

NEW PHARMACEUTICAL PREPARATION FOR PAIN MANAGEMENT

Field of the invention

The present invention relates to a novel pharmaceutical preparation for use as a
5 local anaesthetic for topical administration, to the use of said preparation and to a
process for preparing said preparation.

Background of the invention

10

EMLA[®] cream is the only product on the market giving anaesthesia of intact skin.
EMLA[®] cream is administered to the skin under occlusion for 60 minutes. In
order to obtain faster onset of anaesthesia other local anaesthetic agents and
vehicle systems have been tested (Refs. Freeman, et al., *Pediatr. Anaesthesia*
15 1993:3, 129). For tetracaine, an old well-known topical anaesthetic agent, there
are several patent applications for different formulations, among them a cream and
a patch (Refs. Woolfson and McCafferty, WO 88/09169 and Smith & Nephew, EP
0175609 respectively). WO88/09169 discloses an onset time of approximately 30
minutes for the tetracaine cream and EP 0175609 discloses an onset time of
20 approximately 30-45 minutes for the tetracaine patch. None of these formulations
are so far on the market.

Prior art

25

WO 93/19736 discloses a pharmaceutical composition containing a defined lipid
system of at least two lipid components, at least one being amphiphatic and polar
and one being non polar, and wherein the active agent is lidocaine. The problem
that has been solved according to WO 93/19736 is the considerable difficulties in
30 overcoming the poor bioabsorption of lidocaine.

WO 94/00127 discloses application of lipids and lipid formulations for the treatment of skin and mucous membrane diseases or disorders displaying epidermal hyperproliferation and disruptions of the barrier function.

- 5 EP 455528 discloses cosmetic and dermopharmaceutical compositions containing vesicles of a mixture of phospholipids and glycolipids.

FR 2692781 discloses a cosmetic composition containing sphingomyelin from milk or fish.

10

JP 05163153 discloses a sphingolipid composition, solving the problem with malodour and discoloration, giving a stable composition during storage. The sphingolipid composition can be used as a cosmetic base and in the field of medicines, e.g. as emulsion stabilisers, percutaneous promoting agents etc. The lipids are extracted from the brain.

15

Outline of the invention

- 20 Fig. 1 is a graph showing the solid fat content in the triacylglycerol used, as determined by solid-phase NMR.

We have now surprisingly found that the problem mentioned above, namely to obtain faster onset of anaesthesia, can be solved by the new pharmaceutical preparation according to the present invention. The object of the invention is thus to provide a novel, clinically and pharmaceutically acceptable preparation for dermal pain management.

25

By using the pharmaceutical preparation according to the invention it is possible to achieve pain relief with a faster onset of anaesthesia than what is possible to

30

- achieve with classical topical anaesthetic agents. Another advantage with the pharmaceutical preparation according to the invention is that it is less skin irritating than topical local anaesthetic formulations according to the prior art. Still another advantage is that the hydrolysis of ester compounds such as tetracaine
- 5 seem to be much slower in the lipid vehicle than in conventional formulations.

The pharmaceutical preparation according to the present invention comprises the following ingredients:

- 10 a) one or more local anaesthetic agents
b) one or more polar lipids
c) a triacylglycerol
d) optionally water

- 15 The amount of the local anaesthetic is in the range 1-40 %, preferably 5-10 %. The amount of the polar lipid is in the range 1-40 %, preferably 1-10 %. The amount of the triacylglycerol is in the range 60-95 %, preferably 50-85 %. The amount of water is 0-95 %, preferably 0-20%. All percentages are given as the percentage by weight of the total weight of the pharmaceutical
- 20 preparation.

The local anaesthetic can be selected from tetracaine, lidocaine, prilocaine, mepivacaine, lidocaine/prilocaine, tetracaine/lidocaine, and other local anaesthetics and combinations thereof.

- 25 The polar lipid is preferably a sphingolipid. The sphingolipid can be ceramides, monohexosylceramides, dihexosylceramides, sphingomyelins, lysosphingomyelins, sphingosines or other suitable sphingolipids, or mixtures thereof. The polar lipid can also advantageously be a galactolipid. Preferable galactolipids are
- 30 digalactosyldiacylglycerols and monogalactosylglycerols.

Preferably the sphingolipid is sphingomyelin or products derived from sphingomyelin. The sphingomyelin content is preferably established by chromatographic methods.

- 5 The sphingolipid can be extracted from mammals milk, preferably bovine milk, brain, egg yolk or erythrocytes from animal blood, preferably sheep. The sphingolipid may be synthetic or semisynthetic.

10 The sphingolipid being the polar lipid is related to the composition and structure of human skin lipids, specifically in the epidermis layer. Ceramides, a main lipid component of this layer, is believed to form an extracellular barrier between the cells of the epidermis. The ceramides are further believed to originate from biological processes inside the cells and may be a result of biochemical degradation of sphingomyelins or hexosylceramides. The use of a sphingolipid in
15 the pharmaceutical preparation is therefore preferred from the point of view of biocompatibility of the preparation with the natural barrier of the epidermis.

The polar lipid can also be a galactolipid. The galactolipid can be extracted from almost any kind of glycolipid containing plant material. Preferred plant materials
20 are seeds and kernels from grains and cereals, for instance wheat, rye, oats, corn, rice, millet and sesame. Oat groats as well as wheat gluten have a high lipid concentration and are therefore of advantage to use in the process of preparation.

The triacylglycerol used according to the present invention is preferably selected
25 from palm oil or other natural oils with a similar solid fat content or melting range. When palm oil is selected, commercial palm oil is fractionated to specific mixtures of suitable triacylglycerols, based on the combination of mainly palmitic, oleic and stearic esters of glycerol respectively. It is important for the triacylglycerol mixture utilized in the pharmaceutical preparation to be very pure
30 and free from other glycerides, such as mono- and diacylglycerols. Such purity is

preferably confirmed by established chromatographic methods, for example thin-layer chromatography or high-performance liquid chromatography. It is further important that the triacylglycerol mixture fulfils the bulk quality requirements for use in pharmaceutical preparations. Such requirements are for example oxidation status and content of free fatty acids. The triacylglycerol may also be synthetic or semisynthetic.

The triacylglycerol fractions are defined by the percentage solid fat content, determined by solid-phase NMR as described in IUPAC method no. 2150, 7th edition. Thus, in the temperature range 25-35 °C, the fractions should contain 0-50 % (w/w) solid fat, preferably 0-30 % (w/w). see figure 1.

The pharmaceutical preparation according to the invention is administered topically, and can be administered in form of an ointment, a cream or included in a patch.

20 Pharmaceutical preparations

The pharmaceutical preparations according to the present invention were prepared by melting the triacylglycerol in an open water bath at a temperature range of 40-70 °C. Thereafter the local anaesthetic and the polar lipid were weighed in a vial. The triacylglycerol was melted and transferred to the vial and the mixture was dispersed with an Ystral homogenizer at approximately 1000 rpm and at a temperature range of 40-70 °C for 2-4 minutes. The amount of water required to form a topically applicable formulation was added at room temperature and the formulation was mixed carefully.

30.

The following examples describe in detail the pharmaceutical preparations according to the invention.

5 The sphingolipids used in examples 1-7 and 11 were sphingolipids purified from bovine milk containing 60 - 80 % sphingomyelins. The sphingolipids used in example 10 were sphingolipids purified from egg yolk, containing approximately 98 % sphingomyelin. The galactolipid used in example 9 was purified from oats.

10

Example 1

Tetracaine	5%
Sphingolipids from milk	14.3 %
Palm oil fraction	80.7%

15

Example 2

Tetracaine	25%
Sphingolipids from milk	4%
Palmolein	70%
Water	1%

20

Example 3

Tetracaine	25%
Sphingolipids from milk	3%
Palmolein	52%
Water	20%

25

5	<u>Example 4</u>	
	Tetracaine	25%
	Sphingolipids from milk	4%
	Palmstearin	70%
	Water	1%
10	<u>Example 5</u>	
	Tetracaine	25%
	Sphingolipids from milk	3%
	Palmstearin	52%
	Water	20%
15	<u>Example 6</u>	
	Tetracaine	10%
	Sphingolipids from milk	1%
	Palmstearin	69%
	Water	20%
20	<u>Example 7</u>	
	Tetracaine	10%
	Sphingolipids from milk	10%
	Palmolein	79%
	Water	1%
25	<u>Example 8</u>	
	Tetracaine	5%
	Palm oil fraction	95%

Example 9

5	Tetracaine	25%
	Galactolipids	22%
	Palmstearin	33%
	Water	20%

Example 10

10	Tetracaine	5%
	Sphingolipids from egg yolk	14.3%
	Palm oil fraction	80.7%

Example 11

15	Tetracaine	25%
	Sphingolipids from milk	22%
	Palmstearin	33 %
	Water	20 %

20

Biological studies

Topical anaesthesia/analgesia during occlusion of intact skin in the guinea-pig was studied with lipid formulations of local anaesthetics, as a modification of the method originally described by Edith Bulbring and Isabella Wajda in J Pharmacol Exp Ther 1945: 85: 78-84.

Male guinea-pigs (Dunkin-Hartley strain), weight range of 300-400 g, were used. The hair was removed from the back of the animal with a depilatory (Opilca[®] Hans Schwarzkopf GmbH, Hamburg, Germany). The hairless and smooth skin

was washed with soap and water and the animal was kept in a cage under a desk lamp about two hours before experimentation. On pricking the back of the animal with a cannula (22G) (Kifa (with no point)) or a von Frey filament (4.74) (Semmes-Weinstein pressure aesthesiometer), a twitching of the skin was elicited.

- 5 A circular piece of gauze (one up to eight layers) saturated with test formulation in a thin plastic cup (4.5 cm^2) was applied to the middle of the back. The cup was then covered with Self-adhesive (Fixomull[®] BDF Beiersdorf AG Hamburg, Germany) and the occlusion was finally protected with an elastic bandage. At the end of the application period the treated area was wiped with a tissue and then
- 10 examined for signs of local irritation. The skin which had been in contact with the formulation was pricked with a cannula or a von Frey filament under constant pressure six times at different places and the presence or absence of the twitching response in the skin of treated area was noted. This procedure was repeated at regular intervals of five minutes.

15

The number of pricks not eliciting a response gave an indication of the degree of sensory anaesthesia or analgesia. Groups of two, three or six animals were used for each test formulation.

20

Results

For all formulations discussed below, the percentage of anaesthesia/analgesia was measured after 15 minutes of contact time under occlusion in the guinea-pig.

25

In spite of the short contact time, 15 minutes, a waterfree formulation of 5 % of the active drug, tetracaine in sphingolipids and palm oil gave approximately 80 % of anaesthesia/analgesia with a duration of more than 90 minutes (Example 1).

30

When compared to a similar formulation of 5 % of tetracaine in palm oil without sphingolipids, the percentage of anaesthesia/analgesia was approximately 40 % and the duration was less than 30 minutes (Example 8). From these examples it is obvious that the presence of sphingolipids reduce the onset time and extend the duration of anaesthesia/analgesia.

Further increasing the concentration of tetracaine to 10 % in the presence of sphingolipids, a more saturated triacylglycerol fraction, palmstearin and 20 % of water, resulted in 100 % of anaesthesia/analgesia with a long duration (Example 6).

Further increasing the content of tetracaine to 25 % and sphingolipids to 3 % in palmstearin, also resulted in a high percentage of anaesthesia/ analgesia (Example 4). Addition of 20 % of water to the pharmaceutical preparation according to Example 4, resulted in 100 % of anaesthesia/ analgesia with a long duration. To optimize the effect of the drug, the saturation of the triacylglycerol fraction and the water content of the formulation were important parameters.

Using a more well-defined sphingolipid from egg yolk containing approximately 98 % of sphingomyelins did not alter the effect of the active drug dramatically (Example 10). The sphingolipids used in the other examples were all extracted from bovine milk, containing approximately 60-80 % sphingomyelins. The difference between sphingolipids from bovine milk and egg yolk is obviously not a critical parameter for the effect of the active drug.

A comparison between formulations containing either sphingolipids or galactolipids gave approximately the same percentage of anaesthesia/analgesia at a constant amount of the active drug. The function of the polar lipids is dual, meaning that the polar lipids reduce the onset time and extend the duration of

anaesthesia/analgesia, and are also acting as dispersing agent or stabilizer of the formulation.

5 The best mode of performing the invention is at present considered to be the pharmaceutical preparation according to Example 6.

Conclusion

10 The polar lipids have at least two functions in formulations intended for dermal pain management. They reduce the onset time and extend the duration of anaesthesia, as well as being non-irritating. They are also efficient stabilizers or dispersing agents for pharmaceutical formulations intended for dermal anaesthesia.

CLAIMS


1. A pharmaceutical preparation comprising
 - 5 a) 1-40 % of one or more local anaesthetic agents
 - b) 1-40 % of one or more polar lipids
 - c) 60-95 % of a triacylglycerol
 - d) 0-95 % water
- 10 and wherein the percentages are the percentages by weight of the total weight of the pharmaceutical preparation.
2. A pharmaceutical preparation according to claim 1, wherein the polar lipid is a sphingolipid.
- 15 3. A pharmaceutical preparation according to claim 1, wherein the polar lipid is a galactolipid.
4. A pharmaceutical preparation according to claim 1, wherein the polar lipids are a mixture of sphingolipids and galactolipids.
- 20 5. A pharmaceutical preparation according to claim 2, wherein the sphingolipid is a sphingolipid mixture with at least 1 % sphingomyelin.
- 25 6. A pharmaceutical preparation according to claim 3, wherein the galactolipid is a galactolipid mixture with at least 1 % digalactosyldiacylglycerol.
7. A pharmaceutical preparation according to claim 1, wherein the triacylglycerol is defined by the percentage solid fat content, determined by solid-phase NMR in the temperature range 25-35 °C.
- 30

8. A pharmaceutical preparation according to claim 7, wherein the triacylglycerol is a palm oil fraction.
- 5 9. A pharmaceutical preparation according to claim 7, wherein the triacylglycerol is palmolein.
10. A pharmaceutical preparation according to claim 7, wherein the triacylglycerol is palmstearin.
- 10 11. A pharmaceutical preparation according to claim 1, wherein the amount of the local anaesthetic is 5-10 %, the amount of the polar lipid is 1-10 %, the amount of the triacylglycerol is 50-85 % and the amount of water is 0-20 %.
- 15 12. A pharmaceutical preparation according to claim 1, containing
- | | |
|--------------------------------|------|
| Tetracaine | 10% |
| Sphingolipids from bovine milk | 1% |
| Palmstearin | 69 % |
| 20 Water | 20% |
13. A pharmaceutical preparation according to claim 1 for use in therapy.
14. A pharmaceutical preparation according to claim 1, for use as a local
25 anaesthetic preparation.
15. A pharmaceutical preparation according to claim 1, for topical administration.

16. A pharmaceutical preparation according to claim 1, wherein said preparation is in form of an ointment or a cream.
17. A pharmaceutical preparation according to claim 1, wherein said preparation is incorporated in a patch.
18. Use of a pharmaceutical preparation according to claim 1, for the manufacture of a medicament with local anaesthetic effect.
19. A process for the preparation of a pharmaceutical preparation according to claim 1, wherein the triacylglycerol is melted at a temperature of 40-70°C, one or more local anaesthetic agents and one or more polar lipids are weighed in a vial, the melted triacylglycerol is added and transferred to the vial, the mixture is dispersed with a homogenizer at 40-70°C for 2-4 minutes and the amount of water required to form a topically applicable formulation is added and mixed.
20. A method for the relieving of pain, wherein a pharmaceutical preparation according to claim 1 is administered to a patient in need of pain relief.

1 / 1

Fig. 1



RTMP'S
COPY

970 0253-9



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 31/70, 9/127, 31/19, 31/245	A1	(11) International Publication Number: WO 96/01637 (43) International Publication Date: 25 January 1996 (25.01.96)
(21) International Application Number: PCT/SE95/00760 (22) International Filing Date: 21 June 1995 (21.06.95) (30) Priority Data: 9402453-6 12 July 1994 (12.07.94) SE (71) Applicant (for all designated States except US): ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): BRODIN, Arne [SE/SE]; Kämpevägen 24, S-151 54 Södertälje (SE). CARLSSON, Anders [SE/SE]; St. Göransgatan 80, S-112 38 Stock- holm (SE). HERSLÖF, Bengt [SE/SE]; Brunbärsvägen 2, S-114 21 Stockholm (SE). NICKLASSON, Martin [SE/SE]; Bränningestrandsvägen 72, S-151 39 Södertälje (SE). RY- DHAG, Lisbeth [SE/SE]; P1 1570 Hälleberga, S-150 23 Enhörna (SE). (74) Agent: ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>
(54) Title: NEW PHARMACEUTICAL PREPARATION FOR PAIN MANAGEMENT		
(57) Abstract A new pharmaceutical preparation comprising one or more local anaesthetic agents, a polar lipid, a triacylglycerol and optionally water. The new pharmaceutical preparation is excellent for topical treatment of pain.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Larvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

9700253-9