(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number
WO 01/18175 A1

(51) International Patent Classification[7]: C12N 5/06, 15/12, C07K 14/715

(21) International Application Number: PCT/US00/23978

(22) International Filing Date:
1 September 2000 (01.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/152,311 3 September 1999 (03.09.1999) US

(71) Applicant *(for all designated States except US)*: APPLIED RESEARCH SYSTEMS ARS HOLDING N.V. [NL/NL]; Pietermaai 15, NL-Curacao (AN).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: CUNNINGHAM, Mark [US/US]; Apartment 606, 103 Fox Road, Waltham, MA 02451 (US). IHEY, Tom [US/US]; 39 Pheasant Hill Road, Franklin, MA 02038 (US).

(74) Agents: BROWDY AND NEIMARK, P.L.L.C. et al.; Suite 300, 624 Ninth Street N.W., Washington, DC 20001-5303 (US).


(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:
With international search report.

[Continued on next page]

(54) Title: METHOD FOR PRODUCING A HETEROLOGOUS SECRETED PROTEIN FROM CHINESE HAMSTER OVARY CELLS GROWN ON MICROCARRIERS



(57) Abstract: The present invention relates to a method for producing a heterologous secreted protein from transformed CHO cells grown on microcarriers in a serum-free cell culture medium. This method avoids a production lag phase when switching from a serum-containing to a serum-free environment, a problem observed in the mammalian adherent cell culture production of secreted proteins.

WO 01/18175 A1

— *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.