

# ANTENATAL SCREENING FOR CHROMOSOMAL ABNORMALITIES

This invention relates to a method for antenatal screening for chromosomal abnormalities and to an  
5 assay kit for performing the method and in particular to a method and kit for antenatal screening for Down's Syndrome.

The risk of Down's Syndrome and some other  
10 chromosomal abnormalities in a foetus is known to increase with the age of the mother and it is this knowledge which forms the basis for selection of pregnant women for further investigation. Further investigation for instance in the case of Down's  
15 Syndrome involves sampling of the amniotic fluid by amniocentesis, a procedure which itself carries a risk for the mother of the foetus, induction of a miscarriage being a recognised hazard of this procedure.

20

Maternal serum markers for Down's Syndrome are widely used for antenatal screening for this chromosomal abnormality, the most common of these markers being alpha-fetoprotein (AFP), human chorionic  
25 gonadotropin (hCG) - either the intact molecule thereof or its beta subunit - and unconjugated estriol (uE). Disclosures relating to the use of these markers in antenatal screening for Down's Syndrome include US Patent 4,874,693, WO 89/00696 and WO 90/08325. US  
30 Patent 5,100,806 discloses the use of the beta subunit of hCG as a marker in antenatal screening for Edwards Syndrome.

Maternal serum screening is based on selecting a  
35 subgroup of women who are at the greatest risk of giving birth to a child with an abnormality, in whom

the risks of an invasive diagnostic procedure are considered to be outweighed by the risk of the abnormality. The risk is calculated by multiplying the a priori age related risk by the likelihood ratio. The likelihood ratio is calculated from the relative heights of the multivariate Gaussian distribution functions of marker analytes in affected and unaffected pregnancies, corresponding to the value of the individual serum marker concentrations.

Since the concentrations of the various analytes vary normally with gestational age, the analyte concentrations must be corrected. Correction is performed by dividing the concentration of the analyte by the median concentration expected for that particular gestational age in women with unaffected pregnancies. This is termed the multiple of the median (MoM).

The use of two or more markers together in antenatal screening can be advantageous. For example the markers AFP, hCG and uE can be used together. The combination of marker analytes gives significantly more information than is given by any single marker alone, or by the group of markers when used sequentially. The use of likelihood ratios derived from a multivariate combination is an efficient means of deriving information relating to a woman's risk of carrying an affected child.

Present methods of antenatal screening using maternal serum markers which are in regular use, or are likely to come into regular use in the near future, rely on screening during the second trimester

of pregnancy, that is approximately between 14 and 26 weeks.

However maternal serum screening in the first trimester of pregnancy, that is up to 14 weeks, offers  
5 considerable advantages over screening in the second trimester. In the first trimester the bonding between the mother and the foetus is less strong than in the second trimester and hence there is less psychological impact should the foetus be affected. In addition the  
10 methods of pregnancy interruption are much less traumatic and are safer to the mother in the first trimester than are those available in the second trimester, particularly the late second trimester, that is approximately 18 to 25 or 26 weeks. It is in  
15 the late second trimester that several of the present methods for antenatal screening need to be performed.

Thus it is desirable to develop methods of antenatal screening for chromosomal abnormalities  
20 using two or more maternal serum markers together which can readily be performed in the first trimester of pregnancy.

According to the present invention we provide a  
25 method for antenatal screening for chromosomal abnormalities in which maternal blood from a pregnant woman is measured for levels of free beta hCG and at least a second serum marker and/or precursors and metabolites of these markers and the measured levels  
30 of these markers together with the gestational age of the pregnant woman are compared to reference values at various gestational ages of the levels for free beta hCG and the second serum marker in (a) pregnant women carrying foetuses having the abnormality(ies) subject  
35 to the screen and (b) pregnant women carrying normal

foetuses, the comparison being indicative of the risk of the pregnant woman carrying a foetus with an abnormality subject to the screen characterised in that the second serum marker is pregnancy associated plasma protein A (PAPPA) and the screen is carried out by the end of the thirteenth (13th) completed week of pregnancy.

Further according to the present invention we provide an assay kit for performing a method for antenatal screening for chromosomal abnormalities comprising means for assaying a sample of maternal blood for levels of free beta hCG and at least a second serum marker and/or precursors and metabolites of these markers characterised in that the second serum marker is pregnancy associated plasma protein A (PAPPA) and the blood sample is taken by the end of the thirteenth (13th) completed week of pregnancy.

Further according to the present invention we provide an apparatus comprising means adapted for receiving measurements of a pregnant woman's maternal blood levels of free beta hCG and at least a second serum marker and/or precursors and metabolites of these markers and computer means for comparing the measurements of these levels to sets of reference data to determine fetal chromosomal abnormalities characterised in that the second serum marker is pregnancy associated plasma protein A (PAPPA) and the measurements are made on a blood sample taken by the end of the thirteenth (13th) completed week of pregnancy.

The method of the invention can be used for antenatal screening for a wide range of chromosomal abnormalities. The most significant and frequently

occurring of these is Down's Syndrome (Trisomy 21). Other abnormalities which may be screened for using the invention include Edwards Syndrome (Trisomy 18), Patau's Syndrome (Trisomy 13), Turner Syndrome,

5 Monosomy X and Klinefelter's Syndrome. The method of the invention may be used to screen for individual abnormalities or to screen for groups of abnormalities together, for example it could be used to screen for both Down's Syndrome and Edwards Syndrome.

10

The method of the invention can be used to measure other serum markers in blood samples in addition to hCG and PAPP-A. Such other markers include alpha-fetoprotein (AFP), unconjugated estradiol

15 (uE), inhibin (In), progesterone (Pr), 16-alpha-hydroxy-dehydroepiandrosterone sulphate (16-alpha-hydroxy-DHEAS) and dehydroepiandrosterone sulphate (DHEAS) and precursors and metabolites of these markers. Thus one or more of these other markers may  
20 be measured together with the principal markers of the assay kit of the invention. For example the method of the invention may be used with hCG, PAPP-A and AFP as the markers to be measured.

25 Measurements are carried out using the method of the invention on blood samples taken during the first trimester of pregnancy, i.e. up to the end of the thirteenth completed week of gestation. Preferably the measurements are made on blood samples taken in the  
30 period between the beginning of the eighth week and the end of the thirteenth week of gestation (8th to 13th weeks). The woman's measured serum value for the individual serum marker is divided by the expected median value found in women with unaffected  
35 pregnancies at the same gestational age, to derive the

(MoM). The probability that the (MoM) values for the combination of serum markers tested belongs to the multivariate distribution of values found in unaffected pregnancies is calculated. The same calculation is performed by reference to the probability that the individual combination of values forms part of the multivariate distribution found in abnormal pregnancies. The ratio of the two probabilities is termed the likelihood ratio (LR) which indicates the likelihood that an individual woman has an affected pregnancy or not. The degree of separation between the multivariate distributions for affected and unaffected pregnancies changes with gestational age, i. e. there is a continuous change in the manner of calculating probability depending upon the gestational age. This continuous change can be built into the algorithm used in the calculation.

An individual woman has an a priori age related risk which is independent of the maternal serum marker concentrations. The woman's age related risk, by Baye's theorem, is modified by multiplying by the likelihood ratio (LR) obtained previously to derive a combined risk. This combined risk may then be used to counsel the woman regarding the relative risk of the abnormality as opposed to the risk of miscarriage associated with a subsequent diagnostic invasive procedure.

In the first trimester of pregnancy the concentration of the free beta subunit of hCG is higher in women with a Down's Syndrome fetus and lower in women carrying a fetus with Edwards Syndrome (Trisomy 18). Conversely the levels of PAPPa are lower than normal in women carrying children with either

Down's Syndrome or Trisomy 18 during the first trimester of pregnancy. This makes it very suitable to combine measurements of PAPP A and free beta hCG in to a bivariate distribution for both affected and  
5 unaffected pregnancies in order to calculate the risk of a woman carrying an affected child in the first trimester.

The invention is illustrated by the accompanying drawings wherein:

10

Figure 1 is a graph of Frequency against PAPP A MoM on a log scale which shows the frequency distribution of maternal serum PAPP A expressed as MoM in first trimester pregnancies with Down's Syndrome  
15 (curve 2) and without Down's Syndrome (curve 1).

Figure 2 is a graph of Frequency against free beta hCG MoM on a log scale which shows the frequency distribution of maternal serum free beta hCG expressed  
20 as MoM in first trimester pregnancies both with Down's Syndrome (curve 3) and without Down's Syndrome (curve 4).

Figure 3 contains a series of receiver operating  
25 characteristic curve showing the relationship between detection rate (the proportion of women whose combined risk of abnormality is greater than a given cut-off risk) and the screen positive rate (which is the proportion of all women, both affected and unaffected  
30 who have a combined risk greater than a given cut-off risk). Details of individual curves in Figure 3 are given later in Example 2.

The invention is illustrated by the following  
35 Examples:

## EXAMPLE 1

An example of how the combination of PAPPa and free beta hCG may be used to determine an individual woman's risk is shown below, using as an example a woman of 35 years of age.

In the example a number of sequential calculations are performed as follows:

10

a) The a priori age risk

This is determined from epidemiological survey data relating to a woman's risk of giving birth to a Down's Syndrome child and her age at the date of delivery (Cuckle, H.S., Wald, N.J. and Thompson, S.G., British Journal of Obstetrics and Gynaecology (1987), 94, 387-402).

The risk of giving birth is calculated using the following equation:

20

$$\text{Age-related risk} = \frac{1 - 0.000627 + e^{(-16.2395 + 0.286 \text{ age}^*)}}{0.000627 + e^{(-16.2395 + 0.286 \text{ age}^*)}}$$

25

\* = maternal age at date of delivery

b) Calculation of the multiple of the median value

30

During the first trimester of pregnancy, PAPPa and free beta hCG levels change significantly. To allow for this, all assay results from the mother's serum are first transformed to multiples of the medians to relate the actual result for that analyte to what would be

35

expected for that week of gestation for unaffected pregnancies.

5       The ratio of an individual woman's assay result to the expected value for the gestational age ( estimated to the day ) is termed the multiple of the median (MoM) and it is this value which is required for calculation of her risk.

10       Calculation of the likelihood ratio - the statistical method used to calculate this is trivariate Gaussian frequency distribution analysis. A series of constants are required to calculate the likelihood ratio. These are:

15

- the means and standard deviations of the frequency distributions of PAPPA and free beta hCG ( log transformed to ensure Gaussian distribution ) for both Down's syndrome and unaffected pregnancies.

20

- the correlation coefficients between each pair combination of PAPPA and free beta hCG for Down's Syndrome and unaffected pregnancies.

25       The description of the method used to determine the probability of an unaffected and a Down's Syndrome related pregnancy is described below.

#### Probability Determinations

30

For a single assay, the mathematical calculation of the probability is very simple. The probability that a single result is either in the normal range or the  
35   Down's Syndrome range is given by:

$$f(x) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2}$$

5

where  $\mu$  = transformed population mean for unaffected  
or affected pregnancies

$\sigma$  = standard deviation of the transformed  
populations for unaffected or affected  
10 pregnancies:

$x$  = transformed sample value

When a pair of assays is used, the bivariate  
distribution function set out below is used:

15

$$f(x, y) =$$

$$20 \quad \frac{1}{2\pi \sigma_x \sigma_y \sqrt{1-\rho^2}} e^{-\frac{1}{2} \frac{1}{(1-\rho^2)} \left[ \left( \frac{x-\mu_x}{\sigma_x} \right)^2 + \left( \frac{y-\mu_y}{\sigma_y} \right)^2 - 2\rho \frac{(x-\mu_x)}{\sigma_x} \frac{(y-\mu_y)}{\sigma_y} \right]}$$

where  $\mu_x$  = transformed population mean for unaffected or  
25 affected pregnancies for analyte x

$\mu_y$  = transformed population mean for unaffected or  
affected pregnancies for analyte y

$\sigma_x$  = standard deviation of the transformed population  
for unaffected or affected pregnancies for  
30 analyte x

$\sigma_y$  = standard deviation of the transformed population  
for unaffected or affected pregnancies for  
analyte y

$x$  = transformed sample value, analyte x

35  $y$  = transformed sample value, analyte y

$\rho$  = correlation coefficient between transformed  $x$   
and  $y$  for unaffected or affected pregnancies

#### Calculation of combined risk

---

5

The final step is the calculation of the combined risk. This is given by the following equation:

10 Combined Risk = Age related risk x Likelihood ratio

- The statistical distributions used for the univariate or bivariate Gaussian distributions for unaffected or Down's Syndrome pregnancies are shown in  
15 Table 1. These were derived from a study of 16 first trimester Down's Syndrome samples and 350 unaffected control samples. Log Gaussian univariate and bivariate distributions fitted the data well and the statistical distributions in Table 1 therefore refer to MoM values  
20 after natural log transformation.

25

30

35

Table 1

|    |                                          | Unaffected pregnancies | Down's Syndrome affected pregnancies |
|----|------------------------------------------|------------------------|--------------------------------------|
| 5  |                                          |                        |                                      |
|    | Mean free beta hCG                       | 0.0000                 | 0.8190                               |
| 10 | Standard deviation (sd) of free beta hCG | 0.6325                 | 0.6325                               |
|    | Mean PAPP                                | 0.0000                 | -0.8723                              |
|    | sd of PAPP                               | 0.7222                 | 0.7222                               |
| 15 |                                          |                        |                                      |
|    | Correlation PAPP vs free beta hCG        | 0.3622                 | 0.3622                               |

20 All data are expressed as natural logarithms of the multiple of the median (MoM).

The covariances were not statistically significantly different in unaffected and Down's Syndrome affected pregnancies and were therefore pooled as shown above.

25

The a priori age risk for a 35 year old woman ( following Table 1 ) = 1:384.

30 The risk factors for a 35 year old woman with various levels of free beta hCG and PAPP are shown in Table 2. This is divided into two sections ( respectively 2a and 2b ) showing likelihood ratios and final risks.

35

Table 2

## Section 2a - Likelihood ratios

5

|    |                   | PAPPA MoM   |        |         |         |
|----|-------------------|-------------|--------|---------|---------|
|    | Free beta hCG MoM | Not Assayed | 0.5    | 1.0     | 2.0     |
|    |                   |             |        |         |         |
| 10 | Not Assayed       | -           | 0.651  | 2.074   | 6.611   |
|    | 0.5               | 9.558       | 16.268 | 103.712 | 661.170 |
|    | 1.0               | 2.313       | 1.830  | 11.664  | 74.359  |
| 15 | 2.0               | 0.560       | 0.206  | 1.312   | 8.363   |

## Section 2b - Final risk ( Likelihood ratio x age risk )

20

|    |                   | PAPPA MoM   |        |         |          |
|----|-------------------|-------------|--------|---------|----------|
|    | Free beta HCG MoM | Not Assayed | 0.5    | 1.0     | 2.0      |
|    |                   |             |        |         |          |
| 25 | Not Assayed       | -           | 1:250  | 1:796   | 1:2536   |
|    | 0.5               | 1:3670      | 1:6247 | 1:39825 | 1:253889 |
|    | 1.0               | 1:888       | 1:703  | 1:4479  | 1:28554  |
|    | 2.0               | 1:215       | 1:79   | 1:504   | 1:3211   |

30

The results in Table 2 show that if the result of the first test is known the other result is still informative. For example if a 35 year old woman had a PAPPA level of 1.0 MoM, the a priori age-related risk of 1:384 would be modified to 1:3670 at a free beta hCG

35

level of 0.5 MoM and 1:215 at a free beta hCG level of 2.0 MoM.

## EXAMPLE 2

5

To assess the screening performance realistically in its intended clinical setting it is necessary to simulate the performance when applied to the general screened population.

10

5000 sets of values were constructed from the univariate and bivariate log Gaussian distributions for unaffected pregnancies described in Table 1 using MINITAB<sup>(TM)</sup> ( a statistical package ).

15

5000 sets of data corresponding to the distribution for Down's Syndrome affected pregnancies were similarly constructed.

20

For each sample likelihood ratios were calculated as described in Example 1 univariately for PAPPV and free beta hCG and bivariately using both in combination.

25

The age distribution of pregnancies in England and Wales for the years 1986-1988 were used as the reference maternal age distribution.

The simulation was performed as follows:

30

Various risk cut-offs ( from 1:100 to 1:500 ) were examined. For each risk cut-off and for each maternal age ( from 11 to 55 ) the likelihood ratio required to alter the age risk to below that cut-off was calculated. The proportion of likelihood ratios in the 5000 simulated unaffected samples determines the

35

proportion of unaffected women at that age who would be screen positive. The proportion of likelihood ratios in the 5000 simulated Down's Syndrome affected samples determines the proportion of women at that age who would  
5 screen positive.

Multiplying the proportions by the actual numbers of women at that age gives the false-positive rate ( percentage of unaffected women who screen positive ) and  
10 the detection rate ( percentage of Down's Syndrome affected women who screen positive ) at that maternal age.

Summation of the respective numbers over women of  
15 all ages gives the overall false positive and detection rates in the entire general population.

Examination of different risk cut-off values enables the construction of a receiver operating  
20 characteristic curve ( ROC curve ) which relates the false positive rate to the detection rate.

The ROC curve is shown in Figure 3 which shows the relationship between screen positive rate and  
25 detection rate using as markers free beta hCG alone + age (curve 5), PAPPa alone + age (curve 6) and a combination of these two markers + age (curve 7). From Figure 3 it can be seen that when the combination of markers is used the detection rate is increased  
30 considerably over that using either marker individually.

Table 3 shows the actual detection rates at various screen positive rates for different  
35 combinations of serum markers.

TABLE 3

|    | MARKERS USED                 | SCREEN<br>POSITIVE<br>RATE | DETECTION<br>RATE |
|----|------------------------------|----------------------------|-------------------|
| 5  |                              |                            |                   |
|    | Maternal Age                 | 5.0%                       | 29.8%             |
| 10 |                              |                            |                   |
|    | Free B hCG +<br>Maternal Age | 5.0%                       | 49.6%             |
| 15 |                              |                            |                   |
|    | PAPPA + Maternal<br>Age      | 5.0%                       | 45.8%             |
|    | Free B hCG + PAPPA           | 5.0%                       | 77.3%             |
| 20 | + Maternal Age               |                            |                   |

The results in Table 3 show that where free B hCG and PAPPA are combined as serum markers there are significant improvements in detection rates.

30

35



## CLAIMS

5           1. A method for antenatal screening for  
chromosomal abnormalities in which maternal blood from  
a pregnant woman is measured for levels of free beta  
hCG and at least a second serum marker and/or  
precursors and metabolites of these markers and the  
10 measured levels of these markers together with the  
gestational age of the pregnant woman are compared to  
reference values at various gestational ages of the  
levels for free beta hCG and the second serum marker  
in (a) pregnant women carrying foetuses having the  
15 abnormalitie(s) subject to the screen and (b) pregnant  
women carrying normal foetuses, the comparison being  
indicative of the risk of the pregnant woman carrying  
a foetus with an abnormality subject to the screen  
characterised in that the second serum marker is  
20 pregnancy associated plasma protein A (PAPPA) and the  
screen is carried out by the end of the thirteenth  
(13th) completed week of pregnancy.

          2. A method according to claim 1 characterised  
25 in that a chromosomal abnormality which is screened  
for is Down's Syndrome.

          3. A method according to claim 1 or claim 2  
characterised in that chromosomal abnormalities  
30 screened for include at least one of Edwards Syndrome  
(Trisomy 18), Patau's Syndrome (Trisomy 13), Turner  
Syndrome, Monosomy X and Klinefelter's Syndrome.



4. A method according to any one of the preceding claims characterised in that measurement of the additional marker alpha-fetoprotein is included.
- 5 5. A method according to any one of the preceding claims characterised in that measurement of the additional marker unconjugated estriol is included.
- 10 6. A method according to any one of the preceding claims characterised in that measurement of an additional marker which is at least one of inhibin (In), progesterone (Pr), 16-alpha-hydroxy-dehydroepiandrosterone sulphate (16-alpha-hydroxy-  
15 DHEAS) and dehydroepiandrosterone sulphate (DHEAS) and precursors and metabolites of these markers is included.
- 20 7. A method according to any one of the preceding claims characterised in that measurements are made on a blood sample taken in the period between the beginning of the eighth week and the end of the thirteenth week of gestation.
- 25 8. An assay kit for performing a method for antenatal screening for chromosomal abnormalities comprising means for assaying a sample of maternal blood for levels of free beta hCG and at least a second serum marker and/or precursors and metabolites  
30 of these markers characterised in that the second serum marker is pregnancy associated plasma protein A (PAPPA) and the blood sample is taken by the end of the thirteenth (13th) completed week of pregnancy.



9. An assay kit according to claim 8 characterised in that a chromosomal abnormality to be screened for is Down's Syndrome.


5           10. An assay kit according to claim 8 or claim 9 characterised in that it comprises means for measuring at least one additional marker from the group alpha-fetoprotein (AFP), unconjugated estriol (uE), inhibin (In), progesterone (Pr), 16-alpha-hydroxy-  
10 dehydroepiandrosterone sulphate (16-alpha-hydroxy-DHEAS) and dehydroepiandrosterone sulphate (DHEAS) and precursors and metabolites of these markers.


15           11. An apparatus comprising means adapted for receiving measurements of a pregnant woman's maternal blood levels of free beta hCG and at least a second serum marker and/or precursors and metabolites of these markers and computer means for comparing the measurements of these levels to sets of reference data  
20 to determine fetal chromosomal abnormalities characterised in that the second serum marker is pregnancy associated plasma protein A (PAPPA) and the measurements are made on a blood sample taken by the end of the thirteenth (13th) completed week of  
25 pregnancy.

## ANTENATAL SCREENING FOR CHROMOSOMAL ABNORMALITIES

## Abstract

A method for antenatal screening for chromosomal abnormalities in which maternal blood from a pregnant woman is measured for levels of free beta hCG and at least a second serum marker and/or precursors and metabolites of these markers and the measured levels of these markers together with the gestational age of the pregnant woman are compared to reference values at various gestational ages of the levels for free beta hCG and the second serum marker in (a) pregnant women carrying foetuses having the abnormalitie(s) subject to the screen and (b) pregnant women carrying normal foetuses, the comparison being indicative of the risk of the pregnant woman carrying a foetus with an abnormality subject to the screen characterised in that the second serum marker is pregnancy associated plasma protein A (PAPPA) and the screen is carried out by the end of the thirteenth (13th) completed week of pregnancy.

*Fig.1.*

*Fig.2.*

