

Novel potassium channel inhibitors

Technical field

The present invention relates to novel compounds, pharmaceutical compositions
5 comprising such compounds and their use for treating, alleviating or preventing
diseases or disorders relating to the activity of potassium channels.

Background

10 Ion channels are trans-membrane proteins, which catalyse the transport of inorganic
ions across cell membranes. The ion channels participate in very diverse processes
among which is the generation and timing of action potentials, synaptic transmission,
secretion of hormones, and contraction of muscles.

15 All mammalian cells express potassium (K^+) channels in their cell membranes, and the
channels play a dominant role in the regulation of the membrane potential. In nerve
and muscle cells they influence the form of the action potential, regulate the frequency
and firing patterns of action potentials, the release of neurotransmitters as well as the
degree of bronchodilation and vasodilation. In non-excitabile cells K^+ channels regulate
cellular proliferation and migration as well as the secretion of cytokines.

20 From a molecular and functional point of view, the K^+ channels represent the largest
and most diverse group of ion channels. It can be divided into four broad families:

- voltage-activated K^+ channels (K_v),
- inward rectifier K^+ channels (KIR),
- 25 • two-pore K^+ channels (K2P), and
- calcium-activated K^+ channels (K_{Ca}).

In the K_{Ca} channels, two main groups can be distinguished:

- 30 • the calmodulin-dependent families, consisting of the small conductance (SK's
or $K_{Ca2.x}$) and intermediate conductance channels (IK or $K_{Ca3.1}$), and
- the intracellular ligand gated families, consisting of the classic Ca^{2+} - and
voltage-activated big conductance channel (BK, $K_{Ca1.1}$) as well as channels
sensitive to other intracellular ions ($K_{Ca4.x}$; and $K_{Ca5.1}$).

35

K_{Ca}3.1

K_{Ca}3.1 is a Ca²⁺-activated K⁺ channel encoded by the human gene *KCNN4*. The channel is a tetramer consisting of four identical α -subunits creating the transmembrane K⁺ selective pore at their interfaces, and - at the intracellular side - four calmodulins, which bind incoming Ca²⁺ and open the pore for K⁺ efflux. K_{Ca}3.1 is expressed in many immune cells incl. T- and B-lymphocytes, mast cells, neutrophils, and macrophages, as well as in erythrocytes, fibroblasts, epithelia and endothelia, whereas K_{Ca}3.1 is essentially absent from excitable cells, such as heart, smooth, and striated muscles, and neurons. Furthermore, since K_{Ca}3.1 is essentially absent from excitable cells, pharmacological modulation of this channel is not expected to cause cardiovascular and CNS related adverse effects.

K_{Ca}3.1 in immune cells

The role of K_{Ca}3.1 in immune cells is here described for T-cells but is also valid for other immune cells and for fibroblasts. Activated T-cells (including Th0, Th1 and Th2) require sustained high and strictly controlled intracellular Ca²⁺-concentration to orchestrate activation of enzymes and nuclear transcription factors (eg. the Ca²⁺-dependent calcineurine/NFAT system) for control of the immune response. Cytosolic Ca²⁺ is dynamically regulated via intracellular stores, but long-term Ca²⁺-elevation requires influx from the extracellular space. This causes membrane depolarization, which reduces further influx and quickly terminates the process if not counteracted. This is achieved by K_{Ca}3.1 activation and K⁺ efflux keeping the membrane potential negative. Molecular adaptations occur to consolidate the mechanism long-term: The K_{Ca}3.1 channel is phosphorylated by the H-kinase NDPK-B, which increases its maximal activity, and K_{Ca}3.1 expression is upregulated secondary to NFAT activation. Both processes strengthen the hyperpolarizing capacity of Ca²⁺ mediated K_{Ca}3.1 activation.

Efficient maintenance of high-level cytosolic Ca²⁺ homeostasis is beneficial in controlled immune reactions, while it can be severely pathogenic if becoming an uncontrolled autonomous process.

K_{Ca}3.1 in erythrocytes

Erythrocytes travel between lungs, where O₂ is picked up from alveolar air, and all other tissues, where O₂ is delivered for use in oxidative phosphorylation. The gas

exchange occurs in the smallest blood vessels and the erythrocyte needs to be flexible and adapt size to pass the capillary bed.

5 In this process, $K_{Ca3.1}$ is activated by the Ca^{2+} -influx through Piezo1, which is a Ca^{2+} -permeable channel that is turned-on by the mechanical stress to the membrane during passage. K^+ efflux then drives Cl^- and water efflux resulting in a fast and transient shrinkage allowing a smooth passage. Safe on the other side, where the blood vessels widen out again, both channels close and the salt (K^+ , Cl^- , Ca^{2+}) and water gradients are quickly restored by active transport processes, making the erythrocyte ready for the
10 next passage.

Potassium channel modulators

Consequently, compounds acting as potassium channel modulating agents may be very useful in the treatment, alleviation and/or prevention of diseases like inflammatory
15 bowel diseases (IBD), xerocytosis erythrocytes and acute respiratory distress syndrome (ARDS).

WO 2014/001363 discloses tetrazole derivatives functioning as potassium channel modulators, which are suitable for use in treating diseases and disorders relating to the
20 activity of potassium channels.

WO 2013/191984 discloses fused thiazine-3-ones, which are suitable for the treatment of diseases related to $K_{Ca3.1}$.

25 WO 2014/067861 discloses 3,4-disubstituted oxazolidinone derivatives and their use as inhibitors of calcium activated potassium channel.


Strøbæk *et al.* (2013) discloses the $K(Ca)$ 3.1 channel inhibitor 4-[[3-(Trifluoromethyl)-phenyl]methyl]-2H-1,4-benzothiazin-3(4H)-one (NS6180).
30

$K_{Ca3.1}$ is known to play an essential role in diseases such as IBD, hereditary xerocytosis, and ARDS, and thus $K_{Ca3.1}$ is a promising target for treatment of these diseases. Hence, there is a need for provision of $K_{Ca3.1}$ modulators.

Many known potassium channel modulating agents have poor solubility in water. Thus, there is a further need for potassium channel modulators, such as $K_{Ca3.1}$ modulators, which are more soluble in water.

5 Summary

In one aspect, the current invention relates to a compound of formula (XVI):




Formula (XVI)

wherein

- 10 R^{14} is selected from the group consisting of $-C(O)-C_{1-8}$ alkyl; $-C(O)-O-C_{1-8}$ alkyl; $-C_{2-8}$ alkyl; $-H$ and $-S(O)_2-C_{1-8}$ alkyl;
- R^3 is H , C_{1-5} alkyl, or a bond;
- R^4 is H , C_{1-5} alkyl, or a bond;
- 15 R^5 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by $-O-$;
- R^6 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by $-O-$;
- R^7 is H , a bond, $-OH$, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by $-O-$ and/or substituted with $=O$;
- 20 R^8 is H , a bond, $-OH$, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by $-O-$ and/or substituted with $=O$;
- Anyone of R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 optionally is linked together to form a ring;
- A is a phenyl or a pyridinyl, wherein the phenyl or pyridinyl is optionally substituted with one or more substituents R^{13} individually selected from the
- 25 group consisting of halogen, $-CX_3$, $-OCX_3$, $-CHX_2$, $-OCHX_2$, $-CH_2X$, $-OCH_2X$, $-CH_2CX_3$, OCH_2CX_3 , $-C_{1-8}$ alkyl, $-OC_{1-8}$ alkyl, $-C_{3-7}$ cycloalkyl, $-OC_{3-7}$ cycloalkyl, $-CN$, NO_2 , $-SO_2CH_3$, and $-SF_5$; and
- X is halogen;

or a pharmaceutically acceptable salt thereof.

In one aspect, the current invention relates to a compound of formula (I):



Formula (I)

wherein

R¹ is -OC₁₋₈ alkyl, -C₁₋₈ alkyl, optionally substituted with -OH, or H;

10 R² is a bond, -C(O)-, -S(O)₂-, or -C(H)₂-;

R³ is H, C₁₋₅ alkyl, or a bond;

R⁴ is H, C₁₋₅ alkyl, or a bond;

R⁵ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is replaced by -O-;

15 R⁶ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is replaced by -O-;

R⁷ is H, a bond, -OH, or C₁₋₈ alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;

R⁸ is H, a bond, -OH, or C₁₋₈ alkyl, wherein one or more methylene group optionally and 20 individually is replaced by -O- and/or substituted with =O;

Anyone of R³, R⁴, R⁵, R⁶, R⁷, and R⁸ optionally is linked together to form a ring;

R⁹ is -C(H)- or -N-;

R¹⁰ is H or halogen;

R¹¹ is H or halogen;

25 R¹² is -CX₃, -OCX₃, H or halogen; and

X is halogen;

or a pharmaceutically acceptable salt thereof.


In a further aspect, the present invention relates to a pharmaceutical composition comprising the compound as disclosed herein.

Compounds of the present invention has a high solubility in aqueous medium. Furthermore, compounds of the present invention are active as potassium channel modulators. They are therefore of great interest for the treatment, alleviation and/or prevention of diseases related to potassium channels. Hence, the present invention also relates to the use of a compound as disclosed herein as a medicament. In one aspect, the compound as disclosed herein is used in the treatment of inflammatory bowel disease (IBD). In another aspect, the compound as disclosed herein is used in the treatment of hereditary xerocytosis. In yet another aspect, the compound as disclosed herein is used in the treatment of acute respiratory distress syndrome (ARDS).

Detailed description

Compounds

In one aspect, the present invention relates to a compound of formula (VII):




Formula (VII)

wherein

- R^1 is $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{1-8}$ alkyl, optionally substituted with $-\text{OH}$, or H ;
- R^2 is a bond, $-\text{C}(\text{O})-$, $-\text{S}(\text{O})_2-$, or $-\text{C}(\text{H})_2-$;
- R^3 is H , C_{1-5} alkyl, or a bond;
- R^4 is H , C_{1-5} alkyl, or a bond;

- R⁵ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is replaced by -O-;
- R⁶ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is replaced by -O-;
- 5 R⁷ is H, a bond, -OH, or C₁₋₈ alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;
- R⁸ is H, a bond, -OH, or C₁₋₈ alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;
- Anyone of R³, R⁴, R⁵, R⁶, R⁷, and R⁸ optionally is linked together to form a ring;
- 10 A is a phenyl or a pyridinyl, wherein the phenyl or pyridinyl is optionally substituted with one or more substituents R¹³ individually selected from the group consisting of halogen, -CX₃, -OCX₃, -CHX₂, -OCHX₂, -CH₂X, -OCH₂X, -CH₂CX₃, OCH₂CX₃, -C₁₋₈ alkyl, -OC₁₋₈ alkyl, -C₃₋₇ cycloalkyl, -OC₃₋₇ cycloalkyl, -CN, NO₂, -SO₂CH₃, and -SF₅; and
- 15 X is halogen;
- or a pharmaceutically acceptable salt thereof.

In one embodiment, the compound is of formula (XVI):



Formula (XVI)

- wherein
- R¹⁴ is selected from the group consisting of -C(O)-C₁₋₈ alkyl; -C(O)-O-C₁₋₈ alkyl; -C₂₋₈ alkyl; -H and -S(O)₂-C₁₋₈ alkyl;
- 25 R³ is H, C₁₋₅ alkyl, or a bond;
- R⁴ is H, C₁₋₅ alkyl, or a bond;
- R⁵ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is replaced by -O-;

R^6 is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by -O-;

R^7 is H, a bond, -OH, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;

5 R^8 is H, a bond, -OH, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;

Anyone of R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 optionally is linked together to form a ring;


A is a phenyl or a pyridinyl, wherein the phenyl or pyridinyl is optionally substituted with one or more substituents R^{13} individually selected from the group consisting of halogen, - CX_3 , - OCX_3 , - CHX_2 , - $OCHX_2$, - CH_2X , - OCH_2X , - CH_2CX_3 , OCH_2CX_3 , - C_{1-8} alkyl, - OC_{1-8} alkyl, - C_{3-7} cycloalkyl, - OC_{3-7} cycloalkyl, -
10 CN , NO_2 , - SO_2CH_3 , and - SF_5 ; and

X is halogen;

or a pharmaceutically acceptable salt thereof.

15

In one embodiment, A is a moiety of formula (IX):



Formula (IX)

wherein


20 R^9 is -C(H)-, -N-, or -C(R^{13})-;

R^{13} is individually selected from the group consisting of halogen, - CX_3 , - OCX_3 , - CHX_2 , - $OCHX_2$, - CH_2X , - OCH_2X , - CH_2CX_3 , OCH_2CX_3 , - C_{1-8} alkyl, - OC_{1-8} alkyl, - C_{3-7} cycloalkyl, - OC_{3-7} cycloalkyl, - CN , NO_2 , - SO_2CH_3 , and - SF_5 ;

n is an integer of 0 to 4; and

25 X is halogen.

Thus, in one embodiment, the compound is of formula (VIII):



Formula (VIII)

wherein

R¹⁴ is selected from the group consisting of $-\text{C}(\text{O})-\text{C}_{1-8}$ alkyl; $-\text{C}(\text{O})-\text{O}-\text{C}_{1-8}$ alkyl;
 5 $-\text{C}_{2-8}$ alkyl; $-\text{H}$ and $-\text{S}(\text{O})_2-\text{C}_{1-8}$ alkyl;

R³ is H, C₁₋₅ alkyl, or a bond;

R⁴ is H, C₁₋₅ alkyl, or a bond;

R⁵ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is
 replaced by $-\text{O}-$;

10 R⁶ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is
 replaced by $-\text{O}-$;

R⁷ is H, a bond, $-\text{OH}$, or C₁₋₈ alkyl, wherein one or more methylene group
 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;

15 R⁸ is H, a bond, $-\text{OH}$, or C₁₋₈ alkyl, wherein one or more methylene group
 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;

Any one of R³, R⁴, R⁵, R⁶, R⁷, and R⁸ optionally is linked together to form a ring;
 and

R⁹ is $-\text{C}(\text{H})-$, $-\text{N}-$, or $-\text{C}(\text{R}^{13})-$;

20 R¹³ is individually selected from the group consisting of halogen, $-\text{CX}_3$, $-\text{OCX}_3$, $-\text{CHX}_2$, $-\text{OCHX}_2$, $-\text{CH}_2\text{X}$, $-\text{OCH}_2\text{X}$, $-\text{CH}_2\text{CX}_3$, OCH_2CX_3 , $-\text{C}_{1-8}$ alkyl, $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{3-7}$ cycloalkyl, $-\text{OC}_{3-7}$ cycloalkyl, $-\text{CN}$, NO_2 , $-\text{SO}_2\text{CH}_3$, and $-\text{SF}_5$;


n is an integer of 0 to 4; and

X is halogen;

or a pharmaceutically acceptable salt thereof.

25

In one embodiment, A is a moiety of formula (X):



Formula (X)


wherein

R^9 is $-C(H)-$, $-N-$, or $-C(R^{13})-$;

5 R^{10} , R^{11} , R^{12} , and R^{13} are individually selected from the group consisting of H, halogen, $-CX_3$, $-OCX_3$, $-CHX_2$, $-OCHX_2$, $-CH_2X$, $-OCH_2X$, $-CH_2CX_3$, OCH_2CX_3 , $-C_{1-8}$ alkyl, $-OC_{1-8}$ alkyl, $-C_{3-7}$ cycloalkyl, $-OC_{3-7}$ cycloalkyl, $-CN$, NO_2 , $-SO_2CH_3$, and $-SF_5$; and
 X is halogen.

10

Thus, in one embodiment, the compound is of formula (I):



Formula (I)

15

wherein

R^1 is $-OC_{1-8}$ alkyl, $-C_{1-8}$ alkyl, optionally substituted with $-OH$, or H;

R^2 is a bond, $-C(O)-$, $-S(O)_2-$, or $-C(H)_2-$;

R^3 is H, C_{1-5} alkyl, or a bond;

R^4 is H, C_{1-5} alkyl, or a bond;

20

R^5 is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by $-O-$;

R^6 is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by -O-;

R^7 is H, a bond, -OH, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;

5 R^8 is H, a bond, -OH, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;

Anyone of R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 optionally is linked together to form a ring;

R^9 is -C(H)-, -N-, or -C(R^{13})-;


10 R^{10} , R^{11} , R^{12} , and R^{13} are individually selected from the group consisting of H, halogen, -CX₃, -OCX₃, -CHX₂, -OCHX₂, -CH₂X, -OCH₂X, -CH₂CX₃, OCH₂CX₃, - C_{1-8} alkyl, -OC₁₋₈ alkyl, -C₃₋₇ cycloalkyl, -OC₃₋₇ cycloalkyl, -CN, NO₂, -SO₂CH₃, and -SF₅; and

X is halogen;

or a pharmaceutically acceptable salt thereof.

15

In one embodiment, the present invention relates to a compound of formula (I):



Formula (I)

20

wherein

R^1 is -OC₁₋₈ alkyl, -C₁₋₈ alkyl, optionally substituted with -OH, or H;

R^2 is a bond, -C(O)-, -S(O)₂-, or -C(H)₂-;

R^3 is H, C_{1-5} alkyl, or a bond;

25 R^4 is H, C_{1-5} alkyl, or a bond;

R^5 is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by -O-;

R^6 is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by -O-;

R^7 is H, a bond, -OH, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;

5 R^8 is H, a bond, -OH, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;

Anyone of R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 optionally is linked together to form a ring;

R^9 is -C(H)- or -N-;

R^{10} is H or halogen;

10 R^{11} is H or halogen;


R^{12} is -CX₃, -OCX₃, H or halogen; and

X is halogen;

or a pharmaceutically acceptable salt thereof.

15

In one embodiment, the compound is of formula (XVII):



Formula (XVII)

wherein

20 R^{14} is selected from the group consisting of -C(O)- C_{1-8} alkyl; -C(O)-O- C_{1-8} alkyl; -C₂₋₈ alkyl; -H and -S(O)₂- C_{1-8} alkyl;

R^3 is H, C_{1-5} alkyl, or a bond;

R^4 is H, C_{1-5} alkyl, or a bond;

25 R^5 is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by -O-;

R^6 is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by -O-;

R⁷ is H, a bond, -OH, or C₁₋₈ alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;
R⁸ is H, a bond, -OH, or C₁₋₈ alkyl, wherein one or more methylene group optionally and individually is replaced by -O- and/or substituted with =O;
5 Anyone of R³, R⁴, R⁵, R⁶, R⁷, and R⁸ optionally is linked together to form a ring;
R⁹ is -C(H)- or -N-;
R¹⁰ is H or halogen;
R¹¹ is H or halogen;
R¹² is -CX₃, -OCX₃, H or halogen; and
10 X is halogen;
or a pharmaceutically acceptable salt thereof.

It is well understood that the term "C₁₋₁₀ alkyl" comprises C₁ alkyl, C₂ alkyl, C₃ alkyls, C₄ alkyls, C₅ alkyls, C₆ alkyls, C₇ alkyls, C₈ alkyls, C₉ alkyls, and C₁₀ alkyl. Said alkyl may
15 be linear, branched and/or cyclic. Thus, said alkyl may be partly cyclic. For example, "C₁-C₆-alkyl" designates an alkyl group containing from 1 to 6 carbon atoms that can be linear or branched such as methyl, ethyl, prop-1-yl, prop-2-yl, *iso*-propyl, *tert*-butyl, but-1-yl, but-2-yl, pent-1-yl, pent-2-yl, pent-3-yl, 2-methylbut-1-yl, 3-methylbut-1-yl), hex-1-yl or 2,3-dimethylbut-1-yl.

20 For example, "C₃-C₇-cycloalkyl" designates a saturated monocyclic carbocyclic ring containing from 3 to 7 carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

25 For example, "C₁-C₆-alkoxy" designates a -O-C₁-C₆-alkyl group such as methoxy, ethoxy, 1-propoxy, 2-propoxy, 1-butoxy, 2-butoxy, 2-methyl-2-propoxy, 1-pentoxy, 3-methyl-1-butoxy, 2-pentoxy, 2-methyl-2-butoxy, 1-hexoxy or 3-hexoxy.

30 In one embodiment, R¹ is -OC₁₋₈ alkyl, such as -OC₁₋₇ alkyl, such as -OC₁₋₆ alkyl, such as -OC₁₋₅ alkyl, such as -OC₁₋₄ alkyl, such as -OC₁₋₃ alkyl, such as -OC₁₋₂ alkyl, such as -OC₁ alkyl. Said alkyl may be linear, branched, cyclic or partly cyclic. In one embodiment, R¹ is -OC₁₋₄ alkyl.

35 In one embodiment, R¹ is -C₁₋₈ alkyl, such as -C₁₋₇ alkyl, such as -C₁₋₆ alkyl, such as -C₁₋₅ alkyl, such as -C₁₋₄ alkyl, such as -C₁₋₃ alkyl, such as -C₁₋₂ alkyl, such as -C₁ alkyl.

Said alkyl may be linear, branched, cyclic or partly cyclic. In one embodiment, said alkyl is substituted with -OH. In one embodiment, R¹ is -C₁₋₄ alkyl. In one embodiment, R¹ is a cyclic alkyl, such as cyclopropyl, or cyclobutyl.

5 In one embodiment, R¹ is H.

In one embodiment, R² is a bond. In one embodiment, R² is -C(O)-. In one embodiment, R² is -S(O)₂-. In one embodiment, R² is or -C(H)₂-.

10 In one embodiment, R² is -C(O)- and R¹ is -OC₁₋₄ alkyl. In one embodiment, R² is -C(O)- and R¹ is -OC₁₋₃ alkyl. In one embodiment, R² is -C(O)- and R¹ is -OCH₃. In one embodiment, R² is -C(O)- and R¹ is -OCH₂CH₃. In one embodiment, R² is -C(O)- and R¹ is -OC₃ alkyl, such as -OCH₂(CH₃)₂ or -O-cyclopropyl.

15 In one embodiment, R² is -C(O)- and R¹ is -C₁₋₃ alkyl. In one embodiment, R² is -C(O)- and R¹ is -C₃ alkyl, such as cyclopropyl.

In one embodiment, R² is a bond and R¹ is C₃₋₄ alkyl, corresponding to R² is -C(H)₂- and R¹ is C₂₋₃ alkyl.

20

In one embodiment, R² is -C(H)₂-, and R¹ is -C₃ alkyl, such as cyclopropyl. In one embodiment, R² is or -C(H)₂-, and R¹ is -C₃ alkyl, such as n-propyl, substituted with -OH.

25 In one embodiment, R² is -C(H)₂-, and R¹ is a cyclic alkyl, such as cyclopropyl, or cyclobutyl.

In one embodiment, R² is a bond, and R¹ is a cyclic alkyl, such as cyclopropyl, or cyclobutyl.

30

In one embodiment, R² is -S(O)₂- and R¹ is -C₁₋₃ alkyl. In one embodiment, R² is -S(O)₂- and R¹ is methyl.

35 In one embodiment, -R¹-R² is not -CH₃, such as when R² is a bond, then R¹ is not C₁ alkyl.

In one embodiment, $-R^2-R^1$ is $-R^{14}$.

In one embodiment, R^{14} is $-C(O)-C_{1-8}$ alkyl. In one embodiment, R^{14} is $-C(O)-C_{1-3}$ alkyl.

In one embodiment, R^{14} is $-C(O)-C_3$ alkyl, such as $-C(O)$ -cyclopropyl.

5

In one embodiment, R^{14} is $-C(O)-O-C_{1-8}$ alkyl. In one embodiment, R^{14} is $-C(O)-OC_{1-3}$ alkyl. In one embodiment, R^{14} is $-C(O)-OCH_3$. In one embodiment, R^{14} is $-C(O)-OCH_2CH_3$. In one embodiment, R^{14} is $-C(O)-OC_3$ alkyl, such as $-OCH_2(CH_3)_2$ or $-O$ -cyclopropyl.

10

In one embodiment, R^{14} is $-C_{2-8}$ alkyl, such as C_{3-4} alkyl. In one embodiment, R^{14} is $-C(H)_2-C_{3-7}$ cycloalkyl, such as $-C(H)_2$ -cyclopropyl or $-C(H)_2$ -cyclobutyl. In one embodiment, R^{14} is $-C_{3-7}$ cycloalkyl, such as $-cyclopropyl$ or $-cyclobutyl$. In one embodiment, R^{14} is $-C_{2-8}$ alkyl, such as C_{3-4} alkyl, substituted with one or more $-OH$.

15

In one embodiment, R^{14} is isopropyl substituted with $-OH$.

In one embodiment, R^{14} is $-H$.

In one embodiment, R^{14} is $-S(O)_2-C_{1-8}$ alkyl. In one embodiment, R^{14} is $-S(O)_2-CH_3$.

20

In one embodiment, R^3 is H . In another embodiment, R^3 is a bond. In one embodiment, R^3 is C_{1-5} alkyl, such as C_{1-4} alkyl, such as C_{1-3} alkyl, such as C_{1-2} alkyl, such as C_1 alkyl. Said alkyl may be linear, branched, cyclic or partly cyclic. In one embodiment, R^3 is C_{1-3} alkyl.

25

In one embodiment, R^4 is H . In another embodiment, R^4 is a bond. In one embodiment, R^4 is C_{1-5} alkyl, such as C_{1-4} alkyl, such as C_{1-3} alkyl, such as C_{1-2} alkyl, such as C_1 alkyl. Said alkyl may be linear, branched, cyclic or partly cyclic. In one embodiment, R^4 is C_{1-3} alkyl.

30

In one embodiment, both R^3 and R^4 are H . In another embodiment, only one of R^3 and R^4 are H , whereas the other is a bond or C_{1-5} alkyl.

In one embodiment, R^5 is H . In one embodiment, R^5 is a bond. In one embodiment, R^5 is C_{1-8} alkyl, such as C_{1-7} alkyl, such as C_{1-6} alkyl, such as C_{1-5} alkyl, such as C_{1-4} alkyl,

35


such as C₁₋₃ alkyl, such as C₁₋₂ alkyl, such as C₁ alkyl. In one embodiment, one of the methylene groups in said alkyl is replaced by -O-, thus forming an ether moiety. In one embodiment, R⁵ is C₁₋₄ alkyl.

5 In one embodiment, R⁶ is H. In one embodiment, R⁶ is a bond. In one embodiment, R⁶ is C₁₋₈ alkyl, such as C₁₋₇ alkyl, such as C₁₋₆ alkyl, such as C₁₋₅ alkyl, such as C₁₋₄ alkyl, such as C₁₋₃ alkyl, such as C₁₋₂ alkyl, such as C₁ alkyl. In one embodiment, one of the methylene groups in said alkyl is replaced by -O-, thus forming an ether moiety. In one
10 embodiment, R⁶ is C₁₋₄ alkyl.


10

In one embodiment, R⁵ and R⁶ are H. In one embodiment R⁵ and R⁶ are -CH₃. In one embodiment, R⁵ and R⁶ are linked together to form a ring. Said ring may be a three-membered ring, a four-membered ring, a five-membered ring, a six-membered ring, or a seven-membered ring. In one embodiment, said ring is a three-membered ring.

15 In another embodiment, only one of R⁵ and R⁶ are H, whereas the other is a bond or C₁₋₈ alkyl. In one embodiment, R⁵ and R⁶ are linked together to form a ring as in formula (II), formula (XV) or formula (XXII):




Formula (II)



Formula (XV)

20




Formula (XXII).

In one embodiment, R^7 is H. In one embodiment, R^7 is a bond. In one embodiment, R^7 is -OH. In one embodiment, R^7 is C_{1-8} alkyl, such as C_{1-7} alkyl, such as C_{1-6} alkyl, such as C_{1-5} alkyl, such as C_{1-4} alkyl, such as C_{1-3} alkyl, such as C_{1-2} alkyl, such as C_1 alkyl. In one embodiment, one or more methylene group of said alkyl is replaced by -O-. In one embodiment, said alkyl is substituted with =O, for example R^7 is acetyl. In one embodiment, R^7 is -C(O)-CH₃. In one embodiment, said alkyl is substituted with =O and one or more methylene group of said alkyl is replaced by -O-, for example R^7 is methoxycarbonyl. In one embodiment, R^7 is -C(O)-O-CH₃. In one embodiment, R^7 is C_{1-4} alkyl. In one embodiment, R^7 is methyl.

In one embodiment, R^8 is H. In one embodiment, R^8 is a bond. In one embodiment, R^8 is -OH. In one embodiment, R^8 is C_{1-8} alkyl, such as C_{1-7} alkyl, such as C_{1-6} alkyl, such as C_{1-5} alkyl, such as C_{1-4} alkyl, such as C_{1-3} alkyl, such as C_{1-2} alkyl, such as C_1 alkyl. In one embodiment, one or more methylene group of said alkyl is replaced by -O-. In one embodiment, said alkyl is substituted with =O, for example R^8 is acetyl. Thus, in one embodiment, R^8 is -C(O)-CH₃. In one embodiment, said alkyl is substituted with =O and one or more methylene group of said alkyl is replaced by -O-, for example R^8 is methoxycarbonyl. Thus, in one embodiment, R^8 is -C(O)-O-CH₃. In one embodiment, R^8 is C_{1-4} alkyl. In one embodiment, R^8 is methyl.

In one embodiment, R^5 or R^6 is linked to R^7 or R^8 to form a ring, such as R^5 is linked to R^7 . In one embodiment, when R^5 is linked to R^7 then R^6 and R^8 are H. In one embodiment, when R^5 is linked to R^7 then R^6 is H and R^8 is methyl. In one embodiment, the ring formed by R^5 or R^6 linked to R^7 or R^8 , such as R^5 is linked to R^7 , is a four-membered ring, a five-membered ring, a six-membered ring, a three-

membered ring or a seven-membered ring. In one embodiment, the ring formed when R^5 or R^6 is linked to R^7 or R^8 , such as when R^5 is linked to R^7 , is an azetidine. In one embodiment, the ring formed when R^5 or R^6 is linked to R^7 or R^8 , such as when R^5 is linked to R^7 , is a pyrrolidine. In one embodiment, the ring formed when R^5 or R^6 is linked to R^7 or R^8 , such as when R^5 is linked to R^7 , is a morpholine. In one embodiment, the ring formed when R^5 or R^6 is linked to R^7 or R^8 , such as when R^5 is linked to R^7 , is a piperidine. In one embodiment, R^5 is linked to R^7 as in formula (III), formula (XI) or formula (XVIII):




15


In one embodiment, the compound is the (S)-enantiomer of formula (III).

20


In one embodiment, the ring formed when R^5 or R^6 is linked to R^7 or R^8 , such as when R^5 is linked to R^7 , is a pyrrolidine such as in formula (IV), formula (XII) or formula (XIX):



Formula (IV)



Formula (XII)



Formula (XIX).


5

In one embodiment, the compound is the (S)-enantiomer of formula (IV). In one embodiment, the compound is of formula (IV), and R^9 is $-\text{C}(\text{H})-$ and R^{10} is $-\text{H}$. In one embodiment, the compound is of formula (IV), and R^9 is $-\text{C}(\text{H})-$, R^{10} is $-\text{H}$ and R^8 is $-\text{H}$ or C_{1-3} alkyl, such as methyl.


10

15


In one embodiment, R^3 or R^4 is linked to R^7 or R^8 to form a ring, such as R^3 is linked to R^7 . In one embodiment, when R^3 is linked to R^7 then R^4 and R^8 are H . In one embodiment, the ring formed by R^3 or R^4 linked to R^7 or R^8 , such as R^3 linked to R^7 , is a four-membered ring, a five-membered ring, a six-membered ring, a three-membered ring or a seven-membered ring. In one embodiment, the ring formed when R^3 or R^4 is linked to R^7 or R^8 , such as when R^5 is linked to R^7 , is a four-membered ring. In one embodiment, the ring formed when R^3 or R^4 is linked to R^7 or R^8 , such as when R^3 is linked to R^7 , is an azetidine. In one embodiment, R^3 is linked to R^7 as in formula (V), formula (XIV) or formula (XXI):



Formula (V)



Formula (XIV)




Formula (XXI).


5

In one embodiment, R³ or R⁴ is linked to R⁵ or R⁶ to form a ring, such as R³ is linked to R⁵. In one embodiment, when R³ is linked to R⁵ then R⁴ and R⁶ are H. In one embodiment, the ring formed by R³ or R⁴ linked to R⁵ or R⁶, such as R³ linked to R⁵, is a five-membered ring, a four-membered ring, a six-membered ring, a three-membered ring or a seven-membered ring. In one embodiment, the ring formed when R³ or R⁴ is linked to R⁵ or R⁶, such as when R³ is linked to R⁵, is a five-membered ring. In one embodiment, the ring formed when R³ or R⁴ is linked to R⁵ or R⁶, such as when R³ is linked to R⁵, is a four-membered ring. In one embodiment, the ring formed when R³ or R⁴ is linked to R⁵ or R⁶, such as when R³ is linked to R⁵, is a cyclopentyl. In one embodiment, R³ is linked to R⁵ as in formula (VI), formula (XIII) or formula (XX):


15



Formula (VI)



Formula (XIII)



Formula (XX).

5

In one embodiment R³ and R⁴ are -H, R⁵ and R⁶ are methyl, and R⁷ and R⁸ are -H.

In one embodiment, no more than five of R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are H.

10 In one embodiment, R⁹ is -C(H)-, thus forming a benzene ring. In one embodiment, R⁹ is -N-, thus forming a pyridine ring. In one embodiment, R⁹ is -C(R¹³)-.

In one embodiment, R¹⁰ is H. In another embodiment, R¹⁰ is a halogen, such as Cl.

15 In one embodiment, R¹¹ is H. In another embodiment, R¹¹ is a halogen, such as F.

In one embodiment, R¹² is -CX₃, wherein X is halogen. In one embodiment, said halogen is F, thus R¹² is -CF₃. In one embodiment, R¹² is -OCX₃, wherein X is halogen. In one embodiment, said halogen is F, thus R¹² is -OCF₃. In one embodiment, R¹² is H.

In one embodiment, R^{12} is halogen. In one embodiment, R^{12} is Cl. In one embodiment, R^{12} is Br.

5 In one embodiment R^{11} is F and R^{12} is $-CF_3$. In one embodiment, R^{10} is H, R^{11} is X, and R^{12} is $-CX_3$, wherein X individually is a halogen. In one embodiment, R^{10} is H, R^{11} is F, and R^{12} is $-CF_3$. In one embodiment, R^9 is $-C(H)-$, R^{10} is H, R^{11} is X, and R^{12} is $-CX_3$, wherein X individually is a halogen. In one embodiment, R^9 is $-C(H)-$, R^{10} is H, R^{11} is F, and R^{12} is $-CF_3$. In one embodiment, R^{10} is H, R^{11} is X, and R^{12} is $-OCX_3$, wherein X individually is a halogen. In one embodiment, R^{10} is H, R^{11} is F, and R^{12} is $-OCF_3$. In one embodiment, R^9 is $-C(H)-$, R^{10} is H, R^{11} is X, and R^{12} is $-OCX_3$, wherein X individually is a halogen. In one embodiment, R^9 is $-C(H)-$, R^{10} is H, R^{11} is F, and R^{12} is $-OCF_3$. In one embodiment, R^9 is $-C(H)-$, R^{10} is H, R^{11} is F and R^{12} is $-CF_3$ or $-OCF_3$.

15 In one embodiment, R^9 is $-N-$, R^{10} is H, R^{11} is X, and R^{12} is $-CX_3$, wherein X individually is a halogen. In one embodiment, R^9 is $-N-$, R^{10} is H, R^{11} is F, and R^{12} is $-CF_3$.

In one embodiment, R^{10} is H and R^{11} is H. In one embodiment, R^9 is $-C(H)-$, R^{10} is H and R^{11} is H. In one embodiment, R^{10} is H, R^{11} is H, and R^{12} is halogen, such as R^{10} is H, R^{11} is H, and R^{12} is Br or Cl. In one embodiment, R^9 is $-C(H)-$, R^{10} is H, R^{11} is H, and R^{12} is halogen, such as R^9 is $-C(H)-$, R^{10} is H, R^{11} is H, and R^{12} is Br or Cl.

25 In one embodiment, R^{10} is H, R^{11} is H, and R^{12} is $-CX_3$, wherein X is halogen, such as R^{10} is H, R^{11} is H, and R^{12} is $-CF_3$. In one embodiment, R^9 is $-C(H)-$, R^{10} is H, R^{11} is H, and R^{12} is $-CX_3$, wherein X is halogen, such as R^{10} is H, R^{11} is H, and R^{12} is $-CF_3$.

In one embodiment, R^{11} is H, R^{10} and R^{12} are individually halogen. In one embodiment, R^{11} is H, R^{10} and R^{12} are Cl. In one embodiment, R^9 is $-C(H)-$, R^{11} is H, R^{10} and R^{12} are individually halogen. In one embodiment, R^9 is $-C(H)-$, R^{11} is H, R^{10} and R^{12} are Cl.

30 In one embodiment, no more than two of R^{10} , R^{11} and R^{12} are H. In one embodiment, no more than one of R^{10} , R^{11} and R^{12} are H.

In one embodiment, R^{10} is halogen when R^{11} and R^{12} are H.

In one embodiment, when R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are H, then no more than two of R¹⁰, R¹¹ and R¹² are H.

In one embodiment, n is 0. In one embodiment, n is 1. In one embodiment, n is 2. In one embodiment, n is 3. In one embodiment, n is 4..

In one embodiment, R¹³ is -CX₃, -OCX₃, H or halogen. In one embodiment, R¹³ is H or halogen.

In one embodiment, the compound is not a compound selected from the group consisting of:

- N1-[1-(3-chlorophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine;
- N1-[1-(3-fluorophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine;
- N1-[1-(4-bromophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine;
- N1-[1-(4-fluorophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine;
- N1-methyl-N1-(1-phenylcyclopropyl)-1,2-ethanediamine;
- N-[1-(4-fluorophenyl)cyclopropyl]-N-methyl-3-azetidamine;
- N-[1-(4-fluorophenyl)cyclopropyl]-N-methyl-3-pyrrolidinamine;
- N-[1-(3-fluorophenyl)cyclopropyl]-N-methyl-3-pyrrolidinamine;
- N-[1-(3-fluorophenyl)cyclopropyl]-N-methyl-3-azetidamine;
- N-methyl-N-(1-phenylcyclopropyl)-3-pyrrolidinamine;
- N-methyl-N-(1-phenylcyclopropyl)-3-azetidamine;
- N-methyl-N-(1-phenylcyclopropyl)-3-piperidinamine;
- N-[1-(4-chlorophenyl)cyclopropyl]-N-methyl-3-azetidamine;
- N-[1-(3-chlorophenyl)cyclopropyl]-N-methyl-3-azetidamine; and
- N1-[1-(3-bromophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine.

In one embodiment, the compound is Methyl N-(2-(dimethylamino)ethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate.

In one embodiment, the compound is Methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate.

In one embodiment, the compound is Methyl ((1-aminocyclopropyl)methyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate.

In one embodiment, the compound is Methyl 3-((1-(4-fluoro-3-(trifluoromethyl) phenyl) cyclopropyl) (methoxycarbonyl)amino)azetidine-1-carboxylate.

- In one embodiment, the compound is Methyl azetidin-3-yl(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate.
- In one embodiment, the compound is N-(cyclopropylmethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)azetidin-3-amine.
- 5 In one embodiment, the compound is Methyl (2-aminoethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate.
- In one embodiment, the compound is N1-(cyclopropyl methyl)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-2-methyl propane-1,2-diamine.
- In one embodiment, the compound is N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)cyclopropanecarboxamide.
- 10 In one embodiment, the compound is N1-cyclopropyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine.
- In one embodiment, the compound is Methyl (2-amino-2-methylpropyl)(1-(3,5-dichlorophenyl)cyclopropyl)carbamate.
- 15 In one embodiment, the compound is Methyl (2-amino-2-methylpropyl)(1-(3-bromophenyl)cyclopropyl)carbamate.
- In one embodiment, the compound is Methyl (2-amino-2-methylpropyl)(1-(3-chlorophenyl)cyclopropyl)carbamate.
- In one embodiment, the compound is 1-((azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methyl propan-2-ol.
- 20 In one embodiment, the compound is Methyl (2-amino-2-methylpropyl)(1-(3-(trifluoromethyl)phenyl)cyclopropyl)carbamate.
- In one embodiment, the compound is 1-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)amino)-2-methylpropan-2-ol.
- 25 In one embodiment, the compound is Methyl (2-amino-2-methylpropyl)(1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl) carbamate.
- In one embodiment, the compound is N1-cyclobutyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine.
- In one embodiment, the compound is Ethyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate.
- 30 In one embodiment, the compound is Methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate.
- In one embodiment, the compound is N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methanesulfonamide.

In one embodiment, the compound is Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.

In one embodiment, the compound is Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.

5 In one embodiment, the compound is Methyl (2-acetamido-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate.

In one embodiment, the compound is N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-2-methylpropane-1,2-diamine.

10 In one embodiment, the compound is Methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate.

In one embodiment, the compound is Methyl (S)-(1-(4-fluoro-3-(trifluoro methyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl) carbamate.

In one embodiment, the compound is Methyl (S)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.

15 In one embodiment, the compound is Methyl (R)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.

In one embodiment, the compound is Methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)carbamate.

20 In one embodiment, the compound is Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl)carbamate.

In one embodiment, the compound is Methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate.

In one embodiment, the compound is Methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(2-(hydroxyamino)-2-methylpropyl)carbamate.

25 In one embodiment, the compound is Ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.

In one embodiment, the compound is Ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl) carbamate.

30 In one embodiment, the compound is (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methanesulfonamide.

In one embodiment, the compound is (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-((1-methylpyrrolidin-2-yl)methyl)methanesulfonamide.

- In one embodiment, the compound is Ethyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.
- In one embodiment, the compound is Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate.
- 5 In one embodiment, the compound is Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate.
- In one embodiment, the compound is (R)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methane sulphonamide.
- In one embodiment, the compound is Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate.
- 10 In one embodiment, the compound is Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate.
- In one embodiment, the compound is Isopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.
- 15 In one embodiment, the compound is Cyclopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.
- In one embodiment, the compound is Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methyl pyrrolidin-2-yl)methyl)carbamate.
- In one embodiment, the compound is N-((1-amino cyclopropyl)methyl)-1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine.
- 20 In one embodiment, the compound is Cyclopropylmethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate.
- In one embodiment, the compound is N-((1-aminocyclopropyl)methyl)-N-(1-(4-fluoro-3-(trifluoromethyl)-phenyl)cyclopropyl)methanesulfonamide.
- 25 In one embodiment, the compound is Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methylpyrrolidin-2-yl)methyl)carbamate.
- In one embodiment, the compound is (1S, 2S)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-cyclopentane-1,2-diamine.
- In one embodiment, the compound is (1R,2S)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-cyclopentane-1,2-diamine.
- 30 In one embodiment, the compound is Methyl (S)-(azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)-cyclopropyl)carbamate.

Solubility of compounds

One of the advantages of the compounds of the present invention is that they are more soluble than many other compounds known to modulate potassium channels such as $K_{Ca3.1}$. The compounds tested in Example [54] have a solubility in pH 7.4 phosphate buffer of 400 to 1700 μ M.

Pharmaceutically Acceptable Salts

The chemical compound of the invention may be provided in any form suitable for the intended administration, including pharmaceutically (i.e. physiologically) acceptable salts. Examples of pharmaceutically acceptable addition salts include, without limitation, non-toxic inorganic and organic acid addition salts such as hydrochloride, hydrobromide, nitrate, perchlorate, phosphate, sulphate, formate, acetate, aconate, ascorbate, benzenesulphonate, benzoate, cinnamate, citrate, embonate, enantate, fumarate, glutamate, glycolate, lactate, maleate, malonate, mandelate, methanesulphonate, naphthalene-2-sulphonate, phthalate, salicylate, sorbate, stearate, succinate, tartrate, toluene-p-sulphonate, and the like. Such salts may be formed by procedures well known and described in the art. Other acids such as oxalic acid, which may not be considered pharmaceutically acceptable, may be useful in the preparation of salts useful as intermediates in obtaining a chemical compound of the invention and its pharmaceutically acceptable acid addition salt.

Examples of pharmaceutically acceptable cationic salts of the compound of the invention include, without limitation, the sodium, the potassium, the calcium, the magnesium, the zinc, the aluminium, the lithium, the choline, the lysinium, and the ammonium salt, and the like, of the compound of the invention containing an anionic group. Such cationic salts may be formed by procedures well known and described in the art. In the context of this invention the "onium salts" of *N*-containing compounds are also contemplated as pharmaceutically acceptable salts. Preferred "onium salts" include the alkylonium salts, the cycloalkylonium salts, and the cycloalkylalkylonium salts. In one embodiment, the term "pharmaceutically acceptable salt" of a compound designates any "onium" salts of *N*-containing compounds or any salt of addition of said active principle with a mineral or organic acid among which acetic, hydrochloric, cinnamic, citric, formic, hydrobromic, hydroiodic, hydrofluoric, malonic, methanesulphonic, oxalic, picric, maleic, lactic, nicotinic, phenylacetic, phosphoric, succinic and tartaric acid, ammonium, diethylamine, piperazine, nicotinamide, urea,

sodium, potassium, calcium, magnesium, zinc, lithium, methylamino, dimethylamino, trimethylamino and tris(hydroxymethyl)aminomethane acid.

5 **Preparation of compounds**

Compounds according to the present invention may be prepared according to any conventional methods of chemical synthesis known by the skilled person, e.g. those described in the working examples. The starting materials for the processes described in the present application are known or may readily be prepared by conventional
10 methods known by the skilled artisan from commercially available chemicals.

The end products of the reactions described herein may be isolated by conventional technique such as extraction, crystallisation, distillation, chromatography etc.

15 The compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

20 **Pharmaceutical compositions**

The present invention also relates to a pharmaceutical composition comprising, for example as an active ingredient, a pharmaceutically effective amount of a compound as disclosed herein. In one embodiment, said pharmaceutical composition comprises a therapeutically effective amount of the compound as disclosed herein or a
25 pharmaceutically acceptable salt thereof, together with at least one pharmaceutically acceptable carrier, excipient or diluent.

While a compound as disclosed herein for use in therapy may be administered in the form of the raw chemical compound, it is preferred to introduce the active ingredient,
30 optionally in the form of a pharmaceutically acceptable salt, in a pharmaceutical composition together with one or more adjuvants, excipients, carriers, buffers, diluents, and/or other customary pharmaceutical auxiliaries.

In one embodiment, the invention provides pharmaceutical compositions comprising a
35 compounds disclosed herein or a pharmaceutically acceptable salt thereof, together with one or more pharmaceutically acceptable carriers, and, optionally, other

therapeutic and/or prophylactic ingredients, known and used in the art. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not harmful to the recipient thereof. Pharmaceutical compositions of the invention may be those suitable for oral, rectal, bronchial, nasal, pulmonal, topical
5 (including buccal and sub-lingual), transdermal, vaginal or parenteral (including cutaneous, subcutaneous, intramuscular, intraperitoneal, intravenous, intraarterial, intracerebral, intraocular injection or infusion) administration, or those in a form suitable for administration by inhalation or insufflation, including powders and liquid aerosol administration, or by sustained release systems. Suitable examples of sustained
10 release systems include semipermeable matrices of solid hydrophobic polymers containing the compound of the invention, which matrices may be in form of shaped articles, e.g. films or microcapsules.

A compound as disclosed herein, together with a conventional adjuvant, carrier, or
15 diluent, may thus be placed into the form of pharmaceutical compositions and unit dosages thereof. Such forms include solids, and in particular tablets, filled capsules, powder and pellet forms, and liquids, in particular aqueous or non-aqueous solutions, suspensions, emulsions, elixirs, and capsules filled with the same, all for oral use, suppositories for rectal administration, and sterile injectable solutions for parenteral
20 use. Such pharmaceutical compositions and unit dosage forms thereof may comprise conventional ingredients in conventional proportions, with or without additional active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. A compound as disclosed herein can be administered in a wide
25 variety of oral and parenteral dosage forms. It will be obvious to those skilled in the art that the following dosage forms may comprise, as the active component, either a chemical compound of the invention or a pharmaceutically acceptable salt of a chemical compound of the invention.

30 Unlike many of the other known KCa3.1 inhibitors, the compounds of the present invention has a high solubility in aqueous medium, which makes them suitable for liquid drug administration, such as intravenous or infusion administration.

For preparing pharmaceutical compositions from a compound as disclosed herein,
35 pharmaceutically acceptable carriers can be either solid or liquid. Solid form

preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavouring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

5

The pharmaceutical preparations may be in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packaged tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

10

A therapeutically effective dose refers to that amount of active ingredient, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity, e.g. ED₅₀, may be determined by standard pharmacological procedures in cell cultures or experimental animals. The dose ratio between therapeutic and toxic effects is the therapeutic index and may be expressed by ratio between plasma levels resulting in therapeutic effects and plasma ratios resulting in toxic effects. Pharmaceutical compositions exhibiting large therapeutic indexes are preferred.

15

20

The dose administered must of course be carefully adjusted to the age, weight and condition of the individual being treated, as well as the route of administration, dosage form and regimen, and the result desired, and the exact dosage should of course be determined by the practitioner.

25

The actual dosage depends on the nature and severity of the disease being treated, and is within the discretion of the physician, and may be varied by titration of the dosage to the particular circumstances of this invention to produce the desired therapeutic effect. However, it is presently contemplated that pharmaceutical compositions containing of from about 0.1 to about 10000 mg of active ingredient per individual dose, such as 0.5 to 2000 mg, preferably of from about 1 to about 1000 mg, most preferred of from about 10 to about 500 mg, are suitable for therapeutic treatments. The active ingredient may be administered in one or several doses per day.

30

35

Biological activity

Compounds of the present invention are active as potassium channel modulators. The compounds of the present invention tested in example [55] all inhibit $K_{Ca3.1}$.

5

Method of treatment

Being modulators of potassium ion channels, such as $K_{Ca3.1}$, the compounds of the present invention are of use in the treatment of diseases and disorders of a living body, including human. As used herein, the term "treatment" also includes prevention, and/or
10 alleviation of diseases and disorders. In one aspect, the compound as described herein is for use in medicine. In one aspect, the present invention relates to a method for treatment of IBD, hereditary xerocytosis or ARDS comprising administration of a compound as described herein, or a pharmaceutical composition comprising said compound, to a subject in need thereof.

15

Inflammatory bowel diseases (IBD)

Inflammatory bowel disease (IBD) is a chronic autoimmune disease affecting the gastrointestinal tract with symptoms of abdominal pain, vomiting, diarrhoea, hematochezia, and weight loss. IBD comes in two main forms, ulcerative colitis (UC)
20 and Crohn's disease (CD). UC exclusively affects the colon and rectum, whereas CD may affect the entire gastrointestinal tract. Histologically UC is characterized by extended mucosal inflammation in contrast to CD, where deep punctuate lesions affect all layers of the intestinal wall. It is estimated that approximately 2.5 million patients are diagnosed with IBD (1 million with colitis and 1.5 million with Crohns patients) in the
25 industrialized world (USA, Japan; 5 major EU countries). The incidences are increasing, especially in newly industrialized countries, possibly related to changes in lifestyle.

Currently used anti-IBD drugs are anti-inflammatory (5-ASA's, steroids), generally
30 immune dampening (azathioprine, 6-mercaptopurine), or biological single cytokine/integrine neutralizing agents (eg. infliximab, ustekinumab, vedolizumab). Despite these options and carefully optimized clinical procedures, patients still face rounds of gut-shortening surgeries (many Crohns patients experience at least one surgery in their lifetime), and colitis patients may develop proctitis after colectomy.
35 Suboptimal medical disease control with respect to maintaining long-term remission, to

fight flare ups, and especially avoiding development of irreversible structural changes due to irresolvable gut fibrosis, represents a serious unmet need for IBD patients.

Many of the drugs used to treat IBD today are connected with side effects. For example, side effects of steroids include increased susceptibility to infection; and 5-aminosalicylic acids, such as in the form of *sulphasalazine*, are associated with a significant proportion of non-responders among UC patients, decreased kidney function as well as high and frequent doses, which elicit poor compliance. Drawbacks for TNF-alpha inhibitor infliximab are include high cost, inconvenience of application (injections), waning of efficacy and elicitation of increased risk of infection as a result of their immunosuppressive characteristic; and immunomodulators such as azathioprine, 6-mercaptopurine and methotrexate increase the risk for infections and for some types of cancer, as well as being liver toxic. Thus, there is still a major unmet need for new treatments of inflammatory bowel diseases.

15 *K_{Ca}3.1 as a target for IBD*

T cells play an important role in IBD, and IBD processes (immune cell proliferation, homing, and cytokine release), excessive fibroblast-mediated collagen secretion can lead to fibrosis that causes strictures and intestinal obstructions, and excess water transport across the epithelia can lead to diarrhoea. All these pathological processes can be dampened by K_{Ca}3.1 inhibition.

As demonstrated herein, the compounds of the present invention inhibit K_{Ca}3.1, and thus, in one aspect, the present invention relates to a compound as described herein for use in the treatment, alleviation and/or prevention of inflammatory bowel disease (IBD). In one embodiment, said IBD is colitis, such as ulcerative colitis (UC). In one embodiment, said IBD is Crohn's disease (CD).

Hereditary xerocytosis

Hereditary xerocytosis, also known as dehydrated stomatocytosis, is characterized by increased fragility and haemolysis of erythrocytes, resulting in a fully compensated or mild to severe anaemia. Increased reticulocyte formation (to compensate for erythrocyte loss), ion-overload and jaundice (resulting from the increased break-down of haemoglobin) are characteristic in adults. New-borns may suffer from transient edema/ascites, which in rare cases may develop to life-threatening hydrops fetalis. The disease is very heterogeneous but is classically identified from a combination of

clinical signs, such as fatigue, enlarged spleen, gall stones, thrombosis events, and pulmonary hypertension. Microscopic examination may reveal erythrocytes with abnormal shapes and analysis of haematology parameters reveal shrunken erythrocytes due to salt and water loss. The ethology of hereditary xerocytosis has long been known to differ radically from other hereditary anaemias, such as the haemoglobinopathies (eg. sickle cell anemia and the thalassemia diseases), or glycolytic enzymopathies (eg. glucose-6-phosphate deficiency), in that it is due to a primary membrane permeability defect. The molecular targets involved in this defect have just recently been identified.

K_{Ca}3.1 as a target for hereditary xerocytosis

Recent years of scientific investigations have shown that hereditary xerocytosis is due to gain-of-function mutations in either *Piezo1* or *KCNN4*, the gene encoding K_{Ca}3.1. Both mutations essentially result in the same phenotype: In the case of *Piezo1* mutations Ca²⁺ enters the erythrocyte through the constantly open channel, thus activating K_{Ca}3.1 resulting in permanently dehydrated erythrocytes; in the case of *KCNN4* mutations K_{Ca}3.1 are constitutively open thereby governing erythrocyte dehydration even in the absence of a Ca²⁺-signal from the Piezo1 channel. The clear definition of the genes and mutations responsible for hereditary xerocytosis, allows easy diagnostics of which patients will benefit from the treatment and which should not be treated.

Inhibition of the erythrocyte K_{Ca}3.1 channel will counteract unintentional dehydration and presumably prevent haemolysis of xerocytosis erythrocytes and thereby improve the clinical condition of patients. Importantly, the binding site for K_{Ca}3.1 inhibitors do not overlap with the known gain-of-function mutations in K_{Ca}3.1. This pinpoints K_{Ca}3.1 as a pivotal target for all known causes of hereditary xerocytosis.

The compounds of the present invention inhibit K_{Ca}3.1. Hence, in one aspect, the present invention relates to a compound described herein for use in the treatment, alleviation and/or prevention of hereditary xerocytosis. Hereditary xerocytosis is one of the most frequent variant of hereditary stomatocytoses, a group of rare disorders characterized by a leak of monovalent cations such as K⁺ from the red blood cells (RBCs).

Acute respiratory distress syndrome (ARDS)

Acute respiratory distress syndrome is a serious and often lethal complication to lung infections, as caused for example by SARS, MERS, or Covid-19 virus. The infections can lead to global lung inflammation, which widens the ultrathin barriers between the air-filled alveoli and the blood-filled alveolar vessels and fills-up the alveoli with liquid, and thereby hampers the life-essential oxygen/carbondioxide gas exchange between lung and blood. ARDS is thus a complex condition that involve both components of the immune system as well as the air/blood barrier function. Since there are currently no medical treatments that specifically interfere with ARDS (general immune dampening treatments by steroids are not effective), the only options for patients is medical ventilator treatment at an intensive care unit.


KCa3.1 as a target for ARDS.

Since the KCa3.1 channel is expressed in both the epithelia and endothelia as well as in the inflammatory cells, such as neutrophils, that participate in lung inflammation, inhibition of KCa3.1 can dampen both the basic inflammation and possibly also protect the air/blood barrier function. Experiments with a mouse model of ARDS have recently shown that KCa3.1 knock-out mice have improved gas exchange, and the improvement was also demonstrated by treatment with the classical KCa3.1 inhibitors senicapoc and TRAM-34. In the clinical situation with patients on medical ventilation, oral drug administration is not optimal, whereas intravenous bolus or infusion administration is preferred. Classical KCa3.1 inhibitors like the triarylmethanes (exemplified by senicapoc and TRAM-34) have extremely low water solubility, which makes IV-formulations very challenging. The same drawbacks apply to known KCa3.1 inhibitors based on other chemical scaffolds.

The compounds of the present invention inhibit $K_{Ca3.1}$. Further, the compounds of the present invention has a high solubility in aqueous medium. Hence, the compounds of the present invention are highly suitable for use in treatment of ARDS. Thus, in one aspect, the present invention relates to a compound as described herein for use in the treatment, alleviation and/or prevention of ARDS.


Items

1. A compound of formula (VII):



Formula (VII)

- 5 wherein
- R^1 is $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{1-8}$ alkyl, optionally substituted with $-\text{OH}$, or H ;
- R^2 is a bond, $-\text{C}(\text{O})-$, $-\text{S}(\text{O})_2-$, or $-\text{C}(\text{H})_2-$;
- R^3 is H , C_{1-5} alkyl, or a bond;
- R^4 is H , C_{1-5} alkyl, or a bond;
- 10 R^5 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by $-\text{O}-$;
- R^6 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by $-\text{O}-$;
- R^7 is H , a bond, $-\text{OH}$, or C_{1-8} alkyl, wherein one or more methylene group
- 15 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;
- R^8 is H , a bond, $-\text{OH}$, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;
- Anyone of R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 optionally is linked together to form a ring;
- A is a phenyl or a pyridinyl, wherein the phenyl or pyridinyl is optionally
- 20 substituted with one or more substituents R^{13} individually selected from the group consisting of halogen, $-\text{CX}_3$, $-\text{OCX}_3$, $-\text{CHX}_2$, $-\text{OCHX}_2$, $-\text{CH}_2\text{X}$, $-\text{OCH}_2\text{X}$, $-\text{CH}_2\text{CX}_3$, OCH_2CX_3 , $-\text{C}_{1-8}$ alkyl, $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{3-7}$ cycloalkyl, $-\text{OC}_{3-7}$ cycloalkyl, $-\text{CN}$, NO_2 , $-\text{SO}_2\text{CH}_3$, and $-\text{SF}_5$; and
- X is halogen;
- 25 or a pharmaceutically acceptable salt thereof.
2. The compound according to item 1, wherein the compound is of formula (XVI):



Formula (XVI)

wherein

R¹⁴ is selected from the group consisting of $-\text{C}(\text{O})-\text{C}_{1-8}$ alkyl; $-\text{C}(\text{O})-\text{O}-\text{C}_{1-8}$ alkyl;
 5 $-\text{C}_{2-8}$ alkyl; $-\text{H}$ and $-\text{S}(\text{O})_2-\text{C}_{1-8}$ alkyl;

R³ is H, C₁₋₅ alkyl, or a bond;

R⁴ is H, C₁₋₅ alkyl, or a bond;

R⁵ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is
 replaced by $-\text{O}-$;

10 R⁶ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is
 replaced by $-\text{O}-$;

R⁷ is H, a bond, $-\text{OH}$, or C₁₋₈ alkyl, wherein one or more methylene group
 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;

R⁸ is H, a bond, $-\text{OH}$, or C₁₋₈ alkyl, wherein one or more methylene group


15 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;
 Anyone of R³, R⁴, R⁵, R⁶, R⁷, and R⁸ optionally is linked together to form a ring;

A is a phenyl or a pyridinyl, wherein the phenyl or pyridinyl is optionally
 substituted with one or more substituents R¹³ individually selected from the
 group consisting of halogen, $-\text{CX}_3$, $-\text{OCX}_3$, $-\text{CHX}_2$, $-\text{OCHX}_2$, $-\text{CH}_2\text{X}$, $-\text{OCH}_2\text{X}$, $-\text{CH}_2\text{CX}_3$, OCH_2CX_3 , $-\text{C}_{1-8}$ alkyl, $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{3-7}$ cycloalkyl, $-\text{OC}_{3-7}$ cycloalkyl, $-\text{CN}$, NO_2 , $-\text{SO}_2\text{CH}_3$, and $-\text{SF}_5$; and

20 X is halogen;

or a pharmaceutically acceptable salt thereof.

3. The compound according to any one of the preceding items, wherein the
 25 compound is of formula (VIII):



Formula (VIII)

wherein

R¹⁴ is selected from the group consisting of $-\text{C}(\text{O})-\text{C}_{1-8}$ alkyl; $-\text{C}(\text{O})-\text{O}-\text{C}_{1-8}$ alkyl;
 5 $-\text{C}_{2-8}$ alkyl; $-\text{H}$ and $-\text{S}(\text{O})_2-\text{C}_{1-8}$ alkyl;

R³ is H, C_{1-5} alkyl, or a bond;

R⁴ is H, C_{1-5} alkyl, or a bond;

R⁵ is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is
 replaced by $-\text{O}-$;

10 R⁶ is H, a bond, or C_{1-8} alkyl, wherein one methylene group optionally is
 replaced by $-\text{O}-$;

R⁷ is H, a bond, $-\text{OH}$, or C_{1-8} alkyl, wherein one or more methylene group
 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;

15 R⁸ is H, a bond, $-\text{OH}$, or C_{1-8} alkyl, wherein one or more methylene group
 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;

Any one of R³, R⁴, R⁵, R⁶, R⁷, and R⁸ optionally is linked together to form a ring;
 and

R⁹ is $-\text{C}(\text{H})-$, $-\text{N}-$, or $-\text{C}(\text{R}^{13})-$;


20 R¹³ is individually selected from the group consisting of halogen, $-\text{CX}_3$, $-\text{OCX}_3$, $-\text{CHX}_2$, $-\text{OCHX}_2$, $-\text{CH}_2\text{X}$, $-\text{OCH}_2\text{X}$, $-\text{CH}_2\text{CX}_3$, OCH_2CX_3 , $-\text{C}_{1-8}$ alkyl, $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{3-7}$ cycloalkyl, $-\text{OC}_{3-7}$ cycloalkyl, $-\text{CN}$, NO_2 , $-\text{SO}_2\text{CH}_3$, and $-\text{SF}_5$;

n is an integer of 0 to 4; and

X is halogen;

or a pharmaceutically acceptable salt thereof.


25 4. The compound according to item 1, wherein the compound is of formula (I):



Formula (I)

wherein


- 5 R^1 is $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{1-8}$ alkyl, optionally substituted with $-\text{OH}$, or H ;
 R^2 is a bond, $-\text{C}(\text{O})-$, $-\text{S}(\text{O})_2-$, or $-\text{C}(\text{H})_2-$;
 R^3 is H , C_{1-5} alkyl, or a bond;
 R^4 is H , C_{1-5} alkyl, or a bond;
 R^5 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is
10 replaced by $-\text{O}-$;
 R^6 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is
replaced by $-\text{O}-$;
 R^7 is H , a bond, $-\text{OH}$, or C_{1-8} alkyl, wherein one or more methylene group
optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;
15 R^8 is H , a bond, $-\text{OH}$, or C_{1-8} alkyl, wherein one or more methylene group
optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;
Anyone of R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 optionally is linked together to form a ring;
 R^9 is $-\text{C}(\text{H})-$, $-\text{N}-$, or $-\text{C}(\text{R}^{13})-$;
 R^{10} , R^{11} , R^{12} , and R^{13} are individually selected from the group consisting of H ,
20 halogen, $-\text{CX}_3$, $-\text{OCX}_3$, $-\text{CHX}_2$, $-\text{OCHX}_2$, $-\text{CH}_2\text{X}$, $-\text{OCH}_2\text{X}$, $-\text{CH}_2\text{CX}_3$, OCH_2CX_3 , $-\text{C}_{1-8}$ alkyl, $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{3-7}$ cycloalkyl, $-\text{OC}_{3-7}$ cycloalkyl, $-\text{CN}$, NO_2 , $-\text{SO}_2\text{CH}_3$,
and $-\text{SF}_5$; and
 X is halogen;
or a pharmaceutically acceptable salt thereof.
- 25 5. The compound according to item 1, wherein the compound is of formula (I):



Formula (I)

wherein

- 5 R^1 is $-\text{OC}_{1-8}$ alkyl, $-\text{C}_{1-8}$ alkyl, optionally substituted with $-\text{OH}$, or H ;
 R^2 is a bond, $-\text{C}(\text{O})-$, $-\text{S}(\text{O})_2-$, or $-\text{C}(\text{H})_2-$;
 R^3 is H , C_{1-5} alkyl, or a bond;
 R^4 is H , C_{1-5} alkyl, or a bond;
 R^5 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is
10 replaced by $-\text{O}-$;
 R^6 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is
replaced by $-\text{O}-$;
 R^7 is H , a bond, $-\text{OH}$, or C_{1-8} alkyl, wherein one or more methylene group
optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;
15 R^8 is H , a bond, $-\text{OH}$, or C_{1-8} alkyl, wherein one or more methylene group
optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;
Anyone of R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 optionally is linked together to form a ring;
 R^9 is $-\text{C}(\text{H})-$ or $-\text{N}-$;
 R^{10} is H or halogen;
20 R^{11} is H or halogen;
 R^{12} is $-\text{CX}_3$, $-\text{OCX}_3$, H or halogen; and
 X is halogen;
or a pharmaceutically acceptable salt thereof.
6. The compound according to item 1, wherein the compound is of formula (XVII):



Formula (XVII)

wherein

R¹⁴ is selected from the group consisting of $-\text{C}(\text{O})-\text{C}_{1-8}$ alkyl; $-\text{C}(\text{O})-\text{O}-\text{C}_{1-8}$ alkyl;
 5 $-\text{C}_{2-8}$ alkyl; $-\text{H}$ and $-\text{S}(\text{O})_2-\text{C}_{1-8}$ alkyl;

R³ is H, C₁₋₅ alkyl, or a bond;

R⁴ is H, C₁₋₅ alkyl, or a bond;

R⁵ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is
 replaced by $-\text{O}-$;

10 R⁶ is H, a bond, or C₁₋₈ alkyl, wherein one methylene group optionally is
 replaced by $-\text{O}-$;

R⁷ is H, a bond, $-\text{OH}$, or C₁₋₈ alkyl, wherein one or more methylene group
 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;

R⁸ is H, a bond, $-\text{OH}$, or C₁₋₈ alkyl, wherein one or more methylene group

15 optionally and individually is replaced by $-\text{O}-$ and/or substituted with $=\text{O}$;
 Anyone of R³, R⁴, R⁵, R⁶, R⁷, and R⁸ optionally is linked together to form a ring;

R⁹ is $-\text{C}(\text{H})-$ or $-\text{N}-$;

R¹⁰ is H or halogen;

R¹¹ is H or halogen;


20 R¹² is $-\text{CX}_3$, $-\text{OCX}_3$, H or halogen; and

X is halogen;

or a pharmaceutically acceptable salt thereof.


7. The compound according to any one of the preceding items, wherein R³ and R⁴
 are $-\text{H}$, R⁵ and R⁶ are methyl, and R⁷ and R⁸ are $-\text{H}$.

25 8. The compound according to any one of the preceding items, wherein the
 compound is of formula (XI):




Formula (XI).

9. The compound according to any one of the preceding items, wherein the compound is of formula (XII):




Formula (XII).

10. The compound according to any one of the preceding items, wherein the compound is of formula (XIII):




Formula (XIII).

11. The compound according to any one of the preceding items, wherein the compound is of formula (XIV):




Formula (XIV).

12. The compound according to any one of the preceding items, wherein the compound is of formula (XV):




Formula (XV).

13. The compound according to any one of the preceding items, wherein the compound is of formula (III):




Formula (III).

14. The compound according to any one of the preceding items, wherein the compound is of formula (IV):




Formula (IV).

- 5 15. The compound according to any one of the preceding items, wherein the compound is of formula (VI):



Formula (VI).


- 10 16. The compound according to any one of the preceding items, wherein the compound is of formula (V):



Formula (V).


17. The compound according to any one of the preceding items, wherein the compound is of formula (II):

5



Formula (II).


18. The compound according to any one of the preceding items, wherein the compound is of formula (XVIII):



Formula (XVIII).


19. The compound according to any one of the preceding items, wherein the compound is of formula (XIX):

10




Formula (XIX).

20. The compound according to any one of the preceding items, wherein the compound is of formula (XX):




Formula (XX).

21. The compound according to any one of the preceding items, wherein the compound is of formula (XXI):




Formula (XXI).

22. The compound according to any one of the preceding items, wherein the compound is of formula (XXII):



Formula (XXII).

23. The compound according to any one of the preceding items, wherein A is a moiety of formula (IX):



Formula (IX)

wherein


R^9 is $-C(H)-$, $-N-$, or $-C(R^{13})-$;

R^{13} is individually selected from the group consisting of halogen, $-CX_3$, $-OCX_3$, $-CHX_2$, $-OCHX_2$, $-CH_2X$, $-OCH_2X$, $-CH_2CX_3$, OCH_2CX_3 , $-C_{1-8}$ alkyl, $-OC_{1-8}$ alkyl, $-C_{3-7}$ cycloalkyl, $-OC_{3-7}$ cycloalkyl, $-CN$, NO_2 , $-SO_2CH_3$, and $-SF_5$;

n is an integer of 0 to 4; and

X is halogen.

24. The compound according to any one of the preceding items, wherein A is a moiety of formula (X):



Formula (X)

wherein

R^9 is $-C(H)-$, $-N-$, or $-C(R^{13})-$;

R^{10} , R^{11} , R^{12} , and R^{13} are individually selected from the group consisting of H, halogen, $-CX_3$, $-OCX_3$, $-CHX_2$, $-OCHX_2$, $-CH_2X$, $-OCH_2X$, $-CH_2CX_3$, OCH_2CX_3 , $-C_{1-8}$ alkyl, $-OC_{1-8}$ alkyl, $-C_{3-7}$ cycloalkyl, $-OC_{3-7}$ cycloalkyl, $-CN$, NO_2 , $-SO_2CH_3$, and $-SF_5$; and

5 X is halogen.

25. The compound according to any one of the preceding items, wherein

R^9 is $-C(H)-$ or $-N-$;

R^{10} is H or halogen;

R^{11} is H or halogen;

10 R^{12} is $-CX_3$, $-OCX_3$, H or halogen; and

X is halogen.

26. The compound according to any one of the preceding items, wherein $-R^2-R^1$ is $-R^{14}$, and R^{14} is selected from the group consisting of $-C(O)-C_{1-8}$ alkyl; $-C(O)-O-C_{1-8}$ alkyl; $-C_{2-8}$ alkyl; $-H$ and $-S(O)_2-C_{1-8}$ alkyl.

15 27. The compound according to any one of the preceding items, wherein the compound is the (S)-enantiomer.

28. The compound according to any one of the preceding items, wherein R^1 is $-OC_{1-8}$ alkyl, such as $-OC_{1-7}$ alkyl, such as $-OC_{1-6}$ alkyl, such as $-OC_{1-5}$ alkyl, such as $-OC_{1-4}$ alkyl, such as $-OC_{1-3}$ alkyl, such as $-OC_{1-2}$ alkyl, such as $-OC_1$ alkyl.

20 29. The compound according to any one of the preceding items, wherein R^1 is $-C_{1-8}$ alkyl, such as $-C_{1-7}$ alkyl, such as $-C_{1-6}$ alkyl, such as $-C_{1-5}$ alkyl, such as $-C_{1-4}$ alkyl, such as $-C_{1-3}$ alkyl, such as $-C_{1-2}$ alkyl, such as $-C_1$ alkyl.

30. The compound according to any one of the preceding items, wherein R^1 is $-C_{1-8}$ alkyl substituted with $-OH$.

25 31. The compound according to any one of the preceding items, wherein R^1 is $-H$.

32. The compound according to any one of the preceding items, wherein R^2 is a bond.

33. The compound according to any one of the preceding items, wherein R^2 is $-C(O)-$.

30 34. The compound according to any one of the preceding items, wherein R^2 is $-C(H)_2-$.

35. The compound according to any one of the preceding items, wherein R^2 is $-S(O)_2-$.

35 36. The compound according to any one of the preceding items, wherein R^2 is $-C(O)-$ and R^1 is $-OC_{1-4}$ alkyl.

37. The compound according to any one of the preceding items, wherein R^2 is -C(O)- and R^1 is -OC₁₋₃ alkyl.
38. The compound according to any one of the preceding items, wherein R^2 is a bond and R^1 is C₃₋₄ alkyl.
- 5 39. The compound according to any one of the preceding items, wherein R^{14} is -C(O)-C₁₋₈ alkyl, such as R^{14} is -C(O)-C₁₋₃ alkyl, such as R^{14} is -C(O)-C₃ alkyl, such as -C(O)-cyclopropyl.
40. The compound according to any one of the preceding items, wherein R^{14} is -C(O)-O-C₁₋₈ alkyl, such as, R^{14} is -C(O)-OC₁₋₃ alkyl, such as R^{14} is selected from
10 the group consisting of -C(O)-OCH₃, -C(O)-OCH₂CH₃, -OCH₂(CH₃)₂ and -O-cyclopropyl.
41. The compound according to any one of the preceding items, wherein R^{14} is -C₂₋₈ alkyl, such as C₃₋₄ alkyl.
42. The compound according to any one of the preceding items, wherein R^{14} is -C(H)₂-C₃₋₇ cycloalkyl, such as -C(H)₂-cyclopropyl or -C(H)₂-cyclobutyl. In one
15 43. The compound according to any one of the preceding items, wherein R^{14} is -C₃₋₇ cycloalkyl, such as -cyclopropyl or -cyclobutyl.
44. The compound according to any one of the preceding items, wherein R^{14} is -C₂₋₈ alkyl, such as C₃₋₄ alkyl, substituted with one or more -OH, such as R^{14} is
20 isopropyl substituted with -OH.
45. The compound according to any one of the preceding items, wherein R^{14} is -H.
46. The compound according to any one of the preceding items, wherein R^{14} is -S(O)₂-C₁₋₈ alkyl, such as R^{14} is -S(O)₂-CH₃.
47. The compound according to any one of the preceding items, wherein R^{14} is not
25 -CH₃.
48. The compound according to any one of the preceding items, wherein R^3 is C₁₋₅ alkyl, such as C₁₋₄ alkyl, such as C₁₋₃ alkyl, such as C₁₋₂ alkyl, such as C₁ alkyl.
49. The compound according to any one of the preceding items, wherein R^4 is C₁₋₅ alkyl, such as C₁₋₄ alkyl, such as C₁₋₃ alkyl, such as C₁₋₂ alkyl, such as C₁ alkyl.
- 30 50. The compound according to any one of the preceding items, wherein R^3 and R^4 are H.
51. The compound according to any one of the preceding items, wherein R^5 is C₁₋₈ alkyl, such as C₁₋₇ alkyl, such as C₁₋₆ alkyl, such as C₁₋₅ alkyl, such as C₁₋₄ alkyl, such as C₁₋₃ alkyl, such as C₁₋₂ alkyl, such as C₁ alkyl.

52. The compound according to any one of the preceding items, wherein R^6 is C_{1-8} alkyl, such as C_{1-7} alkyl, such as C_{1-6} alkyl, such as C_{1-5} alkyl, such as C_{1-4} alkyl, such as C_{1-3} alkyl, such as C_{1-2} alkyl, such as C_1 alkyl.
53. The compound according to any one of the preceding items, wherein R^5 and R^6 are H.
54. The compound according to any one of the preceding items, wherein R^5 and R^6 are $-CH_3$.
55. The compound according to any one of the preceding items, wherein R^5 and R^6 are linked together to form a ring.
56. The compound according to any one of the preceding items, wherein R^7 is C_{1-8} alkyl, such as C_{1-7} alkyl, such as C_{1-6} alkyl, such as C_{1-5} alkyl, such as C_{1-4} alkyl, such as C_{1-3} alkyl, such as C_{1-2} alkyl, such as C_1 alkyl.
57. The compound according to any one of the preceding items, wherein R^7 is H.
58. The compound according to any one of the preceding items, wherein R^7 is -
 $C(O)-O-CH_3$ or $-C(O)-CH_3$.
59. The compound according to any one of the preceding items, wherein R^8 is C_{1-8} alkyl, such as C_{1-7} alkyl, such as C_{1-6} alkyl, such as C_{1-5} alkyl, such as C_{1-4} alkyl, such as C_{1-3} alkyl, such as C_{1-2} alkyl, such as C_1 alkyl.
60. The compound according to any one of the preceding items, wherein R^8 is H.
61. The compound according to any one of the preceding items, wherein R^5 or R^6 is linked to R^7 or R^8 to form a ring, such as R^5 is linked to R^7 .
62. The compound according to any one of the preceding items, wherein R^5 is linked to R^7 to form a four-, five- or six membered ring.
63. The compound according to any one of the preceding items, wherein R^3 or R^4 is linked to R^5 or R^6 to form a ring, such as R^3 is linked to R^5 .
64. The compound for use according to any one of the preceding items, wherein R^3 is linked to R^5 to form a four or five membered ring.
65. The compound according to any one of the preceding items, wherein R^3 or R^4 is linked together to R^7 or R^8 to form a ring, such as R^3 is linked to R^7 .
66. The compound for use according to any one of the preceding items, wherein R^3 is linked to R^7 to form a four membered ring.
67. The compound according to any one of the preceding items, wherein R^9 is -
 $C(H)-$.
68. The compound according to any one of the preceding items, wherein R^{10} is H.
69. The compound according to any one of the preceding items, wherein R^{10} is Cl.

70. The compound according to any one of the preceding items, wherein R^{11} is F.
71. The compound according to any one of the preceding items, wherein R^{12} is - CF_3 .
72. The compound according to any one of the preceding items, wherein R^{12} is - OCF_3 .
73. The compound according to any one of the preceding items, wherein R^{12} is Cl or Br.
74. The compound according to any one of the preceding items, wherein R^{11} is F and R^{12} is - CF_3 .
75. The compound according to any one of the preceding items, wherein R^9 is -C(H)-, R^{10} is H, R^{11} is F and R^{12} is - CF_3 or - OCF_3 .
76. The compound according to any one of the preceding items, wherein R^9 is -C(H)-, R^{10} is H, R^{11} is F and R^{12} is - CF_3 .
77. The compound according to any one of the preceding items, wherein R^9 is -C(H)-, R^{10} is H, R^{11} is F and R^{12} is - OCF_3 .
78. The compound according to any one of the preceding items, wherein $-R^1-R^2$ is not $-CH_3$.
79. The compound according to any one of the preceding items, wherein no more than two of R^{10} , R^{11} and R^{12} are H.
80. The compound according to any one of the preceding items, wherein no more than one of R^{10} , R^{11} and R^{12} are H.
81. The compound according to any one of the preceding items, wherein when R^{11} and R^{12} are H, then R^{10} is halogen.
82. The compound according to any one of the preceding items, wherein no more than five of R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are H.
83. The compound according to any one of the preceding items, wherein when R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are H, then no more than two of R^{10} , R^{11} and R^{12} are H.
84. The compound according to any one of the preceding items, wherein the compound is not selected from the group consisting of
- N1-[1-(3-chlorophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine;
- N1-[1-(3-fluorophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine;
- N1-[1-(4-bromophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine;
- N1-[1-(4-fluorophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine;
- N1-methyl-N1-(1-phenylcyclopropyl)-1,2-ethanediamine;
- N-[1-(4-fluorophenyl)cyclopropyl]-N-methyl-3-azetidinamine;

- N-[1-(4-fluorophenyl)cyclopropyl]-N-methyl-3-pyrrolidinamine;
 N-[1-(3-fluorophenyl)cyclopropyl]-N-methyl-3-pyrrolidinamine;
 N-[1-(3-fluorophenyl)cyclopropyl]-N-methyl-3-azetidinamine;
 N-methyl-N-(1-phenylcyclopropyl)-3-pyrrolidinamine;
 5 N-methyl-N-(1-phenylcyclopropyl)-3-azetidinamine;
 N-methyl-N-(1-phenylcyclopropyl)-3-piperidinamine;
 N-[1-(4-chlorophenyl)cyclopropyl]-N-methyl-3-azetidinamine;
 N-[1-(3-chlorophenyl)cyclopropyl]-N-methyl-3-azetidinamine; and
 N1-[1-(3-bromophenyl)cyclopropyl]-N1-methyl-1,2-ethanediamine.
- 10 85. The compound according to any one of the preceding items, wherein the compound is selected from the group consisting of:
 Methyl N-(2-(dimethylamino)ethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate;
 Methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
 15 Methyl ((1-aminocyclopropyl)methyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate;
 Methyl 3-((1-(4-fluoro-3-(trifluoromethyl) phenyl) cyclopropyl) (methoxycarbonyl)amino)azetidine-1-carboxylate;
 20 Methyl azetidin-3-yl(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
 N-(cyclopropylmethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)azetidin-3-amine;
 Methyl (2-aminoethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
 25 N1-(cyclopropyl methyl)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-2-methyl propane-1,2-diamine;
 N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)cyclopropanecarboxamide;
 N1-cyclopropyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine;
 30 Methyl (2-amino-2-methylpropyl)(1-(3,5-dichlorophenyl)cyclopropyl)carbamate;
 Methyl (2-amino-2-methylpropyl)(1-(3-bromophenyl)cyclopropyl)carbamate;
 Methyl (2-amino-2-methylpropyl)(1-(3-chlorophenyl)cyclopropyl)carbamate;
 1-((azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methyl propan-2-ol;
 35

- Methyl (2-amino-2-methylpropyl)(1-(3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
1-((1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)amino)-2-methylpropan-2-ol;
- 5 Methyl (2-amino-2-methylpropyl)(1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl) carbamate;
N1-cyclobutyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine;
- 10 Ethyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
Methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate;
N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-
- 15 (trifluoromethyl)phenyl)cyclopropyl)methanesulfonamide;
Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 20 Methyl (2-acetamido-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl) cyclopropyl)-2-methylpropane-1,2-diamine;
- 25 Methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
Methyl (S)-(1-(4-fluoro-3-(trifluoro methyl) phenyl)cyclopropyl) ((1-methylpyrrolidin-2-yl)methyl) carbamate;
Methyl (S)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 30 Methyl (R)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
Methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)carbamate;
- 35 Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl)carbamate;

- Methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate;
- Methyl (1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)(2-(hydroxyamino)-2-methylpropyl)carbamate;
- 5 Ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- Ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl) carbamate;
- (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methanesulfonamide;
- 10 (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-((1-methylpyrrolidin-2-yl)methyl)methanesulfonamide;
- Ethyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 15 Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate;
- Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate;
- (R)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methane sulphonamide;
- 20 Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate;
- Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate;
- 25 Isopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- Cyclopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methyl pyrrolidin-2-yl)methyl)carbamate;
- 30 N-((1-amino cyclopropyl)methyl)-1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine;
- Cyclopropylmethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;

N-((1-aminocyclopropyl)methyl)-N-(1-(4-fluoro-3-(trifluoromethyl)-phenyl)cyclopropyl)methanesulfonamide;

Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methylpyrrolidin-2-yl)methyl)carbamate;

5 (1S, 2S)-N1-(1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)-cyclopentane-1,2-diamine;

(1R,2S)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-cyclopentane-1,2-diamine; and

10 Methyl (S)-(azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)-cyclopropyl)carbamate.

86. A pharmaceutical composition comprising the compound according to any one of the preceding items.

87. The compound or pharmaceutical composition according to any one of the preceding items for use in medicine.

15 88. The compound according to any one of items 1 to 50 or pharmaceutical composition according to item 51 for use in the treatment of inflammatory bowel disease (IBD).

89. The compound or pharmaceutical composition for use according to item 88, wherein the IBD is colitis.

20 90. The compound or pharmaceutical composition for use according to item 88, wherein the IBD is ulcerative colitis.

91. The compound or pharmaceutical composition for use according to item 88, wherein the IBD is Crohn's disease.

25 92. The compound according to any one of items 1 to 85 or pharmaceutical composition according to item 86 for use in the treatment of hereditary xerocytosis.

93. The compound according to any one of items 1 to 85 or pharmaceutical composition according to item 86 for use in the treatment of acute respiratory distress syndrome (ARDS).

30 94. A method for treatment of IBD, hereditary xerocytosis or ARDS comprising administration of the compound defined in any one of items 1 to 85 or a composition according to item 86 to a subject in need thereof.

35 95. Use of the compound defined in any one of items 1 to 85 or the composition according to item 86 in the manufacture of a medicament for treatment of IBD, hereditary xerocytosis or ARDS.

Examples

Example [1] - Methyl N-(2-(dimethylamino)ethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate

5 *Step 1*

To a solution of the 4-fluoro-3-(trifluoromethyl)benzonitrile (105.75 mmol) and titanium(IV)isopropoxide (116.33 mmol) in dry diethyl ether was added ethylmagnesium bromide 3M solution in Ether (222.09 mmol) at -78 °C. The resulting yellow solution was stirred for 10 minutes and slowly warmed to rt over 4h. Then boron trifluoride diethyl etherate (211.51 mmol) was added and the reaction mixture was stirred for 24h at rt. The reaction mixture was quenched with 1.5 N HCl solution and extracted with ethyl acetate. The aqueous phase was basified with 10% sodium hydroxide solution and extracted with ethyl acetate. The organic layer was washed with brine and dried over anhydrous sodium sulphate to obtain crude which was purified by flash chromatography using neutral silica gel in 10% TEA in ethyl acetate: Pet-ether as eluent to afford 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropane-1-amine [1.1] (LCMS: MH⁺: 220.1) as product.

20 *Step 2*

To a stirred solution of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] (0.2 g, 0.912 mmol) in dichloromethane (10 mL) was added N,N-diisopropylethylamine (0.35 mL, 2 mmol) followed by methyl chloroformate (0.122 g, 1.29 mmol) at 0 °C and the reaction mixture was stirred at rt for 2 h. The reaction mixture was quenched with water and extracted with ethyl acetate. The organic layer was washed with brine solution and dried over sodium sulphate, filtered and concentrated under reduced pressure to afford the crude product which was purified by flash chromatography using ethyl acetate in hexanes as eluent to afford methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [1.2] as an off white solid (0.19 g, 75%).

30 *Step 3*

To a solution of methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [1.2] (0.08 g, 0.29 mmol) in DMF (3 mL) was added sodium hydride (60%, 0.016 g, 0.32 mmol). The reaction was stirred at 0 °C for 10 minutes then, 2-chloro-N,N-dimethylethan-1-amine hydrochloride (0.037 g, 0.26 mmol) added and the reaction stirred at rt for 1 h. The reaction mixture was poured into ice and extracted with ethyl

acetate. The organic layer was washed with brine solution, dried over sodium sulphate, filtered and concentrated under reduced pressure to afford the crude product, which was purified by column chromatography using ethyl acetate in hexane to afford methyl (2-(dimethylamino)ethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate [1] as yellow gum (0.012 g, 15%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.55 (bs, 1H), 7.45-7.35 (m, 2H), 3.55 (bs, 3H), 3.40-3.30 (m, 2H), 1.70 (bs, 2H), 2.05 (s, 6H), 1.48-1.18 (m, 4H). HRMS calculated for: [C₁₆H₂₀F₄N₂O₂+H]⁺ 349.1534; found: 349.1526 (deviation 2.3).

10 Example [2] - Methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate

Step 1

To a solution of tert-butyl (1-hydroxy-2-methylpropan-2-yl)carbamate (20 g, 105.7 mmol) in dichloromethane (100mL) was added Dess-Martin periodinane (55.4 g, 126.8 mmol) at 0 °C and stirred at room temperature for 16h. The reaction mixture was filtered and the filtrate was extracted with dichloromethane, washed with saturated sodium thiosulphate solution and 10% sodium bicarbonate solution. The organic layer was dried over sodium sulphate, filtered and dried under vacuum to afford crude product, which was purified by flash chromatography using ethyl acetate in pet-ether as solvent to afford tert-butyl (2-methyl-1-oxopropan-2-yl)carbamate [2.1] as a white solid (19.5 g, 99%).

Step 2

25 To a stirred solution of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] (0.05 g, 0.22 mmol) in IPA (5 mL) at 0°C was added tert-butyl (2-methyl-1-oxopropan-2-yl)carbamate [2.1] (0.029 g, 0.16 mmol) and acetic acid (0.05 mL). The reaction was stirred for 1 h at room temperature, then sodium cyanoborohydride (0.016 g, 0.26 mmol) was added at 0°C, then the reaction stirred at room temperature for 2h.

30 Saturated aq. sodium bicarbonate solution was added and the reaction extracted with DCM. The combined organic layers were washed with brine solution, dried over sodium sulfate, filtered and concentrated in vacuo to afford the crude product which was purified by column chromatography using EtOAc in hexane as eluent to afford tert-butyl (1-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [2.2] as a colourless liquid (0.03 g, 34%, LCMS MH⁺ = 391.1).

Step 3

The procedure used in Example [1], Step 2 was adapted such that 16 g of tert-butyl (1-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [2.2] was reacted to afford methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [2.3] (14 g, 76%, LCMS MH⁺ = 449.2).

Step 4

To a stirred solution of methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [2.3] (7.8 g, 17.4 mmol) in dioxane (10 mL) was added dioxane.HCl (20 mL) at 0°C and the reaction stirred at room temperature for 12 h. The reaction was concentrated under reduced pressure and the residue triturated in pentane to afford methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [2] hydrochloride salt as an off-white solid (6.5 g, 97%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.53 (s, 2H), 7.42 (t, *J* = 10.40 Hz, 1H), 3.57 (s, 3H), 3.32 (d, *J* = 16.80 Hz, 2H), 1.65 (bs, 4H), 1.31 (s, 2H), 0.93 (s, 6H). HRMS calculated for: [C₁₆H₂₀F₄N₂O₂+H]⁺ 349.1534; found: 349.1521 (deviation 3.7 ppm).

Example [3] - Methyl ((1-aminocyclopropyl)methyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate

Step 1

To a solution of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] (0.2 g, 0.91 mmol) and 0.16 g of tert-butyl N-(1-formylcyclopropyl)carbamate (0.16 g, 0.91 mmol) in dichloromethane/isopropyl alcohol (3:2, 10 mL) was added sodium triacetoxy borohydride (0.38 g, 1.82 mmol) at room temperature and the reaction stirred for 2 h. The reaction mixture was diluted with water and extracted with DCM. The organic layer was washed with brine solution and dried over sodium sulphate, filtered and concentrated under reduced pressure to afford crude product, which was purified by flash chromatography using ethyl acetate in pet-ether as solvent system to afford tert-butyl (1-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)cyclopropyl)carbamate [3.1] as an off-white solid (0.11 g, 31%, LCMS MH⁺ = 389.2).

Step 2

The procedure used in Example [1], Step 2 was adapted such that 0.11 g of tert-butyl (1-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)cyclopropyl)carbamate [3.1] and 0.073 g of methyl chloroformate was reacted to afford the product methyl ((1-((tert-butoxycarbonyl)amino)cyclopropyl)methyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [3.2] as an off white solid (0.1 g, 80%, LCMS $MH^+ = 447.2$).

Step 3

The procedure used in Example [2], Step 4 was adapted such that 0.1 g of methyl ((1-((tert-butoxycarbonyl)amino)cyclopropyl)methyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [3.2] was reacted to afford methyl ((1-aminocyclopropyl)methyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate [3] as a brown solid (0.055 g, 55%). 1H -NMR (400 MHz, DMSO- d_6): δ 7.60-7.32 (m, 4H), 3.65 (s, 3H), 1.60-1.42 (m, 2H), 1.36-1.28 (m, 2H), 1.30-1.21 (m, 2H), 0.95-0.70 (m, 5H). HRMS calculated for: $[C_{16}H_{18}F_4N_2O_2+H]^+$ 347.1377; found: 347.1369 (deviation 2.5 ppm).

Example [4] - Methyl 3-((1-(4-fluoro-3-(trifluoromethyl) phenyl) cyclopropyl) (methoxycarbonyl)amino)azetidine-1-carboxylate

Step 1

The procedure used in Example [3], Step 1 was adapted such that 0.5 g of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine and 0.117 g of tert-butyl 3-oxoazetidine-1-carboxylate were reacted to afford the product tert-butyl 3-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)azetidine-1-carboxylate [4.1] (0.39 g, crude, LCMS $MH^+ = 375.1$).

Step 2

The procedure used in Example [2], Step 4 was adapted such that 0.3 g of tert-butyl 3-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)azetidine-1-carboxylate [4.1] was reacted to afford the product N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) azetidin-3-amine [4.2] (0.2 g, crude, LCMS $MH^+ = 275.1$).

Step 3

The procedure used in Example [1], Step 2 was adapted such that 0.2 g of N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)azetidin-3-amine and 0.2 mL of methyl carbonochloridate were reacted to afford methyl 3-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)azetidine-1-carboxylate [4] (0.045 g, 30%). MS (M+1)⁺ = 391.1. ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.45 (t, *J* = 8.80 Hz, 2H), 7.30 (d, *J* = 5.20 Hz, 1H), 4.40 (q, *J* = 6.80 Hz, 1H), 4.20-3.98 (m, 4H), 3.61 (s, 3H), 3.52 (s, 3H), 1.40 (s, 4H). HRMS calculated for: [C₁₇H₁₈F₄N₂O₄+H]⁺ 391.1275; found: 391.1264 (deviation 2.8 ppm).

Example [5] - Methyl azetidin-3-yl(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate

Step 1

The procedure used in Example [1], Step 2 was adapted such that 0.3 g of tert-butyl 3-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)azetidine-1-carboxylate [4.1] and 0.226 g of methyl carbonochloridate were reacted to afford the product tert-butyl 3-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)azetidine-1-carboxylate [5.1] (0.3 g, crude, LCMS MH⁺ = 433.1).

Step 2

The procedure used in Example [2], Step 4 was adapted such that 0.25 g of tert-butyl 3-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)azetidine-1-carboxylate [5.1] was reacted to afford methyl azetidin-3-yl(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [5] (0.04 g, 40%). MS (M+1)⁺ = 333.1. ¹H-NMR (400 MHz, CDCl₃): δ 7.32-7.27 (m, 2H), 7.15 (t, *J* = 8.80 Hz, 1H), 4.60-4.50 (m, 1H), 4.50-4.20 (m, 4H), 3.82 (s, 3H), 1.32-1.25 (m, 4H). HRMS calculated for: [C₁₅H₁₆F₄N₂O₂+H]⁺ 333.1221; found: 333.1215 (deviation 1.6 ppm).

Example [6] - N-(cyclopropylmethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)azetidin-3-amine

Step 1

The procedure used in Example [3], Step 1 was adapted such that 0.25 g of tert-butyl 3-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)azetidine-1-carboxylate [4.1] and 0.274 g of cyclopropanecarbaldehyde was reacted to afford the product tert-butyl 3-((cyclopropylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl) cyclopropyl) amino)azetidine-1-carboxylate as colourless gum (0.15 g, crude, LCMS MH⁺ = 429.1).

Step 2

The procedure used in Example [2], Step 4 was adapted such that 0.15 g of tert-butyl 3-((cyclopropylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)azetidine-1-carboxylate [6.1] was reacted to afford N-(cyclopropylmethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)azetidin-3-amine [6] as colourless gum (0.1 g, 87%). MS (M+1)⁺ = 329.1. ¹H-NMR (400 MHz, DMSO-*d*₆): δ 8.53 (s, 1H), 8.38 (s, 1H), 7.67-7.62 (m, 2H), 7.47 (t, *J* = 8.80 Hz, 1H), 3.95-3.82 (m, 3H), 3.79 (t, *J* = 8.00 Hz, 1H), 2.38-2.32 (m, 2H), 1.17 (q, *J* = 42.00 Hz, 2H), 0.96 (q, *J* = 11.60 Hz, 2H), 1.40-1.30 (m, 1H), 0.47-0.42 (m, 2H), 0.13 (q, *J* = 3.20 Hz, 2H). HRMS calculated for: [C₁₇H₂₀F₄N₂+H]⁺ 329.1635; found: 329.1624 (deviation 3.3 ppm).

Example [7] - Methyl (2-aminoethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate

Step 1

The procedure used in Example [3], Step 1 was adapted such that 0.4 g of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] and 0.29 g of tert-butyl (2-oxoethyl)carbamate was reacted to afford the product tert-butyl (2-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)ethyl)carbamate [7.1] as an off white solid (0.4 g, 60%, LCMS MH⁺ = 363.2).

Step 2

The procedure used in Example [1], Step 2 was adapted such that 0.4 g of tert-butyl (2-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)ethyl)carbamate [7.1] and 0.32 g of methyl chloroformate was reacted to afford the product methyl (2-((tert-butoxycarbonyl)amino)ethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [7.2] as an off white solid (0.4 g, crude, LCMS MH⁺ = 421.2).

Step 3

The procedure used in Example [2], Step 4 was adapted such that 0.4 g methyl (2-
((tert-butoxycarbonyl)amino)ethyl)(1-(4-fluoro-3-(trifluoromethyl)
5 phenyl)cyclopropyl)carbamate [7.2] was reacted to afford methyl (2-aminoethyl)(1-(4-
fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [7] as a yellow oil (0.011 g,
04%). ¹H-NMR (400 MHz, CDCl₃): δ 7.4 (bs, 3H), 3.72 (s, 3H), 3.40 (s, 2H), 2.82 (s,
2H), 1.45 (s, 2H), 1.25 (s, 4H). HRMS calculated for: [C₁₄H₁₆F₄N₂O₂+H]⁺ 321.1221;
found: 321.1217 (deviation 1.1 ppm).

10

Example [8] - N1-(cyclopropyl methyl)-N1-(1-(4-fluoro-3-
(trifluoromethyl)phenyl)cyclopropyl)-2-methyl propane-1,2-diamine

Step 1

15 The procedure used in Example [3], Step 1 was adapted such that 0.22 g of tert-butyl
(1-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methyl propan-2-
yl)carbamate and 0.047 g of cyclopropanecarbaldehyde was reacted to afford the
product tert-butyl (1-((cyclopropylmethyl)(1-(4-fluoro-3-
(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [8.1] as a
20 colourless gum (0.2 g, 80%, LCMS MH⁺ = 459.2).

Step 2

The procedure used in Example [2], Step 4 was adapted such that 0.2 g of tert-butyl (1-
((cyclopropylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-
25 methylpropan-2-yl)carbamate [8.1] was reacted to afford N1-(cyclopropyl methyl)-N1-
(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-2-methyl propane-1,2-diamine [8] as
a yellow oil (0.1 g, 66%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.70-7.60 (m, 2H), 7.45 (t, *J*
= 40.00 Hz, 1H), 2.52-2.40 (m, 2H), 2.37 (d, *J* = 6.80 Hz, 2H), 1.40-1.20 (m, 4H), 1.10-
1.00 (m, 2H), 1.00-0.88 (m, 7H), 0.45 (q, *J* = 1.20 Hz, 2H), 0.07 (t, *J* = 3.60 Hz, 2H).
30 HRMS calculated for: [C₁₈H₂₄F₄N₂+H]⁺ 345.1948; found: 345.1937 (deviation 3.3 ppm).

Example [9] - N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-
(trifluoromethyl)phenyl)cyclopropyl)cyclopropanecarboxamide

35 *Step 1*

To a stirred solution of tert-butyl (1-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methyl propan-2-yl)carbamate [2.2] (0.1 g, 0.256 mmol) and DIPEA (0.09 mL, 0.512 mmol) in DCM (5 mL) was added cyclopropanecarbonyl chloride (0.054 g, 0.512 mmol) and the reaction stirred at room temperature for 18 h. The reaction was diluted with water and extracted with DCM. The combined organic layers were washed with brine solution, dried over sodium sulfate, filtered and concentrated under reduced pressure to afford tert-butyl (1-(N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)cyclopropanecarboxamido)-2-methylpropan-2-yl)carbamate [9.1] as a colourless liquid (0.1 g, 80%, LCMS MH^+ = 445.2).

10

Step 2

The procedure used in Example [2], Step 4 was adapted such that 0.1 g of tert-butyl (1-(N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)cyclopropanecarboxamido)-2-methylpropan-2-yl)carbamate [9.1] was reacted to afford the crude product, which was purified by Prep HPLC using 0.2% TFA in acetonitrile to afford N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)cyclopropanecarboxamide [9] as TFA salt as a pale brown solid (0.08 g, 80%). 1H -NMR (400 MHz, $DMSO-d_6$): δ 7.76 (s, 3H), 7.52 (t, J = 11.60 Hz, 1H), 7.44-7.39 (m, 1H), 7.27 (dd, J = 3.20, 8.40 Hz, 1H), 3.88 (d, J = 60.00 Hz, 1H), 3.30 (s, 1H), 2.15-1.95 (m, 1H), 1.95-1.80 (m, 1H), 1.80-1.65 (m, 1H), 1.65-1.50 (m, 1H), 1.50-1.35 (m, 1H), 1.22 (s, 4H), (s, 3H), 0.90-0.80 (m, 2H), 0.80-0.65 (m, 1H), 0.65-0.55 (m, 1H). HRMS calculated for: $[C_{18}H_{22}F_4N_2O+H]^+$ 359.1741; found: 359.1733 (deviation 2.2 ppm).

20

25 Example [10] - N1-cyclopropyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine

Step 1

The procedure used in Example [1], Step 1 was adapted such that 1.5 g of 3-(trifluoromethyl)benzonitrile was reacted to afford 1-(3-(trifluoromethyl)phenyl)cyclopropan-1-amine [10.1] (0.5 g, 30%, LCMS MH^+ = 202.1).

30

Step 2

The procedure used in Example [3], Step 1 was adapted such that 0.275 g of 1-(3-(trifluoromethyl)phenyl)cyclopropan-1-amine [10.1] was reacted to afford tert-butyl (2-

35

methyl-1-((1-(3-(trifluoromethyl)phenyl)cyclopropyl)amino)propan-2-yl)carbamate [10.2] as an off-white solid (0.37 g, 73%, LCMS MH^+ = 373.2).

Step 3

5 The procedure used in Example [2], Step 2 was adapted such that 0.270 g of tert-butyl (2-methyl-1-((1-(3-(trifluoromethyl)phenyl)cyclopropyl)amino)propan-2-yl)carbamate [2.2] was reacted with [(1-ethoxycyclopropyl)oxy]trimethylsilane to afford tert-butyl (1-(cyclopropyl(1-(3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [10.3] as an off-white gum (0.1 g, 36%, LCMS MH^+ = 413.4).

10

Step 4

The procedure used in Example [2], Step 4 was adapted such that 0.06 g tert-butyl(1-(cyclopropyl(1-(3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [10.3] was reacted to afford N1-cyclopropyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine [10] as an off-white solid (0.025 g, 50%, LCMS MH^+ = 313.2). 1H -NMR (400 MHz, DMSO- d_6): δ 7.88 (s, 2H), 7.71-7.59 (m, 4H), 2.91 (s, 2H), 1.57 (m, 1H), 1.25 (d, J = 5.68 Hz, 6H), 1.20-1.00 (m, 2H), 1.00-0.85 (m, 2H), 0.85-0.70 (m, 2H), 0.60-0.40 (m, 2H). HRMS calculated for: $[C_{17}H_{23}F_3N_2+H]^+$ 313.1886; found: 313.1880 (deviation 2.1 ppm).

20

Example [11] - Methyl (2-amino-2-methylpropyl)(1-(3,5-dichlorophenyl)cyclopropyl)carbamate

Step 1

25 The procedure used in Example [1], Step 1 was adapted such that 0.5 g of 3,5-dichlorobenzonitrile was reacted to afford 1-(3,5-dichlorophenyl)cyclopropan-1-amine [11.1] (0.2 g, 34%, LCMS MH^+ = 203.1).

Step 2

30 The procedure used in Example [3], Step 1 was adapted such that 0.2 g of 1-(3,5-dichlorophenyl)cyclopropan-1-amine [11.1] and 0.18 g of tert-butyl (2-methyl-1-oxopropan-2-yl)carbamate [2.1] was reacted to afford the product tert-butyl (1-((1-(3,5-dichlorophenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [11.2] as a brown liquid (0.2 g, 54%, LCMS MH^+ = 374.1).

35

Step 3

The procedure used in Example [1], Step 2 was adapted such that 0.2 g of tert-butyl (1-((1-(3,5-dichlorophenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [11.2] was reacted to afford the product methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(3,5-dichlorophenyl)cyclopropyl)carbamate [11.3] as colourless liquid (0.14 g, 60%, LCMS $MH^+ = 432.2$).

Step 4

The procedure used in Example [2], Step 4 was adapted such that 0.14 g of methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(3,5-dichlorophenyl)cyclopropyl)carbamate [11.3] was reacted to afford methyl (2-amino-2-methylpropyl)(1-(3,5-dichlorophenyl)cyclopropyl)carbamate [11] as an off-white solid (0.037 g, 31%). 1H -NMR (400 MHz, DMSO- d_6): δ 7.80 (bs, 3H), 7.48 (d, $J = 1.20$ Hz, 1H), 7.14 (bs, 1H), 3.65 (s, 3H), 3.56 (d, $J = 2.40$ Hz, 1H), 3.50 (s, 1H), 1.45 (s, 2H), 1.35 (s, 1H), 1.30-1.22 (m, 1H), 1.17 (bs, 6H). HRMS calculated for: $[C_{15}H_{20}Cl_2N_2O_2+H]^+$ 331.0975; found: 331.0966 (deviation 2.7 ppm).

Example [12] - Methyl (2-amino-2-methylpropyl)(1-(3-bromophenyl)cyclopropyl)carbamate

Step 1

The procedure used in Example [1], Step 1 was adapted such that 3 g of 3-bromobenzonitrile was reacted to afford 1-(3-bromophenyl)cyclopropan-1-amine [12.1] (1.3 g, 56%, LCMS $MH^+ = 213.0$).

Step 2

The procedure used in Example [3], Step 1 was adapted such that 1 g of 1-(3-bromophenyl)cyclopropan-1-amine [12.1] and 0.97 g of tert-butyl (2-methyl-1-oxopropan-2-yl)carbamate [2.1] were reacted to afford the product tert-butyl (1-((1-(3-bromophenyl)cyclopropyl) amino)-2-methylpropan-2-yl)carbamate [2.2] as a yellow solid (0.9 g, 67%, LCMS $MH^+ = 384.2$).

Step 3

The procedure used in Example [1], Step 2 was adapted such that 0.5 g of tert-butyl (1-((1-(3-bromophenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [2.2] was

reacted to afford the product methyl (1-(3-bromophenyl)cyclopropyl)(2-((tert-butoxycarbonyl)amino)-2-methylpropyl)carbamate as a colourless liquid (0.4 g, 69%, LCMS $MH^+ = 442.1$).

5 *Step 4*

The procedure used in Example [2], Step 4 was adapted such that 0.1 g of methyl (1-(3-bromophenyl)cyclopropyl)(2-((tert-butoxycarbonyl)amino)-2-methylpropyl)carbamate [12.3] was reacted to afford methyl (2-amino-2-methylpropyl)(1-(3-bromophenyl)cyclopropyl)carbamate [12] as TFA salt (compound was purified by prep HPLC using 0.1% TFA in ACN) as a white solid (0.02 g, 26%). 1H -NMR (400 MHz, DMSO- d_6): δ 7.67 (bs, 2H), 7.42-7.38 (m, 1H), 7.27 (t, $J = 10.40$ Hz, 2H), 7.10 (bs, 1H), 3.62 (s, 3H), 3.52-3.37 (m, 2H), 1.50-1.20 (m, 4H), 1.15 (s, 6H). HRMS calculated for: $[C_{15}H_{21}BrN_2O_2+H]^+$ 341.0859; 343.0839; found: 341.0847; 343.0828 (deviation 3.5; 3.3 ppm).

15

Example [13] - Methyl (2-amino-2-methylpropyl)(1-(3-chlorophenyl)cyclopropyl)carbamate

Step 1

20 The procedure used in Example [1], Step 1 was adapted such that 1 g of 3-bromobenzonitrile was reacted to afford 1-(3-chlorophenyl)cyclopropan-1-amine [13.1] (0.4 g, 32%, LCMS $MH^+ = 168.5$).

Step 2

25 The procedure used in Example [3], Step 1 was adapted such that 0.4 g of 1-(3-chlorophenyl)cyclopropan-1-amine [13.1] and 0.49 g of tert-butyl (2-methyl-1-oxopropan-2-yl)carbamate [2.1] were reacted to afford the product tert-butyl (1-((1-(3-chlorophenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [13.2] as a brown liquid (0.25 g, 31%, LCMS $MH^+ = 339.2$).

30

Step 3

The procedure used in Example [1], Step 2 was adapted such that 0.2 g of tert-butyl (1-((1-(3-chlorophenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [13.2] was reacted to afford methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(3-

chlorophenyl)cyclopropyl)carbamate [13.3] as a colourless liquid (0.17 g, 72%, LCMS $MH^+ = 397.1$).

Step 4

- 5 The procedure used in Example [2], Step 4 was adapted such that 0.17 g of methyl 2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(3-chlorophenyl)cyclopropyl)carbamate [13.3] was reacted to afford methyl 2-amino-2-methylpropyl)(1-(3-chlorophenyl)cyclopropyl)carbamate [13] as TFA salt (compound was purified by Prep HPLC using 0.1% TFA in ACN) as a colourless gum (0.1 g, 78%). 1H -NMR (400 MHz, DMSO- d_6): δ 7.59 (s, 2H), 7.36-7.24 (m, 2H), 7.33 (bs, 2H), 3.62 (s, 3H), 3.50 (bs, 2H), 1.35 (s, 4H), 1.15 (s, 6H). HRMS calculated for: $[C_{15}H_{21}ClN_2O_2+H]^+$ 297.1364; found: 297.1257 (deviation 2.5 ppm).
- 10

- Example [14] - 1-((azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methyl propan-2-ol
- 15

Step 1

- To a stirred solution of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] (1.5 g, 6.84 mmol) and 1-(tert-butoxycarbonyl)azetidine-2-carboxylic acid (1.23 g, 6.15 mmol) in DMF was added HATU (3.12 g, 8.21 mmol), followed by DIPEA (2.2 g, 17.10 mmol) at 0 °C and the reaction mixture was stirred at room temperature for 12h. The reaction mixture was diluted with ethyl acetate, washed with water and brine solution. The organic layer was dried over sodium sulphate, filtered and concentrated under reduced pressure to afford a crude product, which was purified by flash chromatography using ethyl acetate in pet-ether as eluent to afford tert-butyl 2-((1-(4-fluoro-3-(trifluoromethyl)phenyl) cyclopropyl) carbamoyl) azetidine-1-carboxylate [14.1] as a colourless gum (2.3 g, 85%, LCMS $MH^+ = 403.3$)."
- 20
- 25

Step 2

- 30 To a solution of tert-butyl 2-((1-(4-fluoro-3-(trifluoromethyl)phenyl) cyclopropyl)carbamoyl)azetidine-1-carboxylate [14.1] (2.3 g, 5.72 mmol) in tetrahydrofuran was added borane dimethyl sulphide complex (1.8 g, 22.86 mmol) at 0 °C and the reaction refluxed at 70 °C for 2h. The reaction mixture was quenched with methanol at 0 °C until the effervescence stopped. The reaction mixture was concentrated under reduced pressure and the residue was diluted with ethyl acetate
- 35

washed with water and brine solution. The organic layer was dried over sodium sulphate, filtered and concentrated under reduced pressure to afford crude product, which was purified by flash chromatography using ethyl acetate in pet-ether as solvent to afford tert-butyl 2-(((1-(4-fluoro-3-

5 (trifluoromethyl)phenyl)cyclopropyl)amino)methyl)azetidine-1-carboxylate [14.2] as colourless liquid (1.45 g, 55%, LCMS M^+ = 389.4).

Step 3

To a solution of tert-butyl 2-(((1-(4-fluoro-3-

10 (trifluoromethyl)phenyl)cyclopropyl)amino)methyl)azetidine-1-carboxylate [14.2] (0.3 g, 0.77 mmol) and potassium carbonate (0.32 g, 2.31 mmol) in acetonitrile was added methyl bromoacetate (0.35 g, 2.31 mmol) and the reaction mixture was heated at 80 °C for 12h. The reaction mixture was concentrated under reduced pressure and the residue was diluted with ethyl acetate, washed with water and brine solution. The

15 organic layer was dried over sodium sulphate, filtered and concentrated under reduced pressure to afford crude product, which was purified by flash chromatography using ethyl acetate in pet-ether as solvent to afford tert-butyl 2-(((1-(4-fluoro-3-

(trifluoromethyl)phenyl)cyclopropyl)(2-methoxy-2-oxoethyl) amino)methyl)azetidine-1-

20 carboxylate [14.3] as a colourless liquid.(0.25 g, 70%, LCMS MH^+ = 461.2).

Step 4

To a stirred solution of tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethyl)

phenyl)cyclopropyl)(2-methoxy-2-oxoethyl) amino)methyl) azetidine-1-carboxylate

[14.3] (0.2 g, 0.43 mmol) in tetrahydrofuran (5 mL) at 0°C was added

25 methylmagnesium bromide (0.21 g, 1.74 mmol) at 0 °C and the reaction mixture was stirred at same temperature for 2h, then stirred at rt for 1h. The reaction mixture was quenched with aqueous ammonium chloride solution and extracted with ethyl acetate, washed with water and brine solution. The organic layer was dried over sodium

sulphate, filtered and concentrated under reduced pressure to afford the crude product

30 which was purified by column chromatography using ethyl acetate/hexane as eluent to afford tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl) cyclopropyl)(2-hydroxy-2-

methylpropyl)amino) methyl)azetidine-1-carboxylate [14.4] as colourless gum (0.15 g, 75%, LCMS MH^+ = 461.2).

35 Step 5

The procedure used in Example [1], Step 2 was adapted such that 0.12 g of tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(2-hydroxy-2-methylpropyl)amino)methyl)azetidine-1-carboxylate [14.4] was reacted to afford 1-
5 ((azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methylpropan-2-ol [14] as an off-white solid (0.06 g, 66%, LCMS MH^+ = 361.2). 1H -NMR (400 MHz, $CDCl_3$): δ 7.52-7.45 (m, 2H), 7.07 (q, J = 62.80 Hz, 1H), 3.45 (q, J = 32.80 Hz, 1H), 3.01 (s, 1H), 2.60 (s, 2H), 2.58-1.87 (m, 4H), 1.60-1.40 (m, 2H), 1.30-1.22 (m, 2H), 1.20 (s, 6H), 0.95-0.90 (m, 2H). HRMS calculated for: $[C_{18}H_{24}F_4N_2O+H]^+$ 361.1898; found: 361.1889 (deviation 2.4 ppm).

10

Example [15] - Methyl (2-amino-2-methylpropyl)(1-(3-(trifluoromethyl)phenyl)cyclopropyl)carbamate

Step 1

15 The procedure used in Example [3], Step 1 was adapted such that 0.4 g of 1-(3-(trifluoromethyl)phenyl)cyclopropan-1-amine [10.1] and 0.52 g of tert-butyl (2-methyl-1-oxopropan-2-yl)carbamate were reacted to afford the product tert-butyl (2-methyl-1-((1-(3-(trifluoromethyl)phenyl)cyclopropyl)amino)propan-2-yl)carbamate [15.1] (0.35 g, 69%, LCMS MH^+ = 373.3).

20

Step 2

The procedure used in Example [1], Step 2 was adapted such that 0.3 g of tert-butyl (2-methyl-1-((1-(3-(trifluoromethyl)phenyl)cyclopropyl)amino)propan-2-yl)carbamate [15.1] was reacted to afford the product methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [15.2] as colourless
25 liquid (0.2 g, 57%, LCMS MH^+ = 431.4).

Step 3

30 The procedure used in Example [2], Step 4 was adapted such that 0.2 g of methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [15.2] was reacted to afford methyl (2-amino-2-methylpropyl)(1-(3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [15] as an off-white solid (0.03 g, 20%). 1H -NMR (400 MHz, $DMSO-d_6$): δ 7.85 (bs, 2H), 7.57 (t, J = 5.20 Hz, 2H), 7.40 (bs, 2H), 3.63 (s, 3H), 3.53 (s, 1H), 1.50 (bs, 2H), 1.37 (bs, 2H),

1.23 (bs, 1H), 1.18 (s, 6H). HRMS calculated for: $[C_{16}H_{21}F_3N_2O_2+H]^+$ 331.1628; found: 331.1620 (deviation 2.5 ppm).

5 Example [16] - 1-((1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)amino)-2-methylpropan-2-ol

The procedure used in Example [2], Step 2 was adapted such that 0.09 g of 1-((azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) amino)-2-methylpropan-2-ol [14] was reacted to afford 1-((1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)amino)-2-methylpropan-2-ol [16] as colourless gum (0.02 g, 21%, LCMS $MH^+ = 375.2$). 1H -NMR (400 MHz, $CDCl_3$): δ 7.46-7.43 (m, 2H), 7.14 (t, $J = 9.20$ Hz, 1H), 3.52-3.48 (m, 1H), 3.42-3.33 (m, 1H), 3.01-2.96 (m, 1H), 2.80-2.70 (m, 3H), 2.65 (d, $J = 13.20$ Hz, 1H), 2.41 (s, 3H), 2.15-2.12 (m, 1H), 1.80-1.70 (m, 2H), 1.30-1.23 (m, 2H), 1.21-1.19 (m, 3H), 1.90-1.30 (m, 3H), 1.08-0.80 (m, 1H), 0.30-0.20 (m, 1H). HRMS calculated for: $[C_{19}H_{26}F_4N_2O+H]^+$ 375.2054; found: 375.2044 (deviation 2.6 ppm).

20 Example [17] - Methyl (2-amino-2-methylpropyl)(1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl) carbamate

Step 1

The procedure used in Example [3], Step 1 was adapted such that 0.054 g of 1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropan-1-amine [36.6] was reacted to afford tert-butyl (1-((1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [17.1] as an off-white solid (0.052 g, 54%, LCMS $MH^+ = 392.2$).

Step 2

The procedure used in Example [1], Step 2 was adapted such that 0.05 g of tert-butyl (1-((1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [17.1] was reacted to afford methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl) carbamate [17.2] as an off-white solid (0.042 g, 73%, LCMS $MH^+ = 450.2$).

Step 3

The procedure used in Example [2], Step 4 was adapted such that 0.04 g of methyl (2-(((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl)carbamate was reacted to afford methyl (2-amino-2-methylpropyl)(1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl) carbamate [17] as an off-white solid (0.021 g, 61%, LCMS $MH^+ = 350.2$). 1H -NMR (400 MHz, $DMSO-d_6$): δ 8.81 (s, 1H), 7.80 (s, 2H), 7.33 (s, 1H), 3.58 (s, 3H), 3.26-3.25 (m, 2H), 1.58-1.54 (m, 4H), 1.24-1.21 (m, 6H). HRMS calculated for: $[C_{15}H_{19}F_4N_3O_2+H]^+$ 350.1486; found: 350.1483 (deviation 0.9 ppm).

Example [18] - N1-cyclobutyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine

To a stirred solution of 2-(cyclobutyl(1-(3-(trifluoromethyl)phenyl)cyclopropyl)amino)acetonitrile [82] (0.25 g, 0.85 mmol) in toluene (3 mL) was added titanium(IV)isopropoxide (0.26 mL, 0.85 mmol) at $-40^\circ C$ and the reaction stirred for 15 minutes. Methylmagnesium bromide solution (7.2 mmol) was added dropwise over a period of 10 minutes and the reaction subsequently stirred at room temperature for 18 h. The reaction was quenched with ammonium chloride solution and extracted with DCM (3 x 50 mL). The combined organics were concentrated in vacuo and purified by flash column chromatography using 15% EtOAc/pet ether as eluent to afford N1-cyclobutyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine [18] as a brownish gum (0.05 g, 18%, LCMS $MH^+ = 327.1$). 1H -NMR (400 MHz, $DMSO-d_6$): δ 7.60-7.54 (m, 5H), 3.45-3.43 (m, 1H), 2.76 (s, 2H), 1.99-1.93 (m, 4H), 1.56-1.45 (m, 2H), 1.37-1.34 (m, 2H), 1.24-1.22 (m, 5H), 1.10-1.07 (m, 2H). HRMS calculated for: $[C_{18}H_{25}F_3N_2+H]^+$ 327.2042; found: 327.2036 (deviation 2.2 ppm).

Example [19] - Ethyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate

Step 1

The procedure used in Example [1], Step 2 was adapted such that 0.15 g of tert-butyl (1-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [2.2] was reacted with ethyl chloroformate to afford ethyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl) (1-(4-fluoro-3-

(trifluoromethyl)phenyl)cyclopropyl)carbamate [19.1] as a colourless liquid (0.14 g, 79% LCMS MH^+ = 463.2).

Step 2

- 5 The procedure used in Example [2], Step 4 was adapted such that 0.1 g of ethyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [19.1] was reacted to afford ethyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [19] as an off-white solid (0.062 g, 72%, LCMS MH^+ = 363.2). 1H -NMR (400 MHz, DMSO- d_6): δ 7.77 (bs, 3H), 7.63-7.58 (m, 1H), 7.54-7.45 (m, 2H), 4.13-4.08 (m, 2H), 3.55 (bs, 2H), 1.44 (s, 2H), 1.35 (bs, 2H), 2.00-1.50 (m, 6H), 1.50-1.10 (m, 3H). HRMS calculated for: $[C_{17}H_{22}F_4N_2O_2+H]^+$ 363.1690; found: 363.1686 (deviation 1.1 ppm).
- 10

- Example [20] - Methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate
- 15

Step 1

- The procedure used in Example [3], Step 1 was adapted such that 0.25 g of 1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropan-1-amine [9.1] and 0.19 g of tert-butyl (2-methyl-1-oxopropan-2-yl)carbamate [2.1] were reacted to afford the product tert-butyl (1-((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [20.1] as brown gum (0.26 g, 60%, LCMS MH^+ = 407.2).
- 20

Step 2

- 25 The procedure used in Example [1], Step 2 was adapted such that 0.26 g of tert-butyl (1-((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [20.1] was reacted to afford the product methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate [20.2] as colourless liquid (0.18 g, 60%, LCMS MH^+ = 465.2).
- 30

Step 3

- The procedure used in Example [2], Step 4 was adapted such that 0.18 g of methyl (2-((tert-butoxycarbonyl)amino)-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate [20.2] was reacted to afford methyl (2-
- 35

amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate [20] as an off-white solid (0.1 g, 71%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.68 (bs, 2H), 7.46-7.42 (m, 1H), 7.28 (bs, 2H), 3.62 (s, 3H), 3.50 (bs, 2H), 1.41 (bs, 2H), 1.25 (bs, 2H), 1.12 (s, 6H). HRMS calculated for: [C₁₆H₂₀F₄N₂O₃+H]⁺ 365.1483; found: 365.1473 (deviation 2.6 ppm).

Example [21] - N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methanesulfonamide

10 *Step 1*

The procedure used in Example [2], Step 4 was adapted such that 0.3 g of tert-butyl ((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methyl propan-2-yl)carbamate [2.2] was reacted with 0.26 g of sulfuryl dichloride to afford tert-butyl (1-(N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methylsulfonamido)-2-methylpropan-2-yl)carbamate [21.1] (0.12 g, 33%, LCMS MH⁺ = 469.2).

Step 2

The procedure used in Example [2], Step 4 was adapted such that 0.12 g of tert-butyl (1-(N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) methyl sulfonamido)-2-methylpropan-2-yl)carbamate [21.1] was reacted to afford N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methanesulfonamide [21] as an off-white solid (0.048 g, 50%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.77 (dd, *J* = 2.40, 6.40 Hz, 1H), 7.63-7.57 (m, 1H), 7.48-7.40 (m, 1H), 3.16 (s, 2H), 3.11 (s, 3H), 1.75-1.68 (m, 2H), 1.55 (s, 2H), 1.36 (d, *J* = 1.20 Hz, 2H), 0.98 (s, 6H). HRMS calculated for: [C₁₅H₂₀F₄N₂O₂S+H]⁺ 369.1254; found: 369.1242 (deviation 3.2 ppm).

Example [22] - Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-yl)methyl)carbamate

30 *Step 1*

The procedure used in Example [2], Step 1 was adapted such that 2 g of tert-butyl (2R)-2-(hydroxymethyl)pyrrolidine-1-carboxylate was reacted to afford tert-butyl (2R)-2-formylpyrrolidine-1-carboxylate [22.1] as a pale yellow liquid (1.5 g (crude), 79%).

35 *Step 2*

The procedure used in Example [2], Step 2 was adapted such that 1.5 g of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] was reacted with tert-butyl (2R)-2-formylpyrrolidine-1-carboxylate [22.1] to afford tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) amino)methyl)pyrrolidine-1-carboxylate as
5 colourless liquid (1.3 g, 48%, LCMS $MH^+ = 403.2$).

Step 3

The procedure used in Example [1], Step 2 was adapted such that 0.25 g of tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [22.2] and 0.09 g of methyl chloroformate were reacted to afford tert-butyl
10 (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl) amino)methyl)pyrrolidine-1-carboxylate [22.3] as a colourless gum (0.13 g, 52%, LCMS $MH^+ = 361.4$ (Boc-cleaved mass)).

Step 4

To a stirred solution of tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) (methoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate [22.3] (0.12 g, 0.26 mmol) in DCM was added HCl gas in diethyl ether at 0°C and the reaction stirred at room temperature for 16 h. The reaction mixture was
20 concentrated under reduced pressure to afford crude product which was purified by trituration in diethyl ether to afford methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [22] (HCl salt) as a white solid (0.1 g, 97%, LCMS $MH^+ = 361.2$). 1H -NMR (400 MHz, DMSO- d_6): δ 9.30-9.00 (bs, 1H), 8.60-8.20 (bs, 1H), 7.54-7.38 (m, 3H), 3.76-0.75 (m, 1H), 3.62-3.50 (m,
25 4H), 3.30-3.18 (m, 1H), 3.12-3.00 (m, 1H), 2.04-1.94 (m, 1H), 1.90-1.82 (m, 2H), 1.57-1.50 (m, 2H), 1.40-1.30 (m, 2H). HRMS calculated for: $[C_{17}H_{20}F_4N_2O_2+H]^+$ 361.1534; found: 361.1521 (deviation 3.6 ppm).

Example [23] - Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate

30

Step 1

The procedure used in Example [2], Step 1 was adapted such that 3 g of tert-butyl (2S)-2-(hydroxymethyl)pyrrolidine-1-carboxylate was reacted to afford tert-butyl (2S)-2-formylpyrrolidine-1-carboxylate [23.1] as a pale yellow liquid (2 g (crude), 62%).
35

Step 2

The procedure used in Example [2], Step 2 was adapted such that 1 g of 1-[4-fluoro-3-(trifluoromethyl)phenyl]cyclopropan-1-amine [1.1] was reacted with tert-butyl (2S)-2-formylpyrrolidine-1-carboxylate [23.1] to afford tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [23.2] (1.2 g, 66%, LCMS MH⁺ = 403.2).

Step 3

The procedure used in Example [1], Step 2 was adapted such that 1 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [23.2] and 0.21 g of methyl chloroformate were reacted to afford tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate (0.25 g, 73%, LCMS MH⁺ = 461.2).

Step 4

The procedure used in Example [22], Step 4 was adapted such that 0.12 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate [23.3] was reacted to afford methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [23] (HCl salt) as an off-white solid (0.09 g, 90%, LCMS MH⁺ = 361.2). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 9.60-9.40 (bs, 1H), 9.00-8.20 (bs, 1H), 7.49-7.38 (m, 3H), 3.73-3.60 (m, 6H), 3.23-3.07 (m, 2H), 1.98-1.85 (m, 3H), 1.70-1.40 (m, 3H), 1.36-1.29 (m, 2H). HRMS calculated for: [C₁₇H₂₀F₄N₂O₂+H]⁺ 361.1534; found: 361.1524 (deviation 2.6 ppm).

Example [24] - Methyl (2-acetamido-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate

The procedure used in Example [1], Step 2 was adapted such that 0.1 g of methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [2] and 0.022 g of acetyl chloride were reacted to afford methyl (2-acetamido-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [24] as a colourless gum (0.034 g, 30%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.53-7.38 (m, 2H), 7.30 (bs, 3H), 3.73 (s, 2H), 3.61 (s, 3H), 1.52 (s, 3H), 1.43 (bs, 2H), 1.30 (bs, 2H), 1.14

(s, 6H). HRMS calculated for: $[C_{18}H_{22}F_4N_2O_3+H]^+$ 391.1639; found: 391.1637 (deviation 0.7 ppm).

Example [25] - N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-2-methylpropane-1,2-diamine

The procedure used in Example [2], Step 4 was adapted such that 0.1 g of tert-butyl (1-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methylpropan-2-yl)carbamate [2.2] was reacted to afford N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-2-methylpropane-1,2-diamine [25] as a colourless gum (0.05 g, 67%). 1H -NMR (400 MHz, DMSO- d_6): δ 7.74 (dd, J = 2.00, 7.00 Hz, 1H), 7.62-7.56 (m, 1H), 7.43-7.36 (m, 1H), 2.59 (bs, 1H), 2.20 (s, 2H), 1.40 (bs, 2H), 1.00-0.98 (m, 2H), 0.93 (s, 6H). HRMS calculated for: $[C_{14}H_{18}F_4N_2+H]^+$ 290.1479; found: 291.1467 (deviation 4.1 ppm).

Example [26] - Methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate

Step 1

The procedure used in Example [1], Step 2 was adapted such that 0.45 g of tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)azetidine-1-carboxylate [14.2] was reacted to afford tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)azetidine-1-carboxylate [26.1] as a colourless gum (0.125 g, 25%, LCMS MH^+ = 447.1).

Step 2

To a stirred solution of tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)azetidine-1-carboxylate [26.1] (0.25 g, 0.56 mmol) in dichloromethane (6 mL) at 0°C was added trifluoroacetic acid (4 mL, 52 mmol) and the reaction stirred at rt for 12h. The reaction was diluted with sat. aq. sodium bicarbonate solution and extracted with ethyl acetate. The combined organic layers were washed with brine solution and concentrated under reduced pressure to afford the crude product which was purified by Prep HPLC to afford methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [26] trifluoroacetate salt as a white solid, (0.035 g, 19%, LCMS MH^+ = 347.1). 1H -NMR (400 MHz, DMSO- d_6): δ 8.58 (bs, 1H), 7.60-7.45 (m, 2H), 7.38 (s, 1H), 4.43 (bs, 1H), 3.90

(bs, 1H), 3.85-3.75 (m, 2H), 3.66-3.63 (m, 3H), 3.63-3.60 (m, 1H), 2.35-2.10 (m, 2H), 1.50-1.25 (m, 4H). HRMS calculated for: $[C_{16}H_{18}F_4N_2O_2+H]^+$ 347.1377; found: 347.1374 (deviation 0.9 ppm).

5 Example [27] - Methyl (S)-(1-(4-fluoro-3-(trifluoro methyl) phenyl)cyclopropyl) ((1-methylpyrrolidin-2-yl)methyl) carbamate

The procedure used in Example [2], Step 2 was adapted such that 0.08 g of methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [23] was reacted to afford methyl (S)-(1-(4-fluoro-3-(trifluoro methyl) phenyl)cyclopropyl) ((1-methylpyrrolidin-2-yl)methyl) carbamate [27] as a colourless gum (0.055 g, 73%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.49-7.42 (m, 3H), 3.65-3.60 (m, 3H), 3.33-3.26 (m, 2H), 2.89-2.86 (m, 1H), 2.50-2.30 (m, 1H), 2.20 (s, 3H), 2.08-2.06 (m, 1H), 1.23-1.61 (m, 8H). HRMS calculated for: $[C_{18}H_{22}F_4N_2O_2+H]^+$ 375.1690; found: 375.1683 (deviation 1.8 ppm).

Example [28] - Methyl (S)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate

20 *Step 1*

The procedure used in Example [3], Step 1 was adapted such that 0.3 g of 1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropan-1-amine [9.1] was reacted to afford tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [28.1] as a pale brown gum (0.3 g, 54%, LCMS MH⁺ = 419.2).

25

Step 2

The procedure used in Example [1], Step 2 was adapted such that 0.3 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [28.1] was reacted to afford tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate [28.2] as colourless gum (0.3 g, 85%, LCMS MH⁺ = 477.1).

30

Step 3

The procedure used in Example [2], Step 4 was adapted such that 0.15 g of tert-butyl (S)-2-(((1-(4-fluoro-3-

35

(trifluoromethoxy)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate [28.2] was reacted to afford methyl (S)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [28] as a white solid (0.65 g, 60%, LCMS $MH^+ = 377.1$). 1H -NMR (400 MHz, DMSO- d_6): δ 9.20 (bs, 1H), 8.45 (bs, 1H), 7.50-7.42 (m, 1H), 7.25 (bs, 2H), 3.69 (bs, 1H), 3.64 (s, 3H), 3.56 (s, 2H), 3.30-3.20 (m, 1H), 3.12-3.02 (m, 1H), 2.05-1.45 (m, 1H), 1.95-1.28 (m, 2H), 1.60-1.40 (m, 3H), 1.35 (bs, 1H), 1.25 (bs, 1H). HRMS calculated for: $[C_{17}H_{20}F_4N_2O_3+H]^+$ 377.1483; found: 377.1476 (deviation 1.8 ppm).

10 Example [29] - Methyl (R)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate

Step 1

The procedure used in Example [3], Step 1 was adapted such that 0.4 g of 1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropan-1-amine [9.1] was reacted to afford tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [29.1] as a colourless liquid (0.35 g, 49%, LCMS $MH^+ = 419.1$).

Step 2

20 The procedure used in Example [1], Step 2 was adapted such that 0.35 g of tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [29.1] was reacted to afford tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate [29.2] as a pale brown gum (0.2 g, 50%, LCMS $MH^+ = 477.2$).

Step 3

25 The procedure used in Example [2], Step 4 was adapted such that 0.1 g of tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate [29.2] was reacted to afford methyl (R)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [29] (HCl salt) as a white solid (0.05 g, 58%, LCMS $MH^+ = 377.2$). 1H -NMR (400 MHz, DMSO- d_6): δ 9.15 (bs, 1H), 8.25 (bs, 1H), 7.27-7.22 (m, 1H), 7.13 (bs, 2H), 3.80-3.68 (m, 1H), 3.64 (s, 3H), 3.62-3.50 (m, 2H), 3.30-3.20 (m, 1H), 3.12-3.02 (m, 1H), 2.05-1.78 (m, 3H), 1.60-

1.40 (m, 2H), 1.40-1.20 (m, 2H). HRMS calculated for: $[C_{17}H_{20}F_4N_2O_3+H]^+$ 377.1483; found: 377.1472 (deviation 2.9 ppm).

Example [30] - Methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)carbamate

5

The procedure used in Example [2], Step 2 was adapted such that 0.19 g of methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [26] was reacted to afford methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)carbamate [30] as a colourless gum (0.02 g, 10%, LCMS $MH^+ = 361.1$). 1H -NMR (400 MHz, $CDCl_3$): δ 7.28-7.20 (m, 1H), 7.15-7.10 (m, 1H), 7.06-7.03 (m, 1H), 3.69 (s, 3H), 3.45-3.25 (m, 3H), 2.64 (bs, 1H), 2.20 (bs, 3H), 1.95-1.73 (m, 2H), 1.63 (bs, 2H), 1.44 (bs, 1H), 1.30-1.18 (m, 2H). HRMS calculated for: $[C_{17}H_{20}F_4N_2O_2+H]^+$ 361.1534; found: 361.1528 (deviation 1.7 ppm).

15

Example [31] - Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl)carbamate

The procedure used in Example [2], Step 2 was adapted such that 0.02 g of methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [22] was reacted to afford methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl)carbamate [31] as colourless gum (0.013 g, 62%). 1H -NMR (400 MHz, $DMSO-d_6$): δ 7.60-7.30 (m, 3H), 3.59 (s, 3H), 3.35-3.25 (m, 2H), 2.91-2.80 (m, 1H), 2.38 (s, 1H), 2.19 (s, 3H), 2.11-2.00 (m, 1H), 1.70-1.50 (m, 4H), 1.50-1.20 (m, 4H). HRMS calculated for: $[C_{18}H_{22}F_4N_2O_3+H]^+$ 375.1690; found: 375.1679 (deviation 3.1 ppm).

25

Example [32] - Methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate

30

Step 1

The procedure used in Example [14], Step 1 was adapted such that 0.6 g of 1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropan-1-amine [9.1] was reacted to afford tert-butyl 2-((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamoyl)azetidine-1-carboxylate [32.1] as a colourless gum (0.6 g, 56%, LCMS $MH^+ = 419.2$).

35

Step 2

To a stirred solution of tert-butyl 2-((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamoyl)azetidine-1-carboxylate [32.1] (0.6 g, 1.43 mmol) in dry THF (3 mL) was added borane dimethyl sulphide complex (0.46 g, 5.74 mmol) dropwise under N₂ atm. The resultant reaction mixture was slowly warmed to rt, then heated to 60°C for 2h. The reaction mixture was quenched sat. aq. ammonium chloride solution and extracted with ethyl acetate (2 x 30 mL). The combined organic layer was washed with brine, dried over sodium sulphate, filtered and concentrated under reduced pressure to afford the crude product which was purified by flash column chromatography using ethyl acetate/hexane as eluent to afford tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)amino)methyl)azetidine-1-carboxylate [32.2] as a colourless gum (0.2 g, 34%, LCMS MH⁺ = 405.1).

Step 3

The procedure used in Example [1], Step 2 was adapted such that 0.2 g of tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)amino)methyl)azetidine-1-carboxylate [32.2] was reacted to afford tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)azetidine-1-carboxylate [32.3] as a colourless gum (0.2 g, 87%, LCMS MH⁺ = 463.2).

Step 4

The procedure used in Example [26], Step 2 was adapted such that 0.2 g of tert-butyl 2-(((1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)azetidine-1-carboxylate [32.3] was reacted to afford methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate [32] as a colourless gum (0.055 g, 35%, LCMS MH⁺ = 463.2). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.42 (t, *J* = 10.40 Hz, 1H), 7.30-7.05 (m, 2H), 3.91 (s, 1H), 3.57 (s, 3H), 3.50-3.40 (m, 2H), 3.05 (s, 1H), 2.08-1.80 (m, 3H), 1.50-1.28 (m, 2H), 1.23 (s, 2H). HRMS calculated for: [C₁₆H₁₈F₄N₂O₃+H]⁺ 363.1326; found: 363.1321 (deviation 1.4 ppm).

Example [33] - Methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(2-(hydroxyamino)-2-methylpropyl)carbamate

Step 1

To a stirred solution of methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [2] (0.25 g, 0.717 mmol) in DMF (5 mL) was added benzoyl peroxide (0.26 g, 1.076 mmol) and potassium phosphate (0.146 g, 1.076 mmol) at 0 °C. The resultant reaction mixture was slowly warmed to rt and stirred at rt for 12h. The reaction mixture was quenched with water and extracted with ethyl acetate (2*25 mL), the combined organic layer was dried over sodium sulphate and concentrated to afford crude product which was purified by column chromatography using 15% ethyl acetate in hexane as eluent to afford methyl (2-((benzoyloxy)amino)-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [33.1] as an off white solid (0.18 g, 54%).

Step 2

To a stirred solution of methyl (2-((benzoyloxy)amino)-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [33.1] (0.16 g, 0.341 mmol) in methanol (5 mL) was added hydrazine monohydrate (3 mL, 61.72 mmol) at 0 °C. The resultant reaction mixture was slowly warmed to rt and stirred at rt for 12h. The reaction mixture was quenched with water and extracted with ethyl acetate (2*25 mL), the combined organic layer was dried over sodium sulphate and concentrated to afford crude product which was purified by Prep HPLC to afford methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(2-(hydroxyamino)-2-methylpropyl)carbamate [33] as an off white solid (15.7 mg, 13%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.50 (bs, 1H), 7.42 (t, *J* = 8.80 Hz, 2H), 6.87 (s, 1H), 5.05 (bs, 1H), 3.56 (s, 3H), 3.40-3.35 (m, 2H), 1.28-1.26 (m, 2H), 1.22-1.24 (m, 2H), 0.85 (d, *J* = 10.40 Hz, 6H). HRMS calculated for: [C₁₆H₂₀F₄N₂O₃+H]⁺ 365.1483; found: 365.1478 (deviation 1.3 ppm).

Example [34] - Ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-yl)methyl)carbamate

Step 1

The procedure used in Example [1], Step 2 was adapted such that 0.25 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [23.2] was reacted to afford tert-butyl (S)-2-(((ethoxycarbonyl)(1-(4-fluoro-

3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [34.1] as colourless gum (0.16 g, 55%, LCMS MH^+ = 375.2 (boc-cleaved mass)).

Step 2

5 The procedure used in Example [2], Step 4 was adapted such that 0.14 g of tert-butyl (S)-2-(((ethoxycarbonyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl) cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate was reacted to afford ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [34] as white solid (0.1 g, 83%). MS ($M+1$)⁺ = 375.2. ¹H-NMR (400 MHz, DMSO-*d*₆): δ 9.40-9.05 (bs, 1H), 8.60-8.20 (bs, 1H), 7.65-7.52 (m, 1H), 7.49 (t, *J* = 9.20 Hz, 2H), 4.11 (q, *J* = 7.20 Hz, 2H), 3.70-3.65 (m, 1H), 3.65-3.52 (m, 2H), 3.30-3.18 (m, 1H), 3.15-3.00 (m, 1H), 2.05-1.60 (m, 3H), 1.65-1.40 (m, 3H), 1.40-1.10 (m, 5H). HRMS calculated for: [C₁₈H₂₂F₄N₂O₂+H]⁺ 375.1690; found: 375.1686 (deviation 1.2 ppm).

15 Example [35] - Ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl) carbamate

The procedure used in Example [2], Step 2 was adapted such that 0.1 g of ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [34] was reacted to afford ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl) carbamate [35] as colourless gum (0.075 g, 80%, LCMS MH^+ = 389.1). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.52 (d, *J* = 5.20 Hz, 2H), 7.38 (t, *J* = 8.80 Hz, 1H), 4.08 (q, *J* = 7.20 Hz, 2H), 3.30 (d, *J* = 6.00 Hz, 2H), 2.90 (q, *J* = 4.80 Hz, 1H), 2.22 (s, 3H), 2.12 (q, *J* = 8.40 Hz, 2H), 1.70-1.50 (m, 4H), 1.50-1.20 (m, 4H), 1.16 (t, *J* = 6.80 Hz, 3H). HRMS calculated for: [C₁₉H₂₄F₄N₂O₂+H]⁺ 389.1847; found: 389.1843 (deviation 1.0 ppm).

25 Example [36] - (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methanesulfonamide

30 *Step 1*

The procedure used in Example [1], Step 2 was adapted such that 0.8 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [23.2] and 0.69 g of methanesulphonyl chloride were reacted to afford the product tert-butyl (S)-2-((N-(1-(4-fluoro-3-

(trifluoromethyl)phenyl)cyclopropyl)methylsulfonamido)methyl)pyrrolidine-1-carboxylate [36.1] as a colourless gum (0.4 g, 42%).

Step 2

- 5 The procedure used in Example [22], Step 4 was adapted such that 0.4 g of tert-butyl (S)-2-((N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methylsulfonamido)methyl)pyrrolidine-1-carboxylate [36.1] was reacted to afford (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methanesulfonamide [36] as an off-white solid (0.28 g, 80%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 9.50-9.25 (s, 1H), 8.95-8.65 (s, 1H), 7.84-7.78 (m, 2H), 7.46 (t, *J* = 8.80 Hz, 1H), 3.80-3.55 (m, 3H), 3.35-3.00 (m, 2H), 2.92 (s, 3H), 2.10-1.65 (m, 3H), 1.65-1.50 (m, 3H), 1.35-1.26 (m, 2H). HRMS calculated for: [C₁₆H₂₀F₄N₂O₂+H]⁺ 381.1254; found: 381.1250 (deviation 1.3 ppm).
- 10

15 Example [37] - (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-((1-methylpyrrolidin-2-yl)methyl)methanesulfonamide

- The procedure used in Example [2], Step 2 was adapted such that 0.12 g of (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl)methanesulfonamide [36] was reacted to afford (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-((1-methylpyrrolidin-2-yl)methyl)methanesulfonamide [37] as a colourless gum (0.08 g, 67%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.69 (dd, *J* = 2.12, 6.68 Hz, 1H), 7.61-7.60 (m, 1H), 7.47 (t, *J* = 9.04 Hz, 1H), 3.25 (dd, *J* = 40.00, Hz, 1H), 3.13-3.04 (m, 1H), 2.95-2.90 (m, 3H), 2.90-2.85 (m, 1H), 2.40-2.32 (m, 1H), 2.30-2.20 (m, 3H), 2.15-2.00 (m, 1H), 1.70-1.50 (m, 6H), 1.45-1.20 (m, 2H). HRMS calculated for: [C₁₇H₂₂F₄N₂O₂+H]⁺ 395.1411; found: 395.1406 (deviation 1.3 ppm).
- 20
- 25

30 Example [38] - Ethyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate

Step 1

- The procedure used in Example [1], Step 2 was adapted such that 0.25 g of tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl) pyrrolidine-1-carboxylate [22.2] and 0.07 g of ethylchloroformate were reacted to afford tert-butyl
- 35

(R)-2-(((ethoxycarbonyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [38.1] as a colourless gum (0.14 g, 48%, LCMS MH^+ = 375.1 (Boc-cleaved mass)).

5 *Step 2*

The procedure used in Example [22], Step 4 was adapted such that 0.13 g of tert-butyl (R)-2-(((ethoxycarbonyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate was reacted to afford ethyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [38]

10 (HCl salt) as a white solid (0.115 g, 99%, LCMS MH^+ = 375.1). 1H -NMR (400 MHz, DMSO- d_6): δ 9.30-9.10 (bs, 1H), 8.60-8.20 (bs, 1H), 7.58-7.46 (m, 3H), 4.13-4.08 (m, 2H), 3.80-3.50 (m, 3H), 3.30-3.18 (m, 1H), 3.15-3.00 (m, 1H), 2.10-1.70 (m, 3H), 1.59-1.49 (m, 3H), 1.35-1.07 (m, 5H). HRMS calculated for: $[C_{18}H_{22}F_4N_2O_2+H]^+$ 375.1690; found: 375.1688 (deviation 0.5 ppm).

15

Example [39] - Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate

Step 1

20 To an ice cooled solution of (S)-4-(tert-butoxy carbonyl)-morpholine-3-carboxylic acid (0.37 g, 1.642 mmol) in dichloromethane (10 mL) was added triethylamine (0.76 mL, 5.474 mmol) and followed by propylphosphonic anhydride (2.61 g, 4.106 mmol) solution under N_2 atm. The resultant reaction mixture was stirred at 0 °C for 20 min, then 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] (0.3 g, 1.368

25 mmol) was added and the reaction stirred at rt for 5h. The reaction mixture was quenched with water and extracted with ethyl acetate (2*80 mL). The combined organic layer was washed with 10% sodium bicarbonate solution, brine, dried over sodium sulphate, filtered and concentrated under reduced pressure to afford tert-butyl (S)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamoyl)morpholine-4-carboxylate

30 [39.1] as a yellow liquid (0.55 g, 93%, LCMS MH^+ = 433.1).

Step 2

To a cooled solution of tert-butyl (S)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamoyl)morpholine-4-carboxylate [39.1] (0.55 g, 1.271 mmol) in dry THF (5 mL) was added borane dimethyl sulphide complex (9.6 mL,

35

2.543 mmol) dropwise under N₂ atm. The resultant reaction mixture was slowly warmed to rt and stirred at rt for 16h. The reaction mixture was quenched with methanol and the refluxed for 1h and then concentrated under reduced pressure. The obtained residue was diluted with water and extracted with dichloromethane (2*100 mL), the combined organic layer was washed with brine, dried over sodium sulphate, filtered and concentrated under reduced pressure to afford tert-butyl (3R)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)morpholine-4-carboxylate [39.2] as a colourless gum (0.28 g, 52%, LCMS MH⁺ = 419.2).

Step 3

The procedure used in Example [1], Step 2 was adapted such that 0.25 g of tert-butyl (3R)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)morpholine-4-carboxylate [39.2] was reacted to afford the product tert-butyl (R)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)morpholine-4-carboxylate as a Colourless gum (0.2 g, 71%, LCMS MH⁺ = 477.2).

Step 4

The procedure used in Example [1], Step 2 was adapted such that 0.2 g of tert-butyl (R)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)morpholine-4-carboxylate [39.3] was reacted to afford methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate [39] as a white solid (0.15 g, 88%, LCMS MH⁺ = 377.2). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 9.11 (bs, 1H), 7.70-7.30 (m, 3H), 3.85 (d, *J* = 11.20 Hz, 2H), 3.75-3.68 (m, 1H), 3.63 (bs, 3H), 3.60-3.40 (m, 4H), 3.24 (d, *J* = 32.00 Hz, 1H), 3.10-3.00 (m, 1H), 1.53 (bs, 2H), 1.50-1.20 (m, 2H). HRMS calculated for: [C₁₇H₂₀F₄N₂O₃+H]⁺ 377.1483; found: 377.1475 (deviation 2.1 ppm).

Example [40] - Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate

The procedure used in Example [2], Step 2 was adapted such that 0.1 g of methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl)carbamate [39] was reacted to afford methyl (R)-(1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate [40] as a colourless gum (0.08 g, 85%, LCMS

MH⁺ = 391.2). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.54-7.44 (m, 3H), 3.61-3.58 (m, 3H), 3.47-3.39 (m, 2H), 3.34-3.33 (m, 2H), 3.30-3.26 (m, 1H), 3.20-3.10 (m, 1H), 2.67-2.61 (m, 1H), 2.27-2.07 (m, 5H), 1.46-1.25 (m, 4H). HRMS calculated for: [C₁₈H₂₂F₄N₂O₃+H]⁺ 391.1639; found: 391.1633 (deviation 1.7 ppm).

5

Example [41] - (R)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methane sulfonamide

Step 1

10 The procedure used in Example [1], Step 2 was adapted such that 0.25 g of tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl) pyrrolidine-1-carboxylate and 0.11 g of methanesulphonyl chloride were reacted to afford tert-butyl (R)-2-((N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methylsulfonamido)methyl)pyrrolidine-1-carboxylate
15 [41.1] as a colourless gum (0.16 g, 34%, LCMS MH⁺ = 381.1 (Boc-cleaved mass)).

Step 2

The procedure used in Example [22], Step 4 was adapted such that 0.15 g of tert-butyl (R)-2-((N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methylsulfonamido)methyl)pyrrolidine-1-carboxylate
20 [41.1] was reacted to afford (R)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methane sulfonamide [41] (HCl salt) as an off white solid (0.11 g, 84%, LCMS MH⁺ = 381.1). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 9.27 (bs, 1H), 8.67 (bs, 1H), 7.80-7.75 (m, 2H), 7.52 (t, *J* = 9.60 Hz, 2H), 3.65-3.56 (m, 3H), 3.22-3.12 (m, 2H),
25 2.90 (s, 3H), 2.00-1.75 (m, 3H), 1.70-1.50 (m, 3H), 1.30 (bs, 2H). HRMS calculated for: [C₁₆H₂₀F₄N₂O₂S+H]⁺ 381.1254; found: 381.1251 (deviation 0.7 ppm).

Example [42] - Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate

30

Step 1

The procedure used in Example [2], Step 2 was adapted such that 0.8 g of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] was reacted with tert-butyl (3R)-3-formylmorpholine-4-carboxylate to afford tert-butyl (S)-3-(((1-(4-fluoro-3-

(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)morpholine-4-carboxylate [42.1] as a colourless gum (0.9 g, 59%, LCMS $MH^+ = 419.2$).

Step 2

5 The procedure used in Example [1], Step 2 was adapted such that 0.3 g of tert-butyl (S)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)morpholine-4-carboxylate [42.1] was reacted to afford tert-butyl(S)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)morpholine-4-carboxylate [42.2] as a colourless gum (0.17 g, 50%, LCMS $MH^+ = 377.2$).

10

Step 3

The procedure used in Example [2], Step 4 was adapted such that 0.17 g of tert-butyl (S)-3-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)morpholine-4-carboxylate [42.2] was reacted to afford 15 methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate [42] as a white solid (0.135 g, 92%; LCMS $MH^+ = 377.2$). 1H -NMR (400 MHz, DMSO- d_6): δ 9.15 (bs, 2H), 7.50-7.37 (m, 3H), 3.86 (d, $J = 11.60$ Hz, 2H), 3.75-3.40 (m, 8H), 3.25 (d, $J = 40.00$ Hz, 1H), 3.12-2.95 (m, 1H), 1.57 (bs, 2H), 1.31-1.23 (m, 2H). HRMS calculated for: $[C_{17}H_{20}F_4N_2O_3+H]^+$ 377.1483; found: 377.1476 20 (deviation 1.9 ppm).

Example [43] - Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate

25 The procedure used in Example [2], Step 2 was adapted such that 0.1 g of methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl)carbamate [42] was reacted to afford methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate [43] as a white solid (0.075 g, 80%, LCMS $MH^+ = 391.2$). 1H -NMR (400 MHz, DMSO- d_6): δ 7.52-7.42 (m, 3H), 3.70-3.55 (m, 5H), 30 3.53-3.37 (m, 2H), 3.35-3.22 (m, 1H), 3.00-3.02 (m, 1H), 2.68-2.58 (m, 1H), 2.35-2.18 (m, 4H), 2.15-2.05 (m, 1H), 1.55-1.33 (m, 4H). HRMS calculated for: $[C_{18}H_{22}F_4N_2O_3+H]^+$ 391.1639; found: 391.1633 (deviation 1.7 ppm).

Example [44] - Isopropyl (S)-(1-(4-fluoro-3-
35 (trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate

Step 1

The procedure used in Example [45], Step 1 was adapted such that 0.1 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [23.2] and 0.67 g of isopropyl chloroformate were reacted to afford tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(isopropoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate [44.1] as colourless liquid (0.1 g, 83%).

10 *Step 2*

The procedure used in Example [26], Step 2 was adapted such that 0.1 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(isopropoxycarbonyl)amino)methyl)pyrrolidine-1-carboxylate [44.1] was reacted to afford isopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [44] as a pale yellow liquid (0.07 g, 88%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 8.45 (bs, 1H), 7.57 (d, *J* = 2.64 Hz, 1H), 7.48 (t, *J* = 8.24 Hz, 2H), 4.83 (t, *J* = 6.24 Hz, 1H), 3.70-3.48 (m, 3H), 3.28-3.18 (m, 1H), 3.12-3.02 (m, 1H), 1.99 (bs, 1H), 1.92-1.74 (m, 2H), 1.60-1.38 (m, 3H), 1.38-1.24 (m, 2H), 1.22-1.05 (m, 6H). HRMS calculated for: [C₁₉H₂₄F₄N₂O₂+H]⁺ 389.1847; found: 389.1838 (deviation 2.3 ppm).

Example [45] - Cyclopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate

25 *Step 1*

To a stirred solution of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [23.2] (0.15 g, 0.37 mmol) in acetonitrile (4 mL) was added cesium carbonate (0.36 g, 1.12 mmol) and cyclopropane chloroformate (0.09 g, 0.75 mmol). The reaction was stirred at room temperature for 3h, filtered and concentrated to afford the crude product, which was purified by flash column chromatography using ethyl acetate in pet ether as eluent to afford tert-butyl (S)-2-(((cyclopropoxycarbonyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [45.1] as a colourless liquid (0.16 g, 89%, LCMS MH⁺ = 387.1 (boc cleaved mass)).

35

Step 2

The procedure used in Example [26], Step 2 was adapted such that 0.16 g of tert-butyl (S)-2-(((cyclopropoxycarbonyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [45.1] was reacted to afford cyclopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [45] as a yellow liquid (0.1 g, 83%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.43-7.41 (m, 3H), 3.50-3.00 (m, 3H), 2.90-2.70 (m, 2H), 1.80-1.50 (m, 4H), 1.40-1.10 (m, 5H), 0.80-0.30 (m, 4H). HRMS calculated for: [C₁₉H₂₂F₄N₂O₂+H]⁺ 387.1690; found: 387.1688 (deviation 0.5 ppm).

Example [46] - Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methyl pyrrolidin-2-yl)methyl)carbamate

Step 1

The procedure used in Example [2], Step 2 was adapted such that 0.7 g of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] and tert-butyl (2R)-2-formyl-2-methylpyrrolidine-1-carboxylate were reacted to afford tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)-2-methyl pyrrolidine-1-carboxylate [46.1] as a colourless liquid (0.68 g, 51%, LCMS MH⁺ = 417.2).

Step 2

The procedure used in Example [1], Step 2 was adapted such that 0.5 g of tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)-2-methylpyrrolidine-1-carboxylate [46.1] was reacted to afford tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)-2-methylpyrrolidine-1-carboxylate [46.2] as a brown liquid (0.13 g, 22%, LCMS MH⁺ = 375.1(boc-cleaved mass)).

Step 3

The procedure used in Example [22], Step 4 was adapted such that 0.15 g of tert-butyl (R)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)-2-methylpyrrolidine-1-carboxylate [46.2] was reacted to afford methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methyl pyrrolidin-2-yl)methyl)carbamate [46] as

colourless gum (0.125 g, 96%, LCMS $MH^+ = 375.4$). 1H -NMR (400 MHz, DMSO- d_6): δ 8.64 (bs, 1H), 8.40 (bs, 1H), 7.59-7.48 (m, 3H), 3.84 (bs, 1H), 3.68 (bs, 3H), 3.60 (bs, 1H), 3.24-3.22 (m, 3H), 2.00-1.85 (m, 2H), 1.77 (bs, 2H), 1.95 (bs, 3H), 1.32-1.20 (m, 1H), 1.10 (bs, 2H). HRMS calculated for: $[C_{18}H_{22}F_4N_2O_2+H]^+$ 375.1690; found: 375.1684 (deviation 1.8 ppm).

Example [47] - N-((1-amino cyclopropyl)methyl)-1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine

The procedure used in Example [2], Step 4 was adapted such that 0.1 g of tert-butyl (1-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)cyclopropyl)carbamate [3.1] was reacted to afford N-((1-amino cyclopropyl)methyl)-1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [47] as a white solid (0.08 g, 96%, LCMS $MH^+ = 288.1$). 1H -NMR (400 MHz, DMSO- d_6): δ 10.60-10.20 (m, 2H), 8.90-8.65 (m, 2H), 8.20-7.90 (m, 2H), 7.65-7.55 (m, 1H), 3.30-3.10 (m, 2H), 1.75-1.55 (m, 2H), 1.35-1.15 (m, 2H), 1.14-0.80 (m, 4H). HRMS calculated for: $[C_{14}H_{16}F_4N_2+H]^+$ 289.1322; found: 289.1310 (deviation 4.2 ppm).

Example [48] - Cyclopropylmethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate

Step 1

The procedure used in Example [45], Step 1 was adapted such that 0.15 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [23.2] and 0.125 g of cyclopropylmethyl chloroformate were reacted to afford tert-butyl (S)-2-(((cyclopropylmethoxy)carbonyl) (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [48.1] (0.25 g, 80%).

Step 2

The procedure used in Example [26], Step 2 was adapted such that 0.12 g of tert-butyl (S)-2-(((cyclopropylmethoxy)carbonyl) (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)pyrrolidine-1-carboxylate [48.1] was reacted to afford cyclopropylmethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate [48] as a yellow

liquid (0.085 g, 88%). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.52-7.43 (m, 3H), 3.87 (d, *J* = 7.00 Hz, 2H), 3.55-3.45 (m, 3H), 3.00-2.53 (m, 2H), 4.80-1.55 (m, 4H), 1.45-1.20 (m, 4H), 1.00 (s, 1H), 0.45 (s, 2H), 0.20 (s, 2H). HRMS calculated for: [C₂₀H₂₄F₄N₂O₂+H]⁺ 401.1847; found: 401.1845 (deviation 0.3 ppm).

5

Example [49] - N-((1-aminocyclopropyl)methyl)-N-(1-(4-fluoro-3-(trifluoromethyl)-phenyl)cyclopropyl)methanesulfonamide

Step 1

10 The procedure used in Example [1], Step 2 was adapted such that 0.3 g of tert-butyl (1-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)cyclopropyl)carbamate [3.1] and 0.13 g of methanesulfonyl chloride were reacted to afford tert-butyl (1-((N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methylsulfonamido)methyl)cyclopropyl)carbamate
15 [49.1] as a white solid (0.3 g, 83%, LCMS MH⁺ = 367.2).

Step 2

The procedure used in Example [2], Step 4 was adapted such that 0.1 g of tert-butyl (1-((N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methylsulfonamido)methyl)cyclopropyl)carbamate [49.1] was
20 reacted to afford N-((1-aminocyclopropyl)methyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methanesulfonamide [49] as a white solid (0.08 g, 93%, LCMS MH⁺ = 367.2). ¹H-NMR (400 MHz, DMSO-*d*₆): δ 8.41 (bs, 3H), 7.76 (s, 1H), 7.66 (d, *J* = 4.96 Hz, 1H), 7.49 (t, *J* = 8.80 Hz, 1H), 3.53 (s, 2H), 3.01 (s, 3H), 1.67
25 (bs, 2H), 1.33 (s, 2H), 0.92 (d, *J* = 44.32 Hz, 4H). HRMS calculated for: [C₁₅H₁₈F₄N₂O₂S+H]⁺ 367.1098; found: 367.1089 (deviation 2.5 ppm).

Example [50] - Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methylpyrrolidin-2-yl)methyl)carbamate

30

Step 1

The procedure used in Example [2], Step 2 was adapted such that 0.5 g of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] and tert-butyl (2S)-2-formyl-2-methylpyrrolidine-1-carboxylate were reacted to afford tert-butyl (S)-2-(((1-(4-fluoro-3-

(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)-2-methylpyrrolidine-1-carboxylate [50.1] as a colourless liquid (0.65 g, 68%, LCMS $MH^+ = 417.2$).

Step 2

5 The procedure used in Example [1], Step 2 was adapted such that 0.4 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)-2-methylpyrrolidine-1-carboxylate [50.1] was reacted to afford tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)-2-methylpyrrolidine-1-carboxylate [50.2] as a brown liquid (0.32 g, 70%, LCMS $MH^+ =$
10 375.1(boc-cleaved mass)).

Step 3

The procedure used in Example [22], Step 4 was adapted such that 0.3 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)-2-methylpyrrolidine-1-carboxylate [50.2] was reacted to afford methyl (S)-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methylpyrrolidin-2-yl)methyl)carbamate [50] as a white solid (0.25 g, 96%, LCMS $MH^+ = 375.0$). 1H -NMR (400 MHz, DMSO- d_6): δ 8.70 (bs, 1H), 8.46 (bs, 1H), 7.52 (q, $J = 9.68$ Hz, 3H), 3.81 (t, $J = 45.48$ Hz, 4H), 3.22 (d, $J = 6.84$ Hz, 2H), 1.95 (d, $J = 8.36$ Hz, 2H), 1.92 (d, $J = 5.52$ Hz, 2H), 1.44 (s, 3H), 1.27 (bs, 2H), 1.10 (s, 3H). HRMS calculated for: $[C_{18}H_{20}F_4N_2O_2+H]^+$ 375.1690; found: 375.1688 (deviation 0.6 ppm).

Example [51] - (1S, 2S)-N1-(1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)-
25 cyclopentane-1,2-diamine

Step 1

The procedure used in Example [2], Step 2 was adapted such that 0.8 g of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] and tert-butyl N-[(1S)-2-oxocyclopentyl]carbamate were reacted to afford the diastereomers tert-butyl ((1S,2S)-2-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)cyclopentyl)carbamate [51.1] as yellow gum (0.43 g, 58%, LCMS $MH^+ = 403.2$) and tert-butyl ((1S,2R)-2-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)cyclopentyl)carbamate [51.2] as a white solid (0.31 g, 42%, LCMS $MH^+ = 403.2$).

35

Step 2

The procedure used in Example [22], Step 4 was adapted such that 0.04 g of tert-butyl ((1S,2S)-2-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)cyclopentyl)carbamate [51.1] was reacted to afford (1S, 2S)-N1-(1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)cyclopentane-1,2-diamine [51] as a brown gum (0.012 g, 35%, LCMS MH^+ = 303.2). 1H -NMR (400 MHz, DMSO- d_6): δ 7.93 (t, J = 22.80 Hz, 2H), 7.51 (t, J = 9.20 Hz, 1H), 3.54 (d, J = 1.60 Hz, 2H), 3.40 (d, J = 6.80 Hz, 2H), 3.15 (t, J = 1.20 Hz, 1H), 2.00-1.88 (m, 2H), 1.63-1.55 (m, 3H), 1.63-1.55 (m, 5H). HRMS calculated for: $[C_{15}H_{18}F_4N_2+H]^+$ 303.1479; found: 303.1472 (deviation 2.3 ppm).

Example [52] - (1R,2S)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-cyclopentane-1,2-diamine

Step 1

The procedure used in Example [22], Step 4 was adapted such that 0.15 g of tert-butyl ((1S,2R)-2-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)cyclopentyl)carbamate [51.2] was reacted to afford (1R,2S)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)cyclopentane-1,2-diamine [52] as an off-white solid (0.112 g, 88%, LCMS MH^+ = 303.2). 1H -NMR (400 MHz, DMSO- d_6): δ 10.66 (bs, 1H), 8.80-8.30 (m, 1H), 8.30-7.60 (m, 3H), 7.70-7.45 (m, 1H), 7.39-7.14 (m, 1H), 4.50-3.80 (m, 2H), 3.80-3.55 (m, 1H), 2.00-1.50 (m, 6H), 1.50-1.05 (m, 5H). HRMS calculated for: $[C_{15}H_{18}F_4N_2+H]^+$ 303.1479; found: 303.1474 (deviation 1.6 ppm).

Example [53] - Methyl (S)-(azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)-cyclopropyl)carbamate

Step 1

The procedure used in Example [39], Step 1 was adapted such that 1.3 g of 1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine [1.1] was reacted to afford tert-butyl (S)-2-((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamoyl)azetidine-1-carboxylate [53.1] as a brown liquid (2.2 g, 92%, LCMS MH^+ = 303.1(boc-cleaved mass)).

Step 2

The procedure used in Example [32], Step 2 was adapted such that 2.2 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamoyl)azetidine-1-carboxylate [53.1] was reacted to afford tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)azetidine-1-carboxylate [53.2] as colourless liquid (1.2 g, 57%, LCMS $MH^+ = 389.2$).

Step 3

The procedure used in Example [1], Step 2 was adapted such that 0.25 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)methyl)azetidine-1-carboxylate [53.2] was reacted to afford tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)azetidine-1-carboxylate [53.3] as a brown gum (0.26 g, 92%, LCMS $MH^+ = 447.2$).

Step 4

The procedure used in Example [26], Step 2 was adapted such that 0.5 g of tert-butyl (S)-2-(((1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(methoxycarbonyl)amino)methyl)azetidine [53.3] was reacted to afford methyl (S)-(azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate [53] as a brown liquid (0.04 g, 11%, LCMS $MH^+ = 347.0$). 1H -NMR (400 MHz, DMSO- d_6): δ 7.44-7.28 (m, 3H), 3.91 (s, 1H), 3.58-3.33 (m, 5H), 3.05 (s, 2H), 2.00 (t, $J = 43.60$ Hz, 1H), 1.76 (s, 2H), 1.39 (s, 2H), 1.24 (s, 2H). HRMS calculated for: $[C_{16}H_{18}F_4N_2O_2+H]^+$ 347.1377; found: 347.1369 (deviation 2.4 ppm).

Example [54] – Solubility of compounds

The aim of this experiment was to determine solubility of test compounds in 50 mM Phosphate buffer by using HPLC.

Method

Incubation time	16 hr at ~ 25°C
Buffer pH	50 mM potassium phosphate buffer, pH 7.40
Test compound Incubation concentration	1600 μ M

Replicates	n=2
Analysis	HPLC
Standard compounds	Caffeine [Solubility (1400-1900 μ M)], Diethylstilbestrol [Solubility (0 μ M)] and Tamoxifen [Solubility (<20 μ M)]
Deliverables	Solubility of test compound mg/mL

Preparation of Phosphate buffer (pH 7.4):

2.79 g of K_2HPO_4 and 0.54 g of KH_2PO_4 was dissolved in 390 mL of milliQ water. pH was adjusted to 7.4 using 1N HCl/1N NaOH and final volume was made up to 400 mL with milliQ water.

Preparation and dilution of test compound:

80 mM master stock of test compounds was prepared in 100% DMSO. In case of compounds not soluble / less quantity submission, 40/20/10Mm stocks were prepared and used for experiment.

Assay procedure:

- 245 μ L of 50 mM phosphate buffer then 5 μ L each of test compound/standards (80mM) in their respective positions was added to the 1.1 mL 96 well plate.
- DMSO Controls was prepared by taking 245 μ L of 100% DMSO then 5 μ L each of test compound in their respective positions and added to the 1.1 mL 96 well plate
- Plate was incubated with mixing at 1600 RPM for 16 hours at room temperature (~23 °C).
- After incubation, samples were filtered using Millipore plates.
- Filtrates were analysed by HPLC-UV.

Solubility calculation:

Solubility is calculated using the following formula:

$$\text{Solubility } (\mu\text{M}) = \frac{\text{(Sample area in Buffer)}}{\text{(Sample area in DMSO)}} \times 1600$$

5

Results

Compound	Measured Solubility in pH 7.4 phosphate buffer		Compound	Measured Solubility in pH 7.4 phosphate buffer	
	(mg/ml)	(μM) (rounded)		(mg/ml)	(μM) (rounded)
[1]	0.5409	1600	[27]	0.5918	1600
[2]	0.5555	1600	[28]	0.6738	1600
[3]	0.5599	1600	[29]	0.6773	1600
[4]	0.0279	100	[30]	0.5984	1700
[6]	0.6792	1500	[31]	0.5575	1500
[8]	0.5109	1500	[32]	0.5578	1500
[9]	0.582	1200	[33]	0.1375	400
[10]	0.4809	1400	[34]	0.594	1400
[11]	0.5778	1600	[35]	0.5679	1500
[12]	0.3755	800	[36]	0.3218	800
[13]	0.6337	1500	[37]	0.621	1600
[14]	0.7554	1600	[38]	0.6157	1500
[15]	0.526	1400	[42]	0.6118	1500
[16]	0.5974	1600	[43]	0.5996	1500
[17]	0.5636	1500	[44]	0.6148	1600
[19]	0.607	1500	[45]	0.6009	1600
[20]	0.5836	1500	[46]	0.585	1400
[21]	0.5646	1500	[47]	0.4635	1400
[22]	0.6666	1700	[50]	0.5747	1400
[23]	0.653	1600	[51]	0.2392	700
[24]	0.5958	1500	[52]	0.4797	1400
[25]	0.4824	1700	[53]	0.5907	1700
[26]	0.6868	1500	NS6180	0.0009	2.8

Conclusion

It is demonstrated that the tested compounds have solubility in pH 7.4 phosphate buffer of 400 to 1700 μM , whereas NS6180 has a solubility of 2.8 μM .

5 Example [55] – Inhibition of $K_{\text{Ca}3.1}$

Erythrocyte $K_{\text{Ca}3.1}$ assay

Human blood was drawn from healthy human volunteers in a standard heparinized blood sampling vial (Vacutainer, Li/heparin, BD Bioscience, Plymouth, UK). The erythrocytes were packed by centrifugation, and the plasma and buffy coat were removed by aspiration. Erythrocytes were washed three times in the experimental salt solution and then stored at 0°C until use. Blood samples from NMRI mice or from Wistar rats were treated similarly. The methodological principle is outlined in Macey *et al.* (1978) and further described in Strøbæk *et al.* (2013). Activation of the erythrocyte $K_{\text{Ca}3.1}$ channels were obtained by addition of the Ca^{2+} ionophore A23187, which causes synchronized hyperpolarization, which is reported as a CCCP-mediated shift in the unbuffered extracellular pH of the erythrocyte suspension. Standard procedure: 3 mL unbuffered experimental salt solution (in mM: 2 KCl, 154 NaCl, 0.05 CaCl_2) was heated to 37°C with stirring. Packed erythrocytes were added (50 μL , final cytocrit 1.5%), and the extracellular pH (pH_o) followed with a glass/calomel (pHG200-8/REF200, Radiometer, Denmark) electrode pair. CCCP (3 μL , final concentration 20 μM) was added followed by varying concentrations of test compounds (DMSO concentration constant). After pH stabilization at ~ 7.2 , A23187 (3 μL , final concentration 0.33 μM) was added to initiate the experiment. After the peak hyperpolarization was attained, the intracellular pH (pH_i constant during the experiment) was found by haemolysing the erythrocytes via addition of 100 μL of Triton-X100.

The erythrocyte membrane potential, V_m , was calculated according to:

$$V_m = -61.5 \text{ mV} \times (\text{pH}_o - \text{pH}_i)$$

30 and the fractional remaining Ca^{2+} -activated K^+ -conductance at the concentration C of blocker, $fG_K(C)$, was calculated from

$$fG_K(C) = \frac{(V_m(0) - E_K) * (E_{\text{Cl}} - V_m(C))}{(E_{\text{Cl}} - V_m(0)) * (V_m(C) - E_K)}$$

where the K^+ equilibrium potential $E_K = -107$ mV, the Cl^- equilibrium potential $E_{Cl} = -12$ mV and the $V_m(0)$ and $V_m(C)$ are the peak hyperpolarizations in the control and in the presence of a concentration of C of blocker respectively.

- 5 IC_{50} values for compounds were calculated from a plot of $fG_K(C)$ versus C by a fit to the Hill equation, using a custom program written in the IGOR-Pro software (WaveMetrics, Lake Oswego, OR, USA). All IC_{50} -values are reported in μM .

Results

Compound	RBC K (in vitro) Human IC_{50} (μM)	Compound	RBC K (in vitro) Human IC_{50} (μM)
[1]	0.33	[28]	0.087
[2]	0.086	[29]	0.33
[3]	0.16	[30]	0.17
[4]	0.071	[31]	0.081
[5]	0.31	[32]	0.27
[6]	0.19	[33]	0.12
[7]	0.48	[34]	0.052
[8]	0.014	[35]	0.11
[9]	0.35	[36]	0.32
[10]	0.024	[37]	0.14
[11]	0.3	[38]	0.12
[12]	0.4	[39]	0.42
[13]	0.44	[40]	0.27
[14]	0.25	[41]	0.38
[15]	0.14	[42]	0.34
[16]	0.4	[43]	0.29
[17]	0.4	[44]	0.2
[18]	0.018	[45]	0.18
[19]	0.047	[46]	0.1

[20]	0.092	[47]	0.22
[21]	0.075	[48]	0.23
[22]	0.21	[49]	0.39
[23]	0.12	[50]	0.15
[24]	0.31	[51]	0.41
[25]	0.18	[52]	0.093
[26]	0.21	[53]	0.36
[27]	0.3		

Conclusion

It is demonstrated that all the compounds inhibit K_{Ca}3.1.

5

10 References

Macey *et al.*, Biochim. Biophys. Acta **1978**, 22, 512(2), 284-95

Strøbæk *et al.*, Br. J. Pharmacol. 2013, 168(2), 432-444


WO 2014/001363 [Clevexel Pharma; Aniona ApS; Saniona ApS]

WO 2013/191984 [Boehringer Ingelheim]

15 WO 2014/067861 [Hoffmann La Roche]

Claims

1. A compound of formula (XVI):



Formula (XVI)

wherein

R^{14} is selected from the group consisting of $-C(O)-C_{1-8}$ alkyl; $-C(O)-O-C_{1-8}$ alkyl; $-C_{2-8}$ alkyl; $-H$ and $-S(O)_2-C_{1-8}$ alkyl;

R^3 is H , C_{1-5} alkyl, or a bond;

R^4 is H , C_{1-5} alkyl, or a bond;

R^5 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by $-O-$;

R^6 is H , a bond, or C_{1-8} alkyl, wherein one methylene group optionally is replaced by $-O-$;

R^7 is H , a bond, $-OH$, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by $-O-$ and/or substituted with $=O$;

R^8 is H , a bond, $-OH$, or C_{1-8} alkyl, wherein one or more methylene group optionally and individually is replaced by $-O-$ and/or substituted with $=O$;


Any one of R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 optionally is linked together to form a ring;

A is a phenyl or a pyridinyl, wherein the phenyl or pyridinyl is optionally substituted with one or more substituents R^{13} individually selected from the group consisting of halogen, $-CX_3$, $-OCX_3$, $-CHX_2$, $-OCHX_2$, $-CH_2X$, $-OCH_2X$, $-CH_2CX_3$, OCH_2CX_3 , $-C_{1-8}$ alkyl, $-OC_{1-8}$ alkyl, $-C_{3-7}$ cycloalkyl, $-OC_{3-7}$ cycloalkyl, $-CN$, NO_2 , $-SO_2CH_3$, and $-SF_5$; and

X is halogen;

or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1, wherein A is a moiety of formula (IX):




Formula (IX)

wherein

- 5 R^9 is $-C(H)-$, $-N-$, or $-C(R^{13})-$;
 R^{13} is individually selected from the group consisting of halogen, $-CX_3$, $-OCX_3$, $-CHX_2$, $-OCHX_2$, $-CH_2X$, $-OCH_2X$, $-CH_2CX_3$, OCH_2CX_3 , $-C_{1-8}$ alkyl, $-OC_{1-8}$ alkyl, $-C_{3-7}$ cycloalkyl, $-OC_{3-7}$ cycloalkyl, $-CN$, NO_2 , $-SO_2CH_3$, and $-SF_5$;
 n is an integer of 0 to 4; and
 10 X is halogen.

3. The compound according to claim 1, wherein A is a moiety of formula (X):



Formula (X)


- 15 wherein
 R^9 is $-C(H)-$, $-N-$, or $-C(R^{13})-$;
 R^{10} , R^{11} , R^{12} , and R^{13} are individually selected from the group consisting of H, halogen, $-CX_3$, $-OCX_3$, $-CHX_2$, $-OCHX_2$, $-CH_2X$, $-OCH_2X$, $-CH_2CX_3$, OCH_2CX_3 , $-C_{1-8}$ alkyl, $-OC_{1-8}$ alkyl, $-C_{3-7}$ cycloalkyl, $-OC_{3-7}$ cycloalkyl, $-CN$, NO_2 , $-SO_2CH_3$,
 20 and $-SF_5$; and
 X is halogen.

4. The compound according to claim 3, wherein

- R^9 is $-C(H)-$ or $-N-$;
 25 R^{10} is H or halogen;
 R^{11} is H or halogen;


R^{12} is $-CX_3$, $-OCX_3$, H or halogen; and
X is halogen.

5. The compound according to any one of the preceding claims, wherein R^3 and R^4 are -H, R^5 and R^6 are methyl, and R^7 and R^8 are -H.
6. The compound according to any one of claims 1 to 4, wherein the compound is of formula (XVIII):




Formula (XVIII).

7. The compound according to any one of the claims 1 to 4, wherein the compound is of formula (XIX):




Formula (XIX).

8. The compound according to any one of claims 1 to 4, wherein the compound is of formula (XXI):




Formula (XXI).

9. The compound according to any one of claims 1 to 4 claim 1, wherein the
5 compound is of formula (XXII):



Formula (XXII).

10. The compound according to any one of claims 1 to 4, wherein the compound is
10 of formula (XX):



Formula (XX).

11. The compound according to any one of the preceding claims, wherein R¹⁴ is -C(O)-OC₁₋₄ alkyl.
12. The compound according to any one of the preceding claims, wherein R¹⁴ is C₃₋₄ alkyl.
13. The compound according to any one of the preceding claims, wherein R¹² is -CF₃, -OCF₃, or a halogen.
14. The compound according to any one of the preceding claims, wherein R⁹ is -C(H)-, R¹⁰ is H, R¹¹ is F and R¹² is -CF₃.
15. The compound according to any one of the preceding claims, wherein the compound is selected from the group consisting of:
- Methyl N-(2-(dimethylamino)ethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate;
- Methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
- Methyl ((1-aminocyclopropyl)methyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl) carbamate;
- Methyl 3-((1-(4-fluoro-3-(trifluoromethyl) phenyl) cyclopropyl) (methoxycarbonyl)amino)azetidine-1-carboxylate;
- Methyl azetidin-3-yl(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
- N-(cyclopropylmethyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)azetidin-3-amine;
- Methyl (2-aminoethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
- N1-(cyclopropyl methyl)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-2-methyl propane-1,2-diamine;
- N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)cyclopropanecarboxamide;
- N1-cyclopropyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine;
- Methyl (2-amino-2-methylpropyl)(1-(3,5-dichlorophenyl)cyclopropyl)carbamate;
- Methyl (2-amino-2-methylpropyl)(1-(3-bromophenyl)cyclopropyl)carbamate;

- Methyl (2-amino-2-methylpropyl)(1-(3-chlorophenyl)cyclopropyl)carbamate;
- 5 1-((azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)amino)-2-methyl propan-2-ol;
- Methyl (2-amino-2-methylpropyl)(1-(3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
- 10 1-((1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)amino)-2-methylpropan-2-ol;
- Methyl (2-amino-2-methylpropyl)(1-(5-fluoro-4-(trifluoromethyl)pyridin-2-yl)cyclopropyl) carbamate;
- 15 N1-cyclobutyl-2-methyl-N1-(1-(3-(trifluoromethyl)phenyl)cyclopropyl)propane-1,2-diamine;
- Ethyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
- 20 Methyl (2-amino-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate;
- 25 N-(2-amino-2-methylpropyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methanesulfonamide;
- Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 30 Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 35 Methyl (2-acetamido-2-methylpropyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
- N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl) cyclopropyl)-2-methylpropane-1,2-diamine;
- 40 Methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)carbamate;
- Methyl (S)-(1-(4-fluoro-3-(trifluoro methyl) phenyl)cyclopropyl) ((1-methylpyrrolidin-2-yl)methyl) carbamate;
- 45 Methyl (S)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 50 Methyl (R)-(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- Methyl (1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylazetidin-2-yl)methyl)carbamate;

- Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl)carbamate;
- 5 Methyl (azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethoxy)phenyl)cyclopropyl)carbamate;
- Methyl (1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)(2-(hydroxyamino)-2-methylpropyl)carbamate;
- 10 Ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 15 Ethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((1-methylpyrrolidin-2-yl)methyl) carbamate;
- (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methanesulfonamide;
- 20 (S)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-((1-methylpyrrolidin-2-yl)methyl)methanesulfonamide;
- Ethyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 25 Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate;
- Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate;
- 30 (R)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-N-(pyrrolidin-2-ylmethyl) methane sulphonamide;
- 35 Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(morpholin-3-ylmethyl) carbamate;
- Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((4-methylmorpholin-3-yl)methyl)carbamate;
- 40 Isopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 45 Cyclopropyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- Methyl (R)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methyl pyrrolidin-2-yl)methyl)carbamate;
- 50 N-((1-amino cyclopropyl)methyl)-1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropan-1-amine;

- Cyclopropylmethyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)(pyrrolidin-2-ylmethyl)carbamate;
- 5 N-((1-aminocyclopropyl)methyl)-N-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)methanesulfonamide;
- Methyl (S)-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)((2-methylpyrrolidin-2-yl)methyl)carbamate;
- 10 (1S, 2S)-N1-(1-(4-fluoro-3-(trifluoromethyl) phenyl)cyclopropyl)-cyclopentane-1,2-diamine;
- (1R,2S)-N1-(1-(4-fluoro-3-(trifluoromethyl)phenyl)cyclopropyl)-cyclopentane-1,2-diamine; and
- 15 Methyl (S)-(azetidin-2-ylmethyl)(1-(4-fluoro-3-(trifluoromethyl)phenyl)-cyclopropyl)carbamate.
- 20 16. A pharmaceutical composition comprising the compound according to any one of the preceding claims.
17. The compound according to any one of claims 1 to 15 or pharmaceutical composition according claim 16 for use in medicine.
- 25 18. The compound according to any one of claims 1 to 15 or pharmaceutical composition according to claim 16 for use in the treatment of inflammatory bowel disease (IBD), such as ulcerative colitis or Crohn's disease.
- 30 19. The compound according to any one of claims 1 to 15 or pharmaceutical composition according to claim 16 for use in the treatment of hereditary xerocytosis.
- 35 20. The compound according to any one of claims 1 to 15 or pharmaceutical composition according to claim 16 for use in the treatment of acute respiratory distress syndrome (ARDS).