

METAL SALTS AND USES THEREOF

Background

The Liver X Receptor (LXR) is a nuclear receptor transcription factor. It has been found that LXR modulators are useful in the treatment of a variety of diseases, including cancers. There is a need to provide salts of such compounds with improved stability and physical properties.

Summary of the Invention

The invention provides a metal salt of an LXR β agonist. The invention also provides a method of preparing such metal salts of the LXR β agonist, pharmaceutical compositions including the metal salts, and methods of treating cancer with such compositions.

Accordingly, in an aspect, the invention features a metal salt (e.g., a pharmaceutically acceptable metal salt) of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid (Compound 1).

In some embodiments, the metal salt of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid is a multivalent metal salt.

In some embodiments, the metal salt of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid is a zinc salt, e.g., a 2:1 (Compound 1:zinc) salt.

In some embodiments, the metal salt of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid is an aluminum salt, e.g., a 3:1 (Compound 1:aluminum) salt.

In some embodiments, the metal salt of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid is a bismuth salt, e.g., a 3:1 (Compound 1:bismuth) salt.

In some embodiments, the metal salt of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid is a calcium salt, e.g., a 2:1 (Compound 1:calcium) salt.

In some embodiments, the metal salt is amorphous. In some embodiments, the metal salt (e.g., an amorphous zinc salt) has a peak with increased intensity relative to the free acid at about 1590 ± 10 cm^{-1} and a peak with reduced intensity relative to the free acid at about 1710 ± 10 cm^{-1} as measured by Fourier-transform infrared spectroscopy (FTIR). In some embodiments, the metal salt has less than 1% mass loss up to decomposition as measured by thermogravimetric analysis.

In another aspect, the invention features a method of producing a zinc salt (e.g., a 2:1 zinc salt) of Compound 1. This method includes combining Compound 1, or a salt thereof (e.g., a 1:1 sodium salt or the free compound), and a zinc salt (e.g., zinc chloride or zinc acetate) in an amount sufficient to produce the zinc salt of Compound 1.

In some embodiments of the method of producing a zinc salt, the method includes dissolving the Compound 1, or salt thereof, and the zinc salt in a solvent to form a mixture. In some embodiments, the solvent is water. In some embodiments, the solvent is a mixture of organic solvent (e.g., methanol) and water (e.g., 9:1 organic solvent and water by volume). In some embodiments of the method of producing a zinc salt, the method further includes cycling the temperature of the mixture between ambient

temperature and 40 °C. In some embodiments of the method of producing a zinc salt, wherein said cycling is performed for 24 hours.

In another aspect, the invention features a pharmaceutically acceptable zinc salt of Compound 1 produced by any of the foregoing methods.

5 In another aspect, the invention features a method of producing an aluminum salt (e.g., a 3:1 aluminum salt) Compound 1. This method includes combining Compound 1, or a salt thereof (e.g., a 1:1 sodium salt or the free compound), and an aluminum salt (e.g., aluminum sulfate) in an amount sufficient to produce the aluminum salt of Compound 1.

10 In another aspect, the invention features a pharmaceutical composition containing any of the foregoing metal salts and a pharmaceutically acceptable excipient.

In some embodiments, the pharmaceutical composition includes less than 1.5 weight % of sodium. In some embodiments, the pharmaceutical composition is substantially free of the 1:1 sodium salt of Compound 1. In some embodiments, the pharmaceutical composition is in unit dosage form.

15 In another aspect, the invention features a method of treating cancer. This method includes administering an effective amount of any of the foregoing salts or pharmaceutical compositions.

In some embodiments of the method of treating cancer, the subject has a cancer that has failed to respond to a previously administered immunotherapy (e.g., the cancer of the subject has progressed despite treatment with the immunotherapy).

20 In some embodiments of the method of treating cancer, the cancer is resistant to an immunotherapy (e.g., the cancer has been determined to be resistant to immunotherapies such as by genetic markers or the level of MDSCs (e.g., monocytic and/or granulocytic MDSCs) in a sample, or is likely to be resistant, to immunotherapies such as a cancer that has failed to respond to an immunotherapy).

25 In another aspect, the invention features a method of treating cancer that has failed to respond to an immunotherapy in a subject. This method includes administering an effective amount of any of the foregoing salts or pharmaceutical compositions to the subject in combination with an immunotherapy.

In another aspect, the invention features a method of treating cancer that is resistant to immunotherapy in a subject. This method includes administering an effective amount of any of the foregoing salts or pharmaceutical compositions to the subject in combination with an immunotherapy.

30 In some embodiments of any of the methods of treating cancer, the cancer is breast cancer, colon cancer, renal cell cancer, lung cancer (e.g., non-small cell lung cancer), hepatocellular carcinoma, gastric cancer, ovarian cancer, pancreatic cancer, esophageal cancer, prostate cancer, sarcoma, glioblastoma, diffuse large B-cell lymphoma, leukemia, or melanoma. In some embodiments, the cancer is metastatic cancer. In some embodiments of any of the methods of treating cancer, the effective
35 amount is an amount effective to suppress metastatic colonization of the cancer.

In certain embodiments of any of the methods of treating cancer, the cancer is a drug resistant cancer or has failed to respond to a prior therapy (e.g., a cancer resistant to, or a cancer that has failed to respond to prior treatment with, vemurafenib, dacarbazine, a CTLA-4 inhibitor, a PD-1 inhibitor, interferon therapy, a BRAF inhibitor, a MEK inhibitor, radiotherapy, temozolimide, irinotecan, a CAR-T therapy,
40 herceptin, perjeta, tamoxifen, xeloda, docetaxol, platinum agents such as carboplatin, taxanes such as paclitaxel and docetaxel, ALK inhibitors, MET inhibitors, alimta, abraxane, adriamycin, gemcitabine,

avastin, halaven, neratinib, a PARP inhibitor, briarane, an mTOR inhibitor, topotecan, gemzar, a VEGFR2 inhibitor, a folate receptor antagonist, demcizumab, fosbretabulin, or a PDL-1 inhibitor).

In an embodiment of any of the methods of treating cancer, the immunotherapy, when present, is a CTLA-4 inhibitor, a PD-1 inhibitor, a PD-L1 inhibitor, or adoptive T-cell transfer therapy. In some
5 embodiments, the immunotherapy includes a PD-1 inhibitor such as a PD-1 antibody, a PD-L1 inhibitor
such as a PD-L1 antibody, a CTLA-4 inhibitor such as a CTLA-4 antibody, a CSF-1R inhibitor, an IDO
inhibitor, an A1 adenosine inhibitor, an A2A adenosine inhibitor, an A2B adenosine inhibitor, an A3A
adenosine inhibitor, an arginase inhibitor, or an HDAC inhibitor. In some embodiments, the
immunotherapy includes a PD-1 inhibitor (e.g., nivolumab, pembrolizumab, pidilizumab, BMS 936559,
10 and atezolizumab). In some embodiments, the immunotherapy includes a PD-L1 inhibitor (e.g.,
atezolizumab and durvalumab). In some embodiments, the immunotherapy includes a CTLA-4 inhibitor
(e.g., ipilimumab). In some embodiments, the immunotherapy includes a CSF-1R inhibitor (e.g.,
peixidartinib and 4-(2,4-difluoroanilino)-7-ethoxy-6-(4-methylpiperazin-1-yl)quinoline-3-carboxamide). In
some embodiments, the immunotherapy includes an IDO inhibitor (e.g., norharmane, rosmarinic acid, and
15 alpha-methyl-tryptophan). In some embodiments, the immunotherapy includes an A1 adenosine inhibitor
(e.g., 8-cyclopentyl-1,3-dimethylxanthine, 8-cyclopentyl-1,3-dipropylxanthine, 8-phenyl-1,3-
dipropylxanthine, bamifylline, BG-9719, tonapofylline, FK-453, FK-838, rollofylline, or N-0861). In some
embodiments, the immunotherapy includes an A2A adenosine inhibitor (e.g., ATL-4444, istradefylline,
MSX-3, preladenant, SCH-58261, SCH-412348, SCH-442416, 2-butyl-9-methyl-8-(triazol-2-yl)purin-6-
20 amine, VER-6623, VER-6947, VER-7835, viadenant, or ZM-241,385). In some embodiments, the
immunotherapy includes an A2B adenosine inhibitor (e.g., *N*-[5-(1-cyclopropyl-2,6-dioxo-3-propyl-7*H*-
purin-8-yl)pyridin-2-yl]-*N*-ethylpyridine-3-carboxamide, 3-ethyl-1-propyl-8-[1-[[3-
(trifluoromethyl)phenyl]methyl]pyrazol-4-yl]-7*H*-purine-2,6-dione, MRS-1706, MRS-1754, *N*-[2-[[2-phenyl-
6-[4-(3-phenylpropyl)piperazine-1-carbonyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]amino]ethyl]acetamide, PSB-
25 603, PSB-0788, or PSB-1115). In some embodiments, the immunotherapy includes an A3A adenosine
inhibitor (e.g., KF-26777, MRS-545, MRS-1191, MRS-1220, MRS-1334, propyl 6-ethyl-5-
ethylsulfanylicarbonyl-2-phenyl-4-propylpyridine-3-carboxylate, MRS-3777, MRE-3005-F20, MRE-3008-
F20, PSB-11, OT-7999, VUF-5574, and SSR161421). In some embodiments, the immunotherapy
includes an arginase inhibitor (e.g., an arginase antibody, (2*S*)-(+)-amino-5-iodoacetamidopentanoic acid,
30 NG-hydroxy-L-arginine, (2*S*)-(+)-amino-6-iodoacetamidohexanoic acid, or (R)-2-amino-6-borono-2-(2-
(piperidin-1-yl)ethyl)hexanoic acid. In some embodiments, the immunotherapy includes an HDAC
inhibitor (e.g., valproic acid, SAHA, or romidepsin).

In another embodiment of any of the methods of treating cancer, the method further includes
administering to the subject an additional anticancer therapy (e.g., an antiproliferative).

35 In particular embodiments, the antiproliferative is: a chemotherapeutic or cytotoxic agent, a
differentiation-inducing agent (e.g., retinoic acid, vitamin D, cytokines), a hormonal agent, an
immunological agent, or an anti-angiogenic agent. Chemotherapeutic and cytotoxic agents include, but
are not limited to, alkylating agents, cytotoxic antibiotics, antimetabolites, vinca alkaloids, etoposides, and
others (e.g., paclitaxel, taxol, docetaxel, taxotere, cis-platinum). A list of additional compounds having
40 antiproliferative activity can be found in L. Brunton, B. Chabner and B. Knollman (eds). Goodman and

Gilman's The Pharmacological Basis of Therapeutics, Twelfth Edition, 2011, McGraw Hill Companies, New York, NY.

In certain embodiments, the antiproliferative is a PD-1 inhibitor, a VEGF inhibitor, a VEGFR2 inhibitor, a PD-L1 inhibitor, a BRAF inhibitor, a CTLA-4 inhibitor, a MEK inhibitor, an ERK inhibitor, vemurafenib, dacarbazine, trametinib, dabrafenib, durvalumab, an mTOR inhibitor, a CAR-T therapy, abiraterone, enzalutamine, apalutamide, 5-fluorouracil (5-FU), FOLFOX (i.e., folinic acid, 5-fluorouracil, and oxaliplatin), FOLFIRI (i.e., folinic acid, 5-fluorouracil, and irinotecan), herceptin, xeloda, a PD-1 antibody (e.g., pembrolizumab or nivolumab), a PD-L1 antibody, a CTLA-4 antibody (e.g., ipilimumab), ramucirumab, rindopepimut, glembatumumab, vedotin, ANG1005, and/or ANG4043.

In some embodiments of any of the methods of treating cancer, the cancer is a renal cell carcinoma and the antiproliferative is a PD-1 inhibitor, a PD-L1 inhibitor, or an mTOR inhibitor. In other embodiments, the cancer is diffuse large B-cell lymphoma and the antiproliferative is a CAR-T therapy. In certain embodiments, the cancer is prostate cancer and the antiproliferative is abiraterone, enzalutamide, or apalutamide. In some embodiments, the cancer is hepatocellular carcinoma, gastric cancer, or esophageal cancer and the antiproliferative is 5-FU, FOLFOX, FOLFIRI, herceptin, or xeloda. In some embodiments, the cancer is sarcoma and the antiproliferative is gemcitabine. In other embodiments, the cancer is pancreatic cancer and the antiproliferative is irinotecan, cisplatin, abraxane, a taxane (e.g., paclitaxel or docetaxel), or capecitabine.

The methods of treating cancer may further include administering an antiproliferative such as alkylating agents, platinum agents, antimetabolites, topoisomerase inhibitors, antitumor antibiotics, antimitotic agents, aromatase inhibitors, thymidylate synthase inhibitors, DNA antagonists, farnesyltransferase inhibitors, pump inhibitors, histone acetyltransferase inhibitors, metalloproteinase inhibitors, ribonucleoside reductase inhibitors, TNF alpha agonists/antagonists, endothelin A receptor antagonist, retinoic acid receptor agonists, immuno-modulators, hormonal and antihormonal agents, photodynamic agents, tyrosine kinase inhibitors, antisense compounds, corticosteroids, HSP90 inhibitors, proteasome inhibitors (for example, marizomib), CD40 inhibitors, anti-CSI antibodies, FGFR3 inhibitors, VEGF inhibitors, MEK inhibitors, cyclin D1 inhibitors, NF-kB inhibitors, anthracyclines, histone deacetylases, kinesin inhibitors, phosphatase inhibitors, COX2 inhibitors, mTOR inhibitors, calcineurin antagonists, IMiDs, and/or other agents used to treat proliferative diseases.

In some embodiments of any of the methods of treating cancer, the cancer is breast cancer such as triple negative breast cancer, colon cancer, renal cell cancer, lung cancer (e.g., non-small cell lung cancer), hepatocellular carcinoma, gastric cancer, ovarian cancer, pancreatic cancer, esophageal cancer, prostate cancer, sarcoma, glioblastoma, diffuse large B-cell lymphoma, leukemia (e.g., acute myeloid leukemia), or melanoma. In some embodiments of any of the methods of treating cancer, the cancer is melanoma. In some embodiments of any of the methods of treating cancer, the cancer is breast cancer. In some embodiments of any of the foregoing methods, the cancer is renal cell cancer. In some embodiments of any of the methods of treating cancer, the cancer is pancreatic cancer. In some embodiments of any of the methods of treating cancer, the cancer is non-small cell lung cancer. In some embodiments of any of the methods of treating cancer, the cancer is colon cancer. In some embodiments of any of the methods of treating cancer, the cancer is ovarian cancer. In some embodiments of any of the methods of treating cancer, the cancer is glioblastoma. In some embodiments of any of the methods

of treating cancer, the cancer is prostate cancer. In some embodiments of any of the methods of treating cancer, the cancer is diffuse large B-cell lymphoma. In some embodiments, the cancer is leukemia (e.g., acute myeloid leukemia).

In particular embodiments of any of the methods of treating cancer, the cancer is melanoma (e.g., metastatic melanoma) that is resistant to, or has failed to respond to prior treatment with, vemurafenib, dacarbazine, interferon therapy, a CTLA-4 inhibitor, a BRAF inhibitor, a MEK inhibitor, a PD1 inhibitor, a PDL-1 inhibitor, and/or a CAR-T therapy. In some embodiments of any of the methods of treating cancer, the cancer is glioblastoma that is resistant to, or has failed to respond to prior treatment with, temozolimide, radiotherapy, avastin, irinotecan, a VEGFR2 inhibitor, a CAR-T therapy, and/or an mTOR inhibitor. In some embodiments of any of the methods of treating cancer, the cancer is non-small cell lung cancer such as metastatic non-small cell lung cancer (e.g., EGFR-wild type non-small cell lung cancer and/or squamous non-small cell lung cancer) that is resistant to, or has failed to respond to prior treatment with, an EGFR inhibitor, platinum agents (e.g., carboplatin), avastin, an ALK inhibitor, a MET inhibitor, a taxane (e.g., paclitaxel and/or docetaxel), gemzar, alimta, radiotherapy, a PD-1 inhibitor, a PD-L1 inhibitor, and/or a CAR-T therapy. In some embodiments of any of the methods of treating cancer, the cancer is a breast cancer (e.g., triple negative breast cancer) that is resistant to, or has failed to respond to prior treatment with, herceptin, perjeta, tamoxifen, xeloda, docetaxel, carboplatin, paclitaxel, abraxane, adriamycin, gemcitabine, avastin, halaven, neratinib, a PARP inhibitor, a PD-1 inhibitor, a PD-L1 inhibitor, a CAR-T therapy, apalutamide, and/or an mTOR inhibitor. In some embodiments of any of the methods of treating cancer, the cancer is ovarian cancer (e.g., advanced ovarian cancer) that is resistant to, or has failed to respond to prior treatment with, a PARP inhibitor, avastin, platinum agents such as carboplatin, paclitaxel, docetaxel, topotecan, gemzar, a VEGFR2 inhibitor, a folate receptor antagonist, a PD-1 inhibitor, a PD-L1 inhibitor, a CAR-T therapy, demcizumab, and/or fosbretabulin.

In some embodiments of any of the methods of treating cancer, the additional anti-cancer therapy, if present, includes chemotherapy.

In some embodiments of any of the methods of treating cancer, the chemotherapy includes docetaxel. In some embodiments, the method includes administering an effective amount of docetaxel to the subject once every seven days. In some embodiments, the effective amount of docetaxel is at least 28 mg/m². In some embodiments, the effective amount of docetaxel is about 28 mg/m² to about 35 mg/m².

In some embodiments of any of the methods of treating cancer, the additional anti-cancer therapy includes chemotherapy and immunotherapy. In some embodiments of any of the methods of treating cancer, the anti-cancer therapy includes carboplatin or cisplatin, pemetrexed, and pembrolizumab. In some embodiments, the method includes administering to the subject an effective amount of pembrolizumab once every twenty-one days. In some embodiments, the effective amount of pembrolizumab is about 200 mg. In some embodiments, the method includes administering to the subject an effective amount of carboplatin or cisplatin once every twenty-one days. In some embodiments, the effective amount of carboplatin or cisplatin is calculated using the formula: Total dose (mg) = (Target area under the curve) x (subject's glomerular filtration rate+25), wherein the target area under the curve is 4 mg/mL*min to 6 mg/mL*min and the subject's glomerular filtration rate was measured by Cr-EDTA clearance. In some embodiments, the effective amount of carboplatin or cisplatin is about

300 mg/m² to about 360 mg/m². In some embodiments, the method includes administering to the subject an effective amount of pemetrexed once every twenty-one days. In some embodiments, the effective amount of pemetrexed is 500 mg/m². In some embodiments of any of the methods of treating cancer, the method further includes administering to the subject an effective amount of folic acid, vitamin B12, and/or corticosteroids. In some embodiments, the method includes administering to the subject an effective amount of corticosteroids twice per day for three days prior to administration of pemetrexed.

In some embodiments of any of the methods of treating cancer, the method further includes administering to the subject an effective amount of a statin (e.g., rosuvastatin or atorvastatin).

In some embodiments of any of the methods of treating cancer, the method further includes administering to the subject an effective amount of an anti-emetic agent (e.g., ondansetron, granisetron, palonosetron, metoclopramide, haloperidol, dexamethasone, aprepitant, fosaprepitant, lorazepam, dronabinol, prochlorperazine, or chlorpromazine), an anti-diarrheal agent (e.g., an opiate agonist or octreotide), an appetite stimulant (e.g., megestrol acetate, metoclopramide, dronabinol, prednisone, or dexamethasone), a general stimulant, a bisphosphonate (e.g., etidronate, clodronate, tiludronate, pamidronate, neridronate, opladronate, alendronate, ibandronate, risedronate, or zoledronate), a gonadotrophin releasing hormone agonist (e.g., buserelin, histrelin, leuporelin, triptorelin, goserelin, or nafarelin), and/or growth factors (e.g., filgrastim).

In some embodiments of any of the methods of treating cancer, the cancer is resistant to an anti-cancer therapy (e.g., platinum-containing chemotherapy, a PD-1 inhibitor, a PD-L1 inhibitor, a CTLA-4 inhibitor, an antimetabolite, a topoisomerase inhibitor, an angiogenesis inhibitor, a kinase inhibitor, and/or an alkylating agent). In some embodiments of any of the methods of treating cancer, the cancer progressed on or after treatment with an anti-cancer therapy (e.g., platinum-containing chemotherapy, a PD-1 inhibitor, a PD-L1 inhibitor, an angiogenesis inhibitor, a kinase inhibitor, and/or an alkylating agent). In some embodiments of any of the methods of treating cancer, the cancer has been determined to be, or is predicted to be, resistant to an anti-cancer therapy (e.g., a PD-1 inhibitor, a PD-L1 inhibitor, a CTLA-4 inhibitor, a topoisomerase inhibitor, an antimetabolite, an angiogenesis inhibitor, a kinase inhibitor, and/or an alkylating agent).

In some embodiments of any of the methods of treating cancer, the cancer has a PD-L1 expression level of less than 1% when tested in an immunohistochemistry assay (e.g., an immunohistochemistry assay with a tumor proportion score). In some embodiments of any of the methods of treating cancer, the cancer has a PD-L1 expression level of about 1% when tested in an immunohistochemistry assay (e.g., an immunohistochemistry assay with a tumor proportion score). In some embodiments of any of the methods of treating cancer, the cancer has a PD-L1 expression level of about 1% to about 49% (e.g., about 1% to about 20%, about 5% to about 30%, about 15% to about 40%, about 25% to about 49%) when tested in an immunohistochemistry assay (e.g., an immunohistochemistry assay with a tumor proportion score). In some embodiments of any of the methods of treating cancer, the cancer is metastatic and/or locally advanced. In some embodiments of any of the methods of treating cancer, the cancer is unresectable.

40

Definitions

As used herein, the term “administration” refers to the administration of a composition (e.g., a salt or a preparation that includes a salt as described herein) to a subject or system. Administration to an animal subject (e.g., to a human) may be by any appropriate route. For example, in some embodiments, administration may be bronchial (including by bronchial instillation), buccal, enteral, interdermal, intra-arterial, intradermal, intragastric, intramedullary, intramuscular, intranasal, intraperitoneal, intrathecal, intravenous, intraventricular, mucosal, nasal, oral, rectal, subcutaneous, sublingual, topical, tracheal (including by intratracheal instillation), transdermal, vaginal, and vitreal.

By “biological sample” or “sample” is meant a fluid or solid sample from a subject. Biological samples may include cells; nucleic acid, protein, or membrane extracts of cells; or blood or biological fluids including (e.g., plasma, serum, saliva, urine, bile). Solid biological samples include samples taken from feces, the rectum, central nervous system, bone, breast tissue, renal tissue, the uterine cervix, the endometrium, the head or neck, the gallbladder, parotid tissue, the prostate, the brain, the pituitary gland, kidney tissue, muscle, the esophagus, the stomach, the small intestine, the colon, the liver, the spleen, the pancreas, thyroid tissue, heart tissue, lung tissue, the bladder, adipose tissue, lymph node tissue, the uterus, ovarian tissue, adrenal tissue, testis tissue, the tonsils, and the thymus. Fluid biological samples include samples taken from the blood, serum, plasma, pancreatic fluid, CSF, semen, prostate fluid, seminal fluid, urine, saliva, sputum, mucus, bone marrow, lymph, and tears. Samples may be obtained by standard methods including, e.g., venous puncture and surgical biopsy. In certain embodiments, the biological sample is a blood, plasma, or serum sample. In some embodiments, the biological sample is a tumor sample from a biopsy.

The term “cancer” refers to any cancer caused by the proliferation of malignant neoplastic cells, such as tumors, neoplasms, carcinomas, sarcomas, leukemias, and lymphomas.

By “determining the level of a cell type” is meant the detection of a cell type by methods known in the art either directly or indirectly. “Directly determining” means performing a process (e.g., performing an assay or test on a sample or “analyzing a sample” as that term is defined herein) to obtain the physical entity or value. “Indirectly determining” refers to receiving the physical entity or value from another party or source (e.g., a third party laboratory that directly acquired the physical entity or value). Methods to measure cell levels generally include, but are not limited to, flow cytometry and immunohistochemistry. Exemplary methods are provided herein. In some embodiments of any of the foregoing methods, the level of MDSCs and/or activated T-cells may be determined as described in Iclozan et al. *Cancer Immunol. Immunother.* 2013, 62(5): 909-918. In some embodiments of any of the foregoing methods, the level of MDSCs and/or activated T-cells may be determined as described in Kitano et al. *Cancer Immunol. Res.* 2014, 2(8): 812-821.

A cancer “determined to be drug resistant,” as used herein, refers to a cancer that is drug resistant, based on unresponsiveness or decreased responsiveness to a chemotherapeutic agent, or is predicted to be drug resistant based on a prognostic assay (e.g., a gene expression assay).

By a “drug resistant” cancer is meant a cancer that does not respond, or exhibits a decreased response to, one or more chemotherapeutic agents (e.g., any agent described herein).

The term “effective amount” means an amount that is sufficient, when administered to a population suffering from or susceptible to a disease, disorder, and/or condition in accordance with a

therapeutic dosing regimen, to treat the disease, disorder, and/or condition. In some embodiments, a therapeutically effective amount is one that reduces the incidence and/or severity of, and/or delays onset of, one or more symptoms of the disease, disorder, and/or condition. Those of ordinary skill in the art will appreciate that the term "effective amount" does not in fact require successful treatment be achieved in a particular individual. Rather, an effective amount may be that amount that provides a particular desired pharmacological response in a significant number of subjects when administered to patients in need of such treatment. It is specifically understood that particular subjects may, in fact, be "refractory" to an "effective amount." To give but one example, a refractory subject may have a low bioavailability such that clinical efficacy is not obtainable. In some embodiments, reference to an effective amount may be a reference to an amount as measured in one or more specific tissues (e.g., a tissue affected by the disease, disorder or condition) or fluids (e.g., blood, saliva, serum, sweat, tears, urine). Those of ordinary skill in the art will appreciate that, in some embodiments, an effective amount may be formulated and/or administered in a single dose. In some embodiments, an effective amount may be formulated and/or administered in a plurality of doses, for example, as part of a dosing regimen.

As used herein, the term "failed to respond to a prior therapy" or "refractory to a prior therapy," refers to a cancer that progressed despite treatment with the therapy.

By "level" is meant a level of a cell type, as compared to a reference. The reference can be any useful reference, as defined herein. By a "decreased level" or an "increased level" of a cell type is meant a decrease or increase in cell level, as compared to a reference (e.g., a decrease or an increase by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 100%, about 150%, about 200%, about 300%, about 400%, about 500%, or more; a decrease or an increase of more than about 10%, about 15%, about 20%, about 50%, about 75%, about 100%, or about 200%, as compared to a reference; a decrease or an increase by less than about 0.01-fold, about 0.02-fold, about 0.1-fold, about 0.3-fold, about 0.5-fold, about 0.8-fold, or less; or an increase by more than about 1.2-fold, about 1.4-fold, about 1.5-fold, about 1.8-fold, about 2.0-fold, about 3.0-fold, about 3.5-fold, about 4.5-fold, about 5.0-fold, about 10-fold, about 15-fold, about 20-fold, about 30-fold, about 40-fold, about 50-fold, about 100-fold, about 1000-fold, or more). A level of a cell type may be expressed in mass/vol (e.g., g/dL, mg/mL, µg/mL, ng/mL) or percentage relative to total cells in a sample. In some embodiments of any of the foregoing methods, the reference is a sample from a healthy subject such as a subject that does not have cancer. In some embodiments of any of the foregoing methods, the reference is an artificial sample with a level (e.g., a level of MDSCs such as monocytic and/or granulocytic MDSCs or activated T-cells) shown beneficial in the treatment of a disorder.

As used herein, "metastatic nodule" refers to an aggregation of tumor cells in the body at a site other than the site of the original tumor.

As used herein, "metastatic tumor" refers to a tumor or cancer in which the cancer cells forming the tumor have a high potential to or have begun to, metastasize, or spread from one location to another location or locations within a subject, via the lymphatic system or via haematogenous spread, for example, creating secondary tumors within the subject. Such metastatic behavior may be indicative of

malignant tumors. In some cases, metastatic behavior may be associated with an increase in cell migration and/or invasion behavior of the tumor cells.

Examples of cancers that can be defined as metastatic include but are not limited to lung cancer (e.g., non-small cell lung cancer), breast cancer, ovarian cancer, colorectal cancer, biliary tract cancer, bladder cancer, brain cancer including glioblastomas and medullablastomas, cervical cancer, choriocarcinoma, endometrial cancer, esophageal cancer, gastric cancer, hematological neoplasms, multiple myeloma, leukemia, intraepithelial neoplasms, liver cancer, lymphomas, neuroblastomas, oral cancer, pancreatic cancer, prostate cancer, sarcoma, skin cancer including melanoma, basocellular cancer, squamous cell cancer, testicular cancer, stromal tumors, germ cell tumors, thyroid cancer, and renal cancer.

“Non-metastatic cell migration cancer” as used herein refers to cancers that do not migrate via the lymphatic system or via haematogenous spread.

As used herein, the term “pharmaceutical composition” refers to an active compound, or pharmaceutically acceptable salt thereof, formulated together with one or more pharmaceutically acceptable carriers. In some embodiments, active compound or salt is present in unit dose amount appropriate for administration in a therapeutic regimen that shows a statistically significant probability of achieving a predetermined therapeutic effect when administered to a relevant population. In some embodiments, pharmaceutical compositions may be specially formulated for administration in solid or liquid form, including those adapted for the following: oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual, and systemic absorption, boluses, powders, granules, pastes for application to the tongue; parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; topical application, for example, as a cream, ointment, or a controlled-release patch or spray applied to the skin, lungs, or oral cavity; intravaginally or intrarectally, for example, as a pessary, cream, or foam; sublingually; ocularly; transdermally; or nasally, pulmonary, and to other mucosal surfaces.

A “pharmaceutically acceptable excipient,” as used herein, refers any inactive ingredient (for example, a vehicle capable of suspending or dissolving the active compound) having the properties of being nontoxic and non-inflammatory in a subject. Typical excipients include, for example: antiadherents, antioxidants, binders, coatings, compression aids, disintegrants, dyes (colors), emollients, emulsifiers, fillers (diluent), film formers or coatings, flavors, fragrances, glidants (flow enhancers), lubricants, preservatives, printing inks, sorbents, suspending or dispersing agents, sweeteners, or waters of hydration.

The term “pharmaceutically acceptable salt,” as use herein, refers to those salts of the compounds described here that are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and animals without undue toxicity, irritation, allergic response and the like and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, pharmaceutically acceptable salts are described in: Berge et al., *J. Pharmaceutical Sciences* 66:1-19, 1977 and in *Handbook of Pharmaceutical Salts: Properties, Selection, and Use*, (Eds. P.H. Stahl and C.G. Wermuth), Wiley-VCH, 2008. The salts can be prepared in situ

during the final isolation and purification of the compounds described herein or separately by reacting the free base group with a suitable organic acid.

"Progression-free survival" as used herein, refers to the length of time during and after medication or treatment during which the disease being treated (e.g., cancer) does not get worse.

5 "Proliferation" as used in this application involves reproduction or multiplication of similar forms (cells) due to constituting (cellular) elements.

As used herein, "slowing the spread of metastasis" refers to reducing or stopping the formation of new loci; or reducing, stopping, or reversing the tumor load.

10 The term "subject," as used herein, refers to a human or non-human animal (e.g., a mammal such as a non-human primate, horse, cow, or dog).

The term "substantially" refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of interest. One of ordinary skill in the biological arts will understand that biological and chemical phenomena rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term "substantially" is therefore used herein to capture the
15 potential lack of completeness inherent in many biological and chemical phenomena.

A "therapeutic regimen" refers to a dosing regimen whose administration across a relevant population is correlated with a desired or beneficial therapeutic outcome.

The term "treatment" (also "treat" or "treating"), in its broadest sense, refers to any administration of a substance (e.g., provided compositions) that partially or completely alleviates, ameliorates, relieves,
20 inhibits, delays onset of, reduces severity of, and/or reduces incidence of one or more symptoms, features, and/or causes of a particular disease, disorder, and/or condition. In some embodiments, such treatment may be administered to a subject who does not exhibit signs of the relevant disease, disorder and/or condition and/or of a subject who exhibits only early signs of the disease, disorder, and/or condition. Alternatively or additionally, in some embodiments, treatment may be administered to a
25 subject who exhibits one or more established signs of the relevant disease, disorder and/or condition. In some embodiments, treatment may be of a subject who has been diagnosed as suffering from the relevant disease, disorder, and/or condition. In some embodiments, treatment may be of a subject known to have one or more susceptibility factors that are statistically correlated with increased risk of development of the relevant disease, disorder, and/or condition.

30 The term "PD-1 inhibitor," as used herein, refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the PDCD1 gene. Known PD-1 inhibitors include nivolumab, pembrolizumab, pidilizumab, BMS 936559, and atezolizumab.

The term "PD-L1 inhibitor," as used herein, refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the CD274 gene. Known PD-L1
35 inhibitors include atezolizumab and durvalumab.

The term "CTLA-4 inhibitor," as used herein, refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the CTLA4 gene. Known CTLA-4 inhibitors include ipilimumab.

40 The term "CSF-1R inhibitors," as used herein refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the CSF1R gene. Known CSF-1R

inhibitors include pexidartinib and 4-(2,4-difluoroanilino)-7-ethoxy-6-(4-methylpiperazin-1-yl)quinoline-3-carboxamide.

The term "IDO inhibitor," as used herein, refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the IDO1 gene. Known IDO inhibitors include norharmane, rosmarinic acid, and alpha-methyl-tryptophan.

The term "A1 adenosine inhibitor," as used herein, refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the ADORA1 gene. Known A1 adenosine inhibitors include 8-cyclopentyl-1,3-dimethylxanthine, 8-cyclopentyl-1,3-dipropylxanthine, 8-phenyl-1,3-dipropylxanthine, bamifylline, BG-9719, tonapofylline, FK-453, FK-838, rolofylline, and N-0861.

The term "A2A adenosine inhibitor," as used herein, refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the ADORA2A gene. Known A2A adenosine inhibitors include ATL-4444, istradefylline, MSX-3, preladenant, SCH-58261, SCH-412,348, SCH-442,416, 2-butyl-9-methyl-8-(triazol-2-yl)purin-6-amine, VER-6623, VER-6947, VER-7835, viadenant, and ZM-241,385.

The term "A2B adenosine inhibitor," as used herein, refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the ADORA2B gene. Known A2B adenosine inhibitors include *N*-[5-(1-cyclopropyl-2,6-dioxo-3-propyl-7*H*-purin-8-yl)pyridin-2-yl]-*N*-ethylpyridine-3-carboxamide, 3-ethyl-1-propyl-8-[1-[[3-(trifluoromethyl)phenyl]methyl]pyrazol-4-yl]-7*H*-purine-2,6-dione, MRS-1706, MRS-1754, *N*-[2-[[2-phenyl-6-[4-(3-phenylpropyl)piperazine-1-carbonyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]amino]ethyl]acetamide, PSB-603, PSB-0788, and PSB-1115.

The term "A3A adenosine inhibitor," as used herein, refers to a compound such as an antibody capable of inhibiting the activity of the protein that in humans is encoded by the ADORA3 gene. Known A3A adenosine inhibitors include KF-26777, MRS-545, MRS-1191, MRS-1220, MRS-1334, MRS-1523, MRS-3777, MRE-3005-F20, MRE-3008-F20, PSB-11, OT-7999, VUF-5574, and SSR161421.

The term "arginase inhibitor," as used herein, refers to a compound capable of inhibiting the activity of a protein that in humans is encoded by the ARG1 or ARG2 genes. Known arginase inhibitors include (2*S*)-(+)-amino-5-iodoacetamidopentanoic acid, NG-hydroxy-L-arginine, (2*S*)-(+)-amino-6-iodoacetamidohexanoic acid, and (R)-2-amino-6-borono-2-(2-(piperidin-1-yl)ethyl)hexanoic acid.


The term "HDAC inhibitor," as used herein, refers to a compound such as an antibody that is capable of inhibiting the activity of the protein that is a member of the histone deacetylase class of enzymes, e.g., HDAC1, HDAC2, HDAC3, HDAC4, HDAC5, HDAC6, HDAC7, HDAC8, HDAC9, HDAC10, HDAC11, SIRT1, SIRT2, SIRT3, SIRT4, SIRT5, SIRT6, and SIRT7. Known HDAC inhibitors include valproic acid, SAHA, and romidepsin.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present disclosure; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting. All publications, patent applications, patents, sequences, database entries, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

prone to discoloration and/or loss of hydrochloric acid. The inventors also found that extraction of the hydrochloride salt with water led to loss of hydrochloric acid creating further processing difficulties. The hydrochloride salt was further found to undergo apparent esterification in the presence of free alcohol groups, e.g., when in alcoholic solvents or in formulations with excipients with free alcohol groups such as sorbitol, or esterification in the presence of lipophilic esters such as linoleate. These instability issues of the hydrochloric acid salt made it less suitable for processing into a drug product, thus salt screening studies were conducted to find a more suitable salt form.

Subsequent salt screening studies were performed to identify counterions and solvent systems that were conducive to forming salts that possess both improved physical properties and stability. From these salt screening studies, including experiments with over sixty-five different counter ion sources, it was found that the 2:1 (Compound 1:zinc) salt was, surprisingly, one of the minority of salts that formed as a stable solid as opposed to the gels, gums, oils, and semi-solids which formed with the majority of counterion sources. As a reproducible flowable solid, the 2:1 (Compound 1:zinc) salt is much more suitable for scale up and development in a drug product.

2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid (Compound 1) is an LXR β agonist having the structure:



Compound 1

Compound 1-mediated activation of the LXR signaling pathway has been shown to induce expression of ApoE, which functions as a tumor suppressor by virtue of its ability to regulate key features of tumorigenesis. Compound 1 has a higher specificity for the LXR β isoform. These features include suppression of cancer cell invasion (~45% in vitro), inhibition of endothelial recruitment (~50% in vitro) and the reduction of MDSCs in circulation (~40%) and in tumors (> 60%). In in vitro studies, Compound 1 induced ApoE gene expression by 3-fold in cancer cells and up to 40-fold in human peripheral blood mononuclear cells (hPBMCs) compared to control cells. The EC₅₀ of Compound 1 for ApoE induction was 385 nM in cancer cells and 271 nM in hPBMCs.

In syngeneic and human xenograft mouse tumor models of melanoma (harboring different genetic backgrounds), glioblastoma, TNBC, ovarian cancer, and lung cancer, Compound 1 inhibited primary tumor growth by 48-95%. Extent of tumor growth inhibition varied with model. In a mouse model of TNBC metastasis, Compound 1 inhibited metastatic spread of cancer cells by ~ 9-fold. Moreover, the anti-tumor activity of Compound 1 in combination with an anti-PD-1 antibody inhibited tumor growth by >80% in a syngeneic mouse melanoma model that is otherwise not responsive to anti-PD-1 antibody. Furthermore, inhibition of tumor growth in the same syngeneic mouse melanoma model was superior when the mice received combination therapy of Compound 1 and anti-CTLA-4 antibody relative to either

therapy alone. Similarly, in a syngeneic mouse melanoma model, Compound 1 demonstrated superior anti-tumor efficacy in combination with dacarbazine (>80%), compared to either treatment alone. In tumor growth studies, minimum efficacious doses ranged from 25-40 mg/kg/day administered orally (PO) resulting in exposures that ranged from 10,000-50,000 ng-h/mL.

- 5 In safety pharmacology assessments, Compound 1 produced a significant increase, but not inhibition of human ether-à-go-go-related gene (hERG) channel conductance in an in vitro hERG assay. There were no Compound 1-related effects on qualitative electrocardiogram (ECG) parameters (PR or QTc intervals or QRS duration) in dogs, but there was a dose related decrease in mean heart rate at the Day 1 post-dose interval that was significantly different in females following the 150 (stepped down to
- 10 100) mg/kg/day dose. This change was not observed during the recovery period and was not considered adverse. Furthermore, no adverse effects were noted during the neurobehavioral functional observation battery (FOB) or respiratory evaluations in rats. Given the favorable safety profile of Compound 1 at the highest doses tested in the repeat dose toxicity studies, the potential for cardiovascular, respiratory or central nervous system (CNS) system effects is considered low.
- 15 In oral PK studies, Compound 1 was well absorbed in CD-1 mice with a calculated absolute oral bioavailability (%F) often >100%, indicative of possible enterohepatic recycling of parent compound. The time to maximum plasma concentration (T_{max}) was similar for males and females and ranged from 2 to 8 hours. The mean apparent oral half-life (t_{1/2}) ranged from 6.5-8 hours in mice. There was a significant food effect showing that Compound 1 plasma concentrations were >2-fold higher when administered to
- 20 mice in the fed state. In Sprague-Dawley rats, the combined, mean %F (following a 30 mg/kg oral dose of 2-[3-[(3R)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid) was moderate (~31%), and T_{max} was similar for males and females, ranging from 4 to 8 hours. Compound 1 was cleared at a lower rate in female rats compared to males, which resulted in higher systemic exposure to Compound 1 in females at all dose levels tested. The mean apparent oral t_{1/2} in
- 25 female rats was 6.5 hours (not calculable in males). In male Beagle dogs given an oral dose of Compound 1, T_{max} ranged from 4-8 hours. The mean %F was moderate (18-30% depending on dose and formulation) and mean apparent oral t_{1/2} ranged from 5-6.7 hours. In Cynomolgus monkeys, mean %F was low to moderate (6-19% depending on dose and formulation). Following an oral dose, monkeys had a mean T_{max} of 4 hours. Mean oral t_{1/2} ranged from 5.5-8 hours.
- 30 Compound 1 is subjected to phase I and phase II metabolism, which includes oxidation, dealkylation, glucuronidation and combinations thereof. In vitro, Compound 1 is metabolized predominantly by the cytochrome P450 (CYP) isoform CYP3A4, but it is also a substrate for CYP2E1, CYP2C9, CYP2C19, and possibly CYP2J2. While Compound 1 is not a strong inhibitor of any human CYP450s in vitro, it is a moderate inhibitor of CYP2C8 (7.5 μM 50% inhibitory concentration [IC₅₀]) and a weak inhibitor of 2B6
- 35 (15 μM IC₅₀). Compound 1 very weakly inhibited 1A1, 2A6, 2C9, 2C19, 2D6, 2E1, and 3A4 in vitro, but CYP3A time-dependent inhibition (TDI) was demonstrated in vitro using testosterone as the substrate. Induction of CYP3A by Compound 1 was demonstrated in primary cultures of cryopreserved hepatocytes (2 donors) and the potential of Compound 1 to induce CYP2B6 cannot be ruled out (1 concentration in 1/3 donors was induced > 2-fold). Compound 1 did not induce CYP1A2. In efflux transporters,
- 40 Compound 1 does not inhibit P-glycoprotein (P-gp) but does inhibit breast cancer resistance protein (BCRP) transport in vitro (55% at 5 μM). Compound 1 is a potent inhibitor of the uptake transporter

organic anion transporting polypeptide (OATP) 1B1 in vitro (0.099 μ M IC50). Compound 1 also appears to be a moderate inhibitor of OATP1B3 (3.7 μ M IC50). Compound 1 only weakly inhibited OAT1, OAT3, and OCT2 in vitro with inhibition less than 50% at 50 μ M.

Potential risks with Compound 1 in the clinical setting based on animal toxicology studies include elevations in serum cholesterol and TG, neutropenia/leukopenia, nausea and/or vomiting, elevations in liver enzymes, development or worsening of cataracts, cardiac rhythm disturbances and/or reduced cardiac function, harderian gland adenocarcinoma, and/or generalized edema.

Nivolumab

Nivolumab is a fully human immunoglobulin (Ig) G4 monoclonal antibody directed against the negative immunoregulatory human cell surface receptor programmed death-1 (PD-1) with immune checkpoint inhibitory and antineoplastic activities. Nivolumab binds to and blocks the activation of PD-1, an Ig superfamily transmembrane protein, by its ligands programmed cell death ligand 1 (PD-L1), overexpressed on certain cancer cells, and programmed cell death ligand 2 (PD-L2), which is primarily expressed on Antigen Presenting Cells. This results in the activation of T-cells and cell-mediated immune responses against tumor cells or pathogens. Activated PD-1 negatively regulates T-cell activation and plays a key role in tumor evasion from host immunity. The nivolumab dose will be 240 mg administered as an intravenous infusion over 60 minutes on Days 1 and 15 of each 28-day cycle.

There is potential for overlapping toxicities between Compound 1 and nivolumab. Specifically, a DLT of Grade 4 neutropenia has been seen on Compound 1 single agent therapy and myelosuppression may be observed with nivolumab treatment. The pharmacological effects of Compound 1 include modulation of sterol biosynthesis. Consequently, hyperlipidemia has been observed in subjects treated with Compound 1, which has also been reported in subjects treated with nivolumab. Liver function abnormalities have been observed in pre-clinical toxicity studies with Compound 1 and immune-mediated hepatitis has been observed with nivolumab treatment.

Ipilimumab

Ipilimumab is a recombinant human IgG1 kappa monoclonal antibody that binds to the cytotoxic T-lymphocyte-associated protein 4 (CTLA-4). CTLA-4 is a negative regulator of T-cell activity. By binding to CTLA-4, ipilimumab blocks the interaction of CTLA-4 with its ligands, CD80/CD86. Blockade of CTLA-4 has been shown to augment T-cell activation and proliferation, including the activation and proliferation of tumor infiltrating T-effector cells. Inhibition of CTLA-4 signaling can also reduce T-regulatory cell function, which may contribute to a general increase in T cell responsiveness, including the anti-tumor immune response.

In some embodiments, the ipilimumab dose is 3 mg/kg administered as an IV infusion on Day 1 of each 28-day cycle for a maximum of 4 doses.

There is potential for overlapping toxicities between Compound 1 and ipilimumab. Liver function abnormalities have been observed in pre-clinical toxicity studies with Compound 1; immune-mediated hepatitis has been observed with ipilimumab treatment.

Docetaxel

Docetaxel is an antineoplastic agent belonging to the taxoid family. It is prepared by a semi-synthesis beginning with a precursor extracted from the renewable needle biomass of yew plants. The chemical name for docetaxel is (2R,3S)-N-carboxy-3-phenylisoserine,N-*tert*-butyl ester, 13-ester with 5b-20-epoxy-1,2a,4,7b, 10b, 13a-hexahydroxytax-11-en-9-one 4-acetate 2- benzoate, trihydrate.

In some embodiments, the docetaxel is administered as an IV infusion on days 1, 8, and 15 of each 28-day cycle. In some embodiments, the docetaxel dose was 35 mg/m². In some embodiments, 28 mg/m² is the docetaxel dose.

There is potential for overlapping toxicities between Compound 1 and docetaxel. Specifically, a DLT of Grade 4 neutropenia has been seen on Compound 1 single agent therapy and myelosuppression may be observed with docetaxel treatment. Liver function abnormalities have been observed in pre-clinical toxicity studies with Compound 1; hepatotoxicity has been observed with docetaxel treatment.

Pembrolizumab

Pembrolizumab is a programmed death receptor-1 (PD 1)-blocking antibody. Pembrolizumab is a humanized monoclonal IgG4 kappa antibody with an approximate molecular weight of 149 kDa. Pembrolizumab is produced in recombinant Chinese hamster ovary (CHO) cells.

In some embodiments, pembrolizumab is administered as a dose of 200 mg using a 30 minutes IV infusion on Day 1 of each 21 days cycle after all procedures and assessments have been completed and prior to the administration of other drugs, and with a gap of 30 minutes between the administration of next drug.

Carboplatin

The chemical name for carboplatin, USP is platinum, diamine[1,1-cyclobutanedicarboxylato(2-)-O,O']-, (SP-4-2). Carboplatin, USP is a crystalline powder. It is soluble in water at a concentration of approximately 14 mg/mL, and the pH of a 1% solution is 5–7. It is virtually insoluble in ethanol, acetone, and dimethylacetamide. Carboplatin produces predominantly interstrand DNA cross-links rather than DNA-protein cross-links. This effect is apparently cell cycle nonspecific. Carboplatin induce equal numbers of drug-DNA cross-links, causing equivalent lesions and biological effects.

In some embodiments, the initial dose of carboplatin injection is determined by the use of mathematical formulae, which is based on a subject's pre-existing renal function or renal function and desired platelet nadir (as renal excretion is the major route of elimination for carboplatin). The use of dosing formulae, as compared to empirical dose calculation based on body surface area, allows compensation for subject variations in pretreatment renal function that might otherwise result in either underdosing (in subjects with above average renal function) or overdosing (in subjects with impaired renal function).

A simple formula for calculating dosage, based upon a subject's glomerular filtration rate (GFR in mL/min) and carboplatin injection target area under the concentration versus time curve (AUC in mg/mL•min), has been proposed by Calvert. In these studies, GFR was measured by Cr-EDTA clearance. The Calvert formula for carboplatin dosing is as follows:

$$\text{Total Dose (mg)} = (\text{target AUC}) \times (\text{GFR} + 25)$$

Note that with this formula, the total dose of carboplatin is calculated in mg, not mg/m². The target AUC of 4 mg/mL•min to 6 mg/mL•min using single agent carboplatin appears to provide the most appropriate dose range in previously treated subjects. This study also showed a trend between the AUC of single agent carboplatin administered to previously treated subjects and the likelihood of developing toxicity.

5

Pemetrexed

Pemetrexed (for injection) is a folate analog metabolic inhibitor. The drug substance, pemetrexed disodium heptahydrate, has the chemical name L-glutamic acid, *N*-[4-[2-(2-amino-4,7-dihydro-4-oxo-1*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)ethyl]benzoyl]-, disodium salt, heptahydrate with a molecular formula of

10 C₂₀H₁₉N₅Na₂O₈•7H₂O and a molecular weight of 597.49 g/mol.

Pemetrexed exerts its antineoplastic activity by disrupting the folate-dependent metabolic processes essential for cell replication. *In vitro* studies have shown that pemetrexed behaves as a multitargeted antifolate by inhibiting thymidylate synthase (TS), dihydrofolate reductase (DHFR), and glycylamide ribonucleotide formyltransferase (GARFT) which are crucial for the *de novo* biosynthesis of

15 thymidine and purine nucleotides. Polyglutamated metabolites of pemetrexed with prolonged intracellular half-life result in prolonged pemetrexed drug action in malignant cells.

In some embodiments, the pemetrexed dose is 500 mg/m² on Day 1 of each 21-day cycle for a maximum of 4 cycles. In some embodiments, subjects treated with pemetrexed must be instructed to take folic acid and vitamin B12 as a prophylactic measure to reduce treatment-related hematologic and GI

20 toxicity. In some embodiments, subjects may also be prescribed with corticosteroids to take 2 times a day for 3 days, beginning the day before each treatment with pemetrexed.

There is potential for overlapping toxicities between Compound 1, pemetrexed, and carboplatin. Specifically, a DLT of Grade 4 neutropenia has been seen on Compound 1 single agent therapy and myelosuppression may be observed with pemetrexed combination treatment with Compound 1 and

25 carboplatin.

Adverse Events

An adverse event (AE) is any untoward medical occurrence in a subject or clinical investigation subject administered a pharmaceutical product, and which does not necessarily have to have a causal

30 relationship with this treatment. An AE can therefore be any unfavorable and unintended sign (including abnormal laboratory findings), symptom, or disease temporally associated with the use of an investigational product, whether or not related to the investigational product.

Death and progressive disease (PD) are not considered AEs. Death is considered an outcome of one or more primary AEs, and PD is considered a worsening of underlying disease. Preexisting

35 conditions (present before the start of the AE collection period) are considered concurrent medical conditions and not AEs. However, a worsening or complication of such a concurrent condition, the worsening or complication is an AE.

An AE or suspected adverse reaction is considered serious if it results in death; is life threatening, i.e., the subject was at immediate risk of death from the reaction as it occurred but does not include a

40 reaction which hypothetically might have caused death had it occurred in a more severe form; requires in-

subject hospitalization or prolongation of existing hospitalization; results in persistent or significant disability/incapacity; is a congenital anomaly/birth defect; or is an important medical event.

Methods of Treatment

5 The methods described here can be used to treat cancer.

 Treating cancer can result in a reduction in size or volume of a tumor. For example, after treatment, tumor size is reduced by 5% or greater (e.g., 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or greater) relative to its size prior to treatment. Size of a tumor may be measured by any reproducible means of measurement. The size of a tumor may be measured as a diameter of the tumor
10 or by any reproducible means of measurement.

 Treating cancer may further result in a decrease in number of tumors. For example, after treatment, tumor number is reduced by 5% or greater (e.g., 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or greater) relative to number prior to treatment. Number of tumors may be measured by any reproducible means of measurement. The number of tumors may be measured by counting tumors
15 visible to the naked eye or at a specified magnification (e.g., 2x, 3x, 4x, 5x, 10x, or 50x).

 Treating cancer can result in a decrease in number of metastatic nodules in other tissues or organs distant from the primary tumor site. For example, after treatment, the number of metastatic nodules is reduced by 5% or greater (e.g., 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or greater) relative to number prior to treatment. The number of metastatic nodules may be measured by any
20 reproducible means of measurement. The number of metastatic nodules may be measured by counting metastatic nodules visible to the naked eye or at a specified magnification (e.g., 2x, 10x, or 50x).

 Treating cancer can result in an increase in average survival time of a population of subjects treated according to the present invention in comparison to a population of untreated subjects. For example, the average survival time is increased by more than 30 days (more than 60 days, 90 days, or
25 120 days). An increase in average survival time of a population may be measured by any reproducible means. An increase in average survival time of a population may be measured, for example, by calculating for a population the average length of survival following initiation of treatment with the compound of the invention. An increase in average survival time of a population may also be measured, for example, by calculating for a population the average length of survival following completion of a first
30 round of treatment with the compound of the invention.

 Treating cancer can also result in a decrease in the mortality rate of a population of treated subjects in comparison to an untreated population. For example, the mortality rate is decreased by more than 2% (e.g., more than 5%, 10%, or 25%). A decrease in the mortality rate of a population of treated subjects may be measured by any reproducible means, for example, by calculating for a population the
35 average number of disease-related deaths per unit time following initiation of treatment with the compound of the invention. A decrease in the mortality rate of a population may also be measured, for example, by calculating for a population the average number of disease-related deaths per unit time following completion of a first round of treatment with the compound of the invention.

 Treating cancer can also result in an increased average progression-free survival time of a
40 population of treated subjects in comparison to an untreated population. For example, the average progression-free survival time is increased by more than 30 days (more than 60 days, 90 days, or 120

days). An increase in average progression-free survival time of a population may be measured by any reproducible means. An increase in average progression-free survival time of a population may be measured, for example, by calculating for a population the average length of progression-free survival following initiation of treatment with the compound of the invention. An increase in average progression-free survival time of a population may also be measured, for example, by calculating for a population the average length of progression-free survival following completion of a first round of treatment with the compound of the invention.

Compositions

Within the scope of this invention is a composition that contains a suitable carrier and one or more of the therapeutic agents described above.

As described above, the pharmaceutical compositions of the present invention additionally include a pharmaceutically acceptable excipient, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, and lubricants, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various excipients used in formulating pharmaceutical compositions and known techniques for the preparation thereof. Except insofar as any conventional excipient is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated to be within the scope of this invention.

The above-described compositions, in any of the forms described above, can be used for treating cancer, or any other disease or condition described herein. An effective amount refers to the amount of an active compound/agent that is required to confer a therapeutic effect on a treated subject. Effective doses will vary, as recognized by those skilled in the art, depending on the types of diseases treated, route of administration, excipient usage, and the possibility of co-usage with other therapeutic treatment.

A pharmaceutical composition of this invention can be administered parenterally, orally, nasally, rectally, topically, or buccally. The term "parenteral" as used herein refers to subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional, or intracranial injection, as well as any suitable infusion technique.

A sterile injectable composition can be a solution or suspension in a non-toxic parenterally acceptable diluent or solvent.

A composition for oral administration can be any orally acceptable dosage form including capsules, tablets, emulsions and aqueous suspensions, dispersions, and solutions. When aqueous suspensions or emulsions are administered orally, the active ingredient can be suspended or dissolved in an oily phase combined with emulsifying or suspending agents. If desired, certain sweetening, flavoring, or coloring agents can be added.

Pharmaceutical compositions for topical administration according to the described invention can be formulated as solutions, ointments, creams, suspensions, lotions, powders, pastes, gels, sprays, aerosols, or oils. Alternatively, topical formulations can be in the form of patches or dressings impregnated with active ingredient(s), which can optionally include one or more excipients or diluents. In

some preferred embodiments, the topical formulations include a material that would enhance absorption or penetration of the active agent(s) through the skin or other affected areas.

Combination Therapies

5 In some embodiments of the methods described herein, the pharmaceutical composition may further include an additional compound having antiproliferative activity.

 It will also be appreciated that the compounds and pharmaceutical compositions of the present invention can be formulated and employed in combination therapies, that is, the compounds and pharmaceutical compositions can be formulated with or administered concurrently with, prior to, or
10 subsequent to, one or more other desired therapeutics or medical procedures (e.g., surgery and/or radiation therapy). The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder, or they may achieve different effects (e.g., control of
15 any adverse effects).

EXAMPLES

Example 1. Methods

X-ray Powder Diffraction (XRPD)

20 Method 1

 XRPD analysis was carried out on a PANalytical X'pert pro, scanning the samples between 3 and 35 °2θ. The material was gently ground to release any agglomerates and loaded onto a multi-well plate with Mylar polymer film to support the sample. The multi-well plate was then placed into the diffractometer and analyzed using Cu K radiation ($\alpha_1 \lambda = 1.54060 \text{ \AA}$; $\alpha_2 = 1.54443 \text{ \AA}$; $\beta = 1.39225 \text{ \AA}$; $\alpha_1 : \alpha_2 \text{ ratio} = 0.5$) running in transmission mode (step size 0.0130 °2θ) using 40 kV / 40 mA generator settings.
25

Method 2

 X-ray powder diffraction patterns were obtained using a Bruker D8 Advance equipped with a Cu K α radiation source ($\lambda = 1.54 \text{ \AA}$), a 9-position sample holder and a LYNXEYE super speed detector.
30 Samples were placed on zero-background, silicon plate holders.

Polarized Light Microscopy (PLM)

Method 1

 The presence of birefringence was determined using an Olympus BX53 polarizing microscope, equipped with a Motic camera and image capture software (Motic Images Plus 3.0). All images
35 were recorded using the 10 × or the 20 × objective.

Method 2

 Samples were analyzed using an Olympus BX53 polarized light microscope equipped with a PAXcam 3 digital microscope camera.
40

Thermogravimetric/Differential Thermal Analysis (TG/DTA)

Method 1

Approximately 5 mg of material was weighed into an open aluminum pan and loaded into a simultaneous thermogravimetric/differential thermal analyzer (TG/DTA) and held at room temperature.

- 5 The sample was then heated, under a nitrogen purge, at a rate of 10 °C/min from 20 °C to 300 °C during which time the change in sample weight was recorded along with any differential thermal events (DTA).

Method 2

- 10 TGA data were collected using a TA Instruments TGA Q500. Typically, samples (~10 mg) were placed in an open, pre-tared aluminum sample pan and scanned from 25 to 300 °C at a rate of 10 °C/min using a nitrogen purge at 60 mL/min.

Differential Scanning Calorimetry (DSC)

Method 1

- 15 Approximately 5 mg of material was weighed into an aluminum DSC pan and sealed with an aluminum lid. The sample pan was then loaded into a TA Instruments Discovery DSC 2500 differential scanning calorimeter equipped with a RC90 cooler. Once a stable heat-flow response was obtained, the sample and reference were heated/cooled, under a nitrogen purge, by using the following temperature programs and the resulting heat flow response monitored.

Program 1

- 20 • Heated 20 °C to 260 °C at 10 °C/minute held at 260 °C for 3 minutes;
• Cooled 260 °C to 20 °C at 10 °C/minute, held at 20 °C for 3 minutes;
• Heated 20 °C to 260 °C at 10 °C/minute held at 260 °C for 3 minutes;
• Cooled 260 °C to 20 °C at 10 °C/minute, held at 20 °C for 3 minutes.

Program 2

- 25 • Heated 20 °C to 270 °C at 10 °C/minute held at 270 °C for 3 minutes;
• Cooled 270 °C to 20 °C at 10 °C/minute, held at 20 °C for 3 minutes;
• Heated 20 °C to 270 °C at 10 °C/minute held at 270 °C for 3 minutes;
• Cooled 270 °C to 20 °C at 10 °C/minute, held at 20 °C for 3 minutes.

Method 2

- 30 DSC data were collected using a TA Instruments Q10 DSC. Typically, samples (2-8 mg) were placed in unsealed but covered hermetic alodined aluminum sample pans and scanned from 30 to 300 °C at a rate of 10 °C/min under a nitrogen purge of 50 mL/min.

¹H Nuclear Magnetic Resonance Spectroscopy (¹H-NMR)

- 35 Method 1

¹H-NMR spectroscopic experiments were performed on a Bruker AV500 (frequency: 500 MHz). Experiments were performed in CDCl₃ and each sample was prepared to about 10 mM concentration.

Method 2

Samples were prepared by dissolving the compound in deuterated dimethylsulfoxide with 0.05% (v/v) tetramethylsilane (TMS). Spectra were collected at ambient temperature on a Bruker Avance 300 MHz NMR with TopSpin software. The number of scans was 16 for proton NMR.

5

¹⁹F Nuclear Magnetic Resonance Spectroscopy (¹⁹F-NMR)

¹⁹F-NMR spectroscopic experiments were performed on a Bruker AV500 (frequency: 470 MHz). Experiments were performed in CDCl₃ and each sample was prepared to about 10 mM concentration.

10

High Performance Liquid Chromatography-Ultraviolet Detection (HPLC-UV)

The HPLC method uses a C18 column with an acetonitrile/water/trifluoroacetic acid gradient.

Dynamic Vapor Sorption (DVS)

15

Samples are analyzed using an Aquadyne DVS-2 gravimetric water sorption analyzer. The relative humidity is adjusted between 2-95% and the weight of the sample is continuously monitored and recorded.

Karl Fisher (KF)

20

The apparent water content in samples is determined by Karl Fischer titration using a Mettler Toledo DL39 Coulometric KF Titrator. HYDRANAL-Coulomat AD was used as the titrant. About 20 mg of the solid is used for titration. The analytical parameters are shown in Table 1 below:

Table 1

KF Parameter	Value
Speed [%]	40
Mix time [sec]	10
Auto start	No
Blank [µg]	0
Drift [µg/min]	5
Calculation	Ug
Standby	Yes
Initial drift [µg/min]	<10
Initial Potential [mV]	100

25

Fourier-Transform Infrared Spectroscopy (FTIR)

FTIR analysis was carried out on a Thermo Scientific, NICOLET™ IS™ 10 FTIR Spectrometer with Attenuated Total Reflectance (ATR). Sufficient material was placed on the center of the plate of the spectrometer and the spectra were obtained.

Example 2. Salt Screening

Study 1

A first study was conducted to identify salts of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]-methyl-(2,2-diphenylmethyl)amino]butoxy]phenyl]acetic acid. 26 counterion sources were selected for salt screening of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]-methyl-(2,2-diphenylmethyl)amino]butoxy]phenyl]acetic acid. The counterion sources are listed in Table 2 below:

Table 2: List of selected counterion sources for salt screening experiments in Study 1

Counterion Source	No.
Hydrochloric Acid (HCl)	1
Sulfuric acid (H ₂ SO ₄)	2
Benzenesulfonic acid (C ₆ H ₅ O ₃ SH)	3
Phosphoric acid (H ₃ PO ₄)	4
Sodium hydroxide (NaOH)	5
Potassium hydroxide (KOH)	6
Calcium hydroxide (Ca(OH) ₂)	7
Maleic acid	8
Fumaric acid	9
L-Arginine	10
L-Histidine	11
L-Malic acid	12
Diethanolamine	13
Salicylic acid	14
Benzoic acid	15
DL-Mandelic acid	16
Ethylenediamine	17
Succinic acid	18
L-Tartaric acid	19
L-Aspartic acid	20
L-Lysine	21
L-Ascorbic acid	22
Citric acid	23
Naphthalene-2-sulfonic acid	24
1,2-Ethane-disulfonic acid	25
Hydroiodic acid	26

Among the counterion sources investigated, only three (hydrochloride, DL-mandelic acid, and naphthalene sulfonic acid) afforded crystalline salts, and reproduction and scaling up of the three crystalline salts was successful only for the hydrochloride salt.

Subsequent studies revealed that the hydrochloride salt, as well as other salts prepared from strong acids, e.g., H₂SO₄, HBr, p-toluenesulfonic acid, and methanesulfonic acid, were found to be unstable. In particular, these salts react with excipients and/or components of a formulation. This discovery led to investigation into the use of weak acids, e.g., oleic acid, octanoic acid, and acetic acid, as the counterion source, none of which formed salts with 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]-methyl-(2,2-diphenylmethyl)amino]butoxy]phenyl]acetic acid.

Study 2

A second study was conducted to identify salts that afforded both improved processability and stability. The 34 counterion sources and 12 solvent conditions investigated in this study are summarized in Tables 3 and 4 below:

Table 3: List of selected counterion sources for salt screening experiments in Study 2

Counterion Source	
Hydriodic acid	EDTA
Naphthalene-2-sulfonic acid	Ethane sulfonic acid
1,2-Ethane disulfonic acid	Pamoic acid
Saccharin	Mucic acid
L-Arginine	Gentistic acid
N-Methylglucamine	Pyroglutamic acid
2-Hydroxyethanesulfonic acid	Ethylenediamine
Ethanolamine	Benzathine
Procaine hydrochloride salt	2-Diethylaminoethanol
Piperazine	Diethanolamine
Triethanolamine	Ammonium hydroxide
Diethylamine	Tromethamine
Choline hydroxide	Benethamine
L-Lysine monohydrate	Calcium hydroxide
Zinc chloride	Magnesium hydroxide
1,5-Naphthalene disulfonic acid	Aluminum hydroxide
Nitric acid	Diethylaminoethanol

Table 4: List of selected solvent conditions for salt screening experiments in Study 2

Solvent and Solvent Mixtures
Isopropanol (IPA)
Tetrahydrofuran (THF)
Dichloromethane (DCM)
Ethyl acetate (EtOAc)
Heptane (Hep)
Tert-Butylmethyl ether (TBME)
Water (H ₂ O)
1,4-Dioxane
Acetone
THF:H ₂ O (9:1)
Acetone:H ₂ O (9:1)
IPA:H ₂ O (95:5)

An initial solubility assessment in 12 solvents/solvent mixtures using amorphous Compound 1. Dissolution was observed in most of the samples. Salt screening by cooling, anti-solvent addition, maturation, liquid assisted grinding, neat grinding and evaporation were all investigated. Gums isolated were subjected to heat/cool maturation cycles (room temperature / 50 °C), sonication, prolonged storage under vacuum, prolonged storage at 40 °C / 75% RH, trituration with anti-solvent and liquid assisted grinding (LAG) with water. From extensive salt screening, eight hits were obtained. Amorphous hits with a high glass transition temperature (T_g) were identified from zinc chloride, naphthalene-2-sulfonic acid, 1,5-naphthalene disulfonic acid, and ethane sulfonic acid. One crystalline hit was identified from ethane sulfonic acid. Poorly crystalline material was isolated from calcium hydroxide, magnesium hydroxide and aluminum hydroxide.

As limited characterization was performed on the isolated hits, focused screens were conducted on the most promising counterions: calcium hydroxide, aluminum hydroxide, ethane sulfonic acid and naphthalene-2-sulfonic acid. Calcium hydroxide and aluminum hydroxide were selected to assess if the initial hits isolated via counterion exchange were single metal or mixed metal salts. Solids isolated from the additional screening were consistent with the initial hits. From additional screening, these samples were likely to be single metal salts. As these hits were not mixed salts, a limited mixed metal screen was performed using magnesium hydroxide and calcium hydroxide. These attempts were unsuccessful. Ethane sulfonic acid and naphthalene-2-sulfonic acid were selected for further screening. However, the initial hits were not reproduced in the further screening attempts. Amorphous solids were isolated, however, the T_g was lower than the T_g of both the free compound (32 °C) and HCl salt (~70-75 °C).

The zinc salt was isolated using the following procedure:

Compound 1 (~ 250 mg) was dissolved in THF (5 vol) at RT. The solution was dispensed into HPLC vials (75 µl, ca. 15 mg in solution). The solution was held at 50 °C for 5 minutes, observations were recorded. The sample was treated with 1.0 eq. of sodium then the selected equivalent of counter-
 5 ion (0.5 eq. of zinc chloride. The sample was held at 50 °C for 10 minutes then cooled to 5 °C at 0.1 °C/min. The resulting solution was held at 5 °C overnight then stored at -20 °C. After 5 days, the sample remained as a solution and was allowed to evaporate at ambient conditions. A gum was obtained post evaporation and was treated with 10 vol of heptane, followed by sonication for 2 hours. The solid collected was analyzed by XRPD and found to be amorphous.

10 None of the salts isolated from this study were recommended for development due to long, complex isolation procedures and the propensity of Compound 1 to form gums. However, several salts, e.g., zinc and aluminum, gave encouraging results that suitable salts might be possible and prompted further investigation.

15 Study 3

A third salt screening study was conducted. Salt formation reactions were attempted at a 100 mg scale using bases with approximate pK_a greater than 8 or acceptable strong acids with a pK_a below 4.

In a first set of salt screening experiments, the counterion sources shown in Table 5 below were investigated.

20 Table 5: List of selected counterion sources for the first set of salt screening experiments in Study 3

Counterion Source	Molecular Weight (g/mol)	pK_{a1}
Ethane-1,2-disulfonic acid	190.20	-2.1
Naphthalene-1,5-disulfonic acid	332.26	-3.4
Ethanesulfonic acid	110.13	-1.6
Nitric acid	63.02	-1.3
Naphthalene-2-sulfonic acid	208.24	0.17
Zinc hydroxide	99.38	14
Choline	121.18	>11.0
Benzathine	240.35	10.0
Benthamine	211.30	9.4
Ethanolamine	61.80	9.6
Diethanolamine	105.14	9.5
L-Arginine	174.20	9.0

Briefly, 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylmethyl)amino]-butoxy]phenyl]acetic acid was suspended/dissolved in six different solvent systems—isopropanol,
 25 acetone, cyclohexanone, methyl ethyl ketone, ethyl acetate, and THF. The corresponding counterion source (1.0 equivalents) was then added as a solution (where possible) or via neat addition. Solutions were temperature cycled between two specific temperatures (e.g., 25 and 40°C). All experiments were monitored for precipitation/crystallization. In vials where crystallization/precipitation was not evident, cooling to sub-ambient temperatures or iterative anti-solvent addition using a suitable anti-solvent was
 30 performed, with or without cooling to sub-ambient temperatures (e.g. 5°C). In vials where crystallization/precipitation was not evident following cooling and/or addition of anti-solvent, the solvent

was removed by evaporation. Any solids were isolated by filtration and analyzed by XRPD. Oils or gums were predominantly obtained.

A second salt screen was conducted with the counterion sources shown in Table 6 below.

Table 6: List of selected counterion sources for the second set of salt screening experiments in

5 Study 3

Counterion Source	Molecular Weight (g/mol)	pKa ₁
Sodium edisylate	234.16	-2.1
Thiocyanic acid	59.09	-1.3
Diethylsulfosuccinic acid	422.16	-0.8
Sodium napsylate	230.22	0.2
Dichloroacetic acid	128.94	1.3
Sodium isethionate	148.11	1.7
Pamoic acid	388.38	2.5
Hippuric acid	179.18	3.6
L-Lysine	146.19	9.2
Histidine	155.15	9.2
1-(2-Hydroxyethyl)-pyrrolidine	115.18	9.4

The procedure for preparing the salts with the counterion sources of Table 5 is as follows. 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylmethyl)amino]butoxy]-phenyl]acetic acid was produced from its hydrochloride salt as per known procedures (Collins et al. J. Med. Chem., 2002, 45:1963-1966). 100 mg of the free material was transferred into 20 mL vials. Ethyl acetate (1 mL) was added to each vial, followed by addition 1.0 eq. of each counterion source. The samples were temperature cycled between ambient temperature and 40 °C in 4-hour cycles with agitation for 24 hours. Heptane was added as an antisolvent to any clear solutions and the samples were placed in the fridge for 24 hours. All solids thus obtained were analyzed by XRPD and all samples that remained as clear solutions were uncapped and left to evaporate. These counterion sources produced solids identified as the counterion sources or sticky, gummy solids.

Another salt screen was conducted using an aqueous precipitative approach. This salt screen used a metathesis procedure where the 1:1 sodium salt of Compound 1 was prepared and 1.0, 0.5, or 0.3 equivalents of the selected metal counterion was added as its chloride or sulfate salt. Alternatively, free Compound 1 was reacted with the selected metal counterion as its acetate or sulfate salt. These aqueous precipitations produced free flowing white solids from zinc, aluminum, and bismuth.

The counterions shown in Table 7 below were investigated.

Table 7: List of selected counterion sources for the third set of salt screening experiments in Study 3

Counterion Source
Zinc chloride
Bismuth chloride
Zinc acetate
Aluminum sulfate

25 The zinc salt was prepared via metathesis using the following procedure.

Compound 1 (200 mg) was dissolved in methanol (2 mL) to form a mixture. A 1M NaOH solution (336 µL, 1.0 eq.) was then added to the mixture to form the 1:1 sodium salt of Compound 1. Zinc chloride (22.5 mg, 0.5 eq.) was dissolved in water (1 mL), and the resulting solution was added to the mixture, upon which a white solid precipitated out of solution. Additional water (1 mL) was added to the mixture and the reaction vessel was sealed with parafilm, after which the mixture was temperature cycled

between ambient temperature and 40 °C in 4 hour cycles for 24 hours with agitation. The methanol was removed from the mixture via rotary evaporation and water (10 mL) was added to reduce the solubility. The resulting precipitate was isolated via Buchner filtration (Whatman Grade 1 filter paper, Ø = 55 mm) and washed with water (100 mL) to remove any remaining sodium chloride. The solid was dried on the filter paper, yielding a white powder (152.9 mg, 72%).

The material obtained via the above-described salt metathesis procedure was characterized by ICP, XRPD (FIG. 1), PLM (FIG. 2), TG/DTA (FIG. 3), DSC (FIG. 4), ¹H-NMR (FIG. 5), and ¹⁹F-NMR (FIG. 6) following the methods described in Example 1. Characterization of the material yielded the following information:

- The material contained 5.27 wt.% zinc by ICP analysis. Stoichiometric salt is 9.9 wt.%, therefore there is approximately 0.51 eq. zinc (0.5 eq. expected).
- The material contained 0.19 wt.% sodium by ICP analysis. Stoichiometric salt is 3.7 wt.%, therefore approximately 5% of the material is the 1:1 sodium salt, and approximately 95% of the material is the 2:1 zinc salt.
- The material was amorphous by XRPD analysis.
- The material showed some birefringence by PLM analysis, with irregular morphology.
- There was minimal mass loss in the TG trace up to decomposition.
- By DSC, there was an endothermic event with an onset temperature of 51 °C in the first heat cycle, and a sharp event corresponding to glass transition with a midpoint temperature at 52 °C in the second heat cycle.
- The ¹H-NMR spectrum was consistent with Compound 1.
- The ¹⁹F-NMR showed one main peak.
- The purity of the amorphous material, as determined by HPLC analysis, is shown in Table 8 below.

Table 8: Purity and stability of the zinc salt of Compound 1 prepared from zinc chloride

Time (days)	Purity
0	99.3%
7 (stored at 40 °C/ 75% RH)	99.5%
7 (stored at 80 °C)	96.2%

As shown in Table 8, the zinc salt prepared by the method described above is stable after one week of storage at 40 °C/ 75% RH and at 80 °C.

The zinc salt was also prepared using zinc acetate as the counterion source. Briefly, Compound 1 (78 mg) was dissolved in methanol (1.5 mL), and zinc acetate was dissolved in methanol/water (9:1; 1.5 mL), after which the two solutions thus obtained were combined to form a mixture. A white solid immediately formed and precipitated from solution. The sample was then cycled between ambient temperature and 40 °C for approximate 5 hours, after which a small amount of white solid remained undissolved. The solvent was then removed via rotary evaporation, and methanol (5 mL) was added to the solid. The cycling and solvent evaporation steps were repeated, then methanol/water (9:1; 5 mL) was added to the sample, and the cycling and solvent evaporation steps were repeated once more. The

sample was subsequently frozen lyophilized overnight to remove water, after which a white solid was obtained.

The aluminum salt was prepared via metathesis using the following procedure. Compound 1 (200 mg) was dissolved in methanol (2 mL). The sodium salt was prepared by adding a 1M NaOH stock solution (336 μ L, 1.0 eq.). The methanol was removed via rotary evaporation. Aluminum sulfate and water (2 mL) were added and the sample temperature cycled between ambient and 40 °C in 4-hour cycles for 24 hours. The solid was isolated via centrifugation. Water (3 mL) was added and the slurry agitated at ambient temperature for 18 hours. The solid was filtered using a Buchner funnel (Whatman Grade 1 filter paper, $\varnothing$ = 42.5 mm) and washed with water (100 mL) before being dried on the filter paper for 10 minutes. The solid was dried under vacuum at ambient temperature for ca. 2 hours, yielding a white powder (128.7 mg, 61%).

Example 3. FTIR Analysis

Study

Stock solutions of the free compound and zinc salt of Compound 1 in CHCl_3 were prepared. The free form of Compound 1 (50 mg) was diluted to 10 mL with CHCl_3 in a volumetric flask. The zinc salt of Compound 1 (50 mg) was also diluted to 10 mL with CHCl_3 in a separate volumetric flask. A FTIR spectrum was taken of the free form of Compound 1 (see FIG. 7), and of the zinc salt form of Compound 1 (see FIG. 9) according to the FTIR method described in Example 1. FTIR spectra were then taken of samples with increasing concentrations of the zinc salt of Compound 1, such as:

- a 1:1 (w/w) mixture of the free form: zinc salt of Compound 1 (mixture of the free form stock solution (500 μ L) with the zinc salt stock solution (500 μ L)) (see FIG. 8);
- a 1:4 (w/w) mixture of the free form: zinc salt of Compound 1 (mixture of the free form stock solution (200 μ L) with the zinc salt stock solution (800 μ L));
- a 1:9 (w/w) mixture of the free form: zinc salt of Compound 1 (mixture of the free form stock solution (100 μ L) with the zinc salt stock solution (900 μ L));
- and a 5:95 (w/w) mixture of the free form: zinc salt of Compound 1 (mixture of the free form stock solution (50 μ L) with the zinc salt stock solution (950 μ L)).

The spectra were compared with each other as well as FTIR spectra of ZnO (FIG. 10) and ZnCl_2 (FIG. 11).

The free form of Compound 1 exhibited a distinct, characteristic peak at about 1710 cm^{-1} (see FIG. 7) which is not present in the zinc salt spectrum (see FIG. 9). The peak at 1710 cm^{-1} was observed to shrink as the percentage of the free form decreased. Additionally, a broad peak at about 1590 cm^{-1} was observed with low intensity in the free compound, but increased in intensity as the concentration of the zinc salt increased.

The zinc salt is evidenced by the disappearance of the IR peak at 1710 cm^{-1} , presumably, due to coordination of the carboxylate carbonyl with zinc. Furthermore, the zinc salt does not contain peaks found in the IR spectrum of ZnO (characterized by a distinct, large, broad band starting at 605 cm^{-1} to <

500 cm^{-1}) or ZnCl_2 (characterized by a distinct broad peak at 3386 cm^{-1}), thus further demonstrating the zinc salt is not simply a mixture of the free compound and inorganic zinc.

Other Embodiments

5 All literature and similar material cited in this application, including, but not limited to, patents, patent applications, articles, books, treatises, and web pages, regardless of the format of such literature and similar materials, are expressly incorporated by reference in their entirety. In the event that one or more of the incorporated literature and similar materials differs from or contradicts this application, including but not limited to defined terms, term usage, described techniques, or the like, this application
10 controls.

 While the methods have been described in conjunction with various embodiments and examples, it is not intended that the methods be limited to such embodiments or examples. On the contrary, the present disclosure encompasses various alternatives, modifications, and equivalents, as will be appreciated by those of skill in the art.

15 While the methods have been particularly shown and described with reference to specific illustrative embodiments, it should be understood that various changes in form and detail may be made without departing from the spirit and scope of the present disclosure. Therefore, all embodiments that come within the scope and spirit of the present disclosure, and equivalents thereto, are intended to be claimed. The claims, descriptions and diagrams of the methods, systems, and assays of the present
20 disclosure should not be read as limited to the described order of elements unless stated to that effect. This application claims the benefit of U.S. provisional serial no. 62/947,968, filed December 13, 2019, which is incorporated herein in its entirety.

CLAIMS

What is claimed is:

- 5 1. A zinc salt of 2-[3-[(3*R*)-3-[[2-chloro-3-(trifluoromethyl)phenyl]methyl-(2,2-diphenylethyl)amino]butoxy]phenyl]acetic acid (Compound 1).
2. The zinc salt of claim 1, wherein the zinc salt is a 2:1 (Compound 1:zinc) salt.
- 10 3. The zinc salt of claim 1 or 2, wherein the zinc salt is amorphous.
4. The zinc salt of any one of claims 1 to 3 having less than 1% mass loss up to decomposition as measured by thermal gravimetric analysis.
- 15 5. A pharmaceutical composition comprising the zinc salt of any one of claims 1 to 4 and a pharmaceutically acceptable excipient.
6. The pharmaceutical composition of claim 5, wherein the pharmaceutical composition comprises less than 1.5 weight % of sodium.
- 20 7. The pharmaceutical composition of claim 6, wherein the pharmaceutical composition is substantially free of the 1:1 sodium salt of Compound 1.
8. The pharmaceutical composition of any one of claims 5 to 7, wherein the pharmaceutical composition is in unit dosage form.
- 25 9. Use of the zinc salt of any one of claims 1 to 4 or the pharmaceutical composition of any one of claims 5 to 8 in the manufacture of a medicament for treating cancer.
- 30 10. The use of claim 9, wherein the cancer is breast cancer, colon cancer, renal cell cancer, lung cancer, hepatocellular carcinoma, gastric cancer, ovarian cancer, pancreatic cancer, esophageal cancer, prostate cancer, sarcoma, bladder cancer, head and neck cancer, glioblastoma, diffuse large B-cell lymphoma, leukemia, or melanoma.
- 35 11. The use of claim 10, wherein the cancer is metastatic cancer.
12. The use of claim 11, wherein the medicament comprises an amount effective to suppress metastatic colonization of said cancer.
- 40 13. A method of producing a zinc salt of Compound 1, the method comprising combining Compound 1, or a salt thereof, and a zinc salt under conditions sufficient to produce the zinc salt of Compound 1.

14. The method of claim 13, wherein the zinc salt of Compound 1 is a 2:1 (Compound 1:zinc) salt.

5 15. The method of claim 13 or 14, wherein the Compound 1, or salt thereof, is a 1:1 (Compound 1:sodium) salt.

16. The method of any one of claims 13 to 15, wherein the zinc salt is zinc chloride.

10 17. The method of any one of claims 13 to 15, wherein the zinc salt is zinc acetate.

18. The method of claim any one of claims 13 to 17, wherein the method comprises dissolving the Compound 1, or salt thereof, and the zinc salt in a solvent to form a mixture.

15 19. The method of claim 18, wherein the method further comprises cycling the temperature of the mixture between ambient temperature and 40 °C.

20. The method of claim 19, wherein the cycling is performed in 4-hour cycles.

20 21. The method of claim 20, wherein the cycling is performed for 24 hours.

22. The zinc salt of claim 3, wherein the zinc salt has a peak with increased intensity relative to the free acid at about $1590 \pm 10 \text{ cm}^{-1}$ and a peak with reduced intensity relative to the free acid at about $1710 \pm 10 \text{ cm}^{-1}$ as measured by Fourier-transform infrared spectroscopy (FTIR).

25