

CHINESE HAMSTER OVARY CELL LINES

Reference is made to US application serial number 61/098,270 filed September 19, 2008 and US application serial number 61/183,647 filed June 3, 2009, in which inventor Song is inventor.

5 The foregoing applications, and each document cited or referenced in each of the present and foregoing applications, including during the prosecution of each of the foregoing applications ("application and article cited documents"), and any manufacturer's instructions or catalogues for any products cited or mentioned in each of the foregoing applications and articles and in any of the application and article cited documents, are hereby incorporated
10 herein by reference. Furthermore, all documents cited in this text, and all documents cited or reference in documents cited in this text, and any manufacturer's instructions or catalogues for any products cited or mentioned in this text or in any document hereby incorporated into this text, are hereby incorporated herein by reference. Documents incorporated by reference into this text or any teachings therein may be used in the practice of this invention. Documents
15 incorporated by reference into this text are not admitted to be prior art.

FIELD

This invention relates to the fields of biotechnology and molecular biology. The invention in particular relates to Chinese Hamster Ovary cell lines and their use in recombinant protein expression.

20 **BACKGROUND**

Proteins are useful in a variety of diagnostic, pharmacologic, agricultural, nutritional, and research applications. Given the high cost of producing proteins, especially therapeutic proteins, even small increases in the efficiency of production or in the function and stability of a protein can be valuable.

25 The function and stability, and hence the utility, of a protein can be affected by the post-translational addition of sugar residues to the protein to form a glycoprotein. For example, the addition of terminal sialic acid residues to polysaccharides attached to a glycoprotein generally increases the protein's lifetime in the bloodstream and can, in particular cases, also affect solubility, thermal stability, resistance to protease attack, antigenicity, and

specific activity of some glycoproteins. See e. g. Gu and Wang (1998), *Biotechnol. and Bioeng.* 58 (6): 642-48; Morell et al. (1968), *J. Biol. Chem.* 243 (1) : 155-59.

Recombinant glycoprotein proteins and drugs produced by cell lines, such as Chinese hamster ovary (CHO) cells, generally consist of differentially sialylated isoforms. Poorly sialylated isoforms have shorter circulatory half-life and are thus less efficacious.

It is therefore desirable to increase the sialic acid content of a glycoprotein, especially a glycoprotein to be used for pharmacologic applications. Indeed, one of the major research focuses in the biotechnology field has been how to increase sialylation of recombinant protein drugs.

10 SUMMARY

According to a 1st aspect of the present invention, we provide a Chinese Hamster Ovary (CHO) cell which is capable of higher protein sialylation compared to a wild type Chinese Hamster Ovary cell, in which the CHO cell is obtainable by selection with *Ricinus communis* agglutinin I (RCA-I). The CHO cell may be capable of higher sialylation in the presence of functional GnT 1.

The CHO cell may comprise a mutation in the GnT 1 gene.

The mutation in the GnT 1 gene may comprise one or more of the following mutations at the specified position of a GnT 1 sequence (GenBank accession number: AF343963): (a) a C to T transition at position 1015; (b) a G to C transversion at position 1300; (c) an A to C transversion at position 638; (d) a C to G transversion at position 784; (e) a T to A transversion at position 811; (f) an insertion at position 706 resulting in a frame shift from position 236 of the encoded amino acid sequence; (g) a C to T transition at position 1015; (h) a G to A transition at position 246; (i) a G to A transition at position 258; (j) an A to T transversion at position 859.

The GnT 1 gene may comprise a GnT 1 nucleic acid sequence shown in SEQ ID NO: 1.

The CHO cell may express a GnT 1 protein comprising one or more of the following mutations at the specified position of a GnT 1 sequence (GenBank accession number: AF343963): (a) Ala to Pro at position 434; (b) Asp → Ala at position 213; (c) Arg → Gly at position 262; (d) Trp → Arg at position 271; (e) frame shift from position 236 resulting from

an insertion at position 706 of an encoding GnT 1 nucleic acid sequence; (f) Gln→STOP at position 339; (g) Trp→STOP at position 82; (h) Trp→STOP at position 86; (i) Lys→STOP at position 287.

The CHO cell may express a GnT 1 protein as shown in SEQ ID NO: 2.

- 5 The CHO cell may comprise a nucleic acid sequence encoding a protein of interest such as a recombinant protein of interest. The protein of interest may comprise erythropoietin (EPO) or interferon- γ (IFN- γ).

- 10 The CHO cell may comprise a nucleic acid sequence encoding functional GnT 1. The nucleic acid sequence encoding the protein of interest and the nucleic acid sequence encoding functional GnT 1 may be comprised in one expression vector.

The nucleic acid sequence encoding the protein of interest and the nucleic acid sequence encoding functional GnT 1 may be stably transfected into the CHO cell.

There is provided, according to a 2nd aspect of the present invention, a CHO cell line comprising a CHO cell according to the 1st aspect of the invention.

- 15 The CHO cell line may comprise a JW152 cell line (deposited at ATCC under the Budapest Treaty as accession number PTA-9657), a JW80 cell line, a JW36 cell line, a KFC15002 cell line, a KFC15071 cell line, a KFC5008 cell line, a JW152 cell line, a KFC5026 cell line, a KFC20011 cell line or a KFC15047 cell line.

The CHO cell or a CHO cell line may be adapted to suspension culture.

- 20 We provide, according to a 3rd aspect of the present invention, a recombinant protein expressed by a CHO cell or CHO cell line according to the 1st or 2nd aspect of the invention.

The recombinant protein may comprise erythropoietin (EPO) or interferon- γ (IFN- γ).

- 25 The recombinant protein may comprise a pKa of 4 or less, 3.5 or less, 3 or less, 2.5 or less or 2 or less or a Z-number of greater than 150, such as greater than 160, greater than 170, greater than 180, such as greater than 190, greater than 200, greater than 210, such as greater than 220 or greater than 230, or both.

As a 4th aspect of the present invention, there is provided a method of expressing a recombinant protein, the method comprising introducing a nucleic acid encoding the protein

into a CHO cell as set out above, allowing the protein to be expressed from the CHO cell or a descendent thereof, and optionally purifying the protein.

The method may comprise introducing a nucleic acid encoding functional GnT 1 into the CHO cell.

- 5 The method may comprise introducing an expression vector comprising a nucleic acid sequence encoding the protein and a nucleic acid sequence encoding functional GnT 1.

The method may comprise stably transfecting the nucleic acid sequence encoding the protein of interest and the nucleic acid sequence encoding functional GnT 1.

- 10 We provide, according to a 5th aspect of the present invention, a nucleic acid comprising a sequence shown in SEQ ID NO: 1, or a sequence comprising a GnT 1 sequence together with a mutation set out in Column 2 of Table D1, or a variant, homologue, derivative or fragment thereof.

- 15 The present invention, in a 6th aspect, provides a polypeptide comprising a sequence as shown in SEQ ID NO: 2, or a sequence comprising a GnT 1 sequence together with a mutation set out in Column 3 of Table D1, or a variant, homologue, derivative or fragment thereof.

In a 7th aspect of the present invention, there is provided a method of providing a CHO cell or cell line, the method comprising culturing CHO cells in the presence of *Ricinus communis* agglutinin I (RCA-I) and selecting cells which survive the culture.

- 20 The method may comprise culturing CHO cells in the presence of *Ricinus communis* agglutinin I (RCA-I) and selecting cells which survive the culture.

The method may comprise exposing the CHO cells to RCA-I at a concentration of between 0.1µg/ml to 100µg/ml, for example up to 50 µg/ml or up to 20µg/ml, such as 10µg/ml or 5µg/ml.

- 25 The method may comprise exposing the CHO cells to RCA-I for a period of from an hour, a few hours (such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12), overnight, to a few days, such as 2 days or 3 days, such as overnight.

The method may comprise further comprising selecting cells which do not react with RCA-I in an agglutination test.

According to an 8th aspect of the present invention, we provide a CHO cell or CHO cell line obtainable by a method set out above.

5 The practice of the present invention will employ, unless otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA and immunology, which are within the capabilities of a person of ordinary skill in the art. Such techniques are explained in the literature. See, for example, J. Sambrook, E. F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Second Edition, Books 1-3, 10 Cold Spring Harbor Laboratory Press; Ausubel, F. M. et al. (1995 and periodic supplements; *Current Protocols in Molecular Biology*, ch. 9, 13, and 16, John Wiley & Sons, New York, N.Y.); B. Roe, J. Crabtree, and A. Kahn, 1996, *DNA Isolation and Sequencing: Essential Techniques*, John Wiley & Sons; J. M. Polak and James O'D. McGee, 1990, *In Situ Hybridization: Principles and Practice*; Oxford University Press; M. J. Gait (Editor), 1984, 15 *Oligonucleotide Synthesis: A Practical Approach*, Irl Press; D. M. J. Lilley and J. E. Dahlberg, 1992, *Methods of Enzymology: DNA Structure Part A: Synthesis and Physical Analysis of DNA* Methods in Enzymology, Academic Press; Using Antibodies : A Laboratory Manual : Portable Protocol NO. I by Edward Harlow, David Lane, Ed Harlow (1999, Cold Spring Harbor Laboratory Press, ISBN 0-87969-544-7); Antibodies : A Laboratory Manual by 20 Ed Harlow (Editor), David Lane (Editor) (1988, Cold Spring Harbor Laboratory Press, ISBN 0-87969-314-2), 1855. Handbook of Drug Screening, edited by Ramakrishna Seethala, Prabhavathi B. Fernandes (2001, New York, NY, Marcel Dekker, ISBN 0-8247-0562-9); and Lab Ref: A Handbook of Recipes, Reagents, and Other Reference Tools for Use at the Bench, Edited Jane Roskams and Linda Rodgers, 2002, Cold Spring Harbor Laboratory, ISBN 0- 25 87969-630-3. Each of these general texts is herein incorporated by reference.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a photograph showing that transiently expressed erythropoietin (EPO) in JW152 cell line is sensitive to endoglucosidase H (Endo H) treatment. Western blot of transiently expressed EPO: from CHO K-1 cells (Lane 1), from JW152 cells (Lane 2), 30 Endoglucosidase H treated EPO from CHO-K1(Lane 3), Endoglucosidase H treated EPO from JW152 (Lane 4). JW152 expressed EPO is sensitive to Endo H treatment suggesting that

the glycan structure found on the EPO molecule is of the high mannose type. It is hence suspected that gene(s) from the early part of the glycosylation pathway is defective.

Figure 2 is a photograph showing that JW152 cells lack a functional GnT I gene. Recombinant EPO is analyzed by the IEF/Western blot assay. The lane on the far left (CHO WT) shows the EPO expressed by the wild type CHO cells as a control. The second lane shows the EPO produced by the JW152 cells. Incomplete glycosylation in JW152 cells is obvious by comparing the two lanes. For the complementation test, each of several glycosylation related genes (as indicated) is co-transfected with the EPO construct into JW152 cells. Only GnT I is able to complement the genetic defect in JW152 cells, suggesting these cells lack a functional GnT I gene. Another very important observation from this IEF results is that in the presence of GnT I, JW152 cells sialylate EPO much better than the wild type cells.

Figure 3 is a photograph showing that improvement in sialylation after co-transfecting with GnT I is observed for JW152 but not for CHO-K1. The sialylation pattern of EPO produced in CHO-K1, CHO-K1 with co-expression of GnT I. EPO produced in JW152 cells and in JW152 cells with the co-expression of GnT I. The EPO sialylation patterns of CHO-K1 with and without co-expression of GnT I appear to be the same and so, the overexpression of GnT I is not responsible for the betterment in sialylation seen in lane 4. Results suggest that in the presence of GnT I, JW152 cells sialylate recombinant proteins much better than the wild type cells. Treating samples with neuraminidase, which cleaves off sialic acid, results in EPO bands in the acidic region being reduced to the basic region after cleaving off sialic acid, showing that the higher sialylated forms of EPO are indeed focused in the acidic region of the gel.

Figure 4 is a photograph showing that EPO expression in the presence of functional GnT I of previously published Lec 1 mutant which has a GnT I defect is shown to be also highly sialylated. Transient EPO expressed in CHO-K1, Lec 1 and Lec 1 with the restoration of functional GnT I. Lec 1 is previously reported and had been independently isolated using a different lectin by another group headed by Pamela Stanley.

Figure 5 is a photograph showing that all nine CHO glycosylation mutants, resistant to RCA and bearing distinct mutations in GnT I gene, lead to incomplete sialylation. Transiently expressed EPO in JW152, JW36, JW80, KFC5008, KFC5026, KFC15002, KFC15047, KFC15071, KFC 20011, CHO-K1. EPO produced in nine CHO glycosylation

mutants selected with RCA lectin, each bearing a different mutation in the GnT I gene lead to a loss in GnT I function. Incomplete sialylation is observed in these cell lines as compared to CHO wild-type (extreme right). EPO clinical standard is on extreme left. RCA lectin yields mainly GnT I deficient mutants.

5 Figure 6 is a photograph showing that all nine CHO glycosylation mutants, after restoring GnT I function transiently express EPO that is better sialylated than that in CHO-K1.

Figure 7 is a photograph showing that JW152, with restored GnT I function, also sialylates EPO-Fc fusion protein better than CHO-K1. EPO-Fc fusion protein produced in CHO wild-type compared alongside with that produced in JW152 before and after rescue. The
10 results demonstrate that the superior sialylation by rescued JW152 is maintained even with a different model glycoprotein.

Figure 8 is a photograph showing that HPAEC chromatogram showing better sialylated glycans cleaved from EPO-Fc expressed in JW152 cells co-expressing functional GnT I. EPO-Fc transiently expressed in CHO-K1 and JW152, cotransfected with GnT I, is
15 purified using affinity purification via a protein A-bound chromatography column. Glycans on the equal amounts of EPO-Fc are cleaved by treatment with Peptide: N-Glycosidase F (PNGase F), and separated, using High pH Anion Exchange Chromatography (HPAEC), according to their number of sialic acids attached to the glycans. The chromatogram shows distinctly higher peaks for the JW152 sample in the 4S group and lower peaks in the 1S group
20 when comparing with the CHO-K1 sample. Arrows point to an internal control, raffinose. Samples are labeled as shown.

Figure 9 is a photograph showing recombinant EPO expressed by 10 randomly picked clones isolated by RCA-I. Top gel, EPO expressed in 10 different mutant lines. Cells are transfected with a construct that expresses EPO alone. Bottom gel, EPO expressed in the same
25 set of cell lines that are transfected with pEIG. This construct expresses both EPO and GnT I.

Figure 10 is a photograph showing that EPO produced in 10 different JW152 clones that are stably transfected with a construct that expresses both EPO and GnT I. JW152 cells are transfected with a construct called pEIG. In pEIG, a transcription unit containing EPO-IRES-GnT I (EIG) is cloned into pcDNA3.1 downstream of the CMV promoter. After
30 transfection, stable clones are selected and picked. The sialylation patterns of EPO produced by 10 such randomly picked clones are analyzed by the IEF/Western blot assay. The results

show that all the EPO samples produced by different stable clones are highly sialylated. This means that better sialylation is maintained at stable expression of EPO in the presence of functional GnT I.

Figure 11 is a photograph showing that stable expression of EPO in CHO-K1 shows a sialylation profile that is inferior to that of JW152 with restored GnT I function. CHO-K1 cells are transfected with expression vector containing EPO coding sequence and put under selection. Clones are isolated and EPO from nine randomly selected stable clones is analysed with IEF. The results show that all the EPO samples produced by different stable clones vary in overall sialylation but in general are not as well sialylated as those in Figure 10.

Figure 12 is a photograph showing that EPO produced in four JW152 stable cell lines that have been adapted to suspension batch culture in protein free medium. The sialylation pattern observed for stable JW152 cell lines in attached culture is maintained in suspension culture.

SEQUENCE LISTING

SEQ ID NO: 1 is a nucleic acid sequence of a N-acetylglucoaminyltransferase I cDNA from CHO JW152 cells.

SEQ ID NO: 2 is an amino acid sequence of N-acetylglucoaminyltransferase I encoded by SEQ ID NO: 1.

DETAILED DESCRIPTION

Using a cytotoxic lectin, RCA-I, we have isolated a novel CHO mutant cell line, JW152, from CHO-K1 cells.

Recombinant EPO produced by JW152 cells that are stably transfected with EPO and GnT I cDNAs contains the highly sialylated isoforms. Several of these stably transfected lines have been adapted in suspension culture and grown in serum-free medium. The EPO produced by these cells are analyzed by an IEF assay. The results showed that EPO produced in serum-free medium remained highly sialylated.

These results suggest that JW152 cells have the potential to become a host cell line for producing proteins, such as highly sialylated proteins including glycoprotein drugs.

The cell line JW152 was deposited on 11 December 2008 at the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, United States of America under the accession number PTA-9657 as the International Deposit Number under the Budapest Treaty on the International Recognition of the Deposit of Microorganisms
5 for the Purposes of Patent Procedure.

We therefore provide for a Chinese Hamster Ovary (CHO) cell or cell line which is capable of higher protein sialylation compared to a wild type Chinese Hamster Ovary cell.

The CHO cell capable of higher sialylation may be produced by a suitable selection method such as a RCA-1 selection method. Such a method is described in further detail in
10 “CHO Cells and Cell Lines” below. Therefore, RCA-I can be used to isolate CHO glycosylation mutant cells produce highly sialylated recombinant proteins in the presence of GnT I.

Such a selection method is therefore specifically included in the methods and compositions described here.

15 We therefore provide for a Chinese Hamster Ovary (CHO) cell or cell line which is capable of higher protein sialylation compared to a wild type Chinese Hamster Ovary cell, the CHO cell being obtainable by selection with *Ricinus communis* agglutinin I (RCA-I). In general, we provide a RCA-I resistant CHO cell or cell line, such as an RCA-I resistant CHO-K1 cell or cell line.

20 Genetic analysis has revealed a dysfunctional N-Acetyl-glucosaminyltransferase I (GnT I) gene in JW152 cells. Molecular cloning of the GnT I cDNA from the mutant cells identified a point mutation that results in a premature stop codon. As a result, the JW152 cells can only synthesize a truncated version of GnT I protein with only 338 amino acids, rather than the normal protein which contains 447 amino acids.

25 Using RCA-I we have isolated many more CHO mutant lines (about 100 clones). Genetic analyses showed that they all lack functional GnT I gene. Many of them carry a different point mutation in the coding region of GnT I gene, suggesting that they derived from different original clones. Yet, they all dramatically improved sialylation of recombinant proteins in the presence of GnT I. The mutations in the GnT I genes from these further CHO
30 cell lines are shown in Table D1 below.

Accordingly, we provide for a Chinese Hamster Ovary (CHO) cell which is capable of higher protein sialylation in the presence of functional GnT 1 compared to a wild type Chinese Hamster Ovary cell, the CHO cell comprising a mutation in the GnT 1 gene. We further provide for a Chinese Hamster Ovary (CHO) cell or cell line which is capable of higher protein sialylation compared to a wild type Chinese Hamster Ovary cell, the CHO cell being obtainable by selection with *Ricinus communis* agglutinin I (RCA-I) and comprising a mutation in the GnT 1 gene.

The mutation in the GnT 1 gene may comprise any point mutation, deletion, inversion, etc. The mutation in the GnT 1 gene may encode a partially functioning, or non-functioning GnT 1 polypeptide. The mutation in the GnT 1 gene may encode a truncated GnT 1 polypeptide.

We provide for CHO cells and cell lines comprising each of these mutant CHO cell lines and clones. We provide for specific cell lines derived from RCA-I selection and capable of higher sialylation compared to wild type or native CHO cells or parental cells. We provide for mutant CHO nucleic acid and polypeptide sequences comprised in such cells or cell lines, as described in further detail below.

The CHO cell or cell line may comprise a JW152 cell line. It may comprise a JW80 cell line. It may comprise a JW36 cell line. It may comprise a KFC15002 cell line. It may comprise a KFC15071 cell line. It may comprise a KFC5008 cell line. It may comprise a JW152 cell line. It may comprise a KFC5026 cell line. It may comprise a KFC20011 cell line. It may comprise a KFC15047 cell line.

The CHO cell or cell line may be transfected with a nucleic acid encoding a protein of interest, for example a heterologous or recombinant protein. Such a protein might comprise a glycoprotein. The CHO cell or cell line may be transfected or co-transfected with a nucleic acid encoding a functional or full length or wild type GnT 1 sequence.

As noted above, we provide for the nucleic acid themselves, e.g., a nucleic acid encoding a mutant GnT 1 gene, or a fragment, variant, derivative or homologue of such a nucleic acid. The nucleic acid encoding a mutant GnT 1 gene, fragment, variant, derivative or homologue may cause a CHO cell comprising it to be capable of higher sialylation, compared to a CHO cell which does not comprise such a nucleic acid, for example a wild type CHO cell.

The nucleic acid encoding the mutant GnT 1 gene may comprise a sequence shown as SEQ ID NO: 1 or a variant, homologue, derivative or fragment thereof.

SEQ ID NO: 1

The coding region of N-acetylglucoaminyltransferase I (Mgat1, GenBank: AF343963)

- 5 mRNA isolated from the CHO JW152 cells. In these mutant cells, a C to T point mutation at position 1015 was identified (shown in bold):

ATGCTGAAGAAGCAGTCTGCAGGGCTTGTGCTTTGGGGTGCTATCCTCTTTGTGGGCTGGAATGCCCTGCTGCTC
CTCTTCTTCTGGACACGCCCAGCCCCCTGGCAGGCCCCCTCAGATAGTGCTATCGATGATGACCTGCCAGCCTC
ACCCGTGAGGTGTTCCGCCTGGCTGAGGACGCTGAGGTGGAGTTGGAGCGGCAGCGGGGGCTGTTGCAGCAAATC
10 AGGGAGCATCATGCTTTGTGGAGACAGAGGTGGAAAGTGCCCAACCGTGGC^{CC}CTCCAGCCTGGCCCCGTGTGCCT
GCGACCCCTCACCAGCCGTGATCCCCATCCTGGTCATTGCCTGTGACCGCAGCACTGTCCGGCGCTGCTTGGAT
AAGTTGTGCACTATCGGCCCTCAGCTGAGCATTCCCCATCATTGTGAGCCAGGACTGCGGGCAGGAAGAGACA
GCACAGGTGATTGCTTCCCTATGGCAGTGCAGTCACACACATCCGGCAGCCAGACCTGAGTAACATCGCTGTGCCC
CCAGACCACCGCAAGTTCAGGGTTACTACAAGATCGCCAGGCACTACCGCTGGGCACTGGGCCAGATCTTCAAC
15 AAGTTCAAGTTCCCAGCAGCTGTGGTAGTGGAGGACGATCTGGAGGTGGCACCAGACTTCTTTGAGTACTTCCAG
GCCACCTACCCACTGCTGAGAACAGACCCCTCCCTTTGGTGTGTGTCTGCTTGAATGACAATGGCAAGGAGCAG
ATGGTAGACTCAAGCAAACCTGAGCTGCTCTATCGAACAGACTTTTTTCTGGCCTTGGCTGGCTGCTGATGGCT
GAGCTGTGGACAGAGCTGGAGCCCAAGTGGCCCAAGGCCTTCTGGGATGACTGGATGCGCAGACCTGAGCAGCGG
AAGGGGCGGGCCTGTATTTCGTCCAGAAATTTCAAGAACGATGACCTTTGGCCGTAAGGGTGTGAGCCATGGGCAG
20 TTCTTTGATCAGCATCTTAAGTTCATCAAGCTGAACCAGTAGTTCGTGTCTTTCACCCAGTTGGATTTGTCATAC
TTGCAGCGGGAGGCTTATGACCGGGATTTCTTGCCCGTGTCTATAGTGCCCCCTGCTACAGGTGGAGAAAGTG
AGGACCAATGATCAGAAGGAGCTGGGGGAGGTGCGGGTACAGTACACTAGCAGAGACAGCTTCAAGGCCTTTGCT
AAGGCCCTGGGTGTGATGGATGACCTCAAGTCTGGTGTCCCCAGAGCTGGCTACCGGGCGTTGTCACTTTCCAG
TTCAGGGGTCGACGTGTCCACCTGGCACCCCAACCTGGGAAGGCTATGATCCTAGCTGGAATTAG

- 25 We further provide for mutant GnT 1 polypeptides, as well as fragments, variants, derivatives and homologues thereof. The mutant GnT 1 polypeptide, fragment, variant, derivative or homologue may cause a CHO cell comprising it to be capable of higher sialylation, compared to a CHO cell which does not comprise such a polypeptide, for example a wild type CHO cell. The mutant GnT 1 polypeptide may comprise a sequence shown as
30 SEQ ID NO: 2 or a variant, homologue, derivative or fragment thereof.

SEQ ID NO: 2

The N-acetylglucoaminyltransferase I (GnT I) protein encoded by the mutated gene in CHO JW152 cells. As a result of the point mutation (C1015T), JW152 cells only produce a truncated version of GnT I which contains only 338 amino acids rather than the normal
35 protein that contains 447 amino acids. The C-terminal portion in bold is not translated in JW152 cells.

MLKKQSAGLVLVGAILFVGWNAALLLLFFWTRPAPGRPPSDSAIDDDPASLTREVFRLAEDAEEVELERQRGILLQOI
REHHALWRQRWKVPTVAPPAWPRVPATPSPAVIPILVIACDRSTVRRCLDKLLHYRPSAEHFPIIVSQDCGHEET
AQVIASYGSAVTHIRQPDLSNIAVPPDHRKFQGYKARHYRWALGQIFNKFKFPAVVVEDDLEVAPDFFEYFQ
40 ATYPLLRTDPSLWCVSAWNDNGKEQMVDSSKPELLYRTDFFPGLGWLLMAELWTELEPKWPKAFFWDDWMRRPEQR

KGRACIRPEISRTMTFGRKGVSHGQFFDQHLKFIKLNQQFVSFTQLDLSYLQREAYDRDFLARVYSAPLLQVEKV
RTNDQKELGEVRVQYTSRDSFKAFKALGVMDLKSQVPRAGYRGVVTQFRGRRVHLAPPQTWEGYDPSWN

Several of the JW152-pEIG stable lines shown in Figure 2 have been adapted in suspension culture and grown in serum-free medium. The EPO produced in serum-free medium remained highly sialylated. We therefore provide for mutant CHO cells and cell lines derived from RCA-I selection, which have been adapted to suspension culture, or growth in semi-solid medium.

In conclusion, we have developed a novel method to isolate glycosylation mutant cells from CHO cells. All CHO cells that survive RCA-I treatment have very similar characteristics. First, they all lack a functional GnT I gene. Second, they all sialylate their recombinant proteins better than the wild type CHO cells in the presence of GnT I. This feature remains the same both in transient transfection and in stably transfected cells.

The CHO glycosylation mutant cells isolated with RCA-I, such as JW152 cells, can produce recombinant glycoproteins with high degree of sialylation as long as GnT I is present. This method can be used to produce recombinant glycoproteins in which sialic acid content is important for the efficacy. These proteins include EPO, IEF- γ , Factor VIII etc.

CHO CELLS AND CELL LINES

The CHO cells and cell lines described here may be made by any suitable means. For example, the CHO cells and cell lines may be produced by selection using a suitable agglutinating agent, such as agglutinin I. The agglutinin may comprise any suitable agglutinin, such as *Ricinus communis* agglutinin I (RCA-I).

We therefore provide a method of providing a CHO cell or cell line, the method comprising culturing CHO cells in the presence of *Ricinus communis* agglutinin I (RCA-I) and selecting cells which survive the culture.

The CHO cells and cell lines described here may be made by treating a starting or parent cell with *Ricinus communis* agglutinin I (RCA-I) and selecting cells that survive such treatment. Such surviving cells may be further cloned and made into cell lines. The selected cells and cell lines may comprise higher sialylation activity as described in this document. The selected cells and cell lines may comprise mutant GnT 1 genes and polypeptides, as described in this document.

For example, the CHO cells or cell lines described here may be selected by exposing a parent cell line to *Ricinus communis* agglutinin I (RCA-I) at a suitable concentration for a suitable period.

5 The RCA-1 concentration could range from between 0.1 µg/ml to 100 µg/ml, for example up to 50 µg/ml or up to 20 µg/ml. Examples of specific concentrations include 10 µg/ml and 5 µg/ml.

The period of incubation or exposure to RCA-1 could be from an hour, a few hours (such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12), overnight, to a few days, such as 2 days or 3 days.

10 In general, the period and concentration can be adjusted to eliminate the majority of CHO cells, but to enable a small proportion of cells, which are resistant to RCA-I, to survive and form colonies. Within this, the concentration and period of exposure and selection may be varied, but generally, the higher the concentration of RCA-I, the lower the period of exposure is necessary, and *vice versa*.

15 The selection could be done on any suitable starting cell or cell line, but this will generally be a CHO cell or cell line. Any known CHO cell or cell line could be used as a starting point or parent cell, including CHO-K1. Other suitable starting cells could include, but are not limited to the following (ECACC accession numbers in brackets): CHO (85050302), CHO (PROTEIN FREE) (00102307), CHO-K1 (85051005), CHO-K1/SF (93061607), CHO/dhFr- (94060607), CHO/dhFr-AC-free (05011002), RR-CHOKI
20 (92052129).

Following selection, the surviving cells are allowed to grow and form colonies following which they may be picked. The time allowed for this will vary, but will generally be long enough for colonies to grow to a pickable size. Examples of such times are 5 days, 7 days, 9 days, 11 days, 13 days, one week, two weeks, three weeks or more.

25 The picking may be done manually, or it may be automated through use of robots, such as CLONEPIX (Genetix, New Milton, Hampshire, UK). The picked colonies may be further cloned, further screened, characterised and cultured, etc.

The selected cells may be subjected to further tests. For example, they may be subjected to agglutination tests using RCA-I to confirm the mutant cells no longer react with
30 RCA-I.

As a specific example, which is not intended to be limiting, CHO-K1 cells may be cultured, for example in 6-well plates, to confluence. Culture media may be changed to serum-free DMEM. *Ricinus communis* agglutinin I (RCA-I, EY Laboratories) may be added into the media to reach a final concentration of 10µg/gml. This may be incubated with cells
 5 overnight. The serum-free DMEM containing RCA-I may be replaced, for example with fresh DMEM with 10% FBS. Nine days later, colonies of the CHO cells that survive the RCA-I treatment may be picked and cultured, for example in 24 well plates.

These cells may be subjected to further tests, such as agglutination tests using RCA-I to confirm the mutant cells no longer react with RCA-I. We therefore provide for a method of
 10 providing a CHO cell or cell line, the method comprising culturing CHO cells in the presence of *Ricinus communis* agglutinin I (RCA-I), selecting cells which survive the culture and which do not react with RCA-I in an agglutination test.

The RCA-I selected CHO cells and CHO cell lines may be tested for their sialylation behaviour, by for example expressing a protein of interest and determining the degree of
 15 sialylation. This may be done by the methods described in "Sialylation" below.

We therefore provide for a method of providing a CHO cell or cell line, the method comprising culturing CHO cells in the presence of *Ricinus communis* agglutinin I (RCA-I), selecting cells which survive the culture and selecting those cells or cell lines which display high sialylation behaviour, for example high Z-number or low pI of expressed proteins.

20 The GnT 1 gene in such selected cells may be cloned and sequenced, using methods known in the art. The GnT 1 gene may comprise a mutant GnT 1 gene as described here.

We therefore provide for a method of providing a CHO cell or cell line, the method comprising culturing CHO cells in the presence of *Ricinus communis* agglutinin I (RCA-I), selecting cells which survive the culture and selecting those cells or cell lines which comprise
 25 mutant GnT 1 genes as described herein.

MUTANT CHO CELLS AND CELL LINES

We provide for a CHO cell or cell line derived from RCA-I selection, as described above. Such a cell line could include a JW152 cell line, or any of the cell lines set out in Table D1 below, including a JW80 cell line, a JW36 cell line, a KFC15002 cell line, a KFC15071

cell line, a KFC5008 cell line, a JW152 cell line, a KFC5026 cell line, a KFC20011 cell line or a KFC15047 cell line.

PROTEIN EXPRESSION

The CHO cells described here may be used as host cells for expression of any protein of interest. This may be done by means known in the art.

Protein expression in CHO cells and cell lines is well described in the literature, and the skilled person will have little difficulty in using the CHO cells and cell lines described here as hosts for protein expression. Thus, for example, the CHO cells and cell lines may be transfected by means known in the art with expression vectors capable of expressing the protein of interest.

The CHO cells and cell lines may further be capable of expressing wild type or functional GnT 1, for example, the sequence set out in GenBank accession number AF343963. This may be done by transfecting the CHO cells and cell lines with an expression vector encoding GnT 1. This may be on the same or different vector as that which contains the nucleic acid encoding the protein of interest

Any suitable protein may be expressed using the CHO cells described here as host cells. The protein may comprise a heterologous protein. The protein may comprise a recombinant protein. The protein may comprise an engineered protein. The protein may comprise a glycoprotein.

Examples include heterologous proteins of therapeutic or pharmacological interest. Proteins which may be expressed include anti-EGFR mAb, α -glucosidase, laronidase, Ig-CTLA4 fusion, N-acetylgalactosamine-4-sulfatase, luteinizing hormone, anti-VEGF mAb, Factor VIII, anti-IgE mAb, anti-CD11a mAb, α -galactosidase, interferon- β , anti-TNF α mAb, erythropoietin, anti-CD52 mAb, Factor VIII, tissue plasminogen activator, anti-HER2 mAb, TNF α receptor fusion, Factor IX, follicle stimulating hormone, anti-CD20 mAb, interferon- β , β -glucocerebrosidase, deoxyribonuclease I, etc.

For example, we describe the expression of erythropoietin (EPO), interferon- γ and Factor VIII with CHO cells and cell lines described here. We also describe highly sialylated forms or isoforms of erythropoietin (EPO), interferon- γ and Factor VIII expressed from CHO cells and cell lines described here.

SIALIC ACID

The term “sialic acid” is intended to refer to any member of a family of nine-carbon carboxylated sugars.

The most common member of the sialic acid family is N-acetylneuraminic acid (2-keto-5-acetamido-3, 5-dideoxy-D-glycero- D-galactononulopyranos-1-onic acid, often abbreviated as Neu5Ac, NeuAc, or NANA). A second member of the family is N-glycolylneuraminic acid (Neu5Gc or NeuGc), in which the N-acetyl group of NeuAc is hydroxylated. A third sialic acid family member is 2-keto-3-deoxy-nonulosonic acid (KDN). 17

Also included are 9-substituted sialic acids such as a 9-O-C1-C6-acyl-Neu5Ac like 9-O-lactyl- Neu5Ac or 9-O-acetyl-Neu5Ac, 9-deoxy-9-fluoro-Neu5Ac and 9-azido-9-deoxy-Neu5Ac. For review of the sialic acid family, see, e. g. , Varki; Glycobiology 2 1992 ; 25-40 ; Sialic Acids: Chemistry, Metabolism and Function, R. Schauer, Ed.

SIALYLATION

The degree of sialylation of a protein may be measured by various means, such as using Z-numbers or expressing the isoelectric point of the protein.

We therefore provide for protein expression using the CHO cells and cell lines described here, such protein expression comprising highly sialylated forms or isoforms of proteins of interest. Protein expression from CHO cells and CHO cell lines described here enables production of proteins such as glycoproteins, for example erythropoietin (EPO), interferon- γ and Factor VIII, which have high Z-numbers or low pIs, or both.

Z-number

The parameter Z-number provides a measure of how many of the antennae of the carbohydrate moieties in a glycoprotein bear charged residues, such as sialic acid.

To determine Z-number, the carbohydrate moieties are released from the peptide, as above, and labelled, if desired. The mixture is then separated by ion exchange chromatography, allowing the separation of species on the basis of charge. Visualisation of the eluted peaks may be by virtue of a label, as mentioned above, or may be by some other method, such as mass-spectrometry.

A chromatogram is then analysed by integrating the peaks associated with mono- di- tri- and tetra-charged carbohydrate species. The percentage of the total carbohydrate represented by each species can then be used to calculate Z-number according to the following equation : $Z = P'_{\text{mono}} + 2 P'_{\text{di}} + 3 P'_{\text{tri}} + 4 P'_{\text{tetra}}$ wherein Z is Z-number, and P'mono, P'di, P'tri and P'tetra are the percentage of total carbohydrate that is mono-, di-, tri- and tetra-charged respectively.

A high Z-number indicates that a large number of antennae bear charged residues, and that the glycoprotein will therefore be highly charged, and in the case of sialic acid residues, acidic.

The CHO cells described here are capable of expressing proteins with a high degree of sialylation. Thus, for example, the CHO cells may express proteins with high Z-number values such as greater than 150, such as greater than 160, greater than 170, greater than 180, such as greater than 190, greater than 200, greater than 210, such as greater than 220, greater than 230, etc.

15 *Isoelectric Point*

The isoelectric point (pI) of a protein may also be used as a measure of its sialylation. The higher the degree of sialylation, the more acidic it is and the lower its pI.

Proteins expressed by the CHO cells described here have significantly lower pI profiles than their normal counterparts, e.g., native proteins, or counterpart proteins expressed by wild type CHO cells. For example, the proteins expressed by the CHO cells and cell lines described here may have low pIs, such as pIs of 4.5 or less, such as 4.3 or less, 4.1 or less, 4.0 or less, 3.8 or less, 3.6 or less, 3.4 or less, 3.2 or less, 3.0 or less, 2.8 or less, 2.6 or less, etc. The pIs may be of individual protein molecules, or a batch or fraction of them, or an average pI of an expressed lot of protein.

The expressed proteins may be isolated from a mixture of isoforms using a number of methods that will be known to one skilled in the art. For example, isoelectric focussing, chromatofocussing or ion-exchange chromatography may be used to separate the isoforms on the basis of pI. The different fractions can be analysed for sialic acid content, and the desired fractions selected for use.

MUTANT GnT 1 SEQUENCES

We disclose mutant GnT 1 sequences comprising mutant GnT 1 amino acid sequences and mutant GnT1 nucleic acid sequences.

5 Example mutant GnT 1 amino acid sequences include the sequence shown as SEQ ID NO: 2, as well as the sequences comprising the mutations shown in column 3 of Table D1 below.

Example mutant GnT 1 nucleic acid sequences include the sequence shown as SEQ ID NO: 1, as well as the sequences comprising the mutations shown in column 2 of Table D1 below.

10 Table D1 shows the mutations in the GnT 1 sequence of clones isolated from selection with RCA-I, as described in the Examples below.

Corresponding mutations are tabulated alongside showing respective nucleotide and amino acid mutation and possible location of the disruption in secondary structure/interaction or the resulting loss in amino acids in the case of a stop codon mutation.

15 A minimum of 4 bacteria colonies were sequenced to ensure that the mutations found were not due to PCR error.

Mutant CHO Cell Line	DNA Mutation	Polypeptide Mutation (position, mutation)	Comment
JW80	G1300C	Position 434 Ala → Pro	Domain 2 β14
JW36	A638C	Position 213 Asp → Ala	Disruption of DxD Motif, similar to JW98, JW191
KFC15002	C784G	Position 262 Arg → Gly	Domain 1 β6, similar to KFC15008 (aka 15008), KFC7501 (aka 7501) and KFC12008
KFC15071	T811A	Position 271 Trp → Arg	Domain 1 β7
KFC5008	_706C	Insertion at 706 bp Frame shift from 236 aa Asp→STOP	Stop codon generated at 245 a.a.
JW152	C1015T	Position 339 Gln→STOP	108 a.a. missing from C terminal
KFC5026	G246A	Position 82 Trp→STOP	365 a.a. missing from C terminal
KFC20011	G258A	Position 86 Trp→STOP	361 a.a. missing from C terminal
KFC15047	A859T	Position 287 Lys→STOP	160 a.a. missing, similar to KFC15026 and KFC15072

Table D1. Table of mutations found in GnT1 mutants from clones derived from RCA-I screening (Example 10 below). Nine CHO glycosylation mutants with different mutations in the GnT1 gene. Point mutations leading to loss in function were found in four cell lines (JW80, JW36, KFC15002, KFC15071). A point insertion resulting in the generation of a stop codon was found in KFC 5008. Mutations leading to a premature stop codon were also found in another four cell lines. (JW152, KFC5026, KFC20011, KFC15047).

MUTANT GNT 1 POLYPEPTIDES

The CHO cells and cell lines comprise mutant GnT 1 polypeptides.

We therefore provide generally for a mutant GnT 1 polypeptide, together with fragments, homologues, variants and derivatives thereof. These polypeptide sequences may comprise the polypeptide sequences disclosed here, and particularly in the sequence listings.

The mutant GnT 1 polypeptide may comprise one or more changes compared to the wild type GnT 1 sequence. Such mutations may result from stop codons being introduced in the encoding nucleic acid sequence and consequent premature termination of translation of the GnT 1 mRNA.

5 The mutant GnT 1 polypeptide may be shorter than a wild type GnT 1 polypeptide. It may be a truncated version of wild type GnT 1 polypeptide. The length of the mutant GnT 1 polypeptide may be 90% or less, 80% or less, 70% or less, etc than the wild type sequence.

 For example, a mutant GnT 1 polypeptide may be missing 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more C-terminal residues compared to full
10 length or wild type GnT 1 polypeptide.

 The mutant GnT 1 polypeptide may for example comprise a sequence set out in SEQ ID NO: 2. This is the mutant GnT 1 polypeptide sequence from the cell line JW152. The mutant GnT 1 polypeptide may comprise a GnT 1 sequence comprising a mutation set out at column 3 of Table D1 above.

15 It will be understood that the mutant GnT 1 polypeptide sequences disclosed here are not limited to the particular sequences set forth in the sequence listing, or fragments thereof, or sequences obtained from mutant GnT 1 protein, but also include homologous sequences obtained from any source, for example related cellular homologues, homologues from other species and variants or derivatives thereof, provided that they have at least one of the
20 biological activities of mutant GnT 1, as the case may be.

 This disclosure therefore encompasses variants, homologues or derivatives of the amino acid sequences set forth in the sequence listings, as well as variants, homologues or derivatives of the amino acid sequences encoded by the nucleotide sequences disclosed here. Such a sequences is generally referred to as a “mutant GnT 1 sequence”.

25 The length of the mutant GnT 1 polypeptide may be 90% or less, 80% or less, 70% or less, etc than a corresponding wild type sequence.

 For example, a mutant GnT 1 nucleic acid may encode a mutant GnT 1 polypeptide that is missing 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more C-terminal residues.

Biological Activities

In some embodiments, the sequences comprise at least one biological activity of mutant GnT 1, as the case may be. The biological activity may comprise improved ability to express proteins with higher sialylation compared to wild type GnT 1

5 *Homologues*

The polypeptides disclosed include homologous sequences obtained from any source, for example related viral/bacterial proteins, cellular homologues and synthetic peptides, as well as variants or derivatives thereof.

10 In the context of the present document, a homologous sequence or homologue is taken to include an amino acid sequence which is at least 60, 70, 80 or 90% identical, such as at least 95 or 98% identical at the amino acid level over at least 30, such as 50, 70, 90 or 100 amino acids with GnT 1, as the case may be, for example as shown in the sequence listing herein. In the context of this document, a homologous sequence is taken to include an amino acid sequence which is at least 15, 20, 25, 30, 40, 50, 60, 70, 80 or 90% identical, such as at
15 least 95 or 98% identical at the amino acid level, such as over at least 15, 25, 35, 50 or 100, such as 200, 300, 400 or 500 amino acids with the sequence of GnT 1.

Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present document it is possible to express homology in terms of sequence identity. In some embodiments, the
20 sequence identity is determined relative to the entirety of the length the relevant sequence, i.e., over the entire length or full length sequence of the relevant gene, for example.

Homology comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences.

25 % homology may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an “ungapped” alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues (for example less than 50 contiguous amino acids).

Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting “gaps” in the sequence alignment to try to maximise local homology.

However, these more complex methods assign “gap penalties” to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. “Affine gap costs” are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, the default values may be used when using such software for sequence comparisons. For example when using the GCG Wisconsin Bestfit package (see below) the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux *et al.*, 1984, *Nucleic Acids Research* 12:387). Examples of other software than can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel *et al.*, 1999 *ibid* – Chapter 18), FASTA (Atschul *et al.*, 1990, *J. Mol. Biol.*, 403-410) and the GENWORKS suite of comparison tools. Both BLAST and FASTA are available for offline and online searching (see Ausubel *et al.*, 1999 *ibid*, pages 7-58 to 7-60).

Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol

comparison table if supplied (see user manual for further details). The public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62 may be used.

- Once the software has produced an optimal alignment, it is possible to calculate %
 5 homology, such as % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

Variants and Derivatives

- The terms “variant” or “derivative” in relation to the amino acid sequences as described here includes any substitution of, variation of, modification of, replacement of,
 10 deletion of or addition of one (or more) amino acids from or to the sequence. For example, the resultant amino acid sequence retains substantially the same activity as the unmodified sequence, such as having at least the same activity as the mutant GnT 1 polypeptide shown in the sequence listings.

- Polypeptides having the amino acid sequence shown in the Examples, or fragments or
 15 homologues thereof may be modified for use in the methods and compositions described here. Typically, modifications are made that maintain the biological activity of the sequence. Amino acid substitutions may be made, for example from 1, 2 or 3 to 10, 20 or 30 substitutions provided that the modified sequence retains the biological activity of the unmodified sequence. Amino acid substitutions may include the use of non-naturally
 20 occurring analogues, for example to increase blood plasma half-life of a therapeutically administered polypeptide.

- Natural variants of mutant GnT 1 are likely to comprise conservative amino acid substitutions. Conservative substitutions may be defined, for example according to the Table below. Amino acids in the same block in the second column and in the same line in the third
 25 column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar - uncharged	C S T M
		N Q
	Polar - charged	D E
		K R
AROMATIC		H F W Y

Fragments

Polypeptides disclosed here and useful as markers also include fragments of the above mentioned full length polypeptides and variants thereof, including fragments of the sequences set out in the sequence listings.

5 Polypeptides also include fragments of the full length sequence of the mutant GnT 1 polypeptide. Such fragments may comprise at least one epitope. Methods of identifying epitopes are well known in the art. Fragments will typically comprise at least 6 amino acids, such as at least 10, 20, 30, 50 or 100 amino acids.

10 Included are fragments comprising, such as consisting of, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145 or 150, or more residues from a mutant GnT 1 amino acid sequence.

15 Polypeptide fragments of the mutant GnT 1 proteins and allelic and species variants thereof may contain one or more (e.g. 5, 10, 15, or 20) substitutions, deletions or insertions, including conserved substitutions. Where substitutions, deletion and/or insertions occur, for example in different species, such as less than 50%, 40% or 20% of the amino acid residues depicted in the sequence listings are altered.

20 Mutant GnT 1, and fragments, homologues, variants and derivatives, may be made by recombinant means. However, they may also be made by synthetic means using techniques well known to skilled persons such as solid phase synthesis. The proteins may also be produced as fusion proteins, for example to aid in extraction and purification. Examples of fusion protein partners include glutathione-S-transferase (GST), 6xHis, GAL4 (DNA binding

and/or transcriptional activation domains) and β -galactosidase. It may also be convenient to include a proteolytic cleavage site between the fusion protein partner and the protein sequence of interest to allow removal of fusion protein sequences. The fusion protein may be such that it does not hinder the function of the protein of interest sequence. Proteins may also be
5 obtained by purification of cell extracts from animal cells.

The mutant GnT 1 polypeptide, variants, homologues, fragments and derivatives disclosed here may be in a substantially isolated form. It will be understood that such polypeptides may be mixed with carriers or diluents which will not interfere with the intended purpose of the protein and still be regarded as substantially isolated. A mutant GnT 1 variant,
10 homologue, fragment or derivative may also be in a substantially purified form, in which case it will generally comprise the protein in a preparation in which more than 90%, e.g. 95%, 98% or 99% of the protein in the preparation is a protein.

The mutant GnT 1 polypeptides variants, homologues, fragments and derivatives disclosed here may be labelled with a revealing label. The revealing label may be any suitable
15 label which allows the polypeptide, etc to be detected. Suitable labels include radioisotopes, e.g. ^{125}I , enzymes, antibodies, polynucleotides and linkers such as biotin. Labelled polypeptides may be used in diagnostic procedures such as immunoassays to determine the amount of a polypeptide in a sample. Polypeptides or labelled polypeptides may also be used in serological or cell-mediated immune assays for the detection of immune reactivity to said polypeptides in
20 animals and humans using standard protocols.

Mutant GnT 1 polypeptide, variants, homologues, fragments and derivatives disclosed here, optionally labelled, may also be fixed to a solid phase, for example the surface of an immunoassay well or dipstick. Such labelled and/or immobilised polypeptides may be packaged into kits in a suitable container along with suitable reagents, controls, instructions and the like.
25 Such polypeptides and kits may be used in methods of detection of antibodies to the polypeptides or their allelic or species variants by immunoassay.

Immunoassay methods are well known in the art and will generally comprise: (a) providing a polypeptide comprising an epitope bindable by an antibody against said protein; (b) incubating a biological sample with said polypeptide under conditions which allow for the
30 formation of an antibody-antigen complex; and (c) determining whether antibody-antigen complex comprising said polypeptide is formed.

The mutant GnT 1 polypeptides variants, homologues, fragments and derivatives disclosed here may be used in *in vitro* or *in vivo* cell culture systems to study the role of their corresponding genes and homologues thereof in cell function, including their function in disease. For example, truncated or modified polypeptides may be introduced into a cell to
5 disrupt the normal functions which occur in the cell. The polypeptides may be introduced into the cell by *in situ* expression of the polypeptide from a recombinant expression vector (see below). The expression vector optionally carries an inducible promoter to control the expression of the polypeptide.

The use of appropriate host cells, such as insect cells or mammalian cells, is expected
10 to provide for such post-translational modifications (e.g. myristolation, glycosylation, truncation, lipidation and tyrosine, serine or threonine phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products. Such cell culture systems in which the mutant GnT 1 polypeptide, variants, homologues, fragments and derivatives disclosed here are expressed may be used in assay systems to identify candidate
15 substances which interfere with or enhance the functions of the polypeptides in the cell.

MUTANT GN T 1 NUCLEIC ACIDS

The CHO cells and cell lines comprise mutant GnT 1 nucleic acids.

We therefore provide generally for a mutant GnT 1 nucleic acid, together with fragments, homologues, variants and derivatives thereof. These nucleic acid sequences may
20 encode the polypeptide sequences disclosed here, and particularly in the sequence listings.

The polynucleotide may comprise a mutant GnT 1 nucleic acid. The mutant GnT 1 nucleic acid may comprise one or more point mutations in the wild type GnT 1 sequence. Such mutations may result in corresponding changes to the amino acid sequence, or introduce stop codons and premature termination of translation of the GnT 1 mRNA.

25 The mutant GnT 1 nucleic acid may comprise a mutation resulting in a stop codon, which results in a mutant GnT 1 polypeptide being shorter than a wild type GnT 1 polypeptide. The length of the mutant GnT 1 polypeptide may be 90% or less, 80% or less, 70% or less, etc than the wild type sequence.

For example, a mutant GnT 1 nucleic acid may encode a mutant GnT 1 polypeptide that is missing 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more C-terminal residues.

The mutant GnT 1 nucleic acid may for example comprise a sequence set out in SEQ ID NO: 1. This is the mutant GnT 1 nucleic acid sequence from the cell line JW152. The mutant GnT 1 nucleic acid may comprise a GnT 1 sequence comprising a mutation set out at column 2 of Table D1 above.

In particular, we provide for nucleic acids or polynucleotides which encode any of the GnT 1 polypeptides disclosed here. Thus, the term “GnT 1 sequence” should be construed accordingly. However, such a nucleic acid or polynucleotide may comprise a sequence set out as SEQ ID NO: 1, or a sequence encoding a of the corresponding polypeptide, and a fragment, homologue, variant or derivative of such a nucleic acid. The above terms therefore may be taken to refer to these sequences.

As used here in this document, the terms “polynucleotide”, “nucleotide”, and nucleic acid are intended to be synonymous with each other. “Polynucleotide” generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. “Polynucleotides” include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. “Modified” bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, “polynucleotide” embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. “Polynucleotide” also embraces relatively short polynucleotides, often referred to as oligonucleotides.

It will be understood by a skilled person that numerous different polynucleotides and nucleic acids can encode the same polypeptide as a result of the degeneracy of the genetic code. In addition, it is to be understood that skilled persons may, using routine techniques, make nucleotide substitutions that do not affect the polypeptide sequence encoded by the polynucleotides described here to reflect the codon usage of any particular host organism in which the polypeptides are to be expressed.

Mutant GnT 1 Variants, Derivatives and Homologues

The mutant GnT 1 polynucleotides described here may comprise DNA or RNA. They may be single-stranded or double-stranded. They may also be polynucleotides which include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate and phosphorothioate backbones, addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present document, it is to be understood that the polynucleotides described herein may be modified by any method available in the art. Such modifications may be carried out in order to enhance the *in vivo* activity or life span of polynucleotides.

Where the polynucleotide is double-stranded, both strands of the duplex, either individually or in combination, are encompassed by the methods and compositions described here. Where the polynucleotide is single-stranded, it is to be understood that the complementary sequence of that polynucleotide is also included.

The terms "variant", "homologue" or "derivative" in relation to a nucleotide sequence include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleotides from or to the sequence. The resulting sequence may be capable of encoding a polypeptide which is capable of mediating higher sialylation in a CHO cell.

As indicated above, with respect to sequence identity, a "homologue" has for example at least 5% identity, at least 10% identity, at least 15% identity, at least 20% identity, at least 25% identity, at least 30% identity, at least 35% identity, at least 40% identity, at least 45% identity, at least 50% identity, at least 55% identity, at least 60% identity, at least 65% identity, at least 70% identity, at least 75% identity, at least 80% identity, at least 85% identity, at least 90% identity, or at least 95% identity to the relevant sequence shown in the sequence listings.

There may be at least 95% identity, such as at least 96% identity, such as at least 97% identity, such as at least 98% identity, such as at least 99% identity. Nucleotide homology comparisons may be conducted as described above. A sequence comparison program that may be used is the GCG Wisconsin Bestfit program described above. The default scoring matrix has a
 5 match value of 10 for each identical nucleotide and -9 for each mismatch. The default gap creation penalty is -50 and the default gap extension penalty is -3 for each nucleotide.

In some embodiments, a mutant GnT 1 polynucleotide has at least 90% or more sequence identity to a sequence shown as SEQ ID NO: 1. The mutant GnT 1 polynucleotide may have 60% or more, such as 65% or more, 70% or more, 75% or more, 80% or more, 85%
 10 or more, 90% or more, 95% or more, 97% or more or 98% or more sequence identity to a sequence shown as SEQ ID NO: 1.

HYBRIDISATION

We further describe mutant GnT 1 nucleotide sequences that are capable of hybridising selectively to any of the sequences presented herein, or any variant, fragment or
 15 derivative thereof, or to the complement of any of the above. Nucleotide sequences are such as at least 15 nucleotides in length, such as at least 20, 30, 40 or 50 nucleotides in length.

The term “hybridisation” as used herein shall include “the process by which a strand of nucleic acid joins with a complementary strand through base pairing” as well as the process of amplification as carried out in polymerase chain reaction technologies.

20 Polynucleotides capable of selectively hybridising to the nucleotide sequences presented herein, or to their complement, will be generally at least 70%, such as at least 80 or 90% or such as at least 95% or 98% homologous to the corresponding nucleotide sequences presented herein over a region of at least 20, such as at least 25 or 30, for instance at least 40, 60 or 100 or more contiguous nucleotides.

25 The term “selectively hybridisable” means that the polynucleotide used as a probe is used under conditions where a target polynucleotide is found to hybridize to the probe at a level significantly above background. The background hybridization may occur because of other polynucleotides present, for example, in the cDNA or genomic DNA library being screened. In this event, background implies a level of signal generated by interaction between the probe and a
 30 non-specific DNA member of the library which is less than 10-fold, such as less than 100 fold as

intense as the specific interaction observed with the target DNA. The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ^{32}P .

Hybridisation conditions are based on the melting temperature (T_m) of the nucleic acid binding complex, as taught in Berger and Kimmel (1987, Guide to Molecular Cloning
5 Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA), and confer a defined "stringency" as explained below.

Maximum stringency typically occurs at about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe); high stringency at about 5°C to 10°C below T_m ; intermediate stringency at about 10°C to 20°C below T_m ; and low stringency at about 20°C to 25°C below T_m . As will be
10 understood by those of skill in the art, a maximum stringency hybridisation can be used to identify or detect identical polynucleotide sequences while an intermediate (or low) stringency hybridisation can be used to identify or detect similar or related polynucleotide sequences.

In a one aspect, we disclose nucleotide sequences that can hybridise to a mutant GnT 1 nucleic acid, or a fragment, homologue, variant or derivative thereof, under stringent conditions
15 (e.g. 65°C and $0.1\times\text{SSC}$ { $1\times\text{SSC} = 0.15\text{ M NaCl}$, $0.015\text{ M Na}_3\text{ Citrate pH } 7.0$ }).

Where a polynucleotide is double-stranded, both strands of the duplex, either individually or in combination, are encompassed by the present disclosure. Where the polynucleotide is single-stranded, it is to be understood that the complementary sequence of that polynucleotide is also disclosed and encompassed.

Polynucleotides which are not 100% homologous to the sequences disclosed here but fall
20 within the disclosure can be obtained in a number of ways. Other variants of the sequences described herein may be obtained for example by probing DNA libraries made from a range of individuals, for example individuals from different populations. In addition, other viral/bacterial, or cellular homologues particularly cellular homologues found in mammalian cells (e.g. rat,
25 mouse, bovine and primate cells), may be obtained and such homologues and fragments thereof in general will be capable of selectively hybridising to the sequences shown in the sequence listing herein. Such sequences may be obtained by probing cDNA libraries made from or genomic DNA libraries from other animal species, and probing such libraries with probes comprising all or part of SEQ ID NO: 1 under conditions of medium to high stringency. Similar
30 considerations apply to obtaining species homologues and allelic variants of mutant GnT 1.

The polynucleotides described here may be used to produce a primer, e.g. a PCR primer, a primer for an alternative amplification reaction, a probe e.g. labelled with a revealing label by conventional means using radioactive or non-radioactive labels, or the polynucleotides may be cloned into vectors. Such primers, probes and other fragments will be at least 15, such as at least 20, for example at least 25, 30 or 40 nucleotides in length, and are also encompassed by the term polynucleotides as used herein. Fragments may be less than 500, 200, 100, 50 or 20 nucleotides in length.

Polynucleotides such as a DNA polynucleotides and probes may be produced recombinantly, synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques.

In general, primers will be produced by synthetic means, involving a step wise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art.

Longer polynucleotides will generally be produced using recombinant means, for example using PCR (polymerase chain reaction) cloning techniques. This will involve making a pair of primers (e.g. of about 15 to 30 nucleotides) flanking a region of the sequence which it is desired to clone, bringing the primers into contact with mRNA or cDNA obtained from an animal or human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the amplified fragment (e.g. by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector.

EXAMPLES

Example 1. Cell Culture

Chinese hamster ovary-K1 (CHO-K1) cells are originally obtained from Dr. Donald K. MacCallum (University of Michigan Medical School, Ann Arbor, MI).

Parental and mutant CHO cells including JW152 cells are cultured in Dulbecco's Modified Eagle Media (DMEM) supplemented with 10% fetal bovine serum (FBS) at 37°C in a humidified incubator with 5% CO₂.

Lec1.3 cells (Lec 1 cells) are kindly provided by Dr. P. Stanley (Albert Einstein College of Medicine, NY) and cultured in α -MEM (Gibco) supplemented with Proline (40 mg/L) (Invitrogen/Gibco) and 10% FBS.

Example 2. Isolation of RCA-I-Resistant CHO Cells

5 CHO-K1 cells are cultured in 6-well plates to confluence before culture media is changed to serum-free DMEM. *Ricinus communis* agglutinin I (RCA-I, *EY Laboratories*) is added into the media to reach a final concentration of 10 μ g/ml and incubated with cells overnight.

10 Then the serum-free DMEM containing RCA-I is replaced with fresh DMEM with 10% FBS. Nine days later, colonies of the CHO cells that survived the RCA-I treatment are picked and cultured in 24 well plates.

 These cells are then subjected to agglutination tests using RCA-I to confirm the mutant cells no longer react with RCA-I.

Example 3. Expression Constructs

15 The coding regions for human erythropoietin (EPO) and several glycosylation related genes are cloned into the pcDNA3.1 vector (Invitrogen).

 A DNA fragment that contained the open reading frame of human EPO followed by the internal ribosome entry site (IRES) of the Encephalomyocarditis Virus (EMCV) and the coding region for Chinese hamster N-acetylglucosaminyltransferase I (EPO-IRES-GnT I, or
20 EIG in short) is also cloned into pcDNA3.1. The resulting vector that expresses both EPO and GnT I is called pEIG. To ensure efficient translation, a Kozak consensus sequence (GCCACC) is placed upstream of the translation start codon ATG in each construct.

Example 4. Transient Expression of Recombinant Human EPO in Parental and Mutant CHO Cell Lines

25 Unless specified, 1 μ g of DNA construct that expresses EPO and 1 μ g of DNA construct that expresses GnT I or another glycosylation related genes are co-transfected into the wild type or JW-152 mutant cells with Lipofectamine (Invitrogen) according to the manufacturer's protocols.

Two days after transfection, conditioned culture media from the transfected cells are collected. The concentrations of recombinant EPO in each transfection sample are determined by standard ELISA using EPO ELISA kits (Roche).

Example 5. Isoelectric Focusing (IEF) Analysis of EPO Produced by Wild Type CHO Cells and JW-152 CHO Cells

The sialylation patterns of EPO in different samples are analyzed by IEF followed by Western blot as previously described (Schriebl *et al.* 2007, Electrophoresis 28:2100-7). The pH range for IEF is 3 to 10.

Example 6. Molecular Cloning and Sequencing Analysis of GnT I cDNA in RCA-I-Resistant CHO Cells

For each cell line, 1×10^7 cells are pelleted and rinsed in PBS. Total RNA is extracted from the pellet using the RNAqueous kit (Ambion). cDNA is then synthesized through reverse transcription using Moloney Murine Leukemia virus (MMLV) reverse transcriptase (Promega) according to the manufacturer's recommendations

The GnT I amplicon from each cell lines' cDNA is obtained through polymerase chain reaction (PCR) using PFX (Invitrogen). This is then cloned into pcDNA 3.1 expression vector and sequenced. A minimum of four clones from each mutant line are sequenced. All plasmid purifications are carried out using mini or midi-preparation kits from Promega. Constructs are sequenced using ABI Prism 3100 Genetic Analyzer (Applied Biosystems) after cycle sequencing with Big Dye 3.1 (Applied Biosystems).

The results are shown at Table D1 above.

Example 7. Isolation of Stably Transfected CHO Cell Lines

CHO-K1 cells are transfected with a construct that expresses EPO alone whilst mutant JW152 cells are transfected with pEIG to express both EPO and GnT I. The transfected cells are selected with G418 (0.8mg/ml) for two weeks.

Stably transfected cells from the transfected pools are cultured in 96 wells using limiting dilutions. Stable cell lines derived from single transfected cells that expressed EPO are selected using a dot blot analysis and the sialylation patterns of the recombinant EPO expressed by each cell line are analyzed with isoelectric focusing.

Example 8. Results: Isolation and Characterization of RCA-I-Resistant Mutant CHO Cell Lines

RCA-I, a plant lectin that is known specific for β -linked galactose residues on glycoproteins or glycolipids, is used to select glycosylation mutants from CHO K1 cells. Most
5 CHO cells are killed by RCA-I after an overnight incubation.

Nine to ten days later cells that survived RCA-I treatment had grown into single clones. These clones are picked and cultured in 24 well plates. One of the clones is named JW152.

To characterize the genetic defect in this cell line, JW152 cells are transfected with a
10 construct to express human EPO. The recombinant EPO produced by JW152 cells is treated with endoglycosidase H (Endo H).

The results are shown in Figure 1. Figure 1 shows that JW152 expressed EPO is sensitive to Endo H treatment, suggesting that the glycan structure (N-glycans) found on the EPO molecule are of the high mannose type. It is hence suspected that gene(s) from the early
15 part of the glycosylation pathway is defective.

Based on this observation, a complementation test is carried out. In this test, in addition to the EPO construct, JW152 cells are co-transfected with several genes on the N-glycosylation pathway. These genes include glucosidase I, glucosidase II, mannosidase IA, IB, IC, mannosidase II, GnT I and GnT II. EPO samples produced by JW152 cells in these co-
20 transfection experiments are analyzed by isoelectric focusing (IEF) followed by Western blot.

The results are shown in Figure 2. The lane on the far left (CHO WT) shows the EPO expressed by the wild type CHO cells as a control. The lane under CHO-JW152 shows the EPO produced by the JW152 cells, showing the incomplete glycosylation of EPO. The rest of the lanes show the EPO produced in JW152 cells that are co-transfected with a different
25 glycosylation related gene as indicated.

Among the genes tested, only GnT I is able to restore the sialylation pattern of EPO produced in JW152 cells. These results suggest that JW152 cells lack functional GnT I gene.

Another important observation made from this IEF/Western blot assay is that in the presence of GnT I, JW152 cells sialylate EPO much better than the wild type cells (comparing with EPO produced by the wild type cells).

To confirm that this difference is due to a unique feature of JW152 cells rather than the result of over expressing GnT I, three CHO cell lines are analyzed by co-transfecting them with a GnT I construct. The recombinant EPO in conditioned media are collected and analyzed by IEF.

The results are shown in Figure 3. GnT I did not improve EPO sialylation in wild CHO cells (comparing lanes CHO-WT and CHO-WT + GnT I). The EPO sialylation patterns of CHO-K1 with and without co-expression of GnT I appear to be the same and so, the overexpression of GnT I is not responsible for the betterment in sialylation seen in the lane labeled CHO-K1 plus GnT1.

GnT I dramatically improved EPO sialylation in JW152 cells (Lane JW152 + GnT I). In the presence of GnT I, JW152 cells sialylate EPO much better than wild type CHO cells (comparing lanes CHO-WT and CHO-JW152 + GnT I).

Figure 3 shows that, in the presence of GnT I, JW152 cells sialylate recombinant proteins much better than the wild type cells. Treating samples with neuraminidase, which cleaves off sialic acid result in EPO bands in the acidic region being reduced to the basic region after cleaving off sialic acid, shows that the higher sialylated forms of EPO are indeed focused in the acidic region of the gel.

Lec1 cells are previously isolated by P. Stanley and they are known to lack GnT I activity. The IEF pattern of EPO produced by Lec1 cells (Lec1) is similar to that of EPO produced by JW152. EPO produced by Lec1 cells that are co-transfected with GnT I show a similar sialylation pattern with that produced by the wild type CHO cells.

This is shown in Figure 4. Thus, Figure 4 shows that EPO expression in the presence of functional GnT I of previously published Lec 1 mutant which has a GnT I defect is also highly sialylated.

Example 9. Results: Stable Expression of Recombinant EPO in JW152 Cells in the Presence of GnT I

Data presented in Figure 2 and Figure 3 are all from transiently transfected cells. To reveal the sialylation pattern of EPO produced by stably transfected cells, JW152 cells are
5 transfected with pEIG and selected with G418.

After two weeks of selection, single clones are picked and cultured in 24 well plates for two more weeks under the selection pressure of G418. EPO produced by 10 randomly picked such stably transfected clones are analyzed by IEF.

As shown in Figure 10, all the EPO samples produced by different clones are highly
10 sialylated. The sialylation patterns are very similar to that of EPO produced by transiently transfected cells shown in Figure 3.

The results shown in Figure 11 demonstrate that all the EPO samples produced by different stable clones are vary in overall sialylation but in general are not as well sialylated as those in Figure 10.

15 Example 10. Results: All the CHO cells that Survived RCA-I Treatment Have Dysfunctional GnT I Gene and they all Sialylate their Proteins Better when Co-transfected with GnT I

RCA-I is known to specifically bind β -Gal residues. Theoretically, as long as the cells do not express β -Gal as the terminal sugar on their surface glycoproteins, they should survive
20 the RCA-I treatment and be isolated as a mutant clones.

We isolated more than 100 mutant CHO clones using RCA-I and hoped that they should carry genetic mutations in different genes. Surprisingly, when the complementation test described in Figure 2 is carried out, all the clones that survived RCA-I treatment are confirmed to have dysfunctional GnT I genes.

25 Furthermore, as shown in Figure 9, EPO produced in these clones are all highly sialylated when co-transfected with GnT I. The mRNA is isolated from some of the mutant lines and reverse transcribed to cDNA. The coding region for GnT I is amplified by PCR using the cDNA as template and cloned into pcDNA3.1.

Sequencing analyses revealed 9 different point mutations in the GnT I gene in these mutant lines, suggesting that they arrived from different original mutant cells (see Table D1 above). Some of them, like JW152, carry a point mutation that results in a premature stop codon. Others carry a mutation that changes an amino acid residue in the GnT I coding region.

- 5 Nine mutant lines each with a different mutation in the GnT I gene are transfected with pEIG. EPO produced by all nine lines are all highly sialylated when analyzed by IEF (data not shown).

Example 10. Sialylation Patterns in JW152 and Other CHO GnT1 Mutants

- 10 As shown in Figure 5, incomplete sialylation of EPO is observed in nine CHO glycosylation mutants selected with RCA lectin, each bearing a different mutation in the GnT1 gene leading to a loss in GnT1 function, as compared to CHO wild-type.

As shown in Figure 6, EPO produced in nine CHO glycosylation mutants, each bearing a different mutation in the GnT1 gene, have a sialylation pattern that is superior to CHO wild-type after rescue.

- 15 Figure 7 shows the results of an experiment with isoelectric focused samples of a different glycoprotein molecule, EPO-Fc, essentially a erythropoietin fusion protein. The experiment was conducted with a protocol as set out above for EPO expression and IEF analysis, only with an expression vector with a different coding sequence.

- 20 The left hand lane shows expression of EPO-Fc in CHO-K1 cells transfected with an expression vector containing EPO-Fc coding sequence. The middle lane shows expression of EPO-Fc in JW152 cells transfected with an expression vector containing EPO-Fc coding sequence. The right hand lane shows expression of EPO-Fc in JW152 cells co-transfected with an expression vector containing EPO-Fc coding sequence and an expression vector containing functional GnT1 coding sequence. As shown in Figure 7, superior sialylation by
25 rescued JW152 is maintained even with a different model glycoprotein.

The results demonstrate that the superior sialylation by rescued JW152 is maintained even with a different model glycoprotein.

Figure 12 shows that the sialylation pattern observed for stable JW152 cell lines in attached culture is maintained in suspension culture.

In conclusion, we have developed a method to isolate novel CHO glycosylation mutant cells. As long as RCA-I is used to treat CHO cells, the surviving cells will have a genetic defect in their GnT I gene. When co-transfected with GnT I, these RCA-I-resistant CHO cells express recombinant glycoproteins with highly sialylated N-glycans.

5 **Example 11. HPAEC Chromatogram**

Figure 8 is a photograph showing that HPAEC chromatogram showing better sialylated glycans cleaved from EPO-Fc expressed in JW152 cells co-expressing functional GnT I. EPO-Fc transiently expressed in CHO-K1 and JW152, cotransfected with GnT I, is purified using affinity purification via a protein A-bound chromatography column. Glycans on
10 the equal amounts of EPO-Fc are cleaved by treatment with Peptide: N-Glycosidase F (PNGase F), and separated, using High pH Anion Exchange Chromatography (HPAEC), according to their number of sialic acids attached to the glycans.

The chromatogram shows distinctly higher peaks for the JW152 sample in the 4S group and a lower peak in the 1S group when comparing with the CHO-K1 sample.

15 **REFERENCES**

Weikert et al. (1999) Engineering Chinese hamster ovary cells to maximize sialic acid content of recombinant glycoproteins. *Nature Biotechnology* 17:1116.

Each of the applications and patents mentioned in this document, and each document cited or referenced in each of the above applications and patents, including during the
20 prosecution of each of the applications and patents (“application cited documents”) and any manufacturer’s instructions or catalogues for any products cited or mentioned in each of the applications and patents and in any of the application cited documents, are hereby incorporated herein by reference. Furthermore, all documents cited in this text, and all documents cited or referenced in documents cited in this text, and any manufacturer’s
25 instructions or catalogues for any products cited or mentioned in this text, are hereby incorporated herein by reference.

Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific
30 preferred embodiments, it should be understood that the invention as claimed should not be

unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the claims.



A global bioscience nonprofit
organization dedicated to biological
standards and biodiversity

IP, Licensing and Services
10801 University Boulevard
Manassas, Virginia 20110-2209 USA
Telephone: (800) 638-6597
Facsimile: (703) 334-2932
Internet: <http://www.atcc.org>

**BUDAPEST RESTRICTED CERTIFICATE OF DEPOSIT
BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF
THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE
INTERNATIONAL FORM
RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT ISSUED PURSUANT TO RULE 7.3
AND VIABILITY STATEMENT ISSUED PURSUANT TO RULE 10.2**

The American Type Culture Collection (ATCC®) has received your deposit of seeds/strain(s)/strain(s) in connection with the filing of an application for patent. The following information is provided to fulfill Patent Office requirements.

To: Agency for Science, Technology and Research
Dr. Song Zhiwei
20 Biopolis way #06-01
Centros, SG
Singapore 138668

Deposited on Behalf of: Agency for Science, Technology and Research

Date of Receipt of seeds/strain(s) by the ATCC®: December 11, 2008

Identification Reference by Depositor:

ATCC ®Patent Deposit Designation:

Chinese Hamster ovary cells

JW152

PTA-9657

The ATCC® understands that:

1. The deposit of these seeds/strain(s) does not grant ATCC® a license, either express or implied, to infringe the patent, and our release of these seeds/strain(s) to others does not grant them a license, either express or implied, to infringe the patent.
2. If the deposit should die or be destroyed during the effective term of the patent, it shall be your responsibility to replace it with viable material. It is also your responsibility to supply a sufficient quantity for distribution for the deposit term. ATCC® will distribute and maintain the material for 30 years or 5 years following the most recent request for the deposit, whichever is longer. The United States and many other countries are signatory to the Budapest Treaty.

Prior to the issuance of a U.S. Patent, the ATCC® agrees in consideration for a one-time service charge, not to distribute these seeds/strain(s) or any information relating thereto or to their deposit except as instructed by the depositor or relevant patent office. After relevant patent issues we are responsible to release the seeds/strain(s) and they will be made available for distribution to the public without any restrictions. We will inform you of requests for the seeds/strain(s) for 30 years from date of deposit.

Doc ID: 20127
Revision: 1
Effective Date: 11/06/2008



A global bioscience nonprofit
organization dedicated to biological
standards and biodiversity

IP, Licensing and Services
10801 University Boulevard
Manassas, Virginia 20110-2209 USA
Telephone: (800) 638-6597
Facsimile: (703) 334-2932
Internet: <http://www.atcc.org>

The deposit was tested January 8, 2009 and on that date, the seeds/strain(s) were viable
International Depository Authority: American Type Culture Collection (ATCC®), Manassas, VA, USA

Signature of person having authority to represent ATCC®:

January 9, 2009

ATCC® Patent Depository

Date

cc: Dr. Khoo Chong Yee

Doc ID: 20127
Revision: 1
Effective Date: 11/06/2008

CLAIMS

1. A method of expressing a protein of interest from a Chinese Hamster Ovary (CHO) cell, the method comprising:
 - (a) introducing a nucleic acid encoding a protein of interest into a Chinese Hamster Ovary (CHO) cell comprising a mutation in the GnT 1 gene (GenBank Accession Number AF343963) leading to loss of GnT 1 function, which CHO cell further comprises an introduced nucleic acid sequence encoding functional GnT I; and
 - (b) allowing the protein of interest to be expressed from the CHO cell or a descendant thereof; and
 - (c) purifying the protein.
2. A method according to Claim 1, in which the protein of interest has higher sialylation than a protein of interest expressed by a wild-type Chinese Hamster Ovary cell.
3. A method according to Claim 1 or 2, in which the protein of interest has a pKa of 4 or less, 3.5 or less, 3 or less, 2.5 or less or 2 or less or a Z-number of greater than 150, such as greater than 160, greater than 170, greater than 180, such as greater than 190, greater than 200, greater than 210, such as greater than 220 or greater than 230, or both.
4. A method according to Claim 1, 2 or 3, in which the protein of interest comprises a glycoprotein, erythropoietin (EPO), interferon- γ (IFN- γ) or Factor VIII.
5. A method according to Claim 4, in which the nucleic acid sequence encoding the protein of interest and the nucleic acid sequence encoding functional GnT I are comprised in one expression vector, which is preferably stably transfected into the CHO cell.
6. A method according to any preceding claim, in which (i) the mutation in the GnT I gene is selected from the following mutations at the specified position of a GnT I sequence (GenBank accession number: AF343963): (a) a C to T transition at position 1015; (b) a G to C transversion at position 1300; (c) an A to C transversion at position 638; (d) a C to G transversion at position 784; (e) a T to A transversion at position 811; (f) an insertion at position 706 resulting in a frame shift from position 236 of the encoded amino acid sequence; (g) a G to A transition at position 246; (h) a G to A transition at position 258; or (i) an A to T transversion at position 859, or (ii) the Chinese Hamster Ovary (CHO) cell expresses a GnT I

protein comprising one or more of the following mutations at the specified position of a GnT I sequence (GenBank accession number: AF343963): (a) Ala to Pro at position 434; (b) Asp → Ala at position 213; (c) Arg → Gly at position 262; (d) Trp → Arg at position 271; (e) frame shift from position 236 resulting from an insertion at position 706 of an encoding GnT I nucleic acid sequence; (f) Gln→STOP at position 339; (g) Trp→STOP at position 82; (h) Trp→STOP at position 86; or (i) Lys→STOP at position 287.

7. A method according to any preceding claim, in which the Chinese Hamster Ovary (CHO) cell comprises a GnT I nucleic acid sequence shown in SEQ ID NO: 1 or expresses a GnT I protein as shown in SEQ ID NO: 2.

8. A method according to any preceding claim, in which the CHO cell is comprised in a cell line, such as a JW152 cell line (deposited at ATCC under the Budapest Treaty as accession number PTA-9657), which has preferably been adapted to suspension culture or growth in semi-solid medium.

9. A method according to any preceding claim, in which the Chinese Hamster Ovary (CHO) cell comprising a mutation in the GnT I gene is obtainable by selection with *Ricinus communis* agglutinin I (RCA-I).

10. A method according to Claim 9, in which the selection comprises culturing CHO cells in the presence of *Ricinus communis* agglutinin I (RCA-I) and selecting cells which survive the culture, in which for example: (a) the CHO cells are exposed to RCA-I at a concentration of between 0.1µg/ml to 100µg/ml, for example up to 50 µg/ml or up to 20µg/ml, such as 10µg/ml or 5µg/ml; or (b) the CHO cells are exposed RCA-I for a period of from an hour, a few hours (such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12), overnight, to a few days, such as 2 days or 3 days, such as overnight, optionally further comprising selecting cells which do not react with RCA-I in an agglutination test.

11. A CHO cell comprising a mutation in the GnT I gene selected from the following mutations at the specified position of a GnT I sequence (GenBank accession number: AF343963): (a) a C to T transition at position 1015; (b) a G to C transversion at position 1300; (c) an A to C transversion at position 638; (d) a C to G transversion at position 784; (e) a T to A transversion at position 811; (f) an insertion at position 706 resulting in a frame shift from position 236 of the encoded amino acid sequence; (g) a G to A transition at position 246; (h) a

G to A transition at position 258; or (i) an A to T transversion at position 859, in which the CHO cell further comprises a nucleic acid encoding a protein of interest comprising a glycoprotein in an expression vector and a nucleic acid encoding a functional GnT I sequence (GenBank Accession Number AF343963), preferably comprised in one expression vector.

12. A CHO cell according to Claim 11, in which the protein of interest comprises a glycoprotein, erythropoietin (EPO), interferon- γ (IFN- γ) or Factor VIII.

13. A CHO cell according to Claim 11 or 12, in which the CHO cell comprises a JW152 (deposited at ATCC under the Budapest Treaty as accession number PTA-9657).

14. A method of providing a CHO cell or cell line comprising a mutation in the GnT I gene (GenBank accession number: AF343963), the method comprising culturing CHO cells in the presence of *Ricinus communis* agglutinin I (RCA-I), in which a recombinant protein of interest expressed by the resulting CHO cell or cell line in the presence of an introduced nucleic acid encoding functional GnT I comprises a higher sialylation than a protein of interest expressed by a wild-type Chinese Hamster Ovary cell.

15. A method according to Claim 14, in which: (a) the CHO cells are exposed to RCA-I at a concentration of between 0.1 $\mu\text{g/ml}$ to 100 $\mu\text{g/ml}$, for example up to 50 $\mu\text{g/ml}$ or up to 20 $\mu\text{g/ml}$, such as 10 $\mu\text{g/ml}$ or 5 $\mu\text{g/ml}$; or (b) the CHO cells are exposed RCA-I for a period of from an hour, a few hours (such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12), overnight, to a few days, such as 2 days or 3 days, such as overnight.

16. A method according to Claim 14 or 15, further comprising introducing a nucleic acid encoding a protein of interest and a nucleic acid sequence encoding functional GnT I sequence (GenBank Accession Number AF343963), preferably comprised in one expression vector.

SEQUENCE LISTING

<110> Agency for Science, Technology and Research (A*STAR)

<120> Chinese Hamster Ovary Cell Lines

<130> P000041W01

<150> US 61/098,270

<151> 2008-09-19

<150> US 61/183,647

<151> 2009-06-03

<160> 2

<170> PatentIn version 3.3p

<210> 1

<211> 1344

<212> DNA

<213> Cricetulus griseus

<220>

<221> C to T point mutation

<222> (1015)..(1015)

<223> C to T point mutation

<400> 1

atgctgaaga agcagtctgc agggccttggtg ctttgggggtg ctatcctctt tgtgggctgg	60
aatgccctgc tgctcctctt cttctggaca cgcccagccc ctggcaggcc cccctcagat	120
agtgtatcgc atgatgaccc tgccagcctc acccgtgagg tgttccgcct ggctgaggac	180
gctgaggtgg agttggagcg gcagcggggg ctgttgacgc aaatcaggga gcatcatgct	240
ttgtggagac agaggtggaa agtgcccacc gtggcccctc cagcctggcc ccgtgtgcct	300
gcgacccctt caccagccgt gatcccatc ctggtcattg cctgtgaccg cagcactgtc	360
cggcgctgct tggataagtt gttgcactat cggccctcag ctgagcattt ccccatcatt	420
gtcagccagg actgcgggca cgaagagaca gcacagggtca ttgcttccta tggcagtgca	480
gtcacacaca tccggcagcc agacctgagt aacatcgctg tgccccaga ccaccgcaag	540
ttccagggtt actacaagat cgccaggcac taccgctggg cactgggcca gatcttcaac	600
aagttcaagt tcccagcagc tgtggtagtg gaggacgac tggaggtggc accagacttc	660

```

tttgagtact tccaggccac ctaccactg ctgagaacag acccctccct ttggtgtgtg 720
tctgcttgga atgacaatgg caaggagcag atggtagact caagcaaacc tgagctgctc 780
tatcgaacag acttttttcc tggccttggc tggctgctga tggctgagct gtggacagag 840
ctggagccca agtggcccaa ggccttctgg gatgactgga tgcgcagacc tgagcagcgg 900
aagggggcggg cctgtattcg tccagaaatt tcaagaacga tgacctttgg ccgtaagggt 960
gtgagccatg ggcagttctt tgatcagcat cttaagttca tcaagctgaa ccagtagttc 1020
gtgtctttca cccagttgga tttgtcatac ttgcagcggg aggcttatga ccgggatttc 1080
cttgcccgtg tctatagtgc cccctgcta caggtggaga aagtgaggac caatgatcag 1140
aaggagctgg gggaggtgcg ggtacagtac actagcagag acagcttcaa ggcctttgct 1200
aaggccctgg gtgtcatgga tgacctcaag tctggtgtcc ccagagctgg ctaccggggc 1260
gttgtcactt tccagttcag gggtcgacgt gtccacctgg cacccccaca aacctgggaa 1320
ggctatgatc ctagctggaa ttag 1344

```

<210> 2

<211> 338

<212> PRT

<213> *Cricetulus griseus*

<400> 2

```

Met Leu Lys Lys Gln Ser Ala Gly Leu Val Leu Trp Gly Ala Ile Leu
1           5           10           15
Phe Val Gly Trp Asn Ala Leu Leu Leu Leu Phe Phe Trp Thr Arg Pro
          20           25           30
Ala Pro Gly Arg Pro Pro Ser Asp Ser Ala Ile Asp Asp Asp Pro Ala
          35           40           45
Ser Leu Thr Arg Glu Val Phe Arg Leu Ala Glu Asp Ala Glu Val Glu
          50           55           60
Leu Glu Arg Gln Arg Gly Leu Leu Gln Gln Ile Arg Glu His His Ala
65           70           75           80
Leu Trp Arg Gln Arg Trp Lys Val Pro Thr Val Ala Pro Pro Ala Trp
          85           90           95
Pro Arg Val Pro Ala Thr Pro Ser Pro Ala Val Ile Pro Ile Leu Val

```

100	105	110
Ile Ala Cys Asp Arg Ser Thr Val Arg Arg Cys Leu Asp Lys Leu Leu		
115	120	125
His Tyr Arg Pro Ser Ala Glu His Phe Pro Ile Ile Val Ser Gln Asp		
130	135	140
Cys Gly His Glu Glu Thr Ala Gln Val Ile Ala Ser Tyr Gly Ser Ala		
145	150	155
Val Thr His Ile Arg Gln Pro Asp Leu Ser Asn Ile Ala Val Pro Pro		
165	170	175
Asp His Arg Lys Phe Gln Gly Tyr Tyr Lys Ile Ala Arg His Tyr Arg		
180	185	190
Trp Ala Leu Gly Gln Ile Phe Asn Lys Phe Lys Phe Pro Ala Ala Val		
195	200	205
Val Val Glu Asp Asp Leu Glu Val Ala Pro Asp Phe Phe Glu Tyr Phe		
210	215	220
Gln Ala Thr Tyr Pro Leu Leu Arg Thr Asp Pro Ser Leu Trp Cys Val		
225	230	235
Ser Ala Trp Asn Asp Asn Gly Lys Glu Gln Met Val Asp Ser Ser Lys		
245	250	255
Pro Glu Leu Leu Tyr Arg Thr Asp Phe Phe Pro Gly Leu Gly Trp Leu		
260	265	270
Leu Met Ala Glu Leu Trp Thr Glu Leu Glu Pro Lys Trp Pro Lys Ala		
275	280	285
Phe Trp Asp Asp Trp Met Arg Arg Pro Glu Gln Arg Lys Gly Arg Ala		
290	295	300
Cys Ile Arg Pro Glu Ile Ser Arg Thr Met Thr Phe Gly Arg Lys Gly		
305	310	315
Val Ser His Gly Gln Phe Phe Asp Gln His Leu Lys Phe Ile Lys Leu		
325	330	335
Asn Gln		