

Cyclic ether derivatives of pyrazolo[1,5-a]pyrimidine-3-carboxamide

Field of the invention

5 The invention relates to cyclic ether derivatives of pyrazolo[1,5-a]pyrimidine-3-carboxamide of general formula (I) which are inhibitors of phosphodiesterase 2, useful in treating central nervous system diseases and other diseases. In addition, the invention relates to processes for preparing pharmaceutical compositions as well as processes for manufacture the compounds according to the
10 invention.

Background of the invention

Phosphodiesterase 2 (PDE2) inhibitors are promising therapeutic targets for
15 treatment of cognitive impairment in diseases such as Schizophrenia, Alzheimer's disease and depression. Inhibitors of PDE2 have emerged as potential candidates to improve synaptic plasticity and memory function.

Phosphodiesterases (PDE) are expressed in nearly all mammalian cells. To date eleven families of phosphodiesterases have been identified in mammals. It is well
20 established that PDEs are critically involved in cell signalling. Specifically, PDEs are known to inactivate the cyclic nucleotides cAMP and/or cGMP.

PDE2 hydrolyses both, cGMP and cAMP. It is both abundantly expressed in the brain indicating their relevance in CNS function.

The expression of PDE2 in the hippocampus, the cortex and in the striatum indicate
25 an involvement in the mechanism of learning and memory/cognition. This is further supported by the fact that increased levels of both cGMP and cAMP are involved in the process of short and long term potentiation (LTP) forming. Further data support the procognitive effect of PDE2 and a synergistic effect of PDE2 on cognition.

Furthermore, the expression of PDE2 in the nucleus accumbens (part of the striatum),
30 the olfactory bulb, the olfactory tubercle and the amygdala supports additional involvement of PDE2 in the pathophysiology of anxiety and depression. This is supported by in vivo studies.

It is commonly accepted (free drug hypothesis) that unbound or free drug concentration at the site of action is responsible for pharmacological activity in vivo at steady state and, in the absence of active transport, the free drug concentration is the same in any biomembrane.

- 5 For drugs with an intended action in the central nervous system (CNS), it is assumed that unbound drug in interstitial spaces (ISF) in the brain is in direct contact or in equilibrium with the site of action. Because cerebrospinal fluid (CSF) is in direct contact with the brain tissue, it is assumed to readily equilibrate with brain interstitial fluid concentration so that CSF concentration is used as a common surrogate
- 10 measure for drug unbound concentration in pre-clinical pharmacology studies. Accordingly, for compounds with an intended action in the central nervous system it is important that they reach a high CSF concentration and a high CSF to plasma ratio in order to have high pharmacological activity in the CNS.
- 15 At steady state and in the absence of active transport, the unbound brain concentration can also be estimated with the experimentally more accessible unbound plasma concentration by measuring the plasma protein binding (PPB) across species.
- 20 High membrane permeability and absence of active transport process at the BBB (blood brain barrier) together with plasma/brain tissue binding are recognised as the primary determinant of drug disposition within CNS.

- High metabolic stability is desirable in order to achieve significant exposure of a drug
- 25 within the body.

- Several families of PDE2 inhibitors are known. Imidazotriazinones are claimed in WO 2002/068423 for the treatment of e.g. memory deficiency, cognitive disorders, dementia and Alzheimer's disease. Oxindoles are described in WO 2005/041957 for
- 30 the treatment of dementia. Further inhibitors of PDE2 are known from WO 2007/121319 for the treatment of anxiety and depression, from WO 2013/034761, WO 2012/104293 and WO2013/000924 for the treatment of neurological and psychiatric disorders, from WO 2006/072615, WO 2006/072612, WO 2006/024640

and WO 2005/113517 for the treatment of arthritis, cancer, edema and septic shock, from WO 2005/063723 for the treatment of renal and liver failure, liver dysfunction, restless leg syndrome, rheumatic disorders, arthritis, rhinitis, asthma and obesity, from WO 2005/041957 for the treatment of cancer and thrombotic disorders, from
5 WO 2006/102728 for the treatment of angina pectoris and hypertension from WO 2008/043461 for the treatment of cardiovascular disorders, erectile dysfunction, inflammation and renal failure and from WO 2005/061497 for the treatment of e.g. dementia, memory disorders, cancer and osteoporosis.

Benzodiazepine like PDE2 inhibitors are described in WO 2005/063723 for the
10 general treatment of CNS diseases including anxiety, depression, ADHD, neurodegeneration, Alzheimer's disease and psychosis.

Newer PDE2 inhibitor families are described in WO 2015/096651, WO 2015/060368 and WO 2015/012328.

15 Aim of the invention

It has now been found that compounds of the present invention according to general formula (I) are effective inhibitors of phosphodiesterase 2.

Besides the inhibition property toward phosphodiesterase 2 enzymes, the compounds of the present invention provide further advantageous properties such as
20 high selectivity with regard to PDE 10, low plasma protein binding across species, high CSF to plasma ratio, adequate tissue permeability and high metabolic stability.

For example the compounds of the present invention show low plasma protein binding across species and as a consequence high fraction unbound in plasma, high
25 concentration in cerebrospinal fluid (CSF) and have a high CSF to plasma ratio, which translates in lower efficacious doses of the compounds for disease treatment and as a consequence in further potential advantages such as minimization of side effects . Furthermore, compounds of the present inventions show good metabolic stability both in rodents and non rodents species, good membrane permeability with
30 no active transport at the BBB. In addition the compounds of the present invention have very high IC50 values for PDE 10.

Accordingly, one aspect of the invention refers to compounds according to formula (I), or salts thereof as inhibitors of phosphodiesterase 2.

Another aspect of the invention refers to compounds according to formula (I), or pharmaceutically acceptable salts thereof as inhibitors of phosphodiesterase 2 and reaching high concentrations in cerebrospinal fluid (CSF) and/or having high CSF to plasma ratio.

Another aspect of the invention refers to compounds according to formula (I), or pharmaceutically acceptable salts thereof as inhibitors of phosphodiesterase 2 with low plasma protein binding and thus high fraction unbound across species.

Another aspect of the invention refers to compounds according to formula (I), or pharmaceutically acceptable salts thereof as inhibitors of phosphodiesterase 2 and showing good membrane permeability and low to moderate in vitro efflux.

Another aspect of the invention refers to according to formula (I), or pharmaceutically acceptable salts thereof as inhibitors of phosphodiesterase 2 and showing good metabolic stability.

In a further aspect this invention relates to pharmaceutical compositions, containing at least one compound according to formula (I), or pharmaceutically acceptable salts thereof, optionally together with one or more inert carriers and/or diluents.

A further aspect of the present invention relates to compounds according to formula (I), or pharmaceutically acceptable salts thereof or pharmaceutical compositions comprising compounds according to formula (I), or pharmaceutically acceptable salts thereof for the use in the prevention and/or treatment of disorders associated with PDE2 hyperactivity and/or cAMP and/or cGMP hypofunction.

Another aspect of the invention relates to processes of manufacture of the compounds of the present invention.

A further aspect of the present invention relates to compounds according to formula (I), or pharmaceutically acceptable salts thereof or pharmaceutical compositions

comprising compounds according to formula (I), or pharmaceutically acceptable salts thereof for the use in the prevention and/or treatment of diseases or conditions which can be influenced by inhibition of PDE2 hyperactivity and/or cAMP and/or cGMP hypofunction, such as (1) disorders comprising the symptom of cognitive deficiency; (2) organic, including symptomatic, mental disorders, dementia; (3) mental retardation; (4) mood affective disorders; (5) neurotic, stress-related and somatoform disorders including anxiety disorders; (6) behavioural and emotional disorders with onset usually occurring in childhood and adolescence, attention deficit hyperactivity syndrome (ADHD) including Autism spectrum disorders; (7) disorders of psychological development, developmental disorders of scholastic skills; (8) schizophrenia and other psychotic disorders; (9) disorders of adult personality and behaviour; (10) mental and behavioural disorders due to psychoactive substance use; (11) extrapyramidal and movement disorders; (12) episodic and paroxysmal disorders, epilepsy; (13) Systemic atrophies primarily affecting the central nervous system, ataxia; (14) Behavioural syndromes associated with physiological disturbances and physical factors; (15) sexual dysfunction comprising excessive sexual drive; (16) factitious disorders; (17) obsessive-compulsive disorders; (18) depression; (19) neuropsychiatric symptoms (e.g. depressive symptoms in Alzheimer's disease); (20) mixed dementia; (21) cognitive impairment in schizoaffective disorder; (22) cognitive impairment in bipolar disorder and (23) cognitive impairment in major depressive disorder.

In addition, the compounds of the present invention can be used for the treatment, amelioration and / or prevention of cognitive impairment being related to perception, concentration, cognition, learning, attention or memory.

In addition, the compounds of the present invention can be used for the treatment amelioration and / or prevention of cognitive impairment being related to age-associated learning and memory impairments, age-associated memory losses, vascular dementia, craniocerebral trauma, stroke, dementia occurring after strokes (post stroke dementia), post-traumatic dementia, general concentration impairments, concentration impairments in children with learning and memory problems, Alzheimer's disease, Lewy body dementia, dementia with degeneration of the frontal lobes, including Pick's syndrome, Parkinson's disease, progressive nuclear palsy,

dementia with corticobasal degeneration, amyotrophic lateral sclerosis (ALS), Huntington's disease, multiple sclerosis, thalamic degeneration, Creutzfeld-Jacob dementia, HIV dementia, schizophrenia with dementia or Korsakoff's psychosis. In addition, the compounds of the present invention can be used for the treatment of

5 Alzheimer's disease.

In addition compounds of the present invention can be used for the treatment of pain disorders, including but not limited to inflammatory, neuropathic and osteoarthritic pain.

In addition, the compounds of the present invention can be used for the treatment of


10 sleep disorders, bipolar disorder, metabolic syndrome, obesity, diabetes mellitus, hyperglycemia, dyslipidemia, impaired glucose tolerance, or a disease of the testes, brain, small intestine, skeletal muscle, heart, lung, thymus or spleen.

Other aims of the present invention will become apparent to the skilled man directly

15 from the foregoing and following remarks.

Summary


In a first aspect the present disclosure relates to a compound of formula (I)



20 (I)

wherein

A is selected from the group A^a consisting of



wherein above mentioned groups are substituted with one R⁵ and one R⁶;

- R^1 is selected from the group R^{1a} consisting of halogen, C_{1-3} -alkyl- and C_{3-6} -cycloalkyl- wherein the above mentioned C_{1-3} -alkyl-, and C_{3-6} -cycloalkyl- groups may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of halogen, NC- and HO-;
- R^2 is selected from the group R^{2a} consisting of aryl and heteroaryl, wherein the above mentioned aryl and heteroaryl-groups may optionally be substituted with 1 to 5 substituents R^4 ;
- R^3 is selected from the group R^{3a} consisting of H- and C_{1-3} -alkyl-, wherein the above mentioned C_{1-3} -alkyl-groups may optionally be substituted with 1 to 7 substituents independently from each other selected from the group consisting of halogen;
- R^4 is independently from each other selected from the group R^{4a} consisting of halogen, NC-, HO-, C_{1-4} -alkyl- and C_{1-3} -alkyl-O- wherein the above mentioned C_{1-4} -alkyl- and C_{1-3} -alkyl-O-groups may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of HO- and F-;


5


- R^5 is selected from the group R^{5a} consisting of H-, halogen, NC-, HO- and C_{1-3} -alkyl-, wherein the above mentioned C_{1-3} -alkyl-group may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of HO- and F- or R^5 and R^6 together form an group O=;

R^6 is selected from the group R^{6a} consisting of
H-, halogen, NC-, HO- and C_{1-3} -alkyl-,
wherein the above mentioned C_{1-3} -alkyl-group may optionally be substituted
with 1 to 5 substituents independently selected from the group consisting of
HO- and F-
or R^5 and R^6 together form a group O=;
or a salt thereof.


In another aspect, the present disclosure relates to a compound selected from the
group consisting of


5







6d












6h



In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:




5


In another aspect, the present disclosure relates to a compound of formula:




10 In another aspect, the present disclosure relates to a compound of formula:



In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:




5

In another aspect, the present disclosure relates to a compound of formula:




6k

In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:




5


In another aspect, the present disclosure relates to a compound of formula:



In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:




5

In another aspect, the present disclosure relates to a compound of formula:




10 In another aspect, the present disclosure relates to a compound of formula:




6m

In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:




5


In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:




5 In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:



In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:




5


In another aspect, the present disclosure relates to a compound of formula:



In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:




5

In another aspect, the present disclosure relates to a compound of formula:




10

In another aspect, the present disclosure relates to a compound of formula:




In another aspect, the present disclosure relates to a compound of formula:


5



In another aspect, the present disclosure relates to a compound of formula:



In another aspect, the present disclosure relates to a compound of formula:



In another aspect, the present disclosure relates to a pharmaceutically acceptable
5 salt of any of the compounds of the present disclosure.

In another aspect, the present disclosure relates to a compound of the present
disclosure, or a pharmaceutically acceptable salt thereof for use as a medicament.


In another aspect, the present disclosure relates to pharmaceutical compositions
10 containing at least one compound of the present disclosure or a pharmaceutically
acceptable salt thereof together with one or more pharmaceutically acceptable
carrier.

In another aspect, the present disclosure relates to a compound of the present
15 disclosure, or a pharmaceutically acceptable salt thereof or a pharmaceutical
composition of the present disclosure for use in the prevention or treatment of (1)
disorders comprising the symptom of cognitive deficiency; (2) organic, including
symptomatic, mental disorders, dementia; (3) mental retardation; (4) mood [affective]
disorders; (5) neurotic, stress-related and somatoform disorders including anxiety
20 disorders; (6) behavioural and emotional disorders with onset usually occurring in
childhood and adolescence, attention deficit hyperactivity syndrome (ADHD) and
Autism spectrum disorders; (7) disorders of psychological development,
developmental disorders of scholastic skills; (8) schizophrenia and other psychotic
disorders; (9) disorders of adult personality and behaviour; (10) mental and
25 behavioural disorders due to psychoactive substance use; (11) extrapyramidal and
movement disorders; (12) episodic and paroxysmal disorders, epilepsy;

(13) Systemic atrophies primarily affecting the central nervous system, ataxia; (14) Behavioural syndromes associated with physiological disturbances and physical factors; (15) sexual dysfunction comprising excessive sexual drive; (16) factitious disorders; (17) Treatment, amelioration and / or prevention of cognitive impairment being related to perception, concentration, cognition, learning or memory; (18) Treatment amelioration and / or prevention of cognitive impairment being related to age-associated learning and memory impairments; (19) Age-associated memory losses; (20) Vascular dementia; (21) Craniocerebral trauma; (22) Stroke; (23) Dementia occurring after strokes (post stroke dementia); (24) Post-traumatic dementia; (25) General concentration impairments; (26) Concentration impairments in children with learning and memory problems; (27) Alzheimer's disease; (28) Lewy body dementia; (29) Dementia with degeneration of the frontal lobes; including Pick's syndrome; (30) Parkinson's disease; (31) Progressive nuclear palsy; (32) Dementia with corticobasal degeneration; (33) Amyotrophic lateral sclerosis (ALS); (34) Huntington's disease; (35) Multiple sclerosis; (36) Thalamic degeneration; (37) Creutzfeld-Jacob dementia; (38) HIV dementia; (38) Schizophrenia with dementia or Korsakoff's psychosis; (39) Sleep disorders; (40) Bipolar disorder; (41) Metabolic syndrome; (42) Obesity; (43) Diabetis mellitus; (44) Hyperglycemia; (45) Dyslipidemia; (46) Impaired glucose tolerance; (47) Disease of the testes, brain, small intestine, skeletal muscle, heart, lung, thymus or spleen; (48) pain disorders; (49) neuropsychiatric symptoms (e.g. depressive symptoms in Alzheimer's disease); (50) mixed dementia; (51) cognitive impairment in schizoaffective disorder; (52) cognitive impairment in bipolar disorder; (53) cognitive impairment in major depressive disorder and (54) cognitive impairment associated with schizophrenia.

Detailed description


In a first aspect the present invention relates to compounds of general formula (I)



5

wherein

A is selected from the group A^a consisting of



wherein above mentioned groups are substituted with one R^5 and one R^6 ;

R^1 is selected from the group R^{1a} consisting of halogen, C_{1-3} -alkyl- and C_{3-6} -cycloalkyl- wherein the above mentioned C_{1-3} -alkyl-, and C_{3-6} -cycloalkyl- groups may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of halogen, NC- and HO-;

R^2 is selected from the group R^{2a} consisting of aryl and heteroaryl, wherein the above mentioned aryl and heteroaryl-groups may optionally be substituted with 1 to 5 substituents R^4 ;

R^3 is selected from the group R^{3a} consisting of H- and C_{1-3} -alkyl-, wherein the above mentioned C_{1-3} -alkyl-groups may optionally be substituted with 1 to 7 substituents independently from each other selected from the group consisting of halogen;

R^4 is independently from each other selected from the group R^{4a} consisting of halogen, NC-, HO-, C_{1-4} -alkyl- and C_{1-3} -alkyl-O- wherein the above mentioned C_{1-4} -alkyl- and C_{1-3} -alkyl-O-groups may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of HO- and F-;

5

R^5 is selected from the group R^{5a} consisting of H-, halogen, NC-, HO- and C_{1-3} -alkyl-, wherein the above mentioned C_{1-3} -alkyl-group may optionally be

substituted with 1 to 5 substituents independently selected from the group consisting of HO- and F-
or R⁵ and R⁶ together form an group O=;

R⁶ is selected from the group R^{6a} consisting of
H-, halogen, NC-, HO- and C₁₋₃-alkyl-,
wherein the above mentioned C₁₋₃-alkyl-group may optionally be
substituted with 1 to 5 substituents independently selected from the
group consisting of HO- and F-
or R⁵ and R⁶ together form a group O=;


or a salt thereof.

- 5 Unless otherwise stated, the groups, residues, and substituents, particularly R¹, R², R³, R⁴ and R⁵ are defined as above and hereinafter. If residues, substituents, or groups occur several times in a compound they may have the same or different meanings. Some preferred meanings of groups and substituents of the compounds according to the invention will be given hereinafter.

10

In a further embodiment of the present invention


A is selected from the group A^b consisting of



wherein above mentioned groups are substituted with with one R⁵
and one R⁶.


In a further embodiment of the present invention

A is selected from the group A^c consisting of




In a further embodiment of the present invention

A is selected from the group A^d consisting of



In a further embodiment of the present invention

A is selected from the group A^e consisting of



5

In a further embodiment of the present invention

R¹ is selected from the group R^{1b} consisting of

F-, Cl-, C₁₋₃-alkyl- and C₃₋₆-cycloalkyl-,

wherein the above mentioned C₁₋₃-alkyl- and C₃₋₆-cycloalkyl-groups may optionally be substituted with 1 to 3 substituents independently selected from the group consisting of F-.

In a further embodiment of the present invention

R¹ is selected from the group R^{1c} consisting of

F-, H₃C- and cyclopropyl-.

10 In a further embodiment of the present invention

R¹ is selected from the group R^{1d} consisting of

H₃C- and cyclopropyl-.

In a further embodiment of the present invention


R^2 is selected from the group R^{2b} consisting of
quinoliny, phenyl and pyridynyl,
wherein the above mentioned quinoline, phenyl and pyridyl-groups
may optionally be substituted with 1 to 5 substituents R^4 .

In a further embodiment of the present invention

R^2 is selected from the group R^{2c} consisting of
phenyl and pyridyl,
wherein the above mentioned phenyl and pyridyl-groups may
optionally be substituted with 1 to 2 substituents R^4 .


5 In a further embodiment of the present invention

R^2 is selected from the group R^{2d} being




In a further embodiment of the present invention

R^2 is selected from the group R^{2e} being



10 In a further embodiment of the present invention

R^2 is selected from the group R^{2f} being



In a further embodiment of the present invention

R^3 is selected from the group R^{3b} consisting of
H-, H_3C -, F_3C -, F_2HC -, FH_2C - and F_3C -.

In a further embodiment of the present invention

R^3 is selected from the group R^{3c} consisting of
H- and H_3C -.

5

In a further embodiment of the present invention

R^3 is selected from the group R^{3d} being H-.

In a further embodiment of the present invention

R^4 is independently from each other selected from the group R^{4b} consisting of
halogen, C_{1-4} -alkyl- and C_{1-3} -alkyl-O-
wherein the above mentioned C_{1-4} -alkyl- and C_{1-3} -alkyl-O-groups may
optionally be substituted with 1 to 5 substituents independently selected
from the group consisting of HO-, and F-.

10 In a further embodiment of the present invention

R^4 is independently from each other selected from the group R^{4c} consisting of
halogen, C_{1-3} -alkyl-, F_3C -O-, F_2HC -O-, FH_2C -O- and H_3C -O-,
wherein the above mentioned C_{1-3} -alkyl-groups may optionally be
substituted with 1 to 5 F-.

In a further embodiment of the present invention

R^4 is independently from each other selected from the group R^{4d} consisting of
F, Cl, Br, F_3C -, F_2HC -, FH_2C -, H_3C -, F_3C -O-, F_2HC -O-, FH_2C -O- and

H₃C-O-.

In a further embodiment of the present invention

R⁴ is independently from each other selected from the group R^{4e} consisting of F, Cl, F₃C-, F₃C-O- and H₃C-O-.

In a further embodiment of the present invention

R⁴ is independently from each other selected from the group R^{4f} consisting of F and F₃C-.

5

In a further embodiment of the present invention

R⁵ is selected from the group R^{5b} consisting of H-, HO- and C₁₋₂-alkyl-,
wherein the above mentioned C₁₋₂-alkyl-group may optionally be substituted with 1 to 5 F-,
or R⁵ and R⁶ together form an group O=.

In a further embodiment of the present invention

R⁵ is selected from the group R^{5c} consisting of H- and HO-.

10 In a further embodiment of the present invention

R⁵ is selected from the group R^{5d} being HO-.

In a further embodiment of the present invention

R⁶ is selected from the group R^{6b} consisting of H- and C₁₋₂-alkyl-,
wherein the above mentioned C₁₋₂-alkyl-group may optionally be substituted with 1 to 5 F-,
or R⁵/R⁶ together form a group O=.

In a further embodiment of the present invention

R^6 is selected from the group R^{6c} consisting of
H and H_3C- ,

wherein the above mentioned methyl-group may optionally be
substituted with 1 to 3 F-.

In a further embodiment of the present invention

R^6 is selected from the group R^{6d} consisting of
H- and H_3C- .

Each A^x , R^{1x} , R^{2x} , R^{3x} , R^{4x} , R^{5x} and R^{6x} represents a characterized, individual

embodiment for the corresponding substituent as described above. Thus given the
above definitions, individual embodiments of the first aspect of the invention are fully
characterized by the term (A^x , R^{1x} , R^{2x} , R^{3x} , R^{4x} , R^{5x} and R^{6x}), wherein for each index
x an individual figure is given that ranges from "a" to the highest letter given above.
All individual embodiments described by the term in parentheses with full permutation
of the indices x, referring to the definitions above, shall be comprised by the present
invention.

The following Table 1 shows such embodiments E-1 to E-39 of the invention that are
considered preferred. Embodiment E-39, represented by the entries in the last row of
Table 1, is the most preferred embodiment.


Table 1: Embodiments E-1 to E-39 of the invention

	A^x	R^{1x}	R^{2x}	R^{3x}	R^{4x}	R^{5x}	R^{6x}
E-1	A^a	R^{1a}	R^{2a}	R^{3a}	R^{4b}	R^{5a}	R^{6a}
E-2	A^a	R^{1a}	R^{2a}	R^{3b}	R^{4b}	R^{5a}	R^{6a}
E-3	A^a	R^{1b}	R^{2b}	R^{3b}	R^{4c}	R^{5a}	R^{6a}
E-4	A^a	R^{1c}	R^{2b}	R^{3b}	R^{4d}	R^{5a}	R^{6a}
E-5	A^a	R^{1c}	R^{2b}	R^{3c}	R^{4e}	R^{5b}	R^{6b}
E-6	A^b	R^{1b}	R^{2b}	R^{3b}	R^{4b}	R^{5a}	R^{6a}
E-7	A^b	R^{1c}	R^{2b}	R^{3c}	R^{4e}	R^{5b}	R^{6b}
E-8	A^c	R^{1c}	R^{2b}	R^{3b}	R^{4b}	R^{5b}	R^{6b}
E-9	A^c	R^{1c}	R^{2b}	R^{3c}	R^{4e}	R^{5b}	R^{6b}
E-10	A^c	R^{1c}	R^{2c}	R^{3b}	R^{4c}	R^{5b}	R^{6b}
E-11	A^c	R^{1d}	R^{2c}	R^{3c}	R^{4d}	R^{5b}	R^{6b}

E-12	A ^c	R ^{1d}	R ^{2c}	R ^{3d}	R ^{4e}	R ^{5b}	R ^{6b}
E-13	A ^c	R ^{1d}	R ^{2c}	R ^{3d}	R ^{4f}	R ^{5c}	R ^{6c}
E-14	A ^c	R ^{1d}	R ^{2d}	R ^{3b}	R ^{4c}	R ^{5b}	R ^{6b}
E-15	A ^c	R ^{1d}	R ^{2d}	R ^{3c}	R ^{4d}	R ^{5b}	R ^{6b}
E-16	A ^c	R ^{1d}	R ^{2d}	R ^{3c}	R ^{4f}	R ^{5d}	R ^{6d}
E-17	A ^c	R ^{1d}	R ^{2d}	R ^{3d}	R ^{4e}	R ^{5b}	R ^{6b}
E-18	A ^c	R ^{1d}	R ^{2d}	R ^{3d}	R ^{4f}	R ^{5c}	R ^{6c}
E-19	A ^c	R ^{1d}	R ^{2e}	R ^{3b}	R ^{4b}	R ^{5b}	R ^{6b}
E-20	A ^c	R ^{1d}	R ^{2e}	R ^{3b}	R ^{4b}	R ^{5c}	R ^{6c}
E-21	A ^c	R ^{1d}	R ^{2e}	R ^{3c}	R ^{4d}	R ^{5b}	R ^{6b}
E-22	A ^c	R ^{1d}	R ^{2e}	R ^{3c}	R ^{4d}	R ^{5c}	R ^{6c}
E-23	A ^c	R ^{1d}	R ^{2e}	R ^{3c}	R ^{4e}	R ^{5b}	R ^{6b}
E-24	A ^c	R ^{1d}	R ^{2e}	R ^{3c}	R ^{4e}	R ^{5c}	R ^{6c}
E-25	A ^c	R ^{1d}	R ^{2e}	R ^{3d}	R ^{4e}	R ^{5b}	R ^{6b}
E-26	A ^c	R ^{1d}	R ^{2e}	R ^{3d}	R ^{4e}	R ^{5c}	R ^{6c}
E-27	A ^c	R ^{1d}	R ^{2f}	R ^{3c}	-	R ^{5d}	R ^{6d}
E-28	A ^c	R ^{1d}	R ^{2f}	R ^{3d}	-	R ^{5d}	R ^{6d}
E-29	A ^d	R ^{1c}	R ^{2c}	R ^{3b}	R ^{4d}	R ^{5b}	R ^{6b}
E-30	A ^d	R ^{1c}	R ^{2d}	R ^{3b}	R ^{4e}	R ^{5c}	R ^{6c}
E-31	A ^d	R ^{1c}	R ^{2e}	R ^{3c}	R ^{4e}	R ^{5d}	R ^{6d}
E-32	A ^d	R ^{1d}	R ^{2f}	R ^{3c}	-	R ^{5d}	R ^{6d}
E-33	A ^e	R ^{1c}	R ^{2c}	R ^{3b}	R ^{4d}	R ^{5b}	R ^{6b}
E-34	A ^e	R ^{1c}	R ^{2d}	R ^{3b}	R ^{4e}	R ^{5c}	R ^{6c}
E-35	A ^e	R ^{1c}	R ^{2e}	R ^{3c}	R ^{4e}	R ^{5d}	R ^{6d}
E-36	A ^e	R ^{1c}	R ^{2f}	R ^{3b}	-	R ^{5b}	R ^{6b}
E-37	A ^e	R ^{1d}	R ^{2f}	R ^{3c}	-	R ^{5c}	R ^{6c}
E-38	A ^e	R ^{1d}	R ^{2f}	R ^{3c}	-	R ^{5d}	R ^{6d}
E-39	A ^e	R ^{1d}	R ^{2f}	R ^{3d}	-	R ^{5d}	R ^{6d}

Accordingly, for example E-1 covers compounds of formula (I),
wherein

A is selected from the group A^a consisting of



wherein above mentioned groups are substituted with one R⁵ and one R⁶;

- R¹ is selected from the group R^{1a} consisting of
halogen, C₁₋₃-alkyl- and C₃₋₆-cycloalkyl-
wherein the above mentioned C₁₋₃-alkyl-, and C₃₋₆-cycloalkyl- groups
may optionally be substituted with 1 to 5 substituents independently
selected from the group consisting of halogen, NC- and HO-;
- R² is selected from the group R^{2a} consisting of
aryl and heteroaryl,
wherein the above mentioned aryl and heteroaryl-groups may
optionally be substituted with 1 to 5 substituents R⁴;
- R³ is selected from the group R^{3a} consisting of
H- and C₁₋₃-alkyl-,
wherein the above mentioned C₁₋₃-alkyl-groups may optionally be
substituted with 1 to 7 substituents independently from each other
selected from the group consisting of halogen;
- R⁴ is independently from each other selected from the group R^{4b} consisting of
halogen, C₁₋₄-alkyl- and C₁₋₃-alkyl-O-
wherein the above mentioned C₁₋₄-alkyl- and C₁₋₃-alkyl-O-groups may
optionally be substituted with 1 to 5 substituents independently selected
from the group consisting of HO-, and F-;
- 5
R⁵ is selected from the group R^{5a} consisting of
H-, halogen, NC-, HO- and C₁₋₃-alkyl-,
wherein the above mentioned C₁₋₃-alkyl-group may optionally be
substituted with 1 to 5 substituents independently selected from the
group consisting of HO- and F-
or R⁵ and R⁶ together form an group O=;
- R⁶ is selected from the group R^{6a} consisting of
H-, halogen, NC-, HO- and C₁₋₃-alkyl-,


wherein the above mentioned C₁₋₃-alkyl-group may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of HO- and F- or R⁵ and R⁶ together form a group O=;

or a salt thereof.

Accordingly, for example E-5 covers compounds of formula (I),

5 wherein

A is selected from the group A^a consisting of



wherein above mentioned groups are substituted with one R⁵ and one R⁶;

R² is selected from the group R^{2b} consisting of

quinolinyl, phenyl and pyridynyl,

wherein the above mentioned quinoline, phenyl and pyridyl-groups may optionally be substituted with 1 to 5 substituents R⁴;

R³ is selected from the group R^{3c} consisting of

H- and H₃C-;

R⁴ is independently from each other selected from the group R^{4e} consisting of

F, Cl, F₃C-, F₃C-O- and H₃C-O-;

10

R⁵ is selected from the group R^{5b} consisting of

H-, HO- and C₁₋₂-alkyl-,

wherein the above mentioned C₁₋₂-alkyl-group may optionally be substituted with 1 to 5 F-,


or R⁵ and R⁶ together form an group O=:

R⁶ is selected from the group R^{6b} consisting of
H- and C₁₋₂-alkyl-,
wherein the above mentioned C₁₋₂-alkyl-group may optionally be
substituted with 1 to 5 F-,
or R⁵/R⁶ together form a group O=;

or a salt thereof.


5 Accordingly, for example E-39 covers compounds of formula (I),
wherein

A is selected from the group A^e consisting of



R¹ is selected from the group R^{1d} consisting of
H₃C- and cyclopropyl-;

R² is selected from the group R^{2f} being



10

R³ is selected from the group R^{3d} being H-;

R⁵ is selected from the group R^{5d} being
HO-;

R⁶ is selected from the group R^{6d} consisting of













H- and methyl-;













or a salt thereof.













Further preferred are the following compounds listed in Table 2:











5











No.	Structure	No.	Structure
I		II	
III		IV	
V		VI	
VII		VIII	
IX		X	












XI		XII	
XIII		XIV	
XV		XVI	
XVII		XVIII	
XIX		XX	
XXI		XXII	

XXIII		XXIV	
XXV		XXVI	
XXVII		XXVIII	
XXIX		XXX	
XXXI		XXXII	
XXXIII		XXXIV	

XXXV		XXXVI	
XXXVII		XXXVIII	
XXXIX		XL	
XLI		XLII	
XLIII		XLIV	
XLV		XLVI	

XLVII		XLVIII	
XLIX		L	
LI		LII	
LIII		LIV	
LV		LVI	

LVII		LVIII	
LIX		LX	
LXI		LXII	
LXIII		LXIV	
LXV		LXVI	


LXVII		LXVIII	
LXIX		LXX	
LXXI		LXXII	
LXXIII		LXXIV	
LXXV		LXXVI	
LXXVII			

or the salts thereof.

Some terms used above and hereinafter to describe the compounds according to the
5 invention will now be defined more closely.

Terms not specifically defined herein should be given the meanings that would be
given to them by one of skill in the art in light of the disclosure and the context. As
used in the specification, however, unless specified to the contrary, the following
10 terms have the meaning indicated and the following conventions are adhered to.
In the groups, radicals, or moieties defined below, the number of carbon atoms is
often specified preceding the group, for example C₁₋₆-alkyl means an alkyl group or
radical having 1 to 6 carbon atoms. In general, for groups comprising two or more
subgroups, the last named subgroup is the radical attachment point, for example, the
15 substituent "aryl-C₁₋₃-alkyl-" means an aryl group which is bound to a C₁₋₃-alkyl-
group, the latter of which is bound to the core molecule or to the group to which the
substituent is attached.

Within the present invention, the term "core molecule" is defined by the following
structure:



In general, the attachment site of a given residue to another group shall be variable,
i.e. any capable atom, bearing hydrogens to be replaced, within this residue may be
the linking spot to the group being attached, unless otherwise indicated.

In case a compound of the present invention is depicted in form of a chemical name
and as a formula in case of any discrepancy the formula shall prevail.

An asterisk may be used in sub-formulas to indicate the bond or attachment point which is connected to the core molecule, rest of the molecule or to the substituent to which it is bound as defined.

- 5 Unless specifically indicated, throughout the specification and the appended claims, a given chemical formula or name shall encompass tautomers and all stereo, optical and geometrical isomers (e.g. enantiomers, diastereomers, E/Z isomers etc...) and racemates thereof as well as mixtures in different proportions of the separate enantiomers, mixtures of diastereomers, or mixtures of any of the foregoing forms
- 10 where such isomers and enantiomers exist, as well as salts, including pharmaceutically acceptable salts thereof and solvates thereof such as for instance hydrates including solvates of the free compounds or solvates of a salt of the compound.
- 15 The phrase "pharmaceutically acceptable" or "physiologically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, and
- 20 commensurate with a reasonable benefit/risk ratio.

- As used herein, "pharmaceutically acceptable salts" or "physiologically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically
- 25 acceptable salts or physiologically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. For example, such salts include salts from ammonia, L-arginine, betaine, benethamine, benzathine, calcium hydroxide, choline, deanol, diethanolamine (2,2'-iminobis(ethanol)), diethylamine, 2-
- 30 (diethylamino)-ethanol, 2-aminoethanol, ethylenediamine, N-ethyl-glucamine, hydrabamine, 1H-imidazole, lysine, magnesium hydroxide, 4-(2-hydroxyethyl)-morpholine, piperazine, potassium hydroxide, 1-(2-hydroxyethyl)-pyrrolidine, sodium hydroxide, triethanolamine (2,2',2"-nitrilotris(ethanol)), tromethamine, zinc hydroxide,

acetic acid, 2,2-dichloro-acetic acid, adipic acid, alginic acid, ascorbic acid, L-aspartic acid, benzenesulfonic acid, benzoic acid, 2,5-dihydroxybenzoic acid, 4-acetamido-benzoic acid, (+)-camphoric acid, (+)-camphor-10-sulfonic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, decanoic acid, dodecylsulfuric acid, ethane-
5 1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxy-ethanesulfonic acid, ethylenediaminetetraacetic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, D-glucoheptonic acid, D-gluconic acid, D-glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycine, glycolic acid, hexanoic acid, hippuric acid, hydrobromic acid, hydrochloric acid, isobutyric acid, DL-lactic
10 acid, lactobionic acid, lauric acid, lysine, maleic acid, (-)-L-malic acid, malonic acid, DL-mandelic acid, methanesulfonic acid, galactaric acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, nitric acid, octanoic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid (embonic acid), phosphoric acid, propionic acid, (-)-L-pyroglutamic acid, salicylic acid,
15 4-amino-salicylic acid, sebacic acid, stearic acid, succinic acid, sulfuric acid, tannic acid, (+)-L-tartaric acid, thiocyanic acid, p-toluenesulfonic acid and undecylenic acid. Further pharmaceutically acceptable salts can be formed with cations from metals like aluminium, calcium, lithium, magnesium, potassium, sodium, zinc and the like (also see Pharmaceutical salts, Berge, S.M. et al., J. Pharm. Sci., (1977), 66, 1-19).

20 The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a sufficient amount of the appropriate base or
25 acid in water or in an organic diluent like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile, or a mixture thereof.

Salts of other acids than those mentioned above which for example are useful for purifying or isolating the compounds of the present invention (e.g. trifluoro acetate
30 salts) also comprise a part of the invention.

The term "substituted" as used herein means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that

the designated atom's viable valence number is not exceeded, and that the substitution results in a stable compound.

5 The term "partially unsaturated" as used herein means that in the designated group or moiety 1, 2, or more, preferably 1 or 2, double bonds are present. Preferably, as used herein, the term "partially unsaturated" does not cover fully unsaturated groups or moieties.

10 The term "halogen" generally denotes fluorine (F), chlorine (Cl), bromine (Br) and iodine (I).

The term "C_{1-n}-alkyl", wherein n is an integer from 2 to n, either alone or in combination with another radical denotes an acyclic, saturated, branched or linear hydrocarbon radical with 1 to n C atoms. For example the term C₁₋₅-alkyl embraces
15 the radicals H₃C-, H₃C-CH₂-, H₃C-CH₂-CH₂-, H₃C-CH(CH₃)-, H₃C-CH₂-CH₂-CH₂-, H₃C-CH₂-CH(CH₃)-, H₃C-CH(CH₃)-CH₂-, H₃C-C(CH₃)₂-, H₃C-CH₂-CH₂-CH₂-CH₂-, H₃C-CH₂-CH₂-CH(CH₃)-, H₃C-CH₂-CH(CH₃)-CH₂-, H₃C-CH(CH₃)-CH₂-CH₂-, H₃C-CH₂-C(CH₃)₂-, H₃C-C(CH₃)₂-CH₂-, H₃C-CH(CH₃)-CH(CH₃)- and H₃C-CH₂-CH(CH₂CH₃)-.

20


The term "C_{3-n}-cycloalkyl", wherein n is an integer from 4 to n, either alone or in combination with another radical denotes a cyclic, saturated, unbranched hydrocarbon radical with 3 to n C atoms. For example the term C₃₋₇-cycloalkyl includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.


25

The term "aryl" as used herein, either alone or in combination with another radical, denotes a carbocyclic aromatic monocyclic group containing 6 carbon atoms which may be further fused to a second 5- or 6-membered carbocyclic group which may be aromatic, saturated or unsaturated. Aryl includes, but is not limited to, phenyl,
30 indanyl, indenyl, naphthyl, anthracenyl, phenanthrenyl, tetrahydronaphthyl and dihydronaphthyl.

The term "heteroaryl" means a mono- or polycyclic-ring systems containing one or more heteroatoms selected from N, O or S(O)_r, wherein r=0, 1 or 2, consisting of 5 to 14 ring atoms wherein at least one of the heteroatoms is part of an aromatic ring. The term "heteroaryl" is intended to include all the possible isomeric forms.

- 5 In one embodiment the term "heteroaryl" means a mono- or bicyclic-ring system containing one to three heteroatoms selected from N, O or S(O)_r, wherein r=0, 1 or 2, consisting of 5 to 10 ring atoms wherein at least one of the heteroatoms is part of an aromatic ring.
- 10 Thus, the term "heteroaryl" includes the following exemplary structures which are not depicted as radicals as each form may be attached through a covalent bond to any atom so long as appropriate valences are maintained:





Many of the terms given above may be used repeatedly in the definition of a formula
 5 or group and in each case have one of the meanings given above, independently of
 one another.

The compounds according to the invention may be obtained using methods of
 synthesis known in principle. Preferably, the compounds are obtained by the
 10 following methods according to the invention which are described in more detail
 hereinafter.


Preparation

The following Schemes shall illustrate generally how to manufacture the compounds of the present invention by way of example. The abbreviated substituents may be as
5 defined above if not defined otherwise within the context of the schemes .


The preparation process might comprises :

a) Reacting a compound of formula (II)

10



or a derivatives thereof, with a compound of formula (III)



15

Wherein R^1 , R^2 , R^3 , R^5 , R^6 and A are as defined above and L is a suitable leaving group such as halogen atom (e.g. chlorine or bromine) or hydroxyl group.


In case of L= halogen, process a) typically comprises the reaction of a compound of
20 formula (II) with a compound of formula (III) in an appropriate solvent such as acetonitrile or N,N-dimethylformamide in the presence of a base such as TEA or DIPEA at room temperature.

In case of L= OH, process a) typically comprises the reaction of a compound of
25 formula (II) with a compound of formula (III) in an appropriate solvent such as N,N-

dimethylformamide and in the presence of a suitable coupling agent (e.g. HATU or TBTU)

Compounds of formula (III) are either commercially available or can be prepared as
5 described in the following Schemes, following known reported procedures.

Scheme 1:




10

In Scheme 1, Step 1 typically involves reaction of commercially available amino pyrazole derivatives with 2-Bromo-malonaldehyde in the presence of acetic acid in a suitable solvent such as EtOH under heating. In Step 2, the cyclopropyl group is
15 introduced by a cross coupling palladium catalyzed reaction using for example potassium cyclopropyltrifluoroborate, a suitable palladium catalyst such as Palladium(II) acetate and 2-dicyclohexylphosphino- 2',6'-diisopropoxy 1,1'-biphenyl as ligand in an appropriate solvent such as toluene under heating . In Step 3 the ethyl ester is then hydrolyzed under basic conditions using sodium hydroxide or

lithium hydroxide monohydrate in an appropriate solvent such as EtOH or a mixture of THF/water.


Scheme 2:

5



In Scheme 2, Step 1 typically involves reaction of commercially available amino
pyrazole derivative with 1,1,3,3,- tetraethoxy-propane in the presence of hydrochloric
acid in a suitable solvent such as EtOH under heating. Bromination using bromine in
acetic acid as solvent at room temperature provides the bromo derivative and the
cyclopropyl group is then introduced as described in Scheme 1.


Scheme 3:



In Scheme 3, Step 1 typically involves reaction of commercially available amino pyrazole derivative with 1,1,3,3,- tetraethoxy-2-methyl-propane in the presence of hydrochloric acid in a suitable solvent such as EtOH under heating . Basic hydrolysis provides the desired carboxylic acid derivative

Compounds of formula (II) are either commercially available or can be prepared as described in the following Schemes.

Scheme 4



In Scheme 4, R^2 is aryl or heteroaryl


In Scheme 4, Step 1, commercially available ketone is converted in the corresponding 2-methyl-propane-2-sulfinyl-imine using titanium (IV) ethoxide and 2-Methyl-propane-2-sulfinic acid amide, as described in WO 2005087751.

The obtained intermediate is then added dropwise to a previously prepared solution of organo lithium derivatives of the appropriate halogen compounds (R^2X , where X is

bromine or iodine) prepared using for example commercially available solution of tert-butyllithium or n-butyllithium in hexane or pentane at low temperature (-75°C) in a suitable solvent such as toluene or THF. Cleavage of the resulting sulphinic amide by treatment with acid such as a 4N solution of HCl in a suitable solvent such as dioxane provides the desired intermediates amines.

The above described synthesis applies also for the analogues with 5 and 4 membered ring, starting from commercially available cyclopentanone and oxetane-3-one.


10 Scheme 5



In Scheme 5, R^2 is aryl derivatives.

- Step 1 involves a cross coupling Suzuki reaction with commercially available boronic acid or pinacol ester derivatives and the appropriate halogen derivatives ($\text{X} = \text{Br}$ or I) using for example 1,1'-bis(diphenylphosphino)ferrocenedichloro palladium(II) as catalyst, potassium carbonate as base in an appropriate solvent such as a mixture of toluene/water under heating. The epoxidation step is performed using MCPBA as oxidating agent in DCM at room temperature. The desired amino alcohol intermediate is then obtained by opening the epoxide with a modified Ritter procedure using trifluoromethane sulfonic acid and acetonitrile followed by basic hydrolysis of the formed intermediate, in analogy to the procedure described in Tetrahedron Asymmetry, 1996, 5, 1501-1506.
- The relative stereochemistry of the above described aminoalcohols is reported in the Experimental description.

Scheme 6




5 In Scheme 6, R^2 is aryl or heteroaryl .

In Scheme 6, the opening of the epoxide is performed using sodium azide in the presence of ammonium chloride under heating in a suitable solvent, such as dimethyl formamide. After separation of the two regioisomers, (see experimental) , the azide group is then converted into amino group by reduction following well known reported
 10 procedure such as for example using zinc and ammonium formate in a suitable solvent such as methanol at room temperature.

The relative stereochemistry of the above described amino alcohols is reported in the Experimental description.

Scheme 7




5 In Scheme 7, R^2 is aryl or heteroaryl

In Scheme 7, the regioisomeric mixture of azide intermediates, obtained following the approach described in Scheme 6, is reduced under catalytic hydrogenation conditions, using for example Pd/C in a suitable solvent such as ethanol and in the presence of di.tert-butylidicarbonate to obtain the protected amino alcohols

10 derivatives. Oxidation to ketone is performed using Dess Martin periodinane in a suitable solvent such as DCM at room temperature or using Swern's procedure. Formation of the tertiary alcohols is accomplished by addition of methyl magnesium chloride to the carbonyl group at low temperature (-20°C) in a suitable solvent such as THF . The cleavage of the Boc protecting group is performed under acid conditions

15 using for example trifluoroacetic acid in a suitable solvent such as DCM at room temperature.

The regioisomeric ratio of epoxide opening and the relative stereochemistry of the above described amino alcohols are reported in the Experimental description.


Scheme 8

5 In Scheme 8 R^2 is aryl.

In Scheme 8, the desired tetrahydro furan-3-ol- intermediates are obtained by addition of the appropriate lithium derivatives, prepared reacting a suitable halogen compounds (R^2X , X =halogen) with commercially available solution of tert-butyllithium or n-buthyllithium in hexane or pentane at low temperature (-75°C) in a suitable solvent such as toluene or THF, to the carbonyl group. Treatment with pTsOH in toluene under reflux provides the double bond derivatives which are transformed into the desired aminoalcohols following the approaches described in Scheme 5.

10 The relative stereochemistry of the aminoalcohols compounds are reported in the Experimental description.

15

Scheme 9

In Scheme 9, R^2 is aryl or heteroaryl.

20 The desired amino alcohols are obtained following the approaches described above in Scheme 6 and 7.

The regiochemistry ratio of the epoxide opening and the relative stereochemistry of the aminoalcohols compounds are reported in the Experimental description.

Biological Examples

5

In-vitro effect:

The *in-vitro* effect of the active compounds of the invention can be shown with the following biological assays.

10

a) Phosphodiesterase (PDE) 2A and 10 assay with fluorescent substrate

Assay principle:

The PDE reaction cleaves cAMP to AMP. The IMAP system (Molecular Device) using
15 fluorescence polarization (FP) as detection principle was used to measure enzyme activity. A fluorescent labeled cAMP was used as substrate for the reaction, generating a labeled AMP. The fluorescent AMP binds specifically to the large M(III)-based nano-particles which reduces the rotational speed of the substrate and thus increases its polarization.

20

Detailed method:

The inhibition of PDE 2A or 10 enzyme activity was assessed using IMAP-
Phosphodiesterase-cAMP fluorescence labeled substrate (Molecular Devices, Order
No. R7506), IMAP TR-FRET screening express (Molecular Devices, Order No.
25 R8160, the TR-FRET component will not be used) and PDE 2A or PDE10 protein expressed upon baculovirus infection in SF9 cells. The cells were incubated after infection for ~3 days and protein production was confirmed by Western Blot. The cells were collected by centrifugation and the pellet frozen in liquid nitrogen before it was resuspended in PBS containing 1% Triton X-100 and protease inhibitors. After
30 45 min incubation on ice, the cell debris was removed by centrifugation (13.000 rpm, 30 min). Since SF 9 cells do not express cAMP hydrolyzing enzymes to a high extent, no further purification of the protein was needed.

All reactions were performed in 384 well plates, Perkin Elmer black optiplates and IMAP reaction buffer with 0.1% Tween20 (kit component)

Compounds were serial diluted in DMSO. With an intermediate dilution step with
5 reaction buffer DMSO concentration was reduced to achieve 1% DMSO in the assay
reaction. Setup of the assay started with 10µl enzyme (~10ng/well, depending on
prep. batch), 5 µl compound, reaction was started by addition of 5 µl labeled cAMP
(30 nM, final concentration), immediately mixed for 15 seconds on a Eppendorf
mixmate (2000 rpm) followed by an incubation at room temperature for 90 minutes in
10 the dark. Reaction is stopped by adding of 60 µl binding buffer for FP/cAMP (kit
component). After at least 90 min of further incubation (room temperature, dark) the
assay was measured at 485 nm excitation/525 nm emission in an Envision multilabel
reader (PerkinElmer).

15 Each assay plate contained wells with vehicle controls (1% DMSO) for the
measurement of non-inhibited reaction (=100% control) and wells without enzyme as
0% controls.

The analysis of the data was performed by calculation of the percentage of inhibition
20 in the presence of test compound compared to the vehicle control samples (100%
control, no inhibition) and a low control (0% control, no enzyme).

IC50 values are calculated with Assay Explorer or other suited software based on
curve fitting of results of at least 8 different compound concentrations. The compound
concentrations may vary according to the needed range, but typically cover the range
25 between 10µM and 0.1pM.

Table 3a: PDE2A Activity of the examples (Ex) compiled in the experimental part,
based on above described assay (IMAP fluorescent).

Ex.	PDE2 IC₅₀ [nM]	Ex.	PDE2 IC₅₀ [nM]	Ex.	PDE2 IC₅₀ [nM]	Ex.	PDE2 IC₅₀ [nM]
1	14	27	534	53	100	81b	149
2	22	28	834	54	12	82a	1.2
3	266	29	661	55	21	82b	40
4	48	30	12	56	229	83a	5.9
5	258	31	30	57	103	83b	1820
6	199	32	67	58	14	84a	120
7	77	33	83	59	60	84b	2790
8	69	34	27	60	22	85a	130
9	117	35	31	61	19	86a	3.9
10	80	36	180	62	127	86b	2550
11	297	37	192	63	124	87a	1000
12	1650	38	331	64	496	87b	184
13	359	39	568	65	30	88a	32
14	456	40	84	66	25	88b	3030
15	746	41	291	67	50	89a	1000
16	537	42	130	68	24	90a	200
17	39	43	359	69	1740	90b	14
18	129	44	840	70	1250	91a	63
19	519	45	239	71	13	91b	1590
20	172	46	5.7	72	122	92a	93
21	74	47	240	73	1142	92b	712
22	119	48	1.35	74	2530	93a	11
23	232	49	70	75	42	93b	1520
24	754	50	596	80b	75		
25	88	51	59	80a	313		
26	174	52	231	81a	3.4		

Table 3b: PDE10 Activity of the examples (Ex) compiled in the experimental part, based on above described assay (IMAP fluorescent).

Ex.	PDE10 IC₅₀ [nM]	Ex.	PDE10 IC₅₀ [nM]	Ex.	PDE10 IC₅₀ [nM]	Ex.	PDE10 IC₅₀ [nM]
1	10100	27	>10000	53	>10000	81b	>10000
2	>10000	28	>10000	54	9670	82a	>10000
3	>10000	29	5930	55	>10000	82b	8461
4	550	30	>10000	56	>10000	83a	>10000
5	12200	31	>10000	57	>10000	83b	>10000
6	>10000	32	>10000	58	>10000	84a	>10000
7	9110	33	>10000	59	>10000	84b	>10000
8	>10000	34	10800	60	>10000	85a	>10000
9	>10000	35	>10000	61	6650	86a	>10000
10	9820	36	>10000	62	7160	86b	9940
11	1470	37	>10000	63	>10000	87a	>10000
12	>10000	38	6710	64	>10000	87b	>10000
13	9910	39	>10000	65	9760	88a	7560
14	8430	40	5730	66	>10000	88b	>10000
15	>10000	41	7950	67	>10000	89a	>10000
16	>10000	42	5590	68	>10000	90a	8590
17	6940	43	6860	69	>10000	90b	7350
18	8630	44	9680	70	>10000	91a	7700
19	>10000	45	7850	71	>10000	91b	5670
20	>10000	46	>10000	72	>10000	92a	>10000
21	>10000	47	>10000	73	>10000	92b	>10000
22	9920	48	6620	74	>10000	93a	>10000
23	>10000	49	>10000	75	>10000	93b	>10000
24	>10000	50	>10000	80b	5280		
25	9070	51	9040	80a	7760		
26	>10000	52	>10000	81a	>10000		

In-vivo effect:**Animal Experiments and sample analysis (CSF):**

Test compounds were administered to animals (rat) different routes at doses of 10.0 or 5 $\mu\text{mol/kg}$, (both oral and intravenous). CSF samples were carefully collected by puncture of the cisterna magna under anaesthesia. Immediately after CSF sampling, blood was taken by heart puncture and brains were dissected out. Blood was collected in EDTA-coated microvettes and plasma was prepared by centrifugation. Concentration of the test compounds in plasma, CSF or brain homogenate was determined using HPLC-MS-MS.

Table 4 : Plasma, brain and CSF concentration

Ex.	Time(*) (h)	conc plasma (nmol/L)	conc brain (nmol/L)	c(brain)/ c(plasma)	conc CSF (nmol/L)	c(CSF)/ c(plasma)
1	0.5	243	471	1.96	11	0.04
21	0.5	1210	1320	1.17	106	0.09
25	0.5	1040	957	0.92	111	0.12
81a	0.5	2460	1070	0.42	261	0.10
82a	0.5	3320	1180	0.36	157	0.05
83a	0.5	794	449	0.6	61	0.08

(*) Time between administration and CSF sampling

For the skilled in the art it is evident from the experimental results shown above that compounds of the present invention are not only potent phosphodiesterase 2 inhibitors but also reach high CSF concentrations and adequate CSF to plasma ratios.

Plasma Protein Binding (Determination of human and rat plasma protein binding with Equilibrium Dialysis)

This equilibrium dialysis (ED) technique is used to determine the approximate in vitro fractional binding of test compounds to human and rat plasma proteins.

Dianorm Teflon dialysis cells (micro 0.2) are used. Each cell consists of a donor and an acceptor chamber, separated by an ultrathin semipermeable membrane with a 5 kDa molecular weight cutoff.

Stock solutions for each test compound are prepared in DMSO at 1mM and diluted to a final concentration of 1.0 μ M. The subsequent dialysis solutions are prepared in pooled human and rat plasma (with NaEDTA)

Aliquots of 200 μ L dialysis buffer (100 mM potassium phosphate, pH 7.4) are dispensed into the buffer chamber. Aliquots of 200 μ L test compound dialysis solution are dispensed into the plasma chambers. Incubation is carried out for 2 hours under rotation at 37°C.

At the end of the dialysis period, the dialysate is transferred into reaction tubes. The tubes for the buffer fraction contain 0.2ml Acetonitril/water (80/20). Aliquots of 25 μ L of the plasma dialysate are transferred into deep well plates and mixed with 25 μ L Acetonitril/water (80/20), 25 μ L buffer, 25 μ L calibration solution and 25 μ L Internal Standard solution. Protein precipitation is done by adding 200 μ L Acetonitrile.

Aliquots of 50 μ L of the buffer dialysate are transferred into deep well plates and mixed with 25 μ L blank plasma, 25 μ L Internal Standard solution and 200 μ L Acetonitril. Samples are measured on HPLC-MS/MS-Systems and evaluated with Analyst-Software.

Percent bound is calculated with the formula: %bound = (plasma concentration - buffer concentration/ plasma concentration) X 100 and % free is calculated as difference.

Table 4 : PPB (Plasma Protein Binding) of compounds of the present invention in human and rat plasma.

EX	PPB HUM %BINDING	PPB RAT %BINDING	EX	PPB HUM %BINDING	PPB RAT %BINDING
2	91,5	94,7	70	79,4	-
1	96	96,8	71	81,1	-
31	95,3	97,6	51	46,9	-

35	92,4	93,7	53	81,7	-
37	94,4	93,50	88b	44,2	-
17	91,4	90,7	82a	78,7	86,5
25	75,4	83,1	81a	63,0	74,3
21	83,8	87,2	46	67,1	68,7
3	84,3	-	81b	65,3	66,2
68	84,4	-	83a	84,4	82,2
69	90,7	-	86a	89,5	93,5

For the skilled in the art it is evident from the experimental results shown above that compounds of the present invention are not only potent phosphodiesterase 2 inhibitors but also have low plasma protein binding.

5

Assessment of efflux in Madin-Darby canine kidney cells transfected with the human MDR1 gene (MDCK assay)

Apparent permeability coefficients (PE) of the compounds across the MDCK-MDR1 cell monolayers are measured (pH 7.4, 37 °C) in apical-to-basal (AB) and basal-to-apical (BA) transport direction. AB permeability (PEAB) represents drug absorption from the blood into the brain and BA permeability (PEBA) drug efflux from the brain back into the blood via both passive permeability as well as active transport mechanisms mediated by efflux and uptake transporters that are expressed on the MDCK-MDR1 cells, predominantly by the overexpressed human MDR1 P-gp. The compounds are assigned to permeability/absorption classes by comparison of the AB permeabilities with the AB permeabilities of reference compounds with known in vitro permeability and oral absorption in the human. Identical or similar permeabilities in both transport directions indicate passive permeation, vectorial permeability points to additional active transport mechanisms. Higher PEBA than PEAB indicates the involvement of active efflux mediated by MDR1 P-gp. Active transport is concentration- dependently saturable.

MDCK-MDR1 cells (1-2 x 10⁵ cells/1 cm² area) are seeded on filter inserts (Costar transwell polycarbonate or PET filters, 0.4 µm pore size) and cultured (DMEM) for 7

days. Subsequently, the MDR1 expression is boosted by culturing the cells with 5 mM sodium butyrate in full medium for 2 days. Compounds are dissolved in appropriate solvent (like DMSO, 1 -20 mM stock solutions). Stock solutions are diluted with HTP-4 buffer (128.13 mM NaCl, 5.36 mM KCl, 1 mM MgSO₄, 1.8 mM CaCl₂, 4.17 mM NaHCO₃, 1.19 mM Na₂HPO₄ x 7H₂O, 0.41 mM NaH₂PO₄xH₂O, 15 mM HEPES, 20 mM glucose, 0.25 % BSA, pH 7.4) to prepare the transport solutions (0.1 - 300 μ M compound, final DMSO $\leq$ 0.5 %). The transport solution (TL) is applied to the apical or basolateral donor side for measuring A-B or B-A permeability (3 filter replicates), respectively. The receiver side contains the same buffer as the donor side. Samples are collected at the start and end of experiment from the donor and at various time intervals for up to 2 hours also from the receiver side for concentration measurement by HPLC-MS/MS or scintillation counting. Sampled receiver volumes are replaced with fresh receiver solution. Efflux ratio is calculated dividing the Papp (b-a) values by the Papp (a-b) values.

15

Table 5 : Papp (PEBA) and efflux of compounds of the present invention

Ex.	Papp (a-b) mean [10-6 cm/s]	efflux ratio	Ex.	Papp (a-b) mean [10-6 cm/s]	efflux ratio
1	78	0.6	53	15	1.8
2	84	0.6	82a	59	1.0
37	86	0.5	51	11	5.0
17	85	0.6	81a	60	1.4
25	100	0.8	46	60	1.4
21	94	0.6	81b	64	1.2
3	97	0.7	83a	34	1.4
69	25	1.3	84a	31	2.2
70	20	2.1	84b	23	2.2
71	23	1.9			

For the skilled in the art it is evident from the experimental results shown above that compounds of the present invention are not only potent phosphodiesterase 2 inhibitors but also have good membrane permeability and low to moderate in vitro efflux.

5

Metabolic stability

The metabolic stability of the compounds according to the invention has been investigated as follows:

10

The metabolic degradation of the test compound was assayed at 37 °C with pooled liver microsomes from various species. The final incubation volume of 100 µl per time point contains TRIS buffer pH 7.6 at room temperature (0.1 M), magnesium chloride (5 mM), microsomal protein (1 mg/mL for human and dog, 0.5 mg/mL for other species) and the test compound at a final concentration of 1 µM. Following a short preincubation period at 37°C, the reactions were initiated by addition of betanicotinamide adenine dinucleotide phosphate, reduced form (NADPH, 1 mM), and terminated by transferring an aliquot into solvent after different time points. After centrifugation (10000 g, 5 min), an aliquot of the supernatant was assayed by LC10
MS/MS for the amount of parent compound. The half-life was determined by the slope of the semi-logarithmic plot of the concentration-time profile.

20

Table 4: Stability of compounds of the present invention in human liver microsomes.

Ex.	Half-life – t1/2 [min] human	Ex.	Half-life – t1/2 [min] human
1	>130	71	>130
2	120	51	>130
37	>130	53	120
17	>130	88b	>130
25	>130	82a	>130
21	>130	81a	>130
3	72	46	>130
68	>130	81b	>130

69	53	83a	>130
70	63	86a	>130

For the skilled in the art it is evident from the experimental results shown above that compounds of the present invention are not only potent phosphodiesterase 2 inhibitors but also have good metabolic stability.

5

In view of their ability to inhibit the activity of phosphodiesterase 2 activity and their advantageous pharmacokinetics properties the compounds of general formula (I) according to the invention, or the physiologically acceptable salts thereof, are suitable for the treatment and/or preventative treatment of all those diseases or conditions which can be influenced by inhibition of PDE2 hyperactivity and/or cAMP and/or cGMP hypofunction. Therefore, compounds according to the invention, including the physiologically acceptable salts thereof, are particularly suitable for the prevention or treatment of diseases, particularly (1) disorders comprising the symptom of cognitive deficiency; (2) organic, including symptomatic, mental disorders, dementia; (3) mental retardation; (4) mood affective disorders; (5) neurotic, stress-related and somatoform disorders including anxiety disorders; (6) behavioural and emotional disorders with onset usually occurring in childhood and adolescence, attention deficit hyperactivity syndrome (ADHD) including Autism spectrum disorders; (7) disorders of psychological development, developmental disorders of scholastic skills; (8) schizophrenia and other psychotic disorders; (9) disorders of adult personality and behaviour; (10) mental and behavioural disorders due to psychoactive substance use; (11) extrapyramidal and movement disorders; (12) episodic and paroxysmal disorders, epilepsy; (13) Systemic atrophies primarily affecting the central nervous system, ataxia; (14) Behavioural syndromes associated with physiological disturbances and physical factors; (15) sexual dysfunction comprising excessive sexual drive; (16) factitious disorders; (17) obsessive-compulsive disorders; (18) depression; (19) neuropsychiatric symptoms (e.g. depressive symptoms in Alzheimer's disease); (20) mixed dementia; (21) cognitive impairment in schizoaffective disorder; (22) cognitive impairment in bipolar disorder and (23) cognitive impairment in major depressive disorder.

30

In addition, the compounds of the present invention can be used for the treatment, amelioration and / or prevention of cognitive impairment being related to perception, concentration, cognition, learning, attention or memory.

- In addition, the compounds of the present invention can be used for the treatment
5 amelioration and / or prevention of cognitive impairment being related to age-associated learning and memory impairments, age-associated memory losses, vascular dementia, craniocerebral trauma, stroke, dementia occurring after strokes (post stroke dementia), post-traumatic dementia, general concentration impairments, concentration impairments in children with learning and memory problems,
10 Alzheimer's disease, Lewy body dementia, dementia with degeneration of the frontal lobes, including Pick's syndrome, Parkinson's disease, progressive nuclear palsy, dementia with corticobasal degeneration, amyotrophic lateral sclerosis (ALS), Huntington's disease, multiple sclerosis, thalamic degeneration, Creutzfeld-Jacob dementia, HIV dementia, schizophrenia with dementia or Korsakoff's psychosis.
15 In addition, the compounds of the present invention can be used for the treatment of Alzheimer's disease.

In addition compounds of the present invention can be used for the treatment of pain disorders, including but not limited to inflammatory, neuropathic and osteoarthritic pain.

- 20 In addition, the compounds of the present invention can be used for the treatment of sleep disorders, bipolar disorder, metabolic syndrome, obesity, diabetes mellitus, hyperglycemia, dyslipidemia, impaired glucose tolerance, or a disease of the testes, brain, small intestine, skeletal muscle, heart, lung, thymus or spleen.

- Preferably the compounds according to the invention are suitable for the treatment of
25 Alzheimer's Disease and for the treatment schizophrenia.

More preferably the compounds according to the invention are suitable for symptomatic treatment of Alzheimer's Disease and for the treatment of cognitive impairment associated with schizophrenia.

In particular the compounds according to the invention are suitable for symptomatic treatment of prodromal and mild-to-moderate Alzheimer's Disease and for the treatment of cognitive impairment associated with schizophrenia and symptomatic treatment of cognitive impairment associated with schizophrenia.

5

In a further aspect of the present invention the present invention relates to methods for the treatment or prevention of above mentioned diseases and conditions, which method comprises the administration of an effective amount of a compound of general formula (I), or the pharmaceutically acceptable salts thereof, to a human
10 being.

The dose range of the compounds of general formula (I) applicable per day is usually from 0.1 to 1000 mg, preferably from 1 to 500 mg by oral route, in each case administered 1 to 4 times a day.

15 Each dosage unit may conveniently contain from 0.1 to 500 mg, preferably 1 to 100 mg.

The actual pharmaceutically effective amount or therapeutic dosage will of course depend on factors known by those skilled in the art such as age and weight of the
20 patient, route of administration and severity of disease. In any case the combination will be administered at dosages and in a manner which allows a pharmaceutically effective amount to be delivered based upon patient's unique condition.

Suitable preparations for administering the compounds of formula I, including the
25 pharmaceutically acceptable salts thereof, will be apparent to those with ordinary skill in the art and include for example tablets, pills, capsules, suppositories, lozenges, troches, solutions, syrups, elixirs, sachets, injectables, inhalatives, powders, etc.. The content of the pharmaceutically active compound(s) should be in the range from 0.1 to 95 wt.-%, preferably 5.0 to 90 wt.-% of the composition as a whole.

30

Suitable tablets may be obtained, for example, by mixing one or more compounds according to formula I with known excipients, for example inert diluents, carriers,

disintegrants, adjuvants, surfactants, binders and/or lubricants. The tablets may also consist of several layers.

For this purpose, the compounds of formula I prepared according to the invention
5 may be formulated, optionally together with other active substances, together with one or more inert conventional carriers and/or diluents, e.g. with corn starch, lactose, glucose, microcrystalline cellulose, magnesium stearate, citric acid, tartaric acid, water, polyvinylpyrrolidone, water/ethanol, water/glycerol, water/sorbitol, water/polyethylene glycol, propylene glycol, cetylstearyl alcohol,
10 carboxymethylcellulose or fatty substances such as hard fat or suitable mixtures thereof.

The compounds according to the invention may also be used in conjunction with other active substances, particularly for the treatment and/or prevention of the
15 diseases and conditions mentioned above. Other active substances which are suitable for such combinations include, for example, BACE inhibitors; amyloid aggregation inhibitors (e.g. ELND-005); directly or indirectly acting neuroprotective and/or disease-modifying substances; anti-oxidants (e.g. vitamin E or ginkgolide); anti-inflammatory substances (e.g. Cox inhibitors, NSAIDs additionally or exclusively
20 having Abeta lowering properties); HMG-CoA reductase inhibitors (statins); acetylcholinesterase inhibitors (e.g., donepezil, rivastigmine, tacrine, galantamine); NMDA receptor antagonists (e.g. memantine); AMPA receptor agonists; AMPA receptor positive modulators, AMPAkinases, monoamine receptor reuptake inhibitors, substances modulating the concentration or release of neurotransmitters; substances
25 inducing the secretion of growth hormone (e.g., ibutamoren mesylate and capromorelin); CB-1 receptor antagonists or inverse agonists; antibiotics (e.g., minocyclin or rifampicin); PDE2, PDE4, PDE5, PDE9, PDE10 inhibitors, GABAA receptor inverse agonists, GABAA receptor antagonists, nicotinic receptor agonists or partial agonists or positive modulators, alpha4beta2 nicotinic receptor agonists or
30 partial agonists or positive modulators, alpha7 nicotinic receptor agonists or partial agonists or positive modulators; histamine H3 antagonists, 5 HT-4 agonists or partial agonists, 5HT-6 antagonists, alpha2-adrenoreceptor antagonists, calcium antagonists, muscarinic receptor M1 agonists or partial agonists or positive

modulators, muscarinic receptor M2 antagonists, muscarinic receptor M4 antagonists, metabotropic glutamate-receptor 5 positive modulators, glycine transporter 1 inhibitors, antidepressants, such as citalopram, fluoxetine, paroxetine, sertraline and trazodone; anxiolytics, such as lorazepam and oxazepam;

5 antipsychotics, such as aripiprazole, clozapine, haloperidol, olanzapine, quetiapine, risperidone and ziprasidone, and other substances that modulate receptors or enzymes in a manner such that the efficacy and/or safety of the compounds according to the invention is increased and/or unwanted side effects are reduced.

The compounds according to the invention may also be used in combination with
10 immunotherapies (e.g., active immunisation with Abeta or parts thereof or passive immunisation with humanised anti-Abeta antibodies or nanobodies) for the treatment of the above-mentioned diseases and conditions.

The dosage for the combination partners mentioned above is usefully 1/5 of the
15 lowest dose normally recommended up to 1/1 of the normally recommended dose.

Therefore, in another aspect, this invention relates to the use of a compound according to the invention or a pharmaceutically acceptable salt thereof combined with at least one of the active substances described above as a combination partner,
20 for preparing a pharmaceutical composition which is suitable for the treatment or prevention of diseases or conditions which can be affected by inhibitors of phosphodiesterase 2. These are preferably pathologies related to PDE2 hyperactivity and/or cAMP and/or cGMP hypofunction, particularly one of the diseases or conditions listed above, most particularly prodromal and mild-to-moderate
25 Alzheimer's Disease and cognitive impairment associated with schizophrenia.

The use of the compound according to the invention in combination with another active substance may take place simultaneously or at staggered times, but particularly within a short space of time. If they are administered simultaneously, the two active substances are given to the patient together; while if they are used at
30 staggered times the two active substances are given to the patient within a period of less than or equal to 12 hours, but particularly less than or equal to 6 hours.

Consequently, in another aspect, this invention relates to a pharmaceutical composition which comprises a compound according to the invention or a pharmaceutically acceptable salt thereof and at least one of the active substances described above as combination partners, optionally together with one or more inert carriers and/or diluents.

The compound according to the invention may both be present together in one formulation, for example a tablet or capsule, or separately in two identical or different formulations, for example as a so-called kit-of-parts.

Examples

The following examples are intended to illustrate the invention, without restricting its scope.

Chemical Manufacture

Abbreviations:

ACN	acetonitrile
APCI	Atmospheric pressure chemical ionization
d	day
Cy	cyclohexane
DCM	dichloromethane
DIPEA	diisopropylethylamine
DMF	dimethylformamide
ESI	electrospray ionization (in MS)
EtOAc	ethylacetate
EtOH	ethanol
Exp.	Example
GC	gas chromatography
GC-MS	coupled gas chromatography-mass spectrometry
h	hour(s)
HATU	O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium-Hexafluorophosphate

	HCl	hydrochloric acid
	HPLC	high performance liquid chromatography
	HPLC-MS	coupled high performance liquid chromatography-mass spectrometry
5	LC	liquid chromatography
	LC-MS	liquid chromatography – mass spectrometry
	M	molar (mol/L)
10	MeOH	methanol
	min	minute(s)
	MS	mass spectrometry
	NaOH	sodiumhydroxide
	NMP	1-methyl-2-pyrrolidinone
15	NOE	Nuclear Overhauser effect
	PE	petroleum ether
	rt	room temperature
	R _t	retention time (in HPLC)
	HATU	1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-
20		oxid hexafluorophosphate
	TBTU	O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate
	TEA	triethylamine
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
25	TLC	thin-layer chromatography
	UPLC- MS	ultra performance liquid chromatography - mass spectrometry

Analytical Methods:

30

UPLC-MS, HPLC-MS, ,LC-MS :

Method 1 :

Instrument: LC/MS ThermoFinnigan HPLC Surveyor DAD, MSQ single quadrupole

35 Column: Synergi Hydro RP100A, 2,5 µm, 3 x 50 mm

Mobile phase: A = H₂O 90% + 10% CH₃CN + NH₄COOH 10 mM
 B = CH₃CN 90% + H₂O 10% + NH₄COOH 10 mM

Time in min:	%A	%B	Flow rate in mL/min
0.00	100	0	1.2
0.50	100	0	1.2
6.50	0	100	1.2
7.50	0	100	1.2
8.00	100	0	1.2
9.00	100	0	1.2

Detection: UV 254 nm

- 5 Detection: Finnigan MSQ, single quadrupole
 Ion source: APCI+/APCI-
 Scan range: 100-900 amu

Method 2 :

- 10 Instrument: LC/MS Waters Acquity UPLC System DAD, SQD single quadrupole
 Column: BEH C18 1.7 μ m 2.1 x 50 mm, Temp 35 °C
 Mobile phase: A = H₂O 90% + CH₃CN 10% + NH₄COOH 5 mM
 B = CH₃CN 90% + H₂O 10%

Time in min:	%A	%B	Flow rate in mL/min
0.00	100	0	0.7
1.20	0	100	0.7
1.45	0	100	0.7
1.55	100	0	0.7
1.75	100	0	0.7

- 15 Detection: UV 254 nm
 Detection: SQD, single quadrupole
 Ion source: ES+/ ES -
 Scan range: 90-900 amu

Method 3 :

Instrument: LC/MS Waters Alliance 2695 HPLC System DAD, Quattro Micro Triple quadrupole

Column: Atlantis dC18 5 μ m 4,6 x 50 mm, Temp 35°C

- 5 Mobile phase: A = H₂O 90% + 10% CH₃CN + CF₃COOH 0,05%
B = CH₃CN 90% + 10% H₂O

Time in min:	%A	%B	Flow rate in mL/min
0.00	100	0	1.3
0.70	100	0	1.3
4.50	0	100	1.3
5.80	0	100	1.3
6.00	100	0	1.3

Detection: UV 254 nm

Detection: Quattro Micro, triple quadrupole

- 10 Ion source: ES+

Scan range: 90-1000 amu

Method 4:

Instrument: LC/MS Waters Alliance 2695 HPLC System DAD, Quattro Micro Triple quadrupole

- 15

Column: XBridge Phenyl 3.5 μ m 3x 30 mm, Temp 35°C

Mobile phase: A = H₂O 90% + 10% CH₃CN + NH₄HCO₃ 5mM

B = CH₃CN 90% + 10% H₂O

Time in min:	%A	%B	Flow rate in mL/min
0.00	100	0	1.3
4.50	0	100	1.3
5.80	0	100	1.3
6.00	100	0	1.3

- 20 Detection: UV 254 nm

Detection: Quattro Micro, triple quadrupole

Ion source: ES+

Scan range: 90-1000 amu

Method 5:

Instrument: LC/MS Waters Acquity UPLC System DAD, SQD single quadrupole

5 Column: BEH C18 1.7 μ m 2.1 x 50 mm, Temp 35 °C

Mobile phase: A = H₂O 90% + CH₃CN 10% + NH₄HCO₃ 5 mM

B = CH₃CN 90% + H₂O 10%

Time in min:	%A	%B	Flow rate in mL/min
0.00	100	0	0.70
1.20	100	0	0.70
1.45	0	100	0.70
1.55	0	100	0.70
1.75	100	0	0.70

Detection: UV 254 nm

10 Detection: SQD, single quadrupole

Ion source: ES+/ ES -

Scan range: 90-900 amu

Method 6:

15

Instrument: LC/MS Waters Acquity System DAD, SQD single quadrupole

Column: XBridge C18 2.5 μ m 3.0 x 30 mm, Temp 60 °C

Mobile phase: A = H₂O + TFA 0.1%

B = CH₃CN

Time in min:	%A	%B	Flow rate in mL/min
0.00	98	2	1.5
1.3	1	99	1.5
1.5	1	99	1.5
1.6	98	2	1.5

Method 7:

Instrument: LC/MS Waters Acquity System DAD, SQD single quadrupole

Column: XBridge C18 2.5 μ m 3.0 x 30 mm, Temp 60 °C

5 Mobile phase: A = H₂O + NH₄OH 0.1%

B = CH₃CN

Time in min:	%A	%B	Flow rate in mL/min
0.00	95	5	1.5
1.3	1	99	1.5
1.5	1	99	1.5
1.6	95	5	1.5

Method 8-:

10 Instrument: LC/MS Agilent 1100 System DAD

Column: Sunfire C18 2.5 μ m 3.0 x 30 mm, Temp 60 °C

Mobile phase: A = H₂O + TFA 0.1%

B = CH₃CN

Time in min:	%A	%B	Flow rate in mL/min
0.00	98	2.0	2.0
1.2	0.0	100	2.0
1.4	0.0	100	2.0

15

Method 10:

Instrument: LC/MS ThermoFinnigan HPLC Surveyor DAD, LCQFleet Ion Trap

Column: Xselect CSH, 2.5 μ m, 4,6 x 50 mm

Mobile phase: A = H₂O 90% + 10% CH₃CN + HCOOH 0.1%

20 B = CH₃CN 90% + H₂O 10% + HCOOH 0.1%

Time in min:	%A	%B	Flow rate in mL/min
0.00	100	0	1.4
4.00	0	100	1.4

5.30	0	100	1.4
5.50	100	0	1.4
6.00	100	0	1.4

Detection: UV 254 nm

Detection: Finnigan Fleet, Ion Trap

Ion source: ES+

5 Scan range: 100-900 amu

GC/MS Method

10 Method 9:

Instrument: GC/MS Thermo Scientific TRACE GC ULTRA, DSQ II MS single quadrupole

Column: Agilent DB-5MS, 25m x 0.25mm x 0.25 um

Carrier gas: Helium, 1 mL/min constant flow

15 Oven Program: 50 °C, to 100 °C in 10 °C/min, to 200 °C in 20 °C/min, to 320 °C in 30 °C/min (hold 10 min).

Detection: DSQ II MS single quadrupole

Ion source: EI

Scan range: 50- 450 amu

20

Chiral HPLC Methods :

Instrument: HPLC Agilent 1100 (DAD equipped; UV Detection: 230 nm); flow rate: 1 mL/min; column temperature: 25°C.

25 **Method C1**

column: Daicel Chiralpack AD-H; eluent: Hexane:Isopropanol=70:30

Method C2

column: Daicel Chiralpack AD-H; eluent: Hexane:Isopropanol=60:40

Method C3

30 column: Daicel Chiralpack AD-H; eluent: Hexane:Isopropanol=80:20

Method C4

column: Daicel Chiralcel OJ-H; eluent: Hexane:EtOH=80:20

Method C5

column: Daicel Chiralcel OJ-H; eluent: Hexane:EtOH=85:15

5 **Method C6**

column: Daicel Chiralcel OJ-H; eluent: Hexane:EtOH=70:30

Method C7

column: Daicel Chiralcel AS-H; eluent: Hexane:EtOH=75:25

10

NMR equipment :

The ¹H NMR spectra were recorded on a Bruker Avance III (500 MHz) or a Varian 400 (400 MHz) or Varian Mercury (300MHz) instrument using deuterated
15 dimethylsulfoxide (DMSO-d₆) as the solvent with tetramethylsilane (TMS) and residual solvent peak as an internal standard. Chemical shifts are reported in δ values (ppm) relative to TMS.

Purification :

20 The most suitable purification techniques applied for the purification of compounds of the present invention are direct phase silica gel flash chromatography and reverse phase chromatography, unless otherwise specifically stated.

General comment concerning the presentation of the structures

25

Compounds with stereogenic centre(s): The structures depicted in the experimental section will not necessarily show all the stereochemical possibilities of the compounds but only one.


The structural presentation of the compounds in the experimental section will show a
30 stereochemical bond only in case where the absolute stereochemistry is known.

The structural presentation of the compounds in the experimental section with unknown absolute stereochemistry will show a planar bond plus an additional

comment that indicates if the described compound is a racemic mixture, a single stereoisomer and where applicable the relative stereochemistry.


Two examples are given below.

- 5 Example 1 : the presented chemical structure is depicted as:




Racemic mixture


- 10 The added term racemic mixture points to the two stereochemical options and thus the manufactured compounds is a mixture of:



- 15 When racemic mixtures of above depicted structures are separated, the single stereoisomers are depicted as :



Single stereoisomer a



Single stereoisomer b


20

The added term 'single stereoisomer' and the planar bond indicates that the absolute configuration is unknown.


Single stereoisomer a is assigned to the first eluting isomer in chiral HPLC, single stereoisomer b is assigned to the second eluting isomer in chiral HPLC.

25

Example 2 : the presented chemical structure is depicted as:

**TRANS-racemic mixture**



The added term 'TRANS- racemic mixture' points to the two stereochemical options
5 and thus the manufactured compounds is a mixture of:



The same principles applies to 'CIS-racemic mixture' .

10

When racemic mixtures of above depicted structures are separated, the single stereoisomers are depicted as:

**TRANS- single stereoisomer a****TRANS- single stereoisomer b**

15 The added term 'TRANS-single stereoisomer' indicates a relative configuration known (trans) and the planar bond indicates the unknown absolute configuration. The same principles applies to 'CIS-single stereoisomer'.

Single stereoisomer a is assigned to the first eluting isomer in chiral HPLC, single stereoisomer b is assigned to the second eluting isomer in chiral HPLC .


20

Experimental

The following intermediates and examples are intended to illustrate the invention, without restricting its scope.

Intermediates

Intermediate 1:



5

To a solution of 3-amino-4-carbethoxypyrazole (4g, 25.27mmol) in absolute EtOH (40mL), 1,1,3,3-Tetraethoxy-2-methyl-propane (6.34g, 26.53mmol) was added followed by 13.90mL of a 1M solution of HCl in dioxane. The mixture was heated at 80°C overnight. Solvents were evaporated, then DCM and water were added. Phases were separated, organics washed with a saturated solution of NaCl, dried over sodium sulphate and evaporated to obtain 5.17g of the title compound


10

LC-MS (Method 2): $R_t = 0.73$ min

MS (ESI pos): $m/z = 206$ (M+H)⁺

15

Intermediate 2




20

Intermediate 1 (5g) was dissolved in a mixture of THF/water (1:1, 100mL) and stirred at room temperature for 48hrs. The resulting suspension was diluted with water and 70mL of EtOAc were added. Phases were separated, aqueous phases were treated with a 4N solution of HCl (ca 20mL). A white solid formed. The mixture was cooled at 0°C, then the white solid formed collected by filtration and dried under vacuum at 65°C to obtain 3.50g of the title compound.

25

LC-MS (Method 3): $R_t = 1.62$ min

MS (ESI pos): $m/z = 178$ (M+H)⁺

Intermediate 3:

5


To a solution of 2-Bromo-malonaldehyde (9.73 g; 64mmol) in EtOH (100mL) at 70 °C, 3-amino-4-carbethoxypyrazole (10g, 64mmol) and AcOH (100mL) were added and the mixture stirred at 70°C for 1 h. Solvents were evaporated, the residue treated with DCM (100mL) and a 1N solution of NaOH (100mL). Phases were separated, organics washed with a saturated solution of NaCl, dried over sodium sulphate and evaporated. The crude was purified flash chromatography (eluent 10:1 PE/EtOAc) to obtain 15g of the title compound as white solid.

10

LC-MS (Method 2): $R_t = 0.98$ min

MS (ESI pos): $m/z = 271$ ($M+H$)⁺

15

Intermediate 4:


20

Intermediate 3 (5g, 18.5mmol) was suspended in dry toluene (50mL) and 5mL of water were added. To this mixture, potassium cyclopropyltrifluoroborate (4g, 28mmol) was added followed by 2-dicyclohexylphosphino-2',6'-diisopropoxy 1,1'-biphenyl (0.864g, 1.85mmol), palladium acetate (0.208g, 0.93mmol) and potassium

carbonate (7.7g, 55mmol). Mixture was refluxed at 130° C for 3hrs, then cooled to room temperature, filtered over celite and washed with AcOEt and then EtOH. Solvent was evaporated under vacuum and the crude used in the next step without further purification.

- 5 LC-MS (Method 2): $R_t = 0.9$ min
MS (ESI pos): $m/z = 232$ (M+H)⁺

Intermediate 5:




10

Intermediate 4 (4g, 17.5mmol) was suspended in 50 ml of EtOH, 8 ml of 4N NaOH and 30 ml of water and stirred overnight. EtOH was evaporated and a 4 N solution of HCl added. The solid formed was filtered, washed with water and dried under vacuum at 70°C overnight to obtain 3.6g of the title compound.

15

LC-MS (Method 3): $R_t = 2.75$ min
MS (ESI pos): $m/z = 204$ (M+H)⁺

20 Intermediate 6:




Intermediate 6 was prepared as described in WO 2010/007074 starting from commercially available (Z) 3- (diethylamino)-2-fluoroprop-2-enal (1.34mL, 9.0 mmol)

25

and 3-amino-4-carbethoxypyrazole (2.1g, 13.6mmol) to obtain 0.53g of the title compound.

¹H NMR (300 MHz, CDCl₃): δ ppm 1.44-1.39 (t, 3H), 4.47-4.40 (q 2H) 8.57 (s, 1H)
5 8.7 (m, 1H), 8.8 (d, 1H)

Intermediate 7:



10


To a solution of 5-amino-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester (1g, 5.91mmol) in absolute EtOH (25mL), 1,1,3,3-Tetraethoxy-2-methyl-propane (1.4g, 6.2 mmol) was added followed by 1.63mL of a 4N solution of HCl in dioxane. Mixture was heated at 80°C for 5hrs, left at room temperature overnight and then solvents
15 were evaporated to dryness. The violet solid obtained was dissolved in DCM, water was added and the phases separated.

The organic phases were dried over sodium sulfate and concentrated under vacuum to obtain 1.26g of title compound used for next step without further purification.

LC-MS (Method 2): R_t = 0.79 min

20 MS (ESI pos): m/z = 224 (M+H)⁺

Intermediate 8:




25

To a solution of intermediate 7 (1.26g, 5.75mmol) in THF (25mL) and water (25mL) 1.5mL of a 1N solution of sodium hydroxide were added and the mixture heated at heated at 60°C for 2 h. Solvent was evaporated, water was added and 30 ml of a 12N solution of HCl added until pH 2. The solid formed was filtered, washed with
5 water and dried at 70°C under vacuum to obtain 0.9g of title compound as white solid.

LC-MS (Method 1): $R_t = 0.27$ min

MS (APCI): $m/z = 192$ (M+H)⁺

10 Intermediate 9:




To a solution of 5-amino-3-methyl-1H-pyrazole-4-carboxylic acid ethyl ester (4g, 23.64mmol) in absolute EtOH (80mL), 1,1,3,3-tetraethoxypropane (5.96mL, 23.64mmol) and 5.9mL of a 4N solution of HCl in dioxane were added. The resulting mixture was heated at 80°C 3hrs. Solvents were evaporated, the residue was diluted with DCM and water. Organic layer was separated, dried over sodium sulphate and evaporated to obtain the title compound as white solid (3.6g)

LC-MS (Method1): $R_t = 2.63$ min

MS (APCI): $m/z = 206$ (M+H)⁺

Intermediate 10:




To a solution of intermediate 9 (3.6g, 17.54mmol) in AcOH (70mL) bromine (2.26mL)
5 was added dropwise. The mixture was stirred at room temperature overnight then
carefully poured into 500 mL of water and extracted with EtOAc (3x100mL). Organic
phases were collected and washed with 100 mL of a 5% solution of Na₂S₂O₃ and
then with 100 mL of a saturated solution of NaCl, dried over sodium sulphate and
concentrated under vacuum. .

10 Crude was purified by flash chromatography (eluent: DCM/EtOAc; gradient from
100% to 70%) to obtain the title compound as white solid (2.1g)

LC-MS (Method 1): $R_t = 3.52$ min

MS (APCI): $m/z = 284$ (M+H)⁺

15 Intermediate 11:



To a solution of intermediate 10 (2.05g, 7.22mmol) in toluene (40mL) water (4mL)
20 was added followed by potassium cyclopropyltrifluoroborate (1.6g, 10.82mmol) ,
palladium(II) acetate (0.08g, 0.36mmol), dicyclohexylphosphino-2',6'-di-*i*-propoxy di-
1,1'-biphenyl (RUPHOS, 0.34g, 0.72mmol) and potassium carbonate (3g,
21.65mmol). Mixture was heated at 130°C for 3hrs then cooled to room
temperature, filtered over celite and washed with AcOEt. Organic layer was dried


and evaporated to obtain the title compound (1.5g) used for the next step without further purification.

LC-MS (Method 2): $R_t = 0.92$ min

MS (ESI pos): $m/z = 246$ (M+H)⁺

5

Intermediate 12:




- 10 To a solution of intermediate 11 (1.5g, 6.2mmol) in absolute EtOH (30mL) water (10mL) was added followed by 7.7mL of a 8N solution of NaOH. Mixture was stirred at room temperature overnight, then solvent evaporated and a 4N solution of HCl added until pH=1. The solid formed was filtered, washed with water and dried under vacuum at 70°C overnight (1.5g).

15 LC-MS (Method 1): $R_t = 0.6$ min

MS (APCI): $m/z = 218$ (M+H)⁺

Intermediate 13 :









- 20 To a solution of 4-bromo-3-fluorobenzotrifluoride (585 mg, 2.36 mmol) in 15 mL of THF, stirred at -75°C under nitrogen atmosphere, 1,53 mL (2.6 mmol) of a 1.7M solution of tert-butyllithium in pentane were added dropwise. The reaction mixture was stirred at -60°C for 15 minutes then a solution of 2-methyl-propylidene-2-sulfonic acid-(tetrahydro-pyran-4-ylidene)-amide (400 mg, 1.97 mmol; prepared as described in literature: WO2005/87751 A2) in 10 mL of THF was added dropwise. The reaction
- 25







mixture was allowed to reach room temperature and stirred for 1 hr. A saturated ammonium chloride solution was added and the reaction mixtures was extracted with ethyl acetate. The organic phases were collected, dried over sodium sulfate and concentrated under vacuum. The crude obtained was purified by flash


- 5 chromatography (eluent: cyclohexane/AcOEt; gradient from 12 % to 100% of AcOEt). The oil obtained was diluted in 2 mL of 1,4-dioxane, 0.4 mL of a 4 M solution of hydrochloric acid in 1,4-dioxane were added, the reaction mixture was stirred at room temperature for 1 hr and then concentrated under vacuum to obtain 100mg of the title compound as white solid.
- 10 LC-MS (Method 2): $R_t = 0.90$ min
MS (ESI pos): $m/z = 264$ (M+H)⁺

The following Amine Intermediates were prepared in analogy to Intermediate 13 starting from the corresponding commercially available bromo-aryl/heteroaryl or iodo-
15 aryl/heteroaryl derivative:

Starting	Amine intermediate		MS m/z	R_t (min)	Method
4-Chloro- 2-fluoro- iodobenze ne	14		230, 232 (M+H) ⁺	0.79	Method 2
4-Iodo- benzotrifu oride	15		246 (M+H) ⁺	0.66	Method 2

-	16 Commercially available from ENAMINE-BB (Cat. Number EN300-185595)	 ClH	-	-	-
2-Bromoquinoline	17	 ClH	229 (M+H)+	0.69	Method 2
-	18 Commercially available from ENAMINE-BB (Cat. Number EN300-50665)	 ClH	-	-	-
4-Bromo-1-(difluoromethoxy)-2-fluorobenzene	19	 ClH	262 (M+H)+	0.68	Method 2

4-Chloro-3-fluoro-iodobenzene	20		230, 232 (M+H)+	0.77	Method 2
2-Bromo-3-fluoro-5-(trifluoromethyl)pyridine	21		265 (M+H)+	0.83	Method 2
2-Iodo-5-(trifluoromethyl)pyridine	22		247 (M+H)+	0.75	Method 2
5-Iodo-2-(trifluoromethyl)pyridine	23		247 (M+H)+	0.71	Method 2
5-Bromo-3-fluoro-2-(trifluoromethyl)pyridine	24		265 (M+H)+	0.81	Method 2
2-Chloro-4-fluoro-iodobenzene	25		230,232 (M+H)+	0.59	Method 2

Intermediate 26 :

Intermediate 26 was prepared as described for Intermediate 13 starting from

- 5 commercially available 4-bromo-3-fluoro-benzotrifluoride (560 mg, 2.30 mmol) and 2-methyl-propane-2-sulfinic acid [dihydro-pyran-(3Z)-ylidene]-amide (390 mg, 1.92 mmol; prepared in analogy to 2-methyl-propane-2-sulfinic acid-(tetrahydro-pyran-4-ylidene)-amide, described in WO2005/87751 A2) to obtain 120 mg of the title compound, as racemic mixture.


- 10 LC-MS (Method 2): $R_t = 1.00$ min

MS (ESI pos): $m/z = 264$ ($M+H$)⁺

The following Amine Intermediates were prepared in analogy to Intermediate 26 starting from the corresponding commercially available bromo-aryl derivative:

15

Starting	Amine intermediate		MS m/z	R_t (min)	Method
4-Bromo-benzotrifluoride	27	 Racemic mixture	246 ($M+H$) ⁺	0.92	Method 2
1-Bromo-4-(trifluoromethoxy)benzene	28	 Racemic mixture	262 ($M+H$) ⁺	0.95	Method 2

Intermediate 29 :




Intermediate 29 was prepared as described for Intermediate 13 starting from commercially available 4-bromo-3-fluoro-benzotrifluoride (462 ,g, 1.90 mmol) and 2-methyl-propane-2-sulfinic acid [dihydro-furan-(3Z)-ylidene]-amide (300 mg, 1.58 mmol; prepared in analogy to 2-methyl-propane-2-sulfinic acid-(tetrahydro-pyran-4-ylidene)-amide, described in WO2005/87751 A2) to obtain 50 mg of the title compound, as racemic mixture.


LC-MS (Method 2): $R_t = 0.90$ min

MS (ESI pos): $m/z = 250$ ($M+H$)⁺

The following Amine Intermediates were prepared in analogy to Intermediate 29 starting from the corresponding commercially available bromo-aryl/heteroaryl or iodo-aryl/heteroaryl derivative:

Starting	Amine intermediate		MS m/z	R_t (min)	Method
4-Iodobenzo trifluoride	30 Racemic mixture		232 ($M+H$) ⁺	0.83	Method 2
4-Chloro-2-fluoroiodo benzene	31 Racemic mixture		216, 218 ($M+H$) ⁺	0.75	Method 2

2-Bromo-3-fluoro-5-(trifluoromethyl)pyridine	32 Racemic mixture		251 (M+H)+	0.76	Method 2
5-Iodo-2-(trifluoromethyl)pyridine	33 Racemic mixture		233 (M+H)+	0.68	Method 2
2-Iodo-5-(trifluoromethyl)pyridine	34 Racemic mixture		233 (M+H)+	0.70	Method 2

Intermediate 35:Step 1:

3,6-Dihydro-2H-pyran-4-boronic acid pinacol ester (5.62 g, 26.75 mmol), 4-bromo-3-fluorobenzotrifluoride (5.00 g, 20.58 mmol), potassium carbonate (8.53 g, 61.73 mmol) and 1,1'-bis(diphenylphosphino)ferrocenedichloro palladium(II) (753 mg, 1.03 mmol) were suspended in 50 mL of 1,4-dioxane and 10 mL of water. The reaction

10

mixture was refluxed for 3 hrs, solvents were evaporated and the crude was extracted with ethyl acetate (50 mL) and water (50 mL). Organic layer was separated, dried over sodium sulfate and evaporated. The crude obtained was purified by flash chromatography (eluent: cyclohexane/AcOEt; gradient from 40 % to 100% of AcOEt) to obtain 4.0 g of the title compound as clear oil.

GC-MS (Method 9): $R_t = 7.76$ min

MS : $m/z = 246$ (M)⁺

Step 2:

To a solution of 4-(2-Fluoro-4-trifluoromethyl-phenyl)-3,6-dihydro-2H-pyran (obtained as described in Step 1; 7.0 g, 25.18 mmol) in 150 mL of dichloromethane, stirred at 0°C, 3-chloroperoxybenzoic acid (11.3 g, 50.37 mmol) was added portionwise. The reaction mixture was allowed to reach room temperature and stirred overnight. The reaction mixture was cooled to 0°C and the precipitate formed was filtered off. The organic solution was washed twice with an aqueous saturated solution of potassium carbonate, dried over sodium sulfate and concentrated under vacuum. The crude obtained was purified by flash chromatography (eluent: cyclohexane/AcOEt; gradient from 50 % to 100% of AcOEt) to obtain 4.2 g of the title compound.

GC-MS (Method 9): $R_t = 7.68$ min

MS : $m/z = 262$ (M)⁺

Step 3:

To a solution of 6-(2-Fluoro-4-trifluoromethyl-phenyl)-3,7-dioxa-bicyclo[4.1.0]heptane (obtained as described in Step 2; 1.64 g, 6.25 mmol) in 10 mL of acetonitrile, stirred under nitrogen atmosphere at -45°C, trifluoromethane sulfonic acid (1.88 g, 12.5 mmol) was added dropwise. The reaction mixture was allowed to reach room temperature and stirred for 2.5 hrs. 10 mL of water were added, the reaction mixture was warmed to 100°C and acetonitrile was distilled out. The reaction mixture was stirred at 100°C for 5 hrs, then cooled to room temperature and stirred overnight. The reaction mixture was diluted with dichloromethane and phases were separated. The aqueous phase was treated with a 4M solution of NaOH until basic pH and extracted with dichloromethane. The organic phase was dried over sodium sulfate and concentrated under vacuum to give 290 mg of the final compound (crude colorless

oil), 4-Amino-4-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-pyran-3-ol, as racemic mixture (TRANS/CIS diastereoisomeric ratio 85:15, determined by NMR).

The crude was used in the next step without any further purification.


LC-MS (METHOD 2): $R_t = 0.77$ min

5 MS (ESI pos): $m/z = 280$ ($M+H$)⁺

1H NMR (500 MHz, DMSO-d₆) δ 7.59 – 7.47 (m, 3H), 4.74 (d, $J = 5.8$ Hz, 1H), 4.04 (dd, $J = 11.7, 1.5$ Hz, 1H), 3.90 (ddd, $J = 12.5, 11.0, 2.0$ Hz, 1H), 3.74 (d, $J = 5.8$ Hz, 1H), 3.67 (dd, $J = 11.1, 4.1$ Hz, 1H), 3.56 – 3.51 (m, 1H), 2.51 – 2.44 (m, 1H), 2.09 (br s, 1H), 1.56 (m, 1H).

NOE: 2.09 (NH₂): 3.74; 4.04. 4.74 (OH): 3.55; 2.45

The following Amino-alcohol Intermediate were prepared in analogy to Intermediate 34 starting from the corresponding commercially available bromo-heteroaryl:


Starting	Amino-alcohol intermediate		MS m/z	R_t (min)	Method
5-Bromo-2-(trifluoromethyl)pyridine	36 Trans racemate		263 ($M+H$) ⁺	0.90	Method 1

Relative stereochemistry of intermediate 36 assigned by NMR and NOE :

20 1H NMR (500 MHz, DMSO-d₆) δ 8.87 (d, $J = 2.3$ Hz, 1H, 13), 8.10 (ddd, $J = 8.3, 2.4, 0.8$ Hz, 1H), 7.81 (dd, $J = 8.3, 0.8$ Hz, 1H), 4.78 (d, $J = 5.9$ Hz, 1H), 4.05 (dd, $J = 11.6, 1.5$ Hz, 1H), 3.91 (td, $J = 11.7, 2.3$ Hz, 1H), 3.68 (ddd, $J = 11.1, 4.9, 2.2$ Hz, 1H), 3.56 (dd, $J = 11.7, 2.4$ Hz, 1H), 3.47 (d, $J = 5.9$ Hz, 1H), 2.49 – 2.44 (m, 1H), 2.12 (s, 2H), 1.49 (dd, $J = 13.1, 1.9$ Hz, 1H).

NOE : 2.12 (NH₂): 3.47; 3.91;4.05. 4.78 (OH): 3.56; 2.48

Intermediate 37



Step 1:

tert-Butyllithium (21.8 mL, 1.7M in pentane, 37.0 mmol) was added dropwise to 4-bromo-3-fluorobenzotrifluoride (5.00 g, 20.58 mmol) in THF (50 mL) at -70°C. After 1h, 3-oxotetrahydrofuran (1.78 g, 20.58 mmol) in THF was added dropwise. The reaction mixture was warmed to -10 °C mixture and quenched with NH₄Cl saturated solution. Ethyl acetate was added, the organic layer was separated, dried over sodium sulfate and evaporated. The crude obtained was purified by flash chromatography (eluent: cyclohexane/AcOEt; gradient from 0 % to 80% of AcOEt) to obtain 1.6 g of the 3-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-furan-ol.

GC-MS (Method 9): R_t = 7.58 min

MS : m/z = 250 (M)⁺

Step 2:

p-Toluenesulfonic acid monohydrate (1.75 g, 9.19 mmol) was added to 3-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-furan-ol (obtained as described in Step 1; 2.3 g, 9.19 mmol) in toluene (20 mL). After refluxing for 1h, volatiles were evaporated, DCM and water were added, the organic layer was separated, dried over sodium sulfate and evaporated to obtain 2.0 g (77% content) of crude 3-(2-fluoro-4-trifluoromethyl-phenyl)-2,5-dihydro-furan, that was used without further purification.

GC-MS (Method 9): R_t = 7.12-7.21 min

MS : m/z = 232 (M)⁺

Step 3:

To a solution of 3-(2-fluoro-4-trifluoromethyl-phenyl)-2,5-dihydro-furan (obtained as described in Step 2; 2.0 g 77% content, 6,63 mmol) in 50 mL of dichloromethane, stirred at 0°C, 3-chloroperoxybenzoic acid (2.63 g, 15,26 mmol) was added

portionwise. The reaction mixture was allowed to reach room temperature and stirred overnight. The reaction mixture was cooled to 0°C and the precipitate formed was filtered off. The organic solution was washed twice with an aqueous saturated solution of potassium carbonate, dried over sodium sulfate and concentrated under vacuum. The crude obtained was purified by flash chromatography (eluent:

cyclohexane/AcOEt; gradient from 50 % to 100% of AcOEt) to obtain 1.2 g (98% content) of 1-(2-fluoro-4-trifluoromethyl-phenyl)-3,6-dioxa-bicyclo[3.1.0]hexane.

¹H NMR (300 MHz, DMSO-*d*₆): δ ppm 3.91-3.95 (m, 2H), 4.06-4.19 (m, 3H) 7.37 (dd, *J* = 10.2, 1.3 Hz, 1H), δ 7.47 (dd, *J* = 8.4, 1.1 Hz, 1H), 7.59 (m, 1H)

Step 4:

To a solution of 1-(2-fluoro-4-trifluoromethyl-phenyl)-3,6-dioxa-bicyclo[3.1.0]hexane (obtained as described in Step 3; 1.20 g, 98% content, 4,74 mmol) in 20 mL of acetonitrile, stirred under nitrogen atmosphere at -40°C, trifluoromethane sulfonic acid (0.84 mL, 9.48 mmol) was added dropwise. The reaction mixture was allowed to reach room temperature and stirred for 2.5 hrs. 20 mL of water were added, the reaction mixture was warmed to 100°C and acetonitrile was distilled out. The reaction mixture was stirred at 100°C for 20 hrs, then cooled to room temperature and stirred overnight. The reaction mixture was diluted with dichloromethane and phases were separated. The aqueous phase was treated with a 4M solution of NaOH until basic pH and extracted with dichloromethane. The organic phase was dried over sodium sulfate and concentrated under vacuum to give 200 mg of 4-amino-4-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-furan-3-ol, as racemic mixture (TRANS/CIS diastereoisomeric ratio 88/12, determined by NMR).

The crude was used in the next step without any further purification.


LC-MS (METHOD 1): *R*_t = 2.52-3.04 min

MS (ESI pos): *m/z* = 266 (*M*+H)⁺

¹H NMR (400 MHz, DMSO-d₆) δ 7.59 – 7.54 (m, 1H), 7.52 – 7.45 (m, 2H), 5.08 (d, J = 4.8 Hz, 1H), 4.29 (q, J = 3.8 Hz, 1H), 4.25 (dd, J = 8.8, 3.8 Hz, 1H), 4.14 (d, J = 7.6 Hz, 1H), 3.95 (dd, J = 8.0, 2.2 Hz, 1H), 3.65 (d, J = 8.8 Hz, 1H), 2.06 (s, 2H).

5 NOE: 2.06 (NH₂): 3.95; 4.29; 4.25. 5.08 (OH): 4.14; 3.65

Intermediate 38:



10

Step 1:

6-(2-Fluoro-4-trifluoromethyl-phenyl)-3,7-dioxabicyclo[4.1.0]heptane (obtained as described in Step 2 in the preparation of Intermediate 34; 4.20 g, 16.02 mmol), sodium azide (2.08 g, 32.04 mmol) and ammonium chloride (1.72 g, 32.04 mmol) were suspended in 50 mL of methanol and 10 mL of water. The reaction mixture was stirred at reflux for 18 hrs. Solvents were removed, the crude was suspended in water and extracted twice with ethyl acetate. The organic layer was dried over sodium sulfate and concentrated under vacuum to give 4.70 g of the final compound as a mixture of the desired regioisomer 4-Azido-4-(2-fluoro-4-trifluoromethyl-phenyl)-

tetrahydro-pyran-3-ol and the undesired regioisomer 3-azido-4-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-pyran-4-ol in a regioisomeric ratio of 76/24 determined by NMR. The regioisomeric mixture was used in the next step without separation.

5 GC-MS (METHOD 9): $R_t = 9.57$ min

MS : $m/z = 248$ (M)⁺

1H NMR (500 MHz, DMSO-d₆) δ 7.77 – 7.74 (m, 1H), 7.72 (d, $J = 7.9$ Hz, 1H), 7.67 – 7.64 (m, 1H), 5.37 (d, $J = 6.0$ Hz, 1H), 3.93 – 3.89 (m, 1H), 3.85 – 3.80 (m, 1H), 3.78
10 (dd, $J = 12.3, 1.5$ Hz, 1H), 3.74 – 3.66 (m, 2H), 2.65 (ddd, $J = 13.8, 11.9, 4.8$ Hz, 1H), 2.02 – 1.95 (m, 1H).

Step 2:

A mixture of 4-Azido-4-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-pyran-3-ol and
15 3-azido-4-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-pyran-4-ol (obtained as described in Step 1, 2.0 g, 6.55 mmol), Pd/C (300 mg, 2.82 mmol) and di-tert-butylidicarbonate (1.86 g, 8.52 mmol) were suspended in 150 mL of ethanol. The reaction mixture was stirred at room temperature under hydrogen atmosphere (2.5 bar) for 1 hr. The reaction mixture was filtered on a celite pad and the organic
20 solution was concentrated under vacuum. The crude obtained was purified by flash chromatography (eluent: cyclohexane/AcOEt; gradient from 10 % to 100% of AcOEt) to obtain 1.75 g of the title compound (yellow solid) as a mixture of the desired regioisomer [4-(2-fluoro-4-trifluoromethyl-phenyl)-3-hydroxy-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester and the undesired regioisomer [4-(2-Fluoro-4-
25 trifluoromethyl-phenyl)-4-hydroxy-tetrahydro-pyran-3-yl]-carbamic acid tert-butyl ester in a regioisomeric ratio of 85/15 determined by NMR. The regioisomeric mixture was used in the next step without separation.

GC-MS (METHOD 9): $R_t = 10.92$ -10.99 min

MS : $m/z = 323$ (M)⁺

30

Step 3:

A mixture of [4-(2-Fluoro-4-trifluoromethyl-phenyl)-3-hydroxy-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester and 4-(2-Fluoro-4-trifluoromethyl-phenyl)-4-hydroxy-

tetrahydro-pyran-3-yl]-carbamic acid tert-butyl ester (obtained as described in Step 2, 3.7 g, 7.30 mmol) was dissolved in 20 mL of dichloromethane, Dess-Martin periodinane (2.18 g, 9.5 mmol) was added portionwise and the reaction mixture was stirred at room temperature for 2 hrs. The reaction mixture was diluted with
5 dichloromethane, washed with aqueous bicarbonate saturated solution, washed with aqueous sodium bisulfite saturated solution, the organic layer was separated, dried over sodium sulfate and concentrated under vacuum. The crude obtained was purified by flash chromatography (eluent: dichloromethane/AcOEt; gradient from 0% to 70% of AcOEt) to give 2.4 g of the desired compound.

10 LC-MS (METHOD 1): $R_t = 4.23-4.83$ min
MS (ESI pos): $m/z = 278$ (fragment) $(M+H)^+$

Step 4:

15 [4-(2-Fluoro-4-trifluoromethyl-phenyl)-3-oxo-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester (obtained as described in Step 3; 340 mg, 0.9 mmol) was suspended in 10 mL of dry THF. The reaction mixture was stirred at -20°C and 0.29 mL of a 3.4M solution of methylmagnesium bromide in methyl-tetrahydrofuran was added dropwise. The reaction mixture was stirred at -20°C for 1 hr, then quenched with aqueous saturated ammonium chloride solution. Organic layer was separated, dried
20 over sodium sulfate and concentrated under vacuum. The crude obtained was purified by flash chromatography (eluent: dichloromethane/AcOEt; gradient from 0% to 30% of AcOEt) to give 200 mg of the title compound, 4-(2-fluoro-4-trifluoromethyl-phenyl)-3-hydroxy-3-methyl-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester, as racemic mixture (TRANS/CIS diastereoisomeric ratio 82/12, determined by NMR).

25 GC-MS (METHOD 9): $R_t = 11.01$ min
MS : $m/z = 292$ (fragment) $(M)^+$

^1H NMR (500 MHz, DMSO- d_6) δ 7.58 – 7.53 (m, 1H), 7.52 – 7.48 (m, 1H), 7.45 (d, $J = 12.8$ Hz, 1H), 6.92 (s, 1H), 4.78 (s, 1H), 3.88 (d, $J = 12.2$ Hz, 1H), 3.70 (t, $J = 6.7$
30 Hz, 1H), 3.60 (q, $J = 12.8, 12.2$ Hz, 1H), 3.28 – 3.26 (d, $J = 12.2$ Hz, 1H), 2.88 (t, $J = 11.4$ Hz, 1H), 1.33 (s, 9H), 0.9 (s, 3H).

NOE: 6.92 (NH): 0.90; 3.88. 4.78 (OH): 2.88; 3.27

Step 5:


4-(2-Fluoro-4-trifluoromethyl-phenyl)-3-hydroxy-3-methyl-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester (obtained as described in Step 4, as preferred diastereoisomer; 200 mg, 0.51 mmol) was dissolved in 5 mL of dichloromethane.

- 5 Trifluoroacetic acid (0.39 mL, 5.1 mmol) was added, the reaction mixture was stirred at room temperature for 1hr and then concentrated under vacuum. The crude obtained was stripped twice with ethyl ether to give 198 mg of the title compound, 4-amino-4-(2-fluoro-4-trifluoromethyl-phenyl)-3-methyl-tetrahydro-pyran-3-ol trifluoroacetate salt as racemic mixture (TRANS/CIS diastereoisomeric ratio 85/15, determined by NMR).
- 10

LC-MS (METHOD 1): $R_t = 3.35$ min

MS (ESI pos): $m/z = 294$ ($M+H$)⁺

- 15 ¹H NMR (500 MHz, DMSO-d₆) δ 8.61 (s, 3H), 7.96 (t, $J = 8.2$ Hz, 1H), 7.81 (dd, $J = 13.3, 2.0$ Hz, 1H), 7.70 – 7.66 (m, 1H), 5.40 (s, 1H), 3.98 (ddd, $J = 13.3, 10.7, 2.8$ Hz, 1H), 3.92 – 3.85 (m, 1H), 3.62 (d, $J = 12.6$ Hz, 1H), 3.40-3.38 (d, $J = 12.6$ Hz, 1H), 2.99 (ddd, $J = 14.4, 10.7, 5.2$ Hz, 1H), 1.79 (dt, $J = 14.4, 3.0$ Hz, 1H), 1.11 (d, $J = 1.8$ Hz, 3H).

Intermediate 39:

5 Step 1:

Step 1 was performed in analogy to Step 1 in the preparation of Intermediate 35, starting from 2-bromo-3-fluoro-5-(trifluoromethyl)pyridine (5 g, 20.49 mmol) to obtain 2-(3,6-dihydro-2H-pyran-4-yl)-3-fluoromethyl-pyridine (5.7 g).

10 LC-MS (METHOD 2): $R_t = 1.17$ min

MS (ESI pos): $m/z = 248$ ($M+H$)⁺

Step 2:

Step 2 was performed in analogy to Step 2 in the preparation of Intermediate 35, starting from 2-(3,6-Dihydro-2H-pyran-4-yl)-3-fluoromethyl-pyridine (5.7 g, 23.06 mmol) to obtain 2-(3,7-dioxa-bicyclo[4.1.0]hept-6-yl)-3-fluoro-5-trifluoromethyl-pyridine (3.25 g).

LC-MS (METHOD 2): $R_t = 0.95$ min

MS (ESI pos): $m/z = 264$ ($M+H$)⁺

Step 3:

Step 3 was performed in analogy to Step 1 in the preparation of Intermediate 38, starting from 2-(3,7-Dioxa-bicyclo[4.1.0]hept-6-yl)-3-fluoro-5-trifluoromethyl-pyridine (250 mg, 0.95 mmol) to obtain after purification by flash chromatography (eluent: cyclohexane/EtOAc; gradient from 0% to 30% of EtOAc), 4-azido-4-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-tetrahydro-pyran-3-ol (160 mg) as major regioisomer

LC-MS (METHOD 2): $R_t = 1.05$ min

MS (ESI pos): $m/z = 307$ (M+H)⁺

¹H NMR (500 MHz, DMSO-d₆) δ 8.90 (dq, $J = 2.0, 0.9$ Hz, 1H), 8.39 (ddd, $J = 11.5, 1.9, 0.7$ Hz, 1H), 5.42 (s, 1H), 4.01 (s, 1H), 3.93 – 3.86 (m, 1H), 3.78 – 3.64 (m, 3H), 2.73 (ddd, $J = 14.6, 12.6, 4.9$ Hz, 1H), 2.02 (dq, $J = 14.7, 2.0$ Hz, 1H).

The minor regioisomer, 3-Azido-4-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-tetrahydro-pyran-4-ol, was also isolated (40mg).

LC-MS (METHOD 2): $R_t = 1.04$ min

MS (ESI pos): $m/z = 307$ (M+H)⁺

Step 4:

To a solution of 4-azido-4-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-tetrahydro-pyran-3-ol (160 mg, 0.52 mmol) in 5 mL of methanol stirred under nitrogen atmosphere, ammonium formate (165 mg, 2.61 mmol) and zinc (51.2 mg, 0.78 mmol) were added.

The reaction mixture was stirred at room temperature overnight and concentrated. A saturated ammonium chloride water solution was added and the reaction mixture was extracted with dichloromethane. The organic phase was separated, washed with brine, dried over sodium sulfate and concentrated under vacuum to give 115 mg of 4-Amino-4-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-tetrahydro-pyran-3-ol, as TRANS-racemic mixture.


LC-MS (METHOD 5): $R_t = 0.71$ min

MS (ESI pos): $m/z = 281$ (M+H)⁺

1H NMR (500 MHz, DMSO-d₆) δ 8.75 (dq, J = 2.0, 1.0 Hz, 1H), 8.18 – 8.12 (m, 1H),
 4.80 (d, J = 5.6 Hz, 1H), 4.01 (dd, J = 11.7, 1.5 Hz, 1H), 3.89 – 3.83 (m, 1H), 3.78 (dt,
 J = 5.5, 1.9 Hz, 1H), 3.74 – 3.66 (m, 1H), 3.55 (dd, J = 11.7, 1.7 Hz, 1H), 2.71 – 2.61
 5 (m, 1H), 2.09 (s, 2H), 1.65 – 1.58 (m, 1H).

NOE: 2.09 (NH₂): 3.55; 3.70; 3.78 4.80 (OH): 1.61; 3.78

The following intermediate was prepared in analogy to Intermediate 39, starting from
 10 2-Bromo-5-trifluoromethyl-pyridine


Starting	Amino-alcohol intermediate		MS m/z	R _t (min)	Method
2-Bromo-5-trifluoromethyl pyridine	40		263	0.68	METHOD 2

1H NMR (500 MHz, DMSO-d₆) δ 8.89 (dq, J = 2.6, 0.9 Hz, 1H), 8.13 (ddd, J = 8.4,
 2.5, 0.8 Hz, 1H), 7.69 (dt, J = 8.4, 0.8 Hz, 1H), 4.69 (d, J = 5.6 Hz, 1H), 4.05 (dd, J =
 15 11.4, 1.7 Hz, 1H), 3.84 (td, J = 11.3, 2.4 Hz, 1H), 3.72 (ddd, J = 10.9, 4.7, 2.7 Hz,
 1H), 3.60 (ddd, J = 5.7, 2.8, 1.4 Hz, 1H), 3.54 (dd, J = 11.5, 2.8 Hz, 1H), 2.56 – 2.51
 (m, 1H), 2.02 (s, 2H), 1.62 – 1.52 (m, 1H).

NOE: 2.09 (NH₂): 3.55; 3.70; 3.78 4.80 (OH): 1.61; 3.78

20

Intermediate 41:



41

Step 1:

Step 1 was performed in analogy to Step 1 in the preparation of Intermediate 38, starting from 1-(2-fluoro-4-trifluoromethyl-phenyl)-3,6-dioxabicyclo[3.1.0]hexane (750, 3.02 mmol, prepared as described in Step 3 in the preparation of Intermediate 37) to obtain 4-azido-4-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-furan-3-ol as major regioisomer and 4-Azido-3-(2-fluoro-4-trifluoromethyl-phenyl)-tetrahydro-furan-3-ol as minor regioisomer . (900mg, regioisomer ratio 82/18 determined by NMR)

GC-MS (METHOD 9): $R_t = 9.22$ min

MS : $m/z = 190$ (fragment) (M)⁺

¹H NMR (500 MHz, DMSO-d₆) δ 7.82 – 7.77 (m, 1H), 7.66 – 7.60 (m, 2H), 5.72 (d, J = 5.4 Hz, 1H), 4.59 – 4.55 (m, 1H), 4.39 (dd, J = 9.7, 1.8 Hz, 1H), 4.23 (d, J = 9.7 Hz, 1H), 4.18 (dd, J = 9.6, 4.1 Hz, 1H), 3.79 – 3.75 (d, 1H).

Step 2:

Step 2 was performed in analogy to Step 1 in the preparation of Intermediate 38, starting from the regioisomeric mixture , to obtain [3-(2-fluoro-4-trifluoromethyl-

phenyl)-4-hydroxy-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester as preferred regioisomer and 4-(2-Fluoro-4-trifluoromethyl-phenyl)-4-hydroxy-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester (670 mg) (regioisomer ratio 80/20 determined by NMR).

- 5 GC-MS (METHOD 9): R_t = 10.67 min
MS : m/z = 265 (fragment) (M)+

Step 3:

- 10 Step 3 was performed in analogy to Step 3 in the preparation of Intermediate 38, starting from the regioisomeric mixture of [3-(2-fluoro-4-trifluoromethyl-phenyl)-4-hydroxy-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester and 4-(2-Fluoro-4-trifluoromethyl-phenyl)-4-hydroxy-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester (670 mg, 1.47 mmol), to obtain [3-(2-fluoro-4-trifluoromethyl-phenyl)-4-oxo-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester (455 mg).

- 15 GC-MS (METHOD 9): R_t = 10.15 min
MS : m/z = 249 (fragment) (M)+

Step 4:

- 20 Step 4 was performed in analogy to Step 4 in the preparation of Intermediate 38, starting from [3-(2-fluoro-4-trifluoromethyl-phenyl)-4-oxo-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester (455 mg, 1.23) to obtain [3-(2-fluoro-4-trifluoromethyl-phenyl)-4-hydroxy-4-methyl-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester as -racemic mixture (TRANS/CIS diastereoisomeric ratio 91/9, determined by NMR). (365 mg).

- 25 LC-MS (METHOD 10): R_t = 3.46-3.62 min
MS (ESI pos): m/z = 280 (fragment) (M+H)⁺

- 30 ¹H NMR (500 MHz, DMSO-d₆) δ 7.64 (t, J = 7.9 Hz, 1H), 7.51 (d, J = 9.5 Hz, 2H), 7.04 (s, 1H), 4.98 (s, 1H), 4.72 – 4.65 (m, 1H), 4.13 (d, J = 8.4 Hz, 1H), 3.94 (d, J = 8.8 Hz, 1H), 3.60 (d, J = 8.8 Hz, 1H), 1.31 (s, 12H).

NOE: 7.04 (NH): 3.94; 4.13; 1.31 4.98 (OH): 1.31; 4.68 1.31 (Me): 7.04; 4.13; 4.98


Step 5:

Step 5 was performed in analogy to Step 5 in the preparation of Intermediate 38, starting from [3-(2-fluoro-4-trifluoromethyl-phenyl)-4-hydroxy-4-methyl-tetrahydro-furan-3-yl]-carbamic acid tert-butyl ester (365 mg, 1.0 mmol) to obtain 4-amino-4-(2-
5 fluoro-4-trifluoromethyl-phenyl)-3-methyl-tetrahydro-furan-3-ol trifluoroacetate, as racemic mixture (TRANS/CIS diastereoisomeric ratio 90/10, determined by NMR). (378 mg)

LC-MS (METHOD 1): R_t = 2.91-3.19 min

MS (ESI pos): m/z = 280 ($M+H$)⁺

10 ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.67 (s, 3H), 7.88 – 7.79 (m, 2H), 7.70 (dd, J = 8.5, 1.9 Hz, 1H), 5.62 (s, 1H), 4.62 (dd, J = 10.0, 1.1 Hz, 1H), 4.19 (dd, J = 10.0, 1.5 Hz, 1H), 3.98 (d, J = 9.6 Hz, 1H), 3.80 (d, J = 9.6 Hz, 1H), 1.48 (d, J = 1.3 Hz, 3H).

Intermediate 42:

Intermediate 42 was prepared in analogy to Intermediate 35, starting from 4-iodobenzotrifluoride (3 g, 10.7 mmol) to obtain, after chromatographic purification in the third step (eluent: cyclohexane/EtOAc; gradient from 0% to 100% of EtOAc), 105 mg the title compound, as racemic mixture (TRANS/CIS diastereoisomeric ratio 93/7,
20 determined by NMR).

LC-MS (METHOD 5): R_t = 0.77 min


MS (ESI pos): m/z = 262 ($M+H$)⁺

25 ¹H NMR (500 MHz, DMSO-*d*₆) δ 7.71 – 7.67 (m, 2H), 7.65 – 7.61 (m, 2H), 4.58 (d, J = 5.9 Hz, 1H), 4.05 (dd, J = 11.5, 1.5 Hz, 1H), 3.92 – 3.86 (m, 1H), 3.67 (ddd, J = 11.1, 4.9, 2.3 Hz, 1H), 3.57 – 3.52 (m, 1H), 3.48 (dq, J = 5.7, 1.5 Hz, 1H), 2.49 – 2.41 (m, 1H), 1.97 (s, 2H), 1.49 – 1.42 (m, 1H).

NOE 1.97 (NH₂): 3.48; 4.05; 1.47 4.58 (OH): 2.46; 3.55 3.48 (CH): 1.97; 3.89

Intermediate 43:

5



Step 1:

Step 1 was performed in analogy to Step 2 in the preparation of Intermediate 38, starting from 4-Azido-4-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-tetrahydro-pyran-3-ol (1.7g, 4.77mmol) to obtain after filtration on silica, 1.3g of the title compound as TRANS-racemic mixture

LC-MS (METHOD 1): R_t = 3.93 min

MS (APCI): m/z = 381 (M+H)⁺

15

Step 2:

To a solution of oxalylchloride (0.25mL, 2.6mmol) in dry DCM (20mL) at -55⁰C, DMSO (0.37mL, 0.51mmol) was added dropwise. After 20min, a solution of (3-fluoro-5-trifluoromethyl-pyridin-2-yl)-3-hydroxy-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester (1.0g, 2.37mmol) dissolved in 3mL of dry DCM was added dropwise.

- 5 Mixture is stirred for 1h at -70 °C , then TEA was added dropwise, stirred for 1.30hrs at -40 °C and then allowed to reach room temperature. Mixture was stirred at room temperature for 48hrs. Solvent was evaporated, residue diluted with 50 ml of ethyl acetate and washed with 3 x 10 ml of water. Organic phase was separated, dried over sodium sulfate to obtain 0.92g of the desired compound used in the next step
- 10 without further purification.

LC-MS (METHOD 1): R_t = 3.79min

MS (APCI): m/z = 475 (M+H)⁺

Step 3:

- 15 Step 3 was performed in analogy to Step 4 in the preparation of Intermediate 38, starting from 4-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-3-oxo-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester (0.92g, 2.41mmol) to obtain after chromatographic purification (eluent: cyclohexane/EtOAc; gradient from 100% to 40% of EtOAc), 0.190g of CIS stereoisomer and 0.35g of TRANS stereoisomer, both
- 20 as racemic mixture.

LC-MS (METHOD 9): R_t = 10.49min (CIS stereoisomer)

MS (ESI): m/z = 394 (M+H)⁺

- 25 ¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.88 (s, 3 H) 1.22 - 1.38 (m, 9 H) 2.26 - 2.42 (m, 1 H) 2.80 (td, J=13.54, 4.65 Hz, 1 H) 3.25 - 3.29 (m, 1 H) 3.54 - 3.63 (m, 2 H) 3.78 (br dd, J=11.37, 3.42 Hz, 1 H) 5.31 (s, 1 H) 6.40 (br s, 1 H) 8.18 (br d, J=11.61 Hz, 1 H) 8.81 - 8.86 (m, 1 H)

- 30 NOE : 6.40 (NH): 5.31 5.31 (OH): 6.40

LC-MS (METHOD 9): R_t = 10.76min (TRANS stereoisomer)

MS (ESI): m/z = 394 (M+H)⁺

1H NMR (500 MHz, DMSO-d₆) δ ppm 1.03 (d, J=2.93 Hz, 3 H) 1.10 - 1.44 (m, 9 H)
1.96 - 2.03 (m, 1 H) 3.06 - 3.18 (m, 1 H) 3.24 (d, J=11.98 Hz, 1 H) 3.54 - 3.69 (m, 1
5 H) 3.72 - 3.86 (m, 2 H) 4.71 (s, 1 H) 7.01 (br s, 1 H) 8.09 (br d, J=11.74 Hz, 1 H) 8.79
(s, 1 H)

NOE : 7.01 (NH): 1.02; 3.62; 3.77 4.71 (OH): 3.12; 3.24

10 Step 4:

Step 4 was performed in analogy to Step 5 in the preparation of Intermediate 38,
starting from TRANS [(R)-4-(3-Fluoro-5-trifluoromethyl-pyridin-2-yl)-3-hydroxy-3-
methyl-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester (0.350g, 0.89mmol) to
obtain 0.25g of desired intermediate 42 , 4-Amino-4-(3-fluoro-5-trifluoro
15 methyl-pyridin-2-yl)-3-methyl-tetrahydro-pyran-3-ol, as trifluoroacetate salt.


LC-MS (METHOD 1): R_t = 2.90min

MS (APCI): m/z = 295 (M+H)⁺

20 1H NMR (500 MHz, DMSO-d₆) δ ppm 1.00 (s, 3 H) 1.87 - 1.93 (m, 1 H) 3.14 (ddd,
J=14.49, 9.72, 4.65 Hz, 1 H) 3.39 - 3.43 (m, 1 H) 3.66 - 3.69 (m, 1 H) 3.81 - 3.88 (m,
1 H) 4.04 (dt, J=11.86, 4.34 Hz, 1 H) 5.48 (br s, 1 H) 8.43 - 8.48 (m, 1 H) 8.72 (br s, 3
H) 8.94 (s, 1 H)

25 NOE : 8.72 (NH₃⁺): 1.00; 1.90; 3.68; 3.84 1.0 (Me):8.72; 1.89; 3.66

Intermediate 44:




[4-(2-Fluoro-4-trifluoromethyl-phenyl)-3-oxo-tetrahydro-pyran-4-yl]-carbamic acid tert-butyl ester (obtained from Step 3 in the preparation of Intermediate 38; 90 mg, 0.23 mmol) was dissolved in 1 mL of 1,4-dioxane. 1.67 mL of a 1.4M solution of hydrogen chloride in 1,4-dioxane was added, the reaction mixture was stirred at room temperature for 1 hr and then concentrated under vacuum. The crude obtained was stripped twice with ethyl ether to give 70 mg of the title compound, used in the next step without further purification.

LC-MS (METHOD 1): $R_t = 3.70$ min

MS (ESI pos): $m/z = 278$ ($M+H$)⁺

Example compounds::

Example 1:











To a suspension of Intermediate 5 (40.68 mg, 0.20 mmol) in 1.0 mL of dry DMF, HATU (82.47 mg, 1.3 mmol) and DIPEA (0.11 mL 0.67 mmol) were added and the reaction mixture was stirred at room temperature. A solution of Intermediate 13 (50 mg, 0.17 mmol) in 1.0 mL of dry DMF was added, and the reaction mixture was stirred at room temperature overnight. The reaction mixture was treated with basic alumina and concentrated under vacuum. The crude obtained was purified by flash chromatography (eluent: water/acetonitrile; gradient from 10% to 100% of acetonitrile) to give 62 mg of the desired compound.








LC-MS (METHOD 1): $R_t = 4.78$ min








MS (ESI pos): $m/z = 463$ ($M+H$)⁺








5 The following Examples were prepared in analogy to Example 1 and purified applying the most suitable purification technique, starting from the corresponding Acid and the corresponding Amine Intermediates:









Starting Acid	Starting Amine	Ex.	Structure	MS m/z [$M+H$] ⁺	R_t (min)	Method
2	13	2		423	4,53	METHOD 1
6	13	3		427	3,50	METHOD 10
8	14	4		403	0.90	METHOD 6
2	14	5		389	4,28	METHOD 1

5	25	6		415	4,67	METHOD 1
5	14	7		415	4,78	METHOD 1
5	15	8		431	4,98	METHOD 1
2	15	9		405	4,40	METHOD 1
5	16	10		411	0.80	METHOD 7
12	17	11		428	3.21	METHOD 10
2	17	12		388	2.86	METHOD 10
5	18	13		441	0.87	METHOD 8


5	19	14		447	3,54	METHOD 10
2	19	15		421	3,26	METHOD 10
5	20	16		415	4,70	METHOD 1
12	21	17		464	4,93	METHOD 1
8	21	18		438	3,38	METHOD 10
2	21	19		424	3,27	METHOD 10
5	21	20		450	3,59	METHOD 10

12	22	21		446	4,52	METHOD 1
8	22	22		420	3,23	METHOD 10
5	22	23		432	3,45	METHOD 10
2	22	24		406	3,12	METHOD 10
12	23	25		446	4,75	METHOD 1
8	23	26		420	3,13	METHOD 10
5	23	27		432	3,32	METHOD 10

2	23	28		406	3,01	METHOD 10
12	24	29		464	3,70	METHOD 10
5	26	30 Racemic mixture		449	3,87	METHOD 10
2	26	31 Racemic mixture		423	3,60	METHOD 10
5	27	32 Racemic mixture		[M+Na] + 453	4,78	METHOD 3
5	28	33 Racemic mixture		447	4,83	METHOD 3
5	29	34 Racemic mixture		435	4,92	METHOD 1

2	29	35 Racemic mixture		409	3,44	METHOD 10
2	30	36 Racemic mixture		391	3,36	METHOD 10
5	30	37 Racemic mixture		417	4,77	METHOD 1
5	31	38 Racemic mixture		401	4,71	METHOD 1
2	31	39 Racemic mixture		[M+Na] + 397	2,81	METHOD 4
12	32	40 Racemic mixture		450	3,63	METHOD 10
8	32	41 Racemic mixture		424	3,32	METHOD 10
12	33	42 Racemic mixture		432	3,47	METHOD 10

8	33	43 Racemic mixture		406	3,14	METHOD 10
8	34	44 Racemic mixture		406	3,25	METHOD 10
12	34	45 Racemic mixture		432	3,53	METHOD 10


Example 46:

- 5 Example 46 was synthesized in analogy to Example 1, starting from acid Intermediate 2 (34.6 mg, 0.2 mmol) and amino alcohol Intermediate 35 (65 mg, 0.19 mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 100% of acetonitrile), 27 mg of the title compound, as TRANS-racemic mixture.
- 10 LC-MS (METHOD 1): $R_t = 3.70$ min
 MS (ESI pos): $m/z = 439$ ($M+H$)⁺
¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.40 (d, $J=1.00$ Hz, 3 H) 2.59 - 2.73 (m, 2 H) 3.62 - 3.71 (m, 1 H) 3.79 - 3.85 (m, 2 H) 3.82 (br d, $J=11.62$ Hz, 2 H) 4.02 (br d, 1 H) 4.10 (m, 1 H) 5.33 (br s, 1 H) 7.46 - 7.50 (m, 1 H) 7.55 (d, $J=7.99$ Hz, 1 H) 7.73 (t,

J=8.19 Hz, 1 H) 8.34 (s, 1 H) 8.42 (s, 1 H) 8.81 (d, J=2.08 Hz, 1 H) 9.19 (dd, J=1.96, 1.10 Hz, 1 H)

NOE: 8.42 (NH): 4.10; 4.03; 3.65; 3.82 5.33 (OH): 4.10; 3.70

5 **Example 47:**



Further elution from the column in the preparation of Example 46 gave 5 mg of the title compound, as CIS- racemic mixture.

10 LC-MS (METHOD 1): $R_t = 3.97$ min



MS (ESI pos): $m/z = 439$ ($M+H$)⁺

¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.35 - 2.48 (m, 4 H) 3.12 (br d, J=14.18 Hz, 1 H) 3.47 (br t, J=11.86 Hz, 1 H) 3.58 (t, J=10.76 Hz, 1 H) 3.72 (br dd, J=11.62, 2.57 Hz, 1 H) 3.74 - 3.84 (m, 1 H) 4.10 (dd, J=10.03, 5.14 Hz, 1 H) 5.66 (s, 1 H) 7.47 - 7.59 (m, 2 H) 7.61 - 7.67 (m, 1 H) 8.49 (s, 1 H) 8.62 (s, 1 H) 8.76 (d, J=1.96 Hz, 1 H) 9.21 (dd, J=2.08, 1.10 Hz, 1 H)


NOE: 8.62 (NH): 4.10; 5.66; 2.49; 3.78; 3.12 5.66(OH): 8.62; 4.10; 3.78; 3.12


The following Examples were prepared in analogy to Example 46 and Example 47
20 starting from the corresponding acid and amino alcohol Intermediates:


Starting Acid	Starting Amino-alcohol	Ex.	Structure	MS m/z [$M+H$] ⁺	R_t (min)	Method

5	35	48 TRANS- Racemic mixture		465	4.12	METHOD 1
5	35	49 CIS- Racemic mixture		465	4.33	METHOD 1

Relative stereochemistry assigned by NMR:

Example		Relative stereochemistry	¹ H-NMR	NOE
48		TRANS Racemic mixture	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 0.94 - 1.11 (m, 4 H) 2.11 (tt, J=8.47, 5.10 Hz, 1 H) 2.56 - 2.76 (m, 2 H) 3.66 (td, J=11.68, 2.08 Hz, 1 H) 3.81 (br d, J=11.25 Hz, 2 H) 4.03 (d, J=11.74 Hz, 1 H) 4.09 (m, 1 H) 5.33 (br s, 1 H) 7.46 - 7.50 (m, 1 H) 7.54 (d, J=8.03 Hz, 1 H) 7.73 (t, J=7.98 Hz, 1 H)	8.40 (NH): 4.09 5.33(OH): 4.09

			8.33 (s, 1 H) 8.40 (s, 1 H) 8.77 (d, J=2.20 Hz, 1 H) 9.13 (d, J=2.20 Hz, 1 H)	
49		CIS-Racemic mixture	1H NMR (500 MHz, DMSO-d6) δ ppm 0.86 - 1.07 (m, 4 H) 2.06 - 2.16 (m, 1 H) 2.42 - 2.48 (m, 1 H) 3.12 (br d, J=14.18 Hz, 1 H) 3.26 - 3.29 (m, 1 H) 3.47 (br t, J=11.98 Hz, 1 H) 3.58 (t, J=10.76 Hz, 1 H) 3.65 - 3.75 (m, 1 H) 3.78 (dd, J=11.49, 5.14 Hz, 1 H) 4.09 (dd, J=10.27, 5.14 Hz, 1 H) 5.66 (br s, 1 H) 7.48 - 7.60 (m, 2 H) 7.60 - 7.65 (m, 1 H) 8.48 (s, 1 H) 8.58 (s, 1 H) 8.71 (d, J=2.20 Hz, 1 H) 9.14 (d, J=1.71 Hz, 1 H)	8.58 (NH): 5.66; 3.58; 3.47; 3.12 5.66(OH): 8.58; 3.77 4.09 (CH): 2.46

Example 50:

Example 50 was synthesized in analogy to Example 1, starting from acid

- 5 Intermediate 2 (100 mg, 0.56 mmol) and amino alcohol Intermediate 36 (171.4 mg, 0.62 mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 0% to 60% of acetonitrile), 209 mg of the title compound, as TRANS-racemic mixture.

LC-MS (METHOD 1): $R_t = 3.04$ min

- 10 MS (ESI pos): $m/z = 422$ ($M+H$)⁺



¹H NMR (400 MHz, DMSO-d₆) δ ppm 2.41 (s, 3 H) 2.45 - 2.48 (m, 1 H) 2.64 - 2.74 (m, 1 H) 3.63 - 3.75 (m, 1 H) 3.76 - 3.87 (m, 2 H) 3.94 (br d, $J=4.89$ Hz, 1 H) 4.08 (d, $J=11.74$ Hz, 1 H) 5.30 (d, $J=5.67$ Hz, 1 H) 7.83 (d, $J=8.22$ Hz, 1 H) 8.10 (dd, $J=8.31$, 1.66 Hz, 1 H) 8.38 (s, 1 H) 8.44 (s, 1 H) 8.82 (d, $J=1.96$ Hz, 1 H) 8.85 (d, $J=1.76$ Hz, 1 H) 9.20 (dd, $J=1.96$, 0.98 Hz, 1 H)

NOE: 8.44 (NH): 4.08; 3.94; 3.68; 2.48 5.32(OH): 8.58; 3.78, 2.68 3.94 (CH): 8.44; 2.48



The following Examples were prepared in analogy to Example 50 starting from the corresponding acid and amino alcohol Intermediates:

20


Starting Acid	Starting Amino-alcohol	Ex.	Structure	MS m/z [M+H] ⁺	R _t (min)	Method

12	36	51 TRANS- Racemic mixture		462	3.87	METHO D 1
5	36	52 TRANS- Racemic mixture		448	3.58	METHO D 1

Relative stereochemistry assigned by NMR:


Example		Relative stereochemistry	¹ H-NMR
51		TRANS Racemic mixture	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 0.86 - 1.07 (m, 4 H) 2.03 - 2.17 (m, 1 H) 2.47 (s, 3 H) 2.49- 2.5 (m, 1H) 2.68 (td, J=12.57, 4.40 Hz, 1 H) 3.64 - 3.72 (m, 1 H) 3.77 - 3.88 (m, 3 H) 4.10 (d, J=12.13 Hz, 1 H) 5.28 (d, J=5.67 Hz, 1 H) 7.83 (d, J=8.22 Hz, 1 H) 8.08 (d, J=7.73 Hz, 1 H) 8.65 (s, 1 H) 8.71 (d, J=1.96 Hz, 1 H) 8.83 (s, 1 H) 9.03 (d, J=1.96 Hz, 1 H)
52		TRANS Racemic mixture	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 0.91 - 1.09 (m, 4 H) 2.08 - 2.15 (m, 1 H) 2.46 (s, 1 H) 2.59 - 2.78 (m, 1 H) 3.64 - 3.75 (m, 1 H) 3.76 - 3.86 (m, 2 H) 3.92 (br d, J=4.21 Hz, 1 H) 4.08 (d, J=11.93

			Hz, 1 H) 5.29 (d, J=5.77 Hz, 1 H) 7.83 (d, J=8.31 Hz, 1 H) 8.10 (dd, J=8.12, 2.15 Hz, 1 H) 8.37 (s, 1 H) 8.43 (s, 1 H) 8.77 (d, J=2.15 Hz, 1 H) 8.84 (d, J=1.96 Hz, 1 H) 9.14 (d, J=2.15 Hz, 1 H)
--	--	--	--

Example 53:

- 5 Example 53 was synthesized in analogy to Example 1, starting from acid Intermediate 5 (98 mg, 0.48 mmol) and amino alcohol Intermediate 38 (198 mg, 0.48 mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 80% of acetonitrile), 7 mg of the title compound, as CIS-racemic mixture.
- 10 LC-MS (METHOD 1): $R_t = 4.45$ min
 MS (ESI pos): $m/z = 479$ (M+H)⁺
¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.94 - 1.07 (m, 4 H) 1.12 (s, 3 H) 2.12 (tt, J=8.38, 5.20 Hz, 1 H) 2.72 (br t, J=11.98 Hz, 1 H) 2.98 (br d, J=13.69 Hz, 1 H) 3.47 (d, J=11.25 Hz, 1 H) 3.56 (br t, J=11.74 Hz, 1 H) 3.72 - 3.76 (m, 1 H) 3.76 - 3.87 (m, 1 H) 5.38 (s, 1 H) 7.46 - 7.56 (m, 2 H) 7.70 (t, J=8.07 Hz, 1 H) 8.35 (s, 1 H) 8.72 (d, J=2.20 Hz, 1 H) 8.88 - 8.95 (m, 1 H) 9.12 (d, J=2.20 Hz, 1 H)
- 15 NOE: 8.92 (NH): 5.38; 3,80 5.38(OH): 8.92; 3,80

Example 54:



Further elution from the column in the preparation of Example 53 gave 29 mg of the title compound, as racemic mixture (TRANS/CIS diastereoisomeric ratio 92/8

5 determined by NMR).


LC-MS (METHOD 1): $R_t = 4.45$ min

MS (ESI pos): $m/z = 479$ ($M+H$)⁺

1H NMR (500 MHz, DMSO-d₆) δ ppm 0.95 - 1.07 (m, 4 H) 1.10 (d, 3 H) 2.12 (tt, J=8.44, 5.14 Hz, 1 H) 2.51 - 2.56 (m, 1 H) 3.03 (td, J=12.65, 4.28 Hz, 1 H) 3.53 - 3.65 (m, 2 H) 3.74 - 3.76 (m, 1 H) 3.88 (d, J=12.23 Hz, 1 H) 5.10 (s, 1 H) 7.44 - 7.48 (dd, 1 H) 7.53 (d, J=8.27 Hz, 1 H) 7.66 (t, J=8.05 Hz, 1 H) 8.38 (s, 1 H) 8.48 (s, 1 H) 8.79 (d, J=2.20 Hz, 1 H) 9.16 (d, J=2.20 Hz, 1 H)

NOE: 8.48 (NH): 1.09; 3.88; 2.53 5.10(OH): 3.04, 3.57 1.09 (Me):
15 8.48; 3.57; 2.53; 3.88

Example 55:



20 Example 55 was synthesized in analogy to Example 1, starting from acid Intermediate 2 (304 mg, 1.68 mmol) and amino alcohol Intermediate 38 (700 mg,

1.68 mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 100% of acetonitrile), 300 mg of the title compound, as racemic mixture (TRANS/CIS diastereoisomeric ratio 96/4 determined by NMR).

LC-MS (METHOD 1): $R_t = 4.02$ min

5 MS (ESI pos): $m/z = 453$ (M+H)⁺

¹H NMR (500 MHz, DMSO-d₆) δ ppm 1.10 (d, J=1.00 Hz, 3 H) 2.41 (d, J=0.86 Hz, 3 H) 2.52 - 2.58 (m, 1 H) 3.03 (br d, J=4.16 Hz, 1 H) 3.46 - 3.66 (m, 2 H) 3.72 - 3.82 (m, 1 H) 3.88 (d, J=12.23 Hz, 1 H) 5.10 (s, 1 H) 7.46 (br d, J=12.72 Hz, 1 H) 7.50 - 7.57 (m, 1 H) 7.67 (t, J=8.23 Hz, 1 H) 8.35 - 8.39 (m, 1 H) 8.50 (s, 1 H) 8.83 (d, J=1.96 Hz, 1 H) 9.18 - 9.24 (m, 1 H)


10 NOE: 8.50 (NH): 1.10; 3.89; 3.64

5.10(OH): 3.03, 3.58

1.10 (Me):

8.50; 3.56; 3.89

Example 56:



15 Example 56 was synthesized in analogy to Example 1, starting from acid


Intermediate 2 (36 mg, 0.21 mmol) and amino alcohol Intermediate 39 (55 mg, 0.2 mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 65% of acetonitrile), 52 mg of the title compound, as TRANS-racemic mixture.

20 LC-MS (METHOD 1): $R_t = 3.57$ min

MS (ESI pos): $m/z = 440$ (M+H)⁺

¹H NMR (400 MHz, DMSO-d₆) δ ppm 2.40 (s, 3 H) 2.42 - 2.48 (m, 1 H) 2.89 (td, J=13.35, 4.40 Hz, 1 H) 3.65 (br t, J=11.30 Hz, 1 H) 3.79 - 3.89 (m, 2 H) 4.01 (d,

25 J=12.42 Hz, 1 H) 4.09 (s, 1 H) 5.35 (br s, 1 H) 8.11 (d, J=11.84 Hz, 1 H) 8.35 (s, 1 H) 8.46 (s, 1 H) 8.81 (d, J=1.76 Hz, 2 H) 9.20 (s, 1 H)


Example 57:

Example 57 was synthesized in analogy to Example 1, starting from acid Intermediate 5 (42 mg, 0.21 mmol) and amino alcohol Intermediate 39 (55 mg, 0.2 mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 70% of acetonitrile), 48 mg of the title compound, as TRANS-racemic mixture.

LC-MS (METHOD 1): $R_t = 4.00$ min

MS (ESI pos): $m/z = 466$ (M+H)⁺

¹H NMR (400 MHz, DMSO-d₆) δ ppm 0.87 - 1.11 (m, 4 H) 2.07 - 2.15 (m, 1 H) 2.47 (m, 1 H) 2.89 (td, J=13.11, 4.11 Hz, 1 H) 3.64 (br t, J=11.54 Hz, 1 H) 3.78 - 3.89 (m, 2 H) 4.01 (d, J=12.52 Hz, 1 H) 4.09 (br s, 1 H) 5.35 (br s, 1 H) 8.11 (d, J=11.93 Hz, 1 H) 8.34 (s, 1 H) 8.44 (s, 1 H) 8.77 (s, 1 H) 8.81 (s, 1 H) 9.14 (d, J=1.96 Hz, 1 H)

Example 58:

Example 58 was synthesized in analogy to Example 1, starting from acid Intermediate 2 (187 mg, 1.06 mmol) and amino alcohol Intermediate 41 (378 mg, 0.96 mmol) to give, after flash chromatographic purification (eluent:


dichloromethane/MeOH; gradient from 0% to 60% of MeOH), 22 mg of the title compound, as racemic mixture (TRANS/CIS diastereoisomeric ratio 91/9)

LC-MS (METHOD 10): $R_t = 3.16$ min

MS (ESI pos): $m/z = 439$ (M+H)⁺

- 5 ¹H NMR (500 MHz, DMSO-d₆) δ ppm 1.56 (s, 3 H) 2.40 (d, $J=0.73$ Hz, 4 H) 3.78 (d, $J=9.05$ Hz, 1 H) 3.90 (d, $J=9.05$ Hz, 1 H) 4.39 (d, $J=8.80$ Hz, 1 H) 4.90 (d, $J=8.80$ Hz, 1 H) 5.31 (s, 1 H) 7.48 - 7.56 (m, 2 H) 7.71 (t, $J=8.07$ Hz, 1 H) 8.37 (s, 1 H) 8.67 (s, 1 H) 8.82 (d, $J=1.96$ Hz, 1 H) 9.09 - 9.26 (m, 1 H) 9.21 (dd, $J=2.08, 1.10$ Hz, 1 H)
- NOE: 8.67 (NH): 1.56; 3.90; 4.38 5.31(OH): 3.78, 4.90 1.56 (Me):
- 10 8.67

Example 59:



Example 59 was synthesized in analogy to Example 1, starting from acid

- 15 Intermediate 2 (35.6 mg, 0.20 mmol) and amino alcohol Intermediate 42 (50 mg, 0.19 mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 65% of acetonitrile), 42 mg of the title compound, as TRANS-racemic mixture.


LC-MS (METHOD 1): $R_t = 3.77$ min

- 20 MS (ESI pos): $m/z = 421$ (M+H)⁺

¹H NMR (500 MHz, DMSO-d₆) δ ppm 2.40 (d, $J=0.98$ Hz, 3 H) 2.51 - 2.54 (m, 1 H) 2.62 - 2.70 (m, 1 H) 3.67 (td, $J=11.68, 1.83$ Hz, 1 H) 3.75 - 3.82 (m, 2 H) 3.82 - 3.89 (m, 1 H) 4.06 (d, $J=11.49$ Hz, 1 H) 5.14 (br d, $J=4.16$ Hz, 1 H) 7.62 - 7.70 (m, 4 H) 8.35 (s, 1 H) 8.37 (s, 1 H) 8.80 (d, $J=1.96$ Hz, 1 H) 9.19 (dd, $J=2.08, 1.10$ Hz, 1 H)

25

Example 60:




Example 60 was synthesized in analogy to Example 1, starting from acid Intermediate 5 (40.8 mg, 0.2 mmol) and amino alcohol Intermediate 42 (50 mg, 0.19mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 70% of acetonitrile), 39 mg of the title compound, as TRANS-racemic mixture.

LC-MS (METHOD 1): $R_t = 4.07$ min

MS (ESI pos): $m/z = 424$ (M+H)⁺

¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.86 - 1.06 (m, 4 H) 2.07 - 2.14 (m, 1 H) 2.51 - 2.53 (m, 1 H) 2.62 - 2.70 (m, 1 H) 3.66 (td, J=11.74, 1.71 Hz, 1 H) 3.75 - 3.87 (m, 3 H) 4.07 (d, J=11.25 Hz, 1 H) 5.14 (d, J=5.62 Hz, 1 H) 7.62 - 7.70 (m, 4 H) 8.35 (s, 1 H) 8.36 (s, 1 H) 8.76 (d, J=2.20 Hz, 1 H) 9.13 (d, J=2.20 Hz, 1 H)

Example 61:



Example 61 was synthesized in analogy to Example 1, starting from acid Intermediate 12 (82 mg, 0.29 mmol) and amino alcohol Intermediate 39 (66.7 mg, 0.31mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 70% of acetonitrile), 110 mg of the title compound, as TRANS-racemic mixture.


LC-MS (METHOD 1): $R_t = 4.22$ min

MS (ESI pos): $m/z = 480$ ($M+H$)⁺

¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.90 - 1.06 (m, 4 H) 2.09 (tt, $J=8.44$, 5.14 Hz, 1 H) 2.45 (s, 3 H) 2.47 (m, 1 H) 2.88 (td, $J=13.27$, 4.28 Hz, 1 H) 3.59 - 3.67 (m, 1 H)
5 3.77 - 3.88 (m, 2 H) 4.01 - 4.05 (m, 1 H) 4.08 (br d, $J=5.14$ Hz, 1 H) 5.32 (d, $J=5.62$ Hz, 1 H) 8.11 (dd, $J=11.86$, 1.59 Hz, 1 H) 8.69 (s, 1 H) 8.71 (d, $J=2.36$ Hz, 1 H) 8.81 (s, 1 H) 9.03 (d, $J=2.20$ Hz, 1 H)

NOE: 8.69 (NH): 4.08; 2.47 3.64 5.32(OH): 3.82, 2.88 4.08 (CH): 8.69; 2.47

10 **Example 62:**



Example 62 was synthesized in analogy to Example 1, starting from acid

Intermediate 12 (103 mg, 0.6 mmol) and amino alcohol Intermediate 43 (250 mg,


15 0.6mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 70% of acetonitrile), 115 mg of the title compound, as TRANS-racemic mixture.

LC-MS (METHOD 1): $R_t = 4.40$ min

MS (ESI pos): $m/z = 494$ ($M+H$)⁺

20 ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.92 - 1.06 (m, 4 H) 1.21 (d, $J=3.18$ Hz, 3 H) 2.10 (tt, $J=8.47$, 5.23 Hz, 1 H) 2.30 - 2.36 (m, 1 H) 2.47 - 2.49 (m, 3 H) 3.22 - 3.29 (m, 1 H) 3.49 - 3.54 (m, 1 H) 3.62 (br t, $J=10.88$ Hz, 1 H) 3.76 - 3.88 (m, 2 H) 4.96 (s, 1 H) 8.08 (dd, $J=11.98$, 1.47 Hz, 1 H) 8.72 (d, $J=1.96$ Hz, 1 H) 8.78 (s, 1 H) 8.84 (s, 1 H) 9.05 (d, $J=2.20$ Hz, 1 H)

25 NOE: 8.78 (NH): 1.21; 2.34; 3.62 4.96(OH): 3.51, 3.35 1.21 (Me): 8.78; 2.34

Example 63:

Example 63 was synthesized in analogy to Example 1, starting from acid

- 5 Intermediate 5 (45 mg, 0.22 mmol) and amino alcohol Intermediate 40 (55mg, 0.21mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 70% of acetonitrile), 55 mg of the title compound, as TRANS-racemic mixture.


LC-MS (METHOD 1): $R_t = 3.82$ min

- 10 MS (ESI pos): $m/z = 448$ ($M+H$)⁺

¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.91 - 1.08 (m, 4 H) 2.06 - 2.17 (m, 1 H) 2.41 - 2.48 (m, 1 H) 2.76 - 2.84 (m, 1 H) 3.65 - 3.76 (m, 2 H) 3.92 (dt, $J=11.23, 3.58$ Hz, 1 H) 3.95 - 4.00 (m, 1 H) 4.04 (br s, 1 H) 5.24 (d, $J=4.03$ Hz, 1 H) 7.75 (d, $J=8.44$ Hz, 1 H) 8.11 (dd, $J=8.50, 2.02$ Hz, 1 H) 8.33 (s, 1 H) 8.43 (s, 1 H) 8.76 (d, $J=2.08$ Hz, 1 H)

- 15 8.88 (dd, $J=1.53, 0.79$ Hz, 1 H) 9.13 (d, $J=2.20$ Hz, 1 H)

NOE 8.43 (NH): 4.04; 2.44; 3.98 5.24(OH): 3.66, 2.80 4.04 (CH): 8.43; 2.44

Example 64:

- 20 Example 64 was synthesized in analogy to Example 1, starting from acid Intermediate 2 (39mg, 0.22 mmol) and amino alcohol Intermediate 40 (55mg,

0.21mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 65% of acetonitrile), 31 mg of the title compound, as TRANS-racemic mixture.

LC-MS (METHOD 10): R_t = 2.66 min


5 MS (ESI pos): m/z = 422(M+H)⁺

¹H NMR (500 MHz, DMSO-d₆) δ ppm 2.40 (s, 3 H) 2.42 (br s, 1 H) 2.76 - 2.85 (m, 1 H) 3.64 - 3.77 (m, 2 H) 3.90 - 3.95 (m, 1 H) 3.97 (br d, J=12.23 Hz, 1 H) 4.05 (br s, 1 H) 5.24 (d, J=5.56 Hz, 1 H) 7.75 (d, J=8.44 Hz, 1 H) 8.11 (dd, J=8.47, 2.11 Hz, 1 H)
10 8.33 (s, 1 H) 8.45 (s, 1 H) 8.80 (d, J=1.90 Hz, 1 H) 8.88 (s, 1 H) 9.19 (s, 1 H)

NOE: 8.45 (NH): 4.05; 2.44; 3.98; 3.68 5.24(OH): 3.72; 3.97; 2.80

4.05 (CH): 8.45; 2.44

15 **Example 65:**



Example 65 was synthesized in analogy to Example 1, starting from acid Intermediate 12 (78mg, 0.36 mmol) and amino alcohol Intermediate 40 (90mg, 0.34mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 10% to 70% of acetonitrile), 110mg of the title compound, as TRANS-racemic mixture.


LC-MS (METHOD 1): R_t = 4.07 min

MS (ESI pos): m/z = 462(M+H)⁺

¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.90 - 1.07 (m, 4 H) 2.09 (tt, J=8.59, 4.98 Hz, 1 H) 2.42 - 2.47 (m, 4 H) 2.79 (ddd, J=13.66, 11.34, 4.34 Hz, 1 H) 3.64 - 3.76 (m, 2 H) 3.91 (dt, J=11.28, 3.59 Hz, 1 H) 3.99 (d, J=12.35 Hz, 1 H) 3.98 - 3.99 (m, 1 H) 4.02 (br s, 1 H) 5.21 (d, J=4.52 Hz, 1 H) 7.73 (d, J=8.44 Hz, 1 H) 8.12 (dd, J=8.50,
25

2.14 Hz, 1 H) 8.69 (d, $J=10.11$ Hz, 2 H) 8.88 (dd, $J=1.47, 0.73$ Hz, 1 H) 9.02 (d, $J=1.83$ Hz, 1 H)

Example 66:




5

Example 66 was synthesized in analogy to Example 1, starting from acid Intermediate 5 (16.8mg, 0.08mmol) and amino alcohol Intermediate 44 (20mg, 0.08mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 0% to 100% of acetonitrile), 11mg of the title compound.

10 LC-MS (METHOD 1): $R_t = 4.78$ min

MS (ESI pos): $m/z = 463$ ($M+H$)⁺

Example 67:



15 Example 67 was synthesized in analogy to Example 1, starting from acid Intermediate 2 (19.7mg, 0.11mmol) and amino alcohol Intermediate 44 (35mg, 0.11mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 0% to 100% of acetonitrile), 24mg of the title compound.



LC-MS (METHOD 1): $R_t = 4.40$ min



MS (ESI pos): $m/z = 437$ ($M+H$)⁺

Examples 68, 69, 70, 71:


5



Examples 68, 69, 70 and 71 were synthesized in analogy to Example 1, starting from acid Intermediate 5 (150 mg, 0,74 mmol) and amino alcohol Intermediate 37 (230 mg, 0.71 mmol) to give, after flash chromatographic purification (eluent: water/acetonitrile; gradient from 0% to 80% of acetonitrile), 143 mg of mixture of the
10 title compounds, which were obtained as single stereoisomers by chiral HPLC separation.


Ex. #	Structure	MS m/z [$M+H$] ⁺	R_t (min) [LC-MS Method]	R_t (min) [Chiral HPLC Method]
68 CIS single stereoisomer a		451	4.28 METHOD 1	9.66 [C3]
69 CIS single stereoisomer b		451	4.28 METHOD 1	11.30 [C3]

70 TRANS single stereoisomer a		451	4.21 METHOD 1	16.32, [C3]
71 TRANS single stereoisomer b		451	4.23 METHOD 1	22.71 [C3]

Relative stereochemistry assigned by NMR:

Example	Relative stereochemistry	¹ H-NMR	NOE
68 	CIS single stereoisomer a	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 0.94 - 1.06 (m, 4 H) 2.07 - 2.14 (m, 1 H) 3.58 (dd, J=9.29, 5.14 Hz, 1 H) 4.02 (dd, J=9.29, 5.87 Hz, 1 H) 4.14 (d, J=9.78 Hz, 1 H) 4.40 (br s, 1 H) 4.80 (d, J=10.03 Hz, 1 H) 6.34 (br s, 1 H) 7.54 - 7.63 (m, 2 H) 7.63 - 7.68 (m, 1 H) 8.39 (s, 1 H)	9.11 (NH): 6.34; 4.14; 3.58 6.34(OH): 9.11, 4.14 ; 3.58 4.40 (CH): 4.80





			8.72 (d, J=2.20 Hz, 1 H) 9.09 - 9.13 (m, 2 H)	
69		CIS single stereoisomer b	1H NMR (500 MHz, DMSO-d₆) δ ppm 0.94 - 1.06 (m, 4 H) 2.07 - 2.14 (m, 1 H) 3.58 (dd, J=9.29, 5.14 Hz, 1 H) 4.02 (dd, J=9.29, 5.87 Hz, 1 H) 4.14 (d, J=9.78 Hz, 1 H) 4.40 (br s, 1 H) 4.80 (d, J=10.03 Hz, 1 H) 6.34 (br s, 1 H) 7.54 - 7.63 (m, 2 H) 7.63 - 7.68 (m, 1 H) 8.39 (s, 1 H) 8.72 (d, J=2.20 Hz, 1 H) 9.09 - 9.13 (m, 2 H)	9.11 (NH): 6.34; 4.14; 3.58 6.34(OH): 9.11, 4.14 ; 3.58 4.40 (CH): 4.80
70		TRANS single stereoisomer a	1H NMR (500 MHz, DMSO-d₆) δ ppm 0.92 - 1.07 (m, 4 H) 3.71 - 3.80 (m, 1 H) 4.25 (dd, J=9.78, 4.40 Hz, 1 H) 4.35 (d, J=8.80 Hz, 1 H) 4.69 (d, J=8.80 Hz, 1 H) 4.76 (br s, 1 H) 5.64 (br s, 1 H)	8.28 (NH): 4.69; 4.76; 4.25 5.64(OH): 4.35; 3.76 4.76 (CH): 8.28

			7.51 - 7.56 (m, 2 H) 7.73 (t, J=7.82 Hz, 1 H) 8.28 (s, 1 H) 8.36 (s, 1 H) 8.74 (d, J=1.96 Hz, 1 H) 9.09 - 9.11 (m, 1 H)	
71		TRANS single stereoisomer b	1H NMR (500 MHz, DMSO-d6) δ ppm 0.92 - 1.07 (m, 4 H) 3.71 - 3.80 (m, 1 H) 4.25 (dd, J=9.78, 4.40 Hz, 1 H) 4.35 (d, J=8.80 Hz, 1 H) 4.69 (d, J=8.80 Hz, 1 H) 4.76 (br s, 1 H) 5.64 (br s, 1 H) 7.51 - 7.56 (m, 2 H) 7.73 (t, J=7.82 Hz, 1 H) 8.28 (s, 1 H) 8.36 (s, 1 H) 8.74 (d, J=1.96 Hz, 1 H) 9.09 - 9.11 (m, 1 H)	8.28 (NH): 4.69; 4.76; 4.25 5.64(OH): 4.35; 3.76 4.76 (CH): 8.28





Examples 72, 73, 74, 75






Examples 72, 73, 74 and 75 were synthesized in analogy to Example 1, starting from
 5 acid Intermediate 2 (92 mg, 0.52 mmol) and amino alcohol Intermediate 37 (160 mg, 43% content, 0.26 mmol) to give, after two subsequent flash chromatographic purifications (eluent: water/acetonitrile; gradient from 0% to 80% of acetonitrile; eluent: DCM/isopropyl alcohol; gradient from 0% to 30% of isopropyl alcohol), 110





mg of mixture of the title compounds, which were obtained as single stereoisomers by chiral HPLC separation.






Ex. #	Structure	MS m/z [M+H] ⁺	R _t (min) [LC-MS Method]	R _t (min) [Chiral HPLC Method]
72 CIS single stereoisomer a		425	3.84 METHOD 1	12.08 [C3]
73 CIS single stereoisomer b		<u>425</u>	<u>3.85</u> METHOD 1	<u>13.41</u> [C3]
74 TRANS single stereoisomer a		<u>425</u>	<u>3.75</u> METHOD 1	<u>22.54</u> [C3]
75 TRANS single stereoisomer b		<u>425</u>	<u>3.78</u> METHOD 1	<u>26.59</u> [C3]






The following examples were obtained as single stereoisomers by chiral HPLC separation of the corresponding racemic mixture:





Starting Racemic Mixture Ex. #	Ex. #	Structure	MS m/z [M+H] ⁺	R _t (min) [LC-MS Method]	R _t (min) [Chiral HPLC Method]
40	80 a Single stereoisomer a		450	3.89 METHOD 10	5.98 [C1]
40	80b single stereoisomer b		450	3.89 METHOD 10	6.76 [C1]
46 TRANS- Racemic mixture	81a TRANS- single stereoisomer a		439	3.67 METHOD 1	11.19 [C4]
46 TRANS- Racemic mixture	81b TRANS- single stereoisomer b		439	3.67 METHOD 1	13.99 [C4]


48 TRANS- Racemic mixture	82a TRANS- single stereoisomer a		465	3.41 METHOD 10	11.78 [C4]
48 TRANS- Racemic mixture	82b TRANS- single stereoisomer b		465	3.41 METHOD 10	14.22 [C4]
55 TRANS /CIS 96/4 Racemic mixture	83a TRANS- single stereoisomer a		453	METHOD 1	11.71 [C3]
55 TRANS /CIS 96/4 Racemic mixture	83b TRANS- single stereoisomer b		453	4.07 METHOD 1	26.90 [C3]
55 TRANS /CIS 96/4 Racemic mixture	84a CIS- single stereoisomer a		453	4.07 METHOD 1	13.67 [C3]

55 TRANS /CIS 96/4 Racemic mixture	84b CIS- single stereoiso mer b		453	4.07 METHOD 1	18.13 [C3]
54 TRANS/ CIS 92/8 Racemic mixture	85a CIS- single stereoiso mer a		479	4.48 METHOD 1	5.46 [C1]
54 TRANS/ CIS 92/8 Racemic mixture	85b CIS- single stereoiso mer b		479	4.48 METHOD 1	6.73 [C1]
54 TRANS/ CIS 92/8 Racemic mixture	86a TRANS- single stereoiso mer a		479	4.48 METHOD 1	6.08 [C1]


54 TRANS/ CIS 92/8 Racemic mixture	86b TRANS- single stereoiso mer b		479	4.48 METHOD 1	10.75 [C1]
50 TRANS- Racemic mixture	87a TRANS- single stereoiso mer a		422	3.08 METHOD 1	9.97 [C6]
50 TRANS- Racemic mixture	87b TRANS- single stereoiso mer b		422	3.08 METHOD 1	12.98 [C6]
51 TRANS- Racemic mixture	88a TRANS- single stereoiso mer a		462	3.87 METHOD 1	7.71 [C1]
51 TRANS- Racemic mixture	88b TRANS- single stereoiso mer b		462	3.87 METHOD 1	9.49 [C1]



52 TRANS- Racemic mixture	89a TRANS- single stereoisomer a		448	3.60 METHOD 1	11.82 [C6]
52 TRANS- Racemic mixture	89b TRANS- single stereoisomer b		448	3.60 METHOD 1	16 [C6]
61 TRANS- Racemic mixture	90a TRANS single stereoisomer a		480	4.22 METHOD 1	4.9 [C7]
61 TRANS- Racemic mixture	90b TRANS single stereoisomer b		480	4.22 METHOD 1	5.54 [C7]
62 TRANS- Racemic mixture	91a TRANS single stereoisomer a		494	4.4 METHOD 1	6.68 [C1]


62 TRANS- Racemic mixture	91b TRANS single stereoiso mer b		494	4.4 METHOD 1	12.94 [C1]
58 TRANS/ CIS 91/9 Racemic mixture	92a CIS Single stereoiso mer a		439	3.16 METHOD 10	4.76 [C1]
58 TRANS/ CIS 91/9 Racemic mixture	92b CIS Single stereoiso mer b		439	3.16 METHOD 10	8.24 [C1]
58 TRANS/ CIS 91/9 Racemic mixture	93a TRANS Single stereoiso mer a		439	3.16 METHOD 10	7.20 [C1]



58 TRANS/ CIS 91/9 Racemic mixture	93b TRANS Single stereoisomer b		439	3.16 METHOD 10	16.60 [C1]
--	--	---	-----	----------------------	---------------


Relative stereochemistry assigned by NMR:


Example		Relative stereochemistry	¹ H-NMR	NOE
81a		TRANS-single stereoisomer a	¹ H NMR (500 MHz, DMSO-d ₆) δ ppm 2.40 (s, 3 H) 2.60 - 2.71 (m, 2 H) 3.63 - 3.69 (m, 1 H) 3.82 (br d, J=11.98 Hz, 2 H) 4.03 (d, J=11.74 Hz, 1 H) 4.08 - 4.11 (m, 1 H) 5.32 (d, J=5.38 Hz, 1 H) 7.46 - 7.56 (m, 2 H) 7.73 (t, J=7.51 Hz, 1 H) 8.34 (s, 1 H) 8.42 (s, 1 H) 8.81 (d, J=2.20 Hz, 1 H) 9.17 - 9.19 (m, 1 H)	8.42 (NH): 4.08; 4.03; 3.65; 3.82 5.32 (OH): 4.10; 3.70.



82a		TRANS-single stereoisomer a	1H NMR (500 MHz, DMSO-d₆) δ ppm 0.93 - 1.06 (m, 4 H) 2.10 (tt, J=8.38, 5.20 Hz, 1 H) 2.59 - 2.71 (m, 2 H) 3.66 (td, J=11.68, 2.08 Hz, 1 H) 3.81 (br d, J=11.25 Hz, 2 H) 4.03 (d, J=11.86 Hz, 1 H) 4.09 (br d, J=4.65 Hz, 1 H) 5.32 (d, J=5.38 Hz, 1 H) 7.48 (d, J=12.47 Hz, 1 H) 7.54 (dd, J=8.31, 1.22 Hz, 1 H) 7.73 (t, J=8.19 Hz, 1 H) 8.33 (s, 1 H) 8.40 (s, 1 H) 8.77 (d, J=2.20 Hz, 1 H) 9.13 (s, 1 H)	8.40 (NH): 4.09 5.32(OH): 4.09
83a		TRANS-single stereoisomer a	1H NMR (500 MHz, DMSO-d₆) δ ppm 1.10 (d, J=2.57 Hz, 3 H) 2.41 (d, J=0.73 Hz, 3 H) 2.52 - 2.57 (m, 1 H) 3.03 (br d, J=4.40 Hz, 1 H) 3.56 - 3.58 (m, 1 H) 3.60 - 3.65 (m,	8.50 (NH): 1.10; 3.88; 2.53 5.10(OH): 3.57, 3.03 1.10 (Me): 8.50; 3.88; 2.53


			1 H) 3.75 (br dd, J=11.49, 2.57 Hz, 1 H) 3.88 (d, J=12.23 Hz, 1 H) 5.10 (s, 1 H) 7.46 (dd, J=12.78, 1.41 Hz, 1 H) 7.53 (dd, J=8.31, 1.34 Hz, 1 H) 7.67 (t, J=8.25 Hz, 1 H) 8.39 (s, 1 H) 8.50 (s, 1 H) 8.83 (d, J=2.08 Hz, 1 H) 9.23 (dd, J=1.96, 1.10 Hz, 1 H)	
84a		CIS-single stereoisomer a	1H NMR (500 MHz, DMSO-d₆) δ ppm 1.12 (s, 2 H) 2.36 - 2.47 (m, 3 H) 2.67 - 2.84 (m, 1 H) 2.98 (br d, J=14.18 Hz, 1 H) 3.47 (d, J=11.00 Hz, 1 H) 3.56 (br t, J=11.98 Hz, 1 H) 3.75 (br dd, J=11.37, 3.55 Hz, 1 H) 3.81 (d, J=11.25 Hz, 1 H) 5.39 (s, 1 H) 7.46 - 7.55 (m, 2 H) 7.70 (t, J=8.07 Hz, 1 H) 8.35 (s, 1 H) 8.77 8.96 (NH): 5.39; 2.73 5.39(OH): 8.96; 2.73	


			(d, J=1.96 Hz, 1 H) 8.96 (s, 1 H) 9.18 (dd, J=2.08, 1.10 Hz, 1 H)	
84b		CIS-single stereoisomer b	1H NMR (500 MHz, DMSO-d ₆) δ ppm 1.12 (s, 2 H) 2.36 - 2.47 (m, 3 H) 2.67 - 2.84 (m, 1 H) 2.98 (br d, J=14.18 Hz, 1 H) 3.47 (d, J=11.00 Hz, 1 H) 3.56 (br t, J=11.98 Hz, 1 H) 3.75 (br dd, J=11.37, 3.55 Hz, 1 H) 3.81 (d, J=11.25 Hz, 1 H) 5.39 (s, 1 H) 7.46 - 7.55 (m, 2 H) 7.70 (t, J=8.07 Hz, 1 H) 8.35 (s, 1 H) 8.77 (d, J=1.96 Hz, 1 H) 8.96 (s, 1 H) 9.18 (dd, J=2.08, 1.10 Hz, 1 H)	8.96 (NH): 5.39; 2.73 5.39(OH): 8.96; 2.73
83b		TRANS-single stereoisomer b	1H NMR (500 MHz, DMSO-d ₆) δ ppm 1.10 (d, J=2.57 Hz, 3 H) 2.41 (d, J=0.73 Hz, 3 H) 2.52 - 2.57 (m, 1 H) 3.03 (br d, 	8.50 (NH): 1.10; 3.88; 2.53 5.10(OH): 3.57, 3.03 1.10 (Me): 8.50; 3.88;


			J=4.40 Hz, 1 H) 3.56 – 3.58 (m, 1 H) 3.60 - 3.65 (m, 1 H) 3.75 (br dd, J=11.49, 2.57 Hz, 1 H) 3.88 (d, J=12.23 Hz, 1 H) 5.10 (s, 1 H) 7.46 (dd, J=12.78, 1.41 Hz, 1 H) 7.53 (dd, J=8.31, 1.34 Hz, 1 H) 7.67 (t, J=8.25 Hz, 1 H) 8.39 (s, 1 H) 8.50 (s, 1 H) 8.83 (d, J=2.08 Hz, 1 H) 9.23 (dd, J=1.96, 1.10 Hz, 1 H)	2.53
85a		CIS-single stereoisomer a	¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.94 - 1.07 (m, 4 H) 1.12 (s, 3 H) 2.12 (tt, J=8.38, 5.20 Hz, 1 H) 2.72 (br t, J=11.98 Hz, 1 H) 2.98 (br d, J=13.69 Hz, 1 H) 3.47 (d, J=11.25 Hz, 1 H) 3.56 (br t, J=11.74 Hz, 1 H) 3.75 (br dd, J=11.37, 3.30 Hz, 1 H) 3.81 (d,	8.92 (NH): 5.38; 3.81; 3.56 5.38(OH): 8.92; 1.12 (Me): 3.47; 2.72

			J=11.25 Hz, 1 H) 5.38 (s, 1 H) 7.46 - 7.56 (m, 2 H) 7.70 (t, J=8.07 Hz, 1 H) 8.35 (s, 1 H) 8.72 (d, J=2.20 Hz, 1 H) 8.92 (s, 1 H) 9.12 (d, J=2.20 Hz, 1 H)	
86a		TRANS-single stereoisomer a	1H NMR (500 MHz, DMSO-d6) δ ppm 0.95 - 1.07 (m, 5 H) 1.10 (d, J=2.57 Hz, 2 H) 2.12 (tt, J=8.44, 5.14 Hz, 1 H) 2.51 - 2.56 (m, 1 H) 3.03 (td, J=12.65, 4.28 Hz, 1 H) 3.54 - 3.59 (m, 1 H) 3.62 (t, J=11.25 Hz, 1 H) 3.75 (br dd, J=11.25, 2.45 Hz, 1 H) 3.88 (d, J=12.23 Hz, 1 H) 5.10 (s, 1 H) 7.46 (br d, J=12.96 Hz, 1 H) 7.47 - 7.56 (m, 1 H) 7.59 - 7.71 (m, 1 H) 8.35 - 8.40 (m, 1 H) 8.48 (s, 1 H) 8.79 (d, J=2.20 Hz, 1 H) 9.16 (d, J=2.20 Hz,	8.48 (NH): 1.10; 3.88; 2.53 5.10(OH): 3.57; 3.03 1.10 (Me): 8.48; 3.88; 2.53

			1 H)	
85b		CIS-single stereoisomer b	1H NMR (500 MHz, DMSO-d₆) δ ppm 0.94 - 1.07 (m, 4 H) 1.12 (s, 3 H) 2.12 (tt, J=8.38, 5.20 Hz, 1 H) 2.72 (br t, J=11.98 Hz, 1 H) 2.98 (br d, J=13.69 Hz, 1 H) 3.47 (d, J=11.25 Hz, 1 H) 3.56 (br t, J=11.74 Hz, 1 H) 3.75 (br dd, J=11.37, 3.30 Hz, 1 H) 3.81 (d, J=11.25 Hz, 1 H) 5.38 (s, 1 H) 7.46 - 7.56 (m, 2 H) 7.70 (t, J=8.07 Hz, 1 H) 8.35 (s, 1 H) 8.72 (d, J=2.20 Hz, 1 H) 8.92 (s, 1 H) 9.12 (d, J=2.20 Hz, 1 H)	8.92 (NH): 5.38; 3.81; 3.56 5.38(OH): 8.92; 1.12 (Me): 3.47; 2.72
86b		TRANS-single stereoisomer b	1H NMR (500 MHz, DMSO-d₆) δ ppm 0.95 - 1.07 (m, 5 H) 1.10 (d, J=2.57 Hz, 2 H) 2.12 (tt, J=8.44, 5.14 Hz, 1 H) 2.51 - 2.56 (m, 1 H) 3.03 (td, J=12.65,	8.48 (NH): 1.10; 3.88; 2.53 5.10(OH): 3.57; 3.03 1.10 (Me): 8.48; 3.88; 2.53


			4.28 Hz, 1 H) 3.54 - 3.59 (m, 1 H) 3.62 (t, J=11.25 Hz, 1 H) 3.75 (br dd, J=11.25, 2.45 Hz, 1 H) 3.88 (d, J=12.23 Hz, 1 H) 5.10 (s, 1 H) 7.46 (br d, J=12.96 Hz, 1 H) 7.47 - 7.56 (m, 1 H) 7.59 - 7.71 (m, 1 H) 8.35 - 8.40 (m, 1 H) 8.48 (s, 1 H) 8.79 (d, J=2.20 Hz, 1 H) 9.16 (d, J=2.20 Hz, 1 H)	
87a		TRANS-single stereoisomer a	1H NMR (400 MHz, DMSO-d6) δ ppm 2.41 (s, 3 H) 2.47 (m, 1 H) 2.63 - 2.74 (m, 1 H) 3.63 - 3.75 (m, 1 H) 3.76 - 3.87 (m, 2 H) 3.94 (br d, J=4.89 Hz, 1 H) 4.08 (d, J=11.74 Hz, 1 H) 5.30 (d, J=5.67 Hz, 1 H) 7.83 (d, J=8.22 Hz, 1 H) 8.10 (dd, J=8.31, 1.66 Hz, 1 H) 8.38 (s, 1 H) 8.44 (NH): 4.08; 3.94; 2.47 5.30(OH): 3.81, 2.68 3.94 (CH): 8.44; 2.47	

			8.44 (s, 1 H) 8.82 (d, J=1.96 Hz, 1 H) 8.85 (d, J=1.76 Hz, 1 H) 9.20 (dd, J=1.96, 0.98 Hz, 1 H)	
88a		TRANS-single stereoisomer a	¹H NMR (500 MHz, DMSO-d₆) δ ppm 0.87 - 0.99 (m, 2 H) 1.00 - 1.06 (m, 2 H) 2.10 (tt, J=8.47, 5.10 Hz, 1 H) 2.46 (s, 3 H) 2.51 - 2.53 (m, 1 H) 2.64 - 2.71 (m, 1 H) 3.64 - 3.72 (m, 1 H) 3.77 - 3.88 (m, 3 H) 4.10 (d, J=11.49 Hz, 1 H) 5.29 (br s, 1 H) 7.84 (d, J=8.07 Hz, 1 H) 8.08 (dd, J=8.19, 2.08 Hz, 1 H) 8.65 (s, 1 H) 8.71 (d, J=2.20 Hz, 1 H) 8.83 (d, J=2.20 Hz, 1 H) 9.04 (d, J=1.71 Hz, 1 H)	-

89b		TRANS-single stereoisomer b	1H NMR (500 MHz, DMSO-d₆) δ ppm 0.92 - 1.07 (m, 4 H) 2.12 (tt, J=8.47, 5.10 Hz, 1 H) 2.44 - 2.48 (m, 1 H) 2.64 - 2.72 (m, 1 H) 3.65 - 3.73 (m, 1 H) 3.76 - 3.86 (m, 2 H) 3.92 (br d, J=5.87 Hz, 1 H) 4.09 (d, J=11.49 Hz, 1 H) 5.29 (d, J=6.11 Hz, 1 H) 7.83 (d, J=8.07 Hz, 1 H) 8.10 (dd, J=8.19, 2.08 Hz, 1 H) 8.37 (s, 1 H) 8.43 (s, 1 H) 8.77 (d, J=2.20 Hz, 1 H) 8.84 (d, J=2.20 Hz, 1 H) 9.14 (d, J=2.20 Hz, 1 H)	-
-----	--	-----------------------------	--	---


Claims

1. A compound of formula (I)



wherein

A is selected from the group A^a consisting of



wherein above mentioned groups are substituted with one R⁵ and one R⁶;

R¹ is selected from the group R^{1a} consisting of

halogen, C₁₋₃-alkyl- and C₃₋₆-cycloalkyl-

wherein the above mentioned C₁₋₃-alkyl-, and C₃₋₆-cycloalkyl- groups may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of halogen, NC- and HO-;

R² is selected from the group R^{2a} consisting of

aryl and heteroaryl,

wherein the above mentioned aryl and heteroaryl-groups may optionally be substituted with 1 to 5 substituents R⁴;

R³ is selected from the group R^{3a} consisting of

H- and C₁₋₃-alkyl-,

wherein the above mentioned C₁₋₃-alkyl-groups may optionally be substituted with 1 to 7 substituents independently from each other selected from the group consisting of halogen;

R⁴ is independently from each other selected from the group R^{4a} consisting of halogen, NC-, HO-, C₁₋₄-alkyl- and C₁₋₃-alkyl-O-

wherein the above mentioned C₁₋₄-alkyl- and C₁₋₃-alkyl-O-groups may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of HO- and F-;

R⁵ is selected from the group R^{5a} consisting of

H-, halogen, NC-, HO- and C₁₋₃-alkyl-,

wherein the above mentioned C₁₋₃-alkyl-group may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of HO- and F-

or R⁵ and R⁶ together form an group O=;

R⁶ is selected from the group R^{6a} consisting of

H-, halogen, NC-, HO- and C₁₋₃-alkyl-,

wherein the above mentioned C₁₋₃-alkyl-group may optionally be substituted with 1 to 5 substituents independently selected from the group consisting of HO- and F-


or R⁵ and R⁶ together form a group O=;

or a salt thereof.

5

2. A compound according to claim 1, wherein

A is selected from the group A^b consisting of




wherein above mentioned groups are substituted with with one R⁵

and one R⁶.

or a salt thereof.

3. A compound according to claim 1, wherein

A is selected from the group A^c consisting of




or a salt thereof.

5

4. A compound according to any one or more of the claims 1 to 3, wherein

A is selected from the group A^c consisting of



or a salt thereof.

5. A compound according to any one or more of the claims 1 to 4, wherein

R¹ is selected from the group R^{1b} consisting of

F-, Cl-, C₁₋₃-alkyl- and C₃₋₆-cycloalkyl-,

wherein the above mentioned C₁₋₃-alkyl- and C₃₋₆-cycloalkyl-groups may optionally be substituted with 1 to 3 substituents independently selected from the group consisting of F-,

10 or a salt thereof.

6. A compound according to any one or more of the claims 1 to 4, wherein

R¹ is selected from the group R^{1c} consisting of

F-, H₃C- and cyclopropyl-,

or a salt thereof.

15 7. A compound according to any one or more of the claims 1 to 6, wherein

R^2 is selected from the group R^{2b} consisting of
quinolinyl, phenyl and pyridynyl,
wherein the above mentioned quinoline, phenyl and pyridyl-groups
may optionally be substituted with 1 to 5 substituents R^4 ,
or a salt thereof.

8. A compound according to any one or more of the claims 1 to 6, wherein
 R^2 is selected from the group R^{2c} consisting of
phenyl and pyridyl,
wherein the above mentioned phenyl and pyridyl-groups may
optionally be substituted with 1 to 2 substituents R^4 ,
or a salt thereof.

5

9. A compound according to any one or more of the claims 1 to 8, wherein
 R^3 is selected from the group R^{3b} consisting of
 $H-$, H_3C- , F_3C- , F_2HC- , FH_2C- and F_3C- ,
or a salt thereof.

10. A compound according to any one or more of the claims 1 to 9, wherein
 R^4 is independently from each other selected from the group R^{4b} consisting of
halogen, C_{1-4} -alkyl- and C_{1-3} -alkyl-O-
wherein the above mentioned C_{1-4} -alkyl- and C_{1-3} -alkyl-O-groups may
optionally be substituted with 1 to 5 substituents independently selected
from the group consisting of $HO-$, and $F-$,

10 or a salt thereof.

11. A compound according to any one or more of the claims 1 to 10, wherein
 R^5 is selected from the group R^{5b} consisting of
 $H-$, $HO-$ and C_{1-2} -alkyl-,

wherein the above mentioned C_{1-2} -alkyl-group may optionally be
substituted with 1 to 5 $F-$,
or R^5 and R^6 together form an group $O=$,

or a salt thereof.

12. A compound according to any one or more of the claims 1 to 11, wherein

R^6 is selected from the group R^{6b} consisting of

H- and C_{1-2} -alkyl-,


wherein the above mentioned C_{1-2} -alkyl-group may optionally be


substituted with 1 to 5 F-,


or R^5/R^6 together form a group O=,


or a salt thereof.


5 13. A compound selected from the group consisting of



















147a

14. A compound of formula:




15. A compound of formula:




5

16. A compound of formula:




10 17. A compound of formula:




147b

18. A compound of formula:




19. A compound of formula:




5

20. A compound of formula:




147c

21. A compound of formula:




22. A compound of formula:




5


23. A compound of formula:



24. A compound of formula:




25. A compound of formula:




5

26. A compound of formula:




10 27. A compound of formula:




147e

28. A compound of formula:




29. A compound of formula:




5


30. A compound of formula:




31. A compound of formula:



5 32. A compound of formula:




33. A compound of formula:




147g

34. A compound of formula:




35. A compound of formula:




5

36. A compound of formula:




147h

37. A compound of formula:




38. A compound of formula:




5

39. A compound of formula:




10

40. A compound of formula:




41. A compound of formula:


5



42. A compound of formula:



43. A compound of formula:



44. A pharmaceutically acceptable salt of any of the compounds according to claims

5 13 to 43.

45. A compound according to any one or more of claims 1 to 43, or a pharmaceutically acceptable salt thereof for use as a medicament.

46. Pharmaceutical compositions containing at least one compound according to one or more of the claims 1 to 43 or a pharmaceutically acceptable salt thereof together with one or more pharmaceutically acceptable carrier.

47. A compound according to one or more of claims 1 to 43, or a pharmaceutically acceptable salt thereof or a pharmaceutical composition according to claim 46 for use in the prevention or treatment of (1) disorders comprising the symptom of cognitive deficiency; (2) organic, including symptomatic, mental disorders, dementia; (3) mental retardation; (4) mood [affective] disorders; (5) neurotic, stress-related and somatoform disorders including anxiety disorders; (6) behavioural and emotional disorders with onset usually occurring in childhood and adolescence, attention deficit hyperactivity syndrome (ADHD) and Autism spectrum disorders; (7) disorders of psychological development, developmental disorders of scholastic skills; (8) schizophrenia and other psychotic disorders; (9) disorders of adult personality and behaviour; (10) mental and behavioural disorders due to psychoactive substance use; (11) extrapyramidal and movement disorders; (12) episodic and paroxysmal disorders, epilepsy; (13) Systemic atrophies primarily affecting the central nervous system, ataxia; (14) Behavioural syndromes associated with physiological disturbances and physical factors; (15) sexual dysfunction comprising excessive sexual drive; (16) factitious disorders; (17) Treatment, amelioration and / or prevention of cognitive impairment being related to perception, concentration, cognition, learning or memory; (18) Treatment amelioration and / or prevention of cognitive impairment being related to age-associated learning and memory impairments; (19) Age-associated memory losses; (20) Vascular dementia; (21) Craniocerebral trauma; (22) Stroke; (23) Dementia occurring after strokes (post stroke dementia); (24) Post-traumatic dementia; (25) General concentration impairments; (26) Concentration impairments in children with learning and memory

problems; (27) Alzheimer's disease; (28) Lewy body dementia; (29) Dementia with degeneration of the frontal lobes; including Pick's syndrome; (30) Parkinson's disease; (31) Progressive nuclear palsy; (32) Dementia with corticobasal degeneration; (33) Amyotrophic lateral sclerosis (ALS); (34) Huntington's disease; (35) Multiple sclerosis; (36) Thalamic degeneration; (37) Creutzfeld-Jacob dementia; (38) HIV dementia; (38) Schizophrenia with dementia or Korsakoff's psychosis; (39) Sleep disorders; (40) Bipolar disorder; (41) Metabolic syndrome; (42) Obesity; (43) Diabetes mellitus; (44) Hyperglycemia; (45) Dyslipidemia; (46) Impaired glucose tolerance; (47) Disease of the testes, brain, small intestine, skeletal muscle, heart, lung, thymus or spleen; (48) pain disorders; (49) neuropsychiatric symptoms (e.g. depressive symptoms in Alzheimer's disease); (50) mixed dementia; (51) cognitive impairment in schizoaffective disorder; (52) cognitive impairment in bipolar disorder; (53) cognitive impairment in major depressive disorder and (54) cognitive impairment associated with schizophrenia.