





FIG. 1



Key Eligibility Criteria


Key Inclusion Criteria


- Men or women ≥50 years of age with treatment-naïve neovascular AMD (nAMD)
- Active subfoveal CNV, with a total area > 50% of the total lesion area in the study eye
- Presence of intraretinal and/or subretinal fluid in the central subfield on OCT
- BCVA of 78 - 24 letters (Snellen equivalent: 20/32 - 20/320) with decreased vision due to nAMD

Exclusion Criteria

- Causes of CNV other than nAMD in the study eye
- History or clinical evidence of diabetic retinopathy, diabetic macular edema, or any retinal vascular disease other than nAMD in either eye
- Scar, fibrosis, atrophy or RPE tears/rips involving the central subfield in the study eye
- IOP ≥25 mmHg in the study eye despite glaucoma treatment
- Intracocular inflammation/infection in either eye within 12 weeks of the screening visit

FIG. 2








FIG. 5




FIG. 6




FIG. 7





FIG. 8






FIG. 10




FIG. 11




FIG. 12




FIG. 13A




FIG. 13B





FIG. 14A






FIG. 15




FIG. 16




FIG. 17





FIG. 18






FIG. 20






FIG. 21

22
G
L







FIG. 24






FIG. 27




FIG. 28




FIG. 29A




FIG. 29B




FIG. 29C




FIG. 30A




FIG. 30B




FIG. 30C

38/53

Non-Ocular Serious TEAEs ≥0.5% through Week 60

	2q8	HDq12	HDq16	All HD
N (SAF)	336	335	338	673
Number of Patients ≥ 1 AE, n (%)	53 (15.8%)	41 (12.2%)	41 (12.1%)	82 (12.2%)
Angina pectoris	0	0	3 (0.9%)	3 (0.4%)
Cardiac failure congestive	0	2 (0.6%)	0	2 (0.3%)
Chest Pain	0	2 (0.6%)	1 (0.3%)	3 (0.4%)
Cellulitis	0	2 (0.6%)	0	2 (0.3%)
Pneumonia	1 (0.3%)	4 (1.2%)	2 (0.6%)	6 (0.9%)
Pyelonephritis acute	0	0	2 (0.6%)	2 (0.3%)
Urinary tract infection	4 (1.2%)	1 (0.3%)	0	1 (0.1%)
Pelvic fracture	2 (0.6%)	0	0	0
Rib fracture	2 (0.6%)	0	0	0
Upper limb fracture	2 (0.6%)	0	0	0
Hyponatraemia	2 (0.6%)	0	2 (0.6%)	2 (0.3%)
Osteoarthritis	1 (0.3%)	1 (0.3%)	3 (0.9%)	4 (0.6%)
Bladder neoplasm	3 (0.9%)	0	0	0
Cerebral infarction	2 (0.6%)	0	0	0
Cerebrovascular accident	2 (0.6%)	1 (0.3%)	0	1 (0.1%)
Hypertension	1 (0.3%)	2 (0.6%)	0	2 (0.3%)

FIG. 30D




FIG. 30E




FIG. 30F




FIG. 30G




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35





FIG. 36






FIG. 38




FIG. 39




FIG. 40





FIG. 41



Points and error bars are means and standard deviations of observed data
Lines and shaded area are medians and 90% prediction intervals of simulated data

FIG. 42