

Description

Title of Invention: MELK EPITOPE PEPTIDES AND VACCINES CONTAINING THE SAME

Technical Field

- [0001] The present application claims the benefit of U.S. Provisional Application No. 61/085,663, filed on August 1, 2008, the entire contents of which are incorporated by reference herein.
- [0002] The present invention relates to the field of biological science, more specifically to the field of cancer therapy. In particular, the present invention relates to novel peptides that are extremely effective as cancer vaccines, and drugs for treating and preventing tumors.

Background Art

- [0003] It has been demonstrated that CD8 positive CTLs recognize epitope peptides derived from the tumor-associated antigens (TAAs) found on major histocompatibility complex (MHC) class I molecules, and then kill the tumor cells. Since the discovery of the melanoma antigen (MAGE) family as the first example of TAAs, many other TAAs have been discovered, primarily through immunological approaches (Boon T, Int J Cancer 1993 May 8, 54(2): 177-80; Boon T & van der Bruggen P, J Exp Med 1996 Mar 1, 183(3): 725-9). Some of these TAAs are currently undergoing of clinical development as immunotherapeutic targets.
- [0004] Identification of new TAAs, capable of inducing potent and specific anti-tumor immune responses, warrants further development and clinical application of peptide vaccination strategies for various types of cancer (Harris CC, J Natl Cancer Inst 1996 Oct 16, 88(20): 1442-55; Butterfield LH et al., Cancer Res 1999 Jul 1, 59(13): 3134-42; Vissers JL et al., Cancer Res 1999 Nov 1, 59(21): 5554-9; van der Burg SH et al., J Immunol 1996 May 1, 156(9): 3308-14; Tanaka F et al., Cancer Res 1997 Oct 15, 57(20): 4465-8; Fujie T et al., Int J Cancer 1999 Jan 18, 80(2): 169-72; Kikuchi M et al., Int J Cancer 1999 May 5, 81(3): 459-66; Oiso M et al., Int J Cancer 1999 May 5, 81(3): 387-94). To date, there have been several reports of clinical trials using these tumor-associated antigen derived peptides. Unfortunately, only a low objective response rate has been observed in these cancer vaccine trials so far (Belli F et al., J Clin Oncol 2002 Oct 15, 20(20): 4169-80; Coulie PG et al., Immunol Rev 2002 Oct, 188: 33-42; Rosenberg SA et al., Nat Med 2004 Sep, 10(9): 909-15).
- [0005] Recently, algorithms for predicting HLA class I-binding peptide sequences have been developed (Journal of Immunological Methods, (1995), Vol.185, pp.181-190, J. Immunol., (1994), Vol.152, pp.163-175, protein science, (2000), Vol.9,

pp.1838-1846). However, it is difficult to estimate if a predicted epitope peptide can be processed naturally in the target cells and expressed on the target cell surface with HLA molecule. Moreover, the algorithms, for example BIMAS (http://bimas.dcrt.nih.gov/cgi-bin/molbio/ken_parker_comboform) (Parker KC, et al., (1994) *J Immunol.*;152(1):163-75.; Kuzushima K, et al., (2001) *Blood.*;98(6):1872-81.)) can suggest less than rigorous HLA molecule-binding peptides (Bachinsky MM, et. al., *Cancer Immun.* 2005 Mar 22;5:6.). Thus, identifying epitope peptides remains challenging and difficult.

[0006] MELK, maternal embryonic leucine zipper kinase, was previously identified as a new member of the snf1/AMPK serine-threonine kinase family that is involved in mammalian embryonic development (Heyer BS et al., *Dev Dyn.* 1999 Aug 215(4):344-51). The gene was shown to play an important role in stem cell renewal (Nakano I et al., *J Cell Biol.* 2005 Aug 1, 170(3):413-27), cell-cycle progression (Blot J et al., *Dev Biol.* 2002 Jan 15, 241(2):327-38; Seong HA et al., *Biochem J.* 2002 Feb 1, 361(Pt 3):597-604) and pre-mRNA splicing (Vulsteke V et al., *J Biol Chem.* 2004 Mar 5, 279(10):8642-7. Epub 2003 Dec 29). In addition, through gene expression profile analysis using a genome-wide cDNA microarray containing 23,040 genes, MELK was recently shown to be up-regulated in breast cancer (Lin ML et al., *Breast Cancer Res.* 2007; 9 (1):R17, WO2006/016525, WO2008/023841). In fact, MELK is up-regulated in several cancer cells, for example lung, bladder, lymphoma and cervical cancer cells (See WO2004/031413, WO2007/013665, and WO2006/085684, the disclosures of which are incorporated by reference herein). Northern blot analysis on multiple human tissues and cancer cell lines demonstrated that MELK was over-expressed at a significantly high level in a great majority of breast cancers and cell lines, but was not expressed in normal vital organs (heart, liver, lung and kidney) (WO2006/016525). Furthermore, suppression of MELK expression by siRNA was shown to significantly inhibit growth of human breast cancer cells. Accordingly, MELK is considered to be a suitable target for cancer immunotherapy and epitope peptides derived therefrom may be expected to serve as cancer immunotherapeutics effective in the treatment of a wide array of cancer types.

Summary

[0007] The present disclosure is based in part on the discovery of suitable targets of immunotherapy. Because TAAs are generally perceived by the immune system as "self" and therefore often have no innate immunogenicity, the discovery of appropriate targets is of extreme importance. Recognizing that MELK has been identified as up-regulated in cancer tissues of breast cancer, bladder cancer, cervical cancer, cholangiocellular carcinoma, chronic myeloid leukemia (CML), colorectal cancer, en-

dometriosis, esophagus cancer, gastric cancer, liver cancer, non-small cell lung cancer (NSCLC), lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma and small cell lung cancer (SCC), the present disclosure targets this maternal embryonic leucine zipper kinase (MELK) protein (SEQ ID NO: 94) encoded by the gene of GenBank Accession No. NM_014791 (SEQ ID NO: 93) for further analysis. In particular, MELK gene products containing epitope peptides that elicit CTLs specific to the corresponding molecules were selected for study. Peripheral blood mononuclear cells (PBMCs) obtained from a healthy donor were stimulated using HLA-A*2402 or HLA-A*0201 binding candidate peptides derived from MELK. CTLs that specifically recognize HLA-A24 or HLA-A02 positive target cells pulsed with the respective candidate peptides were established, and HLA-A24 or HLA-A02 restricted epitope peptides that can induce potent and specific immune responses against MELK expressed on the surface of tumor cells were identified. These results demonstrate that MELK is strongly immunogenic and the epitopes thereof are effective targets for tumor immunotherapy.

[0007a] In one aspect, there is provided an isolated peptide of less than 15 amino acids, wherein said peptide comprises an amino acid sequence selected from the group consisting of:

- (a) SEQ ID NOs:36, 46, 57, 60 and 62; and
- (b) SEQ ID NOs:36, 46, 57, 60 and 62, in which 1 or 2 amino acids are substituted, inserted, deleted and/or added, and has cytotoxic T lymphocyte (CTL) inducibility.

[0008] Accordingly, it is an object of the present disclosure to provide peptides having CTL inducibility as well as an amino acid sequence selected from the group of SEQ ID NOs: 14, 21, 23, 27, 36, 46, 57, 60 and 62. In addition, the present disclosure contemplates modified peptides, having an amino acid sequence of SEQ ID NOs: 14, 21, 23, 27, 36, 46, 57, 60 or 62, wherein one, two or more amino acids are substituted, deleted, incorporated, and/or added, so long as the modified peptides retain the original CTL inducibility. In another aspect, there is provided an isolated polynucleotide encoding a peptide as described herein.

[0008a] In another aspect, there is provided an *in vitro* method for inducing an antigen-presenting cell with high CTL inducibility, wherein the method comprises the step of contacting an antigen-presenting cell with a peptide as described herein, or the step of introducing a polynucleotide encoding the peptide in an expressible form into an antigen-presenting cell. In another aspect, there is provided an *in vitro* method for inducing CTL by using a peptide as described herein.

[0008b] In another aspect, there is provided an isolated CTL which targets any of the peptides as described herein. In another aspect, there is provided an isolated CTL which is induced by a peptide as described herein.

[0008c] In another aspect, there is provided an isolated antigen-presenting cell that presents a complex formed between a HLA antigen and a peptide as described herein on its surface.

[0009] When administered to a subject, the present peptides are presented on the surface of antigen-expressing cells and then induce CTLs targeting the respective peptides. Therefore, it is an object of the present disclosure to provide antigen-presenting cells which present any of the present peptides, as well as methods for inducing antigen-presenting cells. An anti-tumor immune response is induced by the administration of the present MELK polypeptides or polynucleotide encoding the polypeptides, as well as exosomes and antigen-presenting cells which present the MELK polypeptides. Therefore, it is yet another object of the present disclosure to provide pharmaceutical agents or compositions containing the polypeptides or polynucleotides encoding them, as well as the exosomes and antigen-presenting cells as their active ingredients. The pharmaceutical agents of the present disclosure find use as vaccines.

[0009a] In another aspect, there is provided a pharmaceutical composition for the treatment and/or prophylaxis of cancer, and/or the prevention of a postoperative recurrence thereof, wherein the pharmaceutical composition comprises an active ingredient selected from the group consisting of:

- (a) one or more peptides as described herein;
- (b) one or more polynucleotides encoding the peptide in an expressible form;
- (c) one or more antigen-presenting cells and/or exosomes, which antigen-presenting cells and exosomes present a complex formed between a HLA antigen and a peptide as described herein on its surface;
- (d) one or more CTLs induced against a peptide as described herein; and
- (e) combinations thereof.

[0009b] In another aspect, there is provided a composition for inducing an immune response against a cancer in a subject, wherein said composition comprises an active ingredient selected from the group consisting of:

- (a) one or more peptides as described herein;
- (b) one or more polynucleotides encoding the peptide in an expressible form;

(c) one or more antigen-presenting cells and/or exosomes, which antigen-presenting cells and exosomes present a complex formed between a HLA antigen and a peptide as described herein on its surface;

(d) one or more CTLs induced against a peptide as described herein; and

(e) combinations thereof.

[0009c] In another aspect, there is provided a composition for inducing a CTL, wherein said composition comprises an active ingredient selected from the group consisting of:

(a) one or more peptides as described herein;

(b) one or more polynucleotides encoding the peptide in an expressible form; and

(c) one or more antigen-presenting cells and/or exosomes, which antigen-presenting cells and exosomes present a complex formed between a HLA antigen and a peptide as described herein on its surface.

[0010] It is a further object of the present disclosure to provide methods for the treatment and/or prophylaxis of (i.e., preventing) cancers (tumors), and/or prevention of post-operative recurrence thereof, as well as methods for inducing CTLs, methods for inducing an immune response against tumor-associated endothelia and also anti-tumor immunity, which methods include the step of administering the MELK polypeptides,

polynucleotides encoding MELK polypeptides, exosomes or the antigen-presenting cells presenting MELK polypeptides or the pharmaceutical agents of the disclosure. In addition, the CTLs of the disclosure also find use as vaccines against cancer. Examples of the cancer include, but are not limited to breast cancer, bladder cancer, cervical cancer, cholangiocellular carcinomas, CML, colorectal cancer, endometriosis, esophagus cancer, gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma and SCC.

[0011] It is to be understood that both the foregoing summary and the following detailed description are of exemplified embodiments, and not restrictive of the disclosure or other alternate embodiments of the disclosure.

Brief Description of Drawings

[0012] Various aspects and applications of the present invention will become apparent to the skilled artisan upon consideration of the brief description of the figures and the detailed description of the present invention and its preferred embodiments which follows.

[fig.1] Figure 1 includes a series of photographs, (a) - (i) depicting the results of IFN-gamma ELISPOT assay on CTLs that were induced with peptides derived from MELK. The CTLs in well number #5 stimulated with MELK-A24-9-326 (SEQ ID NO: 14) (a), #3 with MELK-A24-9-78 (SEQ ID NO: 21) (b), #4 with MELK-A24-10-637 (SEQ ID NO: 23) (c), #4 with MELK-A24-10-532 (SEQ ID NO: 27) (d), #5 with MELK-A02-9-138 (SEQ ID NO: 36) (e), #3 with MELK-A02-9-193 (SEQ ID NO: 46) (f), #1 with MELK-A02-9-171 (SEQ ID NO: 57) (g), #2 with MELK-A02-9-71 (SEQ ID NO: 60) (h) and #2 with MELK-A02-9-532 (SEQ ID NO: 62) (i) showed potent IFN-gamma production as compared with the control, respectively. The cells in the wells denoted with a rectangular box were expanded to establish CTL lines. In the figures, "+" indicates that the target cells in the well were pulsed with the appropriate peptide, and "-" indicates that the target cells had not been pulsed with any peptides.

[fig.2] Figure 2 includes a series of line graphs, (a) - (i), depicting the IFN-gamma production of CTL lines stimulated with SEQ ID NO: 14 (a), SEQ ID NO: 21 (b), SEQ ID NO: 23 (c), SEQ ID NO: 27 (d), SEQ ID NO: 36 (e), SEQ ID NO: 46 (f), SEQ ID NO: 57 (g), SEQ ID NO: 60 (h) and SEQ ID NO: 62 (i) with IFN-gamma ELISA assay. CTL lines established by stimulation with each peptide showed potent IFN-gamma production as compared with the control. In the figures, "+" indicates that the target cells were pulsed with the appropriate peptide and "-" indicates that the target cells had not been pulsed with any peptides.

[fig.3] Figure 3 depicts the IFN-gamma production of the CTL clone established by limiting dilution from the CTL line stimulated with SEQ ID NO: 27. The results

depicted herein demonstrate that the CTL clone established by stimulation with SEQ ID NO: 27 showed potent IFN-gamma production as compared with the control. In the figure, "+" indicates that the target cells were pulsed with SEQ ID NO: 27 and "-" indicates that the target cells had not been pulsed with any peptides.

[fig.4]Figure 4 is composed of line graphs (a) and (b) depicting specific CTL activity of the CTL clones established with MELK-A02-9-138 (SEQ ID NO:36) (a) or MELK-A02-9-171 (SEQ ID NO: 57) (b) against target cells that exogenously express MELK and HLA-A*0201. The target cells were prepared by co-transfecting COS7 cells with the full length of both MELK and HLA-A*0201 molecule gene (-closed diamond-), and the control target cells were prepared by transfecting COS7 cells with HLA-A*0201 molecule gene and pulsing with an inappropriate peptide that is different from the peptides with which the CTL clones were established (-open triangle-), or transfecting COS7 cells with MELK gene (-open circle-). The CTL clone established with MELK -A02-9-138 (SEQ ID NO: 36) (a) and MELK -A02-9-171 (SEQ ID NO: 57) (b) showed potent IFN-gamma production against COS7 cells transfected with both MELK and HLA-A*0201 (-closed diamond-) as compared with the controls (-open triangle-, -open circle-).

Description of Embodiments

[0013] Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments of the present invention, the preferred methods, devices, and materials are now described. However, before the present materials and methods are described, it is to be understood that the present invention is not limited to the particular sizes, shapes, dimensions, materials, methodologies, protocols, etc. described herein, as these may vary in accordance with routine experimentation and optimization. It is also to be understood that the terminology used in the description is for the purpose of describing the particular versions or embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

[0014] The disclosure of each publication, patent or patent application mentioned in this specification is specifically incorporated by reference herein in its entirety. However, nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

[0015] In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

[0016] I. Definitions

The words "a", "an", and "the" as used herein mean "at least one" unless otherwise

specifically indicated.

- [0017] The terms "polypeptide", "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is a modified residue, or a non-naturally occurring residue, such as an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers.
- [0018] The term "amino acid" as used herein refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that similarly function to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those modified after translation in cells (e.g., hydroxyproline, gamma-carboxyglutamate, and O-phosphoserine). The phrase "amino acid analog" refers to compounds that have the same basic chemical structure (an alpha carbon bound to a hydrogen, a carboxy group, an amino group, and an R group) as a naturally occurring amino acid but have a modified R group or modified backbones (e.g., homoserine, norleucine, methionine, sulfoxide, methionine methyl sulfonium). The phrase "amino acid mimetic" refers to chemical compounds that have different structures but similar functions to general amino acids.
- [0019] Amino acids may be referred to herein by their commonly known three letter symbols or the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission.
- [0020] The terms "gene", "polynucleotides", "nucleotides" and "nucleic acids" are used interchangeably herein and, unless otherwise specifically indicated, are similarly to the amino acids referred to by their commonly accepted single-letter codes.
- [0021] Unless otherwise defined, the term "cancer" refers to the cancers over-expressing the MELK gene, examples of which include, but are not limited to, breast cancer, bladder cancer, cervical cancer, cholangiocellular carcinoma, chronic myeloid leukemia (CML), colorectal cancer, endometriosis, esophagus cancer, gastric cancer, liver cancer, non-small cell lung cancer (NSCLC), lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma and small cell lung cancer (SCC)s.
- [0022] Unless otherwise defined, the term "cytotoxic T lymphocyte", "cytotoxic T cell" and "CTL" are used interchangeably herein and, unless otherwise specifically indicated, refer to a sub-group of T lymphocytes that are capable of recognizing non-self cells (e.g., tumor cells, virus-infected cells) and inducing the death of such cells.
- [0023] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.
- [0024] II. Peptides

To demonstrate that peptides derived from MELK function as an antigen recognized by cytotoxic T lymphocytes (CTLs), peptides derived from MELK (SEQ ID NO: 93) were analyzed to determine whether they were antigen epitopes restricted by HLA-A24 or HLA-A02 which are commonly encountered HLA alleles (Date Y et al., *Tissue Antigens* 47: 93-101, 1996; Kondo A et al., *J Immunol* 155: 4307-12, 1995; Kubo RT et al., *J Immunol* 152: 3913-24, 1994). Candidates of HLA-A24 or HLA-A02 binding peptides derived from MELK were identified based on their binding affinities to HLA-A24 or HLA-A02. After in vitro stimulation of T-cells by dendritic cells (DCs) loaded with these peptides, CTLs were successfully established using the following peptides:

MELK- A24-9-326 (SEQ ID NO: 14),
 MELK-A24-9-78 (SEQ ID NO: 21),
 MELK-A24-10-637 (SEQ ID NO: 23),
 MELK-A24-10- 532 (SEQ ID NO: 27),
 MELK-A02-9-138 (SEQ ID NO: 36),
 MELK- A02-9-193 (SEQ ID NO: 46),
 MELK- A02-9-171 (SEQ ID NO: 57),
 MELK- A02-9-71 (SEQ ID NO: 60),
 and

MELK- A02- 9-532 (SEQ ID NO: 62),

These established CTLs show potent specific CTL activity against target cells pulsed with respective peptides. These results herein demonstrate that MELK is an antigen recognized by CTL and that the peptides may be epitope peptides of MELK restricted by HLA-A24 or HLA-A02.

[0025] Since the MELK gene is over expressed in most cancer tissues, such as breast cancer, bladder cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophagus cancer, gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma and SCC, it is a good target for immunotherapy. Thus, the present invention provides nonapeptides (peptides consisting of nine amino acid residues) and decapeptides (peptides consisting of ten amino acid residues) corresponding to CTL-recognized epitopes of MELK. Particularly preferred examples of nonapeptides and decapeptides of the present invention include those peptides consisting of the amino acid sequence selected from among SEQ ID NOs: 14, 21, 23, 27, 36, 46, 57, 60 and 62.

[0026] Generally, software programs presently available on the Internet, such as those described in Parker KC et al., *J Immunol* 1994 Jan 1, 152(1): 163-75, can be used to calculate the binding affinities between various peptides and HLA antigens in silico. Binding affinity with HLA antigens can be measured as described, for example, in the references to Parker KC et al., *J Immunol* 1994 Jan 1, 152(1): 163-75; and Kuzushima

K et al., Blood 2001, 98(6): 1872-81. The methods for determining binding affinity is described, for example, in; Journal of Immunological Methods, 1995, 185: 181-190.; Protein Science, 2000, 9: 1838-1846. Thus, the present invention encompasses peptides of MELK which bind with HLA antigens identified using such known programs.

[0027] The nonapeptides and decapeptides of the present invention can be flanked with additional amino acid residues so long as the resulting peptide retains its CTL inducibility. Such peptides having CTL inducibility are typically less than about 40 amino acids, often less than about 20 amino acids, usually less than about 15 amino acids. The particular amino acid sequences flanking the nonapeptide and decapeptides of the present invention (e.g., peptides consisting of the amino acid sequence selected from among SEQ ID NOs: 14, 21, 23, 27, 36, 46, 57, 60 and 62 is not limited and can be composed of any kind of amino acids so long as it does not impair the CTL inducibility of the original peptide. Thus, the present invention also provides peptides having CTL inducibility and including the amino acid sequence selected from among SEQ ID NOs: 14, 21, 23, 27, 36, 46, 57, 60 and 62.

[0028] In general, the modification of one, two, or more amino acids in a protein will not influence the function of the protein, and in some cases will even enhance the desired function of the original protein. In fact, modified peptides (i.e., peptides composed of an amino acid sequence in which one, two or several amino acid residues have been modified (i.e., substituted, deleted, added or inserted as compared to an original reference sequence) have been known to retain the biological activity of the original peptide (Mark et al., Proc Natl Acad Sci USA 1984, 81: 5662-6; Zoller and Smith, Nucleic Acids Res 1982, 10: 6487-500; Dalbadie-McFarland et al., Proc Natl Acad Sci USA 1982, 79: 6409-13). Thus, in one embodiment, the peptides of the present invention may have both CTL inducibility and an amino acid sequence selected from among SEQ ID NOs: 14, 21, 23, 27, 36, 46, 57, 60 and 62 wherein one, two or even more amino acids are added, inserted, deleted and/or substituted.

[0029] Those skilled in the art recognize that individual modification to an amino acid sequence which alters a single amino acid or a small percentage of amino acids tend to result in the conservation of the properties of the original amino acid side-chain. As such, they are often referred to as "conservative substitutions" or "conservative modifications", wherein the alteration of a protein results in a modified protein having a function analogous to the original protein. Conservative substitution tables providing functionally similar amino acids are well known in the art. Examples of amino acid side chain characteristics that are desirable to conserve include, for example, hydrophobic amino acids (A, I, L, M, F, P, W, Y, V), hydrophilic amino acids (R, D, N, C, E, Q, G, H, K, S, T), and side chains having the following functional groups or

characteristics in common: an aliphatic side-chain (G, A, V, L, I, P); a hydroxyl group containing side-chain (S, T, Y); a sulfur atom containing side-chain (C, M); a carboxylic acid and amide containing side-chain (D, N, E, Q); a base containing side-chain (R, K, H); and an aromatic containing side-chain (H, F, Y, W). In addition, the following eight groups each contain amino acids that are accepted in the art as conservative substitutions for one another:

- 1) Alanine (A), Glycine (G);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W);
- 7) Serine (S), Threonine (T); and
- 8) Cysteine (C), Methionine (M) (see, e.g., Creighton, *Proteins* 1984).

[0030] Such conservatively modified peptides are also considered to be peptides of the present invention. However, peptides of the present invention are not restricted thereto and can include non-conservative modifications, so long as the modified peptide retains the CTL inducibility of the original peptide. Furthermore, modified peptides should not exclude CTL inducible peptides of polymorphic variants, interspecies homologues, and alleles of MELK.

[0031] To retain the requisite CTL inducibility one can modify (insert, delete, add, and/or substitute) a small number (for example, 1, 2 or several) or a small percentage of amino acids. Herein, the term "several" means 5 or fewer amino acids, for example, 4 or 3 or fewer. The percentage of amino acids to be modified is preferably 20% or less, more preferably 15% or less, even more preferably 10% or less or 1 to 5%.

[0032] Homology analysis of preferred peptides of the present invention, MELK-A24-9-326 (SEQ ID NO: 14), MELK-A24-9-78 (SEQ ID NO: 21), MELK-A24-10-637 (SEQ ID NO: 23), MELK-A24-10-532 (SEQ ID NO: 27), MELK-A02-9-138 (SEQ ID NO: 36), MELK-A02-9-193 (SEQ ID NO: 46), MELK-A02-9-171 (SEQ ID NO: 57), MELK-A02-9-71 (SEQ ID NO: 60) and MELK-A02-9-532 (SEQ ID NO: 62) confirmed that these peptides do not have significant homology with peptides derived from any other known human gene products. Thus, the possibility of these peptides generating unknown or undesired immune responses when used for immunotherapy is significantly lowered. Accordingly, these peptides are expected to be highly useful for eliciting immunity in tumor patients against MELK on cancer cells, such as breast cancer, bladder cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophagus cancer, gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal

carcinoma and SCC.

- [0033] When used in the context of immunotherapy, peptides of the present invention should be presented on the surface of a cell or exosome, preferably as a complex with an HLA antigen. Therefore, one is preferable to select peptides that not only induce CTLs but also possess high binding affinity to the HLA antigen. To that end, the peptides can be modified by substitution, insertion, deletion, and/or addition of the amino acid residues to yield a modified peptide having improved binding affinity. In addition to peptides that are naturally displayed, since the regularity of the sequences of peptides displayed by binding to HLA antigens is already known (J Immunol 1994, 152: 3913; Immunogenetics 1995, 41: 178; J Immunol 1994, 155: 4307), modifications based on such regularity can be introduced into the immunogenic peptides of the present invention. For example, it may be desirable to substitute the second amino acid from the N-terminus with phenylalanine, tyrosine, methionine, or tryptophan, and/or the amino acid at the C-terminus with phenylalanine, leucine, isoleucine, tryptophan, or methionine in order to increase the HLA-A24 binding. Thus, peptides having the amino acid sequences selected from among SEQ ID NO: 14, SEQ ID NO: 21, SEQ ID NO: 23 and SEQ ID NO: 27 wherein the second amino acid from the N-terminus of the amino acid sequence of the SEQ ID NOs is substituted with phenylalanine, tyrosine, methionine, or tryptophan, and/or wherein the C-terminus of the amino acid sequence of the SEQ ID NOs is substituted with phenylalanine, leucine, isoleucine, tryptophan, or methionine are encompassed by the present invention. On the other hand, peptides possessing high HLA-A02 binding affinity have their second amino acid from the N-terminus substituted with leucine or methionine, and/or the amino acid at C-terminus substituted with valine or leucine. Thus, peptides having the amino acid sequences selected from among SEQ ID NO: 36, SEQ ID NO: 46, SEQ ID NO: 57, SEQ ID NO: 60 and SEQ ID NO: 62 wherein the second amino acid from the N-terminus is substituted with leucine or methionine, and/or wherein the C-terminus is substituted with valine or leucine are encompassed by the present invention. Substitutions can be introduced not only at the terminal amino acids but also at the position of potential TCR recognition of peptides. Several studies have demonstrated that amino acid substitutions in a peptide can be equal to or better than the original, for example CAP1, p53₍₂₆₄₋₂₇₂₎, Her-2/neu₍₃₆₉₋₃₇₇₎ or gp100₍₂₀₉₋₂₁₇₎ (Zaremba et al. Cancer Res. 57, 4570-4577, 1997, T. K. Hoffmann et al. J Immunol. (2002) Feb 1;168(3):1338-47., S. O. Dionne et al. Cancer Immunol immunother. (2003) 52: 199-206 and S. O. Dionne et al. Cancer Immunology, Immunotherapy (2004) 53, 307-314).
- [0034] The present invention also contemplates the addition of one to two amino acids to the N and/or C-terminus of the described peptides. Such modified peptides having high HLA antigen binding affinity and retained CTL inducibility are also included in the

present invention.

- [0035] However, when the peptide sequence is identical to a portion of the amino acid sequence of an endogenous or exogenous protein having a different function, side effects such as autoimmune disorders and/or allergic symptoms against specific substances may be induced. Therefore, it is preferable to first perform homology searches using available databases to avoid situations in which the sequence of the peptide matches the amino acid sequence of another protein. When it becomes clear from the homology searches that there exists not even a peptide with 1 or 2 amino acid differences as compared to the objective peptide, the objective peptide can be modified in order to increase its binding affinity with HLA antigens, and/or increase its CTL inducibility without any danger of such side effects.
- [0036] Although peptides having high binding affinity to the HLA antigens as described above are expected to be highly effective, the candidate peptides, which are selected according to the presence of high binding affinity as an indicator, are further examined for the presence of CTL inducibility. Herein, the phrase "CTL inducibility" indicates the ability of the peptide to induce cytotoxic lymphocytes (CTLs) when presented on antigen-presenting cells. Further, "CTL inducibility" includes the ability of the peptide to induce CTL activation, CTL proliferation, promote CTL lysis of target cells, and to increase CTL IFN-gamma production.
- [0037] Confirmation of CTL inducibility is accomplished by inducing antigen-presenting cells carrying human MHC antigens (for example, B-lymphocytes, macrophages, and dendritic cells (DCs)) by stimulating with the peptides, mixing the induced antigen-presenting cells with CD8-positive cells, and then measuring the IFN-gamma produced and released by CTL against the target cells. Preferably, antigen-presenting cells are DCs derived from human peripheral blood mononuclear leukocytes. As the reaction system, transgenic animals that have been produced to express a human HLA antigen (for example, those described in BenMohamed L, Krishnan R, Longmate J, Auye C, Low L, Primus J, Diamond DJ, *Hum Immunol* 2000 Aug, 61(8): 764-79, Related Articles, Books, Linkout Induction of CTL response by a minimal epitope vaccine in HLA A*0201/DR1 transgenic mice: dependence on HLA class II restricted T(H) response) can be used. For example, the target cells can be radio-labeled with ^{51}Cr and such, and cytotoxic activity can be calculated from radioactivity released from the target cells. Alternatively, CTL inducibility can be assessed by measuring IFN-gamma produced and released by CTL in the presence of antigen-presenting cells (APCs) that carry immobilized peptides, and visualizing the inhibition zone on the media using anti-IFN-gamma monoclonal antibodies.
- [0038] As a result of assessing the CTL inducibility of the peptides as described above, it was discovered that those peptides predicted having high binding affinity to an HLA

antigen did not necessarily have high CTL inducibility. However, of those peptides identified and assessed, nonapeptides or decapeptides having an amino acid sequence selected from among SEQ ID NOs: 14, 21, 23, 27, 36, 46, 57, 60 and 62 were found to exhibit particularly high CTL inducibility. Thus, these peptides are exemplified as preferred embodiments of the present invention.

- [0039] In addition to the above-described modifications, the peptides of the present invention can also be linked to other substances, so long as the resulting linked peptide retains the requisite CTL inducibility of the original peptide. Examples of suitable substances include, but are not limited to: peptides, lipids, sugar and sugar chains, acetyl groups, natural and synthetic polymers, etc. The peptides can contain modifications such as glycosylation, side chain oxidation, or phosphorylation, etc., provided the modifications do not destroy the biological activity of the original peptide. These kinds of modifications can be performed to confer additional functions (e.g., targeting function, and delivery function) or to stabilize the polypeptide.
- [0040] For example, to increase the in vivo stability of a polypeptide, it is known in the art to introduce D-amino acids, amino acid mimetics or unnatural amino acids; this concept can also be adapted to the present polypeptides. The stability of a polypeptide can be assayed in a number of ways. For instance, peptidases and various biological media, such as human plasma and serum, can be used to test stability (see, e.g., Verhoef et al., *Eur J Drug Metab Pharmacokin* 1986, 11: 291-302).
- [0041] Further, the peptides of the present invention may be linked to other peptides via spacers or linkers. Examples of other peptides include, but are not limited to, CTL inducible peptides derived from other TAAs. Alternatively, two or more peptides of the present invention may be linked via spacers or linkers. The peptides linked via spacers or linkers may be the same or different each other. Spacers or linkers are not specifically limited, but are preferably peptides, more preferably peptides having one or more cleavage sites which are capable of being cleaved by enzymes such as peptidases, proteases and proteasomes. Examples of linkers or spacers include, but are not limited to: AAY (P. M. Daftarian et al., *J Trans Med* 2007, 5:26), AAA, NKRK (R. P. M. Suttmüller et al., *J Immunol.* 2000, 165: 7308-7315) or, one to several lysine residues (S. Ota et al., *Can Res.* 62, 1471-1476, K. S. Kawamura et al., *J Immunol.* 2002, 168: 5709-5715). The peptides of the present invention encompass those peptides linked to other peptides via spacers or linkers.
- [0042] The peptides of the present invention may be existed on the surface of a cell carrying human MHC antigens (e.g. antigen presenting cell) or an exosome as complexes in combination with MHC molecules and then induce CTLs. The cells and the exosomes can be prepared by well-known methods in the art, for example, the cells may be prepared by contacting with the peptides of the present invention, and the exosomes

may be prepared by collecting an exosome-containing fraction from the cells contacted with the peptides of the present invention (see, e.g., Japanese Patent Application Kohyo Publications Nos. Hei 11-510507 and WO99/03499). The peptide of the present invention encompasses those peptides existed on the surface of a cell or an exosome as complexes in combination with MHC molecules.

[0043] Herein, the peptides of the present invention can also be described as "MELK peptide(s)" or "MELK polypeptide(s)".

[0044] III. Preparation of MELK peptides

The peptides of the present invention can be prepared using well known techniques. For example, the peptides can be prepared synthetically, using recombinant DNA technology or chemical synthesis. Peptide of the invention can be synthesized individually or as longer polypeptides composed of two or more peptides. The peptides can then be isolated i.e., purified or isolated so as to be substantially free of other naturally occurring host cell proteins and fragments thereof, or any other chemical substances.

[0045] A peptide of the present invention can be obtained through chemical synthesis based on the selected amino acid sequence. Examples of conventional peptide synthesis methods that can be adapted to the synthesis include, but are not limited to:

- (i) Peptide Synthesis, Interscience, New York, 1966;
- (ii) The Proteins, Vol. 2, Academic Press, New York, 1976;
- (iii) Peptide Synthesis (in Japanese), Maruzen Co., 1975;
- (iv) Basics and Experiment of Peptide Synthesis (in Japanese), Maruzen Co., 1985;
- (v) Development of Pharmaceuticals (second volume) (in Japanese), Vol. 14 (peptide synthesis), Hirokawa, 1991;
- (vi) WO99/67288; and
- (vii) Barany G. & Merrifield R.B., Peptides Vol. 2, "Solid Phase Peptide Synthesis", Academic Press, New York, 1980, 100-118.

[0046] Alternatively, the present peptides can be obtained adapting any known genetic engineering methods for producing peptides (e.g., Morrison J, J Bacteriology 1977, 132: 349-51; Clark-Curtiss & Curtiss, Methods in Enzymology (eds. Wu et al.) 1983, 101: 347-62). For example, first, a suitable vector harboring a polynucleotide encoding the objective peptide in an expressible form (e.g., downstream of a regulatory sequence corresponding to a promoter sequence) is prepared and transformed into a suitable host cell. The host cell is then cultured to produce the peptide of interest. The peptide can also be produced in vitro adopting an in vitro translation system.

[0047] IV. Polynucleotides

The present invention also provides a polynucleotide which encodes any of the aforementioned peptides of the present invention. These include polynucleotides derived from the natural occurring MELK gene (GenBank Accession No. NM_014791 (SEQ

ID NO: 93)) as well as those having a conservatively modified nucleotide sequence thereof. Herein, the phrase "conservatively modified nucleotide sequence" refers to sequences which encode identical or essentially identical amino acid sequences. Due to the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG, and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a peptide also describes every possible silent variation of the nucleic acid. One of ordinary skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid that encodes a peptide is implicitly described in each disclosed sequence.

[0048] The polynucleotide of the present invention can be composed of DNA, RNA, and derivatives thereof. A DNA is suitably composed of bases such as A, T, C, and G, and T is replaced by U in an RNA.

[0049] The polynucleotide of the present invention can encode multiple peptides of the present invention, with or without intervening amino acid sequences in between. For example, the intervening amino acid sequence can provide a cleavage site (e.g., enzyme recognition sequence) of the polynucleotide or the translated peptides. Furthermore, the polynucleotide can include any additional sequences to the coding sequence encoding the peptide of the present invention. For example, the polynucleotide can be a recombinant polynucleotide that includes regulatory sequences required for the expression of the peptide or can be an expression vector (plasmid) with marker genes and such. In general, such recombinant polynucleotides can be prepared by the manipulation of polynucleotides through conventional recombinant techniques using, for example, polymerases and endonucleases.

[0050] Both recombinant and chemical synthesis techniques can be used to produce the polynucleotides of the present invention. For example, a polynucleotide can be produced by insertion into an appropriate vector, which can be expressed when transfected into a competent cell. Alternatively, a polynucleotide can be amplified using PCR techniques or expression in suitable hosts (see, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, 1989). Alternatively, a polynucleotide can be synthesized using the solid phase techniques, as described in Beaucage SL & Iyer RP, *Tetrahedron* 1992, 48: 2223-311; Matthes et al., *EMBO J* 1984, 3: 801-5.

[0051] Vectors containing the polynucleotide of the present invention and host cells harboring the vectors are also included in the present invention.

[0052] V. Exosomes

The present invention further provides intracellular vesicles called exosomes, which present complexes formed between the peptides of this invention and HLA antigens on their surface. Exosomes can be prepared, for example by using the methods detailed in Japanese Patent Application Kohyo Publications Nos. Hei 11-510507 and WO99/03499, and can be prepared using APCs obtained from patients who are subject to treatment and/or prevention. The exosomes of this invention can be inoculated as vaccines, in a fashion similar to the peptides of this invention.

[0053] The type of HLA antigens contained in the complexes must match that of the subject requiring treatment and/or prevention. For example, in the Japanese population, HLA-A24, particularly HLA-A2402, is prevalent and therefore would be appropriate for treatment of a Japanese patient. The use of the A24 type or the A02 type that is highly expressed among the Japanese and Caucasian is favorable for obtaining effective results, and subtypes such as A2402 or A0201 also find use. Typically, in the clinic, the type of HLA antigen of the patient requiring treatment is investigated in advance, which enables the appropriate selection of peptides having high levels of binding affinity to the particular antigen, or having CTL inducibility by antigen presentation. Furthermore, in order to obtain peptides having both high binding affinity and CTL inducibility, substitution, insertion and/or addition of 1, 2, or several amino acids can be performed based on the amino acid sequence of the naturally occurring MELK partial peptide.

[0054] When using an exosome including the A24 type HLA antigen, the peptides having the amino acid sequences selected from among SEQ ID NOs: 14, 21, 23 and 27 are useful. On the other hand, when using an exosome including A02 type HLA antigen, the peptides having the sequences selected from among SEQ ID NO: 36, 46, 57, 60 and 62 are preferred.

[0055] VI. Antigen-presenting cells (APCs)

The present invention also provides isolated APCs that present complexes formed between HLA antigens and the peptides of this invention on its surface. The APCs that are obtained by contacting the peptides of this invention, or introducing the nucleotides encoding the peptides of this invention in an expressible form can be derived from patients who are subject to treatment and/or prevention, and can be administered as vaccines by themselves or in combination with other drugs including the peptides of this invention, exosomes, or cytotoxic T cells.

[0056] The APCs are not limited to a particular kind of cells and include dendritic cells (DCs), Langerhans cells, macrophages, B cells, and activated T cells, which are known

to present proteinaceous antigens on their cell surface so as to be recognized by lymphocytes. Since DC is a representative APC having the strongest CTL inducing action among APCs, DCs find use as the APCs of the present invention.

[0057] For example, an APC can be obtained by inducing DCs from peripheral blood monocytes and then contacting (stimulating) them with the peptides of this invention in vitro, ex vivo or in vivo. When the peptides of this invention are administered to the subjects, APCs that present the peptides of this invention are induced in the body of the subject. The phrase "inducing APC" includes contacting (stimulating) a cell with the peptides of this invention, or nucleotides encoding the peptides of this invention to present complexes formed between HLA antigens and the peptides of this invention on cell's surface. After introducing the peptides of this invention to the APCs to allow the APCs to present the peptides, the APCs can be administered to the subject as a vaccine. For example, the ex vivo administration can include the steps of:

- a: collecting APCs from a first subject,
- b: contacting with the APCs of step a, with the peptide and
- c: administering the peptide-loaded APCs to a second subject.

[0058] The first subject and the second subject can be the same individual, or may be different individuals. Alternatively, according to the present invention, use of the peptides of the present invention for manufacturing a pharmaceutical composition inducing antigen-presenting cells is provided. In addition, the present invention provides a method or process for manufacturing a pharmaceutical composition inducing antigen-presenting cells, wherein the method or the process includes the step of admixing or formulating the peptide of the present invention with a pharmaceutically acceptable carrier. Further, the present invention also provides the peptides of the present invention for inducing antigen-presenting cells. The APCs obtained by step (b) can be administered to the subject as a vaccine.

[0059] According to an aspect of the present invention, the APCs of the present invention have a high level of CTL inducibility. In the term of "high level of CTL inducibility", the high level is relative to the level of that by APCs contacted with no peptide or peptides which can not induce the CTL. Such APCs having a high level of CTL inducibility can be prepared by a method which includes the step of transferring genes containing polynucleotides that encode the peptides of this invention to APCs in vitro. The introduced genes can be in the form of DNAs or RNAs. Examples of methods for introduction include, without particular limitations, various methods conventionally performed in this field, such as lipofection, electroporation, and calcium phosphate method can be used. More specifically, it can be performed as described in Cancer Res 1996, 56: 5672-7; J Immunol 1998, 161: 5607-13; J Exp Med 1996, 184: 465-72; Published Japanese Translation of International Publication No. 2000-509281. By

transferring the gene into APCs, the gene undergoes transcription, translation, and such in the cell, and then the obtained protein is processed by MHC Class I or Class II, and proceeds through a presentation pathway to present peptides.

[0060] VII. Cytotoxic T cells (CTLs)

A cytotoxic T cell induced against any of the peptides of the present invention strengthens the immune response targeting tumor-associated endothelia in vivo and thus can be used as vaccines, in a fashion similar to the peptides per se. Thus, the present invention also provides isolated cytotoxic T cells that are specifically induced or activated by any of the present peptides.

[0061] Such cytotoxic T cells can be obtained by (1) administering the peptide of the present invention to a subject and then collecting cytotoxic T cells from the subject, or (2) contacting (stimulating) subject-derived APCs and CD8-positive cells, or peripheral blood mononuclear leukocytes in vitro with the peptides of the present invention and then isolating cytotoxic T cells.

[0062] The cytotoxic T cells, which have been induced by stimulation with APCs that present the peptides of this invention, can be derived from patients who are subject to treatment and/or prevention, and can be administered by themselves or in combination with other drugs including the peptides of this invention or exosomes for the purpose of regulating effects. The obtained cytotoxic T cells act specifically against target cells presenting the peptides of this invention, or for example, the same peptides used for induction. In the other word, the cytotoxic T cells can recognize (i.e., binding to) a complex formed between a HLA antigen and the peptide of the present invention on a target cell surface with the T cell receptor and then attack the target cell to induce the death of the target cell. The target cells can be cells that endogenously express MELK, or cells that are transfected with the MELK gene; and cells that present a peptide of this invention on the cell surface due to stimulation by the peptide can also serve as targets of activated CTL attack.

[0063] VIII. T cell receptor (TCR)

The present invention also provides a composition containing nucleic acids sequence encoding polypeptides that are capable of forming a subunit of a T cell receptor (TCR), and methods of using the same. The TCR subunits have the ability to form TCRs that confer specificity to T cells against tumor cells presenting MELK. By using the known methods in the art, the nucleic acid sequence of alpha- and beta- chains of the TCR expressed in the CTL induced with a peptide of the present invention can be identified (WO2007/032255 and Morgan et al., J Immunol, 171, 3288 (2003)). The derivative TCRs can bind to the MELK peptide displaying on the target cells with high avidity, and optionally mediate efficient killing of target cells presenting the MELK peptide in vivo and in vitro.

- [0064] The nucleic acids sequence encoding the TCR subunits can be incorporated into suitable vectors e.g. retroviral vectors. These vectors are well known in the art. The nucleic acids or the vectors containing them usefully can be transferred into a T cell, for example, a T cell from a patient. Advantageously, the invention provides an off-the-shelf composition allowing rapid modification of a patient's own T cells (or those of another mammal) to rapidly and easily produce modified T cells having excellent cancer cell killing properties.
- [0065] Also, the present invention provides CTLs which are prepared by transduction with the nucleic acids encoding the TCR subunits polypeptides that bind to the MELK peptide e.g. SEQ ID NOs: 14, 21, 23, 27, 36, 46, 57, 60 and 62 in the context of HLA-A24 or HLA-A02. The transduced CTLs are capable of homing to cancer cells in vivo, and can be expanded by well known culturing methods in vitro (e.g., Kawakami et al., J Immunol., 142, 3452-3461 (1989)). The T cells of the invention can be used to form an immunogenic composition useful in treating or the prevention of cancer in a patient in need of therapy or protection (WO2006/031221).
- [0066] IX. Pharmaceutical agents or compositions
- Prevention and prophylaxis include any activity which reduces the burden of mortality or morbidity from disease. Prevention and prophylaxis can occur "at primary, secondary and tertiary prevention levels." While primary prevention and prophylaxis avoid the development of a disease, secondary and tertiary levels of prevention and prophylaxis encompass activities aimed at the prevention and prophylaxis of the progression of a disease and the emergence of symptoms as well as reducing the negative impact of an already established disease by restoring function and reducing disease-related complications. Alternatively, prevention and prophylaxis include a wide range of prophylactic therapies aimed at alleviating the severity of the particular disorder, e.g. reducing the proliferation and metastasis of tumors.
- [0067] Treating and/or the prophylaxis of cancer or tumor, and/or the prevention of post-operative recurrence thereof includes any of the following steps, such as surgical removal of cancer cells, inhibition of the growth of cancerous cells, involution or regression of a tumor, induction of remission and suppression of occurrence of cancer, tumor regression, and reduction or inhibition of metastasis. Effectively treating and/or the prophylaxis of cancer decreases mortality and improves the prognosis of individuals having cancer, decreases the levels of tumor markers in the blood, and alleviates detectable symptoms accompanying cancer. For example, reduction or improvement of symptoms constitutes effectively treating and/or the prophylaxis includes 10%, 20%, 30% or more reduction, or stable disease.
- [0068] Since MELK expression is up-regulated in several cancers as compared with normal tissue, the peptides of this invention or polynucleotides encoding such peptides can be

used for the treatment and/or for the prophylaxis of cancer or tumor, and/or prevention of postoperative recurrence thereof. Thus, the present invention provides a pharmaceutical agent or composition for treating and/or for the prophylaxis of cancer or tumor, and/or prevention of postoperative recurrence thereof, which includes one or more of the peptides of this invention, or polynucleotides encoding the peptides as an active ingredient. Alternatively, the present peptides can be expressed on the surface of any of the foregoing exosomes or cells, such as APCs for the use as pharmaceutical agents or compositions. In addition, the aforementioned cytotoxic T cells which target any of the peptides of the present invention can also be used as the active ingredient of the present pharmaceutical agents or compositions. In the context of the present invention, the phrase "targeting a peptide" refers to recognizing (i.e., binding to) a complex formed between a HLA antigen and a peptide on a target cell surface with the T cell receptor, and then attacking the target cell to induce the death of the target cell.

[0069] In another embodiment, the present invention also provides the use of an active ingredient selected from among:

- (a) a peptide of the present invention,
 - (b) a nucleic acid encoding such a peptide as disclosed herein in an expressible form,
 - (c) an APC or an exosome presenting a peptide of the present invention on its surface, and
 - (d) a cytotoxic T cell of the present invention
- in manufacturing a pharmaceutical composition or agent for treating cancer or tumor.

[0070] Alternatively, the present invention further provides an active ingredient selected from among:

- (a) a peptide of the present invention,
 - (b) a nucleic acid encoding such a peptide as disclosed herein in an expressible form,
 - (c) an APC or an exosome presenting a peptide of the present invention on its surface, and
 - (d) a cytotoxic T cells of the present invention
- for use in for treating cancer or tumor.

[0071] Alternatively, the present invention further provides a method or process for manufacturing a pharmaceutical composition or agent for treating cancer or tumor, wherein the method or process includes the step of formulating a pharmaceutically or physiologically acceptable carrier with an active ingredient selected from among:

- (a) a peptide of the present invention,
- (b) a nucleic acid encoding such a peptide as disclosed herein in an expressible form,
- (c) an APC or an exosome presenting a peptide of the present invention on its surface, and
- (d) a cytotoxic T cells of the present invention

as active ingredients.

[0072] In another embodiment, the present invention also provides a method or process for manufacturing a pharmaceutical composition or agent for treating cancer or tumor, wherein the method or process includes the step of admixing an active ingredient with a pharmaceutically or physiologically acceptable carrier, wherein the active ingredient is selected from among:

- (a) a peptide of the present invention,
- (b) a nucleic acid encoding such a peptide as disclosed herein in an expressible form,
- (c) an APC or an exosome presenting a peptide of the present invention on its surface, and
- (d) a cytotoxic T cells of the present invention.

[0073] Alternatively, the pharmaceutical composition or agent of the present invention may be used for either or both the prophylaxis of cancer or tumor and prevention of post-operative recurrence thereof.

[0074] The present pharmaceutical agents or compositions find use as a vaccine. In the context of the present invention, the phrase "vaccine" (also referred to as an "immunogenic composition") refers to a substance that has the function to induce anti-tumor immunity upon inoculation into animals.

[0075] The pharmaceutical agents or compositions of the present invention can be used to treat and/or prevent cancers or tumors, and/or prevention of postoperative recurrence thereof in subjects or patients including human and any other mammal including, but not limited to, mouse, rat, guinea-pig, rabbit, cat, dog, sheep, goat, pig, cattle, horse, monkey, baboon, and chimpanzee, particularly a commercially important animal or a domesticated animal.

[0076] According to the present invention, polypeptides having an amino acid sequence selected from among SEQ ID NOs: 14, 21, 23 and 27 or polypeptides having an amino acid sequence selected from among SEQ ID NOs: 36, 46, 57, 60 and 62 have been found to be HLA-A24 or HLA-A02 restricted epitope peptides or candidates, respectively, that can induce potent and specific immune response. Therefore, the present pharmaceutical agents or compositions which include any of these polypeptides having the amino acid sequences selected from among of SEQ ID NOs: 14, 21, 23 and 27 are particularly suited for the administration to subjects whose HLA antigen is HLA-A24. On the one hand, the present pharmaceutical agents or compositions which contain any of these polypeptides having the amino acid sequences of SEQ ID NOs: 36, 46, 57, 60 and 62 are particularly suited for the administration to subjects whose HLA antigen is HLA-A02. The same applies to pharmaceutical agents which include polynucleotides encoding any of these polypeptides.

[0077] Cancers or tumors to be treated by the pharmaceutical agents or compositions of the

present invention are not limited and include all kinds of cancers or tumors wherein MELK is involved, including for example, bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophagus cancer, gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma and SCC.

[0078] The present pharmaceutical agents or compositions can contain in addition to the aforementioned active ingredients, other peptides which have the ability to induce CTLs against cancerous cells, other polynucleotides encoding the other peptides, other cells that present the other peptides, or such. Herein, the other peptides that have the ability to induce CTLs against cancerous cells are exemplified by cancer specific antigens (e.g., identified TAAs), but are not limited thereto.

[0079] If needed, the pharmaceutical agents or compositions of the present invention can optionally include other therapeutic substances as an active ingredient, so long as the substance does not inhibit the antitumoral effect of the active ingredient, e.g., any of the present peptides. For example, formulations can include anti-inflammatory agents, pain killers, chemotherapeutics, and the like. In addition to including other therapeutic substances in the medicament itself, the medicaments of the present invention can also be administered sequentially or concurrently with the one or more other pharmacologic agents. The amounts of medicament and pharmacologic agent depend, for example, on what type of pharmacologic agent(s) is/are used, the disease being treated, and the scheduling and routes of administration.

[0080] It should be understood that in addition to the ingredients particularly mentioned herein, the pharmaceutical agents or compositions of this invention can include other agents conventional in the art having regard to the type of formulation in question.

[0081] In one embodiment of the present invention, the present pharmaceutical agents or compositions can be included in articles of manufacture and kits containing materials useful for treating the pathological conditions of the disease to be treated, e.g, cancer. The article of manufacture can include a container of any of the present pharmaceutical agents or compositions with a label. Suitable containers include bottles, vials, and test tubes. The containers can be formed from a variety of materials, such as glass or plastic. The label on the container should indicate the agent is used for treating or prevention of one or more conditions of the disease. The label can also indicate directions for administration and so on.

[0082] In addition to the container described above, a kit including a pharmaceutical agent or composition of the present invention can optionally further include a second container housing a pharmaceutically-acceptable diluent. It can further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

[0083] The pharmaceutical agents or compositions can, if desired, be presented in a pack or dispenser device which can contain one or more unit dosage forms containing the active ingredient. The pack can, for example, include metal or plastic foil, such as a blister pack. The pack or dispenser device can be accompanied by instructions for administration.

[0084] (1) Pharmaceutical agents or compositions containing the peptides as the active ingredient

The peptides of this invention can be administered directly as a pharmaceutical agent or composition, or if necessary, that has been formulated by conventional formulation methods. In the latter case, in addition to the peptides of this invention, carriers, excipients, and such that are ordinarily used for drugs can be included as appropriate without particular limitations. Examples of such carriers are sterilized water, physiological saline, phosphate buffer, culture fluid and such. Furthermore, the pharmaceutical agents or compositions can contain as necessary, stabilizers, suspensions, preservatives, surfactants and such. The pharmaceutical agents or compositions of this invention can be used for anticancer purposes.

[0085] The peptides of this invention can be prepared as a combination, composed of two or more of peptides of the present invention, to induce CTLs in vivo. The peptide combination can take the form of a cocktail or can be conjugated to each other using standard techniques. For example, the peptides can be chemically linked or expressed as a single fusion polypeptide sequence. The peptides in the combination can be the same or different. By administering the peptides of this invention, the peptides are presented at a high density by the HLA antigens on APCs, then CTLs that specifically react toward the complex formed between the displayed peptide and the HLA antigen are induced. Alternatively, APCs that present any of the peptides of this invention on their cell surface, which may be obtained by stimulating APCs (e.g., DCs) derived from a subject with the peptides of this invention, may be administered to the subject, and as a result, CTLs are induced in the subject and aggressiveness towards the cancer cells, such as bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophagus cancer, gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma and SCC cells, can be increased.

[0086] The pharmaceutical agents or compositions for the treatment and/or prevention of cancer or tumor, which include a peptide of this invention as the active ingredient, can also include an adjuvant known to effectively cellular immunity. Alternatively, the pharmaceutical agents or compositions can be administered with other active ingredients or administered by formulation into granules. An adjuvant refers to a compound that enhances the immune response against the protein when administered

together (or successively) with the protein having immunological activity. Adjuvants contemplated herein include those described in the literature (Clin Microbiol Rev 1994, 7: 277-89). Examples of suitable adjuvants include aluminum phosphate, aluminum hydroxide, alum, cholera toxin, salmonella toxin, and such, but are not limited thereto.

- [0087] Furthermore, liposome formulations, granular formulations in which the peptide is bound to few-micrometers diameter beads, and formulations in which a lipid is bound to the peptide may be conveniently used.
- [0088] In some embodiments, the pharmaceutical agents or compositions of the present invention may further include a component which primes CTLs. Lipids have been identified as agents capable of priming CTLs in vivo against viral antigens. For example, palmitic acid residues can be attached to the epsilon- and alpha-amino groups of a lysine residue and then linked to a peptide of the present invention. The lipidated peptide can then be administered either directly in a micelle or particle, incorporated into a liposome, or emulsified in an adjuvant. As another example of lipid priming of CTLs responses, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine (P3CSS) can be used to prime CTL when covalently attached to an appropriate peptide (see, e.g., Deres et al., Nature 1989, 342: 561-4).
- [0089] The method of administration can be oral, intradermal, subcutaneous, intravenous injection, or such, and systemic administration or local administration to the vicinity of the targeted sites. The administration can be performed by single administration or boosted by multiple administrations. The dose of the peptides of this invention can be adjusted appropriately according to the disease to be treated, age of the patient, weight, method of administration, and such, and is ordinarily 0.001 mg to 1000 mg, for example, 0.001 mg to 1000 mg, for example, 0.1 mg to 10 mg, and can be administered once in a few days to few months. One skilled in the art can appropriately select a suitable dose.
- [0090] (2) Pharmaceutical agents or compositions containing polynucleotides as the active ingredient
- The pharmaceutical agents or compositions of the present invention can also contain nucleic acids encoding the peptides disclosed herein in an expressible form. Herein, the phrase "in an expressible form" means that the polynucleotide, when introduced into a cell, will be expressed in vivo as a polypeptide that induces anti-tumor immunity. In an exemplified embodiment, the nucleic acid sequence of the polynucleotide of interest includes regulatory elements necessary for expression of the polynucleotide. The polynucleotide(s) can be equipped so to achieve stable insertion into the genome of the target cell (see, e.g., Thomas KR & Capecchi MR, Cell 1987, 51: 503-12 for a description of homologous recombination cassette vectors). See, e.g., Wolff et al.,

Science 1990, 247: 1465-8; U.S. Patent Nos. 5,580,859; 5,589,466; 5,804,566; 5,739,118; 5,736,524; 5,679,647; and WO 98/04720. Examples of DNA-based delivery technologies include "naked DNA", facilitated (bupivacaine, polymers, peptide-mediated) delivery, cationic lipid complexes, and particle-mediated ("gene gun") or pressure-mediated delivery (see, e.g., U.S. Patent No. 5,922,687).

- [0091] The peptides of the present invention can also be expressed by viral or bacterial vectors. Examples of expression vectors include attenuated viral hosts, such as vaccinia or fowlpox. This approach involves the use of vaccinia virus, e.g., as a vector to express nucleotide sequences that encode the peptide. Upon introduction into a host, the recombinant vaccinia virus expresses the immunogenic peptide, and thereby elicits an immune response. Vaccinia vectors and methods useful in immunization protocols are described in, e.g., U.S. Patent No. 4,722,848. Examples of another vector include BCG (Bacille Calmette Guerin). BCG vectors are described in Stover et al., *Nature* 1991, 351: 456-60. A wide variety of other vectors useful for therapeutic administration or immunization e.g., adeno and adeno-associated virus vectors, retroviral vectors, *Salmonella typhi* vectors, detoxified anthrax toxin vectors, and the like, will be apparent. See, e.g., Shata et al., *Mol Med Today* 2000, 6: 66-71; Shedlock et al., *J Leukoc Biol* 2000, 68: 793-806; Hipp et al., *In Vivo* 2000, 14: 571-85.
- [0092] Delivery of a polynucleotide into a subject can be either direct, in which case the subject is directly exposed to a polynucleotide-carrying vector, or indirect, in which case, cells are first transformed with the polynucleotide of interest *in vitro*, then the cells are transplanted into the subject. These two approaches are known, respectively, as *in vivo* and *ex vivo* gene therapies.
- [0093] For general reviews of the methods of gene therapy, see Goldspiel et al., *Clinical Pharmacy* 1993, 12: 488-505; Wu and Wu, *Biotherapy* 1991, 3: 87-95; Tolstoshev, *Ann Rev Pharmacol Toxicol* 1993, 33: 573-96; Mulligan, *Science* 1993, 260: 926-32; Morgan & Anderson, *Ann Rev Biochem* 1993, 62: 191-217; Trends in Biotechnology 1993, 11(5): 155-215). Methods commonly known in the art of recombinant DNA technology which can also be used for the present invention are described in eds. Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, 1993; and Krieger, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY, 1990.
- [0094] The method of administration can be oral, intradermal, subcutaneous, intravenous injection, or such, and systemic administration or local administration to the vicinity of the targeted sites finds use. The administration can be performed by single administration or boosted by multiple administrations. The dose of the polynucleotide in the suitable carrier or cells transformed with the polynucleotide encoding the peptides of this invention can be adjusted appropriately according to the disease to be treated, age

of the patient, weight, method of administration, and such, and is ordinarily 0.001 mg to 1000 mg, for example, 0.001 mg to 1000 mg, for example, 0.1 mg to 10 mg, and can be administered once every a few days to once every few months. One skilled in the art can appropriately select the suitable dose.

[0095] X. Methods using the peptides, exosomes, APCs and CTLs

The peptides of the present invention and polynucleotides encoding such peptides can be used for inducing APCs and CTLs. The exosomes and APCs of the present invention can be also used for inducing CTLs. The peptides, polynucleotides, exosomes and APCs can be used in combination with any other compounds so long as the compounds do not inhibit their CTL inducibility. Thus, any of the aforementioned pharmaceutical agents or compositions of the present invention can be used for inducing CTLs, and in addition thereto, those including the peptides and polynucleotides can be also be used for inducing APCs as discussed below.

[0096] (1) Method of inducing antigen-presenting cells (APCs)

The present invention provides methods of inducing APCs using the peptides of this invention or polynucleotides encoding the peptides. The induction of APCs can be performed as described above in section "VI. Antigen-presenting cells". This invention also provides a method for inducing APCs having a high level of CTL inducibility, the induction of which has been also mentioned under the item of "VI. Antigen-presenting cells", supra.

[0097] Preferably, the methods for inducing APCs include at least one step selected from among:

- a: contacting APCs with the peptides of the present invention, and
- b: introducing the polypeptides of the present invention in an expressible form into APCs.

[0098] Such methods for inducing APCs are preferably performed in vitro or ex vivo. When the methods performed in vitro or ex vivo, APCs to be induced may be obtained from a subject to be treated or others whose HLA antigens are the same as the subject.

[0099] (2) Method of inducing CTLs

Furthermore, the present invention provides methods for inducing CTLs using the peptides of this invention, polynucleotides encoding the peptides, or exosomes or APCs presenting the peptides.

[0100] The present invention also provides methods for inducing CTLs using a polynucleotide encoding a polypeptide that is capable of forming a T cell receptor (TCR) subunit recognizing (i.e., binding to) a complex of the peptides of the present invention and HLA antigens on a cell surface. Preferably, the methods for inducing CTLs include at least one step selected from among:

- a: contacting a CD8-positive T cell with an antigen-presenting cell and/or an

exosome that presents on its surface a complex of an HLA antigen and a peptide of the present invention, and

b: introducing a polynucleotide encoding a polypeptide that is capable of forming a TCR subunit recognizing a complex of a peptide of the present invention and an HLA antigen into a CD8 positive T cell.

[0101] When the peptides of the present invention are administered to a subject, CTLs are induced in the body of the subject, and the strength of the immune response targeting the tumor-associated endothelia is enhanced. Alternatively, the peptides and polynucleotides encoding the peptides can be used for an ex vivo therapeutic method, in which subject-derived APCs and CD8-positive cells, or peripheral blood mononuclear leukocytes are contacted (stimulated) with the peptides of this invention in vitro, and after inducing CTLs, the activated CTL cells are returned to the subject. For example, the method can include steps of:

a: collecting APCs from subject,

b: contacting with the APCs of step a, with the peptide,

c: mixing the APCs of step b with CD8⁺ T cells, and co-culturing for inducing CTLs, and

d: collecting CD8⁺ T cells from the co-culture of step c.

[0102] Alternatively, according to the present invention, use of the peptides of this invention for manufacturing a pharmaceutical agent or composition inducing CTLs is provided. In addition, the present invention provides a method or process for manufacturing a pharmaceutical agent or composition inducing CTLs, wherein the method includes the step of admixing or formulating the peptide of the present invention with a pharmaceutically acceptable carrier. Further, the present invention also provides the peptide of the present invention for inducing CTLs. The CD8⁺ T cells having cytotoxic activity obtained by step d can be administered to the subject as a vaccine. The APCs to be mixed with the CD8⁺ T cells in above step c can also be prepared by transferring genes coding for the present peptides into the APCs as detailed above in section "VI. Antigen-presenting cells"; but are not limited thereto. Accordingly, any APCs or exosomes which effectively presents the present peptides to the T cells can be used for the present method.

[0103] The following examples are presented to illustrate the present invention and to assist one of ordinary skill in making and using the same. The examples are not intended in any way to otherwise limit the scope of the invention.

Examples

[0104] Materials and Methods

Cell lines

A24 lymphoblastoid cell line (A24LCL) was established by transformation with Epstein-bar virus into HLA-A24 positive human B lymphocyte. T2 (HLA-A2), human B-lymphoblastoid cell line, and COS7 were purchased from ATCC.

[0105] Candidate selection of peptides derived from MELK

9-mer and 10-mer peptides derived from MELK that bind to HLA-A*2402 and HLA-A*0201 molecules were predicted using binding prediction software "BIMAS" (http://www-bimas.cit.nih.gov/molbio/hla_bind), which algorithms had been described by Parker KC et al. (J Immunol 1994, 152(1): 163-75) and Kuzushima K et al. (Blood 2001, 98(6): 1872-81). These peptides were synthesized by Sigma (Sapporo, Japan) or Biosynthesis Inc. (Lewisville, TX) according to a standard solid phase synthesis method and purified by reversed phase high performance liquid chromatography (HPLC). The purity (>90%) and the identity of the peptides were determined by analytical HPLC and mass spectrometry analysis, respectively. Peptides were dissolved in dimethylsulfoxide (DMSO) at 20 mg/ml and stored at -80 degrees C.

[0106] In vitro CTL Induction

Monocyte-derived dendritic cells (DCs) were used as antigen-presenting cells (APCs) to induce cytotoxic T lymphocyte (CTL) responses against peptides presented on human leukocyte antigen (HLA). DCs were generated in vitro as described elsewhere (Nakahara S et al., Cancer Res 2003 Jul 15, 63(14): 4112-8). Specifically, peripheral blood mononuclear cells (PBMCs) isolated from a normal volunteer (HLA-A*2402 or HLA-A*0201 positive) by Ficoll-Plaque (Pharmacia) solution were separated by adherence to a plastic tissue culture dish (Becton Dickinson) so as to enrich them as the monocyte fraction. The monocyte-enriched population was cultured in the presence of 1000 U/ml of granulocyte-macrophage colony-stimulating factor (GM-CSF) (R&D System) and 1000 U/ml of interleukin (IL)-4 (R&D System) in AIM-V Medium (Invitrogen) containing 2% heat-inactivated autologous serum (AS). After 7 days of culture, the cytokine-induced DCs were pulsed with 20 microgram/ml of each of the synthesized peptides in the presence of 3 microgram/ml of beta2-microglobulin for 3 hrs at 37 degrees C in AIM-V Medium. The generated cells appeared to express DC-associated molecules, such as CD80, CD83, CD86 and HLA class II, on their cell surfaces (data not shown). These peptide-pulsed DCs were then inactivated by Mitomycin C (MMC) (30 microgram/ml for 30 min) and mixed at a 1:20 ratio with autologous CD8+ T cells, obtained by positive selection with CD8 Positive Isolation Kit (Dyna). These cultures were set up in 48-well plates (Corning); each well contained 1.5×10^4 peptide-pulsed DCs, 3×10^5 CD8+ T cells and 10 ng/ml of IL-7 (R&D System) in 0.5 ml of AIM-V/2% AS medium. Three days later, these cultures were supplemented with IL-2 (CHIRON) to a final concentration of 20 IU/ml. On days 7 and 14, the T cells were further stimulated with the autologous peptide-

pulsed DCs. The DCs were prepared each time by the same way described above. CTL was tested against peptide-pulsed A24LCL or T2 cells after the 3rd round of peptide stimulation on day 21 (Tanaka H et al., Br J Cancer 2001 Jan 5, 84(1): 94-9; Umano Y et al., Br J Cancer 2001 Apr 20, 84(8): 1052-7; Uchida N et al., Clin Cancer Res 2004 Dec 15, 10(24): 8577-86; Suda T et al., Cancer Sci 2006 May, 97(5): 411-9; Watanabe T et al., Cancer Sci 2005 Aug, 96(8): 498-506).

[0107] CTL Expansion Procedure

CTLs were expanded in culture using the method similar to the one described by Riddell et al. (Walter EA et al., N Engl J Med 1995 Oct 19, 333(16): 1038-44; Riddell SR et al., Nat Med 1996 Feb, 2(2): 216-23). A total of 5×10^4 CTLs were suspended in 25 ml of AIM-V/5% AS medium with 2 kinds of human B-lymphoblastoid cell lines, inactivated by MMC, in the presence of 40 ng/ml of anti-CD3 monoclonal antibody (Pharmingen). One day after initiating the cultures, 120 IU/ml of IL-2 were added to the cultures. The cultures were fed with fresh AIM-V/5% AS medium containing 30 IU/ml of IL-2 on days 5, 8 and 11 (Tanaka H et al., Br J Cancer 2001 Jan 5, 84(1): 94-9; Umano Y et al., Br J Cancer 2001 Apr 20, 84(8): 1052-7; Uchida N et al., Clin Cancer Res 2004 Dec 15, 10(24): 8577-86; Suda T et al., Cancer Sci 2006 May, 97(5): 411-9; Watanabe T et al., Cancer Sci 2005 Aug, 96(8): 498-506).

[0108] Establishment of CTL clones

The dilutions were made to have 0.3, 1, and 3 CTLs/well in 96 round-bottomed micro titer plate (Nalge Nunc International). CTLs were cultured with 1×10^4 cells/well of 2 kinds of human B-lymphoblastoid cell lines, 30ng/ml of anti-CD3 antibody, and 125 U/ml of IL-2 in a total of 150 microliter/well of AIM-V Medium containing 5%AS. 50 microliter/well of IL-2 were added to the medium 10 days later so to reach a final concentration of 125 U/ml IL-2. CTL activity was tested on the 14th day, and CTL clones were expanded using the same method as described above (Uchida N et al., Clin Cancer Res 2004 Dec 15, 10(24): 8577-86; Suda T et al., Cancer Sci 2006 May, 97(5): 411-9; Watanabe T et al., Cancer Sci 2005 Aug, 96(8): 498-506).

[0109] Specific CTL activity

To examine specific CTL activity, interferon (IFN)-gamma enzyme-linked immunospot (ELISPOT) assay and IFN-gamma enzyme-linked immunosorbent assay (ELISA) were performed. Specifically, peptide-pulsed A24LCL or T2 (1×10^4 /well) was prepared as stimulator cells. Cultured cells in 48 wells were used as responder cells. IFN-gamma ELISPOT assay and IFN-gamma ELISA assay were performed under manufacture procedure.

[0110] Results

Over expression in cancers

The global gene expression profile data obtained from various cancers using cDNA-

microarray revealed that MELK (GenBank Accession No. NM_014791; SEQ ID No.93) expression was elevated. MELK expression was validly elevated in 29 out of 29 bladder cancer, 31 out of 34 breast cancer, 14 out of 15 cervical cancer, 11 out of 11 cholangiocellular carcinoma, 10 out of 13 CML, 12 out of 15 colorectal cancer, 2 out of 2 endometriosis, 19 out of 42 esophagus cancer, 5 out of 6 gastric cancer, 4 out of 4 liver cancer, 11 out of 11 NSCLC, 13 out of 14 lymphoma, 14 out of 18 osteosarcoma, 3 out of 6 ovarian cancer, 2 out of 2 pancreatic cancer, 18 out of 21 prostate cancer, 5 out of 6 renal carcinoma and 15 out of 15 small cell lung cancer as compared with corresponding normal tissue. (Table 1).

[0111] [Table 1]

Ratio of cases observed up-regulation of *MELK* in cancerous tissue as compared with normal corresponding tissue

Cancer	Ratio
Bladder Cancer	29/29
Breast Cancer	31/34
Cervical Cancer	14/15
Cholangiocellular Carcinoma	11/11
CML	10/13
Colorectal cancer	12/15
Endometriosis	2/2
Esophagus cancer	19/42
Gastric cancer	5/6
Liver cancer	4/4
non-small cell lung cancer	11/11
Lymphoma	13/14
Osteosarcoma	14/18
Ovarian cancer	3/6
Pancreatic cancer	2/2
Prostate cancer	18/21
Renal Carcinoma	5/6
Small cell Lung Cancer	15/15

[0112] Prediction of HLA-A24 and HLA-A2 binding peptides derived from MELK

Table 2 shows the HLA-A24 and HLA-A2 binding peptides of MELK in order of highest binding affinity. A total of 34 peptides with potential HLA-A24 binding ability were selected and examined to determine the epitope peptides (Table 2a), and a total of 58 peptides having potential HLA-A2 binding ability were similarly selected and examined to determine the epitope peptides (Table 2b and c).

[0113]

[Table 2a]

HLA-A24 binding 9mer and 10mer peptides derived from MELK

Start Position	Amino acid sequence	Binding Score	SEQ ID NO.
199	LYVLMCGFL	300	1
96	DYIISQDRL	300	2
560	HYNVTITRL	300	3
373	DYDWCEDDL	200	4
9	KYYELHETI	144	5
87	EYCPGGELF	120	6
637	VYKRLVEDI	60	7
610	QFELEVCQL	30	8
588	DFVQKGYTL	30	9
526	VFGSLERGL	24	10
567	RLVNPDQLL	14.4	11
603	DFGKVTMQF	14	12
522	KGAKVFGSL	13.44	13
326	RGKPVRLRL	13.44	14
450	KNQHKREIL	12	15
230	KWLSPSSIL	12	16
395	KYWTESNGV	12	17
502	RCRSVELDL	11.2	18
145	KLKLIDFGL	11.2	19
574	LLNEIMSIL	10.08	20
78	TANKIFMVL	10.08	21
225	KYDVPKWLS	10	22
637	VYKRLVEDIL	280	23
309	QYDHLTATYL	200	24
142	EYHKLKLIDF	100	25
139	LFDEYHKLKL	26.4	26
532	RGLDKVITVL	20.16	27
230	KWLSPSSILL	12	28
55	KTEIEALKNL	12	29
295	RNNRQTMEDL	12	30
223	RGKYDVPKWL	11.2	31
632	KGDAWVYKRL	11.2	32
266	DYNYPVEWQS	10.5	33
463	RYTTPSKARN	10	34

[0114]

[Table 2b]

HLA-A2 binding 9mer peptides derived from MELK

Start Position	Amino acid sequence	Binding Score	SEQ ID NO.
238	LLLQQMLQV	1006.209	35
138	LLFDEYHKL	826.372	36
263	IMQDYNYPV	350.117	37
103	RLSEEETRV	285.163	38
147	KLIDFGLCA	270.905	39
206	FLPFDDDNV	156.77	40
574	LLNEIMSI	140.409	41
220	KIMRGKYDV	123.846	42
115	QIVSAVAYV	120.977	43
308	WQYDHLTAT	104.878	44
231	WLSPSSILL	98.267	45
193	WSMGILLYV	97.752	46
573	QLLNEIMSI	88.783	47
317	YLLLLAKKA	84.555	48
306	SLWQYDHLT	61.852	49
558	KLHYNVTTT	59.989	50
256	NLLNHPWIM	56.523	51
81	KIFMVLEYC	54.833	52
31	ILTGEMVAI	40.792	53
46	TLGSDLPRI	23.995	54
607	VTMQFELEV	23.089	55
279	FIHLDDDCV	21.556	56
171	SLAYAAPEL	21.362	57
567	RLVNPQQLL	21.362	58
425	NVYTPKSAV	19.475	59
71	QLYHVLETA	17.917	60
407	SLTPALCRT	17.14	61
532	RGLDKVITV	15.841	62
145	KLKLIDFGL	15.178	63
194	SMGILLYVL	14.549	64
457	ILTPNRYT	13.935	65
236	SILLQQML	10.868	66
299	QTMEDLISL	10.352	67

[0115]

[Table 2c]

HLA-A2 binding 10mer peptides derived from MELK

Start Position	Amino acid sequence	Binding Score	SEQ ID NO.
237	ILLQQMLQV	1006.209	68
439	FMFPEPKTPV	854.949	69
581	ILPKKHVDFV	732.901	70
231	WLSPPSILL	226.014	71
262	WIMQDYNYPV	162.769	72
609	MQFELEVCQL	128.47	73
137	NLLFDEYHKL	118.561	74
114	RQIVSAVAYV	89.205	75
111	VVFRQIVSAV	88.043	76
578	IMSILPKKHV	85.394	77
198	LLYVLMCGFL	83.091	78
606	KVTMQFELEV	80.941	79
573	QLLNEIMSIL	74.536	80
640	RLVEDILSSC	46.848	81
306	SLWQYDHLTA	41.234	82
76	LETANKIFMV	30.67	83
617	QLQKPDVVGI	23.995	84
103	RLSEETRVV	23.383	85
411	ALCRTPANKL	21.362	86
273	WQSKNPFIHL	18.93	87
16	TIGTGGFAKV	18.17	88
243	MLQVDPKKRI	17.736	89
271	VEWQSKNPFI	15.743	90
636	WVYKRLVEDI	15.144	91
457	ILTPNR YTT	12.668	92

Start position indicates the number of amino acid residue from the N-terminal of MELK. Binding score is derived from "BIMAS".

[0116] CTL induction with the predicted peptides from MELK restricted with HLA-A*2402 or HLA-A*0201 and establishment for CTL lines stimulated with MELK derived peptides

CTLs for those peptides derived from MELK were generated according to the protocols as described in "Materials and Methods". Peptide specific CTL activity was determined by IFN-gamma ELISPOT assay (Figure 1a-i). It showed that #5 stimulated with MELK-A24-9-326 (SEQ ID NO: 14) (a), #3 with MELK-A24-9-78 (SEQ ID NO: 21) (b), #4 with MELK-A24-10-637 (SEQ ID NO: 23) (c), #4 with MELK-A24-10-532 (SEQ ID NO: 27) (d), #5 with MELK-A02-9-138 (SEQ ID NO: 36) (e), #3 with MELK-A02-9-193 (SEQ ID NO: 46) (f), #1 with MELK-A02-9-171 (SEQ ID NO: 57) (g), #2 with MELK-A02-9-71 (SEQ ID NO: 60) (h) and #2 with MELK-A02-9-532 (SEQ ID NO: 62) (i) demonstrated potent IFN-gamma production as

compared to the control wells. Furthermore, the cells in the positive well number #5 stimulated with SEQ ID NO: 14, #3 with SEQ ID NO: 21, #4 with SEQ ID NO: 23, #4 with SEQ ID NO: 27, #5 with SEQ ID NO: 36, #3 with SEQ ID NO: 46, #1 with SEQ ID NO: 57, #2 with SEQ ID NO: 60 and #2 with SEQ ID NO: 62 were expanded and established CTL lines. CTL activity of those CTL lines was determined by IFN-gamma ELISA assay (Figure 2a-i). All CTL lines demonstrated potent IFN-gamma production against the target cells pulsed with corresponding peptide as compared to target cells without peptide pulse. On the other hand, no CTL lines could be established by stimulation with other peptides shown in Table 2, despite those peptide had possible binding activity with HLA-A*2402 or HLA-A*0201 (data not shown). Consequently, only nine peptides out of 92 candidate peptide derived from MELK could induce potent CTL lines.

[0117] Establishment of CTL clones against MELK specific peptides

CTL clones were established by limiting dilution from CTL lines as described in "Materials and Methods", and IFN-gamma production from CTL clones against target cells pulsed peptide were determined by IFN-gamma ELISA assay. Potent IFN-gamma productions were determined from CTL clones stimulated with SEQ ID NO: 27 in Figure 3.

[0118] Specific CTL activity against target cells presenting the peptide endogenously processed from MELK with HLA-A*0201

The established CTL lines raised against these peptides were examined for their ability to recognize target cells presenting the candidate peptide endogenously processed from MELK with HLA-A*0201 molecule. Specific CTL activity against COS7 cells which transfected with both the full length of MELK and HLA-A*0201 molecule gene (a specific model for the target cells that express MELK and HLA-A*0201 gene) was tested using the CTL lines raised by corresponding peptide as the effector cells. COS7 cells transfected with either full length of MELK genes or HLA-A*0201 were prepared as control. In Figure 4, the CTLs stimulated with SEQ ID NO: 36 (a) and SEQ ID NO: 57 (b) showed potent CTL activity against COS7 cells expressing both MELK and HLA-A02. On the other hand, no significant specific CTL activity was detected against the controls. Thus, these data clearly demonstrate that the peptides having the amino acid sequences of SEQ ID NO: 36 and SEQ ID NO: 57 are naturally presented on the target cells with HLA-A*0201 molecule and are recognized by the CTLs. These results indicated that those two peptides derived from MELK may be available to apply the cancer vaccines for patients with MELK expressing tumors.

[0119] Homology analysis of antigen peptides

The CTLs stimulated with MELK-A24-9-326 (SEQ ID NO: 14), MELK-A24-9-78 (SEQ ID NO: 21), MELK-A24-10-637 (SEQ ID NO: 23), MELK-A24-10-532 (SEQ

ID NO: 27), MELK-A02-9-138 (SEQ ID NO: 36), MELK-A02-9-193 (SEQ ID NO: 46), MELK-A02-9-171 (SEQ ID NO: 57), MELK-A02-9-71 (SEQ ID NO: 60) and MELK-A02-9-532 (SEQ ID NO: 62) showed significant and specific CTL activity. This result may be due to the fact that the sequences of MELK-A24-9-326 (SEQ ID NO: 14), MELK-A24-9-78 (SEQ ID NO: 21), MELK-A24-10-637 (SEQ ID NO: 23), MELK-A24-10-532 (SEQ ID NO: 27), MELK-A02-9-138 (SEQ ID NO: 36), MELK-A02-9-193 (SEQ ID NO: 46), MELK-A02-9-171 (SEQ ID NO: 57), MELK-A02-9-71 (SEQ ID NO: 60) and MELK-A02-9-532 (SEQ ID NO: 62) are homologous to peptides derived from other molecules that are known to sensitize the human immune system. To exclude this possibility, homology analyses were performed for these peptide sequences using as queries the BLAST algorithm (<http://www.ncbi.nlm.nih.gov/blast/blast.cgi>) which revealed no sequence with significant homology. The results of homology analyses indicate that the sequences of MELK-A24-9-326 (SEQ ID NO: 14), MELK-A24-9-78 (SEQ ID NO: 21), MELK-A24-10-637 (SEQ ID NO: 23), MELK-A24-10-532 (SEQ ID NO: 27), MELK-A02-9-138 (SEQ ID NO: 36), MELK-A02-9-193 (SEQ ID NO: 46), MELK-A02-9-171 (SEQ ID NO: 57), MELK-A02-9-71 (SEQ ID NO: 60) and MELK-A02-9-532 (SEQ ID NO: 62) are unique and thus, there is little possibility, to our best knowledge, that these molecules raise unintended immunologic responses to some unrelated molecules.

- [0120] In conclusion, novel HLA-A24 and HLA-A02 epitope peptides derived from MELK were identified and demonstrated to be applicable for cancer immunotherapy.

Industrial Applicability

- [0121] The present invention describes new TAAs, particularly those derived from MELK which induce potent and specific anti-tumor immune responses and have applicability to a wide array of cancer types. Such TAAs warrant further development as peptide vaccines against diseases associated with MELK, e.g. cancers such as breast cancer, bladder cancer, cervical cancer, cholangiocellular carcinoma, CML, colorectal cancer, endometriosis, esophagus cancer, gastric cancer, liver cancer, NSCLC, lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma and SCC.
- [0122] While the invention is herein described in detail and with reference to specific embodiments thereof, it is to be understood that the foregoing description is exemplary and explanatory in nature and is intended to illustrate the invention and its preferred embodiments. Through routine experimentation, one skilled in the art will readily recognize that various changes and modifications can be made therein without departing from the spirit and scope of the invention, the metes and bounds of which are

defined by the appended claims.

CLAIMS:

1. An isolated peptide of less than 15 amino acids, wherein said peptide comprises an amino acid sequence selected from the group consisting of:
 - (a) SEQ ID NOs:36, 46, 57, 60 and 62; and
 - (b) SEQ ID NOs:36, 46, 57, 60 and 62, in which 1 or 2 amino acids are substituted, inserted, deleted and/or added, and has cytotoxic T lymphocyte (CTL) inducibility.
2. The peptide as set forth in Claim 1, wherein the peptide is a nonapeptide or a decapeptide.
3. The peptide as set forth in Claim 1 or 2, wherein the peptide has one or both of the following characteristics:
 - (a) the second amino acid from the N-terminus of the amino acid sequence of SEQ ID NO: 36, 46, 57, 60 or 62 is substituted with an amino acid selected from the group of leucine and methionine; and
 - (b) the C-terminal amino acid of the amino acid sequence of SEQ ID NO: 36, 46, 57, 60 or 62 is substituted with an amino acid selected from the group of valine and leucine.
4. The peptide as set forth in Claim 1 or 2, wherein said peptide consists of an amino acid sequence selected from the group consisting of SEQ ID NOs:36, 46, 57, 60 and 62.
5. An isolated polynucleotide encoding a peptide as set forth in any one of Claims 1 to 4.
6. A pharmaceutical composition for the treatment and/or prophylaxis of cancer, and/or the prevention of a postoperative recurrence thereof, wherein the pharmaceutical composition comprises an active ingredient selected from the group consisting of:
 - (a) one or more peptides as set forth in any one of Claims 1 to 4;
 - (b) one or more polynucleotides encoding the peptide in an expressible form;

- (c) one or more antigen-presenting cells and/or exosomes, which antigen-presenting cells and exosomes present a complex formed between a HLA antigen and a peptide as set forth in any one of Claims 1 to 4 on its surface;
- (d) one or more CTLs induced against a peptide as set forth in any one of Claims 1 to 4; and
- (e) combinations thereof.

7. The pharmaceutical composition as set forth in Claim 6, wherein the pharmaceutical composition is formulated for the administration to a subject whose HLA antigen is HLA-A02.

8. The pharmaceutical composition as set forth in Claim 6, wherein the pharmaceutical composition is formulated for the treatment of a cancer selected from the group consisting of bladder cancer, breast cancer, cervical cancer, cholangiocellular carcinoma, chronic myeloid leukemia (CML), colorectal cancer, endometriosis, esophagus cancer, gastric cancer, liver cancer, non-small cell lung cancer (NSCLC), lymphoma, osteosarcoma, ovarian cancer, pancreatic cancer, prostate cancer, renal carcinoma and small cell lung cancer.

9. The pharmaceutical composition of Claim 6, wherein said pharmaceutical composition is formulated as a vaccine.

10. An *in vitro* method for inducing an antigen-presenting cell with high CTL inducibility, wherein the method comprises the step of contacting an antigen-presenting cell with a peptide as set forth in any one of Claims 1 to 4, or the step of introducing a polynucleotide encoding the peptide in an expressible form into an antigen-presenting cell.

11. An *in vitro* method for inducing CTL by using a peptide as set forth in any one of Claims 1 to 4.

12. The method for inducing CTL as set forth in Claim 11, wherein said method comprises the step selected from the group consisting of:
 - (a) contacting a CD8-positive T cell with an antigen-presenting cell and/or an exosome which present a complex formed between an HLA antigen and a peptide as set forth in any one of Claims 1 to 4 on its surface; and
 - (b) introducing a polynucleotide encoding a polypeptide which is capable of forming a TCR subunit recognizing a complex formed between an HLA antigen and a peptide as set forth in any one of Claims 1 to 4 into a CD8-positive T cell.
13. An isolated CTL which targets any of the peptides of Claims 1 to 4.
14. An isolated CTL which is induced by a peptide as set forth in any one of Claims 1 to 4.
15. The CTL as set forth in Claims 13 or 14, wherein said CTL is capable of recognizing a complex formed between an HLA antigen and a peptide as set forth in any one of Claims 1 to 4 on a cell.
16. The CTL as set forth in any one of Claims 13 to 15, which is induced by the method as set forth in Claim 12.
17. An isolated antigen-presenting cell that presents a complex formed between a HLA antigen and a peptide as set forth in any one of Claims 1 to 4 on its surface.
18. The antigen-presenting cell as set forth in Claim 17, which is induced by the method as set forth in Claim 10.
19. The antigen-presenting cell as set forth in Claim 17 or 18, wherein the HLA antigen is HLA-A02.

20. A composition for inducing an immune response against a cancer in a subject, wherein said composition comprises an active ingredient selected from the group consisting of:

- (a) one or more peptides as set forth in any one of Claims 1 to 4;
- (b) one or more polynucleotides encoding the peptide in an expressible form;
- (c) one or more antigen-presenting cells and/or exosomes, which antigen-presenting cells and exosomes present a complex formed between a HLA antigen and a peptide as set forth in any one of Claims 1 to 4 on its surface;
- (d) one or more CTLs induced against a peptide as set forth in any one of Claims 1 to 4; and
- (e) combinations thereof.

21. A composition for inducing a CTL, wherein said composition comprises an active ingredient selected from the group consisting of:

- (a) one or more peptides as set forth in any one of Claims 1 to 4;
 - (b) one or more polynucleotides encoding the peptide in an expressible form;
- and
- (c) one or more antigen-presenting cells and/or exosomes, which antigen-presenting cells and exosomes present a complex formed between a HLA antigen and a peptide as set forth in any one of Claims 1 to 4 on its surface.