

**NEW ISOINDOLINE OR ISOQUINOLINE COMPOUNDS, A PROCESS FOR THEIR
PREPARATION AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM**

The present invention relates to new isoindoline or isoquinoline compounds, to a process for their preparation and to pharmaceutical compositions containing them.

The compounds of the present invention are new and have very valuable pharmacological characteristics in the field of apoptosis and cancerology.

5 Apoptosis, or programmed cell death, is a physiological process that is crucial for embryonic development and maintenance of tissue homeostasis.


Apoptotic-type cell death involves morphological changes such as condensation of the nucleus, DNA fragmentation and also biochemical phenomena such as the activation of caspases which cause damage to key structural components of the cell, so inducing its
10 disassembly and death. Regulation of the process of apoptosis is complex and involves the activation or repression of several intracellular signalling pathways (Cory S. *et al.*, Nature Review Cancer, 2002, 2, 647-656).

Deregulation of apoptosis is involved in certain pathologies. Increased apoptosis is associated with neurodegenerative diseases such as Parkinson's disease, Alzheimer's
15 disease and ischaemia. Conversely, deficits in the implementation of apoptosis play a significant role in the development of cancers and their chemoresistance, in auto-immune diseases, inflammatory diseases and viral infections. Accordingly, absence of apoptosis is one of the phenotypic signatures of cancer (Hanahan D. *et al.*, Cell 2000, 100, 57-70).

The anti-apoptotic proteins of the Bcl-2 family are associated with numerous pathologies.
20 The involvement of proteins of the Bcl-2 family is described in numerous types of cancer, such as colon cancer, breast cancer, small-cell lung cancer, non-small-cell lung cancer, bladder cancer, ovarian cancer, prostate cancer, chronic lymphoid leukaemia, follicular lymphoma, myeloma, etc. Overexpression of the anti-apoptotic proteins of the Bcl-2 family is involved in tumorigenesis, in resistance to chemotherapy and in the clinical
25 prognosis of patients affected by cancer. There is, therefore, a therapeutic need for compounds that inhibit the anti-apoptotic activity of the proteins of the Bcl-2 family.

In addition to being new, the compounds of the present invention have pro-apoptotic properties making it possible to use them in pathologies involving a defect in apoptosis, such as, for example, in the treatment of cancer and of auto-immune and immune system diseases.

5 The present invention relates more especially to compounds of formula (I):



wherein:

- ♦ Het represents a heteroaryl group,
- ♦ T represents a hydrogen atom, a linear or branched (C₁-C₆)alkyl group optionally substituted by one to three halogen atoms, an alkyl(C₁-C₄)-NR₁R₂ group or an alkyl(C₁-C₄)-OR₆ group,
- ♦ R₁ and R₂ independently of one another represent a hydrogen atom or a linear or branched (C₁-C₆)alkyl group, or R₁ and R₂ form with the nitrogen atom carrying them a heterocycloalkyl group,
- ♦ R₃ represents a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl group, a (C₃-C₁₀)cycloalkyl-(C₁-C₆)alkyl group wherein the alkyl group may be linear or branched, a heterocycloalkyl group, an aryl group or a heteroaryl group, it being understood that one or more carbon atoms of the groups defined hereinbefore, or carbon atoms of their possible substituents, may be deuterated,

- ◆ R_4 represents an aryl, heteroaryl, cycloalkyl or linear or branched (C_1-C_6) alkyl group, it being understood that one or more carbon atoms of the groups defined hereinbefore, or carbon atoms of their possible substituents, may be deuterated,
- ◆ R_5 represents a hydrogen atom or a halogen atom, a linear or branched (C_1-C_6) alkyl group, or a linear or branched (C_1-C_6) alkoxy group,
- ◆ R_6 represents a hydrogen atom or a linear or branched (C_1-C_6) alkyl group,
- ◆ R_7 represents a group selected from R'_7 , R'_7-CO- , $R'_7-O-CO-$, $NR'_7R''_7-CO-$, R'_7-SO_2- , $R'_7-NR''_7-SO_2-$ wherein R'_7 and R''_7 independently of one another represent a hydrogen atom, a linear or branched (C_1-C_6) alkyl group, a linear or branched (C_2-C_6) alkenyl group, a linear or branched (C_2-C_6) alkynyl group, a cycloalkyl, a heterocycloalkyl, an aryl group, or a heteroaryl,
- ◆ R_8 and R_9 represent, independently of one another, an oxo group or a halogen atom,
- ◆ p and p' are, independently of one another, integers equal to 0, 1, 2, 3 or 4,
- ◆ q and q' are, independently of one another, integers equal to 1, 2 or 3,

it being understood that when compound of formula (I) contains a hydroxy group, this latter group may be optionally substituted by one of the following groups: $-PO(OM)(OM')$, $-PO(OM)(O^+M_1)$, $-PO(O^+M_1)(O^+M_2)$, $-PO(O^-)(O^-)M_3^{2+}$, $-PO(OM)(O[CH_2CH_2O]_nCH_3)$, or $-PO(O^+M_1)(O[CH_2CH_2O]_nCH_3)$, wherein M and M' independently of one another represent a hydrogen atom, a linear or branched (C_1-C_6) alkyl group, a linear or branched (C_2-C_6) alkenyl group, a linear or branched (C_2-C_6) alkynyl group, a cycloalkyl or a heterocycloalkyl, both of them being composed of from 5 to 6 ring members, while M_1^+ and M_2^+ independently of one another represent a pharmaceutically acceptable monovalent cation, M_3^{2+} represents a pharmaceutically acceptable divalent cation and n is an integer comprised between 1 to 5,

it also being understood that:

- "aryl" means a phenyl, naphthyl, biphenyl or indenyl group,
- "heteroaryl" means any mono- or bi-cyclic group composed of from 5 to 10 ring members, having at least one aromatic moiety and containing from 1 to 4 hetero atoms selected from oxygen, sulphur and nitrogen (including quaternary nitrogens),
- "cycloalkyl" means any mono- or bi-cyclic non-aromatic carbocyclic group


containing from 3 to 10 ring members, which may include fused, bridged or spiro ring systems,

- "heterocycloalkyl" means any mono- or bi-cyclic non-aromatic carbocyclic group, composed of from 3 to 10 ring members, and containing from one to 3 hetero atoms selected from oxygen, sulphur, SO, SO₂ and nitrogen, it being understood that bicyclic group may be fused or spiro type,

it being possible for the aryl, heteroaryl, cycloalkyl and heterocycloalkyl groups so defined and the alkyl, alkenyl, alkynyl, alkoxy, to be substituted by from 1 to 3 groups selected from: optionally substituted linear or branched (C₁-C₆)alkyl ; optionally substituted linear or branched (C₂-C₆)alkenyl group; optionally substituted linear or branched (C₂-C₆)alkynyl group; (C₃-C₆)spiro; optionally substituted linear or branched (C₁-C₆)alkoxy; (C₁-C₆)alkyl-S-; hydroxyl; oxo (or *N*-oxide where appropriate); nitro; cyano; -COOR'; -OCOR'; -NR'R"; R'CONR''-; NR'R''CO-; linear or branched (C₁-C₆)polyhaloalkyl; trifluoromethoxy; (C₁-C₆)alkylsulphonyl; halogen; optionally substituted aryl; optionally substituted heteroaryl; optionally substituted aryloxy; optionally substituted arylthio; optionally substituted cycloalkyl or optionally substituted heterocycloalkyl, it being understood that R' and R" independently of one another represent a hydrogen atom, an optionally substituted linear or branched (C₁-C₆)alkyl group or an aryl group,

their enantiomers and diastereoisomers, and addition salts thereof with a pharmaceutically acceptable acid or base.

More preferably, the present invention relates to compounds of formula (IA):



wherein:

- ◆ W represents a C-A₃ group or a nitrogen atom,
 - ◆ X, Y and Z represent a carbon atom or a nitrogen atom, it being understood that only one of them represent a nitrogen atom, while the two others represent carbon atoms,
 - ◆ A₁, A₂ and A₃ independently of one another represent a hydrogen atom or a halogen atom, a linear or branched (C₁-C₆)polyhaloalkyl group, a linear or branched (C₁-C₆)alkyl group or a cycloalkyl group,
- or A₃ represents a hydrogen atom (when W represents a C-A₃ group) while A₁ and A₂ form together with the atoms carrying them an optionally substituted aromatic or non-aromatic ring Cy, composed of 5, 6 or 7 ring members, which may contain from one to 4 hetero atoms selected independently from oxygen, sulphur and nitrogen, it being understood that the nitrogen in question may be substituted by a group representing a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or a group -C(O)-O-Alk wherein Alk is a linear or branched (C₁-C₆)alkyl group,
- or W represents a nitrogen atom while A₁ and A₂ form together with the atoms carrying them an optionally substituted aromatic or non-aromatic ring Cy, composed of 5, 6 or 7 ring members, which may contain from one to 4 hetero atoms selected independently from oxygen, sulphur and nitrogen, it being

understood that the nitrogen in question may be substituted by a group representing a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or a group -C(O)-O-Alk wherein Alk is a linear or branched (C₁-C₆)alkyl group,

- 5 ♦ T represents a hydrogen atom, a linear or branched (C₁-C₆)alkyl group optionally substituted by one to three halogen atoms, an alkyl(C₁-C₄)-NR₁R₂ group or an alkyl(C₁-C₄)-OR₆ group,
- ♦ R₁ and R₂ independently of one another represent a hydrogen atom or a linear or branched (C₁-C₆)alkyl group,
 or R₁ and R₂ form with the nitrogen atom carrying them a heterocycloalkyl,
- 10 ♦ R₃ represents a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl group, a (C₃-C₁₀)cycloalkyl-(C₁-C₆)alkyl group wherein the alkyl group may be linear or branched, a heterocycloalkyl group, an aryl group or a heteroaryl group, it being understood that one or more carbon atoms of the groups defined
- 15 hereinbefore, or carbon atoms of their possible substituents, may be deuterated,
- ♦ R₄ represents an aryl, heteroaryl, cycloalkyl or linear or branched (C₁-C₆)alkyl group, it being understood that one or more carbon atoms of the groups defined hereinbefore, or carbon atoms of their possible substituents, may be deuterated,
- ♦ R₅ represents a hydrogen atom or a halogen atom, a linear or branched (C₁-C₆)alkyl group, or a linear or branched (C₁-C₆)alkoxy group,
- 20 ♦ R₆ represents a hydrogen atom or a linear or branched (C₁-C₆)alkyl group,
- ♦ R₇ represents a group selected from R'₇, R'₇-CO-, R'₇-O-CO-, NR'₇R''₇-CO-, R'₇-SO₂-, R'₇-NR''₇-SO₂- wherein R'₇ and R''₇ independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group, a linear or
- 25 branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl, a heterocycloalkyl, an aryl group, or a heteroaryl,
- ♦ R₈ and R₉ represent, independently of one another, an oxo group or a halogen atom,
- ♦ p and p' are, independently of one another, integers equal to 0, 1, 2, 3 or 4,
- ♦ q and q' are, independently of one another, integers equal to 1, 2 or 3,
- 30 it being understood that when compound of formula (I) contains a hydroxy group, this latter group may be optionally substituted by one of the following groups:

$-\text{PO}(\text{OM})(\text{OM}')$, $-\text{PO}(\text{OM})(\text{O}^-\text{M}_1^+)$, $-\text{PO}(\text{O}^-\text{M}_1^+)(\text{O}^-\text{M}_2^+)$, $-\text{PO}(\text{O}^-)(\text{O}^-)\text{M}_3^{2+}$,
 $-\text{PO}(\text{OM})(\text{O}[\text{CH}_2\text{CH}_2\text{O}]_n\text{CH}_3)$, or $-\text{PO}(\text{O}^-\text{M}_1^+)(\text{O}[\text{CH}_2\text{CH}_2\text{O}]_n\text{CH}_3)$, wherein M and M'
independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl
group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl
group, a cycloalkyl or a heterocycloalkyl, both of them being composed of from 5 to 6 ring
members, while M₁⁺ and M₂⁺ independently of one another represent a pharmaceutically
acceptable monovalent cation, M₃²⁺ represents a pharmaceutically acceptable divalent
cation and n is an integer comprised between 1 to 5,

it also being understood that:

- "aryl" means a phenyl, naphthyl, biphenyl or indenyl group,
- "heteroaryl" means any mono- or bi-cyclic group composed of from 5 to 10 ring
members, having at least one aromatic moiety and containing from 1 to 4 hetero
atoms selected from oxygen, sulphur and nitrogen (including quaternary nitrogens),
- "cycloalkyl" means any mono- or bi-cyclic non-aromatic carbocyclic group
containing from 3 to 10 ring members, which may include fused, bridged or spiro
ring systems,
- "heterocycloalkyl" means any mono- or bi-cyclic non-aromatic carbocyclic group,
composed of from 3 to 10 ring members, and containing from one to 3 hetero atoms
selected from oxygen, sulphur, SO, SO₂ and nitrogen, it being understood that
bicyclic group may be fused or spiro type,


it being possible for the aryl, heteroaryl, cycloalkyl and heterocycloalkyl groups so defined
and the alkyl, alkenyl, alkynyl, alkoxy, to be substituted by from 1 to 3 groups selected
from: optionally substituted linear or branched (C₁-C₆)alkyl ; optionally substituted linear
or branched (C₂-C₆)alkenyl group; optionally substituted linear or branched (C₂-C₆)alkynyl
group ; (C₃-C₆)spiro; optionally substituted linear or branched (C₁-C₆)alkoxy;
(C₁-C₆)alkyl-S-; hydroxyl; oxo (or N-oxide where appropriate); nitro; cyano; -COOR';
-OCOR'; -NR'R"; R'CONR''-; NR'R''CO-; linear or branched (C₁-C₆)polyhaloalkyl;
trifluoromethoxy; (C₁-C₆)alkylsulphonyl; halogen; optionally substituted aryl; optionally
substituted heteroaryl; optionally substituted aryloxy; optionally substituted arylthio;
optionally substituted cycloalkyl or optionally substituted heterocycloalkyl, it being

understood that R' and R'' independently of one another represent a hydrogen atom, an optionally substituted linear or branched (C₁-C₆)alkyl group or an aryl group,

it being possible for the Cy moiety defined in formula (IA) to be substituted by from 1 to 3 groups selected from linear or branched (C₁-C₆)alkyl, linear or branched (C₁-C₆)polyhaloalkyl, hydroxy, linear or branched (C₁-C₆)alkoxy, COOH, NR₁'R₁'' and halogen, it being understood that R₁' and R₁'' have the same definitions than the groups R' and R'' mentioned hereinbefore,


their enantiomers and diastereoisomers, and addition salts thereof with a pharmaceutically acceptable acid or base.

- 10 Compounds of formula (IA-1), their enantiomers and diastereoisomers, and addition salts thereof with a pharmaceutically acceptable acid or base, are more preferred:



wherein X, Y, W, A₁, A₂, R₃, R₄, R₅, R₇, R₈, R₉, T, p, p', q and q' are as defined for formula (IA).

- 15 Compounds of formula (IA-2), their enantiomers and diastereoisomers, and addition salts thereof with a pharmaceutically acceptable acid or base, are even more preferred:



wherein A_1 , A_2 , R_3 , R_4 , R_7 and T are as defined for formula (IA).

Among the pharmaceutically acceptable acids there may be mentioned, without implying any limitation, hydrochloric acid, hydrobromic acid, sulphuric acid, phosphonic acid, acetic acid, trifluoroacetic acid, lactic acid, pyruvic acid, malonic acid, succinic acid, glutaric acid, fumaric acid, tartaric acid, maleic acid, citric acid, ascorbic acid, oxalic acid, methanesulphonic acid, camphoric acid etc.

Among the pharmaceutically acceptable bases there may be mentioned, without implying any limitation, sodium hydroxide, potassium hydroxide, triethylamine, *tert*-butylamine etc.

In a preferred embodiment of the invention, R_4 represents a phenyl substituted at the *para* position by a group of formula $-\text{OPO}(\text{OM})(\text{OM}')$, $-\text{OPO}(\text{OM})(\text{O}^-\text{M}_1^+)$, $-\text{OPO}(\text{O}^-\text{M}_1^+)(\text{O}^-\text{M}_2^+)$, $-\text{OPO}(\text{O}^-)(\text{O}^-)\text{M}_3^{2+}$, $-\text{OPO}(\text{OM})(\text{O}[\text{CH}_2\text{CH}_2\text{O}]_n\text{CH}_3)$, or $-\text{OPO}(\text{O}^-\text{M}_1^+)(\text{O}[\text{CH}_2\text{CH}_2\text{O}]_n\text{CH}_3)$, wherein M and M' independently of one another represent a hydrogen atom, a linear or branched $(\text{C}_1\text{-C}_6)$ alkyl group, a linear or branched $(\text{C}_2\text{-C}_6)$ alkenyl group, a linear or branched $(\text{C}_2\text{-C}_6)$ alkynyl group, a cycloalkyl or a heterocycloalkyl, both of them being composed of from 5 to 6 ring members, while M_1^+ and M_2^+ independently of one another represent a pharmaceutically acceptable monovalent cation, M_3^{2+} represents a pharmaceutically acceptable divalent cation and n is an integer comprised between 1 to 5, it being understood that the phenyl group may be optionally substituted by one or more halogen atoms.

The group Het advantageously represents one of the following groups: tetrahydroindolizine, indolizine or 1,2-dimethyl-1*H*-pyrrole. The group 1,2-dimethyl-1*H*-pyrrole is more especially preferred. Alternatively, the group 5,6,7,8-tetrahydroindolizine is more preferred.

5 In the preferred compounds of the invention, $q=1$ and $q'=1$. Alternatively, $q=2$ and $q'=1$.

Preferably, T represents a hydrogen atom, a methyl group, a (morpholin-4-yl)alkyl group, a dimethylaminomethyl group, or an (alkylpiperazin-1-yl)alkyl group. More preferably, T represents a methyl group, a (morpholin-4-yl)methyl group, a 3-(morpholin-4-yl)propyl group or a (4-methylpiperazin-1-yl)methyl group. Even more preferably, T represents a
10 methyl group. Alternatively, T represents more preferably a (morpholin-4-yl)methyl group.

In preferred compounds of the invention, R_3 represents a linear or branched (C_1 - C_6)alkyl group, an aryl group or a heteroaryl group. More specifically, R_3 is a group selected from phenyl, methyl, ethyl, propyl, butyl, 1-methyl-1*H*-pyrrolo[2,3-*b*]pyridine or 5-cyano-1,2-dimethyl-1*H*-pyrrole. More preferably, R_3 represents a methyl, propyl or butyl group. Even
15 more preferably, R_3 represents a methyl group.

Preferably, R_4 represents a linear or branched (C_1 - C_6)alkyl group (more especially a butyl) or an aryl group. R_4 represents advantageously a phenyl group optionally substituted.

Preference is given to the R_4 group being a 4-hydroxyphenyl.

Advantageously, R_7 represents a group R'_7 -CO- or R'_7 -O-CO-.

20 In preferred compounds of the invention, R'_7 represents an aryl optionally substituted, a cycloalkyl optionally substituted or an alkyl optionally substituted. More preferably, R'_7 represents an optionally substituted naphthalene, phenyl or indole group. The most preferred substitutions for the phenyl group are alkyl optionally substituted, cyano, alkynyl, halogen, alkoxy optionally substituted, or -NR'R''. Even more preferably

preferred substitutions for the phenyl group are methyl, ethyl, methoxy, chloro, bromo, cyano, 2-dimethylaminoethylamino, ethynyl, 2-dimethylaminoethoxy, 2-(dimethylamino)ethyl(methyl)amino, dimethylcarbamoylethyl.

In a preferred embodiment, $p=p'=0$.

5 More particularly, the invention relates to the following compounds:

- *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydro isoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide;
- Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydro isoquinoline-2-carboxylate;
- Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydro isoquinoline-2-carboxylate;
- 15 • Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride;
- Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride;
- 20 • 4-Methylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydro indolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride;
- 2-Methylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydro indolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 25 • 4-Methoxyphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydro indolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 30 • 4-Chlorophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydro

indolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;

- 5

• 4-Ethylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 10

• 4-Cyanophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;

• 2-Methoxyphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 15

• 4-Methylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 20

• 4-Chlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;

• 4-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 25

• 4-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;

• 4-{[2-(Dimethylamino)ethyl]amino}phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 30

• Phenyl 6-{4-[(4-hydroxyphenyl)(propyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;


• Phenyl 6-{4-[butyl(4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;

- Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- *N*-Butyl-*N*-(4-hydroxyphenyl)-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide;
- 3-[2-(Dimethylamino)ethoxy]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-Ethynylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- Naphthalen-2-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 1*H*-Indol-5-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 3-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-Bromophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- (1*r*,4*r*)-4-{[(*tert*-Butoxy)carbonyl]amino}cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;

isoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;


- Cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate.

- 5 The invention relates also to a process for the preparation of compounds of formula (I) characterised in that there is used as starting material the compound of formula (II):




wherein Het, R₃, R₄, R₈, R₉, p, p', q and q' are as defined for formula (I) and Alk represents a linear or branched (C₁-C₆) alkyl group,

- 10 the ester function of which compound of formula (II) is hydrolysed to yield the corresponding carboxylic acid which is then subjected to peptide coupling with a compound of formula (III):



wherein R₅ and T are as defined for formula (I),

to yield the compound of formula (IV):




5 wherein Het, R₃, R₄, R₅, R₈, R₉, T, p, p', q and q' are as defined for formula (I),

compound of formula (IV) which is deprotected and subjected to an acylation or a sulfonylation reaction, and which can optionally be subjected to the action of a pyrophosphate or a phosphonate derivative in basic conditions, to yield the compound of formula (I),

which compound of formula (I) may be purified according to a conventional separation technique, which is converted, if desired, into its addition salts with a pharmaceutically acceptable acid or base and which is optionally separated into its isomers according to a conventional separation technique,


- 5 it being understood that at any moment considered appropriate during the course of the process described above, some groups (hydroxy, amino...) of the starting reagents or of the synthesis intermediates can be protected and subsequently deprotected, as required by the synthesis.

- 10 More specifically, the invention relates also to a process for the preparation of compounds of formula (IA-1) characterised in that there is used as starting material the compound of formula (V):




wherein X, Y, W, A₁ and A₂ are as defined for formula (IA) and Alk represents a linear or branched (C₁-C₆) alkyl group,

- 15 the ester function of which compound of formula (V) is hydrolysed to yield the carboxylic acid or the corresponding carboxylate, which may be converted into an acid derivative such as acyl chloride or the corresponding anhydride before being coupled with an amine NHR₃R₄ wherein R₃ and R₄ have the same meanings as for formula (IA), to yield the compound of formula (VI):



wherein X, Y, W, A₁, A₂, R₃ and R₄ are as defined for formula (IA),

compound of formula (VI) which is then halogenated and further converted into the corresponding borohydride derivative of formula (VII) :




5

wherein X, Y, W, A₁, A₂, R₃ and R₄ are as defined for formula (IA), and R_{B1} and R_{B2} represent a hydrogen, a linear or branched (C₁-C₆) alkyl group, or R_{B1} and R_{B2} form with the oxygen atoms carrying them an optionally methylated ring,


10

which compound of formula (VII) is further subjected to coupling with a compound of formula (VIII):



wherein R_8 , R_9 , p , p' , q and q' are as defined for formula (IA), and Alk' represents a linear or branched (C_1 - C_6) alkyl group, and L represents a leaving group selected from halogen or sulfonate,


5 to yield the compound of formula (IIA-1) :



wherein X , Y , W , A_1 , A_2 , R_3 , R_4 , R_8 , R_9 , p , p' , q , and q' are as defined for formula (IA) and Alk' is as defined previously,


the ester function of which compound of formula (IIA-1) is hydrolysed to yield the

corresponding carboxylic acid which is then subjected to peptide coupling with a compound of formula (III) :




wherein R₅ and T are as defined for formula (IA),

5 to yield the compound of formula (IVA-1) :



wherein X, Y, W, A₁, A₂, R₃, R₄, R₅, R₈, R₉, T, p, p', q, q' are as defined for formula (IA),

compound of formula (IVA-1) which is deprotected and subjected to an acylation or a sulfonylation reaction, and which can optionally be subjected to the action of a pyrophosphate or a phosphonate derivative in basic conditions, to yield the compound of formula (IA-1):



wherein X, Y, W, A₁, A₂, R₃, R₄, R₅, R₇, R₈, R₉, T, p, p', q and q' are as defined for formula (IA),

which compound of formula (IA-1) may be purified according to a conventional separation technique, which is converted, if desired, into its addition salts with a pharmaceutically acceptable acid or base and which is optionally separated into its isomers according to a conventional separation technique,

it being understood that at any moment considered appropriate during the course of the process described above, some groups (hydroxy, amino...) of the starting reagents or of the synthesis intermediates can be protected and subsequently deprotected, as required by the synthesis.

More particularly, when one of the groups R₃ or R₄ of the amine NHR₃R₄ is substituted by a hydroxy function, the latter may be subjected beforehand to a protection reaction prior to coupling with the carboxylic acid formed from the compound of formula (V), or with a corresponding acid derivative thereof, the resulting protected function subsequently undergoes a deprotection reaction in the last stage of the process.

The compounds of formulae (III), (V), (VIII) and the amine NHR_3R_4 are either commercially available or can be obtained by the person skilled in the art using conventional chemical reactions described in the literature.

5 Pharmacological study of the compounds of the invention has shown that they have pro-apoptotic properties. The ability to reactivate the apoptotic process in cancerous cells is of major therapeutic interest in the treatment of cancers and of auto-immune and immune system diseases.

10 More especially, the compounds according to the invention will be useful in the treatment of chemo- or radio-resistant cancers, and in malignant haemopathies and small-cell lung cancer.

15 Among the cancer treatments envisaged there may be mentioned, without implying any limitation, treatment of cancers of the bladder, brain, breast and uterus, chronic lymphoid leukaemias, cancer of the colon, oesophagus and liver, lymphoblastic leukaemias, non-Hodgkin lymphomas, melanomas, malignant haemopathies, myelomas, ovarian cancer, non-small-cell lung cancer, prostate cancer and small-cell lung cancer. Among non-Hodgkin lymphomas, there may be mentioned preferably follicular lymphomas, mantle cell lymphomas, diffuse large B-cell lymphomas, small lymphocytic lymphomas and marginal zone B-cell lymphomas.

20 The present invention relates also to pharmaceutical compositions comprising at least one compound of formula (I) in combination with one or more pharmaceutically acceptable excipients.

25 Among the pharmaceutical compositions according to the invention there may be mentioned more especially those that are suitable for oral, parenteral, nasal, per- or trans-cutaneous, rectal, perlingual, ocular or respiratory administration, especially tablets or dragées, sublingual tablets, sachets, paquets, capsules, glossettes, lozenges, suppositories, creams, ointments, dermal gels, and drinkable or injectable ampoules.

The dosage varies according to the sex, age and weight of the patient, the administration route, the nature of the therapeutic indication, or of any associated treatments, and ranges from 0.01 mg to 1 g per 24 hours in one or more administrations.

Furthermore, the present invention relates also to the association of a compound of formula (I) with an anticancer agent selected from genotoxic agents, mitotic poisons, anti-metabolites, proteasome inhibitors, kinase inhibitors and antibodies, and also to pharmaceutical compositions comprising that type of association and their use in the manufacture of medicaments for use in the treatment of cancer.

The compounds of the invention may also be used in association with radiotherapy in the treatment of cancer.

Finally, the compounds of the invention may be linked to monoclonal antibodies or fragments thereof or linked to scaffold proteins that can be related or not to monoclonal antibodies.

Antibody fragments must be understood as fragments of Fv, scFv, Fab, F(ab')₂, F(ab'), scFv-Fc type or diabodies, which generally have the same specificity of binding as the antibody from which they are descended. According to the present invention, antibody fragments of the invention can be obtained starting from antibodies by methods such as digestion by enzymes, such as pepsin or papain, and/or by cleavage of the disulfide bridges by chemical reduction. In another manner, the antibody fragments comprised in the present invention can be obtained by techniques of genetic recombination likewise well known to the person skilled in the art or else by peptide synthesis by means of, for example, automatic peptide synthesizers such as those supplied by the company Applied Biosystems, etc.

Scaffold proteins that can be related or not to monoclonal antibodies are understood to mean a protein that contains or not an immunoglobulin fold and that yields a binding capacity similar to a monoclonal antibody. The man skilled in the art knows how to select the protein scaffold. More particularly, it is known that, to be selected, such a scaffold should display several features as follow (Skerra A., J. Mol. Recogn., 13, 2000, 167-187): phylogenetically good conservation, robust architecture with a well-known three-

dimensional molecular organization (such as, for example, crystallography or NMR), small size, no or only a low degree of post-translational modifications, easy to produce, express and purify. Such a protein scaffold can be, but without limitation, a structure selected from the group consisting in fibronectin and preferentially the tenth fibronectin type III domain (FNfn10), lipocalin, anticalin (Skerra A., J. Biotechnol., 2001, 74(4):257-75), the protein Z derivative from the domain B of staphylococcal protein A, thioredoxin A or any protein with a repeated domain such as an "ankyrin repeat" (Kohl et al., PNAS, 2003, vol.100, No.4, 1700-1705), "armadillo repeat", "leucine-rich repeat" or "tetratricopeptide repeat". There could also be mentioned a scaffold derivative from toxins (such as, for example, scorpion, insect, plant or mollusc toxins) or protein inhibitors of neuronal nitric oxide synthase (PIN).

The following Preparations and Examples illustrate the invention but do not limit it in any way.

General Procedures

All reagents obtained from commercial sources were used without further purification. Anhydrous solvents were obtained from commercial sources and used without further drying. Flash chromatography was performed on ISCO CombiFlash Rf 200i with pre-packed silica-gel cartridges (Siliasep™ F60 (40-63µm, 60Å). Thin layer chromatography was conducted with 5 x 10 cm plates coated with Merck Type 60 F₂₅₄ silica-gel. Microwave heating was performed with a CEM Discover® SP instrument.

Analytical LC-MS

The compounds of the present invention were characterized by high performance liquid chromatography-mass spectroscopy (HPLC-MS) on either an Agilent HP1200 Rapid Resolution Mass detector 6140 multi mode source m/z range 150 to 1000 amu or an Agilent HP1100 Mass detector 1946D ESI source m/z range 150 to 1000 amu. The conditions and methods listed below are identical for both machines.

Detection: UV detection at 230, 254 and 270 nm.

Injection Volume: 2µL

Mobile Phases: A - Water + 10 mMol / ammonium formate + 0.08% (v/v) formic acid at pH ca 3.5.

B - 95% Acetonitrile + 5% A + 0.08% (v/v) formic acid

5 **Method A (3.75 min; either positive (pos) or positive and negative (pos / neg) ionisation)**

Column: Gemini 5µm, C18, 30 mm x 4.6mm (Phenomenex).

Temperature: 35°C.

Gradient:

Time (min)	Solvent A (%)	Solvent B (%)	Flow (mL/min)
0	95	5	2
0.25	95	5	2
2.50	95	5	2
2.55	5	95	3
3.60	5	95	3
3.65	5	95	2
3.70	95	5	2
3.75	95	5	2

Method B (1.9 min; either positive (pos) or positive and negative (pos / neg) ionisation)

10 Column: Gemini 5µm, C18, 30 mm x 4.6mm (Phenomenex).

Temperature: 35°C.

Gradient:

Time (min)	Solvent A (%)	Solvent B (%)	Flow (mL/min)
0	95	5	1.1
0.12	95	5	1.1
1.30	5	95	1.1
1.35	5	95	1.7
1.85	5	95	1.7
1.90	5	95	1.1
1.95	95	5	1.1

Preparative HPLC

Some compounds of the invention were purified by preparative HPLC. These were performed on a Waters FractionLynx MS autopurification system, with a Gemini® 5 µm C18(2), 100 mm × 20 mm i.d. column from Phenomenex, running at a flow rate of 20 cm³min⁻¹ with UV diode array detection (210–400 nm) and mass-directed collection. Gradients used for each compound are shown in Table 1.

At pH 4: solvent A = 10 mM ammonium acetate in HPLC grade water + 0.08% v/v formic acid. Solvent B = 95% v/v HPLC grade acetonitrile + 5% v/v solvent A + 0.08% v/v formic acid.

At pH 9: solvent A = 10 mM ammonium acetate in HPLC grade water + 0.08% v/v ammonia solution. Solvent B = 95% v/v HPLC grade acetonitrile + 5% v/v solvent A + 0.08% v/v ammonia solution.

The mass spectrometer was a Waters Micromass ZQ2000 spectrometer, operating in positive or negative ion electrospray ionisation modes, with a molecular weight scan range of 150 to 1000.

Preparation 1a: 2-*tert*-Butyl 7-methyl 6-[(nonafluorobutanesulfonyl)oxy]-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

Step A: *tert*-Butyl 7-formyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

tert-Butyl 6-hydroxy-3,4-dihydro-1*H*-isoquinoline-2-carboxylate (15.0 g, 0.06 mol) is dissolved in 96 mL of 7% dimethoxymagnesium in methanol (0.06 mol) solution and stirred for 30 minutes at room temperature. The reaction mixture is then concentrated and co-evaporated with toluene to afford a powder, which is then suspended in toluene (300 mL). Paraformaldehyde (6.32 g, 0.21 mol) is added and the suspension heated to reflux. A further 6.32 g of paraformaldehyde (0.21 mol) is added and allowed to stir for 48 h. The reaction mixture is cooled to room temperature, diluted with ethyl acetate and washed with 1 N HCl. The organic phase is separated and the aqueous washed with ethyl acetate. The organics are combined, filtered through a celite pad and the filtrate washed with brine, dried over magnesium sulphate, filtered and concentrated *in vacuo*. The crude material is taken up in dichloromethane, loaded onto isolute and purified on CombiFlash (220 g silica, dichloromethane to 3% methanol/dichloromethane) to afford the undesired regioisomer cleanly as an oil. The remaining fractions are combined, concentrated to a solid, and re-purified on CombiFlash (220 g silica, iso-hexane to 30% ethyl acetate/ iso-hexane). The product is obtained as a 2:1 ratio in favour of desired regioisomer.

LC/MS (C₁₅H₁₉NO₄) 177 [M-Boc+H]⁺; RT 1.29 (Method B)

Step B: 2-*tert*-Butyl 7-methyl 6-hydroxy-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

To a stirred solution of 6.4 g of a mixture of regioisomers of the aldehyde obtained in Step A (approx. 4.22 g, 15.23 mmol) in methanol (250 mL) is added sodium cyanide (0.75 g, 0.02 mol). After 1 minute, activated manganese dioxide (10.03 g, 115.4 mmol) is added and the reaction is stirred at ambient temperature overnight. The reaction mixture is filtered through a celite pad and concentrated. The crude material is purified on

CombiFlash (220 g silica, iso-hexane to 20% ethyl acetate/ iso-hexane) and dried *in vacuo* to afford a solid.

LC/MS ($C_{16}H_{21}NO_5$) 208 $[M-Boc+H]^+$; RT 1.40 (Method B)

Step C: 2-tert-Butyl 7-methyl 6-[(nonafluorobutanesulfonyl)oxy]-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

To a solution of the phenol obtained in Step B (5.97 g, 19.41 mmol) in dichloromethane (10 mL) cooled to 0 °C is added triethylamine (8.1 mL, 58.23 mmol), followed by nonafluorobutane sulphonyl fluoride (10.46 mL, 58.23 mmol). After complete addition, the mixture is left to warm to room temperature and stirred for 1 day, after which time a further 10.46 mL of the sulphonylating reagent is added and stirring is maintained for a further day. 20.92 ml is subsequently added and the reaction stirred for a further 4-5 days. The reaction mixture is cooled to 0 °C and water (100 mL) is added. The mixture is extracted with dichloromethane and the organic layers dried over magnesium sulphate, filtered and concentrated *in vacuo*. The crude material is taken up in dichloromethane, loaded onto isolate and purified on CombiFlash (120 g silica, iso-hexane to 10% ethyl acetate/ iso-hexane), and dried *in vacuo* to afford an oil which slowly crystallises to a solid.

LC/MS ($C_{20}H_{20}NO_7F_9S$) no ionisation; RT 1.62 (Method B)

Preparation 1b: 2-tert-Butyl 7-methyl 6-(trifluoromethanesulfonyloxy)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

To a solution of the phenol from Step B of Preparation 1a (2.06 g, 6.70 mmol) and pyridine (1.27 mL, 20.1 mmol) in dichloromethane (75 mL) is added triflic anhydride (1.69 mL, 10.05 mmol) at -10 – 0 °C over 30 min. The mixture is stirred at 0 °C for 2 h, then ice-water (50 mL) is added and the mixture is acidified with dilute aqueous hydrochloric acid to give pH 2-3. The mixture is extracted with ethyl acetate and the organic extracts are washed sequentially with with brine and saturated aqueous copper sulfate, dried over magnesium sulphate, and concentrated *in vacuo* to afford the product.

Preparation 1c: *tert*-Butyl 7-formyl-6-(trifluoromethanesulfonyloxy)-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of the product from Step A of Preparation 1a (3.1 g, 11.2 mmol) in anhydrous dichloromethane (50 mL), cooled in an ice-bath, is added pyridine (4.52 mL, 56 mmol) followed by slow addition of triflic anhydride (2.75 mL, 16.8 mmol). The mixture is allowed to warm to ambient temperature and left stirring for *ca* 16 h.

The reaction is diluted with ice-cold water, acidified with dilute HCl to ~ pH 3, and then extracted into ethyl acetate, and washed with saturated aqueous copper sulphate solution.

The organic phase is dried over magnesium sulphate, filtered and evaporated. The crude material is purified by column chromatography on silica, eluting with a gradient of iso-hexane to 25% ethyl acetate/iso-hexane to afford the desired product as a mixture with the corresponding regioisomer (*tert*-butyl 5-formyl-6-(trifluoromethanesulfonyloxy)-1,2,3,4-tetrahydroisoquinoline-2-carboxylate).

Preparation 1d: 2-*tert*-Butyl 5-methyl 6-bromo-2,3-dihydro-1*H*-isoindole-2,5-dicarboxylate

Step A: 5-Bromobenzene-1,2,4-tricarboxylic acid

Bromotrimethyl benzene (40.7 g, 205 mmol) was added to a mixture of water (3.25 L), potassium permanganate (232 g, 1.468 mol) and sodium carbonate (28.5 g, 206 mmol). The mixture was stirred at reflux for 60 h. Ethanol (820 mL) was added dropwise, and the resultant mixture was filtered hot through celite, then allowed to cool to ambient temperature. The filtrate was acidified with concentrated aqueous HCl, and the organic solvent was removed *in vacuo*. The solid product was isolated by filtration.

Step B: 6-Bromo-1,3-dioxo-2,3-dihydro-1*H*-isoindole-5-carboxylic acid

The product from Step A (42.8 g, 148 mmol) and ammonium bromide (43.5 g, 444 mmol) were finely powdered, and homogenised. The resulting solid mixture was heated at 230 – 240 °C for 2 h while carefully mixing every 15 min. The mixture was allowed to cool to ambient temperature, then added to water (230 mL), acidified to pH 1-2 with concentrated

aqueous HCl, and extracted with multiple portions of ethyl acetate. The combined organic extracts were evaporated, and the resultant solid was triturated with a minimal amount of ethyl acetate, and then subsequently with ethanol to afford the desired product as a solid.

Step C: (4-Methoxyphenyl)methyl 6-bromo-2-[(4-methoxyphenyl)methyl]-1,3-dioxo-2,3-dihydro-1H-isoindole-5-carboxylate

The product from Step B (40 g, 148 mmol), 1-(chloromethyl)-4-methoxybenzene (48.7 g, 311 mmol), and potassium carbonate (61.3 g, 444 mmol) were added to DMF (760 mL) and the mixture was heated at 45 °C for *ca* 16 h. Water (1140 mL) was added, and the resultant precipitate was isolated by filtration, dissolved in chloroform (1100 mL), and dried (sodium sulphate). Evaporation afforded the desired product as a solid.

Step D: {6-Bromo-2-[(4-methoxyphenyl)methyl]-2,3-dihydro-1H-isoindol-5-yl}methanol

The product from Step C (55.4 g, 108.5 mmol) was suspended in dry THF (165 mL) and borane-THF (1M in THF; 542 mL, 542 mmol) was added dropwise under argon. After 30 min the mixture was heated to 60 °C for 6 h. Methanol (244.5 mL) was added dropwise (liberating gas) and the resultant solution was evaporated under reduced pressure to a volume of 300 mL. Methanol (170 mL) and 10% aqueous HCl (220 mL) were added and the resultant mixture was heated at 70 °C for 3.5 h. The solution was evaporated under reduced pressure, and the residue was partitioned between saturated aqueous potassium carbonate and dichloromethane. The aqueous phase was extracted with dichloromethane, and the combined organic extracts were concentrated *in vacuo* and purified by flash column chromatography, eluting with a gradient of 5 – 60% ethyl acetate / hexane to afford the desired product.

Step E: tert-Butyl 5-bromo-6-(hydroxymethyl)-2,3-dihydro-1H-isoindole-2-carboxylate

A solution of the compound from Step D (11.3 g, 32.5 mmol) in trifluoroacetic acid (145 mL) was heated as rapidly as possible at 200 °C under microwave irradiation for 7 min. The trifluoroacetic acid was removed under vacuum, 10% aqueous sodium hydroxide (185 mL) was added, and the resultant mixture was carefully triturated / homogenised, then extracted with several portions of dichloromethane. The combined organic extracts were evaporated, then dissolved in a 1:1 mixture of methanol/dichloromethane (210 mL) and di-

tert-butyl dicarbonate (1.2 eq) was added. After 40 min, the solvent was removed *in vacuo* and the crude material was purified by flash column chromatography on silica, eluting with 1:1 hexane/ethyl acetate to afford the product.

Step F: tert-Butyl 5-bromo-6-formyl-2,3-dihydro-1H-isoindole-2-carboxylate

5 The product from Step E (5 g, 15.2 mmol) was dissolved in dichloromethane (63.5), and Dess-Martin periodinane (8.4 g, 19.8 mmol) was added portion-wise. After 1 h, the mixture was concentrated *in vacuo* and the crude material was purified by flash column chromatography on silica, eluting with a gradient of 2%-15% ethyl acetate/hexane to afford the desired product.

10 **Step G: 2-tert-Butyl 5-methyl 6-bromo-2,3-dihydro-1H-isoindole-2,5-dicarboxylate**

The compound from Step F (1.63 g, 5.0 mmol), manganese dioxide (8.69 g, 100 mmol), sodium cyanide (1.47 g, 30 mmol) and acetic acid (601 mg, 10.0 mmol) were suspended in methanol (50 mL) and the mixture was stirred for *ca* 16 h. The reaction was filtered through celite, concentrated *in vacuo*, and the residue partitioned between ethyl acetate and
15 water. The organic phase was concentrated *in vacuo* and purified by flash column chromatography on silica, eluting with a gradient of 0% - 10% ethyl acetate/hexane to afford the desired product.

Preparation 1e: tert-Butyl 5-bromo-6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxylate

20 **Step A: 6-Bromo-2-[(tert-butoxy)carbonyl]-2,3-dihydro-1H-isoindole-5-carboxylic acid**

To a solution of the product of Preparation 1d (1 g, 2.81 mmol) in dioxane (5 mL) is added a solution of lithium hydroxide monohydrate (236 mg, 5.62 mmol) in water (5 mL), and the reaction is heated to 90 °C. After 40 min, the reaction is allowed to cool, and is diluted with water and acidified to ~ pH 2 with 2N aqueous HCl. The mixture is extracted with
25 ethyl acetate, and the organic extracts are dried and evaporated under reduced to pressure to afford the product as a solid.

LC/MS (C₁₄H₁₆BrNO₄) 342 [M+H]⁺; RT 2.30 (Method A)

Step B: *tert*-Butyl 5-bromo-6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

To a solution of the product from Step A (933 mg, 2.73 mmol) in DMF (10 mL) is added triethylamine (1.33 mL, 9.55 mmol), HBTU (1.04 g, 2.73 mmol) followed by the product of Preparation 2a (832 mg, 2.73 mmol) and the mixture is stirred at ambient temperature for *ca* 16 h. The reaction is diluted with water and the resulting precipitate is filtered and washed with water. The solid is taken-up in ethyl acetate, dried over magnesium sulphate, filtered and evaporated. The crude material is purified by column chromatography over silica gel, eluting with a gradient of iso-hexane to 1:1 ethyl acetate/iso-hexane to afford the product as a foam.

LC/MS (C₂₈H₃₄BrN₃O₄) 556 [M+H]⁺; RT 2.35 (Method A)

Preparation 1f: *tert*-Butyl 5-bromo-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

The procedure is as in Preparation 1e, replacing the product of Preparation 2a used in Step C with (3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline hydrochloride obtained from Preparation 2b.

LC/MS (C₂₄H₂₇BrN₂O₃) 471 [M+H]⁺; RT 2.76 (Method A)

Preparation 1g: 6-Bromo-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid

Step A: 2-(3-Bromo-4-methoxyphenyl)ethanamine

To a solution of 2-(4-methoxyphenyl)ethanamine (25.0 g, 0.165 mol) in glacial acetic acid (300 mL) was added a solution of bromine (31.7 g, 0.198 mol) in glacial acetic acid (200 mL) dropwise. A white precipitate formed immediately, and the reaction was stirred at rt for 48 h. The white solid was filtered off and washed with hexane to obtain the hydrobromide salt. The mother liquor was evaporated and it was washed with a small amount of acetic acid and hexane. These salts were dissolved in water and the pH was

adjusted to 8. The suspension was extracted with DCM and dried, and concentrated under reduced pressure. The crude product was used in the next step without further purification.

Step B: *N*-[2-(3-Bromo-4-methoxyphenyl)ethyl]-2,2,2-trifluoroacetamide

2-(3-bromo-4-methoxyphenyl)ethanamine (600 mg) was dissolved in 6 mL TFAA and 6 mL DCM and the mixture was stirred for 1h. It was cooled in ice bath and 24 mL water was added to the mixture. The organic phase was washed with water several times, dried and concentrated under reduced pressure. The crude product was used in the next step without further purification.

Step C: *1*-(6-Bromo-7-methoxy-3,4-dihydroisoquinolin-2(1*H*)-yl)-2,2,2-trifluoroethanone

To a solution of 40 % sulphuric acid in acetic acid (150 mL) was added *N*-[2-(3-bromo-4-methoxyphenyl)ethyl]-2,2,2-trifluoroacetamide (15.5 g, 52.4 mmol) and paraformaldehyde (2.4 g, 80 mmol). The reaction mixture was stirred at room temperature for 18 h, poured into cold water and extracted with ethyl acetate. The combined organic layer was washed with sodium hydrogen carbonate solution, dried and concentrated under reduced pressure. The crude product was purified with flash chromatography (eluent: n-hexane:EtOAc gradient).

Step D: *1*-(6-Bromo-7-hydroxy-3,4-dihydroisoquinolin-2(1*H*)-yl)-2,2,2-trifluoroethanone

Under argon atmosphere a solution of 1-(6-bromo-7-methoxy-3,4-dihydroisoquinolin-2(1*H*)-yl)-2,2,2-trifluoroethanone (1 mmol) in 20 mL DCM was slowly added to a precooled solution of boron tribromide (1 mmol) in 30 mL DCM at -30 °C. The resulting solution was slowly warmed to -12 °C and stirred for 16 h. The reaction was quenched by adding ice and the product was extracted with EtOAc. The organic phase was washed with water and saturated aqueous NaCl and dried, The crude product was purified with flash chromatography (eluent: n-hexane:DCM gradient).

Step E: *6*-Bromo-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl trifluoromethanesulfonate

1-(6-bromo-7-hydroxy-3,4-dihydroisoquinolin-2(1*H*)-yl)-2,2,2-trifluoroethanone (5.72 g, 17.7 mmol) and 2.2 mL pyridine (2.12 g, 26.5 mmol) was dissolved in DCM (130 mL). At 0°C 21.2 mL (21.2 mmol) trifluoromethanesulfonic anhydride (1M in DCM) was added dropwise and the temperature was left to be warmed to rt. When the reaction reached an appropriate conversion it was diluted with brine. The layers were separated and the organics were dried with Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified via flash chromatography using heptane and ethyl acetate as eluents to obtain 6-bromo-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl trifluoromethanesulfonate.

Step F: 6-Bromo-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid

6-bromo-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-7-yl trifluoromethanesulfonate (912 mg, 2.0 mmol) and 146 mg (dppf)PdCl₂ (0.2 mmol) were dissolved in 15 mL DMF and 5 mL H₂O. 560 µL TEA (404 mg, 4.0 mmol) was added and the mixture was stirred at 75°C under CO atmosphere (1 bar). When the reaction reached an appropriate conversion it was diluted with DCM, extracted with 0.2 M HCl_{aq}. The layers were separated, the organics were dried with Na₂SO₄ filtered and concentrated under reduced pressure. The crude product was purified with flash chromatography using DCM and methanol as eluents to obtain 6-bromo-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid.

Preparation 1ha: 1-{6-Bromo-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-2-yl}-2,2,2-trifluoroethan-1-one

To a solution of the acid from Preparation 1g (2 g, 0.01 mol) in *N,N*-dimethylformamide (15 mL) was added *N,N*-diisopropylethylamine (1.98 mL, 11.36 mmol), the amine from Preparation 2a (1.39 g, 5.96 mmol) and HBTU (2.15 g, 5.68 mmol), and the mixture was stirred at ambient temperature for *ca* 16 h. The reaction was concentrated *in vacuo* then redissolved in ethylacetate, and washed with water and brine. The organic extract was dried over magnesium sulphate, concentrated *in vacuo*, and purified by flash column chromatography (40 g silica, dichloromethane to 3% methanol in dichloromethane).

LC/MS ($C_{26}H_{27}BrF_3N_3O_3$) 566 $[M+H]^+$; RT 1.26 (Method B)

Preparation 1hb: 1-{6-Bromo-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-2-yl}-2,2,2-trifluoroethan-1-one

The procedure is as in Preparation 1ha, replacing the amine from Preparation 2a with the amine from Preparation 2b.

LC/MS ($C_{22}H_{20}BrF_3N_2O_2$) 481 $[M+H]^+$; RT 1.46 (Method B)

Preparation 2a: (3*S*)-3-(Morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride

Step A: Benzyl (3*S*)-3-(4-morpholinylcarbonyl)-3,4-dihydro-2(1*H*)-isoquinoline carboxylate

To a solution of 5 g of (3*S*)-2-[(benzyloxy)carbonyl]-1,2,3,4-tetrahydro-3-isoquinoline-carboxylic acid (16 mmol) in 160 mL of dichloromethane there are added 1.5 mL of morpholine (17.6 mmol) and then 9 mL of *N,N,N*-triethylamine (64 mmol), 3.3 g of 1-ethyl-3-(3'-dimethylaminopropyl)-carbodiimide (EDC) (19.2 mmol) and 2.6 g of hydroxybenzotriazole (HOBt) (19.2 mmol). The reaction mixture is stirred at ambient temperature overnight and is then poured onto a solution of ammonium chloride and extracted with ethyl acetate. The organic phase is subsequently dried over magnesium sulphate and then filtered and evaporated to dryness. The crude product so obtained is then purified by chromatography over silica gel (dichloromethane/methanol gradient). The product is obtained in the form of a foam.

^1H -NMR: δ (400 MHz; $dmsO-d_6$; 353°K): 7.30 (m, 5H benzyl); 7.15 (m, 4H aromatic); 5.2-5.0 (m, 3H, 2H benzyl, 1H dihydroisoquinoline); 4.75-4.5 (2d, 2H dihydroisoquinoline); 3.55-3.3 (m, 8H morpholine); 3.15-2.9 (2dd, 2H dihydroisoquinoline)

IR: ν : $>C=O$: 1694; 1650 cm^{-1}

Step B: *Benzyl (3S)-3-(4-morpholinylmethyl)-3,4-dihydro-2(1H)-isoquinoline carboxylate*

To a solution of 5.3 g of the product obtained in Step A (13.9 mmol) in 278 mL of tetrahydrofuran there are added 14 mL of $\text{BH}_3\text{Me}_2\text{S}$ (27.8 mmol) at ambient temperature.

5 The whole is heated for 4 hours at 80°C. It is allowed to return to ambient temperature, and then there are added 7 mL (14 mmol) of $\text{BH}_3\text{Me}_2\text{S}$. The reaction mixture is again heated at 80°C for 2 hours. The tetrahydrofuran is subsequently evaporated off, and there are then added slowly methanol and then 5.6 mL of 5N hydrochloric acid (27.8 mmol). The mixture is stirred at ambient temperature overnight and then at 80°C for one hour. There is
10 subsequently added a saturated NaHCO_3 solution to the reaction mixture at 0°C until a pH of 8 is reached, and then extraction with ethyl acetate is carried out. The organic phase is subsequently dried over magnesium sulphate and then filtered and evaporated to dryness. The title product is obtained in the form of an oil.

15 **$^1\text{H-NMR}$:** δ (400 MHz; $\text{dms}\text{-d}_6$; 353°K): 7.43-7.30 (unresolved peak, 5H benzyl); 7.19 (m, 4H aromatic); 5.16 (m, 2H, 2H benzyl); 4.79-4.29 (d, 2H dihydroisoquinoline); 4.58 (m, 1H dihydroisoquinoline); 3.50 (m, 4H morpholine); 3.02-2.80 (dd, 2H dihydroisoquinoline); 2.42-2.28 (unresolved peak, 5H, 4H morpholine, 1H morpholine); 2.15 (dd, 1H morpholine)

20 **IR:** ν : $>\text{CH}$: 2810 cm^{-1} ; ν : $>\text{C}=\text{O}$: 1694 cm^{-1} ; ν : $>\text{C}-\text{O}-\text{C}<$: 1114 cm^{-1} ; ν : $>\text{CH-Ar}$: 751; 697 cm^{-1}

Step C: *(3S)-3-(4-Morpholinylmethyl)-1,2,3,4-tetrahydroisoquinoline*

To a solution of 4.9 g of the compound of Step B (13.4 mmol) in 67 mL of ethanol there is added 0.980 g of palladium dihydroxide (20% by mass) at ambient temperature. The reaction mixture is placed under 1.2 bar of hydrogen at ambient temperature for 4 hours. It
25 is subsequently passed over a Whatman filter, and then the palladium is rinsed several times with ethanol. The filtrate is evaporated to dryness. The title product is obtained in the form of an oil.

$^1\text{H-NMR}$: δ (400 MHz; $\text{dms}\text{-d}_6$; 300°K): 7.12-7.0 (unresolved peak, 4H aromatic); 3.92 (s, 2H tetrahydroisoquinoline); 3.60 (t, 4H morpholine); 2.98 (m, 1H

tetrahydroisoquinoline); 2.68 (dd, 1H tetrahydroisoquinoline); 2.5-2.3 (unresolved peak, 8H, 1H tetrahydroisoquinoline, 6H morpholine, 1H NH)

IR: ν : >NH: 3322 cm^{-1} ; ν : >C-O-C<: 1115 cm^{-1} ; ν : >CH-Ar: 742 cm^{-1}

Step D: *(3S)-3-(4-Morpholinylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride*

5 The free base obtained in Step C is dissolved in a minimum volume of dichloromethane. To the stirred solution at room temperature is added 1M HCl solution in diethyl ether (3 equiv). The reaction is stirred for 15 minutes, after which time diethyl ether is added. The resulting precipitate is filtered, washed with diethyl ether, then dried under vacuum to afford the product.

10 **Preparation 2b:** *(3R)-3-Methyl-1,2,3,4-tetrahydroisoquinoline*

Step A: *3-Methyl-1,2,3,4-tetrahydroisoquinoline*

To a solution of 3-methylisoquinoline and $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (1.2 eq) in methanol (3 mL /mmol), cooled to 0°C, is added sodium borohydride (12 eq) portion-wise over 1 h. The reaction mixture is stirred at ambient temperature for 1 h, then quenched by addition of
15 water. The solvent is removed under reduced pressure, and the residue is partitioned between ethyl acetate and water. The organic phase is dried over sodium sulphate and evaporated to afford the crude product.

Step B: *(3R)-3-Methyl-1,2,3,4-tetrahydroisoquinoline*

The product from Step A (4.5 g) is dissolved in a 1:9 isopropanol/heptane mixture (55
20 mL), and the resultant solution is repeat-injected onto an IC Column (50×500mm, 20 μm particle size), eluting with 5:95 mixture of 2-propanol/heptane containing 0.05% diethylamine, with a flow rate of 50 mL/min at ambient temperature. Under these conditions the (*R*)-isomer eluted as the second fraction.

25 **Preparation 2c:** *tert-Butyl [(3S)-1,2,3,4-tetrahydroisoquinolin-3-ylmethyl] carbamate*

Step A: *Benzyl (3S)-3-(hydroxymethyl)-3,4-dihydro-1H-isoquinoline-2-carboxylate*

This compound is obtained using a protocol from the literature (R. B. Kawthekar *et al* *South Africa Journal of Chemistry* 63, 195, **2009**) starting from 15 g of (3*S*)-1,2,3,4-tetrahydroisoquinolin-3-ylmethanol (91.9 mmol) in the presence of benzyl chloroformate and triethylamine dissolved in dichloromethane. After purification on silica gel (gradient petroleum ether/ AcOEt), the title product is obtained in the form of an oil.

¹H NMR: δ (300 MHz; DMSO-d₆; 300K): 7.33 (m, 5H, aromatic Hs, O-benzyl); 7.15 (s, 4H, aromatic Hs, H tetrahydroisoquinoline); 5.13 (s, 2H, **CH₂**-Ph); 4.73 (d, 1H, H tetrahydroisoquinoline); 4.47 (m, H, CH₂OH); 4.36 (m, 1H, H tetrahydroisoquinoline); 4.28 (d, 1H, H tetrahydroisoquinoline); 3.39 (dd, 1H, **CH₂**OH); 3.23 (dd, 1H, **CH₂**OH); 2.93 (dd, 1H, H tetrahydroisoquinoline); 2.86 (dd, 1H, H tetrahydroisoquinoline).

IR: ν OH: 3416 cm⁻¹; ν <C=O 1694 cm⁻¹; ν aromatic >C-H: 754 cm⁻¹

Step B: Benzyl (3*S*)-3-(azidomethyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

This compound is obtained using a protocol from the literature (D. Pagé *et al* *J. Med. Chem.*, 44, 2387, **2001**) starting from 23 g of the compound obtained in Step A (77.3 mmol) in the presence of diphenylphosphoryl azide and triphenylphosphine dissolved in THF. After purification on silica gel (gradient petroleum ether / AcOEt), the title product is obtained in the form of an oil.

¹H NMR: δ (400 MHz; DMSO-d₆; 300K): 7.36 (m, 5H, aromatic Hs, O-benzyl); 7.19 (m, 4H, aromatic Hs, H tetrahydroisoquinoline); 5.16 (s, 2H, **CH₂**-Ph); 4.76 (d, 1H, H tetrahydroisoquinoline); 4.53 (m, 1H, H tetrahydroisoquinoline); 4.30 (m, 1H, H tetrahydroisoquinoline); 3.28 (m, 2H, **CH₂**N₃); 3.06 (dd, 1H, H tetrahydroisoquinoline); 2.78 (dd, 1H, H tetrahydroisoquinoline)

IR: ν N₃: 2095 cm⁻¹; ν <C=O: 1694 cm⁻¹; ν aromatic >C-H: 754 cm⁻¹

Step C: Benzyl (3*S*)-3-(aminomethyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

To a solution of 20.9 g (64.5 mmol) of the azido compound obtained in Step B in 650 mL of THF there are successively added 25.5 g (97.2 mmol) of triphenylphosphine and 157 mL of water. The complete batch is heated at reflux for 2 hours 30 minutes. The reaction mixture is then concentrated to dryness and the residual oil is then taken up in isopropyl

ether. A white precipitate appears, which is filtered off and washed with isopropyl ether. The filtrate is then concentrated to dryness and is then purified by chromatography on silica gel (gradient CH₂Cl₂/MeOH). The title product is obtained in the form of an oil.

¹H NMR: δ (400 MHz; DMSO-d₆; 300K): 7.40 (m, 5H, aromatic Hs, O-benzyl); 7.20 (m, 4H, aromatic Hs, H tetrahydroisoquinoline); 5.15 (s, 2H, CH₂-Ph); 4.75-4.3 (m, 2H, H tetrahydroisoquinoline); 4.30 (d, 1H, H tetrahydroisoquinoline); 2.90 (m, 2H, CH₂NH₂); 2.45 (m, 2H, H tetrahydroisoquinoline); 1.40 (m, 2H, NH₂)

IR: ν NH₂: 3400-3300 cm⁻¹; ν <C=O: 1688 cm⁻¹

Step D: Benzyl (3S)-3-[(tert-butoxycarbonyl)amino]methyl-3,4-dihydroisoquinoline-2(1H)-carboxylate

To a solution of 18.4 g (62.1 mmol) of the compound obtained in Step C in 630 mL of dichloromethane there are successively added 17.5 mL (124 mmol) of triethylamine and, in portions, 14.9 g (68.3 mmol) of di-*tert*-butyl dicarbonate. The complete batch is stirred at ambient temperature for 2 hours. The reaction mixture is then concentrated, and then ethyl acetate is added. The organic phase is successively washed with 1M HCl solution, saturated NaCl solution, saturated NaHCO₃ solution and then saturated NaCl solution. After drying, concentrating to dryness and purification by chromatography on silica gel (gradient petroleum ether / AcOEt), the title product is obtained in the form of an oil.

¹H NMR: δ (400 MHz; DMSO-d₆; 300K): 7.35 (m, 5H, aromatic Hs, O-benzyl); 7.15 (m, 4H, aromatic Hs, H tetrahydroisoquinoline); 6.51 (m, 1H, NHBoc); 5.12 (s, 2H, CH₂-Ph); 4.76 (d, 1H, H tetrahydroisoquinoline); 4.51 (m, 1H, H tetrahydroisoquinoline); 4.36 (d, 1H, H tetrahydroisoquinoline); 2.95 (m, 3H, H tetrahydroisoquinoline + CH₂NHBoc); 2.71 (d, 1H, H tetrahydroisoquinoline); 1.34 (s, 9H, NHBoc)

IR: ν NH: 3351 cm⁻¹; ν <C=O: 1686 cm⁻¹

Step E: *tert*-Butyl [(3S)-1,2,3,4-tetrahydroisoquinolin-3-ylmethyl]carbamate

To a solution of 21 g (53 mmol) of the compound obtained in Step D in 600 mL of ethyl acetate there are added 2.1 g of palladium-on-carbon 10 %. The complete batch is stirred at ambient temperature under 1.3 bar of dihydrogen pressure for 5 hours. The reaction

mixture is then filtered, and then concentrated to dryness. The title product is obtained in the form of a solid.

¹H NMR: δ (400 MHz; DMSO-d₆; 300K): 7.15 (m, 4H, aromatic Hs, H tetrahydroisoquinoline); 6.85 (t, 1H, **NHBoc**); 3.90 (m, 2H, H tetrahydroisoquinoline); 3.00 (m, 2H, **CH₂NHBoc**); 2.80 (m, 1H, H tetrahydroisoquinoline); 2.65 (dd, 1H, H tetrahydroisoquinoline); 2.40 (dd, 1H, H tetrahydroisoquinoline); 1.40 (s, 9H, **NHBoc**)

IR: ν NH: 3386-3205 cm⁻¹ (NH amide); ν <C=O: 1688 cm⁻¹; ν NH: 1526 cm⁻¹ (NH amine)

Preparation 2d: (3S)-3-[2-(Morpholin-4-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride

Step A: *tert*-Butyl (3S)-3-(2-morpholino-2-oxo-ethyl)-3,4-dihydro-1H-isoquinoline-2-carboxylate

To a solution of 3 g (10.30 mmol) of [(3S)-2-(*tert*-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinolin-3-yl]acetic acid in 100 mL of dichloromethane there are added, dropwise, 1.10 mL (11.32 mmol) of morpholine and, dropwise throughout, 4.3 mL (30.9 mmol) of triethylamine, 2.20 g (12.40 mmol) of 1,2-dichloromethane and 1.70 g (1.68 mmol) of hydroxybenzotriazole. The complete batch is stirred at ambient temperature for 15 hours. The reaction mixture is then diluted with dichloromethane and washed successively with 1M HCl solution, saturated NaHCO₃ solution and then saturated NaCl solution until neutral. The organic phase is then dried over MgSO₄, filtered and concentrated to dryness. After purification by column chromatography on silica gel (dichloromethane /MeOH), the title product is obtained in the form of an oil.

¹H NMR: δ (400 MHz; dmso-d₆; 300K): 7.20-7.10 (m, 4H, aromatic Hs, tetrahydroisoquinoline); 4.70 (m, 1H, aliphatic Hs, CH tetrahydroisoquinoline); 4.75-4.20 (2m, 2H, aliphatic Hs, CH₂ alpha to tetrahydroisoquinoline N); 3.60 (m, 8H, aliphatic Hs, morpholine); 3.00 and 2.70 (2 dd, 2H, aliphatic H, tetrahydroisoquinoline); 2.50-2.20 (2d, 2H, aliphatic Hs, CH₂CO); 1.40 (s, 9H, ^tBu)

IR: ν C=O: 1687; 1625 cm⁻¹

Step B: *1-(Morpholin-4-yl)-2-[(3S)-1,2,3,4-tetrahydroisoquinolin-3-yl]ethanone hydrochloride*

To a solution of 2.88 g (7.18 mmol) of the compound obtained in Step A in 16 mL of dichloromethane there are added, dropwise, 80 mL (80 mmol) of 1M ethereal HCl solution. The complete batch is stirred at ambient temperature for 15 hours, the suspension is then filtered and the precipitate is washed with ether. After drying, the title product is obtained in the form of a solid.

¹H NMR: δ (400 MHz; dms_o-d₆; 300K): 9.80-9.50 (m, 2H, NH₂⁺); 7.30-7.10 (m, 4H, aromatic Hs, tetrahydroisoquinoline); 4.30 (m, 2H, aliphatic Hs, CH₂ alpha to tetrahydroisoquinoline N); 3.80 (m, 1H, aliphatic Hs, CH tetrahydroisoquinoline); 3.70-3.40 (2m, 8H, aliphatic Hs, morpholine); 3.15 and 2.8 (m, 4H, aliphatic H, CH₂ tetrahydroisoquinoline and CH₂CO)

IR: ν -NH₂⁺: 2800-1900 cm⁻¹; ν C=O: 1620 cm⁻¹

Step C: *(3S)-3-[2-(Morpholin-4-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline hydrochloride*

A solution of 2.2 g (7.44 mmol) of the compound obtained in Step B in 22 mL of MTBE and 5 mL of dichloromethane is prepared. After cooling in an ice bath at 0°C, there are then added, dropwise, 15 mL (15 mmol) of a 1M solution of LiAlH₄ in tetrahydrofuran. The complete batch is then stirred at ambient temperature for 6 hours. It is placed at 0°C, and there is then added, dropwise, 1 mL of 5M NaOH solution. The complete batch is stirred at ambient temperature for 45 minutes. The solid is then filtered off and washed with MTBE and then with dichloromethane, and the filtrate is concentrated to dryness. The oil thereby obtained is diluted with dichloromethane and there are added, dropwise, 6.3 mL of 1M ethereal HCl solution. The complete batch is stirred at ambient temperature for 1 hour and then filtered. The crystals thereby obtained are washed with ethyl ether. After drying, the title product is obtained in the form of a solid.

¹H NMR: δ (400 MHz; dms_o-d₆; 300K): 11.35 + 9.80 (2m, 2H, NH₂⁺); 10.00 (m, H, NH⁺); 7.20 (m, 4H, aromatic Hs, tetrahydroisoquinoline); 4.30 (s, 2H, aliphatic Hs, CH₂ alpha to tetrahydroisoquinoline N); 4.00 + 3.85 (2m, 4H, aliphatic Hs, CH₂ alpha to morpholine N); 3.70 (m, 1H, aliphatic Hs, CH tetrahydroisoquinoline); 3.55-3.30 (m, 4H,

aliphatic Hs, CH alpha to morpholine O and CH₂-Morpholine); 3.15 (dd, 1H, aliphatic H, CH₂ tetrahydroisoquinoline); 3.10 (m, 2H, aliphatic H, CH alpha to morpholine O); 2.90 (dd, 1H, aliphatic H, CH₂ tetrahydroisoquinoline); 2.30 + 2.15 (2m, 2H, aliphatic H, CH₂-tetrahydroisoquinoline)

5 **IR:** ν NH⁺ / -NH₂⁺: between 3500 and 2250 cm⁻¹; ν C=C: weak 1593 cm⁻¹; ν aromatic C-H: 765 cm⁻¹

Preparation 2e: *tert*-Butyl {2-[(3*S*)-1,2,3,4-tetrahydroisoquinolin-3-yl]ethyl} carbamate

Step A: *Benzyl (3S)-3-(2-hydroxyethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate*

10 The title compound is obtained starting from (3*S*)-2-[(benzyloxy)carbonyl]-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, based on a protocol from the literature (Jinlong Jiang *et al Bioorganic & Medicinal Chemistry Letters*, 14, 1795, **2004**).

Step B: *Benzyl (3S)-3-{2-[(methanesulphonyl)oxy]ethyl}-3,4-dihydroisoquinoline-2(1H)-carboxylate*

15 To a solution of 10.6 g of the compound of Step A (35.6 mmol) in 350 mL of anhydrous CH₂Cl₂, placed at 0°C, there are successively added triethylamine 10.1 mL (71.2 mmol) and then, dropwise, methanesulphonyl chloride 3.1 mL (39 mmol). The reaction mixture is then stirred at ambient temperature for 2 hours. Hydrolysis is then carried out by slowly adding water. The product is extracted several times with CH₂Cl₂. The organic phases are
20 then combined and successively washed with 1N HCl solution, saturated NaCl solution, saturated NaHCO₃ solution and saturated NaCl solution until neutral. They are then dried over MgSO₄ and concentrated to dryness. After purification by chromatography on silica gel (gradient petroleum ether/ AcOEt), the expected product is obtained in the form of a foam.

25 LC/MS: $m/z = (M + H)^+ = 375$

Step C: *Benzyl (3S)-3-(cyanomethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate*

To a solution of 15.4 g of the compound obtained in Step B (41.02 mmol) in 250 mL of

anhydrous DMSO there are added 22 g (449 mmol) of sodium cyanide. The complete batch is then heated at 60°C for 12 hours. It is allowed to cool and then the reaction mixture is diluted by adding ethyl acetate. Hydrolysis is then carried out using saturated NaHCO₃ solution. After again extracting twice with ethyl acetate, the organic phases are combined, washed with H₂O, dried over MgSO₄ and concentrated to dryness. After purification by chromatography on silica gel (hexane/ AcOEt 7/3), the expected product is obtained in the form of an oil.

LC/MS: $m/z = [M+H]^+ = 307.1$

Step D: Benzyl (3S)-3-(2-aminoethyl)-3,4-dihydroisoquinoline-2(1H)-carboxylate

To a solution of 15.4 g of the compound obtained in Step C (50.3 mmol) in 300 mL of anhydrous THF, placed at 0°C, there is added, dropwise, a 1N solution of BH₃-THF. The reaction mixture is allowed to gradually come back to ambient temperature and then the complete batch is stirred for 14 hours. The reaction mixture is then hydrolysed by slowly adding saturated NH₄Cl solution. After extracting twice with ethyl acetate, the organic phases are combined and dried over MgSO₄. After concentrating to dryness, the expected product is obtained in the form of a foam which is used directly, without purification, in the next, protection step.

Step E: Benzyl (3S)-3-{2-[(tert-butoxycarbonyl)amino]ethyl}-3,4-dihydroisoquinoline-2(1H)-carboxylate

To a solution of 15.6 g of the compound obtained in Step D (50.3 mmol) in 670 mL of CH₂Cl₂, there are successively added 13.2 g (60.36 mmol) of Boc₂O in portions, 14 mL (100.6 mmol) of triethylamine, and DMAP in a catalytic amount. The complete batch is stirred at ambient temperature for 5 hours. The reaction mixture is then hydrolysed with water and extracted twice with CH₂Cl₂. The organic phases are combined, washed with water and dried over MgSO₄. After concentrating to dryness and purification by chromatography on silica gel (gradient heptane/AcOEt), the expected product is obtained in the form of an oil.

LC/MS: $m/z = (M + H)^+ = 411$

Step F: tert-Butyl {2-[(3S)-1,2,3,4-tetrahydroisoquinolin-3-yl]ethyl}carbamate

To a solution of 10.4 g of the compound obtained in Step E (25.5 mmol) in 210 mL of anhydrous MeOH there are added 2.71 g (2.55 mmol) of Pd/C 10%. The complete batch is degassed for 30 minutes and then stirred under a hydrogen atmosphere for 16 hours. The reaction mixture is then filtered and concentrated to dryness. The expected product is obtained in the form of a solid which is taken up in a mixture of pentane/Et₂O (90/10), triturated and filtered. After drying, the product is obtained in the form of a solid.

¹H NMR: δ (400 MHz; dms^o-d₆; 300K): 7.1-6.98 (m, 4H, aromatic Hs, tetrahydroisoquinoline); 6.83 (m, 1H, CH₂NHBoc); 3.85 (s, 2H, aliphatic Hs, tetrahydroisoquinoline); 3.09 (q, 2H, CH₂NHBoc); 2.73 (m, 1H, aliphatic Hs, tetrahydroisoquinoline); 2.70 and 2.39 (2m, 2H, aliphatic Hs, tetrahydroisoquinoline); 1.63 (m, 2H, aliphatic Hs); 1.38 (s, 9H, NHCOOtBu)

IR: ν: >NH: 3378, -3201 cm⁻¹ (amine, amide); ν: >C=O: 1683 cm⁻¹ (amide); ν: >NH: 1524 cm⁻¹ (amide); ν: >C=O: 1168 cm⁻¹

LC/MS: m/z = [M+H]⁺ = 277

Preparation 2f: (3R)-3-[3-(Morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline dihydrochloride

Step A: {(3S)-2-[(4-Methylphenyl)sulphonyl]-1,2,3,4-tetrahydroisoquinolin-3-yl}methyl 4-methylbenzenesulphonate

The procedure is the same as that of Step A of Preparation 1'.

Step B: tert-Butyl 2-[(3R)-2-[(4-methylphenyl)sulphonyl]-1,2,3,4-tetrahydroisoquinolin-3-yl]methyl-3-(morpholin-4-yl)-3-oxopropanoate

To a suspension of 1 g of NaH (60 %) (25.08 mmol) in 30 mL of MTBE there are added, dropwise, a solution of 5 g of tert-butyl 3-morpholino-3-oxopropanoate (21.81 mmol) in 20 mL of anhydrous MTBE. This suspension stirred at ambient temperature for 1 hour and

then the compound obtained in Step A is added in the form of a powder. The batch is stirred at 60°C for 30 hours. 100 mL of saturated ammonium chloride solution are added. The resulting solution is extracted with dichloromethane. The organic phase is then dried over MgSO₄, filtered and concentrated to dryness. After purification by column chromatography over silica gel (dichloromethane/MeOH), the expected product is obtained in the form of an oil.

NMR ¹H (500 MHz, dms_o-d₆) δ ppm : 7.63/7.59 (2d, 2 H), 7.3/7.26 (2d, 2 H), 7.13 (m, 2 H), 7.09/6.97 (2t, 2 H), 4.64/4.55/4.36/4.28 (2AB, 2 H), 4.25/4.11 (2m, 1 H), 3.81 (m, 1 H), 3.73-3.48 (m, 4 H), 3.57-3.32 (m, 4 H), 2.51 (m, 2 H), 2.32/2.31 (2s, 3 H), 1.88/1.79 (2m, 2 H), 1.39/1.38 (2s, 9 H)

IR (ATR) cm⁻¹: ν : >C=O : 1731 (ester); ν : >C=O : 1644 (amide); ν : -SO₂ : 1334-1156; ν : >C-O-C< : 1115; γ : >CH-Ar : 815-746-709

Step C: 2-((3*R*)-2-[(4-Methylphenyl)sulphonyl]-1,2,3,4-tetrahydroisoquinolin-3-yl)methyl)-3-(morpholin-4-yl)-3-oxopropanoic acid

To a solution of 9.5 g (17.97 mmol) of the compound obtained in Step B in 40 mL of dioxane there are added, dropwise, 20 mL of a 4M solution of HCl in dioxane. The batch is stirred at ambient temperature for 48 hours and then the solution is concentrated to dryness. After drying, the expected product is obtained in the form of an oil.

NMR ¹H (400 MHz, dms_o-d₆) δ ppm : 12.75 (m, 1 H), 7.6 (2*d, 2 H), 7.3 (2*d, 2 H), 7.1/6.95 (2*m, 4 H), 4.7-4.2 (d, 2 H), 4.25/4.12 (2*m, 1 H), 3.9-3.3 (m, 9 H), 2.55 (d, 2 H), 2.3 (2*s, 3 H), 1.8 (t, 2 H)

IR (ATR) cm⁻¹ : ν : -OH : 3500 à 2000; ν : >C=O : 1727 (acide) ; ν : >C=O : 1634 (amide) ; ν : -SO₂ : 1330-1155

Step D: 3-((3*R*)-2-[(4-Methylphenyl)sulphonyl]-1,2,3,4-tetrahydroisoquinolin-3-yl)-1-(morpholin-4-yl)propan-1-one

To a solution of 7.80 g (16.51 mmol) of the compound obtained in Step C in 100 mL of DMSO there are added 1.16 g (19.83 mmol) of solid sodium chloride and then, dropwise, 5 mL of water. The batch is stirred at 130°C for 1 hour and then the solution is

concentrated to $\frac{3}{4}$. The reaction mixture is then diluted with dichloromethane and washed successively with saturated lithium chloride solution and then with saturated NaCl solution. The organic phase is then dried over MgSO_4 , filtered and concentrated to dryness. After purification by column chromatography over silica gel (cyclohexane/ethyl acetate), the expected product is obtained in the form of an oil.

NMR ^1H (400 MHz, $\text{dms}\text{-d}_6$) δ ppm : 7.65 (d, 2 H), 7.3 (d, 2 H), 7.15/7 (2 m, 4 H), 4.6 (d, 1 H), 4.25 (d, 1 H), 4.2 (m, 1 H), 3.5 (m, 4 H), 3.4 (2 m, 4 H), 2.6 (2 dd, 2 H), 2.35 (s, 3 H), 2.3 (m, 2 H), 1.5 (quad., 2 H)

IR (ATR) cm^{-1} : ν : $>\text{C}=\text{O}$: 1639 ; ν : $-\text{SO}_2$: 1331-1156 ; γ : $>\text{CH}-\text{Ar}$: 815-675

Step E: (3R)-2-[(4-Methylphenyl)sulphonyl]-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline

To a solution of 6.0 g (14.0 mmol) of the compound obtained in Step D in 60 mL of MTBE and 14 mL of dichloromethane there are added 1.06 g (28 mmol) of LAH in portions over 5 minutes. The batch is stirred at ambient temperature for 15 hours. There are added, dropwise, 1.5 mL of water and stirring is carried out for 15 minutes. There are then added, dropwise, 1.5 mL of 5M sodium hydroxide solution and stirring is carried out for 15 minutes. The reaction mixture is then diluted with MTBE and dichloromethane. The suspension is then filtered and the precipitate is washed with MTBE and dichloromethane. The organic phase is then dried over MgSO_4 , filtered and concentrated to dryness. After purification by column chromatography over silica gel (dichloromethane/EtOH/ NH_4OH), the expected product is obtained in the form of an oil.

NMR ^1H (400 MHz, $\text{dms}\text{-d}_6$) δ ppm : 7.68 (d, 2 H), 7.32 (d, 2 H), 7.1 (massif, 4 H), 4.65/4.23 (AB, 2 H), 4.2 (m, 1 H), 3.55 (t, 4 H), 2.7/2.6 (ABx, 2 H), 2.35 (s, 3 H), 2.25 (t, 4 H), 2.2 (t, 2 H), 1.4/1.3 (2m, 4 H)

IR (ATR) cm^{-1} : ν : $-\text{SO}_2$: 1333-1158

Step F: (3R)-3-[3-(Morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline

To a solution of 1.50 g (3.62 mmol) of the compound obtained in Step E in 20 mL of anhydrous methanol there are added 2.0 g (82.3 mmol), in portions, of magnesium

turnings. The batch is stirred in the presence of ultrasound for 96 hours. The reaction mixture is then filtered, the solid is washed several times with methanol, and the filtrate is concentrated to dryness. After purification by column chromatography over silica gel (dichloromethane/EtOH/NH₄OH), the expected product is obtained in the form of an oil.

5 **NMR** ¹H (400 MHz, dmso-d₆) δ ppm : 7.3 (d, 2 H), 7.1 (t, 2 H), 7.1 (d+t, 3 H), 7 (d, 2 H), 3.9 (s, 2 H), 3.55 (t, 4 H), 2.75 (m, 1 H), 2.72/2.45 (dd, 2 H), 2.35 (t, 4 H), 2.25 (t, 2 H), 1.6 (m, 2 H), 1.45 (m, 2 H)

IR (ATR) cm⁻¹ : ν : >NH₂+ /NH+ : 3500-2300 ; ν : >C-O-C< : 1115

High-resolution mass spectroscopy (ESI+/-/FIA/HR) :

10 Formule brute : C₁₆ H₂₄ N₂ O

[M+H]⁺ calculated: 261.1961

[M+H]⁺ measured: 261.1959

Step _____ G: (3R)-3-[3-(Morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline dihydrochloride

15 The free base obtained in Step F is dissolved in a minimum volume of dichloromethane. To the stirred solution at room temperature is added 1M HCl solution in diethyl ether (3 equiv). The reaction is stirred for 15 minutes, after which time diethyl ether is added. The resulting precipitate is filtered, washed with diethyl ether, then dried under vacuum to afford the product.

20 ***Preparation 2g: N,N-Dimethyl[(3S)-1,2,3,4-tetrahydroisoquinolin-3-ylmethyl]amine dihydrochloride***

The procedure is as in the process of Preparation 2a, replacing the morpholine used in Step A by *N,N*-dimethylamine.

Preparation 2h: **(3*S*)-3-[(4-Methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline trihydrochloride**

The procedure is as in the process of Preparation 2a, replacing the morpholine used in Step A by 1-methyl-piperazine and using 4.5 equivalents of 1M HCl solution in diethyl ether in Step D (salification step).

Preparation 3a: **4-[(*tert*-Butyldimethylsilyl)oxy]-*N*-methylaniline**

To a solution of 3.69 g 4-methylamino-phenol (30 mmol) and 3.20 g imidazole (47 mmol) in 65 mL of dichloromethane containing 1% ethanol, 5.88 g *tert*-butyl-dimethylsilyl chloride (39 mmol) is added with rapid stirring at ambient temperature. After 30 minutes the mixture is poured onto 160 mL water. The organic phase is separated and the aqueous phase is extracted with 50 mL dichloromethane. The combined organic phases are subsequently dried over magnesium sulphate and then filtered and evaporated to dryness. The crude product so obtained is then purified by chromatography over silica gel (hexane/ethyl acetate 100:1) to give the product (6.2 g, 87%) as an oil.

Preparation 3b: **4-(Benzyloxy)-*N*-phenylaniline**

Benzyl bromide (1.79 mL, 15.08 mmol) was added to a mixture of 4-hydroxydiphenylamine (2.54 g, 13.71 mmol), cesium carbonate (5.58g, 1.25 equiv) and potassium iodide (283 mg, 0.1 equiv) in acetone (20 mL).

The mixture was stirred and heated at 50°C for 1.25 h. After this time, a further portion of benzyl bromide (0.2 eq) and cesium carbonate (0.2 eq) were added and the mixture was heated for 45 min. The reaction mixture was allowed to cool to ambient temperature, then diluted with ethyl acetate, filtered, and the solvent removed *in vacuo*. The residue was purified by flash chromatography (Combiflash; 120g SilaSep Silica column) eluting with 0-20% ethyl acetate in hexane gradient to afford the desired product as a solid.

LC/MS (C₁₉H₁₇NO) 276 [M+H]⁺; RT 1.48 (Method B)

Preparation 3c: 4-(Benzyloxy)-N-ethylaniline

Step A: *tert*-Butyl-N-[4-(benzyloxy)phenyl]carbamate

To a mixture of *tert*-butyl-N-(4-hydroxyphenyl)carbamate (25 g, 0.12 mol) and potassium carbonate (24.77 g, 0.18 mol) in DMF (400 mL) was added benzyl bromide (22.48 g, 0.13 mmol) and the reaction was heated at 50 °C for 16 h. The reaction was allowed to cool to ambient temperature, and water (100 mL) was added, causing the precipitation of a white solid. Further water (500 mL) was added and the resultant suspension was stirred for 30 min. The solid material was isolated by filtration, washed with water and dried under vacuum. This solid material was then dissolved in dichloromethane (250 mL), dried over magnesium sulfate and concentrated *in vacuo* to afford the desired material (32.05 g, 0.11 mmol) as a white crystalline solid.

LC/MS (C₁₈H₂₁NO₃) 200 [M-Boc+H]⁺; RT 1.45 (Method B)

Step B: *tert*-Butyl-N-[4-(benzyloxy)phenyl]-N-ethylcarbamate

To a cooled solution of the compound obtained in Step A (5 g, 16.7 mmol) in THF (50 mL) was added sodium hydride (1.34 g, 33.4 mmol) portion-wise, and the resultant mixture was allowed to stir for 40 min. Iodoethane (2.69 mL, 33.4 mmol) was added and the reaction was allowed to stir at ambient temperature for 1 h and then at 40 °C for *ca* 16 h. The reaction was cooled, quenched with water and then extracted with ethyl acetate. The organic phase was washed successively with aqueous sodium thiosulfate solution, aqueous sodium bicarbonate solution, and brine. The organic extract was dried over magnesium sulphate, filtered and concentrated *in vacuo*, and then adsorbed onto isolate and purified by chromatography (CombiFlash R_f, 80 g RediSep™ silica cartridge) eluting in a gradient of iso-hexane to 15 % ethyl acetate in iso-hexane to afford the desired product (5.37 g, 16.4 mmol).

LC/MS (C₂₀H₂₅NO₃) 228 [M-Boc+H]⁺; RT 1.52 (Method B)

Step C: 4-(Benzyloxy)-N-ethylaniline

To a solution of compound obtained in Step B (5.37 g, 16.4 mmol, 1 eq) in dichloromethane (50 mL) was added trifluoroacetic acid (5 mL) and the mixture was stirred

at ambient temperature for 2 hours. The reaction mixture was diluted with water and basified with aqueous 1M sodium hydroxide. The organic extract was dried over magnesium sulphate, filtered, concentrated *in vacuo* and then adsorbed onto isolate and purified by chromatography (CombiFlash R_f, 40 g RediSep™ silica cartridge) eluting in a gradient of iso-hexane to 25 % ethyl acetate in iso-hexane to afford the product as a yellow oil (3.24 g, 14.25 mmol).

LC/MS (C₁₅H₁₇NO) 228 [M+H]⁺; RT 1.03 (Method B)

Preparation 3d: 4-(Benzyloxy)-N-propylaniline

The procedure is as in Preparation 3c, replacing iodoethane used in Step B with 1-iodopropane.

LC/MS (C₁₆H₁₉NO) 242 [M+H]⁺; RT 1.21 (Method B)

Preparation 3e: 4-(Benzyloxy)-N-butylaniline

The procedure is as in Preparation 3c, replacing iodoethane used in Step B with 1-iodobutane.

LC/MS (C₁₇H₂₁NO) 256 [M+H]⁺; RT 1.56 (Method B)

Preparation 3f: 4-({4-[(*tert*-Butyldimethylsilyl)oxy]phenyl}amino)-1,5-dimethyl-1H-pyrrole-2-carbonitrile

Step A: 4-Bromo-1,5-dimethyl-1H-pyrrole-2-carbonitrile

A solution of bromine (6.58 mL, 0.13 mol) in acetic acid (60 mL) is added dropwise with the aid of a dropping funnel to a solution of 1,5-dimethyl-1H-pyrrole-2-carbonitrile (15.0 g, 0.12 mol) in acetic acid (300 mL). The whole is stirred at ambient temperature for 24 hours. The reaction mixture is then poured into a beaker containing 300 mL of water. The solid formed is filtered off and rinsed with water. It is then dissolved in

dichloromethane (300 mL) and the organic phase is washed with brine, dried over sodium sulphate, filtered and concentrated *in vacuo* to give the expected product in the form of a solid.

^1H NMR (CDCl_3) δ ppm: 2.25 (s, 3 H), 3.67 (s, 3 H), 6.74 (s, 1 H)

5 **Step B:** 4-({4-[(*tert*-Butyldimethylsilyl)oxy]phenyl}amino)-1,5-dimethyl-1H-pyrrole-2-carbonitrile

A solution of the compound of the preceding step (1.5 g, 7.53 mmol), 4-[(*tert*-butyldimethylsilyl)oxy]aniline (2.02 g, 9.04 mmol), sodium *tert*-butoxide (1.45 g, 15.06 mmol) and 2-di-*tert*-butylphosphino-2',4',6'-triisopropylbiphenyl (0.13 g, 0.30 mmol) in toluene (20 mL) is purged with nitrogen.
10 Tris(dibenzylideneacetone)dipalladium(0) (0.28 g, 0.30 mmol) is added, and then the reaction mixture is heated at 90°C until the reaction is complete (monitored by TLC). Heating is stopped and the mixture is allowed to return to ambient temperature. Water (75 mL) is added and the mixture is extracted with ethyl acetate (3 x 75 mL). The
15 combined organic phases are washed with brine and then concentrated. The crude product is absorbed on silica gel and purified by flash chromatography over silica gel with a mixture of ethyl acetate and heptane (0 to 30%). The product so obtained is dissolved while hot in heptane and is allowed to precipitate with stirring at ambient temperature and then at 0°C. The solid is filtered off and the operation is repeated on the filtrate to give the
20 expected compound in the form of a solid.

^1H NMR (400 MHz, CDCl_3) δ ppm: 0.15 (s, 6 H), 0.97 (s, 9 H), 2.13 (s, 3 H), 3.66 (s, 3 H), 4.68 (br. s, 1 H), 6.49 (d, $J = 8.5$ Hz, 2 H), 6.64 (s, 1 H), 6.66 (d, $J = 8.7$ Hz, 2 H)

^{13}C NMR (100 MHz, CDCl_3) δ ppm: 4.34, 9.72, 18.30, 25.88, 32.94, 101.27, 114.37, 114.70, 116.41, 120.73, 124.52, 131.23, 141.54, 148.27

25 MS (ESI+): $[\text{M}+\text{H}]^+$ measured: 342.3

Preparation 4a: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: 1-Formylpiperidine-2-carboxylic acid

To a mixture of DL-pipecolinic acid (33.45 g, 259 mmol) in formic acid (250 mL) cooled to 0 °C, is added acetic anhydride (171 mL, 1.81 mol) drop-wise over *ca* 60 minutes. The reaction mixture is allowed to warm to room temperature, stirred for *ca* 16 hours and then cooled in an ice-water bath. Water (250 mL) is added, the mixture is stirred for 10 minutes and then concentrated *in vacuo*. Toluene is added and evaporated *in vacuo* (3 x 50 mL) to azeotropically remove water and acetic acid, and then the residue is dissolved in dichloromethane (60 mL), filtered through a hydrophobic frit and the filtrate evaporated *in vacuo* to afford the product as an oil.

Step B: Methyl 5,6,7,8-tetrahydroindolizine-1-carboxylate

To a stirring solution of formylated pipecolinic acid obtained in Step A (13.26g 84.48 mmol) in dichloroethane (100 mL) is added tosyl chloride (17.69 g, 92.8 mmol) followed by methyl- α -chloroacrylate (15.4 mL, 151.86 mmol). Triethylamine (23.52 mL, 168.74 mmol) is then added dropwise. The reaction mixture is stirred for 10 minutes, before heating to reflux. After 3 hours the reaction is cooled to room temperature and a further portion of methyl- α -chloroacrylate (6.0 mL, 59.1 mmol) is added followed by dropwise addition of triethylamine (11 mL, 78.9 mmol) and the reaction is heated at reflux for *ca.* 16 h. The reaction mixture is allowed to cool to room temperature, partitioned between dichloromethane and 1M HCl, filtered through a pad of celite and the phases separated. The organic phase is washed sequentially with 1N HCl, saturated NaHCO₃ solution and then brine. The organic extract is dried over magnesium sulphate, filtered and concentrated *in vacuo* and then adsorbed onto silica gel and purified by chromatography (CombiFlash R_f, 220g RediSep™ silica cartridge) eluting in a gradient of iso-hexane to 30% ethyl acetate in iso-hexane to obtain an oil.

LC/MS (C₁₀H₁₃NO₂) 180 [M+H]⁺; RT 1.13 (Method B)

Step C: 5,6,7,8-Tetrahydroindolizine-1-carboxylic acid

To a solution of the ester obtained in Step B (2 g, 11.2 mmol) in dioxane (15 mL) is added a solution of LiOH (936 mg, 22.3 mmol) in water (15 mL) and the reaction stirred at 100 °C for 5 hours. The reaction is cooled, diluted with water, acidified to ~ pH 2 with 2M HCl

and the resulting precipitate is filtered and washed with water and then dried under vacuum to afford the product as a powder.

LC/MS ($C_9H_{11}NO_2$) 166 $[M+H]^+$; RT 1.72 (Method A)

Step D: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The acid obtained in Step C (1 g, 6.05 mmol) is azeotroped with a minimal volume of toluene and then dissolved in anhydrous dichloromethane (50 mL). This is cooled to -10 °C under nitrogen and oxalyl chloride (2M in dichloromethane, 3 mL, 6.05 mmol) is added dropwise and then stirred for an hour maintaining the temperature at -10 °C. A solution of pyridine (0.73 mL, 9.08 mmol) and 4-(benzyloxy)-*N*-methylaniline (1.42 g, 6.7 mmol) in dichloromethane (3 mL) is added drop-wise to the reaction mixture at -10 °C and then allowed to warm to ambient temperature. Further pyridine (0.24 mL, 3 mmol) is added after 4 hours and stirring is maintained at ambient temperature for *ca.* 16 h. The reaction mixture is loaded onto a pre-packed silica column and purified in a gradient of *iso*-hexane to 40% ethyl acetate/*iso*-hexane to afford the desired product as a powder.

LC/MS ($C_{23}H_{24}N_2O_2$) 361 $[M+H]^+$; RT 2.68 (Method A)

Step E: *N*-[4-(Benzyloxy)phenyl]-3-bromo-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the compound obtained in Step D (3.49 g, 9.68 mmol) in tetrahydrofuran (35 mL), cooled to -78 °C under nitrogen, is added *N*-bromosuccinimide (1 eq) portion-wise and then the resultant mixture is stirred for 1 hour. The reaction is allowed to warm to ambient temperature, then diluted with ethyl acetate and washed with 10% sodium thiosulphate solution, saturated sodium bicarbonate solution and brine. The organic phase is subsequently dried over magnesium sulphate, filtered and concentrated *in vacuo*. The crude product is purified by chromatography over silica gel in a gradient of *iso*-hexane to 1:1 ethyl acetate/*iso*-hexane, then triturated with ether and filtered to afford a powder.

LC/MS ($C_{23}H_{23}N_2O_2Br$) 439 $[M+H]^+$; RT 2.74 (Method A)

Step F: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the bromide obtained in Step E (6.47g, 14.73 mmol) in anhydrous tetrahydrofuran (83 mL), cooled to -78°C, is added *n*-butyl lithium solution in hexanes (2.17 M, 7.47 mL, 16.20 mmol) drop-wise over *ca* 20 mins. After a further 15 minutes, 2-isopropyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (3.61 mL, 17.67 mmol) is added drop-wise and stirring is continued for 15 min. The reaction mixture is quenched at -78°C by the addition of 20 mL saturated ammonium chloride solution, and then allowed to warm to ambient temperature. The reaction mixture is partitioned between water and ethyl acetate and the phases separated. The organic phase is washed with water, saturated NaCl solution, then dried over magnesium sulphate, filtered and the filtrate concentrated *in vacuo* to a solid. The solid is then triturated with diethyl ether, filtered, washed with cold ether and dried *in vacuo* to afford the product as a powder.

LC/MS (C₂₉H₃₅BN₂O₄) 487 [M+H]⁺; RT 1.59 (Method B)

Preparation 4b: N-[4-(Benzyloxy)phenyl]-N,1,2-trimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrole-3-carboxamide

Step A: Ethyl 1,2-dimethyl-1H-pyrrole-3-carboxylate

To a cooled solution of 2-methyl-1H-pyrrole-3-carboxylate (5 g, 32.64 mmol) in THF (50 mL) is added 60% NaH in mineral oil (2.61 g, 65.28 mmol) and the resultant mixture is stirred at 0 °C for 40 minutes. Methyl iodide (4.07 mL, 65.28 mmol) is then added and allowed to stir for 1 hour. The reaction is quenched by the dropwise addition of water (20 mL) and then extracted with ethyl acetate (20 mL x 2). The organic extracts are washed sequentially with 10% aqueous thiosulphate solution and brine, then dried over magnesium sulphate, filtered and concentrated. The crude product is taken-up in dichloromethane and loaded onto isolate for purification on CombiFlash (120 g silica, Hex to 20% EtOAc/Hex) to yield the desired product as an oil.

LC/MS (C₉H₁₃NO₂) 168 [M+H]⁺; RT 1.11 (Method B)

Step B: 1,2-Dimethyl-1H-pyrrole-3-carboxylic acid

To a solution of the product from Step A (5.3 g, 31.7 mmol) in 1,4-dioxane (60 mL) is added 1M LiOH (2.66 g, 63.4 mmol) in water (60 ml) and the reaction stirred at 100 °C for *ca* 16 h. The reaction is allowed to cool to ambient temperature, diluted with water and acidified with 2M HCl. The resulting precipitate is collected by vacuum filtration to afford the desired product. The filtrate is extracted with ethyl acetate and the organic extracts dried over magnesium sulphate, filtered and concentrated to obtain an additional crop of desired product.

LC/MS (C₇H₉NO₂) no ionisation; RT 0.71 (Method B)

Step C: N-[4-(Benzyloxy)phenyl]-N,1,2-trimethyl-1H-pyrrole-3-carboxamide

The acid obtained from Step B (1 g, 7.19 mmol) is dissolved in dichloromethane (20 ml) and to this is added 1-chloro-N,N,2-trimethyl-1-propenylamine (1.15 g, 8.62 mmol) and stirred at ambient temperature for 2 hours. The mixture is concentrated to an oil which is re-dissolved in toluene (50 mL). A solution of 4-(benzyloxy)-N-methylaniline (1.84 g, 8.62 mmol) in toluene (10 ml) is added and the resultant mixture is stirred at reflux for 2 hours. The reaction is allowed to cool to ambient temperature and left to stand for *ca* 16 h. The reaction is partitioned between ethyl acetate and water, separated, and the organics are washed with water, dried over magnesium sulphate, filtered and concentrated. The crude product is purified on CombiFlash (80 g silica, Hex to 60% EtOAc) to obtain the desired product as a solid.

LC/MS (C₂₁H₂₂N₂O₂) 335 [M+H]⁺; RT 1.33 (Method B)

Step D: N-[4-(Benzyloxy)phenyl]-5-bromo-N,1,2-trimethyl-1H-pyrrole-3-carboxamide

The product from Step C (1.74 g, 5.19 mmol) is dissolved in THF (20 ml) and cooled to -78 °C. N-Bromosuccinimide (924 mg, 5.19 mmol) is then added portion-wise until complete addition and then stirred for 15 minutes. The reaction mixture is allowed to warm to ambient temperature, diluted with ethyl acetate and washed sequentially with 10% sodium thiosulfate solution, saturated sodium bicarbonate solution and brine. The organics are then dried over magnesium sulphate, filtered and concentrated. The crude material is taken-up in dichloromethane, loaded onto isolate and purified on CombiFlash (80 g silica, Hex to 70 % EtOAc/Hex) to afford the product as a solid.

LC/MS ($C_{21}H_{21}N_2O_2Br$) 413 $[M+H]^+$; RT 1.42 (Method B)

Step E: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide

The bromide from Step D (1.74 g, 4.21 mmol) and 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (0.94 g, 5.05 mmol) are combined and dissolved in anhydrous THF (20 ml). The reaction mixture is cooled to - 78 °C under nitrogen, followed by the dropwise addition of 2M *n*-BuLi (2.13 ml, 4.25 mmol) over 50 minutes. The reaction is then quenched by the addition of saturated aqueous ammonium chloride solution, and allowed to warm to ambient temperature, diluted in ethyl acetate and then washed with water followed by brine. The organics are dried over magnesium sulphate, filtered and concentrated to an oil, which solidifies on addition of ether. Re-evaporation and drying *in vacuo* affords the desired product.

LC/MS ($C_{27}H_{33}BN_2O_4$) 461 $[M+H]^+$; RT 1.51 (Method B)

Preparation 4c: *N*,1,2-Trimethyl-*N*-phenyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 4b, replacing 4-(benzyloxy)-*N*-methylaniline used in Step C with *N*-methylaniline.

LC/MS ($C_{20}H_{27}BN_2O_3$) 355 $[M+H]^+$; RT 1.39 (Method B)

Preparation 4d: *N,N*-Dibutyl-1,2-dimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 4b, replacing 4-(benzyloxy)-*N*-methylaniline used in Step C with dibutylamine

LC/MS ($C_{21}H_{37}BN_2O_3$) 377 $[M+H]^+$; RT 1.55 (Method B)

Preparation 4e: *N*-{4-[(*tert*-Butyldimethylsilyl)oxy]phenyl}-*N*-methyl-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Preparation 4a, replacing 4-(benzyloxy)-*N*-methylaniline in Step D with 4-[(*tert*-butyldimethylsilyl)oxy]-*N*-methylaniline from Preparation 3a.

5 **Preparation 4f:** *N*-[4-(Benzyloxy)phenyl]-*N*-phenyl-3-(tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Preparation 4a, replacing 4-(benzyloxy)-*N*-methylaniline in Step D with 4-(benzyloxy)-*N*-phenylaniline from Preparation 3b.

10 **Preparation 4g:** *N*-[4-(Benzyloxy)phenyl]-*N*-ethyl-1,2-dimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 4b, replacing 4-(benzyloxy)-*N*-methylaniline used in Step C with 4-(benzyloxy)-*N*-ethylaniline from Preparation 3c.

LC/MS (C₂₈H₃₅BN₂O₄) 475 [M+H]⁺; RT 1.49 (Method B)

15 **Preparation 4h:** *N*-[4-(Benzyloxy)phenyl]-1,2-dimethyl-*N*-propyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 4b, replacing 4-(benzyloxy)-*N*-methylaniline used in Step C with 4-(benzyloxy)-*N*-propylaniline from Preparation 3d.

LC/MS (C₂₉H₃₇BN₂O₄) 489 [M+H]⁺; RT 1.57 (Method B)

20 **Preparation 4i:** *N*-[4-(Benzyloxy)phenyl]-*N*-butyl-1,2-dimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 4b, replacing 4-(benzyloxy)-*N*-methylaniline used in Step C with 4-(benzyloxy)-*N*-butylaniline from Preparation 3e.

LC/MS (C₃₀H₃₉BN₂O₄) 503 [M+H]⁺; RT 1.55 (Method B)

Preparation 5a: 6-(1-{[4-(Benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-2-[(*tert*-butoxy)carbonyl]-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid

Step A: 2-*tert*-Butyl 7-methyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

The boronic ester obtained in Preparation 4a (1.22 g, 2.50 mmol) and the triflate obtained in Preparation 1b (1.1 g, 2.50 mmol) are suspended in 15 mL of anhydrous *N,N*-dimethylformamide and the mixture is degassed (bubbling with N₂) for 45 minutes. Cs₂CO₃ (1.63 g, 5 mmol) and bis(di-*tert*-butyl(4-dimethylaminophenyl)-phosphine)dichloropalladium(II) (88.5 mg, 0.13 mmol) are added and the resultant mixture is sealed and immediately heated in the microwave at 130°C for 30 minutes. The reaction mixture is concentrated and re-dissolved in ethyl acetate, washed with brine, dried over magnesium sulphate, filtered and concentrated. The crude product is purified by flash chromatography on CombiFlash (120 g silica, dichloromethane to 20% methanol/dichloromethane) to afford the desired product.

LC/MS (C₃₉H₄₃N₃O₆) 650 [M+H]⁺; RT 1.54 (Method B)

Step B: 6-(1-{[4-(Benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-2-[(*tert*-butoxy)carbonyl]-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid

To a solution of the ester obtained in Step A (730 mg, 1.12 mmol) in dioxane (10 mL) is added a solution of LiOH (94 mg, 2.24 mmol) in water (5 mL). The reaction is then heated at 90 °C for *ca* 16 h. A further 94 mg of LiOH (2.24 mmol) in water (5 mL) is added and stirred for 1 hour to complete the reaction. The mixture is cooled, diluted with water and acidified to ~ pH 4 with dilute aqueous HCl. The precipitate which forms is stirred for 30

minutes and then filtered off, washed with cold water and dried under vacuum to afford the desired product.

LC/MS ($C_{38}H_{41}N_3O_6$) 636 $[M+H]^+$; RT 1.46 (Method B)

Preparation 5b: 2-*tert*-Butyl 7-lithio 6-(4-{[4-

(benzyloxy)phenyl](methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

Step A: 2-*tert*-Butyl 7-methyl 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

The boronic ester from *Preparation 4b* (1 g, 1.7 mmol) and the nonaflate from *Preparation 1a* (0.94 g, 2.04 mmol) are dissolved in DMF (20 mL) and degassed (bubbling nitrogen) for 20 minutes. Cesium carbonate (1.3 g, 4.04 mmol) and bis(di-*tert*-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (60 mg, 0.085 mmol) are added and the resultant mixture is immediately heated in the microwave at 130 °C for 30 minutes. The DMF is evaporated, and the residue dissolved in ethyl acetate, washed with brine, then dried over magnesium sulphate, filtered and concentrated. The crude material is taken up in dichloromethane, loaded onto isolate and purified on CombiFlash (120 g silica, Hex to 70 % EtOAc/Hex to obtain the product as a foam.

LC/MS ($C_{37}H_{41}N_3O_6$) 624 $[M+H]^+$; RT 1.55 (Method B)

Step B: 2-*tert*-Butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

To a solution of the ester obtained in Step A (2.17 g, 3.48 mmol) in dioxane (20 ml) is added an aqueous solution of LiOH (2 M; 3.48 mL, 6.96 mmol), and the resultant mixture is heated at reflux for *ca* 16 h. The reaction mixture is allowed to cool to ambient temperature and the solids removed by filtration. The solution is concentrated to obtain the desired product as a solid.

LC/MS ($C_{36}H_{39}N_3O_6$) 610 $[M+H]^+$; RT 1.47 (Method B)

Preparation 5c: 2-[(*tert*-Butoxy)carbonyl]-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid

Step A: 2-*tert*-Butyl 7-methyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

The procedure is as in Step A of Procedure 5b, replacing *N*-[4-(benzyloxy)phenyl]-*N*,1,2-trimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide with the product from Procedure 4c.

Step B: 2-[(*tert*-Butoxy)carbonyl]-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid

The procedure is as in Step B from Procedure 5a, replacing 2-*tert*-butyl 7-methyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate with the product from Step A.

Preparation 5d: 2-*tert*-Butyl 7-lithio 6-[4-(diethylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

The procedure is as in Preparation 5b, replacing *N*-[4-(benzyloxy)phenyl]-*N*,1,2-trimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide with the product from Procedure 4d.

Preparation 5e: 2-[(*tert*-Butoxy)carbonyl]-6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid

Step A: *tert*-Butyl 6-[1-({4-[(*tert*-butyldimethylsilyl)oxy]phenyl}(methyl)carbamoyl)-5,6,7,8-tetrahydroindolizin-3-yl]-7-formyl-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The triflate from Preparation 1c (610 mg, 1.49 mmol), the boronic ester from Preparation 4e (910 mg, 1.79 mmol) and potassium carbonate (412 mg, 2.98 mmol) are dissolved in THF/Water and degassed (nitrogen bubbling) for 15 mins. To this is added tetrakis(triphenylphosphine)palladium(0) (5mol%) and the resultant mixture is stirred at room temperature for 45 min. The reaction is diluted with ethyl acetate, washed with water followed by brine, and the organics dried over magnesium sulphate and evaporated under reduced pressure. The crude product thus obtained is purified on CombiFlash (40 g silica; isohexane to ethyl acetate), affording the desired product.

LC/MS ($C_{37}H_{49}N_3O_5Si$) 644 $[M+H]^+$; RT 1.71 (Method B)

Step B: 2-tert-Butyl 7-methyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

The procedure is as in Step B of Preparation 1a, replacing the 2:1 isomer mixture of *tert*-butyl 7-formyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline-2-carboxylate and *tert*-butyl 5-formyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline-2-carboxylate with *tert*-butyl 6-[1-({4-[(*tert*-butyldimethylsilyl)oxy]phenyl}(methyl)carbamoyl)-5,6,7,8-tetrahydroindolizin-3-yl]-7-formyl-1,2,3,4-tetrahydroisoquinoline-2-carboxylate. The reaction was heated at 45 °C external temperature for 24 h to facilitate completion of the observed partial desilylation.

Step C: 2-[(tert-Butoxy)carbonyl]-6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid

The procedure is as in Step B of Preparation 5a, replacing 2-*tert*-butyl 7-methyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate with 2-*tert*-butyl 7-methyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate.

Preparation 5f: 2-tert-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](ethyl)carbamoyl}-1,5-dimethyl-1H-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

The procedure is as in Preparation 5b, replacing *N*-[4-(benzyloxy)phenyl]-*N*,1,2-trimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide with the product from Procedure 4g.

LC/MS ($C_{37}H_{41}N_3O_6$) 624 $[M+H]^+$; RT 1.50 (Method B)

5 **Preparation 5g:** 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](propyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate

10 The procedure is as in Preparation 5b, replacing *N*-[4-(benzyloxy)phenyl]-*N*,1,2-trimethyl-5-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrrole-3-carboxamide with the product from Procedure 4h.

LC/MS ($C_{38}H_{43}N_3O_6$) 638 $[M+H]^+$; RT 1.54 (Method B)

Preparation 6aa: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide, bis-trifluoroacetic acid salt

15 **Step A:** *tert*-Butyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

20 To a solution of the acid obtained in *Preparation 5a* (150 mg, 0.24 mmol) in *N,N*-dimethylformamide (2 mL) is added triethylamine (0.13 mL, 0.96 mmol), HBTU (91 mg, 0.24 mmol) and the (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride obtained in *Preparation 2a* (72 mg, 0.24 mmol). The reaction is stirred at room temperature for 15 minutes and then diluted with water, and the resulting suspension stirred. The product is then extracted with ethyl acetate and the organic extract dried over magnesium sulphate, filtered and concentrated. The crude material is purified on
25 CombiFlash (12 g silica, dichloromethane to 4% methanol/dichloromethane) and then dried *in vacuo* to obtain the product as a glassy solid.

LC/MS (C₅₂H₅₉N₅O₆) 850 [M+H]⁺; RT 1.49 (Method B)

Step B: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide, bis-trifluoroacetic acid salt

The Boc-protected material obtained in Step A (173 mg, 0.2 mmol) is dissolved in dichloromethane (10 mL) and to this is added trifluoroacetic acid (1 mL). The reaction is allowed to stir for 1 hour at ambient temperature and then concentrated *in vacuo*. The residue is triturated in diethyl ether to afford a precipitate which is filtered off and dried under vacuum.

LC/MS (C₄₇H₅₁N₅O₄) 750 [M+H]⁺; RT 1.10 (Method B)

Preparation 6ab: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; trifluoroacetic acid salt

Step A: *tert*-Butyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbonyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a stirring solution of the acid obtained from *Preparation 5a* (0.87 g, 1.36 mmol) in DMF (10 mL) was added diisopropylethylamine (0.47 mL, 2.72 mmol) and (3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline from *Preparation 2b* (210 mg, 1.43 mmol) followed by HBTU (516 mg, 1.36 mmol). The reaction was stirred for 1 h, then diluted with water and extracted with ethyl acetate. The organic extracts were sequentially washed with aqueous sodium bicarbonate and brine, then dried over magnesium sulphate and evaporated *in vacuo*. The crude product was taken-up in dichloromethane, loaded onto isolate and purified on CombiFlash (40 g silica, isohexane to ethyl acetate gradient) to afford the desired product as a foam.

LC/MS (C₄₈H₅₂N₄O₅) 765 [M+H]⁺; RT 1.62 (Method B)

Step B: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; trifluoroacetic acid salt

The Boc-protected product from Step A (0.90 g, 1.18 mmol) was dissolved in dichloromethane (10 mL) and to this was added TFA (1 mL). The reaction was stirred for 1 h, then solvent was removed under reduced pressure, and residual solvent was removed under high vacuum. Addition of ether to the resultant oil induced the oil to solidify. The mixture was stirred for 45 min, then cooled, filtered, and washed with cold ether to afford the desired product. Concentration of the filtrate afforded further solid product. The combined solids were dried under vacuum to afford the product.

LC/MS (C₄₃H₄₄N₄O₃) 665 [M+H]⁺; RT 1.24 (Method B)

Preparation 6ba: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide; bis trifluoroacetic acid salt

Step A: *tert*-Butyl 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The lithium salt from *Preparation 5b* (1.07 g, 1.74 mmol) is taken-up in DMF (10 mL) and to this is added triethylamine (0.97 ml, 6.96 mmol), HBTU (660 mg, 1.74 mmol), and the (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride obtained in *Preparation 2a* (0.58 g, 1.91 mmol) and the resultant mixture is then stirred for 15 minutes at ambient temperature. The reaction is diluted with water and extracted with ethyl acetate. The organic extract is dried over magnesium sulphate, filtered and concentrated. The crude material is taken up in dichloromethane, loaded onto isolate and purified on CombiFlash (40 g silica, dichloromethane to 5% MeOH/dichloromethane) to afford the desired product.

LC/MS (C₅₀H₅₇N₅O₆) 824 [M+H]⁺; RT 1.44 (Method B)

Step B: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide; bis trifluoroacetic acid salt

The Boc-protected product from Step A (705 mg, 0.86 mmol) is dissolved in dichloromethane and to this is added trifluoroacetic acid (1.33 ml). The resultant mixture is then allowed to stir at ambient temperature for *ca* 16 h. The reaction is concentrated *in vacuo* and the residue triturated with ether to afford a precipitate which is filtered and dried under vacuum to yield the desired product.

LC/MS (C₄₅H₄₉N₅O₄) 724 [M+H]⁺; RT 1.07 (Method B)

Preparation 6bb: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide; trifluoroacetic acid salt

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with (3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline from Preparation 2b.

LC/MS (C₄₁H₄₂N₄O₃) 639 [M+H]⁺; RT 1.18 (Method B)

Preparation 6bc: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-[2-(morpholin-4-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide; bis-trifluoroacetic acid salt

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with (3*S*)-3-[2-(morpholin-4-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline dihydrochloride obtained from Preparation 2d.

LC/MS (C₄₆H₅₁N₅O₄) 738 [M+H]⁺; RT 1.07 (Method B)

Preparation 6bd: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{7-[(3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with (3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline dihydrochloride from Preparation 2f. The intermediate was isolated as the free base.

LC/MS (C₄₇H₅₃N₅O₄) 752 [M+H]⁺; RT 1.03 (Method B)

Preparation 6be: *N*-[4-(Benzyloxy)phenyl]-5-{7-[(3*S*)-3-[(dimethylamino)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide; bis-trifluoroacetic acid salt

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with dimethyl[(3*S*)-1,2,3,4-tetrahydroisoquinolin-3-ylmethyl]amine dihydrochloride from Preparation 2g.

LC/MS (C₄₃H₄₇N₅O₃) 682 [M+H]⁺; RT 1.02 (Method B)

Preparation 6bf: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide; tris-trifluoroacetic acid salt

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with (3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline trihydrochloride from Preparation 2h.

LC/MS (C₄₆H₅₂N₆O₃) 737 [M+H]⁺; RT 1.04 (Method B)

Preparation 6ca: *N*,1,2-Trimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-phenyl-1*H*-pyrrole-3-carboxamide; bis-trifluoroacetic acid salt

The procedure is as in Preparation 6ba, replacing 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate in Step A with 2-[(*tert*-butoxy)carbonyl]-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid from Preparation 5c.

LC/MS (C₃₈H₄₃N₅O₃) 618 [M+H]⁺; RT 0.93 (Method B)

Preparation 6cb: *N*,1,2-Trimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-phenyl-1*H*-pyrrole-3-carboxamide; trifluoroacetic acid salt

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with (3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline from Preparation 2b; and replacing 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate in Step A with 2-[(*tert*-butoxy)carbonyl]-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid from Preparation 5c.

Preparation 6cd: *N*,1,2-Trimethyl-5-{7-[(3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-phenyl-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with (3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline dihydrochloride from Preparation 2f; and replacing 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-

yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate in Step A with 2-[(*tert*-Butoxy)carbonyl]-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid from Preparation 5c. The intermediate was isolated as the free base.

5 **LC/MS** (C₄₀H₄₇N₅O₃) 646 [M+H]⁺; RT 0.91 (Method B)

Preparation 6ce: *N*,1,2-trimethyl-5-{7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-phenyl-1*H*-pyrrole-3-carboxamide; tris-trifluoroacetic acid salt

10 The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with (3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline trihydrochloride from Preparation 2h, and replacing 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate in Step A with
15 2-[(*tert*-butoxy)carbonyl]-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid from Preparation 5c.

LC/MS (C₃₉H₄₆N₆O₂) 631 [M+H]⁺; RT 0.91 (Method B)

Preparation 6da: *N,N*-Dibutyl-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide; bis-trifluoroacetic acid salt

20 The procedure is as in Preparation 6ba, replacing 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate in Step A with 2-*tert*-butyl 7-lithio 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate from Preparation 5d.

25 **LC/MS** (C₃₉H₅₃N₅O₃) 640 [M+H]⁺; RT 1.10 (Method B)

Preparation 6db: *N,N*-Dibutyl-1,2-dimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide; trifluoroacetic acid salt

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with (3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline from Preparation 2b; and replacing 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate in Step A with 2-*tert*-butyl 7-lithio 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate from Preparation 5d.

Preparation 6dc: *N,N*-Dibutyl-1,2-dimethyl-5-[7-(1,2,3,4-tetrahydroisoquinoline-2-carbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 6ba, replacing (3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline dihydrochloride in Step A with tetrahydroisoquinoline; and neutralising the product solution in Step B with dilute aqueous sodium hydroxide solution.

LC/MS (C₃₄H₄₄N₄O₂) 541 [M+H]⁺; RT 1.14 (Method B)

Preparation 6e: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide dihydrochloride

Step A: *tert*-Butyl 6-{1-[4-hydroxyphenyl](methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step A of Preparation 6aa, replacing 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-2-[(*tert*-

butoxy)carbonyl]-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid in Step A with 2-[(*tert*-butoxy)carbonyl]-6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-1,2,3,4-tetrahydroisoquinoline-7-carboxylic acid from Preparation 5e.

5 **LC/MS** (C₄₅H₅₃N₅O₆) 760 [M+H]⁺; RT 1.25 (Method B)

Step B: N-(4-Hydroxyphenyl)-N-methyl-3-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide dihydrochloride

10 The compound obtained in Step A (135 mg, 0.18 mmol) is dissolved in ethyl acetate (6.75 mL) and to this concentrated HCl (148 uL, 1.78 mmol) is added and allowed to stir for 30 minutes. Minimal IPA is then added dropwise to encourage precipitation. The solid is then filtered off, washed with ethyl acetate and dried under vacuum overnight to afford a solid.

LC/MS (C₄₀H₄₅N₅O₄) 660 [M+H]⁺; RT 0.89 (Method B)

15 ***Preparation 6f: N-[4-(Benzyloxy)phenyl]-N-ethyl-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide***

20 The procedure is as in Preparation 6ba, replacing 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate in Step A with 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](ethyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate from Preparation 5f; and neutralising the product solution in Step B with dilute aqueous sodium hydroxide solution.

LC/MS (C₄₆H₅₁N₅O₄) 738 [M+H]⁺; RT 1.08 (Method B)

Preparation 6g: *N*-[4-(Benzyloxy)phenyl]-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-propyl-1*H*-pyrrole-3-carboxamide

The procedure is as in Preparation 6ba, replacing 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate in Step A with 2-*tert*-butyl 7-lithio 6-(4-{[4-(benzyloxy)phenyl](propyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-1,2,3,4-tetrahydroisoquinoline-2,7-dicarboxylate from Preparation 5g; and neutralising the product solution in Step B with dilute aqueous sodium hydroxide solution.

LC/MS (C₄₇H₅₃N₅O₄) 752 [M+H]⁺; RT 1.17 (Method B)

Preparation 6h: *N*-[4-(Benzyloxy)phenyl]-*N*-butyl-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

Step A: *N*-[4-(benzyloxy)phenyl]-*N*-butyl-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

A solution of tetrahydrofuran/ water (5:1, 7 mL) was degassed for 20 min. To a mixture of the boronate ester from Preparation 4i (246 mg, 0.49 mmol) and the bromide from Preparation 1ha (332.78 mg, 0.59 mmol) followed by 7mL of the stock solution. This was degassed for a further 5 min and to this solution was added cesium carbonate (319.05 mg, 0.98 mmol, 2 eq) and bis(di-*tert*-butyl(4-dimethylaminophenyl)phosphine) dichloropalladium(ii) (17.33 mg, 0.02 mmol, 0.05 eq) and then heated to 95 °C in the microwave for 40 min. The reaction was diluted in ethyl acetate and washed with brine. The organic extract was dried over magnesium sulfate and concentrated *in vacuo*, and adsorbed onto isolate and purified on CombiFlash (12 g silica, dichloromethane to 4% methanol in dichloromethane).

LC/MS (C₅₀H₅₄F₃N₅O₅) 862 [M+H]⁺; RT 1.53 (Method B)

Step B: *N*-[4-(benzyloxy)phenyl]-*N*-butyl-1,2-dimethyl-5-{7-[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

To a solution of the product from Step A (138 mg, 0.16 mmol) in ethanol (6 mL) was added a solution of potassium carbonate (132.75 mg, 0.96 mmol) in water (1.2 mL) and the mixture was allowed to stir at reflux for 1 h. The reaction was allowed to cool to ambient temperature and the organic solvent was concentrated *in vacuo*. The solution left behind was washed with Ethyl Acetate. The organic phase was washed with water and saturated sodium chloride. The organic extract was dried over magnesium sulfate and concentrated *in vacuo*. The product was used directly in the next step assuming quantitative transformation.

LC/MS (C₄₈H₅₅N₅O₄) 766 [M+H]⁺; RT 1.23 (Method B)

Preparation 7aa: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{6-[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *tert*-Butyl 5-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-6-[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

A mixture of bromide from Preparation 1e (845 mg, 1.45 mmol), the boronic ester from Preparation 4a (886 mg, 1.82 mmol) and cesium carbonate (991 mg, 3.04 mmol) in THF (9.0 mL) and water (3.6 mL) is degassed by purging with nitrogen, followed by bubbling nitrogen through for 5 min. The mixture is heated to 60 °C, at which point the PdCl₂(Ata-Phos)₂ catalyst (54 mg, 0.08 mmol) is added and the reaction is allowed to stir for 3 h.

The mixture is allowed to cool to ambient temperature, then diluted with ethyl acetate, and washed sequentially with water and brine. The organic phase is dried over magnesium sulphate, and concentrated *in vacuo*.

The crude material was purified on by flash column chromatography (40 g silica; dichloromethane to 5 % MeOH/dichloromethane) to afford the product as a glassy solid.

LC/MS (C₅₁H₅₇N₅O₆) 836 [M+H]⁺; RT 1.53 (Method B)

Step B: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a stirred solution of the product from Step A (850 mg, 1.02 mmol) in dichloromethane (20 mL) is added trifluoroacetic acid (5 mL) and the mixture is stirred for 1 h. Water is added, followed by 2N aqueous sodium hydroxide solution until the aqueous phase was basic. The organic phase is then separated, dried over magnesium sulphate, and concentrated *in vacuo* to afford the product as a solid.

LC/MS (C₄₆H₄₉N₅O₄) 736 [M+H]⁺; RT 1.12 (Method B)

Preparation 7ab: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; trifluoroacetic acid salt

Step A: *tert*-Butyl 5-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

A mixture of THF (20 mL) and water (8 mL) is degassed by multiple evacuation / nitrogen purge cycles and bubbling nitrogen through the mix. 20 mL of this mixture is subsequently added *via* syringe to a mixture of the bromide from Preparation 1f (0.88g, 1.87 mmol), the boronic ester from Preparation 4a (1.09 g, 2.25 mmol) and cesium carbonate (1.22 g, 3.74 mmol) under nitrogen. PdCl₂(Ata-Phos)₂ (66 mg, 0.09 mmol) is then added and the reaction is immediately heated under microwave irradiation at 95°C for 20 min.

The reaction mixture is diluted with ethyl acetate (100 mL) and sequentially washed with water (2 x 50 mL), then saturated NaCl (aq) (2 x 50 mL), dried over magnesium sulphate, filtered and concentrated *in vacuo*. The crude product was purified by flash

chromatography (Combiflash, 80g, eluting with gradient 0 to 80% ethyl acetate in hexane) to afford the product as a foam.

LC/MS (C₄₇H₅₀N₄O₅) 751 [M+H]⁺; RT 1.61 (Method B)

Step B: *N-[4-(Benzyloxy)phenyl]-N-methyl-3-{6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; trifluoroacetic acid salt*

To a solution of the product from Step A (152 mg, 0.20 mmol) in dichloromethane (3 mL) is added trifluoroacetic acid (0.7 mL) and the reaction is stirred at ambient temperature for *ca* 16 h. Solvent was removed *in vacuo* and the resulting solid is triturated with ether collected by filtration.

LC/MS (C₄₂H₄₂N₄O₃) 651 [M+H]⁺; RT 2.39 (Method A)

Preparation 7ba: *N-(4-Benzyloxyphenyl)-N,1,2-trimethyl-5-[6-[(3S)-3-(morpholinomethyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]isoindolin-5-yl]pyrrole-3-carboxamide hydrochloride*

Step A: *tert-Butyl 5-bromo-6-[[(3S)-3-(morpholinomethyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]isoindoline-2-carboxylate*

6-Bromo-2-*tert*-butoxycarbonyl-isoindoline-5-carboxylic acid (0.70 g), TBTU (0.823 g) and DMAP (25 mg) was dissolved dichloromethane (35 ml) and diisopropylethylamine (1.75 ml) was added. After 3 minutes stirring at room temperature 4-[[[(3S)-1,2,3,4-tetrahydroisoquinolin-3-yl]methyl]morpholine hydrochloride (0.656 g) was added and stirred for 1h. After the completion of the reaction it was diluted with dichloromethane (100 ml) and washed with water (50 ml), dried over sodium sulfate and evaporated. The crude product was purified by flash chromatography over silicagel (dichloromethane / methanol, gradient 1-10%) to give the title compound.

High-resolution mass (ESI+):

Empirical formula: C₂₈H₃₄BrN₃O₄

$[M+H]^+$ calculated: 556.1813

$[M+H]^+$ measured: 556.1790

IR v: C-H: 2930 cm^{-1} ; $>\text{C}=\text{O}$: 1697, 1635 cm^{-1} ; C-O-C: 1113 cm^{-1}

Step B: *tert-Butyl 5-[4-[(4-benzyloxyphenyl)-methyl-carbamoyl]-1,5-dimethyl-pyrrol-2-yl]-6-[(3S)-3-(morpholinomethyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]isoindoline-2-carboxylate*

The product from Step A (284 mg, 0.51 mmol) and *N*-(4-benzyloxyphenyl)-*N*,1,2-trimethyl-pyrrole-3-carboxamide (205 mg, 0.61 mmol) from Step C of Preparation 4b were dissolved in dimethylacetamide (5 mL) then nitrogen was bubbled through the solution for 5 min. Potassium acetate (0.11 g, 1.12 mmol) and bis(triphenylphosphine)palladium(II) dichloride (20 mg) were added to the mixture then it was heated up to 140 °C then after 20 min stirring water (20 μL) was added. Stirring at 140 °C under nitrogen atmosphere was continued for additional 16 hours. The reaction mixture was allowed to cool to room temperature then evaporated. The residue was partitioned between dichloromethane (50 mL) and water (10 mL). The organic phase was washed with water (10 mL), dried over Na_2SO_4 and evaporated. The crude product was purified by flash chromatography over silicagel (dichloromethane / methanol, gradient 1-10%) to give the title compound.

Step C: *N*-(4-benzyloxyphenyl)-*N*,1,2-trimethyl-5-[6-[(3S)-3-(morpholinomethyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]isoindolin-5-yl]pyrrole-3-carboxamide hydrochloride

The product from Step B (140 mg) was stirred in a 4M HCl in dioxane (15 mL) solution for 30 min at room temperature. After the completion of the reaction all the solvents were removed to yield 170 mg of the title product.

Preparation 7bb: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindol-5-yl}-1H-pyrrole-3-carboxamide

Step A: *tert-Butyl 5-(4-{{4-(benzyloxy)phenyl}}(methyl)carbamoyl}-1,5-dimethyl-1H-pyrrol-2-yl)-6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxylate*

A mixture of *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide from Step C of Preparation 4b (150 mg, 0.45 mmol), the bromide from Preparation 1f (317 mg, 0.67 mmol), potassium acetate (88 mg, 0.9 mmol) in dimethylacetamide (2 mL) is degassed by bubbling with nitrogen gas for 20 min. Bis(triphenylphosphine)palladium dichloride (31.5 mg, 0.04 mmol) is added, and the reaction is heated at 144 °C. Over the next 8 h, 2 further 0.1 eq portions of bis(triphenylphosphine)palladium dichloride are added. The reaction is then left stirring for *ca* 16 h. The reaction is allowed to cool to ambient temperature, and is concentrated *in vacuo*. The residue is partitioned between ethyl acetate and water, separated, and the organic phase is washed with water, dried (magnesium sulphate), and concentrated *in vacuo*. The crude material is purified by flash column chromatography (24 g silica, isohexane to ethyl acetate gradient) to afford the desired product.

LC/MS (C₄₅H₄₈N₄O₅) 725 [M+H]⁺; RT 1.57 (Method B)

Step B: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-1*H*-pyrrole-3-carboxamide

To a solution of the product from Step A (260 mg, 0.36 mmol) in dichloromethane (5 mL) is added trifluoroacetic acid (1 mL) and the mixture is stirred at ambient temperature for *ca* 16 h. The mixture is partitioned between water and dichloromethane, and basified with 1M aqueous sodium hydroxide. The organic phase is dried (magnesium sulphate), evaporated, and purified on CombiFlash (12 g silica, dichloromethane to 5% methanol/dichloromethane) to afford the product as a glassy solid.

LC/MS (C₄₀H₄₀N₄O₃) 625 [M+H]⁺; RT 1.20 (Method B)

Preparation 7fa: *N*-(4-Benzyloxyphenyl)-*N*-methyl-3-[6-[(3*S*)-3-(morpholinomethyl)-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindolin-5-yl]indolizine-1-carboxamide hydrochloride

Step A: *N*-(4-Benzyloxyphenyl)-*N*-methyl-indolizine-1-carboxamide

5 Indolizine-1-carboxylic acid (0.4 g, 5.6 mmol) was dissolved in dichloromethane/THF mixture (40/40 ml) then 1-chloro-*N,N*,2-trimethyl-1-propenylamine (0.8 mL) was added and the mixture was stirred at room temperature for 10 minutes. After the completion of the acyl chloride formation, the mixture was evaporated and redissolved in dry dichloromethane (50 mL). 4-Benzyloxy-*N*-methylaniline hydrochloride (1.67 g, 6.7 mmol)
10 and triethylamine (2.34 mL, 16.75 mmol) were added and the reaction was stirred at ambient temperature for *ca.* 16 h. The solution was washed with saturated aqueous NaHCO₃ solution, and brine, then dried over Na₂SO₄ and evaporated. The crude product was purified by flash chromatography over silicagel (dichloromethane / methanol, gradient 1-10%) to give the desired material.

15 **Step B:** *tert*-Butyl 5-[1-[(4-benzyloxyphenyl)-methyl-carbamoyl]indolizin-3-yl]-6-[(3*S*)-3-(morpholinomethyl)-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindoline-2-carboxylate

The compound of Preparation 1e (278 mg, 0.5 mmol) and the product from Step A (240 mg, 0.70 mmol) were dissolved in dimethylacetamide (5 mL) then nitrogen was bubbled through the solution for 5 min. Potassium acetate (0.11 g, 1.12 mmol) and
20 bis(triphenylphosphine)palladium(II) dichloride (20 mg) were added to the mixture, it was heated to 110 °C then after 20 min stirring water (20 µL) was added. Stirring at 110 °C under a nitrogen atmosphere was continued for additional 5 hours. The reaction mixture was allowed to cool to ambient temperature then evaporated. The residue was dissolved in THF, filtered through a pad of Celite and after removal of solvent by evaporation the crude
25 product was purified by preparative HPLC (H₂O-TFA/acetonitrile; gradient elution). The pH of the appropriate combined fractions was adjusted to 10 by Na₂CO₃ then the acetonitrile was removed under reduced pressure. The precipitated solid was filtered then dried to yield the title compound.

30 **Step C:** *N*-(4-Benzyloxyphenyl)-*N*-methyl-3-[6-[(3*S*)-3-(morpholinomethyl)-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindolin-5-yl]indolizine-1-carboxamide hydrochloride

The product from Step B (167 mg) was stirred in a 4M HCl in dioxane (15 ml) solution for 30 min at room temperature. After the completion of the reaction all the solvents were removed to afford the desired product.

Preparation 7fb: *N*-(4-Benzoyloxyphenyl)-*N*-methyl-3-[6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindolin-5-yl]indolizine-1-carboxamide hydrochloride

Step A: *tert*-Butyl 5-[1-[(4-benzoyloxyphenyl)-methyl-carbamoyl]indolizin-3-yl]-6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindoline-2-carboxylate

The procedure is as in Step B of Preparation 7fa, replacing *tert*-Butyl 5-bromo-6-[(3*S*)-3-(morpholinomethyl)-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindoline-2-carboxylate with the compound of Preparation 1f. The product is purified by preparative HPLC (H₂O-TFA/acetonitrile; gradient elution).

Step B: *N*-(4-benzoyloxyphenyl)-*N*-methyl-3-[6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindolin-5-yl]indolizine-1-carboxamide hydrochloride

The procedure is as in Step C of Preparation 7fa, replacing *tert*-butyl 5-[1-[(4-benzoyloxyphenyl)-methyl-carbamoyl]indolizin-3-yl]-6-[(3*S*)-3-(morpholinomethyl)-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindoline-2-carboxylate with the product from Step A.

The following Examples are obtained using the compounds of the appropriate Preparations described above. When no specific procedure is detailed, the corresponding Example can be obtained by repeating the procedure described for analogues Examples (*i.e.* whose structure is close) detailed elsewhere in the specification, choosing the appropriate starting materials and using the basic knowledge of the man skilled in the art.

Example 1: *tert*-Butyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is described in Step A of Preparation 6e.

5 **LC/MS** (C₄₅H₅₃N₅O₆) 760 [M+H]⁺; RT 1.25 (Method B)

Example 2: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide dihydrochloride

10 The procedure is described in Step B of Preparation 6e.

LC/MS (C₄₀H₄₅N₅O₄) 660 [M+H]⁺; RT 0.89 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₀H₄₅N₅O₄

[M+H]⁺ calculated: 660.3544

15 [M+H]⁺ measured: 660.3522

Example 3: 3-[2-(Benzenesulfonyl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

20 **Step A:** 4-{*N*-methyl 3-[2-(benzenesulfonyl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-5,6,7,8-tetrahydroindolizine-1-amido}phenyl benzenesulfonate

To a solution of the product of Preparation 6e (35 mg, 0.05 mmol) in dichloromethane (1 mL) is added triethylamine (52 µL, 0.375 mmol) and the mixture is cooled to 0 °C. Benzenesulfonyl chloride (17 µL, 0.13 mmol) is added dropwise, and the resultant mixture

is stirred at 0 °C for 1 h. The reaction is diluted with dichloromethane, washed with 1M aqueous sodium hydroxide solution, followed by brine, and dried over magnesium sulphate. The solvent is removed *in vacuo*, and the crude product is purified on CombiFlash (4 g silica; dichloromethane to 5% MeOH/dichloromethane).

5 **LC/MS** (C₅₂H₅₃N₅O₈S₂) 798 [M-C₆H₅SO₂]⁻; RT 1.41 (Method B)

Step B: 3-[2-(Benzenesulfonyl)-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-N-(4-hydroxyphenyl)-N-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

10 The product from Step A is dissolved in methanol (2 mL), potassium hydroxide (10 eq) is added, and the resultant mixture is allowed to stir at ambient temperature for 7 h. Concentration *in vacuo*, and purification on CombiFlash (4 g silica; dichloromethane to 5% methanol/dichloromethane) affords the desired product as a solid.

LC/MS (C₄₆H₄₉N₅O₆S) 800 [M+H]⁺; RT 1.21 (Method B)

High-resolution mass (ESI+):

15 Empirical formula: C₄₆H₄₉N₅O₆S

[M+H]⁺ calculated: 800.3476

[M+H]⁺ measured: 800.3485

Example 4: ***N-(4-Hydroxyphenyl)-N-methyl-3-{7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-phenylmethanesulfonyl-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide***

20 The procedure is as in the process of Example 3, replacing benzenesulfonyl chloride in Step A with phenylmethanesulfonyl chloride.

LC/MS (C₄₇H₅₁N₅O₆S) 814 [M+H]⁺; RT 1.23 (Method B)

High-resolution mass (ESI+):

25 Empirical formula: C₄₇H₅₁N₅O₆S

$[M+H]^+$ calculated: 814.3633

$[M+H]^+$ measured: 814.3602

Example 5: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(naphthalene-2-sulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in the process of Example 3, replacing benzenesulfonyl chloride in Step A with naphthalene-2-sulfonyl chloride.

LC/MS ($C_{50}H_{51}N_5O_6S$) 850 $[M+H]^+$; RT 1.31 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{50}H_{51}N_5O_6S$

$[M+H]^+$ calculated: 850.3633

$[M+H]^+$ measured: 850.3624

Example 6: *tert*-Butyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

A solution of the product from Step A of Preparation 7aa (16 mg, 0.02 mmol) in ethanol (10 mL) is added to 10% Pd/C (catalytic amount) and the mixture is shaken under an atmosphere of hydrogen for *ca* 16 h. The mixture is filtered through celite, subsequently eluting with methanol, and removing the solvents under reduced pressure.

LC/MS ($C_{44}H_{51}N_5O_6$) 746 $[M+H]^+$; RT 1.25 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{44}H_{51}N_5O_6$

$[M+H]^+$ calculated: 746.3912

[M+H]⁺ measured: 746.3909

Example 7: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; dihydrochloride salt

5 Example 6 (45 mg, 0.05 mmol) was dissolved in a minimal amount of methanol and to this was added 4M solution of HCl in 1,4-dioxane (1 mL), and the mixture was allowed to stir at ambient temperature for *ca* 1 hr. The reaction was then diluted with dry diethyl ether (*ca* 5 mL) and the precipitate was collected by filtration and washed with a minimum of cold diethyl ether to afford the product.

10 **LC/MS** (C₃₉H₄₃N₅O₄) 646 [M+H]⁺; RT 0.89 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₃₉H₄₃N₅O₄

[M+H]⁺ calculated: 646.3388

[M+H]⁺ measured: 646.3385

15 **Example 8:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; hydrochloride salt

Step A: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

20

To a solution of product obtained from Preparation 7aa (50 mg, 0.07 mmol) in dichloromethane (2 mL) was added *N,N*-diisopropylethylamine (0.1 mmol) and the mixture was cooled to 0 °C. To the resultant solution was added dropwise 2-phenylacetyl chloride (0.102 mmol) and after 5 min the reaction was diluted with dichloromethane, and

washed sequentially with aqueous 1M sodium hydroxide, and brine. The organic extract was dried over magnesium sulfate, filtered and concentrated *in vacuo*, adsorbed onto isolate, and purified by chromatography (CombiFlash R_f, 40 g RediSep™ silica cartridge) eluting in a gradient of iso-hexane to 25 % ethyl acetate. The residue was then dissolved in ethanol (5 mL) and to this was added 10% Pd/C (catalytic amount) and the mixture stirred under a hydrogen atmosphere for *ca* 16 h. The mixture was filtered through celite, subsequently eluting with methanol, and solvent was removed under reduced pressure. The residue was adsorbed onto isolate and purified by chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge) eluting in a gradient of dichloromethane to 5 % methanol in dichloromethane to afford the desired product.

LC/MS (C₄₇H₄₉N₅O₅) 764 [M+H]⁺; RT 1.18 (Method B)

Step B: N-(4-Hydroxyphenyl)-N-methyl-3-{6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-2,3-dihydro-1H-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; hydrochloride salt

The product from Step A was dissolved in isopropyl alcohol (0.5 mL) and ethereal HCl (1M; 0.13 mL) was added. The mixture was stirred for 30 min, then concentrated *in vacuo*. Trituration of the residue with ether afforded the desired product as a solid.

LC/MS (C₄₇H₄₉N₅O₅) 764 [M+H]⁺; RT 1.16 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₉N₅O₅

[M+H]⁺ calculated: 764.3806

[M+H]⁺ measured: 764.3807

Example 9: N-(4-Hydroxyphenyl)-N-methyl-3-{6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylpropanoyl)-2,3-dihydro-1H-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride

The procedure is as in Example 8, replacing 2-phenylacetyl chloride in Step A with 3-phenylpropanoyl chloride.

LC/MS ($C_{48}H_{51}N_5O_5$) 778 $[M+H]^+$; RT 1.19 (Method B)

High-resolution mass (ESI+):

5 Empirical formula: $C_{48}H_{51}N_5O_5$

$[M+H]^+$ calculated: 778.3963

$[M+H]^+$ measured: 778.3973

Example 10: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Example 7, replacing Example 6 with product from Preparation 7ab.

LC/MS ($C_{35}H_{36}N_4O_3$) 561 $[M+H]^+$; RT 1.04

Example 11: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-propanoyl-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride

Step A: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-propanoyl-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

20 To a solution of the compound from Preparation 7aa (50 mg, 0.068 mmol) in dichloromethane (2 mL) is added DIPEA (17 μ L, 0.10 mmol) and the mixture is cooled to 0 °C. Propionyl chloride (9 μ L, 0.10 mmol) is added dropwise, and after 30 min the mixture is diluted with dichloromethane, washed sequentially with 1M aqueous NaOH and brine, dried over magnesium sulphate, and concentrated *in vacuo*. Purification by flash

column chromatography (4 g silica; dichloromethane to 5% MeOH/dichloromethane) afforded the product as a glassy solid.

LC/MS ($C_{49}H_{53}N_5O_5$) 792 $[M+H]^+$; RT 1.30 (Method B)

Step B: N-(4-Hydroxyphenyl)-N-methyl-3-{6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-propanoyl-2,3-dihydro-1H-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

A solution of the product from Step A (18.7 mg, 0.024 mmol) in ethanol (5 mL) is added to 10% Pd/C (catalytic amount) and the mixture is shaken under an atmosphere of hydrogen for *ca* 16 h. The mixture is filtered through celite, subsequently eluting with methanol, and removing the solvents under reduced pressure. The crude product is purified by flash column chromatography (4 g silica; dichloromethane to 5% MeOH/dichloromethane) to afford the desired product as a glassy solid.

LC/MS ($C_{42}H_{47}N_5O_5$) 702 $[M+H]^+$; RT 1.08 (Method B)

Step C: N-(4-Hydroxyphenyl)-N-methyl-3-{6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-propanoyl-2,3-dihydro-1H-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride

The product from Step B is dissolved in isopropyl alcohol (0.5 mL) and ethereal HCl (1M; 0.13 mL) is added. The mixture is stirred for 30 min, then concentrated *in vacuo*. Trituration of the residue with ether afforded the desired product as a solid.

LC/MS ($C_{42}H_{47}N_5O_5$) 702 $[M+H]^+$; RT 1.10 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{42}H_{47}N_5O_5$

$[M+H]^+$ calculated: 702.3650

$[M+H]^+$ measured: 702.3684

Example 12: 3-{2-Benzoyl-6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride

The procedure is as in Example 8, replacing 2-phenylacetyl chloride in Step A with benzoyl chloride.

LC/MS (C₄₆H₄₇N₅O₅) 750 [M+H]⁺; RT 1.17 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₇N₅O₅

[M+H]⁺ calculated: 750.3650

[M+H]⁺ measured: 750.3648

Example 13: *tert*-Butyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

The procedure is described in Example 6, replacing the product obtained in Step A of Preparation 7aa with the product obtained in Step A of Preparation 7ab.

LC/MS (C₄₀H₄₄N₄O₅) 661 [M+H]⁺; RT 1.42 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₀H₄₄N₄O₅

[M+H]⁺ calculated: 661.3384

[M+H]⁺ measured: 661.3352

Example 14: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 7ab.

LC/MS ($C_{43}H_{42}N_4O_4$) 679 $[M+H]^+$; RT 1.31 (Method B)

High-resolution mass (ESI+):

5 Empirical formula: $C_{43}H_{42}N_4O_4$

$[M+H]^+$ calculated: 679.3279

$[M+H]^+$ measured: 679.3298

Example 15: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylpropanoyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride in Step A with 3-phenylpropanoyl chloride.

LC/MS ($C_{44}H_{44}N_4O_4$) 693 $[M+H]^+$; RT 2.57 (Method A)

15 **High-resolution mass (ESI+):**

Empirical formula: $C_{44}H_{44}N_4O_4$

$[M+H]^+$ calculated: 693.3435

$[M+H]^+$ measured: 693.3441

Example 16: Phenyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate hydrochloride

The procedure is as in Example 8, replacing 2-phenylacetyl chloride in Step A with phenyl chloroformate.

LC/MS ($C_{46}H_{47}N_5O_6$) 766 $[M+H]^+$; RT 1.24 (Method B)

High-resolution mass (ESI+):

5 Empirical formula: $C_{46}H_{47}N_5O_6$

$[M+H]^+$ calculated: 766.3599

$[M+H]^+$ measured: 766.3602

Example 17: *N-tert-Butyl-5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxamide hydrochloride*

Step A: *5-(1-{[4-(Benzyloxy)phenyl](methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl)-N-tert-butyl-6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxamide*

15 To a solution of the compound of Preparation 7aa (30 mg, 0.04 mmol) in dichloromethane (2 mL) is added DIPEA (10 μ L, 0.06 mmol) and the mixture is cooled to 0 °C. tert-Butyl isocyanate (7 μ L, 0.06 mmol) is added and the reaction is stirred for 10 min, then diluted with dichloromethane and washed sequentially with 1M aqueous NaOH and brine. The organic phase is dried over magnesium sulphate, filtered and concentrated *in vacuo*. The
20 crude material is taken-up in dichloromethane, loaded onto isolate and purified on CombiFlash (4 g silica, dichloromethane to 5% methanol/ dichloromethane) to afford the desired product as a glassy solid that was used directly in the next step.

Step B: *N-tert-Butyl-5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxamide*

The product from Step A (25 mg) is dissolved in EtOH and to this is added 10% Pd/C (catalytic amount) and the mixture is allowed to stir under an atmosphere of hydrogen for *ca* 16 h. The reaction is filtered through celite, eluting with methanol and concentrated *in vacuo* to afford the desired product.

5 **LC/MS** (C₄₄H₅₂N₆O₅) 745 [M+H]⁺; RT 2.20 (Method A)

Step C: N-tert-Butyl-5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxamide hydrochloride

10 To a solution of the product from Step B in isopropyl alcohol (0.5 mL) is added ethereal HCl (1M; 0.17 mL, 0.17 mmol) and the mixture is stirred for 30 min. The solvent is removed *in vacuo* and trituration with ether affords the desired product as a solid.

LC/MS (C₄₄H₅₂N₆O₅) 745 [M+H]⁺; RT 1.10 (Method B)

High-resolution mass (ESI+):

15 Empirical formula: C₄₄H₅₂N₆O₅

[M+H]⁺ calculated: 745.4072

[M+H]⁺ measured: 745.4081

Example 18: **3-[2-(Ethanesulfonyl)-6-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindol-5-yl]-N-(4-hydroxyphenyl)-N-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride**

20

The procedure is as in Example 8, replacing 2-phenylacetyl chloride in Step A with ethanesulfonyl chloride.

LC/MS (C₄₁H₄₇N₅O₆S) 738 [M+H]⁺; RT 1.07 (Method B)

High-resolution mass (ESI+):

25 Empirical formula: C₄₁H₄₇N₅O₆S

$[M+H]^+$ calculated: 738.3320

$[M+H]^+$ measured: 738.3316

Example 19: 3-[2-(Benzenesulfonyl)-6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride

The procedure is as in Example 8, replacing 2-phenylacetyl chloride in Step A with benzenesulfonyl chloride.

LC/MS ($C_{45}H_{47}N_5O_6S$) 786 $[M+H]^+$; RT 1.16 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{45}H_{47}N_5O_6S$

$[M+H]^+$ calculated: 786.3320

$[M+H]^+$ measured: 786.3339

Example 20: 3-{2-Cyclopropanecarbonyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A and Step B of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride in Step A with cyclopropanecarbonyl chloride.

LC/MS ($C_{39}H_{40}N_4O_4$) 629.7 $[M+H]^+$; RT 2.41 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{39}H_{40}N_4O_4$

$[M+H]^+$ calculated: 629.3122

$[M+H]^+$ measured: 629.3129

Example 21: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylethanesulfonyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride

5 The procedure is as in Example 8, replacing 2-phenylacetyl chloride in Step A with 2-phenylethane-1-sulfonyl chloride.

LC/MS ($C_{47}H_{51}N_5O_6S$) 814 $[M+H]^+$; RT 1.21 (Method B)

Example 22: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(pyridine-3-sulfonyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride

10

The procedure is as in Example 8, replacing 2-phenylacetyl chloride in Step A with pyridine-3-sulfonyl chloride.

LC/MS ($C_{44}H_{46}N_6O_6S$) 787 $[M+H]^+$; RT 1.09 (Method B)

High-resolution mass (ESI+):

15 Empirical formula: $C_{44}H_{46}N_6O_6S$

$[M+H]^+$ calculated: 787.3272

$[M+H]^+$ measured: 787.3243

Example 23: 3-[2-(2-Benzylpropanoyl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

20

Step A: *N*-[4-(benzyloxy)phenyl]-3-[2-(2-benzylpropanoyl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the product from Preparation 7ab (52 mg, 0.07 mmol) in tetrahydrofuran (3 mL) was added *N,N*-diisopropylethylamine (37 μ L, 0.21 mmol) and HBTU (27 mg, 0.07 mmol), followed by 2-methyl-3-phenylpropanoic acid (17 mg, 0.10 mmol). The reaction was stirred at ambient temperature for *ca* 16 h, then partitioned between ethyl acetate and water, dried over magnesium sulphate, and concentrated *in vacuo*. Purification by flash column chromatography (silica; iso-hexane to ethyl acetate gradient) afforded the desired product.

LC/MS ($C_{52}H_{52}N_4O_4$) 797 $[M+H]^+$; RT 2.91 (Method A)

Step B: 3-[2-(2-Benzylpropanoyl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The product from Step A was dissolved in ethanol (10 mL) and to this was added 10% Pd/C (catalytic amount) and the mixture was allowed to stir under an atmosphere of hydrogen for *ca* 16 h. The reaction was filtered through celite, eluting with methanol and concentrated *in vacuo*. Purification by flash column chromatography (silica; iso-hexane to ethyl acetate gradient) afforded the desired product.

LC/MS ($C_{45}H_{46}N_4O_4$) 707 $[M+H]^+$; RT 2.64 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{45}H_{46}N_4O_4$

$[M+H]^+$ calculated: 707.3592

$[M+H]^+$ measured: 707.3600

Example 24: 3-(2-{2-[(4-Chlorophenyl)methyl]propanoyl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl)-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the product from Example 10 (20 mg, 0.03 mmol) in dichloromethane (3 ml) was added TEA (13 μ l, 0.09 mmol), HATU (12 mg, 0.03 mmol) and the 3-(4-chlorophenyl)-2-methylpropanoic acid (6 mg, 0.03 mmol), and stirred at ambient temperature for *ca* 16 h. The reaction was diluted with dichloromethane and washed with water, dried over magnesium sulphate, filtered and concentrated *in vacuo*. The crude material was purified by column chromatography in a gradient of dichloromethane to 10% methanol/dichloromethane.

LC/MS ($C_{45}H_{45}N_4O_4Cl$) 741 $[M+H]^+$; RT 2.71 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{45}H_{45}N_4O_4Cl$

$[M+H]^+$ calculated: 741.3202

$[M+H]^+$ measured: 741.3246

Example 25: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[(4-methylphenyl)methyl]propanoyl}-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Example 23, replacing 2-methyl-3-phenylpropanoic acid in Step A with 2-methyl-3-(4-methylphenyl)propanoic acid.

LC/MS ($C_{46}H_{48}N_4O_4$) 721 $[M+H]^+$; RT 2.70 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{48}N_4O_4$

$[M+H]^+$ calculated: 721.3748

$[M+H]^+$ measured: 721.3740

Example 26: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[4-(trifluoromethoxy)phenyl]acetyl}-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Example 23, replacing 2-methyl-3-phenylpropanoic acid in Step A with 2-[4-(trifluoromethoxy)phenyl]acetic acid.

LC/MS (C₄₄H₄₁N₄O₅F₃) 763 [M+H]⁺; RT 2.70 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₄H₄₁N₄O₅F₃

[M+H]⁺ calculated: 763.3102

[M+H]⁺ measured: 763.3102

Example 27: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; trifluoroacetic acid salt

Step A: *tert*-Butyl 6-{1-[4-hydroxyphenyl](methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The product from Preparation 6ab (95 mg, 0.12 mmol) is dissolved in ethanol (5 mL) and to this is added 10% Pd/C (catalytic amount) and the mixture is allowed to stir under an atmosphere of hydrogen for *ca* 6 h. The reaction is filtered through celite, eluting with methanol and concentrated *in vacuo* to afford the desired product.

LC/MS (C₄₁H₄₆N₄O₅) 675 [M+H]⁺; RT 1.43 (Method B)

Step B: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide; trifluoroacetic acid salt

The product from Step A is dissolved in dichloromethane (2 mL) and trifluoroacetic acid (0.1 mL) is added. After stirring at for *ca.* 16 h at ambient temperature the reaction is concentrated *in vacuo* and azeotroped with toluene (x3) to afford the desired product as the trifluoroacetic acid salt.

5 **LC/MS** (C₃₆H₃₈N₄O₃) 575 [M+H]⁺; RT 1.00 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₃₆H₃₈N₄O₃

[M+H]⁺ calculated: 575.3017

[M+H]⁺ measured: 575.2998

10 **Example 28:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(pyridin-2-ylmethyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step B of Example 29, replacing the compound from Step A with Example 10, and replacing cyclohexylacetaldehyde with pyridine-2-carbaldehyde.

15 **LC/MS** (C₄₁H₄₁N₅O₃) 652 [M+H]⁺; RT 1.06 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₁N₅O₃

[M+H]⁺ calculated: 652.3282

[M+H]⁺ measured: 652.3269

20 **Example 29:** 3-[2-(2-Cyclohexylethyl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{6-[*(3R)*-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the compound of Step A of Preparation 7ab (150 mg, 0.20 mmol) in dichloromethane (5 mL) is added trifluoroacetic acid (0.5 mL) and the reaction is stirred for *ca* 16 h. The mixture is diluted with dichloromethane, washed with 1N aqueous sodium hydroxide, dried (magnesium sulphate) and condensed under reduced pressure. Purification on CombiFlash (4 g silica, dichloromethane to 10% methanol/dichloromethane) afforded the desired product as a gum.

LC/MS (C₄₂H₄₂N₄O₃) 651 [M+H]⁺; RT 1.2 (Method B)

Step B: *N*-[4-(Benzyloxy)phenyl]-3-[2-(2-cyclohexylethyl)-6-[*(3R)*-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a stirring solution of the product from Step A (57.5 mg, 0.088 mmol) in THF (2 mL) is added cyclohexylacetaldehyde (13.4 mg, 0.11 mmol) followed by sodium triacetoxyborohydride (22.5 mg, 0.106 mmol) and the reaction was stirred for *ca* 16 h. The mixture is concentrated under reduced pressure, and purified on CombiFlash (4 g silica, dichloromethane to 3% methanol/dichloromethane) to afford the desired product as a gum.

LC/MS (C₅₀H₅₆N₄O₃) 761 [M+H]⁺; RT 1.37 (Method B)

Step C: 3-[2-(2-Cyclohexylethyl)-6-[*(3R)*-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the product from Step B (13 mg, 0.019 mmol) in dichloromethane (2 mL), cooled to 0 °C, is added boron trichloride (2M solution in dichloromethane; 0.1 mL, 0.2 mmol). After 5 h the reaction is quenched by addition of methanol, and concentrated *in vacuo*. Purification by preparative HPLC afforded the desired product.

LC/MS (C₄₃H₅₀N₄O₃) 671 [M+H]⁺; RT 1.16 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₃H₅₀N₄O₃

[M+H]⁺ calculated: 671.3956

[M+H]⁺ measured: 671.3955

Example 30: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylpropyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step B of Example 29, replacing the compound from Step A with Example 10, and replacing cyclohexylacetaldehyde with 3-phenylpropanal.

LC/MS (C₄₄H₄₆N₄O₃) 679 [M+H]⁺; RT 1.16 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₄H₄₆N₄O₃

[M+H]⁺ calculated: 679.3643

[M+H]⁺ measured: 679.3612

Example 31: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[2-(pyrazin-2-yl)-1,3-thiazol-4-yl]acetyl}-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *N*-[4-(benzyloxy)phenyl]-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[2-(pyrazin-2-yl)-1,3-thiazol-4-yl]acetyl}-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the product from Preparation 7ab (50 mg, 0.07 mmol) in dichloromethane (3 ml) was added triethylamine (29.2 μL, 0.21 mmol) and HATU (27 mg, 0.07 mmol), followed by 2-[2-(pyrazin-2-yl)-1,3-thiazol-4-yl]acetic acid (14.46 mg, 0.07 mmol), and the mixture was stirred at ambient temperature for *ca* 16 h. The reaction was diluted with dichloromethane and washed with water. The organic phase was dried over magnesium

sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (silica; dichloromethane to 5% methanol in dichloromethane gradient) afforded the desired product.

LC/MS (C₅₁H₄₇N₇O₄S) 854 [M+H]⁺; RT 2.77 (Method A)

5 **Step B:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[2-(pyrazin-2-yl)-1,3-thiazol-4-yl]acetyl}-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

10 A solution of the product from Step A (1 eq, 46 mg) in dichloromethane (5 ml) was cooled to 0 °C and to this was added boron trichloride (1M in dichloromethane, 100 µL). The reaction was allowed to warm to ambient temperature for *ca* 16 h, and was then quenched with methanol and partitioned between dichloromethane and water. The organic extract was dried over magnesium sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (silica; dichloromethane to 10% methanol in dichloromethane gradient) afforded the desired product.

15 LC/MS (C₄₄H₄₁N₇O₄S) 764 [M+H]⁺; RT 2.44 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₄H₄₁N₇O₄S

[M+H]⁺ calculated: 764.3014

[M+H]⁺ measured: 764.2994

20 **Example 32:** 3-[2-(3-Cyclohexylpropanoyl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *N*-[4-(Benzyloxy)phenyl]-3-[2-(3-cyclohexylpropanoyl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

25

To a solution of the compound from Preparation 7ab (52 mg, 0.066 mmol) in THF (3 mL) are added DIPEA (37 μ L, 0.21 mmol) and HBTU (27 mg, 0.07 mmol) followed by 3-cyclohexylpropionic acid (18 μ L, 0.10 mmol) and the mixture is stirred at ambient temperature for *ca* 16 h.

- 5 The reaction is partitioned between water and ethyl acetate, separated, and the organic phase is dried (magnesium sulphate) and concentrated *in vacuo*. Purification by flash column chromatography on silica gel, eluting with a gradient of iso-hexane to ethyl acetate afforded the product as a gum.

LC/MS ($C_{51}H_{56}N_4O_4$) 789 $[M+H]^+$; RT 3.00 (Method A)

- 10 ***Step B:*** 3-[2-(3-Cyclohexylpropanoyl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

A solution of the product from Step A (50 mg, 0.06 mmol) in ethanol (10 mL) is added to Pd/C (catalytic), and the mixture is shaken under an atmosphere of hydrogen for *ca* 16 h.

- 15 The mixture is filtered through celite, eluting with methanol, and then concentrated *in vacuo*. The crude material is purified by flash column chromatography on silica, eluting with a gradient of dichloromethane to 5% methanol/dichloromethane to afford the product as a solid.

LC/MS ($C_{44}H_{50}N_4O_4$) 699 $[M+H]^+$; RT 2.76 (Method A)

- 20 ***High-resolution mass (ESI⁺):***

Empirical formula: $C_{44}H_{50}N_4O_4$

$[M+H]^+$ calculated: 699.3905

$[M+H]^+$ measured: 699.3926

- 25 **Example 33:** 3-{2-Benzyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step B of Example 29, replacing the compound from Step A with the compound from Example 10, and replacing cyclohexylacetaldehyde with benzaldehyde.

LC/MS ($C_{42}H_{42}N_4O_3$) 651 $[M+H]^+$; RT 1.11 (Method B)

5 ***High-resolution mass (ESI+):***

Empirical formula: $C_{42}H_{42}N_4O_3$

$[M+H]^+$ calculated: 651.3330

$[M+H]^+$ measured: 651.3300

Example 34: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-propanoyl-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride in Step A with propanoyl chloride.

15 LC/MS ($C_{39}H_{42}N_4O_4$) 631 $[M+H]^+$; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{39}H_{42}N_4O_4$

$[M+H]^+$ calculated: 631.3279

$[M+H]^+$ measured: 631.3252

20 **Example 35:** 3-{2-Benzoyl-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride in Step A with benzoyl chloride.

LC/MS ($C_{43}H_{42}N_4O_4$) 679 $[M+H]^+$; RT 1.31 (Method B)

5 ***High-resolution mass (ESI+):***

Empirical formula: $C_{43}H_{42}N_4O_4$

$[M+H]^+$ calculated: 679.3279

$[M+H]^+$ measured: 679.3331

10 **Example 36:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6ab.

LC/MS ($C_{44}H_{44}N_4O_4$) 693 $[M+H]^+$; RT 1.32 (Method B)

15 ***High-resolution mass (ESI+):***

Empirical formula: $C_{44}H_{44}N_4O_4$

$[M+H]^+$ calculated: 693.3435

$[M+H]^+$ measured: 693.3447

20 **Example 37:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylpropanoyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride in Step A with 3-phenylpropanoyl chloride.

LC/MS ($C_{45}H_{46}N_4O_4$) 707 $[M+H]^+$; RT 1.36 (Method B)

5 ***High-resolution mass (ESI+):***

Empirical formula: $C_{45}H_{46}N_4O_4$

$[M+H]^+$ calculated: 707.3592

$[M+H]^+$ measured: 707.3557

10 **Example 38:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-phenylmethanesulfonyl-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing 2-phenylacetyl chloride in Step A with phenylmethanesulfonyl chloride.

LC/MS ($C_{46}H_{49}N_5O_6S$) 800 $[M+H]^+$; RT 1.15 (Method B)

15 ***High-resolution mass (ESI+):***

Empirical formula: $C_{46}H_{49}N_5O_6S$

$[M+H]^+$ calculated: 800.3476

$[M+H]^+$ measured: 800.3487

20 **Example 39:** 3-[2-(Benzenesulfonyl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride in Step A with benzenesulfonyl chloride.

LC/MS ($C_{42}H_{42}N_4O_5S$) 715 $[M+H]^+$; RT 1.37 (Method B)

5 ***High-resolution mass (ESI+):***

Empirical formula: $C_{42}H_{42}N_4O_5S$

$[M+H]^+$ calculated: 715.2949

$[M+H]^+$ measured: 715.2937

Example 40: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-
10 tetrahydroisoquinoline-2-carbonyl]-2-phenylmethanesulfonyl-1,2,3,4-
tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride in Step A with phenylmethanesulfonyl chloride.

15 **LC/MS** ($C_{43}H_{44}N_4O_5S$) 729 $[M+H]^+$; RT 1.37 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{43}H_{44}N_4O_5S$

$[M+H]^+$ calculated: 729.3105

$[M+H]^+$ measured: 729.3135

20 **Example 41:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-
tetrahydroisoquinoline-2-carbonyl]-2-(naphthalene-2-sulfonyl)-1,2,3,4-
tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride in Step A with naphthalene-2-sulfonyl chloride.

LC/MS (C₄₆H₄₄N₄O₅S) 765 [M+H]⁺; RT 1.44 (Method B)

5 ***High-resolution mass (ESI⁺):***

Empirical formula: C₄₆H₄₄N₄O₅S

[M+H]⁺ calculated: 765.3105

[M+H]⁺ measured: 765.3140

10 **Example 42:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(pyridin-3-yl)acetyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *N*-[4-(benzyloxy)phenyl]-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(pyridin-3-yl)acetyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

15 To a solution of the product from Preparation 7aa (40 mg, 0.05 mmol) in DMF (2 mL) was added triethylamine (28 µL, 0.2 mmol), and HBTU (19 mg, 0.05 mmol) followed by 2-(pyridin-3-yl)acetic acid hydrochloride (9.4 mg, 0.05 mmol), and the mixture was stirred at ambient temperature for 1 h. The reaction was concentrated *in vacuo* and directly used in the next step.

20 LC/MS (C₅₃H₅₄N₆O₅) no ionisation; RT 1.20 (Method B)

Step B: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(pyridin-3-yl)acetyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

25 The product from Step A (25 mg, 0.03 mmol) was dissolved in dichloromethane (2 mL) and cooled to 0 °C. To this was added boron trichloride (1M; 0.15 mL, 0.15 mmol) and the

mixture was stirred at ambient temperature for 2 h. The reaction was poured onto ice/water and extracted with dichloromethane. The organic extract was washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (4 g silica; dichloromethane to 5% methanol/ dichloromethane) afforded the desired product.

LC/MS ($C_{46}H_{48}N_6O_5$) 765 $[M+H]^+$; RT 0.95 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{48}N_6O_5$

$[M+2H]^{2+}$ calculated: 383.1916

$[M+2H]^{2+}$ measured: 383.1952

Example 43: *tert*-Butyl 5-{1-[(4-hydroxyphenyl)(phenyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

The procedure is described in Steps A and B of Example 44.

LC/MS ($C_{45}H_{46}N_4O_5$) 723 $[M+H]^+$; RT 1.47 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{45}H_{46}N_4O_5$

$[M+H]^+$ calculated: 723.3541

$[M+H]^+$ measured: 723.3546

Example 44: *N*-(4-Hydroxyphenyl)-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-phenyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *tert*-Butyl 5-(1-([4-(benzyloxy)phenyl](phenyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl)-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

A mix of THF (20 mL) and water (8 mL) is degassed by multiple evacuation / nitrogen purge cycles and bubbling nitrogen through. 4 mL of this mixture is added *via* syringe to a mixture of the boronic ester from Preparation 4f (117 mg, 0.28 mmol), the bromide from Preparation 1f (169 mg, 0.36 mmol) and cesium carbonate (182 mg, 0.56 mmol). Nitrogen is bubbled through the mix for another 3 mins. Bis(*di-tert*-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (10 mg, 5 mole%) is added and the reaction is immediately heated at 95 °C under microwave irradiation for 30 mins.

The reaction mixture is diluted with ethyl acetate (30 mL) and washed

sequentially with water (2 x 30 mL) and saturated NaCl (aq) (1 x 50 mL), dried over magnesium sulphate, filtered and evaporated. The crude material is purified by flash chromatography (Combiflash, 12g silica, eluting with gradient 0 to 100% ethyl acetate in hexane) to afford the product as an oil.

LC/MS (C₅₂H₅₂N₄O₅) 813 [M+H]⁺; RT 1.66 (Method B)

Step B: *tert*-Butyl 5-{1-[(4-hydroxyphenyl)(phenyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

A solution of the product from Step A (94 mg, 0.12 mmol) in ethanol (5 mL) is hydrogenated over 10% Pd/C catalyst under an atmosphere of hydrogen for *ca* 16 h. The mixture was filtered through celite, eluting with ethanol, and concentrated *in vacuo*. Purification by flash chromatography on silica gel (CombiFlash Rf, 4g SiO₂ silica column) eluting with 0 to 100% ethyl acetate in hexane afforded the product as a solid.

LC/MS (C₄₅H₄₆N₄O₅) 723 [M+H]⁺; RT 1.47 (Method B)

Step C: *N*-(4-Hydroxyphenyl)-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-phenyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the product from Step B (40 mg, 0.055 mmol) in anhydrous dichloromethane (4 mL) under nitrogen is added trifluoroacetic acid (180 uL, 2.33 mmol) and the reaction is stirred for *ca* 16 h. The mixture is diluted with dichloromethane (25 mL) and washed sequentially with 1M NaOH (aq) solution (20 mL) and saturated NaCl (aq) solution (20 mL). The organic phase is dried over sodium sulphate, filtered and evaporated. The crude material is purified by flash chromatography on silica gel (CombiFlash Rf, 4g, SilaSep column) eluting with 0 to 15% Methanol in dichloromethane to afford a glassy solid. Trituration with diethyl ether and evaporation afforded the desired product as a solid.

10 **LC/MS** ($C_{40}H_{38}N_4O_3$) 623 $[M+H]^+$; RT 1.08 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{40}H_{38}N_4O_3$

$[M+H]^+$ calculated: 623.3017

$[M+H]^+$ measured: 623.3028

15 **Example 45:** **Phenyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate**

The procedure is as in Step A of Example 8, replacing the product from Preparation 7aa with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride with phenyl chloroformate.

20 **LC/MS** ($C_{42}H_{40}N_4O_5$) 681 $[M+H]^+$; RT 1.36 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{42}H_{40}N_4O_5$

$[M+H]^+$ calculated: 681.3071

25 $[M+H]^+$ measured: 681.3057

Example 46: Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step A of Example 8, replacing the product from Preparation 7aa with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride with phenyl chloroformate.

LC/MS ($C_{43}H_{42}N_4O_5$) 695 $[M+H]^+$; RT 1.41 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{43}H_{42}N_4O_5$

$[M+H]^+$ calculated: 695.3228

$[M+H]^+$ measured: 695.3193

Example 47: *tert*-Butyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: *tert*-Butyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is described in Step A of Preparation 6ab.

LC/MS ($C_{48}H_{52}N_4O_5$) 765 $[M+H]^+$; RT 1.62 (Method B)

Step B: *tert*-Butyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

A solution of the product from Step A (50 mg, 0.07 mmol) in ethanol (5 mL) is added to Pd/C (catalytic), and the mixture is shaken under an atmosphere of hydrogen for *ca* 16 h.

The mixture is filtered through celite, eluting with methanol, and then concentrated *in vacuo*. The crude material is purified by flash column chromatography on silica, eluting with a gradient of dichloromethane to 5% methanol/dichloromethane to afford the product as a solid.

5 **LC/MS** ($C_{41}H_{46}N_4O_5$) 675 $[M+H]^+$; RT 1.42 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{41}H_{46}N_4O_5$

$[M+H]^+$ calculated: 675.3541

$[M+H]^+$ measured: 675.3571

10 **Example 48:** *N-tert-Butyl-6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxamide*

15 The procedure is as in Step A of Example 8, replacing the product from Preparation 7aa with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride with 2-isocyanato-2-methylpropane.

LC/MS ($C_{41}H_{47}N_5O_4$) 674 $[M+H]^+$; RT 1.33 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{41}H_{47}N_5O_4$

$[M+H]^+$ calculated: 674.3701

20 $[M+H]^+$ measured: 674.3706

Example 49: *3-{2-[2-(4-Chlorophenoxy)acetyl]-6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-indol-5-yl}-N-(4-hydroxyphenyl)-N-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide*

Step A: *N*-[4-(Benzyloxy)phenyl]-3-{2-[2-(4-chlorophenoxy)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Step A of Example 8, replacing the product from Preparation 7aa with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride with 2-(4-chlorophenoxy)acetyl chloride.

LC/MS (C₅₀H₄₇N₄O₅Cl) 819 [M+H]⁺; RT 1.56 (Method B)

Step B: 3-{2-[2-(4-Chlorophenoxy)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Step B of Example 42, replacing with the product from Step A of Example 42 with the product from Step A of Example 49.

LC/MS (C₄₃H₄₁N₄O₅Cl) 729 [M+H]⁺; RT 1.37 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₁N₄O₅Cl

[M+H]⁺ calculated: 729.2838

[M+H]⁺ measured: 729.2809

Example 50: 3-{2-[2-(3-Chlorophenoxy)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Example 49, replacing 2-(4-chlorophenoxy)acetyl chloride in Step A with 2-(3-chlorophenoxy)acetyl chloride.

LC/MS (C₄₃H₄₁N₄O₅Cl) 729 [M+H]⁺; RT 1.37 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₁N₄O₅Cl

[M+H]⁺ calculated: 729.2838

[M+H]⁺ measured: 729.2871

Example 51: *N*-(4-Hydroxyphenyl)-3-{2-[2-(4-methoxyphenyl)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]}-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₄₄H₄₄N₄O₅

[M+H]⁺ calculated: 709.3392

[M+H]⁺ measured: 709.3382

Example 52: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenoxyacetyl)-2,3-dihydro-1*H*-isoindol-5-yl]}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride in Step A with 2-phenoxyacetyl chloride.

LC/MS (C₄₃H₄₂N₄O₅) 695 [M+H]⁺; RT 1.31 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₂N₄O₅

[M+H]⁺ calculated: 695.3228

[M+H]⁺ measured: 695.3252

Example 53: 3-{2-[2-(4-Cyanophenyl)acetyl]-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

5 The procedure is as in Example 24, replacing the product from Example 10 with the product from Example 27, and replacing 3-(4-chlorophenyl)-2-methylpropanoic acid with 2-(4-cyanophenyl)acetic acid.

LC/MS (C₄₅H₄₃N₅O₄) 718 [M+H]⁺; RT 1.29

Example 54: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(4-methylphenyl)acetyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

10 LC/MS (C₄₅H₄₆N₄O₄) 707 [M+H]⁺; RT 2.65 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₄₆N₄O₄

[M+H]⁺ calculated: 707.3592

15 [M+H]⁺ measured: 707.3556

Example 55: 5-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-*N*-methyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

Step A: *tert*-Butyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

20 To a solution of the compound of Step A of Preparation 7ab (400 mg, 0.53 mmol) in ethanol (10 mL) is added 10% Pd/C (catalytic), and the mixture is stirred under an atmosphere of hydrogen for *ca* 16 h. The mixture is filtered through celite, concentrated

under reduced pressure, and purified by flash column chromatography on silica to afford the product as a glassy solid.

LC/MS (C₄₀H₄₄N₄O₅) 661 [M+H]⁺; RT 1.41 (Method B)

Step B: N-(4-Hydroxyphenyl)-N-methyl-3-{6-[(3R)-3-methyl-1,2,3,4-

5 ***tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindol-5-yl}-5,6,7,8-***
tetrahydroindolizine-1-carboxamide; trifluoroacetic acid salt

To a solution of the product from Step A (300 mg) in dichloromethane (5 mL) is added trifluoroacetic acid (excess) and the mixture is allowed to stir at ambient temperature for *ca* 16 h. The reaction is concentrated *in vacuo* and triturated with ether to afford the product
10 as a solid.

LC/MS (C₃₅H₃₆N₄O₃) 561 [M+H]⁺; RT 1.01 (Method B)

Step C: 5-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-
[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-
carbonyl chloride

15 To a solution of the product from Step B (100 mg, 0.15 mmol) in dichloromethane (2 mL), cooled to 0 °C, is added DIPEA (61 uL, 0.37 mmol) followed by portion-wise addition of triphosgene (44 mg, 0.15 mmol). The reaction is allowed to warm to ambient temperature and stirred for 1 h, then partitioned between dichloromethane and 1M aqueous HCl. The organic phase is dried over magnesium sulphate and concentrated *in vacuo* to afford the
20 product as a mixture of trichloromethyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxylate and 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carbonyl chloride
25 that was used directly in the next step without further purification.

LC/MS (C₃₇H₃₅Cl₃N₄O₅) 721 [M+H]⁺; RT 1.44, (C₃₆H₃₅ClN₄O₄) 623 [M+H]⁺; RT 1.32 (Method B)

Step D: 5-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-N-methyl-6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxamide

To a solution of the product from Step C (50 mg, 0.07 mmol) in acetonitrile (2 mL) is added methylamine (2M in THF; 346 μ L, 0.69 mmol) and DIPEA (61 μ L, 0.31 mmol) and the mixture is stirred for *ca* 16 h. Solvent is removed *in vacuo* and the residue is partitioned between ethyl acetate and water, the organic phase is dried over magnesium sulphate, and the solvent is removed *in vacuo*. The residue is dissolved in 1,4-dioxane (2 mL) and a 1M aqueous solution of NaOH (5 mL) was added dropwise. After 1 h the mixture is diluted with water, acidified with 1M aqueous HCl, and extracted with ethyl acetate. The organic extracts are dried over magnesium sulphate and concentrated under reduced pressure. Purification on CombiFlash (4 g silica, dichloromethane to 5% MeOH/dichloromethane) afforded the desired product as a solid.

LC/MS ($C_{37}H_{39}N_5O_4$) 618 $[M+H]^+$; RT 2.27 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{37}H_{39}N_5O_4$

$[M+H]^+$ calculated: 618.3075

$[M+H]^+$ measured: 618.3069

Example 56: 3-[2-(Ethanesulfonyl)-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-N-(4-hydroxyphenyl)-N-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Step A and Step B of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6ab, and replacing 2-phenylacetyl chloride in Step A with ethanesulfonyl chloride.

LC/MS ($C_{38}H_{42}N_4O_5S$) 667 $[M+H]^+$; RT 1.27 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₃₈H₄₂N₄O₅S

[M+H]⁺ calculated: 667.2949

[M+H]⁺ measured: 667.2935

Example 57: 3-{2-Ethyl-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

LC/MS (C₃₈H₄₂N₄O₃) 603 [M+H]⁺; RT 1.05 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₃₈H₄₂N₄O₃

[M+H]⁺ calculated: 603.3330

[M+H]⁺ measured: 603.3270

Example 58: 4-Aminophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: 4-nitrophenyl 6-(1-[[4-(benzyloxy)phenyl](methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of the product from Preparation 7ab (43 mg, 0.07 mmol) in dichloromethane (5 mL) was added 4-nitrophenyl chloroformate (18 mg, 0.09 mmol) and allowed to stir at ambient temperature *ca* 16 h. The reaction was diluted with dichloromethane and washed with 1N HCl, NaHCO₃ and brine. The organic phase was dried over magnesium sulphate, filtered and concentrated *in vacuo*, and taken forward assuming quantitative transformation.

LC/MS (C₅₀H₄₇N₅O₇) 830 [M+H]⁺; RT 1.58 (Method B)

Step B: 4-Aminophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of the compound of Step A (54 mg, 0.07 mmol) in ethanol (10 mL) is added 10% Pd/C (catalytic), and the mixture is stirred under an atmosphere of hydrogen for *ca* 16 h. The reaction mixture was filtered through celite and concentrated under reduced pressure.

LC/MS (C₄₃H₄₃N₅O₅) 708 [M-H]⁻; RT 1.26 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₃N₅O₅

[M+H]⁺ calculated: 710.3337

[M+H]⁺ measured: 710.3302

Example 59: 4-Acetamidophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

4-Aminophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate [Example 58] (23 mg, 0.03 mmol) is dissolved in dichloromethane (5 mL) followed by the addition of DIPEA (0.17 mL, 1 mmol) and cooled to 0°C. Acetyl chloride (3 µL, 0.04 mmol) is added and the reaction mixture is stirred for *ca* 16 h at ambient temperature. Ammonia in methanol (7 N; 1 mL) is added, and the resultant mixture is washed with saturated NaHCO₃ solution and brine, dried over magnesium sulphate, filtered and concentrated. The product is purified by FlashChrom (24 g silica, dichloromethane to 5% methanol/ dichloromethane).

LC/MS (C₄₅H₄₅N₅O₆) 752 [M+H]⁺; RT 1.25 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₄₅N₅O₆

[M+H]⁺ calculated: 752.3443

[M+H]⁺ measured: 752.3461

Example 60: 5-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-*N*-phenyl-2,3-dihydro-1*H*-isoindole-2-carboxamide

The procedure is as described in Step A and Step B of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride in Step A with phenyl isocyanate.

LC/MS (C₄₂H₄₁N₅O₄) 680 [M+H]⁺; RT 1.29 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₂H₄₁N₅O₄

[M+H]⁺ calculated: 680.3231

[M+H]⁺ measured: 680.3225

Example 61: *N*-Benzyl-5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

The procedure is as described in Step A and Step B of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride in Step A with benzyl isocyanate.

LC/MS (C₄₃H₄₃N₅O₄) 694 [M+H]⁺; RT 1.28 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₃H₄₃N₅O₄

$[M+H]^+$ calculated: 694.3388

$[M+H]^+$ measured: 694.3392

Example 62: Benzyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

A solution of the product obtained from Example 10 (53 mg, 0.08 mmol) in dichloromethane (5 mL) was cooled to 0 °C and to this was added *N,N*-diisopropylethylamine (0.18 mmol) and benzyl chloroformate (14.7 mg, 0.09 mmol). The reaction was stirred at ambient temperature for 15 min, then diluted with dichloromethane and washed successively with 1M aqueous sodium hydroxide, and brine. The organic extract was dried over magnesium sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; gradient of dichloromethane to 5 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₄₃H₄₂N₄O₅) 695 $[M+H]^+$; RT 1.40 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₂N₄O₅

$[M+H]^+$ calculated: 695.3228

$[M+H]^+$ measured: 695.3241

Example 63: Phenyl 5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

The procedure is as in Example 8, replacing the product from Preparation 7aa with the product from Preparation 7bb, and replacing 2-phenylacetyl chloride in Step A with phenyl chloroformate.

LC/MS ($C_{40}H_{38}N_4O_5$) 655 $[M+H]^+$; RT 1.33 (Method B)

5 ***High-resolution mass (ESI+):***

Empirical formula: $C_{40}H_{38}N_4O_5$

$[M+H]^+$ calculated: 655.2915

$[M+H]^+$ measured: 655.2897

10 **Example 64:** ***1H-pyrrolo[2,3-*b*]pyridin-5-ylmethyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate***

Step A: *4-Nitrophenyl 1H-pyrrolo[2,3-*b*]pyridin-5-ylmethyl carbonate*

15 To a solution of 1*H*-pyrrolo[2,3-*b*]pyridin-5-ylmethanol (148 mg, 1 mmol) and *N,N*-diisopropylethylamine (261 μ L, 1.5 mmol) in DCM (10 mL) was added 4-nitrophenyl chloroformate (202 mg, 1 mmol) and the mixture was stirred at ambient temperature. After concentrating *in vacuo*, purification by flash column chromatography (silica; ethyl acetate) afforded the desired material.

LC/MS ($C_{15}H_{11}N_3O_5$) 314 $[M+H]^+$; RT 1.19 (Method B)

20 **Step B:** *1H-pyrrolo[2,3-*b*]pyridin-5-ylmethyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate*

25 To a solution of the product from Example 10 (10 mg, 0.02 mmol) in tetrahydrofuran (1 mL) was added *N,N*-diisopropylethylamine (11 μ L, 0.06 mmol) followed by the product from Step A (6 mg, 0.02 mmol), and the mixture was stirred at ambient temperature for *ca*

16 h. After this time, further *N,N*-diisopropylethylamine (11 μ L, 0.06 mmol) and product from Step A (6 mg, 0.02 mmol) were added to affect full conversion. Purification by flash column chromatography (silica; gradient of iso-hexane to ethyl acetate) followed by evaporation and trituration with diethyl ether afforded the desired product.

5 **LC/MS** ($C_{44}H_{42}N_6O_5$) 735 $[M+H]^+$; RT 2.48 (Method A)

Example 65: 5-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-*N*-[4-(4-methylpiperazin-1-yl)phenyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

10 The procedure is as in Example 8, replacing the product from Preparation 7aa with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride in Step A with 1-(4-isocyanatophenyl)-4-methylpiperazine.

LC/MS ($C_{47}H_{51}N_7O_4$) 776 $[M-H]^-$; RT 2.18 (Method A)

High-resolution mass (ESI+):

15 Empirical formula: $C_{47}H_{51}N_7O_4$

$[M+2H]^{2+}$ calculated: 389.7074

$[M+2H]^{2+}$ measured: 389.7107

Example 66: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylpyrrolidine-1-carbonyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

20

Step A: 3-Phenylpyrrolidine-1-carbonyl chloride

A solution of the 3-phenylpyrrolidine hydrochloride (50 mg, 0.27 mmol) and *N,N*-diisopropylethylamine (0.14 mL, 0.81 mmol) in dichloromethane (2 mL) was cooled to 0°C. To this was added triphosgene (81 mg, 0.27 mmol) and the mixture was stirred at

ambient temperature. The reaction was subsequently diluted with water and the product extracted into dichloromethane, dried over magnesium sulphate and purified by flash column chromatography (silica; gradient of iso-hexane to 10% ethyl acetate in iso-hexane) to afford the desired product.

5 **LC/MS** ($C_{11}H_{12}NOCl$) 210 $[M+H]^+$; RT 2.56 (Method A)

Step B: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylpyrrolidine-1-carbonyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

10 The procedure is as for Example 8, replacing the product from Preparation 7aa with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride with the acid chloride obtained in Step A of Example 66.

LC/MS ($C_{46}H_{47}N_5O_4$) 734 $[M+H]^+$; RT 2.68 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{47}N_5O_4$

15 $[M+H]^+$ calculated: 734.3701

$[M+H]^+$ measured: 734.3673

Example 67: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylpyrrolidine-1-carbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

20 The procedure is as for Step A from Example 11, replacing the product from Preparation 7aa with the product from Example 27, and replacing propionyl chloride with the acid chloride obtained in Step A of Example 66.

LC/MS ($C_{47}H_{49}N_5O_4$) 748 $[M+H]^+$; RT 2.71 (Method A)

High-resolution mass (ESI+):

25 Empirical formula: $C_{47}H_{49}N_5O_4$

$[M+H]^+$ calculated: 748.3857

$[M+H]^+$ measured: 748.3815

Example 68: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 8, replacing the product from Preparation 7aa with the product from Preparation 6bb, and replacing 2-phenylacetyl chloride in Step A with phenyl chloroformate.

LC/MS ($C_{41}H_{40}N_4O_5$) 669 $[M+H]^+$; RT 1.35 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{41}H_{40}N_4O_5$

$[M+H]^+$ calculated: 669.3071

$[M+H]^+$ measured: 669.3045

Example 69: Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride

Step A: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide dihydrochloride

The product from Step A of Preparation 6aa (549 mg, 0.65 mmol) is dissolved in methanol (5 mL), then a solution of HCl in dioxane (4 M; 10 mL) is added and the mixture is allowed to stir for *ca* 16 h. Ether is added, resulting in the precipitation of a solid which is collected by filtration and washed with ether to afford the desired product.

LC/MS ($C_{47}H_{51}N_5O_4$) 750[M+H]⁺; RT 1.09 (Method B)

Step B: Phenyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

5 The product from Step A (100 mg, 0.12 mmol) is dissolved in dichloromethane (5 mL) and to this is added DIPEA (174 μ L, 1 mmol) and the solution cooled to 0°C. Phenyl chloroformate (18 μ L, 0.14 mmol) is added and the mixture is stirred for 30 minutes before diluting with dichloromethane, washing sequentially with 1M NaOH and brine, and then drying over magnesium sulphate. The solvent is removed *in vacuo*, and the residue is
10 taken up in dichloromethane and purified on CombiFlash (12 g silica; dichloromethane to 5% methanol/dichloromethane) to afford the product as a solid.

LC/MS ($C_{54}H_{55}N_5O_6$) 870[M+H]⁺; RT 1.45 (Method B)

Step C: Phenyl 6-{1-[4-(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-
15 1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride

A solution of the product from Step B (96.6 mg, 0.11 mmol) in ethanol (5 mL) is added to a catalytic amount of 10% Pd/C, and the mixture is shaken under an atmosphere of hydrogen gas for *ca* 16 h. The mixture is filtered through celite and evaporated, whereby the residue is dissolved in a minimum amount of isopropyl alcohol. To the resultant
20 solution is added ethereal HCl (1M, 0.5 mL), followed by ether, and the solid product is collected by vacuum filtration.

LC/MS ($C_{47}H_{49}N_5O_6$) 780[M+H]⁺; RT 1.21 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{47}H_{49}N_5O_6$

25 [M+H]⁺ calculated: 780.3756

[M+H]⁺ measured: 780.3791

Example 70: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa.

LC/MS (C₄₈H₅₁N₅O₅) 778 [M+H]⁺; RT 1.14 (Method B)

Example 71: 6-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-*N*-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

The procedure is as in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing 2-phenylacetyl chloride with phenyl isocyanate.

LC/MS (C₄₃H₄₃N₅O₄) 694 [M+H]⁺; RT 1.31 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₃N₅O₄

[M+H]⁺ calculated: 694.3388

[M+H]⁺ measured: 694.3356

Example 72: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylethyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

LC/MS (C₄₃H₄₄N₄O₃) 665 [M+H]⁺; RT 1.10 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₄N₄O₃

$[M+H]^+$ calculated: 665.3486

$[M+H]^+$ measured: 665.3477

Example 73: *N*-Benzyl-6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

The procedure is as in Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing 2-phenylacetyl chloride with benzyl isocyanate.

LC/MS ($C_{44}H_{45}N_5O_4$) 708 $[M+H]^+$; RT 1.3 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{44}H_{45}N_5O_4$

$[M+H]^+$ calculated: 708.3544

$[M+H]^+$ measured: 708.3556

Example 74: 6-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-*N*-methyl-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-*N*-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

Step A: 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl chloride

A solution of the product from Preparation 6ab (50 mg, 0.08 mmol) in dichloromethane (5 mL) was cooled to 0 °C and to this was added *N,N*-diisopropylethylamine (0.3 mmol) followed by triphosgene (22.3 mg, 0.08 mmol), and the mixture was allowed to stir at ambient temperature for 1 h. The reaction was partitioned between dichloromethane and 1M aqueous HCl, and the organic extract was dried over magnesium sulphate. After

concentration *in vacuo*, the material was used directly in the next step assuming quantitative transformation.

LC/MS (C₄₄H₄₃N₄O₄Cl) 727 [M+H]⁺; RT 1.54 (Method B)

Step B: 6-(1-[[4-(benzyloxy)phenyl](methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl]-N-methyl-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-N-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

To a solution of product obtained from Step A (62 mg, 0.08 mmol) in Acetonitrile (5 mL) was added N,N-diisopropylethylamine (0.40 mmol) and N-methylaniline (80.4 mg, 0.75 mmol), and the mixture was stirred at ambient temperature for *ca* 16 h. The reaction mixture was diluted with ethyl acetate and successively washed with aqueous sodium bicarbonate, and brine. The organic extract was dried over magnesium sulfate, filtered, and concentrated *in vacuo*. Purification by flash column chromatography (4 g silica, dichloromethane to 5% methanol in dichloromethane) afforded the desired product.

LC/MS (C₅₁H₅₁N₅O₄) 798 [M+H]⁺; RT 1.56 (Method B)

Step C: 6-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-N-methyl-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-N-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

To a solution of product obtained in Step B (60 mg, 0.08 mmol) in ethanol (5 mL) was added 10% Pd/C (catalytic amount) and the mixture was shaken under an atmosphere of hydrogen for *ca* 16 h. The mixture was filtered through celite, subsequently eluting with methanol, and the solvent was removed *in vacuo*. Purification by flash column chromatography (4 g silica; dichloromethane to 5% methanol in dichloromethane) afforded the desired product as a solid.

LC/MS (C₄₄H₄₅N₅O₄) 708 [M+H]⁺; RT 1.35 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₄H₄₅N₅O₄

[M+H]⁺ calculated: 708.3544

$[M+H]^+$ measured: 708.3543

Example 75: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

5 The procedure is as in Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6bb.

LC/MS ($C_{42}H_{42}N_4O_4$) 667 $[M+H]^+$; RT 1.27 (Method B)

Example 76: *N*-*tert*-Butyl-5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

10

The procedure is as in Example 8, replacing the product from Preparation 7aa with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride with *tert*-butyl isocyanate.

LC/MS ($C_{40}H_{45}N_5O_4$) 660 $[M+H]^+$; RT 1.29 (Method B)

15 *High-resolution mass (ESI+):*

Empirical formula: $C_{40}H_{45}N_5O_4$

$[M+H]^+$ calculated: 660.3544

$[M+H]^+$ measured: 660.3529

Example 77: 6-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-*N*-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

20

The procedure is as in Example 8, replacing the product from Preparation 7aa with the product from Preparation 6aa, and replacing 2-phenylacetyl chloride with phenyl isocyanate.

LC/MS (C₄₇H₅₀N₆O₅) 779 [M+H]⁺; RT 1.12

5 **Example 78:** *N*-Benzyl-6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

10 The procedure is as in Example 8, replacing the product from Preparation 7aa with the product from Preparation 6aa, and replacing 2-phenylacetyl chloride with benzyl isocyanate.

LC/MS (C₄₈H₅₂N₆O₅) 793 [M+H]⁺; RT 1.12

High-resolution mass (ESI⁺):

Empirical formula: C₄₈H₅₂N₆O₅

[M+H]⁺ calculated: 793.4072

15 [M+H]⁺ measured: 793.4067

Example 79: 6-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-*N*-methyl-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

20 The procedure is as in Example 74, replacing *N*-methylaniline in Step B with *N*-benzylmethylamine.

LC/MS (C₃₈H₄₁N₅O₄) 632 [M+H]⁺; RT 1.16 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₃₈H₄₁N₅O₄

[M+H]⁺ calculated: 632.3231

[M+H]⁺ measured: 632.3232

Example 80: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-1*H*-pyrrole-3-carboxamide

Step A: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-1*H*-pyrrole-3-carboxamide

To a solution of the compound from Preparation 7bb (50 mg, 0.08 mmol) in dichloromethane (2 mL), cooled to 0 °C, are added DIPEA (28 uL, 0.16 mmol) and phenylacetyl chloride (13 uL, 0.10 mmol). After stirring for 10 min, the reaction is diluted with dichloromethane, sequentially washed with 1M aqueous NaOH and brine, dried (magnesium sulphate), and concentrated *in vacuo*. The crude material is purified on CombiFlash (4 g silica, dichloromethane to 3% methanol/dichloromethane) to afford the product.

LC/MS (C₄₈H₄₆N₄O₄) 743 [M+H]⁺; RT 1.47 (Method B)

Step B: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-1*H*-pyrrole-3-carboxamide

A solution of the product from Step A (59 mg, 0.08 mmol) in ethanol was added to 10% Pd/C (catalytic) and the mixture was shaken at ambient temperature under an atmosphere of hydrogen for *ca* 16 h. Filtration through celite, evaporation of solvents under vacuum, and purification on CombiFlash (4 g silica, dichloromethane to 5% methanol/dichloromethane) afforded the desired product.

LC/MS (C₄₁H₄₀N₄O₄) 653 [M+H]⁺; RT 1.25 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₀N₄O₄

[M+H]⁺ calculated: 653.3122

[M+H]⁺ measured: 653.3137

Example 81: *N*-*tert*-Butyl-5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-*N*-methyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

The procedure is as in Example 74, replacing the product from Preparation 6ab in Step A with the product from Preparation 7ab, and replacing *N*-methyl aniline in Step B with *N*-*tert*-butylmethylamine.

LC/MS (C₄₁H₄₇N₅O₄) 674 [M+H]⁺; RT 1.38 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₁H₄₇N₅O₄

[M+H]⁺ calculated: 674.3701

[M+H]⁺ measured: 674.3720

Example 82: 5-{1-[(4-Hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-*N*-methyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-*N*-phenyl-2,3-dihydro-1*H*-isoindole-2-carboxamide

The procedure is as in Example 74, replacing the product from Preparation 6ab with the product from Preparation 7ab.

LC/MS (C₄₃H₄₃N₅O₄) 694 [M+H]⁺; RT 1.33 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₃H₄₃N₅O₄

$[M+H]^+$ calculated: 694.3388

$[M+H]^+$ measured: 694.3395

Example 83: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylazetidine-1-carbonyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Example 74, replacing the product from Preparation 7aa with the product from Preparation 7ab, and replacing *N*-methylaniline with 3-phenylazetidine hydrochloride.

LC/MS ($C_{45}H_{45}N_5O_4$) 720 $[M+H]^+$; RT 2.62 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{45}H_{45}N_5O_4$

$[M+H]^+$ calculated: 720.3544

$[M+H]^+$ measured: 720.3536

Example 84: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylazetidine-1-carbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Example 74, replacing the product from Preparation 7aa with the product from Preparation 6ab, and replacing *N*-methylaniline with 3-phenylazetidine hydrochloride.

LC/MS ($C_{46}H_{47}N_5O_4$) 734 $[M+H]^+$; RT 2.66 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{47}N_5O_4$

$[M+H]^+$ calculated: 734.3701

[M+H]⁺ measured: 734.3668

Example 85: *N*-Benzyl-5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-*N*-methyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

5 **Step A:** *N*-Benzyl-5-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-*N*-methyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

10 The procedure is as in Step A and Step B of Example 74, replacing the product from Preparation 6ab in Step A with the product from Preparation 7ab, and replacing *N*-methylaniline in Step B with *N*-benzylmethylaniline.

LC/MS (C₅₁H₅₁N₅O₄) 798 [M+H]⁺; RT 1.55 (Method B)

Step B: *N*-Benzyl-5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-*N*-methyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

15 The product from Step A (46 mg, 0.06 mmol) was dissolved in dichloromethane (5 mL) and cooled to 0 °C. To this was added boron trichloride (1M in dichloromethane; 0.18 mmol) dropwise. The reaction was stirred at ambient temperature for 1 h, then quenched by the addition of methanol (5 mL) and concentrated *in vacuo*. The crude material was partitioned between ethyl acetate and water, and the organic extract was washed with
20 brine, dried (magnesium sulphate) and concentrated *in vacuo*. Purification by flash column chromatography (4 g silica, dichloromethane to 5% methanol in dichloromethane) afforded the desired product.

LC/MS (C₄₄H₄₅N₅O₄) 708 [M+H]⁺; RT 1.36 (Method B)

High-resolution mass (ESI⁺):

25 Empirical formula: C₄₄H₄₅N₅O₄

[M+H]⁺ calculated: 708.3544

[M+H]⁺ measured: 708.3542

Example 86: 3-{2-[2-(2-Chlorophenoxy)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

5 **Step A:** *N*-[4-(Benzyloxy)phenyl]-3-{2-[2-(2-chlorophenoxy)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Step A of Example 11, replacing the product from Preparation 7aa with the product from Preparation 7ab, and replacing propionyl chloride with 2-(2-chlorophenoxy)acetyl chloride.

10

LC/MS (C₅₀H₄₇N₄O₅Cl) 819 [M+H]⁺; RT 1.53 (Method B)

Step B: 3-{2-[2-(2-Chlorophenoxy)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

15 The procedure is as in Step B of Example 85, replacing the product in Step A of Example 85 with the product of Step A in Example 86.

LC/MS (C₄₃H₄₁N₄O₅Cl) 729 [M+H]⁺; RT 1.35 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₃H₄₁N₄O₅Cl

20 [M+H]⁺ calculated: 729.2838

[M+H]⁺ measured: 729.2820

Example 87: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(phenylamino)acetyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Example 23, replacing 2-methyl-3-phenylpropanoic acid in Step A with 2-(phenylamino)acetic acid.

LC/MS ($C_{43}H_{43}N_5O_4$) 694 $[M+H]^+$; RT 2.58 (Method A)

High-resolution mass (ESI+):

5 Empirical formula: $C_{43}H_{43}N_5O_4$

$[M+H]^+$ calculated: 694.3388

$[M+H]^+$ measured: 694.3357

Example 88: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride

The procedure was as described for Example 69, replacing the compound from Preparation 6aa with the compound from Preparation 6ba.

LC/MS ($C_{45}H_{47}N_5O_6$) 754 $[M+H]^+$; RT 1.17 (Method B)

15 **High-resolution mass (ESI+):**

Empirical formula: $C_{45}H_{47}N_5O_6$

$[M+H]^+$ calculated: 754.3599

$[M+H]^+$ measured: 754.3598

Example 89: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

20 The procedure is as in Example 8, replacing the product from Preparation 7aa with the product from Preparation 6aa.

LC/MS (C₄₆H₄₉N₅O₅) 752 [M+H]⁺; RT 1.10 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₉N₅O₅

[M+H]⁺ calculated: 752.3806

5 [M+H]⁺ measured: 752.3797

Example 90: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[(2*S*)-2-phenylpropanoyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

10 The procedure is as described in Example 23, replacing the product of 7ab with the product of Preparation 6ab, and replacing 2-methyl-3-phenylpropanoic acid in Step A with (2*S*)-2-phenylpropanoic acid.

LC/MS (C₄₅H₄₆N₄O₄) 707 [M+H]⁺; RT 2.65 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₄₆N₄O₄

15 [M+H]⁺ calculated: 707.3592

[M+H]⁺ measured: 707.3589

Example 91: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[(2*R*)-2-phenylpropanoyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

20 The procedure is as described in Example 23, replacing the product of 7ab with the product of Preparation 6ab, and replacing 2-methyl-3-phenylpropanoic acid in Step A with (2*R*)-2-phenylpropanoic acid.

LC/MS (C₄₅H₄₆N₄O₄) 707 [M+H]⁺; RT 2.66 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₄₆N₄O₄

[M+H]⁺ calculated: 707.3592

[M+H]⁺ measured: 707.3589

5 **Example 92:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[(phenylcarbamoyl)methyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

10 The procedure is as in Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 7ab, and replacing 2-phenylacetyl chloride in Step A with 2-chloro-*N*-phenylacetamide

LC/MS (C₄₃H₄₃N₅O₄) 694 [M+H]⁺; RT 1.18 (Method B)

Example 93: Benzyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

15 The procedure is as in Example 62, replacing the product from Example 10 with the product from Example 27.

LC/MS (C₄₄H₄₄N₄O₅) 709 [M+H]⁺; RT 1.41 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₄H₄₄N₄O₅

20 [M+H]⁺ calculated: 709.3384

[M+H]⁺ measured: 709.3387

Example 94: Benzyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide, trifluoroacetic acid salt

To a solution of the product from Step A of Preparation 6aa (43 mg, 0.065 mmol) in dichloromethane (5 mL) is added trifluoroacetic acid (0.4 mL), and the mixture is stirred at ambient temperature for *ca* 16 h. The solvent is removed *in vacuo* to afford the desired product which is used directly in the next step without further purification.

LC/MS (C₄₀H₄₅N₅O₄) 660 [M+H]⁺; RT 0.87 (Method B)

Step B: Benzyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The product from Step A (43 mg, 0.06 mmol) is dissolved in dichloromethane (2 mL) and cooled to 0°C. To this is added triethylamine (42 uL, 0.3 mmol) followed by benzyl chloroformate (9 uL) and the mixture is stirred for 15 minutes. The reaction mixture is diluted with dichloromethane and washed sequentially with 1M aqueous NaOH, and brine. The organics are dried over magnesium sulphate, filtered and concentrated and the residue taken-up in methanol. To the methanolic solution is added 1M aqueous NaOH and the mixture is heated for 2 hours at 50°C. The reaction mixture is concentrated *in vacuo*, partitioned between ethyl acetate and brine, and the organics dried over magnesium sulphate. After evaporation, the crude product is purified by flash column chromatography over silica (4 g), eluting with a gradient of dichloromethane to 5% methanol/dichloromethane.

LC/MS (C₄₈H₅₁N₅O₆) 794 [M+H]⁺; RT 1.21 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₈H₅₁N₅O₆

$[M+H]^+$ calculated: 794.3912

$[M+H]^+$ measured: 794.3908

Example 95: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylbutanoyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as described in Example 23, replacing the product of 7ab with the product of Preparation 6ab, and replacing 2-methyl-3-phenylpropanoic acid in Step A with 1-phenylcyclopropane-1-carboxylic acid.

LC/MS ($C_{46}H_{48}N_4O_4$) 721 $[M+H]^+$; RT 2.72 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{48}N_4O_4$

$[M+H]^+$ calculated: 721.3748

$[M+H]^+$ measured: 721.3740

Example 96: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(4-{2-(morpholin-4-yl)ethyl}amino)phenyl]acetyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: Ethyl 2-(4-{2-(morpholin-4-yl)ethyl}amino)phenyl)acetate

To a boiling tube was added ethyl 2-(4-bromophenyl)acetate (317 mg, 1.3 mmol), the 2-(morpholin-4-yl)ethan-1-amine (257 μ L, 1.96 mmol), potassium phosphate tribasic (386 mg, 1.82 mmol) and (2-biphenyl)di-*tert*-butylphosphine (0.1 mol%) followed by toluene (6 mL). The reaction was degassed with nitrogen followed by the addition of bis(dibenzylideneacetone)palladium(0) (0.05 mol%). The reaction was then heated at 100 °C under nitrogen for *ca* 6 h. The reaction was diluted with dichloromethane and washed

with water. The organic extract was dried over magnesium sulphate, filtered and loaded onto a column for purification on silica in a gradient of iso-hexane to ethyl acetate.

LC/MS ($C_{16}H_{24}N_2O_3$) 293 $[M+H]^+$; RT 1.74 (Method A)

Step B: *Sodium 2-(4-{[2-(morpholin-4-yl)ethyl]amino}phenyl)acetate*

- 5 To a solution of the product obtained in Step A (61 mg, 0.21 mmol) in methanol (5 ml) was added 2M NaOH (21 μ L, 0.42 mmol). The reaction was stirred at ambient temperature for *ca* 16 h. The reaction was then filtered through a cotton wool plug and concentrated *in vacuo*, then triturated with ether, filtered and solvents removed *in vacuo*.

LC/MS ($C_{14}H_{20}N_2O_3$) 265 $[M+H]^+$; RT 0.50 (Method A)

- 10 **Step C:** *N-[4-(Benzyloxy)phenyl]-N-methyl-3-{7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(4-{[2-(morpholin-4-yl)ethyl]amino}phenyl)acetyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-5,6,7,8-tetrahydroindolizine-1-carboxamide*

- 15 To a solution of the product from Preparation 6ab (40 mg, 0.05 mmol) in dichloromethane (3 ml) was added *N,N*-diisopropylethylamine (20 μ L, 0.15 mmol) and HBTU (20 mg, 0.5 mmol) followed by the sodium salt from Step B (23 mg, 0.8 mmol). The reaction was stirred at ambient temperature for *ca* 16 h. The reaction was diluted with dichloromethane and washed with water. The organic extract was dried over magnesium sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (silica; 20 dichloromethane to 5% methanol in dichloromethane) followed by trituration with ether afforded the desired product as a cream powder.

LC/MS ($C_{57}H_{62}N_6O_5$) 911 $[M+H]^+$; RT 2.51 (Method A)

- 25 **Step D:** *N-(4-Hydroxyphenyl)-N-methyl-3-{7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(4-{[2-(morpholin-4-yl)ethyl]amino}phenyl)acetyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-5,6,7,8-tetrahydroindolizine-1-carboxamide*

A solution of the product from Step C (68 mg, 0.07 mmol) in ethanol (5 mL) was added to 10% Pd/C (catalytic) and the mixture was shaken at ambient temperature under an

atmosphere of hydrogen for *ca* 16 h. Filtration through celite, evaporation of solvents under vacuum, and purification on CombiFlash (4 g silica, dichloromethane to 5% methanol in dichloromethane) afforded the desired product.

LC/MS (C₅₀H₅₆N₆O₅) 819 [M-H]⁻; RT 2.21 (Method A)

5 **Example 97:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[4-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Example 96, replacing 2-(morpholin-4-yl)ethan-1-amine in Step A with 1-methylpiperazine.

10 **LC/MS** (C₄₉H₅₄N₆O₄) 791 [M+H]⁺; RT 2.20 (Method A)

High-resolution mass (ESI⁺):

Empirical formula: C₄₉H₅₄N₆O₄

[M+H]⁺ calculated: 791.4279

[M+H]⁺ measured: 791.4268

15 **Example 98:** 4-Methylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride

20 The procedure was as described for Example 99, replacing 3-methylphenol in Step B with 4-methylphenol.

LC/MS (C₄₈H₅₁N₅O₆) 794 [M+H]⁺; RT 1.25 (Method B)

Example 99: 3-Methylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl chloride

To a solution of the compound from Preparation 6aa (150 mg, 0.17 mmol) in dichloromethane (5 mL), cooled to 0 °C, is added DIPEA (89 uL, 0.51 mmol) followed by the portion-wise addition of triphosgene (52 mg, 0.17 mmol). After stirring for 1 h, the reaction mixture is partitioned between dichloromethane and 1M aqueous HCl, and the phase is dried over magnesium sulphate, filtered and concentrated to afford a mixture of trichloromethyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate and 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl chloride that is used directly in the next step without further purification, and assuming quantitative transformation.

Step B: 3-Methylphenyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of the product from Step A (30 mg) in acetonitrile (5 ml) is added potassium carbonate (17 mg, 0.123 mmol) and 3-methylphenol (13 uL, 0.123 mmol) and the mixture is heated at 65°C for 2 hours. The reaction is cooled to room temperature, diluted in ethyl acetate and washed with brine. The organics are dried over magnesium sulphate, filtered and concentrated. The crude material is taken-up in dichloromethane, loaded onto isolate and purified on CombiFlash (4 g silica, dichloromethane to 5% methanol/dichloromethane).

Step C: 3-Methylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

5 A solution of the product from Step B (30 mg, 0.03 mmol) in ethanol is added to 10% Pd/C (catalytic) and the mixture is shaken under an atmosphere of hydrogen for *ca* 16 h. Filtration through celite, evaporation, and purification on CombiFlash (4 g silica, dichloromethane to 5% MeOH/dichloromethane) afforded the desired product. The appropriate fractions were combined and concentrated to obtain the desired product.

LC/MS (C₄₈H₅₁N₅O₆) 794 [M+H]⁺; RT 1.25 (Method B)

10 ***High-resolution mass (ESI+):***

Empirical formula: C₄₈H₅₁N₅O₆

[M+H]⁺ calculated: 794.3912

[M+H]⁺ measured: 794.3925

15 **Example 100:** 2-Methylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure was as described for Example 99, replacing 3-methylphenol in Step B with 2-methylphenol.

LC/MS (C₄₈H₅₁N₅O₆) 794 [M+H]⁺; RT 1.25 (Method B)

20 ***High-resolution mass (ESI+):***

Empirical formula: C₄₈H₅₁N₅O₆

[M+H]⁺ calculated: 794.3912

[M+H]⁺ measured: 794.3877

Example 101: *N*-(4-Hydroxyphenyl)-5-{2-[2-(4-methoxyphenyl)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

5 Empirical formula: C₄₂H₄₂N₄O₅

[M+H]⁺ calculated: 683.3235

[M+H]⁺ measured: 683.3213

Example 102: 5-{2-[2-(4-Fluorophenyl)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

10

High-resolution mass (ESI+):

Empirical formula: C₄₁H₃₉FN₄O₄

[M+H]⁺ calculated: 671.3035

[M+H]⁺ measured: 671.3039

Example 103: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenoxyacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-1*H*-pyrrole-3-carboxamide

15

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₀N₄O₅

20 [M+H]⁺ calculated: 669.3079

[M+H]⁺ measured: 669.3084

Example 104: *N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-5-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenoxypropanoyl)-2,3-dihydro-1*H*-isoindol-5-yl}-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

5 Empirical formula: C₄₂H₄₂N₄O₅

[M+H]⁺ calculated: 683.3235

[M+H]⁺ measured: 683.3210

Example 105: 4-Chlorophenyl 5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

High-resolution mass (ESI+):

Empirical formula: C₄₀H₃₇ClN₄O₅

[M+H]⁺ calculated: 689.2532

[M+H]⁺ measured: 689.2539

15 **Example 106:** 4-Fluorophenyl 5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

High-resolution mass (ESI+):

Empirical formula: C₄₀H₃₇FN₄O₅

20 [M+H]⁺ calculated: 673.2828

[M+H]⁺ measured: 673.2832

Example 107: 4-Methylphenyl 5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

High-resolution mass (ESI+):

5 Empirical formula: C₄₁H₄₀N₄O₅

[M+H]⁺ calculated: 669.3079

[M+H]⁺ measured: 669.3065

Example 108: 3-{2-[2-(4-{[2-(Dimethylamino)ethyl]amino}phenyl)acetyl]-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide hydrochloride

The procedure is as in Example 96, replacing 2-(morpholin-4-yl)ethan-1-amine in Step A with (2-aminoethyl)dimethylamine.

LC/MS (C₄₈H₅₄N₆O₄) 777 [M-H]⁻; RT 2.20 (Method A)

15 **Example 109:** *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(1-phenylcyclopropanecarbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *N*-[4-(benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(1-phenylcyclopropanecarbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

20 To a solution of the product from Preparation 6ab (43 mg, 0.06 mmol) in dichloromethane (5 ml) was added Hunig's Base (32 μL, 0.18 mmol) and HBTU (23 mg, 0.06 mmol) followed by 1-phenylcyclopropane-1-carboxylic acid (9 mg, 0.06 mmol), and stirred at ambient temperature for *ca* 16 h. The reaction was diluted with dichloromethane and

washed with water, followed by aqueous 1M HCl and then brine. The organic extract was dried over magnesium sulfate, filtered and loaded onto a column and purified in ethyl acetate to obtain the desired product.

LC/MS ($C_{53}H_{52}N_4O_4$) 809 $[M+H]^+$; RT 2.92 (Method A)

5 ***Step B: N-(4-Hydroxyphenyl)-N-methyl-3-{7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(1-phenylcyclopropanecarbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide***

A solution of the product from Step A (47.6 mg, 0.06 mmol) in anhydrous dichloromethane (5 ml) was cooled to 0 °C and to this was added boron trichloride (1M in DCM, 120 μ L, 0.12 mmol). The reaction was stirred at ambient temperature for *ca* 16 h. The reaction was quenched with methanol (5 mL), and the reaction was diluted with dichloromethane. The organic phase was washed with water, dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by column chromatography in a gradient of dichloromethane to 5% methanol in dichloromethane followed by trituration with ether and dried *in vacuo*.

LC/MS ($C_{46}H_{46}N_4O_4$) 719 $[M+H]^+$; RT 2.65 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{46}N_4O_4$

$[M+H]^+$ calculated: 719.3592

20 $[M+H]^+$ measured: 719.3556

Example 110: N-Ethyl-5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-6-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1H-isoindole-2-carboxamide

High-resolution mass (ESI+):

25 Empirical formula: $C_{36}H_{39}N_5O_4$

$[M+H]^+$ calculated: 606.3082

$[M+H]^+$ measured: 606.3078

Example 111: *N*-Benzyl-5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₁N₅O₄

$[M+H]^+$ calculated: 668.3239

$[M+H]^+$ measured: 668.3203

Example 112: 5-{2-Acetyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₃₅H₃₆N₄O₄

$[M+H]^+$ calculated: 577.2817

$[M+H]^+$ measured: 577.2782

Example 113: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-{4-[(4-methylpiperazin-1-yl)methyl]phenyl}acetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: 2-{4-[(4-methylpiperazin-1-yl)methyl]phenyl}acetic acid

To a suspension of 4(bromomethyl)phenyl acetic acid (100 mg, 0.44 mmol) in acetonitrile (2 ml) was added potassium carbonate (126 mg, 0.88 mmol) and *N*-methylpiperazine (100 μ L, 0.88 mmol), and the reaction stirred at ambient temperature for *ca* 16 h. The reaction was filtered and washed with ethyl acetate. The filtrate was basified with methanolic sodium hydroxide and then concentrated *in vacuo*. The crude material was dissolved in methanol and loaded onto a PE-AX column which had been pre-wetted with dichloromethane. The column was then washed with dichloromethane and methanol, and the product eluted with 10% formic acid/dichloromethane.

LC/MS ($C_{14}H_{20}N_2O_2$) 249 $[M+H]^+$; RT 0.42 (Method A)

Step B: *N*-[4-(benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-{4-[(4-methylpiperazin-1-yl)methyl]phenyl}acetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution of the product from Preparation 6ab (50 mg, 0.08 mmol) in dichloromethane (3 mL) was added triethylamine (56 μ L, 0.4 mmol) and HBTU (30 mg, 0.08 mmol) followed by the acid obtained in Step A (30 mg, 0.1 mmol) in dichloromethane (2 mL). The reaction was stirred at ambient temperature for *ca* 2 h. The reaction was diluted with dichloromethane and washed with water, dried over magnesium sulphate, filtered and concentrated *in vacuo*. The reaction was purified by column chromatography on 10 g silica column in a gradient of dichloromethane to 5% methanol in dichloromethane.

LC/MS ($C_{57}H_{62}N_6O_4$) 448 $[M+2H]^{2+}$; RT 2.51 (Method A)

Step C: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-{4-[(4-methylpiperazin-1-yl)methyl]phenyl}acetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

A solution of the product from Step B (44 mg, 0.05 mmol) in ethanol (10 mL) is added to 10% Pd/C (catalytic) and the mixture is shaken under an atmosphere of hydrogen for *ca* 16 h. Filtration through celite, evaporation, and purification on column chromatography (4 g

silica, dichloromethane to 5% methanol in dichloromethane) afforded the desired product. Trituration with diethyl ether afforded the desired product.

LC/MS ($C_{50}H_{56}N_6O_4$) 803 $[M-H]^-$; RT 2.20 (Method A)

High-resolution mass (ESI+):

5 Empirical formula: $C_{50}H_{56}N_6O_4$

$[M+2H]^{2+}$ calculated: 403.2254

$[M+2H]^{2+}$ measured: 403.2252

Example 114: Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

10

A solution of the product obtained from Preparation 6cb (50 mg, 0.08 mmol) and *N,N*-diisopropylethylamine (70 μ L, 0.4 mmol) was dissolved in dichloromethane (5 mL) and cooled to 0 °C. To this was added phenylchloroformate (11 μ L, 0.09 mmol), and the mixture was stirred at ambient temperature for 15 min. The reaction was diluted with
15 dichloromethane, then washed with aqueous 1M sodium hydroxide, and brine. The organic extract was dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (4 g silica, dichloromethane to 4% methanol in dichloromethane) afforded the desired product.

LC/MS ($C_{41}H_{40}N_4O_4$) 653 $[M+H]^+$; RT 1.48 (Method B)

20

High-resolution mass (ESI+):

Empirical formula: $C_{41}H_{40}N_4O_4$

$[M+H]^+$ calculated: 653.3122

$[M+H]^+$ measured: 653.3159

Example 115: Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 114, replacing the product from Preparation 6cb with the product from Preparation 6ca.

LC/MS (C₄₅H₄₇N₅O₅) 738 [M+H]⁺; RT 1.28 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₄₇N₅O₅

[M+H]⁺ calculated: 738.3650

[M+H]⁺ measured: 738.3655

Example 116: *N*-(4-Hydroxyphenyl)-5-{2-[3-(4-methoxyphenyl)propyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₆N₄O₄

[M+H]⁺ calculated: 683.3599

[M+H]⁺ measured: 683.3608

Example 117: *N*,1,2-trimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-phenyl-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 114, replacing the product from Preparation 6cb with the product from preparation 6ca, and replacing phenyl chloroformate with phenylacetyl chloride.

LC/MS (C₄₆H₄₉N₅O₄) 736 [M+H]⁺; RT 1.23 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₆H₄₉N₅O₄

[M+H]⁺ calculated: 736.3857

5 [M+H]⁺ measured: 736.3828

Example 118: *N*,1,2-Trimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 The procedure is as in Example 114, replacing phenyl chloroformate with phenylacetyl chloride.

LC/MS (C₄₂H₄₂N₄O₃) 651 [M+H]⁺; RT 1.40 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₂H₄₂N₄O₃

[M+H]⁺ calculated: 651.3330

15 [M+H]⁺ measured: 651.3344

Example 119: *N*,1,2-Trimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-phenyl-1*H*-pyrrole-3-carboxamide hydrochloride

20 The product from Preparation 6cb (22 mg, 0.03 mmol) was dissolved in ethyl acetate (10 mL) and washed with saturated sodium bicarbonate (10 mL). The organic phase was separated, dried over magnesium sulfate and concentrated *in vacuo*. The solid was then dissolved in methanol (2 mL) and to this was added 1M HCl in diethyl ether (0.15 mL, 5 eq.), stirred for 30 min, and then concentrated *in vacuo*. The solid was then triturated in a

small amount of diethyl ether and the solids were isolated by filtration. They were washed with cold ether before being dried under vacuum for 1 h.

LC/MS ($C_{34}H_{36}N_4O_2$) 533 $[M+H]^+$; RT 1.08 (Method B)

Example 120: 4-Fluorophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-
5 5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-
tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing 2-phenylacetyl chloride in Step A with 4-fluorophenyl chloroformate.

10 **LC/MS** ($C_{47}H_{48}N_5O_6F$) 798 $[M+H]^+$; RT 1.22 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{47}H_{48}N_5O_6F$

$[M+H]^+$ calculated: 798.3661

$[M+H]^+$ measured: 798.3663

15 **Example 121:** 4-Methoxyphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-
5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-
tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as described in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing 2-phenylacetyl chloride in Step A with 4-methoxyphenyl chloroformate.

20 **LC/MS** ($C_{48}H_{51}N_5O_7$) 810 $[M+H]^+$; RT 1.21 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{48}H_{51}N_5O_7$

$[M+H]^+$ calculated: 810.3861

$[M+H]^+$ measured: 810.3826

Example 122: 4-Chlorophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: 4-Chlorophenyl 6-(1-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-5,6,7,8-tetrahydroindolizin-3-yl)-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step A of Example 11, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing propionyl chloride with 4-chlorophenyl chloroformate.

LC/MS ($C_{54}H_{54}ClN_5O_6$) 904 $[M+H]^+$; RT 1.51 (Method B)

Step B: 4-Chlorophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step B of Example 85, replacing the product of Step A in Example 85 with the product of Step A in Example 122.

LC/MS ($C_{47}H_{48}ClN_5O_6$) 814 $[M+H]^+$; RT 1.27 (Method B)

Example 123: 3-{2-[2-(3-{[2-(Dimethylamino)ethyl]amino}phenyl)acetyl]-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-N-(4-hydroxyphenyl)-N-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Example 96, replacing 2-(morpholin-4-yl)ethan-1-amine in Step A with (2-aminoethyl)dimethylamine, and replacing ethyl 2-(4-bromophenyl)acetate in Step A with methyl 2-(3-bromophenyl)acetate.

LC/MS ($C_{48}H_{54}N_6O_4$) 777 $[M-H]^-$; RT 2.23 (Method A)

5 **Example 124:** 4-(Trifluoromethyl)phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

10 The procedure is as in Example 74, replacing the product from Preparation 6ab in Step A with the product from Preparation 6aa, and replacing *N*-methylaniline in Step B with 4-(trifluoromethyl)phenol.

LC/MS ($C_{48}H_{48}N_5O_6F_3$) 848 $[M+H]^+$; RT 1.31 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{48}H_{48}N_5O_6F_3$

15 $[M+H]^+$ calculated: 848.3629

$[M+H]^+$ measured: 848.3615

20 **Example 125:** 4-(Trifluoromethoxy)phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 74, replacing the product from Preparation 6ab in Step A with the product from Preparation 6aa, and replacing *N*-methylaniline in Step B with 4-(trifluoromethoxy)phenol.

LC/MS ($C_{48}H_{48}N_5O_7F_3$) 864 $[M+H]^+$; RT 1.31 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₈H₄₈N₅O₇F₃

[M+H]⁺ calculated: 864.3579

[M+H]⁺ measured: 864.3576

5 **Example 126:** **4-Ethylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-
5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-
tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate**

10 The procedure is as in Example 74, replacing the product from Preparation 6ab in Step A
with the product from Preparation 6aa, and replacing *N*-methylaniline in Step B with 4-
ethylphenol.

LC/MS (C₄₉H₅₃N₅O₆) 808 [M+H]⁺; RT 1.32 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₉H₅₃N₅O₆

[M+H]⁺ calculated: 808.4069

15 [M+H]⁺ measured: 808.4026

Example 127: **4-Cyanophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-
5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-
tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate**

20 The procedure is as in Example 74, replacing the product from Preparation 6ab in Step A
with the product from Preparation 6aa, and replacing *N*-methylaniline in Step B with 4-
hydroxybenzonitrile.

LC/MS (C₄₈H₄₈N₆O₆) 805 [M+H]⁺; RT 1.18 (Method B)

Example 128: 2-Methoxyphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step A of Example 8, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing 2-phenylacetyl chloride with 2-methoxyphenyl chloroformate.

LC/MS (C₄₈H₅₁N₅O₇) 810 [M+H]⁺; RT 1.20 (Method B)

Example 129: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[4-(morpholin-4-ylmethyl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *N*-[4-(Benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[4-(morpholin-4-ylmethyl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Step A and B of Example 113, replacing *N*-methylpiperazine in Step A with morpholine.

LC/MS (C₅₆H₅₉N₅O₅) 882 [M+H]⁺; RT 2.51 (Method A)

Step B: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[4-(morpholin-4-ylmethyl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

A solution of the product from Step A (49 mg, 0.06 mmol) in anhydrous dichloromethane (5 ml) was cooled to 0 °C and to this was added boron trichloride (1M in DCM, 170 µL, 0.17 mmol). The reaction was stirred at ambient temperature for *ca* 16 h. The reaction was quenched with methanol (5 mL), and the reaction was diluted with dichloromethane. The organic phase was washed with water, dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude product was purified by column chromatography in a gradient of dichloromethane to 7% methanol in dichloromethane followed by trituration with ether and dried *in vacuo*.

LC/MS (C₄₉H₅₃N₅O₅) 792 [M+H]⁺; RT 2.20 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₉H₅₃N₅O₅

[M+2H]²⁺ calculated: 396.7096

5 [M+2H]²⁺ measured: 396.7104

Example 130: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(3-phenylpropanoyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

10 The procedure was as in Example 23, replacing the product from Preparation 7ab in Step A with the product from Preparation 6ba, and replacing 2-methyl-3-phenylpropanoic acid in Step A with 3-phenylpropanoic acid.

LC/MS (C₄₇H₅₁N₅O₅) 766 [M+H]⁺; RT 1.14 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₅₁N₅O₅

15 [M+H]⁺ calculated: 766.3963

[M+H]⁺ measured: 766.3955

Example 131: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[(2*E*)-3-phenylprop-2-enoyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

20 The procedure was as in Example 109, replacing the product from Preparation 6ab with the product from Preparation 6ba, and replacing 2-methyl-3-phenylpropanoic acid with (2*E*)-3-phenylprop-2-enoic acid.

LC/MS (C₄₇H₄₉N₅O₅) 764 [M+H]⁺; RT 1.15 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₉N₅O₅

[M+H]⁺ calculated: 764.3806

[M+H]⁺ measured: 764.3779

5 **Example 132:** **5-[2-(3-Cyclohexylpropanoyl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide**

The procedure was as in Example 23, replacing the product from Preparation 7ab in Step A with the product from Preparation 6ba, and replacing 2-methyl-3-phenylpropanoic acid in
10 Step A with 3-cyclohexylpropanoic acid.

LC/MS (C₄₇H₅₇N₅O₅) 772 [M+H]⁺; RT 1.25 (Method B)

Example 133: **Phenyl 5-{1-[(4-hydroxyphenyl)(phenyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate**

15 ***High-resolution mass (ESI+):***

Empirical formula: C₄₇H₄₃N₄O₅

[M+H]⁺ calculated: 743.3228

[M+H]⁺ measured: 743.3234

Example 134: ***N*-(4-hydroxyphenyl)-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-phenyl-5,6,7,8-tetrahydroindolizine-1-carboxamide**

20

High-resolution mass (ESI+):

Empirical formula: C₄₉H₄₇N₄O₄

[M+H]⁺ calculated: 755.3592

[M+H]⁺ measured: 755.3595

Example 135: *N*-(4-Hydroxyphenyl)-5-{2-[3-(3-methoxyphenyl)propanoyl]-7-
5 [(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-
tetrahydroisoquinolin-6-yl}-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

The procedure was as in Example 23, replacing the product from Preparation 7ab in Step A with the product from Preparation 6ba, and replacing 2-methyl-3-phenylpropanoic acid in Step A with 3-(2-methoxyphenyl)propanoic acid.

10 **LC/MS** (C₄₈H₅₃N₅O₆) 796 [M+H]⁺; RT 1.14 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₈H₅₃N₅O₆

[M+H]⁺ calculated: 796.4069

[M+H]⁺ measured: 796.4068

15 **Example 136:** *N*-(4-Hydroxyphenyl)-5-{2-[(2*E*)-3-(3-methoxyphenyl)prop-2-
enoyl]-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-
1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

20 The procedure was as in Example 109, replacing the product from Preparation 6ab with the product from Preparation 6ba, and replacing 2-methyl-3-phenylpropanoic acid with (2*E*)-3-(2-methoxyphenyl)prop-2-enoic acid.

LC/MS (C₄₈H₅₁N₅O₆) 794 [M+H]⁺; RT 1.16 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₈H₅₁N₅O₆

$[M+H]^+$ calculated: 794.3912

$[M+H]^+$ measured: 794.3908

Example 137: 5- $\{2-[(2E)-3-(3,4-dichlorophenyl)prop-2-enoyl]-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl\}-N-(4-hydroxyphenyl)-N,1,2-trimethyl-1H$ -pyrrole-3-carboxamide

The procedure was as in Example 109, replacing the product from Preparation 6ab with the product from Preparation 6ba, and replacing 2-methyl-3-phenylpropanoic acid with (2E)-3-(3,4-dichlorophenyl)prop-2-enoic acid.

10 **LC/MS** ($C_{47}H_{47}N_5O_5Cl_2$) 832 $[M+H]^+$; RT 1.27 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{47}H_{47}N_5O_5Cl_2$

$[M+H]^+$ calculated: 832.3027

$[M+H]^+$ measured: 832.3000

15 **Example 138:** 3- $\{2-[3-(4-\{2-(Dimethylamino)ethyl\}amino)phenyl]propanoyl]-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl\}-N-(4-hydroxyphenyl)-N-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide$

20 The procedure is as described in Example 96, replacing 2-(morpholin-4-yl)ethan-1-amine in Step A with (2-aminoethyl)dimethylamine, and replacing ethyl 2-(4-bromophenyl)acetate in Step A with methyl 3-(4-bromophenyl)propanoate.

LC/MS ($C_{49}H_{56}N_6O_4$) 791 $[M-H]^-$; RT 2.22 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₉H₅₆N₆O₄

[M+H]⁺ calculated: 793.4436

[M+H]⁺ measured: 793.4411

Example 139: 4-Methylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 11, replacing the product from Preparation 7aa with the Product from Preparation 6ba, and replacing propionyl chloride with 4-methylphenyl chloroformate.

LC/MS (C₄₆H₄₉N₅O₆) 768 [M+H]⁺; RT 1.22 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₉N₅O₆

[M+H]⁺ calculated: 768.3756

[M+H]⁺ measured: 768.3723

Example 140: 4-Chlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: 4-Chlorophenyl 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1H-pyrrol-2-yl)-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step A of Example 11, replacing the product from Preparation 7aa in Step A with the product from Preparation 6ba, and replacing propionyl chloride with 4-chlorophenyl chloroformate.

LC/MS ($C_{52}H_{52}N_5O_6Cl$) 878 $[M+H]^+$; RT 1.48 (Method B)

Step B: 4-Chlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

- 5 The procedure is as in Step B of Example 85, replacing product in Step A of Example 85 with product of Step A in Example 140.

LC/MS ($C_{45}H_{46}N_5O_6Cl$) 788 $[M+H]^+$; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{45}H_{46}N_5O_6Cl$

10 $[M+H]^+$ calculated: 788.3209

$[M+H]^+$ measured: 788.3191

Example 141: 4-Carbamoylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

15

The procedure is as in Example 74, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing *N*-methylaniline with 4-hydroxybenzamide.

LC/MS ($C_{48}H_{50}N_6O_7$) 823 $[M+H]^+$; RT 1.06 (Method B)

20 ***High-resolution mass (ESI+):***

Empirical formula: $C_{48}H_{50}N_6O_7$

$[M+H]^+$ calculated: 823.3814

$[M+H]^+$ measured: 823.3788

Example 142: 4-(Dimethylcarbamoyl)phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

- 5 The procedure is as in Example 74, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing *N*-methylaniline with 4-hydroxy-*N,N*-dimethylbenzamide.

LC/MS ($C_{50}H_{54}N_6O_7$) 851 $[M+H]^+$; RT 1.11 (Method B)

High-resolution mass (ESI+):

- 10 Empirical formula: $C_{50}H_{54}N_6O_7$

$[M+H]^+$ calculated: 851.4127

$[M+H]^+$ measured: 851.4094

Example 143: 4-(1*H*-imidazol-1-yl)phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

15

The procedure is as in Example 74, replacing the product from Preparation 7aa in Step A with the product from Preparation 6aa, and replacing *N*-methylaniline with 4-(1*H*-imidazol-1-yl)phenol.

- 20 **LC/MS** ($C_{50}H_{51}N_7O_6$) 846 $[M+H]^+$; RT 1.03 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{50}H_{51}N_7O_6$

$[M+2H]^{2+}$ calculated: 423.7023

$[M+2H]^{2+}$ measured: 423.7023

Example 144: 3-[2-(2-{4-[(Dimethylamino)methyl]phenyl}acetyl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

The procedure is as in Example 113, replacing replacing *N*-methylpiperazine in Step A with dimethylamine.

LC/MS (C₄₇H₅₁N₅O₄) 748 [M-H]⁻; RT 2.18 (Method A)

High-resolution mass (ESI⁺):

Empirical formula: C₄₇H₅₁N₅O₄

[M+H]⁺ calculated: 750.4014

[M+H]⁺ measured: 750.3984

Example 145: 5-{2-[2-(4-Cyanophenyl)acetyl]-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 109, replacing the product from Preparation 6ab in Step A with the product from Preparation 6bb, and replacing 1-phenylcyclopropane-1-carboxylic acid in Step A with 2-(4-cyanophenyl)acetic acid.

LC/MS (C₄₃H₄₁N₅O₄) 692 [M+H]⁺; RT 1.24 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₃H₄₁N₅O₄

[M+H]⁺ calculated: 692.3231

[M+H]⁺ measured: 692.3199

Example 146: 4-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 573, replacing 3,4-dichlorophenol in Step B with 4-hydroxybenzonitrile.

LC/MS (C₄₆H₄₆N₆O₆) 779 [M+H]⁺; RT 1.14 (Method B)

Example 147: 4-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 573, replacing the product from Preparation 6ba with the product from Preparation 6bb, and replacing 3,4-dichlorophenol in Step B with 4-hydroxybenzonitrile.

LC/MS (C₄₂H₃₉N₅O₅) 694 [M+H]⁺; RT 1.29 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₂H₃₉N₅O₅

[M+H]⁺ calculated: 694.3024

[M+H]⁺ measured: 694.3044

Example 148: 4-Cyano-2-methoxyphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 573, replacing the product from Preparation 6ba with the product from Preparation 7aa, and replacing 3,4-dichlorophenol in Step B with 4-hydroxybenzonitrile.

LC/MS (C₄₉H₅₀N₆O₇) 835 [M+H]⁺; RT 1.19 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₉H₅₀N₆O₇

[M+H]⁺ calculated: 835.3814

5 [M+H]⁺ measured: 835.3833

Example 149: 5-{2-[2-(4-{[2-(Dimethylamino)ethyl]amino}phenyl)acetyl]-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 The procedure is as in Example 96, replacing 2-(morpholin-4-yl)ethan-1-amine in Step A with (2-aminoethyl)dimethylamine, and replacing the product from Preparation 6ba in Step C with the product obtained from Preparation 6cb.

LC/MS (C₄₆H₅₂N₆O₃) 737 [M+H]⁺; RT 1.13 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₆H₅₂N₆O₃

15 [M+H]⁺ calculated: 737.4174

[M+H]⁺ measured: 737.4173

Example 150: 5-{2-[2-(4-Cyanophenyl)acetyl]-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

20 The procedure is as in Example 109, replacing the product from Preparation 6ab in Step A with the product from Preparation 6cb, and replacing 1-phenylcyclopropane-1-carboxylic acid in Step A with 2-(4-cyanophenyl)acetic acid.

LC/MS (C₄₃H₄₁N₅O₃) 676 [M+H]⁺; RT 1.36 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₁N₅O₃

[M+H]⁺ calculated: 676.3282

[M+H]⁺ measured: 676.3249

5 **Example 151:** 5-{2-[2-(4-Cyanophenyl)acetyl]-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 The procedure is as in Example 109, replacing the product from Preparation 6ab in Step A with the product from Preparation 6ca, and replacing 1-phenylcyclopropane-1-carboxylic acid in Step A with 2-(4-cyanophenyl)acetic acid.

LC/MS (C₄₇H₄₈N₆O₄) 761 [M+H]⁺; RT 1.19 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₈N₆O₄

[M+H]⁺ calculated: 761.3810

15 [M+H]⁺ measured: 761.3811

Example 152: 4-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

20 The procedure is as in Step A and B Example 573, replacing 3,4-dichlorophenol in Step B with 4-hydroxybenzonitrile.

LC/MS (C₄₆H₄₆N₆O₅) 763 [M+H]⁺; RT 1.26 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₆N₆O₅

[M+H]⁺ calculated: 763.3602

[M+H]⁺ measured: 763.3572

Example 153: 4-{{2-(Dimethylamino)ethyl}amino}phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 489, replacing the product from Preparation 6cb with the product from Preparation 6ab.

LC/MS (C₄₇H₅₂N₆O₅) 779 [M-H]⁻; RT 2.27 (Method A)

High-resolution mass (ESI⁺):

Empirical formula: C₄₇H₅₂N₆O₅

[M+2H]²⁺ calculated: 391.2072

[M+2H]²⁺ measured: 391.2081

Example 154: 3-[2-(2-{4-[2-(Dimethylamino)ethoxy]phenyl}acetyl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: methyl 2-{4-[2-(dimethylamino)ethoxy]phenyl}acetate

To a solution of methyl 4-hydroxyphenylacetate (50 mg, 0.3 mmol) in acetone (5 mL) was added cesium carbonate (196 mg, 0.6 mmol) followed by 2-dimethylaminoethyl chloride hydrochloride (45 mg, 0.31 mmol) and the reaction stirred at ambient temperature for *ca* 16 h. To this was added sodium iodide (45 mg, 0.3 mmol) and the mixture was heated at 60 °C for *ca* 48 h. The reaction was cooled to ambient temperature, diluted with dichloromethane and washed with water, dried over magnesium sulphate, filtered and

concentrated *in vacuo* and loaded onto a pre-packed 10 g silica column and purified by column chromatography in a gradient of dichloromethane to 10% methanol in dichloromethane to afford the desired product.

LC/MS (C₁₃H₁₉NO₃) 238 [M+H]⁺; RT 1.49 (Method A)

5 **Step B: sodium 2-{4-[2-(dimethylamino)ethoxy]phenyl}acetate**

To a solution of methyl 2-{4-[2-(dimethylamino)ethoxy]phenyl}acetate (19 mg, 0.08 mmol) in methanol (3 mL) was added sodium hydroxide (2M, 80 µL, 0.16 mmol) and the reaction stirred at ambient temperature for *ca* 48 h. The reaction was concentrated *in vacuo* to afford the desired product and taken on assuming quantitative transformation.

10 LC/MS (C₁₂H₁₇NO₃) 224 [M+H]⁺; RT 0.53 (Method A)

Step C: N-[4-(benzyloxy)phenyl]-3-[2-(2-{4-[2-(dimethylamino)ethoxy]phenyl}acetyl)-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-N-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

15 To a suspension of the product from Step B (16 mg, 0.06 mmol) in dichloromethane (5 mL) was added the product from Preparation 6ab (50 mg, 0.06 mmol) followed by triethylamine (27 µL, 0.19 mmol) and HBTU (24 mg, 0.06 mmol) and the reaction stirred at ambient temperature for *ca* 16 h. The reaction was diluted with dichloromethane and washed with water, dried over magnesium sulphate, filtered and concentrated *in vacuo* and loaded onto a pre-packed 10 g silica column in dichloromethane and purified by column chromatography in a gradient of dichloromethane to 10% methanol in dichloromethane

20

LC/MS (C₅₅H₅₉N₅O₅) 435.5 [M+2H]²⁺; RT 2.54 (Method A)

Step D: 3-[2-(2-{4-[2-(Dimethylamino)ethoxy]phenyl}acetyl)-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-N-(4-hydroxyphenyl)-N-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide

25 To a solution of the product obtained from Step C (51.8 mg, 0.04 mmol, 1 eq) in ethanol (5 mL) was added 10% Pd/C (catalytic). The reaction was stirred under a hydrogen atmosphere for *ca* 48. The reaction was filtered through a celite cartridge and washed with

methanol and concentrated *in vacuo* and purified by reverse phase preparative HPLC (H₂O-TFA/acetonitrile; gradient elution).

LC/MS (C₄₈H₅₃N₅O₅) 778 [M-H]⁻; RT 2.20 (Method A)

Example 155: 4-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 786, replacing the product from Preparation 6cb in Step A with the product from Preparation 6ab, and 4-hydroxybenzonitrile in Step B with 3-(4-hydroxyphenyl)-*N,N*-dimethylpropanamide..

LC/MS (C₄₈H₅₁N₅O₆) 794 [M+H]⁺; RT 2.57 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₈H₅₁N₅O₆

[M+H]⁺ calculated: 794.3912

[M+H]⁺ measured: 794.3950

Example 156: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenoxyacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₉N₅O₆

[M+H]⁺ calculated: 780.3763

[M+H]⁺ measured: 780.3759

Example 157: 1,1-Dioxo-1 λ^6 -thiolan-3-yl 5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

High-resolution mass (ESI+):

5 Empirical formula: C₃₈H₄₀N₄O₇S

[M+H]⁺ calculated: 697.2698

[M+H]⁺ measured: 697.2696

Example 158: *N*-(4-Hydroxyphenyl)-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenoxyacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-{1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Example 159: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenoxyacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}-1*H*-pyrrole-3-carboxamide

15 **High-resolution mass (ESI+):**

Empirical formula: C₄₅H₄₇N₅O₆

[M+H]⁺ calculated: 754.3606

[M+H]⁺ measured: 754.3574

Example 160: 5-{2-[2-(Benzyloxy)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

20 **High-resolution mass (ESI+):**

Empirical formula: C₄₂H₄₂N₄O₅

[M+H]⁺ calculated: 683.3235

[M+H]⁺ measured: 683.3238

Example 161: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-(1,2,3,4-tetrahydroisoquinoline-2-carbonyl)-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The product from Preparation 6dc (81 mg, 0.15 mmol, 1 eq) in dichloromethane (5 mL) was cooled to 0°C. To this was added *N,N*-diisopropylethylamine (52 µL, 0.3 mmol) followed by phenyl chloroformate (21 µL, 0.16 mmol) and the mixture was allowed to stir for 15 min. The reaction was diluted with dichloromethane and washed sequentially with 1M aqueous NaOH, and brine. The organic extracts were dried over magnesium sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 12 g RediSep™ silica cartridge; gradient of dichloromethane to 5 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₄₁H₄₈N₄O₄) 661 [M+H]⁺; RT 1.57 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₁H₄₈N₄O₄

[M+H]⁺ calculated: 661.3748

[M+H]⁺ measured: 661.3769

Example 162: *N,N*-Dibutyl-1,2-dimethyl-5-[2-(2-phenylacetyl)-7-(1,2,3,4-tetrahydroisoquinoline-2-carbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 161, replacing phenyl chloroformate with phenylacetyl chloride.

LC/MS ($C_{42}H_{50}N_4O_3$) 659 $[M+H]^+$; RT 1.52 (Method B)

High-resolution mass (ESI+):

5 Empirical formula: $C_{42}H_{50}N_4O_3$

$[M+H]^+$ calculated: 659.3956

$[M+H]^+$ measured: 659.3969

Example 163: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-
[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-
10 tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 161, replacing the product from Preparation 6dc with product from Preparation 6db.

LC/MS ($C_{42}H_{50}N_4O_4$) 675 $[M+H]^+$; RT 1.52 (Method B)

High-resolution mass (ESI+):

15 Empirical formula: $C_{42}H_{50}N_4O_4$

$[M+H]^+$ calculated: 675.3905

$[M+H]^+$ measured: 675.3900

Example 164: *N,N*-Dibutyl-1,2-dimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-
tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-
20 6-yl}-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 161, replacing the product from Preparation 6dc with product from Preparation 6db, and replacing phenyl chloroformate with phenylacetyl chloride.

LC/MS (C₄₃H₅₂N₄O₃) 673 [M+H]⁺; RT 1.51 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₅₂N₄O₃

[M+H]⁺ calculated: 673.4112

5 [M+H]⁺ measured: 673.4087

Example 165: *N,N*-Dibutyl-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

10 To a solution of the product of Preparation 6da (112 mg, 0.13 mmol) in dichloromethane (5 mL) was added *N,N*-diisopropylethylamine (112 µL, 0.64 mmol) followed by phenylacetyl chloride (17 µL, 0.14 mmol) and the mixture was allowed to stir for 15 min. The reaction was diluted with dichloromethane and washed sequentially with 1M aqueous sodium hydroxide, and brine. The organic extracts were dried over magnesium sulphate, filtered and concentrated. Purification by flash column chromatography (CombiFlash R_f,
15 12 g RediSep™ silica cartridge) eluting with a gradient of dichloromethane to 5 % methanol in dichloromethane afforded the desired product.

LC/MS (C₄₇H₅₉N₅O₄) 758 [M+H]⁺; RT 1.40 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₅₉N₅O₄

20 [M+H]⁺ calculated: 758.4640

[M+H]⁺ measured: 758.4648

Example 166: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenoxyacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}indolizine-1-carboxamide

High-resolution mass (ESI+):

5 Empirical formula: C₄₇H₄₅N₅O₆

[M+H]⁺ calculated: 776.3450

[M+H]⁺ measured: 776.3430

Example 167: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenoxyacetyl)-2,3-dihydro-1*H*-isoindol-5-yl}indolizine-1-carboxamide

10

High-resolution mass (ESI+):

Empirical formula: C₄₃H₃₈N₄O₅

[M+H]⁺ calculated: 691.2922

[M+H]⁺ measured: 691.2913

Example 168: 4-Methylphenyl 5-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxylate

15

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₂N₄O₅

20 [M+H]⁺ calculated: 695.3235

[M+H]⁺ measured: 695.3230

Example 169: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[3-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

Step A: *methyl 2-[3-(4-methylpiperazin-1-yl)phenyl]acetate*

- 5 To a boiling tube was added methyl 2-(3-bromophenyl)acetate (250 mg, 1.09 mmol), 1-methylpiperazine (0.18 mL, 1.64 mmol), di-*tert*-butyl(2-phenylphenyl)phosphane, (32.57 mg, 0.11 mmol) and potassium phosphate tribasic (463.32 mg, 2.18 mmol) which were then suspended in toluene (5 mL) and degassed by sparging with nitrogen for 5 min. Tris(Dibenzylideneacetone)Dipalladium(0) (49.97 mg, 0.05 mmol) was then added to the
- 10 reaction and stirred at 90 °C under nitrogen for *ca* 16 h. The reaction was filtered and solids washed with dichloromethane and concentrated *in vacuo*, and then loaded onto a pre-packed 10 g silica column in dichloromethane and purified by column chromatography in a gradient of dichloromethane to 5% methanol in dichloromethane.

LC/MS (C₁₄H₂₀N₂O₂) 249 [M+H]⁺; RT 1.56 (Method B)

15 **Step B:** *sodium 2-[3-(4-methylpiperazin-1-yl)phenyl]acetate*

To a solution of the product from Step A (39.3 mg, 0.16 mmol) in methanol (5 mL) was added sodium hydroxide (2M, 0.16 mL, 0.32 mmol) and the reaction stirred at ambient temperature for *ca* 16 h. The reaction was concentrated *in vacuo* and triturated with diethyl ether to afford the desired product.

20 LC/MS (C₁₃H₁₈N₂O₂) 235 [M+H]⁺; RT 0.69 (Method B)

Step C : *N*-[4-(benzyloxy)phenyl]-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[3-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

- 25 To a solution of the product from Preparation 6ab (50 mg, 0.06 mmol) in dichloromethane (3 mL) was added HBTU (24.35 mg, 0.06 mmol) and triethylamine (26.79 µL, 0.19 mmol) followed by the product from Step B (24.68 mg, 0.1 mmol) and the reaction stirred at ambient temperature for *ca* 16 h. The reaction was diluted with dichloromethane and washed with water and brine. The organic extract was dried over magnesium sulphate, filtered and concentrated *in vacuo*, and loaded onto a pre-packed 5 g silica column in

dichloromethane and purified by column chromatography in a gradient of dichloromethane in 5% methanol in dichloromethane.

LC/MS (C₅₆H₆₀N₆O₄) 881 [M+H]⁺; RT 2.52 (Method B)

Step D : *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[3-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

To a solution the product from Step C (45 mg, 0.05 mmol) in ethanol (10 mL) was added 10% Pd/C (catalytic), and stirred under a hydrogen atmosphere at ambient temperature for ca 16 h. The reaction was filtered through a celite cartridge, which was washed with methanol and the filtrate was concentrated *in vacuo*, and loaded onto a pre-packed 5 g silica column in dichloromethane and purified by column chromatography in a gradient of dichloromethane to 5% methanol in dichloromethane.

LC/MS (C₄₉H₅₄N₆O₄) 791 [M+H]⁺; RT 2.21 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₉H₅₄N₆O₄

[M+2H]²⁺ calculated: 396.2176

[M+2H]²⁺ measured: 396.2176

Example 170: 5-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-*N*-methyl-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindole-2-carboxamide

High-resolution mass (ESI⁺):

Empirical formula: C₃₅H₃₇N₅O₄

[M+H]⁺ calculated: 592.2926

[M+H]⁺ measured: 592.2925

Example 171: *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(phenylsulfanyl)acetyl]-2,3-dihydro-1*H*-isoindol-5-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide

High-resolution mass (ESI+):

5 Empirical formula: C₄₃H₄₂N₄O₄S

[M+H]⁺ calculated: 711.3007

[M+H]⁺ measured: 711.3001

Example 172: 5-{2-[2-(2*H*-1,3-Benzodioxol-5-yl)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₄₂H₄₀N₄O₆

[M+H]⁺ calculated: 697.3028

[M+H]⁺ measured: 697.2994

Example 173: 5-(2-[(2*S*)-2-Amino-2-phenylacetyl]-6-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2,3-dihydro-1*H*-isoindol-5-yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₁N₅O₄

20 [M+H]⁺ calculated: 668.3239

[M+H]⁺ measured: 668.3238

Example 174: 5-(2-[(2*R*)-2-Amino-2-phenylacetyl]-6-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2,3-dihydro-1*H*-isoindol-5-yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

5 Empirical formula: C₄₁H₄₁N₅O₄

[M+H]⁺ calculated: 668.3239

[M+H]⁺ measured: 668.3247

Example 175: *N*-(4-Hydroxyphenyl)-5-(2-[(2*R*)-2-hydroxy-2-phenylacetyl]-6-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2,3-dihydro-1*H*-isoindol-5-yl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₀N₄O₅

[M+H]⁺ calculated: 669.3079

[M+H]⁺ measured: 669.3063

15 **Example 176:** 5-{2-[2-(1-Piperidyl)acetyl]-6-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2,3-dihydro-1*H*-isoindol-5-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

High-resolution mass (ESI+):

Empirical formula: C₄₀H₄₅N₅O₄

20 [M+H]⁺ calculated: 660.3552

[M+H]⁺ measured: 660.3512

Example 177: 5-[2-[2-(1,3-Benzodioxol-5-yl)acetyl]-6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindolin-5-yl]-*N*,1,2-trimethyl-*N*-(1-methylpyrrolo[2,3-*b*]pyridin-5-yl)pyrrole-3-carboxamide

Example 178: 5-[2-[2-(1,3-Benzodioxol-5-yl)acetyl]-6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindolin-5-yl]-*N*-(4-hydroxyphenyl)-1,2-dimethyl-*N*-(1-methyl-1*H*-pyrazol-4-yl)pyrrole-3-carboxamide

Example 179: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]-2-[(2*R*)-2-phenylpropanoyl]isoindolin-5-yl]pyrrole-3-carboxamide

Example 180: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]-2-[2-(3-thienyl)acetyl]isoindolin-5-yl]pyrrole-3-carboxamide

Example 181: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]-2-[(2*S*)-2-phenylpropanoyl]isoindolin-5-yl]pyrrole-3-carboxamide

Example 182: 5-[2-[3-(4-Chlorophenoxy)propanoyl]-6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]isoindolin-5-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethylpyrrole-3-carboxamide

Example 183: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[6-[(3*R*)-3-methyl-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]-2-(2-pyrrol-1-ylacetyl)isoindolin-5-yl]pyrrole-3-carboxamide

Example 184: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 185: Phenyl 6-{4-[ethyl(4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with product from Preparation 6f, and replacing phenylacetyl chloride in Step A with phenyl chloroformate.

LC/MS (C₄₆H₄₉N₅O₆) 768 [M+H]⁺; RT 1.21 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₉N₅O₆

[M+H]⁺ calculated: 768.3756

[M+H]⁺ measured: 768.3741

Example 186: Phenyl 6-{4-[(4-hydroxyphenyl)(propyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 777, replacing the product from Preparation 6be in Step A with the product from Preparation 6g, and replacing phenylacetyl chloride in Step A with phenyl chloroformate.

LC/MS (C₄₇H₅₁N₅O₆) 782 [M+H]⁺; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₅₁N₅O₆

[M+H]⁺ calculated: 782.3912

[M+H]⁺ measured: 782.3927

Example 187: Phenyl 6-{4-[butyl(4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 777, replacing the product from Preparation 6be in Step A with the product from Preparation 6h, and replacing phenylacetyl chloride in Step A with phenyl chloroformate.

LC/MS (C₄₈H₅₃N₅O₆) 796 [M+H]⁺; RT 1.29 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₈H₅₃N₅O₆

[M+2H]²⁺ calculated: 398.7071

[M+2H]²⁺ measured: 398.7075

Example 188: Phenyl 6-{4-[(4-hydroxyphenyl)(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 189: Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 190: Phenyl 6-{4-[ethyl(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 191: Phenyl 6-{1,5-dimethyl-4-[phenyl(propyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-
7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 192:** Phenyl 6-{4-[butyl(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 193: Phenyl 6-[4-(diphenylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-
{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 194:** Phenyl 6-{4-[butyl(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 195:** Phenyl 6-{4-[butyl(ethyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 196: Phenyl 6-{4-[butyl(propyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 197: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 165, replacing phenylacetyl chloride with phenyl chloroformate.

LC/MS (C₄₆H₅₇N₅O₅) 760 [M+H]⁺; RT 1.40 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₆H₅₇N₅O₅

[M+H]⁺ calculated: 760.4432

[M+H]⁺ measured: 760.4449

Example 198: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 199: Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 200: Phenyl 6-{4-[(4-hydroxyphenyl)(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 201: Phenyl 6-[4-(diphenylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 202: Phenyl 6-{4-[butyl(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-
2(1*H*)-carboxylate

5 **Example 203:** Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-
carboxylate

Example 204: Phenyl 7-(3,4-dihydroisoquinolin-2(1*H*)-ylcarbonyl)-6-{4-[(4-
hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 205:** Phenyl 7-(3,4-dihydroisoquinolin-2(1*H*)-ylcarbonyl)-6-{1,5-dimethyl-4-
[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-3,4-dihydroisoquinoline-2(1*H*)-
carboxylate

15 **Example 206:** Phenyl 7-(3,4-dihydroisoquinolin-2(1*H*)-ylcarbonyl)-6-{4-[(4-
hydroxyphenyl)(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 207: Phenyl 7-(3,4-dihydroisoquinolin-2(1*H*)-ylcarbonyl)-6-[4-
(diphenylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-3,4-dihydroisoquinoline-2(1*H*)-
carboxylate

Example 208: Phenyl 6-{4-[butyl(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-(3,4-dihydroisoquinolin-2(1*H*)-ylcarbonyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 209: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-(3,4-dihydroisoquinolin-2(1*H*)-ylcarbonyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 210: Phenyl 7-[(3*S*)-3-(aminomethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 211: Phenyl 7-[(3*S*)-3-[(dimethylamino)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: Phenyl 6-(4-{4-(benzyloxy)phenyl}(methyl)carbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-[(3*S*)-3-[(dimethylamino)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step A of Example 777, replacing phenylacetyl chloride with phenyl chloroformate.

LC/MS (C₅₀H₅₁N₅O₅) 802 [M+H]⁺; RT 1.34 (Method B)

Step B: Phenyl 7-[(3*S*)-3-[(dimethylamino)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step B of Example 777, but purification was *via* preparative HPLC (H₂O-TFA/acetonitrile; gradient elution).

LC/MS (C₄₃H₄₅N₅O₅) 712 [M+H]⁺; RT 1.13 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₄₅N₅O₅

[M+H]⁺ calculated: 712.3493

[M+H]⁺ measured: 712.3496

5 **Example 212:** Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(pyrrolidin-1-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 213:** Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(piperidin-1-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 214: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

15 **Step A:** Phenyl 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step A of Example 777, replacing the compound from Preparation 6be with the compound from Preparation 6bf, and replacing phenylacetyl chloride with phenyl chloroformate.

20 **LC/MS** (C₅₃H₅₆N₆O₅) 857 [M+H]⁺; RT 1.34 (Method B)

Step B: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step B of Example 777, but purification was *via* preparative HPLC (H₂O-TFA/acetonitrile; gradient elution).

LC/MS (C₄₆H₅₀N₆O₅) 767 [M+H]⁺; RT 1.14 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₅₀N₆O₅

[M+2H]²⁺ calculated: 384.1994

[M+2H]²⁺ measured: 384.1995

Example 215: Phenyl 7-[(3S)-3-[2-(dimethylamino)ethyl]-3,4-dihydroisoquinolin-2(1H)-yl]carbonyl}-6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-3,4-dihydroisoquinoline-2(1H)-carboxylate

Example 216: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-[2-(morpholin-4-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with product from Preparation 6bc, and replacing phenylacetyl chloride in Step A with phenyl chloroformate.

LC/MS (C₄₆H₄₉N₅O₆) 768 [M+H]⁺; RT 1.14 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₉N₅O₆

[M+H]⁺ calculated: 768.3756

[M+H]⁺ measured: 768.3756

Example 217: Phenyl 7-{[(3*S*)-3-(aminomethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 218:** Phenyl 7-{[(3*S*)-3-[(dimethylamino)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 219:** Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(pyrrolidin-1-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 220: Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(piperidin-1-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 221:** Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 222: Phenyl 7-{[(3*S*)-3-[2-(dimethylamino)ethyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 223: Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-[2-(morpholin-4-yl)ethyl]-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 224: Phenyl 7-[[*(3S)*-3-(aminomethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 225: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-[(dimethylamino)methyl]-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 226: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(pyrrolidin-1-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 227: phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(piperidin-1-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 228: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 229: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*S*)-3-[2-(dimethylamino)ethyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 230: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*S*)-3-[2-(morpholin-4-yl)ethyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 231: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 232: *N*-Butyl-*N*-(4-hydroxyphenyl)-1,2-dimethyl-5-{7-{[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 777, replacing the product from Preparation 6be in Step A with the product from Preparation 6h.

LC/MS ($C_{49}H_{55}N_5O_5$) 794 $[M+H]^+$; RT 1.23 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{49}H_{55}N_5O_5$

$[M+2H]^{2+}$ calculated: 397.7174

$[M+2H]^{2+}$ measured: 397.7178

Example 233: *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-[7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 234: *N*,1,2-Trimethyl-5-[7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

5 **Example 235:** *N*-Butyl-1,2-dimethyl-5-[7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 236: 1,2-Dimethyl-5-[7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N,N*-diphenyl-1*H*-pyrrole-3-carboxamide

10 **Example 237:** *N*-Butyl-*N*,1,2-trimethyl-5-[7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

15 **Example 238:** *N,N*-Dibutyl-1,2-dimethyl-5-[7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 239: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 240: *N*,1,2-Trimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 241: *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 242: 1,2-Dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N,N*-diphenyl-1*H*-pyrrole-3-carboxamide

Example 243: *N*-Butyl-*N*,1,2-trimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 244: *N,N*-Dibutyl-1,2-dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 245: 6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-*N*-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6ba, and replacing phenylacetyl chloride with benzyl isocyanate.

LC/MS (C₄₅H₄₈N₆O₅) 753 [M+H]⁺; RT 1.14 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₄₈N₆O₅

[M+H]⁺ calculated: 753.3740

[M+H]⁺ measured: 753.3759

5 **Example 246:** 6-{4-[Butyl(4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-
7-{{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-*N*-
phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

10 **Example 247:** 6-{4-[(4-Hydroxyphenyl)(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-
yl}-7-{{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-*N*-
phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 248: 6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{{{(3*S*)-
3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-*N*-phenyl-3,4-
dihydroisoquinoline-2(1*H*)-carboxamide

15 **Example 249:** 6-{4-[Butyl(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{{{(3*S*)-3-
(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-*N*-phenyl-3,4-
dihydroisoquinoline-2(1*H*)-carboxamide

Example 250: 6-[4-(Diphenylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{{(3*S*)-3-
(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-*N*-phenyl-3,4-
dihydroisoquinoline-2(1*H*)-carboxamide

Example 251: 6-{4-[Butyl(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

5 **Example 252:** 6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 253: 6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-*N*-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

10 The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6bb, and replacing phenylacetyl chloride with benzyl isocyanate.

LC/MS (C₄₁H₄₁N₅O₄) 668 [M+H]⁺; RT 1.31 (Method B)

High-resolution mass (ESI+):

15 Empirical formula: C₄₁H₄₁N₅O₄

[M+H]⁺ calculated: 668.3231

[M+H]⁺ measured: 668.3211

20 **Example 254:** 6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 255: 6-{4-[(4-Hydroxyphenyl)(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 256: 6-[4-(Diphenylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 257: 6-{4-[Butyl(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 258: 6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 259: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(2-phenylethyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 260: Benzyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 261: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-{{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenoxyacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

5 **Example 262:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-(7-{{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-[2-oxo-2-(phenylamino)ethyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

Example 263: 5-(2-Benzoyl-7-{{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

10 **Example 264:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-(7-{{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-[(2*E*)-3-phenylprop-2-enoyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

15 **Example 265:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-{{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(3-phenylpropanoyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 266: 6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-*N*-methyl-7-{{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 267: 6-{4-[Butyl(4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-*N*-methyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 268: 6-{4-[(4-Hydroxyphenyl)(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-*N*-methyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 269: 6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-*N*-methyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 270: 6-{4-[Butyl(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-*N*-methyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 271: 6-[4-(Diphenylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-*N*-methyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 272: 6-{4-[Butyl(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-*N*-methyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 273: 6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-*N*-methyl-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-
phenyl-3,4-dihydroisoquinoline-2(1*H*)-carboxamide

5 **Example 274:** 6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-
yl}-*N*-methyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-
3,4-dihydroisoquinoline-2(1*H*)-carboxamide

Example 275: 6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-*N*-
methyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-
dihydroisoquinoline-2(1*H*)-carboxamide

10 **Example 276:** 6-{4-[(4-Hydroxyphenyl)(phenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-
yl}-*N*-methyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-
3,4-dihydroisoquinoline-2(1*H*)-carboxamide

15 **Example 277:** 6-[4-(Diphenylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-*N*-methyl-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-
dihydroisoquinoline-2(1*H*)-carboxamide

Example 278: 6-{4-[Butyl(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-*N*-methyl-
7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-
dihydroisoquinoline-2(1*H*)-carboxamide

Example 279: 6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-*N*-methyl-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-*N*-phenyl-3,4-
dihydroisoquinoline-2(1*H*)-carboxamide

Example 280: 5-[2-(Benzylsulfonyl)-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-
5 dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-
hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 281: 5-[2-(Benzylsulfonyl)-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-
dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-
trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 **Example 282:** 5-[2-(Benzylsulfonyl)-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-
dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-
hydroxyphenyl)-1,2-dimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 283: 5-[2-(Benzylsulfonyl)-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-
dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N,N*-
15 dibutyl-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 284: 5-[2-(Benzylsulfonyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-
yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-
trimethyl-1*H*-pyrrole-3-carboxamide

Example 285: 5-[2-(Benzylsulfonyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

5 **Example 286:** 5-[2-(Benzylsulfonyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-1,2-dimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 287: 5-[2-(Benzylsulfonyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N,N*-dibutyl-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

10 **Example 288:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylsulfamoyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

15 **Example 289:** *N*,1,2-Trimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylsulfamoyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 290: *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylsulfamoyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 291: *N,N*-Dibutyl-1,2-dimethyl-5-[7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylsulfamoyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

5 **Example 292:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-(2-[methyl(phenyl)sulfamoyl]-7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

Example 293: *N*,1,2-Trimethyl-5-(2-[methyl(phenyl)sulfamoyl]-7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 **Example 294:** *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-(2-[methyl(phenyl)sulfamoyl]-7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

15 **Example 295:** *N,N*-Dibutyl-1,2-dimethyl-5-(2-[methyl(phenyl)sulfamoyl]-7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

Example 296: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-(7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-[methyl(phenyl)sulfamoyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

Example 297: *N*,1,2-Trimethyl-5-(7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-[methyl(phenyl)sulfamoyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

5 **Example 298:** *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-(7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-[methyl(phenyl)sulfamoyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 299: *N,N*-Dibutyl-1,2-dimethyl-5-(7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-[methyl(phenyl)sulfamoyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

10 **Example 300:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

15 **Example 301:** *N*,1,2-Trimethyl-5-[7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

20 **Example 302:** *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-[7-{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 303: *N,N*-Dibutyl-1,2-dimethyl-5-[7-{{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 304: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-{{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 305: *N*,1,2-Trimethyl-5-[7-{{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 306: *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-[7-{{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 307: *N,N*-Dibutyl-1,2-dimethyl-5-[7-{{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 308: Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-{{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 309: Phenyl 6-{1-[methyl(phenyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-{{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 310:** Phenyl 6-{1-[(4-hydroxyphenyl)(phenyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-{{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 311: Phenyl 6-[1-(diphenylcarbamoyl)-5,6,7,8-tetrahydroindolizin-3-yl]-7-{{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 312:** Phenyl 6-{1-[butyl(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-{{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 313:** Phenyl 6-[1-(dibutylcarbamoyl)-5,6,7,8-tetrahydroindolizin-3-yl]-7-{{{(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 314: Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-{{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 315: Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]indolizin-3-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 316:** Phenyl 6-{1-[methyl(phenyl)carbamoyl]indolizin-3-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 317: Phenyl 6-{1-[(4-hydroxyphenyl)(phenyl)carbamoyl]indolizin-3-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 318:** Phenyl 6-[1-(diphenylcarbamoyl)indolizin-3-yl]-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-
2(1*H*)-carboxylate

15 **Example 319:** Phenyl 6-{1-[butyl(methyl)carbamoyl]indolizin-3-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 320: Phenyl 6-[1-(dibutylcarbamoyl)indolizin-3-yl]-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-
carboxylate

Example 321: Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]indolizin-3-yl}-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-
2(1*H*)-carboxylate

5 **Example 322:** Phenyl 6-{3-[(4-hydroxyphenyl)(methyl)carbamoyl]-1*H*-indol-1-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 323: Phenyl 6-{3-[methyl(phenyl)carbamoyl]-1*H*-indol-1-yl}-7-{[(3*S*)-3-
(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 324:** Phenyl 6-{3-[(4-hydroxyphenyl)(phenyl)carbamoyl]-1*H*-indol-1-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 325:** Phenyl 6-[3-(diphenylcarbamoyl)-1*H*-indol-1-yl]-7-{[(3*S*)-3-(morpholin-
4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-
2(1*H*)-carboxylate

Example 326: Phenyl 6-{3-[butyl(methyl)carbamoyl]-1*H*-indol-1-yl}-7-{[(3*S*)-3-
(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 327: Phenyl 6-[3-(dibutylcarbamoyl)-1*H*-indol-1-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 328: Phenyl 6-{3-[(4-hydroxyphenyl)(methyl)carbamoyl]-1*H*-indol-1-yl}-7-
5 {[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 329: Phenyl 6-{3-[(4-hydroxyphenyl)(methyl)carbamoyl]-1*H*-indazol-1-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 330:** Phenyl 6-{3-[methyl(phenyl)carbamoyl]-1*H*-indazol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 331: Phenyl 6-{3-[(4-hydroxyphenyl)(phenyl)carbamoyl]-1*H*-indazol-1-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
15 dihydroisoquinoline-2(1*H*)-carboxylate

Example 332: Phenyl 6-[3-(diphenylcarbamoyl)-1*H*-indazol-1-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 333: Phenyl 6-{3-[butyl(methyl)carbamoyl]-1*H*-indazol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 334:** Phenyl 6-[3-(dibutylcarbamoyl)-1*H*-indazol-1-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 335: Phenyl 6-{3-[(4-hydroxyphenyl)(methyl)carbamoyl]-1*H*-indazol-1-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 336: Phenyl 6-{4-chloro-3-[(4-hydroxyphenyl)(methyl)carbamoyl]-5-methyl-1*H*-pyrazol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 337:** Phenyl 6-{4-chloro-5-methyl-3-[methyl(phenyl)carbamoyl]-1*H*-pyrazol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 338:** Phenyl 6-{4-chloro-3-[(4-hydroxyphenyl)(phenyl)carbamoyl]-5-methyl-1*H*-pyrazol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 339: Phenyl 6-[4-chloro-3-(diphenylcarbamoyl)-5-methyl-1*H*-pyrazol-1-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 340:** Phenyl 6-{3-[butyl(methyl)carbamoyl]-4-chloro-5-methyl-1*H*-pyrazol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 341: Phenyl 6-[4-chloro-3-(dibutylcarbamoyl)-5-methyl-1*H*-pyrazol-1-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 342:** Phenyl 6-{4-chloro-3-[(4-hydroxyphenyl)(methyl)carbamoyl]-5-methyl-1*H*-pyrazol-1-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 343:** Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-2,3-dimethyl-1*H*-pyrrol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 344: Phenyl 6-{2,3-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 345: Phenyl 6-{4-[(4-hydroxyphenyl)(phenyl)carbamoyl]-2,3-dimethyl-1*H*-pyrrol-1-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 346: Phenyl 6-[4-(diphenylcarbamoyl)-2,3-dimethyl-1*H*-pyrrol-1-yl]-7-
5 {[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 347: Phenyl 6-{4-[butyl(methyl)carbamoyl]-2,3-dimethyl-1*H*-pyrrol-1-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 348:** Phenyl 6-[4-(dibutylcarbamoyl)-2,3-dimethyl-1*H*-pyrrol-1-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 349: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-2,3-dimethyl-1*H*-
pyrrol-1-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
15 dihydroisoquinoline-2(1*H*)-carboxylate

Example 350: Phenyl 6-{2-[(4-hydroxyphenyl)(methyl)carbamoyl]-5-methyl-1,3-thiazol-4-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 351: Phenyl 6-{5-methyl-2-[methyl(phenyl)carbamoyl]-1,3-thiazol-4-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 352:** Phenyl 6-{2-[(4-hydroxyphenyl)(phenyl)carbamoyl]-5-methyl-1,3-
thiazol-4-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-
yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 353: Phenyl 6-[2-(diphenylcarbamoyl)-5-methyl-1,3-thiazol-4-yl]-7-{[(3*S*)-3-
(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 354:** Phenyl 6-{2-[butyl(methyl)carbamoyl]-5-methyl-1,3-thiazol-4-yl}-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 355:** Phenyl 6-[2-(dibutylcarbamoyl)-5-methyl-1,3-thiazol-4-yl]-7-{[(3*S*)-3-
(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 356: Phenyl 6-{2-[(4-hydroxyphenyl)(methyl)carbamoyl]-5-methyl-1,3-
thiazol-4-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

Example 357: Phenyl 5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-6-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,3-dihydro-2*H*-isoindole-2-carboxylate

5 **Example 358:** Phenyl 7-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-8-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,3,4,5-tetrahydro-2*H*-2-benzazepine-2-carboxylate

Example 359: Phenyl 7-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-8-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,4,5-tetrahydro-3*H*-3-benzazepine-3-carboxylate

10 **Example 360:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1-oxo-2-(2-phenylethyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

15 **Example 361:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-[6-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1-oxo-2-(2-phenylethyl)-2,3-dihydro-1*H*-isoindol-5-yl]-1*H*-pyrrole-3-carboxamide

Example 362: Phenyl 1,1-difluoro-6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-5-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,3-dihydro-2*H*-isoindole-2-carboxylate

Example 363: Phenyl 1,1,3,3-tetrafluoro-5-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-6-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,3-dihydro-2*H*-isoindole-2-carboxylate

Example 364: 5-[3,3-Difluoro-6-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1-oxo-2-(2-phenylethyl)-2,3-dihydro-1*H*-isoindol-5-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 365: Phenyl 6-(4-{[2-(dimethylamino)ethyl](4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 366: Phenyl 6-(4-{(4-hydroxyphenyl)[2-(morpholin-4-yl)ethyl]carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 367: *N*-[2-(Dimethylamino)ethyl]-*N*-(4-hydroxyphenyl)-1,2-dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 368: *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-[2-(morpholin-4-yl)ethyl]-1*H*-pyrrole-3-carboxamide

Example 369: Phenyl 6-(4-{[2-(dimethylamino)ethyl](phenyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 370:** Phenyl 6-(1,5-dimethyl-4-{[2-(morpholin-4-yl)ethyl](phenyl)carbamoyl}-1*H*-pyrrol-2-yl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 371: *N*-[2-(Dimethylamino)ethyl]-1,2-dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 **Example 372:** 1,2-Dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-[2-(morpholin-4-yl)ethyl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

15 **Example 373:** Phenyl 6-(4-{butyl[2-(dimethylamino)ethyl]carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 374: Phenyl 6-(4-{butyl[2-(morpholin-4-yl)ethyl]carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 375: *N*-Butyl-*N*-[2-(dimethylamino)ethyl]-1,2-dimethyl-5-[7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

5 **Example 376:** *N*-Butyl-1,2-dimethyl-5-[7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-[2-(morpholin-4-yl)ethyl]-1*H*-pyrrole-3-carboxamide

Example 377: Phenyl 6-{1-[2-(dimethylamino)ethyl]-4-[(4-hydroxyphenyl)(methyl)carbamoyl]-5-methyl-1*H*-pyrrol-2-yl}-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 378:** Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-5-methyl-1-[2-(morpholin-4-yl)ethyl]-1*H*-pyrrol-2-yl}-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 379:** 1-[2-(Dimethylamino)ethyl]-*N*-(4-hydroxyphenyl)-*N*,2-dimethyl-5-[7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 380: *N*-(4-Hydroxyphenyl)-*N*,2-dimethyl-5-[7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1-[2-(morpholin-4-yl)ethyl]-1*H*-pyrrole-3-carboxamide

Example 381: Phenyl 6-{1-[2-(dimethylamino)ethyl]-5-methyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 382: Phenyl 7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-6-{5-methyl-4-[methyl(phenyl)carbamoyl]-1-[2-(morpholin-4-yl)ethyl]-1*H*-pyrrol-2-yl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 383: 1-[2-(Dimethylamino)ethyl]-*N*,2-dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 384: *N*,2-Dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1-[2-(morpholin-4-yl)ethyl]-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 385: Phenyl 6-{4-(dibutylcarbamoyl)-1-[2-(dimethylamino)ethyl]-5-methyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 386: Phenyl 6-{4-(dibutylcarbamoyl)-5-methyl-1-[2-(morpholin-4-yl)ethyl]-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 387: *N,N*-Dibutyl-1-[2-(dimethylamino)ethyl]-2-methyl-5-[7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrole-3-carboxamide

Example 388: *N,N*-Dibutyl-2-methyl-5-[7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1-[2-(morpholin-4-yl)ethyl]-1*H*-pyrrole-3-carboxamide

Example 389: 2-(4-Methylpiperazin-1-yl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 390: 2-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 391: 2-{[2-(Dimethylamino)ethyl]amino}phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 392: 2-[2-(Dimethylamino)ethoxy]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 709, replacing the product from Preparation 6ba in Step A with the product from Preparation 6bb, and replacing benzyl 3-hydroxybenzoate from Step B with 3-[2-(dimethylamino)ethoxy]phenol.

LC/MS ($C_{45}H_{49}N_5O_6$) 754 $[M-H]^-$; RT 2.27 (Method A)

5 **High-resolution mass (ESI+):**

Empirical formula: $C_{45}H_{49}N_5O_6$

$[M+2H]^{2+}$ calculated: 378.6914

$[M+2H]^{2+}$ measured: 378.6926

10 **Example 393:** 2-[2-(Dimethylamino)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 394:** 2-[(Dimethylamino)methyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 395: 2-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

20 **Example 396:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-(7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-[[2-(4-methylpiperazin-1-yl)phenyl]acetyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

Example 397: 5-(2-[(2-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl)acetyl]-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-
tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-
carboxamide

5 **Example 398:** 5-(2-[(2-{[2-(Dimethylamino)ethyl]amino}phenyl)acetyl]-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-
yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

10 **Example 399:** 5-[2-({2-[2-(Dimethylamino)ethoxy]phenyl}acetyl)-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-
hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 400: 5-[2-({2-[2-(Dimethylamino)ethyl]phenyl}acetyl)-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-
hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

15 **Example 401:** 5-[2-({2-[(Dimethylamino)methyl]phenyl}acetyl)-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-
hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 402: 5-[2-({2-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-
yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 403: 2-(4-Methylpiperazin-1-yl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 404:** 2-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 405: 2-{[2-(Dimethylamino)ethyl]amino}phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 406:** 2-[2-(Dimethylamino)ethoxy]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 407:** 2-[2-(Dimethylamino)ethyl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 408: 2-[(Dimethylamino)methyl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 409: 2-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 410:** *N*,1,2-Trimethyl-5-(7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-{[2-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 411: 5-(2-[2-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl]acetyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 **Example 412:** 5-(2-[2-{[2-(Dimethylamino)ethyl]amino}phenyl]acetyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

15 **Example 413:** 5-[2-({2-[2-(Dimethylamino)ethoxy]phenyl}acetyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 414: 5-[2-({2-[2-(Dimethylamino)ethyl]phenyl}acetyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 415: 5-[2-({2-[(Dimethylamino)methyl]phenyl}acetyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

5 **Example 416:** 5-[2-({2-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 417: 2-(4-Methylpiperazin-1-yl)phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 418:** 2-{[2-(Dimethylamino)ethyl](methylamino)}phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 419:** 2-{[2-(Dimethylamino)ethyl]amino}phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 420: 2-[2-(Dimethylamino)ethoxy]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 421: 2-[2-(Dimethylamino)ethyl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 422:** 2-[(Dimethylamino)methyl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 423: 2-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 424:** *N,N*-Dibutyl-1,2-dimethyl-5-(7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-{[2-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

15 **Example 425:** *N,N*-Dibutyl-5-(2-[(2-{[2-(dimethylamino)ethyl](methyl)amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 426: *N,N*-Dibutyl-5-(2-[(2-{[2-(dimethylamino)ethyl]amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 427: *N,N*-Dibutyl-5-[2-({2-[2-(dimethylamino)ethoxy]phenyl}acetyl)-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-
tetrahydroisoquinolin-6-yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

5 **Example 428:** *N,N*-Dibutyl-5-[2-({2-[2-(dimethylamino)ethyl]phenyl}acetyl)-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-
yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 429: *N,N*-Dibutyl-5-[2-({2-[(dimethylamino)methyl]phenyl}acetyl)-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-
yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

10 **Example 430:** *N,N*-Dibutyl-5-[2-({2-[3-(dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-
7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-
tetrahydroisoquinolin-6-yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

15 **Example 431:** 3-(4-Methylpiperazin-1-yl)phenyl 6-{4-[(4-
hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-
carboxylate

20 **Example 432:** 3-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl 6-{4-[(4-
hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-
carboxylate

Example 433: 3-{{2-(Dimethylamino)ethyl}amino}phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 434:** 3-{{2-(Dimethylamino)ethoxy}phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

10 The procedure is as in Example 709, replacing the product from Preparation 6ba in Step A with product from Preparation 6bb, and replacing benzyl 3-hydroxybenzoate from Step B with 3-[2-(dimethylamino)ethoxy]phenol.

LC/MS (C₄₅H₄₉N₅O₆) 754 [M-H]⁻; RT 2.24 (Method A)

High-resolution mass (ESI⁺):

Empirical formula: C₄₅H₄₉N₅O₆

15 [M+2H]²⁺ calculated: 378.6914

[M+2H]²⁺ measured: 378.6918

Example 435: 3-[2-(Dimethylamino)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

20 **Example 436:** 3-[(Dimethylamino)methyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 437: 3-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 438:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-(7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-{[3-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

Example 439: 5-(2-[(3-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl)acetyl]-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

10

Example 440: 5-(2-[(3-{[2-(Dimethylamino)ethyl]amino}phenyl)acetyl]-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

15 **Example 441:** 5-[2-[(3-[2-(Dimethylamino)ethoxy]phenyl)acetyl]-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 442: 5-[2-[(3-[2-(Dimethylamino)ethyl]phenyl)acetyl]-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

20

Example 443: 5-[2-({3-[(Dimethylamino)methyl]phenyl}acetyl)-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

5 **Example 444:** 5-[2-({3-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 445: 3-(4-Methylpiperazin-1-yl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 446:** 3-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 786, replacing 4-hydroxybenzonitrile in Step B with 3-{[2-(dimethylamino)ethyl](methyl)amino}phenol.

15 **LC/MS** (C₄₆H₅₂N₆O₄) no ionisation; RT 2.44 (Method A)

High-resolution mass (ESI⁺):

Empirical formula: C₄₆H₅₂N₆O₄

[M+2H]²⁺ calculated: 377.2098

[M+2H]²⁺ measured: 377.2100

Example 447: 3-[[2-(Dimethylamino)ethyl]amino]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 448: 3-[2-(Dimethylamino)ethoxy]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 786, replacing 4-hydroxybenzonitrile in Step B with 3-[2-(dimethylamino)ethoxy]phenol.

LC/MS (C₄₅H₄₉N₅O₅) 740 [M+H]⁺; RT 1.20 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₅H₄₉N₅O₅

[M+2H]²⁺ calculated: 370.6940

[M+2H]²⁺ measured: 370.6954

Example 449: 3-[2-(Dimethylamino)ethyl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 450: 3-[(Dimethylamino)methyl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 451: 3-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 452:** *N*,1,2-Trimethyl-5-(7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-{[3-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 453: 5-(2-[(3-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 **Example 454:** 5-(2-[(3-{[2-(Dimethylamino)ethyl]amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

15 **Example 455:** 5-[2-({3-[2-(Dimethylamino)ethoxy]phenyl}acetyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 456: 5-[2-({3-[2-(Dimethylamino)ethyl]phenyl}acetyl)-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 457: 5-[2-({3-[(Dimethylamino)methyl]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

5 **Example 458:** 5-[2-({3-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 459: 3-(4-Methylpiperazin-1-yl)phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 460:** 3-{{2-(Dimethylamino)ethyl}(methylamino)phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 461:** 3-{{2-(Dimethylamino)ethyl}amino}phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 462: 3-[2-(Dimethylamino)ethoxy]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 463: 3-[2-(Dimethylamino)ethyl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 464: 3-[(Dimethylamino)methyl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 465: 3-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 466: *N,N*-Dibutyl-1,2-dimethyl-5-(7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-{[3-(4-methylpiperazin-1-yl)phenyl]acetyl})-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

Example 467: *N,N*-Dibutyl-5-(2-[(3-{[2-(dimethylamino)ethyl](methyl)amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl})-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 468: *N,N*-Dibutyl-5-(2-[(3-{[2-(dimethylamino)ethyl]amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl})-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 469: *N,N*-Dibutyl-5-[2-({3-[2-(dimethylamino)ethoxy]phenyl}acetyl)-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-
tetrahydroisoquinolin-6-yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 470: *N,N*-Dibutyl-5-[2-({3-[2-(dimethylamino)ethyl]phenyl}acetyl)-7-
5 {[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-
yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 471: *N,N*-Dibutyl-5-[2-({3-[(dimethylamino)methyl]phenyl}acetyl)-7-
3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-
yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

10 **Example 472:** *N,N*-Dibutyl-5-[2-({3-[3-(dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-
7-
7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-
tetrahydroisoquinolin-6-yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 473: 4-(4-Methylpiperazin-1-yl)phenyl 6-{4-[(4-
hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
15 {[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-
carboxylate

Example 474: 4-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl 6-{4-[(4-
hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-
1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-
20 carboxylate

The procedure is as in Example 709, replacing the product from Preparation 6ba in Step A with the product from Preparation 6bb, and replacing benzyl 3-hydroxybenzoate from Step B with 4-[[2-(dimethylamino)ethyl](methyl)amino]phenol.

LC/MS (C₄₆H₅₂N₆O₅) 767 [M-H]⁻; RT 2.26 (Method A)

5 **Example 475:** 4-[[2-(Dimethylamino)ethyl]amino]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 476:** 4-[2-(Dimethylamino)ethoxy]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

15 The procedure is as in Example 709, replacing the product from Preparation 6ba in Step A with the product from Preparation 6bb, and replacing benzyl 3-hydroxybenzoate from Step B with 4-[2-(dimethylamino)ethoxy]phenol.

LC/MS (C₄₅H₄₉N₅O₆) 754 [M-H]⁻; RT 2.21 (Method A)

High-resolution mass (ESI⁺):

Empirical formula: C₄₅H₄₉N₅O₆

[M+2H]²⁺ calculated: 378.6914

20 [M+2H]²⁺ measured: 378.6926

Example 477: 4-[2-(Dimethylamino)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 478: 4-[(Dimethylamino)methyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 479:** 4-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 480:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-(7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-{[4-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

15 **Example 481:** 5-(2-[(4-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 482: 5-(2-[(4-{[2-(Dimethylamino)ethyl]amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 483: 5-[2-({4-[2-(Dimethylamino)ethoxy]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

5 **Example 484:** 5-[2-({4-[2-(Dimethylamino)ethyl]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 485: 5-[2-({4-[(Dimethylamino)methyl]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

10 **Example 486:** 5-[2-({4-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

15 **Example 487:** 4-(4-Methylpiperazin-1-yl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 488: 4-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

20 **Step A:** 6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl chloride

The procedure is as in Step A of Example 573, replacing the product from Preparation 6ba with the product from Preparation 6cb.

LC/MS ($C_{35}H_{35}ClN_4O_3$) 595 $[M+H]^+$; RT 2.73 (Method A)

Step B: 4- $\{2-(\text{Dimethylamino})\text{ethyl}\}(\text{methyl})\text{amino}\}$ phenyl 6- $\{1,5\text{-dimethyl-4-}$
5 $[\text{methyl}(\text{phenyl})\text{carbamoyl}]\text{-1H-pyrrol-2-yl}\}$ -7- $\{(\text{3R})\text{-3-methyl-1,2,3,4-}$
 $\text{tetrahydroisoquinoline-2-carbonyl}\}\text{-1,2,3,4-tetrahydroisoquinoline-2-carboxylate}$

To a solution of the carbamoyl chloride obtained from Step A (55 mg, 0.09 mmol) in acetonitrile (5 mL) was added potassium carbonate (128 mg, 0.92 mmol) and 4-
(dimethylamino)ethyl(methylamino)phenol (22 mg, 0.11 mmol), and the reaction was
10 stirred at 60 °C for *ca* 16 h. The reaction was allowed to cool to ambient temperature, diluted with dichloromethane and washed with water. The organic extract was dried over magnesium sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (silica; dichloromethane to 10% methanol/dichloromethane) afforded the desired product as a cream solid.

15 LC/MS ($C_{46}H_{52}N_6O_4$) 753 $[M+H]^+$; RT 2.42 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{52}N_6O_4$

$[M+2H]^{2+}$ calculated: 377.2098

$[M+2H]^{2+}$ measured: 377.2102

20 **Example 489:** 4- $\{2-(\text{Dimethylamino})\text{ethyl}\}\text{amino}\}$ phenyl 6- $\{1,5\text{-dimethyl-4-}$
 $[\text{methyl}(\text{phenyl})\text{carbamoyl}]\text{-1H-pyrrol-2-yl}\}$ -7- $\{(\text{3R})\text{-3-methyl-1,2,3,4-}$
 $\text{tetrahydroisoquinoline-2-carbonyl}\}\text{-1,2,3,4-tetrahydroisoquinoline-2-carboxylate}$

Step A: 6- $\{1,5\text{-Dimethyl-4-}[\text{methyl}(\text{phenyl})\text{carbamoyl}]\text{-1H-pyrrol-2-yl}\}$ -7- $\{(\text{3R})\text{-3-methyl-}$
 $1,2,3,4\text{-tetrahydroisoquinoline-2-carbonyl}\}\text{-1,2,3,4-tetrahydroisoquinoline-2-carbonyl}$
25 chloride

The procedure is as in Step A of Example 573, replacing the product from Preparation 6ba with the product from Preparation 6cb.

LC/MS ($C_{35}H_{35}ClN_4O_3$) 595 $[M+H]^+$; RT 2.73 (Method A)

Step B: 4- $\{[(tert\text{-}Butoxy)carbonyl][2\text{-}(\text{dimethylamino})ethyl]amino\}$ phenyl 6- $\{1,5\text{-}$
dimethyl-4- $[methyl(phenyl)carbamoyl]\text{-}1H\text{-}pyrrol\text{-}2\text{-}yl\}\text{-}7\text{-}[(3R)\text{-}3\text{-}methyl\text{-}1,2,3,4\text{-}$
tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Step B of Example 573, replacing the product of Step A in Example 573 with the product of Step A in Example 489, and replacing 3,4-dichlorophenol with *tert*-butyl *N*-[2-(dimethylamino)ethyl]-*N*-(4-hydroxyphenyl)carbamate.

LC/MS ($C_{50}H_{58}N_6O_6$) no ionisation; RT 2.54 (Method A)

Step C: 4- $\{2\text{-}(\text{Dimethylamino})ethyl\}amino\}$ phenyl 6- $\{1,5\text{-}dimethyl\text{-}4\text{-}$
 $[methyl(phenyl)carbamoyl]\text{-}1H\text{-}pyrrol\text{-}2\text{-}yl\}\text{-}7\text{-}[(3R)\text{-}3\text{-}methyl\text{-}1,2,3,4\text{-}$
tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of product obtained in Step B (31 mg, 0.04 mmol, 1 eq) in dichloromethane (5 mL) was added trifluoroacetic acid (0.5 mL) and the mixture was stirred at ambient temperature for *ca* 4 h. The reaction was diluted with water and basified with aqueous 2M sodium hydroxide. The phases were separated and the aqueous phase was extracted with dichloromethane. The combined organic phases were dried over magnesium sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (10 g silica; dichloromethane to 10% methanol in dichloromethane) afforded the desired product as a white powder.

LC/MS ($C_{45}H_{50}N_6O_4$) no ionisation; RT 2.38 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{45}H_{50}N_6O_4$

$[M+2H]^{2+}$ calculated: 370.2020

[M+2H]²⁺ measured: 370.2038

Example 490: 4-[2-(Dimethylamino)ethoxy]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 491:** 4-[2-(Dimethylamino)ethyl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 492:** 4-[(Dimethylamino)methyl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 493: 4-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 494:** *N*,1,2-Trimethyl-5-(7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-{[4-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 495: 5-(2-[(4-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl)acetyl]-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 496: 5-(2-[(4-{[2-(Dimethylamino)ethyl]amino}phenyl)acetyl]-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

5 **Example 497:** 5-[2-({4-[2-(Dimethylamino)ethoxy]phenyl}acetyl)-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 498: 5-[2-({4-[2-(Dimethylamino)ethyl]phenyl}acetyl)-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 **Example 499:** 5-[2-({4-[(Dimethylamino)methyl]phenyl}acetyl)-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

15 **Example 500:** 5-[2-({4-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 501: 4-(4-Methylpiperazin-1-yl)phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 502: 4-{{2-(Dimethylamino)ethyl}(methyl)amino}phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 503:** 4-{{2-(Dimethylamino)ethyl}amino}phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 504: 4-[2-(Dimethylamino)ethoxy]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 505:** 4-[2-(Dimethylamino)ethyl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 506:** 4-[(Dimethylamino)methyl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 507: 4-[3-(Dimethylamino)prop-1-yn-1-yl]phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 508: *N,N*-Dibutyl-1,2-dimethyl-5-(7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-2-{{4-(4-methylpiperazin-1-yl)phenyl}acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1*H*-pyrrole-3-carboxamide

5 **Example 509:** *N,N*-Dibutyl-5-(2-[(4-{{2-(dimethylamino)ethyl}(methyl)amino}phenyl)acetyl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 510: *N,N*-Dibutyl-5-(2-[(4-{{2-(dimethylamino)ethyl}amino}phenyl)acetyl]-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

10 **Example 511:** *N,N*-Dibutyl-5-[2-({4-[2-(dimethylamino)ethoxy]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

15 **Example 512:** *N,N*-Dibutyl-5-[2-({4-[2-(dimethylamino)ethyl]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 513: *N,N*-Dibutyl-5-[2-({4-[(dimethylamino)methyl]phenyl}acetyl)-7-{{(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl}carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 514: *N,N*-Dibutyl-5-[2-({4-[3-(dimethylamino)prop-1-yn-1-yl]phenyl}acetyl)-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 515: Disodium 4-[(1,2-Dimethyl-5-[7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-(phenoxycarbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrol-3-yl]carbonyl(methylamino)phenyl phosphate

Example 516: Disodium 4-[(1,2-dimethyl-5-[7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1*H*-pyrrol-3-yl]carbonyl(methylamino)phenyl phosphate

Example 517: Disodium 4-[methyl({3-[7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-(phenoxycarbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-5,6,7,8-tetrahydroindolizin-1-yl}carbonyl)amino)phenyl phosphate

Example 518: Disodium 4-[methyl({3-[7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-5,6,7,8-tetrahydroindolizin-1-yl}carbonyl)amino)phenyl phosphate

Example 519: Disodium 4-[(1,2-dimethyl-5-[7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-2-(phenoxycarbonyl)-1,2,3,4-

tetrahydroisoquinolin-6-yl]-1*H*-pyrrol-3-yl}carbonyl)(methyl)amino]phenyl
phosphate

Example 520: Disodium 4-[(1,2-dimethyl-5-[7-{[(3*R*)-3-methyl-3,4-
dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-
5 tetrahydroisoquinolin-6-yl]-1*H*-pyrrol-3-yl}carbonyl)(methyl)amino]phenyl
phosphate

Example 521: Disodium 4-[methyl({3-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-
2(1*H*)-yl]carbonyl}-2-(phenoxycarbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-5,6,7,8-
tetrahydroindolizin-1-yl}carbonyl)amino]phenyl phosphate

10 **Example 522:** Disodium 4-[methyl({3-[7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-
2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-5,6,7,8-
tetrahydroindolizin-1-yl}carbonyl)amino]phenyl phosphate

Example 523: Phenyl 6-{1-ethyl-4-[(4-hydroxyphenyl)(methyl)carbamoyl]-5-methyl-
1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-
15 yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 524: Phenyl 6-{1-(2-hydroxyethyl)-4-[(4-hydroxyphenyl)(methyl)carbamoyl]-
5-methyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-
2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 525: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1-(2-methoxyethyl)-5-methyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 526: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-5-methyl-1-(2,2,2-trifluoroethyl)-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 527: Phenyl 6-{1-ethyl-5-methyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 528: Phenyl 6-{1-(2-hydroxyethyl)-5-methyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 529: Phenyl 6-{1-(2-methoxyethyl)-5-methyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 530: Phenyl 6-{5-methyl-4-[methyl(phenyl)carbamoyl]-1-(2,2,2-trifluoroethyl)-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 531: Phenyl 6-[4-(dibutylcarbamoyl)-1-ethyl-5-methyl-1*H*-pyrrol-2-yl]-7-
{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-
dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 532:** Phenyl 6-[4-(dibutylcarbamoyl)-1-(2-hydroxyethyl)-5-methyl-1*H*-
pyrrol-2-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-
yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 533: Phenyl 6-[4-(dibutylcarbamoyl)-1-(2-methoxyethyl)-5-methyl-1*H*-
pyrrol-2-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-
yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 534:** Phenyl 6-[4-(dibutylcarbamoyl)-5-methyl-1-(2,2,2-trifluoroethyl)-1*H*-
pyrrol-2-yl]-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-
yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 535:** 2-Methylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-
dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-
2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 536: 3-Methylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-
dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-
2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 537: 4-Methylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

5 **Example 538:** 2-Chlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 539: 3-Chlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

10 **Example 540:** 4-Chlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

15 **Example 541:** 2-Hydroxyphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 542: 3-Hydroxyphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

20 **Example 543:** 4-Hydroxyphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 544: 2-Methoxyphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 545:** 3-Methoxyphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 546: 4-Methoxyphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 547:** 2-(Methylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 548:** 3-(Methylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 549: 4-(Methylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 550: 2-(Dimethylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 551:** 3-(Dimethylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 552: 4-(Dimethylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 553:** 2-(Acetylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 554:** 3-(Acetylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 555: 4-(Acetylamino)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 556: 2-(Trifluoromethyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 557: 3-(Trifluoromethyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 558: 4-(Trifluoromethyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 559: 2-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 560: 3-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 573, replacing 3,4-dichlorophenol in Step B with 3-hydroxybenzonitrile.

LC/MS (C₄₆H₄₆N₆O₆) 779 [M+H]⁺; RT 1.14 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₆H₄₆N₆O₆

[M+H]⁺ calculated: 779.3552

[M+H]⁺ measured: 779.3515

Example 561: 4-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 562: 2-Ethynylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 563: 3-Ethynylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 564: 4-Ethynylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of the product obtained from Example 784 (53 mg, 0.06 mmol, 1 eq) and triethylamine (10 μ L, 0.1 mmol) in THF (1 mL) were added bis(triphenylphosphine)palladium(II) dichloride (5 mg, 0.01 mmol, 0.12 eq), copper(I) iodide (0.7 mg, 3.84 μ mol), and trimethylsilylacetylene (36 μ L, 0.26 mmol, 4 eq) and the mixture was heated at 60°C for *ca* 16 h. The reaction was allowed to cool to ambient temperature, then evaporated and purified preparative HPLC (H₂O-TFA/acetonitrile; gradient elution) to afford the desired product.

LC/MS (C₄₇H₄₇N₅O₆) 778 [M+H]⁺; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₇N₅O₆

$[M+H]^+$ calculated: 778.3599

$[M+H]^+$ measured: 778.3635

Example 565: 2,3-Dichlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 566: Naphthalen-1-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 567: 1*H*-Indol-7-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 568: 1-Methyl-1*H*-indol-7-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 569: 1*H*-Indol-4-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 570: 1-Methyl-1*H*-indol-4-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 571: 1*H*-Pyrrolo[2,3-*b*]pyridin-4-yl 6-{4-[(4-hydroxyphenyl)(methyl) carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 572: 1-Methyl-1*H*-pyrrolo[2,3-*b*]pyridin-4-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 573: 3,4-Dichlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: 6-(4-{[4-(Benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl chloride

To a solution of the compound from Preparation 6ba (450 mg, 0.47 mmol) and *N,N*-diisopropylethylamine (0.41 mL, 2.36 mmol) in dichloromethane (45 mL), cooled to 0 °C, was added triphosgene (139 mg, 0.47 mmol) and the mixture was stirred for 1 h at ambient temperature. The reaction was diluted with dichloromethane and was washed with 1M aqueous HCl. The organic extract was dried over magnesium sulfate, filtered and concentrated *in vacuo* to afford a yellow solid that was used directly in the next step without further purification, and assuming quantitative transformation.

LC/MS (C₄₆H₄₈ClN₅O₅) 786 [M+H]⁺; RT 1.33 (Method B)

Step B: 3,4-Dichlorophenyl 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of the compound from Step A (41 mg, 0.05 mmol) in acetonitrile (5 mL) was added potassium carbonate (72 mg, 0.52 mmol), DMAP (0.052 mmol) and 3,4-dichlorophenol (0.52 mmol) and the mixture heated at 60 °C for *ca* 16 h. The reaction mixture was diluted with ethyl acetate and was washed with saturated aqueous sodium bicarbonate, and brine. The organic extract was dried over magnesium sulfate and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; dichloromethane to 5 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₅₂H₅₁Cl₂N₅O₆) 912 [M+H]⁺; RT 1.48 (Method B)

Step C: 3,4-Dichlorophenyl 6-{4-[4-hydroxyphenyl](methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of the compound from Step A in dichloromethane (5 mL), cooled to 0 °C, was added drop-wise boron trichloride (1 M in dichloromethane; 5 eq.). The reaction was then allowed to warm to ambient temperature and continue stirring for *ca* 16 h. The reaction was quenched by the addition of water (10 mL) and the phases were separated. The organic phase was washed successively with aqueous sodium bicarbonate, and brine. The organic extract was dried over magnesium sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; dichloromethane to 5 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₄₅H₄₅Cl₂N₅O₆) 822 [M+H]⁺; RT 1.28 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₄₅N₅O₆Cl₂

[M+H]⁺ calculated: 822.2820

[M+H]⁺ measured: 822.2832

Example 574: Naphthalen-2-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is an in Example 573, replacing 3,4-dichlorophenol in Step B with naphthalen-2-ol.

LC/MS (C₄₉H₄₉N₅O₆) 804 [M+H]⁺; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₉H₄₉N₅O₆

[M+H]⁺ calculated: 804.3756

[M+H]⁺ measured: 804.3767

Example 575: 1*H*-Indol-6-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is an in Example 573, replacing 3,4-dichlorophenol in Step B with 3*H*-indol-6-ol.

LC/MS (C₄₇H₄₈N₆O₆) 793 [M+H]⁺; RT 1.15 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₈N₆O₆

[M+H]⁺ calculated: 793.3708

[M+H]⁺ measured: 793.3690

Example 576: 1-Methyl-1*H*-indol-6-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 577: 1*H*-Indol-5-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is an in Example 573, replacing 3,4-dichlorophenol in Step B with 1*H*-indol-5-ol.

LC/MS (C₄₇H₄₈N₆O₆) 793 [M+H]⁺; RT 1.13 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₈N₆O₆

[M+2H]²⁺ calculated: 397.1890

[M+2H]²⁺ measured: 397.1879

Example 578: 1-Methyl-1*H*-indol-5-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 579: 1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is an in Example 573, replacing 3,4-dichlorophenol in Step B with 1*H*-pyrrolo[2,3-*b*]pyridin-5-ol.

LC/MS (C₄₆H₄₇N₇O₆) 794 [M+H]⁺; RT 1.07 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₇N₇O₆

[M+2H]²⁺ calculated: 397.6867

$[M+2H]^{2+}$ measured: 397.6880

Example 580: 1-Methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 581: 1*H*-Pyrrolo[2,3-*b*]pyridin-4-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 582: 1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 583: 2-(Methylsulfanyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 584: 3-(Methylsulfanyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 585: 4-(Methylsulfanyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 586: 2-(Dimethylcarbamoyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

5 **Example 587:** 3-(Dimethylcarbamoyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

10 **Example 588:** 4-(Dimethylcarbamoyl)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 589: 2-Methylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

15 **Example 590:** 3-Methylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 591: 4-Methylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 785, replacing the product from Preparation 6cb with the product from Preparation 6ca.

LC/MS (C₄₆H₄₉N₅O₅) 752 [M+H]⁺; RT 1.34 (Method B)

High-resolution mass (ESI+):

5 Empirical formula: C₄₆H₄₉N₅O₅

[M+H]⁺ calculated: 752.3806

[M+H]⁺ measured: 752.3836

10 **Example 592:** 2-Chlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 593: 3-Chlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

15 **Example 594:** 4-Chlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 595: 2-Hydroxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 596: 3-Hydroxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 597:** 4-Hydroxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 598: 2-Methoxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 599:** 3-Methoxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 600:** 4-Methoxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 601: 2-(Methylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 602: 3-(Methylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

5 **Example 603:** 4-(Methylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 604: 2-(Dimethylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

10 **Example 605:** 3-(Dimethylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 606: 4-(Dimethylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

15

Example 607: 2-(Acetylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 608: 3-(Acetylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 609: 4-(Acetylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 610: 2-(Trifluoromethyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 611: 3-(Trifluoromethyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 612: 4-(Trifluoromethyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 613: 2-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 614: 3-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

5 **Example 615:** 4-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 616: 2-Ethynylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

10 **Example 617:** 3-Ethynylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

15 **Example 618:** 4-Ethynylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 619: 2,3-Dichlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 620: Naphthalen-1-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 621: 1*H*-Indol-7-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 622: 1-Methyl-1*H*-indol-7-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 623: 1*H*-Indol-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 624: 1-Methyl-1*H*-indol-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 625: 1*H*-Pyrrolo[2,3-*b*]pyridin-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 626: 1-Methyl-1*H*-pyrrolo[2,3-*b*]pyridin-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 627: 3,4-Dichlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 628: Naphthalen-2-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 629: 1*H*-Indol-6-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: 6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl chloride and

Trichloromethyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of product from Preparation 6ca (200 mg, 0.32 mmol) and *N,N*-diisopropylethylamine (0.17 mL, 0.97 mmol) in dichloromethane (10 mL), cooled to 0 °C, was added triphosgene (96 mg, 0.32 mmol) and the mixture was stirred at ambient temperature 1 h. The reaction was diluted with dichloromethane and washed with 1M aqueous HCl. The organic extract was dried with magnesium sulfate, filtered and concentrated *in vacuo* to afford the desired products as a mixture.

LC/MS (C₃₉H₄₂ClN₅O₄) 680 [M+H]⁺; RT 1.27, and (C₄₀H₄₂Cl₃N₅O₅) 778 [M+H]⁺; RT 1.36 (Method B)

Step B: *1H-Indol-6-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate*

To a solution of the material from Step A (50 mg, 0.06 mmol) in acetonitrile (5 mL) was added potassium carbonate (44 mg, 0.32 mmol) followed by 1*H*-indol-6-ol (1.5 eq) and the mixture was heated at 60 °C for *ca* 16 h. The reaction was allowed to cool to ambient temperature, and was diluted with ethyl acetate (20 mL) and washed successively with 1M aqueous sodium hydroxide, and brine. The organic extract was dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; dichloromethane to 10 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₄₇H₄₈N₆O₅) 777 [M+H]⁺; RT 1.26 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₈N₆O₅

[M+H]⁺ calculated: 777.3759

[M+H]⁺ measured: 777.3795

Example 630: *1-Methyl-1H-indol-6-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1H)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1H)-carboxylate*

Example 631: *1H-Indol-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate*

The procedure is as in Example 629, replacing 1*H*-indol-6-ol in Step B with 1*H*-indol-5-ol, and purifying *via* preparative HPLC (H₂O-TFA/acetonitrile; gradient elution).

LC/MS (C₄₇H₄₈N₆O₅) 777 [M+H]⁺; RT 1.28 (Method B)

High-resolution mass (ESI+):

5 Empirical formula: C₄₇H₄₈N₆O₅

[M+H]⁺ calculated: 777.3759

[M+H]⁺ measured: 777.3742

10 **Example 632:** 1-Methyl-1*H*-indol-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 633: 1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

15 The procedure is as in Example 629, replacing 1*H*-indol-6-ol in Step B with 1*H*-pyrrolo[2,3-*b*]pyridin-5-ol.

LC/MS (C₄₆H₄₇N₇O₅) 778 [M+H]⁺; RT 1.18 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₇N₇O₅

[M+2H]²⁺ calculated: 389.6892

20 [M+2H]²⁺ measured: 389.6874

Example 634: 1-Methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 629, replacing 1*H*-indol-6-ol in Step B with 1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-ol, and purifying *via* preparative HPLC (H₂O-TFA/acetonitrile; gradient elution).

LC/MS (C₄₇H₄₉N₇O₅) 792 [M+H]⁺; RT 1.25 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₄₉N₇O₅

[M+2H]²⁺ calculated: 396.6970

[M+2H]²⁺ measured: 396.6978

Example 635: 1*H*-Pyrrolo[2,3-*b*]pyridin-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 636: 1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 786, replacing 4-hydroxybenzonitrile in Step B with 1*H*-pyrrolo[2,3-*b*]pyridin-5-ol.

LC/MS (C₄₂H₄₀N₆O₄) 693 [M+H]⁺; RT 1.34 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₂H₄₀N₆O₄

[M+2H]²⁺ calculated: 347.1628

[M+2H]²⁺ measured: 347.1639

Example 637: 2-(Methylsulfanyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 638:** 3-(Methylsulfanyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 639:** 4-(Methylsulfanyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 640: 2-(Dimethylcarbamoyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 641:** 3-(Dimethylcarbamoyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 642: 4-(Dimethylcarbamoyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 643: 2-Methylphenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 644:** 3-Methylphenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 645: 4-Methylphenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 646:** 2-Chlorophenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 647:** 3-Chlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 648: 4-Chlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 649: 2-Hydroxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 650:** 3-Hydroxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 651: 4-Hydroxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 652:** 2-methoxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 653:** 3-Methoxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 654: 4-Methoxyphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 655: 2-(Methylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 656:** 3-(Methylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 657: 4-(Methylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 658:** 2-(Dimethylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 659:** 3-(Dimethylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 660: 4-(Dimethylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 661: 2-(Acetylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 662:** 3-(Acetylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 663: 4-(Acetylamino)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 664:** 2-(Trifluoromethyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 665:** 3-(Trifluoromethyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 666: 4-(Trifluoromethyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 667: 2-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 668:** 3-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 669: 4-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 670:** 2-Ethynylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 671:** 3-Ethynylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 672: 4-Ethynylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 673: 2,3-Dichlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

5 **Example 674:** Naphthalen-1-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 675: 1*H*-Indol-7-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

10 **Example 676:** 1-Methyl-1*H*-indol-7-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

15 **Example 677:** 1*H*-Indol-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 678: 1-Methyl-1*H*-indol-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 679: 1*H*-Pyrrolo[2,3-*b*]pyridin-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 680: 1-Methyl-1*H*-pyrrolo[2,3-*b*]pyridin-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 681: 3,4-Dichlorophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 682: Naphthalen-2-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 683: 1*H*-Indol-6-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 684: 1-Methyl-1*H*-indol-6-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 685: 1*H*-Indol-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 686: 1-Methyl-1*H*-indol-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 687: 1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 688: 1-Methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 689: 1*H*-Pyrrolo[2,3-*b*]pyridin-4-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 690: 1*H*-Pyrrolo[2,3-*b*]pyridin-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 691: 2-(Methylsulfanyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl) carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 692:** 3-(Methylsulfanyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl) carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 693: 4-(Methylsulfanyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl) carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 694:** 2-(Dimethylcarbamoyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl) carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 695:** 3-(Dimethylcarbamoyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl) carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 696: 4-(Dimethylcarbamoyl)phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 697: Phenyl 7-{[(3*S*)-3-(hydroxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 698:** 5-[7-{[(3*S*)-3-(Hydroxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 699: Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(hydroxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 700:** 5-[7-{[(3*S*)-3-(Hydroxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

15 **Example 701:** Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*S*)-3-(hydroxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 702: *N,N*-Dibutyl-5-[7-{[(3*S*)-3-(hydroxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 703: Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(methoxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 704: *N*-(4-Hydroxyphenyl)-5-[7-{[(3*S*)-3-(methoxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 705: Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-{[(3*S*)-3-(methoxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 706: 5-[7-{[(3*S*)-3-(Methoxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*,1,2-trimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 707: Phenyl 6-[4-(dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-{[(3*S*)-3-(methoxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 708: *N,N*-Dibutyl-5-[7-{[(3*S*)-3-(methoxymethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2-(phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-1,2-dimethyl-1*H*-pyrrole-3-carboxamide

Example 709: 3-(6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxyloxy)benzoic acid

Step A: 6-(4-{[4-(Benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1H-pyrrol-2-yl)-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl chloride

To a solution of the product from Preparation 6ba (450 mg, 0.47 mmol, 1 eq) in dichloromethane (45 mL), cooled to 0 °C, was added *N,N*-diisopropylethylamine (0.41 mL, 2.36 mmol, 5 eq) and triphosgene (139 mg, 0.47 mmol, 0.99 eq), and the mixture was stirred for 1 h at ambient temperature. The reaction was diluted with dichloromethane and washed with 1M aqueous HCl. The organic phase was dried over magnesium sulfate and concentrated *in vacuo* to afford a yellow solid that was used directly in the next step without further purification, and assuming quantitative transformation.

LC/MS (C₄₆H₄₈ClN₅O₅) 786 [M+H]⁺; RT 1.33 (Method B)

Step B: 3-[(Benzyloxy)carbonyl]phenyl 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1H-pyrrol-2-yl)-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of product obtained from Step A (41 mg, 0.05 mmol) in Acetonitrile (5 mL) was added potassium carbonate (71.87 mg, 0.52 mmol), DMAP (0.052 mmol) and benzyl 3-hydroxybenzoate (0.52 mmol) and the mixture was heated at 60 °C for 16 h.

The reaction mixture was diluted with ethyl acetate, and washed successively with aqueous sodium bicarbonate, and brine. The organic extracts were dried (MgSO₄) and concentrated *in vacuo*. Purification by flash column chromatography (4 g silica; dichloromethane to 5% MeOH/dichloromethane) afforded the desired product.

LC/MS (C₆₀H₅₉N₅O₈) no ionisation; RT 1.46 (Method B)

Step C: 3-(6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxyloxy)benzoic acid

To a solution of the product obtained in Step B (42 mg, 0.04 mmol, 1 eq) in ethanol (8 mL) was added 10% Pd/C (50 mg) and the mixture was shaken under an atmosphere of hydrogen for *ca* 4 h. The reaction was filtered through celite, subsequently eluting with methanol, and concentrated *in vacuo*. Purification by flash column chromatography (4 g silica; dichloromethane to 5% methanol/ dichloromethane) afforded the desired product as a solid.

LC/MS (C₄₆H₄₇N₅O₈) 798 [M+H]⁺; RT 1.07 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₇N₅O₈

[M+H]⁺ calculated: 798.3497

[M+H]⁺ measured: 798.3438

Example 710: 3-{2-[6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 711: 3-({6-[1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}oxy)benzoic acid

Example 712: 3-{2-[6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 713: 3-[(6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}oxy]benzoic acid

Example 714: 3-(2-{6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl)-2-oxoethyl)benzoic acid

Example 715: 3-(6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyloxy)benzoic acid

The procedure is as in Example 709, replacing the product from Preparation 6bb with the product from Preparation 6bf, in Step A.

LC/MS (C₄₂H₄₀N₄O₇) 711 [M-H]⁻; RT 1.27 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₂H₄₀N₄O₇

[M+2H]²⁺ calculated: 713.2970

[M+2H]²⁺ measured: 713.2962

Example 716: 3-{2-[6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 717: 3-({6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}oxy)benzoic acid

Example 718: 3-{2-[6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]-7-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-
2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 719: 3-[(6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*R*)-3-
methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl)-3,4-dihydroisoquinolin-2(1*H*)-
yl]carbonyloxy]benzoic acid

Example 720: 3-(2-{6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*R*)-3-
methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-
2-oxoethyl)benzoic acid

Example 721: 4-(6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-
pyrrol-2-yl]-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-
carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyloxy)benzoic acid

The procedure is as in Example 709, replacing benzyl 3-hydroxybenzoate from Step B with benzyl 4-hydroxybenzoate.

LC/MS (C₄₆H₄₇N₅O₈) 798 [M+H]⁺; RT 1.07 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₇N₅O₈

[M+H]⁺ calculated: 798.3497

[M+H]⁺ measured: 798.3475

Example 722: 4-{2-[6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 723: 4-({[6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}oxy)benzoic acid

Example 724: 4-{2-[6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 725: 4-({[6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}oxy)benzoic acid

Example 726: 4-(2-{6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl)benzoic acid

Example 727: 4-({[6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]}-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}oxy)benzoic acid

Example 728: 4-{2-[6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 729: 4-({[6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}oxy)benzoic acid

Example 730: 4-{2-[6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 731: 4-({[6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}oxy)benzoic acid

Example 732: 4-(2-{6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]}-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl)benzoic acid

Example 733: 3-(6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]}-7-[(*3S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyloxy)benzoic acid

The procedure is as in Example 709, replacing the product from Preparation 6ba with the product from Preparation 6bf, in Step A.

LC/MS (C₄₇H₅₀N₆O₇) 811 [M+H]⁺; RT 2.16 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₅₀N₆O₇

[M+2H]²⁺ calculated: 406.1943

[M+2H]²⁺ measured: 406.1944

5 **Example 734:** 3-{2-[6-{4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

10 **Example 735:** 3-(6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyloxy)benzoic acid

The procedure is as in Example 709, replacing the product from Preparation 6ba with the product from Preparation 6ce, in Step A.

LC/MS (C₄₇H₅₀N₆O₆) 795 [M+H]⁺; RT 1.4 (Method B)

High-resolution mass (ESI+):

15 Empirical formula: C₄₇H₅₀N₆O₆

[M+2H]²⁺ calculated: 398.1969

[M+2H]²⁺ measured: 398.1985

20 **Example 736:** 3-{2-[6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 737: 3-[(6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyloxy]benzoic acid

Example 738: 3-(2-{6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl}-2-oxoethyl)benzoic acid

Example 739: 4-({6-[4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyloxy)benzoic acid

Example 740: 4-{2-[6-[4-[(4-Hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 741: 4-({6-[1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyloxy)benzoic acid

Example 742: 4-{2-[6-[1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]-2-oxoethyl}benzoic acid

Example 743: 4-[(6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyloxy]benzoic acid

5 **Example 744:** 4-(2-{6-[4-(Dibutylcarbamoyl)-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinolin-2(1*H*)-yl}-2-oxoethyl)benzoic acid

Example 745: 5-(2-Acetyl-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-1,2-dimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 **Example 746:** 5-(2-Acetyl-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-1,2-dimethyl-*N*-phenyl-1*H*-pyrrole-3-carboxamide

15 **Example 747:** *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-(2-methyl-7-[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 748: 5-(2-Acetyl-7-[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl)-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-1,2-dimethyl-*N*-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)-1*H*-pyrrole-3-carboxamide

Example 749: 5-(2-Acetyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-1,2-dimethyl-*N*-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)-1*H*-pyrrole-3-carboxamide

5 **Example 750:** *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-(2-methyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)-1*H*-pyrrole-3-carboxamide

Example 751: 5-(2-Acetyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-1,2-dimethyl-*N*-(pyridin-4-yl)-1*H*-pyrrole-3-carboxamide

10 **Example 752:** 5-(2-Acetyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(4-hydroxyphenyl)-1,2-dimethyl-*N*-(pyridin-4-yl)-1*H*-pyrrole-3-carboxamide

15 **Example 753:** *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-(2-methyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(pyridin-4-yl)-1*H*-pyrrole-3-carboxamide

Example 754: 5-(2-Acetyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-*N,N*-diphenyl-1*H*-pyrrole-3-carboxamide

Example 755: 5-(2-Acetyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-*N,N*-diphenyl-1*H*-pyrrole-3-carboxamide

5 **Example 756:** 1,2-Dimethyl-5-(2-methyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N,N*-diphenyl-1*H*-pyrrole-3-carboxamide

Example 757: 5-(2-Acetyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-*N*-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

10 **Example 758:** 5-(2-Acetyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-*N*-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

15 **Example 759:** 1,2-Dimethyl-5-(2-methyl-7-{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)-*N*-phenyl-1*H*-pyrrole-3-carboxamide

Example 760: 5-(2-Acetyl-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-*N*-phenyl-*N*-(pyridin-4-yl)-1*H*-pyrrole-3-carboxamide

Example 761: 5-(2-Acetyl-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-1,2-dimethyl-*N*-phenyl-*N*-(pyridin-4-yl)-1*H*-pyrrole-3-carboxamide

Example 762: 1,2-Dimethyl-5-(2-methyl-7-[[*(3R)*-3-methyl-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl)-*N*-phenyl-*N*-(pyridin-4-yl)-1*H*-pyrrole-3-carboxamide

Example 763: 4-Methylphenyl 6-{4-[(4-hydroxyphenyl)(1-methyl-1*H*-pyrazol-4-yl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 764: 4-Methylphenyl 6-{1-[(4-hydroxyphenyl)(1-methyl-1*H*-pyrazol-4-yl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 765: 4-Methylphenyl 6-{4-[(4-hydroxyphenyl)(pyridin-4-yl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 766: 4-Methylphenyl 6-{1-[(4-hydroxyphenyl)(pyridin-4-yl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 767: 4-Methylphenyl 6-{4-[(4-hydroxyphenyl)(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[[*(3S)*-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(*1H*)-yl]carbonyl]-3,4-dihydroisoquinoline-2(*1H*)-carboxylate

Example 768: 4-Methylphenyl 6-{1-[(4-hydroxyphenyl)(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

5 **Example 769:** 4-Methylphenyl 6-(1,5-dimethyl-4-[(1-methyl-1*H*-pyrazol-4-yl)[4-(phosphonooxy)phenyl]carbamoyl]-1*H*-pyrrol-2-yl)-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

10 **Example 770:** 4-Methylphenyl 6-(1-[(1-methyl-1*H*-pyrazol-4-yl)[4-(phosphonooxy)phenyl]carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl)-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

15 **Example 771:** 4-Methylphenyl 6-(1,5-dimethyl-4-[[4-(phosphonooxy)phenyl](pyridin-4-yl)carbamoyl]-1*H*-pyrrol-2-yl)-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 772: 4-Methylphenyl 7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-6-(1-[[4-(phosphonooxy)phenyl](pyridin-4-yl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

20 **Example 773:** 4-Methylphenyl 6-(1,5-dimethyl-4-[(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)[4-(phosphonooxy)phenyl]carbamoyl]-1*H*-pyrrol-2-yl)-7-{[(3*S*)-3-(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

25 **Example 774:** 4-Methylphenyl 6-(1-[(1-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)[4-(phosphonooxy)phenyl]carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl)-7-{[(3*S*)-3-

(morpholin-4-ylmethyl)-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl)-3,4-dihydroisoquinoline-2(1*H*)-carboxylate

Example 775: 2-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 709, replacing the product from Preparation 6ba in Step A with the product from Preparation 6bb, and replacing benzyl 3-hydroxybenzoate from Step B with 3-(2-hydroxyphenyl)-*N,N*-dimethylpropanamide.

LC/MS ($C_{46}H_{49}N_5O_6$) 768 $[M+H]^+$; RT 2.51 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{49}N_5O_6$

$[M+2H]^{2+}$ calculated: 384.6914

$[M+2H]^{2+}$ measured: 384.6929

Example 776: 3-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 709, replacing the product from Preparation 6ba in Step A with the product from Preparation 6bb, and replacing benzyl 3-hydroxybenzoate from Step B with 3-(3-hydroxyphenyl)-*N,N*-dimethylpropanamide.

LC/MS ($C_{46}H_{49}N_5O_6$) 768 $[M+H]^+$; RT 2.50 (Method A)

High-resolution mass (ESI+):

Empirical formula: $C_{46}H_{49}N_5O_6$

$[M+2H]^{2+}$ calculated: 384.6914

[M+2H]²⁺ measured: 384.6913

Example 777: 5-{7-[(3*S*)-3-[(Dimethylamino)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

5

Step A: *N*-[4-(Benzyloxy)phenyl]-5-{7-[(3*S*)-3-[(dimethylamino)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

10

A solution of the compound from Preparation 6be (50 mg, 0.05 mmol) and *N,N*-diisopropylethylamine (48 μ L, 0.27 mmol) in dichloromethane (5 mL) was cooled to 0°C. To this was added phenylacetyl chloride (8 μ L, 0.06 mmol) and the reaction was then stirred at ambient temperature for 1 h. The reaction mixture was diluted with dichloromethane, washed with 1M aqueous sodium hydroxide, and then brine. The organic extract was dried over magnesium sulphate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; dichloromethane to 10 % methanol in dichloromethane) afforded the desired product.

15

LC/MS (C₅₁H₅₃N₅O₄) 800 [M+H]⁺; RT 1.26 (Method B)

Step B: 5-{7-[(3*S*)-3-[(dimethylamino)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

20

To a solution of the material from Step A (30 mg, 0.04 mmol) in ethanol (5 mL) was added 10% Pd/C (catalytic amount) and the mixture was shaken under an atmosphere of hydrogen for *ca* 16 h. The mixture is filtered through celite, subsequently eluting with methanol, and concentrated under reduced pressure. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; dichloromethane to 7 % methanol in dichloromethane) afforded the desired product.

25

LC/MS (C₄₄H₄₇N₅O₄) 710 [M+H]⁺; RT 1.07 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₄H₄₇N₅O₄

[M+H]⁺ calculated: 710.3701

[M+H]⁺ measured: 710.3702

5 **Example 778:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6bf.

10 **LC/MS** (C₄₇H₅₂N₆O₄) 765 [M+H]⁺; RT 1.08 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₅₂N₆O₄

[M+H]⁺ calculated: 765.4123

[M+H]⁺ measured: 765.4141

15 **Example 779:** *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-[2-(morpholin-4-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with product from Preparation 6bc.

20 **LC/MS** (C₄₇H₅₁N₅O₅) 766 [M+H]⁺; RT 1.08 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₇H₅₁N₅O₅

[M+H]⁺ calculated: 766.3963

$[M+H]^+$ measured: 766.3982

Example 780: *N*-Ethyl-*N*-(4-hydroxyphenyl)-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

- 5 The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with product from Preparation 6f.

LC/MS ($C_{47}H_{51}N_5O_5$) 766 $[M+H]^+$; RT 1.15 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{47}H_{51}N_5O_5$

- 10 $[M+H]^+$ calculated: 766.3963

$[M+H]^+$ measured: 766.3967

Example 781: *N*-(4-Hydroxyphenyl)-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-propyl-1*H*-pyrrole-3-carboxamide

- 15 The procedure is as in Example 777, replacing the product from Preparation 6be in Step A with the product from Preparation 6g.

LC/MS ($C_{48}H_{53}N_5O_5$) 780 $[M+H]^+$; RT 1.19 (Method B)

High-resolution mass (ESI+):

Empirical formula: $C_{48}H_{53}N_5O_5$

- 20 $[M+H]^+$ calculated: 780.4119

$[M+H]^+$ measured: 780.4101

Example 782: 5-{7-[(3*S*)-3-[(Dimethylamino)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

LC/MS (C₃₈H₄₀F₃N₅O₄) 688 [M+H]⁺; RT 1.07 (Method B)

5 **High-resolution mass (ESI+):**

Empirical formula: C₃₈H₄₀N₅O₄F₃

[M+H]⁺ calculated: 688.3105

[M+H]⁺ measured: 688.3094

Example 783: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(trifluoroacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

LC/MS (C₄₁H₄₅N₆O₄F₃) 743 [M+H]⁺; RT 1.09 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₅N₆O₄F₃

15 [M+2H]²⁺ calculated: 372.1800

[M+2H]²⁺ measured: 372.1801

Example 784: 4-Bromophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

20 **Step A:** 4-Bromophenyl 6-(4-{[4-(benzyloxy)phenyl](methyl)carbamoyl}-1,5-dimethyl-1*H*-pyrrol-2-yl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of the compound from Preparation 6ba (108 mg, 0.11 mmol) and *N,N*-diisopropylethylamine (0.1 mL, 0.57 mmol) in dichloromethane (15 mL), cooled to 0 °C,

was added triphosgene (34 mg, 0.11 mmol) and the mixture was stirred for 1 h at ambient temperature. The reaction was diluted with dichloromethane and washed with 1M aqueous HCl. The organic extract was dried over magnesium sulfate, filtered and concentrated *in vacuo* to afford a yellow solid. To a solution of this solid in acetonitrile (10 mL) was added potassium carbonate (152 mg, 1.1 mmol), DMAP (13 mg, 0.11 mmol) and 4-bromophenol (98 mg, 0.57 mmol) and the mixture was heated at 60 °C for *ca* 16 h. The reaction was diluted with ethyl acetate and washed successively with saturated aqueous sodium bicarbonate, and brine. The organic extract was dried over magnesium sulfate and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; dichloromethane to 5 % methanol in dichloromethane) afforded the desired product (79 mg, 0.09 mmol).

LC/MS (C₅₂H₅₂BrN₅O₆) 922 [M+H]⁺; RT 1.50 (Method B)

Step B: 4-Bromophenyl 6-{4-[4-(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The product from Step A was dissolved in dichloromethane (10 mL) and cooled to 0 °C. To this was added boron trichloride (5 eq.) drop-wise. The reaction was then allowed to warm to ambient temperature for 1 h. The reaction mixture was adsorbed onto isolate and purified by chromatography (CombiFlash R_f, 12 g RediSep™ silica cartridge) eluting in a gradient of dichloromethane to 5 % methanol in dichloromethane to afford the desired product.

LC/MS (C₄₅H₄₆BrN₅O₆) 832 [M+H]⁺; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₄₆N₅O₆Br

[M+2H]²⁺ calculated: 416.6388

[M+2H]²⁺ measured: 416.6374

Example 785: 4-Methylphenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

To a solution of product from Preparation 6cb (50 mg, 0.09 mmol) and *N,N*-diisopropylethylamine (38 μ L, 0.19 mmol) in dichloromethane (5 mL), cooled to 0 °C, was added 4-methylphenyl chloroformate (15 μ L, 0.1 mmol) and the mixture was stirred at ambient temperature for 1 h. The reaction was diluted with dichloromethane, then washed successively with 1M aqueous sodium hydroxide, and brine. The organic extract was dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 12 g RediSep™ silica cartridge; dichloromethane to 5 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₄₂H₄₂N₄O₄) 667 [M+H]⁺; RT 1.50 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₂H₄₂N₄O₄

[M+H]⁺ calculated: 667.3279

[M+H]⁺ measured: 667.3287

Example 786: 4-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: 6-{1,5-Dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl chloride

To a solution of the product from Preparation 6cb (300 mg, 0.56 mmol) and *N,N*-diisopropylethylamine (0.29 mL, 1.69 mmol) in dichloromethane (10 mL), cooled to 0 °C, was added triphosgene (167 mg, 0.56 mmol) and the mixture was allowed to warm to ambient temperature over 1 h. The reaction was diluted with dichloromethane, washed with 1M aqueous HCl, and dried over magnesium sulphate. Concentration *in vacuo*

afforded a yellow solid that was used directly in the next step without further purification, and assuming quantitative transformation.

LC/MS ($C_{35}H_{35}ClN_4O_3$) 595 $[M+H]^+$; RT 1.41 (Method B)

Step B: 4-Cyanophenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1H-pyrrol-2-yl}-7-
5 /[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-
tetrahydroisoquinoline-2-carboxylate

To a solution of the carbamoyl chloride from Step A (50 mg, 0.072 mmol) in acetonitrile (5 mL) was added potassium carbonate (50 mg, 0.36 mmol) and 4-hydroxybenzonitrile (10 mg, 0.086 mmol). After heating at 60 °C for *ca.* 16 h the reaction was allowed to cool to
10 ambient temperature, then diluted with ethyl acetate and washed successively with 1M aqueous sodium hydroxide, and brine. The organics extract was dried over magnesium sulphate, concentrated *in vacuo*, and purified by preparative HPLC to afford the desired product.

LC/MS ($C_{42}H_{39}N_5O_4$) 678 $[M+H]^+$; RT 1.41 (Method B)

15 **High-resolution mass (ESI+):**

Empirical formula: $C_{42}H_{39}N_5O_4$

$[M+H]^+$ calculated: 678.3075

$[M+H]^+$ measured: 678.3086

Example 787: 3-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{1,5-dimethyl-4-
20 [methyl(phenyl)carbamoyl]-1H-pyrrol-2-yl}-7-[(3R)-3-methyl-1,2,3,4-
tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 786, replacing 4-hydroxybenzonitrile in Step B with 3-(3-hydroxyphenyl)-*N,N*-dimethylpropanamide.

LC/MS ($C_{46}H_{49}N_5O_5$) 752 $[M+H]^+$; RT 1.39 (Method B)

25 **High-resolution mass (ESI+):**

Empirical formula: C₄₆H₄₉N₅O₅

[M+2H]²⁺ calculated: 376.6940

[M+2H]²⁺ measured: 376.6930

Example 788: 4-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{1,5-dimethyl-4-
5 [methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-
tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 786, replacing 4-hydroxybenzonitrile in Step B with 3-(4-hydroxyphenyl)-*N,N*-dimethylpropanamide.

LC/MS (C₄₆H₄₉N₅O₅) 752 [M+H]⁺; RT 1.38 (Method B)

10 **High-resolution mass (ESI+):**

Empirical formula: C₄₆H₄₉N₅O₅

[M+2H]²⁺ calculated: 376.6940

[M+2H]²⁺ measured: 376.6926

Example 789: 1*H*-Indol-5-yl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-
15 pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-
tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 786, replacing 4-hydroxybenzonitrile in Step B with 1*H*-indol-5-ol.

LC/MS (C₄₃H₄₁N₅O₄) 692 [M+H]⁺; RT 1.41 (Method B)

20 **High-resolution mass (ESI+):**

Empirical formula: C₄₃H₄₁N₅O₄

[M+H]⁺ calculated: 692.3231

[M+H]⁺ measured: 692.3210

Example 790: Cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

5 The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6bb, and replacing phenylacetyl chloride with cyclohexyl chloroformate.

LC/MS (C₄₁H₄₆N₄O₅) 675 [M+H]⁺; RT 1.45 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₆N₄O₅

10 [M+H]⁺ calculated: 675.3541

[M+H]⁺ measured: 675.3548

Example 791: Cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

15 The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6ba, and replacing phenylacetyl chloride with cyclohexyl chloroformate.

LC/MS (C₄₅H₅₃N₅O₆) no ionisation; RT 1.28 (Method B)

High-resolution mass (ESI+):

20 Empirical formula: C₄₅H₅₃N₅O₆

[M+H]⁺ calculated: 760.4069

[M+H]⁺ measured: 760.4078

Example 792: *N*-Cyclohexyl-6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6bb, and replacing phenylacetyl chloride with cyclohexyl isocyanate.

LC/MS (C₄₁H₄₇N₅O₄) 674 [M+H]⁺; RT 1.34 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₇N₅O₄

[M+H]⁺ calculated: 674.3701

[M+H]⁺ measured: 674.3696

Example 793: *N*-Cyclohexyl-6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxamide

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6ba, and replacing phenylacetyl chloride with cyclohexyl isocyanate.

LC/MS (C₄₅H₅₄N₆O₅) no ionisation; RT 1.18 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₅H₅₄N₆O₅

[M+H]⁺ calculated: 759.4228

[M+H]⁺ measured: 759.4192

Example 794: 5-[2-(2-Cyclohexylacetyl)-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6ba, and replacing phenylacetyl chloride with 2-cyclohexylacetyl chloride.

LC/MS (C₄₆H₅₅N₅O₅) no ionisation; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₅₅N₅O₅

[M+H]⁺ calculated: 758.4276

[M+H]⁺ measured: 758.4247

Example 795: 5-[2-(2-Cyclohexylacetyl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6bb, and replacing phenylacetyl chloride with 2-cyclohexylacetyl chloride.

LC/MS (C₄₂H₄₈N₄O₄) 673 [M+H]⁺; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₂H₄₈N₄O₄

[M+H]⁺ calculated: 673.3748

[M+H]⁺ measured: 673.3741

Example 796: Cyclopentyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6ba, and replacing phenylacetyl chloride with cyclopentyl chloroformate.

LC/MS (C₄₄H₅₁N₅O₆) 746 [M+H]⁺; RT 1.24 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₄H₅₁N₅O₆

[M+H]⁺ calculated: 746.3912

[M+H]⁺ measured: 746.3891

Example 797: (1*R*,2*S*,5*R*)-5-Methyl-2-(propan-2-yl)cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6ba, and replacing phenylacetyl chloride with (1*R*,2*S*,5*R*)-5-methyl-2-(propan-2-yl)cyclohexyl chloroformate.

LC/MS (C₄₉H₆₁N₅O₆) 816 [M+H]⁺; RT 1.47 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₉H₆₁N₅O₆

[M+H]⁺ calculated: 816.4695

[M+H]⁺ measured: 816.4717

Example 798: 5-{2-[2-(Adamantan-1-yl)acetyl]-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-*N*-(4-hydroxyphenyl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6ba, and replacing phenylacetyl chloride with 2-(adamantan-1-yl)acetyl chloride.

LC/MS (C₅₀H₅₉N₅O₅) 810 [M+H]⁺; RT 1.32 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₅₀H₅₉N₅O₅

[M+H]⁺ calculated: 810.4589

[M+H]⁺ measured: 810.4586

Example 799: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(piperidin-1-yl)acetyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

Step A: *N*-[4-(Benzyloxy)phenyl]-*N*,1,2-trimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(piperidin-1-yl)acetyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

To a solution of the compound from Preparation 6bb (50 mg, 0.08 mmol) in DMF (2 mL) was added 2-(1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (30 mg, 0.08 mmol), *N,N*-diisopropylethylamine (41 µL, 0.23 mmol, 3 eq), and 2-(piperidin-1-yl)acetic acid (13 mg, 0.09 mmol), and the mixture was stirred at ambient temperature for *ca.* 16 h. The reaction was concentrated *in vacuo* and partitioned between ethyl acetate and water. The organic phase was washed with brine, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R₆, 4 g RediSep™ silica cartridge; dichloromethane to 10 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₄₈H₅₃N₅O₄) no ionisation; RT 1.29 (Method B)

Step B: *N*-(4-Hydroxyphenyl)-*N*,1,2-trimethyl-5-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-[2-(piperidin-1-yl)acetyl]-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide

To a solution of the product from Step A (30 mg, 0.04 mmol) in ethanol (5 mL) was added 10% Pd/C (catalytic amount) and the mixture was shaken under an atmosphere of hydrogen for *ca* 16 h. The mixture was filtered through celite, eluted with methanol, and the filtrate concentrated *in vacuo*. Purification by preparative HPLC (H₂O-TFA/acetonitrile; gradient elution) afforded the desired product.

LC/MS (C₄₁H₄₇N₅O₄) 674 [M+H]⁺; RT 1.09 (Method B)

High-resolution mass (ESI⁺):

Empirical formula: C₄₁H₄₇N₅O₄

[M+2H]²⁺ calculated: 337.6887

[M+2H]²⁺ measured: 337.6887

Example 800: (1*r*,4*r*)-4-{[(*tert*-Butoxy)carbonyl]amino}cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

Step A: (1*r*,4*r*)-4-{[(*tert*-Butoxy)carbonyl]amino}cyclohexyl chloroformate

Triphosgene (69 mg, 0.23 mmol) was dissolved in dichloromethane (3 mL) and cooled to 0 °C. To this was added a solution of *tert*-butyl *N*-(4-hydroxycyclohexyl)carbamate (100 mg, 0.46 mmol) and triethylamine (0.06 mL, 0.46 mmol, 2 eq) in dichloromethane (1 mL) drop-wise. After stirring for 2 h at ambient temperature, the reaction was concentrated *in vacuo* then re-dissolved / suspended in dichloromethane and filtered. The filtrate (2 mL) was used directly in the next step, assuming a quantitative transformation.

Step B: (1*r*,4*r*)-4- $\{[(tert\text{-}Butoxy)carbonyl]amino\}$ cyclohexyl 6-(4- $\{[4\text{-}(benzyloxy)phenyl](methyl)carbamoyl\}$ -1,5-dimethyl-1*H*-pyrrol-2-yl)-7- $[(3R)\text{-}3\text{-methyl-}1,2,3,4\text{-tetrahydroisoquinoline-}2\text{-carbonyl}]\text{-}1,2,3,4\text{-tetrahydroisoquinoline-}2\text{-carboxylate}$

A solution of product from Preparation 6bb (50 mg, 0.08 mmol) in dichloromethane (3 mL) was cooled to 0 °C. To this was added triethylamine (54 μ L, 0.39 mmol) and the solution of the product from Step A (0.49 mmol) and the mixture was stirred for 30 min at ambient temperature. The reaction was diluted with dichloromethane then washed sequentially with 1M aqueous sodium hydroxide, and brine. The organic phase was dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; dichloromethane to 3 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₅₃H₆₁N₅O₇) 780 [M-Boc+H]⁺; RT 1.60 (Method B)

Step C: (1*r*,4*r*)-4- $\{[(tert\text{-}Butoxy)carbonyl]amino\}$ cyclohexyl 6-(4- $\{[4\text{-}(hydroxyphenyl)(methyl)carbamoyl\}$ -1,5-dimethyl-1*H*-pyrrol-2-yl)-7- $[(3R)\text{-}3\text{-methyl-}1,2,3,4\text{-tetrahydroisoquinoline-}2\text{-carbonyl}]\text{-}1,2,3,4\text{-tetrahydroisoquinoline-}2\text{-carboxylate}$

To a solution of the product from Step B (96 mg, 0.11 mmol) in ethanol (5 mL) was added 10% Pd/C (catalytic amount) and the mixture was shaken under an atmosphere of hydrogen for *ca.* 16 h. The mixture was filtered through celite, eluted with methanol, and the filtrate concentrated *in vacuo*. Purification by flash column chromatography (CombiFlash R_f, 4 g RediSep™ silica cartridge; dichloromethane to 5 % methanol in dichloromethane) afforded the desired product.

LC/MS (C₄₆H₅₅N₅O₇) 790 [M+H]⁺; RT 1.43 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₅₅N₅O₇

[M+H]⁺ calculated: 790.4174

[M+H]⁺ measured: 790.4166

Example 801: (1*r*,4*r*)-4-Aminocyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride

5 The product from Example 800 (48 mg, 0.06 mmol) was dissolved in 4M HCl in 1,4-dioxane (5 mL) and stirred at ambient temperature for *ca* 15 min. The reaction was concentrated *in vacuo* to obtain a solid which was suspended in ether and stirred for 1 h at 0 °C. The solid was filtered, washed with ether, and dried under vacuum to afford the desired product.

10 **LC/MS** (C₄₁H₄₇N₅O₅) 690 [M+H]⁺; RT 1.10 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₁H₄₇N₅O₅

[M+2H]²⁺ calculated: 345.6861

[M+2H]²⁺ measured: 345.6864

15 **Example 802:** (1*r*,4*r*)-4-(Dimethylamino)cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

20 To a solution of the product from Example 801 (20 mg, 0.03 mmol, 1 eq) in methanol (2 mL) was added paraformaldehyde (4 mg, 0.14 mmol), acetic acid (8 µL, 0.14 mmol, 5 eq) and sodium cyanoborohydride (9 mg, 0.14 mmol), and the mixture was stirred at ambient temperature for *ca* 16 h. The reaction was quenched by the addition of water (1 mL) and extracted with dichloromethane. The organic extract was washed sequentially with water, and brine, and then dried over magnesium sulfate, filtered and concentrated *in vacuo*. Purification by preparative HPLC (H₂O-TFA/acetonitrile; gradient elution) afforded
25 the desired product

LC/MS (C₄₃H₅₁N₅O₅) 718 [M+H]⁺; RT 1.10 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₃H₅₁N₅O₅

[M+2H]²⁺ calculated: 359.7018

[M+2H]²⁺ measured: 359.7017

5 **Example 803:** 4-(*N*-Methylacetamido)phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

10 The procedure is as in Example 709, replacing benzyl 3-hydroxybenzoate from Step B with *N*-(4-hydroxyphenyl)-*N*-methylacetamide.

LC/MS (C₄₈H₅₂N₆O₇) 825 [M+H]⁺; RT 1.07 (Method B)

High-resolution mass (ESI+):

Empirical formula: C₄₈H₅₂N₆O₇

[M+2H]²⁺ calculated: 413.2022

15 [M+2H]²⁺ measured: 413.2031

Example 804: 4-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate

20 The procedure is as in Example 709, replacing the product from Preparation 6ba in Step A with the product from Preparation 6bb, and replacing benzyl 3-hydroxybenzoate from Step B with 3-(4-hydroxyphenyl)-*N,N*-dimethylpropanamide.

LC/MS (C₄₆H₄₉N₅O₆) 384 [M+2H]²⁺; RT 2.49 (Method A)

High-resolution mass (ESI+):

Empirical formula: C₄₆H₄₉N₅O₆

[M+2H]²⁺ calculated: 384.6914

[M+2H]²⁺ measured: 384.6933

Example 805: Toly 6-[4-[(5-cyano-1,2-dimethyl-1*H*-pyrrol-3-yl)-(4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl]-7-[(3*S*)-3-(morpholinomethyl)-3,4-dihydro-1*H*-isoquinoline-2-carbonyl]-3,4-dihydro-1*H*-isoquinoline-2-carboxylate hydrochloride

Step A: *Benzyl 7-formyl-6-(4-ethoxycarbonyl-1,5-dimethyl-1H-pyrrol-2-yl)-3,4-dihydro-1H-isoquinoline-2-carboxylate*

To a solution of 1.3 g of ethyl 1,2-dimethyl-1*H*-pyrrole-3-carboxylate (7.97 mmol) in 15 mL of *N,N*-dimethylacetamide there are successively added 2.5 g of benzyl 6-bromo-7-formyl-3,4-dihydro-1*H*-isoquinoline-2-carboxylate (6.64 mmol), 1.3 g of potassium acetate (13.3 mmol), and then the batch is stirred under argon for 20 minutes. 0.2 g of palladium catalyst PdCl₂(PPh₃)₂ (0.33 mmol) is then added. The reaction mixture is then heated at 85°C for 4.5 hours. The mixture is allowed to return to ambient temperature and is then diluted with ethyl acetate. After filtration, the organic phase is washed with saturated aqueous sodium chloride solution, dried over magnesium sulphate, filtered and concentrated to dryness. The crude product thereby obtained is purified by chromatography over silica gel (petroleum ether / ethyl acetate gradient). The title product is obtained in the form of a yellow foam.

¹H NMR (400 MHz, dmso-d₆) δ ppm: 9.78 (s, 1 H), 7.78 (s, 1 H), 7.45-7.3 (m, 5 H), 7.32 (s, 1 H), 6.39 (s, 1 H), 5.14 (s, 2 H), 4.72 (m, 2 H), 4.18 (q, 2 H), 3.68 (m, 2 H), 3.31 (s, 3 H), 2.93 (t, 2 H), 2.55 (s, 3 H), 1.25 (t, 3 H)

Step B: *2-Benzylloxycarbonyl-6-(4-ethoxycarbonyl-1,5-dimethyl-1H-pyrrol-2-yl)-3,4-dihydro-1H-isoquinoline-7-carboxylic acid*

A solution containing 1.3 g of the compound obtained in Step A (2.82 mmol) and 2.4 mL (22.6 mmol) of 2-methyl-2-butene in a mixture composed of 2.5 mL of acetone and 2.5 mL of tetrahydrofuran is prepared. There are added, dropwise at 0°C, 20 mL of an aqueous solution containing a mixture of 0.64 g of NaClO₂ (7.05 mmol) and 1.4 g of NaHPO₄ (9.88 mmol). The batch is then stirred vigorously at ambient temperature for 16 hours. The reaction mixture is then concentrated to remove the acetone and the tetrahydrofuran. Ethyl acetate is added and the organic phase is washed with water and then with saturated aqueous sodium chloride solution, dried over magnesium sulphate, filtered and concentrated to dryness. The yellow foam then obtained is subsequently used without being otherwise purified.

¹H NMR (400 MHz, dmso-d₆) δ ppm: 12.75 (m, 1 H), 7.69 (s, 1 H), 7.44-7.3 (m, 5 H), 7.14 (s, 1 H), 6.2 (s, 1 H), 5.14 (s, 2 H), 4.67 (m, 2 H), 4.15 (q, 2 H), 3.67 (m, 2 H), 3.21 (s, 3 H), 2.86 (m, 2 H), 2.49 (s, 3 H), 1.24 (t, 3 H)

Step C: Benzyl 6-(4-ethoxycarbonyl-1,5-dimethyl-1H-pyrrol-2-yl)-7-[(3S)-3-(morpholinomethyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-3,4-dihydro-1H-isoquinoline-2-carboxylate

To a solution of 1.5 g of the compound obtained in Step B (2.49 mmol) in 13 mL of dichloromethane there are added 0.636 g of 4-[(3S)-1,2,3,4-tetrahydroisoquinolin-3-yl]-methylmorpholine (2.74 mmol), 1.1 mL of *N,N,N*-triethylamine (7.47 mmol), 0.51 g of 1-ethyl-3-(3'-dimethylaminopropyl)-carbodiimide (EDC) (2.99 mmol) and then 0.4 g of hydroxybenzotriazole (HOBT) (2.99 mmol). The reaction mixture is stirred at ambient temperature for 16 hours and is then diluted with a mixture of dichloromethane and saturated sodium hydrogen carbonate solution. After separation of the phases, the organic phase is dried over magnesium sulphate, filtered and evaporated to dryness. The crude product thereby obtained is then purified by chromatography over silica gel (dichloromethane / ethyl acetate gradient). The product is obtained in the form of a white foam.

¹H NMR (500 MHz, dmso-d₆) δ ppm: 7.4-7.3 (m, 5 H), 7.25-7.15 (4*s, 2 H), 7.2-6.9 (m, 4 H), 6.32/6.28/6.1 (3*bs, 1 H), 5.15 (3*s, 2 H), 5.15/4.85/3.7 (3*m, 1 H), 5-4 (m, 2 H),

4.7 (m, 2 H), 4.1 (m, 2 H), 3.78/3.6 (2*m, 2 H), 3.6-3.4 (m, 4 H), 3.45/3.39/3.2 (3*s, 3 H), 3-2.45 (m, 2 H), 2.9 (m, 2 H), 2.5-1.9 (4*m, 4 H), 2.5/2.41/2.05 (4*bs, 3 H), 2.35-1.7 (m, 2 H), 1.2/1.1 (3*t, 3 H)

¹³C NMR (500 MHz, dms_o-d₆) δ ppm: 130/125, 129-126, 128, 110, 66.5, 66.5, 59, 57.5, 53.5, 49.5/44.5/43, 45.5, 44.5/41, 41.5, 31.5, 30/29, 28, 15, 11.5

Step D: *1,2-Dimethyl-5-[7-[(3S)-3-(morpholinomethyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-1H-pyrrole-3-carboxylic acid*

To a solution of 1.1 g of the compound obtained in Step C (1.59 mmol) in 8 ml of ethanol there is added 0.9 ml of aqueous 5N sodium hydroxide solution (4.77 mmol). The reaction mixture is stirred at ambient temperature for 24 hours at 80°C and a second addition of 0.9 ml of aqueous 5N sodium hydroxide solution is made. After being in contact for a period of 24 hours at the same temperature, this operation is carried out a second time. The ethanol is then concentrated and the reaction mixture is diluted with water before adding aqueous 1N hydrochloric acid solution until the pH is 7. After extracting the aqueous phase with dichloromethane, the organic phases are combined and concentrated to dryness to obtain the title product in the form of a yellow foam.

¹H NMR (500 MHz, dms_o-d₆) δ ppm: 7.2-6.9 (m, 4 H), 7.05/7 (2*s, 2 H), 6.31/6.25/6.1 (3*s, 1 H), 5.1/4.85/3.7 (3*m, 1 H), 4.15/4.1 (m, 2 H), 3.91/3.81 (bs+dd, 2 H), 3.6-3.4 (m, 4 H), 3.4/3.35/3.15 (3*s, 3 H), 3-2.9 (m, 2 H), 2.9-1.9 (m, 2 H), 2.75 (m, 2 H), 2.5/2.4/1.98 (3*s, 3 H), 2.5-1.9 (m, 6 H)

¹³C NMR (500 MHz, dms_o-d₆) δ ppm: 130/125, 129-126, 110.5, 66.5, 57/54, 49.5/44.5/43, 47.5, 44, 43.5, 32, 30, 29, 11.5

Step E: *1,2-Dimethyl-5-[2-(4-methylphenoxy)carbonyl-7-[(3S)-3-(morpholinomethyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-3,4-dihydro-1H-isoquinolin-6-yl]-1H-pyrrole-3-carboxylic acid*

To a solution of 0.35 g of the compound obtained in Step D (0.66 mmol) in 1 ml of dioxane there is added, at 0°C, 0.33 ml of aqueous 2N sodium hydroxide solution (0.66 mmol) and, each in several portions, 0.106 ml of p-tolyl chloroformate (0.73 mmol) and 0.36 ml of aqueous 2N sodium hydroxide solution (0.73 mmol). The reaction mixture is stirred for 2 hours at ambient temperature and is then diluted with water and extracted with dichloromethane. The aqueous phase is acidified and then extracted with dichloromethane. The organic phases are combined, dried over magnesium sulphate, filtered and concentrated.

The residue obtained is taken up in 12 ml of methanol and 3.2 ml of aqueous 1N potassium hydroxide solution (3.26 mmol) and then the reaction mixture is stirred at ambient temperature for 20 minutes. After adding aqueous 0.1N hydrochloric acid solution until the pH is 4, the mixture is extracted with dichloromethane. The organic phases are combined, washed with saturated aqueous sodium chloride solution, dried over magnesium sulphate, filtered and concentrated. The residue is then taken up in a mixture of dichloromethane and *iso*-propyl ether. The solid then obtained is filtered off and washed with ether. The title product is obtained in the form of a solid, which is subsequently used without being otherwise purified.

¹H NMR (500 MHz, dmso-d₆) δ ppm: 11.5 (bs, 1 H), 7.28 (bs, 1 H), 7.2-6.9 (m, 4 H), 7.2 (bd, 2 H), 7.02 (bd, 2 H), 6.88 (bs, 1 H), 6.35/6.28/6.12 (3*s, 1 H), 5.11/4.85/3.7 (3*m, 1 H), 4.85-4.65 (m, 2 H), 4.8-4.1 (m, 2 H), 4.2-3.6 (m, 2 H), 3.6-3.4 (m, 4 H), 3.45/3.4/3.18 (3*s, 3 H), 3-1.8 (m, 2 H), 2.97 (ml, 2 H), 2.5/2.41/1.99 (4*s, 3 H), 2.5-1.9 (4*m, 4 H), 2.35/2.18 (2*m, 2 H), 2.25 (bs, 3 H)

¹³C NMR (500 MHz, dmso-d₆) δ ppm: 130, 129, 129-126, 125, 122, 110.5, 66.5, 58, 54, 49.5/44/43, 45.5, 44.5/41.5, 41.5, 32, 30-29, 28, 20.5, 11

Step F: Tolyl 6-[4-[(5-cyano-1,2-dimethyl-1H-pyrrol-3-yl)-(4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1H-pyrrol-2-yl]-7-[(3S)-3-(morpholinomethyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-3,4-dihydro-1H-isoquinoline-2-carboxylate hydrochloride

To a solution of 0.232 g of the compound obtained in Step E (0.35 mmol) in 9 mL of dichloroethane there is added 0.05 mL of 1-chloro-*N,N*,2-trimethyl-prop-1-en-1-amine (0.385 mmol). The reaction mixture is stirred at ambient temperature for 2 hours and there is then added 0.131 g of 4-[4-*tert*-butyl(dimethyl)silyl]oxyanilino]-1,5-dimethyl-1*H*-pyrrole-2-carbonitrile (0.385 mmol, product of Preparation 3f). The batch is stirred at 80°C overnight. The reaction mixture is diluted with a mixture of dichloromethane and saturated aqueous sodium hydrogen carbonate solution. After separation of the phases, the organic phase is dried over magnesium sulphate, filtered and concentrated to dryness. The crude product thereby obtained is subsequently used without being otherwise purified.

To a solution of the compound thereby obtained in 1.8 ml of tetrahydrofuran there is added 0.53 ml of a 1N solution of tetrabutylammonium fluoride in tetrahydrofuran (13.5 mmol). The reaction mixture is stirred at ambient temperature for 1 hour and is then diluted with a mixture of dichloromethane and saturated aqueous sodium hydrogen carbonate solution. The aqueous phase is extracted with dichloromethane and then the organic phases are combined, dried over magnesium sulphate, filtered and concentrated. The crude product is then purified by chromatography over silica gel (dichloromethane / methanol gradient). The solid thereby obtained is converted to the hydrochloride form, dissolved in a mixture of acetonitrile and water, filtered and then lyophilised to isolate the title product.

High-resolution mass (ESI/FIA/HR and MS/MS):

Empirical formula: C₅₂H₅₃N₇O₆

[M+H]⁺ calculated: 872.4130

[M+H]⁺ measured: 872.4130

Elemental microanalysis: (% , theoretical:measured)

%C=68.75:68.07; %H=5.99:5.90; %N=10.79:10.68; %Cl=3.90:3.84

Example 806: *N*-(4-Hydroxyphenyl)-5-(2-[(2*S*)-2-hydroxy-2-phenylacetyl]-6-
{[(3*R*)-3-methyl-3,4-dihydroisoquinolin-2(1*H*)-yl]carbonyl}-2,3-dihydro-1*H*-isoindol-
5-yl)-*N*,1,2-trimethyl-1*H*-pyrrole-3-carboxamide

Example 807: **Cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-[2-(morpholin-4-yl)ethyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate**

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6bc, and replacing phenylacetyl chloride with cyclohexyl chloroformate.

LC/MS (C₄₆H₅₅N₅O₆) 774 [M+H]⁺; RT 1.22 (Method B)

Example 808: **Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate**

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with product from Preparation 6bd; and replacing phenylacetyl chloride in Step A with phenyl chloroformate.

LC/MS (C₄₇H₅₁N₅O₆) 782 [M+H]⁺; RT 1.15 (Method B)

Example 809: **Cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate**

The procedure is as in Example 777, replacing the compound from Preparation 6be in Step A with the compound from Preparation 6bd, and replacing phenylacetyl chloride with cyclohexyl chloroformate.

LC/MS (C₄₇H₅₇N₅O₆) 788 [M+H]⁺; RT 1.21 (Method B)

Example 810: **Phenyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate**

Example 811: **Cyclohexyl 6-{1,5-dimethyl-4-[methyl(phenyl)carbamoyl]-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate**

5 The procedure is as in Example 114, replacing the product from Preparation 6cb with the product from Preparation 6cd, and replacing phenyl chloroformate with cyclohexyl chloroformate.

LC/MS (C₄₇H₅₇N₅O₅) 772 [M+H]⁺; RT 1.33 (Method B)

PHARMACOLOGICAL STUDY

EXAMPLE A: Inhibition of Bcl-2 by the fluorescence polarisation technique

The fluorescence polarisation tests were carried out on microplates (384 wells). The Bcl-2 protein, labelled (histag-Bcl-2 such that Bcl-2 corresponds to the UniProtKB® primary
5 accession number: P10415), at a final concentration of 2.50×10^{-8} M, is mixed with a fluorescent peptide (Fluorescein-REIGAQLRRMADDLNAQY), at a final concentration of 1.00×10^{-8} M in a buffer solution (Hepes 10 mM, NaCl 150 mM, Tween20 0.05%, pH 7.4), in the presence or in the absence of increasing concentrations of test compounds. After incubation for 2 hours, the fluorescence polarisation is measured.

10 The results are expressed in IC_{50} (the concentration of compound that inhibits fluorescence polarisation by 50%) and are presented in Table 1 below.

The results show that the compounds of the invention inhibit interaction between the Bcl-2 protein and the fluorescent peptide described hereinbefore.

EXAMPLE B: *In vitro* cytotoxicity

15 The cytotoxicity studies were carried out on the RS4;11 leukaemia cell line and on H146 small Cell Lung Carcinoma cell line. The cells are distributed onto microplates and exposed to the test compounds for 48 hours. The cell viability is then quantified by a colorimetric assay, the Microculture Tetrazolium Assay (Cancer Res., 1987, 47, 939-942).

20 The results are expressed in IC_{50} (the concentration of compound that inhibits cell viability by 50%) and are presented in Table 1 below.

The results show that the compounds of the invention are cytotoxic.

**Table 1: IC₅₀ of Bcl-2 inhibition (fluorescence polarisation test)
and of cytotoxicity for RS4;11 cells**

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 1	13	140
Example 2	397	> 1880
Example 3	18	236
Example 4	9.2	71
Example 5	43	70
Example 6	95	> 1880
Example 7	72% @3.3µM	1080
Example 8	7.3	98
Example 9	9.0	125
Example 10	127	ND
Example 11	22	763
Example 12	43	759
Example 13	108	729
Example 14	7.9	27
Example 15	19	93
Example 16	6.0	27
Example 17	91	1760
Example 18	38	1290
Example 19	47	774
Example 20	20	> 150
Example 21	14	112
Example 22	70	> 150
Example 23	13	61
Example 24	19	> 150
Example 25	29	144
Example 26	30	86
Example 27	83	ND
Example 28	23	> 150

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 29	28	> 150
Example 30	16	119
Example 31	7.4	113
Example 32	47	141
Example 33	33	460
Example 34	4.9	19
Example 35	4.7	8.5
Example 36	4.8	6.2
Example 37	4.9	18
Example 38	23	532
Example 39	14	106
Example 40	3.6	34
Example 41	26	78
Example 42	38	> 150
Example 43	77	502
Example 44	13	443
Example 45	15	20
Example 46	3.7	2.8
Example 47	12	45
Example 48	8.7	40
Example 49	23	58
Example 50	18	26
Example 51	5.5	23
Example 52	3.8	56
Example 53	2.4	5.0
Example 54	4.0	2.4
Example 55	16	372
Example 56	3.8	59
Example 57	65	1250
Example 58	4.2	3.3
Example 59	3.2	6.8

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 60	5.9	46
Example 61	4.4	85
Example 62	20	113
Example 63	4.4	12
Example 64	27	204
Example 65	6.5	51
Example 66	16	80
Example 67	9.4	51
Example 68	2.9	1.8
Example 69	3.1	3.46
Example 70	5.1	9.8
Example 71	3.7	9.9
Example 72	21	229
Example 73	4.3	36
Example 74	6.1	94
Example 75	4.0	6.2
Example 76	24	> 150
Example 77	4.0	84
Example 78	4.4	142
Example 79	5.6	> 150
Example 80	5.5	71
Example 81	21	> 150
Example 82	17	> 150
Example 83	15	122
Example 84	4.9	33
Example 85	6.1	58
Example 86	5.5	60
Example 87	5.1	43
Example 88	3.5	3.9
Example 89	3.5	24
Example 90	6.2	39

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 91	7.0	73
Example 92	20	105
Example 93	5.1	11
Example 94	5.6	19
Example 95	14	115
Example 96	3.9	9.1
Example 97	4.9	15
Example 98	3.8	1.8
Example 99	5.5	13
Example 100	3.9	6.5
Example 101	2.6	31
Example 102	5.0	75
Example 103	3.6	46
Example 104	6.7	65
Example 105	7.3	31
Example 106	5.3	38
Example 107	3.5	6.9
Example 108	2.7	15
Example 109	8.6	89
Example 110	19	329
Example 111	8.4	64
Example 112	42	394
Example 113	3.6	25
Example 114	3.8	28
Example 115	3.4	24
Example 116	21	181
Example 117	2.5	39
Example 118	3.4	43
Example 119	622	> 150
Example 120	2.5	8.2
Example 121	2.9	2.8

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 122	2.6	5.4
Example 123	5.3	8.3
Example 124	12	50
Example 125	8.7	49
Example 126	5.9	5.2
Example 127	3.3	6.4
Example 128	3.5	8.2
Example 129	3.3	39
Example 130	3.1	36
Example 131	4.0	11
Example 132	4.6	32
Example 133	7.0	8.2
Example 134	4.3	2.0
Example 135	8.4	99
Example 136	6.7	24
Example 137	9.4	47
Example 138	12	114
Example 139	5.1	0.84
Example 140	6.6	4.0
Example 141	5.9	82
Example 142	5.5	53
Example 143	5.5	9.3
Example 144	4.8	82
Example 145	4.2	5.5
Example 146	2.8	6.0
Example 147	3.2	2.3
Example 148	3.0	10
Example 149	11	124
Example 150	4.3	16
Example 151	5.3	19
Example 152	6.5	29

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 153	4.2	3.6
Example 154	3.9	5.9
Example 155	3.2	2.9
Example 156	6.8	121
Example 157	21	> 600
Example 158	4.6	3.5
Example 159	8.5	163
Example 160	6.9	86
Example 161	68	> 150
Example 162	33	> 150
Example 163	27	> 150
Example 164	14	> 150
Example 165	18	> 150
Example 166	6.4	58
Example 167	6.5	20
Example 168	5.7	6.6
Example 169	3.3	3.0
Example 170	12	491
Example 171	5.1	63
Example 172	3.4	28
Example 173	55	> 600
Example 174	5.9	407
Example 175	4.2	90
Example 176	78	ND
Example 185	3.7	4.5
Example 186	2.5	4.5
Example 187	3.0	4.6
Example 197	26	> 150
Example 211	3.4	6.9
Example 214	2.7	4.1

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 216	2.4	2.9
Example 232	2.9	11
Example 245	4.2	92
Example 253	5.2	22
Example 392	2.0	14
Example 434	2.9	1.6
Example 446	4.3	14
Example 448	4.0	8.2
Example 474	4.5	2.6
Example 476	1.8	ND
Example 488	5.3	13
Example 489	3.2	68
Example 560	8.8	70
Example 564	4.5	3.3
Example 573	18	43
Example 574	8.8	11
Example 575	6.3	12
Example 577	5.9	4.7
Example 579	4.4	17
Example 591	2.5	7.0
Example 629	7.2	9.0
Example 631	5.2	8.2
Example 633	6.1	22
Example 634	6.9	21
Example 636	11	34
Example 709	4.8	> 150
Example 715	2.2	> 150
Example 721	4.9	> 150
Example 733	1.5	> 150
Example 735	12	> 150

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 775	2.4	6.3
Example 776	2.2	1.1
Example 777	4.1	30
Example 778	3.1	52
Example 779	2.9	24
Example 780	3.5	17
Example 781	2.6	9.7
Example 782	51	112
Example 783	23	> 150
Example 784	5.1	2.9
Example 785	3.3	7.8
Example 786	8.4	24
Example 787	5.0	6.2
Example 788	4.4	14
Example 789	8.9	14
Example 790	3.4	20
Example 791	2.8	54
Example 792	4.6	21
Example 793	4.4	92
Example 794	4.6	70
Example 795	5.3	32
Example 796	6.2	48
Example 797	21	125
Example 798	9.1	76
Example 799	15	> 150
Example 800	2.0	3.2
Example 801	2.7	> 150
Example 802	4.0	99
Example 803	4.6	37
Example 804	3.6	2.2

	IC ₅₀ (nM) Bcl-2 FP	IC ₅₀ (nM) MTT RS4;11
Example 805	3.1	0.12
Example 806	33	> 150
Example 807	3.9	26
Example 808	2.7	1.6
Example 809	3.1	9.8
Example 811	3.8	94

ND: not determined

For partial inhibitors, the percentage fluorescence polarisation inhibition for a given concentration of the test compound is indicated. Accordingly, 72% @3.3 µM means that 72 % fluorescence polarisation inhibition is observed for a concentration of test compound equal to 3.3 µM.

EXAMPLE C: Induction of caspase activity *in vivo*

The ability of the compounds of the invention to activate caspase 3 is evaluated in a xenograft model of RS4;11 leukaemia cells.

1x10⁷ RS4;11 cells are grafted sub-cutaneously into immunosuppressed mice (SCID strain). 25 to 30 days after the graft, the animals are treated orally with the various compounds. Sixteen hours after treatment, the tumour masses are recovered and lysed, and the caspase 3 activity is measured in the tumour lysates.

This enzymatic measurement is carried out by assaying the appearance of a fluorogenic cleavage product (DEVDase activity, Promega). It is expressed in the form of an activation factor corresponding to the ratio between the two caspase activities: the activity for the treated mice divided by the activity for the control mice.

The results show that the compounds of the invention are capable of inducing apoptosis in RS4;11 tumour cells *in vivo*.

EXAMPLE D: Quantification of the cleaved form of caspase 3 *in vivo*

The ability of the compounds of the invention to activate caspase 3 is evaluated in a xenograft model of RS4;11 leukaemia cells.

1x10⁷ RS4;11 cells are grafted sub-cutaneously into immunosuppressed mice (SCID strain). 25 to 30 days after the graft, the animals are treated orally with the various compounds. After treatment, the tumour masses are recovered and lysed, and the cleaved (activated) form of caspase 3 is quantified in the tumour lysates.

The quantification is carried out using the "Meso Scale Discovery (MSD) ELISA platform" test, which specifically assays the cleaved form of caspase 3. It is expressed in the form of an activation factor corresponding to the ratio between the quantity of cleaved caspase 3 in the treated mice divided by the quantity of cleaved caspase 3 in the control mice.

The results show that the compounds of the invention are capable of inducing apoptosis in RS4;11 tumour cells *in vivo*.

EXAMPLE E: Anti-tumour activity *in vivo*

The anti-tumour activity of the compounds of the invention is evaluated in a xenograft model of RS4;11 leukaemia cells.

1x10⁷ RS4;11 cells are grafted sub-cutaneously into immunosuppressed mice (SCID strain). 25 to 30 days after the graft, when the tumour mass has reached about 150 mm³, the mice are treated orally with the various compounds in two different regimes (daily treatment for five days per week for two weeks, or two treatments weekly for two weeks). The tumour mass is measured twice weekly from the start of treatment.


The results obtained therefore show that the compounds of the invention are capable of inducing significant tumour regression during the treatment period.

EXAMPLE F: Pharmaceutical composition: Tablets

	1000 tablets containing a dose of 5 mg of a compound selected from Examples 1 to 811	5 g
	Wheat starch.....	20 g
	Maize starch	20 g
5	Lactose	30 g
	Magnesium stearate.....	2 g
	Silica.....	1 g
	Hydroxypropylcellulose	2 g

CLAIMS

1. Compound of formula (I) :



wherein:

- ◆ Het represents a heteroaryl group,
- ◆ T represents a hydrogen atom, a linear or branched (C₁-C₆)alkyl group optionally substituted by one to three halogen atoms, an alkyl(C₁-C₄)-NR₁R₂ group or an alkyl(C₁-C₄)-OR₆ group,
- ◆ R₁ and R₂ independently of one another represent a hydrogen atom or a linear or branched (C₁-C₆)alkyl group,
or R₁ and R₂ form with the nitrogen atom carrying them a heterocycloalkyl group,
- ◆ R₃ represents a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl group, a (C₃-C₁₀)cycloalkyl-(C₁-C₆)alkyl group wherein the alkyl group may be linear or branched, a heterocycloalkyl group, an aryl group or a heteroaryl group, it being understood that one or more carbon atoms of the groups defined hereinbefore, or carbon atoms of their possible substituents, may be deuterated,
- ◆ R₄ represents an aryl, heteroaryl, cycloalkyl or linear or branched (C₁-C₆)alkyl group, it being understood that one or more carbon atoms of the groups defined

hereinbefore, or carbon atoms of their possible substituents, may be deuterated,

- ◆ R₅ represents a hydrogen atom or a halogen atom, a linear or branched (C₁-C₆)alkyl group, or a linear or branched (C₁-C₆)alkoxy group,
- ◆ R₆ represents a hydrogen atom or a linear or branched (C₁-C₆)alkyl group,
- ◆ R₇ represents a group selected from R'₇, R'₇-CO-, R'₇-O-CO-, NR'₇R''₇-CO-, R'₇-SO₂-, R'₇-NR''₇-SO₂- wherein R'₇ et R''₇ independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl, a heterocycloalkyl, an aryl group, or a heteroaryl,
- ◆ R₈ and R₉ represent, independently of one another, an oxo group or a halogen atom,
- ◆ p and p' are, independently of one another, integers equal to 0, 1, 2, 3 or 4,
- ◆ q and q' are, independently of one another, integers equal to 1, 2 or 3,

it being understood that when compound of formula (I) contains a hydroxy group, this latter group may be optionally substituted by one of the following groups: -PO(OM)(OM'), -PO(OM)(O⁻M₁⁺), -PO(O⁻M₁⁺)(O⁻M₂⁺), -PO(O⁻)(O⁻)M₃²⁺, -PO(OM)(O[CH₂CH₂O]_nCH₃), or -PO(O⁻M₁⁺)(O[CH₂CH₂O]_nCH₃), wherein M and M' independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl or a heterocycloalkyl, both of them being composed of from 5 to 6 ring members, while M₁⁺ and M₂⁺ independently of one another represent a pharmaceutically acceptable monovalent cation, M₃²⁺ represents a pharmaceutically acceptable divalent cation and n is an integer comprised between 1 to 5,

it also being understood that:

- "aryl" means a phenyl, naphthyl, biphenyl or indenyl group,
- "heteroaryl" means any mono- or bi-cyclic group composed of from 5 to 10 ring members, having at least one aromatic moiety and containing from 1 to 4 hetero atoms selected from oxygen, sulphur and nitrogen (including quaternary nitrogens),
- "cycloalkyl" means any mono- or bi-cyclic non-aromatic carbocyclic group containing from 3 to 10 ring members, which may include fused, bridged or spiro ring systems,

- "heterocycloalkyl" means any mono- or bi-cyclic non-aromatic carbocyclic group, composed of from 3 to 10 ring members, and containing from one to 3 hetero atoms selected from oxygen, sulphur, SO, SO₂ and nitrogen, it being understood that bicyclic group may be fused or spiro type,


it being possible for the aryl, heteroaryl, cycloalkyl and heterocycloalkyl groups so defined and the alkyl, alkenyl, alkynyl, alkoxy, to be substituted by from 1 to 3 groups selected from: linear or branched (C₁-C₆)alkyl ; linear or branched (C₂-C₆)alkenyl group; linear or branched (C₂-C₆)alkynyl group; (C₃-C₆)spiro; linear or branched (C₁-C₆)alkoxy; (C₁-C₆)alkyl-S-; hydroxyl; oxo (or *N*-oxide where appropriate); nitro; cyano; -COOR'; -OCOR'; -NR'R"; R'CONR''-; NR'R''CO-; linear or branched (C₁-C₆)polyhaloalkyl; trifluoromethoxy; (C₁-C₆)alkylsulphonyl; halogen; aryl; heteroaryl; aryloxy; arylthio; cycloalkyl or heterocycloalkyl, it being understood that R' and R" independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or an unsubstituted aryl group,

wherein said linear or branched (C₁-C₆)alkyl groups, said linear or branched (C₂-C₆)alkenyl group, said linear or branched (C₂-C₆)alkynyl group, said linear or branched (C₁-C₆)alkoxy, said aryl, said heteroaryl, said aryloxy, said arylthio, said cycloalkyl, and said heterocycloalkyl are optionally substituted by from 1 to 3 groups selected from the substituents of the aryl, heteroaryl, cycloalkyl and heterocycloalkyl groups and the alkyl, alkenyl, alkynyl, alkoxy defined above, namely:

linear or branched (C₁-C₆)alkyl; linear or branched (C₂-C₆)alkenyl group; linear or branched (C₂-C₆)alkynyl group; (C₃-C₆)spiro; linear or branched (C₁-C₆)alkoxy; (C₁-C₆)alkyl-S-; hydroxyl; oxo (or *N*-oxide where appropriate); nitro; cyano; -COOR'; -OCOR'; -NR'R"; R'CONR''-; NR'R''CO-; linear or branched (C₁-C₆)polyhaloalkyl; trifluoromethoxy; (C₁-C₆)alkylsulphonyl; halogen; aryl; heteroaryl; aryloxy; arylthio; cycloalkyl or heterocycloalkyl, it being understood that R' and R" independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or an unsubstituted aryl group,

its enantiomers and diastereoisomers, and addition salts thereof with a pharmaceutically acceptable acid or base.

2. Compound of formula (IA) according to claim 1:



wherein:

- ◆ W represents a C-A₃ group or a nitrogen atom,
- ◆ X, Y and Z represent a carbon atom or a nitrogen atom, it being understood that only one of them represent a nitrogen atom, while the two others represent carbon atoms,
- ◆ A₁, A₂ and A₃ independently of one another represent a hydrogen atom or a halogen atom, a linear or branched (C₁-C₆)polyhaloalkyl group, a linear or branched (C₁-C₆)alkyl group or a cycloalkyl group,
or A₃ represents a hydrogen atom (when W represents a C-A₃ group) while A₁ and A₂ form together with the atoms carrying them an optionally substituted aromatic or non-aromatic ring Cy, composed of 5, 6 or 7 ring members, which may contain from one to 4 hetero atoms selected independently from oxygen, sulphur and nitrogen, it being understood that the nitrogen in question may be substituted by a group representing a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or a group -C(O)-O-Alk wherein Alk is a linear or branched (C₁-C₆)alkyl group,
or W represents a nitrogen atom while A₁ and A₂ form together with the atoms carrying them an optionally substituted aromatic or non-aromatic ring Cy,

composed of 5, 6 or 7 ring members, which may contain from one to 4 hetero atoms selected independently from oxygen, sulphur and nitrogen, it being understood that the nitrogen in question may be substituted by a group representing a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or a group -C(O)-O-Alk wherein Alk is a linear or branched (C₁-C₆)alkyl group,

- ◆ T represents a hydrogen atom, a linear or branched (C₁-C₆)alkyl group optionally substituted by one to three halogen atoms, an alkyl(C₁-C₄)-NR₁R₂ group or an alkyl(C₁-C₄)-OR₆ group,
- ◆ R₁ and R₂ independently of one another represent a hydrogen atom or a linear or branched (C₁-C₆)alkyl group,
or R₁ and R₂ form with the nitrogen atom carrying them a heterocycloalkyl,
- ◆ R₃ represents a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl group, a (C₃-C₁₀)cycloalkyl-(C₁-C₆)alkyl group wherein the alkyl group may be linear or branched, a heterocycloalkyl group, an aryl group or a heteroaryl group, it being understood that one or more carbon atoms of the groups defined hereinbefore, or carbon atoms of their possible substituents, may be deuterated,
- ◆ R₄ represents an aryl, heteroaryl, cycloalkyl or linear or branched (C₁-C₆)alkyl group, it being understood that one or more carbon atoms of the groups defined hereinbefore, or carbon atoms of their possible substituents, may be deuterated,
- ◆ R₅ represents a hydrogen atom or a halogen atom, a linear or branched (C₁-C₆)alkyl group, or a linear or branched (C₁-C₆)alkoxy group,
- ◆ R₆ represents a hydrogen atom or a linear or branched (C₁-C₆)alkyl group,
- ◆ R₇ represents a group selected from R'₇, R'₇-CO-, R'₇-O-CO-, NR'₇R''₇-CO-, R'₇-SO₂-, R'₇-NR''₇-SO₂- wherein R'₇ et R''₇ independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl, a heterocycloalkyl, an aryl group, or a heteroaryl,
- ◆ R₈ and R₉ represent, independently of one another, an oxo group or a halogen atom,
- ◆ p and p' are, independently of one another, integers equal to 0, 1, 2, 3 or 4,
- ◆ q and q' are, independently of one another, integers equal to 1, 2 or 3,

it being understood that when compound of formula (I) contains a hydroxy group, this latter group may be optionally substituted by one of the following groups: $-\text{PO}(\text{OM})(\text{OM}')$, $-\text{PO}(\text{OM})(\text{O}^-\text{M}_1^+)$, $-\text{PO}(\text{O}^-\text{M}_1^+)(\text{O}^-\text{M}_2^+)$, $-\text{PO}(\text{O}^-)(\text{O}^-)\text{M}_3^{2+}$, $-\text{PO}(\text{OM})(\text{O}[\text{CH}_2\text{CH}_2\text{O}]_n\text{CH}_3)$, or $-\text{PO}(\text{O}^-\text{M}_1^+)(\text{O}[\text{CH}_2\text{CH}_2\text{O}]_n\text{CH}_3)$, wherein M and M' independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl or a heterocycloalkyl, both of them being composed of from 5 to 6 ring members, while M₁⁺ and M₂⁺ independently of one another represent a pharmaceutically acceptable monovalent cation, M₃²⁺ represents a pharmaceutically acceptable divalent cation and n is an integer comprised between 1 to 5,

it also being understood that:

- "aryl" means a phenyl, naphthyl, biphenyl or indenyl group,
- "heteroaryl" means any mono- or bi-cyclic group composed of from 5 to 10 ring members, having at least one aromatic moiety and containing from 1 to 4 hetero atoms selected from oxygen, sulphur and nitrogen (including quaternary nitrogens),
- "cycloalkyl" means any mono- or bi-cyclic non-aromatic carbocyclic group containing from 3 to 10 ring members, which may include fused, bridged or spiro ring systems,
- "heterocycloalkyl" means any mono- or bi-cyclic non-aromatic carbocyclic group, composed of from 3 to 10 ring members, and containing from one to 3 hetero atoms selected from oxygen, sulphur, SO, SO₂ and nitrogen, it being understood that bicyclic group may be fused or spiro type,

it being possible for the aryl, heteroaryl, cycloalkyl and heterocycloalkyl groups so defined and the alkyl, alkenyl, alkynyl, alkoxy, to be substituted by from 1 to 3 groups selected from: linear or branched (C₁-C₆)alkyl; linear or branched (C₂-C₆)alkenyl group; linear or branched (C₂-C₆)alkynyl group; (C₃-C₆)spiro; optionally substituted linear or branched (C₁-C₆)alkoxy;

(C₁-C₆)alkyl-S-; hydroxyl; oxo (or N-oxide where appropriate); nitro; cyano; -COOR'; -OCOR'; -NR'R''; R'CONR''-; NR'R''CO-; linear or branched (C₁-C₆)polyhaloalkyl; trifluoromethoxy; (C₁-C₆)alkylsulphonyl; halogen; aryl; heteroaryl; aryloxy; arylthio;

cycloalkyl or heterocycloalkyl, it being understood that R' and R" independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or an unsubstituted aryl group,


wherein said linear or branched (C₁-C₆)alkyl groups, said linear or branched (C₂-C₆)alkenyl group, said linear or branched (C₂-C₆)alkynyl group, said linear or branched (C₁-C₆)alkoxy, said aryl, said heteroaryl, said aryloxy, said arylthio, said cycloalkyl, and said heterocycloalkyl are optionally substituted by from 1 to 3 groups selected from the substituents of the aryl, heteroaryl, cycloalkyl and heterocycloalkyl groups and the alkyl, alkenyl, alkynyl, alkoxy defined above, namely:

linear or branched (C₁-C₆)alkyl; linear or branched (C₂-C₆)alkenyl group; linear or branched (C₂-C₆)alkynyl group; (C₃-C₆)spiro; linear or branched (C₁-C₆)alkoxy; (C₁-C₆)alkyl-S-; hydroxyl; oxo (or *N*-oxide where appropriate); nitro; cyano; -COOR'; -OCOR'; -NR'R"; R'CONR''-; NR'R''CO-; linear or branched (C₁-C₆)polyhaloalkyl; trifluoromethoxy; (C₁-C₆)alkylsulphonyl; halogen; aryl; heteroaryl; aryloxy; arylthio; cycloalkyl or heterocycloalkyl, it being understood that R' and R" independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group or an unsubstituted aryl group,

it being possible for the Cy moiety defined in formula (I) to be substituted by from 1 to 3 groups selected from linear or branched (C₁-C₆)alkyl, linear or branched (C₁-C₆)polyhaloalkyl, hydroxy, linear or branched (C₁-C₆)alkoxy, COOH, NR₁'R₁" and halogen, it being understood that R₁' and R₁" have the same definitions than the groups R' and R" mentioned hereinbefore,


its enantiomers and diastereoisomers, and addition salts thereof with a pharmaceutically acceptable acid or base.

3. Compound of formula (IA-1) according to claim 1 or 2:



wherein X, Y, W, A₁, A₂, R₃, R₄, R₅, R₇, R₈, R₉, T, p, p', q and q' are as defined in claim 2, its enantiomers and diastereoisomers, and addition salts thereof with a pharmaceutically acceptable acid or base.

4. Compound of formula (IA-2) according to claim 1 to 3:



wherein A₁, A₂, R₃, R₄, R₇ and T are as defined in claim 2,

its enantiomers and diastereoisomers, and addition salts thereof with a pharmaceutically acceptable acid or base.

5. Compound of formula (I) according to any one of claims 1 to 4 wherein R₄ represents a phenyl substituted at the *para* position by a group of formula -OPO(OM)(OM'), -OPO(OM)(O⁻M₁⁺), -OPO(O⁻M₁⁺)(O⁻M₂⁺), -OPO(O⁻)(O⁻)M₃²⁺, -OPO(OM)(O[CH₂CH₂O]_nCH₃), or -OPO(O⁻M₁⁺)(O[CH₂CH₂O]_nCH₃), wherein M and M' independently of one another represent a hydrogen atom, a linear or branched (C₁-C₆)alkyl group, a linear or branched (C₂-C₆)alkenyl group, a linear or branched (C₂-C₆)alkynyl group, a cycloalkyl or a heterocycloalkyl, both of them being composed of from 5 to 6 ring members, while M₁⁺ and M₂⁺ independently of one another represent a pharmaceutically acceptable monovalent cation, M₃²⁺ represents a pharmaceutically acceptable divalent cation and n is an integer comprised between 1 to 5, it being understood that the phenyl group may be optionally substituted by one or more halogen atoms.

6. Compound of formula (I) according to any one of claims 1 to 5 wherein Het represents one of the following groups: 5,6,7,8-tetrahydroindolizine, indolizine or 1,2-dimethyl-1*H*-pyrrole.

7. Compound of formula (I) according to any one of claims 1 to 6 wherein q=1 and q'=1.

8. Compound of formula (I) according to any one of claims 1 to 6 wherein q=2 and q'=1.

9. Compound of formula (I) according to any one of claims 1 to 8 wherein T represents a hydrogen atom, a methyl group, a (morpholin-4-yl)methyl group, a 3-(morpholin-4-yl)propyl group, a dimethylaminomethyl group, or a (4-methylpiperazin-1-yl)methyl group.

10. Compound of formula (I) according to any one of claims 1 to 9 wherein R₃ represents a linear or branched (C₁-C₆)alkyl group, an aryl group or a heteroaryl group.

11. Compound of formula (I) according to claim 1 to 10 wherein R₃ represents a group

selected from phenyl, methyl, ethyl, propyl, butyl, 1-methyl-1*H*-pyrrolo[2,3-*b*]pyridine or 5-cyano-1,2-dimethyl-1*H*-pyrrole.

12. Compound of formula (I) according to any one of claims 1 to 11 wherein R₄ represents a linear or branched (C₁-C₆)alkyl group or an aryl group.

13. Compound of formula (I) according to claim 1 to 12 wherein R₄ represents a phenyl or 4-hydroxyphenyl group.

14. Compound of formula (I) according to any one of claims 1 to 13 wherein R₇ represents a group R'₇-CO- or R'₇-O-CO-.

15. Compound of formula (I) according to any one of claims 1 to 14 wherein R'₇ represents an aryl, a cycloalkyl or an alkyl.

16. Compound of formula (I) according to claim 15 wherein R'₇ represents an naphthalene, phenyl or indole group.

17. Compound of formula (I) according to claim 16 wherein the phenyl group is substituted by a group selected from: an alkyl, cyano, alkynyl, halogen, alkoxy, or -NR'R''.

18. Compound of formula (I) according to claim 16 wherein the phenyl group is substituted by a group selected from : methyl, ethyl, methoxy, chloro, bromo, cyano, 2-dimethylaminoethylamino, ethynyl, 2-dimethylaminoethoxy, 2-(dimethylamino)ethyl(methyl)amino, or dimethylcarbamoylethyl.

19. Compound of formula (I) according to any one of claims 1 to 15 wherein p=p'=0.

20. Compound of formula (I) according to any one of claims 1 to 15 which is

- *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide;
- Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-

- yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3R)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - Phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride;
 - Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride;
 - 4-Methylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate hydrochloride;
 - 2-Methylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 4-Methoxyphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 4-Chlorophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 4-Ethylphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 4-Cyanophenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 2-Methoxyphenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3S)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;

- 4-Methylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-Chlorophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-Cyanophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-{[2-(Dimethylamino)ethyl]amino}phenyl 6-{1-[(4-hydroxyphenyl)(methyl)carbamoyl]-5,6,7,8-tetrahydroindolizin-3-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- Phenyl 6-{4-[(4-hydroxyphenyl)(propyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- Phenyl 6-{4-[butyl(4-hydroxyphenyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- Phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-[(4-methylpiperazin-1-yl)methyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- *N*-Butyl-*N*-(4-hydroxyphenyl)-1,2-dimethyl-5-{7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-(2-phenylacetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-1*H*-pyrrole-3-carboxamide;
- 3-[2-(Dimethylamino)ethoxy]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-{[2-(Dimethylamino)ethyl](methyl)amino}phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-

- tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
- 4-Ethynylphenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - Naphthalen-2-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 1*H*-Indol-5-yl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 3-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 4-Bromophenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*S*)-3-(morpholin-4-ylmethyl)-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - (1*r*,4*r*)-4-[[*tert*-Butoxy)carbonyl]amino}cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate;
 - 4-[2-(Dimethylcarbamoyl)ethyl]phenyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate; or
 - Cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-[3-(morpholin-4-yl)propyl]-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate.


21. Compound which is selected from the following list:

- *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[4-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide,
- *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-

tetrahydroisoquinoline-2-carbonyl]-2-(2-{4-[(4-methylpiperazin-1-yl)methyl]phenyl}acetyl)-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide,


- *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[4-(morpholin-4-ylmethyl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide,
- 3-[2-(2-{4-[(Dimethylamino)methyl]phenyl}acetyl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide,
- 3-[2-(2-{4-[2-(Dimethylamino)ethoxy]phenyl}acetyl)-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinolin-6-yl]-*N*-(4-hydroxyphenyl)-*N*-methyl-5,6,7,8-tetrahydroindolizine-1-carboxamide,
- *N*-(4-Hydroxyphenyl)-*N*-methyl-3-{7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-2-{2-[3-(4-methylpiperazin-1-yl)phenyl]acetyl}-1,2,3,4-tetrahydroisoquinolin-6-yl}-5,6,7,8-tetrahydroindolizine-1-carboxamide,
- (1*r*,4*r*)-4-[[(*tert*-Butoxy)carbonyl]amino}cyclohexyl 6-{4-[(4-hydroxyphenyl)(methyl)carbamoyl]-1,5-dimethyl-1*H*-pyrrol-2-yl}-7-[(3*R*)-3-methyl-1,2,3,4-tetrahydroisoquinoline-2-carbonyl]-1,2,3,4-tetrahydroisoquinoline-2-carboxylate.

22. Process for the preparation of a compound of formula (I) according to claim 1 characterised in that there is used as starting material the compound of formula (II):




wherein Het, R₃, R₄, R₈, R₉, p, p', q and q' are as defined for formula (I) and Alk represents a linear or branched (C₁-C₆) alkyl group,

the ester function of which compound of formula (II) is hydrolysed to yield the corresponding carboxylic acid which is then subjected to peptide coupling with a compound of formula (III) :



wherein R₅ and T are as defined for formula (I),

to yield the compound of formula (IV) :




wherein Het, R₃, R₄, R₅, R₈, R₉, T, p, p', q and q' are as defined for formula (I),

compound of formula (IV) which is deprotected and subjected to an acylation or a sulfonylation reaction, and which can optionally be subjected to the action of a pyrophosphate or a phosphonate derivative in basic conditions, to yield the compound of formula (I),

which compound of formula (I) may be purified according to a conventional separation technique, which is converted, if desired, into its addition salts with a pharmaceutically acceptable acid or base and which is optionally separated into its isomers according to a conventional separation technique,


it being understood that at any moment considered appropriate during the course of the process described above, some groups of the starting reagents or of the synthesis intermediates, can be protected and subsequently deprotected, as required by the synthesis.

23. Process according for the preparation of a compound of formula (IA-1) according to claim 3 characterised in that there is used as starting material the compound of formula (V):




wherein X, Y, W, A₁ and A₂ are as defined for formula (IA) and Alk represents a linear or branched (C₁-C₆) alkyl group,

the ester function of which compound of formula (V) is hydrolysed to yield the carboxylic acid or the corresponding carboxylate, which may be converted into an acid derivative such as acyl chloride or the corresponding anhydride before being coupled with an amine NHR₃R₄ wherein R₃ and R₄ have the same meanings as for formula (IA), to yield the compound of formula (VI):




wherein X, Y, W, A₁, A₂, R₃ and R₄ are as defined for formula (IA),

compound of formula (VI) which is then halogenated and further converted into the corresponding borohydride derivative of formula (VII) :




wherein X, Y, W, A₁, A₂, R₃ and R₄ are as defined for formula (IA), and R_{B1} and R_{B2} represent a hydrogen, a linear or branched (C₁-C₆) alkyl group, or R_{B1} and R_{B2} form with the oxygen atoms carrying them an optionally methylated ring,

which compound of formula (VII) is further subjected to coupling with a compound of formula (VIII):




wherein R₈, R₉, p, p', q and q' are as defined for formula (IA), and Alk' represents a linear or branched (C₁-C₆) alkyl group, and L represents a leaving group selected from halogen or sulfonate,

to yield the compound of formula (IIA-1) :




wherein X, Y, W, A₁, A₂, R₃, R₄, R₈, R₉, p, p', q, and q' are as defined for formula (IA) and Alk' is as defined previously,

the ester function of which compound of formula (IIA-1) is hydrolysed to yield the corresponding carboxylic acid which is then subjected to peptide coupling with a compound of formula (III) :




wherein R₅ and T are as defined for formula (IA),

to yield the compound of formula (IVA-1) :



wherein X, Y, W, A₁, A₂, R₃, R₄, R₅, R₈, R₉, T, p, p', q, q' are as defined for formula (IA),

compound of formula (IVA-1) which is deprotected and subjected to an acylation or a sulfonylation reaction, and which can optionally be subjected to the action of a pyrophosphate or a phosphonate derivative in basic conditions, to yield the compound of formula (IA-1):



wherein X, Y, W, A₁, A₂, R₃, R₄, R₅, R₇, R₈, R₉, T, p, p', q and q' are as defined for formula (IA),

which compound of formula (IA-1) may be purified according to a conventional separation technique, which is converted, if desired, into its addition salts with a pharmaceutically acceptable acid or base and which is optionally separated into its isomers according to a conventional separation technique,

it being understood that at any moment considered appropriate during the course of the process described above, some groups of the starting reagents or of the synthesis intermediates, can be protected and subsequently deprotected, as required by the synthesis.

24. Process according to claim 22 or 23 for the preparation of a compound of formula (I) wherein one of the groups R₃ and R₄ is substituted by a hydroxy function, characterised in that the amine NHR₃R₄ is subjected beforehand to a protection reaction of the hydroxy function prior to coupling with the carboxylic acid formed from the compound of formula (V), or with a corresponding acid derivative thereof, the resulting protected function subsequently undergoes a deprotection reaction at the last stage of the process.

25. Pharmaceutical composition comprising a compound of formula (I) according to any one of claims 1 to 21 or an addition salt thereof with a pharmaceutically acceptable

acid or base in combination with one or more pharmaceutically acceptable excipients.

26. Pharmaceutical composition according to claim 25 for use as pro-apoptotic agents.

27. Pharmaceutical composition according to claim 25 for use in the treatment of cancers and of auto-immune and immune system diseases.

28. Pharmaceutical composition according to claim 25 for use in the treatment of cancers of the bladder, brain, breast and uterus, chronic lymphoid leukaemias, cancer of the colon, oesophagus and liver, lymphoblastic leukaemias, non-Hodgkin lymphomas, melanomas, malignant haemopathies, myelomas, ovarian cancer, non-small-cell lung cancer, prostate cancer and small-cell lung cancer.

29. Use of a pharmaceutical composition according to claim 25 in the manufacture of medicaments for use as pro-apoptotic agents.

30. Use of a pharmaceutical composition according to claim 25 in the manufacture of medicaments for use in the treatment of cancers and of auto-immune and immune system diseases.

31. Use of a pharmaceutical composition according to claim 25 in the manufacture of medicaments for use in the treatment of cancers of the bladder, brain, breast and uterus, chronic lymphoid leukaemias, cancer of the colon, oesophagus and liver, lymphoblastic leukaemias, non-Hodgkin lymphomas, melanomas, malignant haemopathies, myelomas, ovarian cancer, non-small-cell lung cancer, prostate cancer and small-cell lung cancer.

32. Compound of formula (I) according to any one of claims 1 to 21, or an addition salt thereof with a pharmaceutically acceptable acid or base, for use in the treatment of cancers of the bladder, brain, breast and uterus, chronic lymphoid leukaemias, cancer of the colon, oesophagus and liver, lymphoblastic leukaemias, non-Hodgkin lymphomas, melanomas, malignant haemopathies, myelomas, ovarian cancer, non-small-cell lung cancer, prostate cancer and small-cell lung cancer.

33. Use of a compound of formula (I) according to any one of claims 1 to 21, or an addition salt thereof with a pharmaceutically acceptable acid or base, in the manufacture of medicaments for use in the treatment of cancers of the bladder, brain, breast and uterus, chronic lymphoid leukaemias, cancer of the colon, oesophagus and liver, lymphoblastic leukaemias, non-Hodgkin lymphomas, melanomas, malignant haemopathies, myelomas, ovarian cancer, non-small-cell lung cancer, prostate cancer and small-cell lung cancer.

34. Combination of a compound of formula (I) according to any one of claims 1 to 21 with an anti-cancer agent selected from genotoxic agents, mitotic poisons, anti-metabolites, proteasome inhibitors, kinase inhibitors and antibodies.

35. Pharmaceutical composition comprising a combination according to claim 34 in combination with one or more pharmaceutically acceptable excipients.

36. Combination according to claim 34 for use in the treatment of cancers.

37. Use of a combination according to claim 34 in the manufacture of medicaments for use in the treatment of cancers.

38. Compound of formula (I) according to any one of claims 1 to 21 for use in combination with radiotherapy in the treatment of cancers.