
PHARMACEUTICAL COMPOSITIONS CONTAINING 3-AMINOAZETIDINE DERIVATIVES, NOVEL DERIVATIVES AND PREPARATION THEREOF

In an aspect of the present invention, there is provided a pharmaceutical compositions containing, as active ingredient, at least one compound of formula:



(I)

or one of its pharmaceutically acceptable salts, to the novel derivatives of formula (I), to their pharmaceutically acceptable salts and to their preparation.

in which

R_1 represents a radical $-NHCOR_4$ or $-N(R_5)-Y-R_6$,

Y is CO or SO_2,

R_2 and R_3, which are identical or different, represent either an aromatic chosen from phenyl, naphthyl and indenyl, these aromatics being unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, formyl, hydroxyl, trifluoromethyl, trifluoromethoxy, -CO-alk, cyano, -COOH, -COOalk, $-CONR_7R_8$, $-CO-NH-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkyl-

sulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, hydroxyalkyl or -alk-NR_7R_8 radicals; or a heteroaromatic chosen from benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzothienyl, pyrimidinyl, furyl, imidazolyl, isochromanyl, isoquinolyl, pyrrolyl, pyridyl, quinolyl, 1,2,3,4-tetrahydroisoquinolyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, cyano, -COOH, -COOalk, -CO-NH-NR_9R_{10}, -$CONR_7R_8$, -alk-NR_9R_{10}, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl or hydroxyalkyl radical,

R_4 represents a radical -alk-SO_2-R_{11}, -alk-SO_2-CH=CH-R_{11}, Het substituted with -SO_2-R_{11} or phenyl substituted with -SO_2-R_{11} or -alk-SO_2-R_{11},

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a phenylalkyl, Het or Ar radical,

R_7 and R_8, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_7 and R_8 together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl

radicals,

R_9 and R_{10}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_9 and R_{10} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

R_{11} represents an alkyl, Ar or Het radical,

Ar represents a phenyl, naphthyl or indenyl radical, these radicals being optionally substituted with one or more halogen atoms or alkyl, alkoxy, cyano, -CO-alk, -COOH, -COOalk, $-CONR_{12}R_{13}$, $-CO-NH-NR_{14}R_{15}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, $-alk-NR_{14}R_{15}$, $-NR_{14}R_{15}$, alkylthioalkyl, formyl, hydroxyl, hydroxyalkyl, Het, -O-alk-NH-cycloalkyl, OCF_3, CF_3, -NH-CO-alk, $-SO_2NH_2$, $-NH-COCH_3$, -NH-COOalk or Het radicals or alternatively, on 2 adjacent carbon atoms, with a dioxymethylene,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl,

oxo, hydroxyl, OCF_3 or CF_3 radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form,

R_{12} and R_{13}, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_{12} and R_{13} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

alk represents an alkyl or alkylene radical,

the alkyl and alkylene radicals and portions and the alkoxy radicals and portions are in the form of a straight or branched chain and contain 1 to 6 carbon

atoms and the cycloalkyl radicals contain 3 to 10 carbon atoms,

an optical isomer of such a compound or one of its pharmaceutically acceptable salts.

The compound of formula (I) for which R_2 and R_3 represent phenyl radicals, R_1 represents a radical $-N(R_5)-Y-R_6$, Y is SO_2, R_5 represents a methyl radical and R_6 represents a phenyl radical is described as a synthesis intermediate in Patent WO 99/01451. The other compounds and their pharmaceutically acceptable salts are novel and as such form parts of the invention.

In formula (I)

R_1 represents a radical $-NHCOR_4$ or $-N(R_5)-Y-R_6$,

Y is CO or SO_2,

R_2 and R_3, which are identical or different, represent either an aromatic chosen from phenyl, naphthyl and indenyl, these aromatics being unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, formyl, hydroxyl, trifluoromethyl, trifluoromethoxy, -CO-alk, cyano, -COOH, -COOalk, $-CONR_7R_8$, $-CO-NH-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, hydroxyalkyl or $-alk-NR_7R_8$ radicals; or a heteroaromatic chosen from benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzothienyl, pyrimidinyl, furyl, imidazolyl, isochromanyl, isoquinolyl, pyrrolyl, pyridyl, quinolyl, 1,2,3,4-tetrahydro-isoquinolyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, cyano, -COOH, -COOalk, $-CO-NH-NR_9R_{10}$, $-CONR_7R_8$, $-alk-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl or hydroxyalkyl radical,

R_4 represents a radical $-alk-SO_2-R_{11}$, $-alk-SO_2-CH=CH-R_{11}$, Het substituted with $-SO_2-R_{11}$ or phenyl substituted with $-SO_2-R_{11}$ or $-alk-SO_2-R_{11}$,

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a phenylalkyl, Het or Ar radical,

R_7 and R_8, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_7 and R_8 together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_9 and R_{10}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_9 and R_{10} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

R_{11} represents an alkyl, Ar or Het radical,

Ar represents a phenyl, naphthyl or indenyl radical, these radicals being optionally substituted with one or more halogen atoms or alkyl, alkoxy, cyano, -CO-alk, -COOH, -COOalk, $-CONR_{12}R_{13}$, $-CO-NH-NR_{14}R_{15}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, $-alk-NR_{14}R_{15}$, $-NR_{14}R_{15}$, alkylthioalkyl, formyl, hydroxyl, hydroxyalkyl, Het, -O-alk-NH-cycloalkyl, OCF_3, CF_3, -NH-CO-alk, $-SO_2NH_2$,

-NH-$COCH_3$, -NH-COOalk or Het radicals, or alternatively, on 2 adjacent carbon atoms, with a dioxymethylene,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo, hydroxyl, OCF_3 or CF_3 radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form,

R_{12} and R_{13}, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_{12} and R_{13} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk,

-CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

alk represents an alkyl or alkylene radical.

In the preceding definitions and in those which follow, unless otherwise stated, the alkyl and alkylene radicals and portions and the alkoxy radicals and portions are in the form of a straight or branched chain and contain 1 to 6 carbon atoms and the cycloalkyl radicals contain 3 to 10 carbon atoms.

Among the alkyl radicals, there may be mentioned the methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, pentyl and hexyl radicals. Among the alkoxy radicals, there may be mentioned the methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy and pentyloxy radicals.

Among the cycloalkyl radicals, there may be mentioned the cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl radicals.


The term halogen comprises chlorine, fluorine, bromine and iodine.

Among the heterocycles represented by Het, the following heterocycles may be mentioned: benzimidazole, benzoxazole, benzothiazole, benzothiophene, cinnoline, thiophene, quinazoline, quinoxaline, quinoline, pyrazole, pyrrole, pyridine, imidazole, indole, isoquinoline, pyrimidine, thiazole, thiadiazole, piperidine, piperazine, pyrrolidine,

triazole, furan, tetrahydroisoquinoline, tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo, hydroxyl, OCF_3 or CF_3 radicals.

The compounds of formula (I) may be provided in the form of enantiomers and of diastereoisomers. These optical isomers and mixtures thereof also form part of the invention.

In another aspect of the invention, there is provided a compound of formula:



(I)

in which

R_1 represents a radical $-NHCOR_4$ or $-N(R_5)-Y-R_6$,

Y is CO or SO_2,

R_2 and R_3, which are identical or different, represent either an aromatic chosen from phenyl, naphthyl and indenyl, these aromatics being unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, formyl, hydroxyl, trifluoromethyl, trifluoromethoxy, -CO-alk, cyano, -COOH, -COOalk, $-CONR_7R_8$, $-CO-NH-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkyl-

sulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, hydroxyalkyl or -alk-NR_7R_8 radicals; or a heteroaromatic chosen from benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzothienyl, pyrimidinyl, furyl, imidazolyl, isochromanyl, isoquinolyl, pyrrolyl, pyridyl, quinolyl, 1,2,3,4-tetrahydro-isoquinolyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, cyano, -COOH, -COOalk, -CO-NH-NR_9R_{10}, -$CONR_7R_8$, -alk-NR_9R_{10}, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl or hydroxyalkyl radical,

R_4 represents a radical -alk-SO_2-R_{11}, -alk-SO_2-CH=CH-R_{11}, Het substituted with -SO_2-R_{11} or phenyl substituted with -SO_2-R_{11} or -alk-SO_2-R_{11},

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a phenylalkyl, Het or Ar radical,

R_7 and R_8, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_7 and R_8 together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_9 and R_{10}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_9 and R_{10} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

R_{11} represents an alkyl, Ar or Het radical,

Ar represents a phenyl, naphthyl or indenyl radical, these radicals being optionally substituted with one or more halogen atoms or alkyl, alkoxy, cyano, -CO-alk, -COOH, -COOalk, $-CONR_{12}R_{13}$, $-CO-NH-NR_{14}R_{15}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, $-alk-NR_{14}R_{15}$, $-NR_{14}R_{15}$, alkylthioalkyl, formyl, hydroxyl, hydroxyalkyl, Het, -O-alk-NH-cycloalkyl, OCF_3, CF_3, -NH-CO-alk, $-SO_2NH_2$, $-NH-COCH_3$, -NH-COOalk or Het radicals, or alternatively, on 2 adjacent carbon atoms, with a dioxymethylene,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo, hydroxyl, OCF_3 or CF_3 radicals, the nitrogen-

containing heterocycles being optionally in their N-oxidized form,

R_{12} and R_{13}, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_{12} and R_{13} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

alk represents an alkyl or alkylene radical,

the alkyl and alkylene radicals and portions and the alkoxy radicals and portions are in the form of a straight or branched chain and contain 1 to 6 carbon atoms and the cycloalkyl radicals contain 3 to 10 carbon atoms,

their optical isomers and their pharmaceutically acceptable salts,

with the exception of the compound for which for which R_2 and R_3 represent phenyl radicals, R_1 represents a radical $-N(R_5)-Y-R_6$, Y is SO_2, R_5 represents a methyl radical and R_6 represents a phenyl radical.

Preferably, the compounds of formula (I) are those for which

R_1 represents a radical $-N(R_5)-Y-R_6$,

Y is SO_2,

R_2 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy, cyano, $-CONR_7R_8$, hydroxyalkyl or $-alk-NR_7R_8$ radicals; or a heteroaromatic chosen from the pyridyl, pyrimidyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, $-CONR_7R_8$, $-alk-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl or hydroxyalkyl radical,

R_3 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy, cyano, $-CONR_7R_8$, hydroxyalkyl or $-alk-NR_7R_8$ radicals; or a

heteroaromatic chosen from the pyridyl, pyrimidyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, $-CONR_7R_8$, $-alk-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl or hydroxyalkyl radical,

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a naphthyl, phenylalkyl, Het or phenyl radical optionally substituted with one or more halogen atoms or alkyl, alkoxy, cyano, -CO-alk, COOalk, $-CONR_{12}R_{13}$, $-alk-NR_{14}R_{15}$, $-NR_{14}R_{15}$, hydroxyl, hydroxyalkyl, Het, OCF_3, CF_3, -NH-CO-alk, $-SO_2NH_2$ or -NH-COOalk radicals, or alternatively, on 2 adjacent carbon atoms, with dioxymethylene,

R_7 and R_8, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_7 and R_8 together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_9 and R_{10}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_9 and R_{10} together form with the nitrogen atom to which they are attached a 3- to 10-membered

saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, oxo or $-CO-NH_2$ radicals,

R_{12} and R_{13}, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_{12} and R_{13} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, oxo, hydroxyalkyl or $-CO-NH_2$ radicals,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl,

oxo or hydroxyl radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form and, preferably, Het represents a heterocycle chosen from the following heterocycles: benzimidazole, benzoxazole, benzothiazole, benzothiophene, thiophene, quinazoline, quinoxaline, quinoline, pyrrole, pyridine, imidazole, indole, isoquinoline, pyrimidine, thiazole, thiadiazole, furan, tetrahydroisoquinoline and tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, oxo, hydroxyl, OCF_3 or CF_3 radicals.

Still more preferably, the compounds of formula (I) are chosen from the following compounds:

R_1 represents a radical $-N(R_5)-Y-R_6$,

Y is SO_2,

R_2 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy or hydroxyalkyl radicals; or a heteroaromatic chosen from pyridyl and pyrimidyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl or trifluoromethoxy radical,

R_3 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy or hydroxyalkyl radicals; or a heteroaromatic chosen from

pyridyl and pyrimidyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl or trifluoromethoxy radical,

R_5 represents a hydrogen atom or an alkyl radical,


R_6 represents a naphthyl, phenylalkyl, Het or phenyl radical optionally substituted with one or more halogen atoms or alkyl, alkoxy, $-NR_{14}R_{15}$, hydroxyl, hydroxyalkyl, OCF_3, CF_3 or $-SO_2NH_2$ radicals, or alternatively, on 2 adjacent carbon atoms, with dioxymethylene,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, oxo, hydroxyalkyl or $-CO-NH_2$ radicals,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo or hydroxyl radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form and, preferably, Het represents a heterocycle chosen


from the following heterocycles: benzimidazole, benzoxazole, benzothiazole, benzothiophene, thiophene, quinoline, pyrrole, pyridine, pyrimidine, thiazole, thiadiazole, furan, tetrahydroisoquinoline and tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, oxo, hydroxyl, OCF_3 or CF_3 radicals.

In still another aspect of the invention, there is provided a process for preparing the compounds of formula (I) as defined in the above aspect for which R_1 represents a radical $-NHCOR_4$, characterized in that an acid R_4COOH for which R_4 has the same meanings as defined in the above aspect is reacted with a derivative of formula:



in which R_2 and R_3 have the same meanings as defined in the above aspect, the product isolated and optionally converted to a pharmaceutically acceptable salt.

The compounds of formula (I) for which R_1 represents a radical $-NHCOR_4$ may be prepared according to the following reaction scheme:



In these formulae, R_2, R_3 and R_4 have the same meanings as in formula (I).

Step a is generally carried out in an inert solvent such as tetrahydrofuran, dioxane, a chlorinated solvent (for example dichloromethane or chloroform), at a temperature of between 15 and 30°C, in the presence of a base such as a trialkylamine (for example


triethylamine or dipropylethylamine) or in pyridine, at a temperature of between 0 and 30°C.

Step b is preferably carried out in methanol, in an autoclave, at a temperature of between 50 and 70°C.

Step c is generally carried out in the presence of a condensing agent used in peptide chemistry, such as a carbodiimide (for example 1-(3-dimethylamiopropyl)-3-ethylcarbodiimide or N,N'-dicyclohexylcarbodiimide) or N,N'carbonyl-diimidazole, in an inert solvent such as an ether (for example tetrahydrofuran or dioxane), an amide (dimethylformamide) or a chlorinated solvent (for example methylene chloride, 1,2-dichloroethane or chloroform) at a temperature of between 0°C and the boiling point of the reaction mixture. It is also possible to use a reactive derivative of the acid such as an acid chloride, optionally in the presence of an acid acceptor such as a nitrogen-containing organic base (for example trialkylamine, pyridine, 1,8-diaza-bicyclo[5.4.0]undec-7-ene or 1,5-diazabicyclo[4.3.0]-non-5-ene), in a solvent as cited above, or a mixture of these solvents, at a temperature of between 0°C and the boiling point of the reaction mixture.

The derivatives R_4COOH are commercially available or may be obtained according to the methods described in R.C. LAROCK, Comprehensive Organic Transformations, VCH editor.

The azetidinols of formula 1 may be obtained by application or adaptation of the methods described by KATRITZKY A.R. et al., J. Heterocycl. Chem., 271 (1994) or DAVE P.R., J. Org. Chem., 61, 5453 (1996) and in the examples. The procedure is generally carried out according to the following reaction scheme:



in these formulae, R_2 and R_3 have the same meanings as in formula (I) and Hal represents a chlorine or bromine atom.

In step A, the procedure is preferably carried out in an inert solvent such as a 1-4C aliphatic alcohol (for example ethanol or methanol), optionally in the presence of an alkali metal hydroxide, at the boiling point of the reaction medium.

In step B, the reduction is generally carried out by means of lithium aluminum hydride, in tetrahydrofuran at the boiling point of the reaction medium.

In step C, the procedure is preferably carried out in an inert solvent such as a 1-4C aliphatic alcohol (for example ethanol or methanol), in the presence of sodium hydrogen carbonate, at a temperature of between 20°C and the boiling point of the reaction medium.

In step D, the procedure is carried out according to the method described by GRISAR M. et al. in J. Med. Chem., 885 (1973). The magnesium compound of the brominated derivative is formed and then the nitrile is caused to react, in an ether such as ethyl ether, at a temperature of between 0°C and the boiling point of the reaction medium. After hydrolysis with an alcohol, the intermediate imine is reduced *in situ* with sodium borohydride at a temperature of between 0°C and the boiling point of the reaction medium.


The derivatives R_2-CO-R_3 are commercially available or may be obtained by application or adaptation of the methods described by KUNDER N.G. et al. J. Chem. Soc. Perkin Trans 1, 2815 (1997); MORENO-MARRAS M., Eur. J. Med. Chem., 23 (5) 477 (1988); SKINNER et al., J. Med. Chem., 14 (6) 546 (1971); HURN N.K., Tet. Lett., 36 (52) 9453 (1995); MEDICI A. et al., Tet. Lett., 24 (28) 2901 (1983); RIECKE R.D. et

al., J. Org. Chem., 62 (20) 6921 (1997); KNABE J. et al., Arch. Pharm., 306 (9) 648 (1973); CONSONNI R. et al., J. Chem. Soc. Perkin Trans 1, 1809 (1996); FR-96-2481 and JP-94-261393.

The derivatives R_3Br are commercially available or may be obtained by application or adaptation of the methods described by BRANDSMA L. et al., Synth. Comm., 20 (11) 1697 and 3153 (1990); LEMAIRE M. et al., Synth. Comm., 24 (1) 95 (1994); GODA H. et al., Synthesis, 9 849 (1992); BAEUERLE P. et al., J. Chem. Soc. Perkin Trans 2, 489 (1993).

The derivatives R_2CN are commercially available or may be obtained by application or adaptation of the methods described by BOUYSSOU P. et al., J. Het. Chem., 29 (4) 895 (1992); SUZUKI N. et al., J. Chem. Soc. Chem. Comm., 1523 (1984); MARBURG S. et al., J. Het. Chem., 17 1333 (1980); PERCEC V. et al., J. Org. Chem., 60 (21) 6895 (1995).

The compounds of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ may be prepared according to the following reaction scheme:



in these formulae, Y, R_2, R_3 and R_6 have the same meanings as in formula (I), Hal represents a halogen atom and, preferably, an iodine, chlorine or bromine atom.

Step a is generally carried out in an inert solvent such as tetrahydrofuran, dioxane or a chlorinated solvent (for example dichloromethane or chloroform), in the presence of an amine such as a trialkylamine (for example triethylamine), at a temperature of between 5°C and 20°C.


Step b is generally carried out in an inert solvent such as tetrahydrofuran, in the presence of sodium hydride, at a temperature of 0°C and the boiling point of the reaction medium.

The Hal-SO_2R_6 derivatives are commercially available or may be obtained by halogenation of the corresponding sulfonic acids, in particular in situ in the presence of chlorosulfonyl isocyanate and alcohol,

in a halogenated solvent (for example dichloromethane or chloroform).

The Hal-CO-R_6 derivatives are commercially available or may be prepared according to the methods described in R. C. LAROCK, Comprehensive Organic Transformations, VCH editor.

The compounds of formula (I) may also be prepared according to the following reaction scheme:



In these formulae, R_1, R_2 and R_3 have the same meanings as in formula (I) and Ph represents a phenyl.

Step a is generally carried out in an alcohol such as methanol, in the presence of sodium borohydride, at a temperature in the region of 20°C.

In step b, the magnesium compound of the brominated derivative is prepared and it is caused to react, in an inert solvent such as ethyl ether or tetrahydrofuran, at a temperature of between 0°C and the boiling point of the reaction medium.

Step c is carried out by means of a halogenating agent such as hydrobromic acid, thionyl bromide, thionyl chloride, a mixture of triphenylphosphine and carbon tetrabromide or tetrachloride, in acetic acid or an inert solvent such as dichloromethane, chloroform, carbon tetrachloride or toluene, at a temperature of between 0°C and the boiling point of the reaction medium.

Step d is carried out by means of hydrogen, in the presence of palladized charcoal, in an alcohol such as methanol, at a temperature in the region of 20°C.

Step e is carried out in an inert solvent such as acetonitrile, in the presence of an alkali metal carbonate (for example potassium carbonate) and potassium iodide, at a temperature of between 20°C and the boiling point of the reaction medium.

The derivatives R_3Br and the derivatives R_2-CHO are commercially available or may be obtained according to the methods described for example by R.C. LAROCK, Comprehensive Organic Transformations, VCH editor.

The compounds of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with a hydroxyl radical may also be prepared by hydrolysis of a corresponding compound of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with an alkoxy radical.

This hydrolysis is generally carried out in an inert solvent such as a chlorinated solvent (for example dichloromethane or chloroform), by means of boron tribromide, at a temperature in the region of 20°C.

The compounds of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with a hydroxy(1C)alkyl radical may also be prepared by the action of diisobutylaluminum hydride on a corresponding compound of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with an alkoxycarbonyl radical.

This reaction is generally carried out in an inert solvent such as toluene, by means of diisopropyl-aluminum hydride, at a temperature of between -50°C and 25°C.

The compounds of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with a 1-pyrrolidinyl radical may also be prepared by the action of pyrrolidine and of a

corresponding compound of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with a fluorine atom.

This reaction is preferably carried out in an inert solvent such as dimethyl sulfoxide, at a temperature of 90°C.

It is understood for persons skilled in the art that, to carry out the processes according to the invention which are described above, it may be necessary to introduce groups protecting amino, hydroxyl and carboxyl functions in order to avoid side reactions. These groups are those which allow removal without affecting the rest of the molecule. As examples of groups protecting the amino function, there may be mentioned tert-butyl or methyl carbamates which may be regenerated using iodotrimethylsilane or allyl using palladium catalysts. As examples of groups protecting the hydroxyl function, there may be mentioned triethylsilyl and tert-butyldimethylsilyl which may be regenerated using tetrabutylammonium fluoride or alternatively asymmetric acetals (methoxymethyl or tetrahydropyranyl for example) with regeneration using hydrochloric acid. As groups protecting carboxyl functions, there may be mentioned esters (allyl or benzyl for example), oxazoles and 2-alkyl-1,3-oxazolines. Other protecting groups which can be used are described by GREENE T.W. et al., Protecting Groups

in Organic Synthesis, second edition, 1991, John Wiley & Sons.

The compounds of formula (I) may be purified by the customary known methods, for example by crystallization, chromatography or extraction.

The enantiomers of the compounds of formula (I) may be obtained by resolution of the racemates for example by chromatography on a chiral column according to PIRCKLE W.H. et al., Asymmetric synthesis, Vol. 1, Academic Press (1983) or by formation of salts or by synthesis from chiral precursors. The diastereoisomers may be prepared according to known conventional methods (crystallization, chromatography or from chiral precursors).

The compounds of formula (I) may be optionally converted to addition salts with an inorganic or organic acid by the action of such an acid in an organic solvent such as an alcohol, a ketone, an ether or a chlorinated solvent. These salts also form part of the invention.

As examples of pharmaceutically acceptable salts, the following salts may be mentioned: benzenesulfonate, hydrobromide, hydrochloride, citrate, ethanesulfonate, fumarate, gluconate, iodate, isethionate, maleate, methanesulfonate, methylene-bis-β-oxynaphthoate, nitrate, oxalate, pamoate, phosphate, salicylate, succinate, sulfate, tartrate, theophyllineacetate and p-toluenesulfonate.

The compounds of formula (I) exhibit advantageous pharmacological properties. These compounds possess a high affinity for the cannabinoid receptors and particularly those of the CB1 type. They are CB1-receptor antagonists and are therefore useful in the treatment and prevention of disorders affecting the central nervous system, the immune system, the cardiovascular or endocrine system, the respiratory system, the gastrointestinal apparatus and reproductive disorders (Hollister, Pharm. Rev.; 38, 1986, 1-20, Reny and Sinha, Prog. Drug Res., 36, 71-114 (1991), Consroe and Sandyk, in Marijuana/Cannabinoids, Neurobiology and Neurophysiology, 459, Murphy L. and Barthe A. Eds, CRC Press, 1992).

Accordingly, these compounds may be used for the treatment or prevention of psychoses including schizophrenia, anxiety disorders, depression, epilepsy, neurodegeneration, cerebellar and spinocerebellar disorders, cognitive disorders, cranial trauma, panic attacks, peripheral neuropathies, glaucomas, migraine, Parkinson's disease, Alzheimer's disease, Huntington's chorea, Raynaud's syndrome, tremor, obsessive-compulsive disorder, senile dementia, thymic disorders, Tourette's syndrome, tardive dyskinesia, bipolar disorders, cancers, movement disorders induced by medicaments, dystonia, endotoxemic shocks, hemorrhagic shocks, hypotension, insomnia, immunological diseases, multiple sclerosis, vomiting, asthma, appetite

disorders (bulimia, anorexia), obesity, memory disorders, in weaning from chronic treatments and alcohol or drug abuse (opioids, barbiturates, cannabis, cocaine, amphetamine, phencyclide, hallucinogens, benzodiazepines for example), as analgesics or potentiators of the analgesic activity of the narcotic and nonnarcotic drugs. They may also be used for the treatment or prevention of intestinal transit.

The affinity of the compounds of formula (I) for the cannabis receptors has been determined according to the method described by KUSTER J.E., STEVENSON J.I., WARD S.J., D'AMBRA T.E., HAYCOCK D.A. in J. Pharmacol. Exp. Ther., 264 1352-1363 (1993).

In this test, the IC_{50} of the compounds of formula (I) is less than or equal to 1000 nM.

Their antagonist activity has been shown by means of the model of hypothermia induced by an agonist of the cannabis receptors (CP-55940) in mice, according to the method described by Pertwee R.G. in Marijuana, Harvey D.J. eds, 84 Oxford IRL Press, 263-277 (1985).

In this test, the ED_{50} of the compounds of formula (I) is less than or equal to 50 mg/kg.

The compounds of formula (I) exhibit low toxicity. Their LD_{50} is greater than 40 mg/kg by the subcutaneous route in mice.

The following examples illustrate the invention.

Example 1

69.3 mm^3 of triethylamine and 110 mg of thien-2-ylsulfonyl chloride are successively added, at room temperature and under an argon atmosphere, to a solution of 61.4 mg of 1-[bis(4-chlorophenyl)methyl]-azetidin-3-ylamine in 3 cm^3 of dichloromethane. After stirring for 68 hours at room temperature, the reaction mixture is introduced into a Bond Elut® SCX cartridge (3 cm^5/500 mg), eluting successively with twice 2 cm^3 of dichloromethane and then twice 2 cm^3 of a 1 M solution of ammonia in methanol. The ammoniacal fractions are pooled and concentrated to dryness under reduced pressure (2.7 kPa). The residue obtained is dissolved in 5 cm^3 of dichloromethane, washed with three times 3 cm^3 of distilled water, dried over magnesium sulfate, filtered and concentrated to dryness under reduced pressure (2.7 kPa). 60 mg of N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}thien-2-ylsulfonamide are thus obtained in the form of a cream-colored foam [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.77 (split t, J = 7 and 2 Hz : 2H); 3.40 (split t, J = 7 and 2 Hz : 2H); 4.06 (mt : 1H); 4.21 (s : 1H); from 4.85 to 5.25 (broad unresolved complex : 1H); 7.06 (t, J = 4.5 Hz : 1H); from 7.15 to 7.35 (mt : 8H); 7.58 (mt : 2H)].

1-[Bis(4-chlorophenyl)methyl]azetidin-3-yl-amine may be obtained in the following manner: 400 cm^3 of a mixture of methanol and liquid ammonia (50/50 by volume) are added to 27 g of 1-[bis(4-chlorophenyl)-

methyl]azetidin-3-yl methylsulfonate contained in an autoclave previously cooled to around -60°C. The reaction medium is then stirred at 60°C for 24 hours and then abandoned in the open air to allow the evaporation of the ammonia and finally concentrated under reduced pressure (2.7 kPa). The residue is taken up in 500 cm^3 of a 0.37 N aqueous sodium hydroxide solution and extracted with four times 500 cm^3 of ethyl ether. The combined organic phases are washed successively with twice 100 cm^3 of distilled water and 100 cm^3 of a saturated sodium chloride solution, dried over magnesium sulfate, filtered and concentrated under reduced pressure (2.7 kPa). The residue obtained is purified by flash chromatography on silica gel [eluant: dichloromethane/methanol (95/5 by volume)]. 14.2 g of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ylamine are obtained in the form of an oil which solidifies into a cream-colored solid.

1-[Bis(4-chlorophenyl)methyl]azetidin-3-yl methylsulfonate may be prepared in the following manner: 3.5 cm^3 of methylsulfonyl chloride are added, under argon over 10 minutes, to a solution of 12 g of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ol in 200 cm^3 of dichloromethane and then the mixture is cooled to +5°C and 3.8 cm^3 of pyridine are poured in over 10 minutes. After stirring for 30 minutes at +5°C and then for 20 hours at 20°C, the reaction mixture is diluted with 100 cm^3 of water and 100 cm^3 of dichloromethane. The

mixture, which is first filtered, is separated after settling. The organic phase is washed with water and then dried over magnesium sulfate, filtered and concentrated to dryness under reduced pressure (2.7 kPa). The oil obtained is chromatographed on a silica gel column (particle size 0.063-0.200 mm, height 40 cm, diameter 3.0 cm), eluting under an argon pressure of 0.5 bar with a mixture of cyclohexane and ethyl acetate (70/30 by volume) and collecting 100-cm^3 fractions. Fractions 4 to 15 are combined and concentrated to dryness under reduced pressure (2.7 kPa). 6.8 g of 1-[bis(4-chlorophenyl)methyl]-azetidin-3-yl ester of methylsulfonic acid are obtained in the form of a yellow oil.

1-[Bis(4-chlorophenyl)methyl]azetidin-3-ol may be prepared according to the procedure described by KATRITZKY A.R. et al., J. Heterocycl. Chem., 271 (1994), starting with 35.5 g of [bis(4-chlorophenyl)methyl]-amine hydrochloride and 11.0 cm^3 of epichlorohydrin. 9.0 g of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ol are isolated.

[Bis(4-chlorophenyl)methyl]amine hydrochloride may be prepared according to the method described by GRISAR M. et al., J. Med. Chem., 885 (1973).

Example 2

By carrying out the operation according to the procedure of Example 1, but starting with 124 mg of

4-methoxyphenylsulfonyl chloride, 12 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-4-methoxyphenylsulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.70 (split t, J = 7 and 2 Hz : 2H); 3.35 (split t, J = 7 and 2 Hz : 2H); 3.85 (s : 3H); 3.94 (mt : 1H); 4.18 (s : 1H); 4.83 (d, J = 9 Hz : 1H); 6.94 (broad d, J = 9 Hz : 2H); 7.22 (s : 8H); 7.75 (broad d, J = 9 Hz : 2H)].

Example 3

By carrying out the operation according to the procedure of Example 1, but starting with 140 mg of 4-acetamidophenylsulfonyl chloride, 13 mg of N-[4-(N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}sulfamoyl)phenyl]acetamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.26 (s : 3H); 2.74 (split t, J = 7 and 2 Hz : 2H); 3.39 (split t, J = 7 and 2 Hz : 2H); 4.01 (mt : 1H); 4.22 (s : 1H); 4.92 (d, J = 9 Hz : 1H); 7.32 (mt : 8H); 7.49 (broad s : 1H); 7.68 (broad d, J = 9 Hz : 2H); 7.81 (broad d, J = 9 Hz : 2H)].

Example 4

By carrying out the operation according to the procedure of Example 1, but starting with 114 mg of 4-methylphenylsulfonyl chloride, 19 mg of N-{1-[bis-(4-chlorophenyl)methyl]azetidin-3-yl}-4-methylphenyl-sulfonamide are obtained in the form of a colorless

lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.42 (s : 3H); 2.71 (split t, J = 7 and 2 Hz : 2H); 3.36 (split t, J = 7 and 2 Hz : 2H); 3.97 (mt : 1H); 4.19 (s : 1H); 4.81 (d, J = 9.5 Hz : 1H); from 7.15 to 7.40 (mt : 10H); 7.71 (broad d, J = 8.5 Hz : 2H)].

Example 5

By carrying out the operation according to the procedure of Example 1, but starting with 142 mg of 3,4-dimethoxyphenylsulfonyl chloride, 10 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-3,4-dimethoxyphenylsulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.72 (broad t, J = 7.5 Hz : 2H); 3.37 (broad t, J = 7.5 Hz : 2H); from 3.85 to 4.00 (mt : 1H); 3.91 (s : 3H); 3.93 (s : 3H); 4.19 (s : 1H); 4.84 (d, J = 9 Hz : 1H); 6.90 (d, J = 8.5 Hz : 1H); 7.23 (mt : 8H); 7.29 (d, J = 2 Hz : 1H); 7.43 (dd, J = 8.5 and 2 Hz : 1H)].

Example 6

By carrying out the operation according to the procedure of Example 1, but starting with 117 mg of 3-fluorophenylsulfonyl chloride, 13.5 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-3-fluorophenyl-sulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (400 MHz, $CDCl_3$, δ in ppm): 2.79 (split t, J = 7 and 2 Hz : 2H); 3.43 (split t, J = 7 and 2 Hz : 2H); 4.05 (unresolved complex : 1H); 4.24 (s : 1H); 4.91 (unresolved complex

: 1H); from 7.20 to 7.40 (mt : 9H); from 7.50 to 7.65 (mt : 2H); 7.67 (broad d, J = 8 Hz : 1H)].

Example 7

By carrying out the operation according to the procedure of Example 1, but starting with 147 mg of 3,4-dichlorophenylsulfonyl chloride, 20 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-3,4-dichlorophenylsulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.77 (split t, J = 7 and 2 Hz : 2H); 3.40 (split t, J = 7 and 2 Hz : 2H); 3.98 (mt : 1H); 4.21 (s : 1H); from 4.85 to 5.15 (unresolved complex : 1H); from 7.20 to 7.35 (mt : 8H); 7.57 (d, J = 8.5 Hz : 1H); 7.65 (dd, J = 8.5 and 2 Hz : 1H); 7.93 (d, J = 2 Hz : 1H)].

Example 8

By carrying out the operation according to the procedure for Example 1, but starting with 121 mg of 3-cyanophenylsulfonyl chloride, 21 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-3-cyanophenylsulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.76 (split t, J = 7 and 2 Hz : 2H); 3.39 (split t, J = 7 and 2 Hz : 2H); 3.99 (mt : 1H); 4.21 (s : 1H); from 4.80 to 5.60 (very broad unresolved complex : 1H); from 7.15 to 7.35 (mt : 8H); 7.65 (t, J = 8 Hz : 1H); 7.86 (broad d, J = 8 Hz : 1H); 8.05 (broad d, J = 8 Hz : 1H); 8.13 (broad s : 1H)].

Example 9

By carrying out the operation according to the procedure of Example 1, but starting with 142 mg of 2,5-dimethoxyphenylsulfonyl chloride, 31 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-2,5-dimethoxyphenylsulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.73 (split t, J = 7 and 2 Hz : 2H); 3.27 (split t, J = 7 and 2 Hz : 2H); 3.80 (s : 3H); from 3.85 to 4.00 (mt : 1H); 3.94 (s : 3H); 4.19 (s : 1H); 5.32 (d, J = 8 Hz : 1H); 6.94 (d, J = 9 Hz : 1H); 7.05 (dd, J = 9 and 3 Hz : 1H); 7.23 (mt : 8H); 7.40 (d, J = 3 Hz : 1H)].

Example 10

By carrying out the operation according to the procedure of Example 1, but starting with 147 mg of 3-trifluoromethylphenylsulfonyl chloride, 8 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-3-trifluoromethylphenylsulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.79 (split t, J = 7 and 2 Hz : 2H); 3.41 (split t, J = 7 and 2 Hz : 2H); 4.03 (mt : 1H); 4.23 (s : 1H); from 4.80 to 5.10 (broad unresolved complex : 1H); from 7.20 to 7.35 (mt : 8H); 7.68 (t, J = 8 Hz : 1H); 7.87 (broad d, J = 8 Hz : 1H); 8.05 (broad d, J = 8 Hz : 1H); 8.15 (broad s : 1H)].

Example 11

By carrying out the operation according to the procedure of Example 1, but starting with 136 mg of naphth-2-ylsulfonyl chloride, 20 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}naphth-2-yl-sulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (400 MHz, $CDCl_3$, δ in ppm): 2.74 (mt : 2H); 3.35 (mt : 2H); 4.02 (mt : 1H); 4.17 (s : 1H); 4.96 (unresolved complex : 1H); from 7.10 to 7.30 (mt : 8H); 7.64 (mt : 2H); 7.78 (dd, J = 7 and 1.5 Hz : 1H); 7.90 to 8.05 (mt : 3H); 8.41 (broad s : 1H)].

Example 12

By carrying out the operation according to the procedure of Example 1, but starting with 136 mg of naphth-1-ylsulfonyl chloride, 52 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}naphth-1-yl-sulfonamide are obtained in the form of a cream-colored foam [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.63 (split t, J = 7 and 2 Hz : 2H); 3.20 (split t, J = 7 and 2 Hz : 2H); 3.90 (mt : 1H); 4.12 (s : 1H); 5.26 (unresolved complex : 1H); 7.16 (mt : 8H); 7.52 (t, J = 8 Hz : 1H); from 7.55 to 7.75 (mt : 2H); 7.95 (d, J = 8.5 Hz : 1H); 8.06 (d, J = 8.5 Hz : 1H); 8.23 (dd, J = 7.5 and 1 Hz : 1H); 8.64 (d, J = 8.5 Hz : 1H)].

Example 13

By carrying out the operation according to the procedure of Example 1, but starting with 128 mg of

3,4-difluorophenylsulfonyl chloride, 7 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-3,4-difluorophenylsulfonamide are obtained in the form of a cream-colored lacquer [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.76 (broad t, J = 7.5 Hz : 2H); 3.39 (broad t, J = 7.5 Hz : 2H); 3.98 (mt : 1H); 4.20 (broad s : 1H); from 4.85 to 5.25 (broad unresolved complex : 1H); from 7.15 to 7.35 (mt : 9H); from 7.55 to 7.75 (mt : 2H)].

Example 14

By carrying out the operation according to the procedure of Example 1, but starting with 108 mg of 1-methylimidazol-4-ylsulfonyl chloride, 22 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-1-methyl-1-*H*-imidazol-4-ylsulfonamide are obtained in the form of a cream-colored foam [^{1}H NMR spectrum (300 MHz, $CDCl_3$, with addition of a few drops of CD_3COOD d4, δ in ppm): 3.22 (mt : 2H); 3.67 (mt : 2H); 3.74 (s : 3H); 4.10 (mt : 1H); 4.65 (broad s : 1H); 7.27 (mt : 8H); 7.47 (broad d, J = 1 Hz : 1H); 7.53 (broad d, J = 1 Hz : 1H)].

Example 15

By carrying out the operation according to the procedure of Example 1, but starting with 152 mg of 4-acetamido-3-chlorophenylsulfonyl chloride, 69 mg of N-[4-(N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}sulfamoyl)-2-chlorophenyl]acetamide are obtained in the form of a cream-colored foam [^{1}H NMR spectrum (300

MHz, $CDCl_3$, δ in ppm): 2.30 (s : 3H); 2.73 (mt : 2H); 3.38 (mt : 2H); 3.97 (mt : 1H); 4.19 (s : 1H); 7.24 (s : 8H); 7.70 (dd, J = 7 and 1.5 Hz : 1H); 7.78 (broad s : 1H); 7.86 (d, J = 1.5 Hz : 1H); 8.61 (d, J = 7 Hz : 1H)].

Example 16

0.79 cm^3 of triethylamine is added, at room temperature under an argon atmosphere, to a solution of 0.7 g of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ylamine in 25 cm^3 of dichloromethane. The mixture is cooled to around 0°C before adding thereto a solution of 1.2 g of pyrid-3-ylsulfonyl chloride in 25 cm^3 of dichloromethane, and then it is stirred at room temperature for 16 hours. The reaction mixture is diluted with 50 cm^3 of dichloromethane and is then washed with twice 25 cm^3 of distilled water. The organic phase is dried over magnesium sulfate, filtered and concentrated to dryness under reduced pressure (2.7 kPa). The residue obtained is purified by flash chromatography on silica gel [eluant: dichloromethane/methanol (97.5/2.5 by volume)]. 0.7 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}pyrid-3-ylsulfonamide is obtained in the form of a cream-colored foam which solidifies in the presence of isopropanol into a cream-colored powder melting at 164°C.

Pyrid-3-ylsulfonyl chloride may be prepared according to the method described by Breant, P. et al., Synthesis, 10, 822-4 (1983).

Example 17

0.214 g of 4-fluorophenylsulfonyl chloride and 0.28 cm^3 of triethylamine are added, at room temperature under an argon atmosphere, to a solution of 0.307 g of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ylamine in 10 cm^3 of dichloromethane. After stirring for 16 hours at room temperature, the reaction mixture is washed with 10 cm^3 of distilled water, dried over magnesium sulfate, filtered and concentrated to dryness under reduced pressure (2.7 kPa). The residue obtained is purified by flash chromatography on silica gel [eluant: dichloromethane/ethyl acetate (100/0 to 95/5 by volume) gradient]. 0.18 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-4-fluorophenylsulfonamide is obtained in the form of a white foam [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.74 (broad t, J = 7.5 Hz : 2H); 2.39 (broad t, J = 7.5 Hz : 2H); 3.98 (mt : 1H); 4.20 (s : 1H); 4.79 (d, J = 9 Hz : 1H); from 7.10 to 7.35 (mt : 10H); 7.86 (mt : 2H)].

Example 18

By carrying out the operation according to the procedure for Example 17, but starting with 0.25 g of quinol-8-ylsulfonyl chloride, 0.36 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}quinol-8-ylsulfonamide is obtained in the form of a white powder [^{1}H NMR

spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.63 (split t, J = 7 and 2 Hz : 2H); 3.16 (split t, J = 7 and 2 Hz : 2H); 3.98 (mt : 1H); 4.11 (s : 1H); 6.77 (d, J = 8 Hz : 1H); 7.15 (mt : 8H); 7.61 (dd, J = 8 and 4 Hz : 1H); 7.64 (dd, J = 8 and 7.5 Hz) : 1H); 8.06 (dd, J = 8 and 1.5 Hz : 1H); 8.30 (dd, J = 8 and 1.5 Hz : 1H); 8.40 (dd, J = 7.5 and 1.5 Hz : 1H); 9.09 (dd, J = 4 and 1.5 Hz : 1H)].

Example 19

By carrying out the operation according to the procedure of Example 17, but starting with 0.14 cm^3 of phenylsulfonyl chloride, 0.35 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}phenylsulfonamide is obtained in the form of a white powder [^1H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.75 (broad t, J = 7.5 Hz : 2H); 3.40 (broad t, J = 7.5 Hz : 2H); 4.03 (mt : 1H); 4.22 (s : 1H); 4.79 (d, J = 10 Hz : 1H); 7.31 (s : 8H); from 7.45 to 7.65 (mt : 3H); 7.87 (broad d, J = 7.5 Hz : 2H)].

Example 20

By carrying out the operation according to the procedure of Example 17, but starting with 0.21 g of (phenylmethyl)sulfonyl chloride, 0.27 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}(phenylmethyl)sulfonamide is obtained in the form of a white powder [^1H NMR spectrum (400 MHz, $CDCl_3$, δ in ppm): 2.76 (split t, J = 7 and 2 Hz : 2H); 3.41 (split t, J = 7 and 2 Hz : 2H); 3.85 (mt : 1H); 4.20 (s : 1H); 4.23 (s

: 2H); 4.46 (d, J = 9 Hz : 1H); from 7.25 to 7.45 (mt : 13H)].

Example 21

By carrying out the operation according to the procedure of Example 17, but starting with 0.42 g of 3,5-difluorophenylsulfonyl chloride in 30 cm^3 of dichloromethane and washing the organic phase with twice 20 cm^3 of distilled water. After purification by flash chromatography on silica gel [eluant: dichloromethane/methanol (100/0 to 95/5 by volume) gradient], 0.1 g of N-{1-[bis(4-chlorophenyl)methyl]-azetidin-3-yl}-3,5-difluorophenylsulfonamide is obtained in the form of a yellow powder [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.77 (split t, J = 7 and 2 Hz : 2H); 3.41 (split t, J = 7 and 2 Hz : 2H); 4.01 (mt : 1H); 4.21 (s : 1H); 4.90 (d, J = 9 Hz : 1H); 7.02 (tt, J = 8.5 and 2.5 Hz : 1H); from 7.20 to 7.35 (mt : 8H); 7.38 (mt : 2H)].

3,5-Difluorophenylsulfonyl chloride may be prepared according to the method described in Patent FR 9615887.

Example 22

By carrying out the operation according to the procedure of Example 21, but starting with 0.21 g of pyrid-2-ylsulfonyl chloride and 0.17 cm^3 of triethylamine and washing the organic phase with twice 30 cm^3 of distilled water. After purification by flash chromatography on silica gel [eluant:

dichloromethane/methanol (100/0 to 98/2 by volume) gradient], 0.3 g of N-{1-[bis(4-chlorophenyl)methyl]-azetidin-3-yl}pyrid-2-sulfonamide is obtained in the form of a white powder [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.78 (split t, J = 7 and 2 Hz : 2H); 3.35 (split t, J = 7 and 2 Hz : 2H); 4.12 (mt : 1H); 4.20 (s: 1H); 5.30 (d, J = 9 Hz : 1H); from 7.15 to 7.35 (mt : 8H); 7.47 (ddd, J = 7.5 and 5 and 1 Hz : 1H); 7.90 (split t, J = 7.5 and 2 Hz; 1H); 7.98 (broad d, J = 7.5 Hz : 1H); 8.65 (broad d, J = 5 Hz : 1H)].

Pyrid-2-ylsulfonyl chloride may be prepared according to the method described by Corey, E.J. et al., J. Org. Chem. (1989), 54(2), 389-93.

Example 23

0.104 cm^3 of pyrrolidine is added, at room temperature, to 0.24 g of N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}-(3,5-difluorophenyl)sulfonamide in solution in 6 cm^3 of dimethyl sulfoxide and then the mixture is heated for 18 hours at 90°C. The reaction mixture is diluted with 30 cm^3 of dichloromethane and washed with three times 30 cm^3 of distilled water. The organic phased is dried over magnesium sulfate, filtered and concentrated to dryness under reduced pressure (2.7 kPa). The residue is purified by flash chromatography on silica gel, eluting with dichloromethane. 50 mg of N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}-(3-fluoro-5-pyrrolidin-1-ylphenyl)sulfonamide are thus obtained in the form of a

white powder [^{1}H NMR spectrum (600 MHz, $CDCl_3$ with addition of a few drops of CD_3COOD d4, δ in ppm): 2.04 (mt : 4H); from 3.20 to 3.35 (mt : 6H); 3.60 (t, J = 8.5 Hz : 2H); 4.14 (mt : 1H); 4.57 (s : 1H); 6.31 (broad d, J = 11.5 Hz : 1H); 6.70 (broad d, J = 8.5 Hz : 1H); 6.72 (broad s : 1H); from 7.20 to 7.35 (mt : 8H)].

Example 24

A solution of 0.26 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-4-fluorophenylsulfonamide in 5 cm^3 of tetrahydrofuran is added, at room temperature under an argon atmosphere, to a suspension of 20.5 mg of 80% sodium hydride in 10 cm^3 of tetrahydrofuran. After stirring for 1 hour at around 20°C, 60 mm^3 of iodomethane are added and then after stirring for an additional 16 hours, the suspension is supplemented with 30 cm^3 of ethyl acetate and 20 cm^3 of distilled water. The organic phase is dried over magnesium sulfate, filtered and concentrated to dryness under reduced pressure (2.7 kPa). The residue is purified by flash chromatography on silica gel [eluant: cyclohexane/ethyl acetate (90/10 by volume)]. 19 mg of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-N-methyl-4-fluorophenylsulfonamide are thus obtained in the form of a white powder [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.69 (s : 3H); 3.02 (split t, J = 7 and 2 Hz : 2H); 3.35 (split t, J = 7 and 2 Hz : 2H);

3.91 (mt : 1H); 4.27 (s : 1H); from 7.15 to 7.35 (mt : 10H); 7.75 (dd, J = 9 and 5 Hz : 2H)].

Example 25

By carrying out the operation according to the procedure of Example 24, but starting with 0.25 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-quinol-8-ylsulfonamide and 18 mg of 80% sodium hydride. After purification by flash chromatography on silica gel [eluant: cyclohexane/ethyl acetate (80/20 by volume)], 70 mg of N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}-N-methylquinol-8-ylsulfonamide are obtained [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): from 3.00 to 3.10 (mt : 2H); 3.05 (s : 3H); 3.35 (mt : 2H); 4.27 (s : 1H); 4.93 (mt : 1H); from 7.15 to 7.35 (mt : 8H); 7.50 (dd, J = 8.5 and 4 Hz : 1H); 7.62 (dd, J = 8 and 8.5 Hz : 1H); 8.03 (dd, J = 8.5 and 1.5 Hz : 1H); 8.22 (dd, J = 8.5 and 1.5 Hz : 1H); 8.48 (dd, J = 8 and 1.5 Hz : 1H); 8.98 (dd, J = 4 and 1.5 Hz : 1H)].

Example 26

By carrying out the operation according to the procedure of Example 24, but starting with 0.21 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-phenylsulfonamide, 17 mg of 80% sodium hydride and introducing iodomethane in two portions at an interval of 3 hours. 80 mg of N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}-N-methylphenylsulfonamide are thus obtained in the form of a white lacquer [^{1}H NMR

spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.70 (s : 3H); 3.03 (broad t, J = 7.5 Hz : 2H); 3.37 (broad t, J = 7.5 Hz : 2H); 3.94 (mt : 1H); 4.28 (s : 1H); from 7.20 to 7.35 (mt : 8H); from 7.45 to 7.65 (mt : 3H); 7.74 (broad d, J = 8 Hz : 2H)].

Example 27

By carrying out the operation according to the procedure of Example 26, but starting with 0.17 g of N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-(phenylmethyl)sulfonamide, 14 mg of 80% sodium hydride and maintaining stirred for 48 hours at 20°C. After purification by flash chromatography on silica gel [eluant: dichloromethane/ethyl acetate (95/5 by volume)], 120 mg of N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}-N-methyl(phenylmethyl)sulfonamide are obtained in the form of a white foam [^1H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.81 (s : 3H); 2.88 (split t, J = 7 and 2 Hz : 2H); 3.16 (split t, J = 7 and 2 Hz : 2H); from 4.10 to 4.25 (mt : 4H); from 7.20 to 7.40 (mt : 13H)].

Example 28

A solution of 0.307 g of 1-[bis(4-chloro-phenyl)methyl]azetidin-3-ylamine is added dropwise, at room temperature, to a solution of 0.412 g of 1,3-benzenedisulfonic acid dichloride and 0.165 cm^3 of triethylamine in 10 cm^3 of acetonitrile. After stirring for 3 hours at room temperature, 0.28 cm^3 of a 20% solution of ammonia is added and the reaction mixture

is left at room temperature. After 18 hours, the mixture is filtered and concentrated to dryness under reduced pressure (2.7 kPa). After chromatography on a silica gel column (particle size 0.06-0.200 mm, height 35 cm, diameter 2 cm), eluting under an argon pressure of 0.9 bar with dichloromethane and then a mixture of dichloromethane + 1% methanol and then a mixture of dichloromethane + 2% methanol by volume and collecting 30-cm^3 fractions, fractions 23 to 34 are combined and concentrated to dryness under reduced pressure (2.7 kPa) to give 90 mg of N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}-3-sulfamoylphenylsulfonamide in the form of a white solid [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.78 (broad t, J = 7 Hz : 2H); 3.35 (broad t, J = 7 Hz : 2H); 4.01 (mt : 1H); 4.24 (s : 1H); 5.27 (unresolved complex : 2H); 5.61 (unresolved complex : 1H); from 7.15 to 7.35 (mt : 8H); 7.67 (t, J = 8 Hz : 1H); 8.04 (broad d, J = 8 Hz : 1H); 8.12 (broad d, J = 8 Hz : 1H); 8.49 (broad s : 1H)].

Example 29

0.031 cm^3 of diisopropylcarbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)-methyl]azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane are added to a solution of 80.1 mg of benzenesulfonylacetic acid, 27 mg of hydroxybenzotriazole in solution in 0.5 cm^3 of dimethylformamide under an inert argon atmosphere, at a

temperature in the region of 23°C. After 17 hours at a temperature in the region of 23°C, the reaction mixture is loaded onto a 3-cm^3 SPE cartridge containing 1 g of SCX phase preconditioned with methanol. After washing with twice 5 cm^3 of methanol and then 4 cm^3 of 0.1 N ammoniacal methanol, the expected product is eluted with 4 cm^3 of 1 N ammoniacal methanol. The fraction containing the expected product is evaporated under an air stream at a temperature in the region of 45°C and then dried under reduced pressure (1 mbar) at a temperature in the region of 40°C. 2-Benzenesulfonyl-N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}acetamide is thus obtained in the form of a white solid [^{1}H NMR spectrum (500 MHz, $CDCl_3$, δ in ppm): 2.96 (mt : 2H); 3.51 (mt : 2H); 4.00 (s : 2H); 4.34 (unresolved complex : 1H); 4.48 (mt : 1H); 7.10 (unresolved complex : 1H); from 7.20 to 7.45 (mt : 8H); 7.57 (t, J = 8 Hz : 2H); 7.70 (t, J = 8 Hz : 1H); 7.90 (d, J = 8 Hz : 2H)].

Example 30

0.031 cm^3 of diisopropylcarbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)-methyl]azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane are added to a solution of 85.7 mg of toluenesulfonylacetic acid, 27 mg of hydroxybenzotriazole in solution in 0.5 cm^3 of dimethylformamide under an inert argon atmosphere, at a temperature in the region of 23°C. After 17 hours at a

temperature in the region of 23°C, the reaction mixture is loaded onto a 3-cm^3 SPE cartridge containing 1 g of SCX phase preconditioned with methanol. After washing with twice 5 cm^3 of methanol and then 4 cm^3 of 0.1 N ammoniacal methanol, the expected product is eluted with 4 cm^3 of 1 N ammoniacal methanol. The fraction containing the expected product is evaporated under an air stream at a temperature in the region of 45°C and then dried under reduced pressure (1 mbar) at a temperature in the region of 40°C. N-{1-[bis(4-chlorophenyl)methyl]-azetidin-3-yl}-2-(toluene-4-sulfonyl)acetamide is obtained in the form of a yellow lacquer [^{1}H NMR spectrum (500 MHz, $CDCl_3$, δ in ppm): 2.85 (t, J = 7 Hz : 2H); 3.07 (s : 3H); 3.48 (t, J = 7 Hz : 2H); 4.24 (s : 1H); 4.49 (mt : 1H); 7.19 (broad d, J = 6 Hz : 1H); from 7.20 to 7.40 (mt : 8H); 8.40 (s : 1H)].

Example 31

By carrying out the operation according to the procedure of Example 30, starting with 85.7 mg of 3-chloro-4-methylsulfonythiophene-2-carboxylic acid, 27 mg of hydroxybenzotriazole in solution in 0.5 cm^3 of dimethylformamide, 0.031 cm^3 of diisopropylcarbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)methyl]-azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane, (3-chloro-4-methylsulfonylthiophene-2-carboxy)-{1-bis(4-chlorophenyl)methyl]azetidin-3-yl}amide is

obtained in the form of a yellow lacquer [^{1}H NMR spectrum (500 MHz, $CDCl_3$, δ in ppm): 2.44 (s : 3H); 2.96 (unresolved complex : 2H); 3.52 (unresolved complex : 2H); 3.98 (s : 2H); 4.35 (unresolved complex : 1H); 4.49 (mt : 1H); from 7.00 to 7.30 (broad unresolved complex : 1H); from 7.20 to 7.45 (mt : 10H); 7.76 (d, J = 8 Hz : 2H)].

Example 32

By carrying out the operation according to the procedure of Example 30, starting with 96.1 mg of 3-(2-phenylethylenesulfonyl)propionic acid, 27 mg of hydroxybenzotriazole in solution in 0.5 cm^3 of dimethylformamide, 0.031 cm^3 of diisopropylcarbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)methyl]-azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane, N-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}-3-(2-phenylethylenesulfonyl)propionamide is obtained in the form of a white foam [^{1}H NMR spectrum (500 MHz, $CDCl_3$, δ in ppm): 2.64 (t, J = 7 Hz : 2H); 2.88 (unresolved complex : 2H); 3.33 (t, J = 7 Hz : 2H); 3.49 (unresolved complex : 2H); 4.29 (unresolved complex : 1H); 4.48 (mt : 1H); from 5.90 to 6.15 (broad unresolved complex : 1H); 6.41 (d, J = 12 Hz : 1H); 7.17 (d, J = 12 Hz : 1H); from 7.20 to 7.35 (mt : 8H); 7.41 (mt : 3H); 7.64 (mt : 2H)].

Example 33

By carrying out the operation according to the procedure of Example 31, starting with 58.5 mg of 4-methylsulfonylbenzoic acid, 26.4 mg of hydroxybenzotriazole in solution in 0.5 cm^3 of dimethylformamide, 0.0302 cm^3 of diisopropylcarbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane, N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}-4-methylsulfonylbenzamide is obtained in the form of a white crystals [^1H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 3.03 (mt : 2H); 3.09 (s : 3H); 3.61 (broad t, J = 7.5 Hz : 2H); 4.35 (s : 1H); 4.73 (mt : 1H); 6.55 (broad d, J = 7.5 Hz : 1H); from 7.20 to 7.35 (mt : 8H); 7.96 (d, J = 8 Hz : 2H); 8.03 (d, J = 8 Hz : 2H)].

Example 34

By carrying out the operation according to the procedure of Example 31, starting with 58.5 mg of 3-phenylsulfonylpropionic acid, 26.4 mg of hydroxybenzotriazole in solution in 0.5 cm^3 of dimethylformamide, 0.0302 cm^3 of diisopropylcarbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane, N-{1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}-4-methanesulfonylbenzamide is obtained in the form of a lacquer [^1H NMR spectrum

(300 MHz, $CDCl_3$, δ in ppm): 2.71 (t, J = 7.5 Hz : 2H); 2.86 (mt : 2H); from 3.40 to 3.55 (mt : 4H); 4.26 (s : 1H); 4.45 (mt : 1H); 6.22 (broad d, J = 7.5 Hz : 1H); from 7.20 to 7.35 (mt : 8H); 7.59 (broad t, J = 7.5 Hz : 2H); 7.69 (tt, J = 7.5 and 1.5 Hz : 1H); 7.93 (broad d, J = 7.5 Hz : 2H)].

Example 35

By carrying out the operation according to the procedure of Example 31, starting with 60.2 mg of 5-methylsulfonylthiophene-2-carboxylic acid, 26.4 mg of hydroxybenzotriazole in solution in 0.5 cm^3 of dimethylformamide, 0.0302 cm^3 of diisopropylcarbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane, (5-methylsulfonylthiophene-2-carboxy)-{1-[bis(4-chlorophenyl)methyl]azetidin-3-yl}amide is obtained in the form of white crystals [^1H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 3.03 (mt : 2H); 3.21 (s : 3H); 3.57 (dd, J = 8 and 7.5 Hz : 2H); 4.34 (s : 1H); 4.67 (mt : 1H); 6.40 (broad d, J = 7.5 Hz : 1H); from 7.20 to 7.35 (mt : 8H); 7.48 (d, J = 4 Hz : 1H); 7.67 (d, J = 4 Hz : 1H)].

Example 36

By carrying out the operation according to the procedure of Example 31, starting with 71.9 mg of 5-methylsulfonyl-3-methyl-4-vinylthiophene-2-carboxylic acid, 26.4 mg of hydroxybenzotriazole in solution in

0.5 cm^3 of dimethylformamide, 0.0302 cm^3 of diisopropylcarbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane, (5-methylsulfonyl-3-methyl-4-vinylthiophene-2-carboxy){1-[bis-(4-chlorophenyl)-methyl]azetidin-3-yl}amide is obtained in the form of a white powder [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.47 (s : 3H); 2.97 (mt : 2H); 3.14 (s : 3H); 3.57 (dd, J = 8 and 7.5 Hz : 2H); 4.32 (s : 1H); 4.65 (mt : 1H); 5.69 (dd, J = 18 and 1 Hz : 1H); 5.77 (dd, J = 12 and 1 Hz : 1H); 6.30 (broad d, J = 7.5 Hz : 1H); 6.96 (dd, J = 18 and 12 Hz : 1H); from 7.20 to 7.35 (mt : 8H)].

Example 37

By carrying out the operation according to the procedure of Example 31, starting with 62.6 mg of 3-methylsulfonylmethylbenzoic acid, 26.4 mg of hydroxybenzotriazole in solution in 0.5 cm^3 of dimethylformamide, 0.0302 cm^3 of diisopropyl-carbodiimide, a solution of 30 mg of 1-[bis(4-chlorophenyl)methyl]azetidin-3-ylamine in 0.5 cm^3 of anhydrous dichloromethane, and 3 cm^3 of anhydrous dichloromethane, (5-methylsulfonyl-3-methyl-4-vinylthiophene-2-carboxy){1-[bis(4-chlorophenyl)-methyl]azetidin-3-yl}amide is obtained in the form of white needles [^{1}H NMR spectrum (300 MHz, $(CD_3)_2SO$ d6 with addition of $CDCl_3$, δ in ppm): 2.84 (s : 3H); 3.02

(broad t, J = 7 Hz : 2H); 3.48 (t, J = 7 Hz : 2H); 4.38 (s : 3H); 4.53 (mt : 1H); 7.21 (d, J = 8 Hz : 4H); 7.34 (d, J = 8 Hz : 4H); 7.40 (t, J = 7.5 Hz : 1H); 7.53 (broad d, J = 7.5 Hz : 1H); 7.84 (broad d, J = 7.5 Hz : 1H); 7.89 (broad s : 1H); 8.54 (d, J = 7 Hz : 1H)].

Example 38

(RS)-N-{1-[(4-chlorophenyl)pyridin-3-yl-methyl]azetidin-3-yl}-3,5-difluorobenzenesulfonamide may be obtained in the following manner: 0.46 g of potassium carbonate and 41 mg of potassium iodide are added to a mixture of 0.3 g of (RS)-3-[bromo-(4-chlorophenyl)methyl]pyridine hydrobromide and 0.28 g of N-azetidin-3-yl-3,5-difluorobenzenesulfonamide hydrochloride in 20 cm^3 of acetonitrile, and then the mixture is heated under reflux for 4 hours. After cooling to a temperature in the region of 20°C, the insoluble matter is removed by filtration and then the mixture is concentrated to dryness under reduced pressure. The residue obtained is taken up in 100 cm^3 of ethyl acetate. The organic phase is washed with twice 50 cm^3 of water, dried over magnesium sulfate in the presence of animal charcoal, filtered on Celite, and then concentrated to dryness under reduced pressure. 230 mg of an orange-colored solid are obtained, which solid is dissolved in a cyclohexane-ethyl acetate mixture (50-50 by volume mixture) and purified by chromatography under pressure on a cartridge of 10 g of silica with the same eluent mixture, with a flow rate

of 6 cm^3/minute. Fractions 22 to 56 are combined and concentrated to dryness under reduced pressure. 100 mg of (RS)-N-{1-[(4-chlorophenyl)pyridin-3-yl-methyl]azetidin-3-yl}-3,5-difluorobenzenesulfonamide are thus obtained in the form of a pale yellow foam melting at 70°C [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.81 (mt : 2H); 3.42 (mt : 2H); 4.03 (mt : 1H); 4.29 (s : 1H); 5.43 (d, J = 9 Hz : 1H); 7.01 (tt, J = 9 and 2.5 Hz : 1H); 7.22 (dd, J = 8 and 5 Hz : 1H); 7.28 (mt : 4H); 7.36 (mt : 2H); 7.62 (broad d, J = 8 Hz : 1H); 8.48 (dd, J = 5 and 1 Hz : 1H); 8.59 (d, J = 1 Hz : 1H)].

(RS)-3-[bromo-(4-chlorophenyl)methyl]pyridine is obtained in the following manner: 3.5 cm^3 of a 48% hydrobromic acid solution in acetic acid and 1 cm^3 of acetyl bromide are added to 1.5 g of (4-chlorophenyl)-pyridin-3-ylmethanol. The amber-colored mixture thus obtained is heated under reflux for 4 hours and then cooled to 20 °C, and concentrated to dryness at 40°C under 2.7 kPa, giving 1.53 g of (RS)-3-[bromo-(4-chlorophenyl)methyl]pyridine (Rf = 75/90, 254 nm, silica plates, reference 1.05719, Merck KGaA, 64271 Darmstadt, Germany).

N-azetidin-3-yl-3,5-difluorobenzene-sulfonamide hydrochloride may be prepared in the following manner: in a 2000 cm^3 hydrogenator, a solution of 7.5 g of N-(1-benzhydrylazetidin-3-yl)-3,5-difluorobenzenesulfonamide in a mixture of 10 cm^3 of

concentrated hydrochloric acid (36% by weight), 1.7 cm^3 of acetic acid and 500 cm^3 of methanol is hydrogenated in the presence of 4.21 g of palladium hydroxide on carbon (20% by weight of catalyst) under 1.7 bar of hydrogen for about 20 hours. The catalyst is removed by filtration on a bed of Celite and then the filtrate is concentrated to dryness under reduced pressure. The residue obtained is beaten with 100 cm^3 of diisopropyl ether for about 16 hours at a temperature in the region of 20°C. The suspension is filtered, and the solid residue is again beaten with 100 cm^3 of diethyl ether at a temperature in the region of 20°C. After filtration, the paste obtained is dried under reduced pressure at a temperature in the region of 40°C. 5.52 g of N-azetidin-3-yl-3,5-difluorobenzenesulfonamide hydrochloride are thus obtained in the form of a white powder.

N-(1-benzhydrylazetidin-3-yl)-3,5-difluorobenzenesulfonamide may be prepared in the following manner: 5.1 g of 3,5-difluorobenzenesulfonyl chloride and then 4.2 cm^3 of triethylamine are successively added to a suspension of 5 g of 1-benzhydrylazetidin-3-ylamine in 80 cm^3 of dichloromethane, at a temperature in the region of 20°C. After stirring for 20 hours at a temperature in the region of 20°C, 50 cm^3 of water are added. The organic phase separated after settling is washed with twice 50 cm^3 of water, dried over magnesium sulfate, and

concentrated to dryness under reduced pressure. 8.99 g of a yellow oil are thus obtained, which oil crystallizes little by little. 4.5 g of this product is purified by chromatography under pressure on 500 g of Amicon silica (diameter of the particles from 0.020 to 0.045 mm), eluting with a methanol-dichloromethane (1-99 by volume) mixture. The fractions containing the desired product are combined and concentrated to dryness under reduced pressure to give 3.58 g of N-(1-benzhydrylazetidin-3-yl)-3,5-difluorobenzene-sulfonamide in the form of a beige powder. The remaining quantity of the preceding yellow oil is purified under the same conditions and provides 3.92 g of N-(1-benzhydrylazetidin-3-yl)-3,5-difluorobenzene-sulfonamide in the form of a beige powder.

1-Benzhydrylazetidin-3-ylamine may be prepared as described in J. Antibiot., 39(9), 1243-1256, 1986.

3,5-Difluorobenzenesulfonyl chloride may be prepared as described in patent: FR2757509.

Example 39

(RS)-N-{1-[(4-chlorophenyl)pyrimidin-5-yl-methyl]azetidin-3-yl}-3,5-difluorobenzenesulfonamide may be obtained by carrying out the procedure as for the preparation of (RS)-N-{1-[(4-chlorophenyl)pyridin-3-ylmethyl]azetidin-3-yl}-3,5-difluorobenzene-sulfonamide: starting with 0.64 g of (RS)-5-[bromo-(4-chlorophenyl)methyl]pyrimidine hydrobromide, 0.5 g

of N-azetidin-3-yl-3,5-difluorobenzenesulfonamide hydrochloride in 20 cm^3 of acetonitril, 1.213 g of potassium carbonate and 379 mg of potassium iodide, 71 mg of (RS)-N-{1-[(4-chlorophenyl)pyrimidin-5-yl-methyl]azetidin-3-yl}-3,5-difluorobenzenesulfonamide are thus obtained in the form of a yellow foam [^{1}H NMR spectrum (300 MHz, $CDCl_3$, δ in ppm): 2.83 (mt : 2H); 3.46 (mt : 2H); 4.03 (mt : 1H); 4.30 (s : 1H); 5.00 (d, J = 9 Hz : 1H); 7.04 (tt, J = 9 and 2.5 Hz : 1H); from 7.20 to 7.35 (mt : 4H); 7.37 (mt : 2H); 8.69 (s : 2H); 9.09 (s : 1H)].

(RS)-5-[bromo-(4-chlorophenyl)methyl]-pyrimidine may be prepared by carrying out the procedure as for the preparation of (RS)-3-[bromo-(4-chlorophenyl)methyl]pyridine using, as raw material, (4-chlorophenyl)pyrimidin-5-ylmethanol.

(4-Chlorophenyl)pyrimidin-5-ylmethanol may be prepared by carrying out the procedure as for (4-chlorophenyl)pyridin-3-ylmethanol, starting with pyrimidine-5-carboxaldehyde and 4-chlorophenylmagnesium bromide.

The medicaments according to the invention consist of a compound of formula (I) or an isomer or a salt of such a compound, in the pure state or in the form of a composition in which it is combined with any other pharmaceutically compatible product which may be inert or physiologically active. The medicaments

according to the invention may be used orally, parenterally, rectally or topically.

As solid compositions for oral administration, tablets, pills, powders (gelatin capsules, sachets) or granules may be used. In these compositions, the active ingredient according to the invention is mixed with one or more inert diluents, such as starch, cellulose, sucrose, lactose or silica, under an argon stream. These compositions may also comprise substances other than diluents, for example one or more lubricants such as magnesium stearate or talc, a coloring, a coating (sugar-coated tablets) or a glaze.

As liquid compositions for oral administration, there may be used pharmaceutically acceptable solutions, suspensions, emulsions, syrups and elixirs containing inert diluents such as water, ethanol, glycerol, vegetable oils or paraffin oil. These compositions may comprise substances other than diluents, for example wetting, sweetening, thickening, flavoring or stabilizing products.

Sterile compositions for parenteral administration may be preferably solutions which are aqueous or nonaqueous, suspensions or emulsions. As solvent or vehicle, there may be used water, propylene glycol, polyethylene glycol, vegetable oils, in particular olive oil, injectable organic esters, for example ethyl oleate or other suitable organic

solvents. These compositions may also contain adjuvants, in particular wetting, isotonizing, emulsifying, dispersing and stabilizing agents. Sterilization may be carried out in several ways, for example by asepticizing filtration, by incorporating sterilizing agents into the composition, by irradiation or by heating. They may also be prepared in the form of sterile solid compositions which may be dissolved at the time of use in sterile water or any other injectable sterile medium.

Compositions for rectal administration are suppositories or rectal capsules which contain, in addition to the active product, excipients such as cocoa butter, semisynthetic glycerides or polyethylene glycols.

Compositions for topical administration may be, for example, creams, lotions, collyria, collutoria, nasal drops or aerosols.

In human therapy, the compounds according to the invention are particularly useful for the treatment and/or prevention of psychoses including schizophrenia, anxiety disorders, depression, epilepsy, neurodegeneration, cerebellar and spinocerebellar disorders, cognitive disorders, cranial trauma, panic attacks, peripheral neuropathies, glaucomas, migraine, Parkinson's disease, Alzheimer's disease, Huntington's chorea, Raynaud's syndrome, tremor, obsessive-compulsive disorder, senile dementia, thymic disorders,

Tourette's syndrome, tardive dyskinesia, bipolar disorders, cancers, movement disorders induced by medicaments, dystonia, endotoxemic shocks, hemorrhagic shocks, hypotension, insomnia, immunological diseases, multiple sclerosis, vomiting, asthma, appetite disorders (bulimia, anorexia), obesity, memory disorders, intestinal transit disorders, in weaning from chronic treatments and alcohol or drug abuse (opioids, barbiturates, cannabis, cocaine, amphetamine, phencyclide, hallucinogens, benzodiazepines for example), as analgesics or potentiators of the analgesic activity of the narcotic and nonnarcotic drugs.

The doses depend on the desired effect, the duration of the treatment and the route of administration used; they are generally between 5 mg and 1000 mg per day orally for an adult with unit doses ranging from 1 mg to 250 mg of active substance.

In general, the doctor will determine the appropriate dosage depending on the age, weight and any other factors specific to the subject to be treated.

The following examples illustrate the compositions according to the invention:

EXAMPLE A

Gelatin capsules containing a dose of 50 mg of active product and having the following composition are prepared according to the usual technique:

- Compound of formula (I) ..... 50 mg
- Cellulose ..... 18 mg
- Lactose ..... 55 mg
- Colloidal silica ..... 1 mg
- Sodium carboxymethylstarch ..... 10 mg
- Talc ..... 10 mg
- Magnesium stearate ..... 1 mg

EXAMPLE B

Tablets containing a dose of 50 mg of active product and having the following composition are prepared according to the usual technique:

- Compound of formula (I) ..... 50 mg
- Lactose ..... 104 mg
- Cellulose ..... 40 mg
- Polyvidone ..... 10 mg
- Sodium carboxymethylstarch ..... 22 mg
- Talc ..... 10 mg
- Magnesium stearate ..... 2 mg
- Colloidal silica ..... 2 mg
- Mixture of hydroxymethylcellulose, glycerin, titanium oxide (72-3.5-24.5) qs 1 finished film-coated tablet containing ..... 245 mg


EXAMPLE C

An injectable solution containing 10 mg of active product and having the following composition is prepared:

- Compound of formula (I)	10 mg
- Benzoic acid	80 mg
- Benzyl alcohol	0.06 ml
- Sodium benzoate	80 mg
- Ethanol, 95%	0.4 ml
- Sodium hydroxide	24 mg
- Propylene glycol	1.6 ml
- Water.....	qs 4 ml

CLAIMS

1. Pharmaceutical composition containing, as active ingredient, a compound of formula:



(I)

in which

R_1 represents a radical $-NHCOR_4$ or $-N(R_5)-Y-R_6$,

Y is CO or SO_2,

R_2 and R_3, which are identical or different, represent either an aromatic chosen from phenyl, naphthyl and indenyl, these aromatics being unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, formyl, hydroxyl, trifluoromethyl, trifluoromethoxy, -CO-alk, cyano, -COOH, -COOalk, $-CONR_7R_8$, $-CO-NH-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, hydroxyalkyl or $-alk-NR_7R_8$ radicals; or a heteroaromatic chosen from benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzothienyl, pyrimidinyl, furyl, imidazolyl, isochromanyl, isoquinolyl, pyrrolyl, pyridyl, quinolyl, 1,2,3,4-tetrahydroisoquinolyl, thiazolyl and thienyl rings, it

being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, cyano, -COOH, -COOalk, -CO-NH-NR_9R_{10}, -$CONR_7R_8$, -alk-NR_9R_{10}, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl or hydroxyalkyl radical,

R_4 represents a radical -alk-SO_2-R_{11}, -alk-SO_2-CH=CH-R_{11}, Het substituted with -SO_2-R_{11} or phenyl substituted with -SO_2-R_{11} or -alk-SO_2-R_{11},

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a phenylalkyl, Het or Ar radical,

R_7 and R_8, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_7 and R_8 together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_9 and R_{10}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_9 and R_{10} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally

substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

R_{11} represents an alkyl, Ar or Het radical,

Ar represents a phenyl, naphthyl or indenyl radical, these radicals being optionally substituted with one or more halogen atoms or alkyl, alkoxy, cyano, -CO-alk, -COOH, -COOalk, $-CONR_{12}R_{13}$, $-CO-NH-NR_{14}R_{15}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, $-alk-NR_{14}R_{15}$, $-NR_{14}R_{15}$, alkylthioalkyl, formyl, hydroxyl, hydroxyalkyl, Het, -O-alk-NH-cycloalkyl, OCF_3, CF_3, -NH-CO-alk, $-SO_2NH_2$, $-NH-COCH_3$, -NH-COOalk or Het radicals or alternatively, on 2 adjacent carbon atoms, with a dioxymethylene,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo, hydroxyl, OCF_3 or CF_3 radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form,

R_{12} and R_{13}, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_{12} and R_{13} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and

being optionally substituted with one or more alkyl radicals,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

alk represents an alkyl or alkylene radical,

the alkyl and alkylene radicals and portions and the alkoxy radicals and portions are in the form of a straight or branched chain and contain 1 to 6 carbon atoms and the cycloalkyl radicals contain 3 to 10 carbon atoms,

an optical isomer of such a compound or one of its pharmaceutically acceptable salts.

2. Composition according to claim 1 for which in the compound of formula (I) Het is chosen from benzimidazole, benzoxazole, benzothiazole, benzothiophene, cinnoline, thiophene, quinazoline, quinoxaline, quinoline, pyrazole, pyrrole, pyridine, imidazole, indole, isoquinoline, pyrimidine, thiazole, thiadiazole, piperidine, piperazine, pyrrolidine,

triazole, furan, tetrahydroisoquinoline, tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo, hydroxyl, OCF_3 or CF_3 radicals.

3. Pharmaceutical composition containing, as active ingredient, at least one compound of formula (I) according to claim 1 in which

R_1 represents a radical $-N(R_5)-Y-R_6$,

Y is SO_2,

R_2 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy, cyano, $-CONR_7R_8$, hydroxyalkyl or $-alk-NR_7R_8$ radicals; or a heteroaromatic chosen from the pyridyl, pyrimidinyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, $-CONR_7R_8$, $-alk-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl or hydroxyalkyl radical,

R_3 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy, cyano, $-CONR_7R_8$, hydroxyalkyl or $-alk-NR_7R_8$ radicals; or a heteroaromatic chosen from the pyridyl, pyrimidinyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or

substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, $-CONR_7R_8$, $-alk-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl or hydroxyalkyl radical,

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a naphthyl, phenylalkyl, Het or phenyl radical optionally substituted with one or more halogen atoms or alkyl, alkoxy, cyano, -CO-alk, COOalk, $-CONR_{12}R_{13}$, $-alk-NR_{14}R_{15}$, $-NR_{14}R_{15}$, hydroxyl, hydroxyalkyl, Het, OCF_3, CF_3, -NH-CO-alk, $-SO_2NH_2$ or -NH-COOalk radicals, or alternatively, on 2 adjacent carbon atoms, with dioxymethylene,

R_7 and R_8, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_7 and R_8 together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_9 and R_{10}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_9 and R_{10} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally

substituted with one or more alkyl, oxo or $-CO-NH_2$ radicals,

R_{12} and R_{13}, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_{12} and R_{13} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, oxo, hydroxyalkyl or $-CO-NH_2$ radicals,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo or hydroxyl radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form and, preferably, Het represents a heterocycle chosen

from the following heterocycles: benzimidazole, benzoxazole, benzothiazole, benzothiophene, thiophene, quinazoline, quinoxaline, quinoline, pyrrole, pyridine, imidazole, indole, isoquinoline, pyrimidine, thiazole, thiadiazole, furan, tetrahydroisoquinoline and tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, oxo, hydroxyl, OCF_3 or CF_3 radicals,

an optical isomer of such a compound or one of its pharmaceutically acceptable salts.

4. Pharmaceutical composition containing, as active ingredient, at least one compound of formula (I) according to claim 1 in which

R_1 represents a radical $-N(R_5)-Y-R_6$,

Y is SO_2,

R_2 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy or hydroxyalkyl radicals; or a heteroaromatic chosen from pyridyl and pyrimidyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl or trifluoromethoxy radical,

R_3 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy or hydroxyalkyl radicals; or a heteroaromatic chosen from

pyridyl and pyrimidyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl or trifluoromethoxy radical,

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a naphthyl, phenylalkyl, Het or phenyl radical optionally substituted with one or more halogen atoms or alkyl, alkoxy, $-NR_{14}R_{15}$, hydroxyl, hydroxyalkyl, OCF_3, CF_3 or $-SO_2NH_2$ radicals, or alternatively, on 2 adjacent carbon atoms, with dioxymethylene,


R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, oxo, hydroxyalkyl or $-CO-NH_2$ radicals,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo or hydroxyl radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form and, preferably, Het represents a heterocycle chosen

from the following heterocycles: benzimidazole, benzoxazole, benzothiazole, benzothiophene, thiophene, quinoline, pyrrole, pyridine, pyrimidine, thiazole, thiadiazole, furan, tetrahydroisoquinoline and tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, oxo, hydroxyl, OCF_3 or CF_3 radicals,

an optical isomer of such a compound or one of its pharmaceutically acceptable salts.

5. Compounds of formula:



(I)

in which

R_1 represents a radical $-NHCOR_4$ or $-N(R_5)-Y-R_6$,

Y is CO or SO_2,

R_2 and R_3, which are identical or different, represent either an aromatic chosen from phenyl, naphthyl and indenyl, these aromatics being unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, formyl, hydroxyl, trifluoromethyl, trifluoromethoxy, -CO-alk, cyano, -COOH, -COOalk, $-CONR_7R_8$, $-CO-NH-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkyl-

sulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, hydroxyalkyl or -alk-NR_7R_8 radicals; or a heteroaromatic chosen from benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzothienyl, pyrimidinyl, furyl, imidazolyl, isochromanyl, isoquinolyl, pyrrolyl, pyridyl, quinolyl, 1,2,3,4-tetrahydro-isoquinolyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, cyano, -COOH, -COOalk, -CO-NH-NR_9R_{10}, -$CONR_7R_8$, -alk-NR_9R_{10}, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, alkylsulfanylalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl or hydroxyalkyl radical,

R_4 represents a radical -alk-SO_2-R_{11}, -alk-SO_2-CH=CH-R_{11}, Het substituted with -SO_2-R_{11} or phenyl substituted with -SO_2-R_{11} or -alk-SO_2-R_{11},

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a phenylalkyl, Het or Ar radical,

R_7 and R_8, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_7 and R_8 together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_9 and R_{10}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_9 and R_{10} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

R_{11} represents an alkyl, Ar or Het radical,

Ar represents a phenyl, naphthyl or indenyl radical, these radicals being optionally substituted with one or more halogen atoms or alkyl, alkoxy, cyano, -CO-alk, -COOH, -COOalk, $-CONR_{12}R_{13}$, $-CO-NH-NR_{14}R_{15}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, $-alk-NR_{14}R_{15}$, $-NR_{14}R_{15}$, alkylthioalkyl, formyl, hydroxyl, hydroxyalkyl, Het, -O-alk-NH-cycloalkyl, OCF_3, CF_3, -NH-CO-alk, $-SO_2NH_2$, $-NH-COCH_3$, -NH-COOalk or Het radicals, or alternatively, on 2 adjacent carbon atoms, with a dioxymethylene,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo, hydroxyl, OCF_3 or CF_3 radicals, the nitrogen-

containing heterocycles being optionally in their N-oxidized form,

R_{12} and R_{13}, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_{12} and R_{13} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, -COOalk, cycloalkyl, alkylcycloalkyl, -alk-O-alk or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, -COalk, -COOalk, -CO-NHalk, -CS-NHalk, oxo, hydroxyalkyl, -alk-O-alk or $-CO-NH_2$ radicals,

alk represents an alkyl or alkylene radical,

the alkyl and alkylene radicals and portions and the alkoxy radicals and portions are in the form of a straight or branched chain and contain 1 to 6 carbon atoms and the cycloalkyl radicals contain 3 to 10 carbon atoms,

their optical isomers and their pharmaceutically acceptable salts,

with the exception of the compound for which for which R_2 and R_3 represent phenyl radicals, R_1 represents a radical $-N(R_5)-Y-R_6$, Y is SO_2, R_5 represents a methyl radical and R_6 represents a phenyl radical.

6. Compounds of formula (I) according to claim 5 for which Het is chosen from benzimidazole, benzoxazole, benzothiazole, benzothiophene, cinnoline, thiophene, quinazoline, quinoxaline, quinoline, pyrazole, pyrrole, pyridine, imidazole, indole, isoquinoline, pyrimidine, thiazole, thiadiazole, piperidine, piperazine, pyrrolidine, triazole, furan, tetrahydroisoquinoline, tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo, hydroxyl, OCF_3 or CF_3 radicals for which R_2 and R_3 represent phenyl radicals, R_1 represents a radical $-N(R_5)-Y-R_6$, Y is SO_2, R_5 represents a methyl radical and R_6 represents a phenyl radical.

7. Compounds of formula (I) according to claim 5 in which

R_1 represents a radical $-N(R_5)-Y-R_6$,

Y is SO_2,

R_2 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy, cyano, $-CONR_7R_8$, hydroxyalkyl or $-alk-NR_7R_8$ radicals; or a

heteroaromatic chosen from the pyridyl, pyrimidinyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, $-CONR_7R_8$, $-alk-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl or hydroxyalkyl radical,

R_3 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy, cyano, $-CONR_7R_8$, hydroxyalkyl or $-alk-NR_7R_8$ radicals; or a heteroaromatic chosen from the pyridyl, pyrimidinyl, thiazolyl and thienyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl, trifluoromethoxy, $-CONR_7R_8$, $-alk-NR_9R_{10}$, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl or hydroxyalkyl radical,

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a naphthyl, phenylalkyl, Het or phenyl radical optionally substituted with one or more halogen atoms or alkyl, alkoxy, cyano, -CO-alk, COOalk, $-CONR_{12}R_{13}$, $-alk-NR_{14}R_{15}$, $-NR_{14}R_{15}$, hydroxyl, hydroxyalkyl, Het, OCF_3, CF_3, -NH-CO-alk, $-SO_2NH_2$ or -NH-COOalk radicals, or alternatively, on 2 adjacent carbon atoms, with dioxymethylene,

R_7 and R_8, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_7

and R_8 together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_9 and R_{10}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_9 and R_{10} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, oxo or $-CO-NH_2$ radicals,

R_{12} and R_{13}, which are identical or different, represent a hydrogen atom or an alkyl radical or alternatively R_{12} and R_{13} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl radicals,

R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen

atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, oxo, hydroxyalkyl or $-CO-NH_2$ radicals,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo or hydroxyl radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form and, preferably, Het represents a heterocycle chosen from the following heterocycles: benzimidazole, benzoxazole, benzothiazole, benzothiophene, thiophene, quinazoline, quinoxaline, quinoline, pyrrole, pyridine, imidazole, indole, isoquinoline, pyrimidine, thiazole, thiadiazole, furan, tetrahydroisoquinoline and tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, oxo, hydroxyl, OCF_3 or CF_3 radicals,

their optical isomers and their pharmaceutically acceptable salts,

with the exception of the compound for which R_2 and R_3 represent phenyl radicals, R_1 represents a radical

$-N(R_5)-Y-R_6$, Y is SO_2, R_5 represents a methyl radical and R_6 represents a phenyl radical.

8. Compounds of formula (I) according to claim 5 in which

R_1 represents a radical $-N(R_5)-Y-R_6$,

Y is SO_2,

R_2 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy or hydroxyalkyl radicals; or a heteroaromatic chosen from pyridyl and pyrimidyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl or trifluoromethoxy radical,

R_3 represents either a phenyl which is unsubstituted or substituted with one or more halogen atoms or alkyl, alkoxy, trifluoromethyl, trifluoromethoxy or hydroxyalkyl radicals; or a heteroaromatic chosen from pyridyl and pyrimidyl rings, it being possible for these heteroaromatics to be unsubstituted or substituted with a halogen atom or an alkyl, alkoxy, hydroxyl, trifluoromethyl or trifluoromethoxy radical,

R_5 represents a hydrogen atom or an alkyl radical,

R_6 represents a naphthyl, phenylalkyl, Het or phenyl radical optionally substituted with one or more halogen atoms or alkyl, alkoxy, $-NR_{14}R_{15}$, hydroxyl, hydroxyalkyl, OCF_3, CF_3 or $-SO_2NH_2$ radicals, or alternatively, on 2 adjacent carbon atoms, with dioxymethylene,


R_{14} and R_{15}, which are identical or different, represent a hydrogen atom or an alkyl, cycloalkyl, alkylcycloalkyl or hydroxyalkyl radical or alternatively R_{14} and R_{15} together form with the nitrogen atom to which they are attached a 3- to 10-membered saturated or unsaturated mono- or bicyclic heterocycle, optionally containing another heteroatom chosen from oxygen, sulfur and nitrogen and being optionally substituted with one or more alkyl, oxo, hydroxyalkyl or $-CO-NH_2$ radicals,

Het represents a 3- to 10-membered unsaturated or saturated mono- or bicyclic heterocycle containing one or more heteroatoms chosen from oxygen, sulfur and nitrogen optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, alkoxycarbonyl, oxo or hydroxyl radicals, the nitrogen-containing heterocycles being optionally in their N-oxidized form and, preferably, Het represents a heterocycle chosen from the following heterocycles: benzimidazole, benzoxazole, benzothiazole, benzothiophene, thiophene, quinoline, pyrrole, pyridine, pyrimidine, thiazole, thiadiazole, furan, tetrahydroisoquinoline and tetrahydroquinoline, these heterocycles being optionally substituted with one or more halogen atoms or alkyl, alkoxy, vinyl, oxo, hydroxyl, OCF_3 or CF_3 radicals,

their optical isomers and their pharmaceutically acceptable salts,


with the exception of the compound for which R_2 and R_3 represent phenyl radicals, R_1 represents a radical $-N(R_5)-Y-R_6$, Y is SO_2, R_5 represents a methyl radical and R_6 represents a phenyl radical.

9. Process for preparing the compounds of formula (I) according to claim 5 for which R_1 represents a radical $-NHCOR_4$, characterized in that an acid R_4COOH for which R_4 has the same meanings as in claim 5 is reacted with a derivative of formula:




in which R_2 and R_3 have the same meanings as in claim 5, the product isolated and optionally converted to a pharmaceutically acceptable salt.

10. Process for preparing the compounds of formula (I) according to claim 5 for which R_1 represents a radical $-N(R_5)-Y-R_6$, characterized in that a derivative of formula:



in which R_2 and R_3 have the same meanings as in claim 5 is reacted with a derivative Hal-Y-R_6 for which Y and R_6 have the same meanings as in claim 5 and Hal represents a halogen atom, optionally followed by a derivative Hal-alk, Hal represents a halogen atom and alk represents an alkyl radical (1-6C in the form of a straight or branched chain) to give the compounds for which R_5 is an alkyl radical, the product isolated and optionally converted to a pharmaceutically acceptable salt.

11. Process for preparing the compounds of formula (I) according to claim 5, characterized in that a derivative R_2-CHBr-R_3 for which R_2 and R_3 have the same meanings as in claim 5 is reacted with a derivative of formula:



in which R_1 has the same meanings as in claim 5, the product isolated and optionally converted to a pharmaceutically acceptable salt.

12. Process for preparing the compounds of formula (I) according to claim 5 for which R_1 represents a radical -N(R_5)-Y-R_6 in which R_6 is a phenyl radical substituted with a hydroxyl radical, characterized in that a corresponding compound of formula (I) for which R_1 represents a radical -N(R_5)-Y-R_6 in which R_6 is a phenyl radical substituted with an alkoxy radical is

hydrolyzed, the product isolated and optionally converted to a pharmaceutically acceptable salt.

13. Process for preparing the compounds of formula (I) according to claim 5 for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with a hydroxy(1C)alkyl radical, characterized in that diisobutylaluminum hydride is reacted with a corresponding compound of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with an alkoxycarbonyl radical, the product isolated and optionally converted to a pharmaceutically acceptable salt.

14. Process for preparing the compounds of formula (I) according to claim 5 for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with a 1-pyrrolidinyl radical, characterized in that pyrrolidine is reacted with a corresponding compound of formula (I) for which R_1 represents a radical $-N(R_5)-Y-R_6$ in which R_6 is a phenyl radical substituted with a fluorine atom, the product isolated and optionally converted to a pharmaceutically acceptable salt.