What is claimed is:

1. An interleukin-4 receptor (IL-4R) antagonist for use in ~~method for enhancing the~~the enhancement of efficacy and/or tolerability of a grass allergen-specific subcutaneous immunotherapy (SCIT) regimen in a subject having a grass allergy, ~~the method comprising administering to the subject~~wherein one or more doses of ~~an~~ the ~~interleukin-4 receptor (~~IL-4R~~)~~ antagonist is administered in combination with the SCIT regimen, wherein at least one dose of the IL-4R antagonist is administered prior to the start of the SCIT regimen, and wherein the IL-4R antagonist is an anti-IL-4R antibody, or an antigen-binding fragment thereof, that comprises the heavy chain complementarity determining regions (HCDRs) of a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO:1 and the light chain complementarity determining regions (LCDRs) of a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO:2.

2. The ~~method~~ IL-4R antagonist of claim 1, wherein the SCIT regimen comprises subcutaneous administration of a grass extract that is from a grass selected from the group consisting of Timothy, Bahia, Bermuda, Johnson, Kentucky bluegrass, Orchard, Redtop, Rye, Sweet Vernal, Meadow Fescue, and combinations thereof.

3. The ~~method~~ IL-4R antagonist of claim 2, wherein the grass extract is from Timothy grass.

4. The ~~method~~ IL-4R antagonist of any one of claims 1 to 3, wherein the SCIT regimen comprises a cluster SCIT regimen.

5. The ~~method~~ IL-4R antagonist of claim 4, wherein the cluster SCIT regimen comprises an up-titration regimen followed by a maintenance regimen, wherein the up-titration regimen comprises administering increasing doses of the grass extract over a period of 4 to 12 weeks and wherein the maintenance regimen comprises administering one or more maintenance doses of the grass extract at the highest dose administered during the up-titration regimen.

6. The IL-4R antagonist~~method~~ of claim 5, wherein the up-titration regimen comprises administering increasing doses of the grass extract over a period of 8 weeks.

7. The IL-4R antagonist~~method~~ of claim 5 or 6, wherein the maintenance regimen comprises administering maintenance doses every 1 to 4 weeks for at least 8 weeks.

8. The IL-4R antagonist~~method~~ of any one of claims 5 to 7, wherein the up-titration regimen comprises up-titrating from a dose of 1 bioequivalent allergy unit (BAU) to a dose of

4,000 BAU and wherein the maintenance regimen comprises administering one or more maintenance doses at 4,000 BAU.

9. The IL-4R antagonist~~method~~ of any one of claims 1 to 8, wherein the IL-4R antagonist is administered at a dose of about 75 mg to about 600 mg.

10. The IL-4R antagonist~~method~~ of any one of claims 1 to 9, wherein the IL-4R antagonist is administered as an initial dose followed by one or more secondary doses, wherein each secondary dose is administered 1 to 4 weeks after the immediately preceding dose.

11. The IL-4R antagonist~~method~~ of claim 10, wherein the initial dose of the IL-4R antagonist is administered from 1 to 7 days before the start of the SCIT regimen.

12. The IL-4R antagonist~~method~~ of claim 10 or 11, wherein the initial dose comprises 600 mg and each secondary dose comprises 300 mg of the IL-4R antagonist.

13. The IL-4R antagonist~~method~~ of any one of claims 10 to 12, wherein each secondary dose is administered 2 weeks after the immediately preceding dose.

14. The IL-4R antagonist~~method~~ of any one of claims 1 to 13, wherein the IL-4R antagonist and the SCIT are not administered to the subject on the same day.

15. The IL-4R antagonist~~method~~ of any one of claims 1 to 14, wherein ~~enhancing the~~the enhancement of efficacy and/or tolerability of the SCIT regimen comprises reducing allergic rhinitis symptoms in the subject.

16. The IL-4R antagonist~~method~~ of any one of claims 1 to 15, wherein ~~enhancing the~~the enhancement of efficacy and/or tolerability of the SCIT regimen comprises:

(a) increasing the amount of serum grass allergen-specific IgG4 (sIgG4) in the subject relative to SCIT monotherapy;

(b) decreasing the amount of serum grass allergen-specific IgE (sIgE) in the subject relative to SCIT monotherapy; and/or

(c) increasing the ratio of sIgG4 to sIgE in the subject relative to SCIT monotherapy.

17. The IL-4R antagonist~~method~~ of any one of claims 1 to 16, wherein administration of the IL-4R antagonist in combination with the SCIT regimen reduces or inhibits the induction of sIgE during the SCIT up-titration regimen and/or the SCIT maintenance regimen.

18. The IL-4R antagonist~~method~~ of any one of claims 1 to 17, wherein administration of the IL-4R antagonist in combination with the SCIT regimen increases the maximum SCIT dose that is tolerated by the subject.

19. The IL-4R antagonist~~method~~ of any one of claims 1 to 18, wherein administration of the IL-4R antagonist in combination with the SCIT regimen reduces the use of epinephrine or an oral steroid as a rescue medication.

20. The IL-4R antagonist~~method~~ of any one of claims 1 to 19, wherein the anti-IL-4R antibody or antigen-binding fragment thereof comprises three HCDRs (HCDR1, HCDR2 and HCDR3) and three LCDRs (LCDR1, LCDR2 and LCDR3), wherein the HCDR1 comprises the amino acid sequence of SEQ ID NO:3; the HCDR2 comprises the amino acid sequence of SEQ ID NO:4; the HCDR3 comprises the amino acid sequence of SEQ ID NO:5; the LCDR1 comprises the amino acid sequence of SEQ ID NO:6; the LCDR2 comprises the amino acid sequence of SEQ ID NO:7; and the LCDR3 comprises the amino acid sequence of SEQ ID NO:8.

21. The IL-4R antagonist~~method~~ of any one of claims 1 to 20, wherein the anti-IL-4R antibody or antigen-binding fragment thereof comprises a HCVR comprising the amino acid sequence of SEQ ID NO:1 and comprises a LCVR comprising the amino acid sequence of SEQ ID NO:2.

22. The IL-4R antagonist~~method~~ of any one of claims 1 to 21, wherein the anti-IL-4R antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO:9 and a light chain comprising the amino acid sequence of SEQ ID NO:10.

23. The IL-4R antagonist~~method~~ of any one of claims 1 to 22, wherein the IL-4R antagonist is dupilumab or a bioequivalent thereof.

24. The IL-4R antagonist~~method~~ of any one of claims 1 to 23, wherein the IL-4R antagonist is contained in a container selected from the group consisting of a glass vial, a syringe, a pre-filled syringe, a pen delivery device, and an autoinjector.

25. The IL-4R antagonist~~method~~ of claim 24, wherein the IL-4R antagonist is contained in a pre-filled syringe.

26. The IL-4R antagonist~~method~~ of claim 25, wherein the pre-filled syringe is a single-dose pre-filled syringe.

27. The IL-4R antagonist~~method~~ of claim 24, wherein the IL-4R antagonist is contained in an autoinjector.

28. The IL-4R antagonist~~method~~ of claim 24, wherein the IL-4R antagonist is contained in a pen delivery device.