

AMIDE DERIVATIVES AS LYSOPHOSPHATIDIC ACID RECEPTOR ANTAGONISTS

The present invention relates to amide derivatives, processes for their preparation, pharmaceutical compositions containing them and their use in therapy, particularly as
5 lysophosphatidic acid receptor antagonists.

Lysophosphatidic acid (LPA) is a bioactive lipid mediator that interacts with G protein-coupled transmembrane LPA receptors to affect fundamental cellular functions including proliferation, differentiation, survival, migration, adhesion, invasion and morphogenesis.

10 The effects of LPA at physiological concentrations are mediated by six high affinity cognate LPA receptors (LPA1-6) and potentially at least three additional putative LPA receptors. All known LPA receptors are type 1, rhodopsin-like G protein-coupled receptors (GPCRs) that differ in their tissue distribution and downstream signalling pathways.

15 LPAR1 was the first high affinity LPA receptor identified. Along with LPAR2 and 3, LPAR 1 is one of the Endothelial Differentiation Gene (EDG) LPA receptors. The LPAR1 gene is widely expressed in adult mice, with presence demonstrated in brain, uterus, testis, lung, small intestine, heart, stomach, kidney, spleen, thymus, placenta and skeletal muscle.

20 Expression is also widespread in humans. LPAR 1 couples with and activates G α i/o, G α q/11 and G α 12/13 G proteins to induce a range of cellular responses. Studies with mice lacking functional LPA1 receptor alleles have implicated LPAR1 in the development of pulmonary fibrosis and neuropathic pain. Intrathecal injection of LPA causes allodynia, hyperalgesia and demyelination of the dorsal root in a LPAR1-dependent manner.

25 LPAR2 has approximately 60% amino acid homology to LPAR1 and in mouse is expressed in kidney, uterus, testis and lung with lower levels of expression in stomach, spleen, thymus brain and heart. In human tissues high expression is detected in testis and leukocytes and moderate expression in prostate, spleen, thymus and pancreas.

30 LPAR2 couples with G α i/o, G α q/11 and G α 12/13 G proteins with signalling associated with processes such as cell survival and cell migration which makes LPAR2 a potential factor for cancer metastasis. In cancer cells, aberrant expression of LPAR2 has been reported, suggesting a tumour promoting role.

Expression of LPAR3 has been seen in human heart, testis, prostate, pancreas, lung, ovary and brain and most abundantly in mouse testis, kidney, lung, small intestine, heart, stomach, spleen, brain and thymus. LPAR3 is able to couple with $G_{\alpha i/o}$, $G_{\alpha q/11}$ to and is also a potential factor for cancer metastasis.


5

LPA receptors 4, 5 and 6 constitute the non-EDG family of LPA receptors. LPAR4 (P2Y9/GPR23) is more closely related to P2Y purinoreceptors. LPAR4 appears to be broadly expressed in humans with particular abundance in ovary, and is present in mouse heart, skin, thymus and ovary. As with other LPA receptors this GPCR can signal via
10 several pathways, namely $G_{\alpha s}$, $G_{\alpha i}$, $G_{\alpha q/11}$ and $G_{\alpha 12/13}$.

Formerly an orphan GPCR, GPR92 was identified as the fifth LPA receptor and renamed LPAR5. Human LPAR5 is located on chromosome 12p13.31 and encodes an approximately 41kDa protein. LPAR5 is broadly expressed in murine tissues such as small
15 intestine, skin, spleen, stomach, thymus, lung, heart, liver, bladder urothelium, dorsal root ganglion and spinal cord. LPA induces neurite retraction and stress fibre formation in LPAR5-expressing cells through coupling to $G_{\alpha 12/13}$ and is also able to activate $G_{\alpha s}$ and $G_{\alpha q/11}$. It has been shown that LPAR5 can be activated by ligands other than LPA, namely endogenous by-products of cholesterol synthesis such as farnesyl pyrophosphate
20 (FPP) and N-arachidonylglycine (NAG). Activation of GPR92/LPAR5 by FPP increases inositol phosphate production, cAMP levels, and Ca^{2+} levels in a concentration-dependent manner. There is evidence of a role for LPAR5 in pain since it is highly expressed in and largely co-localized with TRPV1 in mouse and human dorsal root ganglion, as well as the fact that FPP induces calcium influx in DRG neurons in a LPAR5-specific manner. The
25 targeted deletion of LPAR5 in mice produces an analgesic phenotype, with LPAR5 homozygotic mice being less sensitive to touch and pain stimuli in a number of tests and also resistant to development of neuropathic pain. Additionally, LPAR5 knock-out mice display a urological phenotype consistent with changes in sensory signalling. LPA causes aggregation of human platelets at concentrations that are present in atheromatous lesions;
30 LPAR5 has been shown to have an intrinsic role in this process.

Most recently, P2Y5 was named as the sixth LPA receptor. This member of the purinergic receptor family is more closely related to LPAR4 and 5 than LPA1-3. This receptor has been identified as a critical mediator of human hair growth.

- 5 EP-A-1,553,075 (Ono Pharmaceutical Co., Ltd.) discloses LPA receptor antagonists of general formula



wherein R represents an aliphatic hydrocarbon group which may be substituted, or a cyclic group which may have a substituent(s);

- 10 G represents a bond or a spacer having from 1 to 8 atoms in its principle chain;

T represents $-\text{CH}_2-$, or a spacer having one atom in its principle chain, the principle chain containing a hydrogen bond acceptable group which may have a substituent(s);

J represents a nitrogen atom or a carbon atom;

- B represents an aliphatic hydrocarbon group which may be substituted, or a cyclic group
15 which may have a substituent(s);

K represents (1) a bond, or (2) a spacer having from 1 to 8 atoms in its principle chain which may form a ring together with a substituent of the cyclic group in R, the ring D or a substituent on the ring D;

Q represents (1) a bond, or (2) a spacer having from 1 to 8 atoms in its principle chain which may form a ring together with the cyclic group in R, a substituent of the cyclic group in R or K;

ring D represents a cyclic group which may have an additional substituent(s);

5 L represents a bond, or a spacer having from 1 to 3 atoms in its principle chain;

ring E represents a cyclic group which may have an additional substituent(s);


M represents a bond, or a spacer having from 1 to 8 atoms in its principle chain;

Z represents an acidic group; and

t represents 0 or 1, or a salt thereof.

10

EP-A-2,481,725 (Astellas Pharma Inc.) discloses LPA receptor antagonists of general formula



wherein A is an aryl which may be substituted or a hetero ring group which may be substituted.

B is a 5-membered aromatic hetero ring group which may be substituted,

X is a single bond or $-(CR^{X1}R^{X2})_n-$,

n is 1, 2, 3 or 4,

R^{X1} and R^{X2} are the same or different from each other, and are H, halogen, OH, -O-(lower alkyl which may be substituted), or lower alkyl which may be substituted, or

R^{X1} and R^{X2} are combined with each other to form oxo (=O), or

R^{X1} and R^{X2} are combined with each other to form C₂₋₅ alkylene which may be substituted, in which when n is 2, 3 or 4, R^{X1} may be combined with adjacent R^{X1} to form a new bond,

Y^1 , Y^2 , Y^3 , Y^4 and Y^5 are the same as or different from each other, and are CR^Y or N,

R^Y 's are the same as or different from each other, and are H, OH, halogen, -O-(lower alkyl which may be substituted), -S-(lower alkyl which may be substituted), lower alkyl which may be substituted, lower alkenyl which may be substituted, or cycloalkyl which may be substituted,

m is 1, 2 or 3,

R^3 is H, or lower alkyl which may be substituted,

R^4 is lower alkyl which may be substituted, lower alkenyl which may be substituted, cycloalkyl which may be substituted, aryl which may be substituted, a hetero ring group which may be substituted, or $NR^{101}R^{102}$, or

R^3 and R^4 may be combined with each other to form C₂₋₅ alkylene which may be substituted, and

R^{101} and R^{102} are the same as or different from each other, and are H, OH, -O-(lower alkyl which may be substituted), -C(=O)-(lower alkyl which may be substituted), -C(=O)-O-(lower alkyl which may be substituted), -NH-C(=O)-(lower alkyl which may be substituted), lower alkyl which may be substituted, lower alkenyl which may be substituted, cycloalkyl which may be substituted, aryl which may be substituted, or a hetero ring group which may be substituted, or R^{101} and R^{102} may be combined with nitrogen atoms to which they are bonded to form a nitrogen-containing monocyclic saturated hetero ring group,

in which when R^4 is $NR^{101}R^{102}$, at least one of R^3 , R^{101} and R^{102} is H.


In addition EP-A-1,229,034 describes certain amide derivatives as PDE4 inhibitors for treating respiratory diseases and US 2004/0176446 describes certain amide derivatives for use in treating cardiovascular disorders.

There are currently very few potent LPAR5-selective antagonists. 'LPA5-antagonist 4' has been published by Sanofi-Aventis (Kozian et al, 2012, Bioorg Med Chem Lett., 22(16):5239-43) and is reported to be of low potency against LPAR5 (0.8 μ M).

- 5 It would be desirable to develop compounds that are potent against LPAR5 and/or LPAR1 for use in treating pain disorders and diseases such as atherosclerosis in which these receptors are believed to play a role.

Summary

In one aspect of the present disclosure, there is provided a compound of formula (I)



wherein

R^1 represents a C₅-C₁₀ aryl group substituted by at least one substituent selected from halogen, cyano, nitro, carboxyl, hydroxyl, C₃-C₆ cycloalkyl, C₁-C₆ alkoxy,

C₁-C₆ alkoxy carbonyl, C₁-C₆ alkoxy carbonylamino, C₁-C₆ haloalkoxy, -NR⁵R⁶,

- 15 C₃-C₆ cycloalkylamino, C₁-C₆ alkylthio, C₁-C₆ alkylcarbonyl, C₁-C₆ alkylcarbonyloxy, C₁-C₆ alkylcarbonylamino, sulphonamido, C₁-C₆ alkylsulphonyl,

C₁-C₆ alkylsulphonylamino, -C(O)NR⁷R⁸, C₁-C₆ alkyl which alkyl group may in turn be optionally substituted by at least one halogen, hydroxyl, carboxyl or C₁-C₆ alkoxy carbonyl group, and a saturated or unsaturated 5- to 6-membered ring which may comprise at least
 20 one ring heteroatom selected from nitrogen, oxygen and sulphur, the ring itself being optionally substituted by at least one substituent selected from halogen, hydroxyl, oxo, carboxyl, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy carbonyl and C₁-C₆ hydroxyalkyl;

X represents an oxygen atom or a group -CH₂-, -OCH₂-, -CH₂O-, -CH₂CH₂-, -CH₂NR¹⁷-, -NR¹⁷CH₂-, -CHF- or -CF₂;

m is 0, 1 or 2;

each R² independently represents a halogen atom or a hydroxyl, C₁-C₆ alkyl,

5 difluoromethyl, trifluoromethyl, C₁-C₆ alkoxy, difluoromethoxy, trifluoromethoxy or NR¹⁵R¹⁶ group;

Y represents CR⁹R¹⁰ in which R⁹ and R¹⁰ each independently represent a hydrogen atom or a methyl group;

R³ represents C₁-C₈ alkyl, C₂-C₈ alkenyl, or a saturated or unsaturated 3- to 10-
 10 membered ring system which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur, each of the preceding groups being optionally substituted by at least one substituent selected from halogen, cyano, hydroxyl, C₃-C₆ cycloalkyl, C₁-C₆ alkoxy, C₅-C₆ aryloxy, C₅-C₆arylC₁-C₆alkyloxy, C₁-C₆ alkoxycarbonyl, C₁-C₆ alkoxycarbonylamino, C₁-C₆ alkoxycarbonyloxy, C₁-C₆ haloalkyl,
 15 C₁-C₆ haloalkoxy, -NR¹³R¹⁴, C₃-C₆ cycloalkylamino, C₁-C₆ alkylthio, C₁-C₆ alkylcarbonyl, C₁-C₆ alkylcarbonyloxy, C₁-C₆ alkylcarbonylamino, sulphonamido, C₁-C₆ alkylsulphonyl, C₁-C₆ alkylsulphonylamino, C₁-C₆ alkyl, and a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur, the ring system itself being
 20 optionally substituted by least one substituent selected from halogen, hydroxyl, oxo, carboxyl, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxycarbonyl and C₁-C₆ hydroxyalkyl;

Z represents CR¹¹R¹² in which R¹¹ and R¹² each independently represent a hydrogen atom or a methyl group but cannot both simultaneously represent a methyl group;

25 n is 0, 1 or 2;

each R⁴ independently represents a halogen atom or a hydroxyl, C₁-C₆ alkyl, difluoromethyl, trifluoromethyl, C₁-C₆ alkoxy, difluoromethoxy, trifluoromethoxy or NR¹⁸R¹⁹ group;

R⁵ and R⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or
 5 R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R⁷ and R⁸ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or
 R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

10 R¹³ and R¹⁴ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or
 R¹³ and R¹⁴ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R¹⁵ and R¹⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or
 R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form a 4- to 7-
 15 membered saturated heterocycle;

R¹⁷ represents a hydrogen atom or a C₁-C₆ alkyl group;

R¹⁸ and R¹⁹ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or
 R¹⁸ and R¹⁹ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle; and

20 rings A and B each independently represent a saturated or unsaturated 5- to 10-membered ring system which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur;
 or a pharmaceutically acceptable salt thereof.

25 In a further aspect of the present disclosure, there is provided a compound selected from the group consisting of:

4-((N-Ethyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((N-(2,2-Difluoroethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,
 4-((4-(2-Fluorophenoxy)-*N*-isobutylbenzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclobutylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
 5 4-((4-(2-Fluorophenoxy)-*N*-((3-methoxycyclobutyl)methyl)benzamido)methyl)benzoic
 acid,
 4-((4-(2-Fluorophenoxy)-*N*-isopentylbenzamido)methyl)benzoic acid,
 4-((*N*-(2-Cyclopropylethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-(2-Cyclobutylethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
 10 4-((*N*-Benzyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-(2-Fluorobenzyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
 4-((4-(2-Fluorophenoxy)-*N*-(3-methoxybenzyl)benzamido)methyl)benzoic acid,
 4-((4-(2-Fluorophenoxy)-*N*-(4-methoxybenzyl)benzamido)methyl)benzoic acid,
 4-((4-(2-Fluorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,
 15 4-((4-(2-Fluorophenoxy)-*N*-(3-(2-fluorophenyl)propyl)benzamido)methyl)benzoic
 acid,
 4-((4-(2-Fluorophenoxy)-*N*-(3-(3-fluorophenyl)propyl)benzamido)methyl)benzoic
 acid,
 4-((4-(2-Fluorophenoxy)-*N*-((*trans*-2-phenylcyclopropyl)methyl)benzamido)-
 20 methyl)benzoic acid,
 (*S*)-4-((4-(2-Fluorophenoxy)-*N*-(2-hydroxy-3-phenylpropyl)benzamido)methyl)benzoic
 acid,
 (*R*)-4-((4-(2-Fluorophenoxy)-*N*-(2-hydroxy-3-phenylpropyl)benzamido)-
 methyl)benzoic acid,
 25 4-((4-(2-Fluorophenoxy)-*N*-(4-phenylbutyl)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclobutylmethyl)-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,
 4-((*N*-Benzyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,
 4-((*N*-Phenethyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,
 30 4-((*N*-Phenethyl-4-(2-(trifluoromethyl)phenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-Benzyl-4-(2-cyanophenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-Ethyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-(2,2-Difluoroethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,

- 4-((*N*-Isobutyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,
 4-((4-(2-Methoxyphenoxy)-*N*-(3-phenylpropyl)benzamido)methyl)benzoic acid,
 4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)benzamido)-methyl)benzoic
 5 acid,
 4-((*N*-(Cyclopropylmethyl)-4-(2-ethoxyphenoxy)benzamido)methyl)benzoic acid,
 4-((4-(2-Chlorophenoxy)-*N*-(cyclopropylmethyl)benzamido)methyl)benzoic acid,
 4-((4-(2-Chlorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid,
 10 4-((*N*-(2-Cyclopropylethyl)-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-Benzyl-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(2-fluoro-6-methylphenoxy)benzamido)methyl)benzoic
 acid,
 4-((*N*-(Cyclobutylmethyl)-4-(2-fluoro-6-methylphenoxy)benzamido)-methyl)benzoic
 15 acid,
 4-((4-(2-Chloro-6-fluorophenoxy)-*N*-(cyclopropylmethyl)benzamido)methyl)benzoic
 acid,
 4-((4-(2-Chloro-6-fluorophenoxy)-*N*-(2-cyclopropylethyl)benzamido)methyl)benzoic
 acid,
 20 4-((*N*-Benzyl-4-(2-chloro-6-fluorophenoxy)benzamido)methyl)benzoic acid,
 4-((4-(2,6-Dimethylphenoxy)-*N*-isopentylbenzamido)methyl)benzoic acid,
 4-((*N*-Benzyl-4-(2,6-dimethylphenoxy)benzamido)methyl)benzoic acid,
 4-((4-(2,6-Dimethylphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,
 4-((4-(3-Methoxyphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,
 25 4-((4-(4-Methoxyphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,
 4-((4-(3-Chlorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclobutylmethyl)-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic acid,
 4-((*N*-Benzyl-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic acid,
 30 4-((*N*-(Cyclopropylmethyl)-2-fluoro-4-(2-fluorophenoxy)-benzamido)methyl)benzoic
 acid,
 4-((*N*-Benzyl-4-(2-fluorophenoxy)cyclohexanecarboxamido)methyl)benzoic acid,

trans-4-((*N*-Benzyl-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid,

cis-4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid,

5 *trans*-4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid

2-Fluoro-4-((4-(2-fluorophenoxy)-*N*-(3-methoxybenzyl)benzamido)-methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-propylbenzamido)methyl)benzoic acid,

10 4-((4-(2-Methoxyphenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

4-((4-(2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,

4-((*N*-(2,2-difluoropropyl)-4-(2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

4-((4-(2-methoxyphenoxy)-*N*-(3,3,3-trifluoropropyl)-benzamido)methyl)benzoic acid,

4-((*N*-butyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,

15 4-((*N*-(cyclopropylmethyl)-2-fluoro-4-(2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

4-((*N*-(cyclopropylmethyl)-4-(2-fluoro-6-methoxyphenoxy)-benzamido)methyl)benzoic acid,

20 4-((*N*-(cyclopropylmethyl)-4-(4-fluoro-2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

4-((*N*-(cyclopropylmethyl)-5-(2-methoxyphenoxy)-picolinamido)methyl)benzoic acid,

4-((4-(*o*-Tolyloxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-(3,3,3-trifluoropropyl)benzamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

25 4-((*N*-Ethyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,

4-((*N*-(2,2-Difluoropropyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((*N*-Butyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-ethylbenzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-5-(2-fluorophenoxy)picolinamido)methyl)benzoic acid,

30 4-((4-(2-Chlorophenoxy)-*N*-(2,2-difluoroethyl)benzamido)methyl)benzoic acid,


4-((*N*-(Cyclopropylmethyl)-4-(5-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoic acid,

4-((4-(4-Fluoro-2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,

4-((4-(5-Fluoro-2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(3-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoic
 acid,
 4-((*N*-(Cyclopropylmethyl)-5-(2-fluorophenoxy)pyrimidine-2-
 5 carboxamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-5-(2-methoxyphenoxy)pyrimidine-2-
 carboxamido)methyl)benzoic acid,
 4-((5-(2-Methoxyphenoxy)-*N*-propylpyrimidine-2-carboxamido)methyl)benzoic acid,
 and pharmaceutically acceptable salts thereof.

10

In another aspect of the present disclosure, there is provided a process for the preparation
 of a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined
 herein which comprises hydrolysing a compound of formula (II)



15

wherein L^1 represents a leaving group and R^1 , X , m , R^2 , Y , R^3 , Z , n , R^4 , A and B are as
 defined in formula (I), in the presence of a suitable base;

and optionally thereafter carrying out one or more of the following procedures:

20

- converting a compound of formula (I) into another compound of formula (I)
- removing any protecting groups
- forming a pharmaceutically acceptable salt.

In yet another aspect of the present disclosure, there is provided a pharmaceutical
 composition comprising a compound of formula (I) or a pharmaceutically acceptable salt

thereof as defined herein, in association with a pharmaceutically acceptable adjuvant, diluent or carrier, and optionally one or more other therapeutic agents.


5 In a further aspect of the present disclosure, there is provided a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined herein for use in therapy.

10 In another aspect of the present disclosure, there is provided a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined herein for use in treating pain disorders, atherosclerosis or fibrosis.

15 In yet another aspect of the present disclosure, there is provided a use of a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined herein, in the manufacture of a medicament for treating pain disorders, atherosclerosis or fibrosis.

Detailed Description

In accordance with the present invention, there is therefore provided a compound of formula (I)



wherein

R^1 represents a C_5 - C_{10} aryl group substituted by at least one substituent independently

selected from halogen, cyano, nitro, carboxyl, hydroxyl, C_3 - C_6 cycloalkyl,

C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 alkoxycarbonylamino,

C_1 - C_6 haloalkoxy, $-NR^5R^6$, C_3 - C_6 cycloalkylamino, C_1 - C_6 alkylthio,

C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarbonyloxy, C_1 - C_6 alkylcarbonylamino,

sulphonamido ($-\text{SO}_2\text{NH}_2$), C_1 - C_6 alkylsulphonyl,

C_1 - C_6 alkylsulphonylamino, $-\text{C}(\text{O})\text{NR}^7\text{R}^8$, C_1 - C_6 alkyl which alkyl group may in turn be

optionally substituted by at least one halogen, hydroxyl, carboxyl or C_1 - C_6 alkoxycarbonyl

group, and a saturated or unsaturated 5- to 6-membered ring which may comprise at least one ring heteroatom independently selected from nitrogen, oxygen and sulphur, the ring itself being optionally substituted by at least one substituent independently selected from halogen, hydroxyl, oxo (=O), carboxyl, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy carbonyl and C₁-C₆ hydroxyalkyl;

X represents an oxygen atom or a group -CH₂-, -OCH₂-, -CH₂O-, -CH₂CH₂-, -CH₂NR¹⁷-, -NR¹⁷CH₂-, -CHF- or -CF₂-;

m is 0, 1 or 2;

each R² independently represents a halogen atom or a hydroxyl, C₁-C₆ alkyl, difluoromethyl, trifluoromethyl, C₁-C₆ alkoxy, difluoromethoxy, trifluoromethoxy or NR¹⁵R¹⁶ group;

Y represents CR⁹R¹⁰ in which R⁹ and R¹⁰ each independently represent a hydrogen atom or a methyl group;

R^3 represents C₁-C₈ alkyl, C₂-C₈ alkenyl, or a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom independently selected from nitrogen, oxygen and sulphur, each of the preceding groups being optionally substituted by at least one substituent independently selected from halogen, cyano, hydroxyl, C₃-C₆ cycloalkyl, C₁-C₆ alkoxy, C₅-C₆ aryloxy, C₅-C₆arylC₁-C₆alkyloxy, C₁-C₆ alkoxycarbonyl, C₁-C₆ alkoxycarbonylamino, C₁-C₆ alkoxycarbonyloxy, C₁-C₆ haloalkyl, C₁-C₆ haloalkoxy, -NR¹³R¹⁴, C₃-C₆ cycloalkylamino, C₁-C₆ alkylthio, C₁-C₆ alkylcarbonyl, C₁-C₆ alkylcarbonyloxy, C₁-C₆ alkylcarbonylamino, sulphonamido (-SO₂NH₂), C₁-C₆ alkylsulphonyl, C₁-C₆ alkylsulphonylamino, C₁-C₆ alkyl, and a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom independently selected from nitrogen, oxygen and sulphur, the ring system itself being optionally substituted by least one substituent independently selected from halogen, hydroxyl, oxo

(=O), carboxyl, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxycarbonyl and C₁-C₆ hydroxyalkyl;

Z represents CR¹¹R¹² in which R¹¹ and R¹² each independently represent a hydrogen atom or a methyl group but cannot both simultaneously represent a methyl group;

n is 0, 1 or 2;

each R⁴ independently represents a halogen atom or a hydroxyl, C₁-C₆ alkyl, difluoromethyl, trifluoromethyl, C₁-C₆ alkoxy, difluoromethoxy, trifluoromethoxy or NR¹⁸R¹⁹ group;

R⁵ and R⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or

R⁵ and R⁶ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R⁷ and R⁸ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or

R⁷ and R⁸ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R¹³ and R¹⁴ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or

R¹³ and R¹⁴ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R¹⁵ and R¹⁶ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or

R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R¹⁷ represents a hydrogen atom or a C₁-C₆ alkyl group;

R¹⁸ and R¹⁹ each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or

R¹⁸ and R¹⁹ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle; and

rings A and B each independently represent a saturated or unsaturated 5- to 10-membered ring system which may comprise at least one ring heteroatom independently selected from nitrogen, oxygen and sulphur;
or a pharmaceutically acceptable salt thereof.

Without being bound to any particular theory, the compounds of the invention have been found to possess dual LPAR5/LPAR1 antagonist properties. Such dual antagonist properties would be expected to be particularly beneficial in the treatment of conditions such as fibrosis.

In the context of the present specification, unless otherwise stated, an “alkyl” or “alkenyl” substituent group or an “alkyl” or “alkenyl” moiety in a substituent group may be linear or branched. Examples of C₁-C₈ alkyl groups/moieties include methyl, ethyl, propyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, n-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, n-heptyl and n-octyl. Examples of C₂-C₈ alkenyl groups/moieties include ethenyl, propenyl, 1-butenyl, 2-butenyl, 1-pentenyl, 1-hexenyl, 1,3-butadienyl, 1,3-pentadienyl, 1,4-pentadienyl and 1-hexadienyl.

A “C₁-C₆ haloalkyl” or “C₁-C₆ haloalkoxy” substituent group/moiety will comprise at least one halogen atom, e.g. one, two, three, four or five halogen atoms, examples of which include trifluoromethyl, trifluoromethoxy or pentafluoroethyl.

A “C₁-C₆ hydroxyalkyl” substituent group/moiety will comprise at least one hydroxyl group, e.g. one, two, three or four hydroxyl groups, examples of which include – CH₂OH, –CH₂CH₂OH, –CH₂CH₂CH₂OH, –CH(OH)CH₂OH, –CH(CH₃)OH and –CH(CH₂OH)₂.

A "cycloalkyl" substituent group/moiety is a saturated hydrocarbyl ring containing, for example, from 3 to 8 carbon atoms, examples of which include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

5

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic aromatic hydrocarbon system containing up to 10 carbon atoms, e.g. from 5 to 10 carbon atoms, such as phenyl or naphthyl.

10 In the definition of R^1 , the saturated or unsaturated 5- to 6-membered ring which may comprise at least one ring heteroatom independently selected from nitrogen, oxygen and sulphur may have alicyclic or aromatic properties. An unsaturated ring will be partially or fully unsaturated. Similar comments apply to the saturated or unsaturated 3- to 10-membered ring systems defined in R^3 . In either case, it should be understood that the
15 invention does not encompass any unstable ring structures or any O-O, O-S or S-S bonds and that a substituent, if present, may be attached to any suitable ring atom.

When any of R^5 and R^6 , R^7 and R^8 , R^{13} and R^{14} , R^{15} and R^{16} or R^{18} and R^{19} represents a 4- to 7-membered saturated heterocycle, it should be understood that the heterocycle may
20 contain one or more further heteroatoms (e.g. nitrogen, oxygen or sulphur atoms) in addition to the nitrogen atom to which R^5 and R^6 , R^7 and R^8 , R^{13} and R^{14} , R^{15} and R^{16} or R^{18} and R^{19} are attached.

When any chemical moiety or group in formula (I) is described as being optionally
25 substituted, it will be appreciated that the moiety or group may be either unsubstituted or substituted by one or more of the specified substituents. It will be appreciated that the number and nature of substituents will be selected so as to avoid sterically undesirable combinations.

R¹ represents a C₅-C₁₀ or C₅-C₆ aryl group substituted by at least one substituent, e.g.


one, two, three or four substituents, independently selected from

- (i) halogen (e.g. fluorine, chlorine, bromine or iodine),
- (ii) cyano,
- 5 (iii) nitro,
- (iv) carboxyl,
- (v) hydroxyl,
- (vi) C₃-C₆ or C₅-C₆ cycloalkyl,
- (vii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy,
- 10 (viii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy carbonyl,
- (ix) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy carbonylamino,
- (x) C₁-C₆, or C₁-C₄, or C₁-C₂ haloalkoxy,
- (xi) -NR⁵R⁶,
- (xii) C₃-C₆ or C₅-C₆ cycloalkylamino,
- 15 (xiii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylthio,
- (xiv) C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl carbonyl,
- (xv) C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl carbonyloxy,
- (xvi) C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl carbonylamino,
- (xvii) sulphonamido,
- 20 (xviii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylsulphonyl,
- (xix) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylsulphonylamino,
- (xx) -C(O)NR⁷R⁸,
- (xxi) C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl which alkyl group may in turn be optionally substituted by at least one substituent, e.g. one, two, three or four substituents,
- 25 independently selected from halogen (e.g. fluorine, chlorine, bromine or

iodine), hydroxyl, carboxyl and C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonyl,
and

- (xxii) a saturated or unsaturated 5- to 6-membered ring which may comprise at least one ring heteroatom, e.g. one, two, three or four ring heteroatoms, independently selected from nitrogen, oxygen and sulphur (preferably nitrogen or oxygen), the ring itself being optionally substituted by at least one substituent, e.g. one, two, three or four substituents, independently selected from halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxyl, oxo, carboxyl, cyano, C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl, C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonyl and C₁-C₆, or C₁-C₄, or C₁-C₂ hydroxyalkyl. Examples of saturated or unsaturated 5- to 6-membered rings include hydrocarbyl rings such as phenyl and heterocyclic rings such as pyrrolidinyl, piperidinyl, morpholinyl, tetrahydropyranyl, oxadiazolyl, pyrrolyl, imidazolyl, pyridinyl, pyrazinyl, pyridazinyl, pyrimidinyl, thienyl and furanyl.

The at least one substituent in R¹ is preferably attached in an ortho position relative to the point of attachment of R¹ to X. For example, if R¹ is phenyl, then the preferred carbon atoms to which a substituent is attached are shown by the asterisks in the structure below:



In an embodiment of the invention, R¹ represents a C₅-C₆ aryl (preferably phenyl) group substituted by at least one substituent, e.g. one, two, three or four substituents, independently selected from

- (i) halogen (e.g. fluorine, chlorine, bromine or iodine),
- (ii) cyano,
- (iii) nitro,

- (iv) carboxyl,
- (v) hydroxyl,
- (vi) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy,
- (vii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonyl,
- 5 (viii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonylamino,
- (ix) C₁-C₆, or C₁-C₄, or C₁-C₂ haloalkoxy,
- (x) -NR⁵R⁶,
- (xi) C₃-C₆ or C₅-C₆ cycloalkylamino,
- (xii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylthio,
- 10 (xiii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylcarbonyl,
- (xiv) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylcarbonyloxy,
- (xv) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylcarbonylamino,
- (xvi) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylsulphonyl,
- (xvii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylsulphonylamino,
- 15 (xviii) -C(O)NR⁷R⁸,
- (xix) C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl which alkyl group may in turn be optionally substituted by at least one substituent, e.g. one, two, three or four substituents, independently selected from halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxyl, carboxyl and C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonyl, and
- 20 (xx) a saturated or unsaturated 5- to 6-membered ring which may comprise at least one ring heteroatom, e.g. one, two, three or four ring heteroatoms, independently selected from nitrogen, oxygen and sulphur (preferably nitrogen or oxygen), the ring itself being optionally substituted by at least one substituent, e.g. one, two, three or four substituents, independently selected from halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxyl, oxo, carboxyl, cyano, C₁-C₆, or
- 25

C₁-C₄, or C₁-C₂ alkyl, C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy carbonyl and C₁-C₆,
or C₁-C₄, or C₁-C₂ hydroxyalkyl.

In another embodiment of the invention, R¹ represents a C₅-C₆ aryl group substituted by

one or two substituents independently selected from

- (i) halogen (e.g. fluorine, chlorine, bromine or iodine),
- (ii) cyano,
- (iii) nitro,
- (iv) carboxyl,
- (v) hydroxyl,
- (vi) C₁-C₂ alkoxy,
- (vii) C₁-C₂ alkoxy carbonyl,
- (viii) C₁-C₂ alkoxy carbonylamino,
- (ix) C₁-C₂ haloalkoxy,
- (x) -NR⁵R⁶,
- (xi) C₅-C₆ cycloalkylamino,
- (xii) C₁-C₂ alkylthio,
- (xiii) C₁-C₂ alkylcarbonyl,
- (xiv) C₁-C₂ alkylcarbonyloxy,
- (xv) C₁-C₂ alkylcarbonylamino,
- (xvi) C₁-C₂ alkylsulphonyl,
- (xvii) C₁-C₂ alkylsulphonylamino,
- (xviii) -C(O)NR⁷R⁸,

- (xix) C₁-C₂ alkyl which alkyl group may in turn be optionally substituted by at least one halogen atom, preferably from 1 to 3 halogen atoms (particularly fluorine atoms), and
- (xx) a saturated or unsaturated 5- to 6-membered ring which may comprise at least one ring heteroatom, e.g. one, two, three or four ring heteroatoms, independently selected from nitrogen, oxygen and sulphur (preferably nitrogen or oxygen), the ring itself being optionally substituted by at least one substituent, e.g. one, two, three or four substituents, independently selected from halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxyl, oxo, carboxyl, cyano, C₁-C₂ alkyl, C₁-C₂ alkoxy carbonyl and C₁-C₂ hydroxyalkyl.

In still another embodiment of the invention, R¹ represents a phenyl group substituted by one or two substituents independently selected from

- (i) halogen (e.g. fluorine or chlorine),
- (ii) cyano,
- (iii) C₁-C₂ alkoxy (particularly methoxy), and
- (iv) C₁-C₂ alkyl which alkyl group may in turn be optionally substituted by at least one halogen atom, preferably from 1 to 3 halogen atoms (particularly fluorine atoms).

In yet another embodiment, R¹ represents fluorophenyl (in particular 2-fluorophenyl), chlorophenyl (in particular 2-chlorophenyl or 3-chlorophenyl), difluorophenyl (in particular 2,6-difluorophenyl), chlorofluorophenyl (in particular 2-chloro-6-fluorophenyl), fluoromethylphenyl (in particular 2-fluoro-6-methylphenyl), methylphenyl (in particular 2-methylphenyl), dimethylphenyl (in particular 2,6-dimethylphenyl), cyanophenyl (in particular 2-cyanophenyl), trifluoromethylphenyl (in particular 2-trifluoromethylphenyl), methoxyphenyl (e.g. 2-methoxyphenyl, 3-methoxyphenyl or 4-methoxyphenyl), ethoxyphenyl (in particular 2-ethoxyphenyl), 2-fluoro-6-methoxyphenyl, 3-fluoro-2-methoxyphenyl, 4-fluoro-2-methoxyphenyl or 5-fluoro-2-methoxyphenyl.

In an embodiment of the invention, X represents an oxygen atom or a group -CH₂- , -OCH₂- or -CH₂O-.

5 In another embodiment, X represents an oxygen atom or -CH₂O- (where it will be understood that the oxygen atom is attached to the ring A in formula (I)).

In yet another embodiment, X represents an oxygen atom.

10 In a further embodiment, X represents -CH₂O-.

In a still further embodiment, X represents -CH₂NR¹⁷-, in particular -CH₂NH-.

When m is 1 or 2, each R² independently represents a halogen atom (e.g. fluorine, chlorine, bromine or iodine) or a hydroxyl, C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl, 15 difluoromethyl, trifluoromethyl, C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy, difluoromethoxy, trifluoromethoxy or NR¹⁵R¹⁶ group. Preferably, R² represents a halogen atom (particularly fluorine) or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl (particularly methyl) group.

20 In one embodiment of the invention, m is 1 and R² represents a halogen atom (particularly fluorine).

In a further embodiment, m is 2 and each R² independently represents a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl (particularly methyl) group.

25

In a still further embodiment, m is 0 and the ring A has no R² substituents.

In one aspect, Y represents CH₂ or CH(CH₃), preferably CH₂.

R³ represents C₁-C₈, or C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl, C₂-C₈, or C₂-C₆, or

C₂-C₄ alkenyl, or a saturated or unsaturated 3- to 10-membered (e.g. 3-, 4-, 5- or 6- to 7-, 8-, 9- or 10-membered) ring system which may comprise at least one ring heteroatom (e.g. one, two, three or four ring heteroatoms) independently selected from nitrogen, oxygen and sulphur, each of the preceding groups being optionally substituted by at least one substituent (e.g. one, two, three or four substituents) independently selected from

(i) halogen (e.g. fluorine, chlorine, bromine or iodine),

(ii) cyano,

(iii) hydroxyl,

(iv) C₃-C₆ or C₅-C₆ cycloalkyl,

(v) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy,

(vi) C₅-C₆ aryloxy,

(vii) C₅-C₆arylC₁-C₆, or C₁-C₄, or C₁-C₂ alkyloxy,

(viii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonyl,

(ix) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonylamino,

(x) C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonyloxy,

(xi) C₁-C₆, or C₁-C₄, or C₁-C₂ haloalkyl,

(xii) C₁-C₆, or C₁-C₄, or C₁-C₂ haloalkoxy,

(xiii) -NR¹³R¹⁴,

(xiv) C₃-C₆ or C₅-C₆ cycloalkylamino,

(xv) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylthio,

(xvi) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylcarbonyl,

- (xvii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylcarbonyloxy,
- (xviii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylcarbonylamino,
- (xix) sulphonamido,
- (xx) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylsulphonyl,
- 5 (xxi) C₁-C₆, or C₁-C₄, or C₁-C₂ alkylsulphonylamino,
- (xxii) C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl, and
- (xxiii) a saturated or unsaturated 3- to 10-membered (e.g. 3-, 4-, 5- or 6- to 7-, 8-, 9- or 10-membered) ring system which may comprise at least one ring heteroatom (e.g. one, two, three or four ring heteroatoms) independently selected from
 10 nitrogen, oxygen and sulphur, the ring system itself being optionally substituted by least one substituent (e.g. one, two, three or four substituents) independently selected from halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxyl, oxo, carboxyl, cyano, C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl, C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy, C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxycarbonyl and C₁-C₆, or
 15 C₁-C₄, or C₁-C₂ hydroxyalkyl.

Examples of saturated or unsaturated 3- to 10-membered ring systems that may be used, which may be monocyclic or polycyclic (e.g. bicyclic) in which the two or more rings may be fused, bridged or spiro include one or more (in any combination) of cyclopropyl,
 20 cyclobutyl, cyclopentyl, cyclohexyl, bicyclo[2.2.1]heptyl, cyclopentenyl, cyclohexenyl, phenyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, oxadiazolyl (e.g. 1,2,4-oxadiazolyl), tetrahydrofuranyl, diazabicyclo[2.2.1]hept-2-yl, naphthyl, benzofuranyl, benzothienyl, benzodioxolyl, benzoxazolyl, quinolinyl, oxazolyl, thiadiazolyl (e.g. 1,2,3-thiadiazolyl), 2,3-dihydrobenzofuranyl,
 25 tetrahydropyranyl, pyrazolyl, imidazo[1,2-a]pyridinyl, pyrazinyl, thiazolidinyl, indanyl, thienyl, isoxazolyl, pyridazinyl, pyrrolyl, furanyl, thiazolyl, isothiazolyl, indolyl, isoindolyl, imidazolyl, pyrimidinyl, benzimidazolyl, triazolyl, tetrazolyl and pyridinyl.

Preferred ring systems include phenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and thienyl.

In one aspect of the invention, R^3 represents C_1 - C_4 , or C_1 - C_3 , or C_1 - C_2 alkyl,

C_2 - C_4 alkenyl, or a saturated or unsaturated 3-, 4-, 5- or 6-membered ring system which may comprise at least one ring heteroatom (e.g. one, two, three or four ring heteroatoms) independently selected from nitrogen, oxygen and sulphur, each of the preceding groups being optionally substituted by at least one substituent (e.g. one, two, three or four substituents) independently selected from

(i) halogen (e.g. fluorine, chlorine, bromine or iodine),

(ii) cyano,

(iii) hydroxyl,

(iv) C_3 - C_6 or C_5 - C_6 cycloalkyl,

(v) C_1 - C_2 alkoxy,

(vi) C_5 - C_6 aryloxy,

(vii) C_5 - C_6 aryl C_1 - C_2 alkyloxy,

(viii) C_1 - C_2 alkoxycarbonyl,

(ix) C_1 - C_2 alkoxycarbonylamino,

(x) C_1 - C_2 alkoxycarbonyloxy,

(xi) C_1 - C_2 haloalkyl,

(xii) C_1 - C_2 haloalkoxy,

(xiii) $-NR^{13}R^{14}$,

(xiv) C_3 - C_6 or C_5 - C_6 cycloalkylamino,

(xv) C_1 - C_2 alkylthio,

(xvi) C_1 - C_2 alkylcarbonyl,

(xvii) C_1 - C_2 alkylcarbonyloxy,

- (xviii) C₁-C₂ alkylcarbonylamino,
- (xix) sulphonamido,
- (xx) C₁-C₂ alkylsulphonyl,
- (xxi) C₁-C₂ alkylsulphonylamino,
- 5 (xxii) C₁-C₂ alkyl, and
- (xxiii) a saturated or unsaturated 3-, 4-, 5- or 6-membered ring system which may comprise at least one ring heteroatom (e.g. one, two, three or four ring heteroatoms) independently selected from nitrogen, oxygen and sulphur, the ring system itself being optionally substituted by least one substituent (e.g. one, two, three or four substituents) independently selected from halogen (e.g. 10 fluorine, chlorine, bromine or iodine), hydroxyl, oxo, carboxyl, cyano, C₁-C₂ alkyl, C₁-C₂ alkoxy, C₁-C₂ alkoxycarbonyl and C₁-C₂ hydroxyalkyl.

In a further aspect of the invention, R³ represents C₁-C₄, or C₁-C₃, or C₁-C₂ alkyl,

15 C₂-C₄ alkenyl, or a saturated or unsaturated 3-, 4-, 5- or 6-membered hydrocarbyl ring system (e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or phenyl), each of the preceding groups being optionally substituted by at least one substituent (e.g. one, two, three or four substituents) independently selected from

- (i) halogen (e.g. fluorine or chlorine),
- 20 (ii) hydroxyl,
- (iii) C₃-C₆ cycloalkyl,
- (iv) C₁-C₂ alkoxy,
- (v) C₅-C₆ aryloxy,
- (vi) C₅-C₆arylC₁-C₂ alkyloxy,
- 25 (vii) C₁-C₂ haloalkyl,
- (viii) C₁-C₂ alkyl, and

- (ix) a saturated or unsaturated 3-, 4-, 5- or 6-membered ring system which may comprise at least one ring heteroatom (e.g. one or two ring heteroatoms) independently selected from nitrogen, oxygen and sulphur (e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl and thienyl), the ring system itself being optionally substituted by least one substituent (e.g. one, two, three or four substituents) independently selected from halogen (e.g. fluorine or chlorine), C₁-C₂ alkyl and C₁-C₂ alkoxy.

In a still further aspect, R³ represents C₁-C₄, or C₁-C₃, or C₁-C₂ alkyl, or a saturated or unsaturated 3-, 4-, 5- or 6-membered hydrocarbyl ring system (e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or phenyl), each of the preceding groups being optionally substituted by at least one substituent (e.g. one, two, three or four substituents) independently selected from

- (i) fluorine
- (ii) chlorine,
- (iii) hydroxyl,
- (iv) C₃-C₆ cycloalkyl,
- (v) C₁-C₂ alkoxy,
- (vi) phenoxy,
- (vii) benzyloxy,
- (viii) trifluoromethyl,
- (ix) C₁-C₂ alkyl, and
- (x) a saturated or unsaturated 3-, 4-, 5- or 6-membered ring system which may comprise at least one ring heteroatom (e.g. one or two ring heteroatoms) independently selected from nitrogen, oxygen and sulphur (e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl and thienyl), the ring system itself being optionally substituted by least one substituent (e.g. one, two, three or four substituents) independently selected from halogen (e.g. fluorine or chlorine), C₁-C₂ alkyl and C₁-C₂ alkoxy.

In still another aspect, R^3 represents C_1 - C_4 , or C_1 - C_3 , or C_1 - C_2 alkyl, cyclopropyl, cyclobutyl or phenyl, each of the preceding groups being optionally substituted by at least one substituent (e.g. one, two, three or four substituents) independently selected from

- 5 (i) fluorine,
- (ii) hydroxyl,
- (iii) methoxy,
- (iv) cyclopropyl,
- (v) cyclobutyl, and
- 10 (vi) phenyl optionally substituted by least one halogen (e.g. fluorine or chlorine) atom.

In a particular embodiment of the invention, R^3 represents any one of the following moieties or is selected from a group containing any two or more of such moieties:

15

- (i) methyl,
- (ii) difluoromethyl,
- (iii) trifluoromethyl,
- (iv) ethyl,
- 20 (v) difluoroethyl (e.g. 1,1-difluoroethyl),
- (vi) trifluoroethyl (e.g. 2,2,2-trifluoroethyl),
- (vii) $-CH_2CH_2CH_3$,
- (viii) $-CH(CH_3)_2$,
- (ix) $-CH_2CH(CH_3)_2$,
- 25 (x) cyclopropyl,
- (xi) cyclopropylmethyl,

- (xii) cyclobutyl,
- (xiii) cyclobutylmethyl,
- (xiv) 3-methoxycyclobutyl,
- (xv) phenyl,
- 5 (xvi) 2-fluorophenyl,
- (xvii) 3-methoxyphenyl,
- (xviii) 4-methoxyphenyl,
- (xix) benzyl,
- (xx) phenylethyl,
- 10 (xxi) (2-fluorophenyl)ethyl,
- (xxii) (3-fluorophenyl)ethyl,
- (xxiii) phenylpropyl,
- (xxiv) phenylcyclopropyl,
- (xxv) -CH(OH)CH₂-phenyl.

15

Z represents CH₂ or CH(CH₃), preferably CH₂.

When n is 1 or 2, each R⁴ independently represents a halogen atom (e.g. fluorine, chlorine, bromine or iodine) or a hydroxyl, C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl, difluoromethyl, trifluoromethyl, C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy, difluoromethoxy, trifluoromethoxy or NR¹⁸R¹⁹ group.

20

In one embodiment, n is 1 and R^4 represents a halogen atom (particularly fluorine) or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl (particularly methyl) or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkoxy (particularly methoxy) group.

5 In another embodiment, n is 0 and the ring B has no R^4 substituents.

R^5 and R^6 each independently represent a hydrogen atom or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl group, or R^5 and R^6 together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle (e.g. pyrrolidinyl, morpholinyl,
10 piperazinyl or piperidinyl).

R^7 and R^8 each independently represent a hydrogen atom or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl group, or R^7 and R^8 together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle (e.g. pyrrolidinyl, morpholinyl,
15 piperazinyl or piperidinyl).

R^{13} and R^{14} each independently represent a hydrogen atom or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl group, or R^{13} and R^{14} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle (e.g. pyrrolidinyl, morpholinyl,
20 piperazinyl or piperidinyl).

R^{15} and R^{16} each independently represent a hydrogen atom or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl group, or R^{15} and R^{16} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle (e.g. pyrrolidinyl, morpholinyl,
25 piperazinyl or piperidinyl).


R¹⁷ represents a hydrogen atom or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl (particularly methyl or ethyl) group, especially a hydrogen atom.

R¹⁸ and R¹⁹ each independently represent a hydrogen atom or a C₁-C₆, or C₁-C₄, or C₁-C₂ alkyl group, or R¹⁸ and R¹⁹ together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle (e.g. pyrrolidinyl, morpholinyl, piperazinyl or piperidinyl).






Rings A and B each independently represent a saturated or unsaturated 5-, 6-, 7-, 8-, 9- or 10-membered ring system which may comprise at least one ring heteroatom (e.g. one, two, three or four heteroatoms) independently selected from nitrogen, oxygen and sulphur.

Examples of saturated or unsaturated 5- to 10-membered ring systems that may be used, which may be monocyclic or polycyclic (e.g. bicyclic) in which the two or more rings are fused include one or more (in any combination) of cyclopentyl, cyclohexyl, cyclopentenyl, cyclohexenyl, phenyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, oxadiazolyl (e.g. 1,2,4-oxadiazolyl), tetrahydrofuranyl, naphthyl, benzofuranyl, benzothienyl, benzodioxolyl, benzoxazolyl, quinolinyl, oxazolyl, thiadiazolyl (e.g. 1,2,3-thiadiazolyl), 2,3-dihydrobenzofuranyl, tetrahydropyranyl, pyrazolyl, imidazo[1,2-a]pyridinyl, pyrazinyl, thiazolidinyl, indanyl, thienyl, isoxazolyl, pyridazinyl, pyrrolyl, furanyl, thiazolyl, isothiazolyl, indolyl, isoindolyl, imidazolyl, pyrimidinyl, benzimidazolyl, triazolyl, tetrazolyl and pyridinyl. Preferred ring systems include 5- or 6-membered ring systems such as phenyl, pyridinyl, pyrimidinyl, thienyl and cyclohexyl.

In an embodiment of the invention, ring A is selected from phenyl, pyridinyl, pyrimidinyl, thienyl and cyclohexyl. Examples of ring A showing the points of attachment of R¹-X-, R² and C(O) to the ring are shown below:



In another embodiment of the invention, ring B is selected from phenyl, thienyl and
 5 cyclohexyl. Examples of ring B showing the points of attachment of Z, R^4 and C(O)OH to
 the ring are shown below:



In a further embodiment, rings A and B are both phenyl.

5

In a preferred embodiment of the invention:

R^1 represents a phenyl group substituted by one or two substituents independently selected from halogen, cyano, C_1 - C_2 alkoxy, and C_1 - C_2 alkyl which alkyl group may in turn be optionally substituted by at least one halogen atom;

10

X represents an oxygen atom or $-CH_2O-$;

m is 0 or 1;

if present, R^2 represents a halogen atom;

Y represents CH₂;

R³ represents C₁-C₄ alkyl, or a saturated or unsaturated 3-, 4-, 5- or 6-membered hydrocarbyl ring system, each of the preceding groups being optionally substituted by at least one substituent independently selected from fluorine, hydroxyl, C₁-C₂ alkoxy, and
5 a saturated or unsaturated 3-, 4-, 5- or 6-membered ring system which may comprise at least one ring heteroatom independently selected from nitrogen, oxygen and sulphur, the ring system itself being optionally substituted by least one substituent independently selected from halogen atoms;

Z represents CH₂;

10 n is 0 or 1;

if present, R⁴ represents a halogen atom; and

rings A and B each independently represent a saturated or unsaturated 5- or 6-membered ring system which may comprise at least one ring heteroatom independently selected from nitrogen, oxygen and sulphur.

15 Examples of compounds of the invention include:

4-((N-Ethyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
4-((N-(2,2-Difluoroethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
4-((4-(2-Fluorophenoxy)-N-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,
20 4-((4-(2-Fluorophenoxy)-N-isobutylbenzamido)methyl)benzoic acid,
4-((N-(Cyclopropylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
4-((N-(Cyclobutylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
4-((4-(2-Fluorophenoxy)-N-((3-methoxycyclobutyl)methyl)benzamido)-
methyl)benzoic acid,

25 4-((4-(2-Fluorophenoxy)-N-isopentylbenzamido)methyl)benzoic acid,
4-((N-(2-Cyclopropylethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
4-((N-(2-Cyclobutylethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
4-((N-Benzyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
4-((N-(2-Fluorobenzyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-(3-methoxybenzyl)benzamido)methyl)benzoic acid,
4-((4-(2-Fluorophenoxy)-*N*-(4-methoxybenzyl)benzamido)methyl)benzoic acid,
4-((4-(2-Fluorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,
4-((4-(2-Fluorophenoxy)-*N*-(3-(2-fluorophenyl)propyl)benzamido)methyl)benzoic
5 acid,
4-((4-(2-Fluorophenoxy)-*N*-(3-(3-fluorophenyl)propyl)benzamido)methyl)benzoic
acid,
4-((4-(2-Fluorophenoxy)-*N*-((*trans*-2-phenylcyclopropyl)methyl)benzamido)-
methyl)benzoic acid,
10 (*S*)-4-((4-(2-Fluorophenoxy)-*N*-(2-hydroxy-3-phenylpropyl)-
benzamido)methyl)benzoic acid,
(*R*)-4-((4-(2-Fluorophenoxy)-*N*-(2-hydroxy-3-phenylpropyl)benzamido)-
methyl)benzoic acid,
4-((4-(2-Fluorophenoxy)-*N*-(4-phenylbutyl)benzamido)methyl)benzoic acid,
15 4-((*N*-(Cyclopropylmethyl)-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,
4-((*N*-(Cyclobutylmethyl)-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,
4-((*N*-Benzyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,
4-((*N*-Phenethyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,
4-((*N*-Phenethyl-4-(2-(trifluoromethyl)phenoxy)benzamido)methyl)benzoic acid,
20 4-((*N*-Benzyl-4-(2-cyanophenoxy)benzamido)methyl)benzoic acid,
4-((*N*-Ethyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,
4-((*N*-(2,2-Difluoroethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,
4-((*N*-Isobutyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,
4-((*N*-(Cyclopropylmethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,
25 4-((4-(2-Methoxyphenoxy)-*N*-(3-phenylpropyl)benzamido)methyl)benzoic acid,
4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)benzamido)-
methyl)benzoic acid,
4-((*N*-(Cyclopropylmethyl)-4-(2-ethoxyphenoxy)benzamido)methyl)benzoic acid,
4-((4-(2-Chlorophenoxy)-*N*-(cyclopropylmethyl)benzamido)methyl)benzoic acid,
30 4-((4-(2-Chlorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid,

4-((*N*-(2-Cyclopropylethyl)-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid,

5 4-((*N*-Benzyl-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-(2-fluoro-6-methylphenoxy)benzamido)-methyl)benzoic acid,

4-((*N*-(Cyclobutylmethyl)-4-(2-fluoro-6-methylphenoxy)benzamido)-methyl)benzoic acid,

10 4-((4-(2-Chloro-6-fluorophenoxy)-*N*-(cyclopropylmethyl)benzamido)-methyl)benzoic acid,

4-((4-(2-Chloro-6-fluorophenoxy)-*N*-(2-cyclopropylethyl)benzamido)-methyl)benzoic acid,

4-((*N*-Benzyl-4-(2-chloro-6-fluorophenoxy)benzamido)methyl)benzoic acid,

15 4-((4-(2,6-Dimethylphenoxy)-*N*-isopentylbenzamido)methyl)benzoic acid,

4-((*N*-Benzyl-4-(2,6-dimethylphenoxy)benzamido)methyl)benzoic acid,

4-((4-(2,6-Dimethylphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((4-(3-Methoxyphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((4-(4-Methoxyphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

20 4-((4-(3-Chlorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic acid,

4-((*N*-(Cyclobutylmethyl)-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic acid,

4-((*N*-Benzyl-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic acid,

25 4-((*N*-(Cyclopropylmethyl)-2-fluoro-4-(2-fluorophenoxy)-benzamido)methyl)benzoic acid,

4-((*N*-Benzyl-4-(2-fluorophenoxy)cyclohexanecarboxamido)methyl)benzoic acid,

trans-4-((*N*-Benzyl-4-(2-methoxyphenoxy)cyclohexanecarboxamido)-methyl)benzoic acid,

30 *cis*-4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-

methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid,

trans-4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid

2-Fluoro-4-((4-(2-fluorophenoxy)-*N*-(3-methoxybenzyl)benzamido)methyl)benzoic acid,

5 4-((4-(2-Fluorophenoxy)-*N*-propylbenzamido)methyl)benzoic acid,

4-((4-(2-Methoxyphenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

4-((4-(2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,

4-((*N*-(2,2-difluoropropyl)-4-(2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

4-((4-(2-methoxyphenoxy)-*N*-(3,3,3-trifluoropropyl)-benzamido)methyl)benzoic

10 acid,

4-((*N*-butyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,

4-((*N*-(cyclopropylmethyl)-2-fluoro-4-(2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

4-((*N*-(cyclopropylmethyl)-4-(2-fluoro-6-methoxyphenoxy)-benzamido)methyl)benzoic acid,

15 4-((*N*-(cyclopropylmethyl)-4-(4-fluoro-2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

4-((*N*-(cyclopropylmethyl)-5-(2-methoxyphenoxy)-picolinamido)methyl)benzoic acid,

20 4-((4-(*o*-Tolyloxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-(3,3,3-trifluoropropyl)benzamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

4-((*N*-Ethyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,

4-((*N*-(2,2-Difluoropropyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

25 4-((*N*-Butyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-ethylbenzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-5-(2-fluorophenoxy)picolinamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-(2,2-difluoroethyl)benzamido)methyl)benzoic acid,


4-((*N*-(Cyclopropylmethyl)-4-(5-fluoro-2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

30 4-((4-(4-Fluoro-2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,

4-((4-(5-Fluoro-2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(3-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-5-(2-fluorophenoxy)pyrimidine-2-carboxamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-5-(2-methoxyphenoxy)pyrimidine-2-carboxamido)methyl)benzoic acid,
 4-((5-(2-Methoxyphenoxy)-*N*-propylpyrimidine-2-carboxamido)methyl)benzoic acid,
 and pharmaceutically acceptable salts of any one thereof.

It should be noted that each of the chemical compounds listed above represents a particular and independent aspect of the invention.

The present invention further provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined above which comprises hydrolysing a compound of formula (II)




wherein L^1 represents a leaving group (e.g. alkoxy such as methoxy) and R^1 , X, m, R^2 , Y, R^3 , Z, n, R^4 , A and B are as defined in formula (I), in the presence of a suitable base;

and optionally thereafter carrying out one or more of the following procedures:


- converting a compound of formula (I) into another compound of formula (I)
- removing any protecting groups
- forming a pharmaceutically acceptable salt.

5 The process is conveniently carried out using a base such as lithium hydroxide or sodium hydroxide in a solvent such as 1,4-dioxane or a solvent mixture such as water and tetrahydrofuran, at room temperature or at elevated temperature, e.g. in the range from 20°C to 100°C.

10 Compounds of formula (II) may be prepared by reacting a compound of formula (III)




wherein R^1 , X , m , R^2 and A are as defined in formula (II), with a compound of formula (IV), or a salt (e.g. hydrochloride salt) thereof,




15 wherein L^1 , Y , R^3 , Z , n , R^4 and B are as defined in formula (II). The reaction is conveniently carried out in the presence of an organic solvent such as tetrahydrofuran or *N,N*-dimethylformamide, a base such as diisopropylethylamine and an amide coupling agent such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC),
O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU)
 20 or *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HBTU).

Compounds of formula (III) may be prepared by oxidising the corresponding aldehyde of formula (V)




wherein R^1 , X, m, R^2 and A are as defined in formula (III). Suitable oxidising agents include sodium chlorite.

Compounds of formula (V) in which X represents an oxygen atom or $-\text{CH}_2\text{O}-$ may be prepared by reacting a compound of formula (VI), $R^1-(\text{CH}_2)_p\text{X}'\text{H}$, in which p is 0 or 1, X' is oxygen and R^1 is as defined in formula (V) with a compound of formula (VII)



in which L^2 represents a leaving group (e.g. a halogen atom such as fluorine) and m, R^2 and A are as defined in formula (V).


Compounds of formula (III) in which X represents $-\text{CH}_2-$ may be prepared by reacting a compound of formula (VIII)



in which L^3 represents a leaving group (e.g. a halogen atom such as bromine), P^1 represents a suitable protecting group (e.g. an alkoxy group such as methoxy) and m , R^2 and A are as defined in formula (III), with a compound of formula (IX), $R^1 - B(OH)_2$, in which R^1 is as defined in formula (III), in the presence of a suitable palladium catalyst, e.g.

[1,1'-bis(diphenylphosphino)ferrocene]palladium(II) chloride complex with dichloromethane (see, for example, WO 2012/080221 in the name of Katholieke Universiteit Leuven), followed by ester hydrolysis.

Compounds of formula (III) in which X represents $-CH_2CH_2-$ may be prepared by a Heck Reaction by reacting a compound of formula (X)




in which P^2 is as defined for P^1 above and m , R^2 and A are as defined in formula (III), with a compound of formula (XI), $R^1 - L^4$, in which L^4 represents a leaving group (e.g. a halogen atom such as iodine) and R^1 is as defined in formula (III), in the presence of a suitable palladium catalyst, followed by a hydrogenation reaction and then ester hydrolysis.

Compounds of formula (III) in which X represents $-OCH_2-$ may be prepared by reacting a compound of formula (VIII) as defined above with a compound of formula (XII), $R^1 - X'H$, wherein R^1 and X' are as defined in formula (VI), in the presence of a base such as potassium carbonate and in a solvent such as *N,N*-dimethylformamide (see, for example, WO 2012/007868 in the name of Pfizer Limited), followed by ester hydrolysis.

Compounds of formula (V) in which X represents $-\text{CH}_2\text{NR}^{17}$ may be prepared by reacting a compound of formula (VII) as defined above with a compound of formula (XIII), $\text{R}^1-\text{CH}_2\text{NR}^{17}\text{H}$, in which R^1 and R^{17} are as defined in formula (V), in the presence of a suitable copper catalyst such as copper (I) iodide (see, for example, Kwong *et al.*, Organic Letters, 2002, volume 4, pages 581 to 584).


Compounds of formula (III) in which X represents $-\text{NR}^{17}\text{CH}_2-$ may be prepared by reacting a compound of formula (VIII) as defined above with a compound of formula (XIV), $\text{R}^1-\text{NR}^{17}\text{H}$, wherein R^1 and R^{17} are as defined in formula (III), in the presence of a base such as calcium carbonate and a solvent such as isobutyramide (see, for example, WO 2007/120096 in the names of AstraZeneca AB and Astex Therapeutics Limited), followed by ester hydrolysis.

Compounds of formula (III) in which X represents $-\text{CHF}-$ may be prepared by reacting a compound of formula (XV)




in which P^3 is as defined for P^1 above and m , R^2 and A are as defined in formula (III), with a compound of formula (XI) as defined above in the presence of an organolithium such as phenyllithium, followed by a fluorination reaction using, for example, diethylaminosulfur trifluoride and then ester hydrolysis.

Compounds of formula (III) in which X represents $-\text{CF}_2-$ may be prepared by reacting a compound of formula (VIIa)



in which P^4 is as defined for P^1 above and m , R^2 and A are as defined in formula (VII),
 with a compound of formula (IX) as defined above, in the presence of carbon monoxide
 and a suitable palladium catalyst such as tris-(dibenzylideneacetone)dipalladium (0),
 5 followed by a fluorination reaction using, for example, diethylaminosulfur trifluoride and
 then ester hydrolysis.

Compounds of formula (IV) in which Z is CH_2 or $CH(CH_3)$ may be prepared by reacting a
 compound of formula (XVI), $R^3 - Y - NH_2$, in which R^3 and Y are as defined in formula
 10 (IV), with a compound of formula (XVII)



in which R^{20} represents hydrogen or methyl, and L^1 , n , R^4 and B are as defined in formula
 (IV), in the presence of a reducing agent such as sodium triacetoxyborohydride.

15 Compounds of formulae (V) to (XVII) are either commercially available, are well known
 in the literature or may be prepared using known techniques.

It will be appreciated by those skilled in the art that it may be necessary or desirable at any
 20 stage in the synthesis of the compounds of the invention to protect one or more sensitive
 groups, so as to prevent undesirable side reactions. In particular, it may be necessary or

desirable to protect certain functional groups such as hydroxyl, carboxyl or amino groups. Thus, the preparation of the compounds of formula (I) may involve, at an appropriate stage, the introduction and/or removal of one or more protecting groups.

- 5 The protection and deprotection of functional groups is described in 'Protective Groups in Organic Chemistry', edited by J.W.F. McOmie, Plenum Press (1973) and 'Protective Groups in Organic Synthesis', 3rd edition, T.W. Greene and P.G.M. Wuts, Wiley-Interscience (1999).
- 10 The compounds of formula (I) above may be converted to a pharmaceutically acceptable salt thereof, preferably an acid addition salt such as a hydrochloride, hydrobromide, benzenesulphonate (besylate), saccharin (e.g. monosaccharin), trifluoroacetate, sulphate, nitrate, phosphate, acetate, fumarate, maleate, tartrate, lactate, citrate, pyruvate, succinate, valerate, propanoate, butanoate, malonate, oxalate, 1-hydroxy-2-napthoate (xinafoate),
- 15 methanesulphonate or *p*-toluenesulphonate salt.

In one aspect of the invention, compounds of formula (I) may bear one or more radiolabels. Such radiolabels may be introduced by using radiolabel-containing reagents in the synthesis of the compounds of formula (I), or may be introduced by coupling the

20 compounds of formula (I) to chelating moieties capable of binding to a radioactive metal atom. Such radiolabeled versions of the compounds may be used, for example, in diagnostic imaging studies.

Unless stated otherwise, any atom specified herein may also be an isotope of said atom.

25 For example, the term "hydrogen" encompasses ¹H, ²H and ³H. Similarly carbon atoms are to be understood to include ¹²C, ¹³C and ¹⁴C, nitrogen atoms are to be understood to include ¹⁴N and ¹⁵N, and oxygen atoms are to be understood to include ¹⁶O, ¹⁷O and ¹⁸O.

In a further aspect of the invention, compounds of formula (I) may be isotopically labelled.

30 As used herein, an "isotopically labelled" compound is one in which the abundance of a

particular nuclide at a particular atomic position within the molecule is increased above the level at which it occurs in nature.

Compounds of formula (I) and their salts may be in the form of hydrates or solvates which
5 form an aspect of the present invention. Such solvates may be formed with common organic solvents, including but not limited to, alcoholic solvents e.g. methanol, ethanol or isopropanol.

Where compounds of formula (I) are capable of existing in stereoisomeric forms, it will be
10 understood that the invention encompasses the use of all geometric and optical isomers (including atropisomers) of the compounds of formula (I) and mixtures thereof including racemates. The use of tautomers and mixtures thereof also forms an aspect of the present invention. Enantiomerically pure forms are particularly desired.

15 Compounds of formula (I) and their salts may be amorphous or in a polymorphic form or a mixture of any of these, each of which forms an aspect of the present invention.

The compounds of formula (I) and their pharmaceutically acceptable salts have activity as pharmaceuticals, in particular as modulators (e.g. antagonists) of LPAR5 and, optionally
20 also, as modulators (e.g. antagonists) of LPAR1, and thus may be used in the treatment of: **fibrosis** of organs such as liver, kidney, skin, lung, heart and the like; **liver diseases** (acute hepatitis, chronic hepatitis, liver fibrosis, liver cirrhosis, portal hypertension, regenerative failure, non-alcoholic steato hepatitis (NASH), liver hypofunction, hepatic blood flow disorder, and the like); **cardiovascular disorders** (restenosis, atherosclerosis, heart
25 failure, cardiomyopathy, myocardial infarction, myocardial remodelling, vascular remodelling, hypertension, peripheral arterial occlusive disease, thrombosis, vascular permeability disorders, and the like); **cell proliferative disease** (cancer (solid tumour, solid tumour metastasis, vascular fibroma, myeloma, multiple myeloma, Kaposi's sarcoma, leukaemia, chronic lymphocytic leukaemia (CLL) and the like) and invasive metastasis of
30 cancer cell, and the like); **inflammatory disease** (rheumatoid arthritis, osteoarthritis, psoriasis, nephropathy, pneumonia and the like); **transplant rejection**; **gastrointestinal**

tract disease (irritable bowel syndrome (IBS), inflammatory bowel disease (IBD), abnormal pancreatic secretion, and the like); **renal disease**; **urinary tract –associated disease** (benign prostatic hyperplasia or symptoms associated with neuropathic bladder disease, spinal cord tumour, hernia of intervertebral disk, spinal canal stenosis, symptoms derived from diabetes, lower urinary tract disease (obstruction of lower urinary tract, and the like); **inflammatory disease of lower urinary tract** (dysuria; overactive bladder, interstitial cystitis, frequent urination and the like); **pancreas disease**; **abnormal angiogenesis associated disease** (arterial obstruction and the like); **systemic sclerosis** including localised scleroderma; **brain associated disease** (cerebral infarction, cerebral hemorrhage, and the like); **pain** (neuropathic pain, peripheral neuropathy, lightning pains, phantom pains, shooting pains, acute pain, inflammatory pain, neuralgia, tissue injury pain and the like); **ocular disease** (choroidal neovascularisation, age related macular degeneration (AMD), diabetic retinopathy, proliferative vitreoretinopathy (PVR), cicatricial pemphigoid, glaucoma filtration surgery scarring and the like); **pulmonary disease** (chronic obstructive pulmonary disease, asthma, acute respiratory distress syndrome); **psychiatric disorders**; **neurodegenerative diseases**; **cerebral and peripheral nerve disorders**; **scarring disorders**; and **wound healing**.

Thus, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof as hereinbefore defined for use in therapy, in particular for the treatment of conditions whose development or symptoms are linked to LPAR5 or LPAR1 activity.

The present invention also provides the use of a compound of formula (I) or a pharmaceutically acceptable salt thereof as hereinbefore defined for the preparation of a medicament for the treatment of conditions whose development or symptoms are linked to LPAR5 or LPAR1 activity.

In the context of the present specification, the terms “therapy” and “treatment” also include “prophylaxis” unless there are specific indications to the contrary. The terms “therapeutic”, “therapeutically” and “treating” should be construed accordingly.

Prophylaxis is expected to be particularly relevant to the treatment of persons who have suffered a previous episode of, or are otherwise considered to be at increased risk of, the disorder or condition in question. Persons at risk of developing a particular disorder or condition generally include those having a family history of the disorder or condition, or those who have been identified by genetic testing or screening to be particularly susceptible to developing the disorder or condition or those in the prodromal phase of a disorder.

In particular, the compounds of the invention (including pharmaceutically acceptable salts) may be used in the treatment of pain (such as neuropathic pain), atherosclerosis and fibrosis.

“Pain” means the more or less localised sensation of discomfort, distress or agony resulting from the stimulation of specialised nerve endings. There are many types of pain (as illustrated above) and the goal of treatment of pain is to reduce the degree or severity of pain perceived by a treatment subject.

The invention also provides a method of treating pain (e.g. neuropathic pain), atherosclerosis or fibrosis which comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof as hereinbefore defined.

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated. For example, the daily dosage of the compound of the invention, if inhaled, may be in the range from 0.05 micrograms per kilogram body weight ($\mu\text{g/kg}$) to 100 micrograms per kilogram body weight ($\mu\text{g/kg}$). Alternatively, if the compound is administered orally, then the daily dosage of the compound of the invention may be in the range from 0.01 micrograms per kilogram body weight ($\mu\text{g/kg}$) to 100 milligrams per kilogram body weight (mg/kg).

The compounds of formula (I) and pharmaceutically acceptable salts thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (I) compound/salt (active ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

Therefore the present invention further provides a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof as hereinbefore defined, in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

The invention still further provides a process for the preparation of a pharmaceutical composition of the invention which comprises mixing a compound of formula (I) or a pharmaceutically acceptable salt thereof as hereinbefore defined with a pharmaceutically acceptable adjuvant, diluent or carrier.

Conventional procedures for the selection and preparation of suitable pharmaceutical formulations are described in, for example, "Pharmaceutics - The Science of Dosage Form Design", M. E. Aulton, Churchill Livingstone, 1988.

Pharmaceutically acceptable adjuvants, diluents or carriers that may be used in the pharmaceutical compositions of the invention are those conventionally employed in the field of pharmaceutical formulation, and include, but are not limited to, sugars, sugar alcohols, starches, ion exchangers, alumina, aluminium stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycerine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulphate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene- polyoxypropylene-block polymers, polyethylene glycol and wool fat.

The pharmaceutical compositions of the present invention may be administered orally, topically, parenterally, by inhalation spray, rectally, nasally, buccally, vaginally or via an implanted reservoir. Oral or topical administration is preferred. The pharmaceutical compositions of the invention may contain any conventional non-toxic pharmaceutically acceptable adjuvants, diluents or carriers. The term parenteral as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intra-articular, intrasynovial, intrasternal, intrathecal, intralesional and intracranial injection or infusion techniques.

The pharmaceutical compositions may be in the form of a sterile injectable preparation, for example, as a sterile injectable aqueous or oleaginous suspension. The suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol.

Among the acceptable diluents and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant.

The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, powders, granules, and aqueous suspensions and solutions. These dosage forms are prepared according to techniques well-known in the art of pharmaceutical formulation. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are administered orally, the active ingredient is combined with

emulsifying and suspending agents. If desired, certain sweetening and/or flavouring and/or colouring agents may be added.

The pharmaceutical compositions of the invention may also be administered in the form of suppositories for rectal administration. These compositions can be prepared by mixing the active ingredient with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the active ingredient. Such materials include, but are not limited to, cocoa butter, beeswax and polyethylene glycols.

The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilising or dispersing agents known in the art.

Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.05 to 80 %w, still more preferably from 0.10 to 70 %w, and even more preferably from 0.10 to 50 %w, of active ingredient, all percentages by weight being based on total composition.

The compounds of the invention (that is, compounds of formula (I) and pharmaceutically acceptable salts thereof) may also be administered in conjunction with other compounds used for the treatment of the above conditions.

The invention therefore further relates to combination therapies wherein a compound of the invention or a pharmaceutical composition or formulation comprising a compound of the invention is administered concurrently, sequentially or as a combined preparation with another therapeutic agent or agents, for the treatment of one or more of the conditions previously indicated. Such therapeutic agents may be selected from the following:

(i) **opioid agonists, partial agonists or antagonists** such as, for example, morphine, buprenorphine, alfentanil, fentanyl, pethidine, oxycodone, tramadol and codeine, and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof;

5

(ii) **antidepressants** such as, for example, amitriptyline, amoxapine, bupropion, citalopram, clomipramine, desipramine, doxepin duloxetine, elzasonan, escitalopram, fluvoxamine, fluoxetine, gepirone, imipramine, ipsapirone, maprotiline, nortriptyline, nefazodone, paroxetine, phenelzine, protriptyline, reboxetine, robaizotan, sertraline, sibutramine, thionisoxetine, tranlycypromaine, trazodone, trimipramine, venlafaxine, ketamine, vortioxetine, and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof;

10

(iii) **anxiolytics** including, for example, alnespirone, azapirones, benzodiazepines, barbiturates, and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof. Example anxiolytics include adinazolam, alprazolam, balezepam, bentazepam, bromazepam, brotizolam, buspirone, clonazepam, clorazepate, chlordiazepoxide, cyprazepam, diazepam, diphenhydramine, estazolam, fenobam, flunitrazepam, flurazepam, fosazepam, lorazepam, lormetazepam, meprobamate, midazolam, nitrazepam, oxazepam, prazepam, quazepam, reclazepam, tracazolate, trepipam, temazepam, triazolam, uldazepam, and zolazepam; and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof;

15

20

(iv) **anticonvulsants** including, for example, carbamazepine, valproate, lamotrigine, and gabapentin, and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof;

25

(v) **migraine therapies** including, for example, almotriptan, amantadine, bromocriptine, butalbital, cabergoline, dichloralphenazone, eletriptan, frovatriptan, lisuride, naratriptan, pergolide, pramipexole, rizatriptan, ropinirole, sumatriptan, zolmitriptan, and zomitriptan, and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof;

30

(vi) **stroke therapies** including, for example, abciximab, activase, NXY-059, citicoline, crobenetine, desmoteplase, repinotan, traxoprodil, and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof;

5

(vii) **urinary incontinence therapies** including, for example, darafenacin, falvoxate, oxybutynin, propiverine, robalzotan, solifenacin, and tolterodine, and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof;

10 (viii) **neuropathic pain therapies** including, for example, gabapentin, lidoderm, and pregablin, and equivalents and pharmaceutically active isomer(s) and/or metabolite(s) thereof; and

(ix) **nociceptive pain therapies** such as, for example, celecoxib, etoricoxib, lumiracoxib, rofecoxib, valdecoxib, diclofenac, loxoprofen, naproxen, and paracetamol, and equivalents
15 and pharmaceutically active isomer(s) and/or metabolite(s) thereof.

Such combination products employ the compounds of this invention within the dosage range described herein and the other pharmaceutically active agent within approved dosage
20 ranges.

The present invention will now be further explained by reference to the following illustrative examples. In the illustrative examples, the compounds synthesised are both named and illustrated structurally. Whilst every effort has been made to ensure that the
25 chemical names and the chemical structures are consistent, if any inconsistencies occur the illustrated chemical structure should be taken to be correct, unless the illustrated chemical structure is chemically impossible.

Nuclear magnetic resonance (NMR) spectra were typically recorded at 400 MHz; the
30 chemical shifts (δ) are reported in parts per million. Spectra were recorded using a Bruker

400 Avance instrument fitted with a 5mm BBFO probe or DUL probe. Instrument control was by Bruker TopSpin 2.1 software, unless stated otherwise.

Purity was assessed using UPLC with UV (photodiode array) detection over a wide range of wavelengths, normally 220-450nm, using a Waters Acquity UPLC system equipped with Acquity UPLC BEH or HSS C18 columns (2.1mm id × 50mm long) operated at 50 or 60 °C. Mobile phases typically consisted of MeCN or MeOH mixed with water containing either 0.05% formic acid or 0.025% NH₄OH. Mass spectra were recorded with a Waters SQD single quadrupole mass spectrometer using atmospheric pressure ionisation, unless stated otherwise.

Compounds were purified using normal phase chromatography on silica or alumina, or by reverse-phase chromatographic methods, using Biotage or Isolute KPNH Cartridge, SCX cartridge and SCX-2 solid phase extraction cartridges.

Preparative HPLC was typically performed using an Agilent Technologies 1100 Series system or a Waters autopurification LC/MS system typically using Waters 19mm id × 100mm long C18 columns such as XBridge or SunFire 5µm materials at room temperature. Mobile phases typically consisted of MeCN or MeOH mixed with water containing either 0.1% formic acid or 0.1% NH₄OH, unless stated otherwise.

The compound names were generated using ChemBioDraw Ultra 12.0 (Perkin Elmer Inc.).

Generally:

- (i) Operations were typically carried out at ambient temperature, i.e. in the range 15-25 °C, and under an atmosphere of an inert gas such as nitrogen or argon unless otherwise stated.
- (ii) Inorganic solutions were aqueous unless otherwise stated.
- (iii) Drying of solutions was carried out with magnesium sulfate or sodium sulfate, or using a phase-separating cartridge.
- (iv) Concentration was typically carried out by rotary evaporation under vacuum.

- (v) Column chromatography and high pressure liquid chromatography (HPLC) were performed on silica or reverse-phase C18 silica; unless otherwise stated, preparative HPLC was carried out using MeCN/water (both with 0.1% NH₄OH).
- (vi) Yields, where present, are not necessarily the maximum attainable.
- 5 (vii) In general, the structures of the end-products of the Formula I were confirmed by nuclear magnetic resonance (NMR) and/or mass spectral (MS) techniques; electrospray ionization (in positive or negative mode, ES⁺ or ES⁻) mass spectral data were obtained using a Waters Amity SQD spectrometer and, where appropriate, either positive ion data or negative ion data were collected; NMR chemical shift
- 10 values were measured on the delta scale; the following abbreviations have been used: s, singlet; m, multiplet; br, broad; br s, broad singlet.
- (viii) Intermediates were not necessarily fully purified but their structures and purity were assessed by thin layer chromatography (TLC), HPLC, infra-red (IR) and/or NMR analysis.

15

The following abbreviations are used:


	bp	boiling point
	DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
20	DCM	dichloromethane
	DIPEA	diisopropylethylamine
	DMF	<i>N,N</i> -dimethylformamide
	DMSO	dimethyl sulfoxide
	DMSO- <i>d</i> ₆	deuterated dimethyl sulfoxide
25	eq	equivalent(s)
	Et	ethyl
	Et ₂ O	diethyl ether
	EtOAc	ethyl acetate
	EtOH	ethanol
30	h	hour(s)

	HATU	<i>O</i> -(7-azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
	HBTU	<i>O</i> -(benzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
5	HCl	hydrochloric acid
	IMS	industrial methylated spirit
	iPr	isopropyl
	M	molar
	Me	methyl
10	MeCN	acetonitrile
	MeOH	methanol
	mmHg	millimetres of mercury
	mmol	millimoles
	NaOH	sodium hydroxide
15	petrol	petroleum ether, typically bp 40-60 °C
	RT	room temperature, typically 15-25°C
	THF	tetrahydrofuran

A. Preparation of Acid Intermediates



Intermediate A1: 4-(2-Fluorophenoxy)benzoic acid



Step (i) Potassium carbonate (31.1 g, 225 mmol) was added to a solution of 4-fluorobenzaldehyde (7.96 mL, 75 mmol) and 2-fluorophenol (9 mL, 98 mmol) in DMF (100 mL). The mixture was heated at 100 °C under nitrogen for 24 h. After cooling to RT, the mixture was partitioned between EtOAc (200 mL) and water (150 mL). The aqueous

phase was extracted with EtOAc (200 mL). The combined organic phases were washed with brine (3×50 mL), dried and concentrated. The residue was purified by silica chromatography (0-20% EtOAc/petrol) to give 4-(2-fluorophenoxy)benzaldehyde (crude, 17.7 g) which was used in the next step (ii) without further purification or characterisation.


5

Step (ii) 4-(2-Fluorophenoxy)benzaldehyde (crude, 17.7 g) was dissolved in THF (100 mL), *tert*-butanol (100 mL) and water (50 mL), and the mixture cooled in an ice-water bath. 2-Methyl-2-butene (43.3 mL, 409 mmol), potassium dihydrogen phosphate (27.8 g, 205 mmol) and sodium chlorite (23.12 g, 205 mmol) were added sequentially and the mixture was stirred at RT for 18 h. The mixture was partitioned between water (100 mL) and Et₂O (200 mL), and the aqueous phase extracted with Et₂O (200 mL). The combined organic phases were washed sequentially with saturated sodium thiosulfate solution (100 mL) and brine (100 mL), dried and concentrated. Addition of EtOAc and petrol resulted in a precipitate which was isolated by filtration and air-dried to give the title compound (11.95 g, 63%) as a pale yellow solid.

10

MS ES⁻: 231

Intermediate A2: 4-(2,6-Dimethylphenoxy)benzoic acid



Step (i) Tripotassium phosphate (9.27 g, 43.7 mmol) and palladium(II) acetate (0.490 g, 2.18 mmol) were added to a solution of ethyl 4-bromobenzoate (5 g, 21.8 mmol) and 2,6-dimethylphenol (3.73 g, 30.6 mmol) in toluene (40 mL). The mixture was degassed for about 5 minutes and purged with nitrogen. A solution of di-*tert*-butyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine (1.390 g, 3.27 mmol) in toluene (5 mL) was added. The mixture was degassed for a further 2 minutes, purged with nitrogen, and heated under reflux for 1.5 h. The mixture was partitioned between water and EtOAc (50 mL each), and the organic phase dried and concentrated. The residue was purified by silica chromatography (0-20% EtOAc/petrol) to give ethyl 4-(2,6-dimethylphenoxy)benzoate


20

25

(crude, 2 g) which was used in the next step (ii) without further purification or characterisation.

Step (ii) Ethyl 4-(2,6-dimethylphenoxy)benzoate (crude, 2 g) was dissolved in MeOH (7 mL). NaOH (2 M, 7 mL) was added, and the mixture heated under microwave irradiation at 120 °C for 20 minutes. The mixture was acidified to pH 1 (2 M HCl), diluted with water (50 mL) and extracted with EtOAc (50 mL). The organic phase was dried and concentrated. The residue was dissolved/suspended in EtOAc (50 mL) and the mother liquor concentrated. This process was repeated with Et₂O:EtOAc (4:1, 20 mL) and then Et₂O (20 mL) to give the title compound (crude, 800 mg) which was used in the next step without further purification or characterisation.

Intermediate A3: 4-(2,6-Difluorophenoxy)benzoic acid



Step (i) 4-(2,6-Difluorophenoxy)benzaldehyde was prepared as described for 4-(2-fluorophenoxy)benzaldehyde (Intermediate A1, step (i)) from 4-fluorobenzaldehyde and 2,6-difluorophenol, except with 2 eq potassium carbonate and 1.2 eq of the phenol.


¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.12-7.21 (m, 2H) 7.32-7.50 (m, 3H) 7.90-7.99 (m, 2H) 9.94 (s, 1H)

Step (ii) 4-(2,6-Difluorophenoxy)benzaldehyde (0.500 g, 2.14 mmol) was dissolved in *tert*-butanol (6 mL) and water (3 mL). 2-Methyl-2-butene (2 M in THF, 5.34 mL, 10.67 mmol), potassium dihydrogen phosphate (0.73 g, 5.34 mmol) and sodium chlorite (0.72 g, 2.14 mmol) were added sequentially and the mixture was stirred at RT for 4 h. The mixture was partitioned between water (30 mL) and Et₂O (30 mL). The aqueous phase was acidified with HCl (2 M, 10 mL) and extracted with Et₂O (30 mL). The combined organic phases were washed sequentially with saturated sodium thiosulfate solution (15 mL) and brine (15 mL), dried and concentrated. The residue was loaded onto an anion exchange cartridge.

After washing with MeCN, the product was eluted with 2 M HCl/MeCN then concentrated to give the title compound (0.367 g, 69%) as a white solid.

MS ES⁻: 249


Intermediate A4: 4-(2-Fluoro-6-methylphenoxy)benzoic acid



Step (i) 4-(2-Fluoro-6-methylphenoxy)benzaldehyde was prepared as described for 4-(2-fluorophenoxy)benzaldehyde (Intermediate A1, step (i)) from 4-fluorobenzaldehyde and 2-fluoro-6-methylphenol, and used in the next step (ii) without further purification or characterisation.

Step (ii) The title compound was prepared as described for 4-(2-fluorophenoxy)benzoic acid (Intermediate A1, step (ii)) from 4-(2-fluoro-6-methylphenoxy)benzaldehyde (step (i) above), with recrystallisation of the product from EtOAc/petrol, and used in the next step without further purification or characterisation.

Intermediate A5: 4-(2-Chloro-6-fluorophenoxy)benzoic acid




Step (i) 4-(2-Chloro-6-fluorophenoxy)benzaldehyde was prepared as described for 4-(2-fluorophenoxy)benzaldehyde (Intermediate A1, step (i)) from 4-fluorobenzaldehyde and 2-chloro-6-fluorophenol.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.07-7.17 (m, 2H) 7.40-7.60 (m, 3H) 7.89-7.99 (m, 2H) 9.93 (s, 1H)

Step (ii) The title compound was prepared as described for 4-(2-fluorophenoxy)benzoic acid (Intermediate A1, step (ii)) from 4-(2-chloro-6-fluorophenoxy)benzaldehyde (step (i) above).

MS ES⁻: 265

Intermediate A6: 2-Fluoro-4-(2-fluorophenoxy)benzoic acid




Step (i) 2-Fluoro-4-(2-fluorophenoxy)benzaldehyde was prepared as described for 4-(2-fluorophenoxy)benzaldehyde (Intermediate A1, step (i)) from 2,4-difluorobenzaldehyde and 2-fluorophenol, except with 1 eq of the phenol and heating at 80 °C.

MS ES⁺: 235

Step (ii) The title compound was prepared as described for 4-(2-fluorophenoxy)benzoic acid (Intermediate A1, step (ii)) from 2-fluoro-4-(2-fluorophenoxy)benzaldehyde (step (i) above).

MS ES⁺: 251


Intermediate A7: 4-((2-Fluorobenzyl)oxy)benzoic acid



Step (i) Sodium hydride (60% dispersion in mineral oil, 0.177 g, 4.43 mmol) was added to a mixture of 4-fluorobenzaldehyde (0.437 mL, 4.03 mmol) and (2-fluorophenyl)methanol (0.508 g, 4.03 mmol) in DMF (20 mL). The reaction was stirred under nitrogen at room temperature for 18 h. The mixture was diluted with EtOAc and washed with sodium bicarbonate and then brine. The organic phase was dried and concentrated to give 4-((2-fluorobenzyl)oxy)benzaldehyde as a residue which was used in the next step (ii) without further purification or characterisation.

Step (ii) The title compound was prepared as described for 4-(2,6-difluorophenoxy)benzoic acid (Intermediate A3, step (ii)) from 4-((2-fluorobenzyl)oxy)benzaldehyde (step (i) above) and used in the next step without further purification or characterisation.


Intermediate A8: 2-Fluoro-4-(2-methoxyphenoxy)benzoic acid



Prepared by a two step process analogous to that described for Intermediate A1 in which, in the first step, 2,4-difluorobenzaldehyde was reacted with 2-methoxyphenol to yield 2-fluoro-4-(2-methoxyphenoxy)benzaldehyde which, without further purification or characterisation, was used in the second oxidation step to yield the titled compound.

The intermediates 4-(2-methylphenoxy)benzoic acid and 4-(2-chlorophenoxy)benzoic acid used in the preparation of several of the final compounds is known in the art and is commercially available from chemical suppliers such as Apollo Scientific.

B. Preparation of 4-(Aminomethyl)benzoates and analogues



Intermediates of formula (B) shown in Table 1 below were prepared according to one of the following methods R1 to R6:

Method R1

A solution of the amine (50 mmol) and methyl 4-formylbenzoate (50 mmol) in IMS or EtOH (100 mL) was heated under reflux for 2 h. After cooling, the mixture was concentrated and the residue was dissolved in MeOH (400 mL). The resulting solution was cooled in an ice-water bath and sodium borohydride (40 mmol) was added, typically in several portions over 10 minutes. The mixture was stirred and allowed to warm to RT. After 2 h, the mixture was concentrated. The residue was stirred in water (20 mL). The mixture was acidified HCl (2 M) and extracted with DCM (3 × 20 mL). The aqueous phase was basified with solid NaOH and extracted with DCM (3 × 50 mL). The latter extracts were dried and concentrated to give the intermediate of formula (B).

Method R2

As for Method R1, except that after concentrating the sodium borohydride reaction, the residue was dissolved/suspended in HCl (2 M) or HCl (4 M in 1,4-dioxane). The resulting precipitate was filtered, washed with water, and dried to give the hydrochloride salt of the intermediate of formula (B).

Method R3

A solution of the amine (5 mmol) and methyl 4-formylbenzoate (5 mmol) was stirred in DCM (20 mL) for 2 h at RT. Sodium triacetoxyborohydride (10 mmol) was added, followed by acetic acid (6 mmol). After stirring for 2 h, the mixture was diluted with DCM (20 mL) and washed with NaOH (2 M, 20 mL) or saturated sodium bicarbonate solution (20 mL). The aqueous phase was extracted with DCM (20 mL), and the combined organic phases dried and concentrated to give the intermediate of formula (B).

Method R4


As for Method R1, except that after concentrating the sodium borohydride reaction and adding water and HCl, the mixture was basified with sodium hydrogencarbonate and extracted with EtOAc. The organic phase was dried and concentrated, and the residue was dissolved/suspended in an organic solvent, typically EtOAc or Et₂O. HCl (4 M in MeOH) was added, and the mixture stirred at RT. The resulting precipitate was filtered, and the filter cake air-dried to give the hydrochloride salt of the intermediate of formula (B).

Method R5





As for Method R1, except that (a) in the imine formation, triethylamine (typically 1 eq) was added, and (b) after concentrating the sodium borohydride reaction, the residue was partitioned between (i) sodium hydrogencarbonate or sodium hydroxide and (ii) EtOAc or DCM. The organic phase was dried and concentrated to give the intermediate of formula (B). In some cases, the residue was dissolved/suspended in an organic solvent, typically EtOAc or Et₂O. HCl (4 M in MeOH) was added, and the mixture stirred at RT. The resulting precipitate was filtered, and the filter cake air-dried to give the hydrochloride salt of the intermediate of formula (B). In some cases, the residue was dissolved/suspended in aqueous HCl (2 M). The resulting precipitate was filtered, washed with water, and dried to give the hydrochloride salt of the intermediate of formula (B).










Method R6




As for Method R1, except that after stirring the sodium borohydride reaction for 1 h, the mixture was quenched with HCl (2 M). The resulting precipitate was isolated by filtration and dried under vacuum to give the hydrochloride salt of the intermediate of formula (B).

**TABLE 1**

Inter-mediate	R	Product	Method	Data
B1	CH ₂ CH ₃	methyl 4- ((ethylamino)methyl)benzoate	R1, using 2 M EtNH ₂ in THF (1.8 eq)	MS ES ⁺ : 194

Inter-mediate	R	Product	Method	Data
B2	CH ₂ CHF ₂	methyl 4-(((2,2-difluoroethyl)amino)methyl)benzoate	R1	MS ES ⁺ : 230
B3 (HCl salt)	CH ₂ CF ₃	methyl 4-(((2,2,2-trifluoroethyl)amino)methyl)benzoate hydrochloride	R2, using 5 eq NaBH ₄	MS ES ⁺ : 248
B4 (HCl salt)	CH ₂ iPr	methyl 4-((isobutylamino)methyl)benzoate hydrochloride	R4	n/a
B5 (HCl salt)		methyl 4-(((cyclopropylmethyl)amino)methyl)benzoate hydrochloride	R4	MS ES ⁺ : 220
B6 (HCl salt)		methyl 4-(((cyclobutylmethyl)amino)methyl)benzoate hydrochloride	R5, from Amine 1*	MS ES ⁺ : 234
B7 (HCl salt)		methyl 4-((isopentylamino)methyl)benzoate hydrochloride	R4	MS ES ⁺ : 236
B8 (HCl salt)		methyl 4-(((2-cyclopropylethyl)amino)methyl)benzoate hydrochloride	R4, from Amine 2**	MS ES ⁺ : 234
B9 (HCl salt)		methyl 4-(((2-cyclobutylethyl)amino)methyl)benzoate hydrochloride	R4, from Amine 3***	MS ES ⁺ : 248
B10 (HCl salt)		methyl 4-((benzylamino)methyl)benzoate hydrochloride	R4	MS ES ⁺ : 256

Inter-mediate	R	Product	Method	Data
B11 (HCl salt)		methyl 4-(((2-fluorobenzyl)amino)methyl)benzoate hydrochloride	R2	MS ES ⁺ : 274
B12 (HCl salt)		methyl 4-(((3-methoxybenzyl)amino)methyl)benzoate hydrochloride	R2	MS ES ⁺ : 286
B13 (HCl salt)		methyl 4-(((4-methoxybenzyl)amino)methyl)benzoate hydrochloride	R2	MS ES ⁺ : 286
B14		methyl 4-((phenethylamino)methyl)benzoate	R3	MS ES ⁺ : 270
B15 (HCl salt)		methyl 4-(((3-phenylpropyl)amino)methyl)benzoate hydrochloride	R6	MS ES ⁺ : 284
B16		methyl 4-((((trans-2-phenylcyclopropyl)methyl)amino)methyl)benzoate	R3	MS ES ⁺ : 296
B17 (HCl salt)		methyl 4-(((3-(2-fluorophenyl)propyl)amino)methyl)benzoate hydrochloride	R5	MS ES ⁺ : 302
B18 (HCl salt)		methyl 4-(((3-(3-fluorophenyl)propyl)amino)methyl)benzoate hydrochloride	R5	MS ES ⁺ : 302
B19 (HCl salt)		methyl 4-(((4-phenylbutyl)amino)methyl)benzoate hydrochloride	R4	n/a

Inter-mediate	R	Product	Method	Data
B19a (HCl salt)	CH ₂ CH ₂ CH ₃	methyl 4- ((propylamino)methyl) benzoate hydrochloride	R2	n/a
B19b (HCl salt)		methyl 4-(((2,2- difluoropropyl)amino)methyl) benzoate	R5	MS ES ⁺ : 244
B19c		methyl 4-(((3,3,3- trifluoropropyl)amino)methyl) benzoate	R3	MS ES ⁺ : 262
B19d (HCl salt)		methyl 4- ((butylamino)methyl)benzoate hydrochloride	R2	n/a

***Amine 1: Cyclobutylmethanamine hydrochloride**

A solution of borane in THF (1 M, 296 mL, 296 mmol) was added to a solution of cyclobutanecarbonitrile (20 g, 247 mmol) in THF (60 mL) under argon, and the mixture heated under reflux overnight. The mixture was cooled in an ice-water bath and MeOH (120 mL) added dropwise while keeping the temperature of the mixture below 20 °C. HCl in MeOH (4 M, 300 mL) was added, again keeping the temperature below 20 °C. The resulting solution was heated under reflux for 2.5 h. After cooling, the mixture was concentrated. The residue was diluted with MeOH (100 mL) and concentrated, and this process was repeated. Ether was added to the residue, and the mixture stirred for 30 minutes before filtering to give the title compound (25.9 g, 86%) as a white solid.

¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.70-1.86 (m, 4H) 1.97-2.04 (m, 2H) 2.46-2.61 (m, 1H) 2.75-2.84 (m, 2H) 8.02 (br s, 3H)

****Amine 2: 2-Cyclopropylethanamine**


Sulfuric acid (17.25 mL, 324 mmol) was added dropwise to a suspension of lithium aluminium hydride (24.56 g, 647 mmol) in Et₂O (900 mL) in an ice-water bath under

argon. The mixture was stirred at RT for 1 h. A solution of 2-cyclopropylacetonitrile (17.5 g, 216 mmol) in Et₂O (100 mL) was added and the mixture heated under reflux for 18 h. The mixture was cooled in an ice-water bath and sodium sulfate decahydrate was added until effervescence ceased. The mixture was filtered and concentrated to give the title compound (crude, 18.6 g) which was used without further purification or characterisation.

*****Amine 3: 2-Cyclobutylethanamine**

Prepared as described for Amine 2 from 2-cyclobutylacetonitrile and used in the next step without further purification or characterisation.

Intermediate B20: (S)-methyl 4-(((2-hydroxy-3-phenylpropyl)amino)methyl)benzoate



Step (i) (S,S)-(+)-N,N'-Bis(3,5-di-*tert*-butylsalicylidene)-1,2-cyclohexanediaminocobalt(II) (0.048 g, 0.080 mmol), THF (0.25 mL) and acetic acid (0.018 mL, 0.318 mmol) were added to 2-benzyloxirane (4.27 g, 31.8 mmol). The mixture was cooled in an ice-water bath and water (0.129 mL, 7.16 mmol) was added in one portion. After stirring at RT overnight, the mixture was distilled under vacuum (2 mmHg, bp 60-80 °C) to give (S)-2-benzyloxirane (crude, 2.74 g) which was used in the next step without further purification or characterisation.


Step (ii) (S)-2-Benzyloxirane (step (i) above) (2.74 g, 20.42 mmol) was added to a solution of methyl 4-(aminomethyl)benzoate hydrochloride (7.41 g, 36.8 mmol) and DIPEA (8.92 mL, 51.1 mmol) in MeOH (300 mL). After stirring for 3 days, the reaction mixture was concentrated. The residue was partitioned between saturated sodium hydrogencarbonate solution (200 mL) and DCM (2 × 200 mL). The aqueous phase was extracted with DCM (200 mL). The combined organic phases were concentrated and purified by silica

chromatography (0-10% MeOH/DCM) and then reverse-phase chromatography on C18 silica (eluted with 0-95% MeOH/H₂O with 0.1% NH₄OH) to give the title compound (3.27 g, 53%) as a pale brown gum.

MS ES⁺: 300

5

Intermediate B21: (*R*)-methyl 4-(((2-hydroxy-3-phenylpropyl)amino)methyl)benzoate




Step (i) Prepared as described for (*S*)-2-benzyloxirane (Intermediate B48, step (i)), except using (*R,R*)-(-)-*N,N'*-bis(3,5-di-*tert*-butylsalicylidene)-1,2-cyclohexanediaminocobalt(II) to give (*R*)-2-benzyloxirane which was used in the next step without further purification or characterisation.

Step (ii) Prepared as described for (*S*)-methyl 4-(((2-hydroxy-3-phenylpropyl)amino)methyl)benzoate (Intermediate B20, step (ii)) from (*R*)-2-benzyloxirane (step (i) above) and methyl 4-(aminomethyl)benzoate hydrochloride to give the title compound as a pale brown gum.

MS ES⁺: 300

20 C. Preparation of Intermediate Esters

Intermediates of formula (C) shown in Tables 2 to 4 below were prepared according to one of the following methods E1 to E5:



Method E1

5 Triethylamine (6 mmol), 1-hydroxy-7-azabenzotriazole (2.4 mmol) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) (2.4 mmol) were added to a solution/suspension of the relevant amine or amine hydrochloride (2 mmol) and 4-phenoxybenzoic acid (2 mmol) in THF (10 mL). The reaction was stirred at RT for 18 h. (In cases where the amine hydrochloride was used, an extra 2 mmol triethylamine was

10 added.) The mixture was partitioned between (i) water or saturated sodium hydrogencarbonate solution (10 mL) and (ii) DCM or EtOAc (10 mL). The aqueous phase was extracted with further DCM or EtOAc, and the combined organic phases dried and concentrated. The residue was purified by silica chromatography (typically 0-100% EtOAc/petrol).

Method E2

DIPEA (2.2 mmol) and HBTU (1.1 mmol) were added to a solution of the relevant acid in DMF (5 mL) and the mixture stirred for 5 minutes at RT. The relevant amine or amine hydrochloride (0.95 mmol) and DBU (1.5 mmol) were added. The mixture was stirred for


20 1 h at RT. (In cases where the amine hydrochloride was used, an extra 1 mmol DIPEA was added.) The mixture was partitioned between EtOAc (30 mL) and either water or saturated sodium hydrogencarbonate solution (30 mL). In some cases the aqueous phase was further extracted with EtOAc. The combined organic phases were dried and concentrated. The residue was purified by silica chromatography (typically 0-100% EtOAc/petrol).

Method E3

Thionyl chloride or oxalyl chloride (3-12 mmol) was added to a solution of the acid (3 mmol) in DCM (10 mL) in an ice-water bath. In some cases, several drops of DMF were added. The reaction mixture was stirred at RT for 4 h and then concentrated to yield the crude acid chloride, which was dissolved in THF (3 mL) and added to a solution/suspension of the amine hydrochloride (2 mmol) and potassium carbonate (10 mmol) in THF (10 mL). The reaction was heated under reflux for 18 h. After cooling, the mixture was partitioned between water and EtOAc. The organic phase was dried and concentrated and the residue purified by silica chromatography (0-50% EtOAc/petrol).

Method E4

HATU (1.1 mmol) was added to a solution of the acid (1 mmol) and DIPEA (2.2 mmol) in DMF or THF (2 mL). The mixture was stirred at RT and allowed to stand. After 5 minutes, the relevant amine or amine hydrochloride (1.1 mmol) was added. (In some cases, 0.9 mmol of amine or amine hydrochloride was used.) When the amine hydrochloride was used, DBU (1.5 mmol) was added at this stage. After 10 minutes, the mixture was diluted with EtOAc or DCM (30 mL) and washed with either HCl (1 M, 20 mL) or saturated sodium bicarbonate solution (20 mL) or water (20 mL). In some cases, the aqueous phase was further extracted with EtOAc or DCM. The combined organic phases were washed with water (2 × 20 mL), dried and concentrated. In some cases further purification was carried out, by silica chromatography (typically 0-100% EtOAc/petrol), reverse-phase chromatography on C18 silica (typically eluting with 5-95% MeCN/water) or preparative HPLC.



Method E5

Step (i) Triethylamine (1.351 mL, 9.69 mmol) was added slowly to a mixture of 4-bromobenzoyl chloride (0.851 g, 3.88 mmol) and methyl 4-((phenethylamino)methyl)-benzoate (Intermediate B14) (0.870 g, 3.23 mmol) in DCM (20 mL) at RT under nitrogen and the mixture stirred for 18 h. The mixture was diluted with DCM (30 mL), water (10 mL) and HCl (2 M, 10 mL), and the phases separated. The aqueous phase was extracted with DCM (30 mL) and the combined organic phases washed with water (10 mL), dried and concentrated. The residue was purified by silica chromatography (0-50% EtOAc/petrol) to give methyl 4-((4-bromo-*N*-phenethylbenzamido)methyl)benzoate (1.21 g, 83%) as a pale orange oil.

MS ES⁺: 452, 454

Step (ii) Tripotassium phosphate (2 mmol), palladium(II) acetate (0.1 mmol) and di-*tert*-butyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine (0.15 mmol) were added to a solution of methyl 4-((4-bromo-*N*-phenethylbenzamido)methyl)benzoate (step (i) above) (1 mmol) and the relevant phenol (1.2 mmol) in toluene (5 mL). The mixture was degassed for about 5 minutes, purged with nitrogen and heated either thermally or under microwave irradiation at 120 °C for 20 h. The mixture was filtered through diatomaceous earth, washing through with EtOAc. The filtrate was concentrated and purified either by silica chromatography (typically 0-100% EtOAc/petrol) or preparative HPLC.



















TABLE 2

Inter- mediate	R	Product	Method	Data
C6	CH ₂ CH ₃	methyl 4-((N-ethyl-4-(2-fluorophenoxy)benzamido)methyl)benzoate	E2	MS ES ⁺ : 408
C7	CH ₂ CHF ₂	methyl 4-((N-(2,2-difluoroethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoate	E2	MS ES ⁺ : 444
C8	CH ₂ CF ₃	methyl 4-((4-(2-fluorophenoxy)-N-(2,2,2-trifluoroethyl)benzamido)methyl)benzoate	E3	MS ES ⁺ : 462
C9	CH ₂ iPr	methyl 4-((4-(2-fluorophenoxy)-N-isobutylbenzamido)methyl)benzoate	E1	MS ES ⁺ : 436
C10		methyl 4-((N-(cyclopropylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoate	E1	MS ES ⁺ : 434
C11		methyl 4-((N-(cyclobutylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoate	E4	MS ES ⁺ : 448

Inter- mediate	R	Product	Method	Data
C12		methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -isopentylbenzamido)methyl)benzoate	E1	MS ES ⁺ : 450
C13		methyl 4-((<i>N</i> -(2-cyclopropylethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoate	E1	MS ES ⁺ : 448
C14		methyl 4-((<i>N</i> -(2-cyclobutylethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoate	E1	MS ES ⁺ : 462
C15		methyl 4-((<i>N</i> -benzyl-4-(2-fluorophenoxy)benzamido)methyl)benzoate	E1	MS ES ⁺ : 470
C16		methyl 4-((<i>N</i> -(2-fluorobenzyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoate	E4	MS ES ⁺ : 488
C17		methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -(3-methoxybenzyl)benzamido)methyl)benzoate	E4	n/a
C18		methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -(4-methoxybenzyl)benzamido)methyl)benzoate	E4	n/a

Inter- mediate	R	Product	Method	Data
C19		methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -phenethylbenzamido)methyl)benzoate	E4	MS ES ⁺ : 484
C20		methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -(3-(2-fluorophenyl)propyl)benzamido)methyl)benzoate	E4	MS ES ⁺ : 516
C21		methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -(3-(3-fluorophenyl)propyl)benzamido)methyl)benzoate	E4	MS ES ⁺ : 516
C22		methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -((<i>trans</i> -2-phenylcyclopropyl)methyl)benzamido)methyl)benzoate	E4	MS ES ⁺ : 510
C23		(<i>S</i>)-methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -(2-hydroxy-3-phenylpropyl)benzamido)methyl)benzoate	E4	MS ES ⁺ : 514
C24		(<i>R</i>)-methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -(2-hydroxy-3-phenylpropyl)benzamido)methyl)benzoate	E4	MS ES ⁺ : 514

Inter- mediate	R	Product	Method	Data
C25		methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -(4-phenylbutyl)benzamido) methyl)benzoate	E2	MS ES ⁺ : 512
C25a	CH ₂ CH ₂ CH ₃	methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -propylbenzamido)- methyl)benzoate	E4	MS ES ⁺ : 422
C25b	CH ₂ CH ₂ CF ₃	methyl 4-((4-(2-fluorophenoxy)- <i>N</i> -(3,3,3-trifluoropropyl)benzamido) methyl)benzoate	E4	MS ES ⁺ : n/a
C25c	CH ₂ CF ₂ CH ₃	methyl 4-((<i>N</i> -(2,2-difluoropropyl)-4-(2-fluorophenoxy)benzamido) methyl)benzoate	E3	MS ES ⁺ : n/a
C25d		methyl 4-((<i>N</i> -butyl-4-(2-fluorophenoxy)benzamido) methyl)benzoate	E4	MS ES ⁺ : 436













TABLE 3

Inter-mediate	X	R	Product	Method	Data
C26	Me		methyl 4-((<i>N</i> -(cyclopropylmethyl)-4-(<i>o</i> -tolylloxy)benzamido)methyl)benzoate	E4	MS ES ⁺ : 430
C27	Me		methyl 4-((<i>N</i> -(cyclobutylmethyl)-4-(<i>o</i> -tolylloxy)benzamido)methyl)benzoate	E4	MS ES ⁺ : 444
C28	Me		methyl 4-((<i>N</i> -benzyl-4-(<i>o</i> -tolylloxy)benzamido)methyl)benzoate	E4	MS ES ⁺ : 466
C29	Me		methyl 4-((<i>N</i> -phenethyl-4-(<i>o</i> -tolylloxy)benzamido)methyl)benzoate	E5	MS ES ⁺ : 480
C30	CF ₃		methyl 4-((<i>N</i> -phenethyl-4-(2-(trifluoromethyl)phenoxy)benzamido)-methyl)benzoate	E5	MS ES ⁺ : 534
C31	CN		methyl 4-((<i>N</i> -benzyl-4-(2-cyanophenoxy)benzamido)methyl)-benzoate	E4	MS ES ⁺ : 477
C32	OMe	CH ₂ CH ₃	methyl 4-((<i>N</i> -ethyl-4-(2-methoxyphenoxy)benzamido)methyl)-benzoate	E2	MS ES ⁺ : 420
C33	OMe	CH ₂ CHF ₂	methyl 4-((<i>N</i> -(2,2-difluoroethyl)-4-(2-methoxyphenoxy)benzamido)methyl)-benzoate	E2	MS ES ⁺ : 456

Inter-mediate	X	R	Product	Method	Data
C34	OMe	CH ₂ iPr	methyl 4-((<i>N</i> -isobutyl-4-(2-methoxyphenoxy)benzamido)methyl)-benzoate	E4	MS ES ⁺ : 448
C35	OMe		methyl 4-((<i>N</i> -(cyclopropylmethyl)-4-(2-methoxyphenoxy)benzamido)methyl)-benzoate	E4	MS ES ⁺ : 446
C36	OMe		methyl 4-((4-(2-methoxyphenoxy)- <i>N</i> -(3-phenylpropyl)benzamido)methyl)benzoate	E1	MS ES ⁺ : 510
C37	OMe		methyl 4-((<i>N</i> -(3-(3-fluorophenyl)propyl)-4-(2-methoxyphenoxy)benzamido)methyl)-benzoate	E1	MS ES ⁺ : 528
C38	OEt		methyl 4-((<i>N</i> -(cyclopropylmethyl)-4-(2-ethoxyphenoxy)benzamido)methyl)-benzoate	E5	MS ES ⁺ : 460
C39	Cl		methyl 4-((4-(2-chlorophenoxy)- <i>N</i> -(cyclopropylmethyl)benzamido)methyl)benzoate	E4	MS ES ⁺ : 450
C40	Cl		methyl 4-((4-(2-chlorophenoxy)- <i>N</i> -phenethylbenzamido)methyl)benzoate	E5	MS ES ⁺ : 500









Inter- mediate	X	R	Product	Method	Data
C40a	OM e	CH ₂ CF ₃	methyl 4-((4-(2-methoxyphenoxy)- <i>N</i> -(2,2,2-trifluoroethyl)benzamido)-methyl)benzoate	E3	MS ES ⁺ : n/a
C40b	OM e	CH ₂ CH ₂ CH ₃	methyl 4-((4-(2-methoxyphenoxy)- <i>N</i> -propylbenzamido)methyl)benzoate	E4	MS ES ⁺ : n/a
C40c	OM e	CH ₂ CF ₂ CH ₃	methyl 4-((<i>N</i> -(2,2-difluoropropyl)-4-(2-methoxyphenoxy)benzamido)methyl)-benzoate	E3	MS ES ⁺ : n/a
C40d	OM e	CH ₂ CH ₂ CF ₃	methyl 4-((<i>N</i> -(3,3,3-trifluoropropyl)-4-(2-methoxyphenoxy)benzamido)-methyl)benzoate	E4	MS ES ⁺ : 488
C40e	OM e		methyl 4-((<i>N</i> -butyl-4-(2-methoxyphenoxy)benzamido)methyl)-benzoate	E4	MS ES ⁺ : n/a
C40f	Me	CH ₂ CF ₃	methyl 4-((4-(<i>o</i> -tolylloxy)- <i>N</i> -(2,2,2-trifluoroethyl)benzamido)methyl)-benzoate	E3	MS ES ⁺ : n/a
C40g	Cl	CH ₂ CF ₃	methyl 4-((4-(2-chlorophenoxy)- <i>N</i> -(2,2,2-trifluoroethyl)benzamido)-methyl)benzoate	E3	MS ES ⁺ : n/a











Inter-mediate	X	R	Product	Method	Data
C40h	Me	CH ₂ CH ₃	methyl 4-((<i>N</i> -ethyl-4-(<i>o</i> -tolylloxy)benzamido)methyl)benzoate	E4	MS ES ⁺ : n/a
C40i	Cl	CH ₂ CH ₃	methyl 4-((4-(2-chlorophenoxy)- <i>N</i> -ethylbenzamido)methyl)benzoate	E4	MS ES ⁺ : n/a
C40j	Cl	CH ₂ CHF ₂	methyl 4-((4-(2-chlorophenoxy)- <i>N</i> -(2,2-difluoroethyl)benzamido)methyl)-benzoate	E3	MS ES ⁺ : n/a













5 TABLE 4

Inter-mediate	R ¹	R	Product	Method	Data
C41			methyl 4-((<i>N</i> -(cyclopropylmethyl)-4-(2,6-difluorophenoxy)benzamido)methyl)benzoate	E1	MS ES ⁺ : 452


Inter- mediate	R ^I	R	Product	Method	Data
C42			methyl 4-((<i>N</i> -(2-cyclopropylethyl)-4-(2,6-difluorophenoxy)-benzamido)methyl)benzoate	E1	MS ES ⁺ : 466
C43			methyl 4-((<i>N</i> -benzyl-4-(2,6-difluorophenoxy)benzamido)-methyl)benzoate	E1	MS ES ⁺ : 488
C44			methyl 4-((<i>N</i> -(cyclopropylmethyl)-4-(2-fluoro-6-methylphenoxy)benzamido)-methyl)benzoate	E4	MS ES ⁺ : 448
C45			methyl 4-((<i>N</i> -(cyclobutylmethyl)-4-(2-fluoro-6-methylphenoxy)-benzamido)methyl)benzoate	E4	MS ES ⁺ : 462
C46			methyl 4-((4-(2-chloro-6-fluorophenoxy)- <i>N</i> -(cyclopropylmethyl)benzamido)-methyl)benzoate	E1	MS ES ⁺ : 468

Inter- mediate	R ¹	R	Product	Method	Data
C47			methyl 4-((4-(2-chloro-6-fluorophenoxy)- <i>N</i> -(2-cyclopropylethyl)benzamido)-methyl)benzoate	E1	MS ES ⁺ : 482
C48			methyl 4-((<i>N</i> -benzyl-4-(2-chloro-6-fluorophenoxy)benzamido)methyl)-benzoate	E1	MS ES ⁺ : 504
C49			methyl 4-((4-(2,6-dimethylphenoxy)- <i>N</i> -isopentylbenzamido)methyl)-benzoate	E4	MS ES ⁺ : 460
C50			methyl 4-((<i>N</i> -benzyl-4-(2,6-dimethylphenoxy)benzamido)-methyl)benzoate	E4	MS ES ⁺ : 480
C51			methyl 4-((4-(2,6-dimethylphenoxy)- <i>N</i> -phenethylbenzamido)methyl)-benzoate	E4	MS ES ⁺ : 494

Inter- mediate	R ¹	R	Product	Method	Data
C52			methyl 4-((4-(3-methoxyphenoxy)- <i>N</i> -phenethyl benzamido)methyl)-benzoate	E5	MS ES ⁺ : 496
C53			methyl 4-((4-(4-methoxyphenoxy)- <i>N</i> -phenethyl benzamido)-methyl)benzoate	E5	MS ES ⁺ : 496
C54			methyl 4-((4-(3-chlorophenoxy)- <i>N</i> -phenethyl benzamido)methyl)-benzoate	E5	MS ES ⁺ : 500
C55			methyl 4-((<i>N</i> -(cyclopropylmethyl)-4-((2-fluorobenzyl)oxy) benzamido)-methyl)benzoate	E4	MS ES ⁺ : 448
C56			methyl 4-((<i>N</i> -(cyclobutylmethyl)-4-((2-fluorobenzyl)oxy) benzamido)-methyl)benzoate	E4	MS ES ⁺ : 462

Inter- mediate	R ¹	R	Product	Method	Data
C57			methyl 4-((N-benzyl-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoate	E4	MS ES ⁺ : 484

Intermediate C58: Methyl 4-((4-(2-fluorophenoxy)-N-((3-methoxycyclobutyl)methyl)benzamido)methyl)benzoate (mixture of *cis* and *trans* ring isomers)



Step (i) Prepared using the HATU coupling procedure described in Method E4 above from 3-methoxycyclobutanecarboxylic acid (500 mg, 3.84 mmol) and (4-bromophenyl)methanamine hydrochloride (940 mg, 4.23 mmol) to give *N*-(4-bromobenzyl)-3-methoxycyclobutanecarboxamide (mixture of *cis* and *trans* ring isomers) (880 mg, 77%).

MS ES⁺: 298, 300


Step (ii) Lithium aluminium hydride (1 M solution in THF, 2.62 mL, 2.62 mmol) was added to a solution of *N*-(4-bromobenzyl)-3-methoxycyclobutanecarboxamide from step (i) above (780 mg, 2.62 mmol) in THF (20 mL) at RT. After stirring for 18 h, the mixture was quenched with saturated sodium sulfate dodecahydrate solution (5 mL), filtered and concentrated. The residue was dissolved in DCM (50 mL) and washed with NaOH (2 M, 50 mL). The organic phase was concentrated and the residue purified by silica

chromatography (0-30% EtOAc/petrol) to give *N*-(4-bromobenzyl)-1-(3-methoxycyclobutyl)methanamine (mixture of *cis* and *trans* ring isomers) which was used in the next step without further purification or characterisation.

5 Step (iii) Prepared using the HATU coupling procedure described in Method E4 above from 4-(2-fluorophenoxy)benzoic acid (Intermediate A1) (310 mg, 1.337 mmol) and *N*-(4-bromobenzyl)-1-(3-methoxycyclobutyl)methanamine from step (ii) above (380 mg, 1.337 mmol) to give the desired compound, *N*-(4-bromobenzyl)-4-(2-fluorophenoxy)-*N*-((3-methoxycyclobutyl)methyl)benzamide (mixture of *cis* and *trans* ring isomers), along with
10 the corresponding des-bromo analogue *N*-benzyl-4-(2-fluorophenoxy)-*N*-((3-methoxycyclobutyl)methyl)benzamide (also a mixture of *cis* and *trans* ring isomers). This material (approximately 1:1 mixture of bromo:des-bromo compound, 684 mg in total) was used in the next step without further purification. MS ES⁺: 498, 500

15 Step (iv) A solution of palladium(II) acetate (27.0 mg, 0.120 mmol), 1,3-bis(diphenylphosphino)propane (49.7 mg, 0.120 mmol), and *N*-(4-bromobenzyl)-4-(2-fluorophenoxy)-*N*-((3-methoxycyclobutyl)methyl)benzamide from step (iii) above (300 mg, 0.602 mmol) in MeOH (5 mL) and DMF (3 mL) was degassed and purged with nitrogen. Triethylamine (0.168 mL, 1.204 mmol) was added, the reaction mixture was
20 evacuated and carbon monoxide passed through the solution. After stirring under carbon monoxide at 70°C for 18 h, the mixture was MeOH-evaporated and the residue dissolved in DCM (20 mL). The organic phase was washed with water (20 mL) and concentrated. The resulting residue was purified by silica chromatography (0-30% EtOAc/petrol) to give the title compound (115 mg, 80% based on starting material being 50% pure).
25 MS ES⁺: 478

Intermediate C59: Methyl 4-((*N*-(cyclopropylmethyl)-2-fluoro-4-(2-fluorophenoxy)benzamido)methyl)benzoate




Prepared using the EDC coupling procedure described in Method E1 above from 2-fluoro-4-(2-fluorophenoxy)benzoic acid (Intermediate A6) and methyl 4-

((cyclopropylmethylamino)methyl)benzoate hydrochloride (Intermediate B5).

MS ES⁺: 452

Intermediate C60: Methyl 4-((N-benzyl-4-(2-fluorophenoxy)-cyclohexanecarboxamido)methyl)benzoate (ca. 1:1 mixture of *cis* and *trans* ring isomers)



Step (i) 4-Toluenesulfonyl chloride (18.29 g, 96 mmol) was added in several portions to a solution of ethyl 4-hydroxycyclohexanecarboxylate (ca. 1:1 mixture of *cis* and *trans* ring isomers) (29.5 g, 90 mmol) in pyridine (100 mL) in an ice-water bath, and the mixture stirred, allowing to warm to RT. After the solid had dissolved, the mixture was allowed to stand. After 24 h, the mixture was concentrated and the residue partitioned between water and EtOAc (100 mL each). The organic phase was dried and concentrated to give ethyl 4-(tosyloxy)cyclohexanecarboxylate (ca. 1:1 mixture of *cis* and *trans* ring isomers) (crude, 29.5 g) as a colourless oil which was used in the next step without further purification.

¹H NMR (400 MHz, CDCl₃) δ ppm 1.20-1.30 (m, 3H) 1.44-2.08 (m, 8H) 2.21-2.43 (m, 1H) 2.48 (s, 3H) 4.08-4.20 (m, 2H) 4.40-4.50 and 4.70-4.78 (both m, total 1H) 7.28-7.40 (m, 2H) 7.78-7.86 (m, 2H)

5 Step (ii) 2-Fluorophenol (2.145 mL, 23.25 mmol) was added to a solution/suspension of ethyl 4-(tosyloxy)cyclohexanecarboxylate from step (i) above and cesium carbonate (7.58 g, 23.25 mmol) in DMF (100 mL) and the mixture stirred for 18 h at 80 °C. After cooling, the mixture was diluted with EtOAc (200 mL) and washed with saturated sodium hydrogencarbonate solution (100 mL) and brine (100 mL). The organic phase was dried
10 and concentrated, and the residue purified by silica chromatography (0-100% EtOAc/petrol) to give ethyl 4-(2-fluorophenoxy)cyclohexanecarboxylate (ca. 1:1 mixture of *cis* and *trans* ring isomers) (crude, 3.41 g) which was used without further purification or characterisation.


15 Step (iii) Lithium hydroxide (0.307 g, 12.80 mmol) was added to a solution of ethyl 4-(2-fluorophenoxy)cyclohexanecarboxylate from step (ii) above (crude, 3.41 g) in 1,4-dioxane (30 mL) and water (30 mL), and the mixture heated under microwave irradiation at 100 °C for 15 minutes. The mixture was concentrated. The residue was loaded onto an anion exchange cartridge. After washing with MeCN, the product was eluted with 2 M
20 HCl/MeCN and concentrated to give 4-(2-fluorophenoxy)cyclohexanecarboxylic acid (ca. 1:1 mixture of *cis* and *trans* ring isomers) (crude, 1.0 g) as a pale yellow solid which was used in the next step without further purification.

MS ES⁻: 237

25 Step (iv) The title compound was prepared using the HATU coupling procedure described in Method E4 above, from 4-(2-fluorophenoxy)cyclohexanecarboxylic acid from step (iii) above and methyl 4-((benzylamino)methyl)benzoate hydrochloride (Intermediate B10).

MS ES⁺: 476

30 **Intermediate C61: Methyl 4-((*trans*-*N*-benzyl-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoate**



Step (i) A solution of ethyl 4-hydroxycyclohexanecarboxylate (ca. 1:1 mixture of *cis* and *trans* ring isomers) (5 g, 29 mmol), 2-methoxyphenol (3.96 g, 31.9 mmol), triphenylphosphine (8.38 g, 31.9 mmol) and *trans*-diisopropyl diazene-1,2-dicarboxylate (DIAD) (6.21 mL, 31.9 mmol) in THF (150 mL) was stirred at RT. The mixture was diluted with EtOAc (100 mL) and washed with saturated sodium hydrogencarbonate solution (50 mL) and brine (50 mL). The organic phase was dried and concentrated, and the residue purified by silica chromatography (0-100% EtOAc/petrol) to give ethyl 4-(2-methoxyphenoxy)cyclohexanecarboxylate (ca. 1:1 mixture of *cis* and *trans* ring isomers) (crude, 4.0 g) which was used without further purification or characterisation.

Step (ii) 4-(2-Methoxyphenoxy)cyclohexanecarboxylic acid was prepared as described for 4-(2-fluorophenoxy)cyclohexanecarboxylic acid (Intermediate C60, step (iii)) using ethyl 4-(2-methoxyphenoxy)cyclohexanecarboxylate from step (i) above.


MS ES⁺: 249

Step (iii) The title compound was prepared using the HATU coupling procedure described in Method E4 above from 4-(2-methoxyphenoxy)cyclohexanecarboxylic acid (step (ii) above) and methyl 4-((benzylamino)methyl)benzoate hydrochloride (Intermediate B10).

MS ES⁺: 488

Intermediate C62: Methyl 4-((*cis*-N-(3-(3-fluorophenyl)propyl)-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoate
and


Intermediate C63: Methyl 4-((*trans*-N-(3-(3-fluorophenyl)propyl)-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoate



The title compounds were prepared using the HATU coupling procedure described in Method E4 above, from 4-(2-methoxyphenoxy)cyclohexanecarboxylic acid (ca. 1:1 mixture of *cis* and *trans* ring isomers) (Intermediate C61, step (ii)) and methyl 4-(((3-(3-fluorophenyl)propyl)amino)methyl)benzoate hydrochloride (Intermediate B18), and separated by preparative HPLC.

MS ES⁺: 534 in both cases

Intermediate C64: Methyl 2-fluoro-4-((4-(2-fluorophenoxy)-N-(3-methoxybenzyl)benzamido)methyl)benzoate



Step (i) Potassium carbonate (3.70 g, 26.7 mmol) was added to a solution of 4-cyano-2-fluorobenzoic acid (4 g, 24.3 mmol) in DMF (350 mL). After stirring at RT for 15 minutes, iodomethane (1.66 mL, 26.7 mmol) was added. The flask was stoppered, and the mixture stirred at 40 °C for 2 h. The mixture was concentrated under reduced pressure and the residue partitioned between DCM (50 mL) and brine (50 mL). The organic phase was passed through silica and concentrated to give methyl 4-cyano-2-fluorobenzoate (4.1 g, 94%).

¹H NMR (300 MHz, CDCl₃) δ ppm 3.96 (s, 3H) 7.44-7.48 (m, 1H) 7.49-7.53 (m, 1H) 8.01-8.05 (m, 1H)

Step (ii) Raney nickel (2.0 g) was added to a solution of methyl 4-cyano-2-fluorobenzoate (obtained as described in step (i) above) (10.1 g, 61.6 mmol) in acetic acid (200 mL) and water (100 mL). The mixture was stirred at RT under argon at 20 bar. After 18 h the mixture was filtered through diatomaceous earth, washing through with water (1000 mL). The filtrate was extracted with EtOAc (3 × 300 mL). The combined organic phases were dried and concentrated, adding toluene to assist removal of acetic acid, to give methyl 2-fluoro-4-(hydroxymethyl)benzoate (crude, 6.06 g) which was used in the next step without further purification or characterisation.

Step (iii) Methyl 2-fluoro-4-(hydroxymethyl)benzoate (step (ii) above) (1 g, 5.4 mmol) was dissolved in chloroform (40 mL) and THF (5 mL). Manganese(IV) oxide (2.36 g, 27 mmol) was added and the mixture stirred at RT for 2 h and then at 60 °C for 1 h. Further manganese(IV) oxide (2.36 g, 27 mmol) was added and heating was continued at 60 °C. After 18 h the mixture was filtered through diatomaceous earth to give methyl 2-fluoro-4-formylbenzoate which was used without further purification.

¹H NMR (300 MHz, CDCl₃) δ ppm 3.95 (s, 3H) 7.58-7.62 (m, 1H) 7.63-7.67 (m, 1H) 8.08-8.12 (m, 1H) 10.04 (s, 1H)


Step (iv) Methyl 2-fluoro-4-(((3-methoxybenzyl)amino)methyl)benzoate was prepared using the reductive amination procedure described in Method R3 above from (3-methoxyphenyl)methanamine and methyl 2-fluoro-4-formylbenzoate (step (iii) above).

MS ES⁺: 304

Step (v) Oxalyl chloride (0.30 mL, 3.51 mmol) was added to a solution of 4-(2-fluorophenoxy)benzoic acid (Intermediate A1) (746 mg, 2.34 mmol) in DCM (10 mL) at RT. DMF (1 drop) was added, and the mixture stirred, allowing to warm to RT. After 2 h, the mixture was concentrated. The residue was dissolved in DCM (5 mL) and added to a solution of methyl 2-fluoro-4-(((3-methoxybenzyl)amino)methyl)benzoate (step (iv))

above) (708 mg, 23.4 mmol) and triethylamine (0.49 mL, 3.51 mmol). After 2 h the mixture was partitioned between water (30 mL) and DCM (30 mL). The organic phase was washed sequentially with saturated sodium carbonate solution, citric acid (20% in water) and water, and then dried and concentrated. The residue was purified by silica chromatography (10-30% EtOAc/heptane) to give the title compound which was used in the next step without further purification or characterisation.


Intermediate C65: Methyl 4-((N-(cyclopropylmethyl)-2-fluoro-4-(2-methoxyphenoxy)benzamido)methyl)benzoate



Prepared using the HATU coupling procedure described in Method E4 above, from 2-fluoro-4-(2-methoxyphenoxy)benzoic acid (Intermediate A8) and methyl 4-(((cyclopropylmethyl)amino)methyl)benzoate hydrochloride (Intermediate B5). In this case, the acid, amine hydrochloride and DIPEA were combined in DMF first, and HATU added last.

MS ES⁺: 464

Intermediate C65a: Methyl 4-((4-bromo-N-(cyclopropylmethyl)benzamido)methyl)benzoate




Prepared using a coupling procedure analogous to that described in Method E5 above, from 4-bromobenzoyl chloride and methyl 4-(((cyclopropylmethyl)amino)methyl)benzoate hydrochloride (Intermediate B5).

MS ES⁺: 402, 404

5

Intermediate C65b: Methyl 4-((4-bromo-N-propylbenzamido)methyl)benzoate




Prepared using a coupling procedure analogous to that described in Method E5 above, from 4-bromobenzoyl chloride and methyl 4-((propylamino)methyl)benzoate hydrochloride (Intermediate B19a).

10

MS ES⁺: 390, 392

Intermediate C66: Methyl 4-((N-(cyclopropylmethyl)-4-(2-fluoro-6-methoxyphenoxy)benzamido)methyl)benzoate




15

A mixture of methyl 4-((4-bromo-N-(cyclopropylmethyl)benzamido)methyl)benzoate (Intermediate C65a) (300 mg (0.75 mmol), 2-fluoro-6-methoxyphenol (212 mg, 1.49 mmol) and copper(I) oxide (107 mg, 0.75 mmol) in 2,4,6-trimethylpyridine (3 mL) was heated under microwave irradiation at 220 °C for 2 h. Further 2-fluoro-6-methoxyphenol (212 mg, 1.49 mmol) was added, and the mixture heated under microwave irradiation at 220 °C for another 1 h. The mixture was diluted with water and EtOAc (20 mL each) and filtered through diatomaceous earth. The phases were separated. The aqueous phase was acidified to pH ~3 by addition of HCl (2 M) and extracted with EtOAc (3 × 50 mL). The

20

combined organic phases were dried and concentrated. The residue was purified by silica chromatography (0-60% EtOAc/heptane) to give methyl 4-((*N*-(cyclopropylmethyl)-4-(2-fluoro-6-methoxyphenoxy)benzamido)methyl)benzoate (90 mg, 26%) as a colourless gum.
¹H NMR (400 MHz, CDCl₃) δ ppm -0.12-0.31 (m, 2H) 0.38-0.61 (m, 2H) 0.77-1.15 (m, 1H) 2.99-3.50 (m, 2H) 3.90 (s, 3H), 3.79 (s, 3H) 4.59-5.09 (m, 2H) 6.68-7.58 (m, 9H) 7.93-8.10 (m, 2H)


Intermediate C67: Methyl 4-((*N*-(cyclopropylmethyl)-4-(4-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoate



Prepared as described above for Intermediate C66 from methyl 4-((4-bromo-*N*-(cyclopropylmethyl)benzamido)methyl)benzoate (Intermediate C65a) and 4-fluoro-2-methoxyphenol, except that the mixture was heated under microwave irradiation at 220 °C for 1 h, no additional phenol was required, and slight modifications were made to the workup and purification. In this case the crude reaction mixture was poured into HCl (2 M, 30 mL) and the resulting mixture extracted with EtOAc (3 × 50 mL). The combined organic phases were washed with a mixture of NaOH (1 M) and brine (ca. 1:1, total 30 mL), dried and concentrated, and the residue was purified by silica chromatography (20-40% EtOAc/heptane).

¹H NMR (400 MHz, CDCl₃) δ ppm -0.15-0.30 (m, 2H) 0.40-0.59 (m, 2H) 0.78-1.01 (m, 1H) 2.98-3.45 (m, 2H) 3.78 (s, 3H) 3.91 (s, 3H) 4.64-5.05 (m, 2H) 6.58-7.50 (m, 9H) 7.96-8.10 (m, 2H)


Intermediate C67a: Methyl 4-((4-(4-fluoro-2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoate



Prepared as described above for Intermediate C66, from methyl 4-((4-bromo-*N*-propylbenzamido)methyl)benzoate (Intermediate C65b) and 4-fluoro-2-methoxyphenol, except that the mixture was heated thermally at reflux in air for 18 h, the combined organic phases were washed with HCl (1 M), saturated sodium hydrogencarbonate solution and brine (100 mL each), and the residue was purified by silica chromatography (10-20% EtOAc/heptane).

MS ES⁺: 452


Intermediate C67b: Methyl 4-((*N*-(cyclopropylmethyl)-4-(5-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoate



Prepared as described above for Intermediate C66, from methyl 4-((4-bromo-*N*-(cyclopropylmethyl)benzamido)methyl)benzoate (Intermediate C65a) and 5-fluoro-2-methoxyphenol, except that the combined organic phases were washed with HCl (1 M), NaOH (2 M) and brine (25 mL each).


¹H NMR (400 MHz, CDCl₃) δ ppm -0.10-0.30 (m, 2H) 0.34-0.62 (m, 2H) 0.75-1.20 (m, 1H) 3.00-3.60 (m, 2H) 3.78 (s, 3H) 3.91 (s, 3H) 4.70-5.00 (m, 2H) 6.70-7.01 (m, 5H) 7.20-7.50 (m, 4H) 7.96-8.06 (m, 2H)

Intermediate C67c: Methyl 4-((4-(5-fluoro-2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoate



Prepared as described above for Intermediate C66, from methyl 4-((4-bromo-*N*-propylbenzamido)methyl)benzoate (Intermediate C65b) and 5-fluoro-2-methoxyphenol, except that the mixture was heated under microwave irradiation at 210 °C for 4 h, the combined organic phases were washed with HCl (1 M), saturated sodium hydrogencarbonate solution and brine (50 mL each), the residue was purified by silica chromatography (20-50% EtOAc/heptane), and the material obtained was used in the next step without further purification or characterisation.


Intermediate C67d: Methyl 4-((*N*-(cyclopropylmethyl)-4-(3-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoate



Prepared as described above for Intermediate C66, from methyl 4-((4-bromo-*N*-(cyclopropylmethyl)benzamido)methyl)benzoate (Intermediate C65a) and 3-fluoro-2-methoxyphenol (prepared as described in Synthetic Communications, 1985, 15(1), 61-69), except that the mixture was heated under microwave irradiation at 210 °C for 5 h, the combined organic phases were washed with HCl (1 M), saturated sodium hydrogencarbonate solution and brine (100 mL each), and the residue was purified by preparative HPLC.

MS ES⁺: 464


Intermediate C67e: Methyl 4-((5-bromo-N-(cyclopropylmethyl)picolinamido)methyl)benzoate



Prepared using the coupling procedure described in Method E4 above, from 5-bromopicolinic acid and methyl 4-(((cyclopropylmethyl)amino)methyl)benzoate hydrochloride (Intermediate B5), using 1.3 eq HATU and 6 eq DIPEA.

MS ES⁺: 403, 405


Intermediate C68: Methyl 4-((N-(cyclopropylmethyl)-5-(2-methoxyphenoxy)picolinamido)methyl)benzoate



Prepared as described above for Intermediate C66, from methyl 4-((5-bromo-N-(cyclopropylmethyl)picolinamido)methyl)benzoate (Intermediate C67e) and 2-methoxyphenol, except that 5 eq phenol and 2.5 eq copper(I) oxide were used, the mixture was heated thermally at reflux in air for 40 h, the aqueous phase was adjusted to pH ~7, and the residue was purified by silica chromatography (20-40% EtOAc/heptane).

¹H NMR (400 MHz, CDCl₃) δ ppm -0.10-0.27 (m, 2H) 0.32-0.57 (m, 2H) 0.85-1.35 (m, 1H) 3.27-3.41 (m, 2H) 3.69-3.85 (m, 3H) 3.89 (s, 3H) 4.89-5.05 (m, 2H) 6.87-7.47 (m, 8H) 7.62-7.71 (m, 1H) 7.90-8.06 (m, 2H)


Intermediate C69: Methyl 4-((N-(cyclopropylmethyl)-5-(2-fluorophenoxy)picolinamido)methyl)benzoate



Prepared as described above for Intermediate C68, from methyl 4-((5-bromo-N-(cyclopropylmethyl)picolinamido)methyl)benzoate (Intermediate C67e) and 2-fluorophenol, except that 4 eq phenol was used, the mixture was not filtered through diatomaceous earth, and the residue was purified by silica chromatography (40% EtOAc/heptane).

¹H NMR (400 MHz, CDCl₃) δ ppm -0.50-0.22 (m, 2H) 0.37-0.52 (m, 2H) 0.95-1.10 (m, 1H) 3.28-3.37 (m, 2H) 3.89 (s, 3H) 4.96 (s, 2H) 7.05-7.40 (m, 7H) 7.64-7.74 (m, 1H) 7.91-8.03 (m, 2H), 8.19-8.34 (m, 1H)

Intermediate C70: 5-(2-Fluorophenoxy)pyrimidine-2-carboxylic acid




Step (i) Sodium hydride (60% dispersion in mineral oil, 0.258 g, 6.42 mmol) was added to a solution of 2-fluorophenol (0.492 mL, 5.34 mmol) in pyridine (15 mL) in an ice-water bath. Copper(I) bromide was added, and the mixture heated at 100 °C for 5 h. The mixture was concentrated and the residue triturated with EtOAc and purified by silica chromatography (20% EtOAc/heptane) to give 5-(2-fluorophenoxy)pyrimidine-2-carbonitrile (555 mg, 47%).

¹H NMR (400 MHz, CDCl₃) δ ppm 7.20-7.40 (m, 4H) 8.46 (s, 2H)

Step (ii) 5-(2-Fluorophenoxy)pyrimidine-2-carbonitrile, NaOH (2 M, 10 mL) and IMS (2 mL) were combined and heated at 70 °C for 18 h. After concentration to remove IMS, HCl

(2 M) was added until the pH was 3, and the mixture was extracted with DCM (50 mL). The organic phase was dried and concentrated to give 5-(2-fluorophenoxy)pyrimidine-2-carboxylic acid (250 mg, 48%) as a white solid, which was used in the next step without further purification or characterisation.

Intermediate C71: 5-(2-Methoxyphenoxy)pyrimidine-2-carboxylic acid




Step (i) Prepared as for step (i) of Intermediate C70 to give 5-(2-methoxyphenoxy)pyrimidine-2-carbonitrile (555 mg, 47%).

MS ES⁺: 228

Step (ii) Prepared as for step (ii) of Intermediate C70, from 5-(2-methoxyphenoxy)pyrimidine-2-carbonitrile, to give 5-(2-methoxyphenoxy)pyrimidine-2-carboxylic acid.

¹H NMR (400 MHz, CDCl₃) δ ppm 3.77 (s, 3H) 7.03-7.07 (m, 2H) 7.15-7.19 (m, 1H) 7.28-7.34 (m, 1H) 8.44 (s, 2H)


Intermediate C72: Methyl 4-((N-(cyclopropylmethyl)-5-(2-fluorophenoxy)pyrimidine-2-carboxamido)methyl)benzoate



Prepared using the coupling procedure described in Method E4 above, from 5-(2-fluorophenoxy)pyrimidine-2-carboxylic acid (Intermediate C70) and methyl 4-(((cyclopropylmethyl)amino)methyl)benzoate hydrochloride (Intermediate B5).

MS ES⁺: 436

Intermediate C73: Methyl 4-((*N*-(cyclopropylmethyl)-5-(2-methoxyphenoxy)pyrimidine-2-carboxamido)methyl)benzoate




5

Prepared using the coupling procedure described in Method E4 above, from 5-(2-methoxyphenoxy)pyrimidine-2-carboxylic acid (Intermediate C71) and methyl 4-(((cyclopropylmethyl)amino)methyl)benzoate hydrochloride (Intermediate B5).

MS ES⁺: 448

10

Intermediate C74: Methyl 4-((5-(2-methoxyphenoxy)-*N*-propylpyrimidine-2-carboxamido)methyl)benzoate



15

Prepared using the coupling procedure described in Method E4 above, from 5-(2-methoxyphenoxy)pyrimidine-2-carboxylic acid (Intermediate C71) and methyl 4-((propylamino)methyl)benzoate hydrochloride (Intermediate B19a).

MS ES⁺: 436

20

Preparation of Examples

Compounds derived from the Intermediates C6 to C74 above were prepared according to Method H1 or H2:


Method H1

Lithium hydroxide or lithium hydroxide monohydrate (5 mmol) was added to a solution of the relevant ester (1 mmol) in water (2 mL) and either THF or 1,4-dioxane (4 mL). In some cases, MeOH was added in addition to THF. The mixture was stirred, typically at RT for 18 h, 50 °C for 4 h or 100 °C for 20 minutes. In some cases the reaction was diluted with NaOH (2 M) and extracted with EtOAc or DCM, and the resulting aqueous phase acidified with HCl (2 M) and extracted with EtOAc. In other cases the crude mixture was partitioned between HCl (2 M) and EtOAc. The combined organic phases from the extraction of the acidic aqueous phase were dried and concentrated. In some cases the residue was loaded onto an anion exchange cartridge. After washing with MeCN, the product was eluted with 1 M HCl/MeCN then concentrated. In some cases the material obtained was purified by reverse-phase chromatography on C18 silica (typically eluted with 5-95% MeOH/H₂O with 0.1% NH₄OH). In some cases the crude product was heated under reflux in HCl (4 M), allowed to cool and filtered. In some cases the final product was recrystallised, typically using one or more of the following: water, EtOH, EtOAc, methyl acetate, methyl *tert*-butyl ether, pentane, heptane.

Method H2

Sodium hydroxide (2 M, 6 mmol) was added to a solution of the relevant ester (2 mmol) in THF (3 mL). In some cases, MeOH was added instead of or in addition to THF. The mixture was stirred at RT for 18 h or heated under microwave irradiation at 100 °C for 5 minutes. The mixture was partitioned between HCl (1 M) and EtOAc (30 mL each). The organic phases were dried and concentrated. The residue was loaded onto an anion exchange cartridge. After washing with MeCN, the product was eluted with 4 M HCl/1,4-dioxane then concentrated.

Example 1: **4-((*N*-Ethyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid**




Prepared from the parent methyl ester (Intermediate C6) using Method H1.

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ ppm 0.84-1.36 (m, 3H) 3.05-3.64 (m, 2H) 4.67 (br s, 2H) 6.73-7.09 (m, 2H) 7.09-7.61 (m, 8H) 7.74-8.05 (m, 2H) 12.90 (br s, 1H)

5 MS ES^+ : 394

Example 2: 4-((*N*-(2,2-Difluoroethyl)-4-(2-fluorophenoxy)benzamido)-methyl)benzoic acid




10 Prepared from the parent methyl ester (Intermediate C7) using Method H1.

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ ppm 3.57-3.91 (m, 2H) 4.71 (br s, 2H) 5.95-6.52 (m, 1H) 6.82-7.08 (m, 2H) 7.10-7.57 (m, 8H) 7.79-7.98 (m, 2H) 12.93 (br s, 1H)

MS ES^+ : 430


15 **Example 3:** 4-((4-(2-Fluorophenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)-methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C8) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 4.14-4.36 (m, 2H) 4.76 (s, 2H) 6.92-7.10 (m, 2H)
7.16-7.37 (m, 5H) 7.39-7.55 (m, 3H) 7.84-7.99 (m, 2H) 12.94 (br s, 1H)
MS ES⁺: 448


Example 4: 4-((4-(2-Fluorophenoxy)-*N*-isobutylbenzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C9) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 0.50-1.08 (m, 6H) 1.77-2.17 (m, 1H) 3.00-3.25 (m, 2H) 4.49-4.86 (m, 2H) 6.88-7.13 (m, 2H) 7.13-7.57 (m, 8H) 7.83-8.04 (m, 2H) 12.69-13.02 (m, 1H)
MS ES⁺: 422


Example 5: 4-((*N*-(Cyclopropylmethyl)-4-(2-fluorophenoxy)benzamido)-methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C10) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm -0.13-0.26 (m, 2H) 0.30-0.50 (m, 2H) 0.75-1.06 (m, 1H) 2.97-3.22 (m, 2H) 4.79 (br s, 2H) 6.88-7.12 (m, 2H) 7.12-7.55 (m, 8H) 7.86-7.97 (m, 2H) 12.87 (br s, 1H)
MS ES⁺: 420

Example 6: 4-((N-(Cyclobutylmethyl)-4-(2-fluorophenoxy)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C11) using Method H1.

¹H NMR (400 MHz, CD₃OD) δ ppm 1.37-2.15 (m, 6H) 2.49-2.82 (m, 1H) 3.34-3.58 (m, 2H) 4.60-4.78 (m, 2H) 6.90-7.04 (m, 2H) 7.14-7.29 (m, 5H) 7.33-7.49 (m, 3H) 7.82-8.03 (m, 2H)

MS ES⁺: 434

Example 7: 4-((4-(2-Fluorophenoxy)-N-((3-methoxycyclobutyl)methyl)-benzamido)methyl)benzoic acid (mixture of *cis* and *trans* ring isomers)




Prepared from the parent methyl ester (Intermediate C58) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.64-2.30 (m, 4H) 3.04 (br s, 3H) 3.16-3.70 (m, 4H) 4.65 (br s, 2H) 6.85-7.14 (m, 2H) 7.14-7.59 (m, 8H) 7.82-8.00 (m, 2H)

MS ES⁺:464

Example 8: 4-((4-(2-Fluorophenoxy)-N-isopentylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C12) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 0.48-1.00 (m, 6H) 1.39 (br s, 3H) 3.21 (br s, 2H) 4.49-4.84 (m, 2H) 7.01 (br s, 2H) 7.17-7.57 (m, 8H) 7.88-7.97 (m, 2H) 12.90 (br s, 1H)

MS ES^+ : 436

Example 9: 4-((N-(2-Cyclopropylethyl)-4-(2-fluorophenoxy)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C13) using Method H1.

^1H NMR (400 MHz, CD_3OD) δ ppm -0.22-0.17 (m, 2H) 0.26-0.51 (m, 2H) 0.52-0.79 (m, 1H) 1.35-1.64 (m, 2H) 3.35-3.63 (m, 2H) 4.58-4.79 (m, 2H) 6.90-7.08 (m, 2H) 7.13-7.34 (m, 5H) 7.37-7.54 (m, 3H) 8.06-8.22 (m, 2H)

MS ES^+ : 434

Example 10: 4-((N-(2-Cyclobutylethyl)-4-(2-fluorophenoxy)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C14) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 1.24-2.30 (m, 9H) 2.88-3.40 (m, 2H) 4.48-4.82 (m, 2H) 7.01 (br s, 2H) 7.19-7.54 (m, 8H) 7.88-7.96 (m, 2H) 12.90 (br s, 1H)

MS ES^+ : 448

5 **Example 11:** 4-((*N*-Benzyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C15) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 4.37-4.80 (m, 4H) 6.92-7.06 (m, 2H) 7.11-7.60 (m, 13H) 7.85-7.99 (m, 2H)

10 MS ES^+ : 456

Example 12: 4-((*N*-(2-Fluorobenzyl)-4-(2-fluorophenoxy)benzamido)-methyl)benzoic acid




15 Prepared from the parent methyl ester (Intermediate C16) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 4.45-4.66 (m, 4H) 6.96-7.03 (m, 2H) 7.03-7.45 (m, 10H) 7.45-7.54 (m, 2H) 7.78-7.89 (m, 2H)

MS ES^+ : 474

20 **Example 13:** 4-((4-(2-Fluorophenoxy)-*N*-(3-methoxybenzyl)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C17) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 3.72 (s, 3H) 4.42-4.71 (m, 4H) 6.55-7.10 (m, 5H)
7.16-7.62 (m, 9H) 7.76-7.99 (m, 2H) 12.92 (br s, 1H)

5 MS ES $^+$: 486

Example 14: 4-((4-(2-Fluorophenoxy)-*N*-(4-methoxybenzyl)benzamido)-methyl)benzoic acid




10 Prepared from the parent methyl ester (Intermediate C18) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 3.74 (s, 3H) 4.24-4.80 (m, 4H) 6.83-7.65 (m, 14H)
7.84-8.05 (m, 2H) 12.90 (br s, 1H)

MS ES $^+$: 486

15 **Example 15:** 4-((4-(2-Fluorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C19) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 2.40-3.07 (m, 2H) 3.07-4.11 (m, 2H) 4.11-4.95 (m, 2H) 6.55-7.49 (m, 15H) 7.86-8.05 (m, 2H)

MS ES^+ : 470

Example 16: 4-((4-(2-Fluorophenoxy)-N-(3-(2-fluorophenyl)propyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C20) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 1.66-1.98 (m, 2H) 2.17-2.80 (m, 2H) 3.05-3.52 (m, 2H) 4.44-4.90 (m, 2H) 6.83-7.55 (m, 14H) 7.85-8.01 (m, 2H) 12.90 (br s, 1H)

MS ES^+ : 502

Example 17: 4-((4-(2-Fluorophenoxy)-N-(3-(3-fluorophenyl)propyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C21) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) ppm 1.70-1.95 (m, 2H) 2.30-2.40 (m, 2H) 3.30-3.40 (m, 2H) 4.74 (br s, 2H) 6.80-7.56 (m, 14H) 7.87-7.99 (m, 2H) 12.88 (br s, 1H)

MS ES⁺: 502

Example 18: 4-((4-(2-Fluorophenoxy)-N-((*trans*-2-phenylcyclopropyl)methyl)-benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C22) using Method H1.

¹H NMR (400 MHz, CD₂Cl₂) δ ppm 0.92 (br s, 2H) 1.20-1.35 (m, 1H) 1.52-1.78 (m, 1H) 3.42 (s, 2H) 4.75-4.99 (m, 2H) 6.86-7.10 (m, 4H) 7.10-7.29 (m, 6H) 7.29-7.50 (m, 5H) 8.00-8.12 (m, 2H)

MS ES⁺: 496

Example 19: (*S*)-4-((4-(2-Fluorophenoxy)-N-(2-hydroxy-3-phenylpropyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C23) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 2.60-2.76 (m, 1H) 2.95-3.30 (m, 2H) 3.88-4.17 (m, 1H) 4.65 (br s, 2H) 4.85-4.97 (m, 1H) 5.00-5.29 (m, 1H) 6.85-7.33 (m, 12H) 7.33-7.50 (m, 3H) 7.79-7.90 (m, 2H)

MS ES⁺: 500

Example 20: (*R*)-4-((4-(2-Fluorophenoxy)-N-(2-hydroxy-3-phenylpropyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C24) using Method H1.

^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ ppm 2.55-2.79 (m, 1H) 2.92-3.40 (m, 2H) 3.83-4.17 (m, 1H) 4.72 (br s, 2H) 4.85-5.05 (m, 1H) 5.06-5.23 (m, 1H) 6.83-7.53 (m, 15H) 7.84-7.97 (m, 2H)

MS ES^+ : 500

Example 21: 4-((4-(2-Fluorophenoxy)-*N*-(4-phenylbutyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C25) using Method H1.

^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ ppm 1.29-1.70 (m, 2H) 2.32-2.84 (m, 2H) 3.10-3.55 (m, 2H) 4.62 (br s, 2H) 6.86-7.46 (m, 15H) 7.77-7.95 (m, 2H) 12.57 (br s, 1H)

MS ES^+ : 498

Example 22: 4-((*N*-(Cyclopropylmethyl)-4-(*o*-toloxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C26) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) ppm -0.25 - 0.25 (m, 2H) 0.25 - 0.50 (m, 2H) 0.74 - 1.11 (m, 1H) 2.13 (s, 3H) 2.87 - 3.47 (m, 2H) 4.78 (br s, 2H) 6.75 - 7.62 (m, 10H) 7.83 - 7.98 (m, 2H) 12.87 (br s, 1H)
MS ES^+ : 416

5


Example 23: 4-((*N*-(Cyclobutylmethyl)-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C27) using Method H1.

10 ^1H NMR (400 MHz, $\text{DMSO}-d_6$) ppm 1.23 - 1.99 (m, 7H) 2.15 (s, 3H) 2.40 - 2.74 (m, 2H) 4.65 (br s, 2H) 6.76 - 7.64 (m, 10H) 7.87 - 8.02 (m, 2H) 12.90 (br s, 1H)
MS ES^+ : 430

Example 24: 4-((*N*-Benzyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid




15

Prepared from the parent methyl ester (Intermediate C28) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 2.14 (s, 3H) 4.58 (br s, 4H) 6.62 - 7.04 (m, 3H) 7.07 - 7.60 (m, 12H) 7.84 - 8.00 (m, 2H) 12.90 (br s, 1H)
MS ES^+ : 452

20

Example 25: 4-((*N*-Phenethyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C29) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 2.15 (br s, 3H) 2.72-2.98 (m, 2H) 3.34-3.59 (m, 2H) 4.43-4.88 (m, 2H) 6.82-7.02 (m, 4H) 7.28 (br s, 11H) 7.88-7.96 (m, 2H)

MS ES^+ : 466

Example 26: 4-((N-Phenethyl-4-(2-(trifluoromethyl)phenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C30) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 2.71-2.99 (m, 2H) 3.34-3.59 (m, 2H) 4.43-4.87 (m, 2H) 6.89-7.42 (m, 12H) 7.43-7.54 (m, 1H) 7.64-7.73 (m, 1H) 7.77-7.85 (m, 1H) 7.88-7.98 (m, 2H)

MS ES^+ : 520

Example 27: 4-((N-Benzyl-4-(2-cyanophenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C31) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) ppm 4.40-4.76 (m, 4H) 7.00-7.77 (m, 15H) 7.85-8.00 (m, 2H) 12.97 (br s, 1H)

MS ES⁺: 463

5 **Example 28:** 4-((*N*-Ethyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C32) using Method H1.


¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.01-1.11 (m, 3H) 3.22-3.33 (m, 2H) 3.73 (s, 3H)

10 4.68 (br s, 2H) 6.79-6.88 (m, 2H) 6.96-7.03 (m, 1H) 7.07-7.13 (m, 1H) 7.16-7.28 (m, 2H)

7.40 (br s, 4H) 7.89-7.96 (m, 2H) 12.88 (br s, 1H)

MS ES⁺: 406

15 **Example 29:** 4-((*N*-(2,2-Difluoroethyl)-4-(2-methoxyphenoxy)benzamido)-methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C33) using Method H1.


¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 3.64-3.86 (m, 5H) 4.74 (br s, 2H) 6.07-6.44 (m,

1H) 6.78-6.92 (m, 2H) 6.96-7.03 (m, 1H) 7.07-7.13 (m, 1H) 7.15-7.46 (m, 6H) 7.87-7.97

20 (m, 2H) 12.93 (br s, 1H)

MS ES⁺: 442

Example 30: 4-((*N*-Isobutyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C34) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.58-0.97 (m, 6H) 1.80-2.13 (m, 1H) 3.06-3.19 (m, 2H) 3.72 (s, 3H) 4.68 (s, 2H) 6.77-6.90 (m, 2H) 6.94-7.05 (m, 1H) 7.05-7.13 (m, 1H) 7.16-7.53 (m, 6H) 7.87-7.97 (m, 2H) 12.91 (br s, 1H)

MS ES⁺: 434

Example 31: 4-((*N*-(Cyclopropylmethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C35) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) δ -0.14-0.20 (m, 2H) 0.30-0.44 (m, 2H) 0.81-1.03 (m, 1H) 3.15 (br s, 2H) 3.70 (s, 3H) 4.79 (br s, 2H) 6.78-6.91 (m, 2H) 6.96-7.03 (m, 1H) 7.08-7.14 (m, 1H) 7.16-7.28 (m, 2H) 7.30-7.50 (m, 4H) 7.87-7.96 (m, 2H) 12.88 (s, 1H)

MS ES⁺: 432

Example 32: 4-((4-(2-Methoxyphenoxy)-*N*-(3-phenylpropyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C36) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.83 (br s, 2H) 2.40 (br s, 2H) 3.22 (br s, 2H) 3.73 (s, 3H) 4.70 (br s, 2H) 6.77-6.82 (m, 2H) 6.95-7.44 (m, 13H) 7.88-7.93 (m, 2H) 12.89 (s, 1H)

MS ES⁺: 496

Example 33: 4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)-benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C37) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.84 (br s, 2H) 2.42 (br s, 2H) 3.21 (br s, 2H) 3.73 (s, 3H) 4.70 (br s, 2H) 6.76-6.82 (m, 2H) 6.83-7.47 (m, 12H) 7.86-7.94 (m, 2H) 12.88 (br s, 1H)

MS ES⁺: 514

Example 34: 4-((*N*-(Cyclopropylmethyl)-4-(2-ethoxyphenoxy)-benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C38) using Method H1.

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ ppm -0.24 - 0.60 (m, 4H) 0.69 - 1.41 (m, 4H) 2.87 - 3.31 (m, 2H) 3.80 - 4.15 (m, 2H) 4.76 (br s, 2H) 6.60 - 7.57 (m, 10H) 7.75 - 8.00 (m, 2H) 12.86 (br s, 1H)

MS ES^+ : 446

Example 35: 4-((4-(2-Chlorophenoxy)-*N*-(cyclopropylmethyl)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C39) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) ppm ppm -0.30 - 0.23 (m, 2H) 0.23 - 0.50 (m, 2H) 0.74 - 1.12 (m, 1H) 2.81 - 3.41 (m, 2H) 4.79 (br s, 2H) 6.86 - 7.04 (m, 2H) 7.12 - 7.66 (m, 8H) 7.84 - 7.97 (m, 2H) 12.87 (br s, 1H)

MS ES^+ : 436

Example 36: 4-((4-(2-Chlorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C40) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 2.72 - 2.95 (m, 2H) 3.35 - 3.60 (m, 2H) 4.41 - 4.88 (m, 2H) 6.88 - 7.01 (m, 3H) 7.14 - 7.54 (m, 11H) 7.58 - 7.66 (m, 1H) 7.87 - 7.98 (m, 2H) 12.89 (br s, 1H)

MS ES^+ : 486

Example 37: 4-((N-(Cyclopropylmethyl)-4-(2,6-difluorophenoxy)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C41) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 0.00 (br s, 2H) 0.38 (br s, 2H) 0.92 (br s, 1H) 3.12 (br s, 2H) 4.80 (br s, 2H) 7.01 (br s, 2H) 7.24-7.56 (m, 7H) 7.84-7.99 (m, 2H) 12.88 (br s, 1H)

MS ES $^+$: 438

Example 38: 4-((N-(2-Cyclopropylethyl)-4-(2,6-difluorophenoxy)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C42) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm -0.27-0.10 (m, 2H) 0.18-0.75 (m, 3H) 1.25-1.57 (m, 2H) 3.14-3.47 (m, 2H) 4.52-4.81 (m, 2H) 6.87-7.11 (m, 2H) 7.22-7.56 (m, 7H) 7.84-8.00 (m, 2H) 12.88 (br s, 1H)

MS ES $^+$: 452

Example 39: 4-((N-Benzyl-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C43) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 4.40-4.68 (m, 4H) 6.95-7.04 (m, 2H) 7.11-7.44 (m, 10H) 7.45-7.55 (m, 2H) 7.86-7.94 (m, 2H)

MS ES^+ : 474

Example 40: 4-((N-(Cyclopropylmethyl)-4-(2-fluoro-6-methylphenoxy)-benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C44) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm -0.16-0.24 (m, 2H) 0.26-0.48 (m, 2H) 0.77-1.08 (m, 1H) 2.15 (s, 3H) 2.94-3.32 (m, 2H) 4.78 (br s, 2H) 6.88 (br s, 2H) 7.09-7.28 (m, 3H) 7.42 (br s, 4H) 7.85-7.96 (m, 2H) 12.87 (br s, 1H)

MS ES^+ : 434

Example 41: 4-((N-(Cyclobutylmethyl)-4-(2-fluoro-6-methylphenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C45) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 1.36-2.01 (m, 6H) 2.16 (s, 3H) 3.22-3.42 (m, 3H)
4.64 (br s, 2H) 6.80-6.95 (m, 2H) 7.17-7.29 (m, 3H) 7.32-7.56 (m, 4H) 7.81-8.02 (m, 2H)

12.88 (br s, 1H)

MS ES $^+$: 448

Example 42: 4-((4-(2-Chloro-6-fluorophenoxy)-N-(cyclopropylmethyl)-
benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C46) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 0.00 (br s, 2H) 0.38 (br s, 2H) 0.93 (br s, 1H) 3.13
(br s, 2H) 4.80 (br s, 2H) 6.95 (br s, 2H) 7.24-7.60 (m, 7H) 7.86-7.99 (m, 2H) 12.88 (br s,
1H)

MS ES $^+$: 454

Example 43: 4-((4-(2-Chloro-6-fluorophenoxy)-N-(2-cyclopropylethyl)-
benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C47) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm -0.29 - 0.77 (m, 5H) 1.23 - 1.57 (m, 2H) 3.15 - 3.52 (m, 2H) 4.49 - 4.84 (m, 2H) 6.95 (m, 2H) 7.21 - 7.57 (m, 7H) 7.89 - 7.96 (m, 2H) 12.89 (br s, 1H)

MS ES^+ : 468

Example 44: 4-((*N*-Benzyl-4-(2-chloro-6-fluorophenoxy)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C48) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 4.39 - 4.70 (m, 4H) 6.87 - 6.99 (m, 2H) 7.08 - 7.59 (m, 12H) 7.86 - 7.95 (m, 2H) 12.97 (br s, 1H)

MS ES^+ : 490

Example 45: 4-((4-(2,6-Dimethylphenoxy)-*N*-isopentylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C49) using Method H2.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 0.39 - 1.61 (m, 9H) 2.06 (s, 6H) 2.75 - 3.48 (m, 2H) 4.47 - 4.86 (m, 2H) 6.62 - 6.89 (m, 2H) 7.05 - 7.25 (m, 3H) 7.25 - 7.58 (m, 4H) 7.85 - 8.02 (m, 2H) 12.89 (br s, 1H)

MS ES^+ : 446

Example 46: 4-((*N*-Benzyl-4-(2,6-dimethylphenoxy)benzamido)methyl)benzoic acid




5 Prepared from the parent methyl ester (Intermediate C50) using Method H2.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 2.04 (s, 6H) 4.35-4.79 (m, 4H) 6.68-6.87 (m, 2H)

7.02-7.58 (m, 12H) 7.82-8.02 (m, 2H) 12.89 (br s, 1H)

MS ES $^+$: 466

10 **Example 47:** 4-((4-(2,6-Dimethylphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C51) using Method H2.

^1H NMR (400 MHz, DMSO- d_6) δ ppm 2.07 (s, 6H) 2.62-3.04 (m, 2H) 3.10-3.70 (m, 2H)

15 4.35-4.98 (m, 2H) 6.65-7.63 (m, 14H) 7.85-8.03 (m, 2H)

MS ES $^+$: 480

Example 48: 4-((4-(3-Methoxyphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C52) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 2.72-2.95 (m, 2H) 3.35-3.61 (m, 2H) 3.74 (s, 3H)
4.40-4.88 (m, 2H) 6.54-6.66 (m, 2H) 6.72-6.80 (m, 1H) 6.90-7.11 (m, 3H) 7.13-7.57 (m,
5 9H) 7.86-8.01 (m, 2H) 12.87 (br s, 1H)

MS ES^+ : 482

Example 49: 4-((4-(4-Methoxyphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C53) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 2.70-2.96 (m, 2H) 3.35-3.57 (m, 2H) 3.76 (s, 3H)
4.40-4.87 (m, 2H) 6.85-7.56 (m, 15H) 7.85-8.00 (m, 2H) 12.90 (br s, 1H)

MS ES^+ : 482

Example 50: 4-((4-(3-Chlorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C54) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 2.73-2.98 (m, 2H) 3.36-3.62 (m, 2H) 4.42-4.90 (m, 2H) 6.87-7.57 (m, 15H) 7.93 (br s, 2H) 12.90 (br s, 1H)

MS ES^+ : 486

Example 51: 4-((N-(Cyclopropylmethyl)-4-((2-fluorobenzyl)oxy)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C55) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) ppm ppm -0.22-0.25 (m, 2H) 0.26-0.47 (m, 2H) 0.77-1.08 (m, 1H) 2.73-3.45 (m, 2H) 4.77 (br s, 2H) 5.15 (s, 2H) 6.98-7.63 (m, 10H) 7.85-7.97 (m, 2H) 12.84 (br s, 1H)

MS ES^+ : 434

Example 52: 4-((N-(Cyclobutylmethyl)-4-((2-fluorobenzyl)oxy)benzamido)-methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C56) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) ppm 1.20-2.20 (m, 7H) 2.26-2.74 (m, 2H) 4.64 (s, 2H) 5.18 (s, 2H) 6.97-7.66 (m, 10H) 7.83-8.07 (m, 2H) 12.81 (br s, 1H)

MS ES^+ : 448

Example 53: 4-((*N*-Benzyl-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C57) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) ppm 4.56 (br s, 4H) 5.16 (s, 2H) 7.00-7.61 (m, 15H) 7.87-7.99 (m, 2H) 12.92 (br s, 1H)

MS ES⁺: 470

Example 54: 4-((*N*-(Cyclopropylmethyl)-2-fluoro-4-(2-fluorophenoxy)-benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C59) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm -0.06-0.28 (m, 2H) 0.31-0.52 (m, 2H) 0.73-1.15 (m, 1H) 2.99-3.41 (m, 2H) 4.55-4.97 (m, 2H) 6.73-7.07 (m, 2H) 7.25-7.56 (m, 7H) 7.86-8.02 (m, 2H) 12.91 (br s, 1H)

MS ES⁺: 438

Example 55: 4-((*N*-Benzyl-4-(2-fluorophenoxy)cyclohexanecarboxamido)-methyl)benzoic acid (ca. 1:1 mixture of *cis* and *trans* ring isomers)




Prepared from the parent methyl ester (Intermediate C60) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) ppm 1.27-2.13 (m, 8H) 2.61-2.86 (m, 1H) 4.00-4.40 (m, 1H) 4.45-4.73 (m, 4H) 6.88-7.00 (m, 1H) 7.05-7.44 (m, 10H) 7.85-8.00 (m, 2H) 12.90 (br s, 1H)

MS ES⁺: 462

Example 56: *trans*-4-((*N*-Benzyl-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C61) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) ppm 1.10-2.11 (m, 8H) 2.57-2.78 (m, 1H) 3.73 (s, 3H) 4.11-4.25 (m, 1H) 4.46-4.74 (m, 4H) 6.81-7.04 (m, 4H) 7.15-7.44 (m, 7H) 7.86-8.00 (m, 2H) 12.90 (br s, 1H)

MS ES⁺: 474

Example 57: *cis*-4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C62) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) ppm 1.34-1.98 (m, 10H) 2.39-2.66 (m, 3H) 3.15-3.40 (m, 2H) 3.71-3.87 (m, 3H) 4.37-4.51 (m, 1H) 4.52-4.78 (m, 2H) 6.79-7.12 (m, 7H) 7.19-7.39 (m, 3H) 7.82-8.01 (m, 2H) 12.80 (br s, 1H)

MS ES⁺: 520

Example 58: *trans*-4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C63) using Method H1.

¹H NMR (400 MHz, DMSO-*d*₆) ppm 1.19-2.13 (m, 10H) 2.36-2.65 (m, 3H) 3.15-3.49 (m, 2H) 3.67-3.81 (m, 3H) 4.09-4.24 (m, 1H) 4.51-4.78 (m, 2H) 6.79-7.10 (m, 7H) 7.18-7.40 (m, 3H) 7.81-8.00 (m, 2H) 12.86 (br s, 1H)

MS ES⁺: 520

Example 59: 2-Fluoro-4-((4-(2-fluorophenoxy)-*N*-(3-methoxybenzyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C64) using Method H1.

^1H NMR (300 MHz, CDCl_3) δ ppm 3.79 (s, 3H) 4.34-4.91 (m, 4H) 6.57-7.34 (m, 12H)

7.34-7.61 (m, 2H) 7.90-8.07 (m, 1H)

5 MS ES^+ : 504

Example 60: 4-((4-(2-Fluorophenoxy)-*N*-propylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C25a) using Method H1.

10 ^1H NMR (400 MHz, CDCl_3) δ ppm 0.65-1.06 (m, 3H) 1.48-1.79 (m, 2H) 3.11-3.52 (m, 2H), 4.56-4.85 (m, 2H) 6.87-7.52 (m, 10H) 8.06-8.12 (m, 2H)

MS ES^+ : 408


Example 61: 4-((4-(2-Methoxyphenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C40a) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 3.79 (s, 3H) 3.84-4.12 (m, 2H) 4.77-4.95 (m, 2H) 6.85-7.47 (m, 10H) 8.03-8.12 (m, 2H)
MS ES^+ : 460


5 **Example 62:** 4-((4-(2-Methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C40b) using Method H1.

10 ^1H NMR (400 MHz, CDCl_3) δ ppm 0.65-1.05 (m, 3H) 1.41-1.78 (m, 2H) 3.12-3.52 (m, 2H), 3.80 (s, 3H) 4.55-4.87 (m, 2H) 6.81-7.51 (m, 10H) 8.04-8.12 (m, 2H)
MS ES^+ : 420


Example 63: 4-((*N*-(2,2-Difluoropropyl)-4-(2-methoxyphenoxy)-benzamido)methyl)benzoic acid



15 Prepared from the parent methyl ester (Intermediate C40c) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 1.32-1.84 (m, 3H) 3.45-4.02 (m, 5H) 4.65-5.06 (m, 2H) 6.79-7.50 (m, 10H) 7.99-8.15 (m, 2H)
MS ES^+ : 456

20 **Example 64:** 4-((4-(2-Methoxyphenoxy)-*N*-(3,3,3-trifluoropropyl)-benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C40d) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 2.35-2.67 (m, 2H) 3.50-3.69 (m, 5H) 4.58-4.81 (m, 2H) 6.80-7.48 (m, 10H) 8.06-8.14 (m, 2H)

MS ES^+ : 474

Example 65: 4-((*N*-Butyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C40e) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 0.67-1.01 (m, 3H) 1.01-1.74 (m, 4H) 3.11-3.56 (m, 2H) 4.56-4.90 (m, 2H) 6.80-7.58 (m, 10H) 8.03-8.14 (m, 2H)

MS ES^+ : 434

Example 66: 4-((*N*-(Cyclopropylmethyl)-2-fluoro-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C65) using Method H1.

^1H NMR (400 MHz, DMSO- d_6) δ ppm -0.06 - 0.29 (m, 2H) 0.30 - 0.53 (m, 2H) 0.78 - 1.16 (m, 1H) 2.98 - 3.45 (m, 2H) 3.77 (s, 3H) 4.57 - 4.98 (m, 2H) 6.59 - 6.87 (m, 2H) 6.98 - 7.51 (m, 7H) 7.84 - 8.00 (m, 2H) 12.90 (br s, 1H)

MS ES $^+$: 450

Example 67: 4-((N-(Cyclopropylmethyl)-4-(2-fluoro-6-methoxyphenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C66) using Method H1.

^1H NMR (400 MHz, CDCl $_3$) δ ppm -0.10 - 0.28 (m, 2H) 0.37 - 0.58 (m, 2H) 0.78 - 1.15 (m, 1H) 2.98 - 3.50 (m, 2H) 3.79 (s, 3H) 4.68 - 5.05 (m, 2H) 6.69 - 7.50 (m, 9H) 8.02 - 8.11 (m, 2H)

MS ES $^+$: 450

Example 68: 4-((N-(Cyclopropylmethyl)-4-(4-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C67) using Method H1.

^1H NMR (400 MHz, CDCl $_3$) δ ppm -0.09 - 0.29 (m, 2H) 0.44 - 0.58 (m, 2H) 0.78 - 1.17 (m, 1H) 3.01 - 3.55 (m, 2H) 3.78 (s, 3H) 4.70 - 5.05 (m, 2H) 6.70 - 7.52 (m, 9H) 8.03 - 8.12 (m, 2H)

MS ES $^+$: 450

Example 69: 4-((*N*-(Cyclopropylmethyl)-5-(2-methoxyphenoxy)-picolinamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C68) using Method H1.

¹H NMR (400 MHz, CDCl₃) δ ppm -0.08-0.68 (m, 4H) 0.85-1.33 (m, 1H) 3.16-3.56 (m, 2H) 3.61-4.04 (m, 3H) 4.78-5.19 (m, 2H) 6.80-8.55 (m, 11H)

MS ES⁺: 433

Example 70: 4-((4-(*o*-Tolyloxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C40f) using Method H1.

¹H NMR (400 MHz, CDCl₃) δ ppm 2.18 (s, 3H) 4.02 (br s, 2H) 4.85 (br s, 2H) 6.81-7.58 (m, 10H) 8.00-8.20 (m, 2H)

MS ES⁺: 444

Example 71: 4-((4-(2-Fluorophenoxy)-*N*-(3,3,3-trifluoropropyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C25b) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 2.35-2.65 (m, 2H) 3.53-3.68 (m, 2H) 4.60-4.78 (m, 2H) 6.88-7.45 (m, 10H) 8.06-8.13 (m, 2H)

MS ES^+ : 462

Example 72: 4-((4-(2-Chlorophenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C40g) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 3.81-4.18 (m, 2H) 4.84 (br s, 2H) 6.88-7.57 (m, 10H) 8.01-8.16 (m, 2H)

MS ES^+ : 464


Example 73: 4-((*N*-Ethyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C40h) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 0.92-1.12 (m, 3H) 2.11 (s, 3H) 3.08-3.40 (m, 2H)
4.46-4.78 (m, 2H) 6.74-7.50 (m, 10H) 7.82-7.93 (m, 2H)
MS ES^+ : 390


5 **Example 74:** **4-((N-(2,2-Difluoropropyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid**



Prepared from the parent methyl ester (Intermediate C25c) using Method H1.

10 ^1H NMR (400 MHz, CDCl_3) δ ppm 1.24-1.92 (m, 3H) 3.36-4.08 (m, 2H) 4.57-5.13 (m, 2H) 6.79-7.58 (m, 10H) 7.90-8.20 (m, 2H)
MS ES^+ : 444


Example 75: **4-((N-Butyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid**



15 Prepared from the parent methyl ester (Intermediate C25d) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 0.66-1.77 (m, 7H) 3.08-3.57 (m, 2H) 4.49-4.93 (m, 2H) 6.81-7.57 (m, 10H) 8.04-8.18 (m, 2H)
MS ES^+ : 422

20 **Example 76:** **4-((4-(2-Chlorophenoxy)-N-ethylbenzamido)methyl)benzoic acid**




Prepared from the parent methyl ester (Intermediate C40i) using Method H1.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 0.78-1.23 (m, 3H) 2.98-3.65 (m, 2H) 4.40-4.82 (m, 2H) 6.70-7.70 (m, 10H) 7.77-8.01 (m, 2H)

5 MS ES^+ : 410

Example 77: 4-((N-(Cyclopropylmethyl)-5-(2-fluorophenoxy)picolinamido)methyl)benzoic acid




10 Prepared from the parent methyl ester (Intermediate C69) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm -0.06-0.25 (m, 2H) 0.38-0.56 (m, 2H) 0.77-1.33 (m, 1H) 3.27-3.40 (m, 2H) 4.99 (s, 2H) 7.07-8.44 (m, 11H)

MS ES^+ : 421


15 **Example 78:** 4-((4-(2-Chlorophenoxy)-N-(2,2-difluoroethyl)benzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C40j) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 3.59-3.81 (m, 2H) 4.80 (br s, 2H) 5.05-6.40 (m, 1H)
6.81-7.53 (m, 10H) 8.04-8.17 (m, 2H)
MS ES^+ : 446


Example 79: 4-((N-(Cyclopropylmethyl)-4-(5-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C67b) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ -0.12-0.32 (m, 2H) 0.41-0.58 (m, 2H) 0.78-1.15 (m, 1H)
3.02-3.52 (m, 2H) 3.78 (s, 3H) 4.68-5.08 (m, 2H) 6.69-7.08 (m, 5H) 7.19-7.58 (m, 4H)
8.04-8.12 (m, 2H)
MS ES^+ : 450


Example 80: 4-((4-(4-Fluoro-2-methoxyphenoxy)-N-propylbenzamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C67a) using Method H1.

^1H NMR (300 MHz, CDCl_3) δ ppm 0.65-1.08 (m, 3H) 1.47-1.74 (m, 2H) 3.10-3.50 (m, 2H), 3.77 (s, 3H) 4.55-4.98 (m, 2H) 6.58-7.58 (m, 9H) 7.90-8.23 (m, 2H)
MS ES^+ : 438

Example 81: 4-((4-(5-Fluoro-2-methoxyphenoxy)-N-propylbenzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C67c) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 0.65-1.12 (m, 3H) 1.40-1.84 (m, 2H) 3.03-3.58 (m, 2H), 3.78 (s, 3H) 4.48-5.01 (m, 2H) 6.67-7.12 (m, 5H) 7.20-7.67 (m, 4H) 8.00-8.22 (m, 2H)

MS ES^+ : 438

Example 82: 4-((N-(Cyclopropylmethyl)-4-(3-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C67d) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ -0.09-0.27 (m, 2H) 0.42-0.58 (m, 2H) 0.78-1.15 (m, 1H) 3.03-3.55 (m, 2H) 3.86 (s, 3H) 4.70-5.08 (m, 2H) 6.72-6.87 (m, 1H) 6.87-7.08 (m, 4H) 7.22-7.58 (m, 4H) 8.03-8.12 (m, 2H)

MS ES^+ : 450

Example 83: 4-((N-(Cyclopropylmethyl)-5-(2-fluorophenoxy)pyrimidine-2-carboxamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C72) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ -0.06-0.20 (m, 2H) 0.38-0.57 (m, 2H) 0.91-1.14 (m, 1H) 3.01-3.43 (m, 2H) 4.64-5.06 (m, 2H), 7.12-7.33 (m, 4H) 7.40-7.53 (m, 2H) 8.01-8.10 (m, 2H) 8.38-8.57 (m, 2H)

MS ES^+ : 422

Example 84: 4-((*N*-(Cyclopropylmethyl)-5-(2-methoxyphenoxy)pyrimidine-2-carboxamido)methyl)benzoic acid




Prepared from the parent methyl ester (Intermediate C73) using Method H1.

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ -0.12-0.20 (m, 2H) 0.26-0.42 (m, 2H) 0.76-1.12 (m, 1H) 2.82-3.30 (m, 2H) 3.68-3.76 (m, 3H) 4.35-4.79 (m, 2H), 6.95-7.34 (m, 6H) 7.69-7.82 (m, 2H) 8.39-8.49 (m, 2H)

MS ES^+ : 434

Example 85: 4-((5-(2-Methoxyphenoxy)-*N*-propylpyrimidine-2-carboxamido)methyl)benzoic acid



Prepared from the parent methyl ester (Intermediate C74) using Method H1.

^1H NMR (400 MHz, CDCl_3) δ ppm 0.63-0.99 (m, 3H) 1.48-1.77 (m, 2H) 3.03-3.50 (m, 2H), 3.71-3.86 (m, 3H) 4.49-4.93 (m, 2H) 6.93-7.53 (m, 6H) 7.96-8.14 (m, 2H) 8.28-8.52

MS ES^+ : 422

3. Biological efficacy of compounds of the invention

In Vitro LPA5 Receptor Assay

Engagement of the LPA5 receptor by its ligand, oleoyl-L- α -lysophosphatidic acid (LPA),
5 leads to the release of intracellular stores of calcium into the cytoplasm mediated through a
Gq-driven increase in inositol triphosphate (IP3) levels. Gq is a heterotrimeric G protein
subunit that activates phospholipase C.

The ability of test compounds to prevent LPA-driven intracellular release of stored calcium
10 from RH7777 cells expressing human LPA5 receptors was assayed by stimulating the cells
with LPA in the presence of various test compounds to measure the test compounds'
antagonism, i.e. activity *in vitro*.

Approximately 10,000 cells in normal culture medium (Minimal Essential Media, 10%
15 Foetal Calf Serum) were dispensed per assay well in a black clear bottom poly-D-lysine
coated 384 well plate (Corning). Cells were allowed to settle for thirty minutes at room
temperature before being incubated overnight at 37°C, 5% CO₂. Sixteen to twenty hours
after dispensing, the cells were loaded with a calcium-sensitive fluorescent dye by
replacing the culture medium with assay buffer (1 x Hanks buffered saline, 25mM HEPES,
20 0.1% w/v fatty acid free BSA (bovine serum albumin), pH7.4) containing 1.25mM
probenecid and 1 x Calcium 5 Reagent (Molecular Devices). Cells were incubated at 37°C
for one hour to allow for dye uptake.

To test for antagonist activity, test compounds at a final concentration range between
25 0.32nM - 10 μ M (diluted in assay buffer) were added to the assay wells and allowed to
incubate for fifteen minutes. After incubation with test compounds the assay plate was
placed in a FLIPR Tetra system (Molecular Devices) and LPA (diluted in assay buffer)
was added to give a final concentration equivalent to its determined EC₉₀ against LPA5
receptor. Ligand-dependent changes in intracellular calcium levels were determined by
30 measuring changes in fluorescence of the dye over 515-575nm following excitation over

470-495nm. Readings from control wells that did not contain test compounds enabled percentage inhibition curves to be plotted using a 4-parameter curve fit algorithm and IC₅₀ values to be calculated for each compound.

Results

Example No.	Mean IC₅₀ (nM)	Example No.	Mean IC₅₀ (nM)
		1	840
2	690	3	43
4	63	5	16
6	24	7	31
8	32	9	15
10	48	11	6
12	9	13	3
14	2	15	120
16	1	17	1
18	2	19	9
20	40	21	11
22	24	23	44
24	7	25	29
26	260	27	200
28	45	29	43
30	12	31	5
32	0.4	33	0.4
34	460	35	7
36	24	37	28
38	140	39	11
40	31	41	89
42	17	43	58
44	11	45	99

Example No.	Mean IC ₅₀ (nM)	Example No.	Mean IC ₅₀ (nM)
46	17	47	11
48	3300	49	4600
50	3600	51	53
52	370	53	58
54	26	55	690
56	180	57	410
58	32	59	6
60	60	61	8
62	9	63	10
64	7	65	4
66	5	67	5
68	10	69	8
70	140	71	83
72	42	73	890
74	98	75	22
76	140	77	74
78	170	79	68
80	45	83	350
84	91	85	120

In Vivo Farnesyl Pyrophosphate-induced inflammatory model

Farnesyl pyrophosphate (FPP) is an endogenous ligand for the LPA5 receptor, with a reported *in-vitro* EC₅₀ of 49nM and 3-fold selectivity over other LPA receptors (Williams et al. (2009), 'Unique Ligand Selectivity of the GPR92/LPA5 Lysophosphatidate Receptor Indicates Role in Human Platelet Activation', J Biol. Chem., 284(25):17304–17319). The FPP-induced hyperalgesia model is a pharmacodynamic rat model which involves inducing pain by injecting FPP, an agonist of LPA5 receptor, into the plantar surface of the

hindpaw. Pre-treatment with a LPA5 receptor antagonist prevents the hyperalgesia induced by FPP.

In adult male rats baseline measurements were taken by assessing the withdrawal threshold to a painful pressure stimulus applied to the hindpaw according to the Randall-Selitto method (Randall and Selitto (1957), 'A method for measurement of analgesic activity on inflamed tissue', Arch Int Pharmacodyn Ther., 111(4):409-19). The animals were then administered an intraperitoneal (ip) or oral (po) dose of test compound or vehicle 30 to 120 minutes prior to an intra-plantar injection of FPP (100 μ L at 32nM) or saline vehicle.

Withdrawal responses were reassessed at 30 and 90 minutes after injection of FPP in the ipsilateral paw.

Results

Compound of Example No.	Route	In vivo efficacy (minimum effective dose (MED) mg/kg)
3	po	≤ 10
4	po	≤ 10
4	ip	1
5	po	≤ 10
5	ip	0.3
8	po	100
11	po	≤ 10
24	po	30
29	po	3
31	po	1
40	po	60

In Vitro LPA1 Receptor Assay

Engagement of the LPA1 receptor by its ligand, oleoyl-L- α -lysophosphatidic acid (LPA), leads to the release of intracellular stores of calcium into the cytoplasm mediated through a Gq-driven increase in inositol triphosphate (IP3) levels. Gq is a heterotrimeric G protein subunit that activates phospholipase C.

The ability of test compounds to prevent LPA-driven intracellular release of stored calcium from RH7777 cells expressing human LPA1 receptors was assayed by stimulating the cells with LPA in the presence of various test compounds to measure the test compounds' antagonism, i.e. activity *in vitro*.

Approximately 10,000 cells in normal culture medium (Dulbecco's Modified Eagle Media, 10% Foetal Calf Serum) were dispensed per assay well in a black clear bottom collagen coated 384 well plate (Beckton Dickinson). Cells were allowed to settle for thirty minutes at room temperature before being incubated overnight at 37°C, 5% CO₂. Sixteen to twenty hours after dispensing, the cells were loaded with a calcium-sensitive fluorescent dye by replacing the culture medium with assay buffer (1 x Hanks buffered saline, 25mM HEPES, 0.1% w/v fatty acid free BSA (bovine serum albumin), pH7.4) containing 1.25mM probenecid and 1 x Calcium 5 Reagent (Molecular Devices). Cells were incubated at 37°C for one hour to allow for dye uptake.

To test for antagonist activity, test compounds at a final concentration range between 0.32nM - 10 μ M (diluted in assay buffer) were added to the assay wells and allowed to incubate for twenty five minutes. After incubation with test compounds the assay plate was placed in a FLIPR Tetra system (Molecular Devices) and LPA (diluted in assay buffer) was added to give a final concentration equivalent to its determined EC₈₀ against LPA1 receptor. Ligand-dependent changes in intracellular calcium levels were determined by measuring changes in fluorescence of the dye over 515-575nm following excitation over 470-495nm. Readings from control wells that did not contain test compounds

enabled percentage inhibition curves to be plotted using a 4-parameter curve fit algorithm and IC₅₀ values to be calculated for each compound.

Example No.	Mean IC ₅₀ (nM)	Example No.	Mean IC ₅₀ (nM)
3	35	4	49
5	46	8	46
11	6	13	4
14	2	17	0.4
18	1	19	7
20	12	22	30
24	11	25	14
28	46	29	34
30	12	31	4
32	2	33	0.3
36	24	37	25
38	36	40	160
45	75	46	19
47	22	54	27
58	17		
60	33	61	4
62	5	63	7
64	9	65	5
66	7	67	7
68	13	69	12
70	47	71	59
72	29	73	120
74	40	75	31
76	130	77	86

Example No.	Mean IC ₅₀ (nM)	Example No.	Mean IC ₅₀ (nM)
78	390	79	94
80	110	83	540
84	170	85	180

In vivo Bleomycin induced Scleroderma Model

Experimental outline

- 5 Adult male C57BL/6 mice, aged 8-10 weeks, were assigned to groups and allowed to acclimatise for a week. On Day -1, the back skin of the animals was shaved. From Day 0, animals were administered with 100 µl of a 1 mg/ml solution of bleomycin sulphate in phosphate buffered saline (PBS) by intradermal injection in the shaved skin. Injections were repeated in a one square centimetre area every other day for four weeks. A group of
- 10 ten animals (Control group) was injected with an identical volume of PBS. Treatments were given according to the administration schedule shown in the table below. Animals were weighed at the beginning of the experiment on Day 0 and then twice weekly throughout the study. At the end of the experiment, on Day 28, animals were culled and samples of back skin were taken for further analysis. All groups were n=10; the vehicle
- 15 was distilled water and the administration volume was 10 ml/kg for oral gavage (p.o.); administration volume was 25 µl for topical application.

Group	Substance	Dose	Route	Administration Frequency	Skin sensitisation
1	Vehicle	n/a	p.o.	BID, Day 0 – Day 28	PBS, i.d., QAD, Day 0 – Day 28
2	Vehicle	n/a	p.o.	BID, Day 0 – Day 28	


3	Imatinib mesylate	150 mg/kg	p.o.	BID, Day 0 – Day 28	Bleomycin, i.d., QAD, Day 0 – Day 28
4	Compound of Example 5	30 mg/kg	p.o.	BID, Day 0 – Day 28	
5	Compound of Example 5	30 mg/kg	p.o.	BID, Day 14 – Day 28	
6	“Protopic” (trade mark)	0.1%	topical	BID, Day 0 – Day 28	
7	Betamethasone	0.1%	topical	BID, Day 0 – Day 9 SID, Day 10 – Day 28	

Results

Bodyweights were graphed along with the standard error of the mean (SEM); Skin samples (two punch biopsies per animal, 4 mm in diameter each) samples were hydrolyzed and hydroxyproline content of the samples was determined from analysis in a semi-automated process using a continuous flow analyzer. Autoanalyser readings were calibrated against hydroxyproline standards and collagen content calculated. The bodyweight loss observed in the Example 5 compound- treated groups did not differ from the bodyweight loss observed in the vehicle-treated group. Some weight loss <10% was observed with bleomycin treatment. Bleomycin administration induced a highly significant increase in skin collagen content when compared to the control, saline-injected, group ($p < 0.01$). Imatinib and Betamethasone 0.1% induced a non-significant reduction in skin collagen content when compared to the vehicle-treated group. The Example 5 compound administered from Day 0 and from Day 14 induced a significant reduction in skin collagen content when compared to the vehicle-treated group ($p < 0.05$). Protopic 0.1% induced a highly significant reduction in skin collagen content when compared to the vehicle-treated group ($p < 0.001$). Thus, the compound of Example 5 demonstrated an anti-fibrotic effect in an *in vivo* bleomycin-induced scleroderma model as measured by collagen skin content.

CLAIMS

1. A compound of formula (I)



5 wherein

R^1 represents a C₅-C₁₀ aryl group substituted by at least one substituent selected from halogen, cyano, nitro, carboxyl, hydroxyl, C₃-C₆ cycloalkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxycarbonyl, C₁-C₆ alkoxycarbonylamino, C₁-C₆ haloalkoxy, -NR⁵R⁶, C₃-C₆ cycloalkylamino, C₁-C₆ alkylthio, C₁-C₆ alkylcarbonyl, C₁-C₆ alkylcarbonyloxy, C₁-C₆ alkylcarbonylamino, sulphonamido, C₁-C₆ alkylsulphonyl, C₁-C₆ alkylsulphonylamino, -C(O)NR⁷R⁸, C₁-C₆ alkyl which alkyl group may in turn be optionally substituted by at least one halogen, hydroxyl, carboxyl or C₁-C₆ alkoxycarbonyl group, and a saturated or unsaturated 5- to 6-membered ring which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur, the ring itself being optionally substituted by at least one substituent selected from halogen, hydroxyl, oxo, carboxyl, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxycarbonyl and C₁-C₆ hydroxyalkyl;

X represents an oxygen atom or a group -CH₂-, -OCH₂-, -CH₂O-, -CH₂CH₂-, -CH₂NR¹⁷-, -NR¹⁷CH₂-, -CHF- or -CF₂-;

m is 0, 1 or 2;

each R^2 independently represents a halogen atom or a hydroxyl, C_1 - C_6 alkyl, difluoromethyl, trifluoromethyl, C_1 - C_6 alkoxy, difluoromethoxy, trifluoromethoxy or $NR^{15}R^{16}$ group;

Y represents CR^9R^{10} in which R^9 and R^{10} each independently represent a hydrogen atom or a methyl group;

R^3 represents C_1 - C_8 alkyl, C_2 - C_8 alkenyl, or a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur, each of the preceding groups being optionally substituted by at least one substituent selected from halogen, cyano, hydroxyl, C_3 - C_6 cycloalkyl, C_1 - C_6 alkoxy, C_5 - C_6 aryloxy, C_5 - C_6 aryl C_1 - C_6 alkyloxy, C_1 - C_6 alkoxycarbonyl, C_1 - C_6 alkoxycarbonylamino, C_1 - C_6 alkoxycarbonyloxy, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy, $-NR^{13}R^{14}$, C_3 - C_6 cycloalkylamino, C_1 - C_6 alkylthio, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylcarbonyloxy, C_1 - C_6 alkylcarbonylamino, sulphonamido, C_1 - C_6 alkylsulphonyl, C_1 - C_6 alkylsulphonylamino, C_1 - C_6 alkyl, and a saturated or unsaturated 3- to 10-membered ring system which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur, the ring system itself being optionally substituted by least one substituent selected from halogen, hydroxyl, oxo, carboxyl, cyano, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkoxycarbonyl and C_1 - C_6 hydroxyalkyl;

Z represents $CR^{11}R^{12}$ in which R^{11} and R^{12} each independently represent a hydrogen atom or a methyl group but cannot both simultaneously represent a methyl group;

n is 0, 1 or 2;

each R^4 independently represents a halogen atom or a hydroxyl, C_1 - C_6 alkyl, difluoromethyl, trifluoromethyl, C_1 - C_6 alkoxy, difluoromethoxy, trifluoromethoxy or $NR^{18}R^{19}$ group;

R^5 and R^6 each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or R^5 and R^6 together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^7 and R^8 each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or R^7 and R^8 together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{13} and R^{14} each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or R^{13} and R^{14} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{15} and R^{16} each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or R^{15} and R^{16} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle;

R^{17} represents a hydrogen atom or a C₁-C₆ alkyl group;

R^{18} and R^{19} each independently represent a hydrogen atom or a C₁-C₆ alkyl group, or R^{18} and R^{19} together with the nitrogen atom to which they are attached form a 4- to 7-membered saturated heterocycle; and

rings A and B each independently represent a saturated or unsaturated 5- to 10-membered ring system which may comprise at least one ring heteroatom selected from nitrogen, oxygen and sulphur;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1, wherein R^1 represents a C₅-C₁₀ aryl group substituted by one or two substituents independently selected from halogen, cyano, C₁-C₂ alkoxy, and C₁-C₂ alkyl which alkyl group may in turn be optionally substituted by at least one halogen atom.

3. A compound according to claim 1 or claim 2, wherein X represents an oxygen atom or -CH₂O-.
4. A compound according to any one of claims 1 to 3, wherein either m is 0, or, m is 1 or 2
5 and each R² independently represents a halogen atom or a C₁-C₆ alkyl group.
5. A compound according to any one of claims 1 to 4, wherein Y represents CH₂.
6. A compound according to any one of claims 1 to 5, wherein R³ represents a C₁-C₃ alkyl
10 group optionally substituted by one or more substituents as defined in claim 1.
7. A compound according to any one of claims 1 to 5, wherein R³ represents a saturated or unsaturated 3- to 6-membered hydrocarbyl ring system optionally substituted by one or more substituents as defined in claim 1.
15
8. A compound according to claim 6 or claim 7, wherein the optional substituents are selected from fluorine, chlorine, hydroxyl, C₃-C₆ cycloalkyl, C₁-C₂ alkoxy, phenoxy, benzyloxy, trifluoromethyl, C₁-C₂ alkyl, and a saturated or unsaturated 3-, 4-, 5- or 6-membered ring system which may comprise at least one ring heteroatom selected from
20 nitrogen, oxygen and sulphur, the ring system itself being optionally substituted by least one substituent selected from halogen, C₁-C₂ alkyl and C₁-C₂ alkoxy.
9. A compound according to any one of the preceding claims, wherein Z represents CH₂.
- 25 10. A compound according to any one of the preceding claims, wherein either n is 0, or, n is 1 and R⁴ represents a halogen atom or a C₁-C₆ alkyl or C₁-C₆ alkoxy group.

11. A compound according to any one of the preceding claims, wherein rings A and B are independently selected from cyclopentyl, cyclohexyl, cyclopentenyl, cyclohexenyl, phenyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, oxadiazolyl (e.g. 1,2,4-oxadiazolyl), tetrahydrofuranyl, naphthyl, benzofuranyl, benzothienyl, benzodioxolyl, benzoxazolyl, quinoliny, oxazolyl, thiadiazolyl (e.g. 1,2,3-thiadiazolyl), 2,3-dihydrobenzofuranyl, tetrahydropyranyl, pyrazolyl, imidazo[1,2-a]pyridinyl, pyrazinyl, thiazolidinyl, indanyl, thienyl, isoxazolyl, pyridazinyl, pyrrolyl, furanyl, thiazolyl, isothiazolyl, indolyl, isoindolyl, imidazolyl, pyrimidinyl, benzimidazolyl, triazolyl, tetrazolyl and pyridinyl.

12. A compound according to any one of claims 1 to 10, wherein rings A and B each represent phenyl.

13. A compound selected from the group consisting of:

- 4-((*N*-Ethyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
- 4-((*N*-(2,2-Difluoroethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
- 4-((4-(2-Fluorophenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,
- 4-((4-(2-Fluorophenoxy)-*N*-isobutylbenzamido)methyl)benzoic acid,
- 4-((*N*-(Cyclopropylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
- 4-((*N*-(Cyclobutylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
- 4-((4-(2-Fluorophenoxy)-*N*-((3-methoxycyclobutyl)methyl)benzamido)-methyl)benzoic acid,
- 4-((4-(2-Fluorophenoxy)-*N*-isopentylbenzamido)methyl)benzoic acid,
- 4-((*N*-(2-Cyclopropylethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
- 4-((*N*-(2-Cyclobutylethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
- 4-((*N*-Benzyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
- 4-((*N*-(2-Fluorobenzyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,
- 4-((4-(2-Fluorophenoxy)-*N*-(3-methoxybenzyl)benzamido)methyl)benzoic acid,
- 4-((4-(2-Fluorophenoxy)-*N*-(4-methoxybenzyl)benzamido)methyl)benzoic acid,
- 4-((4-(2-Fluorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-(3-(2-fluorophenyl)propyl)benzamido)methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-(3-(3-fluorophenyl)propyl)benzamido)methyl)benzoic acid,

5 4-((4-(2-Fluorophenoxy)-*N*-((*trans*-2-phenylcyclopropyl)methyl)benzamido)-methyl)benzoic acid,

(*S*)-4-((4-(2-Fluorophenoxy)-*N*-(2-hydroxy-3-phenylpropyl)benzamido)methyl)benzoic acid,

(*R*)-4-((4-(2-Fluorophenoxy)-*N*-(2-hydroxy-3-phenylpropyl)benzamido)-
10 methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-(4-phenylbutyl)benzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,

4-((*N*-(Cyclobutylmethyl)-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,

4-((*N*-Benzyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,

15 4-((*N*-Phenethyl-4-(*o*-tolylloxy)benzamido)methyl)benzoic acid,

4-((*N*-Phenethyl-4-(2-(trifluoromethyl)phenoxy)benzamido)methyl)benzoic acid,

4-((*N*-Benzyl-4-(2-cyanophenoxy)benzamido)methyl)benzoic acid,

4-((*N*-Ethyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,

4-((*N*-(2,2-Difluoroethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,

20 4-((*N*-Isobutyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,

4-((4-(2-Methoxyphenoxy)-*N*-(3-phenylpropyl)benzamido)methyl)benzoic acid,

4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)benzamido)-

25 methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-(2-ethoxyphenoxy)benzamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-(cyclopropylmethyl)benzamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic

30 acid,

4-((*N*-(2-Cyclopropylethyl)-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid,

4-((*N*-Benzyl-4-(2,6-difluorophenoxy)benzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-(2-fluoro-6-methylphenoxy)benzamido)-
methyl)benzoic acid,

4-((*N*-(Cyclobutylmethyl)-4-(2-fluoro-6-methylphenoxy)benzamido)-
methyl)benzoic acid,

4-((4-(2-Chloro-6-fluorophenoxy)-*N*-(cyclopropylmethyl)benzamido)-
methyl)benzoic acid,

4-((4-(2-Chloro-6-fluorophenoxy)-*N*-(2-cyclopropylethyl)benzamido)-
methyl)benzoic acid,

4-((*N*-Benzyl-4-(2-chloro-6-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((4-(2,6-Dimethylphenoxy)-*N*-isopentylbenzamido)methyl)benzoic acid,

4-((*N*-Benzyl-4-(2,6-dimethylphenoxy)benzamido)methyl)benzoic acid,

4-((4-(2,6-Dimethylphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((4-(3-Methoxyphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((4-(4-Methoxyphenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((4-(3-Chlorophenoxy)-*N*-phenethylbenzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic
acid,

4-((*N*-(Cyclobutylmethyl)-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic
acid,

4-((*N*-Benzyl-4-((2-fluorobenzyl)oxy)benzamido)methyl)benzoic acid,

4-((*N*-(Cyclopropylmethyl)-2-fluoro-4-(2-fluorophenoxy)-
benzamido)methyl)benzoic acid,

4-((*N*-Benzyl-4-(2-fluorophenoxy)cyclohexanecarboxamido)methyl)benzoic acid,

trans-4-((*N*-Benzyl-4-(2-methoxyphenoxy)cyclohexanecarboxamido)-
methyl)benzoic acid,

cis-4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-

methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid,

trans-4-((*N*-(3-(3-Fluorophenyl)propyl)-4-(2-methoxyphenoxy)cyclohexanecarboxamido)methyl)benzoic acid

2-Fluoro-4-((4-(2-fluorophenoxy)-*N*-(3-methoxybenzyl)benzamido)-methyl)benzoic acid,

5 4-((4-(2-Fluorophenoxy)-*N*-propylbenzamido)methyl)benzoic acid,

4-((4-(2-Methoxyphenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

4-((4-(2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,

4-((*N*-(2,2-difluoropropyl)-4-(2-methoxyphenoxy)-benzamido)methyl)benzoic

10 acid,

4-((4-(2-methoxyphenoxy)-*N*-(3,3,3-trifluoropropyl)-benzamido)methyl)benzoic acid,

4-((*N*-butyl-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid,

4-((*N*-(cyclopropylmethyl)-2-fluoro-4-(2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

15 4-((*N*-(cyclopropylmethyl)-4-(2-fluoro-6-methoxyphenoxy)-benzamido)methyl)benzoic acid,

4-((*N*-(cyclopropylmethyl)-4-(4-fluoro-2-methoxyphenoxy)-benzamido)methyl)benzoic acid,

20 4-((*N*-(cyclopropylmethyl)-5-(2-methoxyphenoxy)-picolinamido)methyl)benzoic acid,

4-((4-(*o*-Tolyloxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

4-((4-(2-Fluorophenoxy)-*N*-(3,3,3-trifluoropropyl)benzamido)methyl)benzoic acid,

25 4-((4-(2-Chlorophenoxy)-*N*-(2,2,2-trifluoroethyl)benzamido)methyl)benzoic acid,

4-((*N*-Ethyl-4-(*o*-tolyloxy)benzamido)methyl)benzoic acid,

4-((*N*-(2,2-Difluoropropyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((*N*-Butyl-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-ethylbenzamido)methyl)benzoic acid,

30 4-((*N*-(Cyclopropylmethyl)-5-(2-fluorophenoxy)picolinamido)methyl)benzoic acid,

4-((4-(2-Chlorophenoxy)-*N*-(2,2-difluoroethyl)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(5-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoic acid,
 4-((4-(4-Fluoro-2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,
 5 4-((4-(5-Fluoro-2-methoxyphenoxy)-*N*-propylbenzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-4-(3-fluoro-2-methoxyphenoxy)benzamido)methyl)benzoic acid,
 4-((*N*-(Cyclopropylmethyl)-5-(2-fluorophenoxy)pyrimidine-2-carboxamido)methyl)benzoic acid,
 10 4-((*N*-(Cyclopropylmethyl)-5-(2-methoxyphenoxy)pyrimidine-2-carboxamido)methyl)benzoic acid,
 4-((5-(2-Methoxyphenoxy)-*N*-propylpyrimidine-2-carboxamido)methyl)benzoic acid,
 and pharmaceutically acceptable salts thereof.

15

14. The compound according to claim 13, wherein the compound is 4-((*N*-(cyclopropylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid.

15. A compound according to claim 13, wherein the compound is

20 4-((*N*-(cyclopropylmethyl)-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid or a pharmaceutically acceptable salt thereof.

16. A compound according to claim 13, wherein the compound is

4-((*N*-(Cyclopropylmethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid.

25

17. A compound according to claim 13, wherein the compound is

4-((*N*-(Cyclopropylmethyl)-4-(2-methoxyphenoxy)benzamido)methyl)benzoic acid
 or a pharmaceutically acceptable salt thereof.


30

18. A compound according to claim 13, wherein the compound is

4-((*N*-(Cyclopropylmethyl)-2-fluoro-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid.

19. A compound according to claim 13, wherein the compound is 4-((N-(Cyclopropylmethyl)-2-fluoro-4-(2-fluorophenoxy)benzamido)methyl)benzoic acid or a pharmaceutically acceptable salt thereof.

20. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined in claim 1 which comprises hydrolysing a compound of formula (II)



wherein L^1 represents a leaving group and R^1 , X, m, R^2 , Y, R^3 , Z, n, R^4 , A and B are as defined in formula (I), in the presence of a suitable base;

and optionally thereafter carrying out one or more of the following procedures:

- converting a compound of formula (I) into another compound of formula (I)
- removing any protecting groups
- forming a pharmaceutically acceptable salt.

21. A pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 19, in association with a pharmaceutically acceptable adjuvant, diluent or carrier, and optionally one or more other therapeutic agents.

22. A compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 19 for use in therapy.

23. A compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 19 for use in treating pain disorders, atherosclerosis or fibrosis.

24. Use of a compound of formula (I) or a pharmaceutically acceptable salt thereof as claimed in any one of claims 1 to 19, in the manufacture of a medicament for treating pain disorders, atherosclerosis or fibrosis.