

[Document Name] Description

[Title of Invention] FUSION PROTEIN FOR IMPROVING PROTEIN
EXPRESSION LEVEL FROM TARGET mRNA

[Technical Field]

[0001]

The present invention relates to fusion proteins for improving protein expression levels from target mRNAs.

[Background Art]

[0002]

Techniques of binding nucleic acid-binding protein factors revealed by a variety of analyses to sequences of interest are established and used in recent years. Use of this sequence-specific binding enables removal of a target DNA sequence or regulation (activation or inactivation) of expression of a protein coding gene present downstream of the target DNA sequence in some extent.

[0003]

While zinc finger nuclease (ZFN), TAL effector nuclease (TALEN), Crispr-cas9, and the like are known as techniques using protein factors which act on DNA, the development of techniques using protein factors which act specifically to RNA is still limited.

[0004]

The present inventors have proposed a method of designing a protein which can specifically bind to a target RNA sequence using the properties of PPR proteins (protein having one or more pentatricopeptide repeat (PPR) motifs), which are proteins mainly found in plants (Patent Literature 1).

[Citation List]

[Patent Literature]

[0005]

[Patent Literature 1]

WO2013/058404

[Summary of Invention]

[Technical Problem]

[0006]

In the disclosure according to Patent Literature 1, the amino acids which function when a PPR motif demonstrates RNA-binding properties were identified, and the relation between the structure of the PPR motif and the target base was revealed, thereby enabling the construction of proteins which have one or more PPR motifs and can bind to RNAs having any sequence and length. However, no method has ever been found which actually regulates target RNAs using the techniques according to Patent Literature 1.

[Solution to Problem]

[0007]

As a result of extensive research on a method of improving a protein expression level from a target mRNA using a PPR protein, the present inventors have found that a fusion protein of a predetermined functional domain and a PPR protein improves the protein expression level from the target mRNA, and have completed the present invention.

[0008]

Specifically, an embodiment of the present invention relates to a fusion protein for improving a protein expression level from a target mRNA, the fusion protein comprising:

(A) one or more functional domains which improve a protein expression level from an mRNA; and

(B) a polypeptide moiety which can bind to a target mRNA in an RNA base-selective or RNA base sequence-specific manner,

wherein polypeptide moiety (B) is a polypeptide moiety comprising one or more PPR motifs, each PPR motif comprising a polypeptide consisting of 30 to 38 amino acids in length and being represented by Formula 1:

[Formula 1]

(Helix A)-X-(Helix B)-L (Formula 1)

where

Helix A is a moiety which consists of 12 amino acids in length and can form an α -helix structure, and is represented by Formula 2:

[Formula 2]



where A_1 to A_{12} each independently represent an amino acid;

X is not present, or is a moiety consisting of 1 to 9 amino acids in length;

Helix B is a moiety which consists of 11 to 13 amino acids in length and can form an α -helix structure;

L is a moiety consisting of 2 to 7 amino acids in length and represented by Formula 3:

[Formula 3]



where the amino acids are numbered from the C-terminal as "i" (-1), "ii" (-2), ... and

L_{iii} to L_{vii} may not be present, and

a combination of three amino acids A_1 , A_4 , and L_{ii} or a combination of two amino acids A_4 and L_{ii} corresponds to a base or base sequence of the target mRNA.

[0009]

In an embodiment according to the present invention, polypeptide moiety (B) comprises 2 to 30 PPR motifs, and

the plurality of PPR motifs is arranged so as to specifically bind to the base sequence of the target mRNA.

[0010]

Moreover, in an embodiment according to the present invention, polypeptide moiety (B) comprises 5 to 25 PPR motifs.

[0011]

Moreover, in an embodiment according to the present invention, one or more functional domains (A) each bind to an N-terminal side and/or a C-terminal side of polypeptide moiety (B).

[0012]

Moreover, in an embodiment according to the present invention, one or more functional domains (A) are selected from the group consisting of a domain which guides ribosome to the mRNA, a domain associated with initiation or promotion of translation of the mRNA, a domain associated with nuclear export of the mRNA, a domain associated with binding to an endoplasmic reticulum membrane, a domain containing an endoplasmic reticulum retention signal (ER retention signal) sequence, and a domain containing an endoplasmic reticulum signal sequence.

[0013]

Moreover, in an embodiment according to the present invention, the domain which guides ribosome to the mRNA is a domain containing all or functional part of a

polypeptide selected from the group consisting of DENR (Density-regulated protein), MCT-1 (Malignant T-cell amplified sequence 1), TPT1 (Translationally-controlled tumor protein), and Lerepo4 (Zinc finger CCCH-domain),

the domain associated with initiation or promotion of translation of the mRNA is a domain containing all or functional part of a polypeptide selected from the group consisting of eIF4E and eIF4G,

the domain associated with nuclear export of the mRNA is a domain containing all or functional part of SLBP (Stem-loop binding protein),

the domain associated with binding to an endoplasmic reticulum membrane is a domain containing all or functional part of a polypeptide selected from the group consisting of SEC61B, TRAP-alpha (Translocon associated protein alpha), SR-alpha, Dial (Cytochrome b5 reductase 3), and p180,

the endoplasmic reticulum retention signal (ER retention signal) sequence is a signal sequence containing a KDEL (KEEL) sequence, or

the endoplasmic reticulum signal sequence is a signal sequence containing MGWSCIIILFLVATATGAHS (SEQ ID NO: 22).

[0014]

Moreover, in an embodiment according to the present invention, the combination of the three amino acids A₁, A₄, and L_{ii} in each of the PPR motifs is:

(valine, threonine, asparagine), (phenylalanine, serine, asparagine), (phenylalanine, threonine, asparagine), (isoleucine, asparagine, aspartic acid), or (threonine, threonine, asparagine) in order of (A₁, A₄, L_{ii}) if a target base for the PPR motif is A (adenine);

(glutamic acid, glycine, aspartic acid), (valine, threonine, aspartic acid), (lysine, threonine, aspartic acid), or (leucine, threonine, aspartic acid) in order of (A₁, A₄, L_{ii}) if the target base for the PPR motif is G (guanine);

(valine, asparagine, aspartic acid), (isoleucine, asparagine, asparagine), (isoleucine, asparagine, aspartic acid), (isoleucine, methionine, aspartic acid), (phenylalanine, proline, aspartic acid), or (tyrosine, proline, aspartic acid) in order of (A₁, A₄, L_{ii}) if the target base for the PPR motif is U (uracil); or

(valine, asparagine, asparagine), (isoleucine, asparagine, asparagine), (valine, asparagine, serine), or (isoleucine, methionine, aspartic acid) in order of (A₁, A₄, L_{ii}) if the target base for the PPR motif is C (cytosine).

[0015]

Moreover, in an embodiment according to the present invention, the combination of the two amino acids A₄ and L_{ii} in each of the PPR motifs is:

(threonine, asparagine), (serine, asparagine), or (glycine, asparagine) in order of (A₄, L_{ii}) if a target base for the PPR motif is A (adenine);

(threonine, aspartic acid) or (glycine, aspartic acid) in order of (A₄, L_{ii}) if the target base for the PPR motif is G (guanine);

(asparagine, aspartic acid), (proline, aspartic acid), (methionine, aspartic acid), or (valine, threonine) in order of (A₄, L_{ii}) if the target base for the PPR motif is U (uracil); or

(asparagine, asparagine), (asparagine, serine), or (leucine, aspartic acid) in order of (A₄, L_{ii}) if the target base for the PPR motif is C (cytosine).

[0016]

Another embodiment according to the present invention relates to a nucleic acid which encodes the fusion protein according to the present invention.

[0017]

Still another embodiment according to the present invention relates to a vector (preferably an expression vector) comprising the nucleic acid according to the present invention.

[0018]

Further still another embodiment according to the present invention relates to a method of improving a protein expression level from a target mRNA within a cell, the method comprising:

a step of providing the fusion protein according to the present invention or the vector according to the present invention; and

a step of introducing the fusion protein or the vector into the cell.

[0019]

Moreover, in an embodiment according to the present invention, the cell is a eukaryotic cell.

[0020]

Moreover, in an embodiment according to the present invention, the cell is an animal cell.

[0021]

Moreover, in an embodiment according to the present invention, the animal cell is a human cell.

[0022]

Inventions having any combination of one or more features of the present invention described above are also included in the scope of the present invention.

[Brief Description of Drawings]

[0023]

[Figure 1] Figure 1 illustrates a schematic view of an effector plasmid and a reporter plasmid used in Examples, and a schematic view of an experimental outline. Figure 1A illustrates a schematic view of the effector plasmid and the reporter plasmid used in Examples. A fusion protein of PPR motifs and eIF4G expresses from the

effector plasmid. In Examples, a CRR4 protein was used, whose target sequence is well researched. From the reporter plasmid, renilla luciferase (RLuc) and firefly luciferase (FLuc) are transcribed in the form of a dicistronic mRNA. A PPR-binding sequence (here, CRR4-binding sequence) was inserted into a site on the 5' end of FLuc. Figure 1B illustrates a schematic view of an experimental outline of Examples. Irrespective of the presence/absence of the PPR-binding sequence, RLuc is translated at a similar level. For this reason, the activity value of RLuc can be treated as a control in transfection in this reporter system. The translation of FLuc is started only when PPR-eIF4G binds to the PPR-binding sequence and translation factors can be attracted by the effects of eIF4G. In contrast, the translation of FLuc remains at a low level if the PPR-binding sequence is not present and thus, PPR-eIF4G cannot bind to the PPR-binding sequence.

[Figure 2] Figure 2 illustrates an experimental procedure of a reporter assay using HEK293T cells.

[Figure 3] Figure 3 shows the experimental results of Example 1. The activation of sequence-specific translation depends on CRR4-eIF4G and the PPR-binding sequence. This experiment was performed using an effector plasmid, into which CRR4-Flag (without translation activating factor, in white) or CRR4-eIF4G (with translation activating factor, in gray) was

inserted, and a reporter vector with or without an inserted PPR-binding sequence. From the results, it was verified that specific translation activity increased 2.75 times in the presence of both PPR-eIF4G and the PPR-binding sequence. The value represents the average and the standard deviation (N = 3).

[Figure 4] Figure 4 illustrates an outline of the experiment in Example 2.

[Figure 5] Figure 5 illustrates the experimental results in Example 2 and the functions of the domains.

[Figure 6] Figure 6 illustrates the experimental results in Example 2 and the functions of the domains.

[Description of Embodiment]

[0024]

[PPR motifs and PPR proteins]

Unless otherwise specified, the term "PPR motif" used in the present invention indicates a polypeptide which is composed of 30 to 38 amino acids and has an amino acid sequence having an E value equal to or less than a predetermined value (desirably E-03), the E value being obtained at PF01535 in Pfam and PS51375 at Prosite during the analysis of the amino acid sequence with a protein domain search program on the Web. The position number of an amino acid forming the PPR motif defined in the present invention is substantially as defined as PF01535 while it corresponds to the number obtained by

subtracting 2 from the location of the amino acid in PS51375 (for example, position 1 in the present invention corresponds to position 3 in PS51375). Note that the term "ii" (-2)-th amino acid refers to the second amino acid from the tail end (C-terminal side) of the amino acids forming one PPR motif or the amino acid close to the N-terminal by two amino acids from the first amino acid of the next PPR motif (that is, -2 amino acid). If the next PPR motif is not clearly identified, the forward amino acid by two amino acids from the first amino acid of the next helix structure is defined as "ii". See <http://pfam.sanger.ac.uk/> for Pfam and <http://www.expasy.org/prosite/> for Prosite.

[0025]


Although the conserved amino acid sequence of the PPR motif has low conservation properties at the amino acid level, two α -helices are well conserved on the secondary structure. Although a typical PPR motif is composed of 35 amino acids, its length is variable from 30 to 38 amino acids.

[0026]

More specifically, the term PPR motif used in the present invention is composed of a polypeptide having 30 to 38 amino acids in length and being represented by Formula 1:

[0027]

[Formula 4]



where

Helix A is a moiety which consists of 12 amino acids in length and can form an α -helix structure, and is represented by Formula 2:

[0028]

[Formula 5]



where A_1 to A_{12} each independently represent an amino acid;

X is not present, or is a moiety consisting of 1 to 9 amino acids in length;

Helix B is a moiety which consists of 11 to 13 amino acids in length and can form the α -helix structure; and

L is a moiety consisting of 2 to 7 amino acids in length and represented by Formula 3:

[0029]

[Formula 6]



where the amino acids are numbered from the C-terminal side as "i" (-1), "ii" (-2), ... and

L_{iii} to L_{vii} may not be present.

[0030]

Unless otherwise specified, the term "PPR protein" used in the present invention indicates a PPR protein comprising one or more PPR motifs described above, preferably two or more PPR motifs described above. Unless otherwise specified, the term "protein" used herein generally indicates substances consisting of polypeptides (chains of several amino acids bound through peptide bond), also including those consisting of relatively low molecular weight polypeptides. The term "amino acid" used in the present invention may indicate a usual amino acid molecule, or otherwise may indicate an amino acid residue forming a peptide chain in some cases. Persons skilled in the art clearly understand from contexts which case the term indicates.

[0031]

Unless otherwise specified, the "selective" used in the present invention about the binding properties of the PPR motif to the RNA bases indicates that the binding activity of a PPR motif to one of the RNA bases is higher than the binding activity thereof to other bases. Persons skilled in the art can plan the experiment for this selectivity and verify it, and can also determine through calculation.

[0032]

Unless otherwise specified, the term "RNA base" used in the present invention indicates a base of a ribonucleotide forming an RNA, specifically adenine (A),

guanine (G), cytosine (C), or uracil (U). Note that although the PPR protein can have selectivity to the base in the RNA, it does not bind to a nucleic acid monomer.

[0033]

PPR protein is present in many plants, and 500 proteins, about 5000 motifs can be found in *Arabidopsis thaliana*. PPR motifs and PPR proteins having a variety of amino acid sequences are also present in many land plants such as *Oryza*, *Populus*, and *Selaginella tamariscina*. In the present invention, PPR motifs and PPR proteins present in the natural world may be used, or PPR motifs and PPR proteins designed based on the method disclosed in WO2013/058404, for example, may be used. Specifically, desired PPR motifs and PPR proteins can be designed based on the following information disclosed in WO2013/058404.

[0034]

(I) Information on the position of the amino acid essential for selective binding

The combination (A1, A4, Lii) of three, i.e., 1st, 4th, and "ii" (-1)-th amino acids of a PPR motif or the combination (A4, Lii) of two, i.e., 4th and "ii" (-1)-th amino acids is essential for selective binding to the RNA base, and the target RNA base for binding can be determined by these combinations.

[0035]

The present invention can use the findings about the combination of three amino acids A1, A4, and Lii, and/or the combination of two amino acids A4 and Lii disclosed in WO2013/058404.

[0036]

(II) Information about the correspondence of the combination of three amino acids A1, A4, and Lii to RNA bases

(3-1) If the combination of three amino acids A1, A4, and Lii is valine, asparagine, and aspartic acid in this order, the PPR motif has a selective RNA base binding ability as follows: binding to U is the strongest, and binding to C is the second strongest, followed by binding to A or G.

(3-2) If the combination of three amino acids A1, A4, and Lii is valine, threonine, and asparagine in this order, the PPR motif has a selective RNA base binding ability as follows: binding to A is the strongest, and binding to G is the second strongest, followed by binding to C without binding to U.

(3-3) If the combination of three amino acids A1, A4, and Lii is valine, asparagine, and asparagine in this order, the PPR motif has a selective RNA base binding ability as follows: binding to C is the strongest and binding to A or U is the second strongest, without binding to G.

(3-4) If the combination of three amino acids A1, A4, and Lii is glutamic acid, glycine, and aspartic acid in this

order, the PPR motif has a selective RNA base binding ability as follows: binding to G is strong, without binding to A, U, or C.

(3-5) If the combination of three amino acids A1, A4, and Lii is isoleucine, asparagine, and asparagine in this order, the PPR motif has a selective RNA base binding ability as follows: binding to C is the strongest, and binding to U is the second strongest, followed by binding to A, without binding to G.

(3-6) If the combination of three amino acids A1, A4, and Lii is valine, threonine, and aspartic acid in this order, the PPR motif has a selective RNA base binding ability as follows: binding to G is the strongest and binding to U is the second strongest, without binding to A or C.

(3-7) If the combination of three amino acids A1, A4, and Lii is lysine, threonine, and aspartic acid in this order, the PPR motif has a selective RNA base binding ability as follows: binding to G is the strongest and binding to A is the second strongest, without binding to U or C.

(3-8) If the combination of three amino acids A1, A4, and Lii is phenylalanine, serine, and asparagine in this order, the PPR motif has a selective RNA base binding ability as follows: binding to A is the strongest, and binding to C is the second strongest, followed by binding to G and U.

(3-9) If the combination of three amino acids A1, A4, and Lii is valine, asparagine, and serine in this order, the

PPR motif has a selective RNA base binding ability as follows: binding to C is the strongest and binding to U is the second strongest, without binding to A or G.

(3-10) If the combination of three amino acids A1, A4, and Lii is phenylalanine, threonine, and asparagine in this order, the PPR motif has a selective RNA base binding ability as follows: binding to A is strong, without binding to G, U, or C.

(3-11) If the combination of three amino acids A1, A4, and Lii is isoleucine, asparagine, aspartic acid in this order, the PPR motif has a selective RNA base binding ability as follows: binding to U is the strongest and binding to A is the second strongest, without binding to G or C.

(3-12) If the combination of three amino acids A1, A4, and Lii is threonine, threonine, and asparagine in this order, the PPR motif has a selective RNA base binding ability as follows: binding to A is strong, without binding to G, U, or C.

(3-13) If the combination of three amino acids A1, A4, and Lii is isoleucine, methionine, and aspartic acid in this order, the PPR motif has a selective RNA base binding ability as follows: binding to U is the strongest and binding to C is the second strongest, without binding to A or G.

(3-14) If the combination of three amino acids A1, A4, and Lii is phenylalanine, proline, and aspartic acid in

this order, the PPR motif has a selective RNA base binding ability as follows: binding to U is the strongest and binding to C is the second strongest, without binding to A or G.

(3-15) If the combination of three amino acids A1, A4, and Lii is tyrosine, proline, and aspartic acid in this order, the PPR motif has a selective RNA base binding ability as follows: binding to U is strong, without binding to A, G, or C.

(3-16) If the combination of three amino acids A1, A4, and Lii is leucine, threonine, and aspartic acid in this order, the PPR motif has a selective RNA base binding ability as follows: binding to G is strong, without binding to A, U, or C.

[0037]

(II) Information about the correspondence of the combination of two amino acids A4 and Lii to the RNA bases

(2-1) If A4 and Lii in this order are asparagine and aspartic acid, the PPR motif has a selective RNA base binding ability as follows: binding to U is the strongest, and binding to C is the second strongest, followed by binding to A and G.

(2-2) If A4 and Lii in this order are asparagine and asparagine, the PPR motif has a selective RNA base binding ability as follows: binding to C is the strongest,

binding to U is the second strongest, followed by binding to A and G.

(2-3) If A4 and Lii in this order are threonine and asparagine, the PPR motif has a selective RNA base binding ability with strong binding to A and weak binding to G, U, and C.

(2-4) If A4 and Lii in this order are threonine and aspartic acid, the PPR motif has a selective RNA base binding ability with strong binding to G and weak binding to A, U, and C.

(2-5) If A4 and Lii in this order are serine and asparagine, the PPR motif has a selective RNA base binding ability as follows: binding to A is the strongest and binding to G, U, and C is the second strongest.

(2-6) If A4 and Lii in this order are glycine and aspartic acid, the PPR motif has a selective RNA base binding ability as follows: binding to G is the strongest, and binding to U is the second strongest, followed by binding to A, without binding to C.

(2-7) If A4 and Lii in this order are asparagine and serine, the PPR motif has a selective RNA base binding ability as follows: binding to C is the strongest, and binding to U is the second strongest, followed by binding to A and G.

(2-8) If A4 and Lii in this order are proline and aspartic acid, the PPR motif has a selective RNA base binding ability as follows: binding to U is the strongest,

and binding to G, C, and C is the second strongest, without binding to A.

(2-9) If A4 and Lii in this order are glycine and asparagine, the PPR motif has a selective RNA base binding ability as follows: binding to A is the strongest, and binding to G is the second strongest, without binding to C or U.

(2-10) If A4 and Lii in this order are methionine and aspartic acid, the PPR motif has a selective RNA base binding ability with strong binding to U and weak binding to A, G, and C.

(2-11) If A4 and Lii in this order are leucine and aspartic acid, the PPR motif has a selective RNA base binding ability as follows: binding to C is the strongest, and binding to U is the second strongest, without binding to A or G.

(2-12) If A4 and Lii in this order are valine and threonine, the PPR motif has a selective RNA base binding ability as follows: binding to U is the strongest, and binding to A is the second strongest, without binding to G or C.

[0038]

[Use of PPR motifs and PPR proteins]

Identification and design:

One PPR motif can recognize a specific base of an RNA. According to the present invention, PPR motifs selective to A, U, G, or C can be selected or designed by

disposing appropriate amino acids in specific positions of a PPR motif. Furthermore, a protein containing an appropriate series of such PPR motifs can recognize its corresponding specific sequence. Moreover, according to the findings described above, a PPR motif which can selectively bind to a desired RNA base and a protein having a plurality of PPR motifs which can sequence-specifically bind to a desired RNA can be designed. In design, the sequence information of a naturally occurring PPR motif may be referred with respect to moieties other than the amino acids disposed in the important positions of the PPR motif. Alternatively, a PPR motif may be designed by using a naturally occurring PPR motif as a whole and replacing only the amino acids in the important positions with other amino acids. The repetition number of the PPR motif can be appropriately determined according to the target sequence; for example, the repetition number can be 2 or more, or 2 to 30.

[0039]

The PPR motif or PPR protein thus designed can be prepared by a method well known to persons skilled in the art. For example, a nucleic acid sequence encoding an amino acid sequence of the designed PPR motif or PPR protein can be determined from the amino acid sequence, and may be cloned to prepare a transformant (such as an expression vector) which produces a desired PPR motif or PPR protein.

[0040]

Preparation and use of fusion protein:

The present invention relates to a fusion protein of the PPR motif or PPR protein described above (i.e., a polypeptide which can bind RNA base-selectively or RNA base sequence-specifically to the target mRNA) and one or more functional domains which improve a protein expression level from an mRNA.

[0041]

The "functional domain which improves a protein expression level from an mRNA" which can be used in the present invention may be all or functional part of a functional domain of a known protein which directly or indirectly promotes the translation of the mRNA, for example. More specifically, the functional domain which can be used in the present invention may be a domain which guides ribosome to the mRNA, a domain associated with initiation or promotion of translation of the mRNA, a domain associated with nuclear export of the mRNA, a domain associated with binding to an endoplasmic reticulum membrane, a domain containing an endoplasmic reticulum retention signal (ER retention signal) sequence, or a domain containing an endoplasmic reticulum signal sequence, for example.

[0042]

More specifically, the domain which guides ribosome to the mRNA may be a domain containing all or functional

part of a polypeptide selected from the group consisting of DENR (Density-regulated protein), MCT-1 (Malignant T-cell amplified sequence 1), TPT1 (Translationally-controlled tumor protein), and Lerepo4 (Zinc finger CCCH-domain). The domain associated with initiation or promotion of translation of the mRNA may be a domain containing all or functional part of a polypeptide selected from the group consisting of eIF4E and eIF4G. The domain associated with nuclear export of the mRNA may be a domain containing all or functional part of SLBP (Stem-loop binding protein). The domain associated with binding to an endoplasmic reticulum membrane may be a domain containing all or functional part of a polypeptide selected from the group consisting of SEC61B, TRAP-alpha (Translocon associated protein alpha), SR-alpha, Dial (Cytochrome b5 reductase 3), and p180. The endoplasmic reticulum retention signal (ER retention signal) sequence may be a signal sequence containing a KDEL (KEEL) sequence. The endoplasmic reticulum signal sequence may be a signal sequence containing MGWSCIIILFLVATATGAHS (SEQ ID NO: 22).

[0043]

In the fusion protein according to the present invention, the functional domain may be fused to the N-terminal side of the PPR protein, may be fused to the C-terminal side of the PPR protein, or may be fused to both of the N-terminal side and the C-terminal side thereof.

Moreover, the fusion protein according to the present invention may include several functional domains (for example, 2 to 5 functional domains). Furthermore, in the fusion protein according to the present invention, the functional domain and the PPR protein may be indirectly fused via a linker, for example.

[0044]

The present invention also relates to a nucleic acid encoding the fusion protein described above, and a vector (such as an expression vector) comprising the nucleic acid. The expression vector herein refers to, for example, a vector comprising a DNA having a promoter sequence, a DNA encoding a desired protein, and a DNA having a terminator sequence, in this order from upstream. The expression vector may not have these DNAs in this order as long as it demonstrates desired functions. A variety of expression vectors which can be usually used by persons skilled in the art can be used in the present invention.

[0045]

Because the fusion protein according to the present invention uses the RNA translation mechanism of eukaryotes, it can function in cells of eukaryotes (such as animals, plants, microorganisms (e.g., yeasts), and protists). The fusion protein according to the present invention can function within animal cells (in vitro or in vivo) in particular. Examples of animal cells into

which the fusion protein according to the present invention or a vector which expresses the fusion protein according to the present invention can be introduced can include cells derived from human, monkey, pig, cow, horse, dog, cat, mouse, and rat. Examples of cultured cells into which the fusion protein according to the present invention or a vector which expresses the fusion protein according to the present invention can be introduced can include, but should not be limited to, Chinese hamster ovarian (CHO) cells, COS-1 cells, COS-7 cells, VERO (ATCC CCL-81) cells, BHK cells, dog kidney-derived MDCK cells, hamster AV-12-664 cells, HeLa cells, WI38 cells, 293 cells, 293T cells, and PER.C6 cells.

[0046]

The terms used herein excluding those particularly defined are used for illustration of the specific embodiments, and are not intended to be limitative to the invention.

[0047]

The term "comprise" used herein, unless contexts clearly require different understandings, is intended to express that a described entry (such as a member, a step, a component, or a number) is present, and is intended not to exclude the presence of other entries (such as a member, a step, a component, or a number).

[0048]

Unless otherwise defined, all the terms used herein (including technical terms and scientific terms) have the same meanings as those broadly understood by persons skilled in the art to which the present invention belongs. Unless otherwise clearly defined, the terms used herein should be interpreted as having the meanings consistent to those herein and its related technical field, and should not be interpreted as idealized or excessively formal meanings.

[0049]

Hereinafter, the present invention will be described more in detail with reference to Examples. However, the present invention can be implemented with a variety of aspects, and should not be construed as limitative to Examples described below.

[Examples]

[0050]

Example 1: Improvement in protein expression level from target mRNA by fusion protein of PPR motif and eIF4G

[0051]

Materials

(Equipment)

- Basic facility for molecular biological experiment (for construction of plasmids, for example)

- Inverted microscope (DM IL S40, Leica Microsystems, Wetzlar, Germany)

- CO₂ incubator (KM-CC17RH2, Panasonic Healthcare, Tokyo, Japan)

- Clean bench (MHE-S1300A2, Panasonic Healthcare, Tokyo, Japan)

- Aspirator (SP-30, Air Liquide Medical Systems, Bovezzo BS, Italy)

- Centrifuge (swing rotor) (LC-200, Tomy Seiko, Tokyo, Japan)

- Ultra-low temperature freezer (-80°C) (MDF-C8V, Panasonic Healthcare, Tokyo, Japan)

- plate reader (EnSight Kaleido, PerkinElmer, Waltham, MA, USA)

[0052]

(Cell culturing)

- HEK293T cell line (see note 1)

- Dulbecco's modified Eagle's culture medium (DMEM, glucose-rich) (see note 2)

- 100× penicillin-streptomycin solution

- Fetal bovine serum (FBS) (see note 3)

- EDTA-NaCl solution: 10 mM EDTA and 0.85% (w/v) NaCl, pH adjusted to 7.2 to 7.4, autoclave sterilized, stored at room temperature

- 100 × 20 mm cell culture petri dish (Greiner bio one, Frickenhausen, Germany)

- 10 mL disposable sterilized pipette

- 15 mL and 50 mL plastic centrifuge tubes

- 1.8 mL cryotube (Nunc; Thermo Fisher Scientific, Waltham, MA, USA)

- Freeze container (Nalgene; Thermo Fisher Scientific, Waltham, MA, USA)

- Bambanker (Lymphotec, Tokyo, Japan)

[0053]

(Transfection)

- Effector plasmid: pcDNA3.1 (Thermo Fisher Scientific, Waltham, MA, USA) was used as a basic vector. A fusion gene of PPR and eIF4G is inserted into an expression cassette (100 ng/ μ L) (see note 4).

- Reporter plasmid: pcDNA3.1 (Thermo Fisher Scientific, Waltham, MA, USA) was used as a basic vector. Luciferase genes are inserted into an expression cassette, and a PPR-binding sequence is inserted into its 5'-UTR (100 ng/ μ L).

- 96-well plate coated with poly-L-lysine (AGC Techno glass, Shizuoka, Japan)

- 1x phosphate-buffered saline, PBS(-): 1.47 mM KH_2PO_4 , 8.1 mM Na_2HPO_4 , 137 mM NaCl, and 2.7 mM KCl. pH adjusted to 7.4, autoclave sterilized, stored at room temperature

- Hemocytometer (for counting the number of cells) (Improved Neubauer Type Cell counter plate, Watson, Hyogo, Japan)

- Transfection reagent (HilyMax, Dojindo Molecular Technologies, Kumamoto, Japan)

[0054]

(Luciferase assay)

- Dual-Glo Luciferase Assay System (Promega, Madison, WI, USA.)

- 96-well luminometer plate (PerkinElmer, Waltham, MA, USA).

[0055]

Experimental method

(Construction of vector)

The reporter assay requires an effector plasmid and a reporter plasmid. These two plasmids both are constructed based on pcDNA3.1. The effector plasmid includes a fusion gene encoding a PPR protein and a partial domain of human eIF4G (SEQ ID NO: 1) (Figure 1A). The PPR protein moiety used was CRR4 (SEQ ID NO: 2). The reporter plasmid includes two open reading frames (ORFs), specifically, renilla luciferase (RLuc) and firefly luciferase (FLuc), which are dicistronically transcribed (Figure 1A). The RLuc gene is located on the side of the 5'-end of the FLuc gene, and was used as a control of gene expression. The PPR-binding region is inserted into the 5'-UTR of the ORF of FLuc, and consists of three repetitions of a CRR4-recognizing sequence (5'-UAUCUUGUCUUUA-3') (SEQ ID NO: 3) interrupted with four-base sequences (ATCG and GATC). To express both of the fused effector gene and the reporter gene, a cytomegalovirus promoter (CMV) and a bovine growth

hormone gene-derived polyadenylation signal were used. For a control experiment, an effector plasmid having no eIF4G was constructed by fusing a FLAG epitope tag to the PPR. A control reporter plasmid without a PPR-binding region was also constructed.

[0056]

The outline of the procedures from cell culturing to the reporter assay in Examples is shown in Figure 2.

[0057]

(Cell culturing from frozen stock)

This step is aseptically performed. All the tools are preliminarily antisepticized with 70% ethanol.

1. A 9 mL DMEM culture medium is placed into a 15 mL centrifuge tube (sterilized).

2. 1 mL of frozen HEK293T cells in a cryotube is incubated within a water bath at 37°C to quickly melt the cells.

3. The cells are placed into the 15 mL centrifuge tube containing 9 mL DMEM.

4. The centrifuge tube is centrifuged at room temperature and 1100 ×g for two minutes, and the supernatant is removed.

5. The cells are resuspended in 10 mL DMEM (FBS is added such that the final concentration is 10%).

6. The suspended cells are transferred into a 100 mm petri dish. The petri dish was left to stand in an incubator at 37°C and under a 5% CO₂ condition. If the

culturing was started from the frozen stock, the cultured cells were subcultured after 24 hours.

[0058]

To keep the cells healthy (see note 5), the cell density on the surface of the petri dish is maintained between 10% and 80%. The passage is basically performed every three days (two times a week), or is performed according to the growth rate of the cells. Furthermore, to keep the number of passages small, cells are freshly cultured from the frozen stock once a month. Keeping the number of passages small and thus keeping the cells healthy are important for efficient DNA transfection.

[0059]

(Passages to maintain cells)

1. New 100 mm petri dishes are provided as required. 8 mL DMEM and 1 mL FBS are preliminarily placed onto each of the petri dishes.

2. The culture medium on a petri dish containing the cultured cells is removed with an aspirator (see note 6).

3. 2 mL EDTA-NaCl solution is gently added onto adhering cells on the surface of the petri dish so as not to peel off the cells. The petri dish is turned around to evenly distribute the solution across the entire surface of the petri dish. The EDTA-NaCl solution is removed with an aspirator. The petri dish is tapped to peel off the cells.

4. 10 mL DMEM is added to the cells in the petri dish, and the cells are suspended by gently pipetting.

5. 1 mL suspended cells (10% cultured cells) are added to the petri dishes preliminarily provided and each containing 9 mL culture medium. Each of the petri dishes is turned around to distribute the cells across the entire surface thereof.

[0060]

(Freeze storage of cells)

A frozen stock is prepared with Bambanker reagent and cultured cells in a logarithmic growth phase at a cell density up to 50%. Use of Bambanker provides a high recovery rate and facilitates long-term storage.

1. The cells on the second day since the passage are peeled off according to the procedure for passage. 5 to 10 mL DMEM is added, and the cells are recovered in a 50 mL centrifuge tube.

2. The centrifuge tube is centrifuged at room temperature and 1100 xg for two minutes, and the supernatant is removed.

3. 1 mL Bambanker per petri dish is added to suspend the cells.

4. The suspended cells are quickly dispensed into cryotubes, and the cryotubes are covered with their lids.

5. The cryotubes are placed in a dedicated freeze container, and are left to stand at -80°C for 12 hours (see note 7).

6. The cryotubes are transferred into a standard sample box, and are stored at -80°C or in liquid nitrogen.
[0061]

(Transient gene introduction (transfection))

1. Before starting transfection, petri dishes each containing the cells on the second day since the passage are provided as required, and the cells are checked whether they are healthy (normal) or not (see note 8). About 96 assays can be performed with one petri dish as an estimate.

2. The cells on the second day since the passage are peeled off according to the procedure for passage, and the suspended cells are transferred into a 50 mL centrifuge tube.

3. The centrifuge tube is centrifuged at room temperature and $1100 \times g$ for two minutes, and the supernatant is removed.

4. Cell clusters are completely dispersed in 10 mL DMEM (FBS is added such that the final concentration is 10%).

5. The number of cells is counted with a hemocytometer and an inverted microscope. The cells are suspended in an appropriate amount of DMEM (FBS is added such that the final concentration is 10%) such that the number of cells is $1 \text{ to } 2 \times 10^5$ cells/mL.

6. A 96-well plate is provided. $200 \mu\text{L}$ ($2 \text{ to } 4 \times 10^4$ cells/mL) per well of suspended cultured cells is placed

into each well, and the plate is left to stand overnight in an incubator at 37°C under a 5% CO₂ condition. One well is used for one assay.

7. On the next day, the culture medium is carefully removed from each well, and is replaced with 100 µL of new DMEM (FBS is added such that the final concentration is 10%).

8. 400 ng effector plasmid (4 µL of 100 ng/µL) and 100 ng reporter plasmid (1 µL of 100 ng/µL) are placed into a single well on a new 96-well PCR plate (or a 0.2 mL tube).

9. For one assay, 1 µL HilyMAX is diluted with 10 µL serum-free DMEM.

10. 11 µL diluted solution is placed into each of the wells containing the plasmids. The solution is well mixed with the plasmids by pipetting.

11. The solution is left to stand at room temperature for 15 minutes. The total amount of the mixture is placed into the wells containing the cultured cells. The plate is left to stand in an incubator at 37°C under a 5% CO₂ condition for 24 hours.

[0062]

(Luciferase assay)

The dual luciferase assay is performed using Dual-Glo Luciferase Assay System according to the usage instruction from the manufacturer except for a few modifications.

1. After 24 hours from the transfection, the culture medium of each well is replaced with 40 μ L 1 $\times$ PBS(-).

2. 40 μ L of Dual-Glo luciferase reagent is placed into each well, and is well mixed with the culture medium by pipetting.

3. The mixture is left to stand at room temperature for 10 minutes, and the total amount thereof is transferred into a 96-well luminometer plate.

4. The light emission from firefly luciferase related with expression of FLuc gene is measured with a plate reader.

5. A Stop & Glo substrate is 100-fold diluted with a Dual-Glo Stop & Glo buffer. 40 μ L of the diluted solution is added into each well.

6. The plate is left to stand at least at room temperature for 10 minutes, and then the light emission from renilla luciferase related with expression of RLuc gene is measured.

[0063]

(Data analysis)

1. The value of FLuc/RLuc is calculated to correct a difference in transfection efficiency between the assays or experimental errors.

2. An increase in activity of reporter gene expression is determined in the presence of the PPR-binding region and in the absence thereof by dividing an experimental value obtained using the plasmid according

to the present invention (plasmid encoding a fusion protein of CRR4 and a translation activation domain eIF4G) by an experimental value obtained using a control plasmid (plasmid encoding a fusion protein of CRR4 and FLAG-tag).

[0064]

Experimental results

The results of the luciferase assay are shown in Figure 3. As shown in Figure 3, 2.75-fold translation activity was specifically verified in the presence of both of the PPR-eIF4G and the PPR-binding sequence. That is, it is demonstrated that the fusion protein of the PPR protein and the functional domain which improve a protein expression level from an mRNA improves the protein expression level from the target mRNA.

[0065]

Notes

(Note 1) HEK293T is a human fetus-derived kidney cell line which expresses an SV40 large T antigen. The cell line is readily cultured, and can be transfected with high efficiency by a variety of methods. HEK293T cells are available from RIKEN BRC (ja.brc.riken.jp) or ATCC (www.atcc.org).

(Note 2) 1x penicillin-streptomycin solution is added to DMEM to avoid contamination with microorganisms.

(Note 3) Before use, FBS is inactivated at 56°C for 30 minutes, and is stored at 4°C.

(Note 4) The purity of the plasmid is significantly important to the transfection efficiency. The plasmid should be isolated using a kit of a transfection grade.

(Note 5) A daily growth rate is an index indicating that the cells are healthy. To avoid suppression of cell growth, the cells should be always cultured in a sufficient space under a sufficient nutritional condition.

(Note 6) HEK293T cells should be gently treated when the culture medium is replaced because the cells are readily peeled off from the culturing petri dish.

(Note 7) The dedicated freeze container is a box whose freezing speed can be adjusted (about -1°C per minute at -80°C), and enables the cells to be freeze stored in a non-programmable -80°C freezer.

(Note 8) In transfection, cells are used at a culture density of 50 to 80%. However, an appropriate cell density depends on the transfection reagent.

Additionally, the ratio of the transfection reagent (μL) to the plasmid DNA (μg) should be also optimized according to the usage instructions from the manufacturer.

The procedure described herein is optimized for a condition where a 96-well plate, HEK293T cells, and HilyMAX as a transfection reagent are used.

[0066]

Example 2: Improvement in protein expression level from target mRNA by fusion protein of PPR and another functional domain

[0067]

In the case where useful substances are produced using cells, the amounts of protein synthesized by endogenous genes and exogenous genes should be precisely controlled. The final amount of the synthesized protein is determined by the insertion positions of genes, the mRNA transcription amount, post-transcriptional regulation (regulation at an RNA level), post-translational modification, and the like. For these reasons, the present inventors have devised a method of enhancing the translation of mRNAs taking advantage of the fact that a PPR protein sequence-specifically binds to a target RNA molecule (Figure 4). In the translation of mRNAs in eukaryotes, an mRNA undergoes action of a translation initiation factor (eukaryotic initiation factor; eIF). As a result, the ribosome is recruited near the translation starting point, and then the translation of the mRNA is started. In other words, the present inventors have considered that the translation of the mRNA can be artificially enhanced if the ribosome can only be recruited onto the mRNA. Moreover, the translation of an mRNA into a protein is usually performed in the ER. For this reason, the present inventors have considered that the translation of the mRNA can be enhanced by intentionally localizing the target mRNA into the ER.

[0068]

Verification by experiment

To verify the idea above, a reporter assay system using animal cultured cells (HEK293T) was prepared (the experiment was performed by the same method as that in Example 1 except that different functional domains were used). The system was constructed using CRR4 protein (one of *Arabidopsis thaliana* PPR proteins), which is known to bind to a specific RNA sequence (UAUCUUGUCUUUA) (SEQ ID NO: 3). First, a fusion protein expression vector (effector plasmid) of CRR4 and a candidate protein functional domain was prepared. The selected candidate domains were (a) eIF proteins (eIF4E and eIF4G), (b) ribosome-bound proteins (DENR, MCT-1, TPT1, and Lerepo4), (c) translational regulation factors (SLBPs) of Histone which promote transport of the transcribed mRNA from the nucleus to the cytoplasm, (d) ER anchor proteins (SEC61B, TRAP-alpha, SR-alpha, Dial, and p180), (e) ER retention signal (KDEL), and (f) ER signal peptide. The fusion proteins were cloned so as to express in the form of HA-CRR4-XX or XX-CRR4-HA (HA: epitope tag (SEQ ID NO: 4); XX: candidate domain).

[0069]

The reporter plasmid included an expression cassette where renilla luciferase (RLuc) and firefly luciferase (FLuc) are transcribed in the form of a dicistronic mRNA under the control of a CMV promoter. Three PPR-binding

sequences (UAUCUUGUCUUUA) (SEQ ID NO: 3) are inserted into a site on the 5'-end of Fluc.

[0070]

The effector plasmid and the reporter plasmid were transfected into HEK293T cells, and the intensities of light emission from RLUC and FLUC were measured. The intensity of light emission from RLUC was treated as a transfection control, and the value of the intensity of light emission from FLUC/the intensity of light emission from RLUC was treated as a translation activity amount.

[0071]

Results

The results shown in Figures 5 and 6 were examined using the following indices (A) and (B).

(A) Comparison between the absence of and the presence of the target

The comparison shows an amount of sequence-specific change in translation.

(B) Comparison to the presence of the target and the absence of the effector (empty) (black dashed line)

The comparison shows an amount of change in translation caused by addition of the domain.

[0072]

1. eIF4E was fused to the C-terminal side of CRR4.

(A) 2.7 times

(B) 1.6 times

2. eIF4G was fused to the C-terminal side of CRR4.

(A) 4.5 times

(B) 3.3 times

3. DENR was fused to the N-terminal side of CRR4.

(A) 1.7 times

(B) 1.3 times

4. DENR was fused to the C-terminal side of CRR4.

(A) 2.4 times

(B) 1.7 times

5. MCT-1 was fused to the N-terminal side of CRR4.

(A) 1.3 times

(B) 1.0 time

6. MCT-1 was fused to the C-terminal side of CRR4.

(A) 2.0 times

(B) 1.2 times

7. TPT-1 was fused to the N-terminal side of CRR4.

(A) 1.4 times

(B) 1.0 time

8. TPT-1 was fused to the C-terminal side of CRR4.

(A) 2.4 times

(B) 1.9 times

9. Lerepo4 was fused to the N-terminal side of CRR4.

(A) 3.0 times

(B) 1.8 times

10. Lerepo4 was fused to the C-terminal side of CRR4.

(A) 3.3 times

(B) 2.6 times

11. SLBP was fused to the C-terminal side of CRR4.

(A) 4.1 times

(B) 3.3 times

12. Sec61B was fused to the C-terminal side of CRR4.

(A) 1.6 times

(B) 1.6 times

13. Sec61BTM was fused to the C-terminal side of CRR4.

(A) 2.4 times

(B) 1.9 times

14. TRAP-alpha was fused to the C-terminal side of CRR4.

(A) 3.5 times

(B) 4.5 times

15. TRAPTM was fused to the C-terminal side of CRR4.

(A) 2.3 times

(B) 1.6 times

16. SR-alpha was fused to the N-terminal side of CRR4.

(A) 1.7 times

(B) 1.5 times

17. DialTM was fused to the N-terminal side of CRR4.

(A) 1.8 times

(B) 1.2 times

18. P180TM2R was fused to the N-terminal side of CRR4.

(A) 2.1 times

(B) 1.5 times

19. P180TMH was fused to the N-terminal side of CRR4.

(A) 2.3 times

(B) 2.5 times

20. P180TM2 was fused to the N-terminal side of CRR4.

(A) 3.0 times

(B) 2.1 times

21. KDEL was fused to the C-terminal side of CRR4.

(A) 1.8 times

(B) 1.4 times

22. KEEL was fused to the C-terminal side of CRR4.

(A) 2.3 times

(B) 2.1 times

23. Signal peptide (SP) was fused to the N-terminal side of CRR4.

(A) 1.4 times

(B) 2.0 times

[0073]

As shown above, an increase in translation was found in all the functional domains in both of the indices (A) and the targets (B). Namely, it was clearly shown that the fusion protein according to the present invention can enhance the translation of the target mRNA.

[0074]

The amino acid sequences of the functional domains used in Examples are listed below:

[Table 1-1]

Domain	Sequence
eIF4E	MATVEPETTPTPNPPTTEEEKTESNQEVANPEHYIKHPLQNRWALW FFKNDKSKTWQANLRLISKFDTVEDFWALYNHIQLSSNLMPGCDYS LFKDGIEPMLEDEKNKRGGRWLITLNKQQRSDLDLRFWLETLLCLI GESFDDYSDDVCGAVVNVRAKGDKIAIWTTECENREAVTHIGRVYK ERLGLPPKIVIGYQSHADTATKSGSTTKNRFVVGRY (SEQ ID NO: 5)
eIF4G	EEKKRYDREFLLGFQFIFASMQKPEGLPHISDVVLDKANKTFLRPL DPTRLQGINCGPDFTPSFANLGRITLSTRGPPRGGPGGELPRGPQA GLGPRRSQQGPRKEPRKIIATVLMTEDIKLNKAEKAWKPSSKRTAA DKDRGEEDADGSKTQDLFRRVRSILNKLTQPMFQQMLKQVTQLAID TEERLKGVIDLIFEKAISEPNFSVAYANMCRCLMALKVPTTEKPTV TVNFRKLLLNRCQKEFEKDKDDDEVFEKKQKEMDEAATAEERGRK EELEEARDIARRRSLGNIKFIGELFKLKMTEAIMHDCVVKLLKNH DEESLECLCRLLTTIGKDLDFEKAKPRMDQYFNQMEKIIKEKKTSS RIRFMLQDVLDLGRSNWVPRRGDQGPKTIDQIHKEAEMEEREHIK VQQLMAKGS DKRRGGPPGPPI SRGLPLVDDGGWNTVPI SKGSRPID TSRLTKITKPGSIDSNQLFAPGGRLSWGKGSSGGSGAKPSDAASE AARPATSTLNRFSALQQA VPTESDNR RVQRSSLSRERGEKAGDR GDRLE SERGGDRGDR LDRARTPATKRSFSKEVEERSRERPSQPEG LRKAASLTEDRDRGRDAVKREAALPPVSPLKAALSEEELEKSKAI IEEYLHLNDMKEAVQCQVELASPSLLFIFVRHGVSTLERSAIARE HMGQLLHQLLCAGHLSTAQYYQGLYEILELAEDMEIDI PHVWLYLA ELVTPILQEGGVPMGELFREITKPLRPLGKAASLLEILGLLCKSM GPKKVGT LWREAGLSWKEFLPEGQDIGAFVAEQKVEYTLGEESEAP GQRALPSEELNRQLEKLLKEGSSNQRVFDWIEANLSEQQIVSNTLV RALMTAVCYSAII FETPLRVDVAVLKARAKLLQKYLCEQKELQAL YALQALVVTLEQPPNLLRMFFDALYDEDVVKEDAFYSWESSKDPAE QQGKGVALKSVTAFFKWLREAEESDH (SEQ ID NO: 1)
DENR	MAADISESSGADCKGDPRNSAKLDADYPLRVLYCGVCSLPTEYCEY MPDVAKCRQWLEKNFPNEFAKLTVENSPKQEAGISEGQGTAGEEEEE KKKQKRGGRGQIKQKKKTVPOKVITAKIPRAKKKYVTRVCGLATFE IDLKEAQRFFAQKFSCGASVTGEDEII IQGDFTDDIIDVIQEKWPE VDDDSIEDLGEVKK (SEQ ID NO: 6)
MCT-1	MFKKFDEKENVSNCIQLKTSVIKGIKNQLIEQFPGIEPWLNQIMPK KDPVKIVRCHEHIEILTVNGELLFFRQREGPFYPTLRLLHKYPFIL PHQQVDKGAIKFVLSGANIMCPGLTSPGAKLYPAAVDITIVAIMAEG

[Table 1-2]

	KQHALCVGVMKMSAEDIEKVNKGIGIENIHYLNDGLWHMKT (SEQ ID NO: 7)
TPT-1	MIYRDLISHDEMFSDIYKIREIADGLCLEVEGKMVSRT EGNIDDS LIGGNASAEGPEGEGETESTVITGVDIVMNHHLQETSFTKEAYKKYI KDYMKSIKGGLEEQRPVVKPFMTGAAEQIKHILANFKNYQFFIGE NMNPDGMVALLDYREDGVTPYMIFFKDGLEMEKC (SEQ ID NO: 8)
Lerepo4	PPKKQAQAGGSKKAEQKKKEKIEEDKTGFLKNKKGAKQOKFIKAVT HQVKFGQONPRQVAQSEAEKKLKKDDKKELQELNELFKPVVAAQK ISKGADPKSVVCAFFKQGQCTKGDCKKFSHDLTLERKCEKRSVYID ARDEELEKDTMDNWDEKKLEEVVNKKHGEAEKKPKTQIVCKHFLE AIENNKYGFVWVCPGGGDI CMYRHALPPGFVLKKDKKKEEKEDEIS LEDLIERERSALGPNVTKITLESFLAWKKRKRQEKIDKLEQDMERR KADFKAGKALVISGREVFEPPELVNDDDEEADDTRYTQGTGGDEV DDSVSVNDIDLSLYIPRDVDETGITVASLERFSTYTSKDKENKLSE ASGGRAENGERSDLEEDNEREGTENGAI DAVPVDEKSFHWRGFG (SEQ ID NO: 9)
SLBP	ACRPRSPRHQSRCDGASPPSPARWSLGRKRRADGRRWRPEDAEE AEHRGAERRPESFTTPEGPKPRSRCSDWASAVEEDEMRTVRNEMA RYKRKLLINDFGRERKSSSGSSDSKESMSTVPADFETDESVLMMRQ KQINYGKNTIAYDRIYKEVPRHLRQPGIHPKTPNKFKKYSRRSWDQ QIKLWKVALHFWDPPAEBCDLQEIHPVDLESAESSSEPQTSSQDD FDVYSGTPTKVRHMSQVEDEFDLEACLTEPLRDFSAMS (SEQ ID NO: 10)
Sec61B	PGPTPSGTNVGSSGRSPSKAVAARAAGSTVRQRKNASCGRSAGRT TSAGTGGMWRFYTEDSPGLKVGVPVVLVMSLLFIASVFMLHIWGY TRS (SEQ ID NO: 11)
Sec61B-TM	VGPVPVLVMSLLFIASVFMLHIW (SEQ ID NO: 12)
TRAP-alpha	RLLPRLLLLLLVFPATVLFRRGGPRGLLAVAQDLTEDEETVEDSII EDEDDEAEVEEDEPTDLVEDKEEEDVSGEPEASPSADTTILFVKGE DFPANNIVKFLVGFTNKGTEDFIVESLDASFRYPQDYQFYIQNFTA LPLNTVVPQRQATFEYSFIPAEPMGGRPFGLVINLNYKDLNGNVF QDAVFNQTVTVIEREDGLDGETIFMYMFLAGLGLLVIVGLHQLLES RKRKRPIQKVEMGTSSQNDVDMSWIPQETLNQINKASPRRLPRKRA QKRSGSDE (SEQ ID NO: 13)
TRAP-TM	TIFMYMFLAGLGLLVIVGLHQLL (SEQ ID NO: 14)
SR-alpha	LDFFTIFSKGGLVLWCFOGVSDSCTGPVNALIRSVLLQVGFQKILT

[Table 1-3]

	LTYVDKLIDDVHRLFRDKYRTEIQQQSALSLLNGTFDFQNDFLRL REAEBSKIRAPTTMKKFEDSEKAKKPVRSMIETRGEKPKKAKNS KKKGAKKEGSDGPLATSKPVPABKSGLPVGPENGVELSKEELIRRK REEFIQKHGRGMEKSNKSTKSDAPKEKGKKAPRVWELGGCANKEVL DYSTPTTNGTPEAALSEDINLIRGTGSGGQLQDLDCSSSDDEGAAQ NSTKPSATKGTLGGMFGMLKGLVGSKSLSRDMESVLDKMRDHLIA KNVAADIAVQLCESVANKLEBKVMGTFSTVTSTVKQALQESLVQIL QPQRRVDMLRDIMDAQRRQRPYVVTFCGVNGVGKSTNLAKISFWLL ENGFSVLIAACDTFRAGAVEQLRTHTRRLSALHPPEKHGGRTMVQL FEKGYGKDAAGIAMEAIAFARNQGFDVVLVDTAGRMQDNAPLMTAL AKLITVNTFPLVLFVGEALVGNEAVDQLVKFNRALADHSMAQTPLR IDGIVLTKFDTIDDKVGAAISMTYITSKPIVFVGTGQTYCDLRLN AKAVVAALMKA (SEQ ID NO: 15)
DiaTM	STLGHMVLFPVWFLYSLL (SEQ ID NO: 16)
P180TMR2	DIYDTQTLGVVVFGGFMVVSAGIFLVSTFSPMKETSYEEALANQRK EMAKTHHQKVEKKKKEKTVEKKGKTKKKBEKPNGKIPDHDPAPNVT VLLREPVRAPAVAVAPTVPVQPPIIIVAPVATVPAMPQEKCLASSPKDK KKKEKKVAKVEPAVSSVNSIQVLTSKAAILETAPKEGRNTDVAQS PEAPKQEAAPAKKSGSKKKGPPDADGPLYLPYKTLVSTVGSMVFNE GEAQRLEILSEKAGIIQDTHKATQKGDPV(SEQ ID NO: 17)
P180TMH	LGVVVFGGFMVVSAGIFLVSTF (SEQ ID NO: 18)
P180TM2	DIYDTQTLGVVVFGGFMVVSAGIFLVSTF (SEQ ID NO: 19)
KDEL	KDEL (SEQ ID NO: 20)
KEEL	KEEL (SEQ ID NO: 21)
ER signal peptide	MGWSCIIILFLVATATGAHS (SEQ ID NO: 22)

[Document Name] Claims

[Claim 1]

A fusion protein for improving a protein expression level from a target mRNA, the fusion protein comprising:

(A) one or more functional domains which improve a protein expression level from an mRNA; and

(B) a polypeptide moiety which can bind to a target mRNA in an RNA base-selective or RNA base sequence-specific manner,

wherein polypeptide moiety (B) is a polypeptide moiety comprising one or more PPR motifs, each PPR motif comprising a polypeptide consisting of 30 to 38 amino acids in length and being represented by Formula 1:

[Formula 1]

(Helix A)-X-(Helix B)-L (Formula 1)

where

Helix A is a moiety which consists of 12 amino acids in length and can form an α -helix structure, and is represented by Formula 2:

[Formula 2]

A₁-A₂-A₃-A₄-A₅-A₆-A₇-A₈-A₉-A₁₀-A₁₁-A₁₂ (Formula 2)

where A₁ to A₁₂ each independently represent an amino acid;

X is not present, or is a moiety consisting of 1 to 9 amino acids in length;

Helix B is a moiety which consists of 11 to 13 amino acids in length and can form an α -helix structure;

L is a moiety consisting of 2 to 7 amino acids in length and represented by Formula 3:

[Formula 3]



where the amino acids are numbered from the C-terminal as "i" (-1), "ii" (-2), ... and

L_{iii} to L_{vii} may not be present, and

a combination of three amino acids A_1 , A_4 , and L_{ii} or a combination of two amino acids A_4 and L_{ii} corresponds to a base or base sequence of the target mRNA.

[Claim 2]

The fusion protein according to claim 1, wherein polypeptide moiety (B) comprises 2 to 30 PPR motifs, and the plurality of PPR motifs is arranged so as to specifically bind to the base sequence of the target mRNA.

[Claim 3]

The fusion protein according to claim 2, polypeptide moiety (B) comprises 5 to 25 PPR motifs.

[Claim 4]

The fusion protein according to any one of claims 1 to 3, wherein one or more functional domains (A) each

bind to an N-terminal and/or a C-terminal of polypeptide moiety (B).

[Claim 5]

The fusion protein according to any one of claims 1 to 4, wherein one or more functional domains (A) are selected from the group consisting of a domain which guides ribosome to the mRNA, a domain associated with initiation or promotion of translation of the mRNA, a domain associated with nuclear export of the mRNA, a domain associated with binding to an endoplasmic reticulum membrane, a domain containing an endoplasmic reticulum retention signal (ER retention signal) sequence, and a domain containing an endoplasmic reticulum signal sequence.

[Claim 6]

The fusion protein according to claim 5,

wherein the domain which guides ribosome to the mRNA is a domain containing all or functional part of a polypeptide selected from the group consisting of DENR (Density-regulated protein), MCT-1 (Malignant T-cell amplified sequence 1), TPT1 (Translationally-controlled tumor protein), and Lerepo4 (Zinc finger CCCH-domain),

the domain associated with initiation or promotion of translation of the mRNA is a domain containing all or functional part of a polypeptide selected from the group consisting of eIF4E and eIF4G,

the domain associated with nuclear export of the mRNA is a domain containing all or functional part of SLBP (Stem-loop binding protein),

the domain associated with binding to an endoplasmic reticulum membrane is a domain containing all or functional part of a polypeptide selected from the group consisting of SEC61B, TRAP-alpha (Translocon associated protein alpha), SR-alpha, Dial (Cytochrome b5 reductase 3), and p180,

the endoplasmic reticulum retention signal (ER retention signal) sequence is a signal sequence containing a KDEL (KEEL) sequence, or

the endoplasmic reticulum signal sequence is a signal sequence containing MGWSCIIILFLVATATGAHS.

[Claim 7]

The fusion protein according to any one of claims 1 to 6, wherein the combination of the three amino acids A₁, A₄, and L_{ii} in each of the PPR motifs is:

(valine, threonine, asparagine), (phenylalanine, serine, asparagine), (phenylalanine, threonine, asparagine), (isoleucine, asparagine, aspartic acid), or (threonine, threonine, asparagine) in order of (A₁, A₄, L_{ii}) if a target base for the PPR motif is A (adenine);

(glutamic acid, glycine, aspartic acid), (valine, threonine, aspartic acid), (lysine, threonine, aspartic acid), or (leucine, threonine, aspartic acid) in order of

(A₁, A₄, L_{ii}) if the target base for the PPR motif is G (guanine);

(valine, asparagine, aspartic acid), (isoleucine, asparagine, asparagine), (isoleucine, asparagine, aspartic acid), (isoleucine, methionine, aspartic acid), (phenylalanine, proline, aspartic acid), or (tyrosine, proline, aspartic acid) in order of (A₁, A₄, L_{ii}) if the target base for the PPR motif is U (uracil); or

(valine, asparagine, asparagine), (isoleucine, asparagine, asparagine), (valine, asparagine, serine), or (isoleucine, methionine, aspartic acid) in order of (A₁, A₄, L_{ii}) if the target base for the PPR motif is C (cytosine).

[Claim 8]

The fusion protein according to any one of claims 1 to 6, wherein the combination of the two amino acids A₄ and L_{ii} in each of the PPR motifs is:

(threonine, asparagine), (serine, asparagine), or (glycine, asparagine) in order of (A₄, L_{ii}) if a target base for the PPR motif is A (adenine);

(threonine, aspartic acid) or (glycine, aspartic acid) in order of (A₄, L_{ii}) if the target base for the PPR motif is G (guanine);

(asparagine, aspartic acid), (proline, aspartic acid), (methionine, aspartic acid), or (valine, threonine) in order of (A₄, L_{ii}) if the target base for the PPR motif is U (uracil); or

(asparagine, asparagine), (asparagine, serine), or (leucine, aspartic acid) in order of (A₄, L_{ii}) if the target base for the PPR motif is C (cytosine).

[Claim 9]

A nucleic acid encoding the fusion protein according to any one of claims 1 to 8.

[Claim 10]

A vector comprising the nucleic acid according to claim 9.

[Claim 11]

The vector according to claim 10, wherein the vector is an expression vector.

[Claim 12]

A method of improving a protein expression level from a target mRNA within a cell, the method comprising:

a step of providing the fusion protein according to any one of claims 1 to 8 or the vector according to claim 10 or 11; and

a step of introducing the fusion protein or the vector into a cell.

[Claim 13]

The method according to claim 12,
wherein the cell is a eukaryotic cell.

[Claim 14]

The method according to claim 13,
wherein the cell is an animal cell.

[Claim 15]

The method according to claim 14,
wherein the animal cell is a human cell.