[Claim 41]

An instruction, a manual, an appendix, or a product label, indicating that methyl 3-[(4S)-8-bromo-1-methyl-6-(2-pyridinyl)-4H-imidazo[1,2-a][1,4]benzodiazepin-4-yl]propanoate benzenesulfonate can be used in combination with a narcotic analgesic and optionally a muscle relaxant for the induction of general anesthesia by intravenous administration at a dosing rate of about 6 mg or about 12 mg mg/hour/kg in a patient who needs a treatment with the compound.

[Claim 42]

A method which is a sedative method including intravenously administering an effective amount of methyl 3-[(4S)-8-bromo-1-methyl-6-(2-pyridinyl)-4H-imidazo[1,2-a][1,4]benzodiazepin-4-yl]propanoate or a salt thereof, which is an active ingredient, to a patient who needs a treatment with the compound, wherein the dosing rate of the active ingredient by intravenous administration is about 0.3 mg to about 40 mg/hour/kg in a patient who needs a treatment with the compound.

[Claim 43]

Methyl

[Claim 41]

An instruction, a manual, an appendix, or a product label, indicating that methyl 3-[(4S)-8-bromo-1-methyl-6-(2-pyridinyl)-4H-imidazo[1,2-a][1,4]benzodiazepin-4-yl]propanoate benzenesulfonate can be used in combination with a narcotic analgesic and optionally a muscle relaxant for the induction of general anesthesia by intravenous administration at a dosing rate of about 6 mg or about 12 mg mg/hour/kg in a patient who needs a treatment with the compound.

[Claim 42]

The use of an effective amount of methyl 3-[(4S)-8-bromo-1-methyl-6-(2-pyridinyl)-4H-imidazo[1,2-a][1,4]benzodiazepin-4-yl]propanoate or a salt thereof in the manufacture of a sedative to treat the state of consciousness of a patient intravenously, wherein the dosing rate of the active ingredient by intravenous administration is about 0.3 mg to about 40 mg/hour/kg.

[Claim 43]

Methyl