
CLAIMS

1. Composition comprising at least one 7β-hydroxycholesterol derivative and a lipid vehicle, wherein said lipid vehicle comprises a vegetable oil or a mixture of vegetable oils, in which the 7β-hydroxycholesterol derivative corresponds to formula (I)



(I)

in which:

- A represents

a -C(O)R_1 group in which R_1 is a saturated heterocycle comprising 5 members and including 2oxygen atoms, substituted with at least one linear or branched C_1-C_6 alkyl;

- B represents a -C(O)R_4 group, in which R_4 is a linear or branched C_1-C_6, alkyl.

2. Composition according to claim 1, in which said vehicle comprises at least one lipid other than a phospholipid.


3. Composition according to claims 1 or 2, characterized in that it is in a form suitable for administration by the oral route.

4. Composition according to any one of claims 1 to 3, characterized in that the compound of formula (I) is in solution in said lipid vehicle.

5. Composition according to any one of claims 1 to 4, in which said lipid vehicle comprises a vegetable oil selected from argan oil, avocado oil, linseed oil, sunflower oil, palm oil, cabbage palm oil, coconut oil, grapeseed oil, black mustard oil, poppy seed oil, shea butter oil, sweet almond oil, soya oil, peanut oil, cottonseed oil, sesame oil, olive oil, maize oil, cocoa oil, castor oil, Moringa oil (or Ben oil), rapeseed oil, annatto oil, wheatgerm oil, safflower oil, walnut oil, hazelnut oil, turnip rape oil or mixtures thereof.

6. Composition according to any one of claims 1 to 5, in which said lipid vehicle is argan oil.

7. Composition according to any one of claims 1 to 6, in which the 7β-hydroxycholesterol derivative is a compound of formula (I) in which A is a -C(O)R_1 group in which R_1 is a 2,2-dimethyl-1,3-dioxolane group and B is an acetyl group, of formula



.

8. Composition according to any one of claims 1 to 7, in which the content of compound of formula (I) is from 0.1 to 3.5% (w/v) of the total volume of the composition.

9. Composition according to any one of claims 1 to 8, characterized in that the content of compound of formula (I) is from 1 to 35 mg/ml, preferably 30 mg/ml.

10. Composition according to any one of claims 1 to 9, characterized in that the content of lipid vehicle is from 90 to 99% (v/v) of the total volume of the composition.

11. Pharmaceutical composition comprising a composition according to any one of claims 1 to 10.

12. Pharmaceutical composition according to claim 11, characterized in that it is in a form suitable for administration by the oral route.

13. Method for preparing a composition according to any one of claims 1 to 8, comprising the following steps:

- preparing a mixture containing the 7β-hydroxycholesterol derivative of formula (I) as defined in claim 1 or 7, a lipid vehicle and a co-solvent,

- optionally adding an antioxidant and

- if applicable, evaporating the co-solvent.

14. Method according to claim 13, wherein the lipid vehicle is as defined in any one of claim 2 or 5 to 6.

15. Composition according to any one of claims 1 to 12, for use in the treatment of a neoplastic pathology or a malignant haemopathy.

16. Composition for use according to claim 15, in which the neoplastic pathology is glioblastoma multiforme.