
CLAIMS

1. Compound of general formula **(I)** or a pharmaceutically acceptable salt or derivative thereof



(I)

wherein R_2 and R_3 , independently from each other, are selected from:

a) -C_1-C_6 alkyl, with the proviso that R_2 and R_3 are not both methyl,
b) -$(CH_2)_n$-aryl, 4-halo-benzyl or 4-halo-phenyl ,
c) -$(CH_2)_n$-cycloalkyl,
d) -$(CH_2)_n$-heterocycle,
e) -$(CH_2)_n$-OR_6,
f) -$(CH_2)_n$-CN,
g) -$(CH_2)_n$-NR_4R_5,
h) taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle containing from one to three heteroatoms selected from the group consisting of oxygen and sulphur, or
i) taken together with the nitrogen atom to which they are bound to form: (i) a 5- to 8-membered heteromonocycle which is fused to one or two saturated or unsaturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur; or (ii) a halo and/or hydroxy substituted dihydroquinoline;

wherein R_4 and R_5, independently from each other, are selected from:

a) -H,
b) -C_1-C_6 alkyl,
c) -$(CH_2)_n$-aryl,
d) -$(CH_2)_n$-cycloalkyl,
e) -$(CH_2)_n$-heterocycle,
f) -$(CH_2)_n$-OR_6,

g) $-(CH_2)_n-CN$,

h) taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur, or

j) taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle which is fused to one or two saturated or unsaturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur; or

wherein R_6 is selected from:

a) -H,

b) $-C_1-C_6$ alkyl,

c) $-(CH_2)_n$-aryl,

d) $-(CH_2)_n$-cycloalkyl, or

e) $-(CH_2)_n$-heterocycle,;

and wherein **n** is an integer from 0 to 6.

2. Compound according to claim 1,

wherein R_2 and R_3, independently from each other, are selected from:

a) $-C_1-C_6$ alkyl, with the proviso that R_2 and R_3 are not both methyl,

b) $-(CH_2)_n$-aryl, 4-halo-benzyl or 4-halo-phenyl,

c) $-(CH_2)_n$-cycloalkyl,

d) $-(CH_2)_n$-heterocycle,

e) $-(CH_2)_n-OR_6$,

f) $-(CH_2)_n-CN$,

g) taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle containing from one to three heteroatoms selected from the group consisting of oxygen and sulphur, or

h) taken together with the nitrogen atom to which they are bound to form: (i) a 5- to 8-membered heteromonocycle which is fused to one or two a unsaturated or saturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur; or (ii) a halo and/or hydroxy substituted dihydroquinoline;

and/or wherein R_6 is H.

3. Compound according to claim 1,

wherein R_2 and R_3, independently from each other, are selected from:

a) $-C_1$-C_6 alkyl, with the proviso that R_2 and R_3 are not both methyl,

b) $-(CH_2)_n$-aryl, 4-halo-benzyl or 4-halo-phenyl or

c) taken together with the nitrogen atom to which they are bound to form: (i) a 5- to 8-membered heteromonocycle which is fused to one or two a unsaturated or saturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur; or (ii) a halo and/or hydroxy substituted dihydroquinoline.

4. Compound according to claim 1, wherein R_2 and R_3 taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle which is fused to one or two unsaturated or saturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur.

5. Compound according to claim 1, wherein **n** is an integer from 0 to 2.

6. Compound selected from the group consisting of:

8-Hydroxy-N-methyl-N-(4-(2-phenylpropan-2-yl)benzyl)quinoline-2-carboxamide;

N-Benzyl-8-hydroxy-N-methylquinoline-7-carboxamide;

(8-Hydroxyquinolin-7-yl)(isoindolin-2-yl)methanone;

(8-Hydroxyquinolin-7-yl)(morpholino)methanone;

(8-Hydroxyquinolin-7-yl)(piperidin-1-yl)methanone;

8-Hydroxy-N-methyl-N-phenethylquinoline-7-carboxamide;

(8-Hydroxyquinolin-7-yl)(indolin-1-yl)methanone;

N-(Furan-2-ylmethyl)-8-hydroxy-N-methylquinoline-7-carboxamide;

(3,4-Dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;

N-(4-Bromobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;

8-Hydroxy-N-(4-methoxyphenyl)-N-methylquinoline-7-carboxamide;

(6,7-Dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)(8-hydroxyquinolin-7-yl)methanone;

8-Hydroxy-N-methyl-N-(tetrahydro-2H-pyran-4-yl)quinoline-7-carboxamide;


8-Hydroxy-N-methyl-N-phenylquinoline-7-carboxamide;

N-(4-Chlorophenyl)-8-hydroxy-N-methylquinoline-7-carboxamide;

N-Ethyl-8-hydroxy-N-phenylquinoline-7-carboxamide;
N-Cyclohexyl-8-hydroxy-N-methylquinoline-7-carboxamide;
8-Hydroxy-N-methyl-N-(1-methylpiperidin-4-yl)quinoline-7-carboxamide;
8-Hydroxy-N-methyl-N-(1-methylpyrrolidin-3-yl)quinoline-7-carboxamide;
N-(2-Cyanoethyl)-N-(furan-2-ylmethyl)-8-hydroxyquinoline-7-carboxamide;
N-(2-Cyanoethyl)-8-hydroxy-N-((tetrahydrofuran-2-yl)methyl)quinoline-7-carboxamide;
N-Ethyl-8-hydroxy-N-methylquinoline-7-carboxamide;
8-Hydroxy-N-methyl-N-propylquinoline-7-carboxamide;
(3,4-Dihydroisoquinolin-2(1H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(5-Bromoindolin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(6-methoxy-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Hydroxyquinolin-7-yl)(5-nitroindolin-1-yl)methanone;
8-Hydroxy-N-phenyl-N-propylquinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(octahydroquinolin-1(2H)-yl)methanone;
N-(4-Fluorobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
N-(3-Bromobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(4-(trifluoromethyl)piperidin-1-yl)methanone;
(4-*tert*-Butylpiperidin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
(S)-8-Hydroxy-N-methyl-N-(1-phenylethyl)quinoline-7-carboxamide;
N-Benzyl-8-hydroxy-N-(2-hydroxyethyl)quinoline-7-carboxamide;
(3,3-Dimethylpiperidin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
N-(2-Bromobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(4-phenylpiperidin-1-yl)methanone;
((4aS,8S,8aR)-8-Hydroxy-octahydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(2-methylpiperidin-1-yl)methanone;
(8-Hydroxyquinolin-7-yl)(2-phenylpiperidin-1-yl)methanone;
(1,1-Dioxo-thiomorpholin-4-yl)-(8-hydroxy-quinolin-7-yl)-methanone;
(8-Hydroxyquinolin-7-yl)(4-methylpiperidin-1-yl)methanone;
(R)-8-Hydroxy-N-(1-phenylethyl)quinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(2-methylpyrrolidin-1-yl)methanone;
(2,5-Dimethylpyrrolidin-1-yl)(8-hydroxyquinolin-7-yl)methanone;

(8-Hydroxyquinolin-7-yl)(3-phenylpyrrolidin-1-yl)methanone;
(3-(Dimethylamino)pyrrolidin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(3-methylpiperidin-1-yl)methanone;
(8-Hydroxyquinolin-7-yl)(pyrrolidin-1-yl)methanone;
(1,4-Dioxa-8-aza-spiro[4.5]dec-8-yl)-(8-hydroxy-quinolin-7-yl)-methanone;
(8-Hydroxyquinolin-7-yl)(6-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Hydroxyquinolin-7-yl)(6-nitroindolin-1-yl)methanone;
(8-Hydroxyquinolin-7-yl)(7-(trifluoromethyl)-3,4-dihydroquinolin-1(2H)-yl)methanone;
(5-Bromo-8-hydroxyquinolin-7-yl)(isoindolin-2-yl)methanone;
(Hexahydro-1H-isoindol-2(3H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(2-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Hydroxyquinolin-7-yl)(3-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Hydroxyquinolin-7-yl)(4-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Fluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(6-isopropyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(6-Chloro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(7-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(6-Bromo-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone.
(8-Hydroxyquinolin-7-yl)(octahydroisoquinolin-2(1H)-yl)methanone;
N-(4-Bromophenyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
(Decahydro-1H-carbazol-9(9aH)-yl)(8-hydroxyquinolin-7-yl)methanone;
N-(4-Chlorobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
(5,7-Difluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(7-Fluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(5-Fluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(6-Fluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(5-Chloro-8-hydroxy-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(2H-Benzo[b][1,4]oxazin-4(3H)-yl)(8-hydroxyquinolin-7-yl)methanone.

7. A compound of formula



8. Compound according to any of the claims from 1 to 7, for use as a medicament.

9. Compound according to any of the claims from 1 to 7, for use as antifungal agent.

10. Compound according to any of the claims from 1 to 7, for use in the treatment and/or prevention of fungal infections.


11. Compound for use according to claim 10, wherein said fungal infection is from Tricophyton Rubrum, Tricophyton Mentagrophytes, Aspergillus Niger, Scopulariopsis Brevicaulis or Candida, such as Candida Albicans or Candida Glabrata.

12. Compound for use according to claim 10, wherein the recipient of said treatment or prevention is a mammalian, preferably a human.

13. A pharmaceutical formulation containing at least a compound according to any one of claims from 1 to 7, together with at least a pharmaceutically acceptable carrier, excipient and/or adjuvant.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Compound of general formula **(I)** or a pharmaceutically acceptable salt or derivative thereof



(I)

wherein R_2 and R_3 , independently from each other, are selected from:

a) -C_1-C_6 alkyl, with the proviso that R_2 and R_3 are not both methyl,

b) -$(CH_2)_n$-aryl, 4-halo-benzyl or 4-halo-phenyl ,

c) -$(CH_2)_n$-cycloalkyl,

d) -$(CH_2)_n$-heterocycle,

e) -$(CH_2)_n$-OR_6,

f) -$(CH_2)_n$-CN,

g) -$(CH_2)_n$-NR_4R_5,

h) taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle containing from one to three heteroatoms selected from the group consisting of oxygen and sulphur, or

i) taken together with the nitrogen atom to which they are bound to form: (i) a 5- to 8-membered heteromonocycle which is fused to one or two saturated or unsaturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur; or (ii) a halo and/or hydroxy substituted dihydroquinoline;

wherein R_4 and R_5, independently from each other, are selected from:

a. -H,

b. -C_1-C_6 alkyl,

c. -$(CH_2)_n$-aryl,

d. -$(CH_2)_n$-cycloalkyl,

e. -$(CH_2)_n$-heterocycle,

f. -$(CH_2)_n$-OR_6,

g. -$(CH_2)_n$-CN,

h. taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur, or

j) taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle which is fused to one or two saturated or unsaturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur; or

wherein $\mathbf{R_6}$ is selected from:

a) -H,

b) -C_1-C_6 alkyl,

c) -$(CH_2)_n$-aryl,

d) -$(CH_2)_n$-cycloalkyl, or

e) -$(CH_2)_n$-heterocycle,;

and wherein **n** is an integer from 0 to 6.

2. Compound according to claim 1,

wherein $\mathbf{R_2}$ and $\mathbf{R_3}$, independently from each other, are selected from:

a) -C_1-C_6 alkyl, with the proviso that R_2 and R_3 are not both methyl,

b) -$(CH_2)_n$-aryl, 4-halo-benzyl or 4-halo-phenyl,

c) -$(CH_2)_n$-cycloalkyl,

d) -$(CH_2)_n$-heterocycle,

e) -$(CH_2)_n$-OR_6,

f) -$(CH_2)_n$-CN,

g) taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle containing from one to three

heteroatoms selected from the group consisting of oxygen and sulphur, or

h) taken together with the nitrogen atom to which they are bound to form: (i) a 5- to 8-membered heteromonocycle which is fused to one or two a unsaturated or saturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur; or (ii) a halo and/or hydroxy substituted dihydroquinoline;

and/or wherein R_6 is H.

3. Compound according to claim 1,

wherein R_2 and R_3, independently from each other, are selected from:

a) -C_1-C_6 alkyl, with the proviso that R_2 and R_3 are not both methyl,

b) -$(CH_2)_n$-aryl, 4-halo-benzyl or 4-halo-phenyl or

c) taken together with the nitrogen atom to which they are bound to form: (i) a 5- to 8-membered heteromonocycle which is fused to one or two a unsaturated or saturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur; or (ii) a halo and/or hydroxy substituted dihydroquinoline.

4. Compound according to claim 1, wherein R_2 and R_3 taken together with the nitrogen atom to which they are bound to form a 5- to 8-membered heteromonocycle which is fused to one or two unsaturated or saturated rings or to other heterocycles containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur.

5. Compound according to claim 1, wherein **n** is an integer from 0 to 2.

6. Compound selected from the group consisting of:


8-Hydroxy-N-methyl-N-(4-(2-phenylpropan-2-yl)benzyl)quinoline-2-carboxamide;

N-Benzyl-8-hydroxy-N-methylquinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(isoindolin-2-yl)methanone;
(8-Hydroxyquinolin-7-yl)(morpholino)methanone;
(8-Hydroxyquinolin-7-yl)(piperidin-1-yl)methanone;
8-Hydroxy-N-methyl-N-phenethylquinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(indolin-1-yl)methanone;
N-(Furan-2-ylmethyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
(3,4-Dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
N-(4-Bromobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
8-Hydroxy-N-(4-methoxyphenyl)-N-methylquinoline-7-carboxamide;
(6,7-Dimethoxy-3,4-dihydroisoquinolin-2(1H)-yl)(8-hydroxyquinolin-7-yl)methanone;
8-Hydroxy-N-methyl-N-(tetrahydro-2H-pyran-4-yl)quinoline-7-carboxamide;
8-Hydroxy-N-methyl-N-phenylquinoline-7-carboxamide;
N-(4-Chlorophenyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
N-Ethyl-8-hydroxy-N-phenylquinoline-7-carboxamide;
N-Cyclohexyl-8-hydroxy-N-methylquinoline-7-carboxamide;
8-Hydroxy-N-methyl-N-(1-methylpiperidin-4-yl)quinoline-7-carboxamide;
8-Hydroxy-N-methyl-N-(1-methylpyrrolidin-3-yl)quinoline-7-carboxamide;
N-(2-Cyanoethyl)-N-(furan-2-ylmethyl)-8-hydroxyquinoline-7-carboxamide;
N-(2-Cyanoethyl)-8-hydroxy-N-((tetrahydrofuran-2-yl)methyl)quinoline-7-carboxamide;
N-Ethyl-8-hydroxy-N-methylquinoline-7-carboxamide;
8-Hydroxy-N-methyl-N-propylquinoline-7-carboxamide;
(3,4-Dihydroisoquinolin-2(1H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(5-Bromoindolin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(6-methoxy-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Hydroxyquinolin-7-yl)(5-nitroindolin-1-yl)methanone;
8-Hydroxy-N-phenyl-N-propylquinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(octahydroquinolin-1(2H)-yl)methanone;
N-(4-Fluorobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
N-(3-Bromobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;

(8-Hydroxyquinolin-7-yl)(4-(trifluoromethyl)piperidin-1-yl)methanone;
(4-*tert*-Butylpiperidin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
(S)-8-Hydroxy-N-methyl-N-(1-phenylethyl)quinoline-7-carboxamide;
N-Benzyl-8-hydroxy-N-(2-hydroxyethyl)quinoline-7-carboxamide;
(3,3-Dimethylpiperidin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
N-(2-Bromobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(4-phenylpiperidin-1-yl)methanone;
((4aS,8S,8aR)-8-Hydroxy-octahydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(2-methylpiperidin-1-yl)methanone;
(8-Hydroxyquinolin-7-yl)(2-phenylpiperidin-1-yl)methanone;
(1,1-Dioxo-thiomorpholin-4-yl)-(8-hydroxy-quinolin-7-yl)-methanone;
(8-Hydroxyquinolin-7-yl)(4-methylpiperidin-1-yl)methanone;
(R)-8-Hydroxy-N-(1-phenylethyl)quinoline-7-carboxamide;
(8-Hydroxyquinolin-7-yl)(2-methylpyrrolidin-1-yl)methanone;
(2,5-Dimethylpyrrolidin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(3-phenylpyrrolidin-1-yl)methanone;
(3-(Dimethylamino)pyrrolidin-1-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(3-methylpiperidin-1-yl)methanone;
(8-Hydroxyquinolin-7-yl)(pyrrolidin-1-yl)methanone;
(1,4-Dioxa-8-aza-spiro[4.5]dec-8-yl)-(8-hydroxy-quinolin-7-yl)-methanone;
(8-Hydroxyquinolin-7-yl)(6-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Hydroxyquinolin-7-yl)(6-nitroindolin-1-yl)methanone;
(8-Hydroxyquinolin-7-yl)(7-(trifluoromethyl)-3,4-dihydroquinolin-1(2H)-yl)methanone;
(5-Bromo-8-hydroxyquinolin-7-yl)(isoindolin-2-yl)methanone;
(Hexahydro-1H-isoindol-2(3H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(2-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Hydroxyquinolin-7-yl)(3-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Hydroxyquinolin-7-yl)(4-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(8-Fluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(6-isopropyl-3,4-dihydroquinolin-1(2H)-yl)methanone;

(6-Chloro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(7-methyl-3,4-dihydroquinolin-1(2H)-yl)methanone;
(6-Bromo-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(8-Hydroxyquinolin-7-yl)(octahydroisoquinolin-2(1H)-yl)methanone;
N-(4-Bromophenyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
(Decahydro-1H-carbazol-9(9aH)-yl)(8-hydroxyquinolin-7-yl)methanone;
N-(4-Chlorobenzyl)-8-hydroxy-N-methylquinoline-7-carboxamide;
(5,7-Difluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(7-Fluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(5-Fluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(6-Fluoro-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(5-Chloro-8-hydroxy-3,4-dihydroquinolin-1(2H)-yl)(8-hydroxyquinolin-7-yl)methanone;
(2H-Benzo[b][1,4]oxazin-4(3H)-yl)(8-hydroxyquinolin-7-yl)methanone.

7. A compound of formula



8. A medicament comprising a compound according to any of the claims from 1 to 7.

9. An antifungal agent comprising a compound according to any of the claims from 1 to 7.

10. Use of a compound according to any one of claims 1 to 7 in the manufacture of a medicament for the treatment and/or prevention of fungal infections.

11. A pharmaceutical formulation containing at least a compound according to any one of claims from 1 to 7, together with at least a pharmaceutically acceptable carrier, excipient and/or adjuvant.