

HEPARIN-DERIVED POLYSACCHARIDE MIXTURES,
PREPARATION THEREOF AND PHARMACEUTICAL COMPOSITIONS
CONTAINING SAME

5 The present invention relates to mixtures of polysaccharides derived from heparin, their method of preparation and pharmaceutical compositions containing them.

10 Heparin is a mixture of sulfated mucopolysaccharides of animal origin which is used in particular for its anticoagulant and antithrombotic properties.

Heparin nevertheless has disadvantages which limit the
15 conditions for its use. In particular, its high anticoagulant activity (anti-IIa activity) can cause hemorrhages.


Low-molecular-weight heparins obtained by basic
20 depolymerization of heparin esters have been proposed (EP40144); however, these products still have a high anti-IIa activity.

Very-low-molecular-weight heparins have also been
25 described in US 6384021. However, the anti-Xa activity values obtained in the examples described do not exceed 120 IU, and the anti-Xa/anti-IIa ratio obtained is between 15 and 50.

30 In WO-0208295, very-low-molecular-weight heparins are prepared by a method different from US 6384021 and exhibit an activity between 100 and 150 IU with anti-Xa/anti-IIa ratios which are also very high for certain examples of application.

35

A constant need however exists in this class of medicament to improve the anti-Xa activities, in particular in obtaining activities greater than



Δ Ila

II_s

Is

5 The hexasaccharide Δ Ila-II_s-Is contained in the mixture of oligosaccharides may be a sequence which has as high affinity for ARIII and characterized by an aXa activity greater than 740 U/mg.

10 The mixture of oligosaccharides is in the form of an alkali or alkaline-earth metal salt.

15 In a second aspect, there is provided a method for preparing mixtures of oligosaccharides as defined above, during which the quaternary ammonium salt of the benzyl ester of heparin is depolymerized in an organic medium in the presence of a strong organic base with a pKa greater than 20, characterized in that the organic used consists of the family of phosphazenes, in particular in
20 a dichloromethane solution, contains a percentage of water of less than 0.6%.

25 In a third aspect, there is provided a method for preparing oligosaccharides as defined above from heparins, in which the following operations are carried out: a) transalification of sodium heparin by the action of benzethonium chloride, b) esterification of benzethonium heparinate obtained by the action of benzyl

150 IU/mg, and the anti-Xa/anti-IIa ratio, and therefore to develop novel generations of heparin derivatives.

- 5 One of the objectives of the invention is therefore to improve the anti-Xa activity and the anti-Xa/anti-IIa ratio by modifying the methods described in the prior art, in particular by controlling the percentage of water during the depolymerization step. The heparins
10 thus obtained thus exhibit an excellent antithrombotic activity and possess an aXa activity similar to that of heparin while reducing the risks of hemorrhage with a very low aIIa activity. Likewise, the products of the invention exhibit half-life periods which are markedly
15 greater than that of heparin.

The subject of the invention is therefore novel mixtures of polysaccharides derived from heparin possessing a more selective activity toward activated
20 factor X (factor Xa) and toward activated factor II (factor IIa) than heparin.

It is understood that the mixtures of polysaccharides having a mean molecular weight of 1500 to 3000 Da can
25 be termed as oligosaccharides.

In a first aspect, there is provided mixtures of sulfated oligosaccharides having the general structure of the constituent polysaccharides of
30 heparin and having the following characteristics:

- they have a mean molecular weight of 1500 to 3000 daltons, an anti-Xa activity of 120 to 200 IU/mg, an anti-IIa activity of less than 10 IU/mg and an anti-Xa activity/anti-IIa activity ratio of greater than 30,
- 35 - the constituent oligosaccharides of the mixtures contain 2 to 26 saccharide units, have a 4,5-unsaturated uronic acid 2-O-sulfate unit at one of their ends, and contain the hexasaccharide AIIa-IIs-Is of formula:

chloride, c) transalification of the benzy lester obtained and obtaining of the quaternary ammonium salt, d) depolymerization of the quaternary ammonium salt of the benzyl ester of heparin by the method as defined above, e) conversion of the quaternary ammonium salt to a sodium salt, f) optionally saponification of the heparin by the action of a base such as sodium hydroxide, g) optionally purification in particular by the action of an oxidizing agent such as hydrogen peroxide.

In a fourth aspect, there is provided, as a medicament, the oligosaccharides as defined above.

In a fifth aspect, there is provided, as a medicament having an antithrombotic activity, the oligosaccharides as defined above.

In a sixth aspect, there is provided a pharmaceutical composition containing at least one medicament as defined above and one or more pharmaceutically inert excipients or vehicles of additives.

In a seventh aspect, there is provided a mixture of polysaccharides as defined above, which can be obtained by the method as defined above.

As alkali or alkaline-earth metal salt, the sodium, potassium, calcium and magnesium salts are preferred.

The mean molecular weight is determined by high-pressure liquid chromatography using two columns in series, for example those marketed under the name TSK

G3000 XL and TSK G2000 XL. The detection is carried out by refractometry. The eluent used is lithium nitrate and the flow rate is 0.6 ml/min. The system is calibrated with standards prepared by fractionation of enoxaparin (AVENTIS) by chromatography on agarose-polyacrylamide gel (IBF). This preparation is carried out according to the technique described by Barrowcliffe et al., *Thromb. Res.*, 12, 27-36 (1977-78) or D.A. Lane et al., *Thromb. Res.*, 12, 257-271 (1977-78). The results are calculated with the GPC6 software (Perkin Elmer).

The anti-Xa activity is measured by the amidolytic method on a chromogenic substrate described by Teien et al., Thromb. Res., 10, 399-410 (1977), with, as
5 standard, the first international standard for low-molecular-weight heparins.

The anti-IIa activity is measured by the technique described by Anderson L.O. et al., Thromb. Res., 15,
10 531-541 (1979), with, as standard, the first international standard for low-molecular-weight heparins.

The hexasaccharide fraction preferably represents from 15 to 25% of the mixture of oligosaccharides.

15 Preferably, the mixtures as disclosed contain from 8 to 15% of the hexasaccharide Δ IIa-II_s-Is in the hexasaccharide fraction of the mixture of oligosaccharides.

20 The percentage of the hexasaccharide fraction may be analytically determined by high-pressure liquid chromatography on TSK G3000 XL and TSK G2000 XL columns or alternatively by preparative separation of the hexa-
25 saccharide fraction. The mixture is in this case chromatographed on columns filled with a polyacrylamide agarose type gel such as that marketed under the trademark Ultrogel ACA202^R (Biosepra). The mixture is eluted with a sodium hydrogen carbonate solution.
30 Preferably, the sodium hydrogen carbonate solution is a 0.1 mol/l to 1 mol/l solution. Still more preferably, the separation is carried out at a concentration of 1 mol/l. The determination is carried out by UV spectrometry (254 nm). After fractionation, the hexa-
35 saccharide fraction in solution in sodium hydrogen carbonate is neutralized with glacial acetic acid. The solution is then concentrated under reduced pressure so as to obtain a sodium acetate concentration greater than 30% by weight. The hexasaccharide fraction is

precipitated by adding from 3 to 5 volumes of methanol. The hexasaccharide fraction is recovered by filtration on No. 3 sintered glass. The hexasaccharide mixture obtained may be analyzed by HPLC (High-Performance
5 Liquid Chromatography) in order to determine the content of hexasaccharide Δ IIa-II $\bar{s}$ -Is. Hexasaccharide Δ IIa-II $\bar{s}$ -Is may be isolated by preparative HPLC chromatography or by affinity chromatography on an antithrombin III sepharose column according to the
10 techniques used by persons skilled in the art (M. Hook, I. Bjork, J. Hopwood and U. Lindahl, *F.E.B.S letters*, vol 656(1) (1976)).

Most particularly, the mixtures may have an anti-Xa
15 activity of between 150 IU/mg and 200 IU/mg.

Preferably, the mixtures may have an anti-IIa activity of less than 5 IU/mg, and most particularly of 0.5 to 3.5 IU/mg. The examples of applications
20 described below demonstrate values of between 1.1 and 1.6 IU/mg when the preferred characteristics of the process are used.

Preferably, the mixtures exhibit an anti-Xa activity/anti-IIa activity ratio greater than 50 and
25 most particularly greater than 100.

Preferably, the mixtures have a mean molecular weight of between 2000 and 3000 Daltons, and most
30 particularly a mean molecular weight of between 2400 and 2650 Da.

The mixtures as defined above may have an anti-Xa activity of between 150 and 200 IU/mg, an anti-IIa
35 activity of between 0.5 and 3.5 IU/mg and a mean molecular weight of between 2400 and 2650 Da.

The mixtures of oligosaccharides may be prepared by depolymerization of a quaternary ammonium salt of the benzyl ester of heparin in an organic medium, by means of a strong organic base with a pKa preferably greater than 20 (properties preferably similar to the family of phosphazenes defined for example according to R. Schwesinger et al., Angew. Chem. Int. Ed. Engl. 26, 1167-1169 (1987), R. Schwesinger et al., Angew. Chem. 105, 1420 (1993)), conversion of the quaternary ammonium salt of the benzyl ester of the depolymerized heparin to a sodium salt, saponification of the residual esters and optionally purification. The method disclosed may repeat the main steps of the method as described in WO 0208295 while adding an essential characteristic which makes it possible to obtain the mixtures of oligosaccharides with the physicochemical characteristics and the activities described above.

Indeed, in order to obtain the specific mixtures of oligosaccharides, it may be necessary to control the selectivity of the base by a very precise control of the water content of the mixture during the depolymerization step.

The method disclosed may be characterized by a control of the high selectivity of the base during the depolymerization. It makes it possible to depolymerize the heparin while preserving as much as possible the sequences with an affinity for ATIII such as the hexasaccharide Δ IIa-II_s-Is described. This critical step of the method makes it possible to obtain

the polysaccharides.

This characteristic of the method can result in unexpected aXa activities in terms of the mean molecular weight of the mixtures of oligosaccharides (150 IU/mg < aXa <

200 IU/mg for a mean molecular weight of between 2000 Da and 3000 Da). This selectivity may be due to the very particular physicochemical characteristics of the phosphazene bases which have a pKa greater than 20, a very high steric hindrance and a weak nucleophilicity.

This effect may be fully expressed when the reaction medium is anhydrous. On the other hand, when the water content of the reaction medium increases, a drastic reduction in the selectivity of the depolymerization can be observed. The preservation of the sequences with affinity for ATIII decreases and the consequence may be a large drop in the aXa activity. In the presence of a low quantity of water, the phosphazene base may become protonated and the reactive species may become a quaternary ammonium hydroxide. In this case, the very high steric hindrance and weak nucleophilicity properties may be lost and greatly influence the quality of the product obtained. When depolymerization trials are carried out with a measured and controlled water content, this effect can be seen to be fully expressed.

The following table summarizes the impact of the water content on the selectivity of the depolymerization (only this parameter is variable in the trials: the stoichiometry of the reagents, the dilutions, the temperatures remain constant according to the criteria of persons skilled in the art. The base used is the phosphazene base: 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine).

Water content %	0.05%	0.1%	0.2%	0.3%	0.4%	0.57%	1.8%	2.5%
aXa IU/mg	192	177	161	132	122	120	105	99
aIIa IU/mg	1.3	1.5	1.4	1.4	1.3	1.4	3.1	13.4
aXa/aIIa	148	118	115	94	94	85.7	34	7.4

For an optimum selectivity and a maximum preservation of the sequences with affinity for ATIII, it is preferable to carry out the procedure at water contents of less than 0.6% and most particularly less than 0.3% when 1 molar equivalent of phosphazene base is used relative to the benzyl ester of heparin, benzethonium salt.

10

Disclosed herein is the step of depolymerization of the quaternary ammonium salt of the benzyl ester of heparin obtained according to methods known to persons skilled in the art, characterized in that a base of the family of phosphazenes, in particular in dichloromethane solution containing a percentage of water of less than 0.6%, is used. Preferably, this percentage of water should be chosen less than 0.3% and most particularly less than 0.2%.

20


Advantageously, the strong base/ester mol ratio is between 0.2 and 5, preferably between 0.6 and 2 and most particularly between 0.8 and 1.2. The use of the equimolar ratio therefore forms part of the preferred embodiments of the invention.

25

Other aprotic solvents known to persons skilled in the art may be used, such as THF or DMF.

The quaternary ammonium salt of the benzyl ester of heparin is preferably the benzethonium, cetylpyridinium or cetyltrimethylammonium salt.

- 5 The bases of the family of phosphazenes are preferably those of formula:




- 10 in which the radicals R1 to R7 are identical or different and represent alkyl radicals.

In the preceding formulae, the alkyl radicals contain 1 to 6 carbon atoms in the form of a straight or branched chain.

15 Disclosed herein is a method for preparing the mixtures of oligosaccharides according to the invention comprising the following steps

- 20 a) transalification of sodium heparin by the action of benzethonium chloride,
b) esterification of benzethonium heparinate by the action of benzyl chloride,
c) transalification of the benzyl ester obtained to a
25 quaternary ammonium salt,
d) depolymerization of the quaternary ammonium salt of the benzyl ester of heparin by the method as defined above,
e) conversion of the quaternary ammonium salt to a
30 sodium salt,
f) optionally saponification of the heparin by the action of a base such as sodium hydroxide,
g) optionally purification in particular by the action of an oxidizing agent such as hydrogen peroxide.

The following reaction scheme is as illustrated below.



5

The conversion of the quaternary ammonium salt of the benzyl ester of the depolymerized heparin to a sodium salt (step e) is generally carried out by treating the reaction medium with an alcoholic solution of sodium acetate and preferably with a 10% solution of sodium acetate in methanol (weight/volume), at a temperature of between 15 and 25°C. The equivalent by weight of acetate added is preferably 3 times greater than the mass of quaternary ammonium salt of the benzyl ester of heparin used in the depolymerization reaction.

15

The saponification (step f) is generally carried out by means of an alkali metal hydroxide such as sodium hydroxide, potassium hydroxide or lithium hydroxide, in an aqueous medium, at a temperature of between 0 and 20°C and preferably between 0 and 10°C. 1 to 5 molar equivalents of alkali metal hydroxide will be generally used. Preferably, the saponification will be carried out in the presence of 1 to 2 molar equivalents of alkali metal hydroxide.

10

The final product may be optionally purified (step g) by any known method of purifying depolymerized heparins (for example EP 0037319B1). Preferably, the purification is carried out by means of hydrogen peroxide, in an aqueous medium, at a temperature of 10 to 50°C. Preferably, this operation is carried out between 20 and 40°C.

15

The quaternary ammonium salt of the benzyl ester of heparin may be prepared according to the following reaction scheme:

20

a) conversion of the heparin to the form of a sodium salt by means of benzethonium chloride in order to obtain benzethonium heparinate (transalification),

25

b) esterification of the benzethonium salt obtained above by means of benzyl chloride and treatment with an alcoholic solution of sodium acetate in order to obtain the sodium salt of the benzyl ester of heparin,

30

c) transalification of the sodium salt of the benzyl ester of heparin to a quaternary ammonium salt and preferably to a benzethonium, cetylpyridinium or cetyltrimethylammonium salt.

The reaction of step a) may be carried out by the action of benzethonium chloride in excess, on sodium heparin, at a temperature in the region of 15 to 25°C. Advantageously, the salt/sodium heparin molar ratio is between 3 and 4.

35

The starting heparin used is preferably a pig heparin. The latter may be purified beforehand in order to reduce its dermatan sulfate level according to the method described in patent FR2663639.

5

The esterification of step b) is preferably carried out in a chlorinated organic solvent (for example chloroform or methylene chloride), at a temperature of between 25 and 45°C and preferably between 30 and 40°C.

10 The ester in the form of a benzethonium salt is then recovered in the form of a sodium salt by precipitation by means of sodium acetate at 10% by weight in an alcohol such as methanol. 1 to 1.2 volumes of alcohol are generally used per volume of reaction medium. The
15 quantity of benzyl chloride and the reaction time are adjusted in order to obtain a degree of esterification of between 50 and 100% and preferably between 70 and 90%. Preferably, 0.5 to 1.5 parts by weight of benzyl chloride are used for 1 part by weight of benzethonium
20 salt of heparin. Likewise, preferably the reaction time will be between 10 and 35 hours.

The transalification step c) may be carried out by means of a quaternary ammonium chloride and preferably by means
25 of benzethonium chloride, cetylpyridinium chloride or cetyltrimethylammonium chloride, in an aqueous medium, at a temperature of between 10 and 25°C. Advantageously, the quaternary ammonium chloride/sodium salt of the benzyl ester of heparin mol ratio is between 2
30 and 3.

The mixtures ~~according~~
a sodium salt, may be converted to another alkali or alkaline-earth metal salt. The passage from one salt to
35 another is optionally achieved using the method described in patent FR 73 13 580.

~~The mixtures according to the invention disclosed~~ are not toxic and may thus be used as medicaments.

The mixtures of oligosaccharides may be used as antithrombotic agents. In particular, they are useful for the treatment or prevention of venous and arterial thromboses, deep vein thrombosis, pulmonary embolism, unstable angina, myocardial infarction, cardiac ischemia, occlusive diseases of the peripheral arteries and atrial fibrillation. They are also useful in the prevention and treatment of the proliferation of smooth muscle cells, atherosclerosis and arteriosclerosis, for the treatment and prevention of cancer by modulating angio-genesis and growth factors, and for the treatment and prevention of diabetic disorders such as diabetic retinopathies and nephropathies.

Disclosed herein is the pharmaceutical compositions containing, as active ingredient, a mixture of formula (I), optionally in combination with one or more inert excipients.

The pharmaceutical compositions are for example solutions which can be injected by the subcutaneous or intravenous route. They are also useful for administration by the pulmonary route (inhalation) or by the oral route.

The dosage may vary according to the age, weight and state of health of the patient. For an adult, it is in general between 20 and 100 mg per day by the intramuscular or subcutaneous route.

The following examples illustrate the invention without however limiting it.

Example A: PREPARATION OF THE BENZETHONIUM SALT OF BENZYL OF HEPARINATE

Benzethonium heparinate

A solution of 25 g of benzethonium chloride in 125 ml of water is added to a solution of 10 g of heparin in the form of a sodium salt in 100 ml of water. The
5 product is filtered, washed with water and dried.

Benzyl ester of heparin (sodium salt)


16 ml of benzyl chloride are added to a solution of 20 g of benzethonium heparinate in 80 ml of methylene
10 chloride. The solution is heated at a temperature of 30°C for 12 hours. 108 ml of a 10% solution of sodium acetate in methanol are then added, the mixture is filtered, washed with methanol and dried. 7.6 g of benzyl ester of heparin are thus obtained in the form
15 of a sodium salt whose degree of esterification is 77%.

Benzyl ester of heparin (benzethonium salt)

36 g (0.0549 mol) of benzyl ester of heparin (sodium salt) and 540 ml of distilled water are introduced into
20 a 2-litre Erlenmeyer flask A. After homogenization at a temperature of about 20°C, a pale yellow solution is obtained. A solution of 64.45 g (0.1438 mol) of benzethonium chloride and 450 ml of water is prepared, with magnetic stirring, in a 1-litre Erlenmeyer flask
25 B. The solution in Erlenmeyer B is poured over about 35 minutes into the solution in Erlenmeyer A, with stirring. The formation of an abundant white precipitate is observed. The Erlenmeyer B is rinsed with 200 ml of distilled water and the wash water is
30 introduced into the Erlenmeyer A. The stirring is then stopped and the suspension is allowed to settle for 12 hours. Once this time has elapsed, the clear portion of the supernatant is removed and discarded. 560 ml of water are added to the sedimented precipitate (slurry
35 appearance) and the mixture is stirred for 20 minutes. The precipitate is allowed to resediment for about 30 minutes. The supernatant is removed and discarded (560 ml). This operation of washing with about 560 ml of distilled water is repeated twice on the sedimented

precipitate. In the last washing operation, the precipitate is left in suspension and filtered through a No. 3 sintered glass. The cake is then washed with 4 times 200 ml of distilled water. The wet white solid is drained and then dried under reduced pressure (2.7 kPa) at a temperature in the region of 60°C. After drying for 12 hours, 87.5 g of benzyl heparinate, benzethonium salt, are obtained. The yield obtained is 94.9%.

10 **Example B: DESCRIPTION OF THE HEXASACCHARIDE ATIII**
(Δ IIa-II $\underline{s}$ -Is)



15 Proton spectrum in D₂O, 500 MHz, T=298 K, δ in ppm :
1.97 (3H, s), 3.18 (1H, dd, 10 and 3Hz), 3.30 (1H, t, 8Hz), 3.37 (1H, dd, 10 and 3Hz), 3.60 (2H, m), between 3.65 and 3.85 (6H, m), 3.87 (2H, m), 3.95 (1H, d, 8Hz), 4.03 (1H, d, 8Hz), between 4.05 and 4.13 (4H, m),
20 between 4.16 and 4.45 (8H, m), 4.52 (1H, d, 8Hz), 4.67 (1H, m), 5.06 (1H, d, 6Hz), 5.10 (1H, d, 3Hz), 5.33 (1H, d, 4Hz), 5.36 (1H, d, 3Hz), 5.46 (1H, d, 3Hz), 5.72 (1H, d, 4Hz).

Decasodium salt of 4-deoxy- α -L-threo-hexenepyransyluronic
25 uronic acid-(1 $\rightarrow$ 4)-2-deoxy-2-acetamido-6-O-sulfo- α -D-glucopyranosyl-(1 $\rightarrow$ 4)- β -D-glucopyranosyluronic acid-(1 $\rightarrow$ 4)-2-deoxy-2-sulfamido-3,6-di-O-sulfo- α -D-glucopyranosyl-(1 $\rightarrow$ 4)-2-O-sulfo- α -L-idopyranosyluronic acid-(1 $\rightarrow$ 4)-2-deoxy-2-sulfamido-6-O-sulfo- α -D-glucopyranose.

. Examples 1 to 7 and 12 illustrate the influence of the water content on the selectivity of the polymerization reaction and the aXa and aIIa activity of the products obtained.

5 . Examples 8 to 10 illustrate the influence of the number of base equivalents on the aXa and aIIa activity of the product obtained (with a water content of 0.1%)

. Example 11 illustrates the use of a phosphazene base other than 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine:

Use of tert-butyliminotri(pyrrolidino)phosphorane

EXAMPLE 1

Depolymerization and conversion to the sodium salt (0.1% water):

70 ml of dichloromethane are placed in an Erlenmeyer flask A. 10 g (0.006 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A are added slowly with stirring. The water content of the reaction medium is adjusted to 0.1%. The solution is heated to 40°C under nitrogen. After total dissolution, the solution is cooled to a temperature in the region of 20°C, followed by addition of 1.75 ml (0.006 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine. The resulting mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 30 g of anhydrous sodium acetate in 300 ml of methanol is prepared in an Erlenmeyer flask B. After total dissolution, 5 g of Hyflo supercel Celite are added to the solution. The reaction mixture in the Erlenmeyer flask A is poured over 1 minute 30 seconds into the methanolic solution of sodium acetate at a temperature in the region of 5°C, with magnetic stirring. After stirring for 5 minutes, the suspension is left to settle for 1 hour 30 minutes. The clear part of the supernatant is separated out and discarded (220 ml). 220 ml of methanol are added to the sedimented precipitate and

the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 1 hour 20 minutes. The supernatant is separated out and discarded (250 ml). 250 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 100 ml of methanol. The pale yellow wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 18 hours, 2.51 g of the sodium salt of depolymerized heparin in Celite (5 g) are obtained. The yield obtained is 64%.

15 **Saponification:**

2.5 g (0.0038 mol) of the crude depolymerized heparin, sodium salt in Celite (5 g) obtained above, and 17 ml of water are placed in a 50 ml Erlenmeyer flask. The suspension is filtered through a No. 3 sintered glass and rinsed with twice 5 ml of water. The filtrate obtained is placed in a 150 ml Erlenmeyer flask. 0.4 ml (0.004 mol) of 30% caustic soda is introduced with magnetic stirring, at a temperature in the region of 5°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl and 3 g of sodium chloride are added. After dissolution, 21 ml of methanol are added to the reaction medium. After stirring for 15 minutes, 44 ml of methanol are added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 45 minutes at a temperature in the region of 5°C. The supernatant is then separated out and discarded (90 ml). 90 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 20 minutes. The supernatant is separated out and discarded (80 ml). 80 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then

filtered through a No. 3 sintered glass. The cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After
5 drying for 18 hours, 1.31 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 66%.

Purification:

10 1.3 g of crude depolymerized heparin obtained above and 13 ml of distilled water are placed in a 50 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is
15 filtered through a 0.45 μ m membrane and 0.07 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding 1N HCl, followed by addition of 2 g of sodium chloride.
20 The solution is then filtered through a 0.45 μ m membrane and 14 ml of methanol are then poured in. The solution is then cooled to 10°C and stirred for about 15 minutes. 36 ml of methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and
25 the suspension is left to sediment for about 15 minutes. The supernatant is then separated out and discarded (50 ml). 50 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for
30 about 25 minutes. The supernatant is separated out and discarded (50 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced
35 pressure (6 kPa) at a temperature in the region of 40°C. After drying for 18 hours, 1.13 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 87%.

The characteristics of the depolymerized heparin thus obtained are as follows:

mean molecular weight: 2600 Daltons

anti-Xa activity: 177 IU/mg

5 anti-IIa activity: 1.5 IU/mg

anti-Xa activity/anti-IIa activity ratio: 118

EXAMPLE 2

10 Depolymerization and conversion to the sodium salt (0.2%)

70 ml of dichloromethane are placed in a Erlenmeyer flask A. 10 g (0.006 mol) of the benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A are added
15 slowly, with stirring and under nitrogen pressure. The water content of the reaction medium is adjusted to 0.2%. After total dissolution, 1.75 ml (0.006 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The mixture is
20 stirred for 24 hours at a temperature in the region of 20°C. During this time, a solution of 30 g of anhydrous sodium acetate in 300 ml of methanol is prepared in a Erlenmeyer flask B. After total dissolution, 5 g of Hyflo supercel Celite are added to the solution. The
25 reaction mixture in the Erlenmeyer flask A is poured over 1 minute 30 seconds into the methanolic solution of sodium acetate with magnetic stirring, at a temperature in the region of 5°C. After stirring for 5 minutes, the suspension is left to settle for
30 2 hours. The clear part of the supernatant is separated out and discarded (220 ml). 220 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 2 hours. The supernatant is
35 separated out and discarded (230 ml). 230 ml of methanol are added to the sedimented precipitate. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 150 ml of methanol. The pale yellow wet solid is

drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 18 hours, 2.63 g of crude depolymerized heparin in Celite (5 g) are obtained. The yield obtained is 67%.

5

Saponification:

2.5 g (0.0038 mol) of the sodium salt of depolymerized heparin in Celite (5 g) obtained above and 18 ml of water are placed in 50 ml Erlenmeyer flask. The suspension is filtered through a No. 3 sintered glass and rinsed with twice 5 ml of water. The filtrate obtained is placed in a 150 ml Erlenmeyer flask. 0.4 ml (0.004 mol) of 30% caustic soda is added with magnetic stirring, at a temperature in the region of 5°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl and 2 g of sodium chloride are added. 14 ml of methanol are poured into the reaction mixture. After stirring for 15 minutes, 36 ml of methanol are added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 45 minutes at a temperature in the region of 5°C. The supernatant is then separated out and discarded (80 ml). 80 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 48 hours, 2.3 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 65%.

Purification:

1.4 g of crude depolymerized heparin obtained above and 15 ml of distilled water are placed in a 50 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is

filtered through a 0.45 μ m membrane and 0.07 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 40°C, the mixture is cooled to a temperature in the region of 20°C and then neutralized by addition of 1N HCl. 2 g of sodium chloride are added to the reaction medium. The solution is then filtered through a 0.45 μ m membrane and 14 ml of methanol are then poured in. The solution is then cooled to 10°C and stirred for 15 minutes. 36 ml of methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 15 minutes. The supernatant is then separated out and discarded (40 ml). 40 ml of methanol are added to the sedimented precipitate (slurry appearance) and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 20 minutes. The supernatant is separated out and discarded (50 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 18 hours, 1.2 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 86%.

The characteristics of the depolymerized heparin thus obtained are as follows:

mean molecular weight: 2650 daltons

anti-Xa activity: 161 IU/mg

anti-IIa activity: 1.4 IU/mg

anti-Xa activity/anti-IIa activity ratio: 115

EXAMPLE 3

Depolymerization and conversion to the sodium salt (0.3% water)

70 ml of dichloromethane are placed in a Erlenmeyer flask A. 10 g (0.006 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A are added with

stirring and under nitrogen pressure. The water content of the reaction medium is adjusted to 0.3%. After total dissolution, 1.75 ml (0.006 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 30 g of anhydrous sodium acetate in 300 ml of methanol is prepared in a Erlenmeyer flask B. After total dissolution, 5 g of Hyflo supercel Celite are added to the solution. The reaction mixture in the Erlenmeyer flask A is poured over 1 minute 30 seconds into the methanolic sodium acetate solution at a temperature in the region of 5°C, with magnetic stirring. After stirring for 5 minutes, the suspension is left to settle for 1 hour 10 minutes. The clear part of the supernatant is separated out and discarded (220 ml). 220 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 100 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 18 hours, 2.57 g of crude depolymerized heparin are obtained, as the sodium salt in Celite (5 g). The yield obtained is 66%.

Saponification:

2.5 g (0.0038 mol) of the sodium salt of crude depolymerized heparin in Celite (5 g) obtained above and 18 ml of water are placed in 50 ml Erlenmeyer flask. The suspension is filtered through a No. 3 sintered glass and rinsed with twice 5 ml of water. The filtrate obtained is placed in a 150 ml Erlenmeyer flask. 0.4 ml (0.004 mol) of 30% caustic soda is added with magnetic stirring, at a temperature in the region of 5°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl and 3 g of sodium chloride are added. 15 ml of methanol

are poured into the reaction mixture. After stirring for 15 minutes, 36 ml of methanol are added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 1 hour. The supernatant is then separated out and discarded (70 ml). 70 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 48 hours, 1.42 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 62%.

15

Purification:

1.4 g of crude depolymerized heparin obtained above and 14 ml of distilled water are placed in a 50 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μ m membrane and 0.07 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding 1N HCl, followed by addition of 2 g of sodium chloride. After dissolution, the solution is then filtered through a 0.45 μ m membrane and 14 ml of methanol are then poured in. The filtrate is then cooled to 10°C and stirred for 15 minutes. 36 ml of methanol are then added, followed by stirring for about 1 hour. Stirring is then stopped and the suspension is left to sediment for 40 minutes. The supernatant is then separated out and discarded (50 ml). 50 ml of methanol are added to the sedimented precipitate (slurry appearance) and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 25 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then

washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 18 hours, 1.24 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 89%.

The characteristics of the depolymerized heparin thus obtained are as follows:

mean molecular weight: 2400 daltons

anti-Xa activity: 132 IU/mg

10 anti-IIa activity: 1.4 IU/mg

anti-Xa activity/anti-IIa activity ratio: 94

EXAMPLE 4

Depolymerization and conversion to the sodium salt (0.4% water):

15 70 ml of dichloromethane are placed in a Erlenmeyer flask A. 10 g (0.006 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A, are added slowly, with stirring. The water content of the reaction medium is adjusted to 0.4%. The solution is heated to 30°C under nitrogen. After total dissolution, the mixture is cooled to a temperature in the region of 20°C, followed by addition of 1.75 ml (0.006 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 30 g of anhydrous sodium acetate in 300 ml of methanol is prepared in a Erlenmeyer flask B. After total dissolution, 5 g of Hyflo supercel Celite are added to the solution. The reaction mixture in the Erlenmeyer flask A is poured over 1 minute 30 seconds into the methanolic sodium acetate solution at a temperature in the region of 5°C, with magnetic stirring. After stirring for 5 minutes, the suspension is left to settle for 2 hours. The clear part of the supernatant is separated out and discarded (80 ml). 80 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes.

The precipitate is left to resediment for about 1 hour. The supernatant is separated out and discarded (80 ml). 80 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes.

5 The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 150 ml of methanol. The pale yellow wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After

10 drying for 18 hours, 3.25 g of crude depolymerized heparin are obtained, as the sodium salt in Celite (5 g). The yield obtained is 83%.

Saponification:

15 3.1 g (0.0018 mol) of the sodium salt of crude depolymerized heparin in Celite (10 g) obtained above and 21 ml of water are placed in 50 ml Erlenmeyer flask. The suspension is filtered through a No. 3 sintered glass and rinsed with twice 6 ml of water. The

20 filtrate obtained is placed in a 150 ml Erlenmeyer flask. 0.7 ml (0.007 mol) of 30% caustic soda is added with magnetic stirring, at a temperature in the region of 5°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl

25 and 4 g of sodium chloride are added. 28 ml of methanol are poured into the reaction mixture. After stirring for 15 minutes, 72 ml of methanol are added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 1 hour. The

30 supernatant is then separated out and discarded (90 ml). 90 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 20 minutes. The supernatant is separated out and

35 discarded (90 ml). 90 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 50 ml of methanol. The wet

solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 48 hours, 1.9 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 67%.

Purification:

1.9 g of crude depolymerized heparin obtained above and 19 ml of distilled water are placed in a 50 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μ m membrane and 0.1 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding 1N HCl, followed by addition of 2 g of sodium chloride. The solution is then filtered through a 0.45 μ m membrane and 14 ml of methanol are then poured in, and the mixture is stirred for 15 minutes. 36 ml of methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 15 minutes. The supernatant is then separated out and discarded (40 ml). 40 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 20 minutes. The supernatant is separated out and discarded (50 ml). 500 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 20 minutes. The suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 72 hours, 1.56 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 82%.

The characteristics of the depolymerized heparin thus obtained are as follows:

mean molecular weight: 2350 daltons

anti-Xa activity: 122 IU/mg

5 anti-IIa activity: 1.3 IU/mg

anti-Xa activity/anti-IIa activity ratio: 94

EXAMPLE 5

10 Depolymerization and conversion to the sodium salt (0.57% water):

140 ml of dichloromethane are placed in a Erlenmeyer flask A. 20 g (0.019 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A, are added slowly, 15 with stirring. The water content of the reaction medium is adjusted to 0.57%. After total dissolution, 3.5 ml (0.012 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 60 g 20 of anhydrous sodium acetate in 600 ml of methanol is prepared in a Erlenmeyer flask B. After total dissolution, 10 g of Hyflo supercel Celite are added to the solution. The reaction mixture in the Erlenmeyer 25 flask A is poured over 1 minute 30 seconds into the methanolic sodium acetate solution at a temperature in the region of 4°C, with magnetic stirring. After stirring for 5 minutes, the suspension is left to settle for 30 minutes. The clear part of the 30 supernatant is separated out and discarded (400 ml). 400 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 1 hour. The supernatant is separated out and discarded 35 (420 ml). 420 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 200 ml of methanol. The pale yellow wet solid is

drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 50°C. After drying for 18 hours, 6.66 g of crude depolymerized heparin are obtained, as the sodium salt in Celite (10 g). The
5 yield obtained is 85%.

Saponification:

6.66 g (0.0101 mol) of the sodium salt of crude depolymerized heparin in Celite (10 g) obtained above
10 and 47 ml of water are placed in a 50 ml Erlenmeyer flask. The suspension is filtered through a No. 3 sintered glass and rinsed with twice 15 ml of water. The filtrate obtained is placed in a 150 ml Erlenmeyer flask. 1.1 ml (0.011 mol) of 30% caustic soda are added
15 with magnetic stirring, at a temperature in the region of 5°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl and 9.5 g of sodium chloride are added. 66 ml of methanol are added to the reaction medium. After
20 stirring for 15 minutes, 171 ml of methanol are added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for $\frac{3}{4}$ hour at a temperature in the region of 5°C. The supernatant is then separated out and discarded
25 (160 ml). 160 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 20 minutes. The supernatant is separated out and discarded (180 ml). 180 ml of methanol are added to the
30 sedimented precipitate and the mixture is stirred for 5 minutes. The suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with twice 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a
35 temperature in the region of 40°C. After drying for 18 hours, 4.53 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 74%.

Purification:

4.53 g of crude depolymerized heparin obtained above and 45 ml of distilled water are placed in a 100 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by
5 addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μm membrane and 0.25 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding
10 1N HCl, followed by addition of 5.5 g of sodium chloride. The solution is then filtered through a 0.45 μm membrane and 38 ml of methanol are then poured in, at a temperature in the region of 10°C. The solution is then brought to 20°C and stirred for
15 15 minutes. 100 ml of methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 20 minutes. The supernatant is then separated out and discarded (90 ml). 90 ml of methanol are added to the sedimented
20 precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 25 minutes. The supernatant is separated out and discarded (100 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The white
25 cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 50°C. After drying for 18 hours, 3.7 g of pure depolymerized heparin (sodium salt) are obtained. The
30 yield obtained is 82%.

The characteristics of the depolymerized heparin thus obtained are as follows:

mean molecular weight: 2200 daltons

anti-Xa activity: 120 IU/mg

35 anti-IIa activity: 1.4 IU/mg

anti-Xa activity/anti-IIa activity ratio: 86

EXAMPLE 6

Depolymerization and conversion to the sodium salt (1.8% water):

5 140 ml of dichloromethane are placed in a Erlenmeyer flask A. 20 g (0.019 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A, are added slowly, with stirring. The water content of the reaction medium
10 is adjusted to 1.8%. After total dissolution, 3.5 ml (0.012 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 60 g
15 of anhydrous sodium acetate in 600 ml of methanol is prepared in a Erlenmeyer flask B. After total dissolution, 10 g of Hyflo supercel Celite are added to the solution. The reaction mixture in the Erlenmeyer flask A is poured over 1 minute 30 seconds into the
20 methanolic sodium acetate solution at a temperature in the region of 4°C, with magnetic stirring. After stirring for 5 minutes, the suspension is left to settle for 30 minutes. The clear part of the supernatant is separated out and discarded (400 ml).
25 400 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 1 hour. The supernatant is separated out and discarded (420 ml). 420 ml of methanol are added to the
30 sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 200 ml of methanol. The pale yellow wet solid is drained and then dried under
35 reduced pressure (6 kPa) at a temperature in the region of 50°C. After drying for 18 hours, 7.54 g of crude depolymerized heparin are obtained, as the sodium salt in Celite (10 g). The yield obtained is 96%.

Saponification:

7.54 g (0.0101 mol) of the sodium salt of crude depolymerized heparin in Celite (10 g) obtained above and 53 ml of water are placed in a 50 ml Erlenmeyer flask. The solution is filtered through a No. 3 sintered glass and rinsed with twice 15 ml of water. The filtrate obtained is placed in a 150 ml Erlenmeyer flask. 1.25 ml (0.012 mol) of 30% caustic soda are added with magnetic stirring, at a temperature in the region of 4°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl and 10.5 g of sodium chloride are added. 70 ml of methanol are added to the reaction medium. After stirring for 15 minutes, 190 ml of methanol are added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for $\frac{3}{4}$ hour at a temperature in the region of 4°C. The supernatant is then separated out and discarded (180 ml). 180 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 20 minutes. The supernatant is separated out and discarded (180 ml). 180 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with twice 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 50°C. After drying for 18 hours, 5.53 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 80%.

Purification:

5.53 g of crude depolymerized heparin obtained above and 55 ml of distilled water are placed in a 100 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is

filtered through a 0.45 μ m membrane and 0.31 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding 5 1N HCl, followed by addition of 7 g of sodium chloride. The solution is then filtered through a 0.45 μ m membrane and 49 ml of methanol are then poured in, at a temperature in the region of 10°C. The solution is then brought to 20°C and stirred for 15 minutes. 126 ml of 10 methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 20 minutes. The supernatant is then separated out and discarded (105 ml). 105 ml of methanol are added to the sedimented precipitate and 15 the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 25 minutes. The supernatant is separated out and discarded (110 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then 20 washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 55°C. After drying for 18 hours, 4.53 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 82%. 25 The characteristics of the depolymerized heparin thus obtained are as follows:

mean molecular weight: 2600 daltons
anti-Xa activity: 105 IU/mg
anti-IIa activity: 3.1 IU/mg
30 anti-Xa activity/anti-IIa activity ratio: 34

EXAMPLE 7

Depolymerization and conversion to the sodium salt (2.5% water):

35 140 ml of dichloromethane are placed in a Erlenmeyer flask A. 20 g (0.019 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A, are added slowly, with stirring. The water content of the reaction medium

is adjusted to 2.5%. After total dissolution, 3.5 ml (0.012 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 60 g of anhydrous sodium acetate in 600 ml of methanol is prepared in a Erlenmeyer flask B. After total dissolution, 10 g of Hyflo supercel Celite are added to the solution. The reaction mixture in the Erlenmeyer flask A is poured over 1 minute 30 seconds into the methanolic sodium acetate solution at a temperature in the region of 4°C, with magnetic stirring. After stirring for 5 minutes, the suspension is left to settle for 1 hour. The clear part of the supernatant is separated out and discarded (400 ml). 400 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 30 minutes. The clear part of the supernatant is separated out and discarded (400 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 200 ml of methanol. The pale yellow wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 50°C. After drying for 18 hours, 7.78 g of crude depolymerized heparin are obtained, as the sodium salt in Celite (10 g). The yield obtained is 99.6%.

Saponification:

7.78 g (0.0119 mol) of the sodium salt of crude depolymerized heparin in Celite (10 g) obtained above and 79 ml of water are placed in a 50 ml Erlenmeyer flask. The suspension is filtered through a No. 3 sintered glass and rinsed with twice 15 ml of water. The filtrate obtained is placed in a 150 ml Erlenmeyer flask. 1.3 ml (0.012 mol) of 30% caustic soda are added with magnetic stirring, at a temperature in the region of 4°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl.

and 10 g of sodium chloride are added. 60 ml of methanol are added to the reaction medium. After stirring for 15 minutes, 190 ml of methanol are added, followed by stirring for 1 hour. Stirring is then
5 stopped and the suspension is left to sediment for $\frac{3}{4}$ hour at a temperature in the region of 4°C. The supernatant is then separated out and discarded (180 ml). 180 ml of methanol are added to the sedimented precipitate and the mixture is stirred for
10 5 minutes. The precipitate is left to resediment for about 20 minutes. The supernatant is separated out and discarded (180 ml). 180 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then
15 filtered through a No. 3 sintered glass. The cake obtained is then washed with twice 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 50°C. After drying for 18 hours, 5.87 g of crude
20 depolymerized heparin (sodium salt) are obtained. The yield obtained is 82%.

Purification:

5.87 g of crude depolymerized heparin obtained above
25 and 59 ml of distilled water are placed in a 100 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μ m membrane and 0.34 ml of
30 aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding 1N HCl, followed by addition of 7 g of sodium chloride. The solution is then filtered through a 0.45 μ m
35 membrane and 49 ml of methanol are then poured in, at a temperature in the region of 10°C. The solution is then brought to 20°C and stirred for 15 minutes. 126 ml of methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is

left to sediment for 20 minutes. The supernatant is then separated out and discarded (105 ml). 105 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 25 minutes. The supernatant is separated out and discarded (110 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 55°C. After drying for 18 hours, 5.21 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 89%. The characteristics of the depolymerized heparin thus obtained are as follows:

mean molecular weight: 3550 daltons
anti-Xa activity: 99 IU/mg
anti-IIa activity: 13.4 IU/mg
anti-Xa activity/anti-IIa activity ratio: 7.4

20

EXAMPLE 8

Depolymerization and conversion to the sodium salt (0.5 equivalent of base):

140 ml of dichloromethane are placed in a Erlenmeyer flask A. 20 g (0.019 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A, are added slowly, with stirring. After total dissolution at a temperature in the region of 30°C and cooling to a temperature in the region of 20°C, 1.75 ml (0.006 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 60 g of anhydrous sodium acetate in 600 ml of methanol is prepared in a Erlenmeyer flask B. The reaction mixture in the Erlenmeyer flask A is poured into the methanolic sodium acetate solution at a temperature in the region of 4°C, with magnetic stirring. After stirring for 1 hour, the

suspension is left to settle for 2 hours. The clear part of the supernatant is separated out and discarded (420 ml). 420 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 30 minutes. The precipitate is left to resediment for about 18 hours. The clear part of the supernatant is separated out and discarded (400 ml). 400 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 30 minutes. The supernatant 10 is separated out and discarded (400 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with twice 100 ml of methanol. The pale yellow wet solid is drained and then dried under reduced pressure 15 (6 kPa) at a temperature in the region of 60°C. After drying, 5.7 g of crude depolymerized heparin are obtained, as the sodium salt. The yield obtained is 73%.

20 **Saponification:**

5.7 g (0.0086 mol) of the sodium salt of crude depolymerized heparin obtained above and 53 ml of water are placed in a 100 ml Erlenmeyer flask. 0.93 ml (0.009 mol) of 30% caustic soda is added with magnetic 25 stirring, at a temperature in the region of 4°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl and 6 g of sodium chloride are added. 42 ml of methanol are added to the reaction medium. After stirring for 15 minutes, 30 108 ml of methanol are added, followed by stirring for 30 minutes. Stirring is then stopped and the suspension is left to sediment for 30 minutes at a temperature in the region of 4°C. The supernatant is then separated out and discarded (180 ml). 180 ml of methanol are 35 added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 30 minutes. The supernatant is separated out and discarded (170 ml). 170 ml of methanol are added to the sedimented precipitate and

the mixture is stirred for 30 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with twice 30 ml of methanol. The wet solid is drained and then
5 dried under reduced pressure at a temperature in the region of 60°C. After drying, 3.5 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 67.4%.

10 **Purification:**

3.5 g of crude depolymerized heparin obtained above and 35 ml of distilled water are placed in a 100 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.6 ± 0.1 by
15 addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μ m membrane and 0.18 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding
20 1N HCl, followed by addition of 3.6 g of sodium chloride. The solution is then filtered through a 0.45 μ m membrane and 27 ml of methanol are then poured in, at a temperature in the region of 10°C. The solution is then brought to 20°C and stirred for
25 15 minutes. 65 ml of methanol are then added, followed by stirring for 30 minutes. Stirring is then stopped and the suspension is left to sediment for 30 minutes. The supernatant is then separated out and discarded (80 ml). 80 ml of methanol are added to the sedimented
30 precipitate and the mixture is stirred for 30 minutes. The precipitate is left to resediment for about 30 minutes. The supernatant is separated out and discarded (70 ml). 70 ml of methanol are added to the sedimented precipitate and the mixture is stirred for
35 30 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then washed with twice 30 ml of methanol. The wet solid is drained and then dried under reduced pressure at a temperature in the region of 60°C. After

drying, 2.8 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 80%.

The characteristics of the depolymerized heparin thus obtained are as follows:

- 5 mean molecular weight: 2900 daltons
- anti-Xa activity: 146.1 IU/mg
- anti-IIa activity: 5.1 IU/mg
- anti-Xa activity/anti-IIa activity ratio: 28.6

10 **EXAMPLE 9**

Depolymerization and conversion to the sodium salt (0.6 equivalent of base):

- 280 ml of dichloromethane are placed in a three-necked flask A. 40 g (0.024 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A, are added slowly, with stirring. After total dissolution at a temperature in the region of 30°C and cooling to a temperature in the region of 20°C, 4.2 ml (0.014 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 60 g of anhydrous sodium acetate in 600 ml of methanol is prepared in a Erlenmeyer flask B. Half of the reaction mixture in the three-necked flask A is poured into the methanolic sodium acetate solution at a temperature in the region of 4°C, with magnetic stirring. After stirring for 1 hour, the suspension is left to settle. The clear part of the supernatant is separated out and discarded (310 ml). 310 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 1 hour. The precipitate is left to resediment for about 18 hours. The clear part of the supernatant is separated out and discarded (400 ml). 400 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 1 hour. The supernatant is separated out and discarded (300 ml). The precipitate in suspension is then filtered through a No. 3 sintered

glass. The cake obtained is then washed with twice
100 ml of methanol. The pale yellow wet solid is
drained and then dried under reduced pressure at a
temperature in the region of 60°C. After drying, 6 g of
5 crude depolymerized heparin are obtained, as the sodium
salt. The yield obtained is 77%.

Saponification:

6 g (0.0091 mol) of the sodium salt of crude
10 depolymerized heparin obtained above and 56 ml of water
are placed in a 250 ml Erlenmeyer flask. 1 ml
(0.010 mol) of 30% caustic soda is added with magnetic
stirring, at a temperature in the region of 4°C. After
addition, the mixture is stirred for 2 hours. The
15 solution is neutralized by adding 1N HCl and 6.4 g of
sodium chloride are added. 45 ml of methanol are added
to the reaction medium. After stirring for 15 minutes,
115 ml of methanol are added, followed by stirring for
30 minutes. Stirring is then stopped and the suspension
20 is left to sediment for 30 minutes at a temperature in
the region of 4°C. The supernatant is then separated
out and discarded (170 ml). 170 ml of methanol are
added to the sedimented precipitate and the mixture is
stirred for 30 minutes. The precipitate is left to
25 resediment for about 30 minutes. The supernatant is
separated out and discarded (140 ml). 140 ml of
methanol are added to the sedimented precipitate and
the mixture is stirred for 30 minutes. The precipitate
in suspension is then filtered through a No. 3 sintered
30 glass. The cake obtained is then washed with twice
30 ml of methanol. The wet solid is drained and then
dried under reduced pressure at a temperature in the
region of 60°C. After drying, 3.6 g of crude
depolymerized heparin (sodium salt) are obtained. The
35 yield obtained is 65.3%.

Purification:

3.5 g of crude depolymerized heparin obtained above and
35 ml of distilled water are placed in a 100 ml

Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.6 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μ m membrane and 0.18 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding 1N HCl, followed by addition of 3.5 g of sodium chloride. The solution is then filtered through a 0.45 μ m membrane and 25 ml of methanol are then poured in, at a temperature in the region of 10°C. The solution is then brought to 20°C and stirred for 15 minutes. 63 ml of methanol are then added, followed by stirring for 30 minutes. Stirring is then stopped and the suspension is left to sediment for 30 minutes. The supernatant is then separated out and discarded (70 ml). 70 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 30 minutes. The precipitate is left to resediment for a few minutes. The suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then washed with twice 30 ml of methanol. The wet solid is drained and then dried under reduced pressure at a temperature in the region of 60°C. After drying, 2.5 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 71.4%.
The characteristics of the depolymerized heparin thus obtained are as follows:
mean molecular weight: 2600 daltons
anti-Xa activity: 150.5 IU/mg
anti-IIa activity: 3.2 IU/mg
anti-Xa activity/anti-IIa activity ratio: 47

EXAMPLE 10

Depolymerization and conversion to the sodium salt (0.8 equivalent of base):

70 ml of dichloromethane are placed in a Erlenmeyer flask A. 10 g (0.006 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt)

obtained as described in example A, are added slowly, with stirring. After total dissolution at a temperature in the region of 30°C and cooling to a temperature in the region of 20°C, 1.38 ml (0.004 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 30 g of anhydrous sodium acetate in 300 ml of methanol is prepared in a Erlenmeyer flask B. The reaction mixture in the Erlenmeyer flask A is poured over 1 minute 15 seconds into the methanolic sodium acetate solution at a temperature in the region of 4°C, with magnetic stirring. After stirring for 5 minutes, the suspension is left to settle for 1 hour. The clear part of the supernatant is separated out and discarded (190 ml). 190 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 30 minutes. The clear part of the supernatant is separated out and discarded (190 ml). 190 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 30 minutes. The supernatant is separated out and discarded (190 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 150 ml of methanol. The pale yellow wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 18 hours, 3.05 g of crude depolymerized heparin are obtained, as the sodium salt. The yield obtained is 80%.

Saponification:

3.05 g (0.0048 mol) of the sodium salt of crude depolymerized heparin obtained above and 21 ml of water are placed in a 100 ml Erlenmeyer flask. The solution is filtered through a No. 3 sintered glass and rinsed with twice 6 ml of water. The filtrate obtained is placed in a 150 ml Erlenmeyer flask. 0.6 ml (0.006 mol)

of 30% caustic soda is added with magnetic stirring, at a temperature in the region of 4°C. After addition, the mixture is stirred for 2 hours. The solution is neutralized by adding 1N HCl and 4 g of sodium chloride are added. 28 ml of methanol are added to the reaction medium. After stirring for 15 minutes, 72 ml of methanol are added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 30 minutes at a temperature in the region of 4°C. The supernatant is then separated out and discarded (80 ml). 80 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 30 minutes. The supernatant is separated out and discarded (80 ml). 80 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 30 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying, 1.6 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 57%.

25 **Purification:**

1.6 g of crude depolymerized heparin obtained above and 16 ml of distilled water are placed in a 50 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.6 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μ m membrane and 0.08 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding 1N HCl, followed by addition of 2 g of sodium chloride. The solution is then filtered through a 0.45 μ m membrane and 14 ml of methanol are then poured in, at a temperature in the region of 10°C. The solution is then brought to 20°C and stirred for 15 minutes. 36 ml of

- methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 30 minutes. The supernatant is then separated out and discarded (50 ml). 50 ml of
- 5 methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 30 minutes. The supernatant is separated out and discarded (50 ml). The precipitate in suspension is then filtered through a
- 10 No. 3 sintered glass. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying, 1.25 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 78%.
- 15 The characteristics of the depolymerized heparin thus obtained are as follows:
- mean molecular weight: 2400 daltons
anti-Xa activity: 154.3 IU/mg
anti-IIa activity: 1.6 IU/mg
- 20 anti-Xa activity/anti-IIa activity ratio: 96.4

EXAMPLE 11

Depolymerization and conversion to the sodium salt:

- 140 ml of dichloromethane are placed in a Erlenmeyer
- 25 flask A. 20 g (0.019 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A, are added slowly, with stirring. After total dissolution at a temperature in the region of 40°C and cooling to a temperature in
- 30 the region of 20°C, 3.5 ml (0.011 mol) of tert-butyliminotri(pyrrolidino)phosphorane are added. The mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 60 g of anhydrous sodium acetate in 600 ml of methanol is
- 35 prepared in a Erlenmeyer flask B. After total dissolution, 10 g of Hyflo supercel Celite are added to the solution. The reaction mixture in the Erlenmeyer flask A is poured over 1 minute 30 seconds into the methanolic sodium acetate solution at a temperature in

the region of 4°C, with magnetic stirring. After stirring for 5 minutes, the suspension is left to settle for 30 minutes. The clear part of the supernatant is separated out and discarded (400 ml).
5 400 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 1 hour 20 minutes. The supernatant is separated out and discarded (250 ml). 250 ml of methanol are added to the
10 sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 200 ml of methanol. The pale yellow wet solid is drained and then dried under
15 reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying, 5.39 g of depolymerized heparin (benzyl ester, sodium salt) are obtained. The yield obtained is 69%.

20 **Saponification:**

5 g (0.0076 mol) of the depolymerized heparin (benzyl ester, sodium salt) obtained above and 35 ml of water are placed in a 50 ml Erlenmeyer flask. The solution is filtered through a No. 3 sintered glass and rinsed with
25 twice 10 ml of water. The filtrate obtained is placed in a 250 ml Erlenmeyer flask. 1 ml (0.01 mol) of 30% caustic soda is added with magnetic stirring, at a temperature in the region of 4°C. After addition, the mixture is stirred for 2 hours. The solution is
30 neutralized by adding 1N HCl and 6 g of sodium chloride are added. 42 ml of methanol are added to the reaction medium. After stirring for 15 minutes, 104 ml of methanol are added, followed by stirring for 15 minutes. Stirring is then stopped and the suspension is
35 left to sediment for 1 hour at a temperature in the region of 4°C. The supernatant is then separated out and discarded (140 ml). 140 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment

for about 45 minutes. The supernatant is separated out and discarded (160 ml). 160 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate in suspension is then
5 filtered through a No. 3 sintered glass. The cake obtained is then washed with 100 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 40°C. After drying for 48 hours, 2.7 g of crude
10 depolymerized heparin (sodium salt) are obtained. The yield obtained is 59%.

Purification:

2.6 g of crude depolymerized heparin obtained above and
15 25 ml of distilled water are placed in a 50 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μ m membrane and 0.15 ml of
20 aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding 1N HCl, followed by addition of 3 g of sodium chloride. The solution is then filtered through a 0.45 μ m
25 membrane and 21 ml of methanol are then poured in, at a temperature in the region of 10°C. The solution is then brought to 20°C and stirred for 15 minutes. 54 ml of methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is
30 left to sediment for 20 minutes. The supernatant is then separated out and discarded (50 ml). 50 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 20 minutes. The
35 supernatant is separated out and discarded (50 ml). The precipitate in suspension is then filtered through a No. 3 sintered glass. The white cake obtained is then washed with 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a

temperature in the region of 40°C. After drying for 18 hours, 2.35 g of pure depolymerized heparin (sodium salt) are obtained. The yield obtained is 90%.

The characteristics of the depolymerized heparin thus

5 obtained are as follows:

mean molecular weight: 2400 daltons

anti-Xa activity: 167.5 IU/mg

anti-IIa activity: 1.1 IU/mg

anti-Xa activity/anti-IIa activity ratio: 152

10

EXAMPLE 12

Depolymerization and conversion to the sodium salt (0.05% water):

140 ml of dichloromethane are placed in a Erlenmeyer
15 flask. 20 g (0.019 mol) of benzyl ester of heparin (degree of esterification: 75%, benzethonium salt) obtained as described in example A, are added slowly, with stirring. 20 g of 4 A molecular sieve are added to the reaction medium and the water content is brought to
20 0.05%, while stirring slowly for 48 h. The supernatant is transferred under an inert atmosphere to a Erlenmeyer flask A. After total dissolution, 3.5 ml (0.012 mol) of 2-tert-butylimino-2-diethylamino-1,3-dimethylperhydro-1,2,3-diazaphosphorine are added. The
25 mixture is stirred at a temperature in the region of 20°C for 24 hours. During this time, a solution of 60 g of anhydrous sodium acetate in 600 ml of methanol is prepared in a Erlenmeyer flask B. After total dissolution, 10 g of Hyflo supercel Celite are added to
30 the solution. The reaction mixture in the Erlenmeyer flask A is poured over about 2 minutes into the methanolic sodium acetate solution at a temperature in the region of 4°C, with magnetic stirring. After stirring for 15 minutes, the suspension is left to
35 settle for 1 hour. The clear part of the supernatant is separated out and discarded (420 ml). 420 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 15 minutes. The precipitate is left to resediment for about 1 hour. The supernatant

is separated out and discarded (450 ml). 450 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 15 minutes. The suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with 200 ml of methanol. The pale yellow wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 50°C. After drying for 16 hours, 5.36 g of crude depolymerized heparin are obtained, as the sodium salt in Celite (10 g). The yield obtained is 68.6%.

Saponification:

5.36 g (0.00817 mol) of the sodium salt of crude depolymerized heparin in Celite (10 g) obtained above and 50 ml of water are placed in a 50 ml Erlenmeyer flask. The suspension is filtered through a No. 3 sintered glass and rinsed with 4 times 15 ml of water. The filtrate obtained is placed in a 150 ml Erlenmeyer flask. 1.01 ml (0.0122 mol) of 35% caustic soda are added with magnetic stirring, at a temperature in the region of 4°C. After addition, the mixture is stirred for 3 hours. The solution is neutralized by adding 1N HCl and 11 g of sodium chloride are added. 77 ml of methanol are added to the reaction medium. After stirring for 15 minutes, 200 ml of methanol are added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 1 hour at a temperature in the region of 4°C. The supernatant is then separated out and discarded (240 ml). 240 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 10 minutes. The precipitate is left to resediment for about 30 minutes. The supernatant is separated out and discarded (225 ml). 225 ml of methanol are added to the sedimented precipitate and the mixture is stirred for 10 minutes. The suspension is then filtered through a No. 3 sintered glass. The cake obtained is then washed with twice 50 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a

temperature in the region of 40°C. After drying for 18 hours, 2.65 g of crude depolymerized heparin (sodium salt) are obtained. The yield obtained is 53.7%.

5 **Purification:**

2.65 g of crude depolymerized heparin obtained above and 26.5 ml of distilled water are placed in a 100 ml Erlenmeyer flask. The mixture is brought to 40°C with magnetic stirring. The pH is brought to 9.7 ± 0.1 by
10 addition of 1N sodium hydroxide. The reaction medium is filtered through a 0.45 μ m membrane and 0.25 ml of aqueous 30% hydrogen peroxide solution is added. After stirring for about 2 hours at a temperature in the region of 20°C, the mixture is neutralized by adding
15 1N HCl, followed by addition of 3 g of sodium chloride. The solution is then filtered through a 0.45 μ m membrane and 21 ml of methanol are then poured in, at a temperature in the region of 10°C. The solution is then brought to 20°C and stirred for 15 minutes. 54 ml of
20 methanol are then added, followed by stirring for 1 hour. Stirring is then stopped and the suspension is left to sediment for 45 minutes. The supernatant is then separated out and discarded (46 ml). 46 ml of methanol are added to the sedimented precipitate and
25 the mixture is stirred for 5 minutes. The precipitate is left to resediment for about 30 minutes. The supernatant is separated out and discarded (50 ml). 50 ml of methanol are added and the precipitate in suspension is then filtered through a No. 4 sintered
30 glass. The white cake obtained is then washed with 2 portions of 10 ml of methanol. The wet solid is drained and then dried under reduced pressure (6 kPa) at a temperature in the region of 50°C. After drying for 18 hours, 2.363 g of pure depolymerized heparin (sodium
35 salt) are obtained. The yield obtained is 89.1%.

The characteristics of the depolymerized heparin thus obtained are as follows:

mean molecular weight: 2500 Daltons

anti-Xa activity: 192 IU/mg

anti-IIa activity: 1.3 IU/mg

anti-Xa activity/anti-IIa activity ratio: 148


CLAIMS

1) A mixture of sulfated oligosaccharides having the general structure of the constituent polysaccharides of heparin and having the following characteristics:

- they have a mean molecular weight of 1500 to 3000 daltons, an anti-Xa activity of 120 to 200 IU/mg, an anti-IIa activity of less than 10 IU/mg and an anti-Xa activity/anti-IIa activity ratio of greater than 30,

the constituent oligosaccharides of the mixtures

- contain 2 to 26 saccharide units,
- have a 4,5-unsaturated uronic acid 2-O-sulfate unit at one of their ends,
- contain the hexasaccharide Δ IIa-II_s-Is of formula:



and are in the form of an alkali or alkaline-earth metal salt.

2) The mixture of oligosaccharides as claimed in claim 1, characterized in that the alkali or alkaline-earth metal salt is chosen from sodium, potassium, calcium and magnesium salts.

3) The mixture of oligosaccharides as claimed in claim 1 or 2, characterized in that it contains a

hexasaccharide fraction representing from 15 to 25% of the mixture of oligosaccharides.

4) The mixture of oligosaccharides as claimed in claim 3, characterized in that it contains from 8 to 15% of the hexasaccharide Δ IIa-II $\bar{s}$ -Is in the hexasaccharide fraction of the mixture of oligosaccharides.

5) The mixture of oligosaccharides as claimed in any one of claims 1 to 4, characterized in that it exhibits an anti-Xa activity of between 150 IU/mg and 200 IU/mg.

6) The mixture of oligosaccharides as claimed in any one of claims 1 to 5, characterized in that it has an anti-IIa activity of less than 5 IU/mg and in particular of 0.5 to 3.5 IU/mg.

7) The mixture of oligosaccharides as claimed in any one of claims 1 to 6, characterized in that it exhibits an anti-Xa activity/anti-IIa activity ratio greater than 50 and in particular greater than 100.


8) The mixture of oligosaccharides as claimed in any one of claims 1 to 7, characterized in that it has a mean molecular weight of between 2000 and 3000 Daltons and in particular from 2400 to 2650 Da.

9) The mixture of oligosaccharides as claimed in any one of claims 1 to 8, characterized in that it exhibits an anti-Xa activity of between 150 and 200 IU/mg, an anti-IIa activity of between 0.5 and 3.5 IU/mg, and a mean molecular weight of between 2400 and 2650 Da.

10) A method for preparing mixtures of oligosaccharides as claimed in any one of claims 1 to 9, during which the quaternary ammonium salt of the benzyl ester of heparin is depolymerized in an organic medium in the presence of a strong organic base with a pKa greater than 20, characterized in that the organic base

used consists of the family of phosphazenes, in particular in a dichloromethane solution, contains a percentage of water of less than 0.6%.

- 5 **11)** The method as claimed in claim 10 for preparing mixtures of oligosaccharides as claimed in claim 1, characterized in that the water content is less than 0.3% and preferably less than 0.2%.
- 10 **12)** The method of preparation as claimed in claim 10 or 11, characterized in that the quaternary ammonium salt of the benzyl ester of heparin is the benzethonium, cetylpyridinium or cetyltrimethylammonium salt.
- 15 **13)** The method of preparation as claimed in claim 10 or 11, characterized in that the bases of the family of phosphazenes are those of formula:



20

in which the radicals R_1 to R_7 are identical or different and represent linear or branched alkyl radicals containing from 1 to 6 carbon atoms.

- 25 **14)** The method of preparation as claimed in claim 10 or 11, characterized in that the strong base/ester mol ratio is between 0.2 and 5, and preferably between 0.6 and 2.
- 30 **15)** A method for preparing oligosaccharides as claimed in any one of claims 1 to 9 from heparins, in which the following operations are carried out:
- a) transalification of sodium heparin by the action of benzethonium chloride,

- b) esterification of benzethonium heparinate obtained by the action of benzyl chloride,
- c) transalification of the benzyl ester obtained and obtaining of the quaternary ammonium salt,
- 5 d) depolymerization of the quaternary ammonium salt of the benzyl ester of heparin by the method as defined in claim 10 or 11,
- e) conversion of the quaternary ammonium salt to a sodium salt,
- 10 f) optionally saponification of the heparin by the action of a base such as sodium hydroxide,
- g) optionally purification in particular by the action of an oxidizing agent such as hydrogen peroxide.

15 **16)** The method as claimed in claim 15, characterized in that the reaction of step a) is carried out by the action of benzethonium chloride in excess, on sodium heparin, at a temperature in the region of 15 to 25°C, with a salt/ sodium heparin molar ratio of between 3
20 and 4.

17) The method as claimed in claim 15, characterized in that the esterification of step b) is carried out in a chlorinated organic solvent such as chloroform or
25 methylene chloride, at a temperature of between 25 and 45°C, preferably between 30 and 40°C, and the ester in the form of a sodium salt is then recovered by precipitation by means of sodium acetate at 10% by weight in an alcohol such as methanol in a proportion
30 of 1 to 1.2 volumes of alcohol per volume of reaction medium.

18) The method as claimed in claim 15 or 17, characterized in that the degree of esterification of
35 the quaternary ammonium salt of the benzyl ester of heparin is between 50 and 100%, and preferably between 70 and 90%.

19) The method as claimed in claim 15, 17 or 18, characterized in that 0.5 to 1.5 parts by weight of benzyl chloride per 1 part by weight of benzethonium salt of heparin are used with a reaction time which
5 will be between 10 and 35 h.

20) The method as claimed in claim 15, characterized in that the transalification of step c) is carried out by means of a quaternary ammonium chloride, and preferably
10 by means of benzethonium chloride, cetylpyridinium chloride or cetyltrimethylammonium chloride, in an aqueous medium at a temperature between 10 and 25°C.

21) The method as claimed in claim 20, characterized in
15 that the quaternary ammonium chloride/sodium salt of the benzyl ester of heparin mol ratio is between 2 and 3.

22) The method as claimed in claim 15, characterized in
20 that the conversion to a sodium salt of the quaternary ammonium salt of the benzyl ester of depolymerized heparin (step e) is carried out by treating the reaction medium with an alcoholic solution of sodium acetate, and preferably with a 10% solution of sodium
25 acetate in methanol (weight/volume), at a temperature between 15 and 25°C.

23) The method as claimed in claim 15, characterized in that the saponification (step f) is carried out by
30 means of an alkali metal hydroxide such as sodium hydroxide, potassium hydroxide or lithium hydroxide, in an aqueous medium, at a temperature between 0 and 20°C, and preferably 0 and 10°C.

35 24) The method as claimed in claim 23, characterized in that 1 to 5 molar equivalents of alkali metal hydroxide and preferably 1 to 2 molar equivalents of sodium hydroxide are used.

25) The method as claimed in claim 15, characterized in that the purification (step g) is carried out by means of hydrogen peroxide, in an aqueous medium, at a temperature of 10 to 50°C and preferably between 20 and 40°C.

26) As a medicament, the oligosaccharides as claimed in any one of claims 1 to 9.

27) As a medicament having an antithrombotic activity, the oligosaccharides as claimed in any one of claims 1 to 9.

28) The medicaments as claimed in claim 26 or 27, for the prevention or treatment of venous and arterial thromboses, deep vein thrombosis, pulmonary embolism, unstable angina, myocardial infarction, cardiac ischemia, occlusive diseases of the peripheral arteries and atrial fibrillation, the proliferation of smooth muscle cells, atherosclerosis and arteriosclerosis, cancer by modulating angiogenesis and growth factors, and diabetic disorders such as diabetic retinopathies and nephropathies.

29) A pharmaceutical composition containing at least one medicament as defined in claim 26 and one or more pharmaceutically inert excipients or vehicles or additives.

30) The pharmaceutical composition as claimed in claim 29, characterized in that it consists of solutions for injection by the subcutaneous or intravenous route.

31) The pharmaceutical composition as claimed in claim 29, characterized in that it consists of a formulation for inhalation intended for the pulmonary route.

32) The pharmaceutical composition as claimed in claim 29, characterized in that it consists of a formulation for administration intended for the oral route.

- 5 33) A mixture of polysaccharides as defined in any one of claims 1 to 9, which can be obtained by the method as defined in claim 15.