

**METHODS OF USING A G PROTEIN-COUPLED RECEPTOR TO IDENTIFY
PEPTIDE YY (PYY) SECRETAGOGUES AND COMPOUNDS USEFUL IN THE
TREATMENT OF CONDITIONS MODULATED BY PYY**

5 **CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims priority to U.S. Provisional Application No. 61/123,273, filed on April 7, 2008, the disclosure of which is incorporated by reference in its entirety in this application.

10 **FIELD OF THE INVENTION**

The present invention relates to methods of using GPR119 receptor to identify peptide YY (PYY) secretagogues and compounds useful in the treatment of conditions modulated by PYY, such as conditions modulated by stimulation of NPY Y2 receptor (Y2R). Agonists of GPR119 receptor are useful as therapeutic agents for treating or preventing a condition
15 modulated by PYY, such as a condition modulated by stimulation of Y2R. Conditions modulated by PYY such as may be a condition modulated by stimulation of Y2R include bone-related conditions, metabolic disorders, angiogenesis-related conditions, ischemia-related conditions, convulsive disorders, malabsorptive disorders, cancers, and inflammatory disorders.

20 **BACKGROUND OF THE INVENTION**

The following discussion is intended to facilitate the understanding of the invention, but is not intended nor admitted to be prior art to the invention.

A. Osteoporosis

Osteoporosis is a disabling disease characterized by the loss of bone mass and
25 microarchitectural deterioration of skeletal structure leading to compromised bone strength, which predisposes a patient to increased risk of fragility fractures. Osteoporosis affects more than 75 million people in Europe, Japan and the United States, and causes more than 2.3 million fractures in Europe and the United States alone. In the United States, osteoporosis affects at least 25% of all post-menopausal white women, and the proportion rises to 70% in women older
30 than 80 years. One in three women older than 50 years will have an osteoporotic fracture that causes a considerable social and financial burden on society. The disease is not limited to women; older men also can be affected. By 2050, the worldwide incidence of hip fracture in men is projected to increase by 310% and 240% in women. The combined lifetime risk for hip, forearm, and vertebral fractures presenting clinically is around 40%, equivalent to the risk for
35 cardiovascular disease. Osteoporotic fractures therefore cause substantial mortality, morbidity, and economic cost. With an ageing population, the number of osteoporotic fractures and their costs will at least double in the next 50 years unless effective preventive strategies are developed. (See, e.g., Atik et al., Clin Orthop Relat Res (2006) 443:19-24; Raisz, J Clin Invest

(2005) 115:3318-3325; and World Health Organization Technical Report Series 921 (2003), Prevention and Management of Osteoporosis.)

B. Obesity and Diabetes

Obesity is a major public health concern because of its increasing prevalence and associated health risks. Moreover, obesity may affect a person's quality of life through limited mobility and decreased physical endurance as well as through social, academic and job discrimination.

Obesity and being overweight are generally defined by body mass index (BMI), which is correlated with total body fat and serves as a measure of the risk of certain diseases. BMI is calculated by weight in kilograms divided by height in meters squared (kg/m^2). Overweight is typically defined as a BMI of 25-29.9 kg/m^2 , and obesity is typically defined as a BMI of 30 kg/m^2 or higher. See, e.g., National Heart, Lung, and Blood Institute, Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults, The Evidence Report, Washington, DC: U.S. Department of Health and Human Services, NIH publication no. 98-4083 (1998).

Recent studies have found that obesity and its associated health risks are not limited to adults, but also affect children and adolescents to a startling degree. According to the Center for Disease Control, the percentage of children and adolescents who are defined as overweight has more than doubled since the early 1970s, and about 15 percent of children and adolescents are now overweight. Risk factors for heart disease, such as high cholesterol and high blood pressure, occur with increased frequency in overweight children and adolescents compared with normal-weight subjects of similar age. Also, type 2 diabetes, previously considered an adult disease, has increased dramatically in children and adolescents. Overweight conditions and obesity are closely linked to type 2 diabetes. It has recently been estimated that overweight adolescents have a 70% chance of becoming overweight or obese adults. The probability increases to about 80% if at least one parent is overweight or obese. The most immediate consequence of being overweight as perceived by children themselves is social discrimination.

Overweight or obese individuals are at increased risk for ailments such as hypertension, dyslipidemia, type 2 (non-insulin dependent) diabetes, insulin resistance, glucose intolerance, hyperinsulinemia, coronary heart disease, angine pectoris, congestive heart failure, stroke, gallstones, cholecystitis, cholelithiasis, gout, osteoarthritis, obstructive sleep apnea and respiratory problems, gall bladder disease, certain forms of cancer (e.g., endometrial, breast, prostate, and colon) and psychological disorders (such as depression, eating disorders, distorted body image and low self esteem). The negative health consequences of obesity make it the second leading cause of preventable death in the United States and impart a significant economic and psychosocial effect on society. See, e.g., McGinnis et al, JAMA (1993) 270:2207-2212.

Obesity is now recognized as a chronic disease that requires treatment to reduce its associated health risks. Although weight loss is an important treatment outcome, one of the main goals of obesity management is to improve cardiovascular and metabolic values to reduce obesity-related morbidity and mortality. It has been shown that 5-10% loss of body weight can substantially improve metabolic values, such as blood glucose, blood pressure, and lipid concentrations. Hence, it is believed that a 5-10% reduction in body weight may reduce morbidity and mortality.

Currently available prescription drugs for managing obesity generally reduce weight by decreasing dietary fat absorption, as with orlistat, or by creating an energy deficit by reducing food intake and/or increasing energy expenditure, as seen with sibutramine. The search for alternatives to presently available antiobesity agents has taken several paths one of which has focused on certain gut peptides that have been implicated in modulating satiety such as peptide YY (PYY) (Chaudhri et al, Annu Rev Physiol (2008) 70:239-255).

C. Atherosclerosis

Atherosclerosis is a complex disease characterized by inflammation, lipid accumulation, cell death and fibrosis. Atherosclerosis is characterized by cholesterol deposition and monocyte infiltration into the subendothelial space, resulting in foam cell formation. Thrombosis subsequent to atherosclerosis leads to myocardial infarction and stroke. Atherosclerosis is the leading cause of mortality in many countries, including the United States. (See, e.g., Ruggeri, Nat Med (2002) 8:1227-1234; Arehart et al, Circ Res, 2008, Mar 6 Epub ahead of print.)

D. Inflammatory Bowel Disease (IBD)

Inflammatory bowel disease (IBD) is the general name for diseases that cause inflammation in the intestines and includes, e.g. Crohn's disease, ulcerative colitis, ulcerative proctitis. U.S. medical costs of inflammatory bowel disease for 1990 have been estimated to be \$1.4 to \$1.8 billion. Lost productivity has been estimated to have added an additional \$0.4 to \$0.8 billion, making the estimated cost of inflammatory bowel disease \$1.8 to \$2.6 billion. (See, e.g., Pearson, Nursing Times (2004) 100:86-90; Hay et al, J Clin Gastroenterol (1992) 14:309-317; Keighley et al, Ailment Pharmacol Ther (2003) 18:66-70.)

Enteritis refers to inflammation of the intestine, especially the small intestine, a general condition that can have any of numerous different causes. Enterocolitis refers to inflammation of the small intestine and colon.

Crohn's disease (CD) is an inflammatory process that can affect any portion of the digestive tract, but is most commonly seen in the last part of the small intestine otherwise called the (terminal) ileum and cecum. Altogether this area is also known as the ileocecal region. Other cases may affect one or more of: the colon only, the small bowel only (duodenum, jejunum and/or ileum), the anus, stomach or esophagus. In contrast with ulcerative colitis, CD usually does not affect the rectum, but frequently affects the anus instead. The inflammation extends deep into the lining of

the affected organ. The inflammation can cause pain and can make the intestines empty frequently, resulting in diarrhea. Crohn's disease may also be called enteritis. Granulomatous colitis is another name for Crohn's disease that affects the colon. Ileitis is CD of the ileum which is the third part of the small intestine. Crohn's colitis is CD affecting part or all of the colon.

- 5 Ulcerative colitis (UC) is an inflammatory disease of the large intestine, commonly called the colon. UC causes inflammation and ulceration of the inner lining of the colon and rectum. The inflammation of UC is usually most severe in the rectal area with severity diminishing (at a rate that varies from patient to patient) toward the cecum, where the large and small intestine join. Inflammation of the rectum is called proctitis. Inflammation of the sigmoid colon (located just
10 above the rectum) is called sigmoiditis. Inflammation involving the entire colon is termed pancolitis. The inflammation causes the colon to empty frequently resulting in diarrhea. As the lining of the colon is destroyed ulcers form releasing mucus, pus and blood. Ulcerative proctitis is a form of UC that affects only the rectum.

E. Peptide YY (PYY)

- 15 Peptide YY (PYY) is a 36 amino acid peptide originally isolated in 1980 from porcine intestine (Tatemoto et al, Nature (1980) 285:417-418). PYY is secreted from enteroendocrine L-cells within both the large and small intestine. It has been shown that in rat and human gut concentrations of immunoreactive PYY are low in duodenum and jejunum, high in ileum and colon, and highest in rectum (Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al,
20 Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16). (PYY expression in rat been reported to also extend to alpha cells of the islets of Langerhans and to cells in the medulla oblongata (Ekblad et al, Peptides (2002) 23:251-261; PYY is released into the circulation as PYY₁₋₃₆ and PYY₃₋₃₆ (Eberlein et al, Peptides (1989) 10:797-803). PYY₃₋₃₆ is generated from PYY₁₋₃₆ by cleavage of the N-terminal
25 Tyr and Pro residues by dipeptidyl peptidase IV. PYY₃₋₃₆ is the predominant form of PYY in human postprandial plasma (Grandt et al, Regul Pept (1994) 51:151-159). PYY₁₋₃₆ and PYY₃₋₃₆ have been reported to have comparable agonist activity at NPY Y2 receptor (Y2R), a G protein-coupled receptor (Parker et al, Br J Pharmacol (2008) 153:420-431); however, PYY₃₋₃₆ has been reported to be a high-affinity Y2R selective agonist (Keire et al, Am J Physiol Gastrointest Liver
30 Physiol (2000) 279:G126-G131). PYY was subsequently reported to reduce high-fat food intake in rats after peripheral administration (Okada et al, Endocrinology Supplement (1993) 180) and to cause weight loss in mice after peripheral administration (Morley et al, Life Sciences (1987) 41:2157-2165).

- 35 Peripheral administration of PYY₃₋₃₆ has been reported to markedly reduce food intake and weight gain in rats, to decrease appetite and food intake in humans, and to decrease food intake in mice, but not in Y2R-null mice, which was said to suggest that the food intake effect requires the Y2R. In human studies, infusion of PYY₃₋₃₆ was found to significantly decrease

appetite and reduce food intake by 33% over 24 hours. Infusion of PYY₃₋₃₆ to reach the normal postprandial circulatory concentrations of the peptide led to peak serum levels of PYY₃₋₃₆ within 15 minutes, followed by a rapid decline to basal levels within 30 minutes. It was reported that there was significant inhibition of food intake in the 12-hour period following the PYY₃₋₃₆ infusion, but was essentially no effect on food intake in the 12-hour to 24-hour period. In a rat study, repeated administration of PYY₃₋₃₆ intraperitoneally (injections twice daily for 7 days) reduced cumulative food intake (Batterham et al, Nature (2002) 418:650-654; Renshaw et al, Current Drug Targets (2005) 6:171-179).

Peripheral administration of PYY₃₋₃₆ has been reported to reduce food intake, body weight gain and glycemic indices in diverse rodent models of metabolic diseases of both sexes (Pittner et al, Int J Obes Relat Metab Disord (2004) 28:963-971). It has been reported that blockade of Y2R with the specific antagonist BIIIE-246 attenuates the effect of peripherally administered endogenous and exogenous PYY₃₋₃₆ for reducing food intake (Abbott et al, Brain Res (2005) 1043:139-144). It has been reported that peripheral administration of a novel long-acting selective Y2R polyethylene glycol-conjugated peptide agonist reduces food intake and improves glucose metabolism (glucose disposal, plasma insulin and plasma glucose) in rodents (Ortiz et al, JPET (2007) 323:692-700; Lamb et al, J Med Chem (2007) 50:2264-2268). It has been reported that PYY ablation in mice leads to the development of hyperinsulinemia and obesity (Boey et al, Diabetologia (2006) 49:1360-1370). It has been reported that peripheral administration of a long-acting, potent and highly selective Y2R agonist inhibits food intake and promotes fat metabolism in mice (Balasubramaniam et al, Peptides (2007) 28:235-240).

There is evidence that agents which stimulate PYY synthesis *in vivo* can confer protection against diet-induced and genetic obesity and can improve glucose tolerance (Boey et al, Neuropeptides (2008) 42:19-30).

It has been reported that Y2R agonists such as PYY₁₋₃₆ and PYY₃₋₃₆ can confer protection against epileptic seizures, such as against kainate seizures (El Bahh et al, Eur J Neurosci (2005) 22:1417-1430; Woldbye et al, Neurobiology of Disease (2005) 20:760-772).

It has been reported that Y2R agonists such as PYY₁₋₃₆ and PYY₃₋₃₆ act as proabsorbative (or anti-secretory) hormones, increasing upon intravenous administration the absorption of both water and sodium in various parts of the bowel (Bilchik et al, Gastroenterol (1993) 105:1441-1448; Liu et al, J Surg Res (1995) 58:6-11; Nightingale et al, Gut (1996) 39:267-272; Liu et al, Am Surg (1996) 62:232-236; Balasubramaniam et al, J Med Chem (2000) 43:3420-3427). It has been reported that Y2R agonists such as PYY analogues inhibit secretion and promote absorption and growth in the intestinal epithelium (Balasubramaniam et al, J Med Chem (2000) 43:3420-3427). It has been reported that PYY promotes intestinal growth in normal rats (Gomez et al, Am J Physiol (1995) 268:G71-G81). It has been reported that Y2R agonists such

as PYY₁₋₃₆ and PYY₃₋₃₆ inhibit bowel motility and work to prevent diarrhea (EP1902730; also see Cox, Peptides (2007) 28:345-351).

It has been reported that Y2R agonists such as PYY₁₋₃₆ and PYY₃₋₃₆ can confer protection against inflammatory bowel disease such as ulcerative colitis and Crohn's disease (WO 03/105763). It has been reported that PYY-deficient mice exhibit an osteopenic phenotype, i.e. that PYY can increase bone mass and/or can confer protection against loss of bone mass (e.g., decreases loss of bone mass) (Wortley et al, Gastroenterol (2007) 133:1534-1543). It has been reported that PYY₃₋₃₆ can confer protection in rodent models of pancreatitis (Vona-Davis et al, Peptides (2007) 28:334-338).

It has been reported that angiogenesis is impaired in Y2R-deficient mice (Lee et al, Peptides (2003) 24:99-106), i.e. that agonists of Y2R such as PYY₁₋₃₆ and PYY₃₋₃₆ promote angiogenesis. It has been reported that wound healing is impaired in Y2R-deficient mice (Ekstrand et al, PNAS USA (2003) 100:6033-6038), i.e. that agonists of Y2R such as PYY₁₋₃₆ and PYY₃₋₃₆ promote wound healing. It has been reported that ischemic angiogenesis is impaired in Y2R-deficient mice (Lee et al, J Clin Invest (2003) 111:1853-1862), i.e. that agonists of Y2R such as PYY₁₋₃₆ and PYY₃₋₃₆ promotes revascularization and restoration of function of ischemic tissue. It has been reported that agonists of Y2R such as PYY₁₋₃₆ and PYY₃₋₃₆ mediate increases in collateral-dependent blood flow in a rat model of peripheral arterial disease (Cruze et al, Peptides (2007) 28:269-280).

It has been reported that PYY and Y2R agonists such as PYY₃₋₃₆ can suppress tumor growth in the cases of, e.g., pancreatic cancer such as pancreatic ductal adenocarcinoma, breast cancer such as breast infiltrative ductal adenocarcinoma, colon cancer such as colon adenocarcinoma and Barrett's adenocarcinoma (Liu et al, Surgery (1995) 118:229-236; Liu et al, J Surg Res (1995) 58:707-712; Grise et al, J Surg Res (1999) 82:151-155; Tseng et al, Peptides (2002) 23:389-395; McFadden et al, Am J Surg (2004) 188:516-519).

It has been reported that stimulation of Y2R such as by PYY₃₋₃₆ leads to an increase in plasma adiponectin (Ortiz et al, JPET (2007) 323:692-700). Adiponectin is an adipokine with potent anti-inflammatory properties (Ouchi et al, Clin Chim Acta (2007) 380:24-30; Tilg et al, Nat Rev Immunol (2006) 6:772-783). Adiponectin exerts anti-atherogenic effects by targeting vascular endothelial cells and macrophages and insulin-sensitizing effects, predominantly in muscle and liver (Kubota et al, J Biol Chem (2002) 277:25863-25866; Maeda et al, Nat Med (2002) 8:731-737). Low adiponectin levels have been reported to be associated with atherogenic lipoproteins in dyslipidemia (elevated triglycerides, small dense LDL cholesterol, low HDL cholesterol) (Marso et al, Diabetes Care (2008) Feb 5 Epub ahead of print). Adiponectin has been implicated in high density lipoprotein (HDL) assembly (Oku et al, FEBS Letters (2007) 581:5029-5033). Adiponectin has been found to ameliorate the abnormalities of metabolic syndrome, including insulin resistance, hyperglycemia, and dyslipidemia, in a mouse

model of obesity-linked metabolic syndrome associated with decreased adiponectin levels (Hara et al, Diabetes Care (2006) 29:1357-1362). Adiponectin has been reported to stimulate angiogenesis in response to tissue ischemia (Shibata et al, J Biol Chem (2004) 279:28670-28674). Adiponectin has been reported to prevent cerebral ischemic injury through endothelial nitric oxide synthase-dependent mechanisms (Nishimura et al, Circulation (2008) 117:216-223). Adiponectin has been reported to confer protection against myocardial ischemia-reperfusion injury (Shibata et al, Nat Med (2005) 11:1096-1103; Tao et al, Circulation (2007) 115:1408-1416). Adiponectin has been reported to confer protection against myocardial ischaemia-reperfusion injury via AMP-activated protein kinase, Akt, and nitric oxide (Gonon et al, Cardiovasc Res (2008) 78:116-122). Adiponectin has been reported to confer protection against the development of systolic dysfunction following myocardial infarction, through its abilities to suppress cardiac hypertrophy and interstitial fibrosis, and protect against myocyte and capillary loss (Shibata et al, J Mol Cell Cardiol (2007) 42:1065-1074). Adiponectin has been reported to confer protection against inflammatory lung disease; adiponectin-deficient mice exhibit an emphysema-like phenotype (Summer et al, Am J Physiol Lung Cell Mol Physiol (March 7, 2008)). Adiponectin has been reported to confer protection against allergic airway inflammation and airway hyperresponsiveness such as may be associated with asthma (Shore et al, J Allergy Clin Immunol (2006) 118:389-395). Adiponectin has been suggested to confer protection against pulmonary arterial hypertension by virtue of its insulin-sensitizing effects (Hansmann et al, Circulation (2007) 115:1275-1284). Adiponectin has been reported to ameliorate obesity-related hypertension, with said amelioration of hypertension being associated in part with upregulated prostacyclin expression (Ohashi et al, Hypertension (2006) 47:1108-1116). Adiponectin has been reported to decrease tumor necrosis factor (TNF)- α -induced expression of the adhesion molecules VCAM-1, E-selectin and ICAM-1 in human aortic endothelial cells (HAECs) (Ouchi et al, Circulation (1999) 100:2473-2476) and to inhibit production of TNF- α in macrophages (Yokota et al, Blood (2000) 96:1723-1732). Adiponectin has been reported to confer protection against restenosis after vascular intervention (Matsuda et al, J Biol Chem (2002) 277:37487-37491). The central role of TNF- α in inflammation has been demonstrated by the ability of agents that block the action of TNF- α to treat a range of inflammatory conditions. TNF- α -mediated inflammatory conditions encompass rheumatoid arthritis, inflammatory bowel disease such as Crohn's disease, ankylosing spondylitis, psoriasis, ischemic brain injury, cardiac allograft rejection, asthma, and the like (Bradley, J Pathol (2008) 214:149-160). See, e.g., Yamamoto et al, Clinical Science (2002) 103:137-142; Behre, Scand J Clin Lab Invest (2007) 67:449-458; Guerre-Millo, Diabetes & Metabolism (2008) 34:12-18; Parker et al, Br J Pharmacol (2008) 153:420-431.

F. Treatment of Conditions Modulated by PYY

An attractive alternative approach for treatment of conditions modulated by PYY is to develop an orally active composition for increasing an endogenous level of PYY in an individual, such as an endogenous level of plasma PYY, such as by stimulation of PYY secretion in gut enteroendocrine cells.

5 **G. GPR119**

GPR119 is a G protein-coupled receptor (GPR119; *e.g.*, human GPR119, GenBank® Accession No. AAP72125 and alleles thereof; *e.g.*, mouse GPR119, GenBank® Accession No. AY288423 and alleles thereof). GPR119 stimulation as by an agonist leads to elevation of the level of intracellular cAMP, consistent with GPR119 being coupled to Gs. In the patent
10 literature, GPR119 has been referred to as RUP3 (*e.g.*, WO 00/31258); GPR119 has also been referred to as Glucose-Dependent Insulinotropic Receptor (GDIR).

H. G Protein-Coupled Receptors

Although a number of receptor classes exist in humans, by far the most abundant and therapeutically relevant is represented by the G protein-coupled receptor (GPCR) class. It is
15 estimated that there are some 30,000-40,000 genes within the human genome, and of these, approximately 2% are estimated to code for GPCRs.

GPCRs represent an important area for the development of pharmaceutical products. Drugs active at GPCRs have therapeutic benefit across a broad spectrum of human diseases as diverse as pain, cognitive dysfunction, hypertension, peptic ulcers, rhinitis, and asthma. Of the
20 approximately 500 clinically marketed drugs, greater than 30% are modulators of GPCR function. These drugs exert their activity at approximately 30 well-characterized GPCRs. (See, *e.g.*, Wise et al, Annu Rev Pharmacol Toxicol (2004) 44:43-66.)

GPCRs share a common structural motif, having seven sequences of between 22 to 24 hydrophobic amino acids that form seven alpha helices, each of which spans the membrane
25 (each span is identified by number, *i.e.*, transmembrane-1 (TM-1), transmembrane-2 (TM-2), *etc.*). The transmembrane helices are joined by strands of amino acids between transmembrane-2 and transmembrane-3, transmembrane-4 and transmembrane-5, and transmembrane-6 and transmembrane-7 on the exterior, or “extracellular” side, of the cell membrane (these are referred to as “extracellular” regions 1, 2 and 3 (EC-1, EC-2 and EC-3), respectively). The
30 transmembrane helices are also joined by strands of amino acids between transmembrane-1 and transmembrane-2, transmembrane-3 and transmembrane-4, and transmembrane-5 and transmembrane-6 on the interior, or “intracellular” side, of the cell membrane (these are referred to as “intracellular” regions 1, 2 and 3 (IC-1, IC-2 and IC-3), respectively). The “carboxy” (“C”) terminus of the receptor lies in the intracellular space within the cell, and the “amino”
35 (“N”) terminus of the receptor lies in the extracellular space outside of the cell.

Generally, when a ligand binds with the receptor (often referred to as “activation” of the receptor), there is a change in the conformation of the receptor that facilitates coupling between

the intracellular region and an intracellular "G-protein." It has been reported that GPCRs are "promiscuous" with respect to G proteins, *i.e.*, that a GPCR can interact with more than one G protein. *See*, Kenakin, Life Sciences (1988) 43:1095-1101. Although other G proteins exist, currently, Gq, Gs, Gi, Gz and Go are G proteins that have been identified. Ligand-activated GPCR coupling with the G-protein initiates a signaling cascade process (referred to as "signal transduction"). Under normal conditions, signal transduction ultimately results in cellular activation or cellular inhibition. Although not wishing to be bound to theory, it is thought that the IC-3 loop as well as the carboxy terminus of the receptor interact with the G protein.

There are also promiscuous G proteins, which appear to couple several classes of GPCRs to the phospholipase C pathway, such as G15 or G16 (Offermanns & Simon, J Biol Chem (1995) 270:15175-80), or chimeric G proteins designed to couple a large number of different GPCRs to the same pathway, *e.g.* phospholipase C (Milligan & Rees, Trends in Pharmaceutical Sciences (1999) 20:118-24).

Under physiological conditions, GPCRs exist in the cell membrane in equilibrium between two different conformations: an "inactive" state and an "active" state. A receptor in an inactive state is unable to link to the intracellular signaling transduction pathway to initiate signal transduction leading to a biological response. Changing the receptor conformation to the active state allows linkage to the transduction pathway (via the G-protein) and produces a biological response.

A receptor may be stabilized in an active state by a ligand or a compound such as a drug. Recent discoveries, including but not exclusively limited to modifications to the amino acid sequence of the receptor, provide means other than ligands or drugs to promote and stabilize the receptor in the active state conformation. These means effectively stabilize the receptor in an active state by simulating the effect of a ligand binding to the receptor. Stabilization by such ligand-independent means is termed "constitutive receptor activation."

SUMMARY OF THE INVENTION

The present invention relates to the unexpected discovery by Applicant that administration of a GPR119 agonist to an individual, such as by oral administration, can act to increase plasma total PYY in the individual. The present invention features methods relating to GPR119 for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, and compounds useful for a modulation of a PYY-related activity in an individual. A GPR119 agonist is useful for increasing plasma total PYY in an individual. In certain embodiments, the individual is a human.

Nucleotide sequence encoding human GPR119 polypeptide is given in SEQ ID NO: 1. The amino acid sequence of said encoded human GPR119 polypeptide is given in SEQ ID NO: 2.

In a *first aspect*, the invention features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- 5 (a) contacting a test compound with a host cell or with membrane of a host cell comprising a G protein-coupled receptor, wherein the G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:
- 10 (i) amino acids 1-335 of SEQ ID NO:2;
- (ii) amino acids 2-335 of SEQ ID NO:2;
- (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (i);
- 15 (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO:1;
- (v) a variant of SEQ ID NO: 2 when selected from the group consisting of:
- (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
- 20 (b') an amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
- (vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and
- (vii) a biologically active fragment of any one of (i) to (vi); and
- 25 (b) determining the ability of the test compound to stimulate functionality of the G protein-coupled receptor;

wherein the ability of the test compound to stimulate functionality of the G protein-coupled receptor is indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

30

In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2.

In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of SEQ ID NO: 2.

35 In certain embodiments, the variant of SEQ ID NO: 2 is an allele of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2.
In certain embodiments, the variant of SEQ ID NO: 2 is a mammalian ortholog of SEQ ID NO: 2.

In certain embodiments, the G protein-coupled receptor is recombinant.

5 In certain embodiments, the method is a method for identifying PYY secretagogues.

In certain embodiments, the method comprises identifying an agonist of the receptor.

In certain embodiments, the method comprises identifying a partial agonist of the receptor.

10 In certain embodiments, the method is a method for identifying compounds useful for treatment or prevention of a condition modulated by PYY.

In certain embodiments, the method is a method for identifying compounds useful for a modulation of a PYY-related activity in an individual.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds
15 useful for a modulation of a PYY-related activity in an individual, comprising steps (a) and (b) of this *first aspect*, and further comprising:

(c) optionally synthesizing a compound which stimulates the functionality of the receptor in step (b);

20 (d) contacting a compound which stimulates functionality of the receptor in step (b) in vitro with a vertebrate enteroendocrine cell or with a cell capable of secreting PYY; and

(e) determining whether the compound stimulates PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY;

25 wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate enteroendocrine cell is a mammalian
30 enteroendocrine cell. In certain embodiments, the enteroendocrine cell is an L cell. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from the small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from an L cell-containing region of small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from duodenum,
35 jejunum, ileum, colon or rectum tissue (*see, e.g.,* Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16). In certain embodiments, the enteroendocrine

cell is an enteroendocrine cell line. In certain embodiments, the cell capable of secreting PYY is a cell comprised by or derived from the islets of Langerhans. In certain embodiments, the cell capable of secreting PYY is a neuronal cell. In certain embodiments, the cell capable of secreting PYY is a recombinant cell engineered to be capable of secreting PYY.

5 The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising steps (a) and (b) of this *first aspect*, and further comprising:

- 10 (c) optionally synthesizing a compound which stimulates functionality of the receptor in step (b);
- (d) administering a compound which stimulates functionality of the receptor in step (b) to a vertebrate; and
- (e) determining whether the compound increases a PYY level in the vertebrate;

 wherein the ability of the test compound to increase a PYY level in the vertebrate is further
15 indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

 In certain embodiments, the PYY level is blood or plasma concentration of total PYY. In certain embodiments, the PYY level is blood or plasma concentration of PYY₃₋₃₆.

20 In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

 In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged
25 with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

 In certain embodiments, the human is an individual or a patient enrolled or undergoing treatment in a clinical trial.

30 The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising steps (a) and (b) of this *first aspect*, and further comprising:

- 35 (c) optionally synthesizing a compound which stimulates functionality of the receptor in step (b); and

- (d) determining whether the compound increases a PYY level in a sample, said sample having been obtained from a vertebrate previously administered with a compound which stimulates functionality of the receptor in step (b);

wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY; or compounds useful for a modulation of a PYY-related activity in an individual, comprising steps (a) and (b) of this *first aspect*, and further comprising:

- (c) optionally synthesizing a compound which stimulates functionality of the receptor in step (b);
- (d) optionally providing a compound which stimulates functionality of the receptor in step (b);
- (e) contacting a compound which stimulates functionality of the receptor in step (b) in vitro with a vertebrate enteroendocrine cell or with a cell capable of secreting PYY; and
- (f) determining whether the compound stimulates PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY;

wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate enteroendocrine cell is a mammalian enteroendocrine cell. In certain embodiments, the enteroendocrine cell is an L cell. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from the small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or

derived from tissue obtained from an L cell-containing region of small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from duodenum, jejunum, ileum, colon or rectum tissue (*see, e.g.,* Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16). In certain embodiments, the enteroendocrine cell is an enteroendocrine cell line. In certain embodiments, the cell capable of secreting PYY is a cell comprised by or derived from the islets of Langerhans. In certain embodiments, the cell capable of secreting PYY is a neuronal cell. In certain embodiments, the cell capable of secreting PYY is a recombinant cell engineered to be capable of secreting PYY.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity, comprising steps (a) and (b) of this *first aspect*, and further comprising:

- (c) optionally synthesizing a compound which stimulates functionality of the receptor in step (b);
- (d) optionally providing a compound which stimulates functionality of the receptor in step (b);
- (e) administering a compound which stimulates functionality of the receptor in step (b) to a vertebrate; and
- (f) determining whether the compound increases a PYY level in the vertebrate;

wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the PYY level is blood or plasma concentration of total PYY. In certain embodiments, the PYY level is blood or plasma concentration of PYY₃₋₃₆.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds

useful for a modulation of a PYY-related activity, comprising steps (a) and (b) of this *first aspect*, and further comprising:

- (c) optionally synthesizing a compound which stimulates functionality of the receptor in step (b);
 - 5 (d) optionally providing a compound which stimulates functionality of the receptor in step (b); and
 - (e) determining whether the compound increases a PYY level in a sample, said sample having been obtained from a vertebrate previously administered with a compound which stimulates functionality of the receptor in step (b);
- 10 wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the
15 vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain
embodiments, the human is a participant in a study reviewed by a governmental agency charged
with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In
20 certain embodiments, the governmental agency is the Food and Drug Administration of the
United States.

In certain embodiments, the identified PYY secretagogue, or the identified compound
useful for treatment or prevention of a condition modulated by PYY, or the identified compound
useful for a modulation of a PYY-related activity is an agonist of the receptor. In some
25 embodiments, the the identified PYY secretagogue, or the identified compound useful for
treatment or prevention of a condition modulated by PYY, or the identified compound useful for
a modulation of a PYY-related activity is a partial agonist of the receptor.

In certain embodiments, the G protein-coupled receptor is coupled to a G protein. In
certain embodiments, stimulation of the G protein-coupled receptor increases a level of
30 intracellular cAMP accumulation. In certain embodiments, the G protein is Gs.

In some embodiments, a G protein-coupled receptor of the invention is SEQ ID NO:2 or
an allele thereof. In some embodiments, a G protein-coupled receptor of the invention is an
allele of SEQ ID NO:2. In some embodiments, a G protein-coupled receptor of the invention is
an ortholog of SEQ ID NO:2.

35 In some embodiments, a G protein-coupled receptor of the invention is a G protein-
coupled receptor which specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-
isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In some

embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is a ligand. In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, a G protein-coupled receptor of the invention exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP accumulation. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, stringent hybridization conditions comprise hybridization at 42°C in a solution comprising 50% formamide, 5xSSC (1xSSC = 150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20µg/ml denatured, sheared salmon sperm DNA, followed by washing at 65°C in a solution comprising 0.1xSSC. Hybridization techniques are well known to the skilled artisan.

In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is SEQ ID NO:2 or an allele thereof. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an allele of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an ortholog of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP accumulation. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, the variant of SEQ ID NO:2 is a G protein-coupled receptor having at least about 80% identity to SEQ ID NO:2. In certain embodiments, the variant of SEQ ID NO:2 is a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO:2. In certain embodiments, the variant of SEQ ID NO:2 which is a G protein-coupled receptor is an endogenous G protein-coupled receptor. In certain embodiments, the variant of

SEQ ID NO:2 which is a G protein-coupled receptor is a non-endogenous G protein-coupled receptor.

In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is SEQ ID NO: 2 or an allele thereof. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is SEQ ID NO: 2 or an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In some embodiments, the G protein-coupled receptor is part of a fusion protein comprising a G protein. Techniques for making a GPCR:G fusion construct are well known to the skilled artisan (*see, e.g., International Application WO 02/42461*).

In certain embodiments, the host cell comprises an expression vector, said expression vector comprising a polynucleotide encoding the G protein-coupled receptor. An exemplary suitable expression vector is pCMV. Other suitable expression vectors will be readily apparent to those of ordinary skill in the art, and a wide variety of expression vectors are commercially available (*e.g.*, from Clontech, Palo Alto, CA; Stratagene, La Jolla, CA; and Invitrogen, Carlsbad, CA).

In some embodiments, the host cell is a vertebrate cell. In some embodiments, the host cell is a mammalian cell. In some embodiments, the mammalian host cell is selected from the group consisting of a 293 cell, a 293T cell, a CHO cell, an MCB3901 cell, and a COS-7 cell. In some embodiments, the host cell is a yeast cell. In some embodiments, the host cell is a melanophore cell. Other suitable host cells will be readily apparent to those of ordinary skill in the art, and a wide variety of cell lines are available from the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209.

In certain embodiments, said determining is consistent with the G protein-coupled receptor being a Gs-coupled receptor.

In some embodiments, said determining is consistent with the G protein-coupled receptor being coupled through a promiscuous G protein, such as Gα15 or Gα16, to the phospholipase C pathway. Promiscuous G proteins are well known to the skilled artisan (*see, e.g.*, Offermanns et al., J Biol Chem (1995) 270:15175-15180). In some embodiments, said determining is consistent with the G protein-coupled receptor being coupled through a chimeric G protein, *e.g.* to the phospholipase C pathway. Chimeric G proteins are well known to the skilled artisan (*see, e.g.*, Milligan et al., Trends in Pharmaceutical Sciences (1999) 20:118-124; and WO 02/42461).

In some embodiments, said determining is through the measurement of a level of a second messenger.

In some embodiments, said determining is through the measurement of a level of a second messenger selected from the group consisting of cyclic AMP (cAMP), cyclic GMP (cGMP), inositol 1,4,5-triphosphate (IP₃), diacylglycerol (DAG), MAP kinase activity, MAPK/ERK kinase kinase-1 (MEKK1) activity, and Ca²⁺. In some preferred embodiments, the second messenger is cAMP. In certain embodiments, a level of intracellular cAMP is increased.

In certain embodiments, said determining is carried out using isolated membrane comprising the G protein-coupled receptor.

In certain embodiments, said determining is through the use of a melanophore assay. In certain embodiments, a level of pigment dispersion is increased.

In some embodiments, said determining is through a reporter assay. In some embodiments, said reporter assay is CRE-Luc reporter assay.

In some embodiments, said determining is through the measurement of an activity mediated by increasing a level of intracellular cAMP.

In some embodiments, said determining is through CRE-Luc reporter assay. In certain embodiments, a level of luciferase activity is increased.

5 In some embodiments, said determining is through the measurement of GTP γ S binding to membrane comprising the G protein-coupled receptor. In certain embodiments, said GTP γ S is labeled with [³⁵S]. In certain embodiments, said GTP γ S binding to membrane comprising the GPCR is increased.

10 In some embodiments, the test compound is a small molecule. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a polypeptide. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not an antibody or an antigen-binding fragment thereof. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a lipid. In some embodiments, the test compound is a small molecule, with the proviso that
15 the small molecule is not a polypeptide or a lipid. In some embodiments, the test compound is a polypeptide. In some embodiments, the test compound is a polypeptide, with the proviso that the polypeptide is not an antibody or an antigen-binding fragment thereof. In some embodiments, the test compound is a lipid. In some embodiments, the test compound is not an antibody or an antigen-binding fragment thereof. In some embodiments, the test compound is
20 an antibody or an antigen-binding fragment thereof.

In some embodiments, the method further comprises the step of optionally determining the structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

25 In some embodiments, the method further comprises the step of optionally providing the name or structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

30 In some embodiments, said method further comprises the step of optionally producing or synthesizing the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

35 In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual as a pharmaceutical.

In some embodiments, said method further comprises evaluating the bioavailability of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity.

5 In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual into a pharmaceutical composition.

In certain embodiments, the condition modulated by PYY is a bone-related condition.

10 In certain embodiments, the bone-related condition is a condition characterized by low bone mass. In certain embodiments, the condition characterized by low bone mass is selected from the group consisting of osteopenia, osteoporosis, rheumatoid arthritis, osteoarthritis, periodontal disease, alveolar bone loss, osteotomy bone loss, childhood idiopathic bone loss, Paget's disease, bone loss due to metastatic cancer, osteolytic lesions, curvature of the spine, and loss of height.

15 In certain embodiments, the bone-related condition is a bone fracture. In certain embodiments, the bone fracture is selected from the group consisting of a traumatic bone fracture, a long-term bone fracture, and an osteoporotic bone fracture.

20 In certain embodiments, the bone-related condition is a bone disease. In certain embodiments, the bone disease is selected from the group consisting of osteopenia, osteoporosis, rheumatoid arthritis, osteoarthritis, periodontal disease, alveolar bone loss, osteotomy bone loss, childhood idiopathic bone loss, Paget's disease, bone loss due to metastatic cancer, osteolytic lesions, curvature of the spine, and loss of height.

25 In certain embodiments, the bone-related condition is enhanced bone healing or bone growth, wherein said enhanced bone healing or bone growth is selected from the group consisting of enhanced bone healing following facial reconstruction, maxillary reconstruction, mandibular reconstruction, periodontal disease or tooth extraction, enhanced long bone extension, enhanced prosthetic ingrowth, and increased bone synostosis.

In certain embodiments, the condition modulated by PYY is a metabolic disorder.

30 In certain embodiments, the metabolic disorder is selected from the group consisting of overweight, obesity, hyperphagia, diabetes (inclusive of type 1 diabetes and type 2 diabetes), type 2 diabetes, impaired glucose tolerance, insulin resistance, hyperinsulinemia, dyslipidemia, hypertension and metabolic syndrome.

In certain embodiments, the condition modulated by PYY is an angiogenesis-related condition.

35 In certain embodiments, the angiogenesis-related condition is selected from the group consisting of peripheral arterial disease, wound healing and ischemia.

In certain embodiments, the condition modulated by PYY is an ischemia-related condition.

In certain embodiments, the ischemia-related condition is selected from the group consisting of ischemic stroke, ischemic heart disease, myocardial infarction, ischemia-reperfusion injury, post-myocardial infarction remodeling, heart failure, and congestive heart failure.

In certain embodiments, the condition modulated by PYY is a malabsorptive disorder.

5 In certain embodiments, the malabsorptive disorder is selected from the group consisting of short bowel syndrome, enterocolitis, Crohn's disease, ulcerative colitis, irritable bowel syndrome, cholera, and a disorder accompanied by diarrhea.

In certain embodiments, the condition modulated by PYY is a convulsive disorder.

In certain embodiments, the convulsive disorder is epilepsy.

10 In certain embodiments, the condition modulated by PYY is a cancer.

In certain embodiments, the cancer is selected from the group consisting of pancreatic cancer, breast cancer, colon cancer, and Barrett's adenocarcinoma.

In certain embodiments, the condition modulated by PYY is an inflammatory disorder.

15 In certain embodiments, the inflammatory disorder is selected from the group consisting of atherosclerosis, atherothrombosis, insulin resistance, diabetes (inclusive of type 1 diabetes and type 2 diabetes), type 1 diabetes, ischemic stroke, pancreatitis, restenosis, an inflammatory lung disease, and an inflammatory bowel disease. In certain embodiments, the inflammatory lung disease is selected from the group consisting of asthma, emphysema, and pulmonary arterial hypertension. In certain embodiments, the inflammatory bowel disease is selected from the group consisting of
20 enteritis, enterocolitis, Crohn's disease, granulomatous colitis, ileitis, Crohn's colitis, ulcerative colitis, sigmoiditis, pancolitis, and ulcerative proctitis.

In certain embodiments, the modulation of a PYY-related activity is selected from the group consisting of:

- 25 (a) increasing plasma PYY;
- (b) increasing bone mass;
- (c) decreasing loss of bone mass;
- (d) increasing bone formation;
- (e) decreasing food uptake;
- (f) decreasing body weight;
- 30 (g) increasing satiety;
- (h) increasing glucose disposal;
- (i) decreasing plasma glucose;
- (j) increasing angiogenesis;
- (k) decreasing secretion in intestinal epithelium;
- 35 (l) increasing absorption in intestinal epithelium; and
- (m) increasing plasma adiponectin

The invention additionally features a process comprising performing a method according to this *first aspect*.

In a *second aspect*, the invention features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity, comprising the steps of:

- (a) contacting a compound in vitro with a vertebrate enteroendocrine cell or with a cell capable of secreting PYY, said compound having been identified by a method according to the *first aspect*; and
- (b) determining whether the compound stimulates PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY;

wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a PYY-related condition, or a compound useful for a modulation of a PYY-related activity.

- 15 In certain embodiments, the vertebrate enteroendocrine cell is a mammalian enteroendocrine cell. In certain embodiments, the enteroendocrine cell is an L cell. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from the small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from an L cell-containing region of small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from duodenum, jejunum, ileum, colon or rectum tissue (*see, e.g.,* Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16). In certain embodiments, the enteroendocrine cell is an enteroendocrine cell line. In certain embodiments, the cell capable of secreting PYY is a cell comprised by or derived from the islets of Langerhans. In certain embodiments, the cell capable of secreting PYY is a neuronal cell. In certain embodiments, the cell capable of secreting PYY is a recombinant cell engineered to be capable of secreting PYY.

- 25 The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- (a) administering a compound to a vertebrate, said compound having been identified by a method according to the *first aspect*; and
- (b) determining whether the compound increases a PYY level in the vertebrate;

- 35 wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a PYY-related condition, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the PYY level is blood or plasma concentration of total PYY. In certain embodiments, the PYY level is blood or plasma concentration of PYY₃₋₃₆.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

(a) determining whether the compound increases a PYY level in a sample, said sample having been obtained from a vertebrate previously administered with a compound which stimulates functionality of the receptor in step (b);

wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the PYY level is blood or plasma concentration of total PYY. In certain embodiments, the PYY level is blood or plasma concentration of PYY₃₋₃₆.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

In some embodiments, the test compound is a small molecule. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a polypeptide. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not an antibody or an antigen-binding fragment thereof. In some

embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a lipid. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a polypeptide or a lipid. In some embodiments, the test compound is a polypeptide. In some embodiments, the test compound is a polypeptide, with the proviso that
5 the polypeptide is not an antibody or an antigen-binding fragment thereof. In some embodiments, the test compound is a lipid. In some embodiments, the test compound is not an antibody or an antigen-binding fragment thereof. In some embodiments, the test compound is an antibody or an antigen-binding fragment thereof.

In some embodiments, the method further comprises the step of optionally determining
10 the structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

In some embodiments, the method further comprises the step of optionally providing the name or structure of the PYY secretagogue, the compound useful for treatment or prevention of
15 a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

In some embodiments, said method further comprises the step of optionally producing or synthesizing the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related
20 activity in an individual.

In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual as a pharmaceutical.

In some embodiments, said method further comprises evaluating the bioavailability of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity.
25

In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual into a pharmaceutical composition.
30

The invention additionally features a process comprising performing a method according to this *second aspect*.

In a *third aspect*, the invention features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds
35 useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- (a) contacting a GPR119 agonist in vitro with a vertebrate enteroendocrine cell or with a cell capable of secreting PYY; and
- (b) determining whether the GPR119 agonist stimulates PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY;

5 wherein the ability of the GPR119 agonist to stimulate PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY is indicative of the GPR119 agonist being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

10 In certain embodiments, the vertebrate enteroendocrine cell is a mammalian enteroendocrine cell. In certain embodiments, the enteroendocrine cell is an L cell. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from the small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from an L cell-containing region of small or large intestine. In
15 certain embodiments, the enteroendocrine cell is comprised by or derived from duodenum, jejunum, ileum, colon or rectum tissue (*see, e.g.,* Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16). In certain embodiments, the enteroendocrine cell is an enteroendocrine cell line. In certain embodiments, the cell capable of secreting PYY is
20 a cell comprised by or derived from the islets of Langerhans. In certain embodiments, the cell capable of secreting PYY is a neuronal cell. In certain embodiments, the cell capable of secreting PYY is a recombinant cell engineered to be capable of secreting PYY.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds
25 useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- (a) administering a GPR119 agonist to a vertebrate; and
- (b) determining whether the GPR119 agonist increases a PYY level in the vertebrate;

wherein the ability of the GPR119 agonist to increase a PYY level in the vertebrate is indicative of the GPR119 agonist being a PYY secretagogue, a compound useful for treatment or prevention of a
30 condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the PYY level is blood or plasma concentration of total PYY. In certain embodiments, the PYY level is blood or plasma concentration of PYY₃₋₃₆.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the
35 vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- (a) determining whether the compound increases a PYY level in a sample, said sample having been obtained from a vertebrate previously administered with a GPR119 agonist; administering a GPR119 agonist to a vertebrate; wherein the ability of the GPR119 agonist to increase a PYY level in the vertebrate is indicative of the GPR119 agonist being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the PYY level is blood or plasma concentration of total PYY. In certain embodiments, the PYY level is blood or plasma concentration of PYY₃₋₃₆.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

In certain embodiments, the GPR119 agonist is an agonist of an endogenous GPR119.

In certain embodiments, the GPR119 agonist is an agonist of human GPR119.

In certain embodiments, the GPR119 agonist is a GPR119 partial agonist.

In certain embodiments, the GPR119 agonist is a selective GPR119 agonist.

In certain embodiments, the GPR119 agonist is a small molecule. In some embodiments, the small molecule is not a polypeptide. In some embodiments, the small molecule is not an antibody or an antigen-binding fragment thereof. In some embodiments, the small molecule is not a lipid. In some embodiments, the small molecule is not a polypeptide or a lipid.

In certain embodiments, the GPR119 agonist is orally available.

In certain embodiments, the GPR119 agonist has an EC₅₀ value of less than about 10 μ M, less than about 1 μ M, less than about 100 nM, less than about 75 nM, less than about 50

nM, less than about 25 nM, less than about 15 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, or less than about 1 nM. In certain embodiments, the GPR119 agonist has an EC₅₀ value of less than about 10 μM, less than about 1 μM, less than about 100 nM, less than about 75 nM, less than about 50 nM, less than about 25 nM, less than about 15 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, or less than about 1 nM at human GPR119 having SEQ ID NO: 2. In certain embodiments, the GPR119 agonist has an EC₅₀ value of less than about 10 μM, less than about 1 μM, less than about 100 nM, less than about 75 nM, less than about 50 nM, less than about 25 nM, less than about 15 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, or less than about 1 nM at human GPR119 having SEQ ID NO: 2 in adenylyl cyclase assay (exemplary adenylyl cyclase assay is provided in Example 9, Example 10 and in Example 11, *infra*). In certain embodiments, the GPR119 agonist has an EC₅₀ value of less than about 10 μM, less than about 1 μM, less than about 100 nM, less than about 75 nM, less than about 50 nM, less than about 25 nM, less than about 15 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, or less than about 1 nM at human GPR119 having SEQ ID NO: 2 in melanophore assay (exemplary melanophore assay is provided in Example 12, *infra*).

Exemplary GPR119 agonists are disclosed, e.g., in International Application No. PCT/US2004/001267 (published as WO 04/065380); International Application No. PCT/US2004/005555 (published as WO 04/076413); International Application No. PCT/US2004/022327 (published as WO 05/007647); International Application No. PCT/US2004/022417 (published as WO 05/007658); International Application No. PCT/US2005/019318 (published as WO 2005/121121); International Application No. PCT/GB2004/050046 (published as WO 2005/061489); International Application No. PCT/US06/00567 (published as WO 2006/083491); International Application No. PCT/GB2005/050264 (published as WO 2006/067531); International Application No. PCT/GB2005/050265 (published as WO 2006/067532); International Application No. PCT/GB2005/050266 (published as WO 2006/070208); International Application No. PCT/JP02/09350 (published as WO 03/026661); International Application No. PCT/JP2005/018412 (published as WO 06/040966); International Application No. PCT/JP2005/019000 (published as WO 2006/043490); International Application No. PCT/GB2006/050176 (published as WO 2007/003960); International Application No. PCT/GB2006/050177 (published as WO 2007/003961); International Application No. PCT/GB2006/050178 (published as WO 2007/003962); International Application No. PCT/GB2006/050182 (published as WO 2007/003964); International Application No. PCT/JP02/09350 (published as WO 03/026661); International Application No.

PCT/GB2007/050183 (published as WO 2007/116229); International Application No.
PCT/GB2007/050190 (published as WO 2007/116230); International Application No.
PCT/US2007/073364 (published as WO 2008/008895); International Application No.
PCT/EP2007/058991 (published as WO 2008/025798); International Application No.
5 PCT/EP2007/058993 (published as WO 2008/025799); and International Application No.
PCT/EP2007/058995 (published as WO 2008/025800).

In certain embodiments, the method comprises providing the GPR119 agonist.

In certain embodiments, the GPR119 agonist is identifiable by a method according to the *first aspect*.

10 In certain embodiments, the method comprises carrying out a method according to the *first aspect* to identify the GPR119 agonist.

In certain embodiments, the method comprises having identified the GPR119 agonist by a method according to the *first aspect*.

15 In some embodiments, the method further comprises the step of optionally determining the structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

20 In some embodiments, the method further comprises the step of optionally providing the name or structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

25 In some embodiments, said method further comprises the step of optionally producing or synthesizing the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual as a pharmaceutical.

30 In some embodiments, said method further comprises evaluating the bioavailability of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity.

35 In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual into a pharmaceutical composition.

The invention additionally features a process comprising performing a method according to this *third aspect*.

In a *fourth aspect*, the invention features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- (a) contacting a G protein-coupled receptor with an optionally labeled known ligand for the receptor in the presence or absence of a test compound, wherein a ligand for the receptor is a ligand for GPR119 and wherein the G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:
 - (i) amino acids 1-335 of SEQ ID NO:2;
 - (ii) amino acids 2-335 of SEQ ID NO:2;
 - (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (i);
 - (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO:1;
 - (v) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
 - (b') the amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
 - (vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and
 - (vii) a biologically active fragment of any one of (i) to (vi); and
- (b) detecting the complex between said known ligand and said receptor; and
- (c) determining whether less of said complex is formed in the presence of the test compound than in the absence of the test compound;

wherein said determination is indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2.

In certain embodiments, the receptor comprises the amino acid sequence of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an allele of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is a mammalian ortholog of SEQ ID NO: 2.

5 In certain embodiments, the G protein-coupled receptor is recombinant.

In certain embodiments, the method is a method for identifying PYY secretagogues.

In certain embodiments, the method is a method for identifying compounds useful for treatment or prevention of a condition modulated by PYY.

10 In certain embodiments, the method is a method for identifying compounds useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the ligand for GPR119 is a GPR119 agonist. In certain embodiments, the ligand for GPR119 or the GPR119 agonist is a ligand or agonist of an endogenous vertebrate, mammalian or human GPR119 receptor. In certain embodiments, the ligand for GPR119 or the GPR119 agonist is a ligand or agonist of an endogenous human

15 GPR119 receptor. In certain embodiments, the ligand for GPR119 or the GPR119 agonist is identical to a compound disclosed in, e.g., in International Application No. PCT/US2004/001267 (published as WO 04/065380); International Application No. PCT/US2004/005555 (published as WO 04/076413); International Application No. PCT/US2004/022327 (published as WO 05/007647); International Application No.

20 PCT/US2004/022417 (published as WO 05/007658); International Application No. PCT/US2005/019318 (published as WO 2005/121121); International Application No. PCT/GB2004/050046 (published as WO 2005/061489); International Application No. PCT/US06/00567 (published as WO 2006/083491); International Application No. PCT/GB2005/050264 (published as WO 2006/067531); International Application No.

25 PCT/GB2005/050265 (published as WO 2006/067532); International Application No. PCT/GB2005/050266 (published as WO 2006/070208); International Application No. PCT/JP02/09350 (published as WO 03/026661); International Application No. PCT/JP2005/018412 (published as WO 06/040966); International Application No. PCT/JP2005/019000 (published as WO 2006/043490); International Application No.

30 PCT/GB2006/050176 (published as WO 2007/003960); International Application No. PCT/GB2006/050177 (published as WO 2007/003961); International Application No. PCT/GB2006/050178 (published as WO 2007/003962); International Application No. PCT/GB2006/050182 (published as WO 2007/003964); International Application No. PCT/JP02/09350 (published as WO 03/026661); International Application No.

35 PCT/GB2007/050183 (published as WO 2007/116229); International Application No. PCT/GB2007/050190 (published as WO 2007/116230); International Application No. PCT/US2007/073364 (published as WO 2008/008895); International Application No.

PCT/EP2007/058991 (published as WO 2008/025798); International Application No. PCT/EP2007/058993 (published as WO 2008/025799); or International Application No. PCT/EP2007/058995 (published as WO 2008/025800). In certain embodiments, the ligand for GPR119 or the GPR119 agonist is (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-
5 [1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the ligand for GPR119 is an endogenous ligand of an endogenous vertebrate, mammalian, or human GPR119 receptor.

In certain embodiments, the optionally labeled known ligand is a labeled known ligand. In certain embodiments, the labeled known ligand is a radiolabeled known ligand. Techniques
10 for radiolabeling a compound, such as for labeling a known ligand of a G protein-coupled receptor of the invention, are well known to the skilled artisan. See, e.g., International Application WO 04/065380. Also see, e.g., Example 14, *infra*.

Techniques for detecting the complex between a G protein-coupled receptor and a compound known to be a ligand for the G protein-coupled receptor are well known to the skilled
15 artisan. See, e.g., International Application WO 04/065380. Also see, e.g., Example 15, *infra*.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising steps (a) to (c) of this *fourth aspect*, and further comprising:

- 20 (d) optionally synthesizing a compound in the presence of which less of said complex is formed in step (c);
- (e) contacting a compound in the presence of which less of said complex is formed in step (c) in vitro with a vertebrate enteroendocrine cell or with a cell capable of secreting PYY; and
- 25 (f) determining whether the compound stimulates PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY;

wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a
30 condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate enteroendocrine cell is a mammalian enteroendocrine cell. In certain embodiments, the enteroendocrine cell is an L cell. In certain
35 embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from the small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from an L cell-containing region of small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from duodenum,

jejunum, ileum, colon or rectum tissue (*see, e.g.,* Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16). In certain embodiments, the enteroendocrine cell is an enteroendocrine cell line. In certain embodiments, the cell capable of secreting PYY is a cell comprised by or derived from the islets of Langerhans. In certain embodiments, the cell capable of secreting PYY is a neuronal cell. In certain embodiments, the cell capable of secreting PYY is a recombinant cell engineered to be capable of secreting PYY.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising steps (a) to (c) of this *fourth aspect*, and further comprising:

- (d) optionally synthesizing a compound in the presence of which less of said complex is formed in step (c);
 - (e) administering a compound in the presence of which less of said complex is formed in step (c) to a vertebrate; and
 - (f) determining whether the compound increases a PYY level in the vertebrate;
- wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising steps (a) to (c) of this *fourth aspect*, and further comprising:

- (d) optionally synthesizing a compound in the presence of which less of said complex is formed in step (c); and
- (e) determining whether the compound increases a PYY level in a sample obtained from a vertebrate, said vertebrate having been administered with a compound in the presence of which less of said complex is formed in step (c);

wherein the ability of the test compound to increase a PYY level in the vertebrate is indicative of the test compound being PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

5 In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

 In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In
10 certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

 The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds
15 useful for a modulation of a PYY-related activity in an individual, comprising steps (a) to (c) of this *fourth aspect*, and further comprising:

- (d) optionally synthesizing a compound in the presence of which less of said complex is formed according to step (c);
- (e) optionally providing a compound in the presence of which less of said complex is
20 formed according to step (c);
- (f) contacting a compound in the presence of which less of said complex is formed according to step (c) in vitro with a vertebrate enteroendocrine cell or with a cell capable of secreting PYY; and
- (g) determining whether the compound stimulates PYY secretion from the vertebrate
25 enteroendocrine cell or from the cell capable of secreting PYY;

 wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity
30 in an individual.

 In certain embodiments, the vertebrate enteroendocrine cell is a mammalian enteroendocrine cell. In certain embodiments, the enteroendocrine cell is an L cell. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from the small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or
35 derived from tissue obtained from an L cell-containing region of small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from duodenum, jejunum, ileum, colon or rectum tissue (*see, e.g., Lundberg et al, PNAS USA (1982) 79:4471-*

4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16). In certain embodiments, the enteroendocrine cell is an enteroendocrine cell line. In certain embodiments, the cell capable of secreting PYY is a cell comprised by or derived from the islets of Langerhans. In certain embodiments, the cell
5 capable of secreting PYY is a neuronal cell. In certain embodiments, the cell capable of secreting PYY is a recombinant cell engineered to be capable of secreting PYY.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising steps (a) to (c) of
10 this *fourth aspect*, and further comprising:

- (d) optionally synthesizing a compound in the presence of which less of said complex is formed according to step (c);
- (e) optionally providing a compound in the presence of which less of said complex is formed according to step (c);
- 15 (f) administering a compound in the presence of which less of said complex is formed in step (c) to a vertebrate; and
- (g) determining whether the compound increases a PYY level in the vertebrate;

wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or
20 prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

25 In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

30 The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising steps (a) to (c) of this *fourth aspect*, and further comprising:

- (d) optionally synthesizing a compound in the presence of which less of said complex
35 is formed according to step (c);
- (e) optionally providing a compound in the presence of which less of said complex is formed according to step (c); and

- (f) determining whether the compound increases a PYY level in a sample obtained from a vertebrate, said vertebrate having been administered with a compound in the presence of which less of said complex is formed in step (c);

wherein the ability of the test compound to increase a PYY level in the vertebrate is further

- 5 indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

- In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human
10 mammal. In certain embodiments, the mammal is a non-human mammal.

- In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the
15 United States.

In some embodiments, a G protein-coupled receptor of the invention is SEQ ID NO:2 or an allele thereof. In some embodiments, a G protein-coupled receptor of the invention is an allele of SEQ ID NO:2. In some embodiments, a G protein-coupled receptor of the invention is an ortholog of SEQ ID NO:2.

- 20 In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor which specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-
25 piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is a ligand. In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, a G protein-coupled receptor of the invention exhibits a detectable level of constitutive activity. In some embodiments, the
30 constitutive activity is for increasing a level of intracellular cAMP accumulation. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

- In certain embodiments, stringent hybridization conditions comprise hybridization at 42°C in a solution comprising 50% formamide, 5xSSC (1xSSC = 150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20µg/ml denatured, sheared salmon sperm DNA, followed by washing at 65°C in a solution comprising 0.1xSSC. Hybridization techniques are well known to the skilled artisan.
- 35

In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is SEQ ID NO:2 or an allele thereof. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an allele of SEQ ID NO:2. In certain
5 embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an ortholog of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments,
10 the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is a ligand. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-
15 methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive
20 activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, the variant of SEQ ID NO:2 is a G protein-coupled receptor having at least about 80% identity to SEQ ID NO:2. In certain embodiments, the variant of SEQ ID NO:2 is a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO:2. In certain embodiments, the variant of SEQ ID NO:2 which is a G protein-coupled
25 receptor is an endogenous G protein-coupled receptor. In certain embodiments, the variant of SEQ ID NO:2 which is a G protein-coupled receptor is a non-endogenous G protein-coupled receptor.

In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is SEQ ID NO: 2 or an allele thereof. In certain embodiments, the G
30 protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-
35 1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-

pyrimidin-4-yl}-amine is an agonist. In some embodiments, the the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to
5 undergo pigment dispersion.

In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is SEQ ID NO: 2 or an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is an allele of SEQ ID NO: 2. In certain embodiments, the G
10 protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the G protein-coupled receptor which
15 comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for
20 increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In some embodiments, the G protein-coupled receptor is part of a fusion protein comprising a G protein. Techniques for making a GPCR:G fusion construct are well known to the skilled artisan (*see, e.g.,* International Application WO 02/42461).

25 In certain embodiments, said determining is carried out using a host cell comprising the G protein-coupled receptor. In certain embodiments, the host cell comprises an expression vector, said expression vector comprising a polynucleotide encoding the GPCR. An exemplary suitable expression vector is pCMV. Other suitable expression vectors will be readily apparent to those of ordinary skill in the art, and a wide variety of expression vectors are commercially
30 available (*e.g.,* from Clontech, Palo Alto, CA; Stratagene, La Jolla, CA; and Invitrogen, Carlsbad, CA).

In some embodiments, the host cell is a vertebrate cell. In some embodiments, the host cell is a mammalian cell. In some embodiments, the mammalian host cell is selected from the group consisting of a 293 cell, a 293T cell, a CHO cell, a MCB3901 cell, and a COS-7 cell. In
35 some embodiments, the host cell is a yeast cell. In some embodiments, the host cell is a melanophore cell. Other suitable host cells will be readily apparent to those of ordinary skill in

the art, and a wide variety of cell lines are available from the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209.

In certain embodiments, said determining is carried out using membrane comprising the G protein-coupled receptor.

5 In some embodiments, the test compound is a small molecule. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a polypeptide. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not an antibody or an antigen-binding fragment thereof. In some
10 embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a lipid. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a polypeptide or a lipid. In some embodiments, the test compound is a polypeptide. In some embodiments, the test compound is a polypeptide, with the proviso that the polypeptide is not an antibody or an antigen-binding fragment thereof. In some
15 embodiments, the test compound is a lipid. In some embodiments, the test compound is not an antibody or an antigen-binding fragment thereof. In some embodiments, the test compound is an antibody or an antigen-binding fragment thereof.

 In some embodiments, the method further comprises the step of optionally determining the structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related
20 activity in an individual.

 In some embodiments, the method further comprises the step of optionally providing the name or structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

25 In some embodiments, said method further comprises the step of optionally producing or synthesizing the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

 In some embodiments, the method further comprises the step of optionally determining
30 the structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

 In some embodiments, the method further comprises the step of optionally providing the name or structure of the PYY secretagogue, the compound useful for treatment or prevention of
35 a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

In some embodiments, said method further comprises the step of optionally producing or synthesizing the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

5 In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual as a pharmaceutical.

10 In some embodiments, said method further comprises evaluating the bioavailability of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity.

In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual into a pharmaceutical composition.

15 The invention additionally features a process comprising performing a method according to this *fourth aspect*.

In a *fifth aspect*, the invention features a method of screening test compounds to identify a PYY secretagogue, a compound for treatment or prevention of a condition modulated by PYY, or a compound for a modulation of a PYY-related activity in an individual, which is characterized by using a G protein-coupled receptor comprising an amino acid sequence selected from the group consisting of:

- (a) amino acids 1-335 of SEQ ID NO: 2;
- (b) amino acids 2-335 of SEQ ID NO: 2;
- 25 (c) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (a) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (a);
- (d) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1;
- 30 (e) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (i) the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
 - (ii) the amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
- 35 (f) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and

(g) a biologically active fragment of any one of (a) to (f).

In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2.

In certain embodiments, the receptor comprises the amino acid sequence of SEQ ID
5 NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an allele of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is a mammalian ortholog of SEQ ID NO:
2.

10 In certain embodiments, the G protein-coupled receptor is recombinant.

In certain embodiments, the method comprises identifying an agonist of the receptor.

In certain embodiments, the method comprises identifying a partial agonist of the
receptor.

In some embodiments, said method further comprises the step of formulating the PYY
15 secretagogue, the compound useful for treatment or prevention of a condition modulated by
PYY, or the compound useful for a modulation of a PYY-related activity in an individual as a
pharmaceutical.

In some embodiments, said method further comprises evaluating the bioavailability of
the PYY secretagogue, the compound useful for treatment or prevention of a condition
20 modulated by PYY, or the compound useful for a modulation of a PYY-related activity.

In some embodiments, said method further comprises the step of formulating the PYY
secretagogue, the compound useful for treatment or prevention of a condition modulated by
PYY, or the compound useful for a modulation of a PYY-related activity in an individual into a
pharmaceutical composition.

25 The invention additionally features a process comprising performing a method
according to this *fifth aspect*.

In a *sixth aspect*, the invention features a method comprising, having identified a PYY
secretagogue, a compound for treatment or prevention of a condition modulated by PYY, or a
compound for a modulation of a PYY-related activity in an individual according to the *first*
30 *aspect*, the *second aspect*, the *third aspect*, the *fourth aspect* or the *fifth aspect*, formulating said
PYY secretagogue, said compound for treatment or prevention of a condition modulated by
PYY, or said compound for a modulation of a PYY-related activity in an individual into a
pharmaceutical composition.

In a *seventh aspect*, the invention features use of a G protein-coupled receptor to screen
35 test compounds as PYY secretagogues, compounds for treatment or prevention of a condition
modulated by PYY, or compounds for a modulation of a PYY-related activity in an individual,

wherein the G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:

- (a) amino acids 1-335 of SEQ ID NO: 2;
- (b) amino acids 2-335 of SEQ ID NO: 2;
- 5 (c) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (a) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (a);
- (d) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ
10 ID NO: 1;
- (e) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (i) the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
 - (ii) the amino acid sequence of a G protein-coupled receptor comprising at
15 least 20 contiguous amino acids of SEQ ID NO: 2;
- (f) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and
- (g) a biologically active fragment of any one of any one of (a) to (f).

20 In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2.

In certain embodiments, the receptor comprises the amino acid sequence of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an allele of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2.

25 In certain embodiments, the variant of SEQ ID NO: 2 is a mammalian ortholog of SEQ ID NO: 2.

In certain embodiments, the receptor is recombinant.

In certain embodiments, the test compound is a small molecule.

In certain embodiments, the test compound is a GPR119 agonist.

30 In certain embodiments, the GPR119 agonist is an agonist of an endogenous GPR119.

In certain embodiments, the GPR119 agonist is an agonist of human GPR119.

In certain embodiments, the GPR119 agonist is a GPR119 partial agonist.

In certain embodiments, the GPR119 agonist is a selective GPR119 agonist.

In certain embodiments, the GPR119 agonist is a small molecule.

35 In certain embodiments, the GPR119 agonist is orally available.

In certain embodiments, the GPR119 agonist has an EC₅₀ value of less than about 10 μ M, less than about 1 μ M, less than about 100 nM, less than about 75 nM, less than about 50

nM, less than about 25 nM, less than about 15 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, or less than about 1 nM. In certain embodiments, the GPR119 agonist has an EC₅₀ value of less than about 10 μM, less than about 1 μM, less than about 100 nM, less than about 75 nM, less than about 50 nM, less than about 25 nM, less than about 15 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, or less than about 1 nM at human GPR119 having SEQ ID NO: 2. In certain embodiments, the GPR119 agonist has an EC₅₀ value of less than about 10 μM, less than about 1 μM, less than about 100 nM, less than about 75 nM, less than about 50 nM, less than about 25 nM, less than about 15 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, or less than about 1 nM at human GPR119 having SEQ ID NO: 2 in adenylyl cyclase assay (exemplary adenylyl cyclase assay is provided in Example 9, Example 10 and in Example 11, *infra*). In certain embodiments, the GPR119 agonist has an EC₅₀ value of less than about 10 μM, less than about 1 μM, less than about 100 nM, less than about 75 nM, less than about 50 nM, less than about 25 nM, less than about 15 nM, less than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, or less than about 1 nM at human GPR119 having SEQ ID NO: 2 in melanophore assay (exemplary melanophore assay is provided in Example 12, *infra*).

Exemplary GPR119 agonists are disclosed, e.g., in International Application No. PCT/US2004/001267 (published as WO 04/065380); International Application No. PCT/US2004/005555 (published as WO 04/076413); International Application No. PCT/US2004/022327 (published as WO 05/007647); International Application No. PCT/US2004/022417 (published as WO 05/007658); International Application No. PCT/US2005/019318 (published as WO 2005/121121); International Application No. PCT/GB2004/050046 (published as WO 2005/061489); International Application No. PCT/US06/00567 (published as WO 2006/083491); International Application No. PCT/GB2005/050264 (published as WO 2006/067531); International Application No. PCT/GB2005/050265 (published as WO 2006/067532); International Application No. PCT/GB2005/050266 (published as WO 2006/070208); International Application No. PCT/JP02/09350 (published as WO 03/026661); International Application No. PCT/JP2005/018412 (published as WO 06/040966); International Application No. PCT/JP2005/019000 (published as WO 2006/043490); International Application No. PCT/GB2006/050176 (published as WO 2007/003960); International Application No. PCT/GB2006/050177 (published as WO 2007/003961); International Application No. PCT/GB2006/050178 (published as WO 2007/003962); International Application No. PCT/GB2006/050182 (published as WO 2007/003964); International Application No. PCT/JP02/09350 (published as WO 03/026661); International Application No.

PCT/GB2007/050183 (published as WO 2007/116229); International Application No.
PCT/GB2007/050190 (published as WO 2007/116230); International Application No.
PCT/US2007/073364 (published as WO 2008/008895); International Application No.
PCT/EP2007/058991 (published as WO 2008/025798); International Application No.
5 PCT/EP2007/058993 (published as WO 2008/025799); and International Application No.
PCT/EP2007/058995 (published as WO 2008/025800).

In an *eightth aspect*, the invention features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- 10 (a) contacting a compound which stimulates functionality of a G protein-coupled receptor, wherein said G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:
 - (i) amino acids 1-335 of SEQ ID NO:2;
 - (ii) amino acids 2-335 of SEQ ID NO:2;
 - 15 (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (i);
 - (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO:1;
 - 20 (v) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
 - (b') an amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
 - (vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and
 - (vii) a biologically active fragment of any one of (i) to (vi);
 - 30 in vitro with a vertebrate enteroendocrine cell or with a cell capable of secreting PYY, said compound having been determined or identified by a method according to the *first aspect*; and
- (b) determining whether the compound stimulates PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY;
- 35 wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a

condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2.

5 In certain embodiments, the receptor comprises the amino acid sequence of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an allele of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2.
10 In certain embodiments, the variant of SEQ ID NO: 2 is a mammalian ortholog of SEQ ID NO: 2.

In certain embodiments, the G protein-coupled receptor is recombinant.

In certain embodiments, the vertebrate enteroendocrine cell is a mammalian enteroendocrine cell. In certain embodiments, the enteroendocrine cell is an L cell. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from the
15 small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from an L cell-containing region of small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from duodenum, jejunum, ileum, colon or rectum tissue (*see, e.g.,* Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-
20 261; Ueno et al, Regul Pept (2008) 145:12-16). In certain embodiments, the enteroendocrine cell is an enteroendocrine cell line. In certain embodiments, the cell capable of secreting PYY is a cell comprised by or derived from the islets of Langerhans. In certain embodiments, the cell capable of secreting PYY is a neuronal cell. In certain embodiments, the cell capable of secreting PYY is a recombinant cell engineered to be capable of secreting PYY.

25 The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

(a) administering to a vertebrate a compound which stimulates functionality of a G protein-coupled receptor, wherein said G protein-coupled receptor comprises an amino acid
30 sequence selected from the group consisting of:

- (i) amino acids 1-335 of SEQ ID NO:2;
- (ii) amino acids 2-335 of SEQ ID NO:2;
- (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one
35 or several amino acids in the amino acid sequence defined in (i);

- (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO:1;
 - (v) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
 - (b') an amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
 - (vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and
 - (vii) a biologically active fragment of any one of (i) to (vi);
- said compound having been determined or identified by a method according to the *first aspect*; and
- (b) determining whether the compound increases a PYY level in the vertebrate; wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.
- In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2.
- In certain embodiments, the receptor comprises the amino acid sequence of SEQ ID NO: 2.
- In certain embodiments, the variant of SEQ ID NO: 2 is an allele of SEQ ID NO: 2.
- In certain embodiments, the variant of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2.
- In certain embodiments, the variant of SEQ ID NO: 2 is a mammalian ortholog of SEQ ID NO: 2.
- In certain embodiments, the G protein-coupled receptor is recombinant.
- In certain embodiments, the PYY level is blood or plasma concentration of total PYY.
- In certain embodiments, the PYY level is blood or plasma concentration of PYY₃₋₃₆.
- In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.
- In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In

certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- (a) determining whether the compound increases a PYY level in a sample, said sample having been obtained from a vertebrate previously administered with a compound which stimulates functionality of a G protein-coupled receptor, wherein said G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:
 - (i) amino acids 1-335 of SEQ ID NO:2;
 - (ii) amino acids 2-335 of SEQ ID NO:2;
 - (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (i);
 - (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO:1;
 - (v) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
 - (b') an amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
 - (vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and
 - (vii) a biologically active fragment of any one of (i) to (vi);
 said compound having been determined or identified by a method according to the *first aspect*; and
- wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2.

In certain embodiments, the receptor comprises the amino acid sequence of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an allele of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is a mammalian ortholog of SEQ ID NO: 2.

5 In certain embodiments, the G protein-coupled receptor is recombinant.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

10 In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

15 In some embodiments, a G protein-coupled receptor of the invention is SEQ ID NO:2 or an allele thereof. In some embodiments, a G protein-coupled receptor of the invention is an allele of SEQ ID NO:2. In some embodiments, a G protein-coupled receptor of the invention is an ortholog of SEQ ID NO:2.

20 In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor which specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is a ligand. In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, a G protein-coupled receptor of the invention exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP accumulation. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

30 In certain embodiments, stringent hybridization conditions comprise hybridization at 42°C in a solution comprising 50% formamide, 5xSSC (1xSSC = 150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20µg/ml denatured, sheared salmon sperm DNA, followed by washing at 65°C in a solution comprising 0.1xSSC. Hybridization techniques are well known to the skilled artisan.

35 In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is SEQ ID NO:2 or an allele thereof.

In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an allele of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an ortholog of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is a ligand. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, the variant of SEQ ID NO:2 is a G protein-coupled receptor having at least about 80% identity to SEQ ID NO:2. In certain embodiments, the variant of SEQ ID NO:2 is a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO:2. In certain embodiments, the variant of SEQ ID NO:2 which is a G protein-coupled receptor is an endogenous G protein-coupled receptor. In certain embodiments, the variant of SEQ ID NO:2 which is a G protein-coupled receptor is a non-endogenous G protein-coupled receptor.

In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is SEQ ID NO: 2 or an allele thereof. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 exhibits a detectable level of constitutive

activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

5 In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is SEQ ID NO: 2 or an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which
10 comprises at least 20 contiguous amino acids of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-
15 nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

20 In some embodiments, the G protein-coupled receptor is part of a fusion protein comprising a G protein. Techniques for making a GPCR:G fusion construct are well known to the skilled artisan (*see*, e.g., International Application WO 02/42461).

In some embodiments, the compound which stimulates functionality of a G protein-coupled receptor is a small molecule. In some embodiments, the compound which stimulates
25 functionality of a G protein-coupled receptor is a small molecule, with the proviso that the small molecule is not a polypeptide. In some embodiments, the compound which stimulates functionality of a G protein-coupled receptor is a small molecule, with the proviso that the small molecule is not an antibody or an antigen-binding fragment thereof. In some embodiments, the compound which stimulates functionality of a G protein-coupled receptor is a small molecule,
30 with the proviso that the small molecule is not a lipid. In some embodiments, the compound which stimulates functionality of a G protein-coupled receptor is a small molecule, with the proviso that the small molecule is not a polypeptide or a lipid. In some embodiments, the compound which stimulates functionality of a G protein-coupled receptor is a polypeptide. In some embodiments, the compound which stimulates functionality of a G protein-coupled
35 receptor is a polypeptide, with the proviso that the polypeptide is not an antibody or an antigen-binding fragment thereof. In some embodiments, the compound which stimulates functionality of a G protein-coupled receptor is a lipid. In some embodiments, the compound which

stimulates functionality of a G protein-coupled receptor is not an antibody or an antigen-binding fragment thereof. In some embodiments, the compound which stimulates functionality of a G protein-coupled receptor is an antibody or an antigen-binding fragment thereof.

5 In some embodiments, the method further comprises the step of optionally determining the structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

10 In some embodiments, the method further comprises the step of optionally providing the name or structure of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

15 In some embodiments, said method further comprises the step of optionally producing or synthesizing the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual as a pharmaceutical.

20 In some embodiments, said method further comprises evaluating the bioavailability of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity.

25 In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual into a pharmaceutical composition.

The invention additionally features a process comprising performing a method according to this *eighth aspect*.

30 In a *ninth aspect*, the invention features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

35 (a) contacting a compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound, wherein the G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:

(i) amino acids 1-335 of SEQ ID NO:2;

- (ii) amino acids 2-335 of SEQ ID NO:2;
 - (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (i);
 - 5 (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO:1;
 - (v) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (a') the amino acid sequence of a G protein-coupled receptor having
 - 10 at least about 80% identity to SEQ ID NO: 2; and
 - (b') the amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
 - (vi) the amino acid sequence of a constitutively active version of a G
 - 15 protein-coupled receptor having SEQ ID NO: 2; and
 - (vii) a biologically active fragment of any one of (i) to (vi);
 - in vitro with a vertebrate enteroendocrine cell or with a cell capable of secreting PYY, said compound having been determined or identified by a method according to the *fourth aspect*; and
 - 20 (b) determining whether the compound stimulates PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY;
- wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or from the cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a
- 25 condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the G protein-coupled receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2.

30 In certain embodiments, the receptor comprises the amino acid sequence of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an allele of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2.

In certain embodiments, the variant of SEQ ID NO: 2 is a mammalian ortholog of SEQ ID NO: 2.

35 In certain embodiments, the G protein-coupled receptor is recombinant.

In certain embodiments, the vertebrate enteroendocrine cell is a mammalian enteroendocrine cell. In certain embodiments, the enteroendocrine cell is an L cell. In certain

embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from the small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from tissue obtained from an L cell-containing region of small or large intestine. In certain embodiments, the enteroendocrine cell is comprised by or derived from duodenum, jejunum, ileum, colon or rectum tissue (*see, e.g.,* Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16). In certain embodiments, the enteroendocrine cell is an enteroendocrine cell line. In certain embodiments, the cell capable of secreting PYY is a cell comprised by or derived from the islets of Langerhans. In certain embodiments, the cell capable of secreting PYY is a neuronal cell. In certain embodiments, the cell capable of secreting PYY is a recombinant cell engineered to be capable of secreting PYY.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- 15 (a) administering a compound to a vertebrate, wherein in the presence of said compound less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound, wherein the G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:
 - 20 (i) amino acids 1-335 of SEQ ID NO:2;
 - (ii) amino acids 2-335 of SEQ ID NO:2;
 - (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence in (i);
 - 25 (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO:1;
 - (v) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - 30 (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
 - (b') the amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
 - (vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and
 - 35 (vii) a biologically active fragment of any one of (i) to (vi);

said compound having been determined or identified by a method according to the *fourth aspect*; and

- (b) determining whether the compound increases a PYY level in the vertebrate; wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

The invention additionally features a method for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, comprising the steps of:

- (a) determining whether the compound increases a PYY level in a sample, said sample having been obtained from a vertebrate previous administered with a compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound, wherein the G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:
- (i) amino acids 1-335 of SEQ ID NO:2;
 - (ii) amino acids 2-335 of SEQ ID NO:2;
 - (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (i);
 - (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide hybridizing under stringent conditions stringency to the complement of SEQ ID NO:1;
 - (v) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and

(b') the amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;

(vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and

(vii) a biologically active fragment of any one of (i) to (vi);

said compound having been determined or identified by a method according to the *fourth aspect*; and

wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual.

In certain embodiments, the vertebrate is a mammal. In certain embodiments, the vertebrate is a non-human vertebrate. In certain embodiments, the vertebrate is a non-human mammal. In certain embodiments, the mammal is a non-human mammal.

In certain embodiments, the vertebrate or the mammal is a human. In certain embodiments, the human is a participant in a study reviewed by a governmental agency charged with marketing approval for a drug. In certain embodiments, the study is a clinical trial. In certain embodiments, the governmental agency is the Food and Drug Administration of the United States.

In some embodiments, a G protein-coupled receptor of the invention is SEQ ID NO:2 or an allele thereof. In some embodiments, a G protein-coupled receptor of the invention is an allele of SEQ ID NO:2. In some embodiments, a G protein-coupled receptor of the invention is an ortholog of SEQ ID NO:2.

In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor which specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is a ligand. In some embodiments, a G protein-coupled receptor of the invention is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, a G protein-coupled receptor of the invention exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP accumulation. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, stringent hybridization conditions comprise hybridization at 42°C in a solution comprising 50% formamide, 5xSSC (1xSSC = 150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20µg/ml denatured, sheared salmon sperm DNA, followed by washing at 65°C in a solution comprising 0.1xSSC. Hybridization techniques are well known to the skilled artisan.

In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is SEQ ID NO:2 or an allele thereof. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an allele of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an ortholog of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is a ligand. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, the variant of SEQ ID NO:2 is a G protein-coupled receptor having at least about 80% identity to SEQ ID NO:2. In certain embodiments, the variant of SEQ ID NO:2 is a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO:2. In certain embodiments, the variant of SEQ ID NO:2 which is a G protein-coupled receptor is an endogenous G protein-coupled receptor. In certain embodiments, the variant of SEQ ID NO:2 which is a G protein-coupled receptor is a non-endogenous G protein-coupled receptor.

In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is SEQ ID NO: 2 or an allele thereof. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2. In certain embodiments, the G

protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the the G protein-coupled receptor which has at least about 80% identity to SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is SEQ ID NO: 2 or an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is an allele of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is an ortholog of SEQ ID NO: 2. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist. In some embodiments, the G protein-coupled receptor which comprises at least 20 contiguous amino acids of SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In some embodiments, the G protein-coupled receptor is part of a fusion protein comprising a G protein. Techniques for making a GPCR:G fusion construct are well known to the skilled artisan (*see*, e.g., International Application WO 02/42461).

In some embodiments, the compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound is a small molecule. In some embodiments, the compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound is a small molecule, with the proviso that the small molecule is not a polypeptide. In some embodiments, the compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than

in the absence of the compound is a small molecule, with the proviso that the small molecule is not an antibody or an antigen-binding fragment thereof. In some embodiments, the compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound

5 is a small molecule, with the proviso that the small molecule is not a lipid. In some embodiments, the test compound is a small molecule, with the proviso that the small molecule is not a polypeptide or a lipid. In some embodiments, the test compound is a polypeptide. In some embodiments, the compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the

10 absence of the compound is a polypeptide, with the proviso that the polypeptide is not an antibody or an antigen-binding fragment thereof. In some embodiments, the compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound is a lipid. In some embodiments, the compound in the presence of which less of a complex between a G

15 protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound is not an antibody or an antigen-binding fragment thereof. In some embodiments, the compound in the presence of which less of a complex between a G protein-coupled receptor and an optionally labelled known ligand to the receptor is formed than in the absence of the compound is an antibody or an antigen-binding fragment thereof.

20 In certain embodiments, the method is a method for identifying PYY secretagogues.

In certain embodiments, the method is a method for identifying compounds useful for treatment or prevention of a condition modulated by PYY.

In certain embodiments, the method is a method for identifying compounds useful for a modulation of a PYY-related activity in an individual.

25 In certain embodiments, the ligand for GPR119 is a GPR119 agonist. In certain embodiments, the ligand for GPR119 or the GPR119 agonist is a ligand or agonist of an endogenous vertebrate, mammalian or human GPR119 receptor. In certain embodiments, the ligand for GPR119 or the GPR119 agonist is a ligand or agonist of an endogenous human GPR119 receptor. In certain embodiments, the ligand for GPR119 or the GPR119 agonist is

30 identical to a compound disclosed in, e.g., in International Application No. PCT/US2004/001267 (published as WO 04/065380); International Application No. PCT/US2004/005555 (published as WO 04/076413); International Application No. PCT/US2004/022327 (published as WO 05/007647); International Application No. PCT/US2004/022417 (published as WO 05/007658); International Application No.

35 PCT/US2005/019318 (published as WO 2005/121121); International Application No. PCT/GB2004/050046 (published as WO 2005/061489); International Application No. PCT/US06/00567 (published as WO 2006/083491); International Application No.

- PCT/GB2005/050264 (published as WO 2006/067531); International Application No.
PCT/GB2005/050265 (published as WO 2006/067532); International Application No.
PCT/GB2005/050266 (published as WO 2006/070208); International Application No.
PCT/JP02/09350 (published as WO 03/026661); International Application No.
5 PCT/JP2005/018412 (published as WO 06/040966); International Application No.
PCT/JP2005/019000 (published as WO 2006/043490); International Application No.
PCT/GB2006/050176 (published as WO 2007/003960); International Application No.
PCT/GB2006/050177 (published as WO 2007/003961); International Application No.
PCT/GB2006/050178 (published as WO 2007/003962); International Application No.
10 PCT/GB2006/050182 (published as WO 2007/003964); International Application No.
PCT/JP02/09350 (published as WO 03/026661); International Application No.
PCT/GB2007/050183 (published as WO 2007/116229); International Application No.
PCT/GB2007/050190 (published as WO 2007/116230); International Application No.
PCT/US2007/073364 (published as WO 2008/008895); International Application No.
15 PCT/EP2007/058991 (published as WO 2008/025798); International Application No.
PCT/EP2007/058993 (published as WO 2008/025799); or International Application No.
PCT/EP2007/058995 (published as WO 2008/025800). In certain embodiments, the ligand for
GPR119 or the GPR119 agonist is (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-
[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments,
20 the ligand for GPR119 is an endogenous ligand of an endogenous vertebrate, mammalian, or
human GPR119 receptor.

In certain embodiments, the optionally labeled known ligand is a labeled known ligand.
In certain embodiments, the labeled known ligand is a radiolabeled known ligand. Techniques
for radiolabeling a compound, such as for labeling a known ligand of a G protein-coupled
25 receptor of the invention, are well known to the skilled artisan. *See, e.g.,* International
Application WO 04/065380. Also see, *e.g.,* Example 14, *infra*.

Techniques for detecting the complex between a G protein-coupled receptor and a
compound known to be a ligand of the G protein-coupled receptor are well known to the skilled
artisan. *See, e.g.,* International Application WO 04/065380. Also see, *e.g.,* Example 15, *infra*.

30 In some embodiments, the method further comprises the step of optionally determining
the structure of the PYY secretagogue, the compound useful for treatment or prevention of a
condition modulated by PYY, or the compound useful for a modulation of a PYY-related
activity in an individual.

In some embodiments, the method further comprises the step of optionally providing the
35 name or structure of the PYY secretagogue, the compound useful for treatment or prevention of
a condition modulated by PYY, or the compound useful for a modulation of a PYY-related
activity in an individual.

In some embodiments, said method further comprises the step of optionally producing or synthesizing the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual.

5 In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual as a pharmaceutical.

10 In some embodiments, said method further comprises evaluating the bioavailability of the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity.

In some embodiments, said method further comprises the step of formulating the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual into a
15 pharmaceutical composition.

The invention additionally features a process comprising performing a method according to this *ninth aspect*.

Applicant reserves the right to exclude any one or more candidate compounds from any of the embodiments of the invention. Applicant reserves the right to exclude any one or more
20 modulators from any of the embodiments of the invention. By way of example and not limitation, Applicant reserves the right to exclude any one or more agonists from any of the embodiments of the invention. Applicant reserves the right to exclude any polynucleotide or polypeptide from any of the embodiments of the invention. Applicant additionally reserves the right to exclude any condition modulated by PYY, any bone-related condition, any bone-related
25 condition wherein the bone-related condition is a condition characterized by low bone mass, any bone-related condition wherein the bone-related condition is a bone fracture, any bone-related condition wherein the bone-related condition is a bone disease, any bone-related condition wherein the bone-related condition is enhanced bone healing or bone growth, any metabolic disorder, any angiogenesis-related condition, any ischemia-related condition, any malabsorptive
30 disorder, any convulsive disorder, any cancer, any inflammatory disorder, any inflammatory lung disease, any inflammatory bowel disease, and any modulation of a PYY-related activity in an individual from any of the embodiments of the invention. It is also expressly contemplated that conditions modulated by PYY, bone-related conditions, bone-related conditions wherein the bone-related condition is a condition characterized by low bone mass, bone-related conditions
35 wherein the bone-related condition is a bone fracture, bone-related conditions wherein the bone-related condition is a bone disease, bone-related conditions wherein the bone-related condition is enhanced bone healing or bone growth, metabolic disorders, angiogenesis-related conditions,

ischemia-related conditions, malabsorptive disorders, convulsive disorders, cancers, inflammatory disorders, inflammatory lung diseases, inflammatory bowel diseases, and modulations of a PYY-related activity in an individual of the invention can be included in an embodiment either individually or in any combination.

5

BRIEF DESCRIPTION OF THE DRAWINGS

The invention is illustrated in connection with the figures appended hereto in which:

FIG. 1 shows that GPR119 receptor exhibits constitutive activity for increasing cAMP accumulation in transfected HEK293 cells.

10

FIG. 2 shows that **Compound 1** is an agonist of GPR119 receptor.

FIG. 3 shows an effect of administration of GPR119 agonist on plasma total PYY in wild-type mice.

DETAILED DESCRIPTION OF THE INVENTION

15


The present invention is based, at least in part, on the surprising discovery by Applicant that administration of a GPR119 agonist to an individual, such as by oral administration, can act to increase a plasma total PYY in the individual. The present invention features methods relating to GPR119 for identifying PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, and compounds useful for a modulation of a PYY-related activity in an individual. A GPR119 agonist is useful for increasing plasma total PYY in an individual. In certain embodiments, the individual is a human. GPR119 agonists are useful as pharmaceutical agents for treatment or prevention of a condition modulated by PYY and for a modulation of a PYY-related activity in an individual.

25

The term “ligand”, as used herein, shall mean a molecule (*e.g.*, test compound) that specifically binds to a polypeptide, such as GPR119. A ligand may be, for example, a polypeptide, a lipid, a small molecule, an antibody. **Compound 1** is an exemplary ligand of GPR119 receptor polypeptide (see, Table A, which sets forth the chemical structure and chemical name of Compound 1). Compound 1 is identical to a compound disclosed in International Patent Application No. PCT/US2004/001267 (published as WO 2004/065380). (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine (*see*, Table A) is an exemplary ligand of GPR119 receptor polypeptide. An endogenous ligand is a ligand that is an endogenous, natural ligand for a native polypeptide, such as GPR119. A ligand may be an “antagonist”, “agonist”, “partial agonist”, or “inverse agonist”, or the like.

35

TABLE A



The term “agonist”, as used herein, shall mean an agent (*e.g.*, ligand, test compound) that by virtue of binding to a GPCR activates (*e.g.*, stimulates) the GPCR so as to elicit an intracellular response mediated by the GPCR.

- 5 The term “partial agonist”, as used herein, shall mean an agent (*e.g.*, ligand, test compound) that by virtue of binding to a GPCR activates (*e.g.*, stimulates) the GPCR so as to elicit an intracellular response mediated by the GPCR, albeit to a lesser extent or degree than does a full agonist.

- 10 The term “antagonist” shall mean an agent (*e.g.*, ligand, test compound) that binds, and preferably binds competitively, to a GPCR at about the same site as an agonist or partial agonist but which does not activate an intracellular response initiated by the active form of the GPCR, and can thereby inhibit the intracellular response by agonist or partial agonist. An antagonist typically does not diminish the baseline intracellular response in the absence of an agonist or partial agonist.

- 15 The term “inverse agonist” shall mean an agent (*e.g.*, ligand, test compound) which binds to a GPCR and which inhibits the baseline intracellular response initiated by the active form of the receptor below the normal baseline activity which is observed in the absence of an agonist or partial agonist.

- 20 The term “GPR119 agonist,” as used herein, refers to a compound that binds to GPR119 receptor and acts as an agonist. **Compound 1** is an exemplary GPR119 agonist (see, Table A, which sets forth the chemical structure and chemical name of Compound 1). Compound 1 is identical to a compound disclosed in International Patent Application No. PCT/US2004/001267 (published as WO 2004/065380). (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-

[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an exemplary GPR119 agonist.

The term "selective GPR119 agonist," as used herein, refers to a GPR119 agonist having selectivity for GPR119 receptor over one or more related receptors, such as corticotrophin-releasing factor-1 (CRF-1) receptor. **Compound 1** is an exemplary selective GPR119 agonist (see, Table A, which sets forth the chemical structure and chemical name of Compound 1). Compound 1 is identical to a compound disclosed in International Patent Application No. PCT/US2004/001267 (published as WO 2004/065380). (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an exemplary selective GPR119 agonist.

The term "PYY secretagogue" shall mean an agent (e.g., ligand, test compound) that promotes (e.g., stimulates) secretion of PYY in a cell, e.g. an enteroendocrine cell, or that increases a level of total PYY, e.g. a level of blood or plasma total PYY, on administration to an individual such as a vertebrate or a mammal. In certain embodiments, a PYY secretagogue is a compound suitable for increasing a level of total PYY in an individual, for example a level of blood or plasma total PYY.

As used herein, the term "condition modulated by PYY" refers to a condition modulated by PYY₁₋₃₆ or by a fragment thereof that is capable of modulating a G protein-coupled receptor. In certain embodiments, the term "condition modulated by PYY" refers to a condition modulated by PYY₃₋₃₆. In certain embodiments, the term "condition modulated by PYY" refers to a condition modulated by an NPY Y2 receptor (Y2R) agonist. In certain embodiments, the term "condition modulated by PYY" refers to a condition modulated by stimulation of Y2R. In certain embodiments, the term "condition modulated by PYY₃₋₃₆" refers to a condition modulated by a Y2R agonist. In certain embodiments, the term "condition modulated by PYY₃₋₃₆" refers to a condition modulated by stimulation of Y2R.

As used herein, the term "PYY-related activity" refers to an activity for modulating a level of PYY or to an activity associated with PYY. In certain embodiments, the term "PYY-related activity" refers to an activity for increasing a level of PYY or to an activity associated with PYY.

The term "individual," as used herein, refers to a vertebrate, including but not limited to fish (such as commercially farmed fish, pet fish, etc.), amphibians (such as frogs, toads, pet amphibians, etc.), reptiles (such as snakes, lizards, turtles, pet reptiles, etc.), birds (such as chickens, turkeys, pet birds, etc.) and mammals (such as mice, rats, hamsters, rabbits, pigs, dogs, cats, horses, cows, sheep, goats, non-human primates, non-human mammals, pet non-human mammals, humans, etc.). In certain embodiments, the individual is a fish. In certain embodiments, the individual is an amphibian. In certain embodiments, the individual is a reptile. In certain embodiments, the individual is a bird. In certain embodiments, the individual

is a turkey. Over the past 25 yr, commercial selection pressure for turkeys with larger breast muscle mass has placed increasing demands on skeletal integrity. The increased breast muscle mass, however, has not been accompanied by compensatory changes in the skeleton, with the result that the turkey industry has experienced an increase in leg problems. Long bone fracture in young adult male turkeys has been reported. (See, e.g., Crespo et al, Poultry Sci (2000) 79:602-608.) In certain embodiments, the individual is a mammal. In certain embodiments, the individual is a mouse, a rat, a hamster, a rabbit, a pig, a dog, a cat, a horse, a cow, a sheep, a goat, a non-human primate or a human (which may be included in embodiments of the invention individually or in any combination). In certain embodiments, the individual is a horse. Performance horses, which are horses involved in activities such as racing, pacing and other competitive events, are susceptible to bone fracture. In certain embodiments, the individual is a dog or a cat. In certain embodiments, the individual is a human companion animal (such as a dog, a cat, etc.), a farm animal (such as a cow, a sheep, a goat, a pig, a chicken, etc.), a sports animal (such as a horse, a dog, etc.), a beast of burden (such as a mule, a camel, etc.) or an exotic animal (such as an animal found in a zoo, etc.), which may be included in embodiments of the invention individually or in any combination. In certain embodiments, the individual is a non-human vertebrate. In certain embodiments, the individual is a non-human mammal. In certain embodiments, the individual is a non-human primate (such as a rhesus monkey, a chimpanzee, etc.). In certain embodiments, the individual is a human.

The term "in need of prevention or treatment" as used herein refers to a judgement made by a caregiver (e.g. physician, nurse, nurse practitioner in the case of humans; veterinarian in the case of non-human vertebrates, and in particular embodiment non-human mammals) that an individual requires or will benefit from treatment.

The term "therapeutically effective amount" or "therapeutically effective dose" as used herein refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes one or more of the following:

- (1) Preventing the disease; for example, preventing a disease, condition or disorder in an individual that may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease,
- (2) Inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., arresting further development of the pathology and/or symptomatology), and

(3) Ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology).

5 The term “therapeutic efficacy” as used herein refers to elicitation of the biological or medicinal response in a tissue, system, animal, individual or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes one or more of the following:

(1) Preventing the disease; for example, preventing a disease, condition or disorder in an individual that may be predisposed to the disease, condition or disorder but does not yet
10 experience or display the pathology or symptomatology of the disease,

(2) Inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., arresting further development of the pathology and/or symptomatology), and

15 (3) Ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual that is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology).

20 The term “amount that is effective to prevent” refers to that amount of drug that will prevent or reduce the risk of occurrence of the biological or medical event that is sought to be prevented. In many instances, the amount that is effective to prevent is the same as the therapeutically effective amount.

25 The term “peripheral administration” shall mean administration outside of the central nervous system. Peripheral administration does not include direct administration to the brain. Peripheral administration includes, but is not limited to, intravascular, intramuscular, subcutaneous, inhalation, oral, sublingual, enteral, rectal, transdermal, or intra-nasal administration.

The term “composition” shall mean a material comprising at least one component.

30 The term “active ingredient” shall mean any component that provides pharmacological activity or other direct effect in the diagnosis, cure, mitigation, treatment, or prevention of disease.

The term “pharmaceutical composition” shall mean a composition comprising at least one active ingredient, whereby the composition is amenable to investigation and treatment in a mammal.

35 By “pharmaceutically acceptable” it is meant that the carrier, vehicle, diluent, excipients, and/or salt must be compatible with the other ingredients of the formulation, and not deleterious to the recipient thereof.

The term “dosage form” shall mean the physical form in which a drug is produced and dispensed, such as a tablet, capsule, or an injectable.

By “bone” is intended the dense, semi-rigid, porous, calcified connective tissue forming the major portion of the skeleton of most vertebrates, comprising a dense organic matrix and an inorganic, mineral component. Bone is any of numerous anatomically distinct structures making up the skeleton of a vertebrate.

The terms “bone mass” and “bone mineral density (BMD)” are used interchangeably herein. BMD in humans is usually measured by a standard radiographic technique, dual energy X-ray absorptiometry (DXA). Of the many techniques developed to assess BMD, DXA is the most highly developed technically and the most thoroughly validated biologically. DXA technology, with suitably adapted software, can also be used to reliably assess BMD in animal studies. DXA is used in the diagnosis of osteoporosis, prognosis (fracture prediction), monitoring the natural history of the disorder, and assessing response to treatment.

The term “low bone mass” as used herein refers to any decrease or reduction in bone mineral density (BMD) in an individual, and includes both osteoporosis and osteopenia as defined in proposals by the World Health Organization (WHO). The WHO has defined normal as a value of BMD within one standard deviation of the young adult reference mean (T-score ≥ -1). The WHO has defined osteopenia as a value of BMD more than 1 standard deviation below the young adult mean, but less than 2.5 standard deviations below this value (T-score < -1 and > -2.5). The WHO has characterized osteoporosis as a more severe form of osteopenia, and has defined it by value of BMD 2.5 standard deviations or more below the young adult mean (T-score ≤ -2.5). (See, e.g., World Health Organization Technical Report Series 921 (2003), Prevention and Management of Osteoporosis, the disclosure of which is herein incorporated by reference in its entirety.) More commonly, osteopenia is defined as a T-score of less than -1 and greater than -2, and osteoporosis is defined as a T-score of less than or equal to -2. In certain embodiments of the present invention, the T-score is measured at the hip with DXA.

The term “osteoporosis” as used herein is defined by a value of BMD 2 standard deviations or more below the young adult reference mean (T-score ≤ -2) or refers to a diagnosis made by a caregiver (e.g. physician, nurse, nurse practitioner in the case of humans; veterinarian in the case of non-human vertebrates).

Osteoporosis can be classified as either primary or secondary. (See, e.g., World Health Organization Technical Report Series 921 (2003), Prevention and Management of Osteoporosis.) As used herein, the term “osteoporosis” encompasses primary osteoporosis and secondary osteoporosis. In certain embodiments, osteoporosis is primary osteoporosis. In certain embodiments, osteoporosis is secondary osteoporosis.

“Primary osteoporosis” as used herein is associated with menopause (natural, premature, or surgical), aging, or both. It shall be understood that in the present invention,

primary osteoporosis associated with menopause (natural, premature, or surgical), primary osteoporosis associated with aging, and primary osteoporosis associated with menopause and aging can be included in embodiments individually or in any combination.

5 “Secondary osteoporosis” as used herein refers to osteoporosis which is associated not with menopause or aging but rather with medical conditions or with the use of medications or drugs. An increased risk of osteoporosis is associated with a host of medical conditions, including but not limited to endocrine and metabolic disorders, and malignant disease, and with the use of certain medications and drugs, examples of which are well known to those skilled in the art (see, *e.g.*, World Health Organization Technical Report Series 921 (2003), Prevention and Management of Osteoporosis; Williams Textbook of Endocrinology, 10th Edition; the disclosure of which is herein incorporated by reference in its entirety.) Secondary osteoporosis can also be associated with immobilization. A diagnosis of osteoporosis secondary to a medical condition, to use of a medication or drug, or to immobilization can be made by a caregiver (*e.g.* physician, nurse, nurse practitioner in the case of humans; veterinarian in the case of non-human vertebrates).

By “bone fracture” is intended a complete or incomplete break, rupture or crack of a bone. Diagnosis of fractures normally depends upon clinical examination and radiological findings. In the invention, bone fractures include, but are not limited to, traumatic fractures, long-term fractures, and pathological fractures.

20 “Traumatic fracture” as used herein shall refer to an immediate fracture which involves a supraliminal trauma with a degree of local violence that exceeds the natural elasticity of the bone. It can be accompanied by simultaneous injury to the soft tissues and very often the skin. A traumatic fracture can be closed (the adjacent soft tissue can be injured but the covering soft parts are largely preserved). A traumatic fracture can be open (the broken ends of the bone are freed by extensive soft tissue injury so that pathogens from outside can enter the wound directly).

“Long-term fracture” as used herein shall refer to a chronic fracture, fatigue fracture, stress fracture or spontaneous fracture type I.

30 “Pathological fracture” as used herein shall refer to a spontaneous fracture type II. A pathological fracture arises spontaneously, without adequate trauma to account for it. The bone may have been previously damaged, either by a systemic disease (*e.g.*, osteoporosis, osteodystrophy, or Paget’s osteitis deformans) or by a local bone lesion (*e.g.*, metastasis, radioosteonecrosis, or bone tumor). See, Adler, Claus-Peter, BONE DISEASES, p. 114 (Springer-Verlag, Germany 2000).

35 Fractures also include, but are not limited no, oblique torsion fracture, transverse fracture, comminuted fracture, compression fracture, rib fractures, creeping fracture, and

fractured femoral neck (Adler, Claus-Peter, BONE DISEASES, Springer-Verlag, Germany (2000)).

As used herein, "bone-related condition" refers to a condition that derives benefit from treatment according to the invention, by increasing bone mass or bone growth or by decreasing
5 loss of bone mass, and includes but is not limited to a condition characterized by low bone mass, a bone fracture, a bone disease, and enhancing bone healing or bone growth. It shall be understood that in the present invention, bone-related conditions that can be treated according to the invention can be included in embodiments individually or in any combination.

As used herein, the term "condition characterized by low bone mass" includes but is not
10 limited to osteopenia, osteoporosis, rheumatoid arthritis, osteoarthritis, periodontal disease, alveolar bone loss, osteotomy bone loss, childhood idiopathic bone loss, curvature of the spine and loss of height. In certain embodiments, osteoporosis is primary osteoporosis. In certain embodiments, osteoporosis is secondary osteoporosis. In certain embodiments, secondary
15 osteoporosis is associated with a medical condition. In certain embodiments, secondary osteoporosis is associated with the use of a medication or drug. In certain embodiments, secondary osteoporosis is associated with immobilization. Conditions characterized by low bone mass also include but are not limited to Paget's disease, bone loss due to metastatic cancer, and osteolytic lesions such as those caused by neoplastic disease, radiotherapy, or chemotherapy. Conditions characterized by low bone mass also include but are not limited to
20 long-term complications of osteoporosis such as curvature of the spine, loss of height and prosthetic surgery. It shall be understood that in the present invention, conditions characterized by low bone mass can be included in embodiments individually or in any combination. (See, e.g., World Health Organization Technical Report Series 921 (2003), Prevention and Management of Osteoporosis; Williams Textbook of Endocrinology, 10th Edition, Larsen et al, Eds. (2002), W.B. Saunders Company; and Endocrinology and Metabolism, 4th Edition, Felig et al, Eds. (2001), McGraw-Hill Book Company; the disclosure of each of which is herein
25 incorporated by reference in its entirety.)

As used herein, "bone disease" refers to a disorder or condition relating to abnormality of the bone. Bone diseases that can be treated according to the invention, by increasing bone
30 mass or bone growth or decreasing loss of bone mass, include but are not limited to osteopenia, osteoporosis, rheumatoid arthritis, osteoarthritis, periodontal disease, alveolar bone loss, osteotomy bone loss, childhood idiopathic bone loss, curvature of the spine, and loss of height. In certain embodiments, osteoporosis is primary osteoporosis. In certain embodiments, osteoporosis is secondary osteoporosis. In certain embodiments, secondary osteoporosis is
35 associated with a medical conditions. In certain embodiments, secondary osteoporosis is associated with the use of a medication or drug. In certain embodiments, secondary osteoporosis is associated with immobilization. Bone diseases that can be treated according to

the invention, by increasing bone mass or bone growth, also include but are not limited to Paget's disease and bone loss due to metastatic cancer. Destructive bone disorders that can be treated according to the invention, by increasing bone mass or growth, include but are not limited to osteoporosis, osteoarthritis, and osteolytic lesions such as those caused by neoplastic disease, radiotherapy, or chemotherapy. It shall be understood that in the present invention, bone diseases that can be treated according to the invention, by increasing bone mass or growth, can be included in embodiments individually or in any combination. (See, e.g., World Health Organization Technical Report Series 921 (2003), Prevention and Management of Osteoporosis; Williams Textbook of Endocrinology, 10th Edition, Larsen et al, Eds. (2002), W.B. Saunders Company; and Endocrinology and Metabolism, 4th Edition, Felig et al, Eds. (2001), McGraw-Hill Book Company; the disclosure of each of which is herein incorporated by reference in its entirety.)

The present invention also relates to the other conditions that derive benefit from treatment according to the invention, by increasing bone mass or bone growth or decreasing loss of bone mass, including but not limited to enhanced bone healing following facial reconstruction, maxillary reconstruction, mandibular reconstruction, periodontal disease or tooth extraction, enhanced long bone extension, enhanced prosthetic ingrowth and increased bone synostosis.

As used herein, "metabolic disorder" refers to a disorder relating to abnormality of metabolism. Metabolic disorders that can be treated according to the invention include but are not limited to overweight, obesity, hyperphagia, diabetes (inclusive of type 1 diabetes and type 2 diabetes), type 2 diabetes, impaired glucose tolerance, insulin resistance, hyperinsulinemia, dyslipidemia, hypertension and metabolic syndrome. It shall be understood that in the present invention, metabolic disorders that can be treated according to the invention can be included in embodiments individually or in any combination.

The term "metabolic syndrome" as defined herein, and according to the Adult Treatment Panel III (ATP III; National Institutes of Health: Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III), Executive Summary; Bethesda, Md., National Institutes of Health, National Heart, Lung and Blood Institute, 2001 (NIH pub. No 01-3670), occurs when a person meets three or more of five criteria related to obesity, hypertriglyceridemia, low HDL cholesterol, high blood pressure, and high fasting glucose.

The term "obesity," as used herein, is defined as a body mass index (BMI) of 30.0 or greater, in accordance with the WHO classifications of weight [Kopelman, Nature (2000) 404:635-643; the disclosure of which is herein incorporated by reference in its entirety].

"Impaired glucose tolerance" (IGT) as used herein is intended to indicate that condition associated with insulin-resistance that is intermediate between frank, type 2 diabetes and normal

glucose tolerance (NGT). IGT is diagnosed by a procedure wherein an affected person's postprandial glucose response is determined to be abnormal as assessed by 2-hour postprandial plasma glucose levels. In this test, a measured amount of glucose is given to the patient and blood glucose levels are measured at regular intervals, usually every half hour for the first two hours and
5 every hour thereafter. In a "normal" or non-IGT individual, glucose levels rise during the first two hours to a level less than 140 mg/dl and then drop rapidly. In an IGT individual, the blood glucose levels are higher and the drop-off level is at a slower rate.

"Insulin resistance" as used herein is intended to encompass the usual diagnosis of insulin resistance made by any of a number of methods, including but not restricted to: the intravenous
10 glucose tolerance test or measurement of the fasting insulin level. It is well known that there is an excellent correlation between the height of the fasting insulin level and the degree of insulin resistance. Therefore, one could use elevated fasting insulin levels as a surrogate marker for insulin resistance for the purpose of identifying which normal glucose tolerance (NGT) individuals have insulin resistance. A diagnosis of insulin resistance can also be made using the euglycemic glucose
15 clamp test.

"Dyslipidemia" is intended herein to encompass disorders comprising any one of elevated level of plasma free fatty acids, elevated level of plasma cholesterol, elevated level of LDL-cholesterol, reduced level of HDL-cholesterol, elevated ratio of total cholesterol to HDL cholesterol, and elevated level of plasma triglycerides.

20 As used herein, "angiogenesis-related condition" refers to a condition that derives benefit from increased angiogenesis. Angiogenesis-related conditions that can be treated according to the invention, by increasing angiogenesis, include but are not limited to peripheral arterial disease, wound healing, and ischemia. It shall be understood that in the present invention, angiogenesis-related conditions that can be treated according to the invention can be
25 included in embodiments individually or in any combination.

As used herein, "ischemia" refers to a state of tissue oxygen deprivation or insufficiency.

As used herein, "ischemia-related condition" refers to a condition which relates to ischemia. Ischemia-related conditions that can be treated according to the invention include but
30 are not limited to cerebral ischemia, stroke, ischemic heart disease, myocardial infarction, ischemia-reperfusion injury, post-myocardial infarction remodeling, heart failure, and congestive heart failure. It shall be understood that in the present invention, ischemia-related conditions that can be treated according to the invention can be included in embodiments individually or in any combination.

35 "Stroke" is a cardiovascular disease that affects the blood vessels supplying blood to the brain and is intended herein to include cerebral thrombosis, the most common type of stroke. Cerebral thrombosis occurs when a blood clot (thrombus) forms and blocks blood flow in an

artery bring blood to part of the brain. Blood clots usually form in arteries damaged by atherosclerosis.

As used herein, "ischemic heart disease" refers to a disorder caused by lack of oxygen to the tissues of the heart, in which muscles of the heart are affected and the heart cannot pump properly. Ischemic heart disease is the most common cardiomyopathy in the United States.

As used here, "myocardial infarction" refers to the damage or death of an area of heart muscle because of an inadequate supply of oxygen to that area. Myocardial infarctions are often caused by a blood clot that blocks one of the coronary arteries (the blood vessels that bring oxygen to heart muscle). The clot prevents blood and oxygen from reaching that area of the heart, leading to the death of heart cells in that area.

The loss of myocardial tissue due to myocardial infarction results in a sustained excessive hemodynamic burden placed on the ventricle. Ventricular hypertrophy constitutes one of the principle mechanisms by which the heart compensates for an increased load. However, the capacity for this adaptation to sustain cardiac performance in the face of hemodynamic overload is finite and, when chronically maintained, becomes maladaptive. Gradually, the adaptive hypertrophic phenotype transitions to overt heart failure as the enlarged ventricles progressively dilate and contractile function weakens. The natural history of the adaptive and maladaptive response to myocardial infarction in the heart is referred to as "post-myocardial infarction remodeling".

"Congestive heart failure" refers to a disorder in which the heart loses its ability to pump blood efficiently. Congestive heart failure becomes more prevalent with advancing age. Ischemic heart disease is the most common cause of congestive heart failure, accounting for 60-70% of all cases. An increased venous pressure greater than 12 mmHg is one of the major Framingham criteria for congestive heart failure, as is a reduction in cardiac output equivalent to a circulation time great than 25 seconds.

As used herein, "malabsorptive disorder" refers to a disorder relating to abnormality of secretion and/or absorption in the intestinal epithelium. Malabsorptive disorders that can be treated according to the invention include but are not limited to short bowel syndrome, enterocolitis, Crohn's disease, ulcerative colitis, irritable bowel syndrome, cholera and a disorder accompanied by diarrhea. As used herein, the "disorder accompanied by diarrhea" is not particularly limited as long as it is a disease or a symptom accompanied by diarrhea, and it does not matter whether it is an acute disease or a chronic disease. Examples of such a disease or symptom include diarrhea (caused by a parasite, a bacterium, a virus, physical stimulation, dyspepsia, mental disorder, gastrointestinal inflammation or allergy, etc.), irritable bowel syndrome, ulcerative colitis and the like. It is expressly contemplated that each disorder accompanied by diarrhea is a separate embodiment within the scope of the present invention. It

shall be understood that in the present invention, malabsorptive disorders that can be treated according to the invention can be included in embodiments individually or in any combination.

As used herein, "convulsive disorder" refers to a disorder of an individual in which the subject suffers from convulsion, e.g., convulsions due to epileptic seizure. Convulsive disorders that can be treated according to the invention include epilepsy and non-epileptic seizures, e.g., convulsions due to administration of a convulsant agent to the individual. It shall be understood that in the present invention, convulsive disorders that can be treated according to the invention can be included in embodiments individually or in any combination.

As used herein, "cancer" refers to a malignant neoplasm. Cancers that can be treated according to the invention include but are not limited to pancreatic cancer, breast cancer, colon cancer, and Barrett's adenocarcinoma. It shall be understood that in the present invention, cancers that can be treated according to the invention can be included in embodiments individually or in any combination.

As used herein, "inflammatory disorder" refers to a disorder relating to abnormality of inflammation. Inflammatory disorders that can be treated according to the invention include but are not limited to atherosclerosis, atherothrombosis, insulin resistance, pancreatitis, restenosis, an inflammatory lung disease, and an inflammatory bowel disease. It shall be understood that in the present invention, inflammatory disorders that can be treated according to the invention can be included in embodiments individually or in any combination.

As used herein, "atherothrombosis" refers to thrombus formation in an atherosclerotic vessel.

As used herein, "inflammatory lung disease" refers to a condition relating to abnormality of inflammation in the lung. Inflammatory lung diseases that can be treated according to the invention include but are not limited to asthma, emphysema and pulmonary arterial hypertension (PAH). PAH is a life-threatening disease characterized by a progressive pulmonary vasculopathy leading to right ventricular hypertrophy; right heart failure occurs if left untreated. PAH shall be understood to encompass the following forms of pulmonary arterial hypertension described in the 2003 World Health Organization (WHO) clinical classification of pulmonary arterial hypertension: idiopathic PAH, familial PAH, PAH associated with other condition, and PAH associated with significant venous or capillary involvement. It shall be understood that in the present invention, inflammatory lung diseases that can be treated according to the invention can be included in embodiments individually or in any combination.

As used herein, the "inflammatory bowel disease" refers to a condition relating to abnormality of inflammation in the bowel. Inflammatory bowel diseases that can be treated according to the invention include but are not limited to Crohn's disease as may be enteritis, enterocolitis, Crohn's disease, granulomatous colitis, ileitis or Crohn's colitis and ulcerative colitis as may be ulcerative colitis, sigmoiditis, pancolitis and ulcerative proctitis. It shall be

understood that in the present invention, inflammatory bowel diseases that can be treated according to the invention can be included in embodiments individually or in any combination.

5 The term “atherosclerosis” as used herein refers to a form of vascular disease characterized by the deposition of atheromatous plaques containing cholesterol and lipids on the innermost layer of the walls of large and medium-sized arteries.

“Alzheimer’s disease” is a progressive neurodegenerative disease characterized by significant loss of function in more than one cognitive domain and often accompanied by changes in behavior or personality. Alzheimer’s disease is the most common cause of dementia. The beta-amyloid peptide ($A\beta$) is believed to play a central role in the pathology of Alzheimer’s disease.

10

“Mild cognitive impairment” (MCI) is a state of cognitive impairment, most often of memory only, that is not severe enough to significantly impact occupational or social functioning. However, many people with MCI progress to Alzheimer’s disease, at a rate of 10-15% per year and up to 60% within five years. Growing evidence shows that Alzheimer’s disease pathology begins much earlier than the development of symptoms severe enough to allow a diagnosis of Alzheimer’s criteria. In 2001, the American Academy of Neurology (AAN) published clinical practice parameters for the detection of MCI that will increase awareness, diagnosis, and treatment of both MCI and mild forms of Alzheimer’s disease.

15

As used herein, “prion-associated disease” is intended to include but not be limited to bovine spongiform encephalopathy, Creutzfeldt-Jakob disease, kuru, Gerstmann-Straussler-Scheinker syndrome, and fatal familial insomnia (Collins et al, Lancet (2004) 363:51-61).

20

As used herein, “excitotoxic injury” refers to neuronal cell damage or neuronal cell death caused resulting from substance-induced paroxysmal overactivity, e.g. glutamate-induced paroxysmal activity.

25 “Parkinson’s disease” is a chronic, progressive neurodegenerative disorder characterized by motor symptoms such as tremor, bradykinesia, muscle rigidity, gait dysfunction, and postural instability. Researchers have identified degeneration of dopaminergic neurons in the substantia nigra as the primary pathophysiological mechanism of Parkinson’s disease.

“Adiponectin” is a plasma protein secreted by adipocytes and comprised of an N-terminally disposed collagen-like region and a C-terminal globular region. Reduced levels of adiponectin have been associated with a number of metabolic-related disorders, including atherosclerosis, coronary heart disease, stroke, insulin resistance and type 2 diabetes. The serum adiponectin level for women has been reported to be higher than that for men, for example 13.5 $\mu\text{g/ml}$ versus 7.2 $\mu\text{g/ml}$ in one study (Yamamoto et al, Clin Sci (London) (2002) 103:137-142).

30

35 The term “endogenous” shall mean a material that an individual (for example, and not limitation, a human) naturally produces. By contrast, “non-endogenous” shall mean that which is not naturally produced by an individual (for example, and not limitation, a human).

The term "biologically active fragment" of a G protein-coupled receptor shall mean a fragment of the GPCR having structural and biochemical functions of a naturally occurring GPCR. In certain embodiments, the biologically active fragment couples to a G protein. In certain embodiments, the biologically active fragment binds to a known ligand of the GPCR.

5 The term "primer" is used herein to denote a specific nucleotide sequence which is complementary to a target nucleotide sequence and used to hybridize to the target nucleotide sequence. A primer serves as an initiation point for nucleotide polymerization catalyzed by DNA polymerase, RNA polymerase, or reverse transcriptase.

10 The term "expression vector" shall mean a DNA sequence that is required for the transcription of cloned DNA and translation of the transcribed mRNA in an appropriate host cell recombinant for the expression vector. An appropriately constructed expression vector should contain an origin of replication for autonomous replication in host cells, selectable markers, a limited number of useful restriction enzyme sites, a potential for high copy number, and active promoters. The cloned DNA to be transcribed is operably linked to a constitutively or
15 conditionally active promoter within the expression vector.

 The term "host cell" shall mean a cell capable of having a vector incorporated therein. In the present context, the vector will typically contain nucleic acid encoding a GPCR or GPCR fusion protein in operable connection with a suitable promoter sequence to permit expression of the GPCR or GPCR fusion protein to occur. In particular embodiment, the host cell is a
20 eukaryotic host cell. In certain embodiments, the eukaryotic host cell is a mammalian host cell. In certain embodiments, the eukaryotic host cell is a yeast host cell. In certain embodiments, the eukaryotic host cell is a melanophore host cell.

 The term "contact" or "contacting" shall mean bringing at least two moieties together.

25 The terms "modulate" or "modify" shall be taken to refer to an increase or decrease in the amount, quality, or effect of a particular activity, function or molecule or, when in reference to a condition, shall be taken to refer to a change in the status of said condition. By way of illustration and not limitation, agonists, partial agonists, inverse agonists, and antagonists of a G protein-coupled receptor are modulators of the receptor.

30 The term "small molecule" shall be taken to mean a compound having a molecular weight of less than about 10,000 grams per mole, including a peptide, peptidomimetic, amino acid, amino acid analogue, polynucleotide, polynucleotide analogue, nucleotide, nucleotide analogue, organic compound or inorganic compound (i.e. including a heterorganic compound or organometallic compound), and salts, esters and other pharmaceutically acceptable forms thereof. In certain embodiments, small molecules are organic or inorganic compounds having a
35 molecular weight of less than about 5,000 grams per mole. In certain embodiments, small molecules are organic or inorganic compounds having molecular weight of less than about 1,000 grams per mole. In certain embodiments, small molecules are organic or inorganic compounds

having a molecular weight of less than about 800 grams per mole. In certain embodiments, small molecules are organic or inorganic compounds having a molecular weight of less than about 600 grams per mole. In certain embodiments, small molecules are organic or inorganic compounds having a molecular weight of less than about 500 grams per mole.

5 Amino acid abbreviations used herein are set out in Table B:

TABLE B

ALANINE	ALA	A
ARGININE	ARG	R
ASPARAGINE	ASN	N
ASPARTIC ACID	ASP	D
CYSTEINE	CYS	C
GLUTAMIC ACID	GLU	E
GLUTAMINE	GLN	Q
GLYCINE	GLY	G
HISTIDINE	HIS	H
ISOLEUCINE	ILE	I
LEUCINE	LEU	L
LYSINE	LYS	K
METHIONINE	MET	M
PHENYLALANINE	PHE	F
PROLINE	PRO	P
SERINE	SER	S
THREONINE	THR	T
TRYPTOPHAN	TRP	W
TYROSINE	TYR	Y
VALINE	VAL	V

The term “polypeptide” shall refer to a polymer of amino acids without regard to the length of the polymer. Thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not specify or exclude post-expression
 10 modifications of polypeptides. For example, polypeptides that include the covalent attachment

of glycosyl groups, acetyl groups, phosphate groups, lipid groups and the like are expressly encompassed by the term polypeptide.

5 The term "polynucleotide" shall refer to RNA, DNA, or RNA/DNA hybrid sequence of more than one nucleotide in either single chain or duplex form. The polynucleotides of the invention may be prepared by any known method, including synthetic, recombinant, ex vivo generation, or a combination thereof, as well as utilizing any purification methods known in the art.

The term "antibody" is intended herein to encompass monoclonal antibody and polyclonal antibody.

10 The term "second messenger" shall mean an intracellular response produced as a result of receptor activation. A second messenger can include, for example, inositol 1,4,5-triphosphate (IP₃), diacylglycerol (DAG), cyclic AMP (cAMP), cyclic GMP (cGMP), MAP kinase activity, MAPK/ERK kinase kinase-1 (MEKK1) activity, and Ca²⁺. Second messenger response can be measured for a determination of receptor activation.

15 The term "receptor functionality" shall refer to the normal operation of a receptor to receive a stimulus and moderate an effect in the cell, including, but not limited to regulating gene transcription, regulating the influx or efflux of ions, effecting a catalytic reaction, and/or modulating activity through G-proteins, such as eliciting a second messenger response.

20 The term "stimulate" or "stimulating," in relationship to the term "response" or "functionality of the receptor" shall mean that a response or a functionality of the receptor is increased in the presence of a compound as opposed to in the absence of the compound.

25 The term "inhibit" or "inhibiting," in relationship to the term "response" or "functionality of the receptor" shall mean that a response a functionality of the receptor is decreased or prevented in the presence of a compound as opposed to in the absence of the compound.

The term "compound efficacy" shall mean a measurement of the ability of a compound to inhibit or stimulate receptor functionality, as opposed to receptor binding affinity.

30 The term "test compound," used interchangeably herein with "candidate compound," shall mean a molecule (for example, and not limitation, a chemical compound) which is amenable to a screening technique.

The term "constitutively active" in relationship to a G protein-coupled receptor shall mean that the G protein-coupled receptor exhibits agonist-independent activity.

35 The term "directly identifying" or "directly identified", in relationship to the phrase "test compound," shall mean the screening of a compound against a G protein-coupled receptor in the absence of a known ligand (e.g., a known agonist) to the G protein-coupled receptor.

Where a range of values is provided, it is understood that each intervening value, to the tenth of the lower limit unless the context clearly indicates otherwise, between the upper and

lower limit of that range and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes
 5 one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

A. Introduction

The order of the following sections is set forth for presentational efficiency and is not intended, nor should be construed, as a limitation on the disclosure or the claims to follow.

10 B. Receptor Expression

1. GPCR polypeptides of interest

A GPCR of the invention may comprise an amino acid sequence selected from the group consisting of:

- (a) amino acids 1-335 of SEQ ID NO: 2;
- 15 (b) amino acids 2-335 of SEQ ID NO: 2;
- (c) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (a) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (a);
- (d) the amino acid sequence of a G protein-coupled receptor encoded by a
 20 polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1;
- (e) a variant of SEQ ID NO: 2 when selected from the group consisting of:
 - (i) the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO: 2; and
 - 25 (ii) the amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO: 2;
- (f) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2; and
- (g) a biologically active fragment of any one of any one of (a) to (f).

30

In certain embodiments, a GPCR of the invention comprises the amino acid sequence of SEQ ID NO: 2. In some embodiments, a GPCR of the invention is an allele of SEQ ID NO:2. In some embodiments, a GPCR of the invention is an ortholog of SEQ ID NO:2.

In some embodiments, a GPCR of the invention is recombinant. In some embodiments,
 35 the recombinant GPCR is recombinant human GPR119.

In certain embodiments, a GPCR that may be used in the subject methods exhibits a detectable level of constitutive activity.

In some embodiments, a GPCR of the invention is endogenous. In some embodiments, a GPCR of the invention is a mammalian GPR119. In some embodiments, a GPCR of the invention that is endogenous is a mammalian GPR119.

By way of illustration and not limitation, deletion of an N-terminal methionine residue
5 is envisioned to provide a biologically active fragment that may be used in the subject invention. In certain embodiments, a biologically active fragment of the invention is a fragment optionally fused at its N-terminus to a peptide comprising an N-terminal methionine residue and an HA epitope tag (from hemagglutinin influenza virus) that specifically binds Compound 1. In certain
10 embodiments, a biologically active fragment of the invention is a fragment optionally fused at its N-terminus to a peptide comprising an N-terminal methionine residue and an HA epitope tag (from hemagglutinin influenza virus) that specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, a biologically active fragment of the invention is a fragment
15 optionally fused at its N-terminus to a peptide comprising an N-terminal methionine residue and an HA epitope tag that specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine with an IC₅₀ value in receptor binding assay according to Example 15, *infra*, of less than about 50μM, less than about 25μM, less than about 10μM, less than about 5μM, less than about 1μM, less than about 500nM, less than about 100nM, or less than about 50nM. In certain embodiments, a
20 biologically active fragment of the invention is a fragment optionally fused at its N-terminus to a peptide comprising an N-terminal methionine residue and an HA epitope tag that specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine with an IC₅₀ value in receptor binding assay according to Example 15, *infra*, of less than about 10μM, less than about 5μM, less than about 1μM, less than
25 about 500nM, less than about 100nM, or less than about 50nM. In certain embodiments, a biologically active fragment of the invention is a fragment optionally fused at its N-terminus to a peptide comprising an N-terminal methionine residue and an HA epitope tag for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist having an EC₅₀ value at said fragment optionally
30 fused at its N-terminus to said peptide in whole cell adenylyl cyclase assay according to Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than about 5μM, less than about 1μM, less than about 100nM, less than about 50nM, less than about 25nM, less than about 10nM, less than about 5nM, or less than about 1nM. In certain embodiments, a biologically active fragment of the invention is a fragment optionally fused at its N-terminus to a
35 peptide comprising an N-terminal methionine residue and an HA epitope tag for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist having an EC₅₀ value at said fragment optionally

fused at its N-terminus to said peptide in whole cell adenylyl cyclase assay according to Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than about 100nM, less than about 50nM, less than about 25nM, less than about 10nM, less than about 5nM, or less than about 1nM. In some embodiments, a biologically active fragment of the invention is a fragment
5 optionally fused at its N-terminus to a peptide comprising an N-terminal methionine residue and an HA epitope tag that exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion. In certain embodiments, the fragment is fused at its N-terminus to a peptide consisting essentially
10 of an N-terminal methionine residue and an HA epitope tag. Techniques for fusing a peptide comprising or consisting essentially of an N-terminal methionine residue and an HA epitope tag to the N-terminus of a polypeptide fragment are well known in the art and can be obtained commercially (e.g., Clontech, Mountain View, CA).

An allelic variant of human GPR119 of SEQ ID NO: 2 is envisioned to be within the
15 scope of the invention. Human GPR119 is envisioned to be within the scope of the invention.

A variant which is a vertebrate ortholog of human GPR119 of SEQ ID NO: 2 is envisioned to be within the scope of the invention. A variant which is a mammalian ortholog of human GPR119 of SEQ ID NO: 2 is envisioned to be within the scope of the invention. By way of illustration and not limitation, mouse GPR119 (*e.g.*, GenBank[®] Accession No. AY288423),
20 rat GPR119 (GenBank[®] Accession No. AAN95195), hamster GPR119, dog GPR119, and non-human primate GPR119 are envisioned to be within the scope of the invention.

In certain embodiments, the variant of SEQ ID NO: 2 is a GPCR.

A variant of SEQ ID NO: 2 having at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least
25 about 99% identity to SEQ ID NO: 2 is envisioned to be within the scope of the invention. In certain embodiments, the variant of SEQ ID NO: 2 having at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 2 is a GPCR. In some embodiments, the variant of SEQ ID NO: 2 is an endogenous GPCR. In some embodiments, the variant of
30 SEQ ID NO: 2 is a non-endogenous GPCR. In some embodiments, the variant of SEQ ID NO: 2 that is an endogenous GPCR is a mammalian GPCR. In certain embodiments, the variant of SEQ ID NO: 2 specifically binds Compound 1. In certain embodiments, the variant of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-
[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments,
35 the variant of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine with an IC₅₀ value in receptor binding assay according to Example 15, *infra*, of less than about 50μM, less

than about 25 μ M, less than about 10 μ M, less than about 5 μ M, less than about 1 μ M, less than about 500nM, less than about 100nM, or less than about 50nM. In certain embodiments, the variant of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine with an IC₅₀ value in receptor binding assay according to Example 15, *infra*, of less than about 10 μ M, less than about 5 μ M, less than about 1 μ M, less than about 500nM, less than about 100nM, or less than about 50nM. In certain embodiments, the variant of SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist having an EC₅₀ value at said receptor in whole cell adenylyl cyclase assay according to Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than about 5 μ M, less than about 1 μ M, less than about 100nM, less than about 50nM, less than about 25nM, less than about 10nM, less than about 5nM, or less than about 1nM. In certain embodiments, the variant of SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist having an EC₅₀ value at said receptor in whole cell adenylyl cyclase assay according to Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than about 100nM, less than about 50nM, less than about 25nM, less than about 10nM, less than about 5nM, or less than about 1nM. In some embodiments, the variant of SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion. Percent identity can be determined conventionally using known computer programs.

In certain embodiments, a variant GPCR that may be used in the subject methods has an amino acid sequence having at least about 80%, at least about 85%, at least about 90%, at least about 95%, of at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity to SEQ ID NO: 2. By a variant GPCR having, for example, 95% "identity" to SEQ ID NO: 2 is meant that the amino acid sequence of the variant is identical to amino acids 1-335 of SEQ ID NO: 2 except that it may include up to five amino acid alterations per each 100 amino acids of SEQ ID NO: 2. Thus, to obtain for example an amino acid sequence having at least 95% identity to the amino acid sequence of SEQ ID NO: 2, up to 5% (5 of 100) of the amino acid residues in the sequence may be inserted, deleted, or substituted with another amino acid compared with amino acids 1-335 of SEQ ID NO: 2. These alternations may occur at the amino or carboxy termini or anywhere between those terminal positions, interspersed either subjectly among residues in the sequence or in one or more contiguous groups within the sequence.

In certain embodiments, a variant G protein-coupled receptor that may be used in the subject methods is a G protein-coupled receptor having an amino acid sequence derived from SEQ ID NO: 2 by deletion, substitution, and/or addition of one or several amino acids. In

certain embodiments, a variant G protein-coupled receptor that may be used in the subject methods is a G protein-coupled receptor having an amino acid sequence derived from SEQ ID NO: 2 by no more than 10 conservative amino acid substitutions and/or no more than 3 non-conservative amino acid substitutions in the amino acid sequence of SEQ ID NO: 2. In certain

5 embodiments, arginine, lysine and histidine may conservatively substitute for each other; glutamic acid and aspartic acid may conservatively substitute for each other; glutamine and asparagine may conservatively substitute for each other; leucine, isoleucine and valine may conservatively substitute for each other; phenylalanine, tryptophan and tyrosine may conservatively substitute for each other; and glycine, alanine, serine, threonine and methionine

10 may conservatively substitute for each other. The amino acid substitutions, amino acid deletions, and amino acid additions may be at any position (e.g., the C- or N-terminus, or at internal positions). In some embodiments, the variant is an endogenous G protein-coupled receptor. In some embodiments, the variant is an endogenous vertebrate G protein-coupled receptor. In some embodiments, the variant is an endogenous mammalian G protein-coupled

15 receptor. In some embodiments, the variant is an endogenous human G protein-coupled receptor. In some embodiments, the variant is a non-endogenous G protein-coupled receptor. In some embodiments, the variant exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing intracellular cAMP. In some

20 embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion. In certain embodiments, said G protein-coupled receptor having an amino acid sequence derived from SEQ ID NO: 2 is a G protein-coupled receptor for which Compound 1 is a ligand. In certain embodiments, said G protein-coupled receptor having an amino acid

25 sequence derived from SEQ ID NO: 2 is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is a ligand having an IC₅₀ value in receptor binding assay according to Example 15, *infra*, of less than about 50μM, less than about 25μM, less than about 10μM, less than about 5μM, less than about 1μM, less than about 500nM, less than about 100nM, or less than about 50nM. In certain embodiments, said G protein-coupled receptor having an amino acid sequence derived from SEQ ID NO: 2 is a G protein-coupled receptor for which Compound

30 1 is an agonist. In certain embodiments, said G protein-coupled receptor having an amino acid sequence derived from SEQ ID NO: 2 is a G protein-coupled receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist having an EC₅₀ value in in whole cell adenylyl cyclase assay according to Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than

35 about 5μM, less than about 1μM, less than about 100nM, less than about 50nM, less than about 25nM, less than about 10nM, less than about 5nM, or less than about 1nM.

A variant of SEQ ID NO: 2 that is a G protein-coupled receptor comprising at least 20, at least 30, at least 40, at least 50, at least 75, or at least 100 contiguous amino acids of SEQ ID NO: 2 is envisioned to be within the scope of the invention. In certain embodiments, the variant of SEQ ID NO: 2 comprising at least 20, at least 30, at least 40, at least 50, at least 75, or at least 100 contiguous amino acids of SEQ ID NO: 2 is a GPCR. In some embodiments, the variant of SEQ ID NO: 2 is an endogenous GPCR. In some embodiments, the variant of SEQ ID NO: 2 is a non-endogenous GPCR. In some embodiments, the variant of SEQ ID NO: 2 that is an endogenous GPCR is a mammalian GPCR. In certain embodiments, the variant of SEQ ID NO: 2 specifically binds Compound 1. In certain embodiments, the variant of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the variant of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine with an IC_{50} value in receptor binding assay according to Example 15, *infra*, of less than about $50\mu M$, less than about $25\mu M$, less than about $10\mu M$, less than about $5\mu M$, less than about $1\mu M$, less than about $500nM$, less than about $100nM$, or less than about $50nM$. In certain embodiments, the variant of SEQ ID NO: 2 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine with an IC_{50} value in receptor binding assay according to Example 15, *infra*, of less than about $10\mu M$, less than about $5\mu M$, less than about $1\mu M$, less than about $500nM$, less than about $100nM$, or less than about $50nM$. In certain embodiments, the variant of SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist having an EC_{50} value at said receptor in whole cell adenylyl cyclase assay according to Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than about $5\mu M$, less than about $1\mu M$, less than about $100nM$, less than about $50nM$, less than about $25nM$, less than about $10nM$, less than about $5nM$, or less than about $1nM$. In certain embodiments, the variant of SEQ ID NO: 2 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist having an EC_{50} value at said receptor in whole cell adenylyl cyclase assay according to Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than about $100nM$, less than about $50nM$, less than about $25nM$, less than about $10nM$, less than about $5nM$, or less than about $1nM$. In some embodiments, the variant of SEQ ID NO: 2 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion. In some embodiments, the G protein-coupled receptor comprising at least 20, at least 30, at least 40, at least 50, at least 75, or at least 100 contiguous amino acids of SEQ ID NO: 2 exhibits a detectable level of

constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In some embodiments, a variant GPCR that may be used in the subject methods is a
5 GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1. In some embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an endogenous GPCR. In some embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a non-endogenous GPCR. In some
10 embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 and that is an endogenous GPCR is a mammalian endogenous GPCR. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is SEQ ID NO:2 or an allele thereof. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent
15 conditions to the complement of SEQ ID NO: 1 is an allele of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is an ortholog of SEQ ID NO:2. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 specifically binds Compound 1. In certain embodiments, the GPCR encoded by
20 a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine with an IC_{50} value in receptor binding assay
25 according to Example 15, *infra*, of less than about 50 μ M, less than about 25 μ M, less than about 10 μ M, less than about 5 μ M, less than about 1 μ M, less than about 500 nM, less than about 100 nM, or less than about 50 nM. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 specifically binds (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine with an IC_{50} value in receptor binding assay
30 according to Example 15, *infra*, of less than about 10 μ M, less than about 5 μ M, less than about 1 μ M, less than about 500 nM, less than about 100 nM, or less than about 50 nM. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an
35 agonist having an EC_{50} value at said receptor in whole cell adenylyl cyclase assay according to

Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than about 5 μ M, less than about 1 μ M, less than about 100nM, less than about 50nM, less than about 25nM, less than about 10nM, less than about 5nM, or less than about 1nM. In certain embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 is a receptor for which (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine is an agonist having an EC₅₀ value at said receptor in whole cell adenylyl cyclase assay according to Example 10 or in HTRF[®] assay according to Example 11, *infra*, of less than about 100nM, less than about 50nM, less than about 25nM, less than about 10nM, less than about 5nM, or less than about 1nM. In some embodiments, the GPCR encoded by a polynucleotide hybridizing under stringent conditions to the complement of SEQ ID NO: 1 exhibits a detectable level of constitutive activity. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

Hybridization techniques are well known to the skilled artisan. In some embodiments, stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (1xSSC = 150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 μ g/ml denatured, sheared salmon sperm DNA; followed by washing the filter in 0.1xSSC at about 65°C. In some embodiments, stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (1xSSC = 150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 μ g/ml denatured, sheared salmon sperm DNA; followed by a wash in 0.1xSSC/0.1% SDS (sodium dodecyl sulfate) or in 0.2xSSC/0.1% SDS at about 50°C, at about 55°C, at about 60°C or at about 65°C. In some embodiments, said stringent conditions comprise washing at 65°C with 0.1xSSC. In some embodiments, said stringent conditions comprise washing at about 50°C, at about 55°C, at about 60°C, or at about 65° with 0.1xSSC/0.1% SDS or with 0.2xSSC/0.1% SDS.

In some embodiments, a GPCR that may be used in the subject methods is a non-endogenous, constitutively activated receptor comprising the amino acid sequence of SEQ ID NO: 2, wherein the leucine at amino acid position 224 of SEQ ID NO: 2 is substituted with an amino acid other than leucine. In some embodiments, the amino acid other than leucine is lysine. In some embodiments, the amino acid other than leucine is alanine. In some embodiments, the amino acid other than leucine is arginine. In some embodiments, the amino acid other than leucine is histidine. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, a GPCR of the invention comprises a constitutively active version of a G protein-coupled receptor having SEQ ID NO: 2. In some embodiments, the constitutively active version of the receptor is an endogenous constitutively active version having SEQ ID NO: 2. In some embodiments, the constitutively active version of the receptor is a non-endogenous constitutively active version having a mutation positioned at amino acid position 224 of SEQ ID NO: 2. In some embodiments, the mutated residue has been mutated to a residue other than leucine. In some embodiments, the mutated residue has been mutated to a lysine residue. In some embodiments, the mutated residue has been mutated to an alanine residue. In some embodiments, the mutated residue has been mutated to an arginine residue. In some embodiments, the mutated residue has been mutated to a histidine residue. In some embodiments, the constitutive activity is for increasing a level of intracellular cAMP. In some embodiments, the constitutive activity is for causing melanophore cells to undergo pigment dispersion.

In certain embodiments, a GPCR of the invention forms part of a fusion protein with a G protein.

a. Sequence identity

In certain embodiments, percent identity is evaluated using the Basic Local Alignment Search Tool ("BLAST"), which is well known in the art [See, e.g., Karlin and Altschul, Proc Natl Acad Sci USA (1990) 87:2264-2268; Altschul et al., J Mol Biol (1990) 215:403-410; Altschul et al., Nature Genetics (1993) 3:266-272; and Altschul et al., Nucleic Acids Res (1997) 25:3389-3402; the disclosure of each of which is herein incorporated by reference in its entirety]. The BLAST programs may be used with the default parameters or with modified parameters provided by the user. Preferably, the parameters are default parameters.

In certain embodiments, a preferred method for determining the best overall match between a query sequence (e.g., the amino acid sequence of SEQ ID NO:2) and a sequence to be interrogated, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp App Biosci (1990) 6:237-245; the disclosure of which is herein incorporated by reference in its entirety). In a sequence alignment the query and interrogated sequences are both amino acid sequences. The results of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group=25, Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=247 or the length of the interrogated amino acid sequence, whichever is shorter.

If the interrogated sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, the results, in percent identity, must be manually corrected because the FASTDB program does not account for N- and C-terminal truncations of

the interrogated sequence when calculating global percent identity. For interrogated sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the interrogated sequence, that are not matched/aligned with a corresponding
5 interrogated sequence residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to
10 the N- and C-termini of the interrogated sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query amino acid residues outside the farthest N- and C-terminal residues of the interrogated sequence.

For example, a 90 amino acid residue interrogated sequence is aligned with a 100-
15 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the interrogated sequence and therefore, the FASTDB alignment does not match/align with the first residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB
20 program. If the remaining 90 residues were perfectly matched, the final percent identity would be 90%.

In another example, a 90-residue interrogated sequence is compared with a 100-residue query sequence. This time the deletions are internal so there are no residues at the N- or C-termini of the interrogated sequence, which are not matched/aligned with the query. In this case, the
25 percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected. No other corrections are made for the purposes of the present invention.

b. Fusion proteins

30 In certain embodiments, a polypeptide of interest is a fusion protein, and may contain, for example, an affinity tag domain or a reporter domain. Suitable affinity tags include any amino acid sequence that may be specifically bound to another moiety, usually another polypeptide, most usually an antibody. Suitable affinity tags include epitope tags, for example, the V5 tag, the FLAG tag, the HA tag (from hemagglutinin influenza virus), the myc tag,
35 and the like, as is known in the art. Suitable affinity tags also include domains for which, binding substrates are known, e.g., HIS, GST and MBP tags, as is known in the art, and domains from other proteins for which specific binding partners, e.g., antibodies, particularly monoclonal

antibodies, are available. Suitable affinity tags also include any protein-protein interaction domain, such as a IgG Fc region, which may be specifically bound and detected using a suitable binding partner, e.g. the IgG Fc receptor. It is expressly contemplated that such a fusion protein may contain a heterologous N-terminal domain (e.g., an epitope tag) fused in-frame with a GPCR that has had its N-terminal methionine residue either deleted or substituted with an alternative amino acid.

Suitable reporter domains include any domain that can report the presence of a polypeptide. While it is recognized that an affinity tag may be used to report the presence of a polypeptide using, e.g., a labeled antibody that specifically binds to the tag, light emitting reporter domains are more usually used. Suitable light emitting reporter domains include luciferase (from, e.g., firefly, *Vargula*, *Renilla reniformis* or *Renilla muelleri*), or light emitting variants thereof. Other suitable reporter domains include fluorescent proteins, (from e.g., jellyfish, corals and other coelenterates as such those from *Aequoria*, *Renilla*, *Ptilosarcus*, *Stylatula* species), or light emitting variants thereof. Light emitting variants of these reporter proteins are very well known in the art and may be brighter, dimmer, or have different excitation and/or emission spectra, as compared to a native reporter protein. For example, some variants are altered such that they no longer appear green, and may appear blue, cyan, yellow, enhanced yellow red (termed BFP, CFP, YFP eYFP and RFP, respectively) or have other emission spectra, as is known in the art. Other suitable reporter domains include domains that can report the presence of a polypeptide through a biochemical or color change, such as β -galactosidase, β -glucuronidase, chloramphenicol acetyl transferase, and secreted embryonic alkaline phosphatase.

Also as is known in the art, an affinity tags or a reporter domain may be present at any position in a polypeptide of interest. However, in most embodiments, they are present at the C- or N-terminal end of a polypeptide of interest.

2. Nucleic acids encoding GPCR polypeptides of interest

Since the genetic code and recombinant techniques for manipulating nucleic acid are known, and the amino acid sequences of GPCR polypeptides of interest described as above, the design and production of nucleic acids encoding a GPCR polypeptide of interest is well within the skill of an artisan. In certain embodiments, standard recombinant DNA technology (Ausubel, et al, *Short Protocols in Molecular Biology*, 3rd ed., Wiley & Sons, 1995; Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, (1989) Cold Spring Harbor, N.Y.) methods are used. For example, GPCR coding sequences may be isolated from a library of GPCR coding sequence using any one or a combination of a variety of recombinant methods that do not need to be described herein. Subsequent substitution, deletion, and/or addition of nucleotides in the nucleic acid sequence encoding a protein may also be done using standard recombinant DNA techniques.

For example, site directed mutagenesis and subcloning may be used to introduce/delete/substitute nucleic acid residues in a polynucleotide encoding a polypeptide of interest. In other embodiments, PCR may be used. Nucleic acids encoding a polypeptide of interest may also be made by chemical synthesis entirely from oligonucleotides (e.g., Cello et al., Science (2002) 297:1016-8).

In some embodiments, the codons of the nucleic acids encoding polypeptides of interest are optimized for expression in cells of a particular species, particularly a mammalian, e.g., mouse, rat, hamster, non-human primate, or human, species. In some embodiments, the codons of the nucleic acids encoding polypeptides of interest are optimized for expression in cells of a particular species, particularly an amphibian species.

a. Vectors

The invention further provides vectors (also referred to as "constructs") comprising a subject nucleic acid. In many embodiments of the invention, the subject nucleic acid sequences will be expressed in a host after the sequences have been operably linked to an expression control sequence, including, e.g. a promoter. The subject nucleic acids are also typically placed in an expression vector that can replicate in a host cell either as an episome or as an integral part of the host chromosomal DNA. Commonly, expression vectors will contain selection markers, e.g., tetracycline or neomycin, to permit detection of those cells transformed with the desired DNA sequences (see, e.g., U.S. Pat. No. 4,704,362, which is incorporated herein by reference). Vectors, including single and dual expression cassette vectors are well known in the art (Ausubel, et al, *Short Protocols in Molecular Biology*, 3rd ed., Wiley & Sons, 1995; Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, (1989) Cold Spring Harbor, N.Y.). Suitable vectors include viral vectors, plasmids, cosmids, artificial chromosomes (human artificial chromosomes, bacterial artificial chromosomes, yeast artificial chromosomes, etc.), mini-chromosomes, and the like. Retroviral, adenoviral and adeno-associated viral vectors may be used.

A variety of expression vectors are available to those in the art for purposes of producing a polypeptide of interest in a cell and include expression vectors which are commercially available (e.g., from Invitrogen, Carlsbad, CA; Clontech, Mountain View, CA; Stratagene, La Jolla, CA). Commercially available expression vectors include, by way of non-limiting example, CMV promoter-based vectors. One suitable expression vector is pCMV. The expression vector may be adenoviral. An exemplary adenoviral vector may be purchased as AdEasy™ from Qbiogene (Carlsbad, CA) (He TC et al, Proc Natl Acad Sci USA (1998) 95:2509-2514; and US Patent No. 5,922,576; the disclosure of each of which is herein incorporated by reference in its entirety). Other suitable expression vectors will be readily apparent to those of ordinary skill in the art.

The subject nucleic acids usually comprise an single open reading frame encoding a subject polypeptide of interest, however, in certain embodiments, since the host cell for expression of the polypeptide of interest may be a eukaryotic cell, e.g., a mammalian cell, such as a human cell, the open reading frame may be interrupted by introns. Subject nucleic acid are typically part of a transcriptional unit which may contain, in addition to the subject nucleic acid 3' and 5' untranslated regions (UTRs) which may direct RNA stability, translational efficiency, etc. The subject nucleic acid may also be part of an expression cassette which contains, in addition to the subject nucleic acid a promoter, which directs the transcription and expression of a polypeptide of interest, and a transcriptional terminator.

Eukaryotic promoters can be any promoter that is functional in a eukaryotic host cell, including viral promoters and promoters derived from eukaryotic genes. Exemplary eukaryotic promoters include, but are not limited to, the following: the promoter of the mouse metallothionein I gene sequence (Hamer et al., J. Mol. Appl. Gen. 1:273-288, 1982); the TK promoter of Herpes virus (McKnight, Cell 31:355-365, 1982); the SV40 early promoter (Benoist et al., Nature (London) 290:304-310, 1981); the yeast gall gene sequence promoter (Johnston et al., Proc. Natl. Acad. Sci. (USA) 79:6971-6975, 1982); Silver et al., Proc. Natl. Acad. Sci. (USA) 81:5951-5955, 1984), the CMV promoter, the EF-1 promoter, Ecdysone-responsive promoter(s), tetracycline-responsive promoter, and the like. Viral promoters may be of particular interest as they are generally particularly strong promoters. In certain embodiments, a promoter is used that is a promoter of the target pathogen. Promoters for use in the present invention are selected such that they are functional in the cell type (and/or animal) into which they are being introduced. In certain embodiments, the promoter is a CMV promoter.

In certain embodiments, a subject vector may also provide for expression of a selectable marker. Suitable vectors and selectable markers are well known in the art and discussed in Ausubel, et al, (Short Protocols in Molecular Biology, 3rd ed., Wiley & Sons, 1995) and Sambrook, et al, (Molecular Cloning: A Laboratory Manual, Third Edition, (2001) Cold Spring Harbor, N.Y.). A variety of different genes have been employed as selectable markers, and the particular gene employed in the subject vectors as a selectable marker is chosen primarily as a matter of convenience. Known selectable marker genes include: the thymidine kinase gene, the dihydrofolate reductase gene, the xanthine-guanine phosphoribosyl transferase gene, CAD, the adenosine deaminase gene, the asparagine synthetase gene, the antibiotic resistance genes, e.g. tetr, ampr, Cmr or cat, kanr or neor (aminoglycoside phosphotransferase genes), the hygromycin B phosphotransferase gene, and the like.

As mentioned above, polypeptides of interest may be fusion proteins that contain an affinity domain and/or a reporter domain. Methods for making fusions between a reporter or tag and a GPCR, for example, at the C- or N-terminus of the GPCR, are well within the skill of one of skill in the art (e.g. McLean et al, Mol. Pharma. Mol Pharmacol. 1999 56:1182-91; Ramsay et

al., Br. J. Pharmacology, 2001, 315-323) and will not be described any further. It is expressly contemplated that such a fusion protein may contain a heterologous N-terminal domain (e.g., an epitope tag) fused in-frame with a GPCR that has had its N-terminal methionine residue either deleted or substituted with an alternative amino acid. It is appreciated that a polypeptide of interest may first be made from a native polypeptide and then operably linked to a suitable reporter/tag as described above.

The subject nucleic acids may also contain restriction sites, multiple cloning sites, primer binding sites, ligatable ends, recombination sites etc., usually in order to facilitate the construction of a nucleic acid encoding a polypeptide of interest.

b. Host cells

The invention further provides host cells comprising a vector comprising a subject nucleic acid. Suitable host cells include prokaryotic, e.g., bacterial cells (for example *E. coli*), as well as eukaryotic cells e.g. an animal cell (for example an insect, mammal, fish, amphibian, bird or reptile cell), a plant cell (for example a maize or *Arabidopsis* cell), or a fungal cell (for example a *S. cerevisiae* cell). In certain embodiments, any cell suitable for expression of a polypeptide of interest-encoding nucleic acid may be used as a host cell. Usually, an animal host cell line is used, examples of which are as follows: monkey kidney cells (COS cells), monkey kidney CVI cells transformed by SV40 (COS-7, ATCC CRL 165 1); human embryonic kidney cells (HEK-293 ["293"], Graham et al. J. Gen Virol. 36:59 (1977)); HEK-293T ["293T"] cells; baby hamster kidney cells (BHK, ATCC CCL 10); chinese hamster ovary-cells (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. (USA) 77:4216, (1980); Syrian golden hamster cells MCB3901 (ATCC CRL-9595); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CVI ATCC CCL 70); african green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL 51); TRI cells (Mather et al., Annals N. Y. Acad. Sci 383:44-68 (1982)); NIH/3T3 cells (ATCC CRL-1658); and mouse L cells (ATCC CCL-1).

In certain embodiments, melanophores are used. Melanophores are skin cells found in lower vertebrates. Relevant materials and methods will be followed according to the disclosure of U.S. Patent Number 5,462,856 and U.S. Patent Number 6,051,386. These patent disclosures are herein incorporated by reference in their entirety.

Additional cell lines will become apparent to those of ordinary skill in the art, and a wide variety of cell lines are available from the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209.

C. Screening of Candidate Compounds

1. Generic GPCR screening assay techniques

When a G protein receptor becomes active, it binds to a G protein (*e.g.*, Gq, Gs, Gi, Gz, Go) and stimulates the binding of GTP to the G protein. The G protein then acts as a GTPase and slowly hydrolyzes the GTP to GDP, whereby the receptor, under normal conditions, becomes deactivated. However, activated receptors continue to exchange GDP to GTP. A non-
5 hydrolyzable analog of GTP, [³⁵S]GTPγS, can be used to monitor enhanced binding to membranes which express activated receptors. It is reported that [³⁵S]GTPγS can be used to monitor G protein coupling to membranes in the absence and presence of ligand. An example of this monitoring, among other examples well-known and available to those in the art, was reported by Traynor and Nahorski in 1995. A preferred use of this assay system is for initial
10 screening of candidate compounds because the system is generically applicable to all G protein-coupled receptors regardless of the particular G protein that interacts with the intracellular domain of the receptor.

2. Specific GPCR screening assay techniques

Once candidate compounds are identified using the “generic” G protein-coupled
15 receptor assay (*i.e.*, an assay to select compounds that are agonists or inverse agonists), in some embodiments further screening to confirm that the compounds have interacted at the receptor site is preferred. For example, a compound identified by the “generic” assay may not bind to the receptor, but may instead merely “uncouple” the G protein from the intracellular domain.

a. Gs, Gz and Gi.

20 Gs stimulates the enzyme adenylyl cyclase. Gi (and Gz and Go), on the other hand, inhibit adenylyl cyclase. Adenylyl cyclase catalyzes the conversion of ATP to cAMP; thus, activated GPCRs that couple the Gs protein are associated with increased cellular levels of cAMP. On the other hand, activated GPCRs that couple Gi (or Gz, Go) protein are associated with decreased cellular levels of cAMP. *See, generally*, “Indirect Mechanisms of Synaptic
25 Transmission,” Chpt. 8, From Neuron To Brain (3rd Ed.) Nichols, J.G. et al eds. Sinauer Associates, Inc. (1992). Thus, assays that detect cAMP can be utilized to determine if a candidate compound is, *e.g.*, an inverse agonist to the receptor (*i.e.*, such a compound would decrease the levels of cAMP). A variety of approaches known in the art for measuring cAMP can be utilized; in some embodiments a preferred approach relies upon the use of anti-cAMP
30 antibodies in an ELISA-based format. Another type of assay that can be utilized is a whole cell second messenger reporter system assay. Promoters on genes drive the expression of the proteins that a particular gene encodes. Cyclic AMP drives gene expression by promoting the binding of a cAMP-responsive DNA binding protein or transcription factor (CREB) that then binds to the promoter at specific sites called cAMP response elements and drives the expression
35 of the gene. Reporter systems can be constructed which have a promoter containing multiple cAMP response elements before the reporter gene, *e.g.*, β-galactosidase or luciferase. Thus, an activated Gs-linked receptor causes the accumulation of cAMP that then activates the gene and

expression of the reporter protein. The reporter protein such as β -galactosidase or luciferase can then be detected using standard biochemical assays (Chen et al. 1995).

b. Go and Gq.

Gq and Go are associated with activation of the enzyme phospholipase C, which in turn hydrolyzes the phospholipid PIP₂, releasing two intracellular messengers: diacylglycerol (DAG) and inositol 1,4,5-triphosphate (IP₃). Increased accumulation of IP₃ is associated with activation of Gq- and Go-associated receptors. *See, generally, "Indirect Mechanisms of Synaptic Transmission,"* Chpt. 8, *From Neuron To Brain* (3rd Ed.) Nichols, J.G. et al eds. Sinauer Associates, Inc. (1992). Assays that detect IP₃ accumulation can be utilized to determine if a candidate compound is, *e.g.*, an inverse agonist to a Gq- or Go-associated receptor (*i.e.*, such a compound would decrease the levels of IP₃). Gq-associated receptors can also been examined using an AP1 reporter assay in that Gq-dependent phospholipase C causes activation of genes containing AP1 elements; thus, activated Gq-associated receptors will evidence an increase in the expression of such genes, whereby inverse agonists thereto will evidence a decrease in such expression, and agonists will evidence an increase in such expression. Commercially available assays for such detection are available.

3. GPCR Fusion Protein

The use of an endogenous, constitutively active GPCR or a non-endogenous, constitutively activated GPCR, for use in screening of candidate compounds for the direct identification of inverse agonists or agonists provides an interesting screening challenge in that, by definition, the receptor is active even in the absence of an endogenous ligand bound thereto. Thus, in order to differentiate between, *e.g.*, the non-endogenous receptor in the presence of a candidate compound and the non-endogenous receptor in the absence of that compound, with an aim of such a differentiation to allow for an understanding as to whether such compound may be an inverse agonist or agonist or have no affect on such a receptor, in some embodiments it is preferred that an approach be utilized that can enhance such differentiation. In some embodiments, a preferred approach is the use of a GPCR Fusion Protein.

Generally, once it is determined that a non-endogenous GPCR has been constitutively activated using the assay techniques set forth above (as well as others known to the art-skilled), it is possible to determine the predominant G protein that couples with the endogenous GPCR. Coupling of the G protein to the GPCR provides a signaling pathway that can be assessed. In some embodiments it is preferred that screening take place using a mammalian or a melanophore expression system, as such a system will be expected to have endogenous G protein therein. Thus, by definition, in such a system, the non-endogenous, constitutively activated GPCR will continuously signal. In some embodiments it is preferred that this signal be enhanced such that in the presence of, *e.g.*, an inverse agonist to the receptor, it is more likely

that it will be able to more readily differentiate, particularly in the context of screening, between the receptor when it is contacted with the inverse agonist.

The GPCR Fusion Protein is intended to enhance the efficacy of G protein coupling with the GPCR. The GPCR Fusion Protein may be preferred for screening with either an
 5 endogenous, constitutively active GPCR or a non-endogenous, constitutively activated GPCR because such an approach increases the signal that is generated in such screening techniques. This is important in facilitating a significant "signal to noise" ratio; such a significant ratio is preferred for the screening of candidate compounds as disclosed herein.

The construction of a construct useful for expression of a GPCR Fusion Protein is
 10 within the purview of those having ordinary skill in the art. Commercially available expression vectors and systems offer a variety of approaches that can fit the particular needs of an investigator. Important criteria in the construction of such a GPCR Fusion Protein construct include but are not limited to, that the GPCR sequence and the G protein sequence both be in-frame (preferably, the sequence for the endogenous GPCR is upstream of the G protein
 15 sequence), and that the "stop" codon of the GPCR be deleted or replaced such that upon expression of the GPCR, the G protein can also be expressed. The GPCR can be linked directly to the G protein, or there can be spacer residues between the two (preferably, no more than about 12, although this number can be readily ascertained by one of ordinary skill in the art). Based upon convenience, it is preferred to use a spacer. In some embodiments, it is preferred
 20 that the G protein that couples to the non-endogenous GPCR will have been identified prior to the creation of the GPCR Fusion Protein construct.

As noted above, activated GPCRs that couple to Gi, Gz and Go are expected to inhibit the formation of cAMP making assays based upon these types of GPCRs challenging (*i.e.*, the cAMP signal decreases upon activation, thus making the direct identification of, *e.g.*, agonists
 25 (which would further decrease this signal) challenging). As will be disclosed herein, it has been ascertained that for these types of receptors, it is possible to create a GPCR Fusion Protein that is not based upon the GPCR's endogenous G protein, in an effort to establish a viable cyclase-based assay. Thus, for example, an endogenous Gi coupled receptor can be fused to a Gs protein—such a fusion construct, upon expression, "drives" or "forces" the endogenous GPCR to couple
 30 with, *e.g.*, Gs rather than the "natural" Gi protein, such that a cyclase-based assay can be established. Thus, for Gi, Gz and Go coupled receptors, in some embodiments it is preferred that when a GPCR Fusion Protein is used and the assay is based upon detection of adenylyl cyclase activity, that the fusion construct be established with Gs (or an equivalent G protein that stimulates the formation of the enzyme adenylyl cyclase).

TABLE C

	Effect of cAMP Production upon	Effect of IP₃ Accumulation upon Activation	Effect of cAMP Production	Effect on IP₃
--	---	--	--	---------------------------------

G protein	Activation of GPCR (<i>i.e.</i> , constitutive activation or agonist binding)	of GPCR (<i>i.e.</i> , constitutive activation or agonist binding)	upon contact with an Inverse Agonist	Accumulation upon contact with an Inverse Agonist
Gs	Increase	N/A	Decrease	N/A
Gi	Decrease	N/A	Increase	N/A
Gz	Decrease	N/A	Increase	N/A
Go	Decrease	Increase	Increase	Decrease
Gq	N/A	Increase	N/A	Decrease

Equally effective is a G Protein Fusion construct that utilizes a Gq Protein fused with a Gs, Gi, Gz or Go Protein. In some embodiments a preferred fusion construct can be accomplished with a Gq Protein wherein the first six (6) amino acids of the G-protein α -subunit ("G α q") is deleted and the last five (5) amino acids at the C-terminal end of G α q is replaced with the corresponding amino acids of the G α of the G protein of interest. For example, a fusion construct can have a Gq (6 amino acid deletion) fused with a Gi Protein, resulting in a "Gq/Gi Fusion Construct". This fusion construct will force the endogenous Gi coupled receptor to couple to its non-endogenous G protein, Gq, such that the second messenger, for example, inositol triphosphate or diacylglycerol, can be measured *in lieu* of cAMP production.

4. Co-transfection of a Target Gi Coupled GPCR with a Signal-Enhancer Gs Coupled GPCR (cAMP Based Assays)

A Gi coupled receptor is known to inhibit adenylyl cyclase, and, therefore, decreases the level of cAMP production, which can make the assessment of cAMP levels challenging. In certain embodiments, an effective technique in measuring the decrease in production of cAMP as an indication of activation of a receptor that predominantly couples Gi upon activation can be accomplished by co-transfecting a signal enhancer, *e.g.*, a non-endogenous, constitutively activated receptor that predominantly couples with Gs upon activation (*e.g.*, TSHR-A623I; see *infra*), with the Gi linked GPCR. As is apparent, activation of a Gs coupled receptor can be determined based upon an increase in production of cAMP. Activation of a Gi coupled receptor leads to a decrease in production cAMP. Thus, the co-transfection approach is intended to advantageously exploit these "opposite" affects. For example, co-transfection of a non-endogenous, constitutively activated Gs coupled receptor (the "signal enhancer") with expression vector alone provides a baseline cAMP signal (*i.e.*, although the Gi coupled receptor will decrease cAMP levels, this "decrease" will be relative to the substantial increase in cAMP levels established by constitutively activated Gs coupled signal enhancer). By then co-transfecting the signal enhancer with the "target receptor", an inverse agonist of the Gi coupled

target receptor will increase the measured cAMP signal, while an agonist of the Gi coupled target receptor will decrease this signal.

Candidate compounds that are directly identified using this approach should be assessed independently to ensure that these do not target the signal enhancing receptor (this can be done prior to or after screening against the co-transfected receptors).

Composition/Formulation and Methods of Treatment

A compound of the invention, e.g. a GPR119 agonist, a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual, can be formulated into pharmaceutical compositions and medicaments for use in accordance with the present invention using techniques well known in the art. Proper formulation is dependent on the route of administration chosen. In certain embodiments, said administration is to a non-human vertebrate or to a non-human mammal.

As relates to therapies of the present invention, namely therapies relating to a GPR119 agonist, the compounds according to the invention can be administered in any suitable way. Suitable routes of administration include oral, nasal, rectal, transmucosal, transdermal, or intestinal administration, parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, intrapulmonary (inhaled) or intraocular injections using methods known in the art. Other suitable routes of administration are aerosol and depot formulation. Sustained release formulations, particularly depot, of the invented medicaments are expressly contemplated. In certain preferred embodiments, the compounds according to the present invention are administered orally. The compounds according to the present invention can be made up in solid or liquid form, such as tablets, capsules, powders, syrups, elixirs and the like, aerosols, sterile solutions, suspensions or emulsions, and the like. In certain embodiments, the GPR119 agonist is administered orally.

In certain preferred embodiments, the compounds according to the present invention, e.g. GPR119 agonists, are administered orally. In certain preferred embodiments, the compounds according to the present invention, e.g. GPR119 agonists, are administered in a manner that allows for stimulation of PYY secretion in gut enteroendocrine cells. In certain preferred embodiments, the compounds according to the present invention, e.g. GPR119 agonists, are administered orally in a manner that allows for stimulation of PYY secretion in gut enteroendocrine cells.

In certain preferred embodiments, the compounds according to the present invention, e.g. PYY secretagogues, compounds useful for treatment or prevention of a condition modulated by PYY, or compounds useful for a modulation of a PYY-related activity in an individual, are administered orally. In certain preferred embodiments, the compounds according to the present

invention, e.g. GPR119 agonists, are administered in a manner that allows for stimulation of PYY secretion in gut enteroendocrine cells. In certain preferred embodiments, the compounds according to the present invention, e.g. GPR119 agonists, are administered orally in a manner that allows for stimulation of PYY secretion in gut enteroendocrine cells.

5 Formulations for oral administration may be in the form of aqueous solutions and suspensions, in addition to solid tablet and capsule formulations. The aqueous solutions and suspensions may be prepared from sterile powders or granules. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants
10 are well and widely known in the art.

Pharmaceutical compositions of the GPR119 agonist may be prepared by methods well known in the art, e.g., by means of conventional mixing, dissolving, granulation, dragee-making, levigating, emulsifying, encapsulating, entrapping, lyophilizing processes or spray drying.

15 Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Suitable pharmaceutically acceptable carriers are available to those in the art (*see*, e.g., Remington: The Science and Practice of Pharmacy, (Gennaro et al., eds.), 20th Edition, 2000, Lippincott Williams & Wilkins; and Handbook of
20 Pharmaceutical Excipients (Rowe et al., eds), 4th Edition, 2003, Pharmaceutical Press; the disclosure of each of which is herein incorporated by reference in its entirety). Proper formulation is dependent upon the route of administration chosen. The term "carrier" material or "excipient" material herein means any substance, not itself a therapeutic agent, used as a
25 carrier and/or diluent and/or adjuvant, or vehicle for delivery of a therapeutic agent to a subject or added to a pharmaceutical composition to improve its handling or storage properties or to permit or facilitate formation of a dose unit of the composition into a discrete article such as a capsule or tablet suitable for oral administration. Excipients can include, by way of illustration and not limitation, diluents, disintegrants, binding agents, adhesives, wetting agents, polymers,
30 lubricants, glidants, substances added to mask or counteract a disagreeable taste or odor, flavors, dyes, fragrances, and substances added to improved appearance of the composition. Acceptable excipients include stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, magnesium carbonate, talc, gelatin, acacia gum, sodium alginate, pectin, dextrin, mannitol, sorbitol, lactose, sucrose, starches, gelatin, cellulosic
35 materials, such as cellulose esters of alkanolic acids and cellulose alkyl esters, low melting wax cocoa butter or powder, polymers, such as polyvinyl-pyrrolidone, polyvinyl alcohol, and

polyethylene glycols, and other pharmaceutically acceptable materials. The components of the pharmaceutical composition can be encapsulated or tableted for convenient administration.

Pharmaceutically acceptable refers to those properties and/or substances which are acceptable to the patient from a pharmacological/toxicological point of view and to the manufacturing pharmaceutical chemist from a physical/chemical point of view regarding
5 composition, formulation, stability, patient acceptance and bioavailability.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used which may optionally contain gum Arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable
10 organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical compositions which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with a filler such
15 as lactose, a binder such as starch, and/or a lubricant such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, liquid polyethylene glycols, cremophor, capmul, medium or long chain mono-, di- or triglycerides. Stabilizers may be added in these
20 formulations, also.

Additionally, a GPR119 agonist may be delivered using a sustained-release system. Various sustained-release materials have been established and are well known to those skilled in the art. Sustained-release tablets or capsules are particularly preferred. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. The
25 dosage form may also be coated by the techniques described in the U.S. Pat. Nos. 4,256,108, 4,166,452, and 4,265,874 to form osmotic therapeutic tablets for controlled release.

It is expressly contemplated that therapies of the present invention, namely therapies relating to a GPR119 agonist, may be administered or provided alone or in combination with one or more other pharmaceutically or physiologically acceptable compound. In one aspect of
30 the present invention, the other pharmaceutically or physiologically acceptable compound is not a GPR119 agonist. In one aspect of the present invention, the other pharmaceutically or physiologically acceptable compound is a pharmaceutical agent selected from the group consisting of calcium, vitamin D, estrogen, tibolone, selective estrogen receptor modulator (SERM; e.g., raloxifene, tamoxifen), biphosphonate (e.g., etidronate, alendronate, risedronate),
35 calcitonin, 1 α -hydroxylated metabolite of vitamin D, fluoride, thiazide, anabolic steroid, ipriflavone, vitamin K, parathyroid hormone (PTH), strontium, statin, osteoprotegerin, EP4 receptor selective agonist, cannabinoid receptor type 2 (CB2) selective agonist, and p38 MAP

kinase inhibitor. (See, e.g., World Health Organization Technical Report Series 921 (2003), Prevention and Management of Osteoporosis.)

In one aspect, the present invention features a composition comprising or consisting essentially of an amount of a GPR119 agonist according to the present invention. In one aspect, the present invention features a pharmaceutical composition comprising or consisting essentially of an amount of a GPR119 agonist according to the present invention and at least one pharmaceutically acceptable carrier.

In one aspect, the present invention features a composition comprising or consisting essentially of an amount of a GPR119 agonist according to the present invention. In one aspect, the present invention features a pharmaceutical composition comprising or consisting essentially of an amount of a GPR119 agonist according to the present invention and at least one pharmaceutically acceptable carrier. The present invention also relates to a dosage form of the composition or of the pharmaceutical composition wherein the GPR119 agonist, the PYY secretagogue, the compound useful for treatment or prevention of a condition modulated by PYY, or the compound useful for a modulation of a PYY-related activity in an individual is in an amount sufficient to increase a PYY level in an individual, such as a plasma total PYY level or a plasma PYY₃₋₃₆ level. The present invention also relates to a dosage form of the composition or of the pharmaceutical composition wherein the GPR119 agonist is in an amount sufficient to give an effect in treating or preventing a condition characterized by low bone mass, such as osteoporosis, and/or in increasing bone mass in an individual.

In one aspect, the present invention features a composition comprising or consisting essentially of an amount of a GPR119 agonist according to the present invention. In one aspect, the present invention features a pharmaceutical composition comprising or consisting essentially of an amount of a GPR119 agonist according to the present invention and at least one pharmaceutically acceptable carrier. The present invention also relates to a dosage form of the composition or of the pharmaceutical composition wherein the GPR119 agonist is in an amount sufficient to give an effect in increasing a PYY level in an individual. The present invention also relates to a dosage form of the composition or of the pharmaceutical composition wherein the GPR119 agonist is in an amount sufficient to give an effect in increasing a total PYY (inclusive of PYY₁₋₃₆ and PYY₃₋₃₆) level in an individual. The present invention also relates to a dosage form of the composition or of the pharmaceutical composition wherein the GPR119 agonist is in an amount sufficient to give an effect in increasing a PYY₃₋₃₆ level in an individual.

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an amount to achieve their intended purpose. In some embodiments, a pharmaceutical composition of the present invention is understood to be useful for treatment or prevention of a condition modulated by PYY. Conditions modulated by PYY are according to the present invention. In some embodiments, a

pharmaceutical composition of the present invention is understood to be useful for increasing a PYY level in an individual. As relates to the present invention, determination of the amount of a GPR119 agonist sufficient to achieve an intended purpose according to the invention is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

The data obtained from animal studies, including but not limited to studies using mice, rats, rabbits, pigs, and non-human primates, can be used in formulating a range of dosage for use in humans. In general, one skilled in the art understands how to extrapolate *in vivo* data obtained in an animal model system to another, such as a human. In some circumstances, these extrapolations may merely be based on the weight of the animal model in comparison to another, such as a human; in other circumstances, these extrapolations are not simply based on weights but rather incorporate a variety of factors. Representative factors include the type, age, weight, sex, diet and medical condition of the patient, the severity of the disease, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetic and toxicology profiles of the particular compound employed, whether a drug delivery system is utilized, on whether an acute or chronic disease state is being treated or prophylaxis is conducted or on whether further active compounds are administered in addition to the compounds of the present invention and as part of a drug combination. The dosage regimen for treating a disease condition with the compounds and/or compositions of this invention is selected in accordance with a variety factors as cited above. Thus, the actual dosage regimen employed may vary widely and therefore may deviate from a preferred dosage regimen and one skilled in the art will recognize that dosage and dosage regimen outside these typical ranges can be tested and, where appropriate, may be used in the methods of this invention.

An exemplary animal model system is the rat ovariectomy (OVX) bone loss model. The ovariectomized rat is an excellent preclinical animal model that correctly emulates the important clinical feature of the estrogen depleted human skeleton and the response of therapeutic agents. In this model, a therapeutic efficacy is achieved when the bone loss associated with ovariectomy is partially or completely prevented. (See, e.g., Bollag et al, Mol Cell Endocrinol (2001) 177:35-41; and Jee et al, J Musculoskel Neuron Interact (2001) 1:193-207.) In certain embodiments, therapeutic efficacy is achieved when the bone loss associated with ovariectomy is at least about 10% prevented, at least about 20% prevented, at least about 30% prevented, at least about 40% prevented, at least about 50% prevented, at least about 60% prevented, at least about 70% prevented, at least about 75% prevented, at least about 80% prevented, at least about 85% prevented, at least about 90% prevented, at least about 95% prevented, or 100% prevented.

An additional exemplary animal model system is increase of a blood PYY level after glucose challenge in mice. In certain embodiments, the blood PYY level is a plasma PYY level.

In certain embodiments, the PYY level is a glucose-independent PYY level. In certain embodiments, the PYY level is a glucose-dependent PYY level. In certain embodiments, the PYY is total PYY, inclusive of PYY₁₋₃₆ and PYY₃₋₃₆. In certain embodiments, the PYY is PYY₃₋₃₆. Kits for measuring total PYY or PYY₃₋₃₆ in rodents or humans are commercially available, such as from Phoenix Pharmaceuticals, Inc. (Burlingame, CA). Also, PYY₁₋₃₆ and PYY₃₋₃₆ can be independently measured by biochemical means (Eberlein et al, Peptides (1989) 10:797-803; Grandt et al, Regul Pept (1994) 51:151-159; Cruze et al, Peptides (2007) 28:269-280). In certain embodiments, a PYY-related activity of the invention is a PYY₃₋₃₆-related activity. In certain embodiments, therapeutic efficacy is achieved when a blood or plasma PYY level is increased by at least about 10%, at least about 25%, at least about 50%, at least about 100%, at least about 150%, at least about 200%, at least about 300%, at least about 400%, at least about 500%, at least about 600%, at least about 700%, at least about 800%, at least about 900%, or at least about 1000%.

It is expressly contemplated that a GPR119 agonist can be administered on a daily or regular basis to achieve an increased level of PYY in an individual. In certain embodiments, a GPR119 agonist is administered on a daily or regular basis to achieve an increased blood (e.g., plasma) level of PYY in an individual. In certain embodiments, a GPR119 agonist is administered on a daily or regular basis to achieve an increased blood (e.g., plasma) level of PYY in an individual that is at least about 10%, at least about 25%, at least about 50%, at least about 100%, at least about 150%, at least about 200%, at least about 300%, at least about 400%, at least about 500%, at least about 600%, at least about 700%, at least about 800%, at least about 900%, or at least about 1000% a normal blood (e.g., plasma) level of PYY (e.g., a normal pre-meal plasma PYY level (e.g., plasma total PYY level) or a plasma PYY level (e.g., a plasma total PYY level) between the normal pre-meal and post-meal plasma PYY levels) in an individual or the blood (e.g., plasma) level of PYY (e.g., plasma total PYY) in the individual prior to treatment. It is noted that these ranges of increased blood (e.g., plasma) level of PYY (e.g., plasma total PYY) are exemplary ranges and are not meant to be limiting to the invention.

It is expressly contemplated that a GPR119 agonist can be administered on a daily or regular basis to achieve an increased level of PYY in an individual. In certain embodiments, a GPR119 agonist is administered on a daily or regular basis to achieve an increased blood (e.g., plasma) level of PYY in an individual. In certain embodiments, a GPR119 agonist is administered on a daily or regular basis to achieve an increased blood (e.g., plasma) level of PYY in an individual within a concentration range of 0.1 ng/ml to 10 ng/ml or 100 ng/ml, 0.2 ng/ml to 10 ng/ml or 100 ng/ml, 0.3 ng/ml to 10 ng/ml or 100 ng/ml, 0.4 ng/ml to 10 ng/ml or 100 ng/ml, 0.5 ng/ml to 10 ng/ml or 100 ng/ml, 0.6 ng/ml to 10 ng/ml or 100 ng/ml, 0.7 ng/ml to 10 ng/ml or 100 ng/ml, 0.8 ng/ml to 10 ng/ml or 100 ng/ml, 0.9 ng/ml to 10 ng/ml or 100 ng/ml, 1 ng/ml to 10 ng/ml or 100 ng/ml, 2 ng/ml to 10 ng/ml or 100 ng/ml, 3 ng/ml to 10 ng/ml or 100

ng/ml, or 5 ng/ml to 10 ng/ml or 100 ng/ml. It is noted that these ranges of blood (e.g., plasma) PYY concentration (e.g., plasma total PYY concentration) are exemplary ranges and are not meant to be limiting to the invention.

5 Dosage amount and interval may be adjusted in order to provide an intended therapeutic effect. It will be appreciated that the exact dosage of a GPR119 agonist in accordance with the present invention will vary depending on the GPR119 agonist, its potency, the mode of administration, the age and weight of the patient and the severity of the condition to be treated. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. By way of illustration and not limitation, an
10 amount of a GPR119 agonist in accordance with the present invention is an amount less than about 0.001 mg/kg body weight, less than about 0.005 mg/kg body weight, less than about 0.01 mg/kg body weight, less than about 0.05 mg/kg body weight, less than about 0.1 mg/kg body weight, less than about 0.5 mg/kg body weight, less than about 1 mg/kg body weight, less than about 5 mg/kg body weight, less than about 10 mg/kg body weight, less than about 50 mg/kg
15 body weight, or less than about 100 mg/kg body weight.

A preferred dosage range for the amount of a compound of the invention (e.g. a GPR119 agonist, a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity in an individual), which can be administered on a daily or regular basis to achieve desired results
20 is 0.1-100 mg/kg body mass. Other preferred dosage range is 0.1-30 mg/kg body mass. Other preferred dosage range is 0.1-10 mg/kg body mass. Other preferred dosage range is 0.1-3.0 mg/kg body mass. Exemplary preferred dosage is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg. Of course, these daily dosages can be delivered or administered in small amounts periodically during the course
25 of a day. It is noted that these dosage ranges are only preferred ranges and are not meant to be limiting to the invention. It is noted that these dosages are only preferred dosages and are not meant to be limiting to the invention.

Dosage amount and interval may be adjusted individually to provide plasma levels of a GPR119 agonist according to the present invention which achieve an intended therapeutic
30 effect. Dosage intervals can also be determined using the value for a selected range of GPR119 agonist concentration so as to achieve the intended therapeutic effect. A GPR119 agonist should be administered using a regimen that maintains plasma levels within the selected range of GPR119 agonist concentration for 10-90% of the time, preferably between 30-99% of the time, and most preferably between 50-90% of the time. In cases of local administration or selective
35 uptake, the range of GPR119 agonist concentration providing the intended therapeutic effect may not be related to plasma concentration.

The amount of composition administered will, of course, be dependent on the individual being treated, on the individual's weight, the severity of the affliction, the manner of administration, and the judgement of the prescribing physician.

5 In one aspect, the present invention accordingly features a method of treating or preventing a condition characterized by low bone mass, such as osteoporosis, or of increasing bone mass comprising administering to an individual in need thereof a therapeutically effective amount of a composition comprising or consisting essentially of an amount of a GPR119 agonist according to the present invention. In certain embodiments, the composition is a pharmaceutical composition.

10 In one aspect, the present invention relates to a method of treating or preventing a condition characterized by low bone mass, such as osteoporosis, or of increasing bone mass comprising administering to an individual in need thereof a therapeutically effective amount of a composition comprising or consisting essentially of an amount of a GPR119 agonist according to the present invention. In a related aspect, the present invention features said method wherein
15 the GPR119 agonist is administered in an amount sufficient to give an effect in increasing a PYY level in the individual. In certain embodiments, the composition is a pharmaceutical composition.

Therapies of the present invention, such as therapies relating to a GPR119 agonist, are useful in treatment or prevention of a condition modulated by PYY.

20 In certain embodiments, the condition modulated by PYY is a bone-related condition.

In certain embodiments, the bone-related condition is a condition characterized by low bone mass. In certain embodiments, the condition characterized by low bone mass is selected from the group consisting of osteopenia, osteoporosis, rheumatoid arthritis, osteoarthritis, periodontal disease, alveolar bone loss, osteotomy bone loss, childhood idiopathic bone loss, Paget's disease,
25 bone loss due to metastatic cancer, osteolytic lesions, curvature of the spine, and loss of height.

In certain embodiments, the bone-related condition is a bone fracture. In certain embodiments, the bone fracture is selected from the group consisting of a traumatic bone fracture, a long-term bone fracture, and an osteoporotic bone fracture.

In certain embodiments, the bone-related condition is a bone disease. In certain
30 embodiments, the bone disease is selected from the group consisting of osteopenia, osteoporosis, rheumatoid arthritis, osteoarthritis, periodontal disease, alveolar bone loss, osteotomy bone loss, childhood idiopathic bone loss, Paget's disease, bone loss due to metastatic cancer, osteolytic lesions, curvature of the spine, and loss of height.

In certain embodiments, the bone-related condition is enhanced bone healing or bone
35 growth, wherein said enhanced bone healing or bone growth is selected from the group consisting of enhanced bone healing following facial reconstruction, maxillary reconstruction, mandibular

reconstruction, periodontal disease or tooth extraction, enhanced long bone extension, enhanced prosthetic ingrowth, and increased bone synostosis.

In certain embodiments, the condition modulated by PYY is a metabolic disorder.

5 In certain embodiments, the metabolic disorder is selected from the group consisting of overweight, obesity, hyperphagia, diabetes (inclusive of type 1 diabetes and type 2 diabetes), type 1 diabetes, type 2 diabetes, impaired glucose tolerance, insulin resistance, hyperinsulinemia, dyslipidemia, hypertension and metabolic syndrome.

In certain embodiments, the condition modulated by PYY is an angiogenesis-related condition.

10 In certain embodiments, the angiogenesis-related condition is selected from the group consisting of peripheral arterial disease, wound healing and ischemia.

In certain embodiments, the condition modulated by PYY is an ischemia-related condition.

15 In certain embodiments, the ischemia-related condition is selected from the group consisting of cerebral ischemia, stroke, ischemic heart disease, myocardial infarction, ischemia-reperfusion injury, post-myocardial infarction remodeling, heart failure, and congestive heart failure.

In certain embodiments, the condition modulated by PYY is a malabsorptive disorder.

20 In certain embodiments, the malabsorptive disorder is selected from the group consisting of short bowel syndrome, enterocolitis, Crohn's disease, ulcerative colitis, irritable bowel syndrome, cholera, and a disorder accompanied by diarrhea.

In certain embodiments, the condition modulated by PYY is a convulsive disorder.

In certain embodiments, the convulsive disorder is epilepsy.

In certain embodiments, the condition modulated by PYY is a cancer.

25 In certain embodiments, the cancer is selected from the group consisting of pancreatic cancer, breast cancer, colon cancer, and Barrett's adenocarcinoma.

In certain embodiments, the condition modulated by PYY is an inflammatory disorder.

30 In certain embodiments, the inflammatory disorder is selected from the group consisting of atherosclerosis, atherothrombosis, psoriasis, psoriatic arthritis, rheumatoid arthritis, osteoarthritis, insulin resistance, diabetes (inclusive of type 1 diabetes and type 2 diabetes), type 1 diabetes, type 2 diabetes, metabolic syndrome, obesity, lower than normal HDL-cholesterol, hypertension, hyperlipidemia, ischemic heart disease, myocardial infarction, congestive heart failure, osteoporosis, higher than normal osteoclastogenesis, pancreatitis, interstitial nephritis, acute transplant rejection, chronic hepatitis, hepatic steatosis, multiple sclerosis, systemic lupus erythematosus (SLE), restenosis, ischemia-reperfusion injury, allergic inflammation, Parkinson's
35 disease, prion-associated disease, excitotoxic injury, Alzheimer's disease, mild cognitive impairment (MCI), disseminated intravascular coagulation, septic shock, an inflammatory lung disease, and an inflammatory bowel disease. In certain embodiments, the inflammatory lung

disease is selected from the group consisting of asthma, emphysema, interstitial lung disease, idiopathic pulmonary fibrosis, bronchiolitis obliterans syndrome, chronic obstructive pulmonary disease (COPD), and pulmonary arterial hypertension. In certain embodiments, the inflammatory bowel disease is selected from the group consisting of enteritis, enterocolitis, Crohn's disease, 5 granulomatous colitis, ileitis, Crohn's colitis, ulcerative colitis, sigmoiditis, pancolitis, and ulcerative proctitis.

In certain embodiments, the inflammatory disorder is selected from the group consisting of atherosclerosis, atherothrombosis, insulin resistance, pancreatitis, restenosis, an inflammatory lung disease, and an inflammatory bowel disease. In certain embodiments, the inflammatory lung 10 disease is selected from the group consisting of asthma, emphysema, and pulmonary arterial hypertension. In certain embodiments, the inflammatory bowel disease is selected from the group consisting of enteritis, enterocolitis, Crohn's disease, granulomatous colitis, ileitis, Crohn's colitis, ulcerative colitis, sigmoiditis, pancolitis, and ulcerative proctitis.

Conditions characterized by low bone mass include but are not limited to osteopenia, 15 osteoporosis, rheumatoid arthritis, osteoarthritis, periodontal disease, alveolar bone loss, osteotomy bone loss, childhood idiopathic bone loss, Paget's disease, bone loss due to metastatic cancer, osteolytic lesions, curvature of the spine, and loss of height. In certain embodiments, the condition characterized by low bone mass is osteoporosis. In certain embodiments, the condition characterized by low bone mass is osteoporosis. In certain 20 embodiments, osteoporosis is primary osteoporosis. In certain embodiments, osteoporosis is secondary osteoporosis. Conditions characterized by low bone mass also include but are not limited to long-term complications of osteoporosis such as curvature of the spine, loss of height and prosthetic surgery. It is understood that conditions characterized by low bone mass can be included in embodiments individually or in any combination. In certain embodiments, the 25 condition characterized by low bone mass is primary osteoporosis.

In certain embodiments, the individual in need of increased bone mass has a bone mineral density (BMD) of greater than 1 (T-score < -1), greater than or equal to 1.5 (T-score ≤ -1.5), greater than or equal to 2 (T-score ≤ -2) or greater than or equal to 2.5 (T-score ≤ -2.5) standard deviations below the young adult reference mean. In certain embodiments, the 30 individual in need of increased bone mass is in need of treatment of bone fracture. In certain embodiments, the individual in need of treatment of a bone fracture has a traumatic bone fracture, a long-term bone fracture, or an osteoporotic bone fracture. In certain embodiments, the individual is in need of treatment for a bone disease. In certain embodiments, the individual in need of treatment for a bone disease has osteopenia, osteoporosis, rheumatoid arthritis, 35 osteoarthritis, periodontal disease, alveolar bone loss, osteotomy bone loss, childhood idiopathic bone loss, Paget's disease, bone loss due to metastatic cancer, osteolytic lesions, curvature of the spine, or loss of height. In certain embodiments, the individual in need of treatment for a bone

disease has osteoporosis. In certain embodiments, osteoporosis is primary osteoporosis. In certain embodiments, osteoporosis is secondary osteoporosis. Destructive bone disorders that can be treated according to the invention include but are not limited to osteoporosis, osteoarthritis, and osteolytic lesions such as those caused by neoplastic disease, radiotherapy, or chemotherapy. In certain embodiments, osteoporosis is primary osteoporosis. In certain
5 embodiments, osteoporosis is secondary osteoporosis.

Therapies of the present invention, such as therapies relating to a GPR119 agonist are additionally useful in enhancing bone healing following facial reconstruction, maxillary reconstruction, mandibular reconstruction, periodontal disease or tooth extraction, enhancing
10 long bone extension, enhancing prosthetic ingrowth or increasing bone synostosis in an individual.

Therapies of the present invention, such as therapies relating to a GPR119 agonist, are useful in a modulation of a PYY-related activity in an individual. In certain embodiments, the modulation of a PYY-related activity is selected from the group consisting of:

- 15 (a) increasing plasma PYY;
- (b) increasing bone mass;
- (c) decreasing loss of bone mass;
- (d) increasing bone formation;
- (e) decreasing food uptake;
- 20 (f) decreasing body weight;
- (g) increasing satiety;
- (h) increasing glucose disposal;
- (i) decreasing plasma glucose;
- (j) increasing angiogenesis;
- 25 (k) decreasing secretion in intestinal epithelium;
- (l) increasing absorption in intestinal epithelium; and
- (m) increasing plasma adiponectin

In certain embodiments, the individual is a vertebrate. In certain embodiments, the individual that is a vertebrate is a fish, an amphibian, a reptile, a bird or a mammal. In certain
30 embodiments, the individual or vertebrate is a mammal. In certain embodiments, the individual or vertebrate that is a mammal is a mouse, a rat, a hamster, a rabbit, a pig, a dog, a cat, a horse, a cow, a sheep, a goat, a non-human mammal, a non-human primate or a human. In certain embodiments, the individual is a human. In certain embodiments, the human is a post-menopausal woman or a man over the age of 50.

35 Without further elaboration, it is believed that one skilled in the art can, using the preceding description, practice the present invention to its fullest extent. The foregoing detailed description is given for clearness of understanding only, and no unnecessary limitation should

be understood therefrom, as modifications within the scope of the invention may become apparent to those skilled in the art.

Throughout this application, various publications, patents and patent applications are cited. The disclosures of these publications, patents and patent applications referenced in this application are herein incorporated by reference in their entirety into the present disclosure. Citation herein by Applicant of a publication, patent, or patent application is not an admission by Applicant of said publication, patent, or patent application as prior art.

10

EXAMPLES

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, practice the present invention to its fullest extent. The following detailed examples are to be construed as merely illustrative, and not limitations of the preceding disclosure in any way whatsoever. Those skilled in the art will promptly recognize appropriate variations from the procedures.

15

EXAMPLE 1

GPR119 RECEPTOR EXHIBITS CONSTITUTIVE ACTIVITY FOR INCREASING cAMP ACCUMULATION IN TRANSFECTED HEK293 CELLS

GPR119 receptor was shown to exhibit constitutive activity in cAMP assay. cAMP measurements were done with a Flash Plate adenylyl cyclase kit (NEN Life Science Products, Boston, MA) according to the supplier's protocol. Briefly, HEK293 cells were transfected with either empty vector DNA ("Vector") or human GPR119 receptor expression plasmid DNA ("GPR119") using Lipofectamine (Invitrogen, Carlsbad, CA). After 24 h, transfected cells were harvested in GIBCO (Gaithersburg, MD) cell dissociation buffer (Catalog no. 13151-014) and resuspended in assay buffer (50% 1X PBS/50% stimulation buffer). Incubation was carried out with 10^5 cells per well for 60 min at room temperature. After another 2-h incubation with tracer in detection buffer, plates were counted in a Wallac (Gaithersburg, MD) MicroBeta scintillation counter. Values of cAMP per well were extrapolated from a standard cAMP curve that was included on each assay plate. Results are presented in *Figure 1*. From the results shown in *Figure 1*, it is apparent that GPR119 receptor exhibits constitutive activity for increasing cAMP accumulation in transfected HEK293 cells.

20

25

30

EXAMPLE 2

COMPOUND 1 IS AN AGONIST OF GPR119 RECEPTOR AS EVIDENCED, E.G., BY STIMULATION OF cAMP ACCUMULATION IN GPR119-TRANSFECTED HEK293 CELLS

35

Compound 1 (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine) was shown to be an agonist of GPR119 receptor in cAMP assay. cAMP measurements were done with a Flash Plate adenylyl cyclase kit (NEN Life Science Products, Boston, MA) according to the supplier's protocol. Briefly, HEK293 cells were transfected with either empty vector DNA ("Vector") or human GPR119 receptor expression plasmid DNA ("GPR119") using Lipofectamine (Invitrogen, Carlsbad, CA). After 24 h, transfected cells were harvested in GIBCO (Gaithersburg, MD) cell dissociation buffer (Catalog no. 13151-014) and resuspended in assay buffer (50% 1X PBS/50% stimulation buffer). **Compound 1** was incubated with 10^5 cells per well for 60 min at room temperature. After another 2-h incubation with tracer in detection buffer, plates were counted in a Wallac (Gaithersburg, MD) MicroBeta scintillation counter. Values of cAMP per well were extrapolated from a standard cAMP curve that was included on each assay plate. Results are presented in **Figure 2**. From the results shown in **Figure 2**, it is apparent that **Compound 1** is an agonist of GPR119 receptor.

15

EXAMPLE 3

EFFECT OF ADMINISTRATION OF GPR119 AGONIST ON PLASMA PYY IN WILD-TYPE MICE

C57BL/6 male mice (approximately 8 weeks of age; obtained from Harlan, Indianapolis, IN) were fasted for 18 hours, and randomly assigned into eight groups with n=6 for each group. Mice were administered per orally with vehicle (80% polyethylene glycol 400/10% Tween 80/10% ethanol) or with a GPR119 agonist (**Compound 1**; (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine) at 10 mg/kg, as indicated in **Figure 3**. Thirty minutes after treatment, a glucose bolus at 3g/kg was delivered per orally, and blood was collected at 0 minute (no glucose bolus), 2 minutes, 5 minutes, and 10 minutes after glucose bolus. Plasma total PYY (inclusive of PYY₁₋₃₆ and PYY₃₋₃₆) levels were determined by using a PYY EIA kit purchased from Phoenix Pharmaceuticals, Inc. (Peptide YY (PYY) (Rat, Mouse, Porcine, Canine) EIA kit, Catalog No. EK-059-03) according to instructions provided in the kit. From the results shown in **Figure 3**, it is apparent that administration of the GPR119 agonist (**Compound 1**) increased plasma total PYY in the the mice. From the results shown in **Figure 3**, it is apparent that the GPR119 agonist (**Compound 1**) stimulated plasma total PYY in the mice. **Compound 1** is identical to a compound disclosed in International Patent Application No. PCT/US2004/001267 (published as WO 2004/065380).

25
30

EXAMPLE 4

EFFECT OF ADMINISTRATION OF GPR119 AGONIST ON PLASMA PYY IN GPR119-DEFICIENT (KNOCKOUT) MICE COMPARED TO WILD-TYPE MICE

35

GPR119-deficient male mice (Chu et al, Endocrinology (2007) 148:2601-2609) and wild-type littermates are fasted for 18 hours. Mice are administered per orally with vehicle (80% polyethylene glycol 400/10% ethanol/10% Tween 80) or with a GPR119 agonist in accordance with the present invention (**Compound 1**; (2-Fluoro-4-methanesulfonyl-phenyl)-{6-[4-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-yl]-5-nitro-pyrimidin-4-yl}-amine) at 20 mg/kg. Thirty minutes after treatment, blood (100 microliter) is collected via retro orbital vein of the eye (time 0 minutes) followed by a glucose bolus at 3g/kg (per orally). Five minutes after delivering glucose, another blood sample (100 microliter) is collected (time 5 minutes). Plasma are collected after centrifugation and plasma total PYY levels are determined by using PYY EIA kit purchased from Phoenix Pharmaceuticals, Inc. (Peptide YY (PYY) (Rat, Mouse, Porcine, Canine) EIA kit, Catalog No. EK-059-03) according to instructions provided in the kit. It can be further shown that functional GPR119 receptor is necessary for the administered GPR119 agonist to increase plasma total PYY in the mice. **Compound 1** can be shown to stimulate plasma total PYY in the wild-type mice but not in the GPR119-deficient mice. **Compound 1** is identical to a compound disclosed in International Patent Application No. PCT/US2004/001267 (published as WO 2004/065380).

EXAMPLE 5

EFFECT OF ADMINISTRATION OF GPR119 AGONIST ON BONE MASS IN OVARECTOMIZED RATS

A GPR119 agonist in accordance with the present invention can be shown to be effective in treating or preventing a condition characterized by low bone mass, such as osteoporosis, and/or in increasing bone mass in an individual using the *in vivo* ovariectomized (OVX) rat model described below (see, e.g., Bollag et al, Mol Cell Endocrinol (2001) 177:35-41).

Twenty virgin female OVX and 20 virgin non-OVX Sprague-Dawley rats (150-175 g), age 8 weeks, are purchased from Harlan Sprague-Dawley, Inc. (Indianapolis, IN). Animals are fed ad libitum on a normal commercial pellet diet, Teklab Rodent diet (1.46% calcium), with free access to water. The rats are randomly divided into four weight-matched experimental groups and selected to receive per orally vehicle or a GPR119 agonist in accordance with the present invention. Treatment is continued on a daily basis for 6 weeks.

1. **Control.** Ten non-OVX rats are administered per orally vehicle.
2. **Control + Treatment.** Ten non-OVX rats are administered per orally GPR119 agonist.
3. **OVX.** Ten OVX rats are administered per orally vehicle.
4. **OVX + Treatment.** Ten OVX rats are administered per orally GPR119 agonist.

The rats are weighed daily and length measured at baseline and again at 6 weeks. Dual energy X-ray absorptiometry (DXA) using a Hologic QDR 1000/W (Waltham, MA) is performed on all

animals prior to initiation of treatment and at 6 weeks, and data is analyzed using the software Rat Whole Body version 5.53. Bone mineral density (BMD) is determined at the spine.

The percent change in vertebral bone density after 6 weeks of treatment is determined. It is shown that administration of a GPR119 agonist attenuates the negative effects of ovariectomy on vertebral bone density. Attenuation of the negative effects of ovariectomy on vertebral bone density is indicative of the treatment having efficacy in treating or preventing a condition characterized by low bone mass, such as osteoporosis, and/or in increasing bone mass in an individual.

10

EXAMPLE 6

EFFECT OF ADMINISTRATION OF GPR119 AGONIST ON BONE FRACTURE HEALING

A GPR119 agonist in accordance with the present invention can be shown to be effective in treatment of bone fracture using the *in vivo* assay described below.

Fracture Technique

Sprague-Dawley rats at 3 months of age are anesthetized with Ketamine. A 1 cm incision is made on the anteromedial aspect of the proximal part of the right tibia or femur. The following describes the tibial surgical technique. The incision is carried through to the bone, and a 1 mm hole is drilled 4 mm proximal to the distal aspect of the tibial tuberosity 2 mm medial to the anterior ridge. Intramedullary nailing is performed with a 0.8 mm stainless steel tube (maximum load 36.3 N, maximum stiffness 61.8 N/mm, tested under the same conditions as the bones). No reaming of the medullary canal is performed. A standardized closed fracture is produced 2 mm above the tibiofibular junction by three-point bending using specially designed adjustable forceps with blunt jaws. To minimize soft tissue damage, care is taken not to displace the fracture. The skin is closed with monofilament nylon sutures. The operation is performed under sterile conditions. Radiographs of all fractures are taken immediately after nailing, and rats with fractures outside the specified diaphyseal area or with displaced nails are excluded. The remaining animals are divided randomly into the following groups with 10-12 animals per each subgroup per time point for testing the fracture healing. The rats are administered on a daily basis per orally with vehicle or with a GPR119 agonist. The GPR119 agonist is used at an amount between 0.001 mg/kg body weight and 100 mg/kg body weight. Treatment is continued for 10, 20, 40 and 80 days.

At 10, 20, 40 and 80 days, 10-12 rats from each group are anesthetized with Ketamine and sacrificed by exsanguination. Both tibiofibular bones are removed by dissection and all soft tissue is stripped. Bones from 5-6 rats for each group are stored in 70% ethanol for histological analysis, and bones from another 5-6 rats for each group are stored in a buffered Ringer's solution (+4°C, pH 7.4) for radiographs and biomechanical testing which is performed.

Histological Analysis

The methods for histological analysis of fractured bone have been previously published by Mosekilde and Bak (Bone (1993) 14:19-27). Briefly, the fracture site is sawed 8 mm to each side of the fracture line, embedded undecalcified in methymethacrylate, and cut frontals sections on a Reichert-Jung Polycut microtome in 8 μ m thick. Masson-Trichome stained mid-frontal sections (including both tibia and fibula) are used for visualization of the cellular and tissue response to fracture healing with and without treatment. Sirius red stained sections are used to demonstrate the characteristics of the callus structure and to differentiate between woven bone and lamellar bone at the fracture site. The following measurements are performed: (1) fracture gap—measured as the shortest distance between the cortical bone ends in the fracture, (2) callus length and callus diameter, (3) total bone volume area of callus, (4) bony tissue per tissue area inside the callus area, (5) fibrous tissue in the callus, and (6) cartilage area in the callus.

Biomechanical analysis

The methods for biomechanical analysis have been previously published by Bak and Andreassen (Calcif Tissue Int (1989) 45:292-297). Briefly, radiographs of all fractures are taken prior to the biomechanical test. The mechanical properties of the healing fractures are analyzed by a destructive three- or four-point bending procedure. Maximum load, stiffness, energy at maximum load, deflection at maximum load, and maximum stress are determined.

20

EXAMPLE 7**FULL-LENGTH CLONING OF ENDOGENOUS HUMAN GPR119**

Polynucleotide encoding endogenous human GPR119 was cloned by PCR using the GPR119 specific primers
 5'-GTCCTGCCACTTCGAGACATGG-3' (SEQ ID NO:3; sense, ATG as initiation codon)
 5'-GAAACTTCTCTGCCCTTACCGTC-3' (SEQ ID NO:4; antisense, 3' of stop codon)
 and human genomic DNA as template. TaqPlus Precision™ DNA polymerase (Stratagene) was used for amplification by the following cycle with step 2 to step 4 repeated 35 times:
 94°C, 3 minutes; 94°C, 1 minute; 58°C, 1 minute; 72°C, 2 minutes; 72°C, 10 minutes.
 A 1.0 Kb PCR fragment of predicted size was isolated and cloned into the pCRII-TOPO™ vector (Invitrogen) and completely sequenced using the T7 DNA sequenase kit (Amersham). See, SEQ ID NO:1 for nucleic acid sequence and SEQ ID NO:2 for the deduced amino acid sequence.

30

EXAMPLE 8**RECEPTOR EXPRESSION**

Although a variety of cells are available to the art for the expression of G protein-coupled receptors, it is most preferred that eukaryotic cells be utilized. In certain embodiments, mammalian cells or melanophores are utilized. The following are illustrative; those of ordinary

35

skill in the art are credited with the ability to determine those techniques that are preferentially beneficial for the needs of the artisan. See, *e.g.*, Example 12, *infra*, as it relates to melanophores.

a. Transient Transfection

5 On day one, 6×10^6 / 10 cm dish of 293 cells are plated out. On day two, two reaction tubes are prepared (the proportions to follow for each tube are per plate): tube A is prepared by mixing 4 μ g DNA (*e.g.*, pCMV vector; pCMV vector with receptor cDNA, etc.) in 0.5 ml serum free DMEM (Gibco BRL); tube B is prepared by mixing 24 μ l lipofectamine (Gibco BRL) in 0.5ml serum free DMEM. Tubes A and B are admixed by inversions (several times), followed
10 by incubation at room temperature for 30-45min. The admixture is referred to as the "transfection mixture". Plated 293 cells are washed with 1XPBS, followed by addition of 5 ml serum free DMEM. 1 ml of the transfection mixture is added to the cells, followed by incubation for 4hrs at 37°C/5% CO₂. The transfection mixture is removed by aspiration, followed by the addition of 10ml of DMEM/10% Fetal Bovine Serum. Cells are incubated at
15 37°C/5% CO₂. After 48hr incubation, cells are harvested and utilized for analysis.

b. Stable Cell Lines

Approximately 12×10^6 293 cells are plated on a 15cm tissue culture plate. Grown in DME High Glucose Medium containing ten percent fetal bovine serum and one percent sodium pyruvate, L-glutamine, and antibiotics. Twenty-four hours following plating of 293 cells (or to
20 ~80% confluency), the cells are transfected using 12 μ g of DNA (*e.g.*, pCMV-neo^r vector with receptor cDNA). The 12 μ g of DNA is combined with 60 μ l of lipofectamine and 2ml of DME High Glucose Medium without serum. The medium is aspirated from the plates and the cells are washed once with medium without serum. The DNA, lipofectamine, and medium mixture are added to the plate along with 10ml of medium without serum. Following incubation at 37°C for
25 four to five hours, the medium is aspirated and 25ml of medium containing serum is added. Twenty-four hours following transfection, the medium is aspirated again, and fresh medium with serum is added. Forty-eight hours following transfection, the medium is aspirated and medium with serum is added containing geneticin (G418 drug) at a final concentration of approximately 12×10^6 293 cells are plated on a 15cm tissue culture plate. Grown in DME High
30 Glucose Medium containing ten percent fetal bovine serum and one percent sodium pyruvate, L-glutamine, and antibiotics. Twenty-four hours following plating of 293 cells (or to ~80% confluency), the cells are transfected using 12 μ g of DNA (*e.g.*, pCMV vector with receptor cDNA). The 12 μ g of DNA is combined with 60 μ l of lipofectamine and 2ml of DME High Glucose Medium without serum. The medium is aspirated from the plates and the cells are
35 washed once with medium without serum. The DNA, lipofectamine, and medium mixture are added to the plate along with 10ml of medium without serum. Following incubation at 37°C for four to five hours, the medium is aspirated and 25ml of medium containing serum is added.

Twenty-four hours following transfection, the medium is aspirated again, and fresh medium with serum is added. Forty-eight hours following transfection, the medium is aspirated and medium with serum is added containing geneticin (G418 drug) at a final concentration of 500µg/ml. The transfected cells now undergo selection for positively transfected cells containing the G418 resistance gene. The medium is replaced every four to five days as selection occurs. During selection, cells are grown to create stable pools, or split for stable clonal selection.

EXAMPLE 9

10 ASSAYS FOR SCREENING CANDIDATE COMPOUNDS AS, E.G., GPR119 AGONISTS

A variety of approaches are available for screening candidate compounds as, e.g., GPR119 agonists. The following are illustrative; those of ordinary skill in the art are credited with the ability to determine those techniques that are preferentially beneficial for the needs of the artisan. Assays for screening compounds as agonists of a G protein-coupled receptor are well known to the skilled artisan (*see*, e.g., International Application WO 02/42461).

1. Membrane Binding Assays: [³⁵S]GTPγS Assay

When a G protein-coupled receptor is in its active state, either as a result of ligand binding or constitutive activation, the receptor couples to a G protein and stimulates the release of GDP and subsequent binding of GTP to the G protein. The alpha subunit of the G protein-receptor complex acts as a GTPase and slowly hydrolyzes the GTP to GDP, at which point the receptor normally is deactivated. Activated receptors continue to exchange GDP for GTP. The non-hydrolyzable GTP analog, [³⁵S]GTPγS, can be utilized to demonstrate enhanced binding of [³⁵S]GTPγS to membranes expressing activated receptors. The advantage of using [³⁵S]GTPγS binding to measure activation is that: (a) it is generically applicable to all G protein-coupled receptors; (b) it is proximal at the membrane surface making it less likely to pick-up molecules which affect the intracellular cascade.

The assay utilizes the ability of G protein coupled receptors to stimulate [³⁵S]GTPγS binding to membranes expressing the relevant receptors. The assay is generic and has application to drug discovery at all G protein-coupled receptors.

30 Membrane Preparation

In some embodiments, membranes comprising a G protein-coupled receptor of the invention and for use in the identification of candidate compounds as, e.g., agonists of the receptor, are preferably prepared as follows:

a. Materials

35 “Membrane Scrape Buffer” is comprised of 20mM HEPES and 10mM EDTA, pH 7.4; “Membrane Wash Buffer” is comprised of 20 mM HEPES and 0.1 mM EDTA, pH 7.4; “Binding Buffer” is comprised of 20mM HEPES, 100 mM NaCl, and 10 mM MgCl₂, pH 7.4.

b. Procedure

All materials will be kept on ice throughout the procedure. Firstly, the media will be aspirated from a confluent monolayer of cells, followed by rinse with 10ml cold PBS, followed by aspiration. Thereafter, 5ml of Membrane Scrape Buffer will be added to scrape cells; this will be followed by transfer of cellular extract into 50ml centrifuge tubes (centrifuged at 20,000 rpm for 17 minutes at 4°C). Thereafter, the supernatant will be aspirated and the pellet will be resuspended in 30ml Membrane Wash Buffer followed by centrifuge at 20,000 rpm for 17 minutes at 4°C. The supernatant will then be aspirated and the pellet resuspended in Binding Buffer. This will then be homogenized using a Brinkman Polytron™ homogenizer (15-20 second bursts until the all material is in suspension). This is referred to herein as "Membrane Protein".

Bradford Protein Assay

Following the homogenization, protein concentration of the membranes will be determined using the Bradford Protein Assay (protein can be diluted to about 1.5mg/ml, aliquoted and frozen (-80°C) for later use; when frozen, protocol for use will be as follows: on the day of the assay, frozen Membrane Protein is thawed at room temperature, followed by vortex and then homogenized with a Polytron at about 12 x 1,000 rpm for about 5-10 seconds; it is noted that for multiple preparations, the homogenizer should be thoroughly cleaned between homogenization of different preparations).

a. Materials

Binding Buffer (as per above); Bradford Dye Reagent; Bradford Protein Standard will be utilized, following manufacturer instructions (Biorad, cat. no. 500-0006).

b. Procedure

Duplicate tubes will be prepared, one including the membrane, and one as a control "blank". Each contained 800µl Binding Buffer. Thereafter, 10µl of Bradford Protein Standard (1mg/ml) will be added to each tube, and 10µl of membrane Protein will then be added to just one tube (not the blank). Thereafter, 200µl of Bradford Dye Reagent will be added to each tube, followed by vortex of each. After five (5) minutes, the tubes will be re-vortexed and the material therein will be transferred to cuvettes. The cuvettes will then be read using a CECIL 3041 spectrophotometer, at wavelength 595.

Identification Assay**a. Materials**

GDP Buffer consists of 37.5 ml Binding Buffer and 2mg GDP (Sigma, cat. no. G-7127), followed by a series of dilutions in Binding Buffer to obtain 0.2 µM GDP (final concentration of GDP in each well was 0.1 µM GDP); each well comprising a candidate compound, has a final volume of 200µl consisting of 100µl GDP Buffer (final concentration, 0.1µM GDP), 50µl

Membrane Protein in Binding Buffer, and 50µl [³⁵S]GTPγS (0.6 nM) in Binding Buffer (2.5 µl [³⁵S]GTPγS per 10ml Binding Buffer).

b. Procedure

Candidate compounds will be preferably screened using a 96-well plate format (these
5 can be frozen at -80°C). Membrane Protein (or membranes with expression vector excluding
the Target GPCR, as control), will be homogenized briefly until in suspension. Protein
concentration will then be determined using the Bradford Protein Assay set forth above.
Membrane Protein (and control) will then be diluted to 0.25mg/ml in Binding Buffer (final assay
concentration, 12.5µg/well). Thereafter, 100 µl GDP Buffer was added to each well of a Wallac
10 Scintistrip™ (Wallac). A 5ul pin-tool will then be used to transfer 5 µl of a candidate compound
into such well (*i.e.*, 5µl in total assay volume of 200 µl is a 1:40 ratio such that the final
screening concentration of the candidate compound is 10µM). Again, to avoid contamination,
after each transfer step the pin tool should be rinsed in three reservoirs comprising water (1X),
ethanol (1X) and water (2X) – excess liquid should be shaken from the tool after each rinse and
15 dried with paper and kimwipes. Thereafter, 50 µl of Membrane Protein will be added to each
well (a control well comprising membranes without the Target GPCR was also utilized), and
pre-incubated for 5-10 minutes at room temperature. Thereafter, 50µl of [³⁵S]GTPγS (0.6 nM)
in Binding Buffer will be added to each well, followed by incubation on a shaker for 60 minutes
at room temperature (again, in this example, plates were covered with foil). The assay will then
20 be stopped by spinning of the plates at 4000 RPM for 15 minutes at 22°C. The plates will then
be aspirated with an 8 channel manifold and sealed with plate covers. The plates will then be
read on a Wallac 1450 using setting “Prot. #37” (as per manufacturer’s instructions).

2. Adenylyl Cyclase Assay

A Flash Plate™ Adenylyl Cyclase kit (New England Nuclear; Cat. No. SMP004A)
25 designed for cell-based assays can be modified for use with crude plasma membranes. The Flash
Plate wells can contain a scintillant coating which also contains a specific antibody recognizing
cAMP. The cAMP generated in the wells can be quantitated by a direct competition for binding
of radioactive cAMP tracer to the cAMP antibody. The following serves as a brief protocol for
the measurement of changes in cAMP levels in cells that express the receptors.

30 In certain embodiments, a modified Flash Plate™ Adenylyl Cyclase kit (New England
Nuclear; Cat. No. SMP004A) is utilized for identification of candidate compounds as, *e.g.*,
GPR119 agonists in accordance with the following protocol.

Cells transfected with a G protein-coupled receptor of the invention are harvested
approximately three days after transfection. Membranes are prepared by homogenization of
35 suspended cells in buffer containing 20mM HEPES, pH 7.4 and 10mM MgCl₂.
Homogenization is performed on ice using a Brinkman Polytron™ for approximately 10
seconds. The resulting homogenate is centrifuged at 49,000 X g for 15 minutes at 4°C. The

resulting pellet is then resuspended in buffer containing 20mM HEPES, pH 7.4 and 0.1 mM EDTA, homogenized for 10 seconds, followed by centrifugation at 49,000 x g for 15 minutes at 4°C. The resulting pellet is then stored at -80°C until utilized. On the day of direct identification screening, the membrane pellet is slowly thawed at room temperature, resuspended in buffer containing 20mM HEPES, pH 7.4 and 10mM MgCl₂, to yield a final protein concentration of 0.60mg/ml (the resuspended membranes are placed on ice until use).

cAMP standards and Detection Buffer (comprising 2 µCi of tracer ([¹²⁵I]cAMP (100 µl) to 11 ml Detection Buffer)) are prepared and maintained in accordance with the manufacturer's instructions. Assay Buffer was prepared fresh for screening and contained 20mM HEPES, pH 7.4, 10mM MgCl₂, 20mM phosphocreatine (Sigma), 0.1 units/ml creatine phosphokinase (Sigma), 50 µM GTP (Sigma), and 0.2 mM ATP (Sigma); Assay Buffer was then stored on ice until utilized.

Candidate compounds are added, preferably, to *e.g.* 96-well plate wells (3µl/well; 12µM final assay concentration), together with 40 µl Membrane Protein (30µg/well) and 50µl of Assay Buffer. This admixture was then incubated for 30 minutes at room temperature, with gentle shaking.

Following the incubation, 100µl of Detection Buffer is added to each well, followed by incubation for 2-24 hours. Plates are then counted in a Wallac MicroBeta™ plate reader using "Prot. #31" (as per manufacturer's instructions).

3. CRE-Luc Reporter Assay

293 and 293T cells are plated-out on 96 well plates at a density of 2 x 10⁴ cells per well and were transfected using Lipofectamine Reagent (BRL) the following day according to manufacturer instructions. A DNA/lipid mixture is prepared for each 6-well transfection as follows: 260ng of plasmid DNA in 100µl of DMEM is gently mixed with 2µl of lipid in 100µl of DMEM (the 260ng of plasmid DNA consists of 200ng of a 8xCRE-Luc reporter plasmid, 50ng of pCMV comprising a G protein-coupled receptor of the invention or pCMV alone, and 10ng of a GPRS expression plasmid (GPRS in pcDNA3 (Invitrogen)). The 8XCRE-Luc reporter plasmid was prepared as follows: vector SRIF-β-gal was obtained by cloning the rat somatostatin promoter (-71/+51) at BglV-HindIII site in the pβgal-Basic Vector (Clontech). Eight (8) copies of cAMP response element were obtained by PCR from an adenovirus template AdpCF126CCRE8 [see, Suzuki et al., Hum Gene Ther (1996) 7:1883-1893; the disclosure of which is herein incorporated by reference in its entirety) and cloned into the SRIF-β-gal vector at the Kpn-BglV site, resulting in the 8xCRE-β-gal reporter vector. The 8xCRE-Luc reporter plasmid was generated by replacing the beta-galactosidase gene in the 8xCRE-β-gal reporter vector with the luciferase gene obtained from the pGL3-basic vector (Promega) at the HindIII-BamHI site. Following 30 min. incubation at room temperature, the DNA/lipid mixture is diluted with 400 µl of DMEM and 100µl of the diluted mixture is added to each well. 100 µl of

DMEM with 10% FCS are added to each well after a 4hr incubation in a cell culture incubator. The following day the transfected cells are changed with 200 μ l/well of DMEM with 10% FCS. Eight (8) hours later, the wells are changed to 100 μ l /well of DMEM without phenol red, after one wash with PBS. Luciferase activity is measured the next day using the LucLite™ reporter gene assay kit (Packard) following manufacturer instructions and read on a 1450 MicroBeta™ scintillation and luminescence counter (Wallac).

10

EXAMPLE 10

WHOLE CELL ADENYLYL CYCLASE ASSAY FOR, E.G., GPR119 AGONIST ACTIVITY

Cyclic AMP measurements are done with a Flash Plate™ Adenylyl Cyclase kit (New England Nuclear) according to the supplier's protocol. HEK293 cells are plated in 15-cm tissue culture dish at 12×10^6 cells per dish in regular growth medium (DMEM/10%FBS). On the next day, 10 μ g of either empty vector DNA or expression plasmid DNA are transfected into cells with lipofectamine (Invitrogen, Carlsbad, CA) according to manufacturer's protocol. After 24 hours in culture, transfected cells are harvested in GIBCO cell dissociation buffer (Cat #13151-014), pelleted by centrifugation for 5 minutes at 1,100 rpm, and carefully re-suspended into an appropriate volume of Assay Buffer (50% 1xPBS and 50% Stimulation Buffer) to give a final cell count at 2×10^6 cells/ml. Test compounds are prepared in 50 μ l Assay Buffer at desired assay concentration where indicated, and pipetted into wells of the 96-well Flash Plate. The cell suspension prepared above was then added (50 μ l per well). After an incubation time of 60 minutes at room temperature, 100 μ l of Detection Mix containing tracer [^{125} I] -cAMP is then added to the wells. Plates are incubated for additional 2 hours followed by counting in a Wallac MicroBeta scintillation counter. Values of cAMP/well are extrapolated from a standard cAMP curve which is included on each assay plate.

An increase in cAMP level in GPR119-transfected HEK293 cells over that in HEK293 cells transfected with empty vector is indicative of a test compound being a compound that stimulates GPR119 receptor functionality.

EXAMPLE 11

HOMOGENEOUS TIME-RESOLVED FLUORESCENCE (HTRF®) ASSAY FOR DIRECT CAMP MEASUREMENT

Compounds are screened for compounds which stimulate GPR119 receptor, e.g. for agonists of GPR119 receptor, using HTRF® assay for direct cAMP measurement (Gabriel et al, ASSAY and Drug Development Technologies, 1:291-303, 2003) and recombinant CHO-K1 cells

stably transfected with GPR119 receptor. Compounds are screened for agonists of GPR119 receptor using HTRF[®] assay for direct cAMP measurement (Gabriel et al, ASSAY and Drug Development Technologies, 1:291-303, 2003) and recombinant CHO-K1 cells stably transfected with GPR119 receptor. CHO-K1 cells are obtained from ATCC[®] (Manassas, VA; Catalog # CCL-5 61). An agonist of GPR119 receptor is detected in HTRF[®] assay for direct cAMP measurement as a compound which increases cAMP concentration. HTRF[®] assay also is used to determine EC₅₀ values for GPR119 receptor agonists.

Principle of the assay: HTRF[®] assay kit is purchased from Cisbio-US, Inc. (Bedford, MA; Catalog # 62AM4PEC). The HTRF[®] assay supported by the kit is a competitive 10 immunoassay between endogenous cAMP produced by the CHO-K1 cells and tracer cAMP labeled with the dye d2. The tracer binding is visualized by a monoclonal anti-cAMP antibody labeled with Cryptate. The specific signal (i.e., fluorescence resonance energy transfer, FRET) is inversely proportional to the concentration of unlabeled cAMP in the standard or sample.

Standard curve: The fluorescence ratio (665 nm/620 nm) of the standards (0.17 to 712 15 nM cAMP) included in the assay is calculated and used to generate a cAMP standard curve according to the kit manufacturer's instructions. The fluorescence ratio of the samples (test compound or compound buffer) is calculated and used to deduce respective cAMP concentrations by reference to the cAMP standard curve.

Setup of the assay: HTRF[®] assay is carried out using a two-step protocol essentially 20 according to the kit manufacturer's instructions, in 20 µl total volume per well in 384-well plate format (ProxiPlates; PerkinElmer, Fremont, CA; catalog # 6008280). To each of the experimental wells is transferred 3000 recombinant CHO-K1 cells in 5 µl assay buffer (phosphate buffered saline containing calcium chloride and magnesium chloride (Invitrogen, Carlsbad, CA; catalog # 14040) supplemented with IBMX (100 µM) and rolipram (10 µM) (phosphodiesterase inhibitors; Sigma- 25 Aldrich, St. Louis, MO; catalog # I5879 and catalog # R6520, respectively) and 0.1% bovine serum albumin (BSA) fraction V (Sigma-Aldrich; catalog # A3059)), followed by test compound in 5 µl assay buffer or 5 µl assay buffer. The plate is then incubated at room temperature for 1 hour. To each well is then added 5 µl cAMP-d2 conjugate in lysis buffer and 5 µl Cryptate conjugate in lysis buffer according to the kit manufacturer's instructions. The plate is then further incubated at room 30 temperature for 1 hour, after which the assay plate is read.

Assay readout: HTRF[®] readout is accomplished using a PHERAstar (BMG LABTECH Inc., Durham, NC) or EnVision[™] (PerkinElmer, Fremont CA) microplate reader.

EXAMPLE 12

35 MELANOPHORE ASSAY FOR, E.G., GPR119 AGONIST ACTIVITY

Melanophores are maintained in culture as reported by Potenza et al [Pigment Cell Research (1992) 5:372-378] and transfected with an expression vector encoding a GPR119

receptor (GPR119; *e.g.*, human GPR119, GenBank® Accession No. AAP72125 and alleles thereof) using electroporation. Following electroporation, the transfected cells are plated into 96 well plates for the assay. The cells are then allowed to grow for 48 hours in order to both recover from the electroporation procedure and attain maximal receptor expression levels.

5 On the assay day, the growth medium on the cells is replaced with serum-free buffer containing 10nM melatonin. The melatonin acts via an endogenous Gi-coupled GPCR in the melanophores to lower intracellular cAMP levels. In response to lowered cAMP levels, the melanophores translocate their pigment to the center of the cell. The net effect of this is a significant decrease in the absorbance reading of the cell monolayer in the well, measured at
10 600-650nm.

 After a 1-hour incubation in melatonin, the cells become completely pigment-aggregated. At this point a baseline absorbance reading is collected. Serial dilutions of test compounds are then added to the plate, and compounds having GPR119 agonist activity produce increases in intracellular cAMP levels. In response to these increased cAMP levels, the
15 melanophores translocate their pigment back into the cell periphery. After one hour, stimulated cells are fully pigment-dispersed. The cell monolayer in the dispersed state absorbs much more light in the 600-650nm range. The measured increase in absorbance compared to the baseline reading allows one to quantitate the degree of receptor stimulation and plot a dose-response curve.

20 Materials and methods relating to melanophore assay are found in U.S. Pat. Nos. 5,462,856 and 6,051,386, the disclosure of each of which is herein incorporated by reference in its entirety.

 An increase in pigment dispersion in GPR119-transfected melanophores over over that in melanophores transfected with empty vector is indicative of a test compound being a
25 compound that stimulates GPR119 receptor functionality.

 Other assays for identifying a compound as a GPR119 agonist will be readily apparent to the skilled artisan (see, *e.g.*, Example 9, *supra*).

EXAMPLE 13

30 YEAST REPORTER ASSAY FOR, E.G., GPR119 AGONIST ACTIVITY

 The yeast cell-based reporter assays have previously been described in the literature (*e.g.*, see Miret et al, J Biol Chem (2002) 277:6881-6887; Campbell et al, Bioorg Med Chem Lett (1999) 9:2413-2418; King et al, Science (1990) 250:121-123; WO 99/14344; WO 00/12704; and US 6,100,042). Briefly, yeast cells have been engineered such that the endogenous yeast G-alpha
35 (GPA1) has been deleted and replaced with G-protein chimeras constructed using multiple techniques. Additionally, the endogenous yeast alpha-cell GPCR, Ste3 has been deleted to allow for a homologous expression of a mammalian GPCR of choice. In the yeast, elements of the

pheromone signaling transduction pathway, which are conserved in eukaryotic cells (for example, the mitogen-activated protein kinase pathway), drive the expression of Fus1. By placing β -galactosidase (LacZ) under the control of the Fus1 promoter (Fus1p), a system has been developed whereby receptor activation leads to an enzymatic readout.

5 Yeast cells are transformed by an adaptation of the lithium acetate method described by Agatep et al (Agatep et al, 1998, Transformation of *Saccharomyces cerevisiae* by the lithium acetate/single-stranded carrier DNA/polyethylene glycol (LiAc/ss-DNA/PEG) protocol. Technical Tips Online, Trends Journals, Elsevier). Briefly, yeast cells are grown overnight on yeast tryptone plates (YT). Carrier single-stranded DNA (10 μ g), 2 μ g of each of two Fus1p-LacZ reporter
10 plasmids (one with URA selection marker and one with TRP), 2 μ g of GPR119 (e.g., human receptor) in yeast expression vector (2 μ g origin of replication) and a lithium acetate/ polyethylene glycol/ TE buffer is pipetted into an Eppendorf tube. The yeast expression plasmid containing the receptor/ no receptor control has a LEU marker. Yeast cells are inoculated into this mixture and the reaction proceeds at 30°C for 60min. The yeast cells are then heat-shocked at 42°C for 15min. The
15 cells are then washed and spread on selection plates. The selection plates are synthetic defined yeast media minus LEU, URA and TRP (SD-LUT). After incubating at 30°C for 2-3 days, colonies that grow on the selection plates are then tested in the LacZ assay.

In order to perform fluorimetric enzyme assays for β -galactosidase, yeast cells carrying the subject GPR119 receptor are grown overnight in liquid SD-LUT medium to an unsaturated
20 concentration (i.e. the cells are still dividing and have not yet reached stationary phase). They are diluted in fresh medium to an optimal assay concentration and 90 μ l of yeast cells are added to 96-well black polystyrene plates (Costar). Test compounds, dissolved in DMSO and diluted in a 10% DMSO solution to 10X concentration, are added to the plates and the plates placed at 30°C for 4h. After 4h, the substrate for the β -galactosidase is added to each well. In these experiments,
25 Fluorescein di (β -D-galactopyranoside) is used (FDG), a substrate for the enzyme that releases fluorescein, allowing a fluorimetric read-out. 20 μ l per well of 500 μ M FDG/2.5% Triton X100 is added (the detergent is necessary to render the cells permeable). After incubation of the cells with the substrate for 60min, 20 μ l per well of 1M sodium carbonate is added to terminate the reaction and enhance the fluorescent signal. The plates are then read in a fluorimeter at 485/535nm.

30 An increase in fluorescent signal in GPR119-transformed yeast cells over that in yeast cells transformed with empty vector is indicative of a test compound being a compound that stimulates GPR119 receptor functionality (e.g., a compound that is an agonist or partial agonist of GPR119). In certain embodiments, compounds of the invention give an increase in fluorescent signal above that of the background signal (the signal obtained in the presence of vehicle alone).

35

EXAMPLE 14

RADIOLABELED COMPOUND

In certain embodiments, a compound known to be a ligand of a G protein-coupled receptor of the invention is radiolabeled. A radiolabeled compound as described herein can be used in a screening assay to identify/evaluate compounds. In general terms, a newly synthesized or identified compound (i.e., test compound) can be evaluated for its ability to reduce binding of the radiolabeled known ligand to the receptor, by its ability to reduce formation of the complex between the radiolabeled known ligand and the receptor. Suitable radionuclides that may be incorporated in compounds of the present invention include but are not limited to ^3H (also written as T), ^{11}C , ^{14}C , ^{18}F , ^{125}I , ^{82}Br , ^{123}I , ^{124}I , ^{125}I , ^{131}I , ^{75}Br , ^{76}Br , ^{15}O , ^{13}N , ^{35}S and ^{77}Br . Compounds that incorporate ^3H , ^{14}C , ^{125}I , ^{131}I , ^{35}S or ^{82}Br will generally be most useful.

It is understood that a "radiolabelled" compound" is a compound that has incorporated at least one radionuclide. In some embodiments, the radionuclide is selected from the group consisting of ^3H , ^{14}C , ^{125}I , ^{35}S and ^{82}Br . In some embodiments, the radionuclide ^3H or ^{14}C . Moreover, it should be understood that all of the atoms represented in the compounds known to be ligands of a G protein-coupled receptor of the invention can be either the most commonly occurring isotope of such atoms or the more scarce radioisotope or nonradioactive isotope.

Synthetic methods for incorporating radioisotopes into organic compounds including those applicable to those compounds known to be ligands of a G protein-coupled receptor of the invention are well known in the art and include incorporating activity levels of tritium into target molecules include: A. Catalytic Reduction with Tritium Gas - This procedure normally yields high specific activity products and requires halogenated or unsaturated precursors. B. Reduction with Sodium Borohydride [^3H] - This procedure is rather inexpensive and requires precursors containing reducible functional groups such as aldehydes, ketones, lactones, esters, and the like. C. Reduction with Lithium Aluminum Hydride [^3H] - This procedure offers products at almost theoretical specific activities. It also requires precursors containing reducible functional groups such as aldehydes, ketones, lactones, esters, and the like. D. Tritium Gas Exposure Labeling - This procedure involves exposing precursors containing exchangeable protons to tritium gas in the presence of a suitable catalyst. E. N-Methylation using Methyl Iodide [^3H] - This procedure is usually employed to prepare O-methyl or N-methyl (^3H) products by treating appropriate precursors with high specific activity methyl iodide (^3H). This method in general allows for high specific activity, such as about 80-87 Ci/mmol.

Synthetic methods for incorporating activity levels of ^{125}I into target molecules include: A. Sandmeyer and like reactions - This procedure transforms an aryl or heteroaryl amine into a diazonium salt, such as a tetrafluoroborate salt, and subsequently to ^{125}I labelled compound using Na^{125}I . A represented procedure was reported by Zhu, D.-G. and co-workers in *J. Org. Chem.* **2002**, *67*, 943-948. B. Ortho ^{125}I iodination of phenols - This procedure allows for the

incorporation of ^{125}I at the ortho position of a phenol as reported by Collier, T. L. and co-workers in *J. Labelled Compd Radiopharm.* **1999**, 42, S264-S266. C. Aryl and heteroaryl bromide exchange with ^{125}I – This method is generally a two step process. The first step is the conversion of the aryl or heteroaryl bromide to the corresponding tri-alkyltin intermediate using
5 for example, a Pd catalyzed reaction (i.e. $\text{Pd}(\text{Ph}_3\text{P})_4$) or through an aryl or heteroaryl lithium, in the presence of a tri-alkyltinhalide or hexaalkylditin [e.g., $(\text{CH}_3)_3\text{SnSn}(\text{CH}_3)_3$]. A represented procedure was reported by Bas, M.-D. and co-workers in *J. Labelled Compd Radiopharm.* **2001**, 44, S280-S282.

The foregoing techniques are intended to be illustrative and not limiting. Other
10 techniques for radiolabeling a compound known to be a ligand of a G protein-coupled receptor of the invention are well known to the skilled artisan.

15

EXAMPLE 15

RECEPTOR BINDING ASSAY

A test compound can be evaluated for its ability to reduce formation of the complex between a compound known to be a ligand of a G protein-coupled receptor of the invention and the receptor. In certain embodiments, the known ligand is radiolabeled. The radiolabeled
20 known ligand can be used in a screening assay to identify/evaluate compounds. In general terms, a newly synthesized or identified compound (i.e., test compound) can be evaluated for its ability to reduce binding of the radiolabeled known ligand to the receptor, by its ability to reduce formation of the complex between the radiolabeled known ligand and the receptor.

A level of specific binding of the radiolabeled known ligand in the presence of the test
25 compound less than a level of specific binding of the radiolabeled known ligand in the absence of the test compound is indicative of less of the complex between said radiolabeled known ligand and said receptor being formed in the presence of the test compound than in the absence of the test compound.

Assay Protocol for Detecting the Complex Between a Compound Known to be a Ligand of a 30 G Protein-Coupled Receptor of the Invention and the Receptor

A. Preparation of the Receptor

293 cells are transiently transfected with 10 ug expression vector comprising a polynucleotide encoding a G protein-coupled receptor of the invention using 60 ul Lipofectamine (per 15-cm dish). The transiently transfected cells are grown in the dish for 24
35 hours (75% confluency) with a media change and removed with 10 ml/dish of Hepes-EDTA buffer (20mM Hepes + 10mM EDTA, pH 7.4). The cells are then centrifuged in a Beckman Coulter centrifuge for 20 minutes, 17,000 rpm (JA-25.50 rotor). Subsequently, the pellet is

resuspended in 20mM Hepes + 1mM EDTA, pH 7.4 and homogenized with a 50-ml Dounce homogenizer and again centrifuged. After removing the supernatant, the pellets are stored at -80°C, until used in binding assay. When used in the assay, membranes are thawed on ice for 20 minutes and then 10 mL of incubation buffer (20 mM Hepes, 1mM MgCl₂, 100mM NaCl, pH 7.4) added. The membranes are then vortexed to resuspend the crude membrane pellet and homogenized with a Brinkmann PT-3100 Polytron homogenizer for 15 seconds at setting 6. The concentration of membrane protein is determined using the BRL Bradford protein assay.

B. Binding Assay

For total binding, a total volume of 50ul of appropriately diluted membranes (diluted in assay buffer containing 50mM Tris HCl (pH 7.4), 10mM MgCl₂, and 1mM EDTA; 5-50ug protein) is added to 96-well polypropylene microtiter plates followed by addition of 100ul of assay buffer and 50ul of a radiolabeled known ligand. For nonspecific binding, 50ul of assay buffer is added instead of 100ul and an additional 50ul of 10uM said known ligand which is not radiolabeled is added before 50ul of said radiolabeled known ligand is added. Plates are then incubated at room temperature for 60-120 minutes. The binding reaction is terminated by filtering assay plates through a Microplate Devices GF/C Unifilter filtration plate with a Brandell 96-well plate harvester followed by washing with cold 50mM Tris HCl, pH 7.4 containing 0.9% NaCl. Then, the bottom of the filtration plate are sealed, 50ul of Optiphas Supermix is added to each well, the top of the plates are sealed, and plates are counted in a Trilux MicroBeta scintillation counter. For determining whether less of the complex between said radiolabeled known ligand and said receptor is formed in the presence of a test compound, instead of adding 100ul of assay buffer, 100ul of appropriately diluted said test compound is added to appropriate wells followed by addition of 50ul of said radiolabeled known ligand.

C. Calculations

The test compounds are initially assayed at 10, 1 and 0.1 μ M and then at a range of concentrations chosen such that the middle dose would cause about 50% inhibition of binding of the radiolabeled known ligand (i.e., IC₅₀). Specific binding in the absence of test compound (B₀) is the difference of total binding (B_T) minus non-specific binding (NSB) and similarly specific binding (in the presence of test compound) (B) is the difference of displacement binding (B_D) minus non-specific binding (NSB). IC₅₀ is determined from an inhibition response curve, logit-log plot of % B/B₀ vs concentration of test compound.

K_i is calculated by the Cheng and Prustoff transformation:

$$K_i = IC_{50} / (1 + [L]/K_D)$$

where [L] is the concentration of radiolabeled known ligand used in the assay and K_D is the dissociation constant of the radiolabeled known ligand determined independently under the same binding conditions.

EXAMPLE 16**EFFECT OF GPR119 AGONIST ON PYY SECRETION IN ENTEROENDOCRINE CELL OR IN CELLS IN TISSUE DERIVED FROM AN L CELL-CONTAINING REGION OF INTESTINE OR IN CELLS CAPABLE OF SECRETING PYY**

5 A GPR119 agonist in accordance with the present invention can be shown to stimulate PYY secretion in an enteroendocrine cell, e.g. an enteroendocrine cell line, or in cells such as enteroendocrine cells in tissue derived from an L cell-containing region of intestine [e.g., duodenum, jejunum, ileum, colon or rectum tissue; *see*, e.g., Lundberg et al, PNAS USA (1982) 79:4471-4475; Adrian et al, Gastroenterol (1985) 89:1070-1077; Ekblad et al, Peptides (2002) 10 23:251-261; Ueno et al, Regul Pept (2008) 145:12-16]] or in cells capable of secreting PYY using the *in vitro* assay described here. Enteroendocrine cells or cells such as enteroendocrine cells in tissue derived from an L cell-containing region of intestine are plated in 24-well plates on day one in complete culture medium (DMEM/10%FBS). On day two the culture medium is replaced with a low glucose medium (DMEM/3mM Glucose/10%FBS). On day three cells are 15 washed twice with 1xPBS. The washed cells are stimulated with vehicle or with GPR119 agonist at various concentrations (*e.g.*, in the range of 1nM to 20uM) or with forskolin (1uM) as a positive control in serum free DMEM with 15mM glucose for one hour at 37°C and 5%CO₂ in a tissue culture incubator. The supernatants are then collected and clarified by centrifugation at 500g and 4°C for 5 minutes. Total PYY released into the supernatant is determined using a 20 PYY EIA kit purchased from Phoenix Pharmaceuticals, Inc. (Peptide YY (PYY) (Rat, Mouse, Porcine, Canine) EIA kit, Catalog No. EK-059-03) according to instructions provided in the kit.

It is expressly contemplated that cells capable of secreting PYY include but are not limited to an enteroendocrine cell, as may be an enteroendocrine cell line, a cell such as may be an enteroendocrine cell in tissue derived from an L cell-containing region of intestine, an alpha cell of 25 the islets of Langerhans, a cell line derived from the islets of Langerhans such as may be RINm5F (Bargsten, Arch Histol Cytol (2004) 67:79-94), a cell such as a neuronal cell as may be in tissue from the medulla oblongata, a neuronal cell line, and a recombinant cell.

EXAMPLE 17**30 *IN VIVO* EFFECTS OF A GPR119 AGONIST ON ATHEROGENESIS IN MICE**

Male apoE^{-/-} mice (Jackson Laboratory, Bar Harbor, ME) are fed a normal chow. The apolipoprotein E-deficient (apoE^{-/-}) mouse is an established animal model of atherosclerosis, developing extensive atherosclerotic lesions on a chow diet [Zhang et al, Science (1992) 258:468-471]. At the age of 12 weeks, a GPR119 agonist or vehicle alone is administered per 35 daily oral administration. A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

The mice are anesthetized with an intraperitoneal injection of pentobarbital (50 mg/kg), and the hearts, which contain the aortic sinus and aortic arch, are harvested 14 days after the administration of the GPR119 agonist. Hearts are also harvested from unmanipulated mice at the start of the experiment. Ten mice are used for each of the three experimental groups.

5 The frozen cross-sections (10 μ M thick) of aortic sinus embedded in Optimal Cutting Temperature (OCT; Sakura Finetechnical Co., Ltd) compound after overnight fixation in 10% formalin are mounted on slides. For the analysis of plaque size, 3 sections (100 μ M apart) from each mouse were stained with Oil Red O. The lesion size and the diameter of lipid droplets in the lesions are quantified with an image analyzing computer software, and the mean values are
10 determined. The mean value for the group administered with the GPR119 agonist is compared with the mean value for the group administered with vehicle alone.

Data are presented as means $\pm$ SEM and are analyzed by Student's t test or the Mann-Whitney U test, depending on their distribution pattern. A value of $P < 0.05$ is considered statistically significant. *See, e.g., Okamoto et al, Circulation (2002) 106:2767-2770.*

15 These results can demonstrate that the GPR119 agonist is an inhibitor of atherogenesis.

It is expressly contemplated that in other embodiment a GPR119 agonist can be shown to be an inhibitor of atherogenesis in the ApoE^{-/-} mice using a non-invasive *in vivo* technique [see, e.g., Fayad et al, Circulation (1998) 98:1541-1547]. It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that
20 administration of the agonist is intraperitoneal or intravenous. It is expressly contemplated that in other embodiment, treatment begins other than at 12 weeks of age, either earlier or later than at 12 weeks of age. It is expressly contemplated that in other embodiment treatment continues for less than or more than 14 days. It is expressly contemplated that administration may be other than daily.

25

EXAMPLE 18

***IN VIVO* EFFECTS OF A GPR119 AGONIST ON CHRONIC INFLAMMATORY ARTHRITIS IN MICE**

Both male and female MRL-*lpr* mice (Jackson Laboratory, Bar Harbor, ME) are used at 13-14 weeks of age. MRL-*lpr* mice spontaneously develop a chronic inflammatory arthritis with similar characteristics to human rheumatoid arthritis including cell infiltration, pannus
30 formation, bone and cartilage breakdown, and the presence of serum rheumatoid factor. The disease normally develops towards the end of the animal's life span; however, injection with complete Freund's adjuvant (CFA) initiates early onset and increases the severity of arthritis [Rakay et al, J Immunol (1993) 151:5081-5087].

35 On Day 0 of each experiment, all groups of mice are injected with CFA intradermally into a thoracic and an inguinal site with 0.05 ml CFA supplemented to 10 mg/ml with heat inactivated *Mycobacterium tuberculosis* H37 Ra (Difco, Detroit, MI). Immediately, a GPR119

agonist or vehicle alone is administered per daily oral administration. A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

5 Treatment is continued for 30 days. For quantifying swelling, ankle widths are measured with a micrometer. The statistical comparison of paired sets of ankle width measurements is carried out using the Student's *t* test.

Histopathological Analysis

At day 30 after CFA priming, the hind paws are fixed in buffered formalin. After decalcification in 10% formic acid for 48 hours, the tissues are processed for paraffin
10 embedding. Serial sections of the tarso-metatarsal joints are cut to a thickness of 5 mm and stained with hematoxylin and eosin. Sections are examined by an individual without knowledge of the experimental protocol. A minimum of 10 sections/joint are assessed and scored to provide a semiquantitative measure of subsynovial inflammation (0, normal; 1 focal inflammatory infiltrates; 2, inflammatory infiltrate that dominates the cellular histology),
15 synovial hyperplasia (0, normal; 1, a continuous, minimum three-layer thick, synovial lining seen in one joint; 2, minimum three-layer thick, synovial lining detected in several joints), pannus formation and cartilage erosion (0, normal; 1, pannus partially covered cartilage surfaces without evident cartilage loss; 2, pannus connected to evident cartilage loss), bone destruction (0, normal; 1, detectable destruction of bone by the pannus or osteoclast activity; 2, the pannus
20 or osteoclast activity has destroyed a significant part of the bone), and finally, overall pathology is the overall assessment derived by the summation of the values for these criteria [see, e.g., Gong et al, J Exp Med (1997) 186:131-137]. Statistical analysis of the histopathology indices is done using the Student's *t* test.

These results can demonstrate that the GPR119 agonist is an inhibitor of a chronic
25 inflammatory arthritis, for example that the GPR119 agonist is an inhibitor of rheumatoid arthritis.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is intraperitoneal or intravenous. It is expressly contemplated that in other embodiment, treatment begins other than at 13-14
30 weeks of age, either earlier or later than at 13-14 weeks of age. It is expressly contemplated that in other embodiment treatment continues for less than or more than 30 days. It is expressly contemplated that administration may be other than daily.

EXAMPLE 19

35 REDUCTION OF COLON INJURY OF ANIMAL MODEL FOR INFLAMMATORY BOWEL DISEASE

A GPR119 agonist can be shown to confer protection from inflammatory bowel disease using the animal model essentially as described in WO 03/105763.

Methods

Animal preparation:

10-11 weeks old male HSD rats, ranging 250-300 grams, are housed in a 12:12 light:dark cycle, and allowed ad libitum access to a standard rodent diet (Taklad L M 485, Madison, WI) and
5 water. The animals are fasted for 24 hours before the experiment.

Procedure:

A simple and reproducible rat model of chronic colonic inflammation has been previously described by Morris et al (Gastroenterology (1989) 96:795-803). It exhibits a relatively long duration of inflammation and ulceration, affording an opportunity to study the pathophysiology
10 of colonic inflammatory disease in a specifically controlled fashion, and to evaluate new treatments potentially applicable to inflammatory bowel disease in humans.

Rats are anesthetized with 3% isoflurane and placed on a regulated heating pad set at 37°C. A gavage needle is inserted rectally into the colon 7 cm. The hapten trinitrobenzenesulfonic acid (TNBS) dissolved in 50% ethanol (v/v) is delivered into the lumen
15 of the colon through the gavage needle at a dose of 30 mg/kg, in a total volume of 0.4-0.6 mL. Control groups receive saline solution (NaCl 0.9%) intracolonicly.

Four days after induction of colitis, the colon is resected from anesthetized rats, which are then euthanized by decapitation. Weights of excised colon and spleen are measured, and the colon photographed for scoring of gross morphologic damage according to the a rating scale (0
20 = no damage, 1 = localized hyperemia but no ulcers, 2 = linear ulcers with no significant inflammation, 3 = linear ulcer with inflammation at one site, 4 = two or more sites of ulceration and/or inflammation, 5 = two or more major sites of inflammation and ulceration or one major site of inflammation and ulceration extending >1 cm along the length of the colon) before division into 3 parts. The segment of colon 1 cm from the cecum is fixed in 10% formalin.
25 Other segments are immediately frozen on dry ice.

Inflammation is defined as regions of hyperemia and bowel wall thickening. Colons are observed independently by several observers blinded to experimental treatment.

For this assay, 5 treatment groups are set up as follows:

Group A (n=3). intracolonic saline with oral treatment of saline;
30 **Group B** (n=8). oral treatment of saline prior to induction of colitis by 30 mg/kg TNBS;
Group C (n=10). Experimental group—oral treatment of GPR119 agonist (at a dose of, e.g., 30 mg/kg) twice daily for 2 days before, 2 days after, and 30 min before induction of colitis by 30 mg/kg TNBS.

Group D (n=9). Positive control—intramuscular injection of Depo-Medrol 600 µg/kg
35 60 min prior to induction of colitis by 30 mg/kg TNBS;

Group E (n=10). Positive control, 5-amino salicylic acid (Sigma) suspension at 200 mg/kg intracolonic administration in 0.5 mL saline, 60 min before induction of colitis and daily thereafter.

A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is subcutaneous or intravenous.

10

EXAMPLE 20

EFFECT OF GPR119 AGONIST ON KAINATE SEIZURES IN MICE

A GPR119 agonist can be shown to confer protection from kainate seizures in mice using the animal model essentially as described in Woldbye et al (Neurobiology of Disease (2005) 20:760-772).

15

Male NMRI mice (Taconic M&B, DK; 23-30 g) are administered per orally with GPR119 agonist at doses of, e.g., 0.2, 1, 10, 20 or 30 mg/kg (n = 7-8) in vehicle containing 0.05% bovine serum albumin in 10 mM phosphate-buffered saline (PBS, 0.13 M NaCl, 7 mM Na₂HPO₄, 3 mM NaH₂PO₄). A control group receives only vehicle by oral administration (n = 28). Five minutes later, all animals receive a subcutaneous injection of kainate dissolved in isotonic saline and adjusted to pH 7.4 (30 mg/kg; #2020, Ocean Produce International, Canada); seizure severity is scored according to a modified rating scale of Marsh et al (PNAS USA (1999) 96:13518-13523): 0 = no seizure activity, 1 = staring or facial movements, 2 = head nodding or isolated twitches, 3 = motor seizure with forelimb clonus, 4 = motor seizure with rearing, 5 = motor seizure with loss of posture or status epilepticus (at least 10 min of continuous motor seizure activity), 6 = death. The seizure score and latencies to first motor seizure (when scored as at least 3) and motor seizure with loss of posture are determined for each treatment group by an observed blinded to the identity of the mice.

25

30

A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is intraperitoneal or intravenous. It is expressly contemplated that the subcutaneous injection of kainate may alternatively be carried out at a time after administration of the GPR119 agonist which is greater than 5 minutes.

35

EXAMPLE 21**EFFECT OF GPR119 AGONIST ON FOOD INTAKE IN LEAN MICE**

A GPR119 agonist can be shown to decrease food intake using lean fasted-refed mice essentially as described in Ortiz et al (JPET (2007) 323:692-700).

5 Lean C57BL/6 mice (Taconic Farms, Germantown, NY) are acclimated for a minimum of a week with controlled temperature and humidity on a 12-h light/dark cycle. Mice are housed in pairs in cages with a grid floor with food and water (standard chow pellet diet) continuously available. A study includes 20 mice per treatment group. The mice are fasted overnight (18 h) with water available during the dark phase, and they are dosed orally with GPR119 agonist (at a
10 dose of, e.g., 30 mg/kg) in USP saline or only vehicle. Prew weighed food is provided 30 min after dosing, and cumulative food intake is measured over 72 h (e.g., at 1 h, at 2h, at 4 h, at 24 h, at 48 h, and at 72 h).

A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30
15 mg/kg and 100 mg/kg.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is intraperitoneal or intravenous.

EXAMPLE 22**EFFECT OF GPR119 AGONIST ON BODY WEIGHT AND PLASMA ADIPONECTIN IN DIET-
20 INDUCED OBESE (DIO) MICE**

A GPR119 agonist can be shown to provide a reduction in body weight and an increase in plasma adiponectin using the diet-induced obese (DIO) mouse essentially as described in Ortiz et al (JPET (2007) 323:692-700).

Male C57BL/6 mice (Taconic Farms) are fed a high-fat diet containing 45% calories
25 from fat (rodent diet D12451; Research Diets, Inc., New Brunswick, NJ) for 20 to 22 weeks, and they have an average body weight over about 5 SDs greater than mice fed a standard chow pellet diet (5% calories from fat). A study includes 10 mice per treatment group. Mice are kept in standard animal rooms under controlled temperature and humidity and a 12/12-h light dark cycle. Water and food are continuously available. Animals are adapted to the grid floors for 4
30 days, and they are dosed with vehicle (USP saline) for a further 4 days before recording of 2 days of baseline body weight and 24-h food and water consumption. Mice are assigned to treatment groups based upon their body weight such that the initial mean and standard error of the mean of body weight is similar. Animals are administered GPR119 agonist (at a dose of, e.g., 30 mg/kg) in USP saline or only vehicle orally every day before the dark phase, and body weight
35 and food and water consumption are measured. Animals are dosed daily for 40 days.

Blood is collected on day 40 by retro-orbital bleeding for adiponectin analysis; plasma adiponectin is measured using a Mouse Adiponectin ELISA kit (Catalog No. EZMADP-60K; Millipore Corp., Billerica, MA) according to manufacturer's instructions.

A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is
5 selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is subcutaneous or intravenous.

10

EXAMPLE 23

EFFECT OF GPR119 AGONIST ON PLASMA GLUCOSE, PLASMA INSULIN, AND GLUCOSE DISPOSAL IN DIET-INDUCED OBESE (DIO) MICE

A GPR119 agonist can be shown to provide favorable improvements in metabolic parameters of diet-induced obese (DIO) mice associated with risk for type 2 diabetes, such as
15 glucose disposal (to increase glucose disposal in fasted mice, e.g. glucose disposal following an OGTT), plasma glucose level (to decrease blood glucose in fed mice), and plasma insulin level (to increase plasma insulin in fasted mice), essentially as described in Ortiz et al (JPET (2007) 323:692-700).

Male C57BL/6 mice (Taconic Farms) are fed a high-fat diet containing 45% calories
20 from fat (rodent diet D12451; Research Diets, Inc., New Brunswick, NJ) for 20 to 22 weeks, and they have an average body weight over 5 SDs greater than mice fed a standard chow pellet diet (5% calories from fat). A study includes 10 mice per treatment group. Mice are kept in standard animal rooms under controlled temperature and humidity and a 12/12-h light dark cycle. Water and food are continuously available. Animals are adapted to the grid floors for 4
25 days, and they are dosed with vehicle (USP saline) for a further 4 days before recording of 2 days of baseline body weight and 24-h food and water consumption. Mice are assigned to treatment groups based upon their body weight such that the initial mean and standard error of the mean of body weight is similar.

Animals are orally administered GPR119 agonist (at a dose of, e.g., 30 mg/kg) in USP
30 saline or only vehicle every day before the dark phase. Animals are dosed daily for 11 days. Blood samples are collected from the tail of fed mice and analyzed with an Ascensia glucose meter (Bayer Healthcare, Tarrytown, NY) on day 10 of treatment. Food is removed 5 h into the feeding phase, and the animals are fasted for 14 h before blood collection using retro-orbital bleeding for insulin analysis. Oral glucose tolerance test (OGTT) is performed 2 h after retro-
35 orbital bleeding following oral administration of 2 g/kg glucose, and tail blood glucose is measured at 30-min intervals for 3 h. Insulin is measured by enzyme-linked immunosorbent assay (EIA) (Alpco Diagnostics, Windham, NH).

A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is subcutaneous or intravenous. It is expressly contemplated in other embodiment, that intraperitoneal glucose tolerance test (IPGTT) is performed 2 h after retro-orbital bleeding following intraperitoneal administration of 2 g/kg glucose.

EXAMPLE 24

EFFECT OF GPR119 AGONIST ON COLLATERAL-DEPENDENT BLOOD FLOW IN A RAT MODEL OF PERIPHERAL ARTERIAL INSUFFICIENCY

A GPR119 agonist can be shown to increase collateral-dependent blood flow using a rat model of acute onset peripheral arterial insufficiency that is characteristic of the large vessel occlusive disease that often presents with symptoms of intermittent claudication, essentially as described in Cruze et al (Peptides (2007) 28:269-280).

Peripheral arterial insufficiency is established in adult male Sprague Dawley rats (325-350 g; Taconic Farms, Inc.) with bilateral occlusion of the femoral arteries distal to the inguinal ligament. In the same surgical procedure, a catheter connected to a 14 days osmotic pump (Alzet[®]) is inserted into the iliac artery for infusion near the site of collateral expansion for systemic delivery of vascular growth factor-A (residues 1-165) (VEGF₁₆₅) at a dose of 15 µg/kg/day; GPR119 agonist (at a dose of, e.g., 30 mg/kg) and vehicle alone (phosphate buffered saline, 10% sodium citrate, 1.6% glycerol, 0.008% Tween 20) are administered per daily oral administration. All agents are supplied and administered in a manner blinded to the investigators. Blood flow is measured on day 16 to eliminate any acute effects of GPR119 agonist or VEGF₁₆₅ on blood flow. Muscle blood flow is determined in a blinded manner, utilizing radiolabeled microspheres during treadmill running, as described in Yang et al (Am J Physiol Heart Circ Phys (2000) 278:H1966-H1973). The dose of VEGF₁₆₅ used (15 µg/kg/day) yields a near-maximal increase in collateral-dependent blood flow in this model (Cruze et al, Peptides (2007) 28:269-280).

A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is by intravenous infusion at a dose of, e.g., 30 mg/kg/day. A preferred infusion dose of the GPR119 agonist is 0.1-100 mg/kg/day. Other preferred infusion dose is selected from the group consisting of 0.1

mg/kg/day, 0.3 mg/kg/day, 1.0 mg/kg/day, 3.0 mg/kg/day, 10 mg/kg/day, 30 mg/kg/day and 100 mg/kg/day.

EXAMPLE 25

5 EFFECT OF GPR119 AGONIST ON VASOACTIVE INTESTINAL PEPTIDE (VIP)-INDUCED JEJUNAL SECRETIONS IN *IN VIVO* RAT MODEL

A GPR119 agonist can be shown to decrease secretion in intestinal epithelium using an *in vivo* rat model of vasoactive intestinal peptide (VIP)-induced jejunal secretions, essentially as described by Balasubramaniam et al (J Med Chem (2000) 43:3420-3427) and Souli et al
10 (Peptides (1997) 18:551-557). Balasubramaniam et al (J Med Chem (2000) 43:3420-3427) also describe analogues of PYY having activity in this assay.

Male Wistar rats (Charles River Laboratories, Inc., Wilmington, MA) are anesthetized, and a saphenous vein is cannulated to infuse drugs. The abdomen is opened, a jejunal loop (~20 cm long) is delimited by two ligatures. At time zero, 2 mL of 0.9% saline prewarmed to
15 37°C is instilled into the jejunal loop. The loop is then returned into the abdomen and closed. VIP (30 µg/kg/hour) is infused through the saphenous vein at the rate of 2.5 mL/hour for 30 minutes starting at time zero. GPR119 agonist (at a dose of, e.g., 30 mg/kg) or saline (controls) are orally administered as bolus doses 15 min before starting VIP infusions. At the end of the experiment rats are sacrificed, and the jejunal loop excised. The loop is then measured and
20 weighed before and after removing the instilled fluid. The net water flux expressed as µL/cm/30 min is calculated using the formula $(F-E-2000)/L$, where F and E are the weights (mg) of the loop before and after emptying the remaining fluid, L is the length of the loop in cm, and 2000 is the volume in µL of the fluid instilled initially. Net absorption is indicated by a negative value and net secretion by a positive value. Each dose is investigated in 6-8 animals.

25 A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is subcutaneous or intravenous.

30

EXAMPLE 26

EFFECT OF GPR119 AGONIST ON INTESTINAL ABSORPTION IN *IN VIVO* DOG MODEL

A GPR119 agonist can be shown to increase absorption in intestinal epithelium using an *in vivo* dog model, essentially as described by Balasubramaniam et al (J Med Chem (2000) 43:3420-3427). Balasubramaniam et al (J Med Chem (2000) 43:3420-3427) also describe
35 analogues of PYY having activity in this assay.

In vivo effects of GPR119 agonist on water and electrolyte absorption by the jejunum, ileum, and colon under basal conditions are investigated in awake dogs with jejunal, ileal and/or colonic Thiry-Vella loops essentially as described in greater detail in Bilchik et al (Gastroenterology (1993) 105:1441-1448) and Bilchik et al (Am J Surg (1994) 167:570-574).

5 Each experiment consists of a 90-min basal period followed by a 150-min experimental period. The small bowel segment is perfused through the proximal cannula at 2 mL/min using a roller pump. The perfusate, pH 7.4 maintained at 37°C, contained (in nmol/L) 140 Na⁺, 5.2 K⁺, 119.8 Cl⁻, 25 HCO₃⁻, 1.2 Ca²⁺, 1.2 Mg²⁺, 2.4 HPO₄²⁻, 0.4 H₂PO₄⁻, 10 glucose and 10 μC [¹⁴C]poly(ethylene glycol) in 5 g/L poly(ethylene glycol). After a 20-min washout period,

10 perfusate is collected every 15 min for determination of Na⁺, Cl⁻ and water contents.

GPR119 agonist is administered per orally at a dose of, e.g., 30 mg/kg at time zero.

A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

15 It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is by intraluminal infusion at a dose of, e.g., 30 mg/kg/hour starting at time zero. A preferred infusion dose of the GPR119 agonist is 0.1-100 mg/kg/hour. Other preferred infusion dose is selected from the group consisting of 0.1 mg/kg/hour, 0.3 mg/kg/hour, 1.0 mg/kg/hour, 3.0 mg/kg/hour, 10 mg/kg/hour,

20 30 mg/kg/hour and 100 mg/kg/hour.

EXAMPLE 27

INHIBITION OF DIARRHEA OF CASTOR OIL-INDUCED DEFECATION AND DIARRHEAL SYMPTOM IN MICE

25 A GPR119 agonist can be shown to inhibit diarrheal symptom using a castor oil-induced defecation and diarrheal symptom mouse model essentially as described in EP 1902730.

C57BL/6 mice are used. Mice are individually housed in cages with a floor mat in an animal room with a controlled environment (temperature: 23°C, relative humidity: 55%, light hours: from 7:00 till 19:00) with ad libitum access to rodent chow (CE-2, Nippon CLEA) and

30 drinking water.

GPR119 agonist (at a dose of, e.g., 30 mg/kg), conventional antidiarrheal agent loperamide (1 mg/kg), conventional antidiarrheal agent atropine (10 mg/kg), or only vehicle are administered per oral administration, intraperitoneal administration, intraperitoneal administration, and oral administration, respectively, to the mice, and at 15 minutes after the administration, castor oil (0.3

35 ml/animal) is orally administered to the mice. The fecal weight and diarrhea score (normal: 0, soft feces: 1, watery feces: 2) is measured in a time-dependent manner (1, 2, and 4 hours after

administration). At the same time, the same measurement is carried out for mice to which only a vehicle is administered and castor oil is not administered.

A preferred dose of the GPR119 agonist is 0.1-100 mg/kg. Other preferred dose is selected from the group consisting of 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3.0 mg/kg, 10 mg/kg, 30 mg/kg and 100 mg/kg.

It is expressly contemplated that in other embodiment, administration of the agonist is other than oral, for example that administration of the agonist is intraperitoneal or intravenous.

EXAMPLE 28

10 ORAL BIOAVAILABILITY

Physicochemico analytical approaches for directly assessing oral bioavailability of a compound of the invention, such as a GPR119 agonist, a PYY secretagogue, a compound useful for treatment or prevention of a condition modulated by PYY, or a compound useful for a modulation of a PYY-related activity, are well known to those of ordinary skill in the art and may be used (*see*, e.g., without limitation: Wong PC et al., *Cardiovasc Drug Rev* (2002) 20:137-52; and Buchan P et al., *Headache* (2002) Suppl 2:S54-62; the disclosure of each of which is hereby incorporated by reference in its entirety). By way of further illustration and not limitation, said alternative analytical approaches may comprise liquid chromatography-tandem mass spectrometry (Chavez-Eng CM et al., *J Chromatogr B Analyt Technol Biomed Life Sci* (2002) 767:117-29; Jetter A et al., *Clin Pharmacol Ther* (2002) 71:21-9; Zimmerman JJ et al., *J Clin Pharmacol* (1999) 39:1155-61; and Barrish A et al., *Rapid Commun Mass Spectrom* (1996) 10:1033-7; the disclosure of each of which is hereby incorporated by reference in its entirety). Recently, positron emission tomography (PET) has been successfully used to obtain direct measurements of drug distribution, including oral bioavailability, in the mammalian body following oral administration of the drug (Gulyas et al., *Eur J Nucl Med Mol Imaging* (2002) 29:1031-8; the disclosure of which is hereby incorporated by reference in its entirety).

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations, or modifications, as come within the scope of the following claims and its equivalents.

What is claimed is:

1. Use of a G protein-coupled receptor for identifying a peptide YY (PYY) secretagogue, in a
5 method comprising the steps of:
- (a) contacting a test compound with a host cell or with membrane of a host cell
comprising said G protein-coupled receptor wherein the G protein-coupled receptor comprises an
amino acid sequence selected from the group consisting of:
- 10 (i) amino acids 1-335 of SEQ ID NO:2;
(ii) amino acids 2-335 of SEQ ID NO:2;
(iii) an amino acid sequence comprising the amino acid sequence of a
polypeptide derived from (i) by substitution, deletion or addition of one
or several amino acids in the amino acid sequence defined in (i);
15 (iv) the amino acid sequence of a G protein-coupled receptor encoded by a
polynucleotide capable of hybridizing under stringent conditions to the
complement of SEQ ID NO:1;
(v) the amino acid sequence of a variant of SEQ ID NO:2 when selected
from the group consisting of:
- 20 (a') the amino acid sequence of a G protein-coupled receptor having
at least about 80% identity to SEQ ID NO:2; and
(b') the amino acid sequence of a G protein-coupled receptor
comprising at least 20 contiguous amino acids of SEQ ID
NO:2;
25 (vi) the amino acid sequence of a constitutively active version of a G
protein-coupled receptor having SEQ ID NO:2; and
(vii) a biologically active fragment of any one of (i) to (vi); and
(b) determining the ability of the test compound to stimulate functionality of the
receptor;
wherein the ability of the test compound to stimulate functionality of the receptor is
30 indicative of the test compound being a PYY secretagogue.
2. Use according to claim 1, and further comprising:
- (c) contacting a compound which stimulates functionality of the receptor in step (b) in vitro
with a vertebrate enteroendocrine cell or a cell capable of secreting PYY; and
35 (d) determining whether the compound stimulates PYY secretion from the vertebrate
enteroendocrine cell or the cell capable of secreting PYY;

wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue.

- 5 3. Use according to claim 1, and further comprising:
 (c) determining whether the compound increases a PYY level in a sample, said sample having been obtained from a vertebrate previously administered with a compound which stimulates functionality of the receptor in step (b);
 wherein the ability of the test compound to increase a PYY level in the vertebrate is further
10 indicative of the test compound being a PYY secretagogue.
4. Use according to claim 1, and further comprising:
 (c) administering a compound which stimulates functionality of the receptor in step (b) to a vertebrate; and
15 (d) determining whether the compound increases a PYY level in the vertebrate;
 wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue.
5. Use according to claim 3 or 4, wherein the vertebrate is a non-human vertebrate.
20 6. Use according to any one of claims 1 to 5, wherein the receptor is recombinant.
7. Use according to any one of claims 1 to 6, wherein the host cell comprises an expression vector, said expression vector comprising a polynucleotide encoding the G protein-coupled
25 receptor.
8. Use according to any one of claims 1 to 7, wherein said determining is through the measurement of the level of a second messenger.
- 30 9. Use according to claim 8, wherein the second messenger is selected from the group consisting of cyclic AMP (cAMP), cyclic GMP (cGMP), inositol 1,4,5-triphosphate (IP₃), diacylglycerol (DAG), MAP kinase activity, MAPK/ERK kinase kinase-1 (MEKK1) activity, and Ca²⁺.
- 35 10. Use according to claim 9, wherein a level of cAMP is increased.

11. Use according to any one of claims 1 to 7, wherein said determining is through the use of a Melanophore assay, through the measurement of GTP γ S binding to a membrane comprising said GPCR, or through a reporter assay.
- 5 12. Use according to any one of claims 1 to 11, wherein the host cell is a mammalian host cell, or wherein the host cell is a yeast host cell, or wherein the host cell is a melanophore host cell.
13. Use according to any one of claims 1 to 12, wherein the test compound is a small molecule.
- 10 14. Use according to any one of claims 1 to 13, wherein the receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO:2.
- 15 15. Use according to any one of claims 1 to 14, wherein the receptor comprises the amino acid sequence of SEQ ID NO:2.
16. A method for identifying a peptide YY (PYY) secretagogue, comprising the steps of:
- (a) contacting a test compound in vitro with a host cell or with membrane of a host cell comprising said G protein-coupled receptor wherein the G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:
- 20 (i) amino acids 1-335 of SEQ ID NO:2;
- (ii) amino acids 2-335 of SEQ ID NO:2;
- (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (i);
- 25 (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide capable of hybridizing under stringent conditions to the complement of SEQ ID NO:1;
- (v) the amino acid sequence of a variant of SEQ ID NO:2 when selected from the group consisting of:
- 30 (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO:2; and
- (b') the amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO:2;
- (vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO:2; and
- 35 (vii) a biologically active fragment of any one of (i) to (vi); and
- (b) determining the ability of the test compound to stimulate functionality of the receptor;

wherein the ability of the test compound to stimulate functionality of the receptor is indicative of the test compound being a PYY secretagogue.

17. A method for identifying a PYY secretagogue, comprising the step of:

- 5 (a) determining whether a PYY level is increased in a sample as compared with a vertebrate reference sample not previously administered with a GPR119 agonist, said sample having been obtained from a vertebrate previously administered with a GPR119 agonist;

wherein the ability of the GPR119 agonist to increase a PYY level in the sample is indicative of the GPR119 agonist being a PYY secretagogue.

10

18. A method for identifying a PYY secretagogue, comprising the steps of:

(a) administering a GPR119 agonist to a vertebrate; and

(b) determining whether the GPR119 agonist increases a PYY level in the vertebrate;

- 15 wherein the ability of the GPR119 agonist to increase a PYY level in the vertebrate is indicative of the GPR119 agonist being a PYY secretagogue.

19. A method for identifying a PYY secretagogue, comprising the steps of:

(a) contacting a GPR119 agonist in vitro with a vertebrate enteroendocrine cell or a cell capable of secreting PYY; and

- 20 (b) determining whether the GPR119 agonist stimulates PYY secretion from the vertebrate enteroendocrine cell or the cell capable of secreting PYY;

wherein the ability of the GPR119 agonist to stimulate PYY secretion from the vertebrate enteroendocrine cell or cell capable of secreting PYY is indicative of the GPR119 agonist being a PYY secretagogue.

25

20. The method of claim 17, 18, or 19 wherein the vertebrate is a non-human vertebrate.

21. Use of a G protein-coupled receptor for identifying a PYY secretagogue, in a method comprising the steps of:

- 30 (a) contacting a G protein-coupled receptor with an optionally labeled known ligand for the receptor in the presence or absence of a test compound, wherein a ligand for the receptor is a ligand for GPR119 and wherein the G protein-coupled receptor comprises an amino acid sequence selected from the group consisting of:

- 35 (i) amino acids 1-335 of SEQ ID NO:2;
(ii) amino acids 2-335 of SEQ ID NO:2;

- (iii) an amino acid sequence comprising the amino acid sequence of a polypeptide derived from (i) by substitution, deletion or addition of one or several amino acids in the amino acid sequence defined in (i);
- (iv) the amino acid sequence of a G protein-coupled receptor encoded by a polynucleotide capable of hybridizing under stringent conditions to the complement of SEQ ID NO:1;
- (v) the amino acid sequence of a variant of SEQ ID NO:2 when selected from the group consisting of:
- (a') the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO:2; and
- (b') the amino acid sequence of a G protein-coupled receptor comprising at least 20 contiguous amino acids of SEQ ID NO:2;
- (vi) the amino acid sequence of a constitutively active version of a G protein-coupled receptor having SEQ ID NO:2; and
- (vii) a biologically active fragment of any one of (i) to (vi); and
- (b) detecting the complex between said known ligand and said receptor; and
- (c) determining whether less of said complex is formed in the presence of the test compound than in the absence of the test compound;
- wherein said determination is indicative of the test compound being a PYY secretagogue.

22. Use according to claim 21 wherein said use further comprises
- (d) contacting a compound in the presence of which less of said complex is formed in step (c) in vitro with a vertebrate enteroendocrine cell or a cell capable of secreting PYY; and
- (e) determining whether the compound stimulates PYY secretion from the vertebrate enteroendocrine cell or the cell capable of secreting PYY;
- wherein the ability of the test compound to stimulate PYY secretion from the vertebrate enteroendocrine cell or cell capable of secreting PYY is further indicative of the test compound being a PYY secretagogue.

23. Use according to claim 21 wherein said method further comprises
- (d) determining whether the compound increases a PYY level in a sample obtained from a vertebrate, said vertebrate having been administered with a compound in the presence of which less of said complex is formed in step (c) ;
- wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue.

24. Use according to claim 21 wherein said method further comprises:
(d) administering a compound in the presence of which less of said complex is formed in step (c) to a vertebrate; and
5 (e) determining whether the compound increases a PYY level in the vertebrate;
wherein the ability of the test compound to increase a PYY level in the vertebrate is further indicative of the test compound being a PYY secretagogue.
25. Use according to claim 21 or 22, wherein the vertebrate is a non-human vertebrate.
10
26. Use according to any one of claims 21 to 25, wherein the receptor is recombinant.
27. Use according to any one of claims 21 to 26, wherein the host cell comprises an expression vector, said expression vector comprising a polynucleotide encoding the G protein-coupled
15 receptor.
28. Use according to any one of claims 21 to 27, wherein the ligand for GPR119 is a GPR119 agonist.
- 20 29. The method according to any one of claims 16 to 20 or the use according to claim 28, wherein the GPR119 agonist is an agonist of human GPR119.
30. The method according to any one of claims 16 to 20 or 29 or the use according to claim 28 or claim 29 wherein said GPR119 agonist is a selective GPR119 agonist.
25
31. The method according to any one of claims 16 to 20, 29 or 30 or the use according to any one of claims 28 to 30, wherein the GPR119 agonist has an EC₅₀ of less than a value selected from the group consisting of 10 μ M, 1 μ M and 100 nM.
- 30 32. The method according to any one of claims 17 to 20 or 29 to 31, or the use according to any one of claims 28 to 31, wherein the GPR119 agonist is a small molecule.
33. The method according to any one of claims 17 to 20 or 29 to 32, or the use according to any one of claims 28 to 32, wherein the GPR119 agonist is orally bioavailable.
35
34. Use according to any one of claims 21 to 33, or the method according to claim 16 wherein the test compound is a small molecule.

35. Use or method according to any one of claims 21 to 34, wherein the receptor comprises the amino acid sequence of a G protein-coupled receptor having at least about 80% identity to SEQ ID NO:2.
- 5
36. Use or method according to any one of claims 21 to 35, wherein the receptor comprises the amino acid sequence of SEQ ID NO:2.
37. The use of a GPR119 agonist in the treatment or prevention of a condition modulated by peptide YY (PYY).
- 10
38. The use of a GPR119 agonist in the treatment or prevention of a condition modulated by stimulation of a NPY Y2 receptor (Y2R).
39. A method of treating or preventing a condition modulated by peptide YY (PYY), the method comprising administering a pharmaceutical composition comprising a GPR119 agonist to an individual in need thereof.
- 15
40. A method of treating a condition ameliorated by stimulation of a NPY Y2 receptor (Y2R), the method comprising administering a pharmaceutical composition comprising a GPR119 agonist to an individual in need thereof.
- 20
41. The method of claim 39 or 40 wherein the GPR119 agonist is administered in a manner or amount capable of stimulating PYY expression in an enteroendocrine cell of the individual.
- 25
42. The method according to any one of claims 16 to 20 or 29 to 33, further comprising producing or synthesizing the identified PYY secretagogue.
43. The method according to any one of claims 16 to 20, 29 to 33, or 42, further comprising producing or synthesizing the identified PYY secretagogue as a composition.
- 30
44. The method according to claim 16, wherein the test compound is a small molecule.
45. The method according to any one of claims 13, 16, 32, 34, or 44 wherein the small molecule has a weight of less than 10,000 grams per mole.
- 35

46. The method according to any one of claims 13, 16, 32, 34, or 44 wherein the small molecule has a weight of less than 5,000 grams per mole.
47. A dosage form comprising a GPR119 agonist, wherein the GPR119 agonist is present in an amount sufficient to increase PYY levels in a subject.
48. The dosage form of claim 47, wherein the amount is sufficient to increase PYY levels in an enteroendocrine cell in a subject.
49. The use or method according to claims 38-41, wherein the treating or preventing comprises an affect selected from the group comprising:
- (a) increasing plasma PYY;
 - (b) increasing bone mass;
 - (c) decreasing loss of bone mass;
 - (d) increasing bone formation;
 - (e) decreasing food uptake;
 - (f) decreasing body weight;
 - (g) increasing satiety;
 - (h) increasing glucose disposal;
 - (i) decreasing plasma glucose;
 - (j) increasing angiogenesis;
 - (k) decreasing secretion in intestinal epithelium;
 - (l) increasing absorption in intestinal epithelium; and
 - (m) increasing plasma adiponectin.
50. A GPR119 agonist for use in a method for the treatment of a condition, wherein the condition may be treated by increasing PYY levels in an individual.