

4-Aminotetrahydrobenzisoazole or -isothiazole Compounds

The present invention relates to novel 4-aminotetrahydrobenzisoazoles or -isothiazoles having GABA-uptake inhibiting activity and thus being useful in the treatment of analgesia, psychosis, convulsions, anxiety or muscular and movement disorders, such as spastics disorders or symptoms in Huntington's disease or Parkinsonism. The anticonvulsant activity especially provides usefulness as broad spectrum anti-epileptic agents.

Background of the invention

The neutral amino acid, 4-aminobutanoic acid (GABA), is an inhibitory transmitter in the central nervous system. There is considerable direct and indirect evidence that impaired operation of GABA-mediated inhibitory synapses may be an important causative factor in seizure disorders (P. Krogsgaard-Larsen et al., *Epilepsy Res.* 1987, 1, 77-93) making GABA-ergic drugs potential antiepileptic therapeutic agents.

Furthermore, enhancement of GABA-ergic activity may be useful in the treatment of anxiety, pain, muscular and movement disorders and mental and emotional disorders (W. Löscher, *Eur. J. Pharmacol.*, 1985, 110, 103-108).

20

While direct stimulation of GABA receptors by agonists does not seem to represent the most suitable therapeutic approach to epileptic diseases (R.G. Fariello et al., Eds., *Neurotransmitters, Seizures, and Epilepsy II*, 1984, New York, Raven Press; B. Meldrum and R. Horton, *Eur. J. Pharmacol.* 1980, 61, 231-237; Krogsgaard-Larsen et al., *J. Med. Chem.* 1994, 37, 2489-2505.), GABA neurotransmission may be facilitated by manipulation of the GABA uptake mechanisms. Pharmacological inhibition of the neuronal and/or glial GABA transport, assumed to be responsible for the termination of GABA neurotransmission processes, provides a mechanism for sustaining levels of synaptically released GABA in the synapses and thereby increasing GABA-mediated transmission (P. Krogsgaard-Larsen et al., *J. Med. Chem.* 1994, 37, 2489-2505).

30

The strategies for such pharmacological interventions may be: 1) effective blockade

of both neuronal and glial GABA uptake, or 2) selective blockade of the uptake of GABA into glial cells in order to increase the amount of GABA taken up by the neuronal carrier with subsequent elevation of the GABA concentration in nerve terminals. There is evidence suggesting that glia-selective GABA uptake inhibitors may have particular interest as antiepileptic agents (E. Falch et al., *Drug Design and Delivery*, 1987, 2, 9-21; Falch et al. *Drug Dev. Res.*, 1990, 21, 169-188).


Classical GABA uptake inhibitors are nipecotic acid, guvacine and THPO. Oral active N-substituted derivatives of nipecotic acid and guvacine are described in F.E. Ali et al., *J. Med. Chem.* **1985**, 28, 553-560; U.S. patent 4,383,999 and U.S. patent 4,514,414 to SmithKline Beckmann Corporation; EP 236342 and EP 231996 to Novo Industri A/S and H.S. White et al., *Eur. J. Pharmacol.* **1993**, 236, 147-149.

With regard to convulsion, especially epilepsy, in spite of the fact that antiepileptic
15 drugs are available, many patients fail to experience seizure control. Consequently,
it is an object of the present invention to provide new GABA'ergic drugs effective in
the treatment of diseases associated with GABA neurotransmission, in particular
seizure control.

20 Summary of the invention


It has now been found that a class of novel 4-aminotetrahydrobenzisoxazoles or -isothiazoles inhibits neuronal and/or glial GABA-uptake.

25 Accordingly, the present invention relates to novel 4-aminotetrahydrobenzisoxazole
or -isothiazole compounds having general formula Ia or Ib:



wherein R¹ and R² are independently selected from the group consisting of:


A) hydrogen, cycloalkyl, phenyl, or a group




where R⁷, R⁸ and R⁹ are independently selected from the group con-

sisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy-lower
 5 alkyl, cycloalkyl, cycloalkyl-lower alkyl, phenyl, phenyl-lower alkyl, phenoxy-
 lower alkyl and heteroaryl selected from 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, imida-
 zolyl, oxazolyl, pyrazolyl, pyrimidinyl, pyrrolyl, thiazolyl, 1,2,4-triazolyl, 2-pyridyl,
 3-pyridyl or 4-pyridyl, any phenyl or heteroaryl group present optionally being
 substituted with one or two substituents selected from halogen, lower alkyl,
 10 lower alkoxy, hydroxy, nitro, lower alkylthio, lower alkylsulfonyl, lower alkyl- or di-
 (lower)alkylamino, cyano, trifluoromethyl, trifluoromethylthio, trifluoromethylsul-
 fonyloxy and phenyl which again may be substituted with halogen, methyl,
 methoxy or trifluoromethyl; and any alkyl group present being optionally
 substituted with one to three hydroxy groups which again are optionally
 15 esterified with a C₂₋₁₈ carboxylic acid;

B) a group of general formula Y-(CH₂)_r-(CHR¹¹)_s-(CH₂)_t wherein Y is selected
 from the following groups (1) - (5):



20



wherein U is CHR^{10b}, NR^{10b}, O or S, U¹ is NR^{10b}, O or S; p is 0 or 1; q is 0 or 1;
 V is C or N and the dotted line represents a bond when V is C and no bond
 when V is N;

25 A is O, S, CH₂, (CH₂)₂, CH=CH-CH₂, (CH₂)₃, CH=CH or O-CH₂;




R^a and R^b each represent one or more substituents selected from halogen, lower alkyl, lower alkoxy, hydroxy, nitro, lower alkylthio, lower alkylsulfonyl, lower alkyl- or di(lower alkyl)amino, cyano, trifluoromethyl, trifluoromethylsulfonyloxy and trifluoromethylthio;

- 5 r and t are independently 0, 1, 2 or 3, s is 0 or 1, provided that when Y is a group (1) wherein U is NR^{10b} , O or S or a group (4), then $r + s + t$ is at least 2; and when Y is a group (3) or a group (5) where V is N, then $r + s + t$ is at least 1; R^{7b} , R^{8b} and R^{9b} are as defined for R^7 , R^8 and R^9 in A) provided that they are not at the same time selected from hydrogen, lower alkyl, lower alkenyl and lower alkynyl;

10 R^{10b} and R^{11} are independently hydrogen, lower alkyl, lower alkenyl or lower alkynyl; and

- C) a group of general formula $Y^c-(CH_2)_n-W-(CH_2)_m-$ wherein n is 1, 2 or 3, m is 2 or 3; W is O or S; and Y^c is a group (1) - (5) as defined in B) provided that n may not be 1, when Y^c is a group (1) or (4) wherein U or U^1 , respectively, is NR^{10b} , S or O;

- D) a group of general formula $Y^d-O-(CH_2)_k-\overset{\overset{R^{11d}}{|}}{CH}-(CH_2)_l-$ wherein k is 0, 1, 2 or 3; l is 0, 1, 2 or 3; R^{11d} is as defined for R^{11} in B) above; and Y is selected from the groups (2) and (5) as defined in B) above and the following groups (6) - (10):



Wherein p , q , R^a , R^b , and A are as defined in B) and R^{7d} - R^{11d} are as defined for R^{7b} - R^{10b} and R^{11} , respectively, under B) or

R¹ and R² together designate alkylene thereby forming a 4-8 membered nitrogen containing ring; or

- 5 one of R¹ and R² is a group R^{2'}OCO wherein R^{2'} is phenyl, or heteroaryl as defined in A) above or phenyl or such heteroaryl substituted with one or more substituents selected from halogen, lower alkyl, lower alkoxy, hydroxy, nitro, lower alkylthio, lower alkylsulfonyl, lower alkyl- or di(lower)alkylamino, cyano, trifluoromethyl, trifluoromethylthio, trifluoromethylsulfonyloxy, phenyl and phenyl substituted with halogen,
10 methyl, methoxy or trifluoromethyl;

R³ - R⁶ are independently selected from hydrogen, hydroxy and lower alkyl, any alkyl group optionally being substituted with one or two hydroxy groups;

- 15 X is oxygen or sulfur;

P is hydrogen or a group ZR wherein

Z is CO, CS, SO₂ or CR^tR^u, R^t and R^u being hydrogen, hydroxy or lower alkyl, and

if Z is CO or CS, then R is selected from the groups consisting of:

- 20 i) hydrogen, C₁-C₁₈ alkyl, C₂-C₁₈ alkenyl, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkenyl or C₄-C₂₆ cycloalk(en)yl-alk(en)yl, optionally substituted with one or two hydroxy groups, or phenyl optionally substituted with one or more substituents selected from the group consisting of halogen, trifluoromethyl, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₅ acyloxy, or cyano; or
25 ii) QR^v, wherein Q is O or S and R^v is selected from the substituents defined for R under i) above; and
iii) NR^xR^y, wherein R^x and R^y independently are selected from the substituents defined for R under i) above or R^x and R^y are combined to form a four to eight membered heterocyclic ring containing from one to three nitrogen atoms and
30 from zero to three oxygen or sulfur atoms; or
if Z is CR^tR^u, R is selected from the groups consisting of:
iv) a group QR^v as defined in ii);
v) a group NR^xR^y as defined in iii); or

- vi) a group OC(O)R_z, SC(O)R_z, OC(S)R_z or SC(S)R_z wherein R_z is selected from the substituents defined for R under i) above;
- if Z is SO₂, R is selected from group i) defined above;
- provided that P may not be hydrogen, when R¹ to R⁶ are all hydrogen, X is oxygen
- 5 and the compound exists as a racemic mixture;
- or a pharmaceutically acceptable salt thereof.

- The compounds of the invention have been found to inhibit neuronal and/or glial GABA-uptake, some of the compounds predominantly inhibiting the glial uptake.
- 10 Thus the compounds are useful in the treatment of diseases associated with GABA neurotransmission, e.g. as analgesic, antipsychotic, anticonvulsant, or anxiolytic drugs or as drugs for the treatment of muscular and movement disorders, such as spastic disorders or symptoms in Huntington's disease or Parkinsonism.
- 15 In another aspect the invention provides a pharmaceutical composition comprising at least one novel 4-aminotetrahydrobenzisoxazoles or -isothiazoles of Formula I in a therapeutically effective amount together with a pharmaceutically acceptable carrier and or diluent.
- 20 In a further aspect the present invention provides the use of a 4-aminotetrahydrobenzisoxazoles or -isothiazoles of Formula I for the manufacture of a pharmaceutical preparation for the treatment of the above mentioned disorders and diseases.

- Also, the present invention provides a method for the preparation of the 4-amino-
- 25 tetrahydrobenzisoxazoles or -isothiazoles of Formula I.

Detailed Description of the Invention

- The compounds of general Formula I exist as optical isomers thereof and such
- 30 optical isomers as well as any mixture thereof, including the racemic mixtures, are also embraced by the invention.

In the present context, the term lower alkyl designates straight or branched C₁₋₄


alkyl, such as methyl, ethyl, propyl, isopropyl, butyl and tert.butyl. Similarly, lower alkenyl and lower alkynyl designate such groups having up to 4 carbon atoms and having at least one double or tripple bond, respectively. Lower alkoxy, lower alkylthio, lower alkylsulfonyl, lower alkylamino, lower dialkylamino, etc. similarly designate such groups wherein the alkyl moiety is a lower alkyl group as defined above.

The term cycloalkyl designates a saturated carbocyclic ring having 3-7 carbon atoms, inclusive, and the term halogen designates F, Cl, Br or I.

10 Alk(en/yn)yl means that the group may be an alkyl, alkenyl or alkynyl group.

Heteroaryl means 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, imidazolyl, oxazolyl, pyrazolyl, pyrimidinyl, pyrrolyl, thiazolyl, 1,2,4-triazolyl, 2-pyridyl, 3-pyridyl or 4-pyridyl.

15 The compounds wherein P is a group ZR are prodrugs for the compounds wherein P is H. When P is H the compound exists as tautomeric forms as follows:



20 So, when reacted with the proper reactant in order to introduce the group ZR, the said group is either introduced at the O-atom exo to the isoxazole/isothiazole ring or at the ring N-atom depending on the reaction conditions. In the following, compounds wherein P is H, are for the sake of convenience named as having the first mentioned form, i.e. as having formula 1a. Similar the compounds wherein R¹ or R² is a group R^{2'}OCO are prodrugs for the corresponding compounds wherein R¹ or R² is hydrogen.

The pharmaceutically acceptable acid addition salts of the compounds used in the invention are salts formed with non-toxic organic or inorganic acids. Exemplary of such organic salts are those with maleic, fumaric, benzoic, ascorbic, embonic, suc-

cinic, oxalic, bis-methylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-amino-benzoic, glutamic, benzene sulfonic and theophylline acetic acids, as well as the 8-halotheophyllines, for example
 5 8-bromo-theophylline. Exemplary of such inorganic salts are those with hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric and nitric acids.

It is preferred that R^1 is a group as defined in

A) above, namely lower alk(en/yn)yl optionally substituted with hydroxy which may
 10 be esterified with a C_{2-22} carboxylic acid, preferably with a C_{2-4} carboxylic acid; or phenyl or phenyl-lower alkyl, optionally substituted with halogen, lower alkyl, lower alkoxy or trifluoromethyl; R^1 being more preferably methyl, ethyl or phenylbutyl;

B) above wherein Y is a group of formula (1), (2) or (3), wherein R^{7b} and R^{8b} are phenyl, heteroaryl or substituted phenyl or heteroaryl, most preferably phenyl,
 15 phenyl substituted with halogen, lower alkyl, lower alkoxy or trifluoromethyl, 2-thienyl, 2-thienyl substituted with methyl, pyrrolyl or pyrrolyl substituted with methyl or ethyl. In particular Y a group of formula (1) wherein U is CH_2 or O or a group of formula (2), or Y is a group of formula (4) or (5), wherein p and q are 0 and A is sulfur or $-CH_2CH_2-$ and R^a and R^b are hydrogen.

20 In formula (1), (2) or (3) R^{9b} is hydrogen is preferably hydrogen or lower alkyl, especially hydrogen and R^{10b} is hydrogen, and in formula (1), (2), (3), (4) and (5) s is preferably 0 and r+t is 0-5, more preferably 1-3, most preferably 2.

C) above where Y^c is a group of formula (1) where U is CH_2 , a group of formula (2), (3) or (5) in which formulas R^{7b} and R^{8b} are phenyl, heteroaryl or substituted phenyl
 25 or heteroaryl, most preferably phenyl, phenyl substituted with halogen, lower alkyl, lower alkoxy or trifluoromethyl, 2-thienyl, 2-thienyl substituted with methyl, pyrrolyl or pyrrolyl substituted with methyl or ethyl. R^{9b} is preferably hydrogen or lower alkyl, especially hydrogen, R^{10b} is hydrogen, and n+m is preferably 0-5, more preferably 1-3, most preferably 2. or

30 D) above where Y is a group of formula (6), (7) or (8), in which formulas R^{7d} and R^{8d} are independently phenyl, phenyl substituted with halogen, lower alkyl, trifluoromethyl or lower alkoxy, in particular chloro, fluoro, methyl, trifluoromethyl, or methoxy, 2-thienyl, 2-thienyl substituted with lower alkyl; most preferably methyl,

pyrrolyl or pyrrolyl substituted with methyl or ethyl. $R^{9d} - R^{11d}$ are preferably hydrogen and $k+l$ is 0-4, more preferably 0-2, most preferably 1.

R^2 is preferably hydrogen, lower alk(en/yn)yl or a group $R^2'OCO$. Most preferably
5 R^2 is hydrogen or lower alkyl, in particular hydrogen or methyl.

Preferably R^3 is hydrogen or lower alkyl, $R^4 - R^6$ are hydrogen and X is oxygen.

P is preferably hydrogen or a group ZR wherein Z is CR^tR^u , R^t and R^u being
10 hydrogen, hydroxy or lower alkyl, most preferably hydrogen or methyl, and R is a group $OC(O)R^z$, $SC(O)R^z$, $OC(S)R^z$ or $SC(S)R^z$ wherein R^z is lower alk(en)yl, in particular *t*-butyl, or optionally substituted phenyl; or wherein Z is CO or CS and R is NR^xR^y , wherein R^x and R^y are hydrogen or lower alkyl or R^x and R^y form a heterocyclic ring having one or two N-atoms and being optionally mono- or disubsti-
15 tuted with oxo.

Preferably the compound has the structure of formula Ia.

In particular it is preferred that the group P is different from hydrogen when R^1 is a
20 preferred group selected from A) as defined above, especially a lower alk(en/yn)yl group, and R^2 is hydrogen or lower alk(en/yn)yl.

When R^1 and/or R^2 is a group as defined under B), C), or D) it is preferred that P is hydrogen.

25

In a preferred subclass of the compounds of the invention R^1 is a group as defined in A) namely lower alk(en/yn)yl optionally substituted with hydroxy which may be esterified with a C_{2-22} carboxylic acid, or phenyl or phenyl-lower alkyl, optionally substituted with halogen, lower alkyl, lower alkoxy or trifluoromethyl, R^1 being most
30 preferably methyl, ethyl or phenylbutyl;

R^2 is hydrogen or lower alk(en/yn)yl, in particular hydrogen or methyl;

R^3 is hydrogen or lower alkyl, in particular hydrogen or methyl, $R^4 - R^6$ are hydrogen and X is oxygen;

P is a group ZR wherein Z is CH₂ or CH(CH₃), and R is a group OC(O)R_z, wherein R_z is lower alkyl or optionally substituted phenyl or heteroaryl; or wherein Z is CO and R is NR_xR_y, wherein R_x and R_y are hydrogen or lower alkyl or R_x and R_y form a heterocycle.

5

In a further preferred subclass of the compounds of the invention R¹ is a group as defined in B) above wherein s is 0 and r+t is 0-4, in particular 1-3, and Y^b is a group of formula (1) or (2) wherein R^{7b} and R^{8b} are independently phenyl, phenyl substituted with halogen, lower alkyl, trifluoromethyl or lower alkoxy, in particular chloro, fluoro, methyl, trifluoromethyl, or methoxy, 2-thienyl, 2-thienyl substituted with lower alkyl, most preferably methyl, pyrrolyl or pyrrolyl substituted with methyl or ethyl, R^{9b} - R^{10b} are hydrogen and U is CH₂ or O or Y^b is a group of formula (5) where p and q are 0 and A is sulfur or -CH₂CH₂- and R^a and R^b are hydrogen; and R², R³, R⁴-R⁶ and X are as defined in the preferred subclass above and P is hydrogen.

According to another preferred subclass of the compounds of the invention R¹ is a group as defined in C) above where Y is a group of formula (1) where U is CH₂, or a group of formula (3) or (5), in which formulas R^{7b} and R^{8b} are independently phenyl, phenyl substituted with halogen, lower alkyl, trifluoromethyl or lower alkoxy, in particular chloro, fluoro, methyl, trifluoromethyl, or methoxy, 2-thienyl, 2-thienyl substituted with lower alkyl, most preferably methyl, pyrrolyl or pyrrolyl substituted with methyl or ethyl, R^{9b} and R^{10b} are hydrogen, W is O, n is 1 - 3, most preferably 1, and m is 2 - 4, most preferably 2 or 3; and R², R³, R⁴-R⁶ and X are as defined in the preferred subclass above and P is hydrogen.

According to another preferred subclass of the compounds of the invention R¹ is a group as defined in D) above where Y is a group of formula (6), (7) or (8), in which formulas R^{7d} and R^{8d} are independently phenyl, phenyl substituted with halogen, lower alkyl, trifluoromethyl or lower alkoxy, in particular chloro, fluoro, methyl, trifluoromethyl or methoxy, 2-thienyl, 2-thienyl substituted with lower alkyl, most preferably methyl, pyrrolyl, or pyrrolyl substituted with methyl or ethyl, R^{9d}, R^{10d} and

R^{11d} are hydrogen, W is O, k+l is 0-3, most preferably 1 or 2; and R², R³, R⁴-R⁶ and X are as defined in the preferred subclass above and P is hydrogen.

- 5 The pharmaceutical compositions of this invention or those which are manufactured in accordance with this invention may be administered by any suitable route, for example orally in the form of tablets, capsules, powders, syrups, etc., or parenterally in the form of solutions for injection. For preparing such compositions methods well known in the art may be used, and any pharmaceutically acceptable carriers,
- 10 diluents, excipients, or other additive usually used in the art may be used.


Conveniently, the compounds of the invention are administered in unit dosage form containing said compounds in an amount of about 0.1 to 1000 mg.

- 15 The total daily dose is usually in the range of about 0.5 - 5000 mg, and most preferably about 1.0 to 500 mg of the active compound of the invention.


The method of the invention for preparing the novel compounds of formula I comprises:

20

- a) In order to obtain a compound of formula I wherein R² is hydrogen, removing the acyl protection group of a compound of formula II or III :



II



III


25

wherein R¹, R³ - R⁶, X and ZR are as defined above and Ac is an acyl protection group;


- b) Alkylating an amine of formula HNR¹R² wherein R¹ and R² are as defined above

12

with a ketone of formula IV or V:



IV




V


where R⁴-R⁶, X and ZR are as defined above, in the presence of a reducing agent;

5

c) In order to obtain a compound of formula I wherein P is H, removing the B group of a compound of formula VI or VII:




VI




VII

10 wherein R¹ - R⁶ and X are as defined above and B is lower alkyl, phenyl lower alkyl or a group ZR as defined above;

d) Reducing the double bond of a Schiff base, oxime, or oxime ether of the following formula VIII or IX:



VIII




IX

wherein R¹, R⁴ - R⁶, X and ZR are as defined above.

e) Arylating a compound of formula X

20

13



wherein R¹, R³ - R⁶ and X are as defined above with an acylation agent of the formula R^{2'}-O-CO-hal, wherein hal is Cl or Br and R^{2'} is as defined above.

5

The starting material for the preparation of isoxazole intermediates of formulas II to X is 4,5,6,7-tetrahydro-1,2-benzisoxazol-3-ol, which is conveniently prepared according to the method described by R. Jaquier et al., *Bull. Soc. Chim. Fr.* 1970, 5, 1978-1985. Starting material for the corresponding thioisoxazoles is cyclohexanone-2-
10 carboxamide. Details for the conversion of these intermediates to compounds of formula I are given in the Experimental Section.

The acyl protection group in method a) is conveniently removed by aqueous hydrolysis catalyzed by bases (sodium or potassium hydroxide or potassium carbonate)
15 or acids (hydrochloric or hydrobromic), hydrogenation of benzyloxy-carbonyl or 1,1,1-trichloroethyloxycarbonyl groups, or anhydrous acid catalyzed deprotection of e.g. the t-Boc protection group. Hydrogenation either involves catalytic hydrogenation in a Parr apparatus, using Pd as catalyst, or hydrogenation in presence of metals such as zinc in aqueous acidic solution such as diluted acetic acid.

20

Amines used in the reductive alkylation in method b) are commercially available or prepared according to well established literature methods for example as described in F.E. Ali et al., *J. Med. Chem.* 1985, 28, 553-560; U.S. patent 4,383,999 and U.S. patent 4,514,414 to SmithKline Beckmann Corporation; EP 236342 and EP 231996
25 to Novo Industri A/S and H.S. White et al., *Eur. J. Pharmacol.* 1993, 236, 147-149. NaBH₄ or NaCNBH₃, preferably in the presence of a dehydrating agent such as a molecular sieve, may be used as reducing agent in a protic solvent such as methanol, ethanol, water or mixtures thereof. Proper salts of the amines are used to obtain optimal pH conditions.

The protecting groups B in method c) are effectively removed by base or acid catalyzed hydrolysis. If B is an O-alkyl group such as methoxy or ethoxy it might be conveniently removed by treatment with strong acid (*e.g.* 48% hydrobromic acid in
5 glacial acetic acid) at elevated temperatures.

Schiff bases, oximes, or oxime ethers in method d) are conveniently reduced to the corresponding amine derivatives by catalytic hydrogenation using *e.g.* Pd or Pt as catalysts, or by reduction with amalgamated aluminum or with LiAlH_4 or AlH_3 .

10

Experimental Section

Melting points were determined on a Büchi SMP-20 apparatus and are uncorrected. Mass spectra were obtained on a Quattro MS-MS system from VG Biotech, Fisons Instruments. The MS-MS system was connected to an HP 1050 modular HPLC
15 system. A volume of 20-50 μl of the sample (10 $\mu\text{g/ml}$) dissolved in a mixture of 1% acetic acid in acetonitril/water 1:1 was introduced via the autosampler at a flow of 30 $\mu\text{l/min}$ into the Electrospray Source. Spectra were obtained at two standard sets of operating conditions. One set to obtain molecular weight information (MH^+) (21 eV) and the other set to induce fragmentation patterns (70 eV). The background
20 was subtracted. The relative intensities of the ions are obtained from the fragmentation pattern. When no intensity is indicated for the Molecular Ion (MH^+) this ion was only present under the first set of operating conditions. ^1H NMR spectra were recorded of all novel compounds at 250 MHz on a Bruker AC 250 spectrometer, at 200 MHz on a Bruker AC 200 F spectrometer, or as otherwise stated in
25 the experiments. Deuterated chloroform (99,8%D) or dimethylsulfoxide (99,9%D) were used as solvents. TMS was used as internal reference standard. Chemical shift values are expressed in ppm-values. The following abbreviations are used for multiplicity of NMR signals: s=singlet, d=doublet, t=triplet, q=quartet, qui=quintet, h=heptet, dd=double doublet, dt=double triplet, dq=double quartet, tt=triplet of triplets,
30 m=multiplet. NMR signals corresponding to acidic protons are generally omitted. Content of water in crystalline compounds was determined by Karl Fischer titration. Standard workup procedures refer to extraction with the indicated organic solvent from proper aqueous solutions, drying of combined organic extracts (anhydrous MgSO_4 or Na_2SO_4), filtering and evaporation of the solvent in vacuo. For column

chromatography silica gel of type Kieselgel 60, 230-400 mesh ASTM was used.

Example 1

3-Ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, 1a

- 5 The starting material 4,5,6,7-tetrahydro-1,2-benzisoxazol-3-ol was prepared according to a literature procedure (R.Jaquier et al., *Bull.Soc.Chim.Fr.* 1970, 5, 1978-1985). To a solution of this isoxazole derivative (100 g) in acetone (3 L) was added potassium carbonate (200 g). After heating at 50 °C for 45 minutes a solution of bromoethane (170 mL) in acetone (300 mL) was added dropwise during 1.5 hours. The
- 10 mixture was stirred overnight at 50 °C. After cooling, inorganic salts were filtered off and acetone evaporated *in vacuo*. The remaining mixture of O- and N-alkylated product was separated by column chromatography on silica gel (eluted with ethyl acetate/heptane 40:60). Evaporation of the solvents afforded 65 g of the title product as a viscous oil. ¹H NMR (CDCl₃): δ 1.40 (t, 3H), 1.70-1.85 (m, 4H), 2.25-
- 15 2.30 (m, 2H), 2.50-2.60 (m, 2H), 4.30 (q, 2H).

Example 2

3-Ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol-4-one, 2a (Method d))

- To a solution of 3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 1a (35 g) in glacial
- 20 acetic acid (500 mL) at 10 °C was added concentrated H₂SO₄ (29 mL). A solution of sodium dichromate (71 g) in glacial acetic acid (300 mL) kept at 40-45 °C was added dropwise during one hour at 20-25 °C. The mixture was stirred for another 3 hours at 25-30 °C. The reaction mixture was poured onto ice and diethyl ether (3 L). pH was adjusted to >10 by addition of concentrated aqueous NaOH. The organic
- 25 phase was separated and worked up. The remaining crude title compound was purified by column chromatography on silica gel (eluted with ethyl acetate / heptane 1:1). Pure 3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol-4-one was washed with diisopropyl ether and dried. Yield 27 g. Mp 98-99 °C.

30 Example 3

(R,S)-4-amino-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, 3a

To a solution of 3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol-4-one 2a (8 g) in ethanol (400 mL) were added hydroxylammonium hydrochloride (20 g), Na₂CO₃ (16

g), and water (600 mL). The mixture was heated at reflux temperature for 3 hours. The solvents were partly evaporated (2/3). The remaining mixture was left at room temperature for 0.5 hours and the precipitated crystalline product was finally filtered off, washed with water and dried. Yield of the oxime derivative 7 g. Mp 216-218 °C.

- 5 To a solution of HgCl_2 (200 g) in water (4 L) was added aluminium foil (125 g) cut into small pieces (0.5 x 0.5 cm). These pieces were left for about 1 minute and subsequently filtered off and washed with ethanol. The oxime derivative (25 g) was suspended in methanol (2 L) and water (500 mL) and the aluminium foil was added. The resulting mixture was stirred for 5 days. The precipitates were filtered off and
10 washed with methanol. The combined methanol solutions were evaporated *in vacuo*. The remaining crude product was stirred with diethyl ether. Undissolved starting material (oxime) was filtered off and the diethyl ether was evaporated *in vacuo* leaving the title compound as a viscous oil. Yield 23 g. ^1H NMR (CDCl_3) δ 1.40 (t, 3H), 1.40-1.1.75 (m, 4H), 1.90-2.05 (m, 2H), 2.50-2.60 (m, 2H), 3.90 (broad t, 1H),
15 4.30 (q, 2H).

Example 4

4-amino-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 4a enantiomer A

- 20 (*R,S*)-4-Amino-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 3a (17.5 g) was dissolved in dichloromethane (500 mL) and triethylamine (30 mL) was added at 5 °C. A solution of (*R*)-(-)- α -methoxyphenylacetyl chloride (22 g) in dichloromethane (80 mL) was added dropwise at 5-10 °C. The resulting mixture was stirred for 2 hours at room temperature. Water (2 L) was added and the organic phase was separated,
25 washed with diluted hydrochloric acid and worked up as above. Yield 34 g. The diastereomers of this mixture were separated by preparative HPLC on silica gel (eluted with heptane/ethyl acetate 3:2). Yield of the less polar diastereomer B 13 g as an oil. Yield of the more polar diastereomer A 11.5 g. Mp 96-97 °C. ^1H NMR (CDCl_3) δ 1.30 (t, 3H), 1.80-2.05 (m, 4H), 2.45-2.75 (m, 2H), 3.30 (s, 3H), 4.25 (q, 2H), 4.60
30 (s, 1H), 4.95 (dt, 1H), 6.85 (broad d, 1H), 7.30-7.45 (m, 5H). The diastereomer A (3.7 g) was dissolved in 48 % aqueous hydrobromic acid (175 mL) and water (175 mL). This solution was refluxed for 1.25 hours. The solvent was evaporated *in vacuo*. Dichloromethane and water were added. The organic phase was separated,

washed with water and finally discarded. The combined aqueous phases were evaporated *in vacuo*. Ethanol/ether 1:1 were added, the precipitates were filtered off, and the solvents evaporated leaving the impure crude hydrobromide salt 4a (enantiomer A) as an oil. Yield 4.0 g. Purification is shown below in Examples 5 and 6.

- 5 The crude hydrobromic salt of enantiomer B, 4b was isolated correspondingly from diastereomer B.

Example 5

4-(*tert*-butyloxycarbonylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 5a
10 enantiomer A.

A mixture of crude 4-amino-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 4a, enantiomer A from Example 4 (4.0 g) was dissolved in a 1:1 mixture of water/dioxane. After cooling to 10 °C, a solution of NaOH (1.2 g in 12 mL of water) was added. Di-*tert*-butyl dicarbonate (3.5 g) dissolved in dioxane (12 mL) was
15 added at 15-20 °C. The mixture was stirred at room temperature for 1.5 hours. Water (120 mL) was added and pH was adjusted >10 by adding a small amount of NaOH. After stirring for additionally 30 minutes, diethyl ether (200 mL) was added. The organic phase was separated and discarded. By addition of KHSO₄ pH was adjusted to 3-4 and the aqueous phase was extracted with diethyl ether (2 x 100
20 mL). The combined organic phases were worked up as above leaving the crude Boc-protected title compound. The pure title compound 5a was obtained by column chromatography on silica gel (eluted with heptane/ethyl acetate/ethanol 7:3:1 Yield 0.9 g. Mp: 135 °C. ¹H NMR (CDCl₃) δ 1.45 (s, 9H), 1.75-2.10 (m, 4H), 2.45-2.70 (m, 2H), 4.50 (dt, 1H), 5.05 (broad d, 1H).

25

The corresponding enantiomer B was prepared in a similar way:

4-(*tert*-butyloxycarbonylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, 5b
enantiomer B. Mp 134 °C. The NMR spectrum was identical to the spectrum above for compound 5a.

30

Example 6

(+)-4-amino-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrochloride 6a (=s)-
isomer.

The boc-protected enantiomer A, 5a (0.9 g) obtained in Example 5 was dissolved in a saturated solution of HCl in ether (200 mL) and stirred at room-temperature for 1.25 hours. The solvent was evaporated and the remaining oil was dissolved in a 1:1 mixture of ethanol in ether. The precipitated crystalline title compound was
5 filtered off. Yield : 0.5 g. Mp 209-210 °C. $[\alpha]_D = +19.4^\circ$ (c=1.0 M, methanol). ^1H NMR (DMSO- d_6) δ 1.75-2.05 (m, 4H), 2.55-2.65 (m, 2H), 4.25 (broad t, 1H), 8.40 (broad s, 4H). MS m/z (%): 155 (MH $^+$, 49%), 138 (100%), 113 (16%), 65 (58%).

The (-)-isomer (or (R)-isomer) was prepared in a corresponding way:

10 (-)-(R)-4-amino-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrochloride 6b. Mp 209-210 °C. $[\alpha]_D = -20.0^\circ$ (c=1.0 M, methanol). ^1H NMR and MS spectra were the same as for the (+)-enantiomer above.

Compound 8a was resolved in a corresponding way via HPLC separation of the diastereomeric (R)-(-)- α -methoxyphenylacetyl carboxamide derivatives of compound

15 7a. A modification of the method in Example 4 was used to split off protecting groups.

(+)-3-hydroxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide
6c

20 To the most polar diastereomer (9 g) separated via HPLC chromatography was dissolved in dry THF a solution of 1M lithiumtriethylborohydride in dry THF (80 mL) was added dropwise during 20 minutes at 0-5 °C. The mixture was further stirred overnight at roomtemperature. The mixture was poured onto ice (500 g) and pH was adjusted to 2 by addition of concentrated hydrochloric acid. To remove THF the
25 mixture was evaporated *in vacuo*. To the remaining aqueous solution was extracted twice with ethyl acetate (50 mL). The remaining aqueous solution was made alkaline by addition of concentrated NaOH solution (pH = 11). Ethyl acetate (100 mL) was added and the organic phase was subsequently separated and worked up as above. All of the thus isolated 3-ethoxy-4-methylamino-4,5,6,7-tetrahydro-1,2-
30 benzisoxazole enantiomer (4.6 g) was dissolved in a mixture of 33% Hbr in glacial acetic acid (150 mL). The mixture was heated to 90 °C and stirred for an hour. The solution was evaporated *in vacuo*. The remaining viscous oil was stirred with a mixture of ethanol / diethyl ether 1:1. The precipitated hydrobromic acid salt was

filtered off and dried overnight *in vacuo*. Yield 4.2 g. Mp 207-209 °C. $[\alpha]_D = +5.6^\circ$ (c=1.0 M, methanol). ¹H NMR and MS spectra were identical to spectra of the racemic mixture, compound 8a. Enantiomeric purity by HPLC determination : ee > 99.

- 5 (-)-3-hydroxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 6d was prepared analogously from the other diastereomeric carboxamide derivative. Mp 208-209 °C. $[\alpha]_D = -5.9^\circ$ (c=1.0 M, methanol). Enantiomeric purity by HPLC determination : ee > 99.

10 Example 7

(*R,S*)-3-Ethoxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole 7a

- To a solution of 3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol-4-one 2a (4.5 g) in methanol (100 mL) were added methylamine hydrochloride (15 g), a 33% solution of methylamine (5 mL) in ethanol, and molecular sieves (3 Å) powder. Finally
15 sodium cyanoborohydride was added (7 g). The mixture was stirred overnight. Inorganic salts were filtered off and the solvents evaporated *in vacuo*. Ethyl acetate and water were added and pH was adjusted >10 by addition of concentrated NaOH solution. The organic phase was subsequently worked up as above. The crude title compound was used without further purification. Yield 4.8 g. ¹H NMR (CDCl₃) δ 1.40
20 (t, 3H), 1.65-1.80 (m, 3H), 1.85-2.00 (m, 1H), 2.05 (s, 1H), 2.45 (s, 3H), 2.50-2.60 (m, 2H), 3.60 (t, 1H), 4.30 (q, 2H).

The following 3-ethoxyisoxazoles were synthesized in a similar way :

- (*R,S*)-4-[4,4-Bis(4-fluorophenyl)butan-1-ylamino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-
25 benzisoxazole 7b. ¹H NMR (CDCl₃) δ 1.35 (t, 3H), 1.40-1.55 (m, 3H), 1.60-1.85 (m, 3H), 1.90-2.10 (m, 3H), 2.45-2.60 (m, 2H), 2.65 (t, 2H), 3.70 (t, 1H), 3.85 (t, 1H), 4.30 (q, 2H), 6.95 (t, 4H), 7.15 (dd, 4H).

- (*R,S*)-3-Ethoxy-4-(2-hydroxyethylamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole 7c
Mp 72-74 °C. ¹H NMR (CDCl₃) δ 1.40 (t, 3H), 1.65-2.00 (m, 4H), 2.45 (s, 2H), 2.45-
30 2.70 (m, 2H), 2.85 (t, 2H), 3.60-3.70 (m, 2H), 3.75 (t, 1H), 4.30 (q, 2H).

(*R,S*)-3-Ethoxy-4-(1-pyrrolidinyl)-4,5,6,7-tetrahydro-1,2-benzisoxazole 7d ¹H NMR (CDCl₃) δ 1.40 (t, 3H), 1.40-1.55 (m, 2H), 1.70-1.90 (m, 6H), 2.00-2.25 (m, 2H),

2.50-2.80 (m, 4H), 3.25 (t, 1H), 4.30 (dq, 2H).

(*R,S*)-4-(4,4-diphenylbutan-1-ylamino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, 7e. Prepared via the butylamine 30a and isolated as an oil. ¹H NMR (CDCl₃) δ 1.35 (t, 3H), 1.40-1.50 (m, 2H), 1.60 (s, 1H), 1.60-2.00 (m, 6H), 2.00-2.20 (m, 2H), 2.45-2.60 (m, 2H), 2.65 (t, 2H), 3.65 (t, 1H), 3.85 (t, 1H), 4.30 (q, 2H), 7.10-7.30 (m, 10H).

(*R,S*)-4-(4-phenylbutan-1-ylamino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 7f. Isolated as an oil. ¹H NMR (CDCl₃) δ 1.40 (t, 3H), 1.60 (s, 1H), 1.50-2.00 (m, 8H), 2.50-2.70 (m, 6H), 3.70 (t, 1H), 4.30 (q, 2H), 7.15-7.35 (m, 5H).

10 (*R,S*)-4-(3,3-diphenylpropan-1-ylamino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 7g. Isolated as an oil. ¹H NMR (CDCl₃) δ 1.40 (t, 3H), 1.60 (s, 1H), 1.60-1.80 (m, 4H), 1.85-2.00 (m, 1H), 2.25 (q, 2H), 2.50-2.65 (m, 3H), 3.65 (t, 1H), 4.10 (t, 1H), 4.25 (q, 2H), 7.15-7.35 (m, 10H).

15 (*R,S*)-4-[N-[3-(10,11-dihydrodibenzo[*a,d*]cyclohept-5-ylidene)propane-1-yl]amino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, 7h. Prepared via the 3-(10,11-dihydrodibenzo[*a,d*]cyclohept-5-ylidene)propylamine, hydrochloride (prepared as described in J. Org. Chem. (1962), Vol. 27, 4134-37) and isolated as an oil. ¹H NMR (CDCl₃) δ 1.40 (t, 3H), 1.50-2.05 (m, 6H), 2.35 (q, 2H), 2.40-2.65 (m, 2H), 2.65-2.85 (m, 2H), 2.85-3.50 (m, 3H), 3.65 (t, 1H), 4.25 (q, 2H), 5.90 (t, 1H), 7.00-7.07 (m, 1H), 7.07-7.21 (m, 6H), 7.21-7.33 (m, 1H).

Example 8A

(*R,S*)-3-Hydroxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 8a (Method c).

25 A solution of (*R,S*)-3-Ethoxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole 7a (3.8 g) in a 33% solution of HBr in glacial acetic acid (150 mL) was heated at 60-75 °C for one hour. The solvents were evaporated *in vacuo* and a 1:1 mixture of ethanol/ether was added. The crystalline hydrobromic salt 8a was filtered off. Yield : 3.6 g. Mp: 184-186 °C. ¹H NMR (DMSO-d₆) δ 1.75-2.15 (m, 4H), 2.65 (s, 3H), 2.60-2.70 (m, 2H), 4.20 (broad signal, 1H), 8.60 (broad s, 1H).

The following compounds were synthesized correspondingly:

(*R,S*)-4-[4,4-Bis(4-fluorophenyl)butan-1-ylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide **8b** Mp 205-206 °C. ¹H NMR (DMSO-*d*₆) δ 1.45-1.60 (m, 2H), 1.70-2.10 (m, 6H), 2.55-2.70 (m, 2H), 3.05 (t, 2H), 4.00 (t, 1H), 4.20 (broad s, 1H), 7.15 (t, 4H), 7.35 (dd, 4H). MS *m/z* (%): 399 (MH⁺, 4 %), 138 (100 %), 67 (84 %).

(*R,S*)-4-(2-Acetyloxyethylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide **8c**. Mp 164-165 °C. ¹H NMR (DMSO-*d*₆) δ 1.75-1.95 (m, 2H), 2.00-2.20 (m, 2H), 2.10 (s, 3H), 2.55-2.70 (m, 2H), 3.30 (t, 2H), 4.20-4.40 (m, 3H). MS *m/z* (%): 241 (MH⁺, 17 %), 138 (59 %), 67 (100 %), 41 (84 %).

10 (*R,S*)-3-Hydroxy-4-(1-pyrrolidiny)-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide **8d** Mp 209-210 °C. ¹H NMR (DMSO-*d*₆) δ 1.70-2.30 (m, 8H), 2.60-2.75 (m, 2H), 3.10-3.70 (m, 4H), 4.35 (broad s, 1H) MS *m/z* (%): 209 (MH⁺, 5 %), 138 (35 %), 72 (56%), 67 (100%), 41 (79 %).

(*R,S*)-4-(2-propen-1-ylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, oxalate (acetone) **8f**-This compound was purified via the 4-*tert*-butyloxy-carbonylamino derivative, which was deprotected as described in Example 8B. Mp 182-183 °C. ¹H NMR (DMSO-*d*₆) δ 1.75-2.20 (m, 4H), 2.60-2.75 (m, 2H), 3.65 (d, 2H), 4.15 (broad s, 1H), 5.40 (d, 1H), 5.45 (d, 1H), 5.85-6.00 (m, 1H), 7.70 (broad signal, 3H). MS *m/z* (%): 195 (MH⁺, 9 %), 138 (57 %), 67 (100 %), 41 (92 %).

20 (*R,S*)-4-(4,4-diphenylbutan-1-ylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide **8g**. Mp 221-222 °C (ethanol). ¹H NMR (DMSO-*d*₆) δ 1.45-1.65 (m, 2H), 1.70-2.10 (m, 6H), 2.55-2.75 (m, 2H), 3.05 (t, 2H), 3.95 (t, 1H), 4.20 (broad s, 1H), 7.15-7.40 (m, 10H). MS *m/z* (%): 363 (MH⁺, 100 %), 138 (89 %).

(*R,S*)-4-(4-phenylbut-1-ylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide **8h**. Mp 202-204 °C (ethanol). ¹H NMR (DMSO-*d*₆) δ 1.55-1.65 (m, 4H), 1.70-2.15 (m, 4H), 2.50-2.70 (m, 4H), 3.00 (broad t, 2H), 4.25 (broad s, 1H), 7.15-7.35 (m, 5H). MS *m/z* (%): 287 (MH⁺, 6 %), 138 (100 %), 91 (42%), 67 (63%).

(*R,S*)-4-(3,3-diphenylpropan-1-ylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide **8i**. Mp 218-220 °C (ethanol). ¹H NMR (DMSO-*d*₆) δ 1.70-2.10 (m, 4H), 2.40 (t, 2H), 2.55-2.70 (m, 2H), 2.95 (t, 2H), 4.05 (t, 1H), 4.25 (broad s, 1H), 7.15-7.40 (m, 10H). MS *m/z* (%): 349 (MH⁺, 5 %), 138 (100 %), 67 (30%).

The following compounds were prepared in a similar manner except that the title

compounds were crystallized from acetone (8j, 8k, 8l, 8m, 8n, 8p) or diethyl ether (8o).

(*R,S*)-4-[N-[3-(10,11-dihydrodibenzo[*a,d*]cyclohept-5-ylidene)propane-1-yl]amino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide, 8j. Mp 228-230 °C
5 (dec.). ¹H NMR (DMSO-*d*₆) δ 1.55-2.00 (m,4H), 2.15-2.65 (m,4H), 2.65-3.00 (m,4H), 3.10-3.40 (m,2H), 3.77-3.87 (m,1H), 5.80 (t,1H), 7.00-7.28 (m,8H), MS *m/z* (%): 387 (MH⁺, 5%), 233 (7%), 138 (41%), 43 (100%)

(*R,S*)-4-[N-[3-(10,11-dihydrodibenzo[*a,d*]cyclohept-5-ylidene)propane-1-yl]methylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide, 8k. Mp 215-
10 217 °C (dec.). ¹H NMR (DMSO-*d*₆) δ 1.70-2.10 (m,4H), 2.40-2.95 (m,6H), 3.05-3.44 (m,7H), 4.34-4.43 (m,1H), 5.80 (t,1H), 7.05-7.30 (m,8H), MS *m/z* (%): 401 (MH⁺, 26%), 265 (66%), 233 (30%), 138 (84%), 43 (100%)

(*R,S*)-4-[N-3-(Phenothiazin-10-yl)propane-1-yl]amino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide, 8l. Mp 187-189 °C. ¹H NMR (DMSO-*d*₆) δ 1.65-
15 2.12 (m,6H), 2.50-2.71 (m,2H), 3.00-3.16 (m,2H), 3.95 (t,2H), 4.09-4.20 (m,1H), 6.97 (dd,2H), 7.07 (d,2H), 7.10-7.26 (m,4H), MS *m/z* (%): 394 (MH⁺, 3%), 256 (7%), 138 (18%), 43 (100%)

(*R,S*)-4-[N-[4,4-Di-(2-tolyl)butan-1-yl]-methylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide, 8m. Mp 193-195 °C (dec.). ¹H NMR (DMSO-*d*₆) δ
20 1.65-2.20 (m,8H), 2.26 (s,6H), 2.55-2.80 (m,5H), 3.15-3.35 (m,2H), 4.25 (t,1H), 4.36-4.47 (m,1H), 7.03-7.21 (m,8H), MS *m/z* (%): 405 (MH⁺, 4%), 268 (27%), 138 (30%), 43 (100%)

(*R,S*)-4-[N-[4,4-Di-(2-tolyl)butan-1-yl]amino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide, 8n. Mp 217-219 °C (dec.). ¹H NMR (DMSO-*d*₆) δ
25 1.55-2.18 (m,8H), 2.25 (d,6H), 2.52-2.75 (m,2H), 3.06 (t,2H), 4.13-4.27 (m,2H), 7.03-7.20 (m,8H), MS *m/z* (%): 391 (MH⁺, 7%), 195 (15%), 145 (80%), 138 (92%), 105 (100%)

(*R,S*)-4-[N-[1,1-Di-(2-tolyl)but-1-en-4-yl]methylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, 8o. Mp 177-179 °C. ¹H NMR (DMSO-*d*₆) δ 1.70-2.14 (m,8H),
30 2.21 (s,3H), 2.36-2.80 (m,6H), 3.15-3.40 (m,2H), 4.35-4.47 (m,1H), 5.74 (t,1H), 6.98-7.30 (m,8H), MS *m/z* (%): 403 (MH⁺, 19%), 266 (40%), 143 (77%), 138 (100%), 105 (49%), 67 (20%)

(*R,S*)-4-[N-1,1-Di-(2-tolyl)but-1-en-4-yl]amino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide, 8p. Mp 209-211 °C (dec.). ¹H NMR (DMSO-*d*₆) δ 1.70-2.15 (m,7H), 2.22 (s,3H), 2.27-2.45 (m,2H), 2.54-2.73 (m,2H), 3.11 (t,2H), 4.16-4.24 (m,1H), 5.73 (t,1H), 6.96-7.26 (m,8H), MS m/z (%): 389 (MH⁺, 5%), 143 (33%), 138 (100%), 105 (29%), 67 (44%).

Example 8B

(*R,S*)-3-Hydroxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 8a (Method c))

- 10 This method is an alternative way to the method in Example 8A of preparing compound 8a and similar derivatives.

A solution of di-*tert*-butyl dicarbonate (3.56 g) in THF (50 mL) was added to a solution of (*R,S*)-3-ethoxy-4-amino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 3a (2.60 g) and potassium carbonate (2.07 g) in water (25 mL). The mixture was stirred at room temperature for 20 h and evaporated. Water (30 mL) was added to the residue and the mixture was extracted with ether (3 x 50 mL). The combined organic extracts were dried and evaporated, and the residue was recrystallized from ether/light petroleum to give (*R,S*)-3-ethoxy-4-(*tert*. butyloxycarbonylamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole (2.69 g): mp 111-113°C. 60% Sodium hydride in mineral oil (425 mg) was added in small portions to a solution of the said compound (1.00 g) and methyl iodide (2.18 mL) in THF (45 mL). The mixture was stirred overnight at room temperature and methanol was added to destroy excess of sodium hydride. After evaporation, water (25 mL) was added to the residue. Extraction with ethyl acetate (3 x 50 mL), drying and evaporation gave (*R,S*)-3ethoxy-4-(N-methyl-*tert*.butyloxycarbonylamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole (1.05 g) as a yellow oil. ¹H NMR (60 MHz, CDCl₃) δ 1.40 (t, 3H), 1.55 (s, 9H), 1.65-2.2 (m, 4H), 2.55 (m, 2H), 2.60 (s, 3H), 4.35 (q, 2H), 5.20 (m, 1 H). A 33 % solution of hydrobromic acid in glacial acetic acid (15 mL) was added to (*R,S*)-3-ethoxy-4-(N-methyl-*tert*-butyloxycarbonylamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole (1.05 g) and the mixture was stirred at 80 °C for 25 min. After evaporation, 33% hydrobromic acid in glacial acetic acid (15 ml) was added to the residue and the mixture was stirred at 80°C for 25 min. Evaporation and recrystallization of the residue (acetonitrile-

ethanol-ether) gave the title compound (807 mg): mp 188-190°C. ¹H NMR (60 MHz, D₂O) δ 2.05 (m, 4H), 2.75 (m, 2H), 2.85 (s, 3H), 4.35 (m, 1 H).

The following compound was prepared in a similar manner by using ethyl iodide in
5 stead of methyl iodide:

(*R,S*)-4-Ethylamino-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide
8e Mp : 188-191 °C (from acetonitrile-ethanol-ether). ¹H NMR (60 MHz, D₂O) δ 1.30
(t, 3H), 2.05 (m, 4H), 2.70 (m, 2H), 3.25 (q, 2H), 4.35 (m, 1 H).

10 Example 9

(*R,S*)-4-(2-Hydroxyethylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole,
hydrobromide 9a

To a solution of (*R,S*)-4-(2-acetyloxyethylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-
benzisoxazole, hydrobromide 8c (1.3 g) in water (50 mL) was added 48% aqueous
15 HBr (7 mL). The mixture was heated at 100 °C for an hour. The solvent was evapo-
rated *in vacuo* and the remaining oil was dissolved in ethanol. Upon evaporation of
a major fraction of the ethanol, the hydrobromide salt of compound 9a crystallized
and was subsequently filtered off and dried. Yield : 0.9 g. Mp 172-173 °C. ¹H NMR
(DMSO-d₆) δ 1.70-1.95 (m, 2H), 1.95-2.30 (m, 2H), 2.55-2.75 (m, 2H), 3.10 (t, 2H),
20 3.70 (q, 2H), 4.25 (broad s, 1H), 5.20 (broad s, 1H), 8.65 (broad s, 2H), 11.95
(broad s, 1H). MS m/z (%): 199 (MH⁺, 16 %), 138 (33 %), 67 (91%), 41 (100 %).

Example 10

(*R,S*)-4-[N-[4,4-Bis(4-fluorophenyl)butan-1-yl]-N-methylamino]-3-ethoxy-4,5,6,7-
25 tetrahydro-1,2-benzisoxazole 10a

To (*R,S*)-3-ethoxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole 7a (1.0 g) in
methyl isobutyl ketone (MIBK) (10 mL) were added bis-4,4-(4-fluorophenyl)-1-butyl-
chloride (2.0 g), potassium carbonate (1.0 g), and potassium iodide (0.5 g). The mix-
ture was refluxed overnight. Inorganic salts were filtered off and MIBK was evapora-
30 ted. The remaining oil was purified by column chromatography on silica gel (eluted
with heptan/ethyl acetate 2:3). Yield of the title compound as an oil 1.6 g. ¹H NMR
(CDCl₃) δ 1.35 (t, 3H), 1.35-1.50 (m, 2H), 1.60-1.75 (m, 3H), 1.90-2.05 (m, 3H), 2.20
(s, 3H), 2.35-2.60 (m, 4H), 3.60 (t, 1H), 3.85 (t, 1H), 4.25 (q, 2H), 6.95 (t, 4H), 7.15

(dd, 4H).

The following 3-ethoxy derivative was prepared in a corresponding manner :

(*R,S*)-4-[*N*-(4,4-Diphenylbut-1-yl)-*N*-methylamino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, **10b**. Prepared via alkylation with 4,4-Diphenyl-1-butyliodide, **29a** and isolated as an oil. ¹H NMR (CDCl₃) δ 1.30 (t, 3H), 1.35-1.50 (m, 2H), 1.60-1.70 (m, 4H), 1.90-2.10 (m, 2H), 2.15 (s, 3H), 2.30-2.50 (m, 4H), 3.55 (t, 1H), 3.85 (t, 1H), 4.25 (q, 2H), 7.10-7.30 (m, 10H).

10 The following compounds were prepared in a similar manner except that the alkylations were performed in acetone without addition of potassium iodide.

(*R,S*)-4-[*N*-{3-(10,11-dihydrodibenzo[*a,d*]cyclohept-5-ylidene)propane-1-yl}]methylamino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, **10c**. Prepared via alkylation with 3-(10,11-dihydrodibenzo[*a,d*]cyclohept-5-ylidene)-1-propylbromide (prepared as described in J. Org. Chem. (1962), Vol. 27, 4134-37) and isolated as an oil. ¹H NMR (CDCl₃) δ 1.30 (t, 3H), 1.60-1.74 (m, 3H), 1.87-2.06 (m, 1H), 2.20 (s, 3H), 2.20-2.35 (m, 2H), 2.45-2.70 (m, 4H), 2.70-3.50 (m, 4H), 3.50-3.63 (m, 1H), 4.24 (q, 2H), 5.88 (t, 1H), 6.98-7.30 (m, 8H)

(*R,S*)-4-[*N*-[4,4-Di-(2-tolyl)butan-1-yl]-methylamino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, **10d**. Prepared via alkylation with 4,4-di-(2-tolyl)-1-butyliodide, **34a** and isolated as an oil. ¹H NMR (CDCl₃) δ 1.28 (t, 3H), 1.42-1.56 (m, 2H), 1.60-1.80 (m, 4H), 1.87-2.05 (m, 3H), 2.20 (s, 3H), 2.26 (dd, 6H), 2.35-2.60 (m, 4H), 3.60 (t, 1H), 4.23 (q, 2H), 7.00-7.17 (m, 8H)

(*R,S*)-4-[*N*-[1,1-Di-(2-tolyl)but-1-en-4-yl]methylamino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, **10e**. Prepared via alkylation with 4,4-di-(2-tolyl)-3-butenyliodide, **36a** and isolated as an oil. ¹H NMR (CDCl₃) δ 1.33 (t, 3H), 1.55-1.80 (m, 4H), 1.90-2.07 (m, 2H), 2.17 (dd, 6H), 2.26 (s, 3H), 2.45-2.67 (m, 4H), 3.57 (t, 1H), 4.26 (q, 2H), 5.81 (t, 1H), 7.00-7.17 (m, 8H)

30 **Example 11 (Method c))**

(*R,S*)-4-[*N*-[4,4-Bis(4-fluorophenyl)butan-1-yl]-*N*-methylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide **11a**

All of the product **10a** (1.6 g) from Example 10 was heated at 80 °C in a 33%

solution of hydrogen bromide in glacial acetic acid (60 mL) for 1 hour. The solvent was evaporated *in vacuo*. The remaining crude product was repeatedly dissolved in ethanol and evaporated *in vacuo*. The remaining viscous oil was dissolved in glacial acetic acid (15 mL) and water (150 mL) was added. After freeze-drying over-
5 night, an amorphous powder of the title compound 11a remained. This powder was stirred with diethyl ether and filtered off. After drying *in vacuo* at 50 °C for 24 hours, the pure amorphous hydrobromide was collected. Yield 1.2. Mp 68-70 °C. ¹H NMR (DMSO-d₆ recorded at 60 °C) δ 1.60-2.15 (m, 8H), 2.55-2.70 (m, 2H), 2.70 (broad s, 3H), 3.20 (broad t, 2H), 4.05 (t, 1H), 4.45 (t, 1H), 7.10 (t, 4H), 7.35 (dd,
10 4H). MS m/z (%): 413 (MH⁺, 10 %), 203 (13 %), 138 (100 %), 109 (12 %), 67 (58 %).

The following 3-hydroxy derivative was prepared in a corresponding manner :
(*R,S*)-4-[-N-(4,4-Diphenylbut-1-yl)-N-methylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-
15 benzisoxazole, hydrobromide 11b. Mp 174-176 °C (ethanol/ diethyl ether 1:1). ¹H NMR (DMSO-d₆) δ 1.50-2.10 (m, 6H), 2.60-2.80 (m, 4H), 3.15-3.40 (m, 2H), 3.30 (s, 3H), 3.95 (t, 1H), 4.40 (broad s, 1H), 7.15-7.45 (m, 10H). MS m/z (%): 377 (MH⁺, 100%), 240(56%), 138 (49%).

20 Example 12

(*R,S*)-4-(*N*-*tert*-Butyloxycarbonyl-*N*-methylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 12a

To a solution of (*R,S*)-3-hydroxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 8a (7.0 g) in a mixture of dioxane (50 mL) and water (80 mL)
25 were added at 10 °C NaOH (1.1 g) and a solution di-*tert*-butyldicarbonate (6.0 g) in dioxane (20 mL). The mixture was stirred at room temperature for 1.5 hours. Water (150 mL) was added. After stirring for additionally 20 minutes diethyl ether (100 mL) was added. The organic phase was separated and discarded. pH of the aqueous phase was adjusted to 4 by addition of KHSO₄. The aqueous phase was repeatedly
30 extracted with diethyl ether (3 x 100 ml). The combined organic phases were worked up as above. Yield 5.3 g. Mp 151-152 °C. ¹H NMR (CDCl₃) δ 1.50 (s, 9H), 1.65-2.10 (m, 4H), 2.55-2.65 (m, 2H), 2.75 (s, 3H), 5.15 (broad s, 1H).

By the same method was synthesized:

(*R,S*)-4-(*tert*-Butyloxycarbonylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 12b. Mp 175-177 °C (from ethyl acetate-light petroleum).

5 Example 13

(*R,S*)-4-(*N-tert*-butyloxycarbonyl-*N*-methylamino)-3-pivaloyloxymethyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 13a

To a suspension of (*R,S*)-4-(*N-tert*-butyloxycarbonyl-*N*-methylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 12a (5 g) in acetone (50 mL) was added cautiously (temperature below 30 °C) potassium *tert*-butoxide (2.5 g). A solution of pivaloyloxymethyl iodide (7.5 g) in acetone (10 mL) was added and the mixture was stirred overnight. Inorganic salts were filtered off and the acetone evaporated. Column chromatography on silica gel (eluted with heptane/ethyl acetate 3:2) yielded 3.9 g of the title compound 13a as an oil. ¹H NMR (CDCl₃) δ 1.30 (s, 9H), 1.50 (s, 9H), 1.50-2.10 (m, 6H), 2.50-2.60 (m, 5H), 5.00-5.30 (broad signal, 1H), 5.85 (dd, 2H).

A more polar fraction from the column chromatography contained 2.1 g of (*R,S*)-4-(*N-tert*-butyloxycarbonyl-*N*-methylamino)-2-pivaloyloxymethyl-4,5,6,7-tetrahydro-1,2-benzisoxazol-3-one 13b. ¹H NMR (CDCl₃) δ 1.20 (s, 9H), 1.50 (s, 9H), 1.60-2.10 (m, 4H), 2.40-2.50 (m, 2H), 2.65 (s, 3H), 5.00-5.10 (broad signal, 1H), 5.75 (s, 2H).

The following compounds were synthesized correspondingly:

(*R,S*)-4-(*tert*-Butyloxycarbonylamino)-3-pivaloyloxymethyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 13c (oil). ¹H-NMR (CDCl₃) δ 1.25 (s, 9H), 1.45 (s, 9H), 1.80-1.95 (m, 4H), 2.50-2.75 (m, 2H), 4.75 (broad s, 2H), 5.90 (s, 2H).

(*R,S*)-4-(*tert*-Butyloxycarbonylamino)-2-pivaloyloxymethyl-4,5,6,7-tetrahydro-1,2-benzisoxazole 13d (oil). ¹H-NMR (CDCl₃) δ 1.20 (s, 9H), 1.45 (s, 9H), 1.80-2.00 (m, 4H), 2.30-2.55 (m, 2H), 4.50-4.90 (m, 2H), 5.75 (s, 2H).

30 Example 14

(*R,S*)-4-methylamino-3-pivaloyloxymethyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hemioxalate 14a (Method a))

To a solution of (*R,S*)-4-(*N-tert*-butyloxycarbonyl-*N*-methylamino)-3-pivaloyloxyme-

thyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 13a (3.7 g) in dichloromethane kept at 20 °C was added trifluoroacetic acid (19 mL). The mixture was stirred at 20 °C for another hour. The solvents were evaporated at room temperature *in vacuo*. The remaining oil was dissolved in diethyl ether (100 mL) and water (100 mL). Potassium carbonate was added to obtain pH>9. The organic phase was separated and worked up as above. Yield of crude title compound as the free base 2.4 g. To a solution of all of the base of 14a in ethanol (10 mL) was added oxalic acid (0.7 g). The precipitated hemioxalate salt was filtered off. Yield 1.2 g. Mp 201-202 °C. ¹H NMR (DMSO-*d*₆) δ 1.15 (s, 9H), 1.65-2.05 (m, 4H), 2.40 (s, 3H), 2.55-2.80 (m, 2H), 3.85 (t, 1H), 5.90 (dd, 2H). MS *m/z* (%): 283 (MH⁺), 138 (30 %), 57 (100 %).

In a corresponding way from (*R,S*)-4-(*N*-*tert*-butyloxycarbonyl-N-methylamino)-2-pivaloyloxymethyl-4,5,6,7-tetrahydro-1,2-benzisoxazol-3-one (13b) was isolated (*R,S*)-4-methylamino-2-pivaloyloxymethyl-4,5,6,7-tetrahydro-1,2-benzisoxazol-3-one, hemioxalate 14b. Mp 177-178 °C (from acetone). ¹H NMR (DMSO-*d*₆) δ 1.15 (s, 9H), 1.60-2.05 (m, 4H), 2.50 (s, 3H), 2.45-2.60 (m, 2H), 3.80 (t, 1H), 5.80 (dd, 2H). MS *m/z* (%): 283 (MH⁺), 123 (52 %), 57 (52 %), 55 (100%).

Similarly, treatment of (*R,S*)-4-(*tert*-Butyloxycarbonylamino)-3-pivaloyloxy-methoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 13c with a 2.5M solution of HCl in ethyl acetate afforded (*R,S*)-4-amino-3-pivaloyloxymethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole hydrochloride 14c. Mp 157-168 °C (from acetonitrile-ether). ¹H-NMR (CDCl₃) δ 1.20 (s,9H), 2.0-2.35 (m,4H), 2.50-2.85 (m,2H), 4.25-4.50 (m,1H), 5.90 (s,2H).

25

The enantiomers of compound 14a were prepared as follows:

From (+)-3-hydroxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide, 6c, the corresponding enantiomer of 4-(*N*-*tert*-butyloxycarbonyl-N-methylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole was prepared by reaction with di-*tert*-butyldicarbonate according to the procedure in Example 12. Further reaction with pivaloyloxymethyl iodide as in Example 13 gave the pure isomer of 4-(*N*-*tert*-butyloxycarbonyl-N-methylamino)-3-pivaloyloxymethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole. As previously described (Example 14) the BOC-protecting group

was splitted off by treatment with trifluoroacetic acid. The pure enantiomer A of compound 14a crystallized from acetone as the hemioxalate salt. Mp 211-213 °C. The optical rotation was $[\alpha]_D = -5.4^\circ$ (C=1, MeOH).

- 5 In a corresponding way from compound 6d the other isomer was prepared :
4-Methylamino-3-pivaloyloxymethyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hemioxalate, enantiomer B, Mp 210-213 °C. The optical rotation was $[\alpha]_D = +5.6^\circ$ (C=1, MeOH).
- 10 To prove the enantiomeric purity both compounds were analysed by ^1H NMR in the presence of R(-)-1-(9-anthryl)-2,2,2-trifluoroethanol. There is a splitting of the singlets from the *tert*-butyl group and the methyl group both into two singlet signals for the racemate. Enantiomer A gave singlets at $\delta(\text{ppm})$ 1.15 (s, 9H) and 2.15 (s, 3H). Enantiomer B gave singlets at $\delta(\text{ppm})$ 1.20 (s, 9H) and 2.25 (s, 3H). No impuri-
15 ties from the other isomer was detected in the two compounds. The detection limit was estimated to be 2 %.

Example 15 (Method a))

(*R,S*)-4-methylamino-2-phenylaminocarbonyl-4,5,6,7-tetrahydro-1,2-benzisoxazol-3-one, hydrochloride 15a
20

To a solution of (*R,S*)-4-(*N-tert*-butoxycarbonyl-*N*-methylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 12a (1.0 g) in dichloromethane (10 mL) was added phenylisocyanate (0.6 mL). The mixture was stirred overnight at room temperature. Dichloromethane was evaporated. (*R,S*)-4-(*N-tert*-butoxycarbonyl-*N*-methylamino)-
25 -2-phenylaminocarbonyl-4,5,6,7-tetrahydro-1,2-benzisoxazol-3-one crystallized from diethyl ether. The said crystallized compound (0.4 g) was suspended in a saturated solution of hydrogen chloride in diethyl ether. After a few minutes of stirring almost all of the compound had dissolved and the hydrochloride salt of the title compound started precipitating. After stirring for 2 hours the precipitated title compound was
30 filtered off and carefully washed with diethyl ether. Yield 0.3 g. Mp 168-170 °C. ^1H NMR (DMSO- d_6) δ 1.65-2.05 (m, 2H), 2.15-2.30 (m, 2H), 2.65 (s, 3H), 2.60-2.80 (m, 2H), 4.20 (broad s, 1H), 7.20 (t, 1H), 7.40 (t, 2H), 7.55 (d, 2H).

The following compounds were prepared in a corresponding way :

- (*R,S*)-2-Methylaminocarbonyl-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisothiazol-3-one, oxalate **15b**. Mp: 153 °C (acetone). ¹H NMR (DMSO-*d*₆) δ 1.75-2.15 (m, 4H),
5 2.60-2.70 (m, 2H), 2.65 (s, 3H), 2.85 (d, 3H), 4.15 (broad s, 1H), 7.80 (q, 1H), 9.15 (broad s, 3H). MS *m/z* (%): 226 (MH⁺), 160 (9%), 138 (57%), 67 (100%), 65 (50%).
(*R,S*)-2-Benzylaminocarbonyl-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisothiazol-3-onehydrochlorid **15c**. Mp: 85-89 °C (diethyl ether). ¹H NMR (DMSO-*d*₆) δ 1.80-
10 2.05 (m, 2H), 2.10-2.30 (m, 2H), 2.60-2.80 (m, 2H), 2.65 (s, 3H), 4.15 (broad s, 1H), 4.45 (d, 2H), 7.20-7.40 (m, 5H), 8.45 (t, 1H), 9.30-9.60 (broad d, 2H). MS *m/z* (%): 302 (MH⁺), 169 (8 %), 138(58%), 91 (97%), 65 (100 %).

Example 16

- (*R,S*)-4-[N-[bis(4-fluorophenyl)methyl-2-oxyethyl]-N-methylamino]-3-hydroxy-
15 4,5,6,7-tetrahydro-1,2-benzisoxazole, sodium salt **16a** (method c))
To a solution of (*R,S*)-4-methylamino-3-pivaloyloxymethyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole **14a** (1.1 g) in MIBK (18 mL) were added potassium carbonate (0.7 g), and the methanesulfonate ester of bis(4-fluorophenyl)methyl-2-oxyethanol (1.8 g). The mixture was refluxed overnight. Inorganic salts were filtered off and MIBK
20 evaporated. Column chromatography afforded pure (*R,S*)-4-[N-[bis(4-fluorophenyl)methyl-2-oxyethyl]-N-methylamino]-3-pivaloyloxymethyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole. Yield 1.4 g. ¹H NMR (CDCl₃) δ 1.20 (s, 9H), 1.60-1.80 (m, 3H), 1.95-2.05 (m, 1H), 2.25 (s, 3H), 2.55 (broad t, 2H), 2.60-2.80 (m, 2H), 3.50 (t, 2H), 3.65 (dt, 1H), 5.35 (s, 1H), 5.90 (dd, 2H), 7.00 (t, 4H), 7.30 (dd, 4H), 7.55 (d, 2H).
25 To the pivaloyloxymethyl protected derivative (0.6 g) in ethanol (7 mL) was added water (1.4 mL) and NaOH powder (0.7 g). The mixture was stirred overnight. Ethanol was evaporated *in vacuo* and water (25 mL) was added. The precipitated crystalline product was filtered off and washed with water. After drying overnight at 70-80 °C *in vacuo* 350 mg of pure sodium salt of the title compound remained. Mp :
30 178-81 °C ¹H NMR (DMSO-*d*₆) δ 1.30-1.90 (m, 2H), 2.15-2.35 (m, 4H), 2.60-2.85 (m, 2H), 3.25 (t, 1H), 3.40 (t, 2H), MS *m/z* (%): 415 (MH⁺, 4 %), 203 (100 %), 183 (63 %), 138 (42 %), 67 (30 %).

The following compounds were prepared in a corresponding way :

(*R,S*)-4-[*N*-[bis(4-chlorophenyl)methyl-2-oxyethyl]-*N*-methylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, sodium salt 16b. Mp : 201-203 °C (water / ethanol). ¹H NMR (DMSO-*d*₆) δ 1.40-1.60 (m, 2H), 1.60-1.70 (m, 1H), 1.75-1.85 (m, 1H), 2.20-2.35 (m, 2H), 2.20 (s, 3H), 2.65-2.90 (m, 2H), 3.35 (t, 1H), 3.35-3.50 (m, 2H), 5.55 (s, 1H), 7.45 (s, 8H). MS *m/z* (%): 447 (MH⁺), 235 (78 %), 165 (57 %), 138 (100 %), 67 (56 %).

(*R,S*)-4-[*N*-(diphenylmethyl-2-oxyethyl)-*N*-methylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrochloride 16c. Mp : 108-113 °C (amorphous) (diethyl ether). ¹H NMR (DMSO-*d*₆) δ 1.70-2.10 (m, 2H), 2.10-2.30 (m, 2H), 2.60-2.75 (m, 2H), 2.80 (broad s, 3H), 3.40-3.60 (m, 2H), 3.80 (broad t, 2H), 4.55 (broad s, 1H), 5.55 (s, 1H), 7.20-7.45 (m, 10H). MS *m/z* (%): 379 (MH⁺, 4%), 167 (100%), 152 (74%), 138 (36%), 67 (32%).

In a similar manner, treatment of (*R,S*)-4-[*N*-(4,4-bis[3-methylthien-2-yl]-3-butenyl)-*N*-methylamino]-3-pivaloyloxymethyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole with NaOH in aqueous ethanol followed by acidification with 4M HCl and extraction with methylene chloride gave (*R,S*)-4-[*N*-(4,4-bis[3-methylthien-2-yl]-3-butenyl)-*N*-methylamino]-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole hydrochloride 16d. Mp 135-138 °C (from acetonitrile-ether). ¹H-NMR (DMSO-*d*₆) δ 1.95 (s,3H), 2.0 (s,3H), 1.95-2.20 (m,4H), 2.55-2.90 (m,6H), 3.35 (s,3H), 4.40-4.50 (m,1H), 6.0 (t,1H), 6.85 (d,1H), 6.95 (d,1H), 7.35 (d,1H), 7.55 (d,1H).

Example 17

(*R,S*)-3-Ethoxy-4-dimethylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 17a

A mixture of (*R,S*)-3-ethoxy-4-amino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 3a (526 mg) and 1 M sodium hydroxide (8 mL) was extracted with methylene chloride (3x10 mL). The combined extracts were dried and evaporated. Formic acid (3 mL), 30 % formaldehyde (3 mL) and sodium formate (3 g) were added to the residue and the mixture was stirred at 60 °C for 3 h and at 100 °C for 20 h. After evaporation, water (20 mL) was added to the residue and pH was adjusted to 10 with 2 M sodium hydroxide. The mixture was extracted with methylene chloride

(3x40 mL) and the combined extracts were dried and evaporated. The residue was dissolved in ether (25 mL) and an excess of 33 % hydrobromic acid in glacial acetic acid was added dropwise to precipitate the title compound (422 mg). Recrystallization from acetonitrile-ether gave an analytical pure compound: mp 158-160 °C. ¹H NMR (60 MHz, D₂O) δ 1.60 (t, 3H), 2.20 (m, 4H), 2.85 (m, 2H), 3.05 (s, 6H), 4.50 (q, 2H), 4.55 (m, 1 H).

Example 18

(*R,S*)-4-Dimethylamino-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide 18a (Method c))

A solution of (*R,S*)-3-ethoxy-4-dimethylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole hydrobromide 17a (250 mg) and 33 % hydrobromic acid in glacial acetic acid (5 mL) was stirred at 80 °C for 25 min. The mixture was evaporated and 33 % hydrobromic acid in glacial acetic acid (5 mL) was added to the residue. After being stirred at 80 °C for 25 min, the mixture was evaporated and the residue recrystallized from acetonitrile-ethanol-ether to give the title compound (199 mg): mp 183-186 °C. The compound crystallized with 0.33 mol of water. ¹H NMR (60 MHz, D₂O) δ 2.35 (m, 4H), 3.00 (m, 2H), 3.25 (s, 6H), 4.70 (m, 1H).

Example 19

(*R,S*)-3-Benzyloxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrochloride 19a

A mixture of (*R,S*)-3-hydroxy-4-(*N*-methyl-*N*-*tert*-butoxycarbonylamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole 12a (1.20 g) and potassium carbonate (1.23 g) in DMF (25 mL) was stirred at 40 °C for 45 min. Benzyl bromide (1.59 mL) was added and stirring at 40 °C was continued for 20 h. The reaction mixture was evaporated and water (25 mL) was added to the residue. Extraction with methylene chloride (3x 50 mL), drying and evaporation gave an oil. Flash chromatography on silica gel (eluent: toluene containing ethyl acetate (0-75%)) eluted (*R,S*)-3-benzyloxy-4-(*N*-methyl-*N*-*tert*-butoxycarbonylamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole (oil, 620 mg). A mixture of the said compound (620 mg), 1M HCl (15 mL) and ethanol (20 mL) was stirred at 45 °C for 80 min. Evaporation and recrystallization of the residue from acetonitrile-ether gave the title compound (460 mg). Mp 156-159 °C.

The following compound was synthesized correspondingly:

(*R,S*)-3-Benzoyloxy-4-amino-4,5,6,7-tetrahydro-1,2-benzioxazole hydrochloride 19b.
Mp 168-170 °C.

5

Example 20

(*R,S*)-3-Hydroxy-4-[*N*-(4,4-diphenyl-3-butenyl)methylamino]-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrochloride 20a (Method c))

A mixture of Compound 19a (442 mg), potassium carbonate (622 mg), sodium
10 iodide (50 mg) and 4,4-diphenyl-3-butenyl bromide (646 mg) in DMF (8 mL) was
stirred at 120 °C for 24 h. A solution of 4,4-diphenyl-3-butenyl bromide (500 mg) in
DMF (3 mL) was added and stirring at 120 °C was continued for 24 h. The reaction
mixture was evaporated and water (20 mL) was added to the residue. Extractions
with ether (3 x 25 mL), drying and evaporation gave an oil. Flash chromatography
15 on silica gel (eluent: toluene-ethyl acetate (0-100%)) gave (*R,S*)-3-benzoyloxy-4-[*N*-
(4,4-diphenyl-3-butenyl)methylamino]-4,5,6,7-tetrahydro-1,2-benzisoxazole (590
mg) as a light-yellow oil. A solution of the said compound (590 mg) in ethanol (16
mL) and concentrated hydrochloric acid (8 mL) was refluxed for 3 days. Evaporation
and recrystallization of the residue from acetone-ethanol-ether gave the title com-
20 pound (212 mg). Mp. 119-120 °C. ¹H NMR (D₂O and DMSO-d₆) δ 1.60-1.98 (m,
4H), 2.35-2.6 (m, 2H), 2.58 (s, 3H), 3.10 (m, 2H), 4.18 (m, 1H), 5.91 (t, 1H), 7.0-7.4
(m, 10H). The ¹H NMR spectrum showed the presence of 0.75 equivalent of
ethanol.

25 The following compounds were synthesized correspondingly:

(*R,S*)-3-Hydroxy-4-[*N*-(4,4-diphenyl-3-butenyl)amino]-4,5,6,7-tetrahydro-1,2-benzioxazole hydrochloride hydrate 20b. Mp 140-143 °C (from acetonitrile). ¹H-NMR (D₂O and DMSO-d₆) δ 1.80-2.20 (m, 4H), 2.40-2.60 (m, 4H), 3.10-3.30 (m, 2H),
4.15-4.30 (m, 1H), 6.10 (t, 1H), 7.10-7.45 (m, 10H).

30 **(*R,S*)-3-Hydroxy-4-[*N*-(4,4-bis[3-methylthien-2-yl]-3-butenyl)amino]-4,5,6,7-tetrahydro-1,2-benzisoxazole hydrochloride 20c.** Mp 188-191 °C (from ethanol-acetonitrile). ¹H-NMR (D₂O and DMSO-d₆) δ 1.85-2.15 (m, 4H), 2.05 (s, 3H), 2.08 (s, 3H), 2.45-2.90 (m, 4H), 3.15-3.30 (m, 2H), 4.20-4.35 (m, 1H), 6.10 (t, 1H), 6.90

(m,1H), 7.0 (m,1H), 7.25 (t,1H), 7.40 (t,1H).

Example 21

2-Benzylamino-3,4,5,6-tetrahydrobenzamide 21a

- 5 A mixture of cyclohexanone-2-carboxamide (U.S. 4169952 to du Pont de Nemours, (1979)) (10.0 g), benzylamine (8.4 g), toluene (35 mL) and molecular sieves (Union Carbide 3A, 2 g) was refluxed for 2 h in a Dean-Stark water separator. The reaction mixture was filtered, and the filtrate was evaporated. The residue was crystallized from light petroleum to give the title compound (16 g). Mp 73-74 °C.

10

Example 22

3-Hydroxy-4,5,6,7-tetrahydro-1,2-benzisothiazole 22a

- To a solution of 2-benzylamino-3,4,5,6-tetrahydrobenzamide 21a (15 g) in glacial acetic acid (100 mL) was added excess hydrogen sulfide at 80 °C for 4 h. The
15 reaction mixture was evaporated and ether was added to the residue which afforded crystallisation. The crystals were dissolved in ethyl acetate (30 mL) and a solution of bromine (8.3 mL) in ethyl acetate (30 mL) was dropwise added at room temperature. The mixture was stirred for 20 h at room temperature and evaporated. Column chromatography on silica gel (eluent: ethyl acetate-ethanol 1:1 containing 1%
20 glacial acetic acid) gave the title compound (3.3 g): mp 157-158 °C.

Example 23

3-Chloro-4,5,6,7-tetrahydro-1,2-benzisothiazole 23a

- A mixture of 3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisothiazole 22a (4.74 g), pyridi-
25 nium hydrochloride (12.7 g), phosphoric acid (2.1 g) and phosphorus oxychloride (25 mL) was stirred at 90 °C for 5 h. The reaction mixture was evaporated and ethyl acetate (130 mL) was added to the residue. A saturated solution of sodium hydrogencarbonate (130 mL) was added and after 10 min of stirring the phases were separated. The aqueous phase was extracted with ethyl acetate (2 x 150 mL) and
30 the combined organic phases were dried and evaporated to give an oil. Column chromatography on silica gel (eluent: toluene-ethyl acetate 1:1) gave the title compound as a yellow oil (2.8 g).

Example 24

3-Chloro-4,5,6,7-tetrahydro-1,2-benzisothiazol-4-one 24a.

A solution of sodium dichromate (4.4 g) in glacial acetic acid (30 mL) was added dropwise over 1 h to a solution of 3-chloro-4,5,6,7-tetrahydro-1,2-benzisothiazole
5 **23a** (2.7 g) and concentrated sulfuric acid (1.8 mL) in glacial acetic acid (80 mL). The reaction mixture was stirred at room temperature for an additional 2 h, and neutralized with a saturated solution of sodium hydrogencarbonate. Extraction with ether (3 x 150 mL), drying and evaporation gave an oil. Column chromatography on silica gel (eluent: toluene-ethyl acetate 1 :1) eluted first 3-chloro-4,5,6,7-tetrahydro-
10 **1,2-benzisothiazol-7-one** (680 mg). The later fractions contained the title compound (780 mg). Mp 84-85 °C.

Example 25

3-Methoxy-4,5,6,7-tetrahydro-1,2-benzisothiazol-4-one 25a

15 A mixture of 3-chloro-4,5,6,7-tetrahydro-1,2-benzisothiazol-4-one **24a** (600 mg) and a solution of sodium (506 mg) in methanol (22 mL) was stirred at 90 °C for 1 h. The reaction mixture was evaporated and water (20 mL) was added to the residue. Extractions with methylene chloride (3 x 30 mL), drying and evaporation gave an oil. Column chromatography on silica gel (eluent: toluene-ethyl acetate 4:1) gave the
20 title compound (251 mg). Mp 45-46 °C.

Example 26 (Method c)

(*R,S*)-4-Amino-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisothiazole, hydrobromide **26a**
To a solution of 3-methoxy-4,5,6,7-tetrahydro-1,2-benzisothiazol-4-one **25a** (185
25 mg) and ammonium acetate (780 mg) in methanol (7 mL) was portionwise added sodium borohydride (44 mg). The mixture was stirred at room temperature for 48 h and acidified with concentrated hydrochloric acid. The mixture was evaporated and water (3 mL) was added to the residue. The aqueous solution was washed with ether (3 x 15 mL) and solid potassium hydroxide was added until pH > 10. Extrac-
30 tion with ether (3 x 15 ml), drying and evaporation gave an oil. The oil was dissolved in ethanol and excess of hydrochloric acid in ethyl acetate was added to precipitate (*R,S*)-3-methoxy-4-amino-4,5,6,7-tetrahydro-1,2-benzisothiazole, hydrochloride (68 mg). A solution of 33 % hydrobromic acid in glacial acetic acid (3 mL) was added to the said hydrochloride (60 mg) and the mixture was stirred at room

temperature for 48 h. Evaporation and recrystallizations of the residue (methanol-ether) gave the title compound (28 mg). Mp 160-165 °C. ¹H NMR (D₂O) δ 1.78-2.04 (m, 3H), 2.05-2.28 (m, 1H), 2.72-2.87 (m, 2H), 4.23-4.39 (m, 1 H).

5 **Example 27**

(R,S)-3-Hydroxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisothiazole, hydrobromide **27a**

(R,S)-3-Methoxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisothiazole was synthesized as described for Compound 7a in Example 7 using Compound 25a (200 mg)
10 in methanol (5 mL), a 33% solution of methylamine (217 µL) in ethanol, molecular sieves (3 Å) powder and sodium cyanoborohydride (234 mg). The resulting oil was dissolved in ether and excess of hydrochloric acid in ethyl acetate was added to precipitate *(R,S)*-3-methoxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisothiazole hydrochloride (172 mg), mp 146-148. A solution of hydrobromic acid in glacial acetic
15 acid (33%, 6mL) was added to 100 mg thereof and the mixture was stirred at room temperature for 48 hours. Evaporation and recrystallization of the residue from methanol-ether gave the title compound (53 mg), mp 192 °C. (dec.). ¹H-NMR (200 MHz, D₂O): δ 1.82-2.24 (m, 4H), 2.72 (s, 3H), 2.68-2.92 (m, 2H), 4.16-4.30 (1H).

20 **Example 28 (method e)**

(R,S)-3-Hydroxy-N-methyl-N-(4-phenyloxycarbonyl)amino-4,5,6,7-tetrahydro-1,2-benzisoxazole **28a**

To a solution of *(R,S)*-3-hydroxy-N-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrobromide (1.0 g) and triethylamine (1.8 mL) in dry dimethylformamide (10
25 mL) kept at 0 °C was added dropwise phenyl chloroformate (0.7 g) in THF (10 mL). After stirring overnight at room temperature water (200 mL) and ethyl acetate (100 mL) were added. PH was adjusted to 1-2 by addition of hydrochloric acid. The organic phase was separated and worked up as above. The title compound crystallized by stirring with diethyl ether. Yield 1.0 g. Mp 147-149 °C. ¹H NMR
30 (CDCl₃) δ 1.70-1.95 (m, 2H), 2.05-2.20 (m, 2H), 2.6-2.7 (m, 2H), 2.85 (s, 1H), 2.95 (s, 2H), 5.35 (broad s, 1H), 7.10-7.40 (m, 5H), 8.05 (broad s, 1H). MS m/z (%): 289 (MH⁺, 10%), 138 (100%), 95 (24%), 67 (77%).

Example 29**4,4-Diphenyl-1-butyliodide, 29a**

To a suspension of magnesium turnings (15 g) in dry THF (75 mL) was added a little bromobenzene (0.5 g). After an initial exothermic reaction had started, the mixture was heated to reflux and a solution of bromobenzene (90 g) in dry THF (200 mL) was added dropwise during 30 minutes. The mixture was heated for additionally 1.5 hours. The mixture was cooled to room temperature and excess Mg was filtered off in an inert atmosphere. A solution of 4-chlorobutyric acid methylester (40 g) in dry THF (160 mL) was added dropwise at 15-25 °C. After further stirring for 30 minutes, the mixture was poured into an aqueous solution of NH₄Cl and ice. Diethyl ether (500 mL) was added. The organic phase was worked-up according to the standard workup procedure. Yield 65 g of crude 4-chloro-1,1-diphenylbutan-1-ol. The crude alcohol (30 g) was dissolved in a mixture of glacial acetic acid (60 mL) and 57 % aqueous iodic acid (60 mL). Red phosphorus (5 g) was added and the mixture was refluxed for 6 hours. After slowly cooling to room temperature, the mixture was poured into water and diethyl ether. The organic phase was worked up following the standard procedure above yielding 39 g of the title butyliodide 29a as an oil, which was used without further purification.

Example 30**4,4-Diphenyl-1-butylamine, hydrochloride 30a**

To a solution of 4,4-Diphenyl-1-butyliodide, 29a (20 g) in dry DMF (150 mL) was added sodium azide (10 g). After reflux for 1.5 hours, the mixture was cooled to room temperature and subsequently poured into diethyl ether and water. The organic phase was worked up following the standard procedure above. Yield of 4,4-diphenyl-1-butylazide 14 g. The crude azide (10 g) was dissolved in ethanol (150 mL), water (10 mL) and glacial acetic acid (10 mL). 2% Palladium on Carbon black was added and the mixture was hydrogenated in a Parr apparatus at 3 ato for 1.5 hours. The catalyst was filtered off and the solvents evaporated *in vacuo*. The remaining visceous oil was dissolved in water and dichloromethane. Aqueous NaOH solution was added to adjust the pH to >11. The organic phase was separated and worked according to the standard procedure above. The hydrochloric salt was prepared by addition of HCl to a solution of the free amino compound in diethyl ether. Yield 3.4 g. Mp 172-175 °C.

The following compounds were prepared in a similar manner except that the amines were not precipitated as hydrochloric salts.

- 5 4,4-Di-(2-tolyl)-1-butylamine, **30b**. Prepared via the 4,4-di-(2-tolyl)-1-butyliodide, **33a** and isolated as an oil. ¹H NMR (CDCl₃) δ 1.20 (s broad, 2H), 1.42-1.58 (m, 2H), 1.87-2.03 (m, 2H), 2.27 (s, 6H), 2.70 (t, 2H), 4.23 (t, 1H), 7.05-7.16 (m, 8H)
- 4,4-Di-(2-tolyl)-3-butenylamine, **30c**. Prepared via the 4,4-di-(2-tolyl)-3-butenyliodide **36a** and isolated as an oil. ¹H NMR (CDCl₃) δ 1.20 (s broad, 2H), 2.11 (s, 3H), 2.18
- 10 (q, 2H), 2.26 (s, 3H), 2.77 (t, 2H), 5.75 (t, 1H), 7.02-7.18 (m, 8H).

Example 31

(*R,S*)-3-Benzoyloxymethyloxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hemioxalate **31a**

- 15 A mixture of benzoylchloride (22 g), paraformaldehyde (6 g), and a few crystals of ZnCl₂ were heated at 100-105 °C for 2.5 hours. Crude benzoyloxymethylchloride was isolated by elution of the reaction mixture through silica gel (eluted with heptane / dichloromethane 1:1). Yield 12 g. All of the crude chloride was dissolved in acetone (100 mL) and sodiumiodide was added. After reflux for 7 hours acetone
- 20 was evaporated *in vacuo* and crude benzoyloxymethyliodide was isolated by extraction with diethyl ether from water containing sodium thiosulfate by standard work up procedure. The pure iodide derivative was isolated by column chromatography on silica gel (eluted with heptane / dichloromethane 1:1). Yield 5 g. To a solution of (*R,S*)-4-(*N-tert*-Butyloxycarbonyl-*N*-methylamino)-3-hydroxy-4,5,6,7-
- 25 tetrahydro-1,2-benzisoxazole **12a** (1.5 g) in acetone (30 mL) in an inert nitrogen atmosphere was added potassium *tert*-butoxide (0.8 g). The mixture was cooled to 10 °C and a solution of benzoyloxymethyliodide (2.5 g) was added. The mixture was stirred overnight at room temperature in the dark. Acetone was evaporated *in vacuo* and the remaining crude product was purified by column chromatography on
- 30 silica gel (eluted with heptane / ethyl acetate 7 : 3). Yield 1.3 g as an oil. The *tert*-BOC protecting group was removed as described above by treatment with trifluoroacetic acid in dichloromethane. After evaporation of the solvents the crude product was dissolved in ice cooled water, pH was adjusted to 9-10 by addition of

diluted aqueous potassium carbonate. Extraction with diethyl ether and work up of the organic phase using the standard procedure above afforded 1 g of crude title product. The hemioxalate salt **31a** crystallized from acetone. Yield 900 mg. Mp 188-189 °C. ¹H NMR (DMSO-d₆) δ 1.60-2.05 (m, 4H), 2.45 (s, 3H), 2.55-2.80 (m, 2H), 3.85 (t, 1H), 6.15 (s, 1H), 7.60 (t, 2H), 7.65-7.75 (m, 1H), 8.00 (dd, 2H). MS m/z (%): 303 (MH⁺), 198 (17%), 105 (100%).

In a corresponding manner were prepared the following 3-substituted derivatives :

(*R,S*)-4-Methylamino-3-(2,4,6-trimethylbenzoyloxymethyloxy)-4,5,6,7-tetrahydro-1,2-benzisoxazole, hemioxalate **31b**. Mp 217 °C (acetone). ¹H NMR (DMSO-d₆) δ 1.60-2.05 (m, 4H), 2.20 (s, 6H), 2.25 (s, 3H), 2.45 (s, 3H), 2.55-2.80 (m, 2H), 3.90 (t, 1H), 6.10 (s, 1H), 6.95 (s, 2H). MS m/z (%): 345 (MH⁺), 240 (100%), 147 (94%).

(*R,S*)-4-Methylamino-3-(2,6-difluorobenzoyloxymethyloxy)-4,5,6,7-tetrahydro-1,2-benzisoxazole, hemioxalate **31c**. Mp 196-197 °C (acetone) ¹H NMR (DMSO-d₆) δ 1.60-2.05 (m, 4H), 2.45 (s, 3H), 2.55-2.80 (m, 2H), 3.85 (t, 1H), 6.15 (s, 2H) 7.30 (t, 2H), 7.70-7.85 (m, 1H). MS m/z (%): 339 (MH⁺), 234 (17%), 141 (100%).

(*R,S*)-4-Methylamino-3-(2-methylbenzoyloxymethyloxy)-4,5,6,7-tetrahydro-1,2-benzisoxazole, hemioxalate **31d**. Mp 195-196 °C. ¹H NMR (DMSO-d₆) δ 1.60-2.05 (m, 4H), 2.40 (s, 3H), 2.50 (s, 3H), 2.55-2.80 (m, 2H), 3.85 (t, 1H), 6.10 (s, 1H), 7.35-7.45 (m, 2H), 7.55 (t, 1H), 7.90 (d, 1H). MS m/z (%): 317 (MH⁺, 2%), 212 (19%), 119 (100%), 91 (74%).

Example 32

(*R,S*)-4-[N-3-(Phenothiazin-10-yl)propane-1-yl]amino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, **32a**

A solution of 3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol-4-one **2a** (654 mg) and 3-(phenothiazin-10-yl)-1-propylamine (prepared as described in EP-A-0-200-450) (1.02 g) in toluene (130 mL) was boiled under reflux (105 °C) for 6 hrs. p-Toluenesulfonic acid, monohydrate (10 mg) was added to the boiling solution which was boiled for additionally 16 hrs. This solution was cooled to 5 °C and was then added to a solution of NaCNBH₄ (635 mg) in methanol (50 mL) at 10 °C. The resulting reaction mixture was stirred for 20 min, before addition of further NaCNBH₄ (500 mg). The

reaction mixture was stirred for additional 10 min at 10 °C. The reaction mixture was poured onto water and the phases were separated. The aqueous phase was extracted with ethyl acetate (3 x 250 mL). The combined organic phases were washed with a saturated aqueous NaCl solution, dried (Na₂SO₄) and evaporated *in vacuo*. The residue was subjected to CC (*n*-heptane/ethyl acetate - 1:1) to give the title compound as an oil (640 mg). ¹H NMR (CDCl₃) δ 1.33 (t,3H), 1.50-2.00 (m,7H), 2.40-2.60 (m,2H), 2.70-2.84 (m,2H), 3.65 (t,1H), 3.97 (t,2H), 4.26 (q,2H), 6.85-6.95 (m,4H), 7.08-7.18 (m,4H)

10 Example 33

2,2-Di-(2-tolyl)-tetrahydrofuran, 33a.

To a suspension of magnesium turnings (33 g) in dry THF (150 mL) was added 2-bromotoluene (4 mL). The reaction mixture was heated to reflux and an exothermic reaction started. The heating mantle was removed and 2-bromotoluene (137 mL) in dry THF (500 mL) was added dropwise over an hour at reflux temperature (exothermic reaction). The resulting reaction mixture was boiled under reflux for additionally 1.5 hrs. The mixture was cooled to room temperature and excess Mg was filtered off in an inert atmosphere. A solution of 4-chlorobutyric acid methylester (56.4 g) in dry THF (200 mL) was added dropwise at 20 °C. The reaction mixture was stirred at room temperature for additionally 1 hr and was then poured into an aqueous solution of NH₄Cl and ice. The organic phase was worked-up according to the standard workup procedures. After evaporation of the organic solvent the residue was suspended in a mixture of *n*-heptane/ethyl acetate = 4/1. Filtration of the resulting crystals afforded 2,2-di-(2-tolyl)-tetrahydrofuran 33a (32.5 g). ¹H NMR (CDCl₃) δ 1.96 (s,6H), 1.96-2.10 (m,2H), 2.57 (t,2H), 4.02 (t,2H), 7.00-7.07 (m,2H), 7.07-7.23 (m,4H), 7.57-7.65 (m,2H)

Example 34

4,4-Di-(2-tolyl)-1-butyliodide, 34a

30 The crude 2,2-di-(2-tolyl)-tetrahydrofuran 33a (28 g) was dissolved in acetic acid (250 mL). 5 % palladium on Carbon black (3 g) was added and the mixture was hydrogenated in a Parr apparatus at 3 ato at 55 °C for 5 hrs. The catalyst was filtered off and the solvent was evaporated *in vacuo*. The remaining oil was

subjected to CC (*n*-heptane/ethyl acetate - 15:1) to give 4,4-di-(2-tolyl)-1-butanol (17 g). A solution of 4,4-di-(2-tolyl)-1-butanol (19 g) in acetic acid (400 mL) was boiled under reflux for 3 hrs. The cooled solution was evaporated *in vacuo* to give 4,4-di-(2-tolyl)-1-butyl acetate (17 g) as an oil. A solution of 4,4-di-(2-tolyl)-1-butyl acetate (9.2 g) in 57 % aqueous iodic acid (150 mL) was boiled under reflux for 3 hrs. The cooled solution was poured into a mixture of ice and water and the aqueous phase was extracted with diethyl ether. The combined organic phases were washed with water and a saturated aqueous solution of NaCl, dried (Na₂SO₄) and evaporated *in vacuo* to give the title compound 34a (11.7 g) as an oil, which was used without further purification. ¹H NMR (CDCl₃) δ 1.80-2.10 (m,4H), 2.28 (s,6H), 3.17 (t,2H), 4.26 (t,1H), 7.12 (s,8H)

Example 35

(*R,S*)-4-[N-[4,4-Di-(2-tolyl)butan-1-yl]amino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol, 35a

The title compound was prepared according to the procedure described in Synlett (1995) 1079-1080 using 4,4-di-(2-tolyl)-1-butylamine 30b (1.7 g), 3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol-4-one 2a (1.0 g), titanium(IV)isopropylate (4.3 mL), NaCNBH₃ (0.6 g) and ethanol (20 mL). The yield of the title compound was 1.0 g which was isolated as an oil. ¹H NMR (CDCl₃) δ 1.32 (t,3H), 1.40-1.85 (m,7H), 1.85-2.05 (m,3H), 2.27 (s,6H), 2.45-2.75 (m,4H), 3.68 (t,1H), 4.27 (q,2H), 7.05-7.15 (m,8H)

(*R,S*)-4-[N-1,1-di-(2-tolyl)but-1-en-4-yl]amino]-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol, 35b

The title compound was prepared according to the procedure described in Synlett (1995) 1079-1080 using 4,4-di-(2-tolyl)-3-butenylamine, 30c (3.3 g), 3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazol-4-one 2a (2.0 g), titanium(IV)isopropylate (8.2 mL), NaCNBH₃ (1.4 g) and ethanol (40 mL). The yield of the title compound was 1.9 g which was isolated as an oil. ¹H NMR (CDCl₃) δ 1.33 (t,3H), 1.40-2.05 (m,5H), 2.10 (s,3H), 2.17-2.30 (m,5H), 2.40-2.70 (m,2H), 2.76 (t,2H), 3.67 (t,1H), 4.27 (q,2H), 5.80 (t,1H), 7.03-7.17 (m,8H)

Example 36**4,4-Di-(2-tolyl)-3-butenyliodide, 36a**

A solution of 2,2-di-(2-tolyl)-tetrahydrofuran 33a (40 g) in 57 % aqueous iodic acid (250 mL) was boiled under reflux for 30 min. The cooled solution was extracted with diethyl ether. The combined organic phases were washed with water and a saturated aqueous solution of NaCl, dried (Na₂SO₄) and evaporated *in vacuo*. The residue was subjected to CC (*n*-heptane/ethyl acetate 15:1) to give the title compound 36a as an oil (44 g). ¹H NMR (CDCl₃) δ 2.10 (s,3H), 2.30 (s,3H), 2.65 (q,2H), 3.19 (t,2H), 5.73 (t,1H), 7.05-7.22 (m,8H)

10

Example 37**(*R,S*)-3-Benzenesulfonyloxy-4-(*N*-*tert*-butyloxycarbonyl-*N*-methyamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole, 37a**

A solution of (*R,S*)-4-(*N*-*tert*-butyloxycarbonyl-*N*-methyamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 12a (9 g), triethylamine (7.5 mL) and dry THF (350 mL) was cooled to 0 °C. Benzenesulfonylchloride (5.2 mL) in dry THF (100 mL) was added dropwise at 0 °C. The resulting reaction mixture was stirred for 3 hrs at 0 °C and for 60 hrs at 22 °C. The formed precipitate was removed by filtration and the organic solvent was evaporated *in vacuo*. The residue was subjected to CC (*n*-heptane/ethyl acetate 1:1) to give the title compound 37a (5.5 g) as an oil. ¹H NMR (CDCl₃) δ 1.40-1.94 (m,11H), 1.94-2.14 (m,2H), 2.60-2.72 (m,5H), 4.95-5.36 (m,1H), 7.58 (t,2H), 7.70 (t,1H), 8.02 (d,2H)

20

Example 38**(*R,S*)-3-Benzenesulfonyloxy-4-methyamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrochloride, 38a**

To a solution of (*R,S*)-3-benzenesulfonyloxy-4-(*N*-*tert*-butyloxycarbonyl-*N*-methyamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole, 37a (5.4 g) in dry diethyl ether (100 mL) was added a saturated solution of HCl (g) in dry diethyl ether (50 mL) and the resulting mixture was stirred for 48 hrs at room temperature. The reaction mixture was evaporated *in vacuo* and the residue was suspended in dry diethyl ether (100 mL). The resulting crystals were collected by filtration and dried to give the title compound 38a (4.0 g). Mp 162-163 °C (dec.). ¹H NMR (CDCl₃) δ 1.75-2.15 (m,2H),

30

2.46-3.04 (m,7H), 4.25 (dd,1H), 7.60 (t,2H), 7.76 (t,1H), 8.03 (d,2H)

Example 39

(*R,S*)-3-Benzenesulfonyloxy-4-[N-methyl-N-(2-propynyl)amino]-4,5,6,7-tetrahydro-
5 1,2-benzisoxazole, 39a

A suspension of (*R,S*)-3-benzenesulfonyloxy-4-methylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole, hydrochloride, 38a (3.5 g) and K₂CO₃ (4.2 g) in methylisobutylketone (200 mL) was heated to 50 °C for 1 hr. 3-Brom-1-propyne (1.8 mL) in methylisobutylketone (50 mL) was added dropwise at 50 °C. The resulting reaction mixture
10 was stirred for additionally 2 hrs at 50 °C and then for 72 hrs at 117 °C (reflux temperature). The cooled mixture was filtrated and the solvent was evaporated *in vacuo*. The residue was subjected to CC (*n*-heptane/ethyl acetate/triethylamine 14:5:1) to give the title compound 39a (0.92 g). Mp 85-87 °C.

¹H NMR (CDCl₃) δ 1.54-2.15 (m,4H), 2.25 (t,1H), 2.33 (s,3H), 2.52-2.77 (m,2H),
15 3.40 (t,2H), 3.79 (t,1H), 7.57 (t,2H), 7.71 (t,1H), 8.05 (dd,2H)

Example 40

(*R,S*)-4-(*N*-*tert*-butyloxycarbonyl-N-methylamino)-3-ethoxymethyloxy-4,5,6,7-tetrahydro-1,2-benzisoxazole, 40a

20 K₂CO₃ (2.2 g) was added to at solution of (*R,S*)-4-(*N*-*tert*-butyloxycarbonyl-N-methylamino)-3-hydroxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 12a (3.5 g) in acetone (200 mL). The resulting suspension was heated to reflux temperature. Ethoxymethylchloride (1.5 g) in acetone (100 mL) was aded dropwise at 65 °C and the resulting reaction mixture was boiled under reflux for 2 hrs. The cooled suspen-
25 sion was filtrated and the organic solvent was evaporated *in vacuo*. The residue was subjected to CC (*n*-heptane/ethyl acetate/methanol 10:10:1) to give the title compound 40a (0.5 g) as an oil. ¹H NMR (CDCl₃) δ 1.24 (t,3H), 1.48 (s,9H), 1.50-2.10 (m,4H), 2.50-2.70 (m,5H), 3.75 (q,2H), 4.90-5.13 (s broad, 1H), 5.20-5.44 (m,2H).

30

Example 41

(*R,S*)-3-Hydroxy-4-benzylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole hydrobromide 41a.

Benzaldehyde (0.31 mL) was added to a solution of (*R,S*)-4-amino-3-ethoxy-4,5,6,7-tetrahydro-1,2-benzisoxazole 3a (from 0.53 g and the hydrobromide) in ethanol (10 mL). The mixture was stirred at room temperature for 20 min and then cooled in an ice-bath. Sodium borohydride (0.15 g) was added and the mixture
5 was stirred at 0 °C for 30 min and at room temperature for 20h. After evaporation, water (10 mL) was added and the mixture was acidified with 4 M HCl. The aqueous solution was washed with ether (2x10 mL) and made basic by addition of 4 M NaOH. Extraction with methylene chloride (3x15 mL), drying and evaporation afforded (*R,S*)-3-ethoxy-benzylamino-4,5,6,7-tetrahydro-1,2-benzisoxazole (0.31 g).
10 The hydrochloride was prepared by adding a solution of HCl in ethyl acetate and precipitating the hydrochloride by addition of ether. Mp 192-194 °C.

Treatment of said compound with HBr as described in Example 8a gave 41a. Mp 202-205 °C. ¹H-NMR (D₂O) δ 1.95-2.25 (m,4H), 2.60-2.90 (m,2H), 4.45 (s,2H),
15 4.35-4.60 (m,1H), 7.55 (s,5H).

The following compound was synthesized correspondingly:

(*R,S*)-3-Hydroxy-4-(2,2-diphenylethylamino)-4,5,6,7-tetrahydro-1,2-benzisoxazole hydrobromide 41b. Mp 180-183 °C. ¹H-NMR (D₂O and DMSO-d₆ 1:1) δ 1.80-2.15
20 (m,4H), 2.45-2.75 (m,2H), 3.70-4.00 (m,2H), 4.30-4.65 (m,2H), 7.50 (s,10H).

Pharmacology

The compounds of the invention were tested in the following well known and recognised test models:
25

GABA Uptake Synaptosomes

The inhibition of GABA Uptake was performed by the method described by Falch et. al, *Drug Dev. Res.*, 1990,21, 169-188. By this method the overall, i.e. neuronal and
30 glial, inhibition of GABA uptake is determined. The results are shown in the following Table I.

Isoniazid Antagonism.

The test is a test for antagonism of isoniazide induced convulsions in mice.

The test substance is given *s.c.* to mice (male NMRI/BOM, body weight 20-25 g) and 30 min later 300 mg isoniazide is given *s.c.* Five mice are used per dose and a control group only receiving isoniazide is included. This dose of isoniazide induces intermittent tonic seizures.

The animals are placed individually in Macrolon type II cages and the time when convulsions first occur is recorded. The experiment is stopped after 90 min. The animals that have not had convulsions within 60 min. are recorded as + (protected). The results, i.e. the number of protected mice pr number of tested mice are stated in fractions as follows: 0/5, 1/5,....., 5/5. The ED₅₀ values, calculated by log probit analysis, are shown in Table I below.

Table I: Inhibition of GABA-uptake in synaptosomes and Isoniazid antagonism.

Comp No	GABA-Uptake Inhibition IC ₅₀ (μM)	Isoniazid ED ₅₀ (μmol/kg)
6a (+)or(S)-form	>300	nt
6b (-)or(R)-form	120	nt
6c (+)-form	42	> 320.
6d (-)-form	>300	nt
8a	65	>320
8b	0.24	110.
8c	120	> 120.
8d	210	> 140.
8e	100	>300
8f	100	320.
8g	0.17*	56.
8h	4.8*	150.
8i	0.14	310.
8j	1.1	NOT DETERM
8j	nt	68.
8l	nt	71.
8m	nt	27.
8n	nt	44.
8o	nt	50.
8p	nt	30.
9a	180	> 140.
11a	0.73	67.
11b	0.41	640.
14a	>300	86.
14b	>300	220.

Comp No	GABA-Uptake Inhibition IC ₅₀ (μM)	Isoniazid ED ₅₀ (μmol/kg)
15a	70	> 120.
15b	80	72.
15c	84*	> 120.
16a	4.5	55.
16b	4.9	81.
16c	0.36	160.
16d	0.31	63.
18a	280	NT
20a	0.37	97.
20b	0.14	nt
20c	0.17	nt
26 a	>300	NT
26a	nt	200.
31a	nt	210.
31b	300*	55.
31c	nt	210.
31d	nt	nt
41a	140	77.
41b	22	> 190.

* = Preliminary Data

Furthermore, some of the compounds of the invention have been tested for inhibition of neuronal and glial GABA-uptake, respectively, by the method described by Falch et. al, *Drug Dev. Res.*, 1990,21, 169-188. These tests showed that some of the 5 compounds predominantly inhibit the glial uptake. Finally, some of the compounds have shown effect in other convulsion models.

It appears from the above table that some of the compounds of the invention which do not inhibit the GABA uptake *in vitro* show effect in the Isoniazide antagonisme 10 test *in vivo* indicating that these compounds are prodrugs.

Accordingly, the compounds of the invention are considered useful in the treatment of diseases associated with GABA neurotransmission, e.g. as analgesic, antipsychotic, anticonvulsant, or anxiolytic drugs or as drugs for the treatment of muscular and movement disorders, such as antispastics or antisymptomatics in Huntington's disease or Parkinsonism.

Formulation Examples

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. Tablets may e.g. be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvants or additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredients.

Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to desired volume, sterilization of the solution and filling in suitable ampules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

Typical examples of recipes for the formulation of the invention are as follows:

1) Tablets containing 5.0 mg of Compound 20a calculated as the free base:

Compound 20a	5.0 mg
Lactose	60 mg
Maize starch	30 mg
Hydroxypropylcellulose	2.4 mg
Microcrystalline cellulose	19.2 mg
Croscarmellose Sodium Type A	2.4 mg
Magnesium stearate	0.84 mg

2) Tablets containing 50 mg of Compound 8b calculated as the free base:


49

	Compound 8b	0.5 mg
	Lactose	46.9 mg
	Maize starch	23.5 mg
	Povidone	1.8 mg
5	Microcrystalline cellulose	14.4 mg
	Croscarmellose Sodium Type A	1.8 mg
	Magnesium stearate	0.63 mg
3)	Syrup containing per millilitre:	
10	Compound 11a	25 mg
	Sorbitol	500 mg
	Hydroxypropylcellulose	15 mg
	Glycerol	50 mg
	Methyl-paraben	1 mg
15	Propyl-paraben	0.1 mg
	Ethanol	0.005 ml
	Flavour	0.05 mg
	Saccharin natrium	0.5 mg
	Water	ad 1 ml
20	4) Solution for injection containing per millilitre:	
	Compound 14a	10 mg
	Sorbitol	5.1 mg
	Acetic acid	0.08 mg
25	Water for injection	ad 1 ml


CLAIMS



- 5 1. An 4-aminotetrahydrobenzisoxazole or -isothiazole compounds having general formula I:



- 10 wherein R¹ and R² are independently selected from the group consisting of:
A) hydrogen, cycloalkyl, phenyl, or a group





where R⁷, R⁸ and R⁹ are independently selected from the group con-

- sisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy-lower alkyl, cycloalkyl, cycloalkyl-lower alkyl, phenyl, phenyl-lower alkyl, phenoxy-lower alkyl and heteroaryl selected from 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, imidazolyl, oxazolyl, pyrazolyl, pyrimidinyl, pyrrolyl, thiazolyl, 1,2,4-triazolyl, 2-pyridyl, 3-pyridyl or 4-pyridyl, any phenyl or heteroaryl group present optionally being substituted with one or two substituents selected from halogen, lower alkyl, lower alkoxy, hydroxy, nitro, lower alkylthio, lower alkylsulfonyl, lower alkyl- or di(lower)alkylamino, cyano, trifluoromethyl, trifluoromethylthio, trifluoromethylsulfonyloxy and phenyl which again may be substituted with halogen, methyl, methoxy or trifluoromethyl; and any alkyl group present being optionally substituted with one to three hydroxy groups which again are optionally esterified with a C₂₋₁₈ carboxylic acid;

- B) a group of general formula $Y-(CH_2)_r-(CHR^{11})_s-(CH_2)_t$ wherein Y is selected**



from the following groups (1) - (5):



5 wherein U is CHR^{10b} , NR^{10b} , O or S, U^1 is NR^{10b} , O or S; p is 0 or 1; q is 0 or 1; V is C or N and the dotted line represents a bond when V is C and no bond when V is N;

A is O, S, CH_2 , $(CH_2)_2$, $CH=CH-CH_2$, $(CH_2)_3$, $CH=CH$ or $O-CH_2$;

10 R^a and R^b each represent one or more substituents selected from halogen, lower alkyl, lower alkoxy, hydroxy, nitro, lower alkylthio, lower alkylsulfonyl, lower alkyl- or di(lower alkyl)amino, cyano, trifluoromethyl, trifluoromethylsulfonyloxy and trifluoromethylthio;

r and t are independently 0, 1, 2 or 3, s is 0 or 1, provided that when Y is a group (1) wherein U is NR^{10b} , O or S or a group (4), then $r + s + t$ is at least 2; and when Y is a group (3) or a group (5) where V is N, then $r + s + t$ is at least 1; 15 R^{7b} , R^{8b} and R^{9b} are as defined for R^7 , R^8 and R^9 in A) provided that they are not at the same time selected from hydrogen, lower alkyl, lower alkenyl and lower alkynyl;



20 R^{10b} and R^{11} are independently hydrogen, lower alkyl, lower alkenyl or lower alkynyl; and

25 C) a group of general formula $Y^c-(CH_2)_n-W-(CH_2)_m-$ wherein n is 1, 2 or 3, m is 2 or 3; W is O or S; and Y^c is a group (1) - (5) as defined in B) provided that n may not be 1, when Y is a group (1) or (4) wherein U or U^1 , respectively, is NR^{10b} , S or O;

D) a group of general formula $Y^d-O-(CH_2)_k-\overset{R^{11d}}{\underset{|}{CH}}-(CH_2)_l-$ wherein k is 0, 1, 2 or 3;



l is 0, 1, 2 or 3; R^{11d} is as defined for R¹¹ in B) above; and
Y is selected from the groups (2) and (5) as defined in B) above and the following groups (6) - (10):



Wherein p, q, R^a, R^b, and A are as defined in B) and R^{7d} - R^{11d} are as defined for R^{7b} - R^{10b} and R¹¹, respectively, under B) or

R¹ and R² together designate alkylene thereby forming a 4-8 membered nitrogen containing ring; or

one of R¹ and R² is a group R^{2'}OCO wherein R^{2'} is phenyl, or heteroaryl as defined in A) above or phenyl or such heteroaryl substituted with one or more substituents selected from halogen, lower alkyl, lower alkoxy, hydroxy, nitro, lower alkylthio, lower alkylsulfonyl, lower alkyl- or di(lower)alkylamino, cyano, trifluoromethyl, trifluoromethylthio, trifluoromethylsulfonyloxy, phenyl and phenyl substituted with halogen, methyl, methoxy or trifluoromethyl;

R³ - R⁶ are independently selected from hydrogen, hydroxy and lower alkyl, any alkyl group optionally being substituted with one or two hydroxy groups;

X is oxygen or sulfur;

P is hydrogen or a group ZR wherein

Z is CO, CS, SO₂ or CR^tR^u, R^t and R^u being hydrogen, hydroxy or lower alkyl, and if Z is CO or CS, then R is selected from the groups consisting of:



- i) hydrogen, C₁-C₁₈ alkyl, C₂-C₁₈ alkenyl, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkenyl or C₄-C₂₆ cycloalk(en)yl-alk(en)yl, optionally substituted with one or two hydroxy groups, or phenyl optionally substituted with one or more substituents selected from the group consisting of halogen, trifluoromethyl, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₅ acyloxy, or cyano; or
- 5 ii) QR^v, wherein Q is O or S and R^v is selected from the substituents defined for R under i) above; and
- iii) NR^xR^y, wherein R^x and R^y independently are selected from the substituents defined for R under i) above or R^x and R^y are combined to form a four to eight
- 10 membered heterocyclic ring containing from one to three nitrogen atoms and from zero to three oxygen or sulfur atoms; or
- if Z is CR¹R^u, R is selected from the groups consisting of:
- iv) a group QR^v as defined in ii);
- v) a group NR^xR^y as defined in iii); or
- 15 vi) a group OC(O)R^z, SC(O)R^z, OC(S)R^z or SC(S)R^z wherein R^z is selected from the substituents defined for R under i) above;
- if Z is SO₂, R is selected from group i) defined above;
- provided that P may not be hydrogen, when R¹ to R⁶ are all hydrogen, X is oxygen and the compound exists as a racemic mixture;
- 20 or a pharmaceutically acceptable salt thereof.

2. A compound of Claim 1, characterised in that X is oxygen.

3. A compound of Claim 2, characterised in that R¹ is lower alk(en/yn)yl optionally substituted with hydroxy which may be esterified with a C₂₋₂₂ carboxylic acid, or phenyl or phenyl-lower alkyl, optionally substituted with halogen, lower alkyl, lower alkoxy or trifluoromethyl, R¹ being preferably methyl, ethyl or phenylbutyl.

25

4. A compound of Claim 3, characterized in that R² is hydrogen, lower alkyl or a group R^{2'}OCO wherein R^{2'} is as defined in Claim 1.

30

5. A compound of Claim 3 or 4, characterized in that P is not hydrogen, P preferably being a group ZR as defined in Claim 1, wherein Z is CH₂ and R is a



group OC(O)R^z. as defined in vi) in Claim 1.

6. A compound of Claim 5, characterized in that R⁴ - R⁶ are hydrogen and R³ is hydrogen or lower alkyl, preferably hydrogen or methyl.

5

7. A compound of Claim 6, characterized in that it is a compound of formula 1a.

8. A compound of Claim 2, characterized in that R¹ is a group as defined under B), C) or D).

10

9. A compound of Claim 8, characterized in that R¹ is a group as defined in B) wherein Y is a group of formula (1), (2) or (3) wherein R^{7b} and R^{8b} are phenyl, heteroaryl or substituted phenyl or heteroaryl, preferably a group of formula (1) or (2) wherein U is CH₂ or O.

15

10. A compound of Claim 9, characterized in that R^{9b} is hydrogen or lower alkyl, preferably hydrogen.

11. A compound of Claim 10, characterised in that R^{7b} and R^{8b} are independently
20 phenyl, phenyl substituted with halogen, lower alkyl, lower alkoxy or trifluoromethyl, 2-thienyl, 2-thienyl substituted with methyl, pyrrolyl or pyrrolyl substituted with methyl or ethyl.

12. A compound of Claim 11, characterised in that Y is a group of formula (1), (2)
25 or (3) wherein U is CH₂ or O and s is 0, and r+t is 0-5, preferably 1-3, most preferably 2.

13. A compound of Claim 12, characterised in that R⁴ - R⁶ are hydrogen and R³ is hydrogen or lower alkyl, preferably hydrogen or methyl.

30

14. A compound of Claim 8, characterized in that R¹ is a group as defined in B) wherein Y is a group of formula (4) or (5).



15. A compound of Claim 14, characterized in that $r+t$ is 0-5, preferably 1-3, most preferably 2.
16. A compound of Claim 15, characterized in that R^1 is a group of formula (5)
5 wherein p and q are 0, A is sulfur or $-\text{CH}_2\text{CH}_2-$ and R^a and R^b are hydrogen.
17. A compound of Claim 16, characterised in that $R^4 - R^6$ are hydrogen and R^3 is hydrogen or lower alkyl, preferably hydrogen or methyl.
- 10 18. A compound of Claim 8, characterized in that R^1 is a group as defined in C), preferably such a group wherein Y^c is a group of formula (1) where U is CH_2 , or a group of formula (2) in which formulas R^{7b} and R^{8b} are phenyl, heteroaryl or substituted phenyl or heteroaryl.
- 15 19. A compound of Claim 18, characterized in that R^{9b} is hydrogen or lower alkyl, preferably hydrogen.
20. A compound of Claim 8, characterized in that R^1 is a group as defined in C) wherein Y^c is a group of formula (5).
- 20 21. A compound of Claim 8, characterized in that R^1 is a group as defined in D) wherein Y^d is a group of formula (2), (5), (6), (7), (8), (9) or (10).
22. A compound of Claim 21, characterized in that Y^d is a group of formula (6), (7)
25 or (8), R^{9d} is hydrogen or lower alkyl, preferably hydrogen, R^{10d} and R^{11d} are hydrogen and R^{7d} and R^{8d} are phenyl, phenyl substituted with halogen, lower alkyl, lower alkoxy or trifluoromethyl, 2-thienyl, 2-thienyl substituted with lower alkyl, pyrrolyl or pyrrolyl substituted with methyl or ethyl.
- 30 23. A compound of Claim 22, characterized in that $k+l$ is 0-4, preferably 1-3.
24. A compound of any of Claims 8 - 23, characterized in that R^2 is hydrogen, lower alkyl or a group $R^{2'}-\text{O}-\text{CO}-$ wherein $R^{2'}$ is as defined in Claim 1.



25. A compound of Claim 24, characterized in that R² is hydrogen or lower alkyl, preferably hydrogen or methyl.
- 5 26. A compound of any of Claims 8 - 25, characterized in that P is hydrogen.
27. A compound of any of Claims 8 - 25, characterized in that it is a compound of formula 1a and that P is a group ZR as defined in Claim 1, wherein Z is CH₂ and R is a group OC(O)R_z as defined in vi) in Claim 1, preferably pivaloyloxy or
10 benzoyloxy.
28. A pharmaceutical composition comprising at least one novel 4-amino-tetrahydro-benzisoxazoles or -isothiazole of any of Claims 1 -27 in a therapeutically effective amount together with a pharmaceutically acceptable carrier and or diluent.
- 15 29. Use of a 4-aminotetrahydrobenzisoxazole or -isothiazole of any of Claims 1 -27 for the manufacture of a pharmaceutical preparation for the treatment of diseases associated with GABA neurotransmission.
- 20 30. Use of a 4-aminotetrahydrobenzisoxazole or -isothiazole of any of Claims 1 -27 for the manufacture of a pharmaceutical preparation for the treatment of analgesia, psychosis, convulsions, epilepsy, anxiety, muscular and movement disorders, spastic disorders or symptoms of Huntington's or Parkinsons disease, preferably convulsions.