


New dihydroquinoline-2-one derivatives

The present invention relates to organic compounds useful for therapy or prophylaxis in a mammal, and in particular to aldosterone synthase (CYP11B2 or CYP11B1) inhibitors for the treatment or prophylaxis of chronic kidney disease, congestive heart failure,

5 hypertension, primary aldosteronism and Cushing syndrom.

The present invention provides novel compounds of formula (I)



wherein

R¹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

10 R² is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R³ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R⁴ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R³ and R⁴ together with the carbon atoms to which they are attached form a double bond;

15 R⁵ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R⁶ is H or R⁸, wherein in case R⁶ is H then at least one of A¹ and A³ is N;

R⁷ is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R¹ and R⁷ together with the nitrogen and carbon atoms to which they are attached form a heterocycloalkyl substituted with one to three substituents independently selected from H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

5 R⁸ is -O_m-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q-NR¹⁵R¹⁶,
-N_mR¹⁷-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q-NR¹⁵R¹⁶ or
-S_m(O)_r-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q-NR¹⁵R¹⁶;

R⁹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹⁰ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹¹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

10 or R¹⁰ and R¹¹ together with the carbon atoms to which they are attached form a double bond;

R¹² is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R¹¹ and R¹² together with the carbon atom to which they are attached form a cycloalkyl or a heterocycloalkyl;

15 R¹³ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹⁴ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R¹⁰ and R¹⁴ together form -(CH₂)_t-;

R¹⁵ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkoxyalkyl, haloalkoxyalkyl or hydroxyalkyl;

20 R¹⁶ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, phenylalkyl, substituted phenylalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, -S(O)₂R¹⁸, -S(O)R¹⁸, -S(O)₂OR¹⁸, -S(O)₂NR¹⁸R¹⁹, -C(O)R¹⁸, -C(O)OR¹⁸ or -C(O)NR¹⁸R¹⁹, wherein substituted phenylalkyl, substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy,

25

haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and
 5 cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl, wherein in case R^{15} is H or alkyl and R^{16} is H or alkyl, then the sum
 10 of n, p and q is at least 1;

or R^{15} and R^{16} together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R^{23} , R^{24} and R^{25} ;

or R^{13} and R^{15} together with the nitrogen and carbon atoms to which they are
 15 attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R^{23} , R^{24} and R^{25} ;

or R^{11} and R^{15} together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R^{23} , R^{24} and R^{25} ;

or R^9 and R^{15} together with the nitrogen and carbon atoms to which they are attached
 20 form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R^{23} , R^{24} and R^{25} ;

R^{17} is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^{18} is alkyl, cycloalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, halohydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkylcarbonyloxyalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, wherein substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted
 25
 30

amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl;

R^{19} is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

A^1 is CR^{20} or N;

A^2 is CR^{21} or N;

A^3 is CR^{22} or N;

R^{20} is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^{21} is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^{22} is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^{23} , R^{24} and R^{25} are each independently selected from H, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, $-S(O)_2R^{18}$, $-C(O)R^{18}$, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl; wherein in case m is zero and R^9 and R^{15} , or R^{11} and R^{15} or R^{13} and R^{15} or R^{15} and R^{16} together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, then at least one of R^{23} , R^{24} , R^{25} and R^{16} is different from H, alkyl, haloalkyl, alkoxy, haloalkoxy, halogen, hydroxy, cyano, amino, substituted amino, aminocarbonyl, substituted

aminocarbonyl, wherein substituted amino and substituted aminocarbonyl are substituted on the nitrogen atom with one to two alkyl, and $-C(O)R^{18}$, $-S(O)_2R^{18}$, wherein R^{18} is alkyl, haloalkyl, hydroxyalkyl, halohydroxyalkyl, alkoxyalkyl or haloalkoxyalkyl;

5 m is zero or 1, wherein in case m is 1, then the sum of n and p is 2, 3 or 4;

 n is zero, 1 or 2;

 p is zero, 1 or 2;

 q is zero, 1 or 2;

 r is zero, 1 or 2, wherein in case m is zero then r is zero;

10 t is zero, 1 or 2;

 or pharmaceutically acceptable salts or esters.

 Herein we describe inhibitors of aldosterone synthase that have the potential to protect from organ/ tissue damage caused by an absolute or relative excess of aldosterone. Hypertension affects about 20% of the adult population in developed countries. In persons
15 60 years and older, this percentage increases to above 60%. Hypertensive subjects display an increased risk of other physiological complications including stroke, myocardial infarction, atrial fibrillation, heart failure, peripheral vascular disease and renal impairment. The renin angiotensin aldosterone system is a pathway that has been linked to hypertension, volume and salt balance and more recently to contribute directly to end
20 organ damage in advanced stages of heart failure or kidney disease. ACE inhibitors and angiotensin receptor blockers (ARBs) are successfully used to improve duration and quality of life of patients. These drugs are not yielding maximum protection. In a relatively large number of patients ACE and ARB's lead to so-called aldosterone breakthrough, a phenomenon where aldosterone levels, after a first initial decline, return to pathological
25 levels. It has been demonstrated that the deleterious consequences of inappropriately increased aldosterone levels (in relation to salt intake/levels) can be minimized by aldosterone blockade with mineralocorticoid receptor antagonists. A direct inhibition of aldosterone synthesis is expected to provide even better protection as it will also reduce non-genomic effects of aldosterone as well.

30 The effects of aldosterone on Na/K transport lead to increased re-absorption of sodium and water and the secretion of potassium in the kidneys. Overall this results in

increased blood volume and, therefore, increased blood pressure. Beyond its role in the regulation of renal sodium re-absorption aldosterone can exert deleterious effects on the kidney, the heart and the vascular system especially in a “high sodium” context. It has been shown that under such conditions aldosterone leads to increased oxidative stress
5 which ultimately may contribute to organ damage. Infusion of aldosterone into renally compromised rats (either by high salt treatment or by unilaterally nephrectomy) induces a wide array of injuries to the kidney including glomerular expansion, podocyte injury, interstitial inflammation, mesangial cell proliferation and fibrosis reflected by proteinuria. More specifically aldosterone was shown to increase the expression of the adhesion
10 molecule ICAM-1 in the kidney. ICAM-1 is critically involved in glomerular inflammation. Similarly, aldosterone was shown to increase the expression of inflammatory cytokines, such as interleukin IL-1b and IL-6, MCP-1 and osteopontin. On a cellular level it was demonstrated that in vascular fibroblasts aldosterone increased the expression of type I collagen mRNA, a mediator of fibrosis. Aldosterone also stimulates
15 type IV collagen accumulation in rat mesangial cells and induces plasminogen activator inhibitor-1 (PAI-1) expression in smooth muscle cells. In summary aldosterone has emerged as a key hormone involved in renal damage. Aldosterone plays an equally important role in mediating cardiovascular risk.

There is ample preclinical evidence that MR-antagonists (spironolactone and
20 eplerenone) improve blood pressure, cardiac and renal function in various pre-clinical models.

More recently preclinical studies highlight the important contribution of CYP11B2 to cardiovascular and renal morbidity and mortality. The CYP11B2 inhibitor FAD286 and the MR antagonist spironolactone were evaluated in a rat model of chronic kidney disease
25 (high angiotensin II exposure; high salt and uni-nephrectomy). Angiotensin II and high salt treatment caused albuminuria, azotemia, renovascular hypertrophy, glomerular injury, increased PAI-1, and osteopontin mRNA expression, as well as tubulointerstitial fibrosis. Both drugs prevented these renal effects and attenuated cardiac and aortic medial hypertrophy. Following 4 weeks of treatment with FAD286, plasma aldosterone was
30 reduced, whereas spironolactone increased aldosterone at 4 and 8 weeks of treatment. Similarly only spironolactone but not FAD286 enhanced angiotensin II and salt-stimulated PAI-1 mRNA expression in the aorta and the heart. In other studies the CYP11B2 inhibitor FAD286 improved blood pressure and cardiovascular function and structure in rats with experimental heart failure. In the same studies FAD286 was shown to improve kidney
35 function and morphology.

Administration of an orally active CYP11B2 inhibitor, LCI699, to patients with primary aldosteronism, lead to the conclusion that it effectively inhibits CYP11B2 in patients with primary aldosteronism resulting in significantly lower circulating aldosterone levels and that it corrected the hypokalemia and mildly decreased blood pressure. The effects on the glucocorticoid axis were consistent with a poor selectivity of the compound and a latent inhibition of cortisol synthesis. Taken together these data support the concept that a CYP11B2 inhibitor can lower inappropriately high aldosterone levels. Achieving good selectivity against CYP11B1 is important to be free of undesired side effects on the HPA axis and will differentiate different CYP11B2 inhibitors.

Objects of the present invention are the compounds of formula (I) and their aforementioned salts and esters and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments containing the said compounds, their pharmaceutically acceptable salts or esters, the use of the said compounds, salts or esters for the treatment or prophylaxis of illnesses, especially in the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom and the use of the said compounds, salts or esters for the production of medicaments for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.

The term "alkoxy" denotes a group of the formula -O-R', wherein R' is an alkyl group. Examples of alkoxy group include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy and tert-butoxy. Particular alkoxy group include methoxy, ethoxy and tert-butoxy.

The term "alkoxyalkoxy" denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group has been replaced by another alkoxy group. Examples of alkoxyalkoxy group include methoxymethoxy, ethoxymethoxy, methoxyethoxy, ethoxyethoxy, methoxypropoxy and ethoxypropoxy. Particular alkoxyalkoxy groups include methoxymethoxy and methoxyethoxy.

The term "alkoxyalkoxyalkyl" denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by an alkoxyalkoxy group. Examples of alkoxyalkoxyalkyl group include methoxymethoxymethyl, ethoxymethoxymethyl, methoxyethoxymethyl, ethoxyethoxymethyl, methoxypropoxymethyl, ethoxypropoxymethyl, methoxymethoxyethyl, ethoxymethoxyethyl, methoxyethoxyethyl, ethoxyethoxyethyl, methoxypropoxyethyl and ethoxypropoxyethyl.

The term “alkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by an alkoxy group. Exemplary alkoxyalkyl groups include methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methoxypropyl and ethoxypropyl. Particular alkoxyalkyl group include methoxymethyl and methoxyethyl. More particular alkoxyalkyl group is methoxymethyl.

The term “alkoxycarbonyl” denotes a group of the formula $-C(O)-R'$, wherein R' is an alkoxy group. Examples of alkoxycarbonyl groups include groups of the formula $-C(O)-R'$, wherein R' is methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy and tert-butoxy. Particular alkoxycarbonyl group is a group of the formula $-C(O)-R'$, wherein R' is methoxy or tert-butoxy.

The term “alkyl” denotes a monovalent linear or branched saturated hydrocarbon group of 1 to 12 carbon atoms. In particular embodiments, alkyl has 1 to 7 carbon atoms, and in more particular embodiments 1 to 4 carbon atoms. Examples of alkyl include methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, and tert-butyl. Particular alkyl groups include methyl, isopropyl or ethyl. More particular alkyl group is methyl.

The term “alkylcarbonyl” of the formula $-C(O)-R'$, wherein R' is an alkyl group. Examples of alkylcarbonyl groups include groups of the formula $-C(O)-R'$, wherein R' is methyl or ethyl.

The term “alkylcarbonylamino” denotes an amino group wherein one of the hydrogen atoms of the $-NH_2$ group is replaced by an alkylcarbonyl group. Examples of alkylcarbonylamino groups include groups wherein R' is methyl or ethyl.

The term “alkylcarbonylaminoalkyl” denotes an aminoalkyl group wherein one of the hydrogen atoms of the $-NH_2$ group is replaced by an alkylcarbonyl group. Examples of alkylcarbonylaminoalkyl groups include groups wherein R' is methyl or ethyl.

The term “alkylcarbonyloxyalkyl” denotes a hydroxyalkyl group wherein the hydrogen atom of the $-OH$ group is replaced by an alkylcarbonyl group. Examples of alkylcarbonyloxyalkyl groups include alkylcarbonyloxymethyl and alkylcarbonyloxyethyl. Particular alkylcarbonyloxyalkyl group is alkylcarbonyloxymethyl. More particular alkylcarbonyloxyalkyl group is methylcarbonyloxymethyl.

The term “alkylcycloalkyl” denotes a cycloalkyl group wherein at least one of the hydrogen atoms of the cycloalkyl group is replaced by an alkyl group. Examples of alkylcycloalkyl include methyl-cyclopropyl, dimethyl-cyclopropyl, methyl-cyclobutyl,

dimethyl-cyclobutyl, methyl-cyclopentyl, dimethyl-cyclopentyl, methyl-cyclohexyl and dimethyl-cyclohexyl. Particular alkylcycloalkyl groups include methyl-cyclopropyl and dimethyl-cyclopropyl.

5 The term “alkylcycloalkylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by an alkylcycloalkyl group. Examples of alkylcycloalkylalkyl include methyl-cyclopropylmethyl, dimethyl-cyclopropylmethyl, methyl-cyclopropylethyl, dimethyl-cyclopropylethyl, methyl-cyclobutylmethyl, dimethyl-cyclobutylmethyl, methyl-cyclobutylethyl, dimethyl-cyclobutylethyl, methyl-cyclopentylmethyl, dimethyl-cyclopentylmethyl, methyl-cyclopentylethyl, dimethyl-cyclopentylethyl, methyl-cyclohexylmethyl, dimethyl-cyclohexylmethyl, methyl-cyclohexylethyl, dimethyl-cyclohexylethyl, methyl-cycloheptylmethyl, dimethyl-cycloheptylmethyl, methyl-cycloheptylethyl, dimethyl-cycloheptylethyl, methyl-cyclooctylmethyl, dimethyl-cyclooctylmethyl, methyl-cyclooctylethyl and dimethyl-cyclooctylethyl.

15 The term “amino” denotes a -NH_2 group.

The term “aminoalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by an amino group. Examples of aminoalkyl include aminomethyl, aminoethyl, aminopropyl, aminomethylpropyl and diaminopropyl.

The term “aminocarbonyl” of the formula -C(O)-NH_2

20 The term “aryl” denotes a monovalent aromatic carbocyclic mono- or bicyclic ring system comprising 6 to 10 carbon ring atoms. Examples of aryl group include phenyl and naphthyl. Particular aryl group is phenyl.

The term “bicyclic ring system” denotes two rings which are fused to each other via a common single or double bond (annelated bicyclic ring system), via a sequence of three or more common atoms (bridged bicyclic ring system) or via a common single atom (spiro bicyclic ring system). Bicyclic ring systems can be saturated, partially unsaturated, unsaturated or aromatic. Bicyclic ring systems can comprise heteroatoms selected from N, O and S.

The term “carbonyl” denotes a -C(O)- group.

30 The term “cyano” denotes a $\text{-C}\equiv\text{N}$ group.

The term “cycloalkoxy” denotes a group of the formula -O-R', wherein R' is a cycloalkyl group. Examples of cycloalkoxy group include cyclopropoxy, cyclobutoxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy and cyclooctyloxy. Particular cycloalkoxy group is cyclopropoxy.

- 5 The term “cycloalkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a cycloalkoxy group. Examples of cycloalkoxyalkyl group include cyclopropoxymethyl, cyclopropoxyethyl, cyclobutoxymethyl, cyclobutoxyethyl, cyclopentyloxymethyl, cyclopentyloxyethyl, cyclohexyloxymethyl, cyclohexyloxyethyl, cycloheptyloxymethyl, cycloheptyloxyethyl, 10 cyclooctyloxymethyl and cyclooctyloxyethyl.

- The term “cycloalkyl” denotes a monovalent saturated monocyclic or bicyclic hydrocarbon group of 3 to 10 ring carbon atoms. In particular embodiments, cycloalkyl denotes a monovalent saturated monocyclic hydrocarbon group of 3 to 8 ring carbon atoms. Bicyclic means consisting of two saturated carbocycles having two carbon atoms in 15 common. Particular cycloalkyl groups are monocyclic. Examples for monocyclic cycloalkyl are cyclopropyl, cyclobutanyl, cyclopentyl, cyclohexyl or cycloheptyl. Examples for bicyclic cycloalkyl are bicyclo[2.2.1]heptanyl or bicyclo[2.2.2]octanyl. Particular monocyclic cycloalkyl group is cyclopropyl.

- The term “cycloalkylalkoxy” denotes an alkoxy group wherein at least one of the 20 hydrogen atoms of the alkoxy group is replaced by a cycloalkyl group. Examples of cycloalkylalkoxy include cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy, cycloheptylmethoxy and cyclooctylmethoxy.

- The term “cycloalkylalkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a cycloalkylalkoxy group. Examples of 25 cycloalkylalkoxyalkyl include cyclopropylmethoxymethyl, cyclopropylmethoxyethyl, cyclobutylmethoxymethyl, cyclobutylmethoxyethyl, cyclopentylmethoxyethyl, cyclopentylmethoxyethyl, cyclohexylmethoxymethyl, cyclohexylmethoxyethyl, cycloheptylmethoxymethyl, cycloheptylmethoxyethyl, cyclooctylmethoxymethyl and cyclooctylmethoxyethyl.

- 30 The term “cycloalkylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a cycloalkyl group. Examples of cycloalkylalkyl include cyclopropylmethyl, cyclopropylethyl, cyclobutylpropyl and cyclopentylbutyl.

The term “cycloalkylcarbonyl” of the formula $-C(O)-R'$, wherein R' is a cycloalkyl group. Examples of cycloalkylcarbonyl groups include groups of the formula $-C(O)-R'$, wherein R' is cyclopropyl.

5 The term “cycloalkylcarbonylamino” denotes an amino group wherein one of the hydrogen atoms of the $-NH_2$ group is replaced by an cycloalkylcarbonyl group. Examples of alkylcarbonylamino groups include groups wherein R' is cyclopropyl.

The term “cycloalkylcarbonylaminoalkyl” denotes an aminoalkyl group wherein one of the hydrogen atoms of the $-NH_2$ group is replaced by an cycloalkylcarbonyl group. Examples of alkylcarbonylaminoalkyl groups include groups wherein R' is cyclopropyl.

10 The term “haloalkoxy” denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group has been replaced by same or different halogen atoms. The term “perhaloalkoxy” denotes an alkoxy group where all hydrogen atoms of the alkoxy group have been replaced by the same or different halogen atoms. Examples of haloalkoxy include fluoromethoxy, difluoromethoxy, trifluoromethoxy, trifluoroethoxy,
15 trifluoromethylethoxy, trifluorodimethylethoxy and pentafluoroethoxy. Particular haloalkoxy groups are trifluoromethoxy and 2,2-difluoroethoxy.

The term “haloalkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a haloalkoxy group. Examples of haloalkoxyalkyl include fluoromethoxymethyl, difluoromethoxymethyl,
20 trifluoromethoxymethyl, fluoroethoxymethyl, difluoroethoxymethyl, trifluoroethoxymethyl, fluoromethoxyethyl, difluoromethoxyethyl, trifluoromethoxyethyl, fluoroethoxyethyl, difluoroethoxyethyl, trifluoroethoxyethyl, fluoromethoxypropyl, difluoromethoxypropyl, trifluoromethoxypropyl, fluoroethoxypropyl, difluoroethoxypropyl and trifluoroethoxypropyl. Particular haloalkoxyalkyl is 2,2-
25 difluoroethoxyethyl

The term “haloalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by same or different halogen atoms. The term “perhaloalkyl” denotes an alkyl group where all hydrogen atoms of the alkyl group have been replaced by the same or different halogen atoms. Examples of haloalkyl include
30 fluoromethyl, difluoromethyl, trifluoromethyl, trifluoroethyl, trifluoromethylethyl and pentafluoroethyl. Particular haloalkyl groups are trifluoromethyl and trifluoroethyl.

The term “halocycloalkyl” denotes a cycloalkyl group wherein at least one of the hydrogen atoms of the cycloalkyl group has been replaced by same or different halogen

atoms, particularly fluoro atoms. Examples of halocycloalkyl groups include fluorocyclopropyl, difluorocyclopropyl, fluorocyclobutyl and difluorocyclobutyl.

The term “halocycloalkylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a halocycloalkyl. Examples of
5 halocycloalkylalkyl groups include fluorocyclopropylmethyl, fluorocyclopropylethyl, difluorocyclopropylmethyl, difluorocyclopropylethyl, fluorocyclobutylmethyl, fluorocyclobutylethyl, difluorocyclobutylmethyl and difluorocyclobutylethyl.

The term “halogen” and “halo” are used interchangeably herein and denote fluoro, chloro, bromo, or iodo. Particular halogens are chloro and fluoro.

10 The term “halohydroxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by same or different halogen atoms and at least one of the hydrogen atoms of the alkyl group has been replaced by hydroxy.

The term “heteroaryl” denotes a monovalent aromatic heterocyclic mono- or bicyclic ring system of 5 to 12 ring atoms, comprising 1, 2, 3 or 4 heteroatoms selected from N, O
15 and S, the remaining ring atoms being carbon. Examples of heteroaryl group include pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl, triazinyl, azepinyl, diazepinyl, isoxazolyl, benzofuranyl, isothiazolyl, benzothienyl, indolyl, isoindolyl, isobenzofuranyl, benzimidazolyl, benzoxazolyl, benzoisoxazolyl,
20 benzothiazolyl, benzoisothiazolyl, benzooxadiazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl and quinoxalinyl. Particular heteroaryl groups include pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl, isoxazolyl and isothiazolyl. More particular heteroaryl groups include
25 imidazolyl, oxazolyl, thiazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl, isoxazolyl and isothiazolyl.

In particular in the definition of R¹⁸, particular heteroaryl groups include imidazolyl, oxazolyl, furanyl, oxadiazolyl, pyridinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl and isoxazolyl. More particularly oxazolyl, pyridinyl and pyrazolyl.

30 The term “heterocycloalkyl” denotes a monovalent saturated or partly unsaturated mono- or bicyclic ring system of 3 to 9 ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. In particular embodiments, heterocycloalkyl is a monovalent saturated monocyclic ring system of 4 to 7

ring atoms, comprising 1, 2, or 3 ring heteroatoms selected from N, O and S, the remaining ring atoms being carbon. Examples for monocyclic saturated heterocycloalkyl are aziridinyl, oxiranyl, azetidiny, oxetanyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydro-thienyl, pyrazolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, 5 piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, thiomorpholinyl, 1,1-dioxo-thiomorpholin-4-yl, azepanyl, diazepanyl, homopiperazinyl, oxazepanyl and thiazinanyl. Examples for bicyclic saturated heterocycloalkyl are 8-aza-bicyclo[3.2.1]octyl, quinuclidinyl, 8-oxa-3-aza-bicyclo[3.2.1]octyl, 9-aza-bicyclo[3.3.1]nonyl, 3-oxa-9-aza-bicyclo[3.3.1]nonyl, 3-thia-9-aza-bicyclo[3.3.1]nonyl 10 and 2,6-diaza-spiro[3.3]heptanyl. Examples for partly unsaturated heterocycloalkyl are dihydrofuryl, imidazolinyl, dihydro-oxazolyl, tetrahydro-pyridinyl, or dihydropyranyl. More particular examples of heterocycloalkyl group are pyrrolidinyl, pyrazolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperazinyl, morpholinyl, thiomorpholinyl, 1,1-dioxo- 15 thiomorpholin-4-yl, azepanyl, diazepanyl, homopiperazinyl, oxazepanyl, thiazinanyl and 2,6-diaza-spiro[3.3]heptanyl. More particular examples of a heterocycloalkyl are pyrrolydiny, piperidinyl, thiomorpholinyl, thiazinanyl and 2,6-diaza-spiro[3.3]heptanyl.

In particular in the definition of R¹⁶, particular heterocycloalkyl group is oxetanyl.

In particular, the heterocycloalkyl formed by R⁹ and R¹⁵ together with the nitrogen 20 and carbon atoms to which they are attached is azetidiny, pyrrolidinyl, piperidinyl or tetrahydro-pyridinyl. More particularly, pyrrolidinyl.

In particular, the heterocycloalkyl formed by R¹¹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached is azetidiny or pyrrolidinyl.

In particular, the heterocycloalkyl formed by R¹⁵ and R¹⁶ together with the nitrogen 25 and carbon atoms to which they are attached is pyrrolidinyl, piperidinyl, thiomorpholinyl, thiazinanyl, isothiazolidinyl, 1,1-dioxo-thiomorpholin-4-yl or 2,6-diaza-spiro[3.3]heptanyl. More particularly, thiomorpholinyl or 1,1-dioxo-thiomorpholin-4-yl. Further particularly, pyrrolidinyl or 1,1-dioxo-thiomorpholin-4-yl.

In particular, the heterocycloalkyl formed by R¹¹ and R¹² together with the nitrogen 30 and carbon atoms to which they are attached is oxetanyl.

The term "hydroxy" denotes a -OH group.

The term "hydroxyalkyl" denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a hydroxy group. Examples of hydroxyalkyl include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxymethylpropyl and dihydroxypropyl. Particular example is hydroxymethyl.

5 The term "oxo" denotes a divalent oxygen atom =O.

The term "phenylalkyl" denotes an alkyl group wherein one of the hydrogen atoms of the alkyl group has been replaced by a phenyl. Examples of phenylalkyl are benzyl and phenylethyl. Particular example of phenylalkyl is benzyl.

10 The term "tetrazolylalkyl" denotes an alkyl group wherein one of the hydrogen atoms of the alkyl group has been replaced by a tetrazolyl. Examples of tetrazolylalkyl are tetrazolymethyl and tetrazolyethyl. Particular example of tetrazolylalkyl is tetrazolymethyl.

15 The term "triazolylalkyl" denotes an alkyl group wherein one of the hydrogen atoms of the alkyl group has been replaced by a triazolyl. Examples of triazolylalkyl are triazolymethyl and triazolylethyl. Particular example of triazolylalkyl is triazolymethyl.

20 The term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. The salts are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, in particular hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, N-acetylcystein and the like. In addition these salts may be prepared by addition of an inorganic base or an organic

25 base to the free acid. Salts derived from an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium salts and the like. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine,

30 diethylamine, triethylamine, tripropylamine, ethanolamine, lysine, arginine, N-ethylpiperidine, piperidine, polyimine resins and the like. Particular pharmaceutically acceptable salts of compounds of formula (I) are the hydrochloride salts, methanesulfonic acid salts and citric acid salts.

"Pharmaceutically acceptable esters" means that compounds of general formula (I) may be derivatised at functional groups to provide derivatives which are capable of conversion back to the parent compounds in vivo. Examples of such compounds include physiologically acceptable and metabolically labile ester derivatives, such as

5 methoxymethyl esters, methylthiomethyl esters and pivaloyloxymethyl esters.

μAdditionally, any physiologically acceptable equivalents of the compounds of general formula (I), similar to the metabolically labile esters, which are capable of producing the parent compounds of general formula (I) in vivo, are within the scope of this invention.

The term "protecting group" (PG) denotes the group which selectively blocks a

10 reactive site in a multifunctional compound such that a chemical reaction can be carried out selectively at another unprotected reactive site in the meaning conventionally associated with it in synthetic chemistry. Protecting groups can be removed at the appropriate point. Exemplary protecting groups are amino-protecting groups, carboxy-protecting groups or hydroxy-protecting groups. Particular protecting groups are the tert-

15 butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), fluorenylmethoxycarbonyl (Fmoc) and benzyl (Bn). Further particular protecting groups are the tert-butoxycarbonyl (Boc) and the fluorenylmethoxycarbonyl (Fmoc). More particular protecting group is the tert-butoxycarbonyl (Boc). The abbreviation μM means microMolar and is equivalent to the symbol μM.

20 The compounds of the present invention can also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the present invention also embraces isotopically-labeled variants of the present invention which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having the atomic mass or mass number different from the predominant atomic mass or mass number usually

25 found in nature for the atom. All isotopes of any particular atom or element as specified are contemplated within the scope of the compounds of the invention, and their uses. Exemplary isotopes that can be incorporated in to compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine, chlorine and iodine, such as ²H ("D"), ³H ("T"), ¹¹C, ¹³C, ¹⁴C, ¹³N, ¹⁵N, ¹⁵O, ¹⁷O, ¹⁸O, ³²P, ³³P, ³⁵S, ¹⁸F, ³⁶Cl, ¹²³I and ¹²⁵I. Certain isotopically

30 labeled compounds of the present invention (e.g., those labeled with ³H or ¹⁴C) are useful in compound and /or substrate tissue distribution assays. Tritiated (³H) and carbon-14 (¹⁴C) isotopes are useful for their ease of preparation and detectability. Further substitution with heavier isotopes such as deuterium (i.e., ²H) may afford certain therapeutic advantages resulting from greater metabolic stability (e.g., increased in vivo half-life or reduced dosage requirements) and hence may

35 be preferred in some circumstances. Positron emitting isotopes such as ¹⁵O, ¹³N, ¹¹C, and ¹⁸F are useful for positron emission tomography (PET) studies to examine substrate receptor

occupancy. Isotopically labeled compounds of the present inventions can generally be prepared by following procedures analogous to those disclosed in the Schemes and/or in the Examples herein below, by substituting a non-isotopically labeled reagent with a isotopically labeled reagent. In particular, compounds of formula (I) wherein one or more H atom have been replaced by a ^2H atom are also an embodiment of this invention.

The compounds of formula (I) can contain several asymmetric centers and can be present in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates.

According to the Cahn-Ingold-Prelog Convention the asymmetric carbon atom can be of the "R" or "S" configuration.

Also an embodiment of the present invention are compounds according to formula (I) as described herein and pharmaceutically acceptable salts or esters thereof, in particular compounds according to formula (I) as described herein and pharmaceutically acceptable salts thereof, more particularly compounds according to formula (I) as described herein.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein

R^1 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^2 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^3 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^4 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^5 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^6 is H or R^8 , wherein in case R^6 is H then at least one of A^1 and A^3 is N;

R^7 is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^8 is $-\text{O}_m-(\text{CR}^9\text{R}^{10})_n-(\text{CR}^{11}\text{R}^{12})_p-(\text{CR}^{13}\text{R}^{14})_q-\text{NR}^{15}\text{R}^{16}$,
 $-\text{N}_m\text{R}^{17}-(\text{CR}^9\text{R}^{10})_n-(\text{CR}^{11}\text{R}^{12})_p-(\text{CR}^{13}\text{R}^{14})_q-\text{NR}^{15}\text{R}^{16}$ or
 $-\text{S}_m(\text{O})_r-(\text{CR}^9\text{R}^{10})_n-(\text{CR}^{11}\text{R}^{12})_p-(\text{CR}^{13}\text{R}^{14})_q-\text{NR}^{15}\text{R}^{16}$;

R^9 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹⁰ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹¹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹² is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

5 or R¹¹ and R¹² together with the carbon atom to which they are attached form a cycloalkyl;

R¹³ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹⁴ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R¹⁰ and R¹⁴ together form -(CH₂)_t;

10 R¹⁵ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkoxyalkyl, haloalkoxyalkyl or hydroxyalkyl;

15 R¹⁶ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, phenylalkyl, substituted phenylalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, -S(O)₂R¹⁸, -S(O)₂OR¹⁸, -S(O)₂NR¹⁸R¹⁹, -C(O)R¹⁸, -C(O)OR¹⁸ or -C(O)NR¹⁸R¹⁹,
 20 wherein substituted phenylalkyl, substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and
 25 cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl, wherein in case R¹⁵ is H or alkyl and R¹⁶ is H or alkyl, then the sum
 30 of n, p and q is at least 1;

or R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R²³, R²⁴ and R²⁵;

5 or R¹³ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

or R¹¹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

10 or R⁹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

R¹⁷ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

15 R¹⁸ is alkyl, cycloalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, halohydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkylcarbonyloxyalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, wherein substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl;

20

25

30

R¹⁹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

A¹ is CR²⁰ or N;

A² is CR²¹ or N;

A³ is CR²² or N;

R²⁰ is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R²¹ is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

5 R²² is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R²³, R²⁴ and R²⁵ are each independently selected from H, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, -S(O)₂R¹⁸, -C(O)R¹⁸, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein
10 substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl;
15 wherein in case m is zero, then at least one of R²³, R²⁴ and R²⁵ is different from H, alkyl, haloalkyl, alkoxy, haloalkoxy, halogen, hydroxy, cyano, amino, substituted amino, aminocarbonyl or substituted aminocarbonyl, wherein
20 substituted amino and substituted aminocarbonyl are substituted on the nitrogen atom with one to two alkyl;

m is zero or 1, wherein in case m is 1, then the sum of n and p is 2, 3 or 4;

25 n is zero, 1 or 2;

p is zero, 1 or 2;

q is zero, 1 or 2;

r is zero, 1 or 2, wherein in case m is zero then r is zero;

t is zero, 1 or 2;

or pharmaceutically acceptable salts or esters.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

A further embodiment of the present invention are compounds according to formula
5 (I) as described herein, wherein R^1 is H or alkyl.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is alkyl.

In a further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is methyl.

10 Another further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^2 is H.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^3 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

Another embodiment of the present invention are compounds according to formula
15 (I) as described herein, wherein R^3 is H.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^4 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^4 is H or alkyl.

20 A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^4 is H.

A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^5 is H or alkyl.

Also an embodiment of the present invention are compounds according to formula
25 (I) as described herein, wherein R^5 is H.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^6 is H and at least one of A^1 and A^3 is N.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^6 is R^8 .

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^7 is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl. A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^7 is H.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 and R^7 together with the nitrogen and carbon atoms to which they are attached form a heterocycloalkyl substituted with one to three substituents independently selected from H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^8 is $-O_m-(CR^9R^{10})_n-(CR^{11}R^{12})_p-(CR^{13}R^{14})_q-NR^{15}R^{16}$.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^9 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^9 is H, alkyl or cycloalkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^9 is H or alkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^9 is H.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{10} is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{10} is H or alkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^{10} is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{11} is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹¹ is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁰ and R¹¹ together with the carbon atoms to which they
5 are attached form a double bond.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹² is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹¹ and R¹² together with the carbon atom to which they
10 are attached form a cycloalkyl.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹³ is H.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁴ is H, alkyl, haloalkyl, cycloalkyl or
15 halocycloalkyl.

A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁴ is H.

Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁰ and R¹⁴ together form -(CH₂)_t-.

20 Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁵ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkoxyalkyl, haloalkoxyalkyl or hydroxyalkyl.

Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁵ is H, alkyl or alkoxyalkyl.

25 Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁵ is H or alkyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁵ is H.

Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{16} is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, phenylalkyl, substituted phenylalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, $-S(O)_2R^{18}$, $-S(O)_2OR^{18}$, $-S(O)_2NR^{18}R^{19}$, $-C(O)R^{18}$, $-C(O)OR^{18}$ or $-C(O)NR^{18}R^{19}$, wherein substituted phenylalkyl, substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl, wherein in case R^{15} is H or alkyl and R^{16} is H or alkyl, then the sum of n, p and q is at least 1.

Also an embodiment of the present invention are compounds according to formula

25 (I) as described herein, wherein R^{16} is H, hydroxyalkyl, phenylalkyl, heterocycloalkyl substituted with alkyl, $-S(O)R^{18}$, $-S(O)_2R^{18}$, $-C(O)R^{18}$ or $-C(O)OR^{18}$, wherein in case R^{15} is H or alkyl and R^{16} is H or alkyl, then the sum of n, p and q is at least 1. Also an embodiment of the p of n, p and q is at least 1.

Also an embodiment of the present invention are compounds according to formula 30 (I) as described herein, wherein R¹⁶ is hydroxyalkyl, -S(O)₂R¹⁸, -C(O)R¹⁸ or -C(O)OR¹⁸.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁶ is H, hydroxyalkyl or phenylalkyl, wherein in case R¹⁵ is H or alkyl and R¹⁶ is H or alkyl, then the sum of n, p and q is at least 1.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{16} is H, hydroxyethyl, hydroxymethylbutyl or benzyl, wherein in case R^{15} is H or alkyl and R^{16} is H or alkyl, then the sum of n, p and q is at least 1.

- 5 Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{16} is hydroxyalkyl or $-S(O)_2R^{18}$.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{16} is hydroxyalkyl.

- 10 Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{16} is hydroxyethyl, hydroxymethylbutyl or benzyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{16} is $-S(O)_2R^{18}$.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{16} is $-C(O)R^{18}$.

- 15 A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{15} and R^{16} together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R^{23} , R^{24} and R^{25} .

- 20 A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{15} and R^{16} together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl wherein substituted heterocycloalkyl is substituted with one to three substituents independently selected from hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxycarbonyl, oxo, triazolylalkyl and
25 substituted aminoalkyl, wherein substituted aminoalkyl is substituted on the nitrogen atom with one to two alkyl.

- A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{15} and R^{16} together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl wherein substituted
30 heterocycloalkyl is substituted with one to three substituents independently selected from hydroxyalkyl and oxo.

A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein the heterocycloalkyl formed by R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached is selected from pyrrolydiny, piperidiny, thiomorpholiny, thiazinany, isothiazolidiny and 2,6-diaza-

5 spiro[3.3]heptany. A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein the heterocycloalkyl formed by R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached is selected from pyrrolydiny, piperidiny, thiomorpholiny, thiazinany and 2,6-diaza-spiro[3.3]heptany.

Another embodiment of the present invention are compounds according to formula
10 (I) as described herein, wherein R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form 2- hydroxymethyl-pyrrolidin-1-yl, 2- hydroxymethyl-5-oxo-pyrrolidin-1-yl or isothiazolidiny substituted on the sulfur atom by two oxo.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁵ and R¹⁶ together with the nitrogen atom to which they
15 are attached form 2- hydroxymethyl-pyrrolidin-1-yl or 2- hydroxymethyl-5-oxo-pyrrolidin-1-yl.

Another particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁸ is alkyl, cycloalkyl, haloalkyl, hydroxyalkyl, alkylcarbonyloxyalkyl, aryl, substituted aryl, heterocycloalkyl, substituted
20 heterocycloalkyl, heteroaryl, substituted heteroaryl, wherein substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl and halogen.

Another particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁸ is alkyl, cycloalkyl, hydroxyalkyl,
25 alkylcarbonyloxyalkyl or heterocycloalkyl substituted with one to three alkoxy.

Another particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁸ is alkyl, cycloalkyl, hydroxyalkyl, alkylcarbonyloxyalkyl or heteroaryl substituted with one to three alkoxy.

A particular embodiment of the present invention are compounds according to
30 formula (I) as described herein, wherein R¹⁸ is alkyl or heteroaryl substituted with one to three substituents independently selected from alkyl and halogen.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁸ is heteroaryl substituted with one to three substituents independently selected from alkyl and halogen.

A particular embodiment of the present invention are compounds according to
5 formula (I) as described herein, wherein R¹⁸ is alkyl.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R¹⁸ is ethyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein A¹ is CR²⁰.

10 The present invention also relates to compounds according to formula (I) as described herein, wherein A² is CR²¹.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein A³ is CR²².

A embodiment of the present invention are compounds according to formula (I) as
15 described herein, wherein R²⁰ is H or halogen.

A embodiment of the present invention are compounds according to formula (I) as described herein, wherein R²⁰ is H.

A embodiment of the present invention are compounds according to formula (I) as described herein, wherein R²¹ is H, alkyl or halogen.

20 A embodiment of the present invention are compounds according to formula (I) as described herein, wherein R²¹ is H or alkyl.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R²¹ is H.

The present invention also relates to compounds according to formula (I) as
25 described herein, wherein R²² is H or halogen.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R²³, R²⁴ and R²⁵ are each independently selected from hydrogen, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxyalkyl, alkoxycarbonyl, oxo,

triazolylalkyl and substituted aminoalkyl, wherein substituted aminoalkyl is substituted on the nitrogen atom with one to two alkyl.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^{22} is H.

5 Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{23} , R^{24} and R^{25} are each independently selected from hydrogen, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxycarbonyl, oxo, triazolylalkyl and substituted aminoalkyl, wherein substituted aminoalkyl is substituted on the nitrogen atom with one to two alkyl.

10 A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein wherein R^{23} , R^{24} and R^{25} are each independently selected from hydrogen, hydroxyalkyl and oxo.

The present invention also relates to compounds according to formula (I) as described herein, wherein at least one of R^{23} , R^{24} and R^{25} is different from hydrogen.

15 The present invention also relates to compounds according to formula (I) as described herein, wherein A^1 is N.

The present invention also relates to compounds according to formula (I) as described herein, wherein A^2 is N.

20 The present invention also relates to compounds according to formula (I) as described herein, wherein A^3 is N.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein m is zero or 1, wherein in case m is 1, then the sum of n, p and q is 2, 3, 4, 5 or 6.

25 A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein m is 1.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein m is zero.

Also a further embodiment of the present invention are compounds according to formula (I) as described herein, wherein n is zero or 1.

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein n is zero.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein p is zero or 1.

5 A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein q is zero or 1.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein q is zero.

10 A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein r is zero and m is zero.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein t is 2.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein t is zero.

15 Particular examples of compounds of formula (I) as described herein are selected from

Ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-amide;

20 Acetic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]carbamoyl]-methyl ester;

2-Hydroxy-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-acetamide;

6-[5-(2-Hydroxy-ethylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

25 6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

6-[5-((R)-2-Hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

- 6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 5 6-[5-((S)-2-Ethylaminomethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-2-Methoxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 10 1-Methyl-6-[5-((S)-2-[1,2,4]triazol-1-ylmethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 6-(5-Benzylamino-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-(2-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 15 Ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(1,4,4-trimethyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 1-Methyl-6-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 25 6-[5-(1,1-Dioxo-1 λ^6 -thiomorpholin-4-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-2-Methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

- (S)-1-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-pyrrolidine-2-carboxylic acid methyl ester;
- 1-Methyl-6-{5-[(S)-2-(2,2,2-trifluoro-ethoxymethyl)-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-3,4-dihydro-1H-quinolin-2-one;
- 5 6-[6-(2-Hydroxy-ethylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-(6-Benzylamino-pyrazin-2-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 7-Fluoro-6-[6-(2-hydroxy-ethylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 10 6-(6-Benzylamino-pyrazin-2-yl)-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[6-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyrazin-2-yl]-2,6-diazaspiro[3.3]heptane-2-carboxylic acid tert-butyl ester ;
- 6-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-2,6-diazaspiro[3.3]heptane-2-carboxylic acid tert-butyl ester ;
- 15 6-[5-((R)-1-Hydroxymethyl-2-methyl-propylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[6-((R)-1-Hydroxymethyl-2-methyl-propylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- Ethanesulfonic acid [5-(5-methyl-6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-pyridin-3-ylmethyl]-amide;
- 20 6-Pyridin-3-yl-3,4-dihydro-1H-[1,5]naphthyridin-2-one;
- 1-Methyl-6-pyridin-3-yl-3,4-dihydro-1H-[1,5]naphthyridin-2-one;
- 1-Methyl-6-pyridin-3-yl-3,4-dihydro-1H-[1,7]naphthyridin-2-one;
- 6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 25 N-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;

- Propane-2-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- {2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-carbamic acid tert-butyl ester;
- 5 3-Methoxy-isoxazole-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Cyclopropanecarboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 6-[5-(2-Amino-ethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- N-{2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-propionamide;
- 6-[5-(1,1-Dioxo-1 λ ⁶-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 15 and pharmaceutically acceptable salts thereof.
- Also particular examples of compounds of formula (I) as described herein are selected from
- (S)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-azetidine-1-carboxylic acid tert-butyl ester;
- 20 (R)-3-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- 3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-azetidine-1-carboxylic acid tert-butyl ester;
- (S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- 25 (S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;

- 4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;
- (S)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- 5 (S)-3-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- (R)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- 10 (R)- 2-Methyl-propane-2-sulfinic acid {3-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-oxetan-3-yl}-amide;
- Ethanesulfonic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- (R)-2-Methyl-propane-2-sulfinic acid {(S or R)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 15 (R) 2-Methyl-propane-2-sulfinic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- {{(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-carbamic acid tert-butyl ester;
- 20 6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 1-Methyl-6-[5-((S)-piperidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 25 1-Methyl-6-[5-(piperidin-4-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-[5-((S)-1-Azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

- 6-[5-((S or R)-1-Amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-[5-((R or S)-1-Amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 5 1-Methyl-6-[5-((S)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 7-Fluoro-1-methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 10 1-Methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-[5-(3-Amino-oxetan-3-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-[5-(Azetidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 15 6-[5-((trans)-4-Amino-cyclohexyloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-[5-(1-Amino-1-methyl-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-(5-Aminomethyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one ;
- 20 6-{6-[6-(3-Methoxy-isoxazole-5-carbonyl)-2,6-diaza-spiro[3.3]hept-2-yl]-pyrazin-2-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 6-[5-((S)-1-Cyclopropanecarbonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 3-Methyl-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-butyramide;

- 3,3,3-Trifluoro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;
- 2-Hydroxy-2-methyl-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;
- 5 5-Methyl-[1,3,4]oxadiazole-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-6-[5-((S)-1-propionyl-piperidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 10 2-Methoxy-pyrimidine-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide,
- 1-Methyl-1H-imidazole-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Trifluoromethyl-furan-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 Pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Pyrimidine-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-1H-pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 1-Methyl-6-[5-(1-propionyl-piperidin-4-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- Pyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 1-Methyl-6-[5-((S)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-1-Cyclopropanecarbonyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

- 3-Methyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 4-Fluoro-2,6-dimethyl-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-benzamide;
- 5 1-Methyl-6-[5-((S)-1-propionyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 3,6-Dichloro-pyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 3-Cyclopropyl-1-(2,2,2-trifluoro-ethyl)-1H-pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1,3-Dimethyl-1H-pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 Pyrimidine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 6-Methoxy-pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 5-Methyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;
- 1-Methyl-1H-pyrazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;
- 25 6-Chloro-pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Chloro-6-methyl-pyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

- 1-Methyl-6-[5-((S)-1-propionyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 5-Cyclopropyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5 2-Methyl-5-trifluoromethyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 2-Methyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 5-Cyclopropyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 2,5-Dimethyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Methyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 3,5-Dimethyl-isoxazole-4-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 6-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 3,6-Dichloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 6-Methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 3-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Fluoro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

- 5-Chloro-3-methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-6-[5-((R)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 5 7-Fluoro-1-methyl-6-[5-((S)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 5-Trifluoromethyl-pyrimidine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Methyl-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 5-Chloro-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Trifluoromethyl-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-6-[5-((R)-1-propionyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-(1-propionyl-azetidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 20 N-{3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-oxetan-3-yl}-propionamide;
- 3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 3,6-Dichloro-pyridazine-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 7-Fluoro-1-methyl-6-[5-((R)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

- 5-Chloro-3-methoxy-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 5 3-Methyl-5-trifluoromethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 10 3-Chloro-pyridine-2-carboxylic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 3-Chloro-pyridine-2-carboxylic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 3-Methyl-pyridine-2-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 20 N-[(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl]-propionamide;
- 5-Trifluoromethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 7-Fluoro-1-methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- (R)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester;

- 1-Methyl-6-[5-((R)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-(5-Aminomethyl-pyridin-3-yl)-3,4-dihydro-1H-quinolin-2-one;
- 5 6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one ;
- Ethanesulfonic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide ;
- 3-Chloro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-benzenesulfonamide;
- 10 6-Methoxy-pyridine-3-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,5-Dimethyl-isoxazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 Cyclopropanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,4-Dichloro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-benzenesulfonamide;
- 1-Methyl-1H-imidazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 6-Chloro-pyridine-3-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-1H-pyrazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 6-[5-((S)-1-Ethanesulfonyl-piperidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-(1-Ethanesulfonyl-piperidin-4-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

- 2,2,2-Trifluoro-ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- C,C,C-Trifluoro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methanesulfonamide;
- 5 6-[5-((S)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-1-Ethanesulfonyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 10 Ethanesulfonic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 6-[5-((S)-1-Ethanesulfonyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((R)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 15 6-[5-((R)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-(1-Ethanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 20 Ethanesulfonic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- Ethanesulfonic acid {(trans)-4-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-amide;
- Ethanesulfonic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 25 Ethanesulfonic acid {(S or R)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 1-Methyl-1H-pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

- (rac)-Ethanesulfonic acid {2-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-propyl}-amide;
- (rac)-Ethanesulfonic acid {cyclopropyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-methyl}-amide;
- 5 (rac)-Ethanesulfonic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 6-[5-(1,1-Dioxo-1λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 10 (rac)-Ethanesulfonic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-propyl}-amide;
- Ethanesulfonic acid ethyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 Ethanesulfonic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- Ethanesulfonic acid isopropyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid (2-ethoxy-ethyl)-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 (rac)-Ethanesulfonic acid methyl-{1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- (rac)-Ethanesulfonic acid ethyl-{1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 25 3,5-Dimethyl-isoxazole-4-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 6-{5-[2-(1,1-Dioxo-1λ6-isothiazolidin-2-yl)-ethoxy]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one;

Ethanesulfonic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

Ethanesulfonic acid {(S or R)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

5 Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

10 Ethanesulfonic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid methyl-{2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

3-Chloro-pyridine-2-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

15 N-Methyl-N-{2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-propionamide;

1-Methyl-1H-pyrazole-4-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

20 3-Methyl-pyridine-2-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide;

3-Chloro-pyridine-2-carboxylic acid methyl-{1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;

25 1-Methyl-1H-pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide;

Ethanesulfonic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

- 3-Chloro-pyridine-2-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5 6-{5-[(3-Ethyl-oxetan-3-ylamino)-methyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- Ethanesulfonic acid [5-(7-chloro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 Ethanesulfonic acid [5-(8-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(8-chloro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(5-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide;
- N-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;
- 20 (rac)-N-{1-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide;
- (S)-3-[5-(2-Oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester ;
- 3-Chloro-pyridine-2-carboxylic acid methyl-[5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide ;
- 25 3-Chloro-pyridine-2-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide ;
- (R)-2-Methyl-propane-2-sulfinic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide ;

- 6-(5-Aminomethyl-4-chloro-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5 3-Chloro-pyridine-2-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- (R)-2-Methyl-propane-2-sulfinic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide ;
- 10 6-(5-Aminomethyl-4-methyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 (R)-2-Methyl-propane-2-sulfinic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide ;
- 6-(5-Aminomethyl-4-methyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 20 3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide;
- 3-Chloro-pyridine-2-carboxylic acid methyl-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 5'-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-5,6-dihydro-4H-[3,3']bipyridinyl-1-carboxylic acid tert-butyl ester;

{2-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-carbamic acid tert-butyl ester;

3-Chloro-pyridine-2-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

5 Ethanesulfonic acid [5-(4-oxo-1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [5-(3-oxo-2,3,6,7-tetrahydro-1H,5H-pyrido[3,2,1-ij]quinolin-9-yl)-pyridin-3-ylmethyl]-amide;

10 3-Chloro-pyridine-2-carboxylic acid [5-(4-oxo-1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide;

3-Chloro-pyridine-2-carboxylic acid methyl-[5-(4-oxo-1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide ;

6-[5-(2-Amino-ethoxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;

15 (R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

6-[5-((R or S)-1-Amino-ethyl)-4-methyl-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;

20 3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

3,5-Dimethyl-isoxazole-4-carboxylic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

Ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

25 3-Chloro-pyridine-2-carboxylic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

N- {(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-methanesulfonamide;

3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;

5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',6'-dihydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid tert-butyl ester;

5 Ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-methyl-amide;

(R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

10 6-[5-((R or S)-1-Amino-ethyl)-4-chloro-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;

3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

(R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-amide;

15 6-[5-((R or S)-1-Amino-ethyl)-4-methyl-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;

6-[5-((S)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

20 1-Methyl-6-{5-[(S)-1-(1-methyl-1H-pyrazole-4-carbonyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,4-dihydro-1H-quinolin-2-one;

(S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid ethyl ester;

3,5-Dimethyl-isoxazole-4-carboxylic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

25 3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-amide;

5-Methyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;

5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',4',5',6'-tetrahydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid tert-butyl ester ;

N-{(R or S)-1-[4-Chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide ;

5 6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-1H-quinolin-2-one;

Ethanesulfonic acid [5-(8-fluoro-1-methyl-2-oxo-1,2-dihydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

N-{(R or S)-1-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-propionamide;

10 N-[4-Chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;

N-{(R or S)-1-[4-Methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide;

15 3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

and pharmaceutically acceptable salts thereof.

Further particular examples of compounds of formula (I) as described herein are selected from

20 6-[5-(2-Hydroxy-ethylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

25 6-[5-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

Ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

1-Methyl-6-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

- 5 6-[5-((R)-1-Hydroxymethyl-2-methyl-propylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

and pharmaceutically acceptable salts thereof.

Also further particular examples of compounds of formula (I) as described herein are selected from

- 10 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

1-Methyl-6-[5-((S)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

- 15 5-Methyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

1-Methyl-1H-pyrazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

- 20 3-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

- 25 7-Fluoro-1-methyl-6-[5-((R)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

3-Chloro-pyridine-2-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;

- 3-Chloro-pyridine-2-carboxylic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 3-Chloro-pyridine-2-carboxylic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 5 Ethanesulfonic acid {(trans)-4-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-amide;
- 1-Methyl-1H-pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 Ethanesulfonic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 6-{5-[2-(1,1-Dioxo-1λ6-isothiazolidin-2-yl)-ethoxy]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- Ethanesulfonic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 15 Ethanesulfonic acid {(S or R)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- Ethanesulfonic acid methyl-{2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;
- Ethanesulfonic acid [5-(4-oxo-1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide;
- 20 N-[(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl]-methanesulfonamide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;
- 25 5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',6'-dihydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid tert-butyl ester;
- Ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-methyl-amide;

3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

1-Methyl-6-{5-[(S)-1-(1-methyl-1H-pyrazole-4-carbonyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,4-dihydro-1H-quinolin-2-one;

5 and pharmaceutically acceptable salts thereof.


Processes for the manufacture of compounds of formula (I) as described herein are an object of the invention.

The preparation of compounds of formula (I) of the present invention may be carried out in sequential or convergent synthetic routes. Syntheses of the invention are shown in
10 the following general schemes. The skills required for carrying out the reaction and purification of the resulting products are known to those persons skilled in the art. In case a mixture of enantiomers or diastereoisomers is produced during a reaction, these enantiomers or diastereoisomers can be separated by methods described herein or known to the man skilled in the art such as e.g. chiral chromatography or crystallization. The
15 substituents and indices used in the following description of the processes have the significance given herein.

The following abbreviations are used in the present text:


AcOH = acetic acid, BOC = t-butyloxycarbonyl, BuLi = butyllithium, CDI = 1,1-carbonyldiimidazole, CH₂Cl₂ = dichloromethane, DBU = 2,3,4,6,7,8,9,10-octahydro-pyrimido[1,2-a]azepine, DCE = 1,2-dichloroethane, DCM = dichloromethane, DIBALH =
20 di-*i*-butylaluminium hydride, DCC = N,N'-dicyclohexylcarbodiimide, DMA = *N,N*-dimethylacetamide, DMAP = 4-dimethylaminopyridine, DMF = *N,N*-dimethylformamide, EDCI = *N*-(3-dimethylaminopropyl)-*N*'-ethylcarbodiimide hydrochloride, EtOAc = ethylacetate, EtOH = ethanol, Et₂O = diethylether, Et₃N = triethylamine, eq = equivalents,
25 HATU = O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate, HPLC = high performance liquid chromatography, HOBT = 1-hydroxybenzo-triazole, Huenig's base = *i*Pr₂NEt = *N*-ethyl diisopropylamine, IPC = in process control, LAH = lithium aluminium hydride, LDA = lithium diisopropylamide, LiBH₄ = lithium borohydride, MeOH = methanol, NaBH₃CN, sodium cyanoborohydride, NaBH₄ = sodium
30 borohydride, NaI = sodium iodide, Red-Al = sodium bis(2-methoxyethoxy) aluminium hydride, RT = room temperature, TBDMSCl = t-butyldimethylsilyl chloride, TFA = trifluoroacetic acid, THF = tetrahydrofuran, quant = quantitative.

Scheme 1a



X is Halogen or OSO_2CF_3

R^{101} and R^{102} e.g. together with the boron atom to which they are attached form




R^{103} stands for substituents as e.g. shown in Schemes 2a and 2b, which allow further transformation into R^6 at a later stage in the synthesis

- Lactam compounds **1** (Scheme 1a) are known or can be prepared by methods described herein or known to the man skilled in the art (see also Scheme 1b for alternative
- 5 syntheses of compounds **5** and **7**); compounds **1** can be alkylated at nitrogen using a base like sodium hydride or sodium or potassium *tert*-butoxide, followed by addition of an alkylating agent of formula $\text{R}^1\text{-X}$, wherein X is halogen, tosylate or mesylate, in a solvent like DMF or THF preferably in a temperature range between 0 °C and about 80 °C giving N-alkylated lactams **2** (step a).
- 10 Reaction of lactams **2** with e.g. 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-5 dioxaborolane) in solvents like dimethylsulfoxide or dioxane in the presence of potassium acetate and catalysts like (1,1'-bis-diphenylphosphino)-ferrocene)palladium-(II)dichloride (1:1 complex with dichloromethane) at temperatures up to about 100 °C gives boronic ester compounds **3** (step b). Condensation of boronic ester compounds **3** with suitable aryl
- 15 halides **4** or **6** (for possible syntheses of aryl halides **4** or **6**, see Schemes 2) can be performed using *Suzuki* conditions, e.g. in the presence of catalysts, such as *tri*-o-tolylphosphine/palladium(II)acetate, *tetrakis*-(triphenylphosphine)-palladium, *bis*(triphenylphosphine)palladium(II)chloride or dichloro[1,1'-bis(diphenylphosphino)-


ferrocene]palladium(II) optionally in the form of a dichloromethane complex (1:1), and in the presence of a base, such as aqueous or non aqueous potassium phosphate, sodium or potassium carbonate, in a solvent, such as dimethylsulfoxide, toluene, ethanol, dioxane, tetrahydrofuran or *N,N*-dimethylformamide, and in an inert atmosphere such as argon or nitrogen, in a temperature range preferably between room temperature and about 130 °C leading to adducts **5** or **7** (steps c). Compounds **7** can be further transformed into compounds of the general formula **5** by methods described in the following Schemes, the examples or by methods well known to persons skilled in the art (step d).

Halo-nitro pyridine compounds **8** (Scheme 1b) with at least one hydrogen substituent R⁷ ortho to the nitro group react with 1-chloro-1-R⁵-methanesulfonyl-4-methylbenzene in solvents like THF and in the presence of a base like *tert*-BuOK in a temperature range between -78 °C and room temperature to give regioisomeric sulfones **9** and **10** (step a). Treatment of sulfones **9** and **10** with a haloacetic acid ester compound in a solvent like *N,N*-dimethylformamide and in the presence of a weak base as *e.g.* sodium or potassium carbonate preferably in a temperature range between room temperature and about 80 °C gives acetic acid ester adducts **11** and **16** (step b). *Suzuki* reactions of adducts **11** and **16** with suitable heteroaryl-boronic acid derivatives under conditions as described for step c (Scheme 1a) gives adducts **12** and **17** containing acrylic ester moieties by concomitant elimination of the 4-methyl-benzene-sulfonyl groups (step c). Catalytic hydrogenation *e.g.* using Pd/C and AcOH in methanol at elevated temperature and with H₂ pressure of about 50-200 psi gives lactam compounds **13** and **18** (step d). Treatment of lactam compounds **13** and **18** with an alkylating agent like an alkyl or cycloalkyl halide, alkyl or cycloalkyl tosylate or an alkyl or cycloalkyl mesylate in a solvent like THF or *N,N*-dimethylformamide in the presence of a base like sodium or potassium hydride preferable around 0 °C gives alkylated lactam compounds **14** and **19** (step e). Alkylated lactam compounds **14** and **19** can be further transformed into compounds of the general formula **15** or **20** by methods described in the following Schemes, the examples or by methods well known to persons skilled in the art (step f).

Scheme 1b



Scheme 2a



X is Halogen or OSO₂CF₃

Schemes 2a and 2b describe possible syntheses of aryl halide compounds **103**, **105**, **110**, **114**, **117** and **121** which correspond to aryl halides **4** and **6** in Scheme 1. The terms linker used in Scheme 2a are defined as follows:

5 Linker^a = -O_m-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q-, -N_mR¹⁷-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q- or -S_m(O)_r-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q-;

Linker^b = -O_m-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-, -N_mR¹⁷-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p- or -S_m(O)_r-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-;

Linker^c = -O_m-(CR⁹R¹⁰)_n-, -N_mR¹⁷-(CR⁹R¹⁰)_n- or -S_m(O)_r-(CR⁹R¹⁰)_n-.

Compounds **101** (Scheme 2a) carrying an aliphatic Linker^a react with amino compounds **102** either per se or after anion formation *e.g.* with sodium hydride in solvents like *N,N'*-dimethylformamide in a temperature range between 0 °C and about 100 °C to form adducts **103** (step a). Compounds **101** in which Linker^a is absent react with amino compounds **102** (R¹⁶ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, phenylalkyl, substituted phenylalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl or substituted heteroaryl) either directly in the presence of a base like Hunig's base in a solvent like butanol and elevated temperature or by using Pd(0)-catalyzed amination reactions (*Buchwald-Hartwig* coupling) under an inert atmosphere such as argon or nitrogen in the presence of a palladium catalyst such as *tris*(dibenzylideneacetone)dipalladium(0) (Pd₂(dba)₃) or palladium(II) acetate (Pd(COOCH₃)₂), a phosphine ligand like triphenylphosphine, *rac*-2,2'-*bis*(diphenylphosphino)-1,1'-binaphthalene (*rac*-BINAP) or (R)-(-)-1-[(S)-2-(dicyclohexylphosphino)ferrocenyl]ethyl-di-*tert*-butylphosphine and a base such as Cs₂CO₃ or KO*tert*-Bu in a solvent like toluene, ethanol or water or mixtures thereof at room temperature or elevated temperatures, whereby heating might be achieved conventionally or by microwave irradiation to give substituted adducts **103** in which Linker^a is absent (step a).

Compounds **101** in which Linker^a is absent react with amino compounds **102** (R¹⁶ is -S(O)₂R¹⁸, -S(O)₂OR¹⁸, -S(O)₂NR¹⁸R¹⁹, -C(O)R¹⁸, -C(O)OR¹⁸ or -C(O)NR¹⁸R¹⁹) in solvents like 1,4-dioxane, in the presence of copper (I) iodide, potassium or cesium carbonate, and a chelating 1,2-diamino compound like *N,N'*-dimethylethylenediamine or trans-1,2-diamino-hexane, at elevated temperatures, preferable with the aide of microwave heating to give substituted adducts **103** in which Linker^a is absent (step a).

Suitable reductive amination procedures as *e.g.* treatment of aldehydes or ketones **104** and suitable amino-moieties with NaBH(OAc)₃ in a one step procedure in a solvent like methanol preferably around room temperature or in a two step procedure by first treatment with titanium (IV) isopropoxide in solvents like methanol or toluene preferably at temperatures between room temperature and the reflux temperature of the solvents followed by reaction with NaBH₄ preferably between 0 °C and room temperature converts aldehydes or ketones **104** into amino compounds **105**; alternatively imines obtained after treatment with titanium (IV) isopropoxide can be evaporated, then be re-dissolved in a solvent like THF and being treated with a Grignard reagent R¹³MgX, preferably between -40 °C and 0 °C leading to amino compounds **105** carrying the specific R¹³ substituent (step b).

Aldehydes and ketones **106** react in *Horner-Emmons* reactions using *e.g.* reagents like dimethyl(methoxycarbonyl)methylphosphonate, optionally carrying an additional R¹³ substituted at the methylene group, and a base like sodium hydride in a solvent like tetrahydrofuran preferable between about 0 °C and the reflux temperature of the solvent to
5 give unsaturated esters **107** (step c). Reduction of the double bond in unsaturated esters **107** can be performed *e.g.* by using a mixture of nickel chloride and sodiumborohydride as reducing agents in solvents like methanol preferably between about 0 °C and room temperature leading to ester compounds **108** (step d). Alpha mono- or di-substituted esters **109** can be synthesized from esters **108**, by treatment with a base like LDA or HMDS in
10 solvents like tetrahydrofuran or 1,2-dimethoxyethane, followed by addition of one or sequentially two different alkyl halides, a reaction preferably performed between -78 °C and room temperature (step e). Esters **109** can be further transformed into amino compounds **110** suitable to function as compounds **4** or **6** in Scheme 1, *e.g.* by amide formation with ammonia in methanol around room temperature, *Hofmann* rearrangement,
15 treatment with sodium hydroxide, and bromine in a solvent like ethanol preferably between about 0 °C and the reflux temperature of the solvent and subsequent introduction of R¹⁵ and R¹⁶ substituents (steps f, g, h) by methods known to the man skilled in the art.

Amino compounds **110** can also be obtained by hydrolysis of esters **109** to the corresponding acids followed by treatment with diphenylphosphoryl azide, TEA in a
20 solvent like toluene preferably at reflux and subsequent treatment with *e.g.* sodium trimethylsilanoate in a solvent like THF preferably around RT followed by subsequent introduction of R¹⁵ and R¹⁶ substituents (steps f, g, h) by methods known to the man skilled in the art.


Di-halogen or di-triflate substituted heteroaromatic compounds **111** (Scheme 2b)
25 react with unsaturated boronic acid derivatives **112** under *Suzuki* conditions as described in Scheme 1, step c, to adducts **113** (step i). Removal of the double bond in compounds **113** *e.g.* by catalytic hydrogenation, followed by additional standard synthetic modifications transforms compounds **113** into synthons **114** (step k).

Reaction of compounds **111** with amino compounds **115**, performed under
30 conditions as described for the transformation of compounds **101** in which Linker^a is absent into compounds **103**, is giving compounds **116** or **117** (step l). Compounds **116** can be transformed by additional standard synthetic modifications into synthons **117** (step m).


Phenols or thiophenols **118** react with alcohols **119** under *Mitsunobu* conditions *e.g.* with triphenylphosphine and di-*tert*-butyl-, diisopropyl-, diethyl-azodicarboxylate or di-(4-

chlorobenzyl)azodicarboxylate as reagents in solvents like toluene, dichloromethane or tetrahydrofuran preferably at ambient temperature to give adducts **120** or **121** (step n). Compounds **120** can be transformed by additional modifications into synthons **121**, such modifications include oxidation at sulfur to the respective sulfoxide or sulfone *e.g.* by using *m*-chloroperbenzoic acid (step o).

Scheme 2b



Also an embodiment of the present invention is a process to prepare a compound of formula (I) as defined above comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III);



wherein A^1 , A^2 , A^3 , R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined in claim 1 and wherein X is halogen or triflate, R^{101} and R^{102} are alkyl, cycloalkyl or together with the boron atom they are attached to form together a borolanyl.

- 5 In particular in the presence of catalysts, such as *tri-o*-tolylphosphine/palladium(II)acetate, *tetrakis*-(triphenylphosphine)-palladium, *bis*(triphenylphosphine)palladium(II)chloride or dichloro[1,1'-*bis*(diphenylphosphino)-ferrocene]palladium(II) optionally in the form of a dichloromethane complex (1:1), and in the presence of a base, such as aqueous or non aqueous potassium phosphate, sodium or
- 10 potassium carbonate, in a solvent, such as dimethylsulfoxide, toluene, ethanol, dioxane, tetrahydrofuran or *N,N*-dimethylformamide, and in an inert atmosphere such as argon or nitrogen, in a temperature range particularly between room temperature and about 130 °, wherein A^1 , A^2 , A^3 , R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined in claim 1 and wherein X is halogen or triflate, R^{103} and R^{104} are alkyl, cycloalkyl or together with the boron atom they
- 15 are attached to form together a borolanyl.

Also an object of the present invention is a compound according to formula (I) as described herein for use as therapeutically active substance.

- Likewise an object of the present invention is a pharmaceutical composition comprising a compound according to formula (I) as described herein and a therapeutically
- 20 inert carrier.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrome.

- The present invention also relates to the use of a compound according to formula (I)
- 25 as described herein for the treatment or prophylaxis of chronic kidney disease.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of congestive heart failure.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of hypertension.

- 5 The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of primary aldosteronism.

A particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.

- 10 Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of chronic kidney disease.

Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of congestive heart failure.

- 15 Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of hypertension.

Also a particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of primary aldosteronism.

- 20 The present invention also relates to the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.

Also an embodiment of the present invention is the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of chronic kidney disease.

- 25 Also an embodiment of the present invention is the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of congestive heart failure.

- 30 Also an embodiment of the present invention is the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of hypertension.

Also an embodiment of the present invention is the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of primary aldosteronism.

Also an object of the invention is a method for the treatment or prophylaxis of
5 chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Also an embodiment of the present invention is a method for the treatment or prophylaxis of chronic kidney disease, which method comprises administering an effective
10 amount of a compound according to formula (I) as described herein.

Also an embodiment of the present invention is a method for the treatment or prophylaxis of congestive heart failure, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Also an embodiment of the present invention is a method for the treatment or
15 prophylaxis of hypertension, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

Also an embodiment of the present invention is a method for the treatment or prophylaxis of primary aldosteronism, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

20 formula (I) as described herein, when manufactured according to any one of the described processes.

Assay procedures

Herein we identified the use of the G-402 cell line as a host cell to ectopically express (transiently or stably) enzymes of the CYP11 family. Specifically we developed
25 stable G-402 cells expressing ectopically human CYP11B1, human CYP11B2, human CYP11A1, cynmolgus CYP11B1 or cynomolgus CYP11B2 enzyme activity. Importantly the identified cell line G-402 expresses co-factors (adrenodoxin and adrenodoxin reductase) important for the activity of the CYP11 family and no relevant enzyme activity of the CYP11 family (in comparison to H295R cells) was detected in these cells. Therefore
30 the G-402 cell line is uniquely suited as a host cell for the ectopic expression of enzymes from the CYP11 family.

G-402 cells can be obtained from ATCC (CRL-1440) and were originally derived from a renal leiomyoblastoma.

The expression plasmids contains the ORF for either human / cyno CYP11B1 or CYP11B2 under the control of a suitable promoter (CMV-promoter) and a suitable resistance marker (neomycin). Using standard techniques the expression plasmid is transfected into G-402 cells and these cells are then selected for expressing the given resistance markers. Individual cell-clones are then selected and assessed for displaying the desired enzymatic activity using 11-Deoxycorticosterone (Cyp11B2) or 11-Deoxycortisol (Cyp11B1) as a substrate.

G-402 cells expressing CYP11 constructs were established as described above and maintained in McCoy's 5a Medium Modified, ATCC Catalog No. 30-2007 containing 10% FCS and 400 µg/ml G418 (Geneticin) at 37 °C under an atmosphere of 5% CO₂/95% air. Cellular enzyme assays were performed in DMEM/F12 medium containing 2.5 % charcoal treated FCS and appropriate concentration of substrate (0.3-10 uM 11-Deoxycorticosterone, 11-Deoxycortisol or Corticosterone). For assaying enzymatic activity, cells were plated onto 96 well plates and incubated for 16h. An aliquot of the supernatant is then transferred and analyzed for the concentration of the expected product (Aldosterone for CYP11B2; Cortisol for CYP11B1). The concentrations of these steroids can be determined using HTRF assays from CisBio analyzing either Aldosterone or Cortisol.

Inhibition of the release of produced steroids can be used as a measure of the respective enzyme inhibition by test compounds added during the cellular enzyme assay. The dose dependent inhibition of enzymatic activity by a compound is calculated by means of plotting added inhibitor concentrations (x-axes) vs. measured steroid/product level (y-axes). The inhibition is then calculated by fitting the following 4-parameter sigmoidal function (Morgan-Mercer-Flodin (MMF) model) to the raw data points using the least squares method:

$$y = \frac{AB + Cx^D}{B + x^D}$$

wherein, A is the maximum y value, B is the EC₅₀ factor determined using XLFit, C is the minimum y value and D is the slope value.

The maximum value A corresponds to the amount of steroid produced in the absence of an inhibitor, the value C corresponds to the amount of steroid detected when the enzyme is fully inhibited.

- EC50 values for compounds claimed herein were tested with the G402-based assay system described. Cyp11B2 enzyme activity was tested in presence of 1 μ M Deoxycorticosterone and variable amounts of inhibitors; Cyp11B1 enzyme activity was tested in presence of 1 μ M Deoxycortisol and variable amounts of inhibitors.

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
1	0.170	3.70
2	0.430	>3
3	0.715	>3
4	0.011	0.35
5	0.019	1.08
6	0.098	3.34
7	0.056	0.57
8	0.049	2.33
9	0.829	>10
10	0.020	0.17
11	0.076	0.41
12	0.002	0.05
13	0.319	17.89

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
14	0.308	1.57
15	0.019	1.20
16	0.011	0.86
17	4.649	>10
18	0.058	0.75
19	0.023	0.80
20	0.003	0.02
21	0.021	0.26
22	0.025	0.10
23	0.032	0.39
24	0.007	0.22
25	0.005	0.11
26	0.018	0.57

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
27	0.007	0.27
28	0.121	0.78
29	0.260	5.56
30	0.170	10.22
31	0.172	>3
32	1.293	>10
33	1.256	>10
34	0.416	29.22
35	0.325	>10
36	0.071	2.57
37	0.044	1.59
38	0.081	1.50
39	0.010	0.12
40	0.015	0.75
41	0.067	5.56
42	0.028	0.16
43	0.040	1.56
44	0.005	0.568

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
45	0.087	0.863
46	0.005	0.298
47	0.003	0.061
48	0.049	0.392
49	0.271	1.003
50	0.003	0.024
51	0.032	1.872
52	0.030	0.139
53	0.008	0.453
54	n.d.	3.855
55	0.682	11.133
56	0.040	0.587
57	0.026	0.393
59	0.033	0.710
60	0.685	n.d.
61	0.508	20.108
62	0.177	8.543
63	0.731	15.994

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
64	0.423	12.328
65	0.315	13.084
66	0.710	n.d.
67	0.160	7.675
68	0.350	33.719
69	0.411	3.492
70	0.202	15.608
71	0.081	2.497
72	0.007	0.101
73	0.017	0.575
74	0.286	n.d.
75	0.008	0.509
76	0.023	0.452
77	0.042	0.223
78	0.121	1.145
79	0.035	0.302
80	0.043	0.493
81	0.172	4.635

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
82	0.008	0.020
83	0.009	0.046
84	0.034	0.526
85	0.003	0.048
86	0.004	0.108
87	0.017	1.066
88	0.001	0.013
89	0.014	0.276
90	0.011	1.497
91	0.043	3.068
92	0.016	0.632
93	0.023	0.613
94	0.023	1.336
95	0.017	1.781
96	0.022	1.676
97	0.006	0.111
98	0.011	0.360
99	0.035	1.117

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
100	0.007	0.140
101	0.002	0.179
102	0.014	1.441
103	0.015	2.927
104	0.010	0.094
105	0.086	2.581
106	0.020	2.683
107	0.002	0.044
108	0.032	1.110
109	0.078	0.712
110	0.006	0.187
111	0.004	0.081
112	0.006	0.151
113	0.075	1.575
114	0.008	0.142
115	0.001	0.014
116	0.003	0.081
117	0.008	0.121

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
118	0.004	0.682
119	0.006	0.075
120	0.000	0.008
121	0.019	1.371
122	0.009	0.535
123	0.062	2.917
124	0.005	0.138
125	0.003	0.154
126	0.007	0.642
127	0.012	0.400
128	0.119	2.286
129	0.030	0.613
130	0.162	10.483
131	0.003	0.440
132	0.029	4.800
133	0.010	0.837
134	0.003	0.094
135	0.006	0.163

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
136	0.047	2.433
137	0.006	0.872
138	0.003	0.267
139	0.018	0.455
140	0.114	6.646
141	0.856	19.889
142	0.003	0.380
143	0.009	0.051
144	0.004	0.460
145	0.175	3.779
151	0.005	0.090
152	0.154	4.752
153	0.375	11.100
154	0.114	1.427
155	0.047	0.589
156	0.155	1.374
157	0.463	3.256
158	0.239	6.429

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
159	0.180	7.933
160	0.113	0.716
161	0.001	0.008
162	0.062	2.853
163	0.061	2.387
164	0.009	0.207
165	0.013	0.494
166	0.022	1.206
167	0.018	0.898
168	0.014	0.273
169	0.035	0.850
170	0.015	0.699
171	0.009	0.420
172	0.001	0.069
173	0.031	0.800
174	0.033	1.225
175	0.007	0.688
176	0.128	n.d.

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
177	0.117	8.464
178	0.010	0.535
179	0.012	0.169
180	0.033	1.891
181	0.025	2.320
182	0.006	0.302
183	0.004	0.268
184	0.227	9.741
185	0.633	9.289
186	0.011	0.199
187	0.027	0.705
188	0.014	0.572
189	0.002	0.093
190	0.007	0.442
191	0.010	0.777
192	0.029	1.981
193	0.022	1.009
194	0.041	1.704

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
195	0.001	0.056
196	0.008	0.514
197	0.004	0.062
198	0.007	0.188
199	0.003	0.028
200	0.001	0.031
201	0.024	0.436
202	0.013	0.170
203	0.083	4.957
204	0.020	2.237
205	0.097	4.025
206	0.010	0.237
207	0.044	2.543
208	0.059	4.202
209	0.162	7.544
210	0.030	1.344
211	0.005	0.119
212	0.051	4.834

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
213	0.041	0.834
214	0.030	0.104
216	0.042	0.879
217	0.029	1.256
218	0.029	1.196
219	0.012	0.557
220	0.002	0.040
222	0.004	1.008
223	0.036	2.855
224	0.006	0.173
225	0.006	0.691
226	0.011	1.767
227	0.004	0.157
228	0.009	0.356
229	0.010	0.656
230	0.001	0.007
231	0.019	0.802
232	0.001	0.016

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
233	0.013	0.219
234	0.027	2.258
235	0.109	4.662
236	0.003	0.077
237	0.008	0.143
238	0.005	0.513
239	0.021	0.401
240	0.250	15.012
241	0.007	0.251
242	0.100	3.686
243	0.001	0.026
244	0.002	0.075
245	0.001	0.072
246	0.013	4.723
247	0.005	0.069
248	0.000	0.007
249	0.014	0.261
250	0.035	1.247

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
251	0.001	0.006
252	0.026	0.468
253	0.193	4.475
254	0.001	0.049
255	0.013	1.325
256	0.008	0.158
257	0.022	0.679
258	0.013	0.291
259	0.001	0.154

Example	EC50 human CYP11B2 μM	EC50 human CYP11B1 μM
260	0.000	0.001
261	0.062	1.222
262	0.084	1.817
263	3.101	n.d.
264	0.024	2.921
265	0.009	0.623
266	0.054	0.623
267	0.001	0.236

Compounds of formula (I) and their pharmaceutically acceptable salts or esters thereof as described herein have EC₅₀ (CYP11B2) values between 0.000001 uM and 1000 uM, particular compounds have EC₅₀ (CYP11B2) values between 0.00005 uM and 500 uM, further particular compounds have EC₅₀ (CYP11B2) values between 0.0005 uM and 5 uM. These results have been obtained by using the described enzymatic assay.

The compounds of formula (I) and their pharmaceutically acceptable salts can be used as medicaments (e.g. in the form of pharmaceutical preparations). The pharmaceutical preparations can be administered internally, such as orally (e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatin capsules, solutions, emulsions or suspensions), nasally (e.g. in the form of nasal sprays) or rectally (e.g. in the form of suppositories). However, the administration can also be effected parentally, such as intramuscularly or intravenously (e.g. in the form of injection solutions).

The compounds of formula (I) and their pharmaceutically acceptable salts can be processed with pharmaceutically inert, inorganic or organic adjuvants for the production of tablets, coated tablets, dragées and hard gelatin capsules. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts etc. can be used, for example, as such

5 adjuvants for tablets, dragées and hard gelatin capsules.

Suitable adjuvants for soft gelatin capsules, are, for example, vegetable oils, waxes, fats, semi-solid substances and liquid polyols, etc.

Suitable adjuvants for the production of solutions and syrups are, for example, water, polyols, saccharose, invert sugar, glucose, etc.

10 Suitable adjuvants for injection solutions are, for example, water, alcohols, polyols, glycerol, vegetable oils, etc.

Suitable adjuvants for suppositories are, for example, natural or hardened oils, waxes, fats, semi-solid or liquid polyols, etc.

Moreover, the pharmaceutical preparations can contain preservatives, solubilizers,

15 viscosity-increasing substances, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

The dosage can vary in wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily

20 dosage of about 0.1 mg to 20 mg per kg body weight, preferably about 0.5 mg to 4 mg per kg body weight (e.g. about 300 mg per person), divided into preferably 1-3 individual doses, which can consist, for example, of the same amounts, should be appropriate. It will, however, be clear that the upper limit given herein can be exceeded when this is shown to be indicated.

25 In accordance with the invention, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis of aldosterone mediated diseases.

The compounds of formula (I) or their pharmaceutically acceptable salts and esters herein display also variable inhibition of CYP11B1. These compounds may be used for the

30 inhibition of CYP11B1 in combination with variable inhibition of CYP11B2. Such compounds may be used for treatment or prophylaxis of conditions displaying excessive cortisol production/levels or both excessive cortisol and aldosterone levels (for ex.

Cushing syndrome, burn trauma patients, depression, post-traumatic stress disorders, chronic stress, corticotrophic adenomas, Morbus Cushing).

In accordance with the invention, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis
5 of cardiovascular conditions (including hypertension and heart failure), renal conditions, liver conditions, vascular conditions, inflammatory conditions, pain, retinopathy, neuropathy (such as peripheral neuropathy), insulinopathy, edema, endothelial dysfunction, baroreceptor dysfunction; fibrotic diseases, depression and the like.

Cardiovascular conditions include congestive heart failure, coronary heart disease,
10 arrhythmia, arterial fibrillation, cardiac lesions, decreased ejection fraction, diastolic and systolic heart dysfunction, fibrinoid necrosis of coronary arteries, heart failure, hypertrophic cardiomyopathy, impaired arterial compliance, impaired diastolic filling, ischemia, left ventricular hypertrophy, myocardial and vascular fibrosis, myocardial infarction, myocardial necrotic lesions, myocardial necrotic lesions cardiac arrhythmias,
15 prevention of sudden cardiac death, restenosis, stroke, vascular damage.

Renal conditions include acute and chronic renal failure, end-stage renal disease, decreased creatinine clearance, decreased glomerular filtration rate, diabetic nephropathy, expansion of reticulated mesangial matrix with or without significant hypercellularity, focal thrombosis of glomerular capillaries, global fibrinoid necrosis, glomerulosclerosis,
20 ischemic lesions, malignant nephrosclerosis (such as ischemic retraction, microalbuminuria, nephropathy, proteinuria, reduced renal blood flow, renal arteriopathy, swelling and proliferation of intracapillary (endothelial and mesangial) and/or extracapillary cells (crescents).

Liver conditions include, but are not limited to, liver cirrhosis, liver ascites, hepatic
25 congestion, nonalcoholic steatohepatitis and the like.

Vascular conditions include, but are not limited to, thrombotic vascular disease (such as mural fibrinoid necrosis, extravasation and fragmentation of red blood cells, and luminal and/or mural thrombosis), proliferative arteriopathy (such as swollen myointimal cells surrounded by mucinous extracellular matrix and nodular thickening),
30 atherosclerosis, decreased vascular compliance (such as stiffness, reduced ventricular compliance and reduced vascular compliance), endothelial dysfunction, and the like.

Inflammatory conditions include, but are not limited to, arthritis (for example, osteoarthritis), inflammatory airways diseases (for example, chronic obstructive pulmonary disease (COPD)), and the like.

Pain includes, but is not limited to, acute pain, chronic pain (for example, arthralgia),
5 and the like.

Edema includes, but is not limited to, peripheral tissue edema, hepatic congestion, splenic congestion, liver ascites, respiratory or lung congestion, and the like.

Insulinopathies include, but are not limited to, insulin resistance, Type I diabetes mellitus, Type II diabetes mellitus, glucose sensitivity, pre-diabetic state, syndrome X, and
10 the like.

Fibrotic diseases include, but are not limited to myocardial and intrarenal fibrosis, renal interstitial fibrosis and liver fibrosis.

Furthermore, the compounds of formula (I) or their pharmaceutically acceptable salts and esters as described herein can also be used for the treatment or prophylaxis of
15 cardiovascular condition selected from the group consisting of hypertension, heart failure (particularly heart failure post myocardial infarction), left ventricular hypertrophy, and stroke.

In another embodiment, the cardiovascular condition is hypertension.

In another embodiment, the cardiovascular condition is heart failure.

20 In another embodiment, the cardiovascular condition is left ventricular hypertrophy.

In another embodiment, the cardiovascular condition is stroke.

In another embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis renal condition.

In another embodiment, the renal condition is nephropathy.

25 In another embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis Type II diabetes mellitus

In another embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis Type I diabetes mellitus

The invention is illustrated hereinafter by Examples, which have no limiting
5 character.


In case the preparative examples are obtained as a mixture of enantiomers, the pure enantiomers can be separated by methods described herein or by methods known to the man skilled in the art, such as e.g. chiral chromatography or crystallization.

Examples


All examples and intermediates were prepared under argon atmosphere if not specified otherwise.

5 Intermediate A-1

1-Methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one




[A] 6-Bromo-1-methyl-3,4-dihydro-1H-quinolin-2-one



To a solution of 6-bromo-3,4-dihydroquinolin-2(1H)-one (5 g, 22.1 mmol) in DMF (100 mL) cooled to 0°C was added potassium *tert*-butoxide (4.96 g, 44.2 mmol) portionwise and the reaction mixture was stirred at 0 °C for 15 min. Then, methyl iodide (4.08 g, 28.8 mmol) was added and the reaction mixture allowed to warm up to room temperature and stirring was continued over night. More MeI (1.25 g, 8.86 mmol) was added and the reaction mixture was heated to 40 °C until completion of the reaction. The mixture was diluted with EtOAc, poured into 100 mL of 1M HCl and the aqueous phase was extracted with EtOAc (2 x 200 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 30% EtOAc-heptane gradient to give the title compound (4.23 g, 80 %) as an off white solid. MS: 240.0, 242.1 (M+H⁺).


[B] 1-Methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one




A flask was charged with 6-bromo-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (3 g, 12.5 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-*bi*(1,3,2-dioxaborolane) (3.81 g, 15.0 mmol), potassium acetate (3.68 g, 37.5 mmol) and dioxane (48 mL). The mixture was purged with Ar, then dichloro[1,1'-*bis*(diphenylphosphino)-ferrocene]palladium(II) dichloromethane complex (1:1) [PdCl₂(DPPF)-CH₂Cl₂ adduct] (457 mg, 0.625 mmol) was added and the resulting mixture was heated to 80 °C over night. The reaction mixture was diluted with EtOAc, filtered through Dicalite and washed with EtOAc (2 x 150 mL). The resulting filtrate was washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 40% EtOAc-heptane gradient to give the title compound (2.63 g, 73 %) as an off white solid. MS: 288.0 (M+H⁺).

Intermediate A-2

6-(5-Amino-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride




[A] 2,2-Dimethyl-*N*-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-propionamide



A microwave vial was charged with *N*-(5-bromopyridin-3-yl)pivalamide (0.2 g, 0.778 mmol), 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) (0.246 g, 0.856 mmol) and DMF (4 mL). The reaction mixture was purged with Argon; then, *bis*(triphenylphosphine)palladium(II)chloride (0.055 g, 0.078 mmol), followed by 1*N* aqueous Na₂CO₃ solution (2.33 mL, 2.33 mmol) were added and the reaction was heated in the microwave at 120 °C for 5 min. The reaction mixture was diluted with EtOAc, filtered through Dicalite and washed with EtOAc (2 x 20 mL). The resulting filtrate was washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.195 g, 74 %) as a white solid. MS: 338.2 (M+H⁺).

[B] 6-(5-Amino-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride




A suspension of 2,2-dimethyl-*N*-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-propionamide (0.195 g, 0.578 mmol) in 1*N* aqueous HCl (5.78 mL) was heated to 90 °C over night. The reaction mixture was evaporated to dryness and the resulting solid material was triturated in MeOH, filtered off and further dried in a the high vacuum to give the title compound (0.115 g, 69 %) as an off white solid as the HCl salt. MS: 254.1 (M+H⁺).


20

Intermediate A-3

6-{5-[(*S*)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




[A] (S)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidine



To a solution of (S)-pyrrolidin-2-yl-methanol (0.69 g, 6.82 mmol) in DCM (3 mL) cooled to 0 °C was added TEA (1.38 g, 13.6 mmol) followed by TBDMS-Cl (1.03 g, 6.82 mmol) in DCM (3 mL). The reaction mixture was then stirred at room temperature over night and poured into NH₄Cl (20 mL). The aqueous layer was extracted with DCM (2 x 50 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (1.11 g, 76%) as a yellow oil. MS: 216.2 (M+H⁺).

10 [B] 3-Bromo-5-[(S)-2-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridine




To a solution of (S)-2-((*tert*-butyldimethylsilyloxy)methyl)pyrrolidine (0.455 g, 2.11 mmol) in toluene (20 mL) were added *tris*(dibenzylideneacetone)dipalladium(0) (Pd₂(dba)₃) (0.039 g, 0.042 mmol) and *rac*-2,2'-bis(diphenylphosphino)-1,1'-binaphthalene (*rac*-BINAP) (0.066 g, 0.106 mmol). The solution was purged with Argon and heated to 85 °C for 10 min. After cooling to room temperature, sodium *tert*-butoxide

(0.406 g, 4.22 mmol) and 3,5-dibromopyridine (0.5 g, 2.11 mmol) were added and the reaction mixture was then heated to 85 °C over night. The mixture was poured into sat. NH₄Cl (20 mL) and the aqueous layer was extracted with DCM (2 x 25 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness.

- 5 The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.412 g, 53 %) as a yellow oil. MS: 371.0, 372.9 (M+H⁺).

[C] 6-{5-[(S)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




10

A microwave vial was charged with 3-bromo-5-[(S)-2-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridine (0.091 g, 0.245 mmol), 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1, 0.077 g, 0.270 mmol) and DMF (1.5 mL). The reaction mixture was purged with Argon;


- 15 then, *bis*(triphenylphosphine)palladium(II)chloride (0.017 g, 0.025 mmol), followed by a 1N Na₂CO₃ aqueous solution (0.980 mL, 0.980 mmol) were added and the reaction mixture was heated in the microwave at 120 °C for 30 min. The reaction mixture was diluted with EtOAc, poured into aq. NaHCO₃ (10 mL) and the aqueous layer was extracted with EtOAc (2 x 40 mL). Combined organics were washed with brine, dried over Na₂SO₄,
20 filtered and evaporated. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.1 g, 90%) as a white solid. MS: 452.1 (M+H⁺).

Intermediate A-4

- 25 **6-{5-[(R)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one**




[A] (R)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidine




In analogy to the procedure described for the preparation of intermediate A-3 [A], (R)-pyrrolidin-2-yl-methanol was reacted with TBDMS-Cl in the presence of TEA to give the
5 title compound as a yellow oil. MS: 216.3 ($M+H^+$).

[B] 3-Bromo-5-[(R)-2-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridine



In analogy to the procedure described for the preparation of intermediate A-3 [B], (R)-2-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidine was reacted with 3,5-dibromopyridine
10 in the presence of $Pd_2(dba)_3$, *rac*-BINAP and sodium *tert*-butoxide to give the title compound as a yellow oil. MS: 371.0, 372.9 ($M+H^+$).


[C] 6-{5-[(R)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




In analogy to the procedure described for the preparation of intermediate A-3 [C], 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1) has been coupled to 3-bromo-5-[(R)-2-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridine to give the title compound as a white solid. MS: 452.1 (M+H⁺).

Intermediate A-5

3-Bromo-5-((S)-2-[1,2,4]triazol-1-ylmethyl-pyrrolidin-1-yl)-pyridine




[A] [(S)-1-(5-Bromo-pyridin-3-yl)-pyrrolidin-2-yl]-methanol



To a solution of 3-bromo-5-[(S)-2-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridine (intermediate A-3 [B], 0.1 g, 0.269 mmol) in MeOH (2 mL) was added 4M HCl in dioxane (0.202 mL, 0.808 mmol) and the reaction mixture was stirred at room temperature for 1h. The mixture was evaporated to dryness, the residue diluted with DCM, poured into aq. NaHCO₃ (10 mL) and extracted with DCM (25 mL). The organic layer


was dried over Na₂SO₄, filtered and evaporated to give the title compound (0.060 g, 87 %) as a yellow oil. MS: 257.0, 259.0 (M+H⁺).

[B] 3-Bromo-5-((S)-2-chloromethyl-pyrrolidin-1-yl)-pyridine




- 5 To a solution of (S)-1-(5-bromopyridin-3-yl)pyrrolidin-2-yl)methanol (0.060 g, 0.233 mmol) in DCM (1 mL) cooled to 0 °C was added thionyl chloride (0.112 g, 0.932 mmol) dropwise. The reaction mixture was stirred at this temperature for 30 min and then allowed to warm up to room temperature and stirring was continued over night. The mixture was diluted with DCM, poured into aq. NaHCO₃ (10 mL) and the aqueous layer was extracted
- 10 with DCM (2 x 20 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (0.055 g, 86 %) as an orange oil. MS: 275.0, 277.0 (M+H⁺).

[C] 3-Bromo-5-((S)-2-[1,2,4]triazol-1-ylmethyl-pyrrolidin-1-yl)-pyridine




- 15 To a solution of NaH (0.011 g, 0.272 mmol) in DMF (1 mL) was added 1H-1,2,4-triazole (0.016 g, 0.236 mmol) and the reaction mixture was stirred at room temperature for 30 min. Then, 3-bromo-5-((S)-2-chloromethyl-pyrrolidin-1-yl)-pyridine (0.05 g, 0.181 mmol) in DMF (0.5 mL) was added to the reaction mixture which was heated to 80 °C over night. The mixture was diluted with DCM, poured into aq. NaHCO₃ (5 mL) and the aqueous
- 20 layer was extracted with DCM (2 x 10 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (0.04 g, 72 %) as a yellow oil. MS: 308.0, 310.0 (M+H⁺).

Intermediate A-6

3-Bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyridine


In analogy to the procedure described for the preparation of intermediate A-3 [B], (S)-2-methoxymethyl-pyrrolidine was reacted with 3,5-dibromopyridine in the presence of
5 Pd₂(dba)₃, *rac*-BINAP and sodium *tert*-butoxide to give the title compound as a yellow oil.
MS: 271.1, 273.1 (M+H⁺).

Intermediate A-7**[2-(5-Bromo-pyridin-3-yloxy)-ethyl]-carbamic acid *tert*-butyl ester**


10

To a solution of 5-bromopyridin-3-ol (0.5 g, 2.87 mmol) in THF (30 mL) cooled to 0 °C were added triphenylphosphine (0.829 g, 3.16 mmol) and *tert*-butyl 2-hydroxyethylcarbamate (0.51 g, 3.16 mmol) in THF (5 mL) followed by *di*-(4-chlorobenzyl)azodicarboxylate (1.16 g, 3.16 mmol), added portionwise, and the reaction
15 mixture was then stirred at room temperature over night. The mixture was diluted with EtOAc, poured into aq. NaHCO₃ (50 mL) and the aqueous layer was extracted with EtOAc (2 x 100 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated. The residue was taken up in diethyl ether (50 mL) and left to stand in the fridge for 2h. After this time, the solid precipitate was filtered off and the resulting filtrate
20 evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 30% EtOAc-heptane gradient to give the title compound (0.57 g, 63%) as a white solid. MS: 317.0 and 319.0 (M+H⁺).

Intermediate A-8

Benzyl-(5-bromo-pyridin-3-yl)-amine

In analogy to the procedure described for the preparation of intermediate A-3 [B], benzylamine was reacted with 3,5-dibromopyridine in the presence of Pd₂(dba)₃, *rac*-BINAP and sodium *tert*-butoxide to give the title compound as a yellow solid. MS: 262.9, 264.9 (M+H⁺).


Intermediate A-9**(S)-1-(5-Bromo-pyridin-3-yl)-5-hydroxymethyl-pyrrolidin-2-one**

10

In a sealed tube, 3,5-dibromopyridine (0.5 g, 2.11 mmol) was combined with (S)-5-(hydroxymethyl)pyrrolidin-2-one (0.243 g, 2.11 mmol), copper (I) iodide (0.040 g, 0.021 mmol), potassium carbonate (0.583 g, 4.22 mmol) and *N,N'*-dimethylethylenediamine (0.037g, 0.042 mmol) in 1,4-dioxane (20 mL). The reaction mixture was heated to 110 °C over night. The mixture was cooled to room temperature, filtered through Dicalite and washed with DCM. The residue was purified by silica gel flash chromatography eluting with a 0 to 10% MeOH-DCM gradient to give the title compound (0.140 g, 25 %) as a light yellow oil. MS: 271.1, 273.1 (M+H⁺).

Intermediate A-10**1-(5-Bromo-pyridin-3-yl)-pyrrolidin-2-one**

20




In analogy to the procedure described for the preparation of intermediate A-9, pyrrolidin-2-one has been coupled to 3,5-dibromopyridine to yield the title compound as a light yellow solid. MS: 241.0, 243.0 ($M+H^+$).

5

Intermediate A-11


Ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide




A flask was charged with 5-bromonicotinaldehyde (2.55 g, 13.7 mmol),
10 ethanesulfonamide (2.99 g, 27.4 mmol) and toluene (250 mL), then titanium isopropoxide (5.84 g, 20.6 mmol) was added dropwise. The reaction mixture was heated to 115 °C over night and then concentrated in vacuo. The residue was taken up in DCM (200 mL) and MeOH (200 mL) and NaBH₄ (1.04 g, 27.4 mmol) was added portionwise at 0 °C. The reaction mixture was stirred at 0 °C for 30 min and then poured into water (100 mL) and
15 the resulting suspension was filtered through a pad of Dicalite and washed with DCM (3 x 100 mL). The aqueous layer was separated and extracted with DCM (2 x 200 mL). Combined organics were dried over Na₂SO₄, filtered and preadsorbed on silica gel. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (3.01 g, 79%) as an orange solid. MS: 279.0,
20 281.0 ($M+H^+$).

Intermediate A-12

1-(5-Bromo-pyridin-3-ylmethyl)-pyrrolidin-2-one




[A] 3-Bromo-5-chloromethyl-pyridine




To a solution of (5-bromopyridin-3-yl)methanol (3 g, 16.0 mmol) in DCM (15 mL) cooled
5 to 0 °C was added thionylchloride (7.59 g, 63.8 mmol) dropwise and the reaction mixture
was stirred at room temperature over night. The mixture was poured onto ice/water (20
mL), basified with NaOH conc. (8 mL) and extracted with EtOAc (2 x 50 mL). Combined
organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was
10 purified by silica gel flash chromatography eluting with a 0 to 40% EtOAc-Heptane
gradient to give the title compound (3.08 g, 93 %) as a white solid. MS: 206.0, 207.9
(M+H⁺).

[B] 1-(5-Bromo-pyridin-3-ylmethyl)-pyrrolidin-2-one




To a suspension of sodium hydride (60% in mineral oil, 0.044 g, 1.09 mmol) in DMF (2
15 mL) was added pyrrolidin-2-one (0.081 g, 0.945 mmol) and the reaction mixture was
stirred at room temperature for 20 min. Then, 3-bromo-5-chloromethyl-pyridine (0.15 g,
0.727 mmol) was added and the resulting suspension was heated at 60 °C over night. The
mixture was quenched with water (2 mL) and extracted with EtOAc (2 x 10 mL).
Combined organics were dried over Na₂SO₄, filtered and concentrated in vacuo. The
20 residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-
DCM gradient to give the title compound (0.217 g, 87 %) as a white solid. MS: 251.1,
257.1 (M+H⁺).

Intermediate A-13

1-(5-Bromo-pyridin-3-ylmethyl)-piperidin-2-one


In analogy to the procedure described for the preparation of intermediates A-12 [B], piperidin-2-one was reacted with 3-bromo-5-chloromethyl-pyridine (intermediate A-12
5 [A]) in the presence of NaH to give the title compound as a colorless oil. MS: 269.2, 271.2 (M+H⁺).

Intermediate A-14**4-(5-Bromo-pyridin-3-ylmethyl)-thiomorpholine 1,1-dioxide**

10

In analogy to the procedure described for the preparation of intermediates A-12 [B], thiomorpholine 1,1-dioxide was reacted with 3-bromo-5-chloromethyl-pyridine (intermediate A-12 [A]) in the presence of NaH to give the title compound as a white solid. MS: 304.9, 307.0 (M+H⁺).

15


Intermediate A-15**3-Bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridine**

In analogy to the procedure described for the preparation of intermediates A-12 [B], (S)-2-(methoxymethyl)pyrrolidine was reacted with 3-bromo-5-chloromethyl-pyridine (intermediate A-12 [A]) in presence of NaH to give the title compound as a yellow oil. MS: 285.0, 286.9 ($M+H^+$).


5

Intermediate A-16

6-{5-[(S)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




10 [A] 3-Bromo-5-[(S)-2-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-ylmethyl]-pyridine



In analogy to the procedure described for the preparation of intermediates A-12 [B], (S)-2-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidine (intermediate A-3 [A]) was reacted
15 with 3-bromo-5-chloromethyl-pyridine (intermediate A-12 [A]) in presence of NaH to give the title compound as a yellow solid. MS: 385.2, 387.2 ($M+H^+$).


[B] 6-{5-[(S)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one



In analogy to the procedure described for the preparation of intermediate A-3 [C], 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) has been coupled to 3-bromo-5-[(*S*)-2-(*tert*-butyl-dimethyl-silyloxymethyl)-pyrrolidin-1-ylmethyl]-pyridine to give the title compound as an off white waxy solid. MS: 352.3 ($M+H^+$).

Intermediate A-17

(*S*)-1-(5-Bromo-pyridin-3-ylmethyl)-pyrrolidine-2-carboxylic acid methyl ester




10

In analogy to the procedure described for the preparation of intermediates A-12 [B], (*S*)-pyrrolidine-2-carboxylic acid methyl ester was reacted with 3-bromo-5-chloromethyl-pyridine (intermediate A-12 [A]) in presence of NaH to give the title compound as a light yellow oil. MS: 299.2, 301.1 ($M+H^+$).


15

Intermediate A-18

6-{5-[(*S*)-2-(*tert*-Butyl-dimethyl-silyloxymethyl)-5-oxo-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




[A] (S)-5-(*tert*-Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one




- In analogy to the procedure described for the preparation of intermediate A-3 [A], (S)-5-hydroxymethyl-pyrrolidin-2-one was reacted with TBDMS-Cl in the presence of TEA to give the title compound as a colorless liquid. MS: 230.3 ($M+H^+$).

[B] (S)-1-(5-Bromo-pyridin-3-ylmethyl)-5-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one



- In analogy to the procedure described for the preparation of intermediates A-12 [B], (S)-5-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one was reacted with 3-bromo-5-chloromethyl-pyridine (intermediate A-12 [A]) in presence of NaH to give the title compound as a colorless oil. MS: 399.2, 401.2 ($M+H^+$).


- [C] 6-{5-[(S)-2-(*tert*-Butyl-dimethyl-silanyloxymethyl)-5-oxo-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one



In analogy to the procedure described for the preparation of intermediate A-3 [C], 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) has been coupled to (S)-1-(5-bromo-pyridin-3-ylmethyl)-5-(*tert*-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one to give the title compound as a brown foam. MS: 480.3 ($M+H^+$).


Intermediate A-19

3-Bromo-5-[(S)-2-(2,2,2-trifluoro-ethoxymethyl)-pyrrolidin-1-ylmethyl]-pyridine



10


[A] (S)-1-Benzyl-2-(2,2,2-trifluoro-ethoxymethyl)-pyrrolidine



To a suspension of sodium hydride (60% in mineral oil, 0.314 g, 7.84 mmol) in THF (20 mL) was added (S)-1-(1-benzyl-pyrrolidin-2-yl)-methanol (1.0 g, 5.23 mmol) and the


reaction mixture was stirred at room temperature for 20 min. Then, 2,2,2-trifluoroethyl methanesulfonate (1.4 g, 7.84 mmol) was added and stirring was continued at room temperature over night. The reaction mixture was quenched with H₂O (10 mL) and extracted with EtOAc (2 x 50 mL). Combined organics were dried over Na₂SO₄, filtered and concentrated in vacuo. The residue was purified by silica gel flash chromatography eluting with a 0 to 5 % MeOH(1% NH₄OH)-DCM gradient to give the title compound (0.706 g, 37 %) as a light yellow liquid. MS: 274.3 (M+H⁺).

[B] (S)-2-(2,2,2-Trifluoro-ethoxymethyl)-pyrrolidine




To a solution of (S)-1-benzyl-2-(2,2,2-trifluoroethoxymethyl)-pyrrolidine (0.5 g, 1.83 mmol) in MeOH (20 mL) was added Hunig's base (1.18 g, 9.15 mmol). The flask was purged three times with Ar and 10 % Pd/C (0.487 g, 0.457 mmol) was added to the mixture. The flask was purged three times with hydrogen and the reaction mixture was stirred at room temperature for 1h. The mixture was filtered through a pad of Celite, washed with MeOH (20 mL) and the resulting solution was concentrated in vacuo to give the title compound (0.250 g, 75%) as a colorless semi-solid. MS: 184.1 (M+H⁺).

[C] 3-Bromo-5-[(S)-2-(2,2,2-trifluoro-ethoxymethyl)-pyrrolidin-1-ylmethyl]-pyridine




In analogy to the procedure described for the preparation of intermediates A-12 [B], (S)-2-(2,2,2-trifluoro-ethoxymethyl)-pyrrolidine was reacted with 3-bromo-5-chloromethylpyridine (intermediate A-12 [A]) in presence of NaH to give the title compound as a yellow oil. MS: 353.1, 355.1 (M+H⁺).

Intermediate A-20**2-(6-Chloro-pyrazin-2-ylamino)-ethanol**

In a sealed tube, 2,6-dichloropyrazine (1 g, 6.71 mmol) was mixed with 2-aminoethanol
5 (0.492 g, 8.05 mmol), Hunig's base (1.13 g, 8.73 mmol) in BuOH (7 mL) and the reaction mixture was heated to 80 °C over night. The mixture was concentrated in vacuo and the residue partitioned between NaHCO₃ (20 mL) and EtOAc (100 mL). The organic phase was separated, washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (0.583 g, 50%) as a light yellow foam. MS: 174.1 (M+H⁺).


10

Intermediate A-21**Benzyl-(6-chloro-pyrazin-2-yl)-amine**


In analogy to the procedure described for the preparation of intermediates A-20, benzyl-
15 amine was reacted with 2,6-dichloropyrazine in presence of Hunig's base to give the title compound as a light yellow foam. MS: 220.1 (M+H⁺).

Intermediate A-22

**7-Fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-
20 quinolin-2-one**




[A] 6-Bromo-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one



In analogy to the procedure described for the preparation of intermediates A-1 [A], 6-bromo-7-fluoro-3,4-dihydro-1H-quinolin-2-one was reacted with methyl iodide in presence of potassium *tert*-butoxide to give the title compound as a white foam. MS: 258.0, 259.9 ($M+H^+$).

[B] 7-Fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one




10

In analogy to the procedure described for the preparation of intermediates A-1 [B], 6-bromo-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one was reacted with 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) in presence of potassium acetate and $PdCl_2(DPPF)-CH_2Cl_2$ to give the title compound as a white solid. MS: 306.1 ($M+H^+$).

15

Intermediate A-23


1,4,4-Trimethyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one



- In analogy to the procedure described for the preparation of intermediates A-1 [B], 6-bromo-1,4,4-trimethyl-3,4-dihydro-1*H*-quinolin-2-one was reacted with 4,4,4',4',5,5,5',5'-octamethyl-2,2'-*bi*(1,3,2-dioxaborolane) in presence of potassium acetate and
- 5 PdCl₂(DPPF)-CH₂Cl₂ to give the title compound as an off-white solid. MS: 316.1 (M+H⁺).

Intermediate A-24

- 6-(6-Chloro-pyrazin-2-yl)-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid *tert*-butyl ester
- 10 ester




- In analogy to the procedure described for the preparation of intermediate A-20, 2,6-diaza-spiro[3.3]heptane-2-carboxylic acid *tert*-butyl ester was reacted with 2,6-dichloropyrazine in presence of Hunig's base to give the title compound as a white solid.

15

Intermediate A-25


- 6-(5-Bromo-pyridin-3-yl)-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid *tert*-butyl ester



- In analogy to the procedure described for the preparation of intermediate A-3 [B], 2,6-diaza-spiro[3.3]heptane-2-carboxylic acid *tert*-butyl ester was reacted with 3,5-dibromopyridine in the presence of $\text{Pd}_2(\text{dba})_3$, *rac*-BINAP and sodium *tert*-butoxide to
- 5 give the title compound as a white solid.

Intermediate A-26


(S)-2-(5-Bromo-pyridin-3-yloxymethyl)-azetidine-1-carboxylic acid *tert*-butyl ester



- 10 In analogy to the procedure described for the preparation of intermediate A-7, 5-bromopyridin-3-ol was reacted with (S)-2-hydroxymethyl-azetidine-1-carboxylic acid *tert*-butyl ester in presence of *di*-(4-chlorobenzyl)azodicarboxylate and triphenylphosphine to give the title compound as an amorphous colorless solid. MS: 343.1 and 345.1 ($\text{M}+\text{H}^+$).

Intermediate A-27

(R)-3-(5-Bromo-pyridin-3-yloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester




In analogy to the procedure described for the preparation of intermediate A-7, 5-bromopyridin-3-ol was reacted with (S)-3-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester in presence of *di*-(4-chlorobenzyl)azodicarboxylate and triphenylphosphine to give the title compound as a white solid. MS: 343.1 and 345.1 ($M+H^+$).

5

Intermediate A-28


3-(5-Bromopyridin-3-yloxy)-azetidine-1-carboxylic acid *tert*-butyl ester



To a solution of 5-bromopyridin-3-ol (0.285 g, 1.64 mmol) in DMF (2.5 mL) was added
10 K_2CO_3 (0.453 g, 3.28 mmol), followed by 3-bromo-azetidine-1-carboxylic acid *tert*-butyl ester (0.425 g, 1.8 mmol) in DMF (0.5 mL) and the reaction mixture was heated to 60 °C and stirred over night. The reaction mixture was diluted with EtOAc, poured into sat. $NaHCO_3$ solution (10 mL) and the aqueous layer was extracted EtOAc (2 x 20 mL). Combined organics were washed with brine, dried over Na_2SO_4 , filtered and evaporated to
15 dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 50% EtOAc-heptane gradient to give the title compound (0.539 g, 100%) as a colorless crystalline solid. MS: 329.1 ($M+H^+$).


Intermediate A-29

(S)-3-(5-Bromo-pyridin-3-yloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester




20

In analogy to the procedure described for the preparation of intermediate A-7, 5-bromopyridin-3-ol was reacted with (R)-3-hydroxy-pyrrolidine-1-carboxylic acid *tert*-butyl ester in presence of *di*-(4-chlorobenzyl)azodicarboxylate and triphenylphosphine to give the title compound as a colorless oil. MS: 343.1 and 345.1 ($M+H^+$).

Intermediate A-30**(S)- 3-(5-Bromopyridin-3-yloxy)-piperidine-1-carboxylic acid *tert*-butyl ester**


- 5 In analogy to the procedure described for the preparation of intermediate A-7, 5-bromopyridin-3-ol was reacted with (R)-3-hydroxy-piperidine-1-carboxylic acid *tert*-butyl ester in presence of *di*-(4-chlorobenzyl)azodicarboxylate and triphenylphosphine to give the title compound as a light yellow oil. MS: 357.1 and 359.1 (M+H⁺).

10 **Intermediate A-31**

4-(5-Bromopyridin-3-yloxy)-piperidine-1-carboxylic acid *tert*-butyl ester

- In analogy to the procedure described for the preparation of intermediate A-7, 5-bromopyridin-3-ol was reacted with 4-hydroxy-piperidine-1-carboxylic acid *tert*-butyl ester in presence of *di*-(4-chlorobenzyl)azodicarboxylate and triphenylphosphine to give the title compound as a light yellow amorphous solid. MS: 357.0 and 359.0 (M+H⁺).
- 15


Intermediate A-32**(S)-2-((5-Bromopyridin-3-yloxy)methyl)pyrrolidine-1-carboxylic acid *tert*-butyl ester**




In analogy to the procedure described for the preparation of intermediate A-7, 5-bromopyridin-3-ol was reacted with (S)-2-(hydroxymethyl)pyrrolidine-1-carboxylic acid *tert*-butyl ester in presence of *di*-(4-chlorobenzyl)azodicarboxylate and triphenylphosphine
 5 to give the title compound as a colorless oil. MS: 357.0 and 359.0 ($M+H^+$).

Intermediate A-33


(R)-N-(3-(5-Bromopyridin-3-yl)oxetan-3-yl)-2-methylpropane-2-sulfinamide




10 [A] (R)-2-Methyl-N-(oxetan-3-ylidene)-propane-2-sulfinamide



To a solution of $Ti(OEt)_4$ (2.02 g, 7.53 mmol) in THF (20 mL) was added (R)-2-methylpropane-2-sulfinamide (0.854 g, 7.05 mmol) followed by oxetan-3-one (0.493 g, 6.84 mmol) in THF (5 mL) and the reaction mixture was heated to 50 °C over night. After
 15 cooling, the reaction mixture was diluted with EtOAc (50 mL) and a saturated solution of brine (20 mL) was slowly added under stirring. After 15 min, the mixture was filtered through a plug of Dicalite and washed with EtOAc. The organic phase was separated, dried over Na_2SO_4 and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 40 % EtOAc-heptane gradient to give the title
 20 compound (0.242 g, 20 %) as a light yellow oil. MS: 176.2 ($M+H^+$).

[B] (R)-N-(3-(5-Bromopyridin-3-yl)oxetan-3-yl)-2-methylpropane-2-sulfinamide

To a solution of 3-bromo-5-iodopyridine (0.14 g, 0.493 mmol) in THF (2 mL) cooled at -78 °C was added nBuLi (462 μ l, 0.74 mmol) dropwise over 5 min. The reaction mixture was stirred at -78 °C for 15 min, then (R)-2-methyl-N-(oxetan-3-ylidene)-propane-2-sulfinamide (0.1 g, 0.567 mmol) in THF (1 mL) was added dropwise. The reaction mixture was stirred at -78 °C for 10 min then allowed to warm up to RT. The mixture was quenched with water (2 mL) and extracted with EtOAc (2 x 10 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 100 % EtOAc-heptane gradient to give the title compound (0.096 g, 58 %) as a light yellow oil. MS: 333.0 and 335.1 (M+H⁺).

Intermediate A-34**6-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroquinolin-2(1H)-one**


In analogy to the procedure described for the preparation of intermediates A-1 [B],

6-bromo-3,4-dihydroquinolin-2(1H)-one was reacted with 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) in presence of potassium acetate and PdCl₂(DPPF)-CH₂Cl₂ to give the title compound as a colorless solid. MS: 274.4 (M+H⁺).

Intermediate A-35**(R)-2-Methyl-propane-2-sulfinic acid [(S and R)-1-(5-bromo-pyridin-3-yl)-ethyl]-**


amide

[A] (R, E)-*N*-(1-(5-Bromopyridin-3-yl)ethylidene)-2-methylpropane-2-sulfinamide




- To a solution of 1-(5-bromopyridin-3-yl)ethanone (1.01 g, 5.04 mmol) in DCM (20 mL) was added (R)-2-methylpropane-2-sulfinamide (0.555 g, 4.58 mmol) followed by $\text{Ti}(\text{OEt})_4$ (1.23 g, 4.58 mmol) dropwise and the reaction mixture was heated to 40 °C over night. After cooling, the solvent was removed under vacuum and the residue taken up in EtOAc (80 mL). This solution was vigorously stirred while a saturated solution of brine (20 mL) was slowly added. After 15 min, the mixture was filtered through a plug of Dicalite and washed with EtOAc. The organic phase was separated, dried over Na_2SO_4 and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 50 % EtOAc-heptane gradient to give the title compound (0.964 g, 69 %) as a yellow oil. MS: 303.1 and 305.0 ($\text{M}+\text{H}^+$).

- [B] (R)-2-Methyl-propane-2-sulfinic acid [(S and R)-1-(5-bromo-pyridin-3-yl)-ethyl]-amide



(S and R)



(R and S)


- To a solution of (R, E)-*N*-(1-(5-bromopyridin-3-yl)ethylidene)-2-methylpropane-2-sulfinamide (0.3 g, 0.989 mmol) in MeOH (20 mL) cooled at 0 °C with an ice bath was added NaBH_4 (0.075 g, 1.98 mmol) portionwise and the reaction mixture was stirred at this temperature for 30 min and then at room temperature for another 30 min. The mixture was quenched with sat. ammonium chloride solution (10 mL), the organic solvent was

evaporated and the resulting aqueous solution was extracted with EtOAc (2 x 25 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (0.293 g, 97 %) as a colorless solid as a 2:1 mixture of both diastereoisomers. MS: 305.0 and 307.1 (M+H⁺).

5

Intermediate A-36


6-(6-(2,6-Diazaspiro[3.3]heptan-2-yl)pyrazin-2-yl)-1-methyl-3,4-dihydroquinolin-2(1H)-one 2,2,2-trifluoroacetate



- 10 To a solution of 6-[6-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyrazin-2-yl]-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid *tert*-butyl ester (example 28, 0.231 g, 0.53 mmol) in DCM (3 mL) was added TFA (0.605 g, 5.3 mmol) and the reaction mixture was stirred at room temperature over night. The mixture was evaporated several times with toluene and then the residue was purified by reverse phase HPLC on a Gemini-NX
- 15 column, eluting with a 20 to 98% MeOH-H₂O (0.05 % TEA) gradient to give the title compound (0.203 g, 99 %) as a yellow solid (TFA salt). MS: 336.3 (M+H⁺).

Intermediate A-37

(rac)-N-(1-(5-Bromopyridin-3-yl)ethyl)ethanesulfonamide




20

In analogy to the procedure described for the preparation of intermediate A-11, 1-(5-bromo-pyridin-3-yl)-ethanone was reacted with ethanesulfonamide in presence of titanium

tetra-isopropoxide and then NaBH₄ to give the title compound as a light yellow oil. MS: 293.1 and 295.3 (M+H⁺).

Intermediate A-38


5 (rac)-N-(1-(5-Bromopyridin-3-yl)propyl)ethanesulfonamide



- A flask was charged with 5-bromonicotinaldehyde (0.5 g, 2.69 mmol), ethanesulfonamide (0.367 g, 3.36 mmol) and toluene (25 mL), then titanium *tetra*-isopropoxide (1.53 g, 5.38 mmol) was added dropwise. The reaction mixture was heated to 110 °C over night and
- 10 then concentrated in vacuo. The residue was dissolved in THF (25 mL) and cooled down to -40 °C. A 3 M solution of ethylmagnesium bromide in ether (2.24 mL, 6.72 mmol) was added dropwise at this temperature and the resulting mixture was slowly warmed up to -20 °C and stirred for 2.5 h. The mixture was poured into a saturated NH₄Cl solution (15 mL), the resulting suspension was filtered and the filtrate was extracted with DCM (2 x 50 mL).
- 15 Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.5 g, 56%) as light yellow oil. MS: 307.1 and 308.9.


20 Intermediate A-39

(rac)-N-((5-Bromopyridin-3-yl)(cyclopropyl)methyl)ethanesulfonamide




In analogy to the procedure described for the preparation of intermediate A-38, 5-bromonicotinaldehyde was reacted with ethanesulfonamide in presence of titanium *tetra*-isopropoxide and then cyclopropylmagnesium bromide (0.5 M in THF) to give the title compound as a yellow oil. MS: 319.0 and 320.9 (M+H⁺).

5


Intermediate A-40**(rac)-N-(1-(5-Bromopyridin-3-yl)-2-methylpropyl)ethanesulfonamide**

10 In analogy to the procedure described for the preparation of intermediate A-38, 5-bromonicotinaldehyde was reacted with ethanesulfonamide in presence of titanium *tetra*-isopropoxide and then isopropylmagnesium chloride (2 M in THF) to give the title compound as a red oil. MS: 321.0 and 323.0 (M+H⁺).

Intermediate A-41**15 N-((5-Bromopyridin-3-yl)methyl)-N-methylethanesulfonamide**

To a solution of ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11, 0.21 g, 0.752 mmol) in DMF (6 mL) cooled at 0 °C with an ice bath was added 60 % NaH in mineral oil (0.039 g, 0.98 mmol) and the mixture was stirred for 15 min. Then, MeI (0.136 g, 0.96 mmol) was added and the reaction was stirred at 0 °C for another 15 min before being quenched with aq. ammonia (2 mL). The mixture was further diluted with brine and then extracted with EtOAc (2 x 15 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (0.209 g, 85 %) as a yellow oil which was used with no further purification. MS: 293.1 and 295.1 (M+H⁺).


20

Intermediate A-42***N-((5-Bromopyridin-3-yl)methyl)-N-ethylethanesulfonamide***

- 5 To a solution of ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11, 0.051 g, 0.182 mmol) in DMF (3 mL) cooled at 0 °C with an ice bath was added 60 % NaH in mineral oil (0.011 g, 0.274 mmol) and the mixture was stirred at room temperature for 20 min. Then, iodoethane (0.045 g, 0.292 mmol) was added and the reaction mixture was stirred for 3 h before being quenched with water (5 mL). The aqueous solution was
- 10 extracted with EtOAc (2 x 15 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (0.051 g, 82 %) as orange oil which was used with no further purification. MS: 307.1 and 309.2 (M+H⁺).


Intermediate A-43

- 15 ***N-((5-Bromopyridin-3-yl)methyl)-N-isopropylethanesulfonamide***



- In analogy to the procedure described for the preparation of intermediate A-42, ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) was reacted with 2-iodopropane in presence of NaH (60 % in mineral oil) to give the title compound as
- 20 white solid after purification by reverse phase HPLC on a Gemini-NX column. MS: 323.1 (M+H⁺).


Intermediate A-44

N-((5-Bromopyridin-3-yl)methyl)-N-2-ethoxy-ethylethanesulfonamide

- In analogy to the procedure described for the preparation of intermediate A-42, ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) was reacted
5 with 2-bromoethyl ethyl ether in presence of NaH (60 % in mineral oil) to give the title compound as yellow oil after purification by reverse phase HPLC on a Gemini-NX column. MS: 351.0 and 353.1 ($M+H^+$).

Intermediate A-45


- 10 ***(rac)-N-(1-(5-Bromopyridin-3-yl)ethyl)-N-methylethanesulfonamide***



- In analogy to the procedure described for the preparation of intermediate A-41, *N*-(1-(5-bromopyridin-3-yl)ethyl)ethanesulfonamide (intermediate A-37) was reacted with MeI in
15 presence of NaH (60 % in mineral oil) to give the title compound as a light yellow solid. MS: 307.1 and 309.2 ($M+H^+$).

Intermediate A-46


(rac)-N-(1-(5-Bromopyridin-3-yl)ethyl)-N-ethylethanesulfonamide



In analogy to the procedure described for the preparation of intermediate A-42, *N*-(1-(5-bromopyridin-3-yl)ethyl)ethanesulfonamide (intermediate A-37) was reacted with iodoethane in presence of NaH (60 % in mineral oil) to give the title compound as a
5 yellow oil. MS: 321.0 and 323.1 ($M+H^+$).

Intermediate A-47


1-Methyl-1*H*-pyrazole-4-carboxylic acid (5-bromo-pyridin-3-ylmethyl)-amide



10 In analogy to the procedure described for the preparation of example 75, coupling of (5-bromopyridin-3-yl)methanamine with 1-methyl-1*H*-pyrazole-4-carboxylic gave the title compound as a colorless solid. MS: 297.1 and 295.0 ($M+H^+$).

Intermediate A-48

15 2-(5-Bromo-pyridin-3-ylmethyl)-isothiazolidine 1,1-dioxide




In analogy to the procedure described for the preparation of intermediate A-12 [B], isothiazolidine 1,1-dioxide was reacted with 3-bromo-5-chloromethyl-pyridine

(intermediate A-12 [A]) in the presence of NaH to give the title compound as a light yellow oil. MS: 292.8 and 290.9 ($M+H^+$).

Intermediate A-49


5 *N*-(2-(5-Bromopyridin-3-yl)propan-2-yl)ethanesulfonamide



A flask was charged with 1-(5-bromopyridin-3-yl)ethanone (0.5 g, 2.5 mmol), ethanesulfonamide (0.341 g, 3.12 mmol) and toluene (25 mL), then titanium *tetra*-isopropoxide (1.42 g, 5 mmol) was added dropwise. The reaction mixture was heated to
10 110 °C over night and then concentrated in vacuo to give the imine intermediate (0.73 g) as a light brown oil. A part of this the imine intermediate (0.4 g) was dissolved in THF (10 mL) and the mixture was cooled down to -78 °C. A 3 M solution of methylmagnesium bromide in ether (1.14 mL, 3.43 mmol) was added dropwise at this temperature and the resulting mixture was stirred at -70 °C for 3 h. It was then allowed to warm up to 0 °C
15 before being quenched with a saturated NH_4Cl solution (10 mL). The resulting suspension was filtered and the filtrate was extracted with DCM (2 x 30 mL). Combined organics were dried over Na_2SO_4 , filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.1 g, 17 %) as a yellow oil which was used with no further purification.
20 MS: 307.1 and 309.1 ($M+H^+$).

Intermediate A-50

***N*-((5-Bromopyridin-3-yl)methyl)-*N*,3,5-trimethylisoxazole-4-carboxamide**




25 In analogy to the procedure described for the preparation of intermediate A-41, 3,5-

dimethyl-isoxazole-4-carboxylic acid (5-bromo-pyridin-3-ylmethyl)-methyl-amide (prepared as described for the preparation of intermediate A-47) was reacted with MeI in presence of NaH (60 % in mineral oil) to give the title compound as a yellow solid. MS: 326.3 and 324.2 ($M+H^+$).


5

Intermediate A-51

3-Bromo-5-[2-(1,1-dioxo-1 λ^6 -isothiazolidin-2-yl)-ethoxy]-pyridine



[A] 2-(2-Chloro-ethyl)-isothiazolidine-1,1-dioxide




10

To a solution of 2-(1,1-dioxo-1 λ^6 -isothiazolidin-2-yl)-ethanol (0.14 g, 0.847 mmol) in DCM (4 mL) was added SOCl_2 (0.151 g, 1.27 mmol) and the reaction mixture was heated to 40 °C for 2 h. The mixture was cooled to room temperature, diluted with DCM, poured into a sat. NaHCO_3 solution (10 mL) and the aqueous layer was extracted with DCM (2 x 20 mL). Combined organics were dried over Na_2SO_4 , filtered and evaporated to dryness to give the title compound (0.046 g, 30 %) as an orange oil which was used with no further purification. MS: 184.1 ($M+H^+$).

15

[B] 3-Bromo-5-[2-(1,1-dioxo-1 λ^6 -isothiazolidin-2-yl)-ethoxy]-pyridine




To a solution of 2-(2-chloro-ethyl)-isothiazolidine-1,1-dioxide (0.046 g, 0.253 mmol) and 5-bromopyridin-3-ol (0.04 g, 0.230 mmol) in DMF (1 mL) was added K_2CO_3 (0.064 g, 0.46 mmol), followed by KI (0.008 g, 0.046 mmol) and the reaction mixture was heated to

20

60 °C and stirred over night. The mixture was diluted with EtOAc, poured into a sat. NaHCO₃ solution (5 mL) and the aqueous layer was extracted with EtOAc (2 x 10 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.015 g, 20 %) as a yellow oil which was used with no further purification. MS: 321.3 and 323.2 (M+H⁺).

Intermediate A-52

***N*-((5-Bromo-4-methylpyridin-3-yl)methyl)ethanesulfonamide**




In analogy to the procedure described for the preparation of intermediate A-11, 5-bromo-4-methyl-pyridine-3-carbaldehyde has been reacted first with ethanesulfonamide, followed by reduction of the thus formed imine with NaBH₄ in MeOH to give the title compound as an off-white solid. MS: 293.1 and 295.3 (M+H⁺).

15

Intermediate A-53


***N*-((5-Bromo-4-chloropyridin-3-yl)methyl)ethanesulfonamide**



20 In analogy to the procedure described for the preparation of intermediate A-11, 5-bromo-4-chloro-pyridine-3-carbaldehyde has been reacted first with ethanesulfonamide, followed by reduction of the thus formed imine with NaBH₄ in MeOH to give the title compound as a light orange solid. MS: 313.2 and 315.1 (M+H⁺).

Intermediate A-54


***tert*-Butyl (*trans*)-4-(5-bromopyridin-3-yloxy)cyclohexylcarbamate**



- In analogy to the procedure described for the preparation of intermediate A-7, 5-bromopyridin-3-ol was reacted with (*cis*)-(4-hydroxy-cyclohexyl)-carbamic acid *tert*-butyl ester in presence of *di*-(4-chlorobenzyl)azodicarboxylate and triphenylphosphine to give the title compound as a colorless solid. MS: 371.3 and 373.3 ($M+H^+$).

Intermediate A-55


- 10 **(*R*)-2-((5-Bromopyridin-3-yloxy)methyl)pyrrolidine-1-carboxylic acid *tert*-butyl ester**



- In analogy to the procedure described for the preparation of intermediate A-7, 5-bromopyridin-3-ol was reacted with (*R*)-2-(hydroxymethyl)pyrrolidine-1-carboxylic acid *tert*-butyl ester in presence of *di*-(4-chlorobenzyl)azodicarboxylate and triphenylphosphine to give the title compound as an orange oil. MS: 357.3 and 359.3 ($M+H^+$).

Intermediate A-56


***N*-((5-Bromopyridin-3-yl)methyl)-3-chloropyridin-2(1H)-one**



In analogy to the procedure described for the preparation of example 75, coupling of (5-bromopyridin-3-yl)methanamine with 3-chloro-pyridine-2-carboxylic acid gave the title compound as off-white solid. MS: 326.2 and 328.2 ($M+H^+$).

5 **Intermediate A-57**


***N*-((5-Bromopyridin-3-yl)methyl)-3-chloro-*N*-methylpicolinamide**



In analogy to the procedure described for the preparation of intermediate A-41, *N*-((5-bromopyridin-3-yl)methyl)-3-chloropicolinamide (intermediate A-56) was reacted with
10 MeI in presence of NaH (60 % in mineral oil) to give the title compound as a dark brown oil. MS: 340.2 and 342.2 ($M+H^+$).

Intermediate A-58


(*R*)-*N*-((5-Bromo-4-chloropyridin-3-yl)methyl)-2-methylpropane-2-sulfinamide




15

(*R*)-2-Methyl-propane-2-sulfinic acid 1-(5-bromo-4-chloro-pyridin-3-yl)-meth-(*E*)-ylideneamide, prepared from (*R*)-2-methylpropane-2-sulfinamide and 5-bromo-4-chloro-pyridine-3-carbaldehyde in analogy to procedure described for the preparation of intermediate A-35 [A], has been reduced with NaBH₄ in MeOH in analogy to the
20 procedure described for the preparation of intermediate A-35 [B] to give the title compound as a yellow oil. MS: 325.2 ($M+H^+$).

Intermediate A-59

(R)-2-Methyl-propane-2-sulfinic acid (5-bromo-4-methyl-pyridin-3-ylmethyl)-amide


- (R)-2-Methyl-propane-2-sulfinic acid 1-(5-bromo-4-methyl-pyridin-3-yl)-meth-(E)-ylideneamide, prepared from (R)-2-methylpropane-2-sulfinamide and 5-bromo-4-methyl-pyridine-3-carbaldehyde in analogy to procedure described for the preparation of intermediate A-35 [A], has been reduced with NaBH₄ in MeOH in analogy to the procedure described for the preparation of intermediate A-35 [B] to give the title compound as colorless solid. MS: 305.3 and 307.4 (M+H⁺).

10 **Intermediate A-60****(R)-2-Methyl-propane-2-sulfinic acid [(R or S)-1-(5-bromo-4-methyl-pyridin-3-yl)-ethyl]-amide**

- (R)-2-Methyl-propane-2-sulfinic acid 1-(5-bromo-4-methyl-pyridin-3-yl)-meth-(E)-ylideneamide, prepared from (R)-2-methylpropane-2-sulfinamide and 5-bromo-4-methyl-pyridine-3-carbaldehyde in analogy to procedure described for the preparation of intermediate A-35 [A], has been reacted with methylmagnesium bromide in analogy to the procedure described for the preparation of intermediate A-49 from the corresponding imine intermediate to give the title compound as light yellow amorphous solid. MS: 319.2 and 321.3 (M+H⁺).

Intermediate A-61

(R)-2-Methyl-propane-2-sulfinic acid [(R or S)-1-(5-bromo-4-chloro-pyridin-3-yl)-ethyl]-amide




- (R)-2-Methyl-propane-2-sulfinic acid 1-(5-bromo-4-chloro-pyridin-3-yl)-meth-(E)-ylideneamide, prepared from (R)-2-methylpropane-2-sulfinamide and 5-bromo-4-chloro-pyridine-3-carbaldehyde in analogy to procedure described for the preparation of intermediate A-35 [A], has been reacted with methylmagnesium bromide in analogy to the procedure described for the preparation of intermediate A-49 from the corresponding imine intermediate to give the title compound as orange oil. MS: 341.2 (M+H⁺).


10

Example 1

Ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-amide




- 15 To a solution of 6-(5-amino-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-2, 0.02 g, 0.079 mmol) in DCM (0.5 mL) was added triethylamine (0.024 g, 0.237 mmol) followed by ethanesulfonyl chloride (0.01 g, 0.079 mmol) and the reaction mixture was stirred at room temperature for 1h. The mixture was diluted with DCM, poured into water (5 mL) and the aqueous layer was extracted with DCM (2 x 20 mL).
- 20 Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 10 % MeOH-DCM gradient to give the title compound (0.007 g, 23 %) as a brown solid. MS: 346.0 (M+H⁺).

Example 2**Acetic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylcarbamoyl]-methyl ester**

5

To a solution of 6-(5-amino-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (intermediate A-2, 0.045 g, 0.155 mmol) in DCM (1.5 mL) was added TEA (0.039 g, 0.388 mmol) followed by 2-chloro-2-oxoethyl acetate (0.017 g, 0.124 mmol) in DCM (1 mL) and the reaction mixture was stirred at room temperature for 2h. The mixture
10 was diluted with DCM, poured into aq. NaHCO₃ (10 mL) and the aqueous layer was extracted with DCM (2 x 25 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5 % MeOH-DCM gradient to give the title compound (0.025 g, 46 %) as a light brown solid. MS: 354.2 (M+H⁺).

15


Example 3**2-Hydroxy-*N*-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-acetamide**

20 To a solution of acetic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylcarbamoyl]-methyl ester (example 2, 0.025 g, 0.071 mmol) in MeOH (1 mL) was


added a 1M aqueous solution of Na_2CO_3 (0.354 mL) and the reaction mixture was stirred at room temperature for 1h. The mixture was evaporated to dryness, the residue dissolved in EtOAc (20 mL) and washed with aq. NaHCO_3 (5 mL). The organic layer was separated, dried over Na_2SO_4 , filtered and evaporated to dryness to give the title compound (0.02 g, 91%) as light brown solid. MS: 312.1 ($\text{M}+\text{H}^+$).

Example 4

6-[5-(2-Hydroxy-ethylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one




10 [A] 6-{5-[2-(*tert*-Butyl-dimethyl-silanyloxy)-ethylamino]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one



To a solution of 6-(5-amino-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (intermediate A-2, 0.062 g, 0.214 mmol) in MeOH (1.5 mL) was added AcOH (0.154 g, 2.57 mmol), followed by 2-(*tert*-butyldimethylsilyloxy)acetaldehyde (0.039 g, 0.225 mmol) and the reaction mixture was stirred at room temperature for 1h. Then, NaBH_3CN (0.027 g, 0.428 mmol) in THF (0.6 mL) was added to the reaction mixture and stirring at room temperature was continued for 2h. The mixture was diluted with EtOAc, poured into aq. NaHCO_3 (10 mL) and the aqueous layer was extracted with EtOAc (2 x 25 mL). Combined organics were washed with brine, dried over Na_2SO_4 , filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5 % MeOH-DCM gradient to give the title compound (0.06 g, 68 %) as a yellow solid. MS: 412.3 ($\text{M}+\text{H}^+$).


[B] 6-[5-(2-Hydroxy-ethylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one






To solution of 6-{5-[2-(*tert*-butyl-dimethyl-silanyloxy)-ethylamino]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (0.06 g, 0.146 mmol) in MeOH (2.5 mL) was
 5 added 4M HCl in dioxane (0.255 ml, 1.02 mmol) and the reaction mixture was stirred at room temperature for 2h. The mixture was concentrated in vacuo, the residue dissolved in EtOAc (20 mL) and washed with aq. NaHCO₃ (5 mL). The organic layer was washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 2 % MeOH(1% NH₄OH)-DCM
 10 gradient to give the title compound (0.01 g, 22 %) as an off white solid. MS: 298.3 (M+H⁺).

The following compounds listed in Table 1 were prepared in analogy to the procedure described for the preparation of example 4 [B] using appropriate starting materials


15 **Table 1**

Ex	Name and Structure	Starting Materials	Aspect	MS (M+H ⁺)
5	6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one 	6-{5-[(S)-2-(<i>tert</i> -Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (intermediate A-3)	Light yellow solid	338.2


Ex	Name and Structure	Starting Materials	Aspect	MS (M+H ⁺)
6	<p>6-[5-((R)-2-Hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	6-{5-[(R)-2-(<i>tert</i> -Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-yl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (intermediate A-4)	Light yellow solid	338.2
7	<p>6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	6-{5-[(S)-2-(<i>tert</i> -Butyl-dimethyl-silanyloxymethyl)-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (intermediate A-16)	Off white waxy solid	352.3
8	<p>6-[5-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	6-{5-[(S)-2-(<i>tert</i> -Butyl-dimethyl-silanyloxymethyl)-5-oxo-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (intermediate A-18)	White Foam	366.2

Example 9

6-[5-((S)-2-Ethylaminomethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one




5 [A] 6-[5-((S)-2-Chloromethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one



In analogy to the procedure described for the preparation of intermediate A-5 [B], 6-[5-((S)-2-hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 5) has been reacted with thionyl chloride to give the title compound as an orange oil. MS: 356.1 (M+H⁺).

10

[B] 6-[5-((S)-2-Ethylaminomethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one




In a sealed tube, a solution of 6-[5-((S)-2-chloromethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one (0.066 g, 0.185 mmol) in CH₃CN (2.5 mL) was mixed with K₂CO₃ (0.064 g, 0.464 mmol), TEA (0.188 g, 1.85 mmol), sodium iodide (0.028 g, 0.185 mmol) and ethanamine hydrochloride (0.151 g, 1.85 mmol). The reaction mixture was then heated to 80 °C over night. The mixture was diluted with EtOAc, poured

15

into aq. NaHCO_3 (10 mL) and the aqueous layer was extracted with EtOAc (2 x 20 mL). Combined organics were washed with brine, dried over Na_2SO_4 , filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography, eluting with a 0 to 15% MeOH(1% NH_4OH)-DCM gradient to give the title compound (0.02 g, 30 %) as an orange gum. MS: 365.2 ($\text{M}+\text{H}^+$).

Example 10

6-[5-((S)-2-Methoxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one



10




In analogy to the procedure described for the preparation of intermediate A-3 [C], 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1) has been coupled to 3-bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-yl)-pyridine (intermediate A-6) to give the title compound as a yellow oil. MS: 352.3 ($\text{M}+\text{H}^+$).




15




The following compounds listed in Table 2 were prepared in analogy to the procedure described for the preparation of intermediate A-3 [C], using the appropriate starting materials.



20



Table 2




Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
11	<p>1-Methyl-6-[5-((S)-2-[1,2,4]triazol-1-ylmethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one</p> 	<p>3-Bromo-5-((S)-2-[1,2,4]triazol-1-ylmethyl-pyrrolidin-1-yl)-pyridine (intermediate A-5) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	Light brown foam	389.1
12	<p>6-(5-Benzylamino-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one</p> 	<p>Benzyl-(5-bromopyridin-3-yl)-amine (intermediate A-8) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	White solid	344.1
13	<p>6-[5-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one</p> 	<p>(S)-1-(5-Bromopyridin-3-yl)-5-hydroxymethyl-pyrrolidin-2-one (intermediate A-9) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	White yellow foam	352.3



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
14	<p>1-Methyl-6-[5-(2-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one</p> 	1-(5-Bromo-pyridin-3-yl)-pyrrolidin-2-one (intermediate A-10) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)	White solid	322.1
15	<p>Ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	Ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)	White foam	360.1
16	<p>Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	Ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-22)	White solid	378.2


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
17	<p>Ethanesulfonic acid [5-(1,4,4-trimethyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p>Ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) and 1,4,4-trimethyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-23)</p>	Off white solid	388.1
18	<p>1-Methyl-6-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one</p> 	<p>1-(5-Bromo-pyridin-3-ylmethyl)-pyrrolidin-2-one (intermediate A-12) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	Brown foam	336.4
19	<p>1-Methyl-6-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one</p> 	<p>1-(5-Bromo-pyridin-3-ylmethyl)-piperidin-2-one (intermediate A-13) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	White solid	350.3

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
20	<p>6-[5-(1,1-Dioxo-1λ⁶-thiomorpholin-4-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one</p> 	<p>4-(5-Bromo-pyridin-3-ylmethyl)-thiomorpholine 1,1-dioxide (intermediate A-14) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	White solid	386.2
21	<p>6-[5-((S)-2-Methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one</p> 	<p>3-Bromo-5-((S)-2-methoxymethyl-pyrrolidin-1-ylmethyl)-pyridine (intermediate A-15) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	Colorless oil	366.3

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
22	<p>(S)-1-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-pyrrolidine-2-carboxylic acid methyl ester</p> 	<p>(S)-1-(5-Bromopyridin-3-ylmethyl)-pyrrolidine-2-carboxylic acid methyl ester (intermediate A-17) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	Brown oil	380.3
23	<p>1-Methyl-6-{5-[(S)-2-(2,2,2-trifluoro-ethoxymethyl)-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-3,4-dihydro-1H-quinolin-2-one</p> 	<p>3-Bromo-5-[(S)-2-(2,2,2-trifluoro-ethoxymethyl)-pyrrolidin-1-ylmethyl]-pyridine (intermediate A-19) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	Colorless waxy solid	434.4


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
24	<p>6-[6-(2-Hydroxy-ethylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	<p>2-(6-Chloro-pyrazin-2-ylamino)-ethanol (intermediate A-20) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Light brown solid	299.2
25	<p>6-(6-Benzylamino-pyrazin-2-yl)-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	<p>Benzyl-(6-chloro-pyrazin-2-yl)-amine (intermediate A-21) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Light yellow solid	345.1
26	<p>7-Fluoro-6-[6-(2-hydroxy-ethylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	<p>2-(6-Chloro-pyrazin-2-ylamino)-ethanol (intermediate A-20) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-22)</p>	White solid	317.1

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
27	<p>6-(6-Benzylamino-pyrazin-2-yl)-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one</p> 	<p>Benzyl-(6-chloro-pyrazin-2-yl)-amine (intermediate A-21) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-22)</p>	Yellow solid	362.9
28	<p>6-[6-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyrazin-2-yl]-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>6-(6-Chloro-pyrazin-2-yl)-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid <i>tert</i>-butyl ester (intermediate 24) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	Off white solid	436.1


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
29	<p>6-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>6-(5-Bromo-pyridin-3-yl)-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid <i>tert</i>-butyl ester (intermediate 25) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Brown waxy solid	435.3

Example 30

5 **6-[5-((*R*)-1-Hydroxymethyl-2-methyl-propylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one**




[A] 6-{5-[(*R*)-1-(*tert*-butyl-dimethyl-silanyloxymethyl)-2-methyl-propylamino]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one



In analogy to the procedures described for the preparation of intermediate A-3 [B] and A-3 [C]: i) (R)-1-(*tert*-butyl-dimethyl-silanyloxymethyl)-2-methyl-propylamine has been reacted with 3,5-dibromopyridine to give (5-bromo-pyridin-3-yl)-[(R)-1-(*tert*-butyl-dimethyl-silanyloxymethyl)-2-methyl-propyl]-amine; ii) subsequent condensation with 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) gave the title compound as light yellow amorphous solid. MS: 454.3 ($M+H^+$).


[B] 6-[5-((R)-1-Hydroxymethyl-2-methyl-propylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one



In a 10 mL round-bottomed flask, 6-{5-[(R)-1-(*tert*-butyl-dimethyl-silanyloxymethyl)-2-methyl-propylamino]-pyridin-3-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (60 mg, 132 μ mol) was dissolved in THF (1.5 mL) to give a light yellow solution and cooled down to 0 °C. TBAF, 1 M sol. in THF (132 μ l, 132 μ mol) was added dropwise. The solution was stirred at 0 °C for 1.5 h and partitioned between aqueous sat. NaHCO_3 solution and AcOEt (3 x). The organic layers were collected, dried over Na_2SO_4 and evaporated in vacuo. The crude material was purified by flash chromatography (silica gel, 20 g, 100% EtOAc followed by 1% MeOH in EtOAc) to give the title compound (22 mg, 49%) as a light yellow foam. MS: 340.2 ($M+H^+$).

Example 31

6-[6-((R)-1-Hydroxymethyl-2-methyl-propylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one




- In analogy to the procedures described for the preparation of intermediate A-20 and A-3
- 5 [C]: i) (R)-2-amino-3-methyl-butan-1-ol has been reacted with 2,6-dichloropyrazine to give (R)-2-(6-chloro-pyrazin-2-ylamino)-3-methyl-butan-1-ol; ii) subsequent condensation with 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1) gave the title compound as a light yellow solid. MS: 341.2 (M+H⁺).


10


Example 32

Ethanesulfonic acid [5-(5-methyl-6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-pyridin-3-ylmethyl]-amide




- 15 [A] 6-Bromo-3-nitro-2-(toluene-4-sulfonylmethyl)-pyridine




and 2-bromo-5-nitro-4-(toluene-4-sulfonylmethyl)-pyridine

To a stirred solution of 2-bromo-5-nitro-pyridine (3.4 g, 25.0 mmol) and 1-chloromethanesulfonyl-4-methyl-benzene (3.4 g, 25 mmol) in THF (100 mL) was added t-BuOK solution (1.0 M in THF, 55 mL) at -78 °C. After stirring for 30 minutes, AcOH (3.0 mL) was added to the above solution and the reaction mixture was allowed to warm up to RT and stirred for additional 20 minutes. After extraction of the reaction mixture with EtOAc, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give a suspension (50 mL), which after filtration gave 2-bromo-5-nitro-4-(toluene-4-sulfonylmethyl)-pyridine as a pale white solid (3.25 g, 35%); the filtrate was concentrated in vacuo to afford 6-bromo-3-nitro-2-(toluene-4-sulfonylmethyl)-pyridine as a light yellow solid (4.2 g, 45%). MS: 371.1 & 373.1 (M+H⁺).

[B] 3-(6-Bromo-3-nitro-pyridin-2-yl)-3-(toluene-4-sulfonyl)-propionic acid methyl ester


A mixture of 6-bromo-3-nitro-2-(toluene-4-sulfonylmethyl)-pyridine (3.3 g, 9.0 mmol), bromo-acetic acid methyl ester (2.1 mL, 22.5 mmol) and K₂CO₃ (8.4 g, 60.8 mmol) were suspended in DMF (18.0 mL) and stirred at 40 °C for 1 hour before pouring into water (50 mL). After extraction with EtOAc, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give 3-(6-bromo-3-nitro-pyridin-2-yl)-3-(toluene-4-sulfonyl)-propionic acid methyl ester (3.9 g, quant.) as a light yellow solid. MS: 443.1 & 445.1 (M+H⁺).

[C] 6-(2-Methoxycarbonyl-vinyl)-5-nitro-[2,3']bipyridinyl-5'-carboxylic acid methyl ester




3-(6-Bromo-3-nitro-pyridin-2-yl)-3-(toluene-4-sulfonyl)-propionic acid methyl ester (1.2 g, 2.7 mmol), Pd(PPh₃)₂Cl₂ (190 mg, 0.27 mmol), Na₂CO₃ (572 mg, 5.4 mmol) and 3-(methoxycarbonyl)pyridine-5-boronic acid pinacol ester (926 mg, 3.5 mmol) were dissolved in 1,4-dioxane (4.0 mL) and the resulting reaction mixture was heated at 120 °C for 3 hours before it was poured into H₂O (50 mL). After extraction with EtOAc, the organic layer was washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo to give 6-(2-methoxycarbonyl-vinyl)-5-nitro-[2,3']bipyridinyl-5'-carboxylic acid methyl ester (926 mg, quant.) as a light yellow solid. MS: 344.1 (M+H⁺).

[D] 5-(6-Oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-nicotinic acid methyl ester



6-(2-Methoxycarbonyl-vinyl)-5-nitro-[2,3']bipyridinyl-5'-carboxylic acid methyl ester (926 mg, 2.7 mmol), 10% Pd/C (300 mg) and AcOH (1.3 mL) were suspended in methanol (300 mL) and the reaction mixture was stirred at 50 °C under 50 psi H₂ pressure for 13 hours. After filtration, the reaction mixture was concentrated in vacuo to give a crude product of 5-(6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-nicotinic acid methyl ester (500 mg, 65.4%) as a light yellow solid. MS: 284.1 (M+H⁺).


[E] 5-(5-Methyl-6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-nicotinic acid methyl ester



To a stirred solution of 5-(6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-nicotinic acid methyl ester (430 mg, 1.52 mmol) in THF (15.0 mL) was added 60% NaH (91 mg, 2.28 mmol) at 0 °C and the reaction mixture was stirred at 2-5 °C for 0.5 h before CH₃I (0.3


mL, 4.56 mmol) was added. After stirring overnight at RT, it was poured into water (5.0 mL). After extraction with EtOAc, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give 5-(5-methyl-6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-nicotinic acid methyl ester (383.7 mg, 85%) as a light yellow solid. MS: 298.1 (M+H⁺).

[F] 6-(5-Hydroxymethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-[1,5]naphthyridin-2-one




To a stirred solution of 5-(5-methyl-6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-nicotinic acid methyl ester (270 mg, 0.91 mmol) in MeOH (10 mL) was added sodium borohydride (300 mg, 8.0 mmol) at RT. After stirring overnight, the reaction mixture was quenched by adding water (5.0 mL). It was then concentrated in vacuo to give a light yellow oil. After extraction with EtOAc, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give 6-(5-hydroxymethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-[1,5]naphthyridin-2-one (190 mg, 77.6%) as a white solid. MS: 270.1 (M+H⁺).

[G] 6-(5-Chloromethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-[1,5]naphthyridin-2-one



At 0 °C, 6-(5-hydroxymethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-[1,5]naphthyridin-2-one (170 mg, 0.63 mmol) in DCM (15 mL) was treated with thionyl chloride (0.32 mL, 4.0 mmol). After the addition, the reaction mixture was allowed to stir at 2-5 °C for 2 hours before it was poured into satd. aq. NaHCO₃ solution (50 mL). After extraction with EtOAc, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo to give 6-(5-chloromethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-[1,5]naphthyridin-2-one (156 mg, 86.3%) as yellow oil. MS: 288.1 (M+H⁺).


[H] Ethanesulfonic acid [5-(5-methyl-6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-pyridin-3-ylmethyl]-amide



To a stirred solution of ethanesulfonic acid amide (118 mg, 1.08 mmol) in DMF (5.0 mL)
 5 was added 60% NaH (35 mg, 0.81 mmol) at RT and the resulting reaction mixture was stirred for 0.5 hour before 6-(5-chloromethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-[1,5]naphthyridin-2-one (156 mg, 0.54 mmol) was added. After continued stirring at RT for additional 2 hours, water (1.0 mL) was added. After extraction with EtOAc, the organic layer was washed with brine, dried over anhy. Na₂SO₄, filtered and concentrated in vacuo
 10 to give a crude product which was purified by prep-HPLC to give title compound (15 mg, 7.7%) as a light yellow solid. MS: 361.1 (M+H⁺).

Example 33

6-Pyridin-3-yl-3,4-dihydro-1H-[1,5]naphthyridin-2-one




15

In analogy to the procedures described for the preparation of example 32 (steps [C] to [D]), the title compound was prepared using 3-(6-bromo-3-nitro-pyridin-2-yl)-3-(toluene-4-sulfonyl)-propionic acid methyl ester and 3-pyridine boronic acid as corresponding starting materials. MS: 226.1(M+H⁺).

20

Example 34

1-Methyl-6-pyridin-3-yl-3,4-dihydro-1H-[1,5]naphthyridin-2-one




In analogy to the procedures described for the preparation of example 32 (step [E]), the title compound was prepared using 6-pyridin-3-yl-3,4-dihydro-1*H*-[1,5]naphthyridin-2-one (example 33) as corresponding starting material. MS: 240.2 ($M+H^+$).

5

Example 35


1-Methyl-6-pyridin-3-yl-3,4-dihydro-1*H*-[1,7]naphthyridin-2-one



10 In analogy to the procedures described for the preparation of example 32 (steps [B] to [E]), the title compound was prepared using 2-bromo-5-nitro-4-(toluene-4-sulfonylmethyl)-pyridine (example 32 [A]) and 3-pyridine boronic acid as corresponding starting materials. MS: 240.1 ($M+H^+$).

Example 36

15 6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




A round-bottomed flask was charged with 5-bromo-pyridin-3-yl-methylamine (0.5 g, 2.67 mmol), 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1, 0.845 g, 2.94 mmol) and EtOH (48 mL). Then, *tetrakis*(triphenylphosphine)palladium(0) (0.093 g, 0.080 mmol), followed by aqueous

5 Na_2CO_3 solution (8 mL, 0.312 g, 2.94 mmol) were added and the reaction mixture heated to 85 °C over night. The mixture was evaporated to dryness, the residue was taken up in EtOAc, filtered through Dicalite and washed with EtOAc (2 x 50 mL). The filtrate was evaporated to dryness. Then, the residue was purified by silica gel flash chromatography eluting with a 0 to 20% MeOH(1% NH_4OH)-DCM gradient to give the title compound

10 (0.7 g, 94 %) as an off white solid. MS: 268.2 ($\text{M}+\text{H}^+$).

Example 37

***N*-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide**




To a solution of 6-(5-aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (example 36, 0.05 g, 0.187 mmol) in dry DMF (1 mL) were added EDCI (0.039 g, 0.206 mmol), Hünig's base (0.060 g, 0.468 mmol) and propionic acid (0.021 g, 0.282 mmol) and the reaction mixture was stirred at room temperature over night. The reaction mixture was

20 diluted with EtOAc, poured into sat. NH_4Cl solution (5 mL) and extracted with EtOAc (2 x 10 mL). Combined organics were dried over Na_2SO_4 , filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 10% MeOH(1% NH_4OH)-DCM gradient to give the title compound (0.029 g, 48 %) as a white solid. MS: 324.4 ($\text{M}+\text{H}^+$).

25

Example 38


Propane-2-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide



To a solution of 6-(5-aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (example 36, 0.05 g, 0.187 mmol) in DCM (1 mL) cooled to 0 °C were added triethylamine (0.038 g, 0.374 mmol) and isopropylsulfonylchloride (0.029 g, 0.206 mmol). The resulting suspension was stirred at 0 °C for 4h and then allowed to warm up to room temperature and stirring was continued for 2h. The reaction mixture was diluted with DCM, poured into H₂O (5 mL) and extracted with DCM (2 x 10 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 10% MeOH(1% NH₄OH)-DCM gradient to give the title compound (0.023 g, 33 %) as a white foam. MS: 374.1 (M+H⁺).

Example 39

{2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-carbamic acid *tert*-butyl ester




In analogy to the procedure described for the preparation of intermediate A-3 [C], 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) has been coupled to [2-(5-bromo-pyridin-3-yloxy)-ethyl]-carbamic acid

tert-butyl ester (intermediate A-7) to give the title compound as a white solid. MS: 398.1 ($M+H^+$).

Example 40


5 **3-Methoxy-isoxazole-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide**



To a solution of 6-(5-aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (example 36, 0.05 g, 0.187 mmol) in dry DCM (1 mL) were added EDCI (0.039 g, 0.206 mmol), hydroxybenzotriazole (0.032 g, 0.206 mmol), Hünig's base (0.060 g, 0.468 mmol) and 3-methoxy-isoxazole-5-carboxylic acid (0.040 g, 0.281 mmol) and the resulting solution was stirred at room temperature over night. The reaction mixture was diluted with EtOAc, poured into sat. NaHCO₃ solution (5 mL) and extracted with EtOAc (2 x 10 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 10% MeOH(1% NH₄OH)-DCM gradient to give the title compound (0.03 g, 41 %) as a white solid. MS: 393.1 ($M+H^+$).

Example 41

20 **Cyclopropanecarboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide**




In analogy to the procedure described for the preparation of example 37, 6-(5-aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (example 36) has been reacted with cyclopropanecarboxylic acid to give the title compound as a white solid.

5 MS: 336.3 ($M+H^+$).

Example 42

6-[5-(2-Amino-ethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride



10


To a solution of {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-carbamic acid *tert*-butyl ester (example 39, 0.06 g, 0.151 mmol) in MeOH (1 mL) was added 4M HCl in dioxane (0.151 mL, 0.604 mmol) and the reaction mixture was stirred at room temperature over night. The resulting suspension was filtered off and the solid material was triturated in diethyl ether, filtered off and further dried in a high vacuum to give the title compound (0.046 g, 91 %) as a white solid. MS: 298.3 ($M+H^+$).

15

Example 43

***N*-{2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-propionamide**


20



In analogy to the procedure described for the preparation of example 37, 6-[5-(2-aminoethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 42) has been reacted with propionic acid to give the title compound as a white solid. MS:
 5 354.3 ($M+H^+$).

Example 44

6-[5-(1,1-Dioxo-1 λ ⁶-[1,2]thiazinan-2-ylmethyl)pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




10

In analogy to the procedure described for the preparation of intermediate A-12 [B] and to the procedure described for the preparation of intermediate A-3 [C], [1,2]thiazinane 1,1-dioxide was reacted with 3-bromo-5-chloromethyl-pyridine (intermediate A-12 [A]) in the presence of NaH to give 2-(5-bromo-pyridin-3-ylmethyl)-[1,2]thiazinane 1,1-dioxide,
 15 which was subsequently reacted with 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) to give the title compound as a off-white solid. MS: 386.2 ($M+H^+$).

Example 45

20 **(S)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)pyridin-3-yloxymethyl]-azetidine-1-carboxylic acid *tert*-butyl ester**





A sealed tube was charged with (S)-2-(5-bromo-pyridin-3-yloxymethyl)-azetidine-1-carboxylic acid *tert*-butyl ester (intermediate A-26, 0.260 g, 0.76 mmol), 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1, 0.239 g, 0.83 mmol) and DMF (3 mL). Then, *bis*(triphenylphosphine)palladium(II)chloride (0.053 g, 0.076 mmol), followed by 1*N* aqueous Na₂CO₃ solution (2.27 mL, 2.27 mmol) were added and the reaction was heated to 110 °C (pre-heated oil bath) for 1h. The mixture was diluted with EtOAc, filtered through Dicalite and washed with EtOAc (20 mL). The resulting filtrate was poured into aq. NaHCO₃ (20 mL) and the aqueous layer was extracted with EtOAc (20 mL). Combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated. The residue was purified by silica gel flash chromatography eluting with a 0 to 100 % EtOAc-heptane gradient to give the title compound (0.277 g, 86%) as a light brown foam. MS: 424.1 (M+H⁺).



15



The following compounds listed in Table 3 were prepared in analogy to the procedure described for the preparation of example 45, using the appropriate starting materials.



Table 3



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
46	<p>(R)-3-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>(R)-3-(5-Bromopyridin-3-yloxy)-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester (intermediate A-27) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-22)</p>	Colorless solid	442.3
47	<p>3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-azetidine-1-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>3-(5-Bromopyridin-3-yloxy)-azetidine-1-carboxylic acid <i>tert</i>-butyl ester (intermediate A-28) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless solid	410.5


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
48	<p>(S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>(S)-3-(5-Bromopyridin-3-yloxy)-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester (intermediate A-29) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless solid	424.1
49	<p>(S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-piperidine-1-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>(S)- 3-(5-Bromopyridin-3-yloxy)-piperidine-1-carboxylic acid <i>tert</i>-butyl ester (intermediate A-30) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless oil	438.3



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
50	<p>4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-piperidine-1-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>4-(5-Bromopyridin-3-yloxy)-piperidine-1-carboxylic acid <i>tert</i>-butyl ester (intermediate A-31) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless amorphous solid	438.3
51	<p>(S)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>(S)-2-((5-Bromopyridin-3-yloxy)methyl)pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester (intermediate A-32) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless solid	438.5

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
52	<p>(S)-3-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>(S)-3-(5-Bromopyridin-3-yloxy)-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester (intermediate A-29) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-22)</p>	Colorless solid	n.d.
53	<p>(R)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester</p> 	<p>(R)-3-(5-Bromopyridin-3-yloxy)-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester (intermediate A-27) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless solid	424.2

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
54	<p>(R)-2-Methyl-propane-2-sulfinic acid {3-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-oxetan-3-yl}-amide</p> 	<p>(R)-N-(3-(5-Bromopyridin-3-yl)oxetan-3-yl)-2-methylpropane-2-sulfinamide (intermediate A-33) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)</p>	Amorphous brown solid	414.4
55	<p>Ethanesulfonic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p>Ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) and 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroquinolin-2(1H)-one (intermediate A-34)</p>	Amorphous colorless solid	346.1

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
56	<p>(R)-2-Methyl-propane-2-sulfinic acid {(S or R)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p>  <p>(S or R)</p>	<p>(R)-2-Methyl-propane-2-sulfinic acid [(S and R)-1-(5-bromo-pyridin-3-yl)-ethyl]-amide (intermediate A-35) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1) followed by HPLC separation.</p>	Amorphous colorless solid	386.1
57	<p>(R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p>  <p>(R or S)</p>	<p>(R)-2-Methyl-propane-2-sulfinic acid [(S and R)-1-(5-bromo-pyridin-3-yl)-ethyl]-amide (intermediate A-35) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1) followed by HPLC separation.</p>	Amorphous colorless solid	386.1

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
58	<p>{(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-carbamic acid <i>tert</i>-butyl ester</p> 	<p><i>tert</i>-Butyl (trans)-4-(5-bromopyridin-3-yloxy)cyclohexylcarbamate (intermediate A-54) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1).</p>	Off-white solid	452.5


Example 59**6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one****5 [A] Methyl 1-(5-bromopyridin-3-yl)cyclopropanecarboxylate**

To a solution of methyl 2-(5-bromopyridin-3-yl)acetate (4 g, 17.4 mmol) in DMF (80 mL), cooled at 0 °C with an ice bath, was added 60 % NaH in mineral oil (0.918 g, 38.3 mmol) and the reaction mixture was stirred at 0 °C for 15 min. Then, a solution of 1,2-

10 dibromoethane (3.27 g, 17.4 mmol) in DMF (16 mL) was added at 0 °C. After the addition, the mixture was stirred at room temperature for 1h. Two other portions of 60 %


NaH in mineral oil (2 x 0.2 g) were added sequentially, until conversion was complete. Then, the reaction mixture was poured into H₂O (50 mL) and extracted with EtOAc (3 x 125 mL). The organic layers were combined, dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (4.764 g, 99 %) as a brown oil, which was used with
5 no further purification. MS: 256.0 and 258.0 (M+H⁺).

[B] Potassium 1-(5-bromopyridin-3-yl)cyclopropanecarboxylate



To a solution of methyl 1-(5-bromopyridin-3-yl)cyclopropanecarboxylate (4.764 g, 18.6 mmol) in THF (190 mL) was added 90 % potassium trimethylsilanolate (2.65 g, 18.6
10 mmol) and the reaction mixture was stirred at room temperature over night. Then, the mixture was filtered and washed with THF to give the desired product (2.668 g) as a light brown solid. The mother liquors were concentrated to around 50 mL of THF. Another portion of 90 % potassium trimethylsilanolate (0.43 g) was added and stirring was continued for 5 h at room temperature. Then, the mixture was filtered and washed again
15 with THF to give a second batch of desired product (0.953 g) which was combined with the first batch to give the title compound (3.621 g, 59.1 %) as a light brown solid. MS: 242.0 and 244.1 (M+H⁺).

[C] 1-(5-(1-Methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)pyridin-3-yl)cyclopropane-carboxylic acid




20

A flask was charged with potassium 1-(5-bromopyridin-3-yl)cyclopropanecarboxylate (0.5 g, 1.52 mmol), 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1, 0.479 g, 1.67 mmol) and DMF (7 mL). Then, *bis*(triphenylphosphine)palladium(II)chloride (0.106 g, 0.152 mmol), followed by 1N
25 aqueous Na₂CO₃ solution (3.64 mL, 2.4 mmol) were added and the reaction was heated to

120 °C (pre-heated oil bath) for 1h. The mixture was evaporated to dryness and the residue purified by silica gel flash chromatography eluting with a 0 to 15 % MeOH-DCM gradient to give the title compound (0.551 g, 100%) as a brown solid. MS: 323.4 (M+H⁺).

[D] 6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one



5

To solution of 1-(5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)pyridin-3-yl)cyclopropanecarboxylic acid (0.551 g, 1.52 mmol) in toluene (20 mL) were added TEA (0.185 g, 1.83 mmol) and diphenylphosphoryl azide (0.419 g, 1.52 mmol) and then the reaction mixture was heated to reflux for 4h. After cooling to 0 °C, a 1 M solution of sodium trimethylsilanoate in THF (3.04 mL, 3.04 mmol) was added and the mixture was stirred for 45 min at room temperature. After quenching with 0.1 M aq. HCl (40 mL), the aqueous solution was washed with ether (2 times), then basified with 1 M aq. NaOH and then extracted with DCM (2 x 100 mL). The combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated to dryness to give the title compound (0.333 g, 71%) as a yellow foam. MS: 294.4 (M+H⁺).



15



The following compounds listed in Table 4 were prepared by treatment of the appropriate starting material with HCl in methanol in analogy to the procedure described for the preparation of example 42 (isolation of the compounds as HCl salts by direct evaporation or by basic extraction followed by silica gel or reverse phase chromatography):



20



Table 4



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
60	<p>1-Methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride</p> 	<p>(S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester (example 48)</p>	Yellow solid	324.3
61	<p>1-Methyl-6-[5-((S)-piperidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one</p> 	<p>(S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-piperidine-1-carboxylic acid <i>tert</i>-butyl ester (example 49)</p>	Colorless amorphous solid	338.3

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
62	<p>1-Methyl-6-[5-(piperidin-4-yloxy)-pyridin-3-yl]-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride</p> 	<p>4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-piperidine-1-carboxylic acid <i>tert</i>-butyl ester (example 50)</p>	<p>Light yellow solid</p>	<p>338.4</p>
63	<p>6-[5-((S)-1-Azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	<p>(S)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-azetidine-1-carboxylic acid <i>tert</i>-butyl ester (example 45)</p>	<p>Light yellow oil</p>	<p>324.2</p>

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
64	<p>6-[5-((S or R)-1-Amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride</p>  <p>(S or R)</p>	(R)-2-Methylpropane-2-sulfinic acid {(S or R)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-yl]-ethyl}-amide (example 56)	Colorless solid	282.1
65	<p>6-[5-((R or S)-1-Amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride</p>  <p>(R or S)</p>	(R)-2-Methylpropane-2-sulfinic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-yl]-ethyl}-amide (example 57)	Colorless solid	282.1


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
66	<p>1-Methyl-6-[5-((S)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride</p> 	(S)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-pyrrolidine-1-carboxylic acid <i>tert</i> -butyl ester (example 51)	Yellow solid	338.2
67	<p>7-Fluoro-1-methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride</p> 	(R)-3-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid <i>tert</i> -butyl ester (example 46)	Light yellow solid	342.3

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
68	<p>1-Methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride</p> 	<p>(R)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid <i>tert</i>-butyl ester (example 53)</p>	Yellow solid	324.5
69	<p>6-[5-(3-Amino-oxetan-3-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride</p> 	<p>(R)-2-Methylpropane-2-sulfinic acid {3-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-oxetan-3-yl}-amide (example 54)</p>	Light yellow solid	310.2


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
70	<p>6-[5-(Azetidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride</p> 	<p>3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-azetidine-1-carboxylic acid <i>tert</i>-butyl ester (example 47)</p>	Yellow solid	310.3
71	<p>6-[5-((<i>trans</i>)-4-Amino-cyclohexyloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride</p> 	<p>{(<i>trans</i>)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-carbamic acid <i>tert</i>-butyl ester (example 58)</p>	Yellow solid	352.5

Example 72

6-[5-(1-Amino-1-methyl-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




[A] Methyl 2-(5-bromopyridin-3-yl)-2-methylpropanoate



To a solution of methyl 2-(5-bromopyridin-3-yl)acetate (0.7 g, 3.04 mmol) in DMF (14
 5 mL) cooled at 0 °C with an ice bath was added 60 % NaH in mineral oil (0.244 g, 6.09
 mmol) and the reaction mixture was stirred for 15 min. Then, a solution of MeI (0.864 g,
 6.09 mmol) in DMF (16 mL) was added dropwise at 0 °C. After the addition, the mixture
 was stirred at room temperature for 1h. An other portion of 60 % NaH in mineral oil
 (0.073 g) was added to the mixture which was stirred at room temperature for 1h. The
 10 reaction mixture was poured into H₂O (15 mL) and the aqueous layer was extracted with
 EtOAc (3 x 50 mL). The organic layers were combined, dried over Na₂SO₄, filtered and
 evaporated to dryness to give the title compound (0.617 g, 75 %) as a brown oil which was
 used with no further purification. MS: 258.2 and 260.3 (M+H⁺).

[B] Potassium 2-(5-bromopyridin-3-yl)-2-methylpropanoate




15

In analogy to the procedure described for the preparation of example 59 [B], methyl 2-(5-
 bromopyridin-3-yl)-2-methylpropanoate (example 72 [A]) was reacted with 90 %
 potassium trimethylsilanolate to give the title compound as a light brown solid. MS: 244.3
 (M+H⁺).

20


[C] 2-Methyl-2-(5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)pyridin-3-yl)propan-

oic acid



In analogy to the procedure described for the preparation of example 59 [C], potassium 2-(5-bromopyridin-3-yl)-2-methylpropanoate (example 72 [B]) was reacted with 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) in presence of *bis*(triphenylphosphine)palladium(II)chloride and Na₂CO₃ (1M aq. solution) to give the title compound as a brown foam. MS: 325.4 (M+H⁺).

[D] 6-[5-(1-Amino-1-methyl-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




10

In analogy to the procedure described for the preparation of example 59 [D], 2-methyl-2-(5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)pyridin-3-yl)propanoic acid (example 72 [C]) was reacted with diphenylphosphoryl azide in presence of TEA and then sodium trimethylsilanoate (1 M in THF) to give the title compound as a light brown amorphous solid. MS: 296.5 (M+H⁺).

15

Example 73


6-(5-Aminomethyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one



In analogy to the procedure described for the preparation of example 45, reaction of 3-aminomethyl-5-bromopyridine and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-22) gave the
 5 title compound as a light yellow solid. MS: 286.7 ($M+H^+$).

Example 74

6-{6-[6-(3-Methoxy-isoxazole-5-carbonyl)-2,6-diaza-spiro[3.3]hept-2-yl]-pyrazin-2-yl}-1-methyl-3,4-dihydro-1*H*-quinolin-2-one




10

To a solution of 6-(6-(2,6-diazaspiro[3.3]heptan-2-yl)pyrazin-2-yl)-1-methyl-3,4-dihydroquinolin-2(1*H*)-one 2,2,2-trifluoroacetate (intermediate A-36, 0.030 g, 0.067 mmol) in DCM (1 mL) were added 3-methoxyisoxazole-5-carboxylic acid (0.014 g, 0.1 mmol) and TBTU (0.024 g, 0.073 mmol) followed by Hünig's base (0.022 g, 0.167 mmol)
 15 and the reaction mixture was stirred at room temperature over night. The reaction mixture was diluted with EtOAc, poured into sat. NaHCO_3 solution (5 mL) and extracted with EtOAc (2 x 10 mL). Combined organics were dried over Na_2SO_4 , filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.007 g, 20 %) as a yellow solid.
 20 MS: 461.3 ($M+H^+$).

Example 75


3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide



To a solution of 6-(5-aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (example 36, 0.047 g, 0.177 mmol) in DMF (1 mL) were added 3,5-dimethylisoxazole-4-carboxylic acid (0.037 g, 0.266 mmol) and TBTU (0.063 g, 0.195 mmol) followed by Hünig's base (0.048 g, 0.372 mmol) and the reaction mixture was stirred at room temperature over night. The mixture was purified directly by reverse phase HPLC on a Gemini-NX column, eluting with a 20 to 98% MeOH-H₂O (0.05 % TEA) gradient to give the title compound (0.041 g, 59 %) as a colorless solid. MS: 391.3 (M+H⁺).

Example 76

6-[5-((S)-1-Cyclopropanecarbonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one



15


To a solution of 1-methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 60, 0.05 g, 0.139 mmol) in dry DCM (1.5 mL) was added cyclopropanecarbonyl chloride (0.015 g, 0.139 mmol) followed by TEA (0.035 g, 0.347 mmol) and the reaction mixture was stirred at room temperature for 1h. Then, the reaction mixture was diluted with EtOAc, poured into sat. NaHCO₃ solution (10 mL) and extracted with EtOAc (2 x 20 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography



20



eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.032 g, 59 %) as a colorless foam. MS: 392.2 ($M+H^+$).



The following compounds listed in Table 5 were prepared by treatment of the appropriate starting materials under the appropriate coupling conditions as described in examples 40, 74, 75 or 76.



Table 5



Ex	Name	Starting Materials	Aspect	MS ($M+H^+$)
			Prep. by analogy to	
77	3-Methyl-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-butyramide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3-methyl-butyric acid	Colorless solid	352.4
			expl. 74	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
78	3,3,3-Trifluoro- <i>N</i> -[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 3,3,3-trifluoropropionic acid	Colorless solid	378.3
			expl. 74	
79	2-Hydroxy-2-methyl- <i>N</i> -[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 2-hydroxy-2-methylpropionic acid	Colorless solid	354.3
			expl. 74	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
80	5-Methyl-[1,3,4]oxadiazole-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 5-methyl-[1,3,4]oxadiazole-2-carboxylic acid	Colorless solid	378.3
			expl. 75	
81	1-Methyl-6-[5-((S)-1-propionyl-piperidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1 <i>H</i> -quinolin-2-one 	1-Methyl-6-[5-((S)-piperidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 61) and propionic acid	Orange amorphous solid	394.1
			expl. 74	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
82	2-Methoxy-pyrimidine-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 2-methoxy-pyrimidine-5-carboxylic acid	Colorless solid	404.3
			expl. 75	
83	1-Methyl-1H-imidazole-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 1-methyl-1H-imidazole-2-carboxylic acid	Colorless solid	376.2
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
84	5-Trifluoromethyl-furan-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 5-trifluoromethyl-furan-3-carboxylic acid	Colorless amorphous solid	430.4
			expl. 75	
85	Pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and pyridazine-3-carboxylic acid	Colorless solid	374.3
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
86	Pyrimidine-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4- tetrahydro-quinolin-6-yl)- pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl- pyridin-3-yl)-1- methyl-3,4-dihydro- 1 <i>H</i> -quinolin-2-one (example 36) and pyrimidine-5- carboxylic acid	Light brown solid	374.3
			expl. 75	
87	1-Methyl-1 <i>H</i> -pyrazole-4- carboxylic acid [5-(1-methyl- 2-oxo-1,2,3,4-tetrahydro- quinolin-6-yl)-pyridin-3- ylmethyl]-amide 	6-(5-Aminomethyl- pyridin-3-yl)-1- methyl-3,4-dihydro- 1 <i>H</i> -quinolin-2-one (example 36) and 1- methyl-1 <i>H</i> - pyrazole-4- carboxylic acid	Colorless solid	376.4
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
88	1-Methyl-6-[5-(1-propionyl- piperidin-4-yloxy)-pyridin-3- yl]-3,4-dihydro-1 <i>H</i> -quinolin- 2-one 	1-Methyl-6-[5-(piperidin-4-yloxy)- pyridin-3-yl]-3,4- dihydro-1 <i>H</i> - quinolin-2-one hydrochloride (example 62) and propionic acid	Orange amorphou s solid	394.1
			expl. 74	
89	Pyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4- tetrahydro-quinolin-6-yl)- pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl- pyridin-3-yl)-1- methyl-3,4-dihydro- 1 <i>H</i> -quinolin-2-one (example 36) and pyridazine-4- carboxylic acid	Colorless amorphou s solid	374.3
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
90	1-Methyl-6-[5-((S)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one 	1-Methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 60) and propionic acid	Light yellow oil	380.3
			expl. 74	
91	6-[5-((S)-1-Cyclopropanecarbonyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one 	6-[5-((S)-1-Azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 63) and cyclopropanecarbonyl chloride	Off-white amorphous solid	392.2
			expl. 76	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
92	3-Methyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3-methyl-isoxazole-4-carboxylic acid	Light brown solid	377.1
			expl. 75	
93	4-Fluoro-2,6-dimethyl-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-benzamide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 4-fluoro-2,6-dimethylbenzoic acid	Colorless solid	418.3
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
94	1-Methyl-6-[5-((S)-1-propionyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one 	6-[5-((S)-1-Azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 63) and propionyl chloride	Light yellow oil	380.3
			expl. 76	
95	3,6-Dichloro-pyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3,6-dichloro-pyridazine-4-carboxylic acid	Colorless amorphous solid	442.3
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
96	3-Cyclopropyl-1-(2,2,2-trifluoro-ethyl)-1 <i>H</i> -pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 3-cyclopropyl-1-(2,2,2-trifluoro-ethyl)-1 <i>H</i> -pyrazole-4-carboxylic acid	Colorless solid	484.4
			expl. 75	
97	Pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and pyridine-2-carboxylic acid	Colorless solid	373.3
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
98	1,3-Dimethyl-1 <i>H</i> -pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 1,3-dimethyl-1 <i>H</i> -pyrazole-4-carboxylic acid	Colorless amorphous solid	390.3
			expl. 75	
99	Pyrimidine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and pyrimidine-2-carboxylic acid	Colorless solid	374.3
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
100	6-Methoxy-pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 6-methoxy-pyridazine-3-carboxylic acid	Colorless solid	404.5
			expl. 75	
101	5-Methyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 5-methyl-isoxazole-4-carboxylic acid	Colorless solid	377.1
			expl. 40	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
102	3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide 	6-[5-(2-Aminoethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 42) and 3,5-dimethyl-isoxazole-4-carboxylic acid	Light yellow solid	421.1
			expl. 75	
103	1-Methyl-1H-pyrazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide 	6-[5-(2-Aminoethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 42) and 1-methyl-1H-pyrazole-4-carboxylic acid	Colorless solid	406.4
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
104	6-Chloro-pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 6-chloro-pyridazine-3-carboxylic acid	Colorless solid	408.3
			expl. 40	
105	3-Chloro-6-methylpyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3-chloro-6-methylpyridazine-4-carboxylic acid	Colorless solid	422.0
			expl. 40	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
106	1-Methyl-6-[5-((S)-1-propionyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one 	1-Methyl-6-[5-((S)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 66) and propionyl chloride	Colorless amorphous solid	394.3
			expl. 76	
107	5-Cyclopropyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 5-cyclopropyl-isoxazole-4-carboxylic acid	Colorless amorphous solid	403.3
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
108	2-Methyl-5-trifluoromethyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 2-methyl-5-trifluoromethyl-oxazole-4-carboxylic acid	Colorless solid	445.3
			expl. 75	
109	2-Methyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 2-methyl-oxazole-4-carboxylic acid	Off-white solid	377.4
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
110	5-Cyclopropyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 5-cyclopropyl-oxazole-4-carboxylic acid	Colorless amorphous solid	403.6
			expl. 75	
111	2,5-Dimethyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 2,5-dimethyl-oxazole-4-carboxylic acid	Colorless solid	391.3
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
112	5-Methyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 5-methyl-oxazole-4-carboxylic acid	Colorless solid	377.4
			expl. 75	
113	3,5-Dimethyl-isoxazole-4-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide 	6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 59) and 3,5-dimethyl-isoxazole-4-carboxylic acid	Colorless solid	417.4
			expl. 40	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
114	6-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 6-chloro-pyridine-2-carboxylic acid	Off-white solid	407.2
			expl. 75	
115	3-Methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3-methyl-pyridine-2-carboxylic acid	Colorless solid	387.2
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
116	3,6-Dichloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3,6-dichloro-pyridine-2-carboxylic acid	Colorless solid	441.2
			expl. 75	
117	6-Methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 6-methyl-pyridine-2-carboxylic acid	Colorless solid	387.2
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
118	3-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3-chloro-pyridine-2-carboxylic acid	Off-white solid	407.2
			expl. 75	
119	3-Fluoro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3-fluoro-pyridine-2-carboxylic acid	Colorless solid	391.2
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
120	5-Chloro-3-methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 5-chloro-3-methyl-pyridine-2-carboxylic acid	Off-white solid	421.1
			expl. 75	
121	1-Methyl-6-[5-((R)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1 <i>H</i> -quinolin-2-one 	1-Methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1 <i>H</i> -quinolin-2-one hydrochloride (example 68) and propionyl chloride	Light yellow solid	380.2
			expl. 76	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
122	7-Fluoro-1-methyl-6-[5-((S)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one 	7-Fluoro-1-methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one (example 146) and propionyl chloride	Yellow solid	398.2
			expl. 76	
123	5-Trifluoromethyl-pyrimidine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 5-trifluoromethyl-pyrimidine-2-carboxylic acid	Colorless solid	442.5
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
124	5-Methyl-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 5-methyl-pyrazine-2-carboxylic acid	Colorless solid	388.3
			expl. 75	
125	5-Chloro-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 5-chloro-pyrazine-2-carboxylic acid	Colorless solid	408.4
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
126	5-Trifluoromethyl-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 5-trifluoromethyl-pyrazine-2-carboxylic acid	Colorless solid	442.4
			expl. 75	
127	3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 73) and 3,5-dimethyl-isoxazole-4-carboxylic acid	Colorless solid	409.5
			expl. 40	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
128	1-Methyl-6-[5-((R)-1-propionyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one 	1-Methyl-6-[5-((R)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 148) and propionyl chloride	Yellow solid	394.5
			expl. 76	
129	1-Methyl-6-[5-(1-propionyl-azetidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one 	6-[5-(Azetidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 70) and propionyl chloride	Colorless amorphous solid	366.5
			expl. 76	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
130	<p><i>N</i>-{3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-oxetan-3-yl}-propionamide</p> 	<p>6-[5-(3-Amino-oxetan-3-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride (example 69) and propionyl chloride</p>	Colorless amorphous solid	366.1
			expl. 76	
131	<p>3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p>6-(5-Aminomethyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one (example 73) and 3-chloro-pyridine-2-carboxylic acid</p>	Off-white solid	425.4
			expl. 75	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
132	3,6-Dichloro-pyridazine-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 73) and 3,6-dichloro-pyridazine-4-carboxylic acid	Colorless amorphous solid	460.3
			expl. 40	
133	7-Fluoro-1-methyl-6-[5-((R)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1 <i>H</i> -quinolin-2-one 	7-Fluoro-1-methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1 <i>H</i> -quinolin-2-one hydrochloride (example 67) and propionyl chloride	Yellow solid	398.5
			expl. 76	



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
134	5-Chloro-3-methoxy- pyrazine-2-carboxylic acid [5- (1-methyl-2-oxo-1,2,3,4- tetrahydro-quinolin-6-yl)- pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl- pyridin-3-yl)-1- methyl-3,4-dihydro- 1 <i>H</i> -quinolin-2-one (example 36) and 5- chloro-3-methoxy- pyrazine-2- carboxylic acid	Yellow amorphou s solid	438.1
			expl. 75	
135	3,5-Dimethyl-isoxazole-4- carboxylic acid {(R or S)-1- [5-(1-methyl-2-oxo-1,2,3,4- tetrahydro-quinolin-6-yl)- pyridin-3-yl]-ethyl}-amide  (R or S)	6-[5-((R or S)-1- Amino-ethyl)- pyridin-3-yl]-1- methyl-3,4-dihydro- 1 <i>H</i> -quinolin-2-one hydrochloride (example 65) and 3,5-dimethyl- isoxazole-4- carboxylic acid	Colorless solid	405.5
			expl. 74	

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
136	3-Methyl-5-trifluoromethyl- isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4- tetrahydro-quinolin-6-yl)- pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl- pyridin-3-yl)-1- methyl-3,4-dihydro- 1 <i>H</i> -quinolin-2-one (example 36) and 3- methyl-5- trifluoromethyl- isoxazole-4- carboxylic acid	Off-white solid	445.5
			expl. 40	
137	3-Chloro-pyridine-2- carboxylic acid {1-[5-(1- methyl-2-oxo-1,2,3,4- tetrahydro-quinolin-6-yl)- pyridin-3-yl]-cyclopropyl}- amide 	6-[5-(1-Amino- cyclopropyl)- pyridin-3-yl]-1- methyl-3,4-dihydro- 1 <i>H</i> -quinolin-2-one (example 59) and 3- chloro-pyridine-2- carboxylic acid	Off-white solid	433.5
			expl. 75	

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
138	3-Chloro-pyridine-2-carboxylic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide 	6-[5-(1-Amino-1-methyl-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 72) and 3-chloro-pyridine-2-carboxylic acid	Off-white amorphous solid	435.5
			expl. 75	
139	3,5-Dimethyl-isoxazole-4-carboxylic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide 	6-[5-(1-Amino-1-methyl-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 72) and 3,5-dimethyl-isoxazole-4-carboxylic acid	Off-white solid	419.5
			expl. 40	


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
140	3-Chloro-pyridine-2-carboxylic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-3,4-dihydro-1H-quinolin-2-one (example 149) and 3-chloro-pyridine-2-carboxylic acid	Light yellow solid	393.5
			expl. 75	
141	3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethyl-pyridin-3-yl)-3,4-dihydro-1H-quinolin-2-one (example 149) and 3,5-dimethyl-isoxazole-4-carboxylic acid	Colorless solid	377.5
			expl. 40	

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
142	3-Chloro-pyridine-2-carboxylic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide 	6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 150) and 3-chloro-pyridine-2-carboxylic acid	Light brown amorphous solid	451.5
			expl. 75	
143	3-Methyl-pyridine-2-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide 	6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 59) and 3-methyl-pyridine-2-carboxylic acid	Colorless solid	413.5
			expl. 75	

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
			Prep. by analogy to	
144	<p><i>N</i>-{((<i>trans</i>)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-propionamide</p> 	<p>6-[5-((<i>trans</i>)-4-Amino-cyclohexyloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one hydrochloride (example 71) and propionyl chloride</p>	Colorless solid	408.5
			expl. 76	
145	<p>5-Trifluoromethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p>6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one (example 36) and 5-trifluoromethyl-isoxazole-4-carboxylic acid</p>	Light brown amorphous solid	431.5
			expl. 40	

Example 146


7-Fluoro-1-methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one



- In analogy to the procedure described for the preparation of example 42, reaction of (S)-3-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid *tert*-butyl ester (example 52) with hydrogen chloride (in dioxane) in methanol gave the title compound as a yellow solid. MS: 342.1 (M+H⁺).

Example 147


- (R)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester**



- In analogy to the procedure described for the preparation of example 45, reaction of (R)-2-((5-bromopyridin-3-yloxy)methyl)pyrrolidine-1-carboxylic acid *tert*-butyl ester (intermediate A-55) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1) gave the title compound as a colorless solid. MS: 438.6 (M+H⁺).

Example 148


- 1-Methyl-6-[5-((R)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride**



In analogy to the procedure described for the preparation of example 42, reaction of (R)-2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-pyrrolidine-1-carboxylic acid *tert*-butyl ester (example 147) with hydrogen chloride (in dioxane) in
5 methanol gave the title compound as a yellow solid. MS: 338.2 (M+H⁺).

Example 149


6-(5-Aminomethyl-pyridin-3-yl)-3,4-dihydro-1H-quinolin-2-one



10 In analogy to the procedure described for the preparation of example 45, reaction of 3-aminomethyl-5-bromopyridine and 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroquinolin-2(1H)-one (intermediate A-34) gave the title compound as a light yellow amorphous solid. MS: 254.4 (M+H⁺).

Example 150


6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one









In analogy to the procedures described for the preparation of example 59, potassium 1-(5-bromopyridin-3-yl)cyclopropanecarboxylate (example 59 [B]) has been reacted with 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-22) and subsequently with diphenylphosphoryl azide and sodium trimethylsilanolate to give the title compound as a light yellow oil. MS: 312.5 ($M+H^+$).




The following compounds listed in Table 6 were prepared in analogy to the procedure described for the preparation of example 38, using the appropriate starting materials.




Table 6




Ex	Name	Starting Materials	Aspect	MS ($M+H^+$)
151	Ethanesulfonic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide 	6-[5-(2-Aminoethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one hydrochloride (example 42) and ethanesulfonyl chloride	Colorless oil	390.1




Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
152	3-Chloro- <i>N</i> -[5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-benzenesulfonamide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 3-chlorobenzenesulfonyl chloride	Light yellow solid	442.2
153	6-Methoxy-pyridine-3-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 6-methoxy-pyridine-3-sulfonyl chloride	Light yellow solid	439.3
154	3,5-Dimethyl-isoxazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide 	6-(5-Aminomethylpyridin-3-yl)-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 36) and 3,5-dimethyl-isoxazole-4-sulfonyl chloride	Yellow solid	427.2




Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
155	<p>Cyclopropanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and cyclopropanesulfonyl chloride	Off-white solid	372.2
156	<p>3,4-Dichloro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-benzenesulfonamide</p> 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 3,4-dichlorobenzenesulfonyl chloride	Light yellow solid	476.1
157	<p>1-Methyl-1H-imidazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 1-methyl-1H-imidazole-4-sulfonyl chloride	Off-white solid	412.3



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
158	<p>6-Chloro-pyridine-3-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 6-chloro-pyridine-3-sulfonyl chloride	Light yellow solid	443.2
159	<p>1-Methyl-1H-pyrazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 1-methyl-1H-pyrazole-4-sulfonyl chloride	Light brown solid	412.2
160	<p>6-[5-((S)-1-Ethanesulfonyl-piperidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one</p> 	1-Methyl-6-[5-((S)-piperidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one (example 61) and ethanesulfonyl chloride	Colorless solid	430.1

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
161	<p>6-[5-(1-Ethanesulfonyl-piperidin-4-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one</p> 	1-Methyl-6-[5-(piperidin-4-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 62) and ethanesulfonyl chloride	Colorless amorphous solid	430.1
162	<p>2,2,2-Trifluoro-ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and 2,2,2-trifluoro-ethanesulfonyl chloride	Colorless solid	414.1
163	<p>C,C,C-Trifluoro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methanesulfonamide</p> 	6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 36) and trifluoromethanesulfonyl chloride	Colorless amorphous solid	400.0

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
164	6-[5-((S)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one 	1-Methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1 <i>H</i> -quinolin-2-one hydrochloride (example 60) and ethanesulfonyl chloride	Colorless amorphous solid	416.1
165	6-[5-((S)-1-Ethanesulfonyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one 	6-[5-((S)-1-Azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 63) and ethanesulfonyl chloride	Colorless solid	416.1
166	Ethanesulfonic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide 	6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1 <i>H</i> -quinolin-2-one (example 59) and ethanesulfonyl chloride	Off-white solid	386.3

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
167	6-[5-((S)-1-Ethanesulfonyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one 	1-Methyl-6-[5-((S)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 66) and ethanesulfonyl chloride	Colorless amorphous solid	430.5
168	6-[5-((R)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one 	7-Fluoro-1-methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 67) and ethanesulfonyl chloride	Off-white solid	434.4
169	6-[5-((R)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one 	1-Methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 68) and ethanesulfonyl chloride	Colorless oil	416.4



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
170	<p>6-[5-(1-Ethanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one</p> 	6-[5-(Azetidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 70) and ethanesulfonyl chloride	Colorless amorphous solid	402.5
171	<p>Ethanesulfonic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide</p> 	6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one (example 150) and ethanesulfonyl chloride	Brown amorphous solid	404.5
172	<p>Ethanesulfonic acid {(trans)-4-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-amide</p> 	6-[5-((trans)-4-Amino-cyclohexyloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 71) and ethanesulfonyl chloride	Yellow amorphous solid	444.4



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
173	<p>Ethanesulfonic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p>  <p>(R or S)</p>	6-[5-((R or S)-1-Amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 65) and ethanesulfonyl chloride	Colorless amorphous solid	374.1
174	<p>Ethanesulfonic acid {(S or R)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p>  <p>(S or R)</p>	6-[5-((S or R)-1-Amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 64) and ethanesulfonyl chloride	Colorless amorphous solid	374.0



The following compounds listed in Table 7 were prepared in analogy to the procedure described for the preparation of example 45, using the appropriate starting materials.



5 **Table 7**



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
----	------	--------------------	--------	---------------------------



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
175	<p>1-Methyl-1<i>H</i>-pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p>1-Methyl-1<i>H</i>-pyrazole-4-carboxylic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-47) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-22)</p>	Colorless solid	394.0
176	<p>(rac)-Ethanesulfonic acid {2-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-propyl}-amide</p> 	<p>(rac)-<i>N</i>-(1-(5-Bromopyridin-3-yl)-2-methylpropyl)ethane sulfonamide (intermediate A-40) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Yellow solid	402.5



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
177	<p>(rac)-Ethanesulfonic acid {cyclopropyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-methyl}-amide</p> 	(rac)-N-((5-Bromopyridin-3-yl)(cyclopropyl)methyl)ethanesulfonamide (intermediate A-39) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)	Colorless amorphous solid	400.3
178	<p>(rac)-Ethanesulfonic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p> 	(rac)-N-(1-(5-Bromopyridin-3-yl)ethyl)ethanesulfonamide (intermediate A-37) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-22)	Colorless amorphous solid	392.1



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
179	<p>6-[5-(1,1-Dioxo-1λ⁶-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	<p>2-(5-Bromo-pyridin-3-ylmethyl)-isothiazolidine 1,1-dioxide (intermediate A-48) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless solid	372.1
180	<p>(rac)-Ethanesulfonic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-propyl}-amide</p> 	<p>(rac)-<i>N</i>-(1-(5-Bromopyridin-3-yl)propyl)ethanesulfonamide (intermediate A-38) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Light yellow amorphous solid	388.0



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
181	<p>Ethanesulfonic acid ethyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p><i>N</i>-((5-Bromopyridin-3-yl)methyl)-<i>N</i>-ethylethanesulfonamide (intermediate A-42) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	<p>Light yellow amorphous solid</p>	388.2
182	<p>Ethanesulfonic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p><i>N</i>-((5-Bromopyridin-3-yl)methyl)-<i>N</i>-methylethanesulfonamide (intermediate A-41) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	<p>Colorless solid</p>	374.1



Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
183	<p>Ethanesulfonic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p> 	<i>N</i> -(2-(5-Bromopyridin-3-yl)propan-2-yl)ethanesulfonamide (intermediate A-49) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1 <i>H</i> -quinolin-2-one (intermediate A-1)	Colorless solid	388.0
184	<p>Ethanesulfonic acid isopropyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<i>N</i> -((5-Bromopyridin-3-yl)methyl)- <i>N</i> -isopropylethanesulfonamide (intermediate A-43) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1 <i>H</i> -quinolin-2-one (intermediate A-1)	Colorless solid	402.2

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
185	<p>Ethanesulfonic acid (2-ethoxy-ethyl)-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p><i>N</i>-((5-Bromopyridin-3-yl)methyl)-<i>N</i>-2-ethoxy-ethylethanesulfonamide (intermediate A-44) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless solid	432.3
186	<p>(rac)-Ethanesulfonic acid methyl-{1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p> 	<p>(rac)-<i>N</i>-(1-(5-Bromopyridin-3-yl)ethyl)-<i>N</i>-methylethanesulfonamide (intermediate A-45) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless amorphous solid	388.4

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
187	<p>(rac)-Ethanesulfonic acid ethyl- {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p> 	(rac)-N-(1-(5-Bromopyridin-3-yl)ethyl)-N-ethylethanesulfonamide (intermediate A-46) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)	Colorless amorphous solid	402.5
188	<p>3,5-Dimethyl-isoxazole-4-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	N-((5-Bromopyridin-3-yl)methyl)-N,3,5-trimethylisoxazole-4-carboxamide (intermediate A-50) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1)	Colorless solid	405.5


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
189	<p>6-{5-[2-(1,1-Dioxo-1λ^6-isothiazolidin-2-yl)-ethoxy]-pyridin-3-yl}-1-methyl-3,4-dihydro-1<i>H</i>-quinolin-2-one</p> 	<p>3-Bromo-5-[2-(1,1-dioxo-1λ^6-isothiazolidin-2-yl)-ethoxy]-pyridine (intermediate A-51) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Light yellow oil	402.4
190	<p>Ethanesulfonic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p>  <p>(R or S)</p>	<p>(rac)-<i>N</i>-(1-(5-Bromopyridin-3-yl)ethyl)ethanesulfonamide (intermediate A-37) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-22) followed by separation with chiral phase HPLC.</p>	Colorless amorphous solid	392.1

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
191	<p>Ethanesulfonic acid {(S or R)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-yl]-ethyl}-amide</p>  <p>(S or R)</p>	<p>(rac)-N-(1-(5-Bromopyridin-3-yl)ethyl)ethanesulfonamide (intermediate A-37) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-22) followed by separation with chiral phase HPLC.</p>	Colorless amorphous solid	392.1
192	<p>Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide</p> 	<p>N-((5-Bromo-4-methylpyridin-3-yl)methyl)ethanesulfonamide (intermediate A-52) and 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-22).</p>	Colorless solid	392.5

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
193	<p>Ethanesulfonic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p><i>N</i>-((5-Bromo-4-chloropyridin-3-yl)methyl)ethanesulfonamide (intermediate A-53) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Colorless solid	394.3
194	<p>Ethanesulfonic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide</p> 	<p><i>N</i>-((5-Bromo-4-methylpyridin-3-yl)methyl)ethanesulfonamide (intermediate A-52) and 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1<i>H</i>-quinolin-2-one (intermediate A-1)</p>	Grey solid	374.4

Example 195

Ethanesulfonic acid methyl-{2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide









To a solution of ethanesulfonic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide (example 151, 0.06 g, 0.154 mmol) in DMF (1.5 mL) cooled at 0 °C with an ice bath was added 60% NaH in mineral oil (0.008 g, 0.2 mmol) and the reaction mixture was stirred for 15 min. Then, MeI (0.026 g, 0.185 mmol) was added and the reaction mixture was stirred at 0 °C for another 30 min. The mixture was quenched with aq. ammonia (1 mL), diluted with brine (5 mL) and then extracted with EtOAc (2 x 20 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.050 g, 80 %) as a light yellow amorphous solid. MS: 404.4 (M+H⁺).


The following compounds listed in Table 8 were prepared in analogy to the procedure described for the preparation of example 195, using the appropriate starting materials.

Table 8

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
196	3-Chloro-pyridine-2-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	3-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 118)	Colorless amorphous solid	421.4
197	N-Methyl-N-{2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-propionamide 	N-{2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-propionamide (example 43)	Colorless amorphous solid	368.3
198	1-Methyl-1H-pyrazole-4-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	1-Methyl-1H-pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 87)	Colorless amorphous solid	390.4

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
199	3-Methyl-pyridine-2-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	3-Methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 115)	Colorless amorphous solid	401.5
200	3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methylamide 	3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 131)	Off-white amorphous solid	439.1
201	3-Chloro-pyridine-2-carboxylic acid methyl-{1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide 	3-Chloro-pyridine-2-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide (example 137)	Colorless amorphous solid	447.4

Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
202	1-Methyl-1 <i>H</i> -pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methylamide 	1-Methyl-1 <i>H</i> -pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 175)	Light brown solid	408.5

Example 203

Ethanesulfonic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide





5

To a solution of ethanesulfonic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 55, 0.681 g, 0.197 mmol) in DMF (1 mL) heated to 65 °C was added *N*-chloro-succinimide (0.034 g, 0.256 mmol) and the reaction mixture was stirred at this temperature over night. The mixture was diluted with EtOAc, poured into water (5 mL) and extracted with EtOAc (2 x 10 mL). The combined organics were washed with brine, dried over Na₂SO₄, filtered and evaporated. The residue was purified by reverse phase HPLC on a Gemini-NX column, eluting with a MeOH-H₂O (0.05 % TEA) gradient to give the title compound (0.025 g, 33 %) as a colorless amorphous solid. MS: 380.4 (M+H⁺).

10


The following compounds listed in Table 9 were prepared in analogy to the procedure described for the preparation of example 203, using the appropriate starting materials.

5 **Table 9**


Ex	Name	Starting Materials	Aspect	MS (M+H ⁺)
204	3-Chloro-pyridine-2-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	3-Chloro-pyridine-2-carboxylic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 140)	Colorless solid	427.4
205	3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide 	3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 141)	Colorless solid	411.4

Example 206

6-{5-[(3-Ethyl-oxetan-3-ylamino)-methyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one




[A] 6-(5-Hydroxymethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one



A microwave vial was charged with (5-bromo-pyridin-3-yl)-methanol (376 mg, 2.0 mmol), 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1) (574 mg, 2.0 mmol) and DMF (4 mL). After purging the reaction mixture with argon, *bis*(triphenylphosphine)palladium(II)chloride (140 mg, 0.2 mmol) and 2 N aq. Na₂CO₃ solution (2.0 mL, 4 mmol) were added and the reaction was heated in the microwave at 120 °C for 50 min. The reaction mixture was diluted with EtOAc, filtered through Dicalite and washed with EtOAc (2 x 20 mL). The resulting filtrate was washed with brine, dried over anhydrous Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (198 mg, 74 %) as a white solid. MS: 269.2 (M+H⁺).


[B] 6-(5-Chloromethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one



At 0 °C, 6-(5-hydroxymethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (190 mg, 0.7 mmol) in DCM (15 mL) was treated slowly with thionyl chloride (0.32 mL, 4.0 mmol). After the addition, the reaction mixture was allowed to stir at 2-5 °C for 2 hours before it was poured into satd. aq. NaHCO₃ solution (50 mL). After extraction with

EtOAc, the organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated *in vacuo* to give title compound (190 mg, 93.5%) as yellow oil. MS: 287.1 ($\text{M}+\text{H}^+$).


5 [C] 6-{5-[(3-Ethyl-oxetan-3-ylamino)-methyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one




To a stirred solution of 6-(5-chloromethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (190 mg, 0.66 mmol) in DMF (5.0 mL) was added K_2CO_3 (276 mg, 2.0 mmol) and 3-ethyl-oxetan-3-ylamine (415 mg, 3.0 mmol) at room temperature and the
 10 resulting reaction mixture was stirred at room temperature for additional 12 hours. After extraction with EtOAc, the organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated *in vacuo* to give a crude product which was purified by prep-HPLC to give title compound (35 mg, 17.2%) as a light yellow solid. MS: 352.1 ($\text{M}+\text{H}^+$).

15 Example 207

Ethanesulfonic acid [5-(7-chloro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide




[A] N-(4-Bromo-3-chloro-phenyl)-3-chloro-propionamide




- To a solution of 4-bromo-3-chloroaniline (6.0 g, 29.1 mmol) and pyridine (3.45 g, 43.6 mmol) in 1,2-dichloroethane (50 mL) was added 3-chloropropionyl chloride (5.53 g, 43.6 mmol) dropwise at 15 °C. After stirring for 2 hours at room temperature, the mixture was washed with water and then hydrochloric acid (2 N, aqueous). The organic layer was dried over anhydrous sodium sulfate. After filtration, the solvent was removed under reduced pressure. The product *N*-(4-bromo-3-chloro-phenyl)-3-chloro-propionamide (8.20 g, yield: 95%) was obtained as an oil. MS: 298.0 (M+H)⁺.

[B] 6-Bromo-7-chloro-3,4-dihydro-1H-quinolin-2-one




- A flame-dried 50-mL flask equipped with a magnetic stirring bar was charged with *N*-(4-bromo-3-chloro-phenyl)-3-chloro-propionamide (1.0 g, 3.36 mmol) and aluminium chloride (0.67 g, 5.04 mmol). In a pre-heated oil bath, the flask was heated at 135~140 °C for 2 hours. After cooling to room temperature, the reaction mixture was treated slowly with ice-water, then extracted with EtOAc. The organic layer was washed with water and brine in sequence and dried over anhydrous Na₂SO₄. After evaporation of the solvent, the residue was purified by recrystallization from EtOAc (2 mL) which gave title compound (0.44 g, 50%) as a solid. MS: 260.0 (M+H)⁺.

[C] 6-Bromo-7-chloro-1-methyl-3,4-dihydro-1H-quinolin-2-one




- Potassium *tert*-butoxide (0.45 g, 4.0 mmol) was added to a solution of 6-bromo-7-chloro-3,4-dihydro-1H-quinolin-2-one (0.52 g, 2.0 mmol) in DMF (5 mL) at 0 °C. Then, the reaction mixture was stirred at 0 °C for 30 min and methyl iodide (0.18 g, 1.29 mmol) was added. The resulting mixture was stirred for 2 hours before water was added. After extraction of the reaction mixture with EtOAc, the organic layer was washed with water and brine in sequence. Then, it was dried over anhydrous Na₂SO₄ and the solvent was removed under reduced pressure to give a crude product (0.49 g, 90%) as a white solid. MS: 274.0 (M+H)⁺.

[D] 7-Chloro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one



A mixture of 6-bromo-7-chloro-1-methyl-3,4-dihydro-1H-quinolin-2-one (0.1 g, 0.36 mmol), *bis*(pinacolato)diboron (0.13 g, 0.55 mmol), 1,1'-*bis*(diphenylphosphino)ferrocenedichloro-palladium (II) (14.7 mg, 0.018 mmol) and potassium acetate (0.11 g, 1.08 mmol) in dioxane (3 mL) was heated in a microwave at 100 °C for 3 hours. The mixture was diluted with EtOAc, washed with water, dried over anhydrous Na₂SO₄, and concentrated *in vacuo*. The residue was purified by flash chromatography (silica gel, 30% ethyl acetate in hexane) to afford the title compound (82 mg, 70%) as a white solid. MS: 322.1 (M+H)⁺.


[E] Ethanesulfonic acid [5-(7-chloro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide




A mixture of 7-chloro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (20 mg, 0.06 mmol), ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11, 26 mg, 0.09 mmol), *bis*(triphenylphosphine)palladium(II) chloride (4.2 mg, 0.006 mmol) and sodium carbonate (2 N aq., 0.06 mL) in DMF (1 mL) was heated in a microwave at 120 °C for 30 minutes. The mixture was diluted with EtOAc, washed with water, dried over anhydrous Na₂SO₄ and concentrated *in vacuo*. The residue was purified by prep-HPLC to afford the title product (7.1 mg, 30%) as a white solid. MS, 394.2 (M+H)⁺.

Example 208

Ethanesulfonic acid [5-(8-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide




[A] 3-Chloro-N-(2-fluoro-phenyl)-propionamide



5

To a solution of 2-fluoroaniline (6.67 g, 60 mmol) and pyridine (5.21 g, 66 mmol) in 1,2-dichloroethane (50 mL) was added 3-chloropropionyl chloride (8.38 g, 66 mmol) dropwise at 15 °C. After stirring at room temperature for 2 hours, the mixture was washed with water and then hydrochloric acid (2 N, aqueous). The organic layer was dried over
10 anhydrous Na₂SO₄, filtered, and concentrated *in vacuo* to afford title compound (10.9 g, yield: 90%) as an oil. MS: 202.1 (M+H)⁺.


[B] 8-Fluoro-3,4-dihydro-1H-quinolin-2-one



A flame-dried 50-mL flask equipped with a magnetic stirring bar was charged with 3-chloro-N-(2-fluoro-phenyl)-propionamide (5.33 g, 26.5 mmol) and aluminium chloride (5.30 g, 39.7 mmol). In a pre-heated oil bath, the flask was heated at 160 °C for 1.5 hours.
15 After cooling to room temperature, the mixture was treated slowly with ice-water and extracted with EtOAc. The organic layer was washed with water and brine in sequence, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. The resulting residue


was then purified by flash chromatography (silica gel, 30% ethyl acetate in hexane) to afford title compound (3.1 g, 70%) as a solid. MS: 166.0 (M+H)⁺.

[C] 6-Bromo-8-fluoro-3,4-dihydro-1H-quinolin-2-one




- 5 At 0 °C, NBS (0.62 g, 3.5mmol) was added dropwise to a solution of 8-fluoro-3,4-dihydro-1H-quinolin-2-one (0.52 g, 3.2 mmol) in DMF (5 mL). The resulting reaction mixture was stirred at room temperature for 12 hours before it was treated with water. The precipitated solid was collected through filtration, washed with ether, and dried in a vacuum to afford title compound (0.65 g, 85%) as a white solid. MS: 244.1 (M+H)⁺.

10 [D] 6-Bromo-8-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one




- A solution of 6-bromo-8-fluoro-3,4-dihydro-1H-quinolin-2-one (0.21 g, 0.86 mmol) in DMF (2 mL) was treated with potassium *tert*-butoxide (0.19 g, 1.72 mmol) at 0 °C. The resulting mixture was stirred at 0 °C for 30 minutes before methyl iodide (0.18 g, 1.29 mmol) was added. After stirring for 2 hours, the reaction mixture was treated with water, extracted with EtOAc, washed with water and brine in sequence, and dried over anhydrous Na₂SO₄. After removal of solvent under reduced pressure, the crude product (0.18 g, 80%) was obtained as a white solid. MS: 258.0 (M+H)⁺.

20 [E] 8-Fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one



A mixture of 6-bromo-8-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (0.18 g, 0.36 mmol), *bis*(pinacolato)diboron (0.25g, 1.05mmol), 1,1'-*bis*(diphenylphosphino)ferrocene-dichloropalladium (II) (28.6 mg, 0.035 mmol) and potassium acetate (0.21 g, 2.1 mmol) in dioxane (3 mL) was heated in a microwave at 100 °C for 3 hours. After dilution with EtOAc, the organic layer was washed with water, dried over anhydrous Na₂SO₄ and concentrated *in vacuo*. The residue was then purified by flash chromatography (silica gel, 30% ethyl acetate in hexane) to afford the title compound (0.15g, 70%) as a white solid. MS: 306.2 (M+H)⁺.


- 10 [F] Ethanesulfonic acid [5-(8-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide




- 15 A mixture of 8-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (20 mg, 0.066 mmol), ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11, 27.4 mg, 0.098 mmol), *bis*(triphenylphosphine)-palladium(II) chloride (4.6 mg, 0.0066 mmol) and aq. sodium carbonate solution (2 N, 0.07 mL) in DMF (1 mL) was heated in a microwave at 120 °C for 30 min. The resulting reaction mixture was then diluted with EtOAc, washed with water, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by prep-HPLC to afford the
20 title compound (12.4 mg, 50%) as a white solid. MS: 378.1 (M+H)⁺.

Example 209

Ethanesulfonic acid [5-(8-chloro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide



[A] *N*-(4-Bromo-2-chloro-phenyl)-3-chloro-propionamide




5

To a solution of 4-bromo-2-chloro-phenylamine (32 g, 0.15 mol) and pyridine (13.45 g, 0.17 mol) in DCM (200 mL) was added 3-chloropropionyl chloride (21.65 g, 0.17 mol) dropwise at 15 °C. After stirring at room temperature for 1 hour, the mixture was washed with water and then hydrochloric acid (2 N, aqueous). The organic layer was dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo* to afford title compound (10.9 g, yield: 90%) as a white solid.

10

[B] 6-Bromo-8-chloro-3,4-dihydro-1*H*-quinolin-2-one




A flame-dried 500-mL flask equipped with a magnetic stirring bar was charged with *N*-(4-bromo-2-chloro-phenyl)-3-chloro-propionamide (29.7 g, 0.1 mol) and aluminium chloride (53.3 g, 0.4 mol). In a pre-heated oil bath, the flask was heated at 140 °C for 1 hour. After cooling to room temperature, the mixture was treated slowly with ice-water and extracted with EtOAc. The organic layer was washed with water and brine in sequence, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. The resulting residue was then purified by flash chromatography (silica gel, 30% ethyl acetate in hexane) to afford the title compound (7.0 g, 27%) as a white solid.

15


20

[C] 6-Bromo-8-chloro-1-methyl-3,4-dihydro-1H-quinolin-2-one




A solution of 6-bromo-8-chloro-3,4-dihydro-1H-quinolin-2-one (7.0 g, 26.9 mmol) in DMF (100 mL) was treated with potassium *tert*-butoxide (6.0 g, 53.8 mmol) at 0 °C
5 portionwise. The resulting mixture was stirred at 0 °C for 30 minutes before methyl iodide (5.0 g, 35.0 mmol) was added. After stirred for 12 hours, the reaction mixture was treated with water, extracted with EtOAc, washed with water and brine in sequence, and dried over anhydrous Na₂SO₄. After removal of solvent under reduced pressure, the crude product (3.3 g, 45%) was obtained as a white solid.

10 [D] 8-Chloro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one



A mixture of 6-bromo-8-chloro-1-methyl-3,4-dihydro-1H-quinolin-2-one (0.23 g, 0.84 mmol), *bis*(pinacolato)diboron (0.255 g, 1.01 mmol), 1,1'-*bis*(diphenylphosphino)-
15 ferrocene-dichloropalladium (II) (30.7 mg, 0.04 mmol) and potassium acetate (0.247 g, 2.52 mmol) in dioxane (5 mL) was heated in a microwave at 80 °C over night. After dilution with EtOAc, the organic layer was washed with water, dried over anhydrous Na₂SO₄ and concentrated *in vacuo*. The residue was then purified by flash chromatography (silica gel, 30% ethyl acetate in hexane) to afford the title compound (0.17g, 63%) as a
20 white solid.


[E] Ethanesulfonic acid [5-(8-chloro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide




Under argon protection, *bis*(triphenylphosphine)palladium(II)chloride (4 mg, 0.05 mmol) followed by 1 N aqueous Na₂CO₃ solution (1 mL), was added to a solution of 8-chloro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (160 mg, 0.5 mmol) and ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11, 158 mg, 0.6 mmol) in DMF (3 mL). The resulting reaction mixture was then heated in a microwave at 110 °C for 45 min. After cooling to room temperature, the reaction mixture was diluted with EtOAc (5 mL) and poured into a satd. aq. solution of NaHCO₃ (10 mL). The aqueous layer was extracted with EtOAc (3 x 5 mL). Combined organics were washed with water and brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo* to give a crude product which was then purified by prep-HPLC to give title compound (65 mg, 45%) as a white solid. MS: 394.1 (M+H)⁺.

Example 210

Ethanesulfonic acid [5-(5-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)-pyridin-3-ylmethyl]-amide




[A] 3-Chloro-N-(3-fluoro-phenyl)-propionamide




To a solution of 3-fluoroaniline (10 mL, 104.02 mmol) in DCM (100 mL) was added pyridine (21 mL, 260.2 mmol) and 3-chloropropionyl chloride (12 mL, 124.4 mmol). The reaction mixture was stirred for 3 hours at room temperature until all starting material had disappeared as shown by LC-MS analysis. The reaction mixture was then diluted with H₂O and extracted with EtOAc. The organic layer was dried over anhydrous Na₂SO₄ and concentrated in *vacuo* to afford the title compound as a solid. It was used in the next step without further purification.

[B] 7-Fluoro-3,4-dihydro-1H-quinolin-2-one (compound A) and 5-fluoro-3,4-dihydro-1H-quinolin-2-one (compound B)



A flame-dried 50-mL flask equipped with a magnetic stirring bar was charged with 3-chloro-*N*-(3-fluoro-phenyl)-propionamide (10 g, 49.6 mmol) and AlCl₃ (23.1 g, 173.6 mmol). On a pre-heated oil bath, the flask was heated at 120~125 °C for 2 hours until LC-MS indicated the reaction was complete. After cooling to room temperature, the mixture was treated slowly with ice-water. After extraction with EtOAc, the combined organic layers were washed with water and brine in sequence. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated in *vacuo* to afford a white solid as a crude mixture of two regioisomeric products (A:B) in a ratio of 5.3 :1. (7.63 g, 93.2%).


[C] 7-Fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one and 5-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one



To an ice cold solution of a mixture of 7-fluoro-3,4-dihydro-1H-quinolin-2-one and 5-fluoro-3,4-dihydro-1H-quinolin-2-one (16.5 g, 0.1 mol) in DMF (200 mL) was added potassium *tert*-butoxide (22.4 g, 0.2 mol) in 2 portions. The reaction mixture was stirred at


- 0 °C for 30 min before MeI (25.4 g, 0.18 mol) was added. After the addition, the reaction mixture was allowed to warm up to RT slowly and stirred at RT over night. It was then diluted with EtOAc (500 mL) and poured into 200 mL of 1 M aq. HCl. After extraction with EtOAc (200 mL, 3x), the combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered and concentrated in vacuo to give the crude title compounds as oil (16.0 g, 89 %). It was used in the next step without further purification.

[D] 6-Bromo-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one and 6-bromo-5-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one



- To an ice cold solution of the mixture of 7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one and 5-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one (16.0 g, 89.4 mmol) in DMF (200 mL) was added NBS (16.0 g, 89.4 mmol). After the addition, the reaction mixture was warmed up to RT and stirred for 3 hours. When LC-MS indicated the completion of the reaction, the mixture was diluted with EtOAc (500 mL) and poured into 500 mL of water.
- The aqueous layer was then extracted with EtOAc (200 mL, 3x). Combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated in vacuo to give the crude title compound as oil (18.0 g, 78 %). It was used in the next step without further purification.

- [E] Ethanesulfonic acid [5-(5-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide




In analogy to the procedure described for the preparation of intermediates A-1 [B], the mixture of 6-bromo-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one and 6-bromo-5-

fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one was reacted with 4,4,4',4',5,5,5',5'-octamethyl-2,2'-*bi*(1,3,2-dioxaborolane) in the presence of potassium acetate and PdCl₂(DPPF)-CH₂Cl₂ and the reaction product was subsequently reacted with ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) in analogy to the procedure described for the preparation of example 45. The crude reaction product has been separated by preparative HPLC to give ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 16) as well as the title compound as a white solid. MS: 378.1 (M+H)⁺.

10 **Example 211**


Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide




To a solution of ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 16, 0.15 g, 0.39 mmol) in DMF (5 mL) was added potassium *tert*-butoxide (67.2 mg, 0.60 mmol) at 0 °C. The resulting mixture was stirred for 30 minutes at 0 °C before methyl iodide (84.6 mg, 0.6 mmol) was added dropwise. After stirring for 2 hours, the mixture was treated with water, and extracted with EtOAc. The combined organic layers were then washed with water and brine in sequence, dried over anhydrous Na₂SO₄ and concentrated *in vacuo*. The residue was then purified by flash chromatography (silica gel, 5% methanol in dichloromethane) to afford the title compound (93 mg, 60%) as a white solid. MS: 392.1(M+H)⁺.

Example 212

N-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide




[A] *N*-(5-Bromo-pyridin-3-ylmethyl)-propionamide



To a solution of propionamide (146 mg, 2.0 mmol) in DMF (10 mL) was added NaH (61
5 mg, 2.4 mmol) at 0 °C and reaction mixture was stirred for 10 min before the addition of 3-
bromo-5-chloromethyl-pyridine (intermediate A-12 [A], 412 mg, 2.0 mmol). The resulting
reaction mixture was stirred at room temperature for 1 hour before water was added. Then,
the reaction mixture was extracted with EtOAc and combined organic layers were dried by
anhydrous Na₂SO₄, filtered and concentrated *in vacuo*. The residue was then purified by
10 flash chromatography to give title compound (400 mg, 82%) as a solid.

[B] *N*-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-
propionamide





To a mixture of 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-
15 dihydro-1*H*-quinolin-2-one (intermediate A-22, 168 g, 0.55 mmol) and *N*-(5-bromo-
pyridin-3-ylmethyl)-propionamide (122 mg, 0.5 mmol) in DMF (3 mL), purged with argon
for 1 min, was added *bis*(triphenylphosphine)palladium (II)chloride (38 mg, 0.054 mmol)
and 1 N aq. Na₂CO₃ (2.5 mL). Then, the resulting reaction mixture was heated in a
microwave at 110 °C for 45 min. After cooling to room temperature, it was diluted with
20 EtOAc (5 mL) and poured into a satd. aq. solution of NaHCO₃ (10 mL). The aqueous layer

was extracted with EtOAc (3 x 5 mL) and combined organics were washed with water and brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo*. The residue was then purified by prep-HPLC to afford the title compound (30 mg, 18%) as a white solid. MS: 342.1(M+H)⁺.

5

Example 213


(rac)-N-{1-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide

10 A] 5-Bromo-N-methoxy-N-methyl-nicotinamide

To a solution of 5-bromo-nicotinic acid (2.02 g, 10 mmol) and *O,N*-dimethylhydroxylamine (670 mg, 11 mmol) in DMF (100 mL) was added HOBt (400 mg, 0.3 mmol), EDCI (2.09 g, 11 mmol) and Et₃N (1.11 g, 11 mmol). The resulting mixture was stirred at room temperature for 1 hour. Water was added to the reaction mixture and it was extracted with EtOAc. The organic layer was washed with brine, dried by anhydrous Na₂SO₄, filtered and concentrated *in vacuo*. The residue was then purified by flash chromatography to give the title compound (2.08 g, 85%) as a solid.

15


[B] 1-(5-Bromo-pyridin-3-yl)-ethanone



20


To a solution of 5-bromo-*N*-methoxy-*N*-methyl-nicotinamide (2.08 g, 8.5 mmol) in THF (20 mL) was added MeMgBr (1.52 g, 12.75 mmol) at -78 °C. After the addition, the reaction mixture was stirred at room temperature for 2 hours before quenching with water. After extraction with EtOAc, the organic layer was washed with brine, dried over
5 anhydrous Na₂SO₄, filtered and concentrated *in vacuo*. The residue was then purified by flash column chromatography to afford the title compound (1.5 g, 88%).

[C] (rac)-1-(5-Bromo-pyridin-3-yl)-ethylamine




To a solution of 1-(5-bromo-pyridin-3-yl)-ethanone (400 mg, 2 mmol) in methanolic
10 ammonia (10 mL) was added Ti(*O-i*Pr)₄ (1.14 g, 4 mmol). The resulting reaction mixture was heated to reflux and stirred over night. After cooling at 0 °C, NaBH₄ was added into the mixture and it was allowed to warm up to room temperature and stirred for 3 hours. Water was added and the reaction mixture was extracted with EtOAc. The organic layer was washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated
15 *in vacuo*. It was directly used in the next step without further purification.

[D] (rac)-*N*-[1-(5-Bromo-pyridin-3-yl)-ethyl]-propionamide



At 0 °C, propionyl chloride (138 mg, 1.5 mmol) was added dropwise to a solution of (rac)-
1-(5-bromo-pyridin-3-yl)-ethylamine (201 mg, 1 mmol) and Et₃N (153 mg, 1.5 mmol) in
20 DCM (10 mL). The reaction mixture was stirred at room temperature for 1 hour before it was concentrated to dryness. The residue was redissolved in EtOAc and washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo*. The residue was purified by flash column chromatography to give the title compound (200 mg, 78%).


25 [E] (rac)-*N*-{1-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide



To a mixture of 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-22, 168 g, 0.55 mmol) and (rac)-*N*-[1-(5-bromo-pyridin-3-yl)-ethyl]-propionamide (125 mg, 0.5 mmol) in DMF (3 mL), purged
 5 with argon for 1 min, was added *bis*(triphenylphosphine)palladium(II)chloride (38 mg, 0.054 mmol) and 1 N aq. Na₂CO₃ (2.5 mL). Then, the resulting reaction mixture was heated in a microwave at 100 °C for 45 min. After cooling to room temperature, it was diluted with EtOAc (5 mL), and poured into satd. aq. solution of NaHCO₃ (10 mL). The aqueous layer was extracted with EtOAc (3 x 5 mL) and combined organics were washed
 10 with water and brine (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated *in vacuo*. The residue was purified by prep-HPLC to give the title compound (35 mg, 20%) as a white solid. MS: 356.3 (M+H)⁺.

Example 214

15 **(S)-3-[5-(2-Oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid *tert*-butyl ester**




In analogy to the procedure described for the preparation of example 45, 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroquinolin-2(1*H*)-one (intermediate A-34)
 20 has been coupled to (S)-3-(5-bromo-pyridin-3-yloxy)-pyrrolidine-1-carboxylic acid *tert*-butyl ester (intermediate A-29) to give the title compound as a dark grey amorphous solid.

MS: 410.6 ($M+H^+$).

Example 215


3-Chloro-pyridine-2-carboxylic acid methyl-[5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide



In analogy to the procedure described for the preparation of example 45, 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,4-dihydroquinolin-2(1*H*)-one (intermediate A-34) has been coupled to *N*-((5-bromopyridin-3-yl)methyl)-3-chloro-*N*-methylpicolinamide (intermediate A-57) to give the title compound as a light brown amorphous solid. MS: 407.5 ($M+H^+$).

Example 216

3-Chloro-pyridine-2-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide




In analogy to the procedure described for the preparation of example 203, 3-chloro-pyridine-2-carboxylic acid methyl-[5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 215) was reacted with *N*-chloro-succinimide to give the title compound as a light brown solid. MS: 441.5 ($M+H^+$).

Example 217

(R)-2-Methyl-propane-2-sulfinic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide

Chiral




- 5 In analogy to the procedure described for the preparation of example 45, 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) has been coupled to (R)-*N*-((5-bromo-4-chloropyridin-3-yl)methyl)-2-methylpropane-2-sulfinamide (intermediate A-58) to give the title compound as an orange solid. MS: 406.4 ($M+H^+$).

10

Example 218


6-(5-Aminomethyl-4-chloro-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride



- 15 In analogy to the procedure described for the preparation of example 42, reaction of (R)-2-methyl-propane-2-sulfinic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 217) with hydrogen chloride (in dioxane) in methanol gave the title compound as a yellow solid. MS: 302.4 ($M+H^+$).

20 **Example 219**


3,5-Dimethyl-isoxazole-4-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide



To a solution of 6-(5-aminomethyl-4-chloro-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 218, 0.05 g, 0.148 mmol) in dry DMF (1 mL) were added EDCI (0.034 g, 0.077 mmol), hydroxybenzotriazole (0.017 g, 0.077 mmol), Hünig's base (0.057 g, 0.443 mmol) and 3,5-dimethyl-isoxazole-4-carboxylic acid (0.021 g, 0.148 mmol) and the resulting solution was stirred at room temperature for 2h. The reaction mixture was diluted with EtOAc, poured into sat. NaHCO₃ solution (10 mL) and extracted with EtOAc (2 x 20 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was purified by silica gel flash chromatography eluting with a 0 to 5% MeOH-DCM gradient to give the title compound (0.03 g, 48 %) as a colorless solid. MS: 425.4 (M+H⁺).

Example 220


3-Chloro-pyridine-2-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide



In analogy to the procedure described for the preparation of example 75, 6-(5-aminomethyl-4-chloro-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 218) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a off-white solid. MS: 441.3 (M+H⁺).

Example 221

(R)-2-Methyl-propane-2-sulfinic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide




- 5 In analogy to the procedure described for the preparation of example 45, 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) has been coupled to (R)-2-methyl-propane-2-sulfinic acid (5-bromo-4-methyl-pyridin-3-ylmethyl)-amide (intermediate A-59) to give the title compound as a colorless amorphous solid. MS: 386.5 (M+H⁺).

10

Example 222


6-(5-Aminomethyl-4-methyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride



- 15 In analogy to the procedure described for the preparation of example 42, reaction of (R)-2-methyl-propane-2-sulfinic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 221) with hydrogen chloride (in dioxane) in methanol gave the title compound as a yellow solid. MS: 282.5 (M+H⁺).

20 **Example 223**


3,5-Dimethyl-isoxazole-4-carboxylic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide



In analogy to the procedure described for the preparation of example 219, 6-(5-aminomethyl-4-methyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 222) has been coupled with 3,5-dimethyl-isoxazole-4-carboxylic acid to give the title compound as a off-white solid. MS: 405.5 (M+H⁺).

Example 224


- 10 **3-Chloro-pyridine-2-carboxylic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide**



- In analogy to the procedure described for the preparation of example 75, 6-(5-aminomethyl-4-methyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 222) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a colorless solid. MS: 421.5 (M+H⁺).
- 15

Example 225


(R)-2-Methyl-propane-2-sulfinic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide



In analogy to the procedure described for the preparation of example 45, 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-22) has been coupled to (R)-2-methyl-propane-2-sulfinic acid (5-bromo-4-methyl-pyridin-3-ylmethyl)-amide (intermediate A-59) to give the title compound as a brown amorphous solid. MS: 404.5 ($M+H^+$).

10 **Example 226**


6-(5-Aminomethyl-4-methyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride



In analogy to the procedure described for the preparation of example 42, reaction of (R)-2-methyl-propane-2-sulfinic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide (example 225) with hydrogen chloride (in dioxane) in methanol gave the title compound as a light brown solid. MS: 300.5 ($M+H^+$).

Example 227


3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide



In analogy to the procedure described for the preparation of example 75, 6-(5-aminomethyl-4-methyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 226) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a colorless solid. MS: 439.4 ($M+H^+$).

Example 228


3-Chloro-pyridine-2-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide



In analogy to the procedure described for the preparation of example 195, 3-chloro-pyridine-2-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 220) has been reacted with sodium hydride and methyl iodide to give the title compound as off-white solid. MS: 455.1 ($M+H^+$).

Example 229


3-Chloro-pyridine-2-carboxylic acid methyl-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide



In analogy to the procedure described for the preparation of example 195, 3-chloro-pyridine-2-carboxylic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide (example 224) has been reacted with sodium hydride and
 5 methyl iodide to give the title compound as light brown amorphous solid. MS: 435.5 ($M+H^+$).

Example 230

**3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-
 10 tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide**




(R or S)

In analogy to the procedures described for the preparation of example 74, 6-[5-((R or S)-
 1-amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride
 15 (example 65) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a colorless solid. MS: 421.4 ($M+H^+$).

Example 231


**5'-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-5,6-dihydro-4*H*-
 20 [3,3']bipyridinyl-1-carboxylic acid *tert*-butyl ester**



- In analogy to the procedure described for the preparation of example 45, 6-(5-bromopyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one (prepared from 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-1) and 3,5-dibromo pyridine also with a procedure as used for the preparation of example 45) has been coupled to 5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-2H-pyridine-1-carboxylic acid *tert*-butyl ester to give the title compound as a amorphous colorless solid. MS: 420.5 (M+H⁺).

10 **Example 232**


{2-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-carbamic acid *tert*-butyl ester



- In analogy to the procedure described for the preparation of example 45, 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1H-quinolin-2-one (intermediate A-22) has been coupled to [2-(5-bromo-pyridin-3-yloxy)-ethyl]-carbamic acid *tert*-butyl ester (intermediate A-7) to give the title compound as a yellow waxy solid. MS: 416.4 (M+H⁺).

20 **Example 233**


3-Chloro-pyridine-2-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide



In analogy to the procedure described for the preparation of example 75, 6-[5-(2-amino-ethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 42) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a light yellow viscous oil. MS: 437.4 ($M+H^+$).

Example 234


- 10 **Ethanesulfonic acid [5-(4-oxo-1,2,5,6-tetrahydro-4*H*-pyrrolo[3,2,1-*ij*]quinolin-8-yl)-pyridin-3-ylmethyl]-amide**



In analogy to the procedure described for the preparation of example 45, 8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydro-1*H*-pyrrolo[3,2,1-*ij*]quinolin-4(2*H*)-one has been coupled to ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) to give the title compound as a colorless amorphous solid. MS: 372.2 ($M+H^+$).

Example 235


- 20 **Ethanesulfonic acid [5-(3-oxo-2,3,6,7-tetrahydro-1*H*,5*H*-pyrido[3,2,1-*ij*]quinolin-9-yl)-pyridin-3-ylmethyl]-amide**



- In analogy to the procedure described for the preparation of example 45, 9-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,6,7-tetrahydropyrido[3,2,1-ij]quinolin-3(5*H*)-one (prepared from 9-bromo-1,2,6,7-tetrahydropyrido[3,2,1-ij]quinolin-3(5*H*)-one and
- 5 4,4,4',4',5,5,5',5'-octamethyl-2,2'-*bi*(1,3,2-dioxaborolane) in analogy to the procedure described for the preparation of intermediate A-1 [B]) has been coupled to ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11) to give the title compound as a off-white amorphous solid. MS: 386.2 ($M+H^+$).

10 **Example 236**


3-Chloro-pyridine-2-carboxylic acid [5-(4-oxo-1,2,5,6-tetrahydro-4*H*-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide



- In analogy to the procedure described for the preparation of example 45, 8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydro-1*H*-pyrrolo[3,2,1-ij]quinolin-4(2*H*)-one
- 15 has been coupled to *N*-((5-bromopyridin-3-yl)methyl)-3-chloropicolinamide (intermediate A-56) to give the title compound as off-white solid. MS: 419.3 ($M+H^+$).

Example 237


- 20 **3-Chloro-pyridine-2-carboxylic acid methyl-[5-(4-oxo-1,2,5,6-tetrahydro-4*H*-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide**



In analogy to the procedure described for the preparation of example 195, N-methylation of 3-chloro-pyridine-2-carboxylic acid [5-(4-oxo-1,2,5,6-tetrahydro-4*H*-pyrrolo[3,2,1-*ij*]quinolin-8-yl)-pyridin-3-ylmethyl]-amide (example 236) gave the title compound as a
 5 light yellow solid. MS: 433.4 ($M+H^+$).

Example 238

6-[5-(2-Amino-ethoxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride




10

In analogy to the procedure described for the preparation of example 42, reaction of {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-carbamic acid *tert*-butyl ester (example 232) with hydrogen chloride (in dioxane) in methanol gave the title compound as light yellow solid. MS: 316.5 ($M+H^+$).

Example 239

(*R*)-2-Methyl-propane-2-sulfinic acid {(*R* or *S*)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide

15




(R or S)

In analogy to the procedure described for the preparation of example 45, reaction of 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) with (R)-2-methyl-propane-2-sulfinic acid [(R or S)-1-(5-bromo-4-methyl-pyridin-3-yl)-ethyl]-amide (intermediate A-60) gave the title compound as a dark brown amorphous solid. MS: 400.5 ($M+H^+$).

Example 240

6-[5-((R or S)-1-Amino-ethyl)-4-methyl-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride




(R or S)

In analogy to the procedure described for the preparation of example 42, reaction of (R)-2-methyl-propane-2-sulfinic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide (example 239) with hydrogen chloride (in dioxane) in methanol gave the title compound as light brown solid. MS: 296.5 ($M+H^+$).

Example 241

3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide




(R or S)

In analogy to the procedure described for the preparation of example 75, 6-[5-((R or S)-1-amino-ethyl)-4-methyl-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 240) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a colorless solid. MS: 435.6 (M+H⁺).

Example 242

3,5-Dimethyl-isoxazole-4-carboxylic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide




(R or S)

In analogy to the procedure described for the preparation of example 219, 6-[5-((R or S)-1-amino-ethyl)-4-methyl-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 240) has been coupled with 3,5-dimethyl-isoxazole-4-carboxylic acid to give the title compound as a colorless amorphous solid. MS: 419.6 (M+H⁺).

Example 243


Ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide



In analogy to the procedure described for the preparation of example 38, 6-[5-(2-amino-ethoxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 238) and ethanesulfonyl chloride gave the title compound as a orange amorphous solid. MS: 408.4 (M+H⁺).

Example 244


10 **3-Chloro-pyridine-2-carboxylic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide**



In analogy to the procedure described for the preparation of example 75, 6-[5-(2-amino-ethoxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 238) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a off-white solid. MS: 455.3 (M+H⁺).

Example 245


20 ***N*-{(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-methanesulfonamide**



In analogy to the procedure described for the preparation of example 38, 6-[5-((*trans*)-4-amino-cyclohexyloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 71) and methanesulfonyl chloride gave the title compound as a
5 light yellow amorphous solid. MS: 430.4 (M+H⁺).

Example 246

3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide




10

In analogy to the procedure described for the preparation of example 219, 6-(5-aminomethyl-4-methyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 226) has been coupled with 3,5-dimethyl-isoxazole-4-carboxylic acid to give the title compound as a colorless solid. MS: 423.6 (M+H⁺).

15

Example 247


5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',6'-dihydro-2'*H*-[3,4']bipyridinyl-1'-carboxylic acid *tert*-butyl ester



- In analogy to the procedure described for the preparation of example 45, 6-(5-bromopyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one (prepared from 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) and 3,5-dibromo pyridine also with a procedure as used for the preparation of example 45) has been coupled to 4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,6-dihydro-2*H*-pyridine-1-carboxylic acid *tert*-butyl ester to give the title compound as light yellow amorphous solid. MS: 420.5 ($M+H^+$).

10 **Example 248**


Ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-methyl-amide



- In analogy to the procedure described for the preparation of example 195, ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide (example 243) has been reacted with sodium hydride and methyl iodide to give the title compound as orange amorphous solid. MS: 422.4 ($M+H^+$).

Example 249

- 20 **(R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide**




(R or S)

In analogy to the procedure described for the preparation of example 45, reaction of 1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (intermediate A-1) with (R)-2-methyl-propane-2-sulfinic acid [(R or S)-1-(5-bromo-4-chloro-pyridin-3-yl)-ethyl]-amide (intermediate A-61) gave the title compound as a colorless amorphous solid. MS: 420.4 ($M+H^+$).

Example 250

6-[5-((R or S)-1-Amino-ethyl)-4-chloro-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride




(R or S)

In analogy to the procedure described for the preparation of example 42, reaction of (R)-2-methyl-propane-2-sulfinic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide (example 249) with hydrogen chloride (in dioxane) in methanol gave the title compound as yellow solid. MS: 316.4 ($M+H^+$).

Example 251

3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide




(R or S)

In analogy to the procedure described for the preparation of example 75, 6-[5-((R or S)-1-amino-ethyl)-4-chloro-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one
 5 hydrochloride (example 250) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a colorless solid. MS: 455.5 ($M+H^+$).

Example 252

10 **(R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-amide**




(R or S)

In analogy to the procedure described for the preparation of example 45, reaction of 7-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-
 15 quinolin-2-one (intermediate A-22) with (R)-2-methyl-propane-2-sulfinic acid [(R or S)-1-(5-bromo-4-methyl-pyridin-3-yl)-ethyl]-amide (intermediate A-60) gave the title compound as a yellow amorphous solid. MS: 418.5 ($M+H^+$).

Example 253

6-[5-((R or S)-1-Amino-ethyl)-4-methyl-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride




(R or S)

- 5 In analogy to the procedure described for the preparation of example 42, reaction of (R)-2-methyl-propane-2-sulfonic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-amide (example 252) with hydrogen chloride (in dioxane) in methanol gave the title compound as light brown solid. MS: 314.5 (M+H⁺).


10

Example 254

6-[5-((S)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one



- 15 In analogy to the procedure described in example 76, 1-methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 60) has been reacted with acetyl chloride to give the title compound as a colorless solid. MS: 366.4 (M+H⁺).


Example 255**1-Methyl-6-{5-[(S)-1-(1-methyl-1H-pyrazole-4-carbonyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,4-dihydro-1H-quinolin-2-one**

5

To a solution of 1-methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (0.04 g, 0.111 mmol, example 60), 1-methyl-1H-pyrazole-4-carboxylic acid (0.021 g, 0.167 mmol) and Hunig's base (0.036 g, 0.278 mmol) in EtOAc (2 mL) was added propylphosphonic acid anhydride (cyclic trimer) solution (50% in EtOAc, 0.177 mL, 0.25 mmol) dropwise and the reaction mixture was stirred at room temperature over night. The mixture was diluted with EtOAc, poured into a sat. NaHCO₃ solution (5 mL) and extracted with EtOAc (2 x 20 mL). Combined organics were dried over Na₂SO₄, filtered and evaporated. The residue was purified by silica gel flash chromatography eluting with a 0 to 10% MeOH-DCM gradient to give the title compound (0.036 g, 75.1%) as a colorless amorphous solid. MS: 432.4 (M+H⁺).

15


Example 256**(S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid ethyl ester**



In analogy to the procedure described in example 76, 1-methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 60) has been reacted with chloroformic acid ethyl ester to give the title compound as a light yellow
 5 amorphous solid. MS: 396.5 (M+H⁺).

Example 257

3,5-Dimethyl-isoxazole-4-carboxylic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide



(R or S)


10

In analogy to the procedure described for the preparation of example 219, 6-[5-((R or S)-1-amino-ethyl)-4-chloro-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride (example 250) has been coupled with 3,5-dimethyl-isoxazole-4-carboxylic acid to give the title compound as a off-white amorphous solid. MS: 439.4 (M+H⁺).

15

Example 258

3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-amide




(R or S)

- In analogy to the procedure described for the preparation of example 75, 6-[5-((R or S)-1-amino-ethyl)-4-methyl-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 253) has been coupled with 3-chloro-pyridine-2-carboxylic acid to give the title compound as a off-white solid. MS: 453.4 (M+H⁺).

Example 259


- 10 **5-Methyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide**



- In analogy to the procedure described for the preparation of example 219, 6-(5-aminomethyl-4-methyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 226) has been coupled with 5-methyl-isoxazole-4-carboxylic acid to give the title compound as a off-white solid. MS: 409.6 (M+H⁺).

Example 260


- 20 **5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',4',5',6'-tetrahydro-2'*H*-[3,4']bipyridinyl-1'-carboxylic acid *tert*-butyl ester**



- 5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',6'-dihydro-2'*H*-[3,4']bipyridinyl-1'-carboxylic acid *tert*-butyl ester (example 247, 0.128 g, 0.305 mmol) in MeOH (50 mL) was run through the H-cube for 2 h (flow rate: 0.5 mL/min; temperature: 60 °C; catalyst: 10 % Pd/C (THS 01111); full hydrogen mode). The resulting solution was evaporated to dryness to give the title compound (0.106 g, 74 %) as a yellow amorphous solid. MS: 422.5 ($M+H^+$).

Example 261


- 10 **N-{(R or S)-1-[4-Chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide**



(R or S)


- In analogy to the procedure described in example 76, 6-[5-((R or S)-1-amino-ethyl)-4-chloro-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 250) has been reacted with propionyl chloride to give the title compound as a colorless solid. MS: 372.5 ($M+H^+$).

Example 262

6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-1*H*-quinolin-2-one

To solution of 1-(5-(1-methyl-2-oxo-1,2,3,4-tetrahydroquinolin-6-yl)pyridin-3-yl)cyclopropanecarboxylic acid (example 59 [C], 0.3 g, 0.931 mmol) in toluene (15 mL) were added TEA (0.141 g, 1.4 mmol) and diphenylphosphoryl azide (0.384 g, 1.4 mmol) and then the reaction mixture was heated to reflux for 4 h. The mixture was concentrated in vacuo, the residue was dissolved in THF (30 mL) and the solution cooled to 0 °C. Then, a solution of KOtBu (0.627 g, 5.58 mmol) in THF (13 mL) was added and the mixture was stirred for 1 h at 0 °C and then over night at room temperature. The mixture was concentrated in vacuo, the residue was preadsorbed on silica gel and purified by silica gel flash chromatography eluting with a 0 to 10% MeOH-DCM gradient to give a yellow amorphous solid (0.104 g). This material was then further purified by reverse phase HPLC on a Gemini-NX column, eluting with a MeOH-H₂O (0.05 % TEA) gradient to give the title compound (0.02 g, 7 %) as a colorless amorphous solid. MS: 292.3 (M+H⁺).

15

Example 263**Ethanesulfonic acid [5-(8-fluoro-1-methyl-2-oxo-1,2-dihydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide**


A mixture of 8-fluoro-1-methyl-6-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,4-dihydro-1*H*-quinolin-2-one (example 208 [E], 20 mg, 0.066 mmol), ethanesulfonic acid (5-bromo-pyridin-3-ylmethyl)-amide (intermediate A-11, 27.4 mg, 0.098 mmol), *bis*(triphenylphosphine)-palladium(II) chloride (4.6 mg, 0.0066 mmol), and aq. sodium

20

carbonate solution (2 N, 0.07 mL) in DMF (1 mL) was heated in a microwave at 120 °C for 30 minutes. After dilution of the reaction mixture with EtOAc, the organic layer was washed with water, dried over anhydrous Na₂SO₄ and concentrated *in vacuo*. The residue was then purified by prep-HPLC to afford the title compound (2.5 mg, 10%) as a white solid. MS: 376.3 (M+H)⁺.

Example 264


N-{(R or S)-1-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-propionamide



In analogy to the procedure described in example 76, 6-[5-((R or S)-1-amino-ethyl)-4-methyl-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 253) has been reacted with propionyl chloride to give the title compound as a colorless amorphous solid. MS: 370.6 (M+H⁺).

Example 265

N-[4-Chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide




In analogy to the procedure described in example 76, 6-(5-aminomethyl-4-chloro-pyridin-3-yl)-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 218) has been reacted with propionyl chloride to give the title compound as a off-white solid. MS: 358.4 (M+H⁺).

5

Example 266


N-{(R or S)-1-[4-Methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide



- 10 In analogy to the procedure described in example 76, 6-[5-((R or S)-1-amino-ethyl)-4-methyl-pyridin-3-yl]-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 240) has been reacted with propionyl chloride to give the title compound as a colorless amorphous solid. MS: 352.5 (M+H⁺).

15 **Example 267**

3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide



In analogy to the procedure described for the preparation of example 219, 6-[5-(2-aminoethoxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1*H*-quinolin-2-one hydrochloride (example 238) has been coupled with 3,5-dimethyl-isoxazole-4-carboxylic acid to give the title compound as a colorless solid. MS: 439.4 (M+H⁺).

Example A

A compound of formula (I) can be used in a manner known per se as the active ingredient for the production of tablets of the following composition:

	<u>Per tablet</u>
5 Active ingredient	200 mg
Microcrystalline cellulose	155 mg
Corn starch	25 mg
Talc	25 mg
Hydroxypropylmethylcellulose	<u>20 mg</u>
10	425 mg


Example B

A compound of formula (I) can be used in a manner known per se as the active ingredient for the production of capsules of the following composition:

	<u>Per capsule</u>
15 Active ingredient	100.0 mg
Corn starch	20.0 mg
Lactose	95.0 mg
Talc	4.5 mg
20 Magnesium stearate	<u>0.5 mg</u>
	220.0 mg

CLAIMS

1. Compounds of formula (I)



wherein

5 R^1 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^2 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^3 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^4 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

10 or R^3 and R^4 together with the carbon atoms to which they are attached form a double bond;

R^5 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^6 is H or R^8 , wherein in case R^6 is H then at least one of A^1 and A^3 is N;

R^7 is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

15 or R^1 and R^7 together with the nitrogen and carbon atoms to which they are attached form a heterocycloalkyl substituted with one to three substituents independently selected from H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^8 is $-O_m-(CR^9R^{10})_n-(CR^{11}R^{12})_p-(CR^{13}R^{14})_q-NR^{15}R^{16}$,
 $-N_mR^{17}-(CR^9R^{10})_n-(CR^{11}R^{12})_p-(CR^{13}R^{14})_q-NR^{15}R^{16}$ or
 $-S_m(O)_r-(CR^9R^{10})_n-(CR^{11}R^{12})_p-(CR^{13}R^{14})_q-NR^{15}R^{16}$;

20 R^9 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^{10} is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^{11} is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R¹⁰ and R¹¹ together with the carbon atoms to which they are attached form a double bond;

R¹² is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

5 or R¹¹ and R¹² together with the carbon atom to which they are attached form a cycloalkyl or a heterocycloalkyl;

R¹³ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹⁴ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R¹⁰ and R¹⁴ together form -(CH₂)_t;

10 R¹⁵ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkoxyalkyl, haloalkoxyalkyl or hydroxyalkyl;

R¹⁶ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, phenylalkyl, substituted phenylalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, -S(O)₂R¹⁸, -S(O)R¹⁸, -S(O)₂OR¹⁸, -S(O)₂NR¹⁸R¹⁹, -C(O)R¹⁸, -C(O)OR¹⁸ or -C(O)NR¹⁸R¹⁹, wherein substituted phenylalkyl, substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl, wherein in case R¹⁵ is H or alkyl and R¹⁶ is H or alkyl, then the sum of n, p and q is at least 1;

15
20
25
30

or R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R²³, R²⁴ and R²⁵;

5 or R¹³ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

or R¹¹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

10 or R⁹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

R¹⁷ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

15 R¹⁸ is alkyl, cycloalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, halohydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkylcarbonyloxyalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, wherein substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl;

20

25

30

R¹⁹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

A¹ is CR²⁰ or N;

A^2 is CR^{21} or N;

A^3 is CR^{22} or N;

R^{20} is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^{21} is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

5 R^{22} is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R^{23} , R^{24} and R^{25} are each independently selected from H, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, $-S(O)_2R^{18}$, $-C(O)R^{18}$, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein

10 substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl;

15 wherein in case m is zero and R^9 and R^{15} , or R^{11} and R^{15} or R^{13} and R^{15} or R^{15} and R^{16} together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, then at least one of R^{23} , R^{24} , R^{25} and R^{16} is different from H, alkyl, haloalkyl, alkoxy, haloalkoxy, halogen, hydroxy, cyano, amino, substituted amino, aminocarbonyl, substituted aminocarbonyl, wherein substituted amino and substituted aminocarbonyl are

20 substituted on the nitrogen atom with one to two alkyl, and $-C(O)R^{18}$, $-S(O)_2R^{18}$, wherein R^{18} is alkyl, haloalkyl, hydroxyalkyl, halohydroxyalkyl, alkoxyalkyl or haloalkoxyalkyl;

25

m is zero or 1, wherein in case m is 1, then the sum of n and p is 2, 3 or 4;

n is zero, 1 or 2;

30 p is zero, 1 or 2;

q is zero, 1 or 2;

r is zero, 1 or 2, wherein in case m is zero then r is zero;

t is zero, 1 or 2;

or pharmaceutically acceptable salts or esters.

2. A compound according to claim 1, wherein

5 R¹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R² is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R³ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R⁴ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R⁵ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

10 R⁶ is H or R⁸, wherein in case R⁶ is H then at least one of A¹ and A³ is N;

R⁷ is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R⁸ is -O_m-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q-NR¹⁵R¹⁶,
-N_mR¹⁷-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q-NR¹⁵R¹⁶ or
-S_m(O)_r-(CR⁹R¹⁰)_n-(CR¹¹R¹²)_p-(CR¹³R¹⁴)_q-NR¹⁵R¹⁶;

15 R⁹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹⁰ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹¹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹² is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R¹¹ and R¹² together with the carbon atom to which they are attached form a
20 cycloalkyl;

R¹³ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R¹⁴ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

or R¹⁰ and R¹⁴ together form -(CH₂)_t-;

R¹⁵ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkoxyalkyl, haloalkoxyalkyl or hydroxyalkyl;

R¹⁶ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, phenylalkyl, substituted phenylalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, -S(O)₂R¹⁸, -S(O)₂OR¹⁸, -S(O)₂NR¹⁸R¹⁹, -C(O)R¹⁸, -C(O)OR¹⁸ or -C(O)NR¹⁸R¹⁹, wherein substituted phenylalkyl, substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl, wherein in case R¹⁵ is H or alkyl and R¹⁶ is H or alkyl, then the sum of n, p and q is at least 1;

or R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R²³, R²⁴ and R²⁵;

or R¹³ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

or R¹¹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

or R⁹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵;

R¹⁷ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

5 R¹⁸ is alkyl, cycloalkyl, haloalkyl, halocycloalkyl, hydroxyalkyl, haloalkoxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkylcarbonyloxyalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, wherein substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, 10 haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein 15 substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl;

R¹⁹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

20 A¹ is CR²⁰ or N;

A² is CR²¹ or N;

A³ is CR²² or N;

R²⁰ is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R²¹ is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

25 R²² is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl;

R²³, R²⁴ and R²⁵ are each independently selected from H, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, 30 haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, -S(O)₂R¹⁸, -C(O)R¹⁸, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino,

5 substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl; wherein in case m is zero, then at least one of R^{23} , R^{24} and R^{25} is different from H, alkyl, haloalkyl, alkoxy, haloalkoxy, halogen, hydroxy, cyano, amino, substituted amino, aminocarbonyl or substituted aminocarbonyl, wherein substituted amino and substituted aminocarbonyl are substituted on the nitrogen atom with one to two alkyl;

m is zero or 1, wherein in case m is 1, then the sum of n and p is 2, 3 or 4;

n is zero, 1 or 2;

15 p is zero, 1 or 2;

q is zero, 1 or 2;

r is zero, 1 or 2, wherein in case m is zero then r is zero;

t is zero, 1 or 2;

or pharmaceutically acceptable salts or esters.

20 3. A compound according to claim 1 or 2, wherein R^1 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

4. A compound according to any one of claims 1 to 3, wherein R^1 is H or alkyl.

5. A compound according to any one of claims 1 to 4, wherein R^1 is alkyl.

6. A compound according to any one of claims 1 to 5, wherein R^1 is methyl.

25 7. A compound according to any one of claims 1 to 6, wherein R^2 is H.

8. A compound according to any one of claims 1 to 7, wherein R^3 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.

9. A compound according to any one of claims 1 to 8, wherein R^3 is H.

10. A compound according to any one of claims 1 to 9, wherein R^4 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.
11. A compound according to any one of claims 1 to 10, wherein R^4 is H or alkyl.
12. A compound according to any one of claims 1 to 11, wherein R^4 is H.
- 5 13. A compound according to any one of claims 1 to 12, wherein R^5 is H or alkyl.
14. A compound according to any one of claims 1 to 13, wherein R^5 is H.
15. A compound according to any one of claims 1 to 14, wherein R^6 is R^8 .
16. A compound according to any one of claims 1 to 12, wherein R^7 is H, halogen, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.
- 10 17. A compound according to any one of claims 1 to 15, wherein R^7 is H.
18. A compound according to any one of claims 1 and 7 to 15, wherein R^1 and R^7 together with the nitrogen and carbon atoms to which they are attached form a heterocycloalkyl substituted with one to three substituents independently selected from H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.
- 15 19. A compound according to any one of claims 1 to 18, wherein R^8 is $-O_m-(CR^9R^{10})_n-(CR^{11}R^{12})_p-(CR^{13}R^{14})_q-NR^{15}R^{16}$.
20. A compound according to any one of claims 1 to 19, wherein R^9 is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.
21. A compound according to any one of claims 1 to 20, wherein R^9 is H, alkyl or
20 cycloalkyl.
22. A compound according to any one of claims 1 to 21, wherein R^9 is H or alkyl.
23. A compound according to any one of claims 1 to 22, wherein R^9 is H.
24. A compound according to any one of claims 1 to 23, wherein R^{10} is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.
- 25 25. A compound according to any one of claims 1 to 24, wherein R^{10} is H or alkyl.
26. A compound according to any one of claims 1 to 25, wherein R^{10} is H.

27. A compound according to any one of claims 1 to 26, wherein R¹¹ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.
28. A compound according to any one of claims 1 to 27, wherein R¹¹ is H.
29. A compound according to any one of claims 1 to 23, wherein R¹⁰ and R¹¹ together
5 with the carbon atoms to which they are attached form a double bond.
30. A compound according to any one of claims 1 to 29, wherein R¹² is H.
31. A compound according to any one of claims 1 to 26, wherein R¹¹ and R¹² together with the carbon atom to which they are attached form a cycloalkyl.
32. A compound according to any one of claims 1 to 31, wherein R¹³ is H.
- 10 33. A compound according to any one of claims 1 to 32, wherein R¹⁴ is H, alkyl, haloalkyl, cycloalkyl or halocycloalkyl.
34. A compound according to any one of claims 1 to 33, wherein R¹⁴ is H.
35. A compound according to any one of claims 1 to 23, 27, 28 and 30 to 32, wherein R¹⁰ and R¹⁴ together form -(CH₂)_t-.
- 15 36. A compound according to any one of claims 1 to 35, wherein R¹⁵ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkoxyalkyl, haloalkoxyalkyl or hydroxyalkyl.
37. A compound according to any one of claims 1 to 36, wherein R¹⁵ is H, alkyl or alkoxyalkyl.
38. A compound according to any one of claims 1 to 37, wherein R¹⁵ is H or alkyl.
- 20 39. A compound according to any one of claims 1 to 38, wherein R¹⁵ is H.
40. A compound according to any one of claims 1 to 19 and 25 to 35, wherein R⁹ and R¹⁵ together with the nitrogen and carbon atoms to which they are attached form a substituted heterocycloalkyl, wherein substituted heterocycloalkyl is substituted with R²³, R²⁴ and R²⁵.
- 25 41. A compound according to any one of claims 1 to 40, wherein R¹⁶ is H, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, phenylalkyl, substituted phenylalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, -S(O)₂R¹⁸, -

- $S(O)_2OR^{18}$, $-S(O)_2NR^{18}R^{19}$, $-C(O)R^{18}$, $-C(O)OR^{18}$ or $-C(O)NR^{18}R^{19}$, wherein substituted phenylalkyl, substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkylcycloalkyl, halocycloalkyl, alkylcycloalkylalkyl, halocycloalkylalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, haloalkoxyalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxycarbonyl, halogen, hydroxy, oxo, cyano, triazolylalkyl, tetrazolylalkyl, amino, substituted amino, aminoalkyl, substituted aminoalkyl, aminocarbonyl, substituted aminocarbonyl, alkylcarbonylamino, cycloalkylcarbonylamino, alkylcarbonylaminoalkyl and cycloalkylcarbonylaminoalkyl, wherein substituted amino, substituted aminoalkyl and substituted aminocarbonyl are substituted on the nitrogen atom with one to two substituents independently selected from alkyl, cycloalkyl, alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkylalkyl, hydroxyalkyl and alkoxyalkyl, wherein in case R^{15} is H or alkyl and R^{16} is H or alkyl, then the sum of n, p and q is at least 1.
42. A compound according to any one of claims 1 to 41, wherein R^{16} is H, hydroxyalkyl, phenylalkyl, heterocycloalkyl substituted with alkyl, $-S(O)R^{18}$, $-S(O)_2R^{18}$, $-C(O)R^{18}$ or $-C(O)OR^{18}$, wherein in case R^{15} is H or alkyl and R^{16} is H or alkyl, then the sum of n, p and q is at least 1.
43. A compound according to any one of claims 1 to 42, wherein R^{16} is H, hydroxyalkyl, phenylalkyl, $-S(O)_2R^{18}$, $-C(O)R^{18}$ or $-C(O)OR^{18}$, wherein in case R^{15} is H or alkyl and R^{16} is H or alkyl, then the sum of n, p and q is at least 1.
44. A compound according to any one of claims 1 to 43, wherein R^{16} is H, hydroxyalkyl or phenylalkyl, wherein in case R^{15} is H or alkyl and R^{16} is H or alkyl, then the sum of n, p and q is at least 1.
45. A compound according to any one of claims 1 to 44, wherein R^{16} is hydroxyalkyl, $-S(O)_2R^{18}$, $-C(O)R^{18}$ or $-C(O)OR^{18}$.
46. A compound according to any one of claims 1 to 45, wherein R^{16} is H, hydroxyethyl, hydroxymethylbutyl or benzyl, wherein in case R^{15} is H or alkyl and R^{16} is H, then the sum of n, p and q is at least 1.
47. A compound according to any one of claims 1 to 43, wherein R^{16} is hydroxyalkyl or $-S(O)_2R^{18}$.

48. A compound according to any one of claims 1 to 44 and 45, wherein R^{16} is hydroxyalkyl.
49. A compound according to any one of claims 1 to 44, wherein R^{16} is hydroxyethyl, hydroxymethylbutyl or benzyl.
- 5 50. A compound according to any one of claims 1 to 43, 45 and 47, wherein R^{16} is $-S(O)_2R^{18}$.
51. A compound according to any one of claims 1 to 43 and 45, wherein R^{16} is $-C(O)R^{18}$.
- 10 52. A compound according to any one of claims 1 to 35, wherein R^{15} and R^{16} together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl or a substituted heteroaryl, wherein substituted heterocycloalkyl and substituted heteroaryl are substituted with R^{23} , R^{24} and R^{25} .
- 15 53. A compound according to any one of claims 1 to 35 and 52, wherein R^{15} and R^{16} together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl wherein substituted heterocycloalkyl is substituted with one to three substituents independently selected from hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxycarbonyl, oxo, triazolylalkyl and substituted aminoalkyl, wherein substituted aminoalkyl is substituted on the nitrogen atom with one to two alkyl.
- 20 54. A compound according to any one of claims 1 to 35, 53 and 53, wherein R^{15} and R^{16} together with the nitrogen atom to which they are attached form a substituted heterocycloalkyl wherein substituted heterocycloalkyl is substituted with one to three substituents independently selected from hydroxyalkyl and oxo.
- 25 55. A compound according to any one of claims 1 to 35 and 53 to 54, wherein the heterocycloalkyl formed by R^{15} and R^{16} together with the nitrogen atom to which they are attached is selected from pyrrolidiny, piperidiny, thiomorpholinyl, thiazinanyl, isothiazolidinyl and 2,6-diaza-spiro[3.3]heptanyl.
- 30 56. A compound according to any one of claims 1 to 35 and 53 to 55, wherein the heterocycloalkyl formed by R^{15} and R^{16} together with the nitrogen atom to which they are attached is selected from pyrrolidiny, piperidiny, thiomorpholinyl, thiazinanyl and 2,6-diaza-spiro[3.3]heptanyl.

57. A compound according to any one of claims 1 to 35 and 53 to 56, wherein R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form 2-hydroxymethyl-pyrrolidin-1-yl, 2-hydroxymethyl-5-oxo-pyrrolidin-1-yl or isothiazolidinyl substituted on the sulfur atom by two oxo.
- 5 58. A compound according to any one of claims 1 to 35 and 53 to 57, wherein R¹⁵ and R¹⁶ together with the nitrogen atom to which they are attached form 2-hydroxymethyl-pyrrolidin-1-yl or 2-hydroxymethyl-5-oxo-pyrrolidin-1-yl.
59. A compound according to any one of claims 1 to 43, 50 and 51, wherein R¹⁸ is alkyl, cycloalkyl, haloalkyl, hydroxyalkyl, alkylcarbonyloxyalkyl, aryl, substituted aryl, heterocycloalkyl, substituted heterocycloalkyl, heteroaryl, substituted heteroaryl, wherein substituted aryl, substituted heterocycloalkyl and substituted heteroaryl are substituted with one to three substituents independently selected from alkyl, hydroxyalkyl, haloalkyl, cycloalkyl and halogen.
- 10
60. A compound according to any one of claims 1 to 53, 50, 51 and 59, wherein R¹⁸ is alkyl, cycloalkyl, hydroxyalkyl, alkylcarbonyloxyalkyl or heterocycloalkyl substituted with one to three alkoxy.
- 15
61. A compound according to any one of claims 1 to 53, 50, 51, 59 and 60, wherein R¹⁸ is alkyl, cycloalkyl, hydroxyalkyl, alkylcarbonyloxyalkyl or heteroaryl substituted with one to three alkoxy.
- 20
62. A compound according to any one of claims 1 to 53, 50, 51, 59 to 61, wherein R¹⁸ is alkyl or heteroaryl substituted with one to three substituents independently selected from alkyl and halogen.
63. A compound according to any one of claims 1 to 53, 50, 51, 59 to 62, wherein R¹⁸ is heteroaryl substituted with one to three substituents independently selected from alkyl and halogen.
- 25
64. A compound according to any one of claims 1 to 53, 50, 51, 59 to 63, wherein R¹⁸ is alkyl.
65. A compound according to any one of claims 1 to 53, 50, 51, 59 to 64, wherein R¹⁸ is ethyl.
- 30
66. A compound according to any one of claims 1 to 65, wherein A¹ is CR²⁰.

67. A compound according to any one of claims 1 to 66, wherein A^2 is CR^{21} .
68. A compound according to any one of claims 1 to 67, wherein A^3 is CR^{22} .
69. A compound according to any one of claims 1 to 68, wherein R^{20} is H or halogen.
70. A compound according to any one of claims 1 to 69, wherein R^{20} is H.
- 5 71. A compound according to any one of claims 1 to 70, wherein R^{21} is H, alkyl or halogen.
72. A compound according to any one of claims 1 to 71, wherein R^{21} is H or alkyl.
73. A compound according to any one of claims 1 to 72, wherein R^{21} is H.
74. A compound according to any one of claims 1 to 73, wherein R^{22} is H or halogen.
- 10 75. A compound according to any one of claims 1 to 74, wherein R^{22} is H.
76. A compound according to any one of claims 1 to 75, wherein R^{23} , R^{24} and R^{25} are each independently selected from hydrogen, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxyalkyl, alkoxycarbonyl, oxo, triazolylalkyl and substituted aminoalkyl, wherein substituted aminoalkyl is substituted on the nitrogen atom with one to two
15 alkyl.
77. A compound according to any one of claims 1 to 76, wherein R^{23} , R^{24} and R^{25} are each independently selected from hydrogen, hydroxyalkyl, alkoxyalkyl, haloalkoxyalkyl, alkoxycarbonyl, oxo, triazolylalkyl and substituted aminoalkyl, wherein substituted aminoalkyl is substituted on the nitrogen atom with one to two
20 alkyl.
78. A compound according to any one of claims 1 to 77, wherein R^{23} , R^{24} and R^{25} are each independently selected from hydrogen, hydroxyalkyl and oxo.
79. A compound according to any one of claims 1 to 78, wherein at least one of R^{23} , R^{24} and R^{25} is different from hydrogen.
- 25 80. A compound according to any one of claims 1 to 65, 67, 68 and 71 to 79, wherein A^1 is N.
81. A compound according to any one of claims 1 to 66, 68 to 70 and 74 to 80, wherein A^2 is N.

82. A compound according to any one of claims 1 to 67, 69 to 73 and 76 to, wherein A^3 is N.
83. A compound according to any one of claims 1 to 82, wherein m is zero or 1, wherein in case m is 1, then the sum of n, p and q is 2, 3, 4, 5 or 6.
- 5 84. A compound according to any one of claims 1 to 83, wherein m is 1.
85. A compound according to any one of claims 1 to 82, wherein m is zero.
86. A compound according to any one of claims 1 to 85, wherein n is zero or 1.
87. A compound according to any one of claims 1 to 86, wherein n is zero.
88. A compound according to any one of claims 1 to 87, wherein p is zero or 1.
- 10 89. A compound according to any one of claims 1 to 88, wherein q is zero or 1.
90. A compound according to any one of claims 1 to 89, wherein q is zero.
91. A compound according to any one of claims 1 to 90, r is zero and m is zero.
92. A compound according to any one of claims 1 to 91, wherein t is 2.
93. A compound according to any one of claims 1 to 91, wherein t is zero.
- 15 94. A compound according to any one of claims 1 to 14, 16 to 18, 66 to 82 and 92 to 93, wherein R^6 is H and at least one of A^1 and A^3 is N.
95. A compound according to any one of claims 1 to 94, selected from
- Ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-amide;
- 20 Acetic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylcarbamoyl]-methyl ester;
- 2-Hydroxy-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-acetamide;
- 6-[5-(2-Hydroxy-ethylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 25

- 6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((R)-2-Hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 5 6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 10 6-[5-((S)-2-Ethylaminomethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-2-Methoxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-((S)-2-[1,2,4]triazol-1-ylmethyl-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 15 6-(5-Benzylamino-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-(2-oxo-pyrrolidin-1-yl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 20 Ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(1,4,4-trimethyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 1-Methyl-6-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-(2-oxo-piperidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

- 6-[5-(1,1-Dioxo-1 λ^6 -thiomorpholin-4-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S)-2-Methoxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 5 (S)-1-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-pyrrolidine-2-carboxylic acid methyl ester;
- 1-Methyl-6-{5-[(S)-2-(2,2,2-trifluoro-ethoxymethyl)-pyrrolidin-1-ylmethyl]-pyridin-3-yl}-3,4-dihydro-1H-quinolin-2-one;
- 10 6-[6-(2-Hydroxy-ethylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-(6-Benzylamino-pyrazin-2-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 7-Fluoro-6-[6-(2-hydroxy-ethylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-(6-Benzylamino-pyrazin-2-yl)-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 15 6-[6-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyrazin-2-yl]-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid tert-butyl ester ;
- 6-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-2,6-diaza-spiro[3.3]heptane-2-carboxylic acid tert-butyl ester ;
- 20 6-[5-((R)-1-Hydroxymethyl-2-methyl-propylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[6-((R)-1-Hydroxymethyl-2-methyl-propylamino)-pyrazin-2-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- Ethanesulfonic acid [5-(5-methyl-6-oxo-5,6,7,8-tetrahydro-[1,5]naphthyridin-2-yl)-pyridin-3-ylmethyl]-amide;
- 25 6-Pyridin-3-yl-3,4-dihydro-1H-[1,5]naphthyridin-2-one;
- 1-Methyl-6-pyridin-3-yl-3,4-dihydro-1H-[1,5]naphthyridin-2-one;
- 1-Methyl-6-pyridin-3-yl-3,4-dihydro-1H-[1,7]naphthyridin-2-one;

- 6-(5-Aminomethyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- N-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;
- 5 Propane-2-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- {2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-carbamic acid tert-butyl ester;
- 3-Methoxy-isoxazole-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 Cyclopropanecarboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 6-[5-(2-Amino-ethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 15 N-{2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-propionamide;
- 6-[5-(1,1-Dioxo-1λ⁶-[1,2]thiazinan-2-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- and pharmaceutically acceptable salts thereof.
96. A compound according to any one of claims 1 to 94, selected from
- 20 (S)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-azetidine-1-carboxylic acid tert-butyl ester;
- (R)-3-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- 25 3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-azetidine-1-carboxylic acid tert-butyl ester;
- (S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester;

- (S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;
- 4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-piperidine-1-carboxylic acid tert-butyl ester;
- 5 (S)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxymethyl]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- (S)-3-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- 10 (R)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- (R)- 2-Methyl-propane-2-sulfinic acid {3-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-oxetan-3-yl}-amide;
- Ethanesulfonic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 (R)-2-Methyl-propane-2-sulfinic acid {(S or R)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- (R) 2-Methyl-propane-2-sulfinic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 20 {(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-carbamic acid tert-butyl ester;
- 6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 25 1-Methyl-6-[5-((S)-piperidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-[5-(piperidin-4-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;

- 6-[5-((S)-1-Azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((S or R)-1-Amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 5 6-[5-((R or S)-1-Amino-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 1-Methyl-6-[5-((S)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 10 7-Fluoro-1-methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 1-Methyl-6-[5-((R)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-[5-(3-Amino-oxetan-3-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 15 6-[5-(Azetidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-[5-((trans)-4-Amino-cyclohexyloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 6-[5-(1-Amino-1-methyl-ethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 20 6-(5-Aminomethyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one ;
- 6-{6-[6-(3-Methoxy-isoxazole-5-carbonyl)-2,6-diaza-spiro[3.3]hept-2-yl]-pyrazin-2-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 6-[5-((S)-1-Cyclopropanecarbonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

- 3-Methyl-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-butyramide;
- 3,3,3-Trifluoro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;
- 5 2-Hydroxy-2-methyl-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;
- 5-Methyl-[1,3,4]oxadiazole-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 1-Methyl-6-[5-((S)-1-propionyl-piperidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 2-Methoxy-pyrimidine-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide,
- 1-Methyl-1H-imidazole-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 5-Trifluoromethyl-furan-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Pyrimidine-5-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 1-Methyl-1H-pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-6-[5-(1-propionyl-piperidin-4-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 25 Pyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-6-[5-((S)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

- 6-[5-((S)-1-Cyclopropanecarbonyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 3-Methyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5 4-Fluoro-2,6-dimethyl-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-benzamide;
- 1-Methyl-6-[5-((S)-1-propionyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 10 3,6-Dichloro-pyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Cyclopropyl-1-(2,2,2-trifluoro-ethyl)-1H-pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 1,3-Dimethyl-1H-pyrazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Pyrimidine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 6-Methoxy-pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Methyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;
- 25 1-Methyl-1H-pyrazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;
- 6-Chloro-pyridazine-3-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

- 3-Chloro-6-methyl-pyridazine-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-6-[5-((S)-1-propionyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 5 5-Cyclopropyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 2-Methyl-5-trifluoromethyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 2-Methyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Cyclopropyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 2,5-Dimethyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 5-Methyl-oxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 20 6-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,6-Dichloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 6-Methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

- 3-Fluoro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Chloro-3-methyl-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5 1-Methyl-6-[5-((R)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 7-Fluoro-1-methyl-6-[5-((S)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 10 5-Trifluoromethyl-pyrimidine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Methyl-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5-Chloro-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 5-Trifluoromethyl-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-6-[5-((R)-1-propionyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 20 1-Methyl-6-[5-(1-propionyl-azetidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- N-{3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-oxetan-3-yl}-propionamide;
- 25 3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,6-Dichloro-pyridazine-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

- 7-Fluoro-1-methyl-6-[5-((R)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;
- 5-Chloro-3-methoxy-pyrazine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5 3,5-Dimethyl-isoxazole-4-carboxylic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 3-Methyl-5-trifluoromethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 3-Chloro-pyridine-2-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 3-Chloro-pyridine-2-carboxylic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 15 3-Chloro-pyridine-2-carboxylic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 3-Chloro-pyridine-2-carboxylic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 3-Methyl-pyridine-2-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- N-[(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl]-propionamide;
- 25 5-Trifluoromethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 7-Fluoro-1-methyl-6-[5-((S)-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

- (R)-2-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy-methyl]-pyrrolidine-1-carboxylic acid tert-butyl ester;
- 1-Methyl-6-[5-((R)-1-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 5 6-(5-Aminomethyl-pyridin-3-yl)-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one ;
- Ethanesulfonic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide ;
- 10 3-Chloro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-benzenesulfonamide;
- 6-Methoxy-pyridine-3-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 3,5-Dimethyl-isoxazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Cyclopropanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3,4-Dichloro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-benzenesulfonamide;
- 20 1-Methyl-1H-imidazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 6-Chloro-pyridine-3-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 1-Methyl-1H-pyrazole-4-sulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 25 6-[5-((S)-1-Ethanesulfonyl-piperidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

- 6-[5-(1-Ethanesulfonyl-piperidin-4-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 2,2,2-Trifluoro-ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5 C,C,C-Trifluoro-N-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methanesulfonamide;
- 6-[5-((S)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 10 6-[5-((S)-1-Ethanesulfonyl-azetidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- Ethanesulfonic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- 6-[5-((S)-1-Ethanesulfonyl-pyrrolidin-2-ylmethoxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 15 6-[5-((R)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 6-[5-((R)-1-Ethanesulfonyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 20 6-[5-(1-Ethanesulfonyl-azetidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- Ethanesulfonic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;
- Ethanesulfonic acid {(trans)-4-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-amide;
- 25 Ethanesulfonic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- Ethanesulfonic acid {(S or R)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

- 1-Methyl-1H-pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- (rac)-Ethanesulfonic acid {2-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-propyl}-amide;
- 5 (rac)-Ethanesulfonic acid {cyclopropyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-methyl}-amide;
- (rac)-Ethanesulfonic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 10 6-[5-(1,1-Dioxo-1λ6-isothiazolidin-2-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- (rac)-Ethanesulfonic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-propyl}-amide;
- Ethanesulfonic acid ethyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 Ethanesulfonic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- Ethanesulfonic acid isopropyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 20 Ethanesulfonic acid (2-ethoxy-ethyl)-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- (rac)-Ethanesulfonic acid methyl-{1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 25 (rac)-Ethanesulfonic acid ethyl-{1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

6-{5-[2-(1,1-Dioxo-1λ6-isothiazolidin-2-yl)-ethoxy]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one;

Ethanesulfonic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

5 Ethanesulfonic acid {(S or R)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;

10 Ethanesulfonic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid methyl-{2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

15 3-Chloro-pyridine-2-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

N-Methyl-N-{2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-propionamide;

20 1-Methyl-1H-pyrazole-4-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

3-Methyl-pyridine-2-carboxylic acid methyl-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide;

25 3-Chloro-pyridine-2-carboxylic acid methyl-{1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;

1-Methyl-1H-pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide;

- Ethanesulfonic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 5 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 6-{5-[(3-Ethyl-oxetan-3-ylamino)-methyl]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 10 Ethanesulfonic acid [5-(7-chloro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(8-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(8-chloro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 Ethanesulfonic acid [5-(5-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide;
- 20 N-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;
- (rac)-N-{1-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide;
- (S)-3-[5-(2-Oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid tert-butyl ester ;
- 25 3-Chloro-pyridine-2-carboxylic acid methyl-[5-(2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide ;
- 3-Chloro-pyridine-2-carboxylic acid [5-(8-chloro-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide ;

- (R)-2-Methyl-propane-2-sulfinic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide ;
- 6-(5-Aminomethyl-4-chloro-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 5 3,5-Dimethyl-isoxazole-4-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 10 (R)-2-Methyl-propane-2-sulfinic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide ;
- 6-(5-Aminomethyl-4-methyl-pyridin-3-yl)-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 15 3-Chloro-pyridine-2-carboxylic acid [4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- (R)-2-Methyl-propane-2-sulfinic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide ;
- 20 6-(5-Aminomethyl-4-methyl-pyridin-3-yl)-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid [4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-methyl-amide;
- 25 3-Chloro-pyridine-2-carboxylic acid methyl-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;
- 3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

5'-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-5,6-dihydro-4H-[3,3']bipyridinyl-1-carboxylic acid tert-butyl ester;

{2-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-carbamic acid tert-butyl ester;

5 3-Chloro-pyridine-2-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

Ethanesulfonic acid [5-(4-oxo-1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide;

10 Ethanesulfonic acid [5-(3-oxo-2,3,6,7-tetrahydro-1H,5H-pyrido[3,2,1-ij]quinolin-9-yl)-pyridin-3-ylmethyl]-amide;

3-Chloro-pyridine-2-carboxylic acid [5-(4-oxo-1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide;

3-Chloro-pyridine-2-carboxylic acid methyl-[5-(4-oxo-1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide ;

15 6-[5-(2-Amino-ethoxy)-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;

(R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

20 6-[5-((R or S)-1-Amino-ethyl)-4-methyl-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;

3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

3,5-Dimethyl-isoxazole-4-carboxylic acid {(R or S)-1-[4-methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

25 Ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

3-Chloro-pyridine-2-carboxylic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

- N- {(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-methanesulfonamide;
- 3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;
- 5 5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',6'-dihydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid tert-butyl ester;
- Ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-methyl-amide;
- 10 (R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 6-[5-((R or S)-1-Amino-ethyl)-4-chloro-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 15 (R)-2-Methyl-propane-2-sulfinic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-amide;
- 6-[5-((R or S)-1-Amino-ethyl)-4-methyl-pyridin-3-yl]-7-fluoro-1-methyl-3,4-dihydro-1H-quinolin-2-one hydrochloride;
- 20 6-[5-((S)-1-Acetyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;
- 1-Methyl-6-{5-[(S)-1-(1-methyl-1H-pyrazole-4-carbonyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,4-dihydro-1H-quinolin-2-one;
- (S)-3-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-pyrrolidine-1-carboxylic acid ethyl ester;
- 25 3,5-Dimethyl-isoxazole-4-carboxylic acid {(R or S)-1-[4-chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;
- 3-Chloro-pyridine-2-carboxylic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-amide;

5-Methyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;

5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',4',5',6'-tetrahydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid tert-butyl ester ;

5 N-{(R or S)-1-[4-Chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide ;

6-[5-(1-Amino-cyclopropyl)-pyridin-3-yl]-1-methyl-1H-quinolin-2-one;

Ethanesulfonic acid [5-(8-fluoro-1-methyl-2-oxo-1,2-dihydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

10 N-{(R or S)-1-[5-(7-Fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-yl]-ethyl}-propionamide;

N-[4-Chloro-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-propionamide;

15 N-{(R or S)-1-[4-Methyl-5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-propionamide;

3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

and pharmaceutically acceptable salts thereof.

97. A compound according to any one of claims 1 to 94 and 95, selected from

20 6-[5-(2-Hydroxy-ethylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-yl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

25 6-[5-((S)-2-Hydroxymethyl-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

6-[5-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

Ethanesulfonic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

Ethanesulfonic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

5 1-Methyl-6-[5-(2-oxo-pyrrolidin-1-ylmethyl)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

6-[5-((R)-1-Hydroxymethyl-2-methyl-propylamino)-pyridin-3-yl]-1-methyl-3,4-dihydro-1H-quinolin-2-one;

and pharmaceutically acceptable salts thereof.

10 98. A compound according to any one of claims 1 to 94 and 96, selected from

3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

1-Methyl-6-[5-((S)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

15 5-Methyl-isoxazole-4-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

20 1-Methyl-1H-pyrazole-4-carboxylic acid {2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

3-Chloro-pyridine-2-carboxylic acid [5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

3-Chloro-pyridine-2-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

25 7-Fluoro-1-methyl-6-[5-((R)-1-propionyl-pyrrolidin-3-yloxy)-pyridin-3-yl]-3,4-dihydro-1H-quinolin-2-one;

3-Chloro-pyridine-2-carboxylic acid {1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;

3-Chloro-pyridine-2-carboxylic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

3-Chloro-pyridine-2-carboxylic acid {1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-cyclopropyl}-amide;

5 Ethanesulfonic acid {(trans)-4-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-amide;

1-Methyl-1H-pyrazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-ylmethyl]-amide;

10 Ethanesulfonic acid {1-methyl-1-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

6-{5-[2-(1,1-Dioxo-1λ6-isothiazolidin-2-yl)-ethoxy]-pyridin-3-yl}-1-methyl-3,4-dihydro-1H-quinolin-2-one;

Ethanesulfonic acid {(R or S)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

15 Ethanesulfonic acid {(S or R)-1-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yl]-ethyl}-amide;

Ethanesulfonic acid methyl-{2-[5-(1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

20 Ethanesulfonic acid [5-(4-oxo-1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolin-8-yl)-pyridin-3-ylmethyl]-amide;

N-{(trans)-4-[5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-cyclohexyl}-methanesulfonamide;

3,5-Dimethyl-isoxazole-4-carboxylic acid [5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-4-methyl-pyridin-3-ylmethyl]-amide;

25 5-(1-Methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-3',6'-dihydro-2'H-[3,4']bipyridinyl-1'-carboxylic acid tert-butyl ester;


Ethanesulfonic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-methyl-amide;

3,5-Dimethyl-isoxazole-4-carboxylic acid {2-[5-(7-fluoro-1-methyl-2-oxo-1,2,3,4-tetrahydro-quinolin-6-yl)-pyridin-3-yloxy]-ethyl}-amide;

1-Methyl-6-{5-[(S)-1-(1-methyl-1H-pyrazole-4-carbonyl)-pyrrolidin-3-yloxy]-pyridin-3-yl}-3,4-dihydro-1H-quinolin-2-one;

5 and pharmaceutically acceptable salts thereof.

99. A process to prepare a compound according to any one of claims 1 to 98 comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III);



10 wherein A^1 , A^2 , A^3 , R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined in claim 1 and wherein X is halogen or triflate, R^{101} and R^{102} are alkyl, cycloalkyl or together with the boron atom they are attached to form together a borolanyl.

100. A compound according to any one of claims 1 to 98 for use as therapeutically active substance.

15 101. A pharmaceutical composition comprising a compound according to any one of claims 1 to 98 and a therapeutically inert carrier.

102 The use of a compound according to any one of claims 1 to 98 for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrome.

20 103. A compound according to any one of claims 1 to 98 for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrome.

104. The use of a compound according to any one of claims 1 to 98 for the preparation of a medicament for the treatment or prophylaxis of chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom.
105. A method for the treatment or prophylaxis chronic kidney disease, congestive heart failure, hypertension, primary aldosteronism and Cushing syndrom, which method comprises administering an effective amount of a compound according to any one of claims 1 to 98.
106. A compound according to any one of claims 1 to 98, when manufactured according to a process of claim 99.
107. The invention as hereinbefore described.