

Pharmaceutical preparation containing an antivirally active dihydroquinazoline derivative

The present invention relates to a pharmaceutical composition, containing {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof.

The invention further relates to methods for the production and use of the composition in methods of treatment and/or prevention of virus infections, as well as its use for the production of medicinal drugs for use in treating and/or preventing virus infections, in particular for use in treating infections with the human cytomegalovirus (HCMV) or another representative of the Herpes viridae group.

{8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid is known, for example, from WO 2004/096778, full disclosure of which is included herein by reference; it was developed by Applicant as a promising candidate for an antivirally active substance, in particular for combating infections caused by the human cytomegalovirus. However, in the course of development it was discovered that problems occurred with the solubility of the substance, and in particular it proved complicated to produce stable formulations for intravenous administration or solid easily soluble compositions for producing solutions used for intravenous administration.

Summary

It is thus an object of the present disclosure to describe a pharmaceutical composition that is used in particular for intravenous administration, that contains {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid, that has long-term stability and can be stored, and that in addition has a substantially physiological pH.

A further object of the present disclosure is to describe a pharmaceutical composition with which it is possible, in a simple and reliable manner, to produce pharmaceutical compositions for intravenous administration which contain

{8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid and which also remain stable for an adequate period of time, e.g. more than 24 hours.

- 5 In a first aspect, there is provided a pharmaceutical composition, particularly for intravenous administration, comprising the following, namely: a) {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof, b) at least one excipient selected from the cyclodextrins, lysine and arginine, and c) water.

10

In a second aspect, there is provided a solid pharmaceutical composition produced by lyophilizing a pharmaceutical preparation as disclosed herein.

- 15 In a third aspect, there is provided a method for producing a pharmaceutical composition as disclosed herein using the following steps, namely: A) Dissolving the at least one excipient in the water, B) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(tri-fluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof to the solution obtained in step A). C) If necessary, add at least one sugar and/or at least one buffer, D) Adjust the pH to the desired value in order
20 to obtain a pharmaceutical composition, and E) Sterile-filter the solution obtained in step D) and fill into suitable containers. F) If necessary, perform a final sterilization of the solution obtained in step E) under heat.

- 25 In a fourth aspect, there is provided a method for producing a pharmaceutical composition as disclosed herein using the following steps, namely: I.) Dissolving the at least one excipient in a part of the water, II) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(tri-fluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof to the solution obtained in step I.), III.) If necessary, adjusting the pH of the solution obtained in step II.) to the desired value to
30 obtain a first solution. IV.) Dissolving at least one sugar and/or a buffer in a part of the water, V.) If necessary, adjust the pH of the solution obtained in step IV.) to the desired value to obtain a second solution. VI.) Mixing the first and second solution to obtain a

pharmaceutical composition. and VII.) Sterile filtration of the solution obtained in step VI.) and filling into suitable containers. VIII.) If necessary, perform a final sterilization of the solution obtained in step VII.) under heat.

5 In a fifth aspect, there is provided a method for producing a pharmaceutical composition as disclosed herein using the following steps, namely: a.) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(tri-fluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt to an aqueous NaOH solution, preferably an aqueous 0.1M NaOH solution to produce a solution or
10 suspension, b.) Adding water to the solution or suspension obtained in step a.), c.) Adding cyclodextrin and NaCl to the solution or suspension obtained in step b.). d.) Sterile filtration of the solution obtained in step c.) and fill into suitable containers. e.) If necessary, perform a final sterilization of the solution obtained in step d.) under heat.

15 In a sixth aspect, there is provided a method for producing a solid pharmaceutical composition as disclosed above comprising the production of a pharmaceutical composition corresponding to a method as disclosed above followed by lyophilization of the pharmaceutical composition obtained in order to obtain a solid pharmaceutical composition.

20

In a seventh aspect. there is provided the pharmaceutical composition as disclosed herein for use in a method of treatment and/or prophylaxis of diseases, in particular of virus infections, preferably human cytomegalovirus (HCMV) infections or infections with another member of the herpes viridae group.

25

In an eighth aspect, there is provided the use of a pharmaceutical composition as disclosed herein for producing a medicament for the treatment and/or prophylaxis of diseases, in particular of virus infections, preferably human cytomegalovirus (HCMV) infections or infections with another member of the herpes viridae group.

30 **Detailed Description**

Within the scope of the invention the term "stability" is understood to mean not only the chemical stability of the constituents of the pharmaceutical composition, but also the

stability of the solution itself. In particular, the composition according to the invention must be stable against precipitation of the constituents.

In this context, the term "stability" means that at 2 °C to 8 °C, or at 25 °C or at 40 °C the
5 pharmaceutical compositions according to the invention contain a minimum proportion of
>90% and preferably >95% {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-
methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid for a storage
period of at least two, preferably at least three, and most preferred at least six weeks, when
said liquid pharmaceutical compositions are measured using one of the HPLC methods 1-
10 3. Said stability of the liquid pharmaceutical compositions is regarded as adequate within
the scope of the invention.

Furthermore, the term "stability" means that, after they have been diluted or reconstituted
to a final concentration of 0.8-10 mg/ml for infusion at 2 °C to 8 °C, the compositions
15 according to the invention contain a minimum proportion of 90%, preferably at least 95%,
of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-
(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid for a period of at least
four hours, preferably at least six hours, and most preferred at least 24 hours in storage
when, after dilution or reconstitution, said liquid pharmaceutical compositions are
20 measured using one of the HPLC methods 1-3. Said stability of the pharmaceutical
compositions after dilution or reconstitution is regarded as adequate within the scope of the
invention.

It has surprisingly been discovered that pharmaceutical compositions, in particular those
25 used for intravenous administration, that contain

{8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid as well as water can be stabilized by adding at least one excipient selected from cyclodextrins, lysine and arginine. It has further been discovered that such compositions can be lyophilized in order to obtain a stable, solid pharmaceutical composition that can be reconstituted in a simple manner for injection purposes, e.g. by adding water, as a result of which, in turn, a stable pharmaceutical composition, e.g. for intravenous administration, can be obtained.

The subject matter of the invention are thus pharmaceutical compositions, in particular for intravenous administration, that possess the following constituents, namely:

- a) {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof,
- b) at least one excipient selected from the cyclodextrins, lysine and arginine, and
- c) water.

In addition, subject matter of the invention are pharmaceutical compositions which are produced by lyophilization of the above-mentioned pharmaceutical composition.

Within the scope of the invention, the term "salts" is understood to mean physiologically acceptable salts of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid. Physiologically acceptable salts of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid comprise acid addition salts of mineral acids, carbonic acids and sulfonic acids, for example of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, toluenesulfonic acid, benzenesulfonic acid, naphthalenedisulfonic acid, acetic acid, trifluoroacetic acid, propionic acid, lactic acid, tartaric acid, malic acid, citric acid, fumaric acid, maleic acid and benzoic acid.

Physiologically acceptable salts of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid also comprise salts of usual bases, such as for example and preferably alkali metal salts (e.g. sodium and potassium salts), alkaline earth metal salts (e.g. calcium and magnesium salts), ammonium salts derived from ammonia or organic amines having 1 to 16 C-atoms, such as for example and preferably monoethylamine, diethylamine, triethylamine, ethyldiisopropylamine, monoethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, dimethylaminoethanol, 2-amino-2-methyl-1,3-propanediol, procaine, dibenzylamine, N-methylmorpholine, ethylene diamine and N-methylpiperidine as well as salts of alkaline amino acids.

Within the scope of the invention the term „solvates“ refers to those forms of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid which form a complex through coordination with solvent molecules. Hydrates are a special form of solvates in which the coordination takes place with water.

As is readily apparent to a person skilled in the art, {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid possesses a stereocentre at the carbon in the 4-position in the dihydroquinazoline ring. Within the scope of the present invention, it is particularly preferable if this carbon possesses the S-configuration.

A cyclodextrin according to the invention is understood to be any modified or non-modified cyclodextrin. In this case, because of the size of the cavity in the ring, preference is given to β -cyclodextrins and in particular to modified β -cyclodextrins such as, for example, hydroxyalkyl- β -cyclodextrins, e.g. hydroxymethyl- β -cyclodextrin, hydroxyethyl- β -cyclodextrin or hydroxypropyl- β -cyclodextrin, alkyl-hydroxyalkyl- β -cyclodextrins, e.g. methyl-hydroxypropyl- β -cyclodextrins or ethyl-hydroxypropyl-cyclodextrins or sulfoalkyl-cyclodextrins.

Within the scope of the invention, the water used for preparation purposes is normally water that is used for injections.

5 Within the scope of the present invention the expression "possess" or "possessing" denotes a non-exhaustive enumeration and, along with the explicitly mentioned components or steps, does not exclude any other components or steps.

10 Within the meaning of the present invention, the expression "consist of" or "consisting of" denotes an exhaustive enumeration and, apart from the explicitly mentioned components or steps, excludes any other components or steps.

15 Within the scope of the present invention, the expression "consist substantially of" or "consisting substantially of" denotes a partially exhaustive enumeration and denotes methods or compositions which, besides the mentioned components and steps, possess only such other components and steps that do not materially modify the character of the composition or of the method according to the invention, or which are present in quantities that do not materially modify the character of the composition or of the method according to the invention.

20 If, within the scope of the present invention, a composition or a method is described using the expression "possess" or "possessing", this explicitly includes compositions or methods that consist of the components or steps mentioned or that substantially consist of the components or steps mentioned.

25 Within the framework of the invention it is preferred if the pharmaceutical composition according to invention further possesses at least one buffer that is preferably selected from the phosphate buffers, the Tris buffers and the citrate buffers.

30 By adding the buffer it can, in particular, be ensured that the composition always possesses a physiological pH. The buffers named are preferred, in particular due to the fact that they are well tolerated.

It is further preferable, within the scope of the invention, if the pharmaceutical composition according to the invention further possesses at least one sugar, preferably selected from the group consisting of glucose, sucrose, lactose, maltose, trehalose, sorbitol and mannitol.

- 5 It has been found that the pharmaceutical composition according to the invention can again be significantly stabilized by adding a sugar, and in particular one of the sugars explicitly mentioned above. Furthermore, it has been found that the addition of a sugar can facilitate the production of a solid composition by lyophilization, as well as renewed reconstitution of such a solid composition in order to produce a pharmaceutical composition in particular
10 for intravenous administration. Also, the addition of the at least one sugar serves to adjust the osmolality of the solution and to suppress any hemolysis that might occur.

- Within the scope of the invention it is furthermore preferred that {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-
15 dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof is present in the pharmaceutical composition in an amount corresponding to 1 to 100 mg, preferably 2 to 50 mg, more preferably 2 to 25 mg, and in particular to 5 to 20 mg of pure active compound per ml of composition.

- 20 For the stability of the solution and also in the interest of simple storage it has proven advantageous if a quantity of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid is present in the above-mentioned ranges.

- 25 It is further preferred, within the scope of the invention, if the composition possesses a pH in the range of 7.5 to 8.5.

- The aforementioned pH range has proved advantageous because it is a pH in the range of a physiological pH. It has furthermore been found that the solubility of the pharmaceutical
30 composition according to the invention is again significantly better in the slightly alkaline range, i.e. in a range greater than 7.0, than at a pH value of 7.0 or less.

It is further preferred, within the scope of the invention, that the at least one excipient is present in the pharmaceutical composition in an amount of 1 to 5 equivalents, preferably of 2 to 5 equivalents and more preferably of 2.5 to 4.5 equivalents in relation to the content of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-

(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid.

It is further preferred within the scope of the invention, that said composition, in relation to the content of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid, possesses 1 to 10 equivalents, preferably 2 to 7 equivalents, and in particular 2.5 to 5 equivalents of cyclodextrin as well as 0 to 2.0 equivalents, preferably 0.5 to 1.5 equivalents, and in particular 0.75 to 0.9 equivalents of NaOH.

Within the framework of the invention the term "equivalents" is understood to mean "molar equivalents".

It has been found that adding less excipient than that given as the lower limit in the above-mentioned ranges causes inadequate stabilization of the solution. Adding amounts of excipient that exceed the aforementioned upper limits is not more advantageous in terms of the stability of the composition. It is furthermore feared that adding larger amounts of excipient will also lead to interactions with the active substance and thus tend to reduce the effectiveness of the composition.

Within the scope of the invention, particular preference is given to a pharmaceutical composition possessing the following constituents based on 100 ml of composition:

- a) 0.25 – 2.0 g, preferably 0.5 – 1.25 g {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof,
- b) 0.25 – 2.5 g, preferably 0.5 g – 1.5 g of arginine,
- c) 1.5 – 9.5 g, preferably 2.0 – 4.75 g of glucose,
- d) 0.5 – 4.0 g, preferably 0.75 – 2.0 g of NaH₂PO₄, and

e) water,

wherein said composition has a pH in the range of 7.5 to 8.5, preferably 7.7 to 8.0.

Furthermore, within the scope of the invention, particular preference is given to a pharmaceutical composition possessing the following constituents based on 100 ml of composition:

- a) 0.5 – 2.5 g, preferably 1.0 – 2.0 g {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof,
- b) 10.0 – 30.0 g, preferably 12.5g – 22.5 g of HP- β -cyclodextrin,
- c) 0.0 - 350 mg, preferably 75 - 225 mg, in particular 100 - 125 mg of NaOH, and
- d) water,

wherein said composition has a pH in the range of 7.5 to 8.5.

In the latter composition the NaOH is used preferably in the form of an approx. 0.1M aqueous solution.

It has been found that pharmaceutical compositions constituted in this way are particularly advantageous both as regards clinical effectiveness and also stability.

The pharmaceutical compositions according to the invention are generally produced by first producing an aqueous solution of the excipient and then adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid to this solution, if necessary followed by the addition of other additives, such as the at least one sugar and/or the at least one buffer. After adding all the constituents, the pH of the pharmaceutical composition is adjusted to the desired value, with particular attention being paid to the fact that when the pH is adjusted from a value in the alkaline range towards the physiological pH value, by adding an acid or a buffer, this adjustment is carried out slowly and carefully to avoid any precipitation of

{8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid resulting from an excessive local reduction in the pH value.

- 5 It is also possible to produce first of all individual solutions, with one solution containing the excipient and {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid and the other solution containing the other excipients, such as for example the at least one sugar and/or the at least one buffer, where in the next step the solutions are adjusted to the desired pH and
10 then mixed with each other.

It is further possible to dissolve, at least partially, the {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid in an aqueous alkaline solution, e.g. a solution of an
15 alkali metal hydroxide, preferably an NaOH solution, and then to add the excipient as well as, if necessary, the other constituents to the solution and, if necessary, to adjust the solution to the desired pH value.

It is further possible to lyophilize the solutions obtained by the above-mentioned methods
20 in order to obtain the solid pharmaceutical compositions according to the invention.

The subject matter of the invention is thus also a method to produce a pharmaceutical composition according to the invention having the following steps:

- 25 A) Dissolving the at least one excipient in the water,
B) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof to the solution obtained in step A),
C) If necessary, add at least one sugar and/or at least one buffer,
30 D) Adjust the pH to the desired value in order to obtain a pharmaceutical composition, and
E) Sterile-filter the solution obtained in step D) and fill into suitable containers.

- F) If necessary, perform a final sterilization of the solution obtained in step E) under heat.

The subject matter of the invention is also a method to produce a pharmaceutical composition according to the invention having the following steps:

- I.) Dissolving the at least one excipient in a part of the water,
II.) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof to the solution obtained in step I.),
III.) If necessary, adjusting the pH of the solution obtained in step II.) to the desired value to obtain a first solution,
IV.) Dissolving at least one sugar and/or a buffer in a part of the water,
V.) If necessary, adjust the pH of the solution obtained in step IV.) to the desired value to obtain a second solution,
VI.) Mixing the first and second solution to obtain a pharmaceutical composition, and
VII.) Sterile filtration of the solution obtained in step VI.) and filling into suitable containers.
VIII.) If necessary, perform a final sterilization of the solution obtained in step VII.) under heat.

The subject matter of the invention is also a method to produce a pharmaceutical composition according to the invention having the following steps:


- a.) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt to an aqueous NaOH solution, preferably an aqueous 0.1M NaOH solution to produce a solution or suspension,
b.) Adding water to the solution or suspension obtained in step a.),
c.) Adding cyclodextrin and NaCl to the solution or suspension obtained in step b.),
d.) Sterile filtration of the solution obtained in step c.) and fill into suitable containers.

e.) If necessary, perform a final sterilization of the solution obtained in step d.) under heat.

The subject matter of the invention is in addition a method to produce a solid pharmaceutical composition, wherein a pharmaceutical composition produced according to the aforementioned methods is lyophilized.


The {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid, or salts, solvates and solvates of the salts thereof, which are used to produce the pharmaceutical compositions according to the invention, are known and can be produced, for example, by the method described in WO 2006/133822.

The production takes place in particular by the saponification of the ester of a compound having the formula (II)




with a base.

The compound having the formula (II) can be produced by reacting a compound having the formula (III)




(III),

with a compound having the formula (IV) in the presence of a base.



(IV),


The compound having the formula (III) can be produced by reacting a compound having
5 the formula (V)



(V),

with phosphorus oxychloride, phosphorus trichloride or phosphorus pentachloride in the
presence of a base.

10 The compound having the formula (V) can be produced by reacting a compound having
the formula (VI)



(VI),

in the first step with acrylic acid methyl ester in the presence of a palladium catalyst and oleum, and in the second step with a base.

Compounds having the formulae (IV) and (VI) are in principle known to a person skilled
5 in the art or can be produced by customary methods known from the literature.

The saponification of the ester of a compound having the formula (II) to form {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid is achieved by reacting a compound having the
10 formula (II) with a base in an inert solvent, in a temperature range from 18 °C up to reflux of the solvent, preferably at 18 to 50 °C, more preferably at 20 to 30 °C, at normal pressure, within a period of, for example, 0.5 to 10 hours, preferably within 1 to 5 hours.

Bases are, for example, alkali hydroxides, such as sodium, lithium or potassium hydroxide,
15 or alkali carbonates, such as cesium carbonate, sodium or potassium carbonate, or alcoholates such as sodium or potassium methanolate, or sodium or potassium ethanolate, where the base may be present in aqueous solution.

Inert solvents are, for example, ethers, such as 1,2-dimethoxyethane, methyl *tert*-butyl
20 ether (MTBE), dioxane, tetrahydrofuran, glycol dimethyl ether or diethylene glycol dimethyl ether, alcohols such as methanol, ethanol, n-propanol, iso-propanol, n-butanol or *tert*-butanol, or water, or mixtures of solvents.

Sodium hydroxide in water and MTBE are preferred.

25 The synthesis of a compound having the formula (II) from a compound having the formula (III) and a compound having the formula (IV), in the presence of a base, takes place in an inert solvent, in a temperature range from 40 °C up to reflux of the solvent, preferably at reflux of the solvent, at normal pressure, within for example 2 to 48 hours, preferably
30 within 4 to 12 hours.

Bases are, for example, amine bases such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), 1-(3-methoxyphenyl)piperazine or triethylamine, or other bases such as potassium tert-butylate.

- 5 Inert solvents are, for example, chlorobenzene or ethers such as 1,2 dimethoxyethane, dioxane, glycol dimethyl ether or diethylene glycol dimethyl ether.

DBU in dioxane is preferred.

- 10 The conversion of a compound having the formula (V) to a compound having the formula (III) takes place by reacting a compound having the formula (V) with phosphorus oxychloride, phosphorus trichloride or phosphorus pentachloride, with phosphorus oxychloride being preferred, in the presence of a base in an inert solvent, in a temperature range from 40 °C up to reflux of the solvent, preferably at reflux of the solvent, at normal pressure,
15 within for example 1 to 48 hours, preferably within 2 to 12 hours.

Bases are, for example, amines such as 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), pyridine or triethylamine, or other bases such as potassium tert-butylate.

- 20 Inert solvents are for example hydrocarbons such as benzene, xylene, toluene or chlorobenzene.

DBU in chlorobenzene is preferred.

- 25 The conversion of a compound having the formula (VI) to a compound having the formula (V) takes place, in the first step, by reacting a compound of the formula (VI) with acrylic acid methyl ester in the presence of a palladium catalyst and oleum in a solvent, in a temperature range from 0 °C to 40 °C, preferably at room temperature, and in the second step by reaction with a base in an inert solvent, in a temperature range from 40 °C up to
30 reflux of the solvent, preferably at reflux of the solvent, at normal pressure, within for example 1 to 48 hours, preferably within 2 to 12 hours.

Palladium catalysts in the first step are, for example, palladium(II) acetate, bis(triphenylphosphine)palladium(II)chloride, tetrakis(triphenylphosphine)palladium(0), bis(tri(o-tolyl)phosphine)palladium-(II)-chloride, or a palladium catalyst produced from bis(acetonitrile)dichloropalladium or palladium(II) acetate and a ligand, for example tris(o-tolyl)phosphine, triphenylphosphine or diphenylphosphino ferrocene.

Solvents in the first step are, for example, organic acids such as acetic acid or propionic acid.

Palladium(II) acetate in acetic acid is preferred.


Bases in the second step are, for example, DBU, triethylamine or diisopropylethylamine.

Inert solvents in the second step are, for example, ethers such as 1,2-dimethoxyethane, dioxane, glycol dimethyl ether or diethylene glycol dimethyl ether, hydrocarbons such as benzene, xylene or toluene, or other solvents such as isobutyronitrile, acetonitrile, acetone, nitrobenzene, dimethylformamide, dimethylacetamide, dimethylsulfoxide or N-methylpyrrolidone.

DBU in acetone is preferred.

The production of the {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid used to produce the pharmaceutical composition according to the invention is described in more detail, by way of example, in the following Synthesis Diagram 1. This synthesis diagram is nothing more than an example and should in no way be understood as restrictive.


Synthesis Diagram 1



5[[Translation key:
Essigsäure = acetic acid]

As already mentioned further above, the {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid is used preferably in the form of the S-enantiomer. This S-enantiomer can be produced as shown, for example, in the following Synthesis Diagram 2.

Synthesis Diagram 2



[Translation key:
Bernsteinsäure = succinic acid]

The {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid, as well as the salts, solvates, and solvates of the salts thereof contained in the pharmaceutical composition exhibit an antiviral effect against representatives of the Herpes viridae group (herpes viruses), above all
5 against the cytomegaloviruses (CMV), in particular against the human cytomegalovirus (HCMV). The pharmaceutical compositions according to the invention are thus suitable for use in methods of treating and/or preventing diseases, especially infections with viruses, in particular the viruses referred to herein and the infectious diseases caused by them. The term "virus infection" is understood to mean not only an infection with a virus but also a
10 disease caused by infection with a virus.

Due to their properties and characteristics the pharmaceutical compositions according to the invention can be used to produce drugs that are suitable for use in methods of preventing and/or treating diseases, in particular virus infections.

15 The following areas of indication can be mentioned, by way of example:

- 1) Treatment and prevention of HCMV infections in AIDS patients (retinitis, pneumonitis, gastrointestinal infections).
- 2) Treatment and prevention of cytomegalovirus infections in bone marrow and organ
20 transplant patients who often contract life-threatening HCMV pneumonitis or encephalitis, as well as gastrointestinal and systemic HCMV infections.
- 3) Treatment and prevention of HCMV infections in neonates and infants.
- 4) Treatment of acute HCMV infection in pregnant women.
- 5) Treatment of HCMV infection in immune-suppressed patients suffering from cancer
25 and undergoing cancer therapy.
- 6) Treatment of HCMV-positive cancer patients with the aim of reducing HCMV-mediated tumour progression (cf. J. Cinatl, et al., *FEMS Microbiology Reviews* 2004, 28, 59-77).

30 The pharmaceutical compositions according to the invention are preferably used to produce drugs which are suitable for use in methods of preventing and/or treating infections

with a representative of the Herpes viridae group, in particular a cytomegalovirus, in particular the human cytomegalovirus.

Due to their pharmacological properties and characteristics, the pharmaceutical compositions according to the invention can be used by themselves and, if needed, also in combination with other active substances, especially antiviral substances such as for example valganciclovir, ganciclovir, valacyclovir, acyclovir, foscarnet, cidofovir and related derivatives in methods of treating and/or preventing virus infections, in particular HCMV infections.

Further subject matter of the present invention is the use of the pharmaceutical compositions according to the invention in a method of treating and/or preventing diseases, preferably virus infections, in particular infections with the human cytomegalovirus (HCMV) or another representative of the Herpes viridae group.

Further subject matter of the present invention is the use of the pharmaceutical compositions according to the invention in a method of treating and/or preventing diseases, in particular the aforementioned diseases.

Further subject matter of the present invention is the use of the pharmaceutical compositions according to the invention to produce a drug for use in methods of treating and/or preventing diseases, in particular the aforementioned diseases.

Further subject matter of the present invention is a method of treating and/or preventing diseases, in particular the aforementioned diseases, using an antivirally effective amount of the pharmaceutical compositions according to the invention.

The term "antivirally effective amount" denotes the pharmaceutical compositions according to the invention in a dose of at least 0.001 mg/kg.

In general, it has proved to be advantageous to administer the pharmaceutical compositions in such a way that about 0.001 to 10 mg per kg, preferably 0.01 to 5 mg per kg body

weight of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} acetic acid is administered.

Nevertheless, it may be necessary to deviate from the stated amounts, namely depending
5 on body weight, individual response to the active substance and the time and interval at
which it is applied. For example, in certain cases it may be sufficient to get by with less
than the aforementioned minimum amount, while in other cases the stated upper limit has
to be exceeded. When administering large amounts it may be recommendable to distribute
these in several individual doses over the course of a day.

10 The invention will now be described in detail on the basis of non-restrictive examples.

Unless otherwise stated, the percentages given in the following tests and examples are
weight percentages, parts are weight proportions, solvent ratios, dilution ratios and
15 concentrations of liquid solutions relate, in each case, to the volume.

List of abbreviations

	ACN	Acetonitrile
	API-ES-pos.	Atmospheric pressure ionization, electrospray, positive (in MS)
	API-ES-neg.	Atmospheric pressure ionization, electrospray, negative (in MS)
5	ca.	circa
	CI, NH ₃	chemical ionization (with ammonia)
	DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene
	DMAP	4-(Dimethylamino)pyridine
	DMSO	Dimethyl sulfoxide
10	ESTD	external standardization
	h	hour(s)
	HPLC	high pressure liquid chromatography
	conc.	concentrated
	min.	minutes
15	MS	mass spectroscopy
	MTBE	Methyl <i>tert</i> -butylether
	NMR	nuclear magnetic resonance spectroscopy
	R _T	retention time (in HPLC)
	VTS	vacuum drying cabinet

20

General HPLC methods:

Method 1 (HPLC): Instrument: HP 1050 with variable wavelength detection; column:
25 Phenomenex Prodigy ODS (3) 100A, 150 mm x 3 mm, 3 µm; Eluent A: (1.0 g KH₂PO₄ +
1.0 mL H₃PO₄) / 1 water, Eluent B: acetonitrile; gradient: 0 min 10% B, 25 min 80% B,
35 min 80% B; flow: 0.5 ml/min; temp.: 45 °C; UV detection: 210 nm.

Method 2 (HPLC): Instrument: HP 1050 with variable wavelength detection; column:
30 Chiral AD-H, 250 mm x 4.6 mm, 5 µm; Eluent A: *n*-heptane + 0.2% diethylamine, Eluent
B: isopropanol + 0.2% diethylamine; gradient: 0 min 12.5% B, 30 min 12.5% B; flow:
1 ml/min; temp.: 25 °C; UV detection: 250 nm.


Method 3 (HPLC): Instrument: HP 1050 with variable wavelength detection; column: Chiral AD-H, 250 mm x 4.6 mm, 5 μ m; Eluent A: *n*-heptane + 0.2% diethylamine, Eluent B: isopropanol + 0.2% diethylamine; gradient: 0 min 25% B, 15 min 25% B; flow: 1 ml/min; temp.: 30 °C; UV detection: 250 nm.

Examples

A) Production of {8-fluoro-2-[4-(3-methoxyphenyl)piperazin-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid

Example 1A

N-(2-fluorophenyl)-*N'*-[2-methoxy-5-(trifluoromethyl)phenyl]urea



2-methoxy-5-trifluoromethylphenyl isocyanate (78 kg) is melted at approx. 35 °C and dissolved in acetonitrile (a total of approx. 270 l), then 2-fluoroaniline (39.9 kg) is added and rinsed with acetonitrile (approx. 25 l). The resulting clear solution is agitated for 4 h at reflux and then cooled to approx. 75 °C. Once this temperature is reached, the solution is inoculated with seed crystals of the desired end product (200 g), agitated for an additional 15 min., and then cooled to 0 °C over the course of 3 h. The resulting crystalline product is isolated by centrifugation, washed with cold acetonitrile (twice using approx. 13 l), and dried at 45 °C in the VTS under purging with nitrogen (approx. 3.5 h). A total of 101.5 kg of *N*-(2-fluorophenyl)-*N'*-[2-methoxy-5-(trifluoromethyl)phenyl]urea is thus obtained as a solid, corresponding to 85.9% of theory.


¹H NMR (300 MHz, d₆-DMSO): δ = 8.93 (s, 1H), 8.84 (s, 1H), 8.52 (d, ³J = 2,3, 2H), 7.55 (d, ²J = 7.7, 1H), 7.38-7.26 (m, 3H), 7.22 (d, ²J = 8.5, 1H), 4.00 (s, 3H) ppm;

MS (API-ES-pos.): m/z = 409 [(M+H)⁺, 100%];

HPLC (Method 1): R_T = 22.4 and 30.6 min.

Example 2A

Methyl-(2Z)-3-[3-fluoro-2-({[2-methoxy-5-(trifluoromethyl)phenyl]carbamoyl}amino)-phenyl]acrylate



5 *N*-(2-fluorophenyl)-*N'*-[2-methoxy-5-(trifluoromethyl)phenyl] urea (51 kg) is dissolved in acetic acid (approx. 430 l) in one reactor in a nitrogen atmosphere. Methyl acrylate (20.1 kg) is added to the resulting solution and the resulting suspension is agitated until further use. Acetic acid (950 l) is placed in a second reactor, oleum (57 kg) is carefully added and palladium (II) acetate (7 kg) is dissolved in the mixture. The suspension formed in the first
10 reactor is then added to the mixture contained in the second reactor over the course of approx. 2 h; the reaction mixture is overflowed with a mixture of 96% nitrogen and 4% oxygen and the resulting reaction mixture is agitated for approx. 18 h at room temperature. Part of the acetic acid (approx. 900 l) is then distilled off; water (approx. 500 l) is added to the remaining reaction mixture over the course of approx. 1 h and the resulting suspension
15 is agitated for 1 h. The resulting particulate matter is filtered off, washed once with a mixture of acetic acid and water (1:1) and twice with water, and finally dried at approx. 30 mbar and 50 °C. A total of 44.8 kg of *methyl*-(2Z)-3-[3-fluoro-2-({[2-methoxy-5-(trifluoromethyl)phenyl]carbamoyl}amino)phenyl]acrylate is thus obtained as a solid, corresponding to 65.0% of theory.

20

¹H NMR (300 MHz, d₆-DMSO): δ = 9.16 (s, 1H), 8.84 (s, 1H), 8.45 (d, 1.7 Hz, 1H), 7.73 (m, 2H), 7.33 (m, 3H), 7.22 (d, 8.6 Hz, 1H), 6.70 (d, 16 Hz, 1H), 3.99 (s, 3H), 3.71 (s, 3H) ppm;

MS (API-ES-pos.): m/z = 429.9 [(M+NH₄)⁺]; 412.9 [(M+H)⁺]

25


HPLC : R_T = 46.4 min.

Instrument: HP 1100 with variable wavelength detection; column: Phenomenex Prodigy ODS (3) 100A, 150 mm x 3 mm, 3 μ m; Eluent A: (1.36 g KH_2PO_4 + 0.7 ml H_3PO_4) / 1 of water, Eluent B: acetonitrile; gradient: 0 min 20% B, 40 min 45% B, 50 min 80% B, 65 min 80% B; flow: 0.5 ml/min; temp.: 55 $^\circ\text{C}$; UV detection: 210 nm.

5

Example 3A

{8-fluoro-3-[2-methoxy-5-(trifluoromethyl)phenyl]-2-oxo-1,2,3,4-tetrahydroquinazoline-4-yl} methyl acetate




- 10 The compound in Example 2A (75 kg) is suspended in acetone (1600 l), and DBU (5.7 kg) is added. The resulting suspension is heated to reflux and agitated for 4 h at reflux. The resulting solution is cooled to a jacket temperature of 55 $^\circ\text{C}$ and filtered through kieselguhr. Part of the solvent (approx. 1125 l) is removed by distillation and the remaining residue is cooled for 2 h to 0 $^\circ\text{C}$. The resulting solid is separated out by centrifugation,
- 15 washed twice using cold acetone (approx. 15 l), and dried overnight at 45 $^\circ\text{C}$ under reduced pressure and purging with nitrogen to constant mass. A total of 58.3 kg of {8-fluoro-3-[2-methoxy-5-(trifluoromethyl)phenyl]-2-oxo-1,2,3,4-tetrahydroquinazoline-4-yl} methyl acetate is thus obtained as a solid, corresponding to 84.1% of theory.

HPLC (Method 1): R_T = 19.4 min.

20

Example 4A

(2*S*,3*S*)-2,3-bis[(4-methylbenzoyl)oxy] succinic acid-{(4*S*)-8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} methyl acetate (1:1 salt) chlorination/amination/crystallization



A solution of {8-fluoro-3-[2-methoxy-5-(trifluoromethyl)phenyl]-2-oxo-1,2,3,4-tetrahydroquinazoline-4-yl} methyl acetate (Example 3A, 129.2 kg) in chlorobenzene (800 l) is heated to reflux and azeotropically dried. Phosphorous oxychloride (144 kg) is added, and the reaction mixture is agitated for 3 h at reflux. Next, DBU (95 kg) and chlorobenzene (45 l) are added and agitated for an additional 9 h at reflux. The reaction mixture is cooled to room temperature, hydrolyzed by adding water, diluted with chlorobenzene (80 l), and neutralized with an aqueous solution of ammonia (25%). The phases are separated, the organic phase is washed with water and the solvent is distilled off. The remaining residue is dissolved in dioxane (170 l). 3-methoxyphenylpiperazine (66 kg), DBU (52 kg), and an additional 90 l of dioxane are added and the reaction mixture is heated for 4 h at reflux. The reaction mixture is cooled to room temperature, added to ethyl acetate (1300 l), washed once with water, 3 times with 0.2 N HCl, and once with an aqueous solution of NaCl, and the solvent is distilled off. The resulting residue is dissolved in ethyl acetate (800 l) and added to a solution of (2*S*,3*S*)-2,3-bis[(4-methylbenzoyl)oxy]-succinic acid (121 kg) in ethyl acetate (600 l). The resulting mixture is agitated for approx. 60 min. at room temperature and then inoculated with (2*S*,3*S*)-2,3-bis[(4-methylbenzoyl)oxy]-succinic acid-{(4*S*)-8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)-phenyl]-3,4-dihydroquinazoline-4-yl} methyl acetate and agitated for 3 days at room temperature. It is then cooled to 0 – 5 °C and agitated for an additional 3 h. The suspension is filtered and the remaining solid is rewashed in batches with ethyl acetate. A total of about 141 kg (calculated as dry weight) of the salt is thus obtained as a solid, corresponding to around 46.2% of theory, in three stages (chlorination, amination, and crystallization) compared to the racemate.

¹H NMR (300 MHz, d₆-DMSO): δ = 7.90 (d, ²J = 7.8, 4H), 7.56 (d, ²J = 8.3, 1H), 7.40 (d, ²J = 7.8, 4H), 7.28-7.05 (m, 4H), 6.91-6.86 (m, 2H), 6.45 (d, ²J = 8.3, 1H), 6.39-6.36 (m, 2H), 5.82 (s, 2H), 4.94 (m, 1H), 4.03 (q, ²J = 7.1, 2H), 3.83 (brs, 3H), 3.69 (s, 3H), 3.64 (s, 3H), 3.47-3.36 (m, 8H and water, 2H), 2.98-2.81 (m, 5H), 2.58-2.52 (m, 1H), 2.41 (s, 6H),
5 1.99 (s, 3H), 1.18 (t, ²J = 7.2, 3H) ppm;
HPLC (Method 1): R_T = 16.6 and 18.5 min.


Example 5A

(2*S*,3*S*)-2,3-bis[(4-methylbenzoyl)oxy]succinic acid-{(4*S*)-8-fluoro-2-[4-(3-methoxy-
10 phenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydro-
quinazoline-4-yl} methyl acetate (1:1 salt) / recrystallization

(2*S*,3*S*)-2,3-bis[(4-methylbenzoyl)oxy]-succinic acid – (S) {(4*S*)-8-fluoro-2-[4-(3-
methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydro-
15 quinazoline-4-yl} acetic acid methyl ester (1:1 salt) (141 kg, calculated as dry weight) is
suspended in ethyl acetate (1400 l) and dissolved by heating to reflux (77 °C). The solution
is filtered and slowly cooled to room temperature, which results in spontaneous
crystallization. The suspension is agitated for 16 h at RT, and then cooled to 0 – 5 °C and
agitated for an additional 3 h. The suspension is filtered and the remaining solid is re-
20 washed with cold ethyl acetate. The crystals are dried for 16 h in a vacuum at around 40
°C. A total of 131.2 kg of the salt is obtained as a solid, corresponding to 93.0% of theory.
HPLC (Method 1): R_T = 16.9 and 18.8 min.;
HPLC (Method 3): 99.9% e.e.

Example 6A

(*S*)-{8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-(2-methoxy-5-trifluoromethyl-
phenyl)-3,4-dihydroquinazoline-4-yl} acetic acid



A mixture of (2*S*,3*S*)-2,3-bis[(4-methylbenzoyl)oxy] succinic acid-{(4*S*)-8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} acetic acid methyl ester (1:1 salt) (30.8 kg), sodium bicarbonate (16.4 kg), and water (315 l) is mixed with MTBE (160 l). The phases are separated and the organic phase is treated with 35 l of an approximately seven-percent aqueous solution of sodium bicarbonate. The phases are separated and the organic phase is added to 125 l of an approximately four-percent aqueous solution of sodium hydroxide. The reaction mixture is heated to reflux, the solution is evaporated to dryness, and the reactor contents are then agitated for an additional 5 h at 55 – 60 °C. The reaction mixture is then added at approx. 22 °C to MTBE (160 l) and water (65 l) and agitated. The phases are separated and the organic phase is extracted with an approximately six-percent aqueous solution of sodium chloride (30 l). The combined aqueous phases are mixed with water (25 l) and MTBE (160 l) and the pH value is adjusted to approx. 6.5 with approx. 1 N of hydrochloric acid. The organic phase is separated out, the solvent is evaporated to dryness, and the residue is dissolved in acetone (approx. 75 l). The solvent is changed to acetone (6 distillations with approx. 130 l each). The final product is then precipitated by adding water, isolated through centrifugation, and dried in a vacuum dryer. A total of 16.5 kg of (*S*)-{8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-(2-methoxy-5-trifluoromethylphenyl)-3,4-dihydroquinazoline-4-yl} acetic acid is thus obtained as an amorphous solid, corresponding to 96.4% of theory.

¹H NMR (300 MHz, d₆-DMSO): δ = 7.53 (d, ²J = 8.4, 1H), 7.41 (brs, 1H), 7.22 (d, ²J = 8.5, 1H), 7.09-7.01 (m, 2H), 6.86 (m, 2H), 6.45 (dd, ²J = 8.2, ³J = 1.8, 1H), 6.39-6.34 (m, 2H), 4.87 (t, ²J = 7.3, 1H), 3.79 (brs, 3H), 3.68 (s, 3H), 3.50-3.38 (m, 4H), 2.96-2.75 (m, 5H), 2.45-2.40 (m, 1H) ppm;

5

MS (API-ES-neg.): m/z = 571 [(M+H), 100%];

HPLC (Method 1): R_T = 15.1 min;

10 HPLC (Method 2): 99.8% e.e.; Pd (ICP): <1 ppm.

B) Exemplary embodiments of pharmaceutical compositions according to the invention

15 Example 1

Production of a pharmaceutical composition using cyclodextrin:

In a three-necked flask, 30.03 g of hydroxypropyl-β-cyclodextrin HP5 (Kleptose HPB, Roquette) are mixed with 68.365 g of water for injection purposes, and 6.6 g of a 1 M
20 solution of sodium hydroxide is added to the mixture. Following the addition of 5.005 g of the compound from Example 6A, the mixture is heated to 50 °C and agitated for 24 h until a clear solution is formed. The solution is sterile filtered (pore diameter 0.22 μm) and transferred under aseptic conditions to sterile 20-ml glass containers. The filled glass containers are sealed with infusion plugs and flange caps.

25

Example 2

Production of a first pharmaceutical composition using arginine as an excipient:

To produce a first stock solution, 262.38 mg of L-arginine are weighed in a 25-ml volumetric flask and then dissolved in approx. 22 ml of water for injection. 504.51 mg of the
30 compound from Example 6A are added to the resulting arginine solution, and the mixture

is agitated for approx. 1 h until a clear solution is obtained. The volume is then topped up with water for injection purposes.

To produce a second stock solution, 40.05 mg of sodium dihydrogen phosphate dihydrate
5 are weighed in a 50 ml volumetric flask and dissolved in approx. 48 ml of water for injection. The mixture is agitated until a clear solution is obtained and the volume is topped up with water for injection.

To produce a solution with a concentration of 10 mg/ml of {8-fluoro-2-[4-(3-
10 methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid, 12.5 ml of the first stock solution are mixed with 10 ml of the second stock solution in a 25-ml volumetric flask and the pH is slowly and carefully adjusted with approx. 200 µl of 1 M HCl. The volume is then topped up with the second stock solution in order to obtain a mixture with a final pH of 7.9.

15 Using this protocol, compositions with varying concentrations of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid can be produced and only the quantity of the first stock solution used must be varied. However, it is important that the pH does not vary too
20 much and particularly that it shifts not towards the acid range.

The solutions thus obtained are sterile filtered (pore diameter 0.22 µm) and transferred under aseptic conditions to sterile containers. The containers are sealed with infusion plugs and flange caps.

Example 3

Production of a second pharmaceutical composition using arginine as an excipient:

To produce a first stock solution, 2.1 g of L-arginine are dissolved in 88.8 g of water for
30 injection purposes. 2 g of the compound from Example 6A are added to the resulting arginine solution, and the mixture is agitated for approx. 1 h until a clear solution is obtained. The pH value of the resulting solution is adjusted to 7.8 by adding 1 M HCl

dropwise; it is important that the HCl be added slowly so that the compound from Example 6A does not precipitate. If necessary, the volume of solution is then topped up to 100 ml.

To produce a second stock solution, 3.1 g of sodium dihydrogen phosphate dihydrate and 8.4 g of glucose are weighed in an appropriate container and dissolved in approx. 74.5 g of water for injection. The mixture is agitated until a clear solution is obtained, and the pH of the resulting solution is adjusted to 7.8 using 1 M NaOH. Finally, if necessary, the volume of solution is topped up to 100 ml.

To produce a solution with a concentration of 10 mg/ml of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid, 50.5 g of the first stock solution are mixed with 53.0 g of the second stock solution and agitated for 5 min.

Using this protocol, compositions with varying concentrations of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid can be produced and only the quantity of the first stock solution used must be varied.

The solutions thus obtained are sterile filtered (pore diameter 0.22 µm) and transferred under aseptic conditions to sterile containers. The containers are sealed with infusion plugs and flange caps.

Example 4

Production of a third pharmaceutical composition using arginine as an excipient:

To produce a first stock solution, 1.05 g of L-arginine and 1 g of the compound from Example 6A are dissolved in approx. 50.0 g of water for injection purposes in a 100-ml volumetric flask and the mixture is agitated until a clear solution is obtained. The pH value of the resulting solution is adjusted to 7.8 by adding 0.1 M HCl dropwise (approx. 43.5 ml); it is important that the HCl be added slowly so that the compound from Example 6A does not precipitate. Finally, the volume of solution is topped up to 100 ml.

To produce a second stock solution, 1.56 g of sodium dihydrogen phosphate dihydrate and 4.18 g of glucose are dissolved in approx. 80.0 g of water for injection purposes in a 100-ml volumetric flask. The mixture is agitated until a clear solution is obtained, and the pH of the resulting solution is adjusted to 7.8 using 1 M NaOH (approx. 9.1 ml). Finally, the volume of solution is topped up to 100 ml.

To produce a solution with a concentration of 5 mg/ml of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid, 50.24 g of the first stock solution are mixed with 51.35 g of the second stock solution and agitated for 5 min.

Using this protocol, compositions with varying concentrations of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid can be produced and only the quantity of the first stock solution used must be varied.

The solutions thus obtained are sterile filtered (pore diameter 0.22 µm) and transferred under aseptic conditions to sterile containers. The containers are sealed with infusion plugs and flange caps.

Example 5

Production of a fourth pharmaceutical composition using arginine as an excipient:

To produce a first stock solution, 2.11 g of L-arginine are mixed with 2.01 g of the compound from Example 6A in a 100-ml volumetric flask and the volume is topped up with water for injection. The pH value of this first stock solution was 9.8.

To produce a second stock solution, 3.12 g of sodium dihydrogen phosphate dihydrate, 8.35 g of glucose, and 0.50 g of NaCl are dissolved in approx. 80.0 g of water for injection in a 100-ml volumetric flask. The mixture is agitated until a clear solution is obtained, and the pH of the resulting solution is adjusted to 6.5 using 1 M NaOH (approx. 9.7 ml). Finally, the volume of solution is topped up to 100 ml.

To produce a solution with a concentration of 10 mg/ml of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid, 50.50 g of the first stock solution are mixed with
5 52.65 g of the second stock solution and agitated for 5 min.

Using this protocol, compositions with varying concentrations of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid can be produced and only the quantity of the first
10 stock solution used must be varied or the pH may need to be adjusted.

The solutions thus obtained are sterile filtered (pore diameter 0.22 μ m) and transferred under aseptic conditions to sterile containers. The containers are sealed with infusion plugs and flange caps.

15

Example 6

Production of a pharmaceutical composition using lysine as an excipient:

To produce a first stock solution, 217.24 mg of lysine are weighed in a 25-ml volumetric
20 flask and then dissolved in 22 ml of water for injection. 500.71 mg of the compound from Example 6A are added to the resulting solution, and the mixture is agitated for approx. 1 h until a clear solution is obtained. The pH of the resulting solution is then adjusted to pH 8 using approx. 460 μ l of 1 M HCl; it is again important to avoid too sharp a local decline of the pH value and the associated precipitation of the compound from Example 6A. The
25 volume is then topped up with water for injection purposes in order to obtain a first stock solution.

To produce a second stock solution, 242.01 mg of sodium dihydrogen phosphate are weighed in a 50-ml volumetric flask and then dissolved in approx. 48 ml of water for
30 injection purposes. The mixture is agitated until a clear solution is obtained. The pH of the resulting solutions is adjusted to a pH value of 8 using approx. 1.825 μ l of 1 M NaOH, and the volume is then topped up with water for injection.

To produce a solution with a concentration of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} acetic acid of 5 mg per ml of solution, 6.5 ml of the first stock solution are filled into a 25-ml volumetric flask and the volume is topped up with the
5 second stock solution to obtain a solution with a final pH of 8.

As previously described in Example 2, for instance, pharmaceutical compositions with other concentrations of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid can also be produced
10 by varying the quantity of the first stock solution.

The resulting solution is sterile filtered (pore diameter 0.22 μ m) and transferred under aseptic conditions to sterile containers.

15 Example 7

Production of a solid pharmaceutical composition which can be reconstituted to produce an infusion solution:

To produce a first stock solution, 261.16 mg of L-arginine are weighed in a 25-ml volumetric flask and then dissolved in 22 ml of water for injection purposes. 502.45 mg of the
20 compound in Example 6A are added to the resulting solution, and the mixture is agitated for approx. 1 h until a clear solution is obtained. The pH of this solution is adjusted to a value of 7.8 using approx. 660 μ l of 1 M HCl; it is again important that the compound from Example 6A does not precipitate. The volume is then topped up with water for
25 injection.

To produce a second stock solution, 240.05 mg of sodium dihydrogen phosphate are weighed in a 50-ml volumetric flask and dissolved in approx. 48 ml of water for injection purposes. The mixture is agitated until a clear solution is obtained. The pH of the resulting
30 solutions is adjusted to a value of 7.8 using approx. 1.850 μ l of 1 M NaOH, and the volume is topped up with water for injection.

To produce a solution with a concentration of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} acetic acid of 10 mg/ml, 12.5 ml of the first stock solution are placed in a 25-ml volumetric flask and the volume is topped up with the second stock solution to obtain a solution with a final pH of 7.8. One ml of the clear, colourless solutions is placed in each of 2-ml glass containers with suitable plugs and lyophilized in an EPSILON 2-4 D freeze dryer (Martin Christ GmbH, Germany) to obtain a colourless powder, which can easily be reconstituted into a solution suitable for intravenous application by adding 1 ml of water.

10

Example 8

Production of a solid pharmaceutical composition which can be reconstituted to produce an infusion solution:

15 To produce a first stock solution, 210.49 g of L-arginine and 9665.8 g of water for injection are mixed and the mixture is agitated until a clear solution is obtained. While agitating the resulting solution, 199.23 g of the compound from Example 6A are added in small portions and the mixture is agitated for at least 30 min until a clear solution is obtained.

20

To produce a second stock solution, 309.19 g of sodium dihydrogen phosphate dihydrate, 827.47 g of glucose, 49.55 g of sodium chloride, and 7900.0 g of water for injection are agitated until a clear solution is obtained. The pH of the resulting solution is adjusted to 6.55 using 1 M NaOH, and 1418.29 g of water for injection purposes—less the amount of NaOH solution used to adjust the pH value—are added to the resulting solution.

25

To produce the desired solution, the second stock solution is added slowly and in small portions to the first stock solution while gently agitating, and the resulting solution is sterile filtered. Appropriate sterile glass containers with suitable plugs are each filled with 15 ml of the clear, colourless solution and lyophilized in a freeze dryer to obtain a colourless powder.

30

The lyophilisate thus obtained can be reconstituted into a solution by adding, for instance, 30 ml of water for injection purposes, and the resulting solution can then be further diluted as necessary for use in infusions.

5 Example 9

Production of a second pharmaceutical composition using cyclodextrin:

2.00 g of the compound from Example 6A are measured in and 30 g of a 0.1 M solution of NaOH are added. The resulting mixture is agitated for 30 min (the compound from
10 Example 6A need not dissolve completely). 57.7 g of water for injection as well as 15.0 g hydroxypropyl- β -cyclodextrin HP5 (Kleptose HPB, Roquette) and 0.31 g of NaCl are added to the resulting mixture and agitated until a clear solution is obtained. The solution is sterile filtered (pore diameter 0.22 μ m) and transferred under aseptic conditions to sterile 20-ml glass containers. The filled glass containers are sealed with infusion plugs and
15 flange caps. The filled glass containers thus obtained may be heat-sterilized.

Example 10

Production of a third pharmaceutical composition using cyclodextrin:

20 2.00 g of the compound from Example 6A are measured in and 30 g of a 0.1 M solution of NaOH are added. The resulting mixture is agitated for 30 min (the compound from Example 6A need not dissolve completely). 54.8 g of water for injection purposes as well as 20.0 g hydroxypropyl- β -cyclodextrin HP5 (Kleptose HPB, Roquette) and 0.205 g of NaCl are added to the resulting mixture and agitated until a clear solution is obtained. The
25 solution is sterile filtered (pore diameter 0.22 μ m) and transferred under aseptic conditions to sterile 20-ml glass containers. The filled glass containers are sealed with infusion plugs and flange caps. The filled glass containers thus obtained may be heat-sterilized.

Example 11

Production of a fourth pharmaceutical composition using cyclodextrin:

0.5 g of the compound from Example 6A are weighed in and added to 8.75 g of a 0.1 M solution of NaOH. The resulting mixture is agitated for 30 min (the compound from Example 6A does not need to be dissolved completely). 12.45 g of water for injection purposes and 5.0 g of 2-O-methyl- β -cyclodextrin (Crysmeb, Roquette) are added to the resulting mixture and agitated until a clear solution is obtained. The pH value of the solution is adjusted to 7.5 using 1M HCl, the solution is sterile filtered (pore diameter 0,22 μ m) and transferred under aseptic conditions to sterile 20-ml glass containers. The filled glass containers are sealed with infusion plugs and flange caps. The filled glass containers thus obtained may be heat-sterilized.

Example 12

Production of a fifth pharmaceutical composition using cyclodextrin:

0.5 g of the compound from Example 6A are measured in and added to 13.125 g of a 0.1 M solution of NaOH. The resulting mixture is agitated for 30 min (the compound from Example 6A does not need to be dissolved completely). 8.075 g of water for injection purposes and 5.0 g of sulphaalkyl ether- β -cyclodextrin (Captisol, CyDex Pharmaceuticals Inc.) are added to the resulting mixture and agitated until a clear solution is obtained. The pH value of the solution is adjusted to 7.5 using 500 μ l of 1M HCl, the solution is sterile filtered (pore diameter 0.22 μ m) and transferred under aseptic conditions to sterile 20-ml glass containers. The filled glass containers are sealed with infusion plugs and flange caps. The filled glass containers thus obtained may be heat-sterilized. Prior to dispensing, the described solution may be diluted with an isotonic solution, e.g. an isotonic infusion solution.

C) Stability measurement

To measure stability, the solutions produced in Examples 1 to 6 were stored for two, three and six weeks at 2 – 8 °C, 25 °C, 40 °C. All solutions demonstrated adequate stability.

Furthermore, the stability of a solution reconstituted from the preparation produced from Example 7 was tested over 24 h at 2 – 8 °C, 25 °C, and 40 °C. The solution was demonstrated to be stable under all conditions over a period of 24 h.

5 **D) Comparative trials for fixed pharmaceutical compositions**

To demonstrate the advantageous properties of the solid pharmaceutical compositions obtained in Example 7 as compared to other solid compositions, the solid substances contained in the solution were mixed and reconstitution trials were then carried out. In
10 none of the investigated cases it was possible to obtain a clear solution.

E) Assessment of physiological efficacy

The in vitro effects of the compositions according to the present invention on the
15 replication of the HCMV (human cytomegalovirus) can be seen in the following antiviral assay:

F) HCMV fluorescence-reduction test

20 The solution from Example 8 is used without further dilution in the test. The composition from Example 6A is used as a 50-millimolar (mM) solution in dimethyl sulphoxide (DMSO). Ganciclovir[®], Foscarnet[®] or Cidofovir[®] can be used as reference compositions. One day before the beginning of the test, 1.5×10^4 human foreskin fibroblasts (NHDF cells)/well are seeded in 200 µl of cell culture medium in Wells B2 – G11 of 96-well plates
25 (black with transparent floor). The wells along the edges of each 96-well plate are filled with 200 µl of medium only in order to prevent edge effects. On the day of the test the cell culture medium in Wells B2 – G11 of each 96-well plate is vacuumed off by a suction device and replaced with 100 µl of virus suspension (multiplicity of infection (MOI): 0.1 – 0.2). The virus used is a recombinant HCMV which has integrated an expression cassette
30 for green fluorescence protein (GFP) in the virus genome (HCMV AD 169 RV-HG [E. M. Borst, K. Wagner, A. Binz, B. Sodeik, and M. Messerle, 2008, *J. Virol.* 82:2065-2078.]). After an incubation time of 2 h at 37 °C and 5% CO₂, the virus inoculate is vacuumed off

by a suction device and all wells, with the exception of the wells in Column 3, are filled with 200 μ l of cell culture medium. Column 2 is not treated further and serves as a virus control. The wells in Column 3 are each filled with 300 μ l of composition or solution of the test substance (the latter diluted in cell culture medium) for duplicate analysis. The concentration of the respective antiviral substance in Column 3 is ~ 27 times as concentrated as the respective anticipated EC_{50} value. The test substance in Column 3 is diluted in 8 steps to a concentration of 1:3 across the 96-well plate by transferring 100 μ L from each columns into its respective right-hand column, where it is mixed with the 200 μ l of cell culture medium already present there. In this way, three antiviral substances are tested in duplicate analyses. The plates are incubated for 7 days at 37 $^{\circ}$ C and 5% CO_2 . Subsequently, all wells on the plate are washed 3 times with PBS (phosphate-buffered saline) and filled with 50 μ l of PBS. The GFP intensity of each well in a 96-well plate is then determined using a fluorescence scanner (FluoBox; Bayer Technology Services GmbH; filter settings: GFP, Ex 480 nm, Em 520 nm). The measured values thus obtained can be used to determine the EC_{50} of an anti-HCMV:

EC_{50} (GFP-RA) = substance concentration in μ M which reduces GFP fluorescence by 50% in comparison to the untreated virus control.

Representative in vitro efficacy data for the compositions according to the present invention are reproduced in Table 1:

Table 1

Virus strain	Example 6A EC_{50} [μM]	Example 8 EC_{50} [μM]	Ganciclovir EC_{50} [μM]
AD169 RV-HG	0.0022 ± 0.0002	0.0026 ± 0.0005	2.5 ± 0.4

Claims

1. A pharmaceutical composition, particularly for intravenous administration,
 5 comprising the following, namely:
 - a) {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} acetic acid or a salt, a solvate or a solvate of a salt thereof,
 - b) at least one excipient selected from modified β -cyclodextrins, in particular the
 10 hydroxyalkyl β -cyclodextrins, the alkylhydroxyalkyl β -cyclodextrins and the sulfoalkyl cyclodextrins, and
 - c) water.
2. The pharmaceutical composition according to claim 1, further comprising at least
 15 one buffer, preferably selected from the phosphate buffers, the Tris buffers and the citrate buffers.
3. The pharmaceutical composition according to claim 1 or 2, further comprising at least one sugar, preferably selected from the group consisting of glucose, sucrose, lactose,
 20 maltose, trehalose, sorbitol and mannitol.
4. The pharmaceutical composition according to any one of claims 1 to 3 characterized in that {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a
 25 solvate of a salt thereof is present in an amount corresponding to 1 to 100 mg, preferably 2 to 50 mg, more preferably 2 to 25 mg, and in particular to 5 to 20 mg of pure active compound per ml of preparation.
5. The pharmaceutical composition according to any one of claims 1 to 4,
 30 characterized in that the preparation has a pH in the range of 7.5 to 8.5.
6. The pharmaceutical composition according to any one of claims 1 to 5, characterized in that the at least one excipient is present in an amount of 1 to 5 equivalents, preferably of 2 to 5 equivalents and more preferably of 2.5 to 4.5 equivalents in relation to

the content of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} acetic acid.

- 5 7. The pharmaceutical composition according to any one of claims 1 to 6, characterized in that 100 ml of said composition comprises the following:
- a) 0.2 - 2.0 g, preferably 0.5 - 1.25 g of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} acetic acid or a salt, a solvate or a solvate of a salt thereof,
 - 10 b) 2.5 g, preferably 0.5 - 1.5 g of cyclodextrin,
 - c) 1.5 - 9.5 g, preferably 2.0 - 4.75 g of glucose,
 - d) 0.6 - 4.0 g, preferably 0.75 - 2.0 g of NaH_2PO_4 , and
 - e) water,

wherein said composition has a pH in the range of 7.5 to 8.5, preferably 7.7 to 8.0.

15

8. The pharmaceutical composition according to claim 1, characterized in that said preparation, in relation to the content of {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl} acetic acid, contains 1 to 10 equivalents, preferably 2 to 7 equivalents, and in particular 2.5 to 5
- 20 equivalents of cyclodextrin as well as 0 to 2.0 equivalents, preferably 0.5 to 1.5 equivalents, and in particular 0.75 to 0.9 equivalents of NaOH.

- 9 The pharmaceutical composition according to any one of claims 1 to 5 or according to claim 8, characterized in that 100 ml of said composition comprises the following:

25

30

- a) 0.5 - 2.5 g, preferably 1.0 - 2.0 g {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof,
 - b) 10.0 - 30.0 g, preferably 12.5g - 22.5 g of HP- β -cyclodextrin,
 - 5 c) 0.0 - 350 mg, preferably 75 - 225 mg, in particular 100 - 125 mg of NaOH, and
 - d) water,
- wherein said preparation has a pH in the range of 7.5 to 8.5.

10. A solid pharmaceutical composition produced by lyophilizing a pharmaceutical
10 preparation according to any one of claims 1 to 9.

11. A method for producing a pharmaceutical composition according to any one of claims 1 to 7 using the following steps, namely:
- A) Dissolving the at least one excipient in the water,
 - 15 B) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt thereof to the solution obtained in step A),
 - C) If necessary, add at least one sugar and/or at least one buffer,
 - D) Adjust the pH to the desired value in order to obtain a pharmaceutical composition,
 - 20 and
 - E) Sterile-filter the solution obtained in step D) and fill into suitable containers.
 - F) If necessary, perform a final sterilization, of the solution obtained in step E) under heat.

- 25 12. A method for producing a pharmaceutical composition according to any one of claims 1 to 7 using the following steps, namely:
- I.) Dissolving the at least one excipient in a part of the water,
 - II.) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a
 - 30 solvate of a salt thereof to the solution obtained in step I.),

- III.) If necessary, adjusting the pH of the solution obtained in step II.) to the desired value to obtain a first solution,
- IV.) Dissolving at least one sugar and/or a buffer in a part of the water,
- V.) If necessary, adjust the pH of the solution obtained in step IV.) to the desired value to obtain a second solution,
- VI.) Mixing the first and second solution to obtain a pharmaceutical composition, and
- VII.) Sterile filtration of the solution obtained in step VI.) and filling into suitable containers.
- VIII.) If necessary, perform a final sterilization of the solution obtained in step VII.) under heat.
13. A method for producing a pharmaceutical composition according to any one of claims 1 to 5 or according to claim 8 or 9 using the following steps, namely:
- a.) Adding {8-fluoro-2-[4-(3-methoxyphenyl)piperazine-1-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazoline-4-yl}acetic acid or a salt, a solvate or a solvate of a salt to an aqueous NaOH solution, preferably an aqueous 0.1M NaOH solution to produce a solution or suspension,
- b.) Adding water to the solution or suspension obtained in step a.),
- c.) Adding cyclodextrin and NaCl to the solution or suspension obtained in step b.),
- d.) Sterile filtration of the solution obtained in step c.) and fill into suitable containers.
- e.) If necessary, perform a final sterilization of the solution obtained in step d.) under heat.
14. A method for producing a solid pharmaceutical composition according to claim 10 comprising the production of a pharmaceutical composition corresponding to a method according to any one of claims 11 to 13 followed by lyophilization of the pharmaceutical composition obtained in order to obtain a solid pharmaceutical composition.

15. The pharmaceutical composition according to any one of claims 1 to 10 for use in a method of treatment and/or prophylaxis of diseases, in particular of virus infections, preferably human cytomegalovirus (HCMV) infections or infections with another member of the herpes viridae group.

5

16. The use of a pharmaceutical composition according to any one of claims 1 to 10 for producing a medicament for the treatment and/or prophylaxis of diseases, in particular of virus infections, preferably human cytomegalovirus (HCMV) infections or infections with another member of the herpes viridae group.