

COMPOUND FOR TREATING GOUT OR HYPERURICEMIA

CROSS-REFERENCE

[0001] This application claims benefit of U.S. Provisional Application No. 62/847,519, filed on May 14, 2019, which is herein incorporated by reference in its entirety.


BACKGROUND

[0002] Hyperuricemia is caused by the overproduction or under-excretion of uric acid, and is considered to be a causative factor of several diseases that significantly impair the quality of life. For example, hyperuricemia is considered the causative factor of gout – the most prevalent form of inflammatory arthritis, characterized by severe pain and tenderness in joints caused by urate crystal accumulation. The identification of a gout/hyperuricemia drug effective in lowering serum uric acid (sUA) with reduced toxicity represents an unmet medical need that would have beneficial impact on patients.

SUMMARY OF THE INVENTION

[0003] In one aspect, described herein is a compound that is (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof.

[0004] In another aspect, described herein is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof:



Formula (I);

wherein:


R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of hydrogen, deuterium, hydroxy, and methoxy; wherein at least one of R₁, R₂, R₃, and R₄ is deuterium; and

R₅ is -CH₂CH₃, -CH₂CH₂(OH), or -CH(OH)CH₂(OH).

[0005] In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein at least two of R₁, R₂, R₃, and R₄ is deuterium. In some

embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein at least three of R_1 , R_2 , R_3 , and R_4 is deuterium. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R_1 , R_2 , R_3 , and R_4 are deuterium. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R_5 is $-\text{CH}_2\text{CH}_3$. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R_5 is $-\text{CH}_2\text{CH}_2(\text{OH})$. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R_5 is $-\text{CH}(\text{OH})\text{CH}_2(\text{OH})$.

[0006] In another aspect, described herein is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof:



Formula (II);

wherein:

R_1 , R_2 , R_3 , and R_4 are each independently selected from a group consisting of hydrogen, deuterium, hydroxy, and methoxy; wherein at least one of R_1 , R_2 , R_3 , and R_4 is deuterium.

[0007] In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein at least two of R_1 , R_2 , R_3 , and R_4 is deuterium. In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein at least three of R_1 , R_2 , R_3 , and R_4 is deuterium. In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein at least one of R_1 , R_2 , R_3 , and R_4 is hydroxy.

[0008] In another aspect, described herein is a method for treating or preventing hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl)-4,5,7- d_3 methanone, or a pharmaceutically acceptable salt or solvate thereof, described herein. In another aspect, described herein is a method for treating or preventing hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II) described herein, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating or preventing hyperuricemia in an individual in need thereof. In some

embodiments is a method for treating hyperuricemia in an individual in need thereof. In some embodiments is a method for preventing hyperuricemia in an individual in need thereof. In some embodiments is a method for treating or preventing gout in an individual in need thereof. In some embodiments is a method for treating gout in an individual in need thereof. In some embodiments is a method for preventing gout in an individual in need thereof. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is from about 3 mg to about 1500 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is from about 3 mg to about 600 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is from about 5 mg to about 300 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is from about 10 mg to about 200 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is from about 10 mg to about 100 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered orally. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is taken with food. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is taken without food. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered to the individual once per day. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered to the individual twice per day. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-

hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered with at least one additional therapeutic agent. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered with a xanthine oxidase inhibitor. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered with a xanthine oxidase inhibitor selected from allopurinol, oxypurinol, febuxostat, topiroxostat, and inositol. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered with a sodium-glucose co-transporter-2 (SGLT2) inhibitor. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered with an SGLT2 inhibitor selected from canagliflozin, dapagliflozin, empagliflozin, empagliflozin/linagliptin, empagliflozin/metformin, and dapagliflozin/metformin. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered with a xanthine oxidase inhibitor and a SGLT2 inhibitor. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, is administered with a xanthine oxidase inhibitor and a SGLT2 inhibitor, wherein the xanthine oxidase inhibitor is selected from allopurinol, oxypurinol, febuxostat, topiroxostat, and inositol, and the SGLT2 inhibitor is selected from canagliflozin, dapagliflozin, empagliflozin, empagliflozin/linagliptin, empagliflozin/metformin, and dapagliflozin/metformin.

[0009] In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients.

[0010] In some embodiments is a pharmaceutical composition comprising a compound of Formula (I) or (II) described herein, or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients.

[0011] In another aspect is a pharmaceutical composition for the treatment or prevention of hyperuricemia or gout, wherein the pharmaceutical composition comprises a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients. In another aspect is a pharmaceutical composition for the treatment or prevention of hyperuricemia or gout, wherein the pharmaceutical composition comprises a therapeutically effective amount of a compound of Formula (I) or (II) described herein, or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients. In some embodiments is a pharmaceutical composition for the treatment or prevention of hyperuricemia in an individual in need thereof. In some embodiments is a pharmaceutical composition for the treatment of hyperuricemia in an individual in need thereof. In some embodiments is a pharmaceutical composition for the prevention of hyperuricemia in an individual in need thereof. In some embodiments is a pharmaceutical composition for the treatment or prevention of gout in an individual in need thereof. In some embodiments is a pharmaceutical composition for the treatment of gout in an individual in need thereof. In some embodiments is a pharmaceutical composition for the prevention of gout in an individual in need thereof. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is from about 3 mg to about 1500 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is from about 3 mg to about 600 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is from about 5 mg to about 300 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is from about 10 mg to about 200 mg. In some embodiments, the therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate

thereof, is from about 10 mg to about 100 mg. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is formulated for oral, intravenous, intramuscular, or subcutaneous administration. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is formulated for oral administration. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is taken with food. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is taken without food. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is administered to the individual once per day. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, is administered to the individual twice per day. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, further comprises at least one additional therapeutic agent. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, further comprises a xanthine oxidase inhibitor. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, further comprises a xanthine oxidase inhibitor selected from allopurinol, oxypurinol, febuxostat, topiroxostat, and inositol. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-

hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, further comprises an SGLT2 inhibitor. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, further comprises an SGLT2 inhibitor selected from canagliflozin, dapagliflozin, empagliflozin, empagliflozin/linagliptin, empagliflozin/metformin, and dapagliflozin/metformin. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, further comprises a xanthine oxidase inhibitor and an SGLT2 inhibitor. In some embodiments, the pharmaceutical composition of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, further comprises a xanthine oxidase inhibitor and an SGLT2 inhibitor, wherein the xanthine oxidase inhibitor is selected from allopurinol, oxypurinol, febuxostat, topiroxostat, and inositol, and the SGLT2 inhibitor is selected from canagliflozin, dapagliflozin, empagliflozin, empagliflozin/linagliptin, empagliflozin/metformin, and dapagliflozin/metformin.


INCORPORATION BY REFERENCE

[0012] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the extent applicable and relevant and to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

DETAILED DESCRIPTION OF THE INVENTION

[0013] Benzbromarone is a uricosuric agent effective in lowering serum uric acid sUA and treating gout. It has been found that therapy using benzbromarone can lead to lowering of sUA even following a single dose and continue to be lowered following multiple doses, and that chronic therapy can bring sUA into target levels of <6 mg/dL. However, in certain patients, benzbromarone is associated with hepatotoxicity. A high proportion of these patients developed acute liver failure leading to death or emergency liver transplantation. As a result, benzbromarone was never approved for use in the United States. In addition, the hepatotoxicity of benzbromarone led to its withdrawal in Europe in 2003. Benzbromarone is converted to reactive metabolites by CYP2C9. Benzbromarone is metabolized to 5,6-


dihydroxybenzbromarone via 6-OH benzbromarone by CYP2C9, followed by the oxidation of 5,6-dihydroxybenzbromarone to a reactive ortho-quinone intermediate. The mechanism of benzbromarone hepatotoxicity is believed to be a result of its hepatic metabolism by CYP2C9 and possible effects of the 6-OH benzbromarone and its further metabolites on mitochondrial function (Iwamura et al., Drug Metabolism and Disposition, 2011, 39, 838-846; Uchida et al., Drug Metab. Pharmacokinet., 2010, 25, 605-610).



[0014] Described herein is (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), an active metabolite of (3,5-dibromo-4-hydroxyphenyl)(2-(1-hydroxyethyl)benzofuran-3-yl-4,5,6,7-*d*₄)methanone.

Compound 1


[0015] In one embodiment is (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone. “Compound 1” or “(3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone” refers to the compound with the following structure:



[0016] In some embodiments, Compound 1 undergoes phase II metabolism. In some embodiments, Compound 1 is glucuronidated. In some embodiments, Compound 1 forms a sulfate conjugate.

Compounds of Formula (I)

[0017] In some embodiments, described herein is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof:



Formula (I);

wherein:

R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of hydrogen, deuterium, hydroxy, and methoxy; wherein at least one of R₁, R₂, R₃, and R₄ is deuterium; and


R₅ is -CH₂CH₃, -CH₂CH₂(OH), or -CH(OH)CH₂(OH).

[0018] In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of hydrogen and deuterium. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of deuterium, hydroxy, and methoxy. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of deuterium and hydroxy. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein at least two of R₁, R₂, R₃, and R₄ is deuterium. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein at least three of R₁, R₂, R₃, and R₄ is deuterium. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are deuterium. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R₅ is -CH₂CH₃. In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R₅ is -CH₂CH₂(OH). In some embodiments is a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, wherein R₅ is -CH(OH)CH₂(OH).

[0019] In some embodiments, a compound of Formula (I) is undergoes phase II metabolism. In some embodiments, a compound of Formula (I) is glucuronidated. In some embodiments, a compound of Formula (I) forms a sulfate conjugate.

Compounds of Formula (II)

[0020] In some embodiments, described herein is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof:



Formula (II);

wherein:

R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of hydrogen, deuterium, hydroxy, and methoxy; wherein at least one of R₁, R₂, R₃, and R₄ is deuterium.

[0021] In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of hydrogen and deuterium. In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of deuterium, hydroxy, and methoxy. In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of deuterium and hydroxy. In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein at least two of R₁, R₂, R₃, and R₄ is deuterium. In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein at least three of R₁, R₂, R₃, and R₄ is deuterium. In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein at least one of R₁, R₂, R₃, and R₄ is hydroxy. In some embodiments is a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are deuterium.

[0022] In some embodiments, a compound of Formula (II) is undergoes phase II metabolism. In some embodiments, a compound of Formula (II) is glucuronidated. In some embodiments, a compound of Formula (II) forms a sulfate conjugate.

[0023] In some embodiments, Compound 1 includes the solvent addition forms (solvates). In some embodiments, a compound of Formula (I) or (II) described herein includes the solvent addition forms (solvates). Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and are formed during the process of product formation or isolation with pharmaceutically acceptable solvents such as water, ethanol, methanol, tert-butyl methyl ether (MTBE), diisopropyl ether (DIPE), ethyl acetate, isopropyl acetate, isopropyl alcohol, methyl isobutyl ketone (MIBK), methyl ethyl ketone (MEK), acetone, nitromethane, tetrahydrofuran


(THF), dichloromethane (DCM), dioxane, heptanes, toluene, anisole, acetonitrile, and the like. In some embodiments, solvates are formed using, but not limited to, Class 3 solvent(s). In some embodiments, solvates are formed using, but not limited to, Class 2 solvent(s). Categories of solvents are defined in, for example, the International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH), "Impurities: Guidelines for Residual Solvents Q3C(R6)," (October 2016). Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol.

[0024] In some embodiments, Compound 1 includes a pharmaceutically acceptable salt. In some embodiments, a compound of Formula (I) or (II) described herein includes a pharmaceutically acceptable salt.

[0025] In other embodiments, Compound 1 is prepared in various forms, including but not limited to, an amorphous phase, crystalline forms, milled forms, and nano-particulate forms. In other embodiments, a compound of Formula (I) or (II) is prepared in various forms, including but not limited to, an amorphous phase, crystalline forms, milled forms, and nano-particulate forms.


[0026] While not intending to be bound by any particular theory, certain solid forms are characterized by physical properties, *e.g.*, stability, solubility, and dissolution rate, appropriate for pharmaceutical and therapeutic dosage forms. Moreover, while not wishing to be bound by any particular theory, certain solid forms are characterized by physical properties (*e.g.*, density, compressibility, hardness, morphology, cleavage, stickiness, solubility, water uptake, electrical properties, thermal behavior, solid-state reactivity, physical stability, and chemical stability) affecting particular processes (*e.g.*, yield, filtration, washing, drying, milling, mixing, tableting, flowability, dissolution, formulation, and lyophilization) which make certain solid forms suitable for the manufacture of a solid dosage form. Such properties can be determined using particular analytical chemical techniques, including solid-state analytical techniques (*e.g.*, X-ray diffraction, microscopy, spectroscopy and thermal analysis), as described herein.


[0027] In some embodiments, is a compound selected from:




; or a pharmaceutically acceptable salt or solvate thereof.

[0028] In some embodiments, is a compound selected from:







[0029] In some embodiments, is a compound with the following structure:



[0030] In some embodiments, is a compound selected from:




[0031] In some embodiments, is a compound selected from:



, and ; or a pharmaceutically acceptable salt or solvate thereof.

[0032] In some embodiments, is a compound selected from:



solvate thereof.

Certain Terminology

[0033] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which the claimed subject matter belongs. It is to be understood that the foregoing general description and the following detailed

description are exemplary and explanatory only and are not restrictive of any subject matter claimed. In this application, the use of the singular includes the plural unless specifically stated otherwise. It must be noted that, as used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. In this application, the use of “or” means “and/or” unless stated otherwise. Furthermore, use of the term “including” as well as other forms, such as “include”, “includes,” and “included,” is not limiting. The term "comprising" (and related terms such as "comprise" or "comprises" or "having" or "including") is not intended to exclude that in other certain embodiments, for example, an embodiment of any composition of matter, composition, method, or process, or the like, described herein, may "consist of" or "consist essentially of" the described features. The term "about" when referring to a number or a numerical range means that the number or numerical range referred to is an approximation within experimental variability (or within statistical experimental error), and thus the number or numerical range may vary between 1% and 15% of the stated number or numerical range.

[0034] The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All documents, or portions of documents, cited in the application including, but not limited to, patents, patent applications, articles, books, manuals, and treatises are hereby expressly incorporated by reference in their entirety.

[0035] The term “acceptable” or “pharmaceutically acceptable”, with respect to a formulation, composition or ingredient, as used herein, means having no persistent detrimental effect on the general health of the subject being treated or does not abrogate the biological activity or properties of the compound, and is relatively nontoxic.

[0036] As used herein, “amelioration” of the symptoms of a particular disease, disorder, or condition by administration of a particular compound or pharmaceutical composition refers to any lessening of severity, delay in onset, slowing of progression, or shortening of duration, whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the compound or composition.

[0037] “Bioavailability” refers to the percentage of Compound 1 dosed that is delivered into the general circulation of the animal or human being studied. The total exposure ($AUC_{(0-\infty)}$) of a drug when administered intravenously is usually defined as 100% bioavailable (F%). “Oral bioavailability” refers to the extent to which Compound 1 is absorbed into the general circulation when the pharmaceutical composition is taken orally as compared to intravenous injection.

[0038] “Blood plasma concentration” refers to the concentration of Compound 1 in the plasma component of blood of a subject. It is understood that the plasma concentration of Compound 1

may vary significantly between subjects, due to variability with respect to metabolism and/or possible interactions with other therapeutic agents. In accordance with one embodiment disclosed herein, the blood plasma concentration of Compound 1 may vary from subject to subject.

Likewise, values such as maximum plasma concentration ($C_{\max}$) or time to reach maximum plasma concentration ($T_{\max}$), or total area under the plasma concentration time curve ($AUC_{(0-\infty)}$) may vary from subject to subject. Due to this variability, the amount necessary to constitute “a therapeutically effective amount” of Compound 1 may vary from subject to subject.

[0039] The terms “co-administration” or the like, as used herein, are meant to encompass administration of the selected therapeutic agents to a single patient, and are intended to include treatment regimens in which the agents are administered by the same or different route of administration or at the same or different time.

[0040] The terms “effective amount” or “therapeutically effective amount,” as used herein, refer to a sufficient amount of an agent or a compound being administered which will relieve to some extent one or more of the symptoms of the disease or condition being treated. The result can be reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an “effective amount” for therapeutic uses is the amount of the composition including a compound as disclosed herein required to provide a clinically significant decrease in disease symptoms without undue adverse side effects. An appropriate “effective amount” in any individual case may be determined using techniques, such as a dose escalation study. The term “therapeutically effective amount” includes, for example, a prophylactically effective amount. An “effective amount” of a compound disclosed herein is an amount effective to achieve a desired pharmacologic effect or therapeutic improvement without undue adverse side effects. It is understood that “an effect amount” or “a therapeutically effective amount” can vary from subject to subject, due to variation in metabolism of Compound 1, age, weight, general condition of the subject, the condition being treated, the severity of the condition being treated, and the judgment of the prescribing physician. By way of example only, therapeutically effective amounts may be determined by a dose escalation clinical trial.

[0041] The terms “enhance” or “enhancing” means to increase or prolong either in potency or duration a desired effect. By way of example, “enhancing” the effect of therapeutic agents refers to the ability to increase or prolong, either in potency or duration, the effect of therapeutic agents on during treatment of a disease, disorder, or condition. An “enhancing-effective amount,” as used herein, refers to an amount adequate to enhance the effect of a therapeutic agent in the treatment of a disease, disorder, or condition. When used in a patient, amounts effective for this use will depend on the severity and course of the disease, disorder, or condition, previous

therapy, the patient's health status and response to the drugs, and the judgment of the treating physician.

[0042] The term “prophylactically effective amount,” as used herein, refers that amount of a composition applied to a patient which will relieve to some extent one or more of the symptoms of a disease, condition or disorder being treated. In such prophylactic applications, such amounts may depend on the patient's state of health, weight, and the like. As an example, one can determine such prophylactically effective amounts by a dose escalation clinical trial.

[0043] "Pharmaceutically acceptable salt" includes both acid and base addition salts. A pharmaceutically acceptable salt of any one of the compounds described herein is intended to encompass any and all pharmaceutically suitable salt forms. Preferred pharmaceutically acceptable salts of the compounds described herein are pharmaceutically acceptable acid addition salts, and pharmaceutically acceptable base addition salts.

[0044] "Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, hydroiodic acid, hydrofluoric acid, phosphorous acid, and the like. Also included are salts that are formed with organic acids such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanoic acids, hydroxy alkanoic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc. and include, for example, acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid, salicylic acid, and the like. Exemplary salts thus include sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, nitrates, phosphates, monohydrogenphosphates, dihydrogenphosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, acetates, trifluoroacetates, propionates, caprylates, isobutyrate, oxalates, malonates, succinate suberates, sebacates, fumarates, maleates, mandelates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, phthalates, benzenesulfonates, toluenesulfonates, phenylacetates, citrates, lactates, malates, tartrates, methanesulfonates, and the like. Also contemplated are salts of amino acids, such as arginates, gluconates, and galacturonates (see, for example, Berge S.M. et al., "Pharmaceutical Salts," *Journal of Pharmaceutical Science*, 66:1-19 (1997)). Acid addition salts of basic compounds are prepared by contacting the free base forms with a sufficient amount of the desired acid to produce the salt.

[0045] "Pharmaceutically acceptable base addition salt" refers to those salts that retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise

undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. In some embodiments, pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Salts derived from inorganic bases include, but are not limited to, sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts, and the like. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, for example, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, diethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, *N,N*-dibenzylethylenediamine, chloroprocaine, hydrabamine, choline, betaine, ethylenediamine, ethylenedianiline, *N*-methylglucamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, *N*-ethylpiperidine, polyamine resins, and the like. See Berge et al., *supra*.

[0046] The term “subject” as used herein, refers to an animal which is the object of treatment, observation or experiment. By way of example only, a subject may be, but is not limited to, a mammal including, but not limited to, a human.

[0047] As used herein, the term “target activity” refers to a biological activity capable of being modulated by a selective modulator. Certain exemplary target activities include, but are not limited to, binding affinity, signal transduction, enzymatic activity, tumor growth, inflammation or inflammation-related processes, and amelioration of one or more symptoms associated with a disease or condition.

[0048] The terms “treat,” “treating” or “treatment”, as used herein, include alleviating, abating or ameliorating a disease or condition symptoms, preventing additional symptoms, ameliorating or preventing the underlying metabolic causes of symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition. The terms “treat,” “treating” or “treatment”, include, but are not limited to, prophylactic and/or therapeutic treatments.

[0049] As used herein, IC_{50} refers to a dosage, concentration or amount of a particular test compound that elicits a dose-dependent response at 50% of maximal expression of a particular response that is induced, provoked or potentiated by the particular test compound.

Pharmaceutical Compositions/Formulations

[0050] Pharmaceutical compositions may be formulated in a conventional manner using one or more physiologically acceptable carriers including excipients and auxiliaries which facilitate

processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions described herein may be found, for example, in *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999), herein incorporated by reference in their entirety.

[0051] A pharmaceutical composition, as used herein, refers to a mixture of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), a compound of Formula (I), or a compound of Formula (II) with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients. The pharmaceutical composition facilitates administration of the compound to a mammal. In practicing the methods of treatment or use provided herein, therapeutically effective amounts of Compound 1, a compound of Formula (I), or a compound of Formula (II) are administered in a pharmaceutical composition to a mammal having a disease, disorder, or condition to be treated. Preferably, the mammal is a human. A therapeutically effective amount can vary widely depending on the severity of the disease, the age and relative health of the subject, the potency of the compound used and other factors. The compounds can be used singly or in combination with one or more therapeutic agents as components of mixtures.

[0052] In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients.

[0053] In some embodiments is a pharmaceutical composition comprising a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients.

[0054] In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, for the treatment or prevention of hyperuricemia or gout, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients. In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-

hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, for the treatment or prevention of hyperuricemia, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients. In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, for the treatment of hyperuricemia, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients. In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, for the prevention of hyperuricemia, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients. In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, for the treatment or prevention of gout, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients. In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, for the treatment of gout, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients. In some embodiments is a pharmaceutical composition comprising (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone (Compound 1), a compound of Formula (I), or a compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof, for the prevention of gout. In some embodiments, the therapeutically effective amount is from about 3 mg to about 1500 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 1250 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 1000 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 750 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 600 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 500 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 400

[illegible]

mg to about 30 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 25 mg.

[0055] The term “pharmaceutical combination” as used herein, means a product that results from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term “fixed combination” means that the active ingredients, e.g. Compound 1, and a co-agent, are both administered to a patient simultaneously in the form of a single entity or dosage. The term “non-fixed combination” means that the active ingredients, e.g. Compound 1, and a co-agent, are administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific intervening time limits, wherein such administration provides effective levels of the two compounds in the body of the patient. The latter also applies to cocktail therapy, e.g. the administration of three or more active ingredients.

[0056] In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) is incorporated into pharmaceutical compositions to provide solid oral dosage forms. In other embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) is used to prepare pharmaceutical compositions other than oral solid dosage forms. The pharmaceutical formulations described herein can be administered to a subject by multiple administration routes, including but not limited to, oral, parenteral (e.g., intravenous, subcutaneous, intramuscular), intranasal, buccal, topical, rectal, or transdermal administration routes. The pharmaceutical formulations described herein include, but are not limited to, aqueous liquid dispersions, self-emulsifying dispersions, solid solutions, liposomal dispersions, aerosols, solid dosage forms, powders, immediate release formulations, controlled release formulations, fast melt formulations, tablets, capsules, pills, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate and controlled release formulations.

[0057] Pharmaceutical compositions including a compound described herein may be manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

Dosage Forms

[0058] The pharmaceutical compositions described herein can be formulated for administration to a mammal via any conventional means including, but not limited to, oral, parenteral (e.g., intravenous, subcutaneous, or intramuscular), buccal, intranasal, rectal, or transdermal administration routes. As used herein, the term “subject” or “individual” is used to mean an

animal, preferably a mammal, including a human or non-human. The terms individual, patient and subject may be used interchangeably.

[0059] Moreover, the pharmaceutical compositions described herein, which include Compound 1, a compound of Formula (I), or a compound of Formula (II), can be formulated into any suitable dosage form, including but not limited to, solid oral dosage forms, controlled release formulations, fast melt formulations, effervescent formulations, tablets, powders, pills, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and controlled release formulations.

[0060] Pharmaceutical preparations for oral use can be obtained by mixing one or more solid excipients with one or more of the compounds described herein, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for example, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methylcellulose, microcrystalline cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose; or others such as: polyvinylpyrrolidone (PVP or povidone) or calcium phosphate. If desired, disintegrating agents may be added, such as the cross-linked croscarmellose sodium, polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

[0061] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration.

[0062] In some embodiments, the solid dosage forms disclosed herein may be in the form of a tablet, (including a suspension tablet, a fast-melt tablet, a bite-disintegration tablet, a rapid-disintegration tablet, an effervescent tablet, or a caplet), a pill, a powder (including a sterile packaged powder, a dispensable powder, or an effervescent powder) a capsule (including both soft or hard capsules, e.g., capsules made from animal-derived gelatin or plant-derived HPMC, or “sprinkle capsules”), solid dispersion, solid solution, bioerodible dosage form, controlled release formulations, pulsatile release dosage forms, multiparticulate dosage forms, pellets, granules, or an aerosol. In other embodiments, the pharmaceutical formulation is in the form of a powder. In

still other embodiments, the pharmaceutical formulation is in the form of a tablet, including but not limited to, a fast-melt tablet. Additionally, pharmaceutical formulations described herein may be administered as a single capsule or in multiple capsule dosage form. In some embodiments, the pharmaceutical formulation is administered in two, or three, or four, capsules or tablets.

[0063] In some embodiments, solid dosage forms, e.g., tablets, effervescent tablets, and capsules, are prepared by mixing particles of Compound 1, a compound of Formula (I), or a compound of Formula (II) with one or more pharmaceutical excipients to form a bulk blend composition.

When referring to these bulk blend compositions as homogeneous, it is meant that the particles of Compound 1, a compound of Formula (I), or a compound of Formula (II) are dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms, such as tablets, pills, and capsules. The individual unit dosages may also include film coatings, which disintegrate upon oral ingestion or upon contact with diluent. These formulations can be manufactured by conventional pharmacological techniques.

[0064] Conventional pharmacological techniques include, e.g., one or a combination of methods: (1) dry mixing, (2) direct compression, (3) milling, (4) dry or non-aqueous granulation, (5) wet granulation, or (6) fusion. See, e.g., Lachman et al., *The Theory and Practice of Industrial Pharmacy* (1986). Other methods include, e.g., spray drying, pan coating, melt granulation, granulation, fluidized bed spray drying or coating (e.g., wurster coating), tangential coating, top spraying, tableting, extruding and the like.

[0065] The pharmaceutical solid dosage forms described herein can include Compound 1, a compound of Formula (I), or a compound of Formula (II), and one or more pharmaceutically acceptable additives such as a compatible carrier, binder, filling agent, suspending agent, flavoring agent, sweetening agent, disintegrating agent, dispersing agent, surfactant, lubricant, colorant, diluent, solubilizer, moistening agent, plasticizer, stabilizer, penetration enhancer, wetting agent, anti-foaming agent, antioxidant, preservative, or one or more combination thereof. In still other aspects, using standard coating procedures, such as those described in *Remington's Pharmaceutical Sciences*, 20th Edition (2000), a film coating is provided around the formulation of Compound 1, a compound of Formula (I), or a compound of Formula (II). In one embodiment, some or all of the particles of the Compound 1, the compound of Formula (I), or the compound of Formula (II) are coated. In another embodiment, some or all of the particles of the Compound 1, the compound of Formula (I), or the compound of Formula (II) are microencapsulated. In still another embodiment, the particles of the Compound 1, the compound of Formula (I), or the compound of Formula (II) are not microencapsulated and are uncoated.

[0066] Suitable carriers for use in the solid dosage forms described herein include, but are not limited to, acacia, gelatin, colloidal silicon dioxide, calcium glycerophosphate, calcium lactate, maltodextrin, glycerine, magnesium silicate, sodium caseinate, soy lecithin, sodium chloride, tricalcium phosphate, dipotassium phosphate, sodium stearyl lactylate, carrageenan, monoglyceride, diglyceride, pregelatinized starch, hydroxypropylmethylcellulose, hydroxypropylmethylcellulose acetate stearate, sucrose, microcrystalline cellulose, lactose, mannitol, and the like.

[0067] Suitable filling agents for use in the solid dosage forms described herein include, but are not limited to, lactose, calcium carbonate, calcium phosphate, dibasic calcium phosphate, calcium sulfate, microcrystalline cellulose, cellulose powder, dextrose, dextrans, dextrans, starches, pregelatinized starch, hydroxypropylmethylcellulose (HPMC), hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose acetate stearate (HPMCAS), sucrose, xylitol, lactitol, mannitol, sorbitol, sodium chloride, polyethylene glycol, and the like.

[0068] In order to release the Compound 1, the compound of Formula (I), or the compound of Formula (II) from a solid dosage form matrix as efficiently as possible, disintegrants are often used in the formulation, especially when the dosage forms are compressed with binder. Disintegrants help rupturing the dosage form matrix by swelling or capillary action when moisture is absorbed into the dosage form. Suitable disintegrants for use in the solid dosage forms described herein include, but are not limited to, natural starch such as corn starch or potato starch, a pregelatinized starch such as National 1551 or Amijel[®], or sodium starch glycolate such as Promogel[®] or Explotab[®], a cellulose such as a wood product, methylcrystalline cellulose, e.g., Avicel[®], Avicel[®] PH101, Avicel[®] PH102, Avicel[®] PH105, Elcema[®] P100, Emcocel[®], Vivacel[®], Ming Tia[®], and Solka-Floc[®], methylcellulose, croscarmellose, or a cross-linked cellulose, such as cross-linked sodium carboxymethylcellulose (Ac-Di-Sol[®]), cross-linked carboxymethylcellulose, or cross-linked croscarmellose, a cross-linked starch such as sodium starch glycolate, a cross-linked polymer such as crospovidone, a cross-linked polyvinylpyrrolidone, alginate such as alginic acid or a salt of alginic acid such as sodium alginate, a clay such as Veegum[®] HV (magnesium aluminum silicate), a gum such as agar, guar, locust bean, Karaya, pectin, or tragacanth, sodium starch glycolate, bentonite, a natural sponge, a surfactant, a resin such as a cation-exchange resin, citrus pulp, sodium lauryl sulfate, sodium lauryl sulfate in combination starch, and the like. In some embodiments provided herein, the disintegrating agent is selected from the group consisting of natural starch, a pregelatinized starch, a sodium starch, methylcrystalline cellulose, methylcellulose, croscarmellose, croscarmellose sodium, cross-

linked sodium carboxymethylcellulose, cross-linked carboxymethylcellulose, cross-linked croscarmellose, cross-linked starch such as sodium starch glycolate, cross-linked polymer such as crospovidone, cross-linked polyvinylpyrrolidone, sodium alginate, a clay, or a gum. In some embodiments provided herein, the disintegrating agent is croscarmellose sodium.

[0069] Binders impart cohesiveness to solid oral dosage form formulations: for powder filled capsule formulation, they aid in plug formation that can be filled into soft or hard shell capsules and for tablet formulation, they ensure the tablet remaining intact after compression and help assure blend uniformity prior to a compression or fill step. Materials suitable for use as binders in the solid dosage forms described herein include, but are not limited to, carboxymethylcellulose, methylcellulose (e.g., Methocel[®]), hydroxypropylmethylcellulose (e.g. Hypromellose USP Pharmaccoat-603, hydroxypropylmethylcellulose acetate stearate (Aquate HS-LF and HS), hydroxyethylcellulose, hydroxypropylcellulose (e.g., Klucel[®]), ethylcellulose (e.g., Ethocel[®]), and microcrystalline cellulose (e.g., Avicel[®]), microcrystalline dextrose, amylose, magnesium aluminum silicate, polysaccharide acids, bentonites, gelatin, polyvinylpyrrolidone/vinyl acetate copolymer, crospovidone, povidone, starch, pregelatinized starch, tragacanth, dextrin, a sugar, such as sucrose (e.g., Dipac[®]), glucose, dextrose, molasses, mannitol, sorbitol, xylitol (e.g., Xylitab[®]), lactose, a natural or synthetic gum such as acacia, tragacanth, ghatti gum, mucilage of isapol husks, starch, polyvinylpyrrolidone (e.g., Povidone[®] CL, Kollidon[®] CL, Polyplasdone[®] XL-10, and Povidone[®] K-12), larch arabogalactan, Veegum[®], polyethylene glycol, waxes, sodium alginate, and the like.

[0070] In general, binder levels of 20-70% are used in powder-filled gelatin capsule formulations. Binder usage level in tablet formulations varies whether direct compression, wet granulation, roller compaction, or usage of other excipients such as fillers which itself can act as moderate binder. Formulators skilled in art can determine the binder level for the formulations, but binder usage level of up to 70% in tablet formulations is common.

[0071] Suitable lubricants or glidants for use in the solid dosage forms described herein include, but are not limited to, stearic acid, calcium hydroxide, talc, corn starch, sodium stearyl fumarate, alkali-metal and alkaline earth metal salts, such as calcium, magnesium, stearic acid, sodium stearates, magnesium stearate, zinc stearate, waxes, Stearowet[®], boric acid, sodium benzoate, sodium acetate, sodium chloride, leucine, a polyethylene glycol or a methoxypolyethylene glycol such as Carbowax[™], PEG 4000, PEG 5000, PEG 6000, propylene glycol, sodium oleate, glyceryl behenate, glyceryl palmitostearate, glyceryl benzoate, magnesium or sodium lauryl sulfate, and the like. In some embodiments provided herein, the lubricant is selected from the group consisting of stearic acid, calcium hydroxide, talc, corn starch, sodium stearyl fumarate,

stearic acid, sodium stearates, magnesium stearate, zinc stearate, and waxes. In some embodiments provided herein, the lubricant is magnesium stearate.

[0072] Suitable diluents for use in the solid dosage forms described herein include, but are not limited to, sugars (including lactose, sucrose, and dextrose), polysaccharides (including dextrans and maltodextrin), polyols (including mannitol, xylitol, and sorbitol), cyclodextrins and the like. In some embodiments provided herein, the diluent is selected from the group consisting of lactose, sucrose, dextrose, dextrans, maltodextrin, mannitol, xylitol, sorbitol, cyclodextrins, calcium phosphate, calcium sulfate, starches, modified starches, microcrystalline cellulose, microcellulose, and talc. In some embodiments provided herein, the diluent is microcrystalline cellulose.

[0073] The term “non water-soluble diluent” represents compounds typically used in the formulation of pharmaceuticals, such as calcium phosphate, calcium sulfate, starches, modified starches, microcrystalline cellulose, microcellulose (e.g., having a density of about 0.45 g/cm³, e.g. Avicel, powdered cellulose), and talc.

[0074] Suitable wetting agents for use in the solid dosage forms described herein include, for example, oleic acid, glyceryl monostearate, sorbitan monooleate, sorbitan monolaurate, triethanolamine oleate, polyoxyethylene sorbitan monooleate, polyoxyethylene sorbitan monolaurate, quaternary ammonium compounds (e.g., Polyquat 10[®]), sodium oleate, sodium lauryl sulfate, magnesium stearate, sodium docusate, triacetin, vitamin E TPGS, and the like.

[0075] Suitable surfactants for use in the solid dosage forms described herein include, for example, sodium lauryl sulfate, sorbitan monooleate, polyoxyethylene sorbitan monooleate, polysorbates, polaxomers, bile salts, glyceryl monostearate, copolymers of ethylene oxide and propylene oxide, e.g., Pluronic[®] (BASF), and the like. In some embodiments provided herein, the surfactant is selected from the group consisting of sodium lauryl sulfate, sorbitan monooleate, polyoxyethylene sorbitan monooleate, polysorbates, polaxomers, bile salts, glyceryl monostearate, copolymers of ethylene oxide and propylene oxide. In some embodiments provided herein, the surfactant is sodium lauryl sulfate.

[0076] Suitable suspending agents for use in the solid dosage forms described here include, but are not limited to, polyvinylpyrrolidone, e.g., polyvinylpyrrolidone K12, polyvinylpyrrolidone K17, polyvinylpyrrolidone K25, or polyvinylpyrrolidone K30, polyethylene glycol, e.g., the polyethylene glycol can have a molecular weight of about 300 to about 6000, or about 3350 to about 4000, or about 7000 to about 5400, vinyl pyrrolidone/vinyl acetate copolymer (S630), sodium carboxymethylcellulose, methylcellulose, hydroxy-propylmethylcellulose, polysorbate-80, hydroxyethylcellulose, sodium alginate, gums, such as, e.g., gum tragacanth and gum acacia,

guar gum, xanthans, including xanthan gum, sugars, cellulose, such as, e.g., sodium carboxymethylcellulose, methylcellulose, sodium carboxymethylcellulose, hydroxypropylmethylcellulose, hydroxyethylcellulose, polysorbate-80, sodium alginate, polyethoxylated sorbitan monolaurate, polyethoxylated sorbitan monolaurate, povidone, and the like.

[0077] Suitable antioxidants for use in the solid dosage forms described herein include, for example, e.g., butylated hydroxytoluene (BHT), sodium ascorbate, and tocopherol.

[0078] It should be appreciated that there is considerable overlap between additives used in the solid dosage forms described herein. Thus, the above-listed additives should be taken as merely exemplary, and not limiting, of the types of additives that can be included in solid dosage forms described herein. The amounts of such additives can be readily determined by one skilled in the art, according to the particular properties desired.

[0079] In other embodiments, one or more layers of the pharmaceutical formulation are plasticized. Illustratively, a plasticizer is generally a high boiling point solid or liquid. Suitable plasticizers can be added from about 0.01% to about 50% by weight (w/w) of the coating composition. Plasticizers include, but are not limited to, diethyl phthalate, citrate esters, polyethylene glycol, glycerol, acetylated glycerides, triacetin, polypropylene glycol, polyethylene glycol, triethyl citrate, dibutyl sebacate, stearic acid, stearyl, stearate, and castor oil.

[0080] Compressed tablets are solid dosage forms prepared by compacting the bulk blend of the formulations described above. In various embodiments, compressed tablets which are designed to dissolve in the mouth will include one or more flavoring agents. In other embodiments, the compressed tablets will include a film surrounding the final compressed tablet. In some embodiments, the film coating can provide a delayed release of Compound 1, compound of Formula (I), or compound of Formula (II) from the formulation. In other embodiments, the film coating aids in patient compliance (e.g., Opadry[®] coatings or sugar coating). Film coatings including Opadry[®] typically range from about 1% to about 3% of the tablet weight. In other embodiments, the compressed tablets include one or more excipients.

[0081] A capsule may be prepared, for example, by placing the bulk blend of the formulation of Compound 1, compound of Formula (I), or compound of Formula (II) inside of a capsule. In some embodiments, the formulations (non-aqueous suspensions and solutions) are placed in a soft gelatin capsule. In some embodiments, the formulations (non-aqueous suspensions and solutions) are placed in a hard shell gelatin capsule. In other embodiments, the formulations are placed in standard gelatin capsules or non-gelatin capsules such as capsules comprising HPMC. In other embodiments, the formulation is placed in a sprinkle capsule, wherein the capsule may

be swallowed whole or the capsule may be opened and the contents sprinkled on food prior to eating. In some embodiments, the therapeutic dose is split into multiple (e.g., two, three, or four) capsules. In some embodiments, the entire dose of the formulation is delivered in a capsule form.

[0082] In various embodiments, the particles of Compound 1, compound of Formula (I), or compound of Formula (II) and one or more excipients are dry blended and compressed into a mass, such as a tablet, having a hardness sufficient to provide a pharmaceutical composition that substantially disintegrates within less than about 30 minutes, less than about 35 minutes, less than about 40 minutes, less than about 45 minutes, less than about 50 minutes, less than about 55 minutes, or less than about 60 minutes, after oral administration, thereby releasing the formulation into the gastrointestinal fluid.

[0083] In another aspect, dosage forms may include microencapsulated formulations. In some embodiments, one or more other compatible materials are present in the microencapsulation material. Exemplary materials include, but are not limited to, pH modifiers, erosion facilitators, anti-foaming agents, antioxidants, flavoring agents, and carrier materials such as binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, and diluents.

[0084] Materials useful for the microencapsulation described herein include materials compatible with Compound 1, compound of Formula (I), or compound of Formula (II) which sufficiently isolate Compound 1, compound of Formula (I), or compound of Formula (II) from other non-compatible excipients. Materials compatible with Compound 1, compound of Formula (I), or compound of Formula (II) are those that delay the release of the compounds *in vivo*.

[0085] Exemplary microencapsulation materials useful for delaying the release of the formulations including compounds described herein, include, but are not limited to, hydroxypropyl cellulose ethers (HPC) such as Klucel[®] or Nisso HPC, low-substituted hydroxypropyl cellulose ethers (L-HPC), hydroxypropyl methyl cellulose ethers (HPMC) such as Seppifilm-LC, Pharmacoat[®], Metolose SR, Methocel[®]-E, Opadry YS, PrimaFlo, Benecel MP824, and Benecel MP843, methylcellulose polymers such as Methocel[®]-A, hydroxypropylmethylcellulose acetate stearate Aqoat (HF-LS, HF-LG, HF-MS) and Metolose[®], Ethylcelluloses (EC) and mixtures thereof such as E461, Ethocel[®], Aqualon[®]-EC, Surelease[®], Polyvinyl alcohol (PVA) such as Opadry AMB, hydroxyethylcelluloses such as Natrosol[®], carboxymethylcelluloses and salts of carboxymethylcelluloses (CMC) such as Aqualon[®]-CMC, polyvinyl alcohol and polyethylene glycol co-polymers such as Kollicoat IR[®], monoglycerides (Myverol), triglycerides (KLX), polyethylene glycols, modified food starch, acrylic polymers and mixtures of acrylic polymers with cellulose ethers such as Eudragit[®] EPO, Eudragit[®] L30D-55,

Eudragit[®] FS 30D Eudragit[®] L100-55, Eudragit[®] L100, Eudragit[®] S100, Eudragit[®] RD100, Eudragit[®] E100, Eudragit[®] L12.5, Eudragit[®] S12.5, Eudragit[®] NE30D, and Eudragit[®] NE 40D, cellulose acetate phthalate, sepiifilms such as mixtures of HPMC and stearic acid, cyclodextrins, and mixtures of these materials.

[0086] In still other embodiments, plasticizers such as polyethylene glycols, e.g., PEG 300, PEG 400, PEG 600, PEG 1450, PEG 3350, and PEG 800, stearic acid, propylene glycol, oleic acid, and triacetin are incorporated into the microencapsulation material. In other embodiments, the microencapsulating material useful for delaying the release of the pharmaceutical compositions is from the USP or the National Formulary (NF). In yet other embodiments, the microencapsulation material is Klucel. In still other embodiments, the microencapsulation material is methocel.

[0087] Microencapsulated Compound 1, compound of Formula (I), or compound of Formula (II) may be formulated by several methods, illustrative examples of which include, e.g., spray drying processes, spinning disk-solvent processes, hot melt processes, spray chilling methods, fluidized bed, electrostatic deposition, centrifugal extrusion, rotational suspension separation, polymerization at liquid-gas or solid-gas interface, pressure extrusion, or spraying solvent extraction bath. In addition to these, several chemical techniques, e.g., complex coacervation, solvent evaporation, polymer-polymer incompatibility, interfacial polymerization in liquid media, in situ polymerization, in-liquid drying, and desolvation in liquid media could also be used. Furthermore, other methods such as roller compaction, extrusion/spheronization, coacervation, or nanoparticle coating may also be used.

[0088] In one embodiment, the particles of Compound 1, compound of Formula (I), or compound of Formula (II) are microencapsulated prior to being formulated into one of the above forms. In still another embodiment, some or most of the particles are coated prior to being further formulated by using standard coating procedures, such as those described in *Remington's Pharmaceutical Sciences*, 20th Edition (2000).

[0089] In other embodiments, the solid dosage formulations of Compound 1, compound of Formula (I), or compound of Formula (II) are plasticized (coated) with one or more layers. Illustratively, a plasticizer is generally a high boiling point solid or liquid. Suitable plasticizers can be added from about 0.01% to about 50% by weight (w/w) of the coating composition. Plasticizers include, but are not limited to, diethyl phthalate, citrate esters, polyethylene glycol, glycerol, acetylated glycerides, triacetin, polypropylene glycol, polyethylene glycol, triethyl citrate, dibutyl sebacate, stearic acid, stearyl, stearate, and castor oil.

[0090] In other embodiments, a powder including the formulations with Compound 1, compound of Formula (I), or compound of Formula (II) may be formulated to include one or more

pharmaceutical excipients and flavors. Such a powder may be prepared, for example, by mixing the formulation and optional pharmaceutical excipients to form a bulk blend composition.

Additional embodiments also include a suspending agent and/or a wetting agent. This bulk blend is uniformly subdivided into unit dosage packaging or multi-dosage packaging units.

[0091] In still other embodiments, effervescent powders are also prepared in accordance with the present disclosure. Effervescent salts have been used to disperse medicines in water for oral administration. Effervescent salts are granules or coarse powders containing a medicinal agent in a dry mixture, usually composed of sodium bicarbonate, citric acid and/or tartaric acid. When salts of the compositions described herein are added to water, the acids and the base react to liberate carbon dioxide gas, thereby causing “effervescence.” Examples of effervescent salts include, e.g., the following ingredients: sodium bicarbonate or a mixture of sodium bicarbonate and sodium carbonate, citric acid and/or tartaric acid. Any acid-base combination that results in the liberation of carbon dioxide can be used in place of the combination of sodium bicarbonate and citric and tartaric acids, as long as the ingredients were suitable for pharmaceutical use and result in a pH of about 6.0 or higher.

[0092] In some embodiments, the solid dosage forms described herein can be formulated as enteric coated delayed release oral dosage forms, i.e., as an oral dosage form of a pharmaceutical composition as described herein which utilizes an enteric coating to affect release in the small intestine of the gastrointestinal tract. The enteric coated dosage form may be a compressed or molded or extruded tablet/mold (coated or uncoated) containing granules, powder, pellets, beads or particles of the active ingredient and/or other composition components, which are themselves coated or uncoated. The enteric coated oral dosage form may also be a capsule (coated or uncoated) containing pellets, beads or granules of the solid carrier or the composition, which are themselves coated or uncoated.

[0093] The term “delayed release” as used herein refers to the delivery so that the release can be accomplished at some generally predictable location in the intestinal tract more distal to that which would have been accomplished if there had been no delayed release alterations. In some embodiments the method for delay of release is coating. Any coatings should be applied to a sufficient thickness such that the entire coating does not dissolve in the gastrointestinal fluids at pH below about 5, but does dissolve at pH about 5 and above. It is expected that any anionic polymer exhibiting a pH-dependent solubility profile can be used as an enteric coating in the methods and compositions described herein to achieve delivery to the lower gastrointestinal tract. In some embodiments the polymers described herein are anionic carboxylic polymers. In other

embodiments, the polymers and compatible mixtures thereof, and some of their properties, include, but are not limited to:

[0094] Shellac, also called purified lac, a refined product obtained from the resinous secretion of an insect. This coating dissolves in media of pH >7;

[0095] Acrylic polymers. The performance of acrylic polymers (primarily their solubility in biological fluids) can vary based on the degree and type of substitution. Examples of suitable acrylic polymers include methacrylic acid copolymers and ammonium methacrylate copolymers. The Eudragit series E, L, S, RL, RS and NE (Rohm Pharma) are available as solubilized in organic solvent, aqueous dispersion, or dry powders. The Eudragit series RL, NE, and RS are insoluble in the gastrointestinal tract but are permeable and are used primarily for colonic targeting. The Eudragit series E dissolve in the stomach. The Eudragit series L, L-30D and S are insoluble in the stomach and dissolve in the intestine;

[0096] Cellulose Derivatives. Examples of suitable cellulose derivatives are ethyl cellulose; and reaction mixtures of partial acetate esters of cellulose with phthalic anhydride. The performance can vary based on the degree and type of substitution. Cellulose acetate phthalate (CAP) dissolves in pH >6. Aquateric (FMC) is an aqueous based system and is a spray dried CAP psuedolatex with particles <1 μm . Other components in Aquateric can include pluronics, Tweens, and acetylated monoglycerides. Other suitable cellulose derivatives include: cellulose acetate trimellitate (Eastman); methylcellulose (Pharmacoat, Methocel); hydroxypropylmethyl cellulose phthalate (HPMCP); hydroxypropylmethyl cellulose succinate (HPMCS); and hydroxypropylmethylcellulose acetate succinate (e.g., AQOAT (Shin Etsu)). The performance can vary based on the degree and type of substitution. For example, HPMCP such as, HP-50, HP-55, HP-55S, or HP-55F grades are suitable. The performance can vary based on the degree and type of substitution. For example, suitable grades of hydroxypropylmethylcellulose acetate succinate include, but are not limited to, AS-LG (LF), which dissolves at pH 5, AS-MG (MF), which dissolves at pH 5.5, and AS-HG (HF), which dissolves at higher pH. These polymers are offered as granules, or as fine powders for aqueous dispersions; Poly Vinyl Acetate Phthalate (PVAP). PVAP dissolves in pH >5, and it is much less permeable to water vapor and gastric fluids.

[0097] In some embodiments, the coating can, and usually does, contain a plasticizer and possibly other coating excipients such as colorants, talc, and/or magnesium stearate. Suitable plasticizers include triethyl citrate (Citroflex 2), triacetin (glyceryl triacetate), acetyl triethyl citrate (Citroflex A2), Carbowax 400 (polyethylene glycol 400), diethyl phthalate, tributyl citrate, acetylated monoglycerides, glycerol, fatty acid esters, propylene glycol, and dibutyl phthalate. In

particular, anionic carboxylic acrylic polymers usually will contain 10-25% by weight of a plasticizer, especially dibutyl phthalate, polyethylene glycol, triethyl citrate and triacetin. Conventional coating techniques such as spray or pan coating are employed to apply coatings. The coating thickness must be sufficient to ensure that the oral dosage form remains intact until the desired site of topical delivery in the intestinal tract is reached.

[0098] Colorants, detackifiers, surfactants, antifoaming agents, lubricants (e.g., carnuba wax or PEG) may be added to the coatings besides plasticizers to solubilize or disperse the coating material, and to improve coating performance and the coated product.

[0099] In other embodiments, the formulations described herein, which include Compound 1, a compound of Formula (I), or a compound of Formula (II) are delivered using a pulsatile dosage form. A pulsatile dosage form is capable of providing one or more immediate release pulses at predetermined time points after a controlled lag time or at specific sites. Other types of controlled release systems may be used. Examples of such delivery systems include, e.g., polymer-based systems, such as polylactic and polyglycolic acid, polyanhydrides and polycaprolactone; porous matrices, nonpolymer-based systems that are lipids, including sterols, such as cholesterol, cholesterol esters and fatty acids, or neutral fats, such as mono-, di- and triglycerides; hydrogel release systems; silastic systems; peptide-based systems; wax coatings, bioerodible dosage forms, compressed tablets using conventional binders and the like. See, e.g., Liberman et al., *Pharmaceutical Dosage Forms*, 2 Ed., Vol. 1, pp. 209-214 (1990); Singh et al., *Encyclopedia of Pharmaceutical Technology*, 2nd Ed., pp. 751-753 (2002); U.S. Pat. Nos. 4,327,725, 4,624,848, 4,968,509, 5,461,140, 5,456,923, 5,516,527, 5,622,721, 5,686,105, 5,700,410, 5,977,175, 6,465,014 and 6,932,983, each of which is specifically incorporated by reference.

[00100] In some embodiments, pharmaceutical formulations are provided that include particles of Compound 1, a compound of Formula (I), or a compound of Formula (II) and at least one dispersing agent or suspending agent for oral administration to a subject. The formulations may be a powder and/or granules for suspension and, upon admixture with water, a substantially uniform suspension is obtained.

[00101] It is to be appreciated that there is overlap between the above-listed additives used in the aqueous dispersions or suspensions described herein, since a given additive is often classified differently by different practitioners in the field, or is commonly used for any of several different functions. Thus, the above-listed additives should be taken as merely exemplary, and not limiting, of the types of additives that can be included in formulations described herein. The amounts of such additives can be readily determined by one skilled in the art, according to the particular properties desired.

Methods

[00102] In some embodiments is a method for treating hyperuricemia or gout comprising administering to the individual in need thereof a therapeutically effective amount of Compound 1, or a pharmaceutically acceptable salt or solvate thereof, described herein. In some embodiments is a method for the treatment of diseases or conditions that would benefit from lowering serum uric acid (sUA) in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating or preventing hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for preventing hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating or preventing hyperuricemia in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating hyperuricemia in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for preventing hyperuricemia in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating or preventing gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for

treating gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for preventing gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating hyperuricemia or gout comprising administering to the individual in need thereof a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof, described herein. In some embodiments is a method for the treatment of diseases or conditions that would benefit from lowering serum uric acid (sUA) in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating or preventing hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for preventing hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating or preventing hyperuricemia in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating hyperuricemia in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for preventing hyperuricemia in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for treating or preventing gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some

embodiments is a method for treating gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments is a method for preventing gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of Formula (I) or (II), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the therapeutically effective amount is from about 3 mg to about 1500 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 1250 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 1000 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 750 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 600 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 500 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 400 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 300 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 250 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 200 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 150 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 100 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 75 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 50 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 45 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 40 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 35 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 30 mg. In some embodiments, the therapeutically effective amount is from about 3 mg to about 25 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 300 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 250 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 200 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 150 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 100 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 75 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 50 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 45 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 40 mg. In some

embodiments, the therapeutically effective amount is from about 5 mg to about 35 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 30 mg. In some embodiments, the therapeutically effective amount is from about 5 mg to about 25 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 200 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 150 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 100 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 75 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 50 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 45 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 40 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 35 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 30 mg. In some embodiments, the therapeutically effective amount is from about 10 mg to about 25 mg.

Methods of Dosing and Treatment Regimens

[00103] In some embodiments, Compound 1 is used in the preparation of medicaments for the treatment of diseases or conditions that would benefit from lowering serum uric acid (sUA). In addition, a method for treating any of the diseases or conditions described herein in an individual in need of such treatment, involves administration of pharmaceutical compositions containing Compound 1, or pharmaceutically acceptable solvate thereof, in therapeutically effective amounts to said individual.

[00104] In some embodiments, compositions containing a compound of Formula (I) or (II) are administered for prophylactic, therapeutic, or maintenance treatment. In some embodiments, compositions containing a compound of Formula (I) or (II) are administered for therapeutic applications. In some embodiments, compositions containing a compound of Formula (I) or (II) are administered for prophylactic applications.

[00105] In some embodiments, compositions containing Compound 1 are administered for prophylactic, therapeutic, or maintenance treatment. In some embodiments, compositions containing Compound 1 are administered for therapeutic applications. In some embodiments, compositions containing Compound 1 are administered for prophylactic applications.

[00106] In some embodiments, compositions containing a compound of Formula (I) or (II) are administered for prophylactic, therapeutic, or maintenance treatment. In some embodiments, compositions containing a compound of Formula (I) or (II) are administered for therapeutic

applications. In some embodiments, compositions containing a compound of Formula (I) or (II) are administered for prophylactic applications.

[00107] In therapeutic applications, the compositions are administered to a patient already suffering from a disease or condition, in an amount sufficient to cure or at least partially arrest the symptoms of the disease or condition. Amounts effective for this use will depend on the severity and course of the disease or condition, previous therapy, the patient's health status, weight, and response to the drugs, and the judgment of the treating physician.

[00108] In prophylactic applications, compositions containing the compounds described herein are administered to a patient susceptible to or otherwise at risk of a particular disease, disorder, or condition. Such an amount is defined to be a "prophylactically effective amount or dose." In this use, the precise amounts also depend on the patient's state of health, weight, and the like. When used in a patient, effective amounts for this use will depend on the severity and course of the disease, disorder, or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician.

[00109] In some embodiments, Compound 1 is administered daily. In some embodiments, Compound 1 is administered every other day. In some embodiments, a compound of Formula (I) or (II) is administered daily. In some embodiments, a compound of Formula (I) or (II) is administered every other day.

[00110] In some embodiments, Compound 1 is administered once per day. In some embodiments, Compound 1 is administered twice per day. In some embodiments, Compound 1 is administered three times per day. In some embodiments, Compound 1 is administered four times per day. In some embodiments, a compound of Formula (I) or (II) is administered once per day. In some embodiments, a compound of Formula (I) or (II) is administered twice per day. In some embodiments, a compound of Formula (I) or (II) is administered three times per day. In some embodiments, a compound of Formula (I) or (II) is administered four times per day.

[00111] In the case wherein the patient's condition does not improve, upon the doctor's discretion the administration of the compounds may be administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disease or condition.

[00112] Once improvement of the patient's conditions has occurred, a maintenance dose is administered, if necessary. Subsequently, the dosage or the frequency of administration, or both, can be reduced, as a function of the symptoms, to a level at which the improved disease, disorder, or condition is retained. Patients can, however, require intermittent treatment on a long-term basis upon any recurrence of symptoms.

[00113] The amount of a given agent that will correspond to such an amount will vary depending upon factors such as the particular compound, disease or condition and its severity, the identity (e.g., weight) of the subject or host in need of treatment, but can nevertheless be determined in a manner recognized in the field according to the particular circumstances surrounding the case, including, e.g., the specific agent being administered, the route of administration, the condition being treated, and the subject or host being treated. In general, however, doses employed for adult human treatment will typically be in the range of about 0.02 - about 5000 mg per day, in some embodiments, about 1 – about 1500 mg per day. The desired dose may conveniently be presented in a single dose or as divided doses administered simultaneously (or over a short period of time) or at appropriate intervals, for example as two, three, four or more sub-doses per day.

[00114] The pharmaceutical composition described herein may be in unit dosage forms suitable for single administration of precise dosages. In unit dosage form, the formulation is divided into unit doses containing appropriate quantities of one or more compound. The unit dosage may be in the form of a package containing discrete quantities of the formulation. Non-limiting examples are packaged tablets or capsules, and powders in vials or ampoules. Aqueous suspension compositions can be packaged in single-dose non-reclosable containers. Alternatively, multiple-dose reclosable containers can be used, in which case it is typical to include a preservative in the composition. By way of example only, formulations for parenteral injection may be presented in unit dosage form, which include, but are not limited to ampoules, or in multi-dose containers, with an added preservative.

[00115] The daily dosages appropriate for the compounds described herein are from about 0.01 mg/kg to about 20 mg/kg. In one embodiment, the daily dosages are from about 0.1 mg/kg to about 10 mg/kg. An indicated daily dosage in the larger mammal, including, but not limited to, humans, is in the range from about 3 mg to about 1500 mg, conveniently administered in a single dose or in divided doses, including, but not limited to, up to four times a day or in extended release form. Suitable unit dosage forms for oral administration include from about 1 to about 500 mg active ingredient. In one embodiment, the unit dosage is about 1 mg, about 5 mg, about 10 mg, about 20 mg, about 50 mg, about 100 mg, about 200 mg, about 250 mg, about 400 mg, or about 500 mg. The foregoing ranges are merely suggestive, as the number of variables in regard to an individual treatment regime is large, and considerable excursions from these recommended values are not uncommon. Such dosages may be altered depending on a number of variables, not limited to the activity of the compound used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the practitioner.

[00116] Toxicity and therapeutic efficacy of such therapeutic regimens can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the determination of the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between the toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD₅₀ and ED₅₀. The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with minimal toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized.

Combination Treatments

[00117] Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, and pharmaceutical compositions thereof, may also be used in combination with other therapeutic agents that are selected for their therapeutic value for the condition to be treated. In general, the compositions described herein and, in embodiments where combinational therapy is employed, other agents do not have to be administered in the same pharmaceutical composition, and may, because of different physical and chemical characteristics, have to be administered by different routes. The determination of the mode of administration and the advisability of administration, where possible, in the same pharmaceutical composition, is well within the knowledge of the clinician. The initial administration can be made according to established protocols recognized in the field, and then, based upon the observed effects, the dosage, modes of administration and times of administration can be modified by the clinician.

[00118] In certain instances, it may be appropriate to administer Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, and pharmaceutical compositions thereof, in combination with another therapeutic agent. By way of example only, if one of the side effects experienced by a patient upon receiving one of the compounds herein, such as Compound 1, is nausea, then it may be appropriate to administer an anti-nausea agent in combination with the initial therapeutic agent. Or, by way of example only, the therapeutic effectiveness of one of the compounds described herein may be enhanced by administration of an adjuvant (i.e., by itself the adjuvant may have minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced). Or, by way of example only, the benefit experienced by a patient may be increased by administering one of the compounds described herein with another therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit. In any case, regardless of the disease,

disorder, or condition being treated, the overall benefit experienced by the patient may simply be additive of the two therapeutic agents or the patient may experience a synergistic benefit.

[00119] In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor, wherein the xanthine oxidase inhibitor is allopurinol, oxypurinol, febuxostat, topiroxostat, or inositol. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor, wherein the xanthine oxidase inhibitor is allopurinol. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor, wherein the xanthine oxidase inhibitor is oxypurinol. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor, wherein the xanthine oxidase inhibitor is febuxostat. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor, wherein the xanthine oxidase inhibitor is topiroxostat. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor, wherein the xanthine oxidase inhibitor is ositol.

[00120] In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, and the xanthine oxidase inhibitor are administered in combination in a single dosage form. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, and the xanthine oxidase inhibitor are administered in combination in separate dosage forms.

[00121] In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with an SGLT2 inhibitor. In some embodiments, Compound 1 is administered in combination with an SGLT2 inhibitor, wherein the SGLT2 inhibitor is canagliflozin, dapagliflozin, empagliflozin, empagliflozin/linagliptin, empagliflozin/metformin, or

dapagliflozin/metformin. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with an SGLT2 inhibitor, wherein the SGLT2 inhibitor is canagliflozin. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with an SGLT2 inhibitor, wherein the SGLT2 inhibitor is dapagliflozin. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with an SGLT2 inhibitor, wherein the SGLT2 inhibitor is empagliflozin. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with an SGLT2 inhibitor, wherein the SGLT2 inhibitor is empagliflozin/linagliptin. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with an SGLT2 inhibitor, wherein the SGLT2 inhibitor is empagliflozin/metformin. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with an SGLT2 inhibitor, wherein the SGLT2 inhibitor is dapagliflozin/metformin.

[00122] In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, and the SGLT2 inhibitor are administered in combination in a single dosage form. In some embodiments, Compound 1 and the SGLT2 inhibitor are administered in combination in separate dosage forms.

[00123] In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor and an SGLT2 inhibitor. In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, is administered in combination with a xanthine oxidase inhibitor and an SGLT2 inhibitor, wherein the xanthine oxidase inhibitor is allopurinol, oxypurinol, febuxostat, topiroxostat, or inositol, and the SGLT2 inhibitor is canagliflozin, dapagliflozin, empagliflozin, empagliflozin/linagliptin, empagliflozin/metformin, or dapagliflozin/metformin.

[00124] In some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, the xanthine oxidase inhibitor, and the SGLT2 inhibitor are administered in combination in a single dosage form. In

some embodiments, Compound 1, a compound of Formula (I), or a compound of Formula (II) described herein, or a pharmaceutical composition thereof, the xanthine oxidase inhibitor, and the SGLT2 inhibitor are administered in combination in separate dosage forms.

[00125] The particular choice of compounds used will depend upon the diagnosis of the attending physicians and their judgment of the condition of the patient and the appropriate treatment protocol. The compounds may be administered concurrently (e.g., simultaneously, essentially simultaneously or within the same treatment protocol) or sequentially, depending upon the nature of the disease, disorder, or condition, the condition of the patient, and the actual choice of compounds used. The determination of the order of administration, and the number of repetitions of administration of each therapeutic agent during a treatment protocol, is well within the knowledge of the physician after evaluation of the disease being treated and the condition of the patient.

[00126] Therapeutically-effective dosages can vary when the drugs are used in treatment combinations. Methods for experimentally determining therapeutically-effective dosages of drugs and other agents for use in combination treatment regimens are described in the literature. For example, the use of metronomic dosing, i.e., providing more frequent, lower doses in order to minimize toxic side effects, has been described extensively in the literature. Combination treatment further includes periodic treatments that start and stop at various times to assist with the clinical management of the patient.

[00127] For combination therapies described herein, dosages of the co-administered compounds will of course vary depending on the type of co-drug employed, on the specific drug employed, on the disease or condition being treated and so forth. In addition, when co-administered with one or more biologically active agents, the compound provided herein may be administered either simultaneously with the biologically active agent(s), or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein in combination with the biologically active agent(s).

[00128] In any case, the multiple therapeutic agents (one of which is Compound 1, a compound of Formula (I), a compound of Formula (II), or a pharmaceutical composition thereof) may be administered in any order or even simultaneously. If simultaneously, the multiple therapeutic agents may be provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills). One of the therapeutic agents may be given in multiple doses, or both may be given as multiple doses. If not simultaneous, the timing between the multiple doses may vary from more than zero weeks to less than four weeks. In addition, the

combination methods, compositions and formulations are not to be limited to the use of only two agents; the use of multiple therapeutic combinations are also envisioned.

[00129] The dosage regimen to treat, prevent, or ameliorate the condition(s) for which relief is sought, can be modified in accordance with a variety of factors. These factors include the disorder or condition from which the subject suffers, as well as the age, weight, sex, diet, and medical condition of the subject. Thus, the dosage regimen actually employed can vary widely and therefore can deviate from the dosage regimens set forth herein.

[00130] The pharmaceutical agents which make up the combination therapy disclosed herein may be a combined dosage form or in separate dosage forms intended for substantially simultaneous administration. The pharmaceutical agents that make up the combination therapy may also be administered sequentially, with either therapeutic compound being administered by a regimen calling for two-step administration. The two-step administration regimen may call for sequential administration of the active agents or spaced-apart administration of the separate active agents. The time period between the multiple administration steps may range from a few minutes to several hours, depending upon the properties of each pharmaceutical agent such as potency, solubility, bioavailability, plasma half-life and kinetic profile of the pharmaceutical agent. Circadian variation of the target molecule concentration may also determine the optimal dose interval.

[00131] In addition, the compounds described herein also may be used in combination with procedures that may provide additional or synergistic benefit to the patient. By way of example only, patients are expected to find therapeutic and/or prophylactic benefit in the methods described herein, wherein pharmaceutical composition of a compound disclosed herein and /or combinations with other therapeutics are combined with genetic testing to determine whether that individual is a carrier of a mutant gene that is correlated with certain diseases or conditions.

[00132] The compounds described herein and combination therapies can be administered before, during, or after the occurrence of a disease or condition, and the timing of administering the composition containing a compound can vary. Thus, for example, the compounds can be used as a prophylactic and can be administered continuously to subjects with a propensity to develop conditions or diseases in order to prevent the occurrence of the disease or condition. The initial administration can be via any route practical, such as, for example, an intravenous injection, a bolus injection, infusion over about 5 minutes to about 5 hours, a pill, a capsule, transdermal patch, buccal delivery, and the like, or combination thereof. A compound is preferably administered as soon as is practicable after the onset of a disease or condition is detected or suspected, and for a length of time necessary for the treatment of the disease or condition. The

length of treatment can vary for each subject, and the length can be determined using specified criteria.

Kits/Articles of Manufacture

[00133] For use in the therapeutic methods of use described herein, kits and articles of manufacture are also described herein. Such kits include a carrier, package, or container that is compartmentalized to receive one or more containers such as vials, tubes, and the like, each of the container(s) comprising one of the separate elements to be used in a method described herein. Suitable containers include, for example, bottles, vials, syringes, and test tubes. In one embodiment, the containers are formed from a variety of materials such as glass or plastic.

[00134] The articles of manufacture provided herein contain packaging materials. Packaging materials for use in packaging pharmaceutical products include, e.g., U.S. Patent No. 5,323,907. Examples of pharmaceutical packaging materials include, but are not limited to, blister packs, bottles, tubes, bags, containers, bottles, and any packaging material suitable for a selected formulation and intended mode of administration and treatment.

[00135] In some embodiments, the compounds or compositions described herein, are presented in a package or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The compound or composition described herein is packaged alone, or packaged with another compound or another ingredient or additive. In some embodiments, the package contains one or more containers filled with one or more of the ingredients of the pharmaceutical compositions. In some embodiments, the package comprises metal or plastic foil, such as a blister pack. In some embodiments, the package or dispenser device is accompanied by instructions for administration, such as instructions for administering the compounds or compositions for treating a neoplastic disease. In some embodiments, the package or dispenser is accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. In some embodiments, such notice, for example, is the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. In some embodiments, compositions include a compound described herein formulated in a compatible pharmaceutical carrier are prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

[00136] For example, the container(s) include Compound 1, a compound of Formula (I), or a compound of Formula (II), optionally in a composition or in combination with another agent as

disclosed herein. Such kits optionally include an identifying description or label or instructions relating to its use in the methods described herein.

[00137] A kit typically includes labels listing contents and/or instructions for use, and package inserts with instructions for use. A set of instructions will also typically be included.

[00138] In one embodiment, a label is on or associated with the container. In one embodiment, a label is on a container when letters, numbers or other characters forming the label are attached, molded or etched into the container itself; a label is associated with a container when it is present within a receptacle or carrier that also holds the container, e.g., as a package insert. In one embodiment, a label is used to indicate that the contents are to be used for a specific therapeutic application. The label also indicates directions for use of the contents, such as in the methods described herein.

[00139] In certain embodiments, the pharmaceutical compositions are presented in a pack or dispenser device which contains one or more unit dosage forms containing a compound provided herein. The pack, for example, contains metal or plastic foil, such as a blister pack. In one embodiment, the pack or dispenser device is accompanied by instructions for administration. In one embodiment, the pack or dispenser is also accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, is the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. In one embodiment, compositions containing a compound provided herein formulated in a compatible pharmaceutical carrier are also prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

EXAMPLES

List of abbreviations

[00140] As used throughout the description of the invention, the following abbreviations, unless otherwise indicated, shall be understood to have the following meanings:


ACN or MeCN	acetonitrile
Bn	benzyl
BOC or Boc	<i>tert</i> -butyl carbamate
<i>t</i> -Bu	<i>tert</i> -butyl
Cy	cyclohexyl
DCE	dichloroethane (ClCH ₂ CH ₂ Cl)

DCM	dichloromethane (CH ₂ Cl ₂)
DIPEA or DIEA	diisopropylethylamine
DMAP	4-(<i>N,N</i> -dimethylamino)pyridine
DMF	dimethylformamide
DMA	<i>N,N</i> -dimethylacetamide
DMSO	dimethylsulfoxide
eq or equiv	equivalent(s)
Et	ethyl
Et ₂ O	diethyl ether
EtOH	ethanol
EtOAc	ethyl acetate
HPLC	high performance liquid chromatography
Me	methyl
MeOH	methanol
MS	mass spectroscopy
GC	gas chromatography
h	hour(s)
KF	Karl Fischer
min	minutes
MsOH	methanesulfonic acid
NMR	nuclear magnetic resonance
RP-HPLC	reverse phase-high performance liquid chromatography
r.t.	room temperature
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TLC	thin layer chromatography
V	volumes


I. Chemical Synthesis

[00141] Unless otherwise noted, reagents and solvents were used as received from commercial suppliers. Anhydrous solvents and oven-dried glassware were used for synthetic transformations sensitive to moisture and/or oxygen. Yields were not optimized. Reaction times are approximate and were not optimized. Column chromatography and thin layer chromatography (TLC) were performed on silica gel unless otherwise noted.

Example 1: Preparation of 2,6-dibromo-4-(2-ethylbenzofuran-3-carbonyl-4,5,6,7-*d*₄)phenyl acetate (Int-7)




Step 1: 2-Hydroxybenzaldehyde-3,4,5,6-*d*₄ (Int-1)



[00142] A solution of phen-*d*₅-ol (1.0 eq), magnesium chloride (1.5 eq), and triethylamine (3.7 eq) in ACN (10 V) was stirred at 20°C for 0.5 h. Formaldehyde (8.0 eq) was added and the reaction mixture was heated at reflux for 3 h. The reaction mixture was cooled to room temperature and 10% HCl solution (10V) was added. The mixture was extracted with EtOAc (3 x 6V). The combined organic layers were washed with brine (6 V), dried with Na₂SO₄, and concentrated to give 2-hydroxybenzaldehyde-3,4,5,6-*d*₄ (Int-1) as a yellow oil.


Step 2: 1-(Benzofuran-2-yl-4,5,6,7-*d*₄)ethan-1-one (Int-2)



[00143] A solution of 2-hydroxybenzaldehyde-3,4,5,6-*d*₄ (Int-1) (1.0 eq), bromopropanone (1.0 eq), and potassium carbonate (3.0 eq) in acetone (14 V) was heated at reflux for 6 h. The


reaction mixture was cooled to room temperature and filtered. The filtrate was concentrated and the crude product was recrystallized (petroleum ether/EtOAc 10:1) to give 1-(benzofuran-2-yl-4,5,6,7-d₄)ethan-1-one (**Int-2**) as a yellow solid.

Step 3: 2-Ethylbenzofuran-4,5,6,7-d₄ (Int-3)




[00144] A solution of 1-(benzofuran-2-yl-4,5,6,7-d₄)ethan-1-one (**Int-2**) (1.0 eq) in diethylene glycol (16 V) was heated at 120°C. N₂H₄·H₂O (2.0 eq) and water (1V) was added. The reaction mixture was heated at 180°C for 10 min and then cooled to 120°C. KOH (2.0 eq) was added and the reaction mixture was heated at 120°C for 6 h. The reaction mixture was cooled, poured into water, and extracted with EtOAc (20 V x 3). The combined organic layers were washed with brine (20 V) and concentrated to give 2-ethylbenzofuran-4,5,6,7-d₄ (**Int-3**) as a colorless oil.

Step 4: (2-Ethylbenzofuran-3-yl-4,5,6,7-d₄)(4-methoxyphenyl)methanone (Int-4)




[00145] A solution of 2-ethylbenzofuran-4,5,6,7-d₄ (**Int-3**) (1.0 eq) and 4-methoxybenzoyl chloride (1.15 eq) in DCM (30 V) was cooled to 0°C and charged with AlCl₃ (1.1 eq). The reaction mixture was stirred for 2 h at 0°C. D₂O (2 V) was added to the mixture dropwise at 5°C and the mixture was stirred for 0.5 h. Water (8 V) was added. The organic layer was separated, washed with brine (10 V), dried with Na₂SO₄, and concentrated under vacuum at 40°C to give (2-ethylbenzofuran-3-yl-4,5,6,7-d₄)(4-methoxyphenyl)methanone (**Int-4**) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.81-7.77 (dd, 2H), 7.12-7.08 (dd, 2H), 3.88(s, 3H), 2.86-2.78 (q, 2H), 1.28-1.23 (t, 3H); LCMS: 285 [M+H]⁺.

Step 5: (2-Ethylbenzofuran-3-yl-4,5,6,7-d₄)(4-hydroxyphenyl)methanone (Int-5)




[00146] To a solution of (2-ethylbenzofuran-3-yl-4,5,6,7-*d*₄)(4-methoxyphenyl)methanone (**Int-4**) (1.0 eq) in DCM (10 V) at 0°C was added BBr₃ (2.2 eq) dropwise at 0-5°C. The reaction mixture was warmed to room temperature and stirred for 14 h. Ice water (10 V) was added and the mixture was stirred for 0.5 h. The organic layer was separated, washed with brine (10 V), dried with Na₂SO₄, and concentrated under vacuum at 40°C to give (2-ethylbenzofuran-3-yl-4,5,6,7-*d*₄)(4-hydroxyphenyl)methanone (**Int-5**) as a brown solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 10.47 (s, 1H), 7.71-7.68 (dd, 2H), 6.92-6.88 (dd, 2H), 2.84-2.78 (q, 2H), 1.28-1.24 (t, 3H); LCMS: 271 [M+H]⁺.

Step 6: (3,5-Dibromo-4-hydroxyphenyl)(2-ethylbenzofuran-3-yl-4,5,6,7-*d*₄)methanone (Int-6)



[00147] To a solution of (2-ethylbenzofuran-3-yl-4,5,6,7-*d*₄)(4-hydroxyphenyl)methanone (**Int-5**) (1.0 eq) in DCM (10 V) at 10°C was added NBS (1.7 eq) dropwise at 0-5°C. The reaction mixture was warmed to 18°C and stirred for 16 h. The reaction mixture was charged with additional NBS (0.14 eq) at 10°C and stirred for 16 h at 18°C. The reaction mixture was charged with additional NBS (0.05 eq) at 10°C and stirred for 3 h at 18°C. Water (15 V) was added and the mixture was stirred for 0.5 h. The organic layer was separated, washed with brine (15 V), dried with Na₂SO₄, and concentrated under vacuum at 40°C to give a yellow solid. The yellow solid was slurried in EtOAc/n-heptane (1 V/10 V) at 60°C for 2 h. The mixture was cooled to 10°C and filtered to give (3,5-dibromo-4-hydroxyphenyl)(2-ethylbenzofuran-3-yl-4,5,6,7-*d*₄)methanone (**Int-6**) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 11.05 (s, 1H), 7.92 (s, 2H), 2.84-2.75 (q, 2H), 1.27-1.20 (t, 3H); LCMS: 429 [M+H]⁺.


Step 7: 2,6-Dibromo-4-(2-ethylbenzofuran-3-carbonyl-4,5,6,7-*d*₄)phenyl acetate (Int-7)




[00148] To a solution of (3,5-dibromo-4-hydroxyphenyl)(2-ethylbenzofuran-3-yl-4,5,6,7-*d*₄)methanone (**Int-6**) (1.0 eq) and triethylamine (2.5 eq) in DCM (10 V) at 0°C was added acetyl

chloride (2.0 eq) dropwise at 0-5°C. The reaction mixture was warmed to 15°C and stirred for 2 h. Water (10 V) was added. The organic layer was separated, washed with brine (10 V), dried with Na₂SO₄, and concentrated under vacuum at 40°C to give a crude solid. The crude solid was decolorized with activated charcoal (0.5 w/w) in EtOAc (10 V) at 50°C for 1 h. The mixture was cooled to 30°C and filtered with kieselguhr aid to remove the activated charcoal. The filtrate was concentrated under vacuum at 40°C. The residue was dissolved in i-PrOH (2 V) and heated at 60°C for 1 h. The solution was cooled to 45°C, charged with seed crystals (0.5% w/w), and stirred for 1 h. The mixture was cooled to 25°C and stirred for 16 h. The mixture was filtered and the solid dried to give 2,6-dibromo-4-(2-ethylbenzofuran-3-carbonyl-4,5,6,7-*d*₄)phenyl acetate (**Int-7**) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.08 (s, 2H), 2.81-2.74 (q, 2H), 2.44 (s, 3H), 1.27-1.22 (t, 3H); LCMS: 471 [M+H]⁺.

Example 2: Preparation of (3,5-dibromo-4-hydroxyphenyl)(2-(1-hydroxyethyl)benzofuran-3-yl-4,5,6,7-*d*₄)methanone (Compound 1)




Step 1: (3,5-Dibromo-4-hydroxyphenyl)(2-ethyl-6-nitrobenzofuran-3-yl-4,5,7-*d*₃)methanone (Int-8)




[00149] To a mixture of 2,6-dibromo-4-(2-ethylbenzofuran-3-carbonyl-4,5,6,7-*d*₄)phenyl acetate (34 g, 72.3 mmol) (**Int-7**) in dichloromethane (250 mL) was added nitronium tetrafluoroborate (11.5 g, 86.8 mmol) slowly at 0 °C. The reaction mixture was stirred at room temperature for overnight. The mixture was quenched with brine and extracted with dichloromethane. The organic phase was washed with brine, dried over sodium sulfate, and evaporated to dryness. The crude product was purified with column chromatography on silica gel to afford (3,5-dibromo-4-(2-ethyl-6-nitrobenzofuran-3-yl-4,5,7-*d*₃)methanone (**Int-8**) (12.5 g, 37 %) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.1 (bs, 1 H), 7.98 (s, 2 H), 2.87 (q, *J*=7.2 Hz, 2 H), 1.3 (t, *J*=7.2 Hz, 3 H).

Step 2: (6-Amino-2-ethylbenzofuran-3-yl-4,5,7-*d*₃)(3,5-dibromo-4-hydroxyphenyl)methanone (Int-9)




[00150] A mixture of (3,5-dibromo-4-(2-ethyl-6-nitrobenzofuran-3-yl-4,5,7-*d*₃)methanone (**Int-8**) (12.5 g, 26.5 mol), iron powder (7.4 g, 132 mmol), conc. hydrochloride (1.2 mL, 13.3 mmol), ethanol (250 mL), and H₂O (50 mL) was heated to reflux for 2 hours. The mixture was filtered to remove the iron powder. The solution was evaporated to dryness. The residue was purified with column chromatography on silica gel to afford (6-amino-2-ethylbenzofuran-3-yl-4,5,7-*d*₃)(3,5-dibromo-4-hydroxyphenyl)methanone (**Int-9**) (9.2 g, 79%) as a yellow solid. ESIMS (*m/z*): 441.9, 439.9, 437.9 (*M*+H)⁺.

Step 3: 3-(3,5-Dibromo-4-hydroxybenzoyl)-2-ethylbenzofuran-6-diazonium-4,5,7-*d*₃ tetrafluoro-borate (Int-10)




[00151] To a mixture of (6-amino-2-ethylbenzo-furan-3-yl-4,5,7- d_3)(3,5-dibromo-4-hydroxyphenyl)methanone (**Int-9**) (9.2 g, 20.8 mmol) and fluoroboric acid (4.2 g, 22.9 mmol) in H_2O (100 mL) was slowly added a solution of sodium nitrite (1.58 g, 22.9 mmol) in H_2O . The reaction mixture was stirred at room temperature overnight. The suspension obtained was then filtered in vacuum. The residue was washed with ice water and dried in vacuum to afford 3-(3,5-dibromo-4-hydroxybenzoyl)-2-ethylbenzofuran-6-diazonium-4,5,7- d_3 tetrafluoroborate (**Int-10**) (11 g, 98%) as a pink solid which was used for next steps without further purification. ESIMS (m/z): 455.9, 453.9, 451.9 ($\text{M}+\text{H}$) $^+$.

Step 4: (3-(3,5-Dibromo-4-hydroxybenzoyl)-2-ethylbenzofuran-6-yl-4,5,7- d_3)boronic acid (Int-11)




[00152] To a mixture of hypodiboric acid (3.65 g, 40.6 mmol) in DMF (160 mL) was added 3-(3,5-dibromo-4-hydroxybenzoyl)-2-ethylbenzofuran-6-diazonium-4,5,7- d_3 tetrafluoroborate (**Int-10**) (11 g, 20.3 mmol) at room temperature. The reaction mixture was stirred at the same temperature for 2 hours. The mixture was quenched with brine and extracted with ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate, and evaporated to dryness. The residue was purified with column chromatography on silica gel to afford (3-(3,5-dibromo-4-hydroxybenzoyl)-2-ethylbenzofuran-6-yl-4,5,7- d_3)boronic acid (**Int-11**) (3.9 g, 41%) as a white solid. ESIMS (m/z): 473.9, 471.9, 469.9 ($\text{M}+\text{H}$) $^+$.

Step 5: (3,5-Dibromo-4-hydroxyphenyl)(2-ethyl-6-hydroxybenzofuran-3-yl-4,5,7- d_3)methanone (Int-12)




[00153] To a mixture of (3-(3,5-dibromo-4-hydroxybenzoyl)-2-ethylbenzofuran-6-yl-4,5,7- d_3)boronic (**Int-11**) (3.9 g, 8.28 mmol), and sodium hydroxide (662 mg, 16.6 mmol) in THF (60 mL) was added hydrogen peroxide (w/w 48%, 1.2 g, 16.6 mmol) dropwise at 0 °C. The mixture was stirred at room temperature for 2 hours. It was quenched with 1M hydrochloride to pH 5-6, extracted with ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate and evaporated to dryness. The residue was purified on silica gel with column chromatography to afford (3,5-dibromo-4-hydroxyphenyl)(2-ethyl-6-hydroxybenzofuran-3-yl-4,5,7- d_3)methanone (**Int-12**) (3.0 g, 83%) as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 10.9 (bs, 1 H), 9.68 (bs, 1 H), 7.91 (s, 2 H), 2.74 (q, $J=7.6$ Hz, 2 H), 1.24 (t, $J=7.6$ Hz, 3 H). ESIMS (m/z): 445.9, 443.9, 441.9 ($\text{M}+\text{H}$) $^+$.

Step 6: 4-(6-Acetoxy-2-ethylbenzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (Int-13)



[00154] To a mixture of (3,5-dibromo-4-hydroxyphenyl)(2-ethyl-6-hydroxybenzofuran-3-yl-4,5,7- d_3)methanone (**Int-12**) (3.0 g, 6.78 mmol), and triethylamine (1.71 g, 17.0 mmol) in dichloromethane (60 mL) was added acetyl chloride (1.06 g, 13.6 mmol) dropwise at 0 °C. The reaction mixture was stirred at room temperature for 2h. The mixture was quenched with brine and extracted with ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate, and evaporated to dryness. The residue was purified on silica gel with column chromatography to afford 4-(6-acetoxy-2-ethylbenzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (**Int-13**) (2.8 g, 78%) as a white solid.


Step 7: 4-(6-Acetoxy-2-(1-bromoethyl)benzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (Int-14)



[00155] A mixture of 4-(6-acetoxy-2-ethylbenzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (**Int-13**) (2.8 g, 5.31 mmol), *N*-bromosuccinimide (945 mg, 5.31 mmol), and 2,2'-azobis(2-methylpropionitrile) (87 mg, 0.53 mmol) in perchloromethane (80 mL) was heated to


reflux overnight. The mixture was evaporated to dryness. The crude was purified with column chromatography on silica gel to afford 4-(6-acetoxy-2-(1-bromoethyl)benzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (**Int-14**) (3.06 g, 95%) as a white solid.

Step 8: 4-(6-Acetoxy-2-(1-hydroxyethyl)benzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (Int-15**)**



[00156] A mixture of 4-(6-acetoxy-2-(1-bromoethyl)benzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (**Int-14**) and silver(I) oxide (1.75 g, 7.57 mmol) in *N,N*-dimethylformamide (40 mL) and H₂O (8 mL) was heated to 70 °C for 6 hours. The mixture was filtered to remove the silver(I) oxide. The filtrate was quenched with brine and extracted with ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate, and evaporated to dryness. The residue was purified on silica gel with column chromatography to afford 4-(6-acetoxy-2-(1-hydroxyethyl)benzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (**Int-15**) (2.3 g, 84 %) as a white solid. ¹H NMR (400 MHz, DMSO- d_6) δ 11.7 (bs, 1 H), 8.14 (bs, 1 H), 5.65 (d, *J*=4.8 Hz, 1 H), 4.80 (dd, *J*=6.8 Hz, 4.8 Hz, 1 H), 2.45 (s, 3 H), 2.33 (s, 3 H), 1.48 (d, *J*=4.8 Hz, 3 H). ESIMS (*m/z*): 527.9, 525.9, 523.9 (*M*+H)⁺.

Step 9: (3,5-Dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7- d_3)methanone (Compound 1**)**



[00157] A mixture of 4-(6-acetoxy-2-(1-hydroxyethyl)benzofuran-3-carbonyl-4,5,7- d_3)-2,6-dibromophenyl acetate (**Int-15**) (2.3 g, 4.24 mmol) and lithium hydroxide hydrate (427 mg, 10.2 mmol) in methanol (40 mL) and H₂O (5 mL) was stirred at room temperature for 2 hours. The mixture was poured into 30 mL of H₂O and evaporated in vacuum to remove methanol. The solution obtained was acidified to pH 5-6 with 1M hydrochloride and extracted with ethyl acetate. The organic phase was washed with brine, dried over sodium sulfate, and evaporated to dryness. The residue was purified with column chromatography on silica gel to afford (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7- d_3)methanone

(Compound 1) (1.06 g, 55%) as a yellow solid. ^1H NMR (400 MHz, DMSO- d_6) δ 11.1 (bs, 1 H), 9.75 (bs, 1 H), 7.92 (s, 2 H), 5.47 (bs, 1 H), 4.77 (d, $J=6.8$ Hz, 1 H), 1.45 (d, $J=6.8$ Hz, 3 H). ESIMS (m/z): 443.9, 441.9, 439.9 (M+H) $^+$.

II. Biological Data

Example 3: *In vitro* Interaction Studies of Compound 1 with the human URAT1 Uptake Transporter

[00158] Uptake experiments were performed using MDCKII cells stably expressing the human URAT1 uptake transporter. Cells were cultured at $37 \pm 1^\circ\text{C}$ in an atmosphere of 95:5 air:CO₂ and were plated onto standard 96-well tissue culture plates at the cell number described in Table 1.

Table 1

Transporter	Control cell line	Cell number/well	Culturing medium	Incubation prior to the assay	Buffer
human URAT1	Mock-transfected MDCKII	1×10^5	DMEM 4.5 g/L glucose	24 h	HBSS w/o Cl $^-$ (pH 7.4)

DMEM: Dulbecco's Modified Eagle's Medium; HBSS: Hank's balanced salt solution; w/o: without

[00159] Before the experiment, the medium was removed and the cells were washed twice with 100 μL of HBSS without Cl $^-$. Uptake experiments were carried out at $37 \pm 1^\circ\text{C}$ in 50 μL of HBSS without Cl $^-$, pH 7.4 containing the probe substrate (20 μM uric acid) and the test article (TA) or solvent. The organic solvent concentration was equal in all wells, and did not exceed 1% (v/v).

[00160] Treatment groups are presented in Table 2.

Table 2

Treatment groups in the 96-well plate format	No. of wells
Compound 1 in assay buffer (0.01, 0.04, 0.12, 0.37, 1.11, 3.33 and 10.0 μM) in transfected cells	3 per TA concentration
1% DMSO control in transfected cells	3
1% DMSO control in control cells	3
Reference inhibitor in assay buffer with 1% DMSO in transfected cells	3
Reference inhibitor in assay buffer with 1% DMSO in control cells	3


[00161] After the experiment, cells were washed twice with 100 μL of ice cold HBSS without Cl $^-$ and lysed with 50 μL of 0.1 M NaOH. Radiolabelled probe substrate transport was determined by measuring an aliquot (35 μL) from each well for liquid scintillation counting.

[00162] Results: Compound 1 was soluble in HBSS buffer at all tested concentrations; the highest tested concentration being 10 μ M. Compound 1 inhibited URAT1 mediated uric acid accumulation with an $IC_{50} = 0.913 \mu$ M.

CLAIMS

We Claim:

1. A compound that is (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof.
2. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof:




Formula (I);


wherein:

R₁, R₂, R₃, and R₄ are each independently selected from a group consisting of hydrogen, deuterium, hydroxy, and methoxy; wherein at least one of R₁, R₂, R₃, and R₄ is deuterium; and


R₅ is -CH₂CH₃, -CH₂CH₂(OH), or -CH(OH)CH₂(OH).

3. The compound of claim 2, or a pharmaceutically acceptable salt or solvate thereof, wherein at least two of R₁, R₂, R₃, and R₄ is deuterium.
4. The compound of claim 2, or a pharmaceutically acceptable salt or solvate thereof, wherein at least three of R₁, R₂, R₃, and R₄ is deuterium.
5. The compound of claim 2, or a pharmaceutically acceptable salt or solvate thereof, wherein R₁, R₂, R₃, and R₄ are deuterium.
6. The compound of any one of claims 2-5, or a pharmaceutically acceptable salt or solvate thereof, wherein R₅ is -CH₂CH₃.
7. The compound of any one of claims 2-5, or a pharmaceutically acceptable salt or solvate thereof, wherein R₅ is -CH₂CH₂(OH).
8. The compound of any one of claims 2-5, or a pharmaceutically acceptable salt or solvate thereof, wherein R₅ is -CH(OH)CH₂(OH).
9. The compound of claim 2, or a pharmaceutically acceptable salt or solvate thereof, selected





10. A compound of Formula (II), or a pharmaceutically acceptable salt or solvate thereof:




Formula (II);


wherein:

R_1 , R_2 , R_3 , and R_4 are each independently selected from a group consisting of hydrogen, deuterium, hydroxy, and methoxy; wherein at least one of R_1 , R_2 , R_3 , and R_4 is deuterium.

11. The compound of claim 10, wherein at least two of R_1 , R_2 , R_3 , and R_4 is deuterium.
12. The compound of claim 10, wherein at least three of R_1 , R_2 , R_3 , and R_4 is deuterium.
13. The compound of any one of claims 10-12, wherein at least one of R_1 , R_2 , R_3 , and R_4 is hydroxy.
14. The compound of claim 10, or a pharmaceutically acceptable salt or solvate thereof,



selected from:



15. A method for treating or preventing hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof.
16. A method for treating or preventing hyperuricemia or gout in an individual in need thereof, comprising administering to the individual a therapeutically effective amount of a compound of any one of claims 1-14, or a pharmaceutically acceptable salt or solvate thereof.
17. The method of claim 15 or claim 16 for treating or preventing hyperuricemia in an individual in need thereof.
18. The method of any one of claims 15-17 for treating hyperuricemia in an individual in need thereof.
19. The method of any one of claims 15-17 for preventing hyperuricemia in an individual in need thereof.
20. The method of claim 15 or claim 16 for treating or preventing gout in an individual in need thereof.
21. The method of claim 20 for treating gout in an individual in need thereof.
22. The method of claim 20 for preventing gout in an individual in need thereof.
23. The method of any one of claims 15-22, wherein the therapeutically effective amount is from about 3 mg to about 1500 mg
24. The method of any one of claims 15-23, wherein the therapeutically effective amount is from about 3 mg to about 600 mg.
25. The method of any one of claims 15-24, wherein the therapeutically effective amount is from about 5 mg to about 300 mg.
26. The method of any one of claims 15-25, wherein the therapeutically effective amount is from about 10 mg to about 200 mg.
27. The method of any one of claims 15-26, wherein the therapeutically effective amount is from about 10 mg to about 100 mg.
28. The method of any one of claims 15-27, wherein (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone is administered orally.

29. The method of claim 28, wherein the therapeutically effective amount is taken with food.
30. The method of claim 28, wherein the therapeutically effective amount is taken without food.
31. The method of any one of claims 15-30, wherein the therapeutically effective amount is administered to the individual once per day.
32. The method of any one of claims 15-31, wherein the therapeutically effective amount is administered to the individual twice per day.
33. The method of any one of claims 15-32, further comprising administering at least one additional therapeutic agent.
34. The method of any one of claims 15-33, further comprising administering a xanthine oxidase inhibitor.
35. The method of claim 34, wherein the xanthine oxidase inhibitor is allopurinol, oxypurinol, febuxostat, topiroxostat, or inositol.
36. The method of any one of claims 15-35, further comprising administering an SGLT2 inhibitor.
37. The method of claim 36, wherein the SGLT2 inhibitor is canagliflozin, dapagliflozin, empagliflozin, empagliflozin/linagliptin, empagliflozin/metformin, or dapagliflozin/metformin.
38. A pharmaceutical composition comprising the compound of claim 1, or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients.
39. A pharmaceutical composition comprising a compound of any one of claims 1-14, or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients.
40. A pharmaceutical composition for the treatment or prevention of hyperuricemia or gout, wherein the pharmaceutical composition comprises a therapeutically effective amount of (3,5-dibromo-4-hydroxyphenyl)(6-hydroxy-2-(1-hydroxyethyl)benzofuran-3-yl-4,5,7-*d*₃)methanone, or a pharmaceutically acceptable salt or solvate thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients.
41. A pharmaceutical composition for the treatment or prevention of hyperuricemia or gout, wherein the pharmaceutical composition comprises a therapeutically effective amount of a compound of any one of claims 1-14, or a pharmaceutically acceptable salt or solvate

- thereof, and at least one inactive ingredient selected from pharmaceutically acceptable carriers, diluents, and excipients.
42. The pharmaceutical composition of claim 40 or claim 41 for the treatment or prevention of hyperuricemia in an individual in need thereof.
 43. The pharmaceutical composition of any one of claims 40-42 for the treatment of hyperuricemia in an individual in need thereof.
 44. The pharmaceutical composition of any one of claims 40-42 for the prevention of hyperuricemia in an individual in need thereof.
 45. The pharmaceutical composition of claim 40 or claim 41 for the treatment or prevention of gout in an individual in need thereof.
 46. The pharmaceutical composition of claim 45 for the treatment of gout in an individual in need thereof.
 47. The pharmaceutical composition of claim 45 for the prevention of gout in an individual in need thereof.
 48. The pharmaceutical composition of any one of claims 40-47, wherein the therapeutically effective amount is from about 3 mg to about 1500 mg.
 49. The pharmaceutical composition of any one of claims 40-48, wherein the therapeutically effective amount is from about 3 mg to about 600 mg.
 50. The pharmaceutical composition of any one of claims 40-49, wherein the therapeutically effective amount is from about 5 mg to about 300 mg.
 51. The pharmaceutical composition of any one of claims 40-50, wherein the therapeutically effective amount is from about 10 mg to about 200 mg.
 52. The pharmaceutical composition of any one of claims 40-51, wherein the therapeutically effective amount is from about 10 mg to about 100 mg.
 53. The pharmaceutical composition of any one of claims 40-52 formulated for oral, intravenous, intramuscular, or subcutaneous administration.
 54. The pharmaceutical composition of any one of claims 40-53 formulated for oral administration.
 55. The pharmaceutical composition of claim 54, wherein the therapeutically effective amount is taken with food.
 56. The pharmaceutical composition of claim 54, wherein the therapeutically effective amount is taken without food.
 57. The pharmaceutical composition of any one of claims 40-56, wherein the therapeutically effective amount is administered once per day.

58. The pharmaceutical composition of any one of claims 40-56, wherein the therapeutically effective amount is administered twice per day.
59. The pharmaceutical composition of any one of claims 40-58, further comprising at least one additional therapeutic agent.
60. The pharmaceutical composition of any one of claims 40-59, further comprising a xanthine oxidase inhibitor.
61. The pharmaceutical composition of claim 60, wherein the xanthine oxidase inhibitor is allopurinol, oxypurinol, febuxostat, topiroxostat, or inositol.
62. The pharmaceutical composition of any one of claims 40-61, further comprising an SGLT2 inhibitor.
63. The pharmaceutical composition of claim 62, wherein the SGLT2 inhibitor is canagliflozin, dapagliflozin, empagliflozin, empagliflozin/linagliptin, empagliflozin/metformin, or dapagliflozin/metformin.