SB	gesehen	erledigt
GT		

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro

(43) Internationales Veröffentlichungsdatum
10. Mai 2002 (10.05.2002)

PCT

(10) Internationale Veröffentlichungsnummer
WO 02/036104 A3

(51) Internationale Patentklassifikation[7]: A61K 9/72, 31/5386

(21) Internationales Aktenzeichen: PCT/EP01/12292

(22) Internationales Anmeldedatum:
24. Oktober 2001 (24.10.2001)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
100 54 036.8 31. Oktober 2000 (31.10.2000) DE

(71) Anmelder *(für alle Bestimmungsstaaten mit Ausnahme von US)*: **BOEHRINGER INGELHEIM PHARMA KG** [DE/DE]; 55216 Ingelheim am Rhein (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder *(nur für US)*: **NIKLAUS-HUMKE, Barbara** [DE/DE]; Henry-Moisand-Strasse 12, 55130 Mainz (DE). **SCHMELZER, Christel** [DE/DE]; Welfenstrasse 14, 55218 Ingelheim am Rhein (DE). **BARTH, Petra** [DE/DE]; Oberer Laubenheimer Weg 17, 55131 Mainz (DE). **DRECHSEL, Karin** [DE/DE]; Wackernheimer Strasse 17, 55218 Ingelheim (DE).

(81) Bestimmungsstaaten *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Bestimmungsstaaten *(regional)*: ARIPO-Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI-Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Veröffentlicht:
— *mit internationalem Recherchenbericht*
— *vor Ablauf der für Änderungen der Ansprüche geltenden Frist; Veröffentlichung wird wiederholt, falls Änderungen eintreffen*

(88) Veröffentlichungsdatum des internationalen Recherchenberichts: 1. August 2002

Zur Erklärung der Zweibuchstaben-Codes und der anderen Abkürzungen wird auf die Erklärungen ("Guidance Notes on Codes and Abbreviations") am Anfang jeder regulären Ausgabe der PCT-Gazette verwiesen.

WO 02/036104 A3

(54) **Title:** INHALATIVE SOLUTION FORMULATION CONTAINING A TIOTROPIUM SALT

(54) **Bezeichnung:** INHALATIVE LÖSUNGSFORMULIERUNG MIT EINEM TIOTROPIUMSALZ

(57) **Abstract:** The invention relates to a propellant-free inhalation formulation of tiotropium bromide or tiotropium bromide monohydrate, which is dissolved in water or in a mixture consisting of water and ethanol, and to propellant-free inhalable aerosols resulting therefrom.

(57) **Zusammenfassung:** Die vorliegende Erfindung betrifft eine treibgasfreie Inhalationsformulierung von in Wasser oder in einem Gemisch aus Ethanol gelöstem Tiotropiumbromid bzw. Tiotropiumbromid-Monohydrat und daraus resultierenden treibgasfreien inhalierbaren Aerosolen.

Inhalative Solution Formulation Containing A Tiotropium Salt

The present invention relates to a propellant-free inhalable formulation of a pharmaceutically acceptable salt of tiotropium dissolved in water or a mixture of water and ethanol and propellant-free inhalable aerosols resulting therefrom. The formulation according to the invention is particularly suitable for administering the active substance by inhalation, especially for treating asthma and COPD.

Tiotropium, chemically (1α,2β,4β,5α,7β)-7-[(hydroxydi-2-thienylacetyl)oxy]-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.1.$0^{2,4}$]nonane, is known as tiotropium bromide from European Patent Application EP 418 716 A1. The bromide salt of tiotropium has the following chemical structure:



(I)

The compound has valuable pharmacological properties and is known by the name tiotropium bromide. Tiotropium and its salts are highly effective anticholinergics and can provide therapeutic benefit in the treatment of asthma or COPD (chronic obstructive pulmonary disease). The monohydrate of tiotropium bromide is also pharmacologically valuable.
Both compounds are a preferred object of the present invention.

The present invention relates to liquid active substance formulations of these compounds which can be administered by inhalation; the liquid formulations according to the invention have to meet high quality standards.

To achieve an optimum distribution of active substances in the lung it makes sense to use a liquid formulation without propellant gases administered using suitable inhalers. Those inhalers which are capable of nebulising a small amount of a liquid formulation

in the dosage needed for therapeutic purposes within a few seconds into an aerosol suitable for therapeutic inhalation are particularly suitable. Within the scope of the invention, preferred nebulisers are those in which an amount of less than 100 microlitres, preferably less than 50 microlitres, most preferably less than 20 microlitres of active substance solution can be nebulised preferably in one puff to form an aerosol having an average particle size of less than 20 microns, preferably less than 10 microns, so that the inhalable part of the aerosol already corresponds to the therapeutically effective quantity.

An apparatus of this kind for the propellant-free administration of a metered amount of a liquid pharmaceutical composition for inhalation is described in detail for example in International Patent Application WO 91/14468 "Atomizing Device and Methods" and also in WO 97/12687, cf. Figures 6a and 6b and the accompanying description. In a nebuliser of this kind a pharmaceutical solution is converted by means of a high pressure of up to 500 bar into an aerosol destined for the lungs, which is sprayed. Within the scope of the present specification reference is expressly made to the entire contents of the literature mentioned above.

In inhalers of this kind the formulations of solutions are stored in a reservoir. It is essential that the active substance formulations used are sufficiently stable when stored and at the same time are such that they can be administered directly, if possible without any further handling, in accordance with their medical purpose. Moreover, they must not contain any ingredients which might interact with the inhaler in such a way as to damage the inhaler or the pharmaceutical quality of the solution or of the aerosol produced.

To nebulise the solution a special nozzle is used as described for example in WO 94/07607 or WO 99/16530. Reference is expressly made here to both these publications.

WO 98/27959 discloses formulations of solutions for the inhaler described above which contain as additive the disodium salt of editic acid (sodium edetate). For aqueous formulations of solutions which are to be converted into inhalable aerosols using the inhaler described above, the specification favours a minimum concentration

of sodium edetate of 50 mg/100 ml, in order to reduce the incidence of spray anomalies. Among the Examples disclosed there is a formulation containing tiotropium bromide. In this formulation the active substance is dissolved in water. The proportion of sodium edetate is again 50 mg / 100 ml.

Surprisingly, it has now been found that formulations of solutions of tiotropium salts in water or a water-ethanol mixture wherein the proportion of the additive sodium edetate is significantly less than 50 mg / 100 ml show a reduction in the scattering of the composition delivered, compared with the formulation containing tiotropium bromide known from the prior art. In addition, the spray quality is very good. Water is preferably used as the solvent. The resulting aerosol has very good properties for administration by inhalation.

Another advantage of the formulation is that, thanks to the absence of or reduction in the additive sodium edetate in the active substance formulation, the pH of the solution formulation can be lowered. Low pH levels are necessary for the long-term stability of the formulation.

It is therefore an aim of the present invention to provide an aqueous active substance formulation containing a pharmaceutically acceptable tiotropium salt which meets the high standards needed in order to be able to achieve optimum nebulisation of a solution using the inhalers mentioned hereinbefore. The active substance formulations according to the invention must be of sufficiently high pharmaceutical quality, i.e. they should be pharmaceutically stable over a storage time of some years, preferably at least one year, more preferably two years.

Another aim is to provide propellant-free formulations of solutions containing tiotropium salts which are nebulised under pressure using an inhaler, the composition delivered by the aerosol produced falling reproducibly within a specified range.

According to the invention, any pharmaceutically acceptable salts of tiotropium may be used for the formulation. When the term tiotropium salt is used within the scope of the present invention, this is to be taken as a reference to tiotropium. According to the invention a reference to tiotropium, which is the free ammonium cation,

corresponds to a reference to tiotropium in the form of a salt (tiotropium salt) which contains an anion as counter-ion. Tiotropium salts which may be used within the scope of the present invention are preferably compounds which contain, in addition to tiotropium as counter-ion (anion), chloride, bromide, iodide, methanesulphonate, para-toluenesulphonate and/or methylsulphate.

Accordingly, a first aspect of the present invention provides a pharmaceutical preparation comprising of

- a tiotropium salt as active substance, in a concentration based on tiotropium of between 0.0005 and 5 % by weight,
- water on its own or a water/ethanol mixture as solvent for the active substance,
- acid for achieving a pH between 2.0 and 3.1,
- a pharmacologically acceptable preservative,
- editic acid or an edetic acid salt in an amount of greater than 0 up to 25 mg /100 ml,
- optionally a pharmacologically acceptable complexing agent and/or stabiliser and/or a pharmacologically acceptable cosolvent and/or other pharmacologically acceptable adjuvants and additives in addition to the preservative.

A second aspect of the present invention provides a pharmaceutical preparation containing water, 0.1 % by weight of tiotropium bromide, 0.01 % by weight of benzalkonium chloride, 0.05 % by weight of sodium edetate, which is adjusted to a pH of 3.0 using hydrochloric acid.

A third aspect of the present invention provides a use of a pharmaceutical preparation as defined above for nebulising in an inhaler which nebulises defined amounts of the pharmaceutical formulation by the application of pressures from 100 to 600 bar through a nozzle having at least one nozzle opening with a depth of 2 to 10 microns and a width of 5 to 15 microns to form an inhalable aerosol.

A fourth aspect of the present invention provides a pharmaceutical preparation as defined above for use as a medicament, particularly for the treatment of asthma and/or COPD.

A fifth aspect of the present invention provides a process for preparing a pharmaceutical preparation as defined above by mixing the individual components.

Within the scope of the present invention tiotropium bromide is preferred as the salt. References to tiotropium bromide within the scope of the present invention must always be taken as references to all possible amorphous and crystalline modifications of tiotropium bromide. These may for example contain molecules of solvent in their crystalline structure. Of all the crystalline modifications of tiotropium bromide those which also contain water (hydrates) are preferred according to the invention. It is particularly preferred within the scope of the present invention to use tiotropium bromide monohydrate. The formulation preferably does not contain any other active substance which is not tiotropium or a pharmaceutically acceptable salt thereof.

In the formulations according to the invention the tiotropium salts are dissolved in a solvent. The solvent may be exclusively water, or it may be a mixture of water and ethanol. The relative proportion of ethanol to water is not limited, but the maximum limit is preferably up to 70 % by volume, particularly up to 60 % by volume and most preferably up to 30 % by volume. The remaining % by volume consist of water. The preferred solvent is water without the addition of ethanol.

According to one embodiment, the formulation preferably contains only a single tiotropium salt . However, the formulation may also contain a mixture of different tiotropium salts and solvates. Preferably, the formulation does not contain any active substance other than tiotropium as hereinbefore defined.

The concentration of the tiotropium salt based on the proportion of tiotropium in the finished pharmaceutical preparation depends on the therapeutic effect sought. For most of the complaints which respond to tiotropium the concentration of tiotropium is between 0.0005 and 5 % by weight, preferably between 0.001 and 3 % by weight. In the case of tiotropium bromide or tiotropium bromide monohydrate the preferred

amount based on tiotropium is 0.0005 to 0.5 % by weight, more preferably 0.0005 to 0.25 % by weight and particularly preferably 0.001 to 0.1 % by weight.

The pH of the formulation according to the invention is between 2.0 and 4.5, preferably between 2.5 and 3.5 and more preferably between 2.7 and 3.3 and particularly preferably between 2.7 and 3.2. Most preferred are pHs with an upper limit of 3.1.

The pH is adjusted by the addition of pharmacologically acceptable acids. Examples of inorganic acids which are preferred for this purpose include: hydrochloric acid, hydrobromic acid, nitric acid, sulphuric acid and/or phosphoric acid.

Examples of particularly suitable organic acids are: ascorbic acid, citric acid, malic acid, tartaric acid, maleic acid, succinic acid, fumaric acid, acetic acid, formic acid and/or propionic acid, etc. Preferred inorganic acids are hydrochloric acid and sulphuric acid. It is also possible to use acids which form an acid addition salt with the active substance.
Of the organic acids, ascorbic acid, fumaric acid and citric acid are preferred. If desired, mixtures of the abovementioned acids may also be used, particularly in the case of acids which have other properties in addition to their acidifying properties, e.g. those which act as flavourings or antioxidants, such as for example citric acid or ascorbic acid.
Hydrochloric acid deserves special mention as an inorganic acid.

If desired, pharmacologically acceptable bases may be used to titrate the pH precisely. Suitable bases include for example alkali metal hydroxides and alkali metal carbonates. The preferred alkali ion is sodium. If bases of this kind are used, care must be taken to ensure that the resulting salts, which are then contained in the finished pharmaceutical formulation, are pharmacologically compatible with the abovementioned acid.

According to another embodiment, there is no need to add editic acid (EDTA) or one of the known salts thereof, sodium edetate, to the present formulation as a stabiliser or complexing agent.

Yet another embodiment contains editic acid and/ or the abovementioned salts thereof.

In a preferred embodiment of this kind the content based on sodium edetate is less than 10 mg / 100 ml. In this case, there is one preferred range from 5 mg/ 100 ml to less than 10 mg/100 ml or another from greater than 0 to 5 mg/100ml.

In another embodiment the content of sodium edetate is 10 to 30 mg / 100 ml, preferably not more than 25 mg/ 100 ml.

In a preferred embodiment this additive is omitted entirely.

The remarks made concerning sodium edetate also apply analogously to other comparable additives which have complexing properties and can be used instead, such as for example nitrilotriacetic acid and the salts thereof.

By complexing agents is preferably meant within the scope of the present invention molecules which are capable of entering into complex bonds. Preferably, these compounds should have the effect of complexing cations, most preferably metal cations.

As well as ethanol, other cosolvents and/or other adjuvants may be added to the formulation according to the invention.

Preferred cosolvents are those which contain hydroxyl groups or other polar groups, for example alcohols - especially isopropylalcohol, glycols - especially propyleneglycol, polyethyleneglycol, polypropyleneglycol, glycolether, glycerol, polyoxyethylene alcohols and polyoxyethylene fatty acid esters, provided that these are not already being used as the solvent or suspension agent.

By adjuvants and additives are meant, in this context, any pharmacologically acceptable and therapeutically useful substance which is not an active substance, but can be formulated together with the active substance in the pharmacologically suitable solvent, in order to improve the qualities of the active substance formulation.

Preferably, these substances have no pharmacological effects or no appreciable or at least no undesirable pharmacological effects in the context of the desired therapy. The adjuvants and additives include, for example, surfactants such as e.g. soya lecithin, oleic acid, sorbitan esters such as sorbitan trioleate, polyvinylpyrrolidone, other stabilisers, complexing agents, antioxidants and/or preservatives which prolong the shelf life of the finished pharmaceutical formulation, flavourings, vitamins and/or other additives known in the art. The additives also include pharmacologically acceptable salts such as sodium chloride, for example.

The preferred excipients include antioxidants such as ascorbic acid, for example, provided that it has not already been used to adjust the pH, vitamin A, vitamin E, tocopherols and similar vitamins or provitamins occurring in the human body.

Preservatives can be added to protect the formulation from contamination with pathogenic bacteria. Suitable preservatives are those known from the prior art, particularly benzalkonium chloride or benzoic acid or benzoates such as sodium benzoate in the concentration known from the prior art.

Preferred formulations contain only benzalkonium chloride and sodium edetate, in addition to the solvent water and the tiotropium salt.
In another preferred embodiment, sodium edetate is omitted.

As already mentioned a number of times, tiotropium bromide is described in EP 418 716 A1 and crystalline tiotropium bromide monohydrate may be obtained using a process which is described in more detail below.

In order to prepare the crystalline monohydrate according to the present invention, the tiotropium bromide obtained by the method disclosed in EP 418 716 A1, for example, first has to be taken up in water, heated, purified with activated charcoal and, after removal of the activated charcoal, the tiotropium bromide-monohydrate is slowly crystallised while cooling slowly.

The following procedure is preferably followed:

In a reaction vessel of suitable dimensions, the solvent is mixed with tiotropium bromide, which has been obtained by the method disclosed in EP 418 716 A1, for example.
For each mol of tiotropium bromide put in, 0.4 to 1.5 kg, preferably 0.6 to 1 kg, most preferably about 0.8 kg of water are used as solvent.
The mixture obtained is heated with stirring, preferably to above 50°C, most preferably to above 60°C. The maximum temperature which can be selected is determined by the boiling point of the solvent used. Preferably, the mixture is heated to a range from 80-90°C.
Activated charcoal, either dry or moistened with water, is added to this solution. Preferably, 10 to 50 g, more preferably 15 to 35 g, most preferably about 25 g of activated charcoal are put in per mol of tiotropium bromide used. If desired the activated charcoal is suspended in water before being added to the solution containing tiotropium bromide. 70 to 200 g, preferably 100 to 160 g, more preferably about 135 g of water are used, per mol of tiotropium bromide put in, in order to suspend the activated charcoal. If the activated charcoal is suspended in water beforehand, before being added to the solution containing tiotropium bromide, it is advisable to rinse again with the same amount of water.
After the activated charcoal has been added, stirring is continued at constant temperature for between 5 and 60 minutes, preferably between 10 and 30 minutes, more preferably for about 15 minutes and the mixture obtained is filtered to remove the activated charcoal. The filter is then rinsed with water. 140 to 400 g, preferably 200 to 320 g, most preferably about 270 g of water are used for this, per mol of tiotropium bromide used.
The filtrate is then slowly cooled, preferably to a temperature of 20-25°C. The cooling preferably takes place at a cooling rate of 1 to 10°C every 10 to 30 minutes, preferably 2 to 8°C every 10 to 30 minutes, more preferably 3 to 5°C every 10 to 20 minutes, most preferably 3 to 5°C about every 20 minutes. If desired, the cooling to 20 to 25°C may be followed by further cooling to below 20°C, more preferably to 10 to 15°C.
After cooling is complete, stirring is continued for between 20 minutes and 3 hours, preferably between 40 minutes and 2 hours, more preferably for about one hour to complete the crystallisation.
The crystals obtained are then isolated by filtering or suction filtering to remove the solvent. If it should prove necessary to subject the crystals obtained to a further washing step, it is advisable to use water or acetone as the washing solvent. 0.1 to 1.0 L, preferably 0.2 to 0.5 L, more preferably about 0.3 L of solvent may be used per mol of tiotropium bromide put in, in order to wash the tiotropium bromide

monohydrate crystals obtained. If necessary the washing step may be repeated. The product obtained is dried *in vacuo* or using circulating heated air until a water content of 2.5 – 4.0 % is obtained.

Accordingly, one embodiment may also relate to formulations of solutions of the type described above using crystalline tiotropium bromide monohydrate which may be obtained by the procedure described above.

The pharmaceutical formulations containing tiotropium salts according to an embodiment are preferably used in an inhaler of the kind described hereinbefore in order to produce the propellant-free aerosols according to the invention. At this point we should once again expressly mention the patent documents described hereinbefore, to which reference is hereby made.

As described at the beginning, a further developed embodiment of the preferred inhaler is disclosed in WO 97/12687 and Figure 6 thereof. This nebuliser (Respimat®) can advantageously be used to produce the inhalable aerosols according to the invention containing a tiotropium salt as active substance. Because of its cylindrical shape and handy size of less than 9 to 15 cm long and 2 to 4 cm wide, the device can be carried anywhere by the patient. The nebuliser sprays a defined volume of the pharmaceutical formulation out through small nozzles at high pressures, so as to produce inhalable aerosols.

The preferred atomiser essentially consists of an upper housing part, a pump housing, a nozzle, a locking clamp, a spring housing, a spring and a storage container, characterised by

- a pump housing fixed in the upper housing part and carrying at one end a nozzle body with the nozzle or nozzle arrangement,
- a hollow piston with valve body,
- a power take-off flange in which the hollow body is fixed and which is located in the upper housing part,
- a locking clamping mechanism located in the upper housing part ,
- a spring housing with the spring located therein, which is rotatably mounted on the upper housing part by means of a rotary bearing,

- a lower housing part which is fitted onto the spring housing in the axial direction.

The hollow piston with valve body corresponds to a device disclosed in WO 97/12687. It projects partially into the cylinder of the pump housing and is disposed to be axially movable in the cylinder. Reference is made particularly to Figures 1-4 - especially Figure 3 - and the associated parts of the description. At the moment of release of the spring the hollow piston with valve body exerts, at its high pressure end, a pressure of 5 to 60 Mpa (about 50 to 600 bar), preferably 10 to 60 Mpa (about 100 to 600 bar) on the fluid, the measured amount of active substance solution. Volumes of 10 to 50 microlitres are preferred, volumes of 10 to 20 microlitres are more preferable, whilst a volume of 15 microlitres per actuation is particularly preferred.

The valve body is preferably mounted at the end of the hollow piston which faces the nozzle body.

The nozzle in the nozzle body is preferably microstructured, i.e. produced by micro-engineering. Microstructured nozzle bodies are disclosed for example in WO-94/07607; reference is hereby made to the contents of this specification, especially Figure 1 and the associated description.
The nozzle body consists for example of two sheets of glass and/or silicon securely fixed together, at least one of which has one or more microstructured channels which connect the nozzle inlet end to the nozzle outlet end. At the nozzle outlet end there is at least one round or non-round opening 2 to 10 microns deep and 5 to 15 microns wide, the depth preferably being 4.5 to 6.5 microns and the length being 7 to 9 microns.
If there is a plurality of nozzle openings, preferably two, the directions of spraying of the nozzles in the nozzle body may run parallel to each other or may be inclined relative to one another in the direction of the nozzle opening. In the case of a nozzle body having at least two nozzle openings at the outlet end, the directions of spraying may be inclined relative to one another at an angle of 20 degrees to 160 degrees, preferably at an angle of 60 to 150 degrees, most preferably 80 to 100°.

The nozzle openings are preferably arranged at a spacing of 10 to 200 microns, more preferably at a spacing of 10 to 100 microns, still more preferably 30 to 70 microns. A spacing of 50 microns is most preferred.
The directions of spraying therefore meet in the region of the nozzle openings.

As already mentioned, the liquid pharmaceutical preparation hits the nozzle body at an entry pressure of up to 600 bar, preferably 200 to 300 bar and is atomised through the nozzle openings into an inhalable aerosol. The preferred particle sizes of the aerosol are up to 20 microns, preferably 3 to 10 microns.

The locking clamping mechanism contains a spring, preferably a cylindrical helical compression spring as a store for the mechanical energy. The spring acts on the power take-off flange as a spring member the movement of which is determined by the position of a locking member. The travel of the power take-off flange is precisely limited by an upper stop and a lower stop. The spring is preferably tensioned via a stepping-up gear, e.g. a helical sliding gear, by an external torque which is generated when the upper housing part is turned relative to the spring housing in the lower housing part. In this case, the upper housing part and the power take-off flange contain a single- or multi-speed spline gear.

The locking member with the engaging locking surfaces is arranged in an annular configuration around the power take-off flange. It consists for example of a ring of plastics or metal which is inherently radially elastically deformable. The ring is arranged in a plane perpendicular to the axis of the atomiser. After the locking of the spring, the locking surfaces of the locking member slide into the path of the power take-off flange and prevent the spring from being released. The locking member is actuated by means of a button. The actuating button is connected or coupled to the locking member. In order to actuate the locking clamping mechanism the actuating button is moved parallel to the annular plane, preferably into the atomiser, and the deformable ring is thereby deformed in the annular plane. Details of the construction of the locking clamping mechanism are described in WO 97/20590.

The lower housing part is pushed axially over the spring housing and covers the bearing, the drive for the spindle and the storage container for the fluid.

When the atomiser is operated, the upper part of the housing is rotated relative to the lower part, the lower part taking the spring housing with it. The spring meanwhile is compressed and biased by means of the helical sliding gear, and the clamping mechanism engages automatically. The angle of rotation is preferably a whole-number fraction of 360 degrees, e.g. 180 degrees. At the same time as the spring is tensioned, the power take-off component in the upper housing part is moved along by a given amount, the hollow piston is pulled back inside the cylinder in the pump housing, as a result of which some of the fluid from the storage container is sucked into the high pressure chamber in front of the nozzle.

If desired, a plurality of replaceable storage containers containing the fluid to be atomised can be inserted in the atomiser one after another and then used. The storage container contains the aqueous aerosol preparation according to the invention.

The atomising process is initiated by gently pressing the actuating button. The clamping mechanism then opens the way for the power take-off component. The biased spring pushes the piston into the cylinder in the pump housing. The fluid emerges from the nozzle of the atomiser in the form of a spray.

Further details of the construction are disclosed in PCT applications WO 97/12683 and WO 97/20590, to which reference is hereby made.

The components of the atomiser (nebuliser) are made of a material suitable for their function. The housing of the atomiser and – if the function allows – other parts as well are preferably made of plastics, e.g. by injection moulding. For medical applications, physiologically acceptable materials are used.

Figures 1a/b, which are identical to Figures 6 a/b of WO 97/12687, show the Respimat® nebuliser with which the aqueous aerosol preparations according to the invention can advantageously be inhaled.

Figure 1 a shows a longitudinal section through the atomiser with the spring under tension, Figure 1 b shows a longitudinal section through the atomiser with the spring released.

The upper housing part (51) contains the pump housing (52), on the end of which is mounted the holder (53) for the atomiser nozzle. In the holder is the nozzle body (54) and a filter (55). The hollow piston (57) fixed in the power take-off flange (56) of the locking clamping mechanism projects partly into the cylinder of the pump housing. At its end the hollow piston carries the valve body (58). The hollow piston is sealed off by the gasket (59). Inside the upper housing part is the stop (60) on which the power take-off flange rests when the spring is relaxed. Located on the power take-off flange is the stop (61) on which the power take-off flange rests when the spring is under tension. After the tensioning of the spring, the locking member (62) slides between the stop (61) and a support (63) in the upper housing part. The actuating button (64) is connected to the locking member. The upper housing part ends in the mouthpiece (65) and is closed off by the removable protective cap (66).

The spring housing (67) with compression spring (68) is rotatably mounted on the upper housing part by means of the snap-fit lugs (69) and rotary bearings. The lower housing part (70) is pushed over the spring housing. Inside the spring housing is the replaceable storage container (71) for the fluid (72) which is to be atomised. The storage container is closed off by the stopper (73), through which the hollow piston projects into the storage container and dips its end into the fluid (supply of active substance solution).

The spindle (74) for the mechanical counter is mounted on the outside of the spring housing. The drive pinion (75) is located at the end of the spindle facing the upper housing part. On the spindle is the slider (76).

The nebuliser described above is suitable for nebulising the aerosol preparations according to the invention to form an aerosol suitable for inhalation.

If the formulation according to the invention is nebulised using the method described above (Respimat®), the mass expelled, in at least 97%, preferably at least 98% of all the actuations of the inhaler (puffs), should correspond to a defined quantity with a range of tolerance of not more than 25%, preferably 20% of this quantity. Preferably, between 5 and 30 mg, more preferably between 5 and 20 mg of formulation are delivered as a defined mass per puff.

However, the formulation according to the invention can also be nebulised using inhalers other than those described above, for example jet-stream inhalers.

Examples

I. Example of the synthesis of tiotropium bromide monohydrate

15.0 kg of tiotropium bromide are added to 25.7 kg of water in a suitable reaction vessel. The mixture is heated to 80-90°C and stirred at constant temperature until a clear solution is formed. Activated charcoal (0.8 kg), moistened with water, is suspended in 4.4 kg of water, this mixture is added to the solution containing tiotropium bromide and rinsed with 4.3 kg of water. The mixture thus obtained is stirred for at least 15 min. at 80-90°C and then filtered through a heated filter into an apparatus which has been preheated to an outer temperature of 70°C. The filter is rinsed with 8.6 kg of water. The contents of the apparatus are cooled to a temperature of 20-25°C at a rate of 3-5°C every 20 minutes. Using cold water the apparatus is cooled further to 10-15°C and crystallisation is completed by stirring for at least another hour. The crystals are isolated using a suction filter drier, the crystal slurry isolated is washed with 9 L of cold water (10-15°C) and cold acetone (10-15°C). The crystals obtained are dried at 25°C for 2 hours in a nitrogen current.

Yield: 13.4 kg of tiotropium bromide monohydrate (86 % of theory).

II. Examples of formulations

100 g of pharmaceutical preparation contain:

Example	Amount of tiotropium bromide, based on tiotropium:	Amount of tiotropium bromide monohydrate, based on tiotropium:	Amount of benzalkonium chloride	Amount of sodium edetate	pH, adjusted with HCl (1N)
1	0.099 g	---	10 mg	25 mg	3.0
2	0.006 g	---	10 mg	25 mg	3.0
3	0.099 g	---	10 mg	10 mg	3.0
4	0.006 g	---	10 mg	10 mg	3.0
5	---	0.099 g	10 mg	25 mg	3.0
6	---	0.006 g	10 mg	25 mg	3.0
7	---	0.099 g	10 mg	10 mg	3.0
8	---	0.006 g	10 mg	10 mg	3.0

The remainder is water.

Claims: 17

1. Pharmaceutical preparation comprising of
   - a tiotropium salt as active substance, in a concentration based on tiotropium of between 0.0005 and 5 % by weight,
   - water on its own or a water/ethanol mixture as solvent for the active substance,
   - acid for achieving a pH between 2.0 and 3.1,
   - a pharmacologically acceptable preservative,
   - editic acid or an edetic acid salt in an amount of greater than 0 up to 25 mg /100 ml,
   - optionally a pharmacologically acceptable complexing agent and/or stabiliser and/or a pharmacologically acceptable cosolvent and/or other pharmacologically acceptable adjuvants and additives in addition to the preservative.

2. Pharmaceutical preparation according to claim 1, characterised in that the active substance is tiotropium bromide.

3. Pharmaceutical preparation according to claim 1, characterised in that the active substance is tiotropium bromide monohydrate.

4. Pharmaceutical preparation according to one of claims 1 to 3, characterised in that the solvent is water on its own.

5. Pharmaceutical preparation according to one of claims 1 to 3, characterised in that the solvent is a water-ethanol mixture with preferably up to 70 vol.% of

ethanol, more preferably up to 60 vol.% of ethanol and most preferably up to 30 vol.% of ethanol.

6. Pharmaceutical preparation according to one of claims 1 to 5, characterised in that it does not contain a complexing agent.

7. Pharmaceutical preparation according to one of claims 1 to 6, characterised in that it does not contain a stabiliser.

8. Pharmaceutical preparation according to one of claims 1 to 5, characterised in that editic acid salt is present in an amount from 5 to less than 10 mg /100 ml.

9. Pharmaceutical preparation according to claim 8, characterised in that the editic acid salt is sodium edetate.

10. Pharmaceutical preparation according to one of claims 1 to 9, characterised in that the pH is between 2.5 and 3.1, preferably between 2.7 and 3.1 and most preferably between 2.7 and 3.0.

11. Pharmaceutical preparation according to one of claims 1 to 10, characterised in that the concentration based on tiotropium is between 0.001 and 3 % by weight, preferably 0.001 to 0.5 % by weight, more preferably 0.001 to 0.25 % by weight and particularly preferably 0.001 to 0.1 % by weight.

12. Pharmaceutical preparation according to one of claims 1 to 11, characterised in that the preparation contains benzalkonium chloride as preservative.

13. Pharmaceutical preparation according to one of claims 1 to 5 and 8 to 12, characterised in that pharmacologically acceptable adjuvants and additives are used in addition to the preservative.

14. Pharmaceutical preparation according to claim 13, characterised in that the preparation contains an antioxidant as adjuvant.

15. Pharmaceutical preparation according to one of claims 1 to 14, characterised in that no cosolvents and/or pharmacologically acceptable adjuvants and additives are used apart from the preservative.

16. Pharmaceutical preparation containing water, 0.1 % by weight of tiotropium bromide, 0.01 % by weight of benzalkonium chloride, 0.05 % by weight of sodium edetate, which is adjusted to a pH of 3.0 using hydrochloric acid.

17. Use of a pharmaceutical preparation according to one of claims 1 to 16 for nebulising in an inhaler which nebulises defined amounts of the pharmaceutical formulation by the application of pressures from 100 to 600 bar through a nozzle having at least one nozzle opening with a depth of 2 to 10 microns and a width of 5 to 15 microns to form an inhalable aerosol.

18. Use according to claim 17, characterised in that the at least one nozzle opening is at least two nozzle openings which are inclined relative to one another in the direction of the nozzle opening at an angle of from 20 degrees to 160 degrees.

19. Use according to claim 17 or 18, characterised in that the defined amounts are 10 to 50 microlitres.

20. Use according to one of claims 17 to 19, characterised in that the mass of formulation delivered in at least 97% of all actuations of the inhaler is between 5 and 30 mg with a range of tolerance of 25%.

21. Use according to claim 20, characterised in that the mass of formulation delivered in at least 97% of all actuations of the inhaler is between 5 and 30 mg with a range of tolerance of 20%.

22. Use according to one of claims 20 or 21, characterised in that the mass delivered is achieved in at least 98% of all actuations of the inhaler.

23. A pharmaceutical preparation according to one of claims 1 to 16 for use as a medicament, particularly for the treatment of asthma and/or COPD.

24. Process for preparing a pharmaceutical preparation according to one of claims 1 to 16 by mixing the individual components.