


ENZYMATIC HYDROLYSIS OF 4-METHYLTHIOBUTYRONITRILE

The present invention relates to the use of a nitrilase as a catalyst for hydrolysing a nitrile group into a carboxylic group. The invention relates more
5 precisely to the use of the nitrilase chosen from nitrilases of microorganisms of the genus *Alcaligenes*, *Rhodococcus* or *Gordona* and more precisely of *Alcaligenes faecalis* filed under the number ATCC 8750, *Rhodococcus* sp. HT 29-7 filed under the number FERM BP-
10 3857 or *Gordona terrae* MA-1 filed under the number FERM BP-4535.

The nitrilase of *Alcaligenes faecalis* is described, for example, in the European and Japanese patents published under the numbers EP 348,901 and
15 JP 03 224 496 respectively, both belonging to the Asahi company. In these patents, and more particularly in the European patent, the nitrilase is used to produce optically active acids from racemic nitriles. The preferred starting nitriles are α -substituted and
20 contain an alkyl chain preferably containing 1 to 3 carbon atoms or an aromatic radical. Example 11 of the abovementioned European patent describes the hydrolysis of racemic mandelonitrile with *Alcaligenes faecalis*; the R-(-)-mandelic acid is obtained in an enantiomeric
25 excess of 91%. European patent 486,289 and its equivalent American patent US 5,326,702 from the company Nitto Chemical describe the use of *Alcaligenes*

sp. BC35-2 (FERM No. 11265 or FERM-BP-3318) to hydrolyse mandelonitrile in the presence of a sulphite; in this case, mandelic acid is obtained in an enantiomeric excess of about 98%.

5 Patent JP 04 341 185 from the company Asahi also describes the preparation and use of the nitrilase from *Alcaligenes faecalis* ATCC 8750 to solve the problems of preparation of optically pure compounds from their racemic nitrile. The enantioselective
10 nitrilase described in the Japanese patent referred to above is more preferably capable of hydrolysing the 2-hydroxynitrile of general formula (I) below into 2-hydroxycarboxylic acid of formula (II) below:



in which R represents an optionally substituted aryl
15 group or an optionally substituted heterocyclic group. It is mentioned in that patent that hydrolysis of the nitrile of mandelic acid by the said nitrilase makes it possible to obtain a 100% enantiomeric excess of R-(-)-mandelic acid. It is mentioned in Example 5 that
20 the said nitrilase may be used to hydrolyse aliphatic nitriles such as valeronitrile, acrylonitrile, 2-halopropionitriles and chloroacetonitrile. Nothing is

specified as regards the enantioselectivity of this hydrolysis on aliphatic nitriles having an asymmetric centre.

The said text also specifies that the
5 nitrilase of *Alcaligenes faecalis* is of optimum activity at a pH between 6.5 and 8.0; the working temperature of the said enzyme should, according to that reference, always be below 45°C.

It appeared to be entirely surprising that
10 the said enzyme of *Alcaligenes faecalis* ATCC 8750, when used to hydrolyse 4-methylthio-2-hydroxybutyronitrile, carried out this hydrolysis without any selectivity from the point of view of the optical purity. This enzyme hydrolyses the said nitrile producing a racemic
15 mixture of the two acids with no preponderance of either of the isomers.

The nitrilase of *Rhodococcus* sp. HT 29-7 filed under the number FERM BP-3857 is described, for example, in the European and American patents published
20 under the numbers EP 610,049 and US 5,296,373 respectively, both belonging to the company Nitto.

In these patents, and particularly in the American patent, the nitrilase is used to produce optically active acids from racemic nitriles bearing a
25 phenyl group. The starting nitriles all contain an aromatic radical; this essentially involves hydrolysis of mandelonitrile or of derivatives thereof substituted on the aromatic ring. Example 1 of the abovementioned

American patent describes the hydrolysis of racemic mandelonitrile with *Rhodococcus* sp. HT 29-7; the R-(-)-mandelic acid is obtained in an enantiomeric excess of 100%. Example 2 describes the hydrolysis of substituted mandelonitrile derivatives on the ring; the enantiomeric excess of mandelic acid derivative is also 100% and this is likewise the case when the hydrolysis is carried out on benzaldehyde nitriles.

European patent 610 049 from the company Nitto Chemical describes the use of *Rhodococcus* sp. HT 29-7 (FERM BP-3857), of *Alcaligenes* sp. BC35-2 (FERM BP-3318) or of *Brevibacterium acetylicum* IAM 1790 to hydrolyse α -hydroxynitriles substituted in the γ position with an aromatic ring into α -hydroxycarboxylic acid substituted with an optically active phenyl group. The acids obtained with *Rhodococcus* sp. HT 29-7 always have a variable enantiomeric excess depending on the starting nitrile and on the presence or absence of phosphoric acid (pH set at about 8.2).

It appeared to be entirely surprising that the said enzyme of *Rhodococcus* sp. HT 29-7 (FERM BP-3318), when used to hydrolyse 4-methylthio-2-hydroxybutyronitrile, carried out this hydrolysis without any enantiomeric selectivity. This enzyme hydrolyses the said nitrile producing a racemic mixture of the two acids without any preponderance of either of the isomers.

The nitrilase of *Gordona terrae* MA-1 filed

under the number FERM BP-4535 is described in European patent 0,610,048 to hydrolyse nitriles bearing a γ -phenyl group and an α -hydroxy group into the corresponding optically active acid. The enantiomeric excess ranges in all the examples between 92 and 100%.

It appeared to be entirely surprising that the said enzyme of *Gordona terrae* MA-1 filed under the number FERM BP-4535, when used to hydrolyse 4-methylthio-2-hydroxybutyronitrile, carried out this hydrolysis without any enantiomeric selectivity. This enzyme hydrolyses the said nitrile producing a racemic mixture of the two acids without any preponderance of either of the isomers.

The present invention thus relates to a process for the preparation of racemic α -substituted 4-methylthiobutyric acids by hydrolysis of racemic α -substituted 4-methylthiobutyronitriles with the nitrilase of one of the following microorganisms: *Alcaligenes faecalis* filed under the number ATCC 8750, *Rhodococcus* sp. HT 29-7 filed under the number FERM BP-3857 or *Gordona terrae* MA-1 filed under the number FERM BP-4535.

According to the process of the invention, the microorganism may be used as it is or immobilized on supports which are well known to those skilled in the art.

Moreover, the genetic information which codes for the enzyme may be transferred from the parent

microorganism (such as *A. faecalis* ATCC8750, etc.) into a microorganism such as *Bacillus subtilis*.

One variant of the process of the invention consists in using, in place of a microorganism, a
5 corresponding amount of its free or immobilized enzyme, this enzyme being totally or partially purified.

Among the α -substituted
4-methylthiobutyronitriles, it is preferred to use
4-methylthio-2-hydroxybutyronitrile, which makes it
10 possible to prepare the hydroxy analogue of racemic methionine. This technique for the preparation of methionine derivatives makes it possible to avoid the formation of large amounts of inorganic side products whose disposal causes ever-increasing problems on
15 account of environmental protection constraints.

In the context of the present invention, these enzymes have a nitrilase activity in a pH range between 4 and 11 and optimum activity at a pH of between 5 and 9. Their optimum working temperature is
20 between 30 and 60°C; it is preferred, however, to use them between 30 and 50°C.

It is preferred to work with a nitrile concentration in the starting solution of between 10 mmol and 400 mmol per litre and preferably between 50
25 and 200 mmol per litre. The concentration of ammonium salt of the acid obtained has little influence on the activity of the enzyme when it is less than 2 mol per litre and it is preferably between 0.1 and 1.5 mol per

litre.

The invention will be described more fully with the aid of the examples which follow, which should not be considered as limiting the invention.

5 EXAMPLE 1

The microorganism *Alcaligenes faecalis* ATCC 8750 is cultured in the following medium:

	ammonium acetate	10 g/l
	yeast extract	5 g/l
10	peptone	5 g/l
	K ₂ HPO ₄	5 g/l
	MgSO ₄ ·7H ₂ O	0.2 g/l
	FeSO ₄ ·7H ₂ O	30 mg/l
	NaCl	1 g/l
15	benzonitrile	0.5 g/l
	pH 7.2	

Culturing is carried out at 30°C in 2 litre conical flasks containing 600 ml of medium. The whole mixture is stirred at 150 revolutions per minute for 30 hours. The cell pellet is recovered, washed with 9 g/l NaCl solution and then frozen.

Operating conditions to hydrolyse the butyronitrile:

	4-methylthio-2-hydroxybutyronitrile	50 mM
25	cells	5 mg/ml
	phosphate buffer	200 mM
	temperature	30°C
	duration	4 hours

Study of the kinetics of hydrolysis of
4-methylthio-2-hydroxybutyronitrile shows a linear
production of 2.8 μmol of hydroxy analogue of
methionine per hour and per mg of dry cells. The
5 enantiomeric excess of the acid formed was measured by
liquid chromatography; it is equal to 0% for a yield of
acid of 80%.

EXAMPLE 2

The stability of the enzyme was evaluated
10 according to the kinetics of production of the hydroxy
analogue of methionine over 180 hours with free cells.

The operating conditions are as follows:

4-methylthio-2-hydroxybutyronitrile	100 mM
cells	10 mg/ml
15 phosphate buffer	300 mM
	5 ml
temperature	25°C

The nitrile is added sequentially by addition
of 100 mmol of nitrile after depletion. The formation
of the corresponding acid is indicated in Table I

duration

1 hour

Nitrile concentration (mM)	Activity ($\mu\text{mol/h}\cdot\text{mg}$ of cell $\cdot\text{s}$)
50	2
100	2
150	2
200	2
500	0

5

Stability of the enzyme as a function of the

10 amount of acid formed

operating conditions:

nitrile 50 mM

cells 2.5 mg/ml

phosphate buffer 100 mM

15

pH 7.0 1 ml

temperature 30°C

duration 2 hours

Acid concentration (mM)	Activity ($\mu\text{mol/h}\cdot\text{mg}$ of cell $\cdot\text{s}$)
0	3.2
100	4.5
200	3.8
300	4.1
400	4
500	4
600	3.4
700	2.8

EXAMPLE 4**Purification of the nitrilase**

The cells of *Alcaligenes faecalis* are cultured for 24 hours at 30°C in the medium described in Example 1.

After centrifuging the culture, the pellet is taken up in pH 7.5 buffer (25 mM trisHCl, 10% (w/v) glycerol). The cell suspension is treated with ultrasound and then centrifuged in order to obtain the crude extract. The crude extract is then treated with ammonium sulphate up to 30% of saturation. The precipitate obtained is resuspended in pH 7.5 buffer and then dialysed against 2 litres of the same buffer overnight.

The solution obtained is then loaded onto a Q Sepharose Fast Flow HR 26/10° anion-exchange column pre-

equilibrated with the pH 7.5 buffer. The activity is then eluted with a gradient of 0 to 1 M NaCl.

The active fractions are then loaded onto a Mono Q HR 5/5[°] anion-exchange column pre-equilibrated with pH 7.5 buffer. The nitrilase is eluted using a gradient of 0 to 1 M NaCl.

Lastly, the fractions containing the activity are combined and 1 M ammonium sulphate is then added. This solution is loaded onto a Phenyl Sepharose HR 5/5[°] hydrophobic-interaction column pre-equilibrated with the pH 7.5 buffer, to which is added 1 M (NH₄)₂SO₄. The activity is then eluted with a gradient of 1 to 0 M ammonium sulphate.

The molecular weight of the protein is determined by gel filtration. It is about 260 kDa. A single band of 43 kDa is observed on SDS-PAGE (purity of 95%).

The kinetics of hydrolysis of the 4-methylthio-2-hydroxybutyronitrile into the hydroxy analogue of methionine with the nitrilase of *A. faecalis* is linear.

Time (hour)	Activity ($\mu\text{mol/h}\cdot\text{mg}$ of prot.)	RY (%)
0	0	1
0.5	150	3
1	171	5
21.5	150	68
24.8	150	77
29	150	87

EXAMPLE 5 Influence of the pH

10

Operating conditions:

nitrile 50 mM

protein 50 $\mu\text{g/ml}$

acetate buffer 100 mM of pH 4 to 5

phosphate buffer 100 mM of pH 6 to 7 1 ml

15

Tris-HCl buffer 100 mM of pH 8 to 9

borate buffer 100 mM of pH 10 to 11

temperature 30°C

duration 1 to 2 hours

pH	Activity ($\mu\text{mol/h}\cdot\text{mg}$ of prot.)
4	0
5	42
6	232
7	272
8	405
9	412
10	158
11	0

10 **EXAMPLE 6 Influence of the temperature**

Operating conditions:

nitrile 50 mM

protein 50 $\mu\text{g/ml}$

phosphate buffer 100 mM

15 of pH 7.0

variable temperatures

from 4°C to 60°C

duration 1 hour

The optimum working temperature of the enzyme is 50°C.

Temperature in °C	Activity ($\mu\text{mol/h}\cdot\text{mg}$ of prot.)
4	45
10	67
20	140
30	272
40	419
50	570
60	333

10 COMPARATIVE EXAMPLE WITH MANDELONITRILE

Operating conditions:

mandelonitrile	7 mM
protein	5 $\mu\text{g/ml}$
phosphate buffer 100 mM, pH 7.0	1 ml
temperature	30°C

Time (minute)	Activity ($\mu\text{mol/h}\cdot\text{mg}$ of prot.)	ee
15	1000	1
60	1030	1

20 The purified nitrilase is thus enantioselective on mandelonitrile but not on 4-methylthio-2-hydroxybutyronitrile.

EXAMPLE 7

Hydrolysis of 4-methylthio-2-hydroxybutyronitrile with *Rhodococcus* HT 29-7 FERM BP-3857

5 The microorganism *Rhodococcus* sp. HT 29-7 FERM BP-3857 is cultured in the following medium:

	glycerol	20 g/l
	yeast extract (DIFCO)	3 g/l
	KH_2PO_4	1 g/l
10	$\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$	4.4 g/l
	Na_2SO_4	2.8 g/l
	$\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$	0.85 g/l
	$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	0.05 g/l
	$\text{MnSO}_4 \cdot \text{H}_2\text{O}$	0.033 g/l
15	$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	0.013 g/l
	$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	0.005 g/l
	benzonitrile	0.5 g/l
	pH 7.5	

The culturing is performed at 30°C in 2 litre
 20 conical flasks containing 600 ml of medium. The whole mixture is stirred at 150 revolutions per minute for 140 hours. The cell pellet is recovered, washed with 9 g/l NaCl solution and then frozen.

Influence of the pH on the hydrolysis:

25 Operating conditions:

4-methylthio-2-hydroxybutyronitrile	100 mM
optical density at 660 nm	15
acetate buffer 100 mM of pH 4 to 5	

phosphate buffer 100 mM of pH 6.0

to 7.0 1 ml

Tris-HCl buffer 100 mM of pH 8.0

to 9.0

5 borate buffer 100 mM of pH 10.0

to 11.0

temperature 30°C

duration 10 hours

10

pH	Yield of acid
4	0%
5	100%
6	100%
7	100%
8	100%
9	100%
10	20%
11	0%

15

EXAMPLE 8 Influence of the concentration of substrate

The stability of the enzyme as a function of
 20 the amount of substrate was also measured under the
 following conditions:

4-methylthio-2-hydroxybutyronitrile cf Table
 of results

optical density at 660 nm 20

25

phosphate buffer 300 mM, pH 7.0 5 ml

temperature

30°C

duration

2 hours

Nitrile concentration (mM)	Yield of acid
50	90%
100	44%
200	22%
300	10%
400	0%
500	0%

EXAMPLE 9 Stability of the enzyme as a function of the amount of acid formed

Operating conditions:

4-methylthio-2-hydroxybutyronitrile 100 mM

optical density at 660 nm 15

phosphate buffer 100 mM, pH 7.0 1 ml

temperature 30°C

duration 6 hours

5

Acid concentration (mM)	Yield of acid
0	100%
200	100%
400	100%
600	100%
800	100%
1000	100%

EXAMPLE 10**10 Enantiomeric excess**

4-methylthio-2-hydroxybutyronitrile 140 mM
 optical density at 660 nm 14
 phosphate buffer 100 mM, pH 7.0 1 ml
 temperature 20°C

15

Incubation time	Yield of acid	Optical purity
2 h	15%	0
6 h	50%	0
24 h	100%	0

20

Study of the kinetics of hydrolysis of the
 4-methylthio-2-hydroxybutyronitrile shows a linear
 production of hydroxy analogue of methionine per hour
 per mg of dry cells. The enantiomeric excess of the
 acid formed was measured by liquid chromatography; it

is equal to 0% for a yield of acid ranging from 15% to 100%.

EXAMPLE 11 Influence of the temperature.

Operating conditions:

5	4-methylthio-2-hydroxybutyronitrile	100 mM
	optical density at 660 nm	15
	phosphate buffer 100 mM, pH 7.0	1 ml
	temperature	see table
	duration	6 hours

10	T (°C)	Yield of acid
	10	70%
	20	100%
	30	100%
15	40	100%
	50	31%
	60	15%

EXAMPLE 12 Hydrolysis of 4-methylthio-2-aminobutyronitrile

Operating conditions:

20	4-methylthio-2-hydroxybutyronitrile	23 mM
	optical density at 660 nm	15
	phosphate buffer 100 mM, pH 7.0	1 ml
	temperature	30°C

Time (hours)	Yield of acid	Enantiomeric excess of the acid
0.5	40%	n.d.
1	51%	0.12
2	62%	n.d.
5	78%	0.15

EXAMPLE 13

Culture conditions used for *Gordona terrae*

MA-1 (FERM BP-4535) are as follows:

10	glycerol	10 g/l
	yeast extract	0.4 g/l
	K_2HPO_4	6.8 g/l
	$Na_2HPO_4 \cdot 12H_2O$	7.1 g/l
	Na_2SO_4	2.8 g/l
15	$MgCl_2 \cdot 6H_2O$	0.4 g/l
	$CaCl_2 \cdot 2H_2O$	40 mg/l
	$MnSO_4 \cdot H_2O$	4 mg/ml
	$FeCl_3$	0.6 mg/l
	$ZnSO_4$	0.3 mg/l
20	benzonitrile	0.5 g/l
	pH 7.2	

Activity of the enzyme

The cultured cells are next washed and then placed in contact with the hydroxymethylthiobutyronitrile in order to assay the activity.

Operating conditions:

[nitrile] = 23 mM;

[cells] = 6.8 g/l;

phosphate buffer 100 mM

pH 7.0, 35°C

5




Figure 3: Hydrolysis of the cyanohydrin of AMTP by *Gordona terrae* MA-1.

Time (hours)

10

The kinetics of hydrolysis of the hydroxymethylthiobutyronitrile are linear. The initial rate is estimated at 13 mmol/h·g dry cells.

EXAMPLE 14 Influence of the concentration of cyanohydrin of AMTP on the nitrilase activity.

15

The influence of the initial concentration of hydroxymethylthiobutyronitrile on the nitrilase activity was determined, and the results are indicated in the following table.

Operating conditions:

20

[cells]=5.1 g/l;

phosphate buffer 100 mM

pH 7.0; 35°C.

The kinetics are effected over 0.5, 1, 2 and 3 hours.

Nitrile concentration (mM)	Activity (mmol/h·g dry cells)
50	14
100	13
200	15
300	0
400	0

Up to 200 mM, the activity varies little with the concentration of substrate.

EXAMPLE 15 Influence of the concentration of hydroxy analogue of methionine on the nitrilase activity.

In this test, we studied the influence of the concentration of ammonium 4-hydroxy-2-methylthiobutanoate on the nitrilase activity.

Operating conditions:

[cells] = 5 g/l;
[nitrile] = 100 mM;
phosphate buffer 100 mM, pH 7.0;
35°C.

The concentration of ammonium carboxylate varies between 0 and 1.5 M.

Acid concentration (mol/l)	Activity (μ mol/h·mg of dry cells)
0	9.4
0.5	14
1	19
1.5	15

5

The concentration of ammonium 4-hydroxy-2-methylthiobutanoate does not at all modify the initial activity of the strain *Gordona terrae* HT29-7.

10 EXAMPLE 16 Influence of the pH and of the temperature.

Operating conditions:

[nitrile]=100 mM;

[cells]=7.5 g/l;

acetate buffer 100 mM, pH 4 to 5;

15

phosphate buffer 100 mM, pH 6 to 7;

Tris-HCl buffer 100 mM, pH 8 to 9

borate buffer 100 mM, pH 10 to 11; 3

30°C; kinetics over 1, 2, 4 and 6 hours.

pH	Activity (mmol/h·g dry cells)
4	0.6
5	2.5
6	7.8
7	9.6
8	15.3
9	18
10	5.7
11	11

10 The optimum pH is at about 8-9 under our
operating conditions.

Influence of the temperature on the nitrilase activity
of *Gordona terrae* MA-1.

Operating conditions:

15 [nitrile]=100 mM;
 [cells]=7.5 g/l;
 phosphate buffer 100 mM, pH 7.0;
 kinetics over 1 hour.

T(°C)	Activity (mmol/h·g dry cells)
10	0.6
20	2.3
30	3.2
40	3.4
50	3.5
60	1.9

The optimum temperature is about 40-50°C.



WO 96/09403

27

PCT/FR95/01196

CLAIMS

1. Process for the preparation of racemic α -substituted 4-methylthiobutyric acids, characterized in that the racemic α -substituted
5 4-methylthiobutyronitriles are hydrolysed by a nitrilase chosen from those of Alcaligenes faecalis, Rhodococcus sp. HT 29-7 or Gordona terrae.
2. Process according to claim 1, characterized in that the racemic α -substituted
10 4-methylthiobutyronitrile is 4-methylthio-2-hydroxybutyronitrile.
3. Process according to claim 1, characterized in that the pH of the medium is between 4 and 11 and preferably between 5 and 9.
- 15 4. Process according to claim 1, characterized in that the hydrolysis temperature is between 30 and 60°C and preferably 30 and 50°C.
5. Process according to claim 1, characterized in that the concentration of
20 α -substituted 4-methylthiobutyronitrile in the starting solution is between 10 and 400 mmol and preferably between 50 and 200 mmol per litre.
6. Process according to claim 1, characterized in that the concentration of
25 α -substituted 4-methylthiobutyric acid in the solution is between 10 and 2000 mmol per litre and preferably between 100 and 1500 mmol per litre.
7. Process according to claim 1,





characterized in that a nitrilase belonging to a
microorganism of the genus *Alcaligenes* is used.

8. Process according to claim 1,
characterized in that a nitrilase belonging to a
5 microorganism of the genus *Rhodococcus* is used.

9. Process according to claim 1,
characterized in that a nitrilase belonging to a
microorganism of the genus *Gordona* is used.

