

Description

POLYPEPTIDE HAVING AMIDASE ACTIVITY AND GENE THEREOF

5 Technical Field

The present invention relates to a polypeptide having amidase activity, a gene thereof, a microorganism or a transformant that is able to produce the above polypeptide thereof, and a method for producing amidase using the above
10 microorganism or the transformant thereof.

Background Art

Amidase (enzyme No. [EC3.5.1.4]) is an enzyme, which hydrolyzes a carboxylic amide to carboxylic acid and ammonia.
15 It has been known that this enzyme has industrially useful activities. For example, this enzyme stereoselectively hydrolyzes a racemic α -amino acid amide to generate an optically active α -amino acid. Such an optically active α -amino acid is a compound that is useful as a synthetic intermediate of
20 pharmaceuticals or a sweetener. It has also been reported that amidase catalyzes hydrolysis of an ester as a substrate (Non-Patent Document 1).

A large number of methods for producing optically active α -amino acids have been known. However, the mass production of

D- α -amino acids via a fermentation method has been considered particularly difficult. Thus, it has been desired that an inexpensive efficient synthetic method be developed. For example, as a method for synthesizing D- α -amino acid via biological means, the following synthetic methods, which involve the use of a microorganism or an enzyme having amidase activity and optical resolution of a racemic α -amino acid amide, have been known.

(1) A method for producing a D- α -amino acid from a racemic α -amino acid amide, using activity for specifically hydrolyzing the D- α -amino acid amide of a microorganism belonging to genus *Rhodococcus* (Patent Document 1).

(2) A method for producing a D- α -amino acid from a racemic α -amino acid amide, using aminopeptidase produced by a microorganism belonging to genus *Achromobacter*, *Corynebacterium*, *Flavobacterium*, *Bacillus*, *Micrococcus*, *Cellulomonas*, *Pseudomonas*, *Protaminobacter*, *Mycobacterium*, *Arthrobacter*, or *Streptomyces* (Patent Document 2).

(3) A method for producing D-alanine from a racemic alanine amide, using amidase produced by a microorganism belonging to genus *Arthrobacter* (Patent Documents 3 and 4).

(4) A method for producing a D- α -amino acid from a racemic α -amino acid amide, using activity for specifically hydrolyzing

the D- α -amino acid amide of a microorganism belonging to genus *Achromobacter* (Patent Document 5).

(5) A method for producing a D- α -amino acid from a racemic α -amino acid amide, using a transformant having a D-amino acid
5 amidase gene derived from a microorganism, which has been modified by genetic engineering (Patent Document 6).

However, in the case of the methods described in (1) to (4) above, since the hydrolytic activity of the microorganisms used is low, it cannot be said that these methods are satisfactory
10 for industrial applications.

In addition, in the case of the method described in (5) above, only D-amino acid amidase derived from bacteria belonging to genus *Ochrobactrum* is described, and bacteria belonging to genus *Arthrobacter* are not described at all.

15 It has been generally known that bacteria belonging to genus *Arthrobacter* produce amidase. The enzyme has been purified and isolated, and the properties thereof have been then clarified (Patent Documents 3 and 4 and Non-Patent Documents 2 and 3).
However, the amino acid sequence of the above enzyme and the
20 nucleotide sequence of a gene encoding the above enzyme have not yet been reported.

[Patent Document 2] Japanese Patent Publication No.
7-106149

[Patent Document 3] U.S. Patent No. 5130240

[Patent Document 4] U.S. Patent No. 5252470

5 [Patent Document 5] Japanese Patent Laid-Open No. 2-234678

[Patent Document 6] Japanese Patent Laid-Open No.
2002-253256

[Non-Patent Document 1] Eur. J. Biochem., 2000, Vol. 267,
p. 2028

10 [Non-Patent Document 2] Bioscience, Biotechnology and
Biochemistry, 1992, Vol. 56, No. 12, p. 1980

[Non-Patent Document 3] Agricultural and Biological
Chemistry, 1982, Vol. 46, No. 5, p. 1175

15 Disclosure of the Invention

Problems to be Solved by the Invention

Taking into consideration the aforementioned circumstances,
it is an object of the present invention to provide a novel
D-amidase, which can be used for the optical resolution of a
20 racemic amino acid amide and a racemic amino acid ester. It is
another object of the present invention to determine the amino
acid sequence of the above D-amidase and the nucleotide sequence
of the gene thereof to provide a microorganism or a transformant

that is able to produce the above enzyme, and a method for producing the above enzyme using them.

Means for Solving the Problems

5 As a result of intensive studies directed towards achieving the aforementioned objects, the present inventors have separated from the soil, bacteria belonging to genus *Arthrobacter*, which produce a novel D-amidase that stereoselectively hydrolyzes a D-amino acid amide and a D-amino acid ester. Thereafter, the
10 inventors have isolated and purified the above amidase from the above bacteria, so that they have achieved isolation of an amidase gene and its expression in host microorganisms, thereby completing the present invention.

That is to say, the present invention relates to a
15 polypeptide described in the following (a) or (b):

(a) a polypeptide, which has the amino acid sequence shown in SEQ ID NO: 1 of the sequence listing; or

(b) a polypeptide, which has an amino acid sequence formed by comprising a substitution, insertion, deletion, and/or addition
20 of one or more amino acids in the amino acid sequence shown in SEQ ID NO: 1 of the sequence listing, and which has amidase activity.

The present invention also relates to DNA encoding the above polypeptide.

The present invention also relates to DNA described in any one of the following (c) to (e):

(c) DNA, which has the nucleotide sequence shown in SEQ ID NO: 3 of the sequence listing;

5 (d) DNA, which hybridizes with DNA having a nucleotide sequence that is complementary to the nucleotide sequence shown in SEQ ID NO: 3 of the sequence listing under stringent conditions, and which encodes a polypeptide having amidase activity; and

(e) DNA, which has a nucleotide sequence resulting from
10 substitution, insertion, deletion, and/or addition of one or more nucleotides in the nucleotide sequence shown in SEQ ID NO: 3 of the sequence listing, and which encodes a polypeptide having amidase activity.

In addition, the present invention relates to a recombinant
15 plasmid, which is obtained by insertion of the above DNA into a vector.

Moreover, the present invention relates to a transformant, which is obtained by transformation of a host microorganism with the above DNA or recombinant plasmid.

20 Furthermore, the present invention relates to a microorganism, which is able to produce the above polypeptide and belongs to genus *Arthrobacter*.

Still further, the present invention relates to a method for producing amidase, which comprises culturing the above

microorganism or the above transformant that is able to produce a polypeptide, accumulating the above polypeptide in the culture, and collecting the above polypeptide.

5 Advantages of the Invention

The present invention comprises the aforementioned structures and enables efficient production of novel amidase. In addition, the present invention enables efficient production of D-amino acid from an amino acid amide or an amino acid ester,
10 using the above amidase, or microorganisms that produce the above amidase.

Brief Description of the Drawings

Figure 1 is a view of a recombinant plasmid pHA001 comprising
15 the amidase gene of the present invention; and

Figure 2 is a view of a recombinant plasmid pHA002 comprising the amidase gene of the present invention.

Best Mode for Carrying Out the Invention

20 The present invention will be described in detail below.

First, the polypeptide of the present invention will be described. The polypeptide of the present invention is a polypeptide, which has amidase activity and enables the

stereoselective hydrolysis of a D-amino acid amide and a D-amino acid ester.

In the present invention, the amidase activity of the polypeptide can be detected and measured by carrying out a
5 reaction at 30°C in a tris(hydroxymethyl)aminomethane (Tris)-HCl buffer solution (pH 8.5) that contains a 50 mM D-phenylalanine amide, and then quantifying the generated phenylalanine by high performance liquid chromatography (HPLC) or the like.

The polypeptide of the present invention can be obtained
10 from a microorganism having amidase activity. The type of such a microorganism is not particularly limited, as long as it produces the above polypeptide. Examples of such a microorganism include microorganisms belonging to genus *Arthrobacter*. Among others, *Arthrobacter* sp. are preferable,
15 and *Arthrobacter* sp. KNK1101J, which has been newly isolated from the soil by the present inventors, is more preferable.

The aforementioned *Arthrobacter* sp. KNK1101J has been deposited with the National Institute of Advanced Industrial Science and Technology, an Independent Administrative
20 Institution under the Ministry of Economy, Trade and Industry, at the AIST (Tsukuba Central 6, Higashi 1-1-1, Tsukuba, Ibaraki, Japan) (accession No. FERM BP-10192: a national deposition on October 22, 2003 has been transferred to an international deposition under the provisions of the Budapest Treaty).

The mycological properties of the *Arthrobacter* sp. KNK1101J will be described below.

1. Form

- 1) Bacillus having a diameter of approximately 0.7 to 0.8 μm
5 and a length of approximately 1.0 to 1.5 μm
- 2) Gram staining: positive
- 3) Motility: no
- 4) Sporulation: no
- 5) Cell polymorphism: no
- 10 6) Colony form on broth agar plate medium: round shape, smooth entire fringe, low convex, lustrous, and light yellow color

2. Cultural properties

- 1) Nutrient agar (manufactured by Oxoid) medium
15 Color: no; luster: no; pigment production: no
- 2) Nutrient broth (manufactured by Oxoid) medium
Surface growth: no; turbidity of medium: yes
- 3) Gelatin stab culture
Growth: no; gelatin change: no
- 20 4) Litmus milk
Coagulation: no; liquefaction: no

3. Physiological test

- 1) Nitrate reduction: -

- 2) Denitrification reaction: -
- 3) MR test: -
- 4) VP test: -
- 5) Indole generation: -
- 5 6) Hydrogen sulfide generation: -
- 7) Hydrolysis of starch: +
- 8) Use of citric acid Koser: +; Christensen: +
- 9) Use of inorganic nitrogen Nitrate: +; ammonium salt: +
- 10) Urease activity: -
- 10 11) Oxidase activity: -
- 12) Catalase activity: +
- 13) β -galactosidase activity: +
- 14) Arginine dihydrolase activity: -
- 15) Lysine decarboxylase activity: -
- 15 16) Tryptophan deaminase activity: -
- 17) Gelatinase activity: -
- 18) Pyrazinamidase activity: +
- 19) Pyrrolidonyl allyl amidase activity: +
- 20) Alkaline phosphatase activity: +
- 20 21) β -glucuronidase activity: -
- 22) N-acetyl- β -glucosaminidase activity: -
- 23) α -glucosidase activity: +
- 24) Esculin (β -glucosidase) activity: +
- 25) Gelatin hydrolytic activity: -

26) Growth range

pH 5: +; pH 9: +; pH 10: +

20°C: +; 25°C: +; 40°C: +W; 45°C: -

27) Growth conditions

5 Aerobic: +; Anaerobic: -

28) O-F test (oxidation/fermentation): -/-

29) Generation of acid and gas from sugars (acid/gas)

L-arabinose: -/-

D-glucose: +/-

10 D-fructose: +/-

Maltose: -/-

Lactose: -/-

D-sorbitol: -/-

Inositol: -/-

15 D-xylose: +/-

D-mannose: +/-

D-galactose: +/-

Sucrose: +/-

Trehalose: +W/-

20 D-mannitol: +/-

Glycerin: +/-

30) Fermentation test

Glucose: -

Ribose: -

Xylose: -

Mannitol: -

Maltose: -

Lactose: -

5 Saccharose: -

Glycogen: -

31) Main menaquinone component: MK-9 (H₂)

32) Fatty acid composition in cell mass

10 Table 1

fatty acid		component ratio(%)
C15 : 0	ANTEISO	60.58
C15 : 0	ISO	11.44
C16 : 0	ISO	10.63
C17 : 0	ANTEISO	7.06
C16 : 0		4.14
C14 : 0	ISO	3.13
C14 : 0		1.88
C17 : 0		1.14

The microorganism that produces the polypeptide of the present invention may be either the wild strain of the aforementioned microorganism, or a mutant strain obtained by mutation and improvement. Such a mutant strain can be obtained by methods known to persons skilled in the art, such as UV irradiation or treatment with agents such as

N-methyl-N'-nitro-N-nitrosoguanidine (NTG) or ethyl methanesulfonate (EMS).

The type of a medium used to culture the microorganism that produces the polypeptide of the present invention is not particularly limited, as long as the above microorganism can grow therein. Ordinary liquid mediums that contain the following nutrients can be used. That is, examples of a carbon source may include: carbohydrates such as glucose or sucrose; alcohols such as ethanol or glycerol; fatty acids such as oleic acid or stearic acid and the esters thereof; and oils such as rapeseed oil or soybean oil. Examples of a nitrogen source may include ammonium sulfate, sodium nitrate, peptone, casamino acid, corn steep liquor, bran, and yeast extract. Examples of inorganic salts may include magnesium sulfate, sodium chloride, calcium carbonate, dipotassium hydrogen phosphate, and potassium dihydrogen phosphate. Examples of other suitable nutrients may include malt extract and meat extract.

Moreover, small quantities of substances for enhancing the production of amidase, which include amides such as an amino acid amide or a fatty acid amide, or esters such as an amino acid ester or a fatty acid ester, can also be added. The concentration of such an amidase production-enhancing substance contained in a medium is between 0.001% by weight and 10% by weight, and preferably between 0.01% by weight and 1% by weight.

The culture can be aerobically carried out within a temperature range between 10°C and 60°C, and preferably between 20°C and 50°C, and a pH range between pH 3 and 11, and preferably between pH 5 and 9. The culture period is between approximately 1 and 5 days. In addition, either a batch culture method or a continuous culture method may be applied.

After completion of the culture, the cells are harvested from the culture broth by centrifugation or the like, and the harvested cells are then disrupted by means such as ultrasonic disruption to obtain a crude enzyme solution. Thereafter, this crude enzyme solution is purified by salting-out, column chromatography, or the like to obtain the polypeptide of the present invention.

As stated above, the polypeptide of the present invention may be either a natural enzyme obtained from microorganisms, or a recombinant enzyme produced utilizing genetic recombination technology. An example of such a natural enzyme is a polypeptide having the amino acid sequence shown in SEQ ID NO: 1 of the sequence listing.

In addition, the polypeptide of the present invention may also be a polypeptide, which has an amino acid sequence resulting from substitution, insertion, deletion, and/or addition of one or more amino acids in the amino acid sequence shown in SEQ ID NO: 1 of the sequence listing, and which has amidase activity.

An "amino acid sequence resulting from deletion, addition, insertion, and/or substitution of one or more amino acids" can be obtained by deleting, adding, inserting, and/or substituting amino acid(s) according to methods known to persons skilled in
5 the art, such as a site-directed mutagenesis. Specifically, such methods are described in publications such as Nucleic Acid Res. 10, 6487 (1982), and Methods in Enzymology 100, 448 (1983).

Moreover, the term "polypeptide, which has amidase activity" is used to mean a polypeptide exhibiting 10% or more
10 of the amidase activity of a polypeptide having the amino acid sequence shown in SEQ ID NO: 1, when measured under the aforementioned activity measurement conditions, preferably 40% or more, more preferably 60% or more, and further more preferably 80% or more.

15 Next, the DNA of the present invention will be described. The DNA of the present invention may be DNA encoding the aforementioned polypeptide. An example is DNA having the nucleotide sequence shown in SEQ ID NO: 3 of the sequence listing.

In addition, the DNA of the present invention may also be
20 either DNA hybridizing with DNA having a nucleotide sequence complementary to the nucleotide sequence shown in SEQ ID NO: 3 of the sequence listing under stringent conditions, or DNA having a nucleotide sequence resulting from substitution, insertion, deletion, and/or addition of one or more nucleotides in the

nucleotide sequence shown in SEQ ID NO: 3. Any type of DNA is included in the DNA of the present invention, as long as it encodes the aforementioned polypeptide having amidase activity.

The expression "DNA hybridizing with DNA having a nucleotide
5 sequence complementary to the nucleotide sequence shown in SEQ
ID NO: 3 of the sequence listing under stringent conditions" is
used herein to mean DNA capable of specifically forming a hybrid
with DNA having nucleotide sequence complementary to the
nucleotide sequence shown in SEQ ID NO: 3 of the sequence listing,
10 when the colony hybridization method, the plaque hybridization
method, the Southern hybridization method, or the like is carried
out.

The term "stringent conditions" is used herein to mean
conditions wherein hybridization is carried out at 65°C in an
15 aqueous solution consisting of 75 mM trisodium citrate, 750 mM
sodium chloride, 0.5% sodium dodecyl sulfate, 0.1% bovine serum
albumin, 0.1% polyvinylpyrrolidone, and 0.1% Ficoll 400
(manufactured by Amersham Biosciences), and wherein washing is
then carried out at 60°C with an aqueous solution consisting of
20 15 mM trisodium citrate, 150 mM sodium chloride, and 0.1% sodium
dodecyl sulfate, for example. Such stringent conditions are
preferably conditions wherein after hybridization has been
carried out under the aforementioned conditions, washing is
carried out at 65°C with an aqueous solution consisting of 15

mM trisodium citrate, 150 mM sodium chloride, and 0.1% sodium dodecyl sulfate. Such stringent conditions are more preferably conditions wherein after hybridization has been carried out under the aforementioned conditions, washing is carried out at 65°C
5 with an aqueous solution consisting of 1.5 mM trisodium citrate, 15 mM sodium chloride, and 0.1% sodium dodecyl sulfate.

Moreover, the term "nucleotide sequence resulting from deletion, addition, insertion, and/or substitution of one or more nucleotides" is used to mean a nucleotide sequence obtained by
10 deletion, addition, insertion, and/or substitution of nucleotides, which can be deleted, added, inserted, and/or substituted by methods known to persons skilled in the art, which are described, for example, in *Tanpaku Kakusan Koso Zokan* (Proteins, Nucleic acids, and Enzymes, Supplement), *Idenshi*
15 *Zofuku PCR Ho* (Gene Amplification PCR Method), 35 (17), 2951-3178 (1990), or *PCR Technology* (1990), edited by Henry A. Erlich, translated by Ikunoshin Kato.

The DNA of the present invention (amidase gene) can be obtained from the aforementioned microorganisms having amidase
20 activity. For example, the following method can be applied to obtain DNA of interest.

First, the amino acid sequence of the N-terminus of amidase that has been purified from microorganisms having amidase activity is determined using a vapor-phase protein sequencer or

the like. Further, protease such as lysyl endopeptidase is allowed to act on the purified amidase, so that the amidase can be digested into a polypeptide with a suitable size, and the obtained polypeptide is then purified by HPLC or the like to
5 determine an internal amino acid sequence according to the aforementioned method. Based on the resultant N-terminal amino acid sequence and internal amino acid sequence, DNA primers are designed and synthesized.

Thereafter, chromosomal DNA is isolated from microorganisms
10 used as origins of amidase. Such chromosomal DNA is obtained by dissolving the cultured cells in a surfactant, cetyltrimethylammonium bromide (CTAB), chloroform, phenol, or the like, treating the cells therein, precipitating the extracted DNA with isopropanol, and then washing with ethanol the
15 precipitate obtained by centrifugation (refer to Current Protocols in Molecular Biology (Greene Publishing Associates and Wiley-Interscience), for example).

Such chromosomal DNA is used as a template, and PCR is carried out using the aforementioned DNA primers to obtain a portion of
20 the gene of interest.

Subsequently, a DNA fragment encoding the N- and C-termini of the obtained partial gene can be obtained by the inverse PCR method (refer to Nucleic Acids Res. 16, 8186 (1988), for example). The nucleotide sequence of the above DNA fragment is determined.

Thereafter, based on the nucleotide sequence of a portion presumably located upstream of the N-terminus of the enzyme and the nucleotide sequence of a portion presumably located downstream of the C-terminus thereof, DNA primers are prepared.

5 Using such DNA primers, PCR is carried out with the obtained chromosomal DNA as a template to obtain a DNA fragment that contains the full-length amidase gene of interest.

Thereafter, the obtained DNA fragment that contains an amidase gene is allowed to bind to vector DNA using T4 DNA ligase,
10 thereby obtaining a recombinant plasmid. Using this plasmid, the nucleotide sequence of the DNA fragment portion containing an amidase gene, which has been inserted into the vector, is analyzed to confirm the presence of nucleotides that encode the N-terminal amino acid sequence and internal amino acid sequence
15 of an amidase enzyme. Moreover, a translation initiation site and a termination codon are confirmed to determine an open reading frame.

Host microorganisms are transformed with the thus obtained DNA or with a plasmid obtained by incorporation of the above DNA
20 into a vector to obtain a transformant.

As a host and a vector, the host-vector system described in "*Kumikae DNA Jikken Shishin* (Recombinant DNA Experimental Guideline)" (edited by Science and Technology Agency, Research and Development Bureau, Life Sciences Division; revised on March

22, 1996) can be used. Examples of a host used herein may include microorganisms belonging to genus *Escherichia*, *Pseudomonas*, *Flavobacterium*, *Bacillus*, *Serratia*, *Corynebacterium*, *Brevibacterium*, *Agrobacterium*, *Acetobacter*, *Gluconobacter*,
5 *Lactobacillus*, *Streptococcus*, or *Streptomyces*.

As a vector, a plasmid, phage, or derivative thereof derived from microorganisms, which is capable of autonomously replicating in the aforementioned host, can be used. Among others, *Escherichia coli* is preferably used as a host
10 microorganism, and a vector capable of autonomously replicating in such a microorganism is preferably used as a vector. Examples of such a vector may include pUC18, pUC19, pBR322, pACYC184, pSC101, pT7Blue, and pUCNT. Moreover, a vector, which is modified to have a strong constitutive promoter for increasing
15 the production amount of enzyme, can also be used.

As an example of a transformant, *Escherichia coli* HB101 is transformed with a recombinant plasmid pHA002 obtained by incorporation of the DNA as obtained above into pUCNT (refer to WO94/03613) to obtain a transformant, *Escherichia coli* HB101
20 (pHA002).

The transformant *Escherichia coli* HB101 (pHA002) obtained by the present invention has been deposited with the National Institute of Advanced Industrial Science and Technology, an Independent Administrative Institution under the Ministry of

Economy, Trade and Industry (under accession No. FERM BP-10193; a national deposition with the original deposition date of January 22, 2004 has been transferred to an international deposition under the provisions of the Budapest Treaty).

5 The recombinant DNA technology used in the present invention is well known in the related field. For example, such a technology is described in Molecular Cloning 2nd Edition (Cold Spring Harbor Laboratory Press, 1989), and Current Protocols in Molecular Biology (Greene Publishing Associates and
10 Wiley-Interscience).

 The microorganism capable of producing the amidase of the present invention (*Arthrobacter* sp. KNK1101J, a mutant thereof, the aforementioned transformant, etc.) is cultured to produce the above enzyme at a high level. Thus, this technique can be
15 used for production of D-amino acid.

 Such a microorganism may be cultured in a common medium. As a medium used for the culture, a common medium that contains nutrients such as a carbon source, a nitrogen source, and inorganic salts, may be used. When a trace amount of organic
20 nutrient such as vitamin or amino acid is added, preferred results can often be obtained. Examples of a carbon source, which is appropriately used, may include: carbohydrates such as glucose or sucrose; organic acids such as acetic acid; and alcohols. Examples of a nitrogen source used herein may include ammonium

salts, ammonia water, ammonia gas, urea, yeast extract, peptone, and corn steep liquor. Examples of inorganic salts used herein may include phosphates, magnesium salts, potassium salts, sodium salts, calcium salts, iron salts, sulfates, and chlorides.

5 The culture may be carried out within a temperature range between 25°C and 40°C. Such a temperature range is particularly preferably between 25°C and 37°C. In addition, the culture may be carried out within a pH range between 4 and 8. Such a pH range is preferably between 5 and 7.5. Moreover, either a batch culture
10 method or a continuous culture method may be applied.

Treatments for inducing the enzyme, such as addition of isopropyl-1-thio- β -D-galactoside (IPTG) or lactose, can also be carried out, as necessary.

When the amidase of the present invention is used for the
15 stereoselective hydrolysis of an amino acid amide or amino acid ester, the culture as obtained above (a culture broth, cells of microorganisms) may be allowed to directly act thereon.

Otherwise, amidase may be isolated and purified from the culture before use.

20

Examples

Specific examples of the present invention will be given below. However, these examples are not intended to limit the scope of the present invention.

(Example 1) Isolation and purification of amidase

The *Arthrobacter* sp. KNK1101J (accession No. FERM BP-10192) was inoculated into 6 ml of medium A (produced by mixing 10 g of meat extract, 5 g of yeast extract, 10 g of polypeptone, and 3 g of sodium chloride in deionized water to total 1 L in volume; pH 6.5 before sterilization), which had been sterilized in a test tube. Thereafter, the strain was subjected to an aerobic shaking culture at 30°C for 24 hours. Thereafter, 2 ml of the culture broth was inoculated into 200 ml of medium A, which had been sterilized in a flask, and it was then subjected to an aerobic shaking culture at 30°C for 48 hours. After completion of the culture, the cells were harvested by centrifugation, and they were then suspended in a 50 mM Tris-HCl buffer solution (pH 8.0) contained 1 mM dithiothreitol (DTT). The cells were then disrupted with ultrasonic wave, followed by centrifugation. Thereafter, ammonium sulfate was added to the supernatant, resulting in a 45% of the saturated solution, and the generated precipitate was then obtained by centrifugation. This fraction was dissolved in a 50 mM Tris-HCl buffer solution (pH 8.0) containing 1 mM DTT and dialyzed in the buffer solution. Thereafter, the resultant solution was applied to DEAE-TOYOPEAL (manufactured by Tosoh Corporation) to conduct column chromatography. After washing with the same above buffer

solution, fractions with activity were collected by elution via concentration gradient from 0 M to 0.5 M sodium chloride with the above buffer solution. Ammonium sulfate was dissolved in such active fractions to a final concentration of 0.8 M, and the
5 resultant was then applied to Phenyl-TOYOPEAL (manufactured by Tosoh Corporation) to conduct column chromatography. Thus, fractions were collected by elution via concentration gradient from 0.8 M to 0 M ammonium sulfate with a 50 mM Tris-HCl buffer solution (pH 8.0) that contained 1 mM DTT. The obtained active
10 fractions were applied to MonoQ HR5/5 (manufactured by Amersham Pharmacia Biotech) to conduct column chromatography. After washing with a 50 mM Tris-HCl buffer solution (pH 8.0) that contained 1 mM DTT, fractions were collected by elution via concentration gradient from 0 M to 0.5 M sodium chloride with
15 the above buffer solution. The obtained active fractions were concentrated with an ultrafiltration membrane (molecular weight cut off: 10,000). Thereafter, the concentrate was applied to Superdex 200 HR 10/30 (manufactured by Amersham Pharmacia Biotech) to conduct gel filtration via FPLC. Fractions were then
20 eluted with a 50 mM Tris-HCl buffer solution (pH 8.0) that contained 1 mM DTT and 0.15 M sodium chloride. The obtained active fractions were analyzed by SDS-polyacrylamide electrophoresis. As a result, amidase was detected in the form

of a single band, and thus the purity of the purified enzyme could be confirmed.

(Example 2) Hydrolysis of amino acid amide with amidase

5 0.1 ml of the purified amidase active fraction obtained in
Example 1 was mixed with 0.1 ml of a 0.2 M Tris-HCl buffer solution
(pH 8.5) that contained 0.5% of the compound shown in Table 2.
The mixture was shaken at 30°C for 15 hours. After completion
of the reaction, solids were eliminated by centrifugation. The
10 yield and optical purity of the generated amino acid were analyzed
by high performance liquid chromatography. The results are
shown in Table 2.

Conditions for analysis by high performance liquid
15 chromatography

[Analysis of yield of amino acid]

Column: COSMOSIL 5C18-AR (4.6 mmφ × 250 mm; manufactured by
Nacalai Tesque, Inc.)

Eluent: 10 mM potassium phosphate buffer solution (pH
20 2)/acetonitrile = 19/1

Flow rate: 0.5 ml/min.

Column temperature: 40°C

Measurement wavelength: 210 nm

[Analysis of optical purity of amino acid]

Column: SUMICHIRAL OA-5000 (4.6 mmφ × 150 mm; manufactured by Sumika Chemical Analysis Service, Ltd.)

Eluent: 2 mM copper sulfate/acetonitrile = 85/15

Flow rate: 0.8 ml/min.

5 Column temperature: 40°C

Measurement wavelength: 254 nm

Table 2

substrate	Generated amino acid		
	yield (mol%)	configuration	optical purity (%ee)
DL-phenylalanine amide hydrochloride	47.2	D	98.5
DL-leucine amide hydrochloride	3.1	D	88.9

10 (Example 3) Hydrolysis of amino acid ester with purified amidase

0.1 ml of the purified amidase active fraction obtained in Example 1 was mixed with 0.1 ml of a 0.2 M 2-(N-morpholino)ethanesulfonic acid-NaOH buffer solution (pH 6.5) that contained a racemic phenylalanine ethyl ester with a concentration of 0.5%. The mixture was shaken at 30°C for 15 hours. After completion of the reaction, solids were eliminated by centrifugation. The generated amino acid was analyzed by the same method as that described in Example 2. As a result, it was

found that D-phenylalanine was obtained in a yield of 29.7 mol % and at an optical purity of 72.8%ee.

(Example 4) Isolation of amidase gene

5 First, the cells were harvested by culturing the
Arthrobacter sp. KNK1101J according to the same method as that
described in Example 1 were dissolved in CTAB, chloroform and
phenol, and were then treated therewith. Thereafter, the
extracted DNA was precipitated with isopropanol, and the
10 precipitate obtained by centrifugation was then washed with
ethanol to prepare chromosomal DNA (refer to Current Protocols
in Molecular Biology (Greene Publishing Associates and
Wiley-Interscience)). Subsequently, the amino acid sequence of
the amino terminus of the amidase purified in Example 1 was
15 determined using a vapor-phase protein sequencer. Thereafter,
lysyl endopeptidase was allowed to act on the amidase purified
in Example 1 in the presence of 4 M urea. The generated
polypeptide fragment was purified by reverse phase HPLC.
Thereafter, the internal amino acid sequence of the amidase was
20 determined by the same above method. Using a DNA primer
(Primer-1; SEQ ID NO: 4 of the sequence listing) designed based
on the N-terminal amino acid sequence and a DNA primer (Primer-2;
SEQ ID NO: 5 of the sequence listing) designed based on the
internal amino acid sequence, PCR was carried out with the

previously obtained chromosomal DNA as a template. As a result, a portion of the amidase gene of interest (referred to as a partial gene) was obtained.

Subsequently, the following operations were carried out to
5 obtain the full-length gene of interest. Based on the nucleotide sequences corresponding to each of the N-terminal portion and C-terminal portion of the enzyme in the aforementioned partial gene, DNA primers directed towards the external direction of the partial gene (Primer-3 shown in SEQ ID NO: 6 of the sequence
10 listing; and Primer-4 shown in SEQ ID NO: 7 of the sequence listing) were synthesized. Using such primers, inverse PCR was carried out. As a template, DNA, which had been obtained by cyclization with T4 DNA ligase of a fragment obtained by decomposition of the previously obtained chromosomal DNA with
15 restriction enzymes, SacII, PvuI, SalI, and XhoI, was used. As a result of the inverse PCR, a DNA fragment, which contained a gene portion located further outside of the previously obtained partial gene, was obtained. The nucleotide sequence of this DNA fragment was determined. Thereafter, there were prepared a DNA
20 primer (Primer 5; SEQ ID NO: 8 of the sequence listing) having a sequence obtained by binding the restriction site of BamHI to the nucleotide sequence of a portion presumably located upstream of the N-terminus of the enzyme, and a DNA primer (Primer 6; SEQ ID NO: 9 of the sequence listing) having a sequence obtained by

binding the restriction site of SacI to the nucleotide sequence of a portion presumably located downstream of the C-terminus of the enzyme. Using such DNA primers, DNA between the sequences was amplified by PCR to obtain a DNA fragment (SEQ ID NO: 2 of the sequence listing) that contained the full-length amidase gene. The nucleotide sequence of a portion of the obtained DNA fragment was analyzed, and as a result, it was confirmed that the full-length amidase gene (SEQ ID NO: 3 of the sequence listing) was contained therein.

10

(Example 5) Production of recombinant plasmid that contains amidase gene and analysis of the gene

Using T4 DNA ligase, the DNA fragment that contained the full-length amidase gene obtained in Example 4 was ligated to a vector plasmid pT7Blue cleaved with the restriction enzyme EcoRV. As a result, a plasmid pHA001 that contained the amidase gene, as shown in the restriction map of Figure 1, was obtained.

Using the obtained plasmid pHA001, the nucleotide sequence of the DNA fragment obtained in Example 4 was analyzed. As a result, the presence of nucleotides encoding the N-terminal amino acid sequence and internal amino acid sequence determined with the purified amidase was confirmed. In addition, a translation initiation site and a termination codon were confirmed to determine an open reading frame. The nucleotide sequence of the

thus obtained DNA fragment that contained the full-length amidase gene is shown in SEQ ID NO: 2 of the sequence listing. The nucleotide sequence of the open reading frame is shown in SEQ ID NO: 3 of the sequence listing. The amino acid sequence
5 estimated from the nucleotide sequence is shown in SEQ ID NO: 1 of the sequence listing.

(Example 6) Production of recombinant plasmid that expresses amidase gene in high volume

10 Primers (Primer-7 shown in SEQ ID NO: 10 of the sequence listing; and Primer-8 shown in SEQ ID NO: 11 thereof) having sequences obtained by binding the restriction sites of NdeI and SacI to the N-terminal and C-terminal portions of the amidase gene obtained in Example 5, respectively, were used. DNA between
15 them was amplified by PCR to obtain the DNA fragment of the open reading frame shown in SEQ ID NO: 3 of the sequence listing.

The obtained DNA fragment was cleaved with the restriction enzymes NdeI and SacI. Using T4 DNA ligase, the cleaved portion was allowed to bind to a vector plasmid pUCNT (refer to
20 WO94/03613), which had been cleaved with the same above enzymes. As a result, the pHA002 shown in the restriction map of Figure 2, which had been designed to express a larger amount of amidase gene than pHA001 does, was obtained.

(Example 7) Production of transformant using recombinant DNA that contains amidase gene

The plasmid pHA002 obtained in Example 6 was mixed with *Escherichia coli* HB101 competent cells to transform the cells.

5 Thereafter, the cells were plated on agar medium B (produced by mixing 10 g of trypton, 5 g of yeast extract, 10 g of sodium chloride, 15 g of agar, and 100 mg of ampicillin, in 1 L of deionized water; pH 7.0 before sterilization; ampicillin was added after completion of the sterilization) to obtain a
10 transformant, *Escherichia coli* HB101 (pHA002), which contained recombinant DNA that contained an amidase gene, in the form of a colony.

The obtained transformant colony was inoculated into 6 ml of medium C (produced by mixing 10 g of trypton, 5 g of yeast
15 extract, 10 g of sodium chloride, and 100 mg ampicillin, in 1 L of deionized water; pH 7.0 before sterilization; ampicillin was added after completion of the sterilization), which had been sterilized in a test tube. Thereafter, the resultant was shaken at 37°C for 24 hour for an aerobic culture. Thereafter, cells
20 were harvested from the obtained culture broth by centrifugation, and it was then suspended in a 50 mM Tris-HCl buffer solution (pH 8.0) that contained 1 mM dithiothreitol (DTT). The cells were then disrupted with ultrasonic wave, and insoluble products derived from the cells were then eliminated by centrifugation

to obtain an amidase solution of the transformant. 0.1 ml of the obtained enzyme solution was mixed with 0.1 ml of a 0.2 M Tris-HCl buffer solution (pH 8.5) that contained 50 mM D-phenylalanine amide hydrochloride. The obtained mixture was
5 reacted at 30°C for 15 hours. As a result, it was confirmed that D-phenylalanine was generated, and that the transformant had amidase activity.

It is to be noted that the obtained transformant, *Escherichia coli* HB101 (pHA002), has been deposited with the National
10 Institute of Advanced Industrial Science and Technology, an Independent Administrative Institution under the Ministry of Economy, Trade and Industry (under accession No. FERM BP-10193; a national deposition with the original deposition date of January 22, 2004 has been transferred to an international
15 deposition under the provisions of the Budapest Treaty).

(Example 8) Hydrolysis of amino acid ester using recombinant bacteria

0.1 ml of the culture broth of *Escherichia coli* HB101
20 (pHA002) obtained in Example 7 was mixed with 0.1 ml of a 0.2 M 2-(N-morpholino)ethanesulfonic acid-NaOH buffer solution (pH 6.5) that contained 1% of racemic phenylalanine ethyl ester. The obtained mixture was shaken at 30°C for 15 hours. After completion of the reaction, solids were eliminated by

centrifugation, and the generated amino acid was analyzed by the same method as that described in Example 2. As a result, it was found that D-phenylalanine was obtained at a yield of 34.7 mol % and an optical purity of 53.6%ee.

CLAIMS

1. A polypeptide described in the following (a) or (b):

(a) a polypeptide, which has the amino acid sequence shown in SEQ ID NO: 1 of the sequence listing; or

5 (b) a polypeptide, which has an amino acid sequence resulting from substitution, insertion, deletion, and/or addition of one or more amino acids in the amino acid sequence shown in SEQ ID NO: 1 of the sequence listing, and which has amidase activity.

10 2. The polypeptide according to claim 1, which is derived from a microorganism belonging to genus *Arthrobacter*.

3. The polypeptide according to claim 2, wherein the microorganism is *Arthrobacter* sp. KNK1101J (FERM BP-10192).

15

4. DNA, which encodes the polypeptide according to any one of claims 1 to 3.

5. DNA described in any one of the following (c) to (e):

20 (c) DNA, which has the nucleotide sequence shown in SEQ ID NO: 3 of the sequence listing;

(d) DNA, which hybridizes with DNA having a nucleotide sequence that is complementary to the nucleotide sequence shown in SEQ

ID NO: 3 of the sequence listing under stringent conditions, and which encodes a polypeptide having amidase activity; and

(e) DNA, which has a nucleotide sequence resulting from substitution, insertion, deletion, and/or addition of one or more
5 nucleotides in the nucleotide sequence shown in SEQ ID NO: 3 of the sequence listing, and which encodes a polypeptide having amidase activity.

6. A recombinant plasmid, which is obtained by insertion of
10 the DNA according to claim 4 or 5 into a vector.

7. The recombinant plasmid according to claim 6, wherein the vector is pUC18, pUC19, pBR322, pACYC184, pSC101, pT7Blue, or pUCNT.

15

8. The recombinant plasmid according to claim 6 or 7, which is pHA002 specified in the restriction map shown in Figure 2.

9. A transformant, which is obtained by transformation of a
20 host microorganism with the recombinant plasmid according to any one of claims 6 to 8.

10. The transformant according to claim 9, wherein the host microorganism is *Escherichia coli*.

11. The transformant according to claim 9, which is *Escherichia coli* HB101 (pHA002) (FERM BP-10193).

5 12. A microorganism, which is able to produce the polypeptide according to claim 1 and belongs to genus *Arthrobacter*.

13. The microorganism according to claim 12, which is *Arthrobacter* sp. KNK1101J (FERM BP-10192) or a mutant thereof.

10

14. A method for producing amidase, comprising culturing a microorganism that is able to produce the polypeptide according to any one of claims 1 to 3, accumulating said polypeptide in the culture, and collecting it.

15

15. The production method according to claim 14, wherein the microorganism is the transformant according to any one of claims 9 to 11.

20 16. The production method according to claim 14, wherein the microorganism is the microorganism according to claim 12 or 13.

SEQUENCE LISTING

<110> KANEKA CORPORATION

<120> POLYPEPTIDE HAVING AMIDASE ACTIVITY AND GENE THEREOF

<130> B040008W001

<150> JP2004-028041

<151> 2004-2-4

<160> 11

<170> PatentIn version 3.1

<210> 1

<211> 388

<212> PRT

<213> Arthrobacter sp.

<400> 1

```

Met Ser Arg Leu Leu Arg Glu His Gly Ile Val Ile Gly Arg Leu Gln
1      5      10      15
Pro Gly Ser Leu Asn Thr Ile Ala Asp Val Ala Gly Val Arg Val Gly
      20      25      30
His Ser Thr Ile Met Arg Gly Ser Gly Pro Leu Ser Ile Gly His Gly
      35      40      45
Pro Val Arg Thr Gly Val Thr Ala Ile Ile Pro His Glu Gly Asp Ile
      50      55      60
Trp Glu Glu Pro Arg Phe Ala Gly Val Phe Ser Leu Asn Gly Ser Gly
65      70      75      80
Glu Trp Ser Gly Thr Ser Phe Val Arg Glu Thr Gly Cys Leu Tyr Gly
      85      90      95
Pro Ile Met Thr Thr Asn Ser His Ser Ile Gly Ser Val Arg Asn Ala
      100      105      110
Val Ile Lys Arg Glu Val Ala Arg Arg Gly Ser Leu Glu Arg Leu Pro
      115      120      125
Leu Val Gly Glu Thr Phe Asp Gly Leu Leu Asn Asp Ile Ser Gly Met
      130      135      140
His Val Lys Asp Glu His Val Ala Glu Ala Ile Asp Ser Ala Ser Ala
145      150      155      160
Asn Val Thr Glu Gly Asn Val Gly Gly Gly Thr Gly Asn Val Cys His
      165      170      175
Gly Phe Lys Gly Gly Ile Gly Ser Ala Ser Arg Val Leu Gln Leu Gly
      180      185      190
Glu Glu Thr Tyr Thr Leu Gly Val Leu Val Gln Ala Asn His Gly Leu
      195      200      205
Arg Asp Glu Phe Gln Val Thr Gly Val Pro Val Gly Arg Leu Ile Ser
      210      215      220
Thr Asp Glu Ile Pro Leu Gly Pro Ser Gly Phe Asp Arg Arg Ser Ser
225      230      235      240
Pro His Lys Asn Ser Ile Leu Val Val Val Ala Thr Asp Ala Pro Leu
      245      250      255
Leu Pro Gly Gln Leu Glu Arg Val Ala His Arg Ser Thr Leu Gly Ile
      260      265      270
Ala Arg Asn Gly Ala Tyr Ala His Asn Leu Ser Gly Asp Leu Ala Leu
      275      280      285
Ala Phe Ser Thr Cys Pro Gln Pro Val Ser Gly Tyr Asp Phe Gly Val
290      295      300
Asp Thr Ser Pro Gly Thr Ile Arg Ala Leu Pro Asn Ala Ala Thr Ala
305      310      315      320

```

Gly Leu Phe Glu Ala Ala Val Glu Ala Thr Glu Glu Ala Ile Val Ser
 325 330 335
 Ala Leu Val His Ala Asp Thr Cys Thr Gly Ile Asp Asp Arg Val Ala
 340 345 350
 Tyr Gly Leu Glu Ala Ala Arg Leu Ala Arg Ser Ile Ser Glu Tyr Arg
 355 360 365
 Gly Thr Gln Leu Tyr Pro Glu Lys Val Ser Asp Ser His Leu Glu Arg
 370 375 380
 Arg Ser Gln Pro
 385

<210> 2
 <211> 1290
 <212> DNA
 <213> Arthrobacter sp.

<220>
 <221> CDS
 <222> (53).. (1219)
 <223>

<400> 2
 agcgcgtcgt ggactgggtg cagaaataca caggcgagcc cgaggacgaa aa atg agc 58
 Met Ser
 1
 cgt ctg ctc cgt gag cac gga atc gtc atc ggt cgt ctc caa ccg ggt 106
 Arg Leu Leu Arg Glu His Gly Ile Val Ile Gly Arg Leu Gln Pro Gly
 5 10 15
 tct ctg aac acc att gca gac gtc gca ggc gtt cgc gta ggc cat tca 154
 Ser Leu Asn Thr Ile Ala Asp Val Ala Gly Val Arg Val Gly His Ser
 20 25 30
 aca atc atg cgc ggt tct ggg ccc ctg tcc atc ggc cat ggc cca gtt 202
 Thr Ile Met Arg Gly Ser Gly Pro Leu Ser Ile Gly His Gly Pro Val
 35 40 45 50
 cgc aca ggg gta aca gcc atc atc cct cac gaa gga gac atc tgg gag 250
 Arg Thr Gly Val Thr Ala Ile Ile Pro His Glu Gly Asp Ile Trp Glu
 55 60 65
 gag cca cgg ttc gcc ggc gtc ttc tcc ctg aat ggc agc ggt gaa tgg 298
 Glu Pro Arg Phe Ala Gly Val Phe Ser Leu Asn Gly Ser Gly Glu Trp
 70 75 80
 agc gga acc tcg ttc gtc agg gag acc ggg tgt ctt tat ggc cct atc 346
 Ser Gly Thr Ser Phe Val Arg Glu Thr Gly Cys Leu Tyr Gly Pro Ile
 85 90 95
 atg acg acg aat tcg cac agc att gga tcg gtc agg aac gcc gtc atc 394
 Met Thr Thr Asn Ser His Ser Ile Gly Ser Val Arg Asn Ala Val Ile
 100 105 110
 aag cgt gaa gta gcc cgg cgg gga agc ctg gag agg ctc cct ctc gtg 442
 Lys Arg Glu Val Ala Arg Arg Gly Ser Leu Glu Arg Leu Pro Leu Val
 115 120 125 130
 gga gaa acc ttt gat ggc cta ctc aat gac atc agc ggc atg cac gtc 490
 Gly Glu Thr Phe Asp Gly Leu Leu Asn Asp Ile Ser Gly Met His Val
 135 140 145
 aag gac gaa cac gtg gcc gag gcc atc gac tcc gcc tcc gca aat gtt 538
 Lys Asp Glu His Val Ala Glu Ala Ile Asp Ser Ala Ser Ala Asn Val
 150 155 160
 acc gaa ggc aat gtt ggc ggt ggg acc gga aat gtt tgt cac ggt ttc 586
 Thr Glu Gly Asn Val Gly Gly Gly Thr Gly Asn Val Cys His Gly Phe
 165 170 175
 aaa ggc ggt att gga agt gcc tcg cgc gtg ttg caa ttg ggc gag gaa 634
 Lys Gly Gly Ile Gly Ser Ala Ser Arg Val Leu Gln Leu Gly Glu Glu
 180 185 190

acc tac act ttg ggg gtt ctc gtc caa gcc aac cac ggc ctt cgt gac 682
Thr Tyr Thr Leu Gly Val Leu Val Gln Ala Asn His Gly Leu Arg Asp
195 200 205 210

gaa ttt cag gtg acg gga gta ccc gtg gga agg ctc ata tct acg gac 730
Glu Phe Gln Val Thr Gly Val Pro Val Gly Arg Leu Ile Ser Thr Asp
215 220 225

gag atc ccc ttg ggg cct tca ggt ttt gat cga agg tct tca cca cac 778
Glu Ile Pro Leu Gly Pro Ser Gly Phe Asp Arg Arg Ser Ser Pro His
230 235 240

aaa aac agt att ctt gtc gtc gtg gca acc gac gcg cct cta cta ccg 826
Lys Asn Ser Ile Leu Val Val Val Ala Thr Asp Ala Pro Leu Leu Pro
245 250 255

ggc caa ctg gaa cgc gtt gct cac cgt tct acc cta ggc att gcc cgt 874
Gly Gln Leu Glu Arg Val Ala His Arg Ser Thr Leu Gly Ile Ala Arg
260 265 270

aat ggt gcc tac gcg cac aat ctc agc ggc gac ctt gca ctt gcg ttc 922
Asn Gly Ala Tyr Ala His Asn Leu Ser Gly Asp Leu Ala Leu Ala Phe
275 280 285 290

tcc acc tgc ccg cag cct gta agc ggt tac gat ttc gga gtg gat aca 970
Ser Thr Cys Pro Gln Pro Val Ser Gly Tyr Asp Phe Gly Val Asp Thr
295 300 305

agt cct ggg acc att cgc gcc ctg ccc aac gcc gca acg gct ggc ctc 1018
Ser Pro Gly Thr Ile Arg Ala Leu Pro Asn Ala Ala Thr Ala Gly Leu
310 315 320

ttc gag gcg gcc gtt gag gcc act gag gaa gcg att gtt tcc gcg ctt 1066
Phe Glu Ala Ala Val Glu Ala Thr Glu Glu Ala Ile Val Ser Ala Leu
325 330 335

gtc cac gcc gac acc tgc acc ggg atc gat gac agg gtt gcc tat ggg 1114
Val His Ala Asp Thr Cys Thr Gly Ile Asp Asp Arg Val Ala Tyr Gly
340 345 350

ttg gag gcg gct cga ctt gct cgt tca att tgc gaa tat cga ggc acc 1162
Leu Glu Ala Ala Arg Leu Ala Arg Ser Ile Ser Glu Tyr Arg Gly Thr
355 360 365 370

cag ctg tat ccg gag aaa gtg tgc gat tcc cat ctt gaa cga agg agc 1210
Gln Leu Tyr Pro Glu Lys Val Ser Asp Ser His Leu Glu Arg Arg Ser
375 380 385

cag ccg tga ccgccgcga gccaaagccaa gcaccacccc gggcaaaggc 1259
Gln Pro

cgggaaacgg tccaacctaa cacgcaacga t 1290

<210> 3
<211> 1167
<212> DNA
<213> *Arthrobacter* sp.

<220>
<221> CDS
<222> (1).. (1167)
<223>

<400> 3
atg agc cgt ctg ctc cgt gag cac gga atc gtc atc ggt cgt ctc caa 48
Met Ser Arg Leu Leu Arg Glu His Gly Ile Val Ile Gly Arg Leu Gln
1 5 10 15

ccg ggt tct ctg aac acc att gca gac gtc gca ggc gtt cgc gta ggc 96
Pro Gly Ser Leu Asn Thr Ile Ala Asp Val Ala Gly Val Arg Val Gly
20 25 30

cat tca aca atc atg cgc ggt tct ggg ccc ctg tcc atc ggc cat ggc 144
His Ser Thr Ile Met Arg Gly Ser Gly Pro Leu Ser Ile Gly His Gly
35 40 45

cca gtt cgc aca ggg gta aca gcc atc atc cct cac gaa gga gac atc 192

Pro Val Arg Thr Gly Val Thr Ala Ile Ile Pro His Glu Gly Asp Ile	
50 55 60	
tgg gag gag cca cgg ttc gcc ggc gtc ttc tcc ctg aat ggc agc ggt	240
Trp Glu Glu Pro Arg Phe Ala Gly Val Phe Ser Leu Asn Gly Ser Gly	
65 70 75 80	
gaa tgg agc gga acc tcg ttc gtc agg gag acc ggg tgt ctt tat ggc	288
Glu Trp Ser Gly Thr Ser Phe Val Arg Glu Thr Gly Cys Leu Tyr Gly	
85 90 95	
cct atc atg acg acg aat tcg cac agc att gga tcg gtc agg aac gcc	336
Pro Ile Met Thr Thr Asn Ser His Ser Ile Gly Ser Val Arg Asn Ala	
100 105 110	
gtc atc aag cgt gaa gta gcc cgg cgg gga agc ctg gag agg ctc cct	384
Val Ile Lys Arg Glu Val Ala Arg Arg Gly Ser Leu Glu Arg Leu Pro	
115 120 125	
ctc gtg gga gaa acc ttt gat ggc cta ctc aat gac atc agc ggc atg	432
Leu Val Gly Glu Thr Phe Asp Gly Leu Leu Asn Asp Ile Ser Gly Met	
130 135 140	
cac gtc aag gac gaa cac gtg gcc gag gcc atc gac tcc gcc tcc gca	480
His Val Lys Asp Glu His Val Ala Glu Ala Ile Asp Ser Ala Ser Ala	
145 150 155 160	
aat gtt acc gaa ggc aat gtt ggc ggt ggg acc gga aat gtt tgt cac	528
Asn Val Thr Glu Gly Asn Val Gly Gly Gly Thr Gly Asn Val Cys His	
165 170 175	
ggt ttc aaa ggc ggt att gga agt gcc tcg cgc gtg ttg caa ttg ggc	576
Gly Phe Lys Gly Gly Ile Gly Ser Ala Ser Arg Val Leu Gln Leu Gly	
180 185 190	
gag gaa acc tac act ttg ggg gtt ctc gtc caa gcc aac cac ggc ctt	624
Glu Glu Thr Tyr Thr Leu Gly Val Leu Val Gln Ala Asn His Gly Leu	
195 200 205	
cgt gac gaa ttt cag gtg acg gga gta ccc gtg gga agg ctc ata tct	672
Arg Asp Glu Phe Gln Val Thr Gly Val Pro Val Gly Arg Leu Ile Ser	
210 215 220	
acg gac gag atc ccc ttg ggg cct tca ggt ttt gat cga agg tct tca	720
Thr Asp Glu Ile Pro Leu Gly Pro Ser Gly Phe Asp Arg Arg Ser Ser	
225 230 235 240	
cca cac aaa aac agt att ctt gtc gtc gtg gca acc gac ggc cct cta	768
Pro His Lys Asn Ser Ile Leu Val Val Val Ala Thr Asp Ala Pro Leu	
245 250 255	
cta ccg ggc caa ctg gaa cgc gtt gct cac cgt tct acc cta ggc att	816
Leu Pro Gly Gln Leu Glu Arg Val Ala His Arg Ser Thr Leu Gly Ile	
260 265 270	
gcc cgt aat ggt gcc tac gcg cac aat ctc agc ggc gac ctt gca ctt	864
Ala Arg Asn Gly Ala Tyr Ala His Asn Leu Ser Gly Asp Leu Ala Leu	
275 280 285	
gcg ttc tcc acc tcg ccg cag cct gta agc ggt tac gat ttc gga gtg	912
Ala Phe Ser Thr Cys Pro Gln Pro Val Ser Gly Tyr Asp Phe Gly Val	
290 295 300	
gat aca agt cct ggg acc att cgc gcc ctg ccc aac gcc gca acg gct	960
Asp Thr Ser Pro Gly Thr Ile Arg Ala Leu Pro Asn Ala Ala Thr Ala	
305 310 315 320	
ggc ctc ttc gag gcg gcc gtt gag gcc act gag gaa gcg att gtt tcc	1008
Gly Leu Phe Glu Ala Ala Val Glu Ala Thr Glu Glu Ala Ile Val Ser	
325 330 335	
gcg ctt gtc cac gcc gac acc tcg acc ggg atc gat gac agg gtt gcc	1056
Ala Leu Val His Ala Asp Thr Cys Thr Gly Ile Asp Asp Arg Val Ala	
340 345 350	
tat ggg ttg gag gcg gct cga ctt gct cgt tca att tcg gaa tat cga	1104
Tyr Gly Leu Glu Ala Ala Arg Leu Ala Arg Ser Ile Ser Glu Tyr Arg	
355 360 365	
ggc acc cag ctg tat ccg gag aaa gtg tcg gat tcc cat ctt gaa cga	1152

Gly Thr Gln Leu Tyr Pro Glu Lys Val Ser Asp Ser His Leu Glu Arg
370 375 380

agg agc cag ccg tga
Arg Ser Gln Pro
385

1167

<210> 4
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer-1

<220>
<221> misc_feature
<222> (2)..(2)
<223> n represents a, t, g or c

<220>
<221> misc_feature
<222> (11)..(11)
<223> n represents a, t, g or c

<400> 4
gngarcaygg nathgt

16

<210> 5
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer-2

<220>
<221> misc_feature
<222> (6)..(6)
<223> n represents a, t, g or c

<220>
<221> misc_feature
<222> (15)..(15)
<223> n represents a, t, g or c

<400> 5
gtrtangtyt cytcncc

17

<210> 6
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer-3

<400> 6
tgccattcag ggagaagac

19

<210> 7
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer-4

<400> 7
cacgtcaagg acgaacac

18

<210> 8
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer-5

<400> 8
taattaggat ccagcgcgtc gtggactg 28

<210> 9
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer-6

<400> 9
ataaccggag ctcatcggtg cgtgttaggt t 31

<210> 10
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer-7

<400> 10
tccigctcat atgagccgtc tgcctcgt 28

<210> 11
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer-8

<400> 11
attctatgag ctacggctg gctccttcg 29