Vaccine

FIELD OF THE INVENTION

This invention relates to rotavirus vaccine formulations. The invention relates to the use of an attenuated rotavirus population from one rotavirus serotype in the prevention of disease associated with rotavirus infection from another rotavirus serotype. In particular the invention relates to the use of an attenuated rotavirus population from G1 serotype in the prevention of disease associated with rotavirus infection from non-G1 serotypes.

BACKGROUND

Acute, infectious diarrhoea is a leading cause of disease and death in many areas of the world. In developing countries, the impact of diarrhoeal disease is staggering. For Asia, Africa and Latin America, it has been estimated that there are between 3-4 billion cases of diarrhoea each year and of those cases about 5-10 million result in death (Walsh, J.A. et al.: N. Engl. J. Med., 301:967-974 (1979)).

Rotaviruses have been recognised as one of the most important causes of severe diarrhoea in infants and young children (Estes, M.K. Rotaviruses and Their Replication in Fields Virology, Third Edition, edited by Fields et al., Raven Publishers, Philadelphia, 1996). It is estimated that rotavirus disease is responsible for over one million deaths annually. Rotavirus-induced illness most commonly affects children between 6 and 24 months of age, and the peak prevalence of the disease generally occurs during the cooler months in temperate climates, and year-round in tropical areas. Rotaviruses are typically transmitted from person to person by the faecal-oral route with an incubation period of from about 1 to about 3 days. Unlike infection in the 6-month to 24-month age group, neonates are generally asymptomatic or have only mild disease. In contrast to the severe disease normally encountered in young children, most adults are protected as a result of previous rotavirus infection so most adult infections are mild or asymptomatic (Offit, P.A. et al. Comp. Ther., 8(8):21-26, 1982).

Rotaviruses are generally spherical, and their name is derived from their distinctive outer and inner or double-shelled capsid structure. Typically, the double-shelled capsid structure of a rotavirus surrounds an inner protein shell or core that contains the genome. The genome of a rotavirus is composed of 11 segments of double-stranded RNA which encode at least 11 distinct viral proteins. Two of these viral proteins

designated as VP4 and VP7 are arranged on the exterior of the double-shelled capsid structure. The inner capsid of the rotavirus presents one protein, which is the rotavirus protein designated VP6. The relative importance of these three particular rotaviral proteins in eliciting the immune response that follows rotavirus infection is not yet clear. Nevertheless, the VP6 protein determines the group and subgroup antigen, and VP4 and VP7 proteins are the determinants of serotype specificity.

To date, at least 14 rotavirus G serotypes and 11 rotavirus P serotypes have been identified (Linhares A.C. & Bresse J.S., Pan. Am. J. Publ. Health 2000, 9, 305-330). Among these, 10 G serotypes and 6 P serotypes have been identified among the human rotavirus.

VP7 protein is a 38,000 MW glycoprotein (34,000 MW when non-glycosylated) which is the translational product of genomic segment 7, 8 or 9, depending on the strain. This protein stimulates formation of the neutralising antibody following rotavirus infection. VP4 protein is a non-glycosylated protein of approximately 88,000 MW which is the translational product of genomic segment 4. This protein also stimulates neutralising antibody following rotavirus infection.

Since VP4 and VP7 proteins are the viral proteins against which neutralising antibodies are directed, they are believed to be prime candidates for development of rotavirus vaccines, affording protection against rotavirus illness.

Natural rotavirus infection during early childhood is known to elicit protective immunity. A live attenuated rotavirus vaccine is thus highly desirable. Preferably this should be an oral vaccine, as this is the natural route of infection of the virus.

Early vaccine development for preventing rotavirus infections began in the 1970s after the discovery of the virus. Initially, attenuated strains from animals and humans were studied and had mixed or disappointing results. More recent efforts have focused on human-animal reassortants that have been more successful.

A rotavirus strain known as 89-12 has been described by Ward; see US Patent Number 5,474,773 and Bernstein, D.L. et al, Vaccine, 16 (4), 381-387, 1998. The 89-12 strain

was isolated from a stool specimen collected from a 14 month-old child with natural rotavirus illness in 1988. According to US Patent Number 5,474,773 the HRV 89-12 human rotavirus was then culture-adapted by 2 passages in primary African Green Monkey Kidney (AGMK) cells and 4 passages in MA-104 cells as described by Ward in J. Clin. Microbiol., 19, 748-753, 1984. It was then plaque purified 3 times in MA-104 cells (to passage 9) and grown after 2 additional passages in these cells. One additional passage was made (passage 12) for deposition with the ATCC under the accession number ATCC VR 2272. The deposited strain is known as 89-12C2.

The 1998 paper in Vaccine by Bernstein et al is referred to below as the Vaccine (1998) paper. The paper describes the safety and immunogenicity of an orally administered live human rotavirus vaccine candidate. This vaccine was obtained from strain 89-12, attenuated by passaging without plaque purification 26 times in primary AGMK cells and then another 7 times in an established AGMK cell line (33 passages in total).

Hereinafter the aforesaid material which has been serially passaged 26 times will be referred to as P26 and the material which has been serially passaged 33 times will be referred to as P33. In general, rotavirus derived by passaging 89-12 n times will be referred to as Pn.

In the examples which follow the P33 material was passaged a further 5 times on Vero cells. This is referred to as P38.

The P26 and P33 isolates described in the Vaccine (1998) paper were not deposited in a culture collection, nor were they analysed to establish their genetic characterisation.

It has now been found that the P26 population described in the literature comprises a mixture of variants. This has been established by genetic characterisation as described hereinbelow (see examples). P26 is therefore not a reliably consistent population for further passages, in particular for the production of vaccine lots. Similarly, P33 comprises a mixture of variants and is not reliably consistent for the production of vaccine lots.

It has been found that the P26 material is a mixture of at least three VP4 gene variants. P33 and P38 are similarly a mixture of two variants. These variants appear to be antigenically different, in terms of neutralising epitopes, to the 89-12C2 strain deposited at the ATCC when evaluating the neutralizing antibody titers of sera from infants vaccinated with P33 against these variants.

Furthermore it has been found that when the P33 material is administered to infants, two identified variants are replicated and excreted. Of 100 vaccinated infants, only 2 showed signs of gastro-enteritis due to rotavirus infection, while 20% of a placebo group were infected. These findings suggest that the identified variants are associated with protection from rotavirus disease.

WO 0112797 discloses a method of separating rotavirus variants and an improved live attenuated rotavirus vaccine derived from a cloned (homogeneous) human rotavirus strain. Also disclosed is an attenuated rotavirus population (isolate), characterised in that it comprises a single variant or substantially a single variant, said variant defined by the nucleotide sequence encoding at least one of the major viral proteins designated as VP4 and VP7. Protective efficacy of such an oral attenuated human rotavirus vaccine against G9 heterologous strain has been reported in Latin American infants (Perez et al. 42nd Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC 2002) 27-30 September 2002, San Diego). The whole contents of WO0112797 are herein incorporated by reference.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is the nucleotide sequence encoding the VP4 protein of P43.
Figure 2 is the nucleotide sequence encoding the VP7 protein of P43.

DETAILED DESCRIPTION OF THE INVENTION

In the present invention we have determined that an attenuated rotavirus population, characterised in that it comprises a single variant or substantially a single variant, said variant defined by a nucleotide sequence encoding at least one of the major viral proteins designated as VP4 and VP7, can be used as a vaccine to provide cross protection against disease caused by rotavirus infection of a different serotype to that used in the vaccine.

In particular the use of a G1 rotavirus population, [for example as deposited at the European Collection of Animal Cell Cultures (ECACC), Vaccine Research and Production Laboratory, Public Health Laboratory Service, Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire, SP4 0JG, United Kingdom on 13 August 1999 under the deposition number 99081301, under the terms of the Budapest Treaty], can be used to prevent disease caused by both G1 and at least one non-G1 rotavirus serotype, such as the G2, G3, G4 and G9 rotavirus serotypes.

Accordingly the present invention relates to use of an attenuated rotavirus population from one rotavirus serotype in the prevention of disease associated with rotavirus infection from another rotavirus serotype, wherein the serotype is preferably defined by reference to the sequence of the rotavirus G protein.

Preferably the rotavirus population is characterised in that it comprises a single variant or substantially a single variant, said variant defined by a nucleotide sequence encoding at least one of the major viral proteins designated as VP4 and VP7.

Preferably the rotavirus population comprises VP4 and/or VP7 viral proteins from ECACC deposit 99081301 suitable to provide a cross protective effect.

Preferably the attenuated rotavirus serotype is G1 and is able to provide cross protection against disease caused by G1 and non-G1 rotavirus serotypes such as serotypes selected from the group consisting of: G2, G3, G4, G5, G6, G7, G8, G9, G10, G11, G12, G13 and G14.

In particular the use of a G1 attenuated rotavirus population, [for example as deposited at the European Collection of Animal Cell Cultures (ECACC), Vaccine Research and Production Laboratory, Public Health Laboratory Service, Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire, SP4 0JG, United Kingdom on 13 August 1999 under the deposition number 99081301, under the terms of the Budapest Treaty], can be used to prevent disease caused by G1 and at least one, preferably at least two, more preferably at least three, most preferably at least four non-G1 rotavirus serotypes selected from the group consisting of: G2, G3, G4, G5, G6, G7,

G8, G9, G10, G11, G12, G13 and G14. In a particularly preferred aspect, an immune response is induced against two or more rotavirus non-G1 serotypes, typically against any serotype selected from the group consisting of: G2, G3, G4, G5, G6, G7, G8, G9, G10, G11, G12, G13 and G14. It is preferred to have heterotypic protection against at least two, at least three, at least four, more preferably at least five or more of these serotypes different to that of the vaccine composition. Most preferably an immune response is induced against at least one, preferably at least two, more preferably at least three, most preferably at least four of the following non-G1 serotypes: G2, G3, G4 and G9, in addition to homotypic (G1) protection.

In particular according to the present invention, there is provided the use of an attenuated rotavirus strain from a G1 serotype in the manufacture of a vaccine composition for the induction of an immune response against a rotavirus which is not G1 or G9. In a preferred aspect an immune response is additionally induced against rotavirus infection by a G9 serotype and/or G1 serotype. In a particular preferred aspect, an immune response is induced against two or more rotavirus serotypes, these serotypes not being G1 or G9. Typically any serotype other than G1 or G9 are selected from the list consisting of: G2, G3, G4, G5, G6, G7, G8, G10, G11, G12, G13 and G14. It is preferred to have heterotypic protection against at least two, at least three, at least four, more preferably at least five or more of these serotypes different to that of the vaccine composition. Most preferably an immune response is induced against all of the following non-G1 serotypes: G2, G3, G4 and G9.

The invention also relates to a method of inducing an immune response against rotavirus infection from a rotavirus serotype, the method comprising administering to a subject a composition comprising an attenuated rotavirus vaccine from a different serotype.

Preferably the method is a method of inducing an immune response against rotavirus non- G9 serotype, the method comprising administering to a subject a composition comprising a rotavirus G1 serotype vaccine comprising a single variant or substantially a single variant as defined herein.

Preferably the rotavirus population within the vaccine composition is of G1P1A P[8] specificity. More preferably the rotavirus population comprises VP4 and/or VP7 viral proteins from ECACC deposit 99081301 suitable to elicit an immune response and, typically, provide a cross protective effect. Preferably the rotavirus vaccine used is the ECACC deposit 99081301, or is derived from that deposit.

In one embodiment the invention excludes the use of the rotavirus strain corresponding to ECACC 99081301, in a method or use as indicated above for inducing an immune response to the G9 serotype alone, although use of such a G1 strain for the induction of immune responses to G9 in combination with induction of immune responses to other G serotypes is generally not excluded.

Suitable the invention relates to G1P8 rotavirus strains in methods or uses as described above.

Preferably the vaccine is used in a 2 dose regime.

Preferably the vaccine provides cross protection against gastro-enteritis. Accordingly, in another aspect there is provided a method according to the invention wherein in composition is up to 60% protective, in a population of vaccinated individuals, against diarrhoea caused by infection of a rotavirus of a different type to that of the attenuated rotavirus present in the composition. In another aspect, there is provided a method according to the invention wherein the composition is up to 81% protective against gastro-enteritis caused by infection of a rotavirus of a different type to that of the attenuated rotavirus present in the composition. In a still further aspect, there is provided a method according to the invention wherein the composition comprises a G1 rotavirus strain which is up to 83% protective in a population of vaccinated individuals against severe gastro-enteritis caused by infection of rotaviruses of at least two non-G1 serotypes.

Preferably the protection rate against diarrhoea and/or gastro-enteritis and/or severe gastro-enteritis achieved in a population of vaccinated individuals infected by a rotavirus of a different type to that of the attenuated rotavirus present in the composition, is between 10 to 90%, more preferably between 20 to 80%, most preferably at least 50%.

The rotavirus vaccine used to give cross protection has the following preferred features:

Preferably the rotavirus population according to the invention is a cloned variant.

By a population comprising a single variant, or substantially a single variant, is meant a rotavirus population which does not contain more than 10%, and preferably less than 5% and most preferably less than 1% of a different variant or variants. Virus populations can be purified to homogeneity or substantial homogeneity by passaging on suitable cell types or by performing a series of one or more cloning steps.

An advantage of the invention is that a population comprising a single variant is more suitable for the formulation of a consistent vaccine lot. Particular variants defined by nucleotide sequences encoding the major viral proteins may also be associated with enhanced efficacy in the prevention of rotavirus infection.

In one preferred aspect, the single or substantially single variant in the rotavirus population of the invention is a variant in which the VP4 gene comprises a nucleotide sequence comprising at least one of the following: an adenine base (A) at position 788, an adenine base (A) at position 802 and a thymine base (T) at position 501 from the start codon.

In a further aspect the single or substantially single variant in the population of the invention is a variant in which the VP7 gene comprises a nucleotide sequence comprising at least one of the following: a thymine (T) at position 605, an adenine (A) at position 897, or a guanine (G) at position 897 from the start codon. Preferably at position 897 there is an adenine (A).

In a preferred aspect the single variant in the population according to the invention has an adenine (A) at positions 788 and 802 and a thymine (T) at position 501 from the start codon in the VP4 gene sequence.

In another preferred aspect the single variant in the population according to the invention has a thymine (T) at position 605 and an adenine/guanine (A/G) at position 897 from the

start codon in the VP7 sequence. Most preferably in the VP7 sequence there is an adenine (A) at position 897.

In a particularly preferred aspect the single variant in the population according to the invention has an adenine (A) at positions 788 and 802 and a thymine (T) at position 501 from the start codon in the VP4 gene sequence, and a thymine (T) at position 605 and an adenine/guanine (A/G) at position 897 from the start codon in the VP7 sequence. Most preferably in the VP7 sequence there is an adenine (A) at position 897.

In another aspect the single variant comprises a nucleotide sequence encoding a VP4 protein wherein the nucleotide sequence is as shown in Figure 1, and/or a nucleotide sequence encoding a VP7 protein wherein the nucleotide sequence is as shown in Figure 2.

Preferred rotavirus populations for use in the present invention may be obtained by a method comprising:
passaging a rotavirus preparation on a suitable cell type;
optionally selecting homogeneous culture using the steps of either:
a) limit dilution; or
b) Individual plaque isolation; and
checking for the presence of a substantially single variant by carrying out a sequence determination of an appropriate region of the VP4 and/or VP7 gene sequence.

Preferably the population is derived from the P33 or P26 strains as described above.

The sequence determination may suitably be carried out by a quantitative or semi-quantitative hybridisation technique such as slot blot hybridisation or plaque hybridisation.

Preferably the selected variant is a variant which is replicated and excreted when the starting rotavirus preparation is administered to a human subject, in particular a child.

The resulting cloned virus population resulting from the method according to the invention may be amplified by further passaging on a suitable cell line.

Suitable cell types for passaging the rotavirus population in the above method include African green monkey kidney (AGMK) cells, which may be established cell lines or primary AGMK cells. Suitable AGMK cell lines include for example Vero (ATCC CCL-81), DBS-FRhL-2 (ATCC CL-160), BSC-1 (ECACC 85011422) and CV-1 (ATCC CCL-70). Also suitable are MA-104 (rhesus monkey) and MRC-5 (human – ATCC CCL-171) cell lines. Vero cells are particularly preferred for amplification purposes. Passaging on Vero cells gives a high virus yield.

Techniques for checking whether there is a single variant in a virus population resulting from the method, and for determining the nature of that single variant involve standard sequencing or hybridisation procedures known in the art and are described hereinbelow.

In a preferred aspect the method of the invention is carried out using an appropriate rotavirus, particularly rotavirus having the characteristics of the 89-12 strain or of a passaged derivative thereof.

A particularly preferred single variant population is P43, which was obtained from P33 (an isolated human rotavirus passages 33 times in culture on appropriate cell types) by a series of end dilution cloning steps followed by passaging the cloned material on Vero cells for amplification.

A P43 population was deposited at the European Collection of Animal Cell Cultures (ECACC), Vaccine Research and Production Laboratory, Public Health Laboratory Service, Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire, SP4 0JG, United Kingdom on 13 August 1999 under the deposition number 99081301, under the terms of the Budapest Treaty.

Although this indicated public availability is the simplest method of obtaining the human rotavirus P43, similar and functionally substantially identical rotaviruses may be produced by these or other methods in view of the teachings of this invention. Such functionally substantially identical rotaviruses are considered to be biologically equivalent to the human rotavirus P43 of this invention and therefore are within the general scope of the present invention. It will therefore be understood that the invention

encompasses rotavirus populations having the characteristics of the P43 variant as described herein.

It will also be understood that the invention encompasses materials derived from the deposited P43 ECACC 99081301 by subjecting it to further processing such as by propagating it by further passaging, cloning, or other procedures using the live virus or by modifying P43 in any way including by genetic engineering techniques or reassortant techniques. Such steps and techniques are well known in the art.

Materials derived from the deposited P43 which are covered by the invention include protein and genetic material. Of particular interest are reassortant rotaviruses which comprise at least one antigen or at least one segment of P43, for example reassortants which comprise a virulent strain of rotavirus in which one or part of one of the 11 genome segments has been replaced by the genome segment or part thereof of P43. Specifically, a rotavirus reassortant in which the segment or partial segment coding for NSP4 is a P43 segment or partial segment, may have useful properties. Reassortant rotaviruses and techniques for preparing them are well known (Foster, R. H. and Wagstaff, A. J. Tetravalent Rotavirus Vaccine, a review. ADIS drug evaluation, BioDrugs, Gev, 9 (2), 155-178, 1998).

Materials of particular interest are progeny of P43 and immunologically active derivatives of P43. Immunologically active derivatives means materials obtained from or with the P43 virus, particularly antigens of the virus, which are capable of eliciting an immune response that is reactive against Rotavirus when injected into a host animal.

In adapting the rotavirus to an appropriate cell line, for example Vero cells, it may be necessary to treat the virus so as to get rid of any potential contaminant such as any adventitious agents that may be present and which would otherwise cause contamination. In the case of ether-sensitive adventitious viruses, this may be done by ether treatment as described hereinbelow. The present invention also relates to inclusion of such ether treatment as an optional step in the overall procedure for obtaining an attenuated live rotavirus or vaccine formulated therewith.

Vaccine comprising single or substantially single rotavirus variants are preferred, however, use of single variants or substantially single variants is not essential. The cross protective rotavirus strain of the present invention may be combined with other rotavirus strains to provide additional protection or cross-protection against rotavirus infection or disease.

Therefore, also within the scope of the invention are admixtures of P43 with other rotavirus variants, for example other cloned variants, or with other viruses in particular other attenuated viruses. Such mixtures are useful in the vaccines of the invention which are described hereinbelow.

The present invention also provides a live attenuated rotavirus vaccine capable of providing cross protection, preferably which comprises a substantially single variant population, admixed with a suitable adjuvant or a pharmaceutical carrier.

Preferably, the rotavirus vaccine according to the invention is a monovalent rotavirus vaccine containing a single rotavirus strain.

The present invention is particularly advantageous in providing a live rotavirus vaccine in which the live attenuated rotavirus is a human rotavirus and does not cause intussusception.

Suitable pharmaceutical carriers for use with the attenuated rotavirus strain according to the invention include those known in the art as being suitable for oral administration, especially to infants. Such carriers include and are not limited to carbohydrates, polyalcohols, amino acids, aluminium hydroxide, magnesium hydroxide, hydroxyapatite, talc, titanium oxide, iron hydroxide, magnesium stearate, carboxymethylcellulose, hydroxypropylmethylcellulose, microcrystalline cellulose, gelatin, vegetal peptone, xanthane, caraghenane, arabic gum, ß-cyclodextrin.

The invention also provides a process for preparing a rotavirus vaccine, for example by freeze drying the virus in the presence of suitable stabilisers or admixing the virus according to the invention with a suitable adjuvant or pharmaceutical carrier.

It may also be advantageous to formulate the virus of the invention in lipid-based vehicles such as virosomes or liposomes, in oil in water emulsions or with carrier particles. Alternatively or in addition immunostimulants such as those known in the art for oral vaccines may be included in the formulation. Such immunostimulants include bacterial toxins, particularly cholera toxin (CT) in the form of the holotoxin (entire molecule) or the B chain only (CTB) and the heat labile enterotoxin of *E. coli* (LT). Mutated LTs (mLTs) which are less likely to convert to their active form than the native LT are described in WO 96/06627, WO 93/13202 and US 5,182,109.

Further immunostimulants which may advantageously be included are saponin derivatives such as QS21 and monophosphoryl lipid A, in particular 3-de-O-acylated monophosphoryl lipid A (3D-MPL). Purified saponins as oral adjuvants are described in WO 98/56415. Saponins and monophosphoryl lipid A may be employed separately or in combination (e.g. WO 94/00153) and may be formulated in adjuvant systems together with other agents. 3D-MPL is a well-known adjuvant manufactured by Ribi Immunochem, Montana and its manufacture is described in GB 2122204.

A general discussion of vehicles and adjuvants for oral immunisation can be found in Vaccine Design, The Subunit and Adjuvant Approach, edited by Powell and Newman, Plenum Press, New York, 1995.

The invention also provides a method for vaccinating human subjects, especially infants, by administering to a subject in need thereof an effective amount of a vaccine composition according to the invention. Preferably the live attenuated vaccine is administered by oral administration.

In a preferred aspect the attenuated rotavirus strain according to the invention is formulated with an antacid to minimise inactivation of the vaccine by acid in the stomach. Suitable antacid components include inorganic antacids for example aluminium hydroxide $Al(OH)_3$ and magnesium hydroxide $Mg(OH)_2$. Commercially available antacids which are suitable for use in the invention include Mylanta (trade mark) which contains aluminium hydroxide and magnesium hydroxide. These are insoluble in water and are given in suspension.

Aluminium hydroxide is a particularly preferred component of a vaccine composition according to the invention as it can provide not only an antacid effect but also an adjuvantation effect.

Also suitable for use as antacids in the vaccine of the invention are organic antacids such as organic acid carboxylate salts. A preferred antacid in the vaccine composition of the invention contains an organic acid carboxylate salt, preferably a salt of citric acid such as sodium citrate or potassium citrate.

A particularly preferred antacid that may be used in the vaccine composition of the present invention is the insoluble inorganic salt, calcium carbonate ($CaCO_3$). The calcium carbonate is able to associate with the rotavirus and the rotavirus activity is maintained during the association with the calcium carbonate.

To prevent sedimentation of calcium carbonate during the filling step, viscous agents are preferably present in the formulation.

Possible viscous agents that may be used include pseudoplastic excipients. A pseudoplastic solution is defined as a solution having higher viscosity on standing compared to its viscosity under agitation. Excipients of this type are natural polymers such as arabic gum, adragante gum, agar-agar, alginates, pectines or semi-synthetic polymers for example: carboxymethylcellulose (Tyloses C®), methylcellulose (Methocels A®, Viscotrans MC®, Tylose MH® and MB®), hydroxypropylcellulose (Klucels®), and hydroxypropylmethylcellulose (Methocels E® and K®, Viscontrans MPHC®). In general those pseudoplastic excipients are used together with thixotropic agents. Alternative viscous agents that may be used are pseudoplastic excipients with low flowing capacity. Those polymers, at a sufficient concentration, give rise to a structural fluid arrangement resulting in a high viscosity solution having low flowing capacity on standing. A certain quantity of energy needs to be given to the system to allow flowing and transfer. External energies (agitation) are needed to destroy temporarily the structural fluid arrangement in order to obtain a fluid solution.
Examples of such polymers are Carbopols® and xanthane gum.

Thixotropic excipents become a gel structure on standing whilst under agitation they form a fluid solution. Examples of thixotropic excipients are: Veegum ®(Magnesium-aluminium silicate) and Avicel RC® (about 89% microcrystalline cellulose and 11% Carboxymethylcellulose Na).

The vaccine composition of the present invention preferably comprises a viscous agent selected from xanthane gum or starch.

Thus the vaccine composition of the present invention is preferably formulated with a combination of calcium carbonate and xanthane gum.

Other components of a composition used in the invention suitably include sugars for example sucrose and/or lactose.

The vaccine composition according to the invention may contain additional components including for example flavourings (particularly for an oral vaccine) and bacteriostatic agents.

Different presentations of the vaccine composition according to the invention are envisaged.

In one preferred embodiment, the vaccine is administered as a liquid formulation. Preferably the liquid formulation is reconstituted prior to administration from at least the following two components:

i) virus component
ii) liquid component.

In this embodiment, the virus component and the liquid component are normally present in separate containers, which may conveniently be separate compartments of a single vessel, or separate vessels which can be connected in such a way that the final vaccine composition is reconstituted without exposing it to the air.

Prior to reconstitution, the virus may be in a dry form or a liquid form. Preferably the virus component is lyophilised. Lyophilised virus is more stable than virus in an aqueous

solution. The lyophilised virus may be suitably reconstituted using a liquid antacid composition to produce a liquid vaccine formulation. Alternatively the lyophilised virus may be reconstituted with water or aqueous solution, in which case the lyophilised virus composition preferably contains an antacid component.

Preferably, the vaccine formulation comprises a virus component formulated with calcium carbonate and xanthane gum in one compartment or vessel and this is reconstituted with water or aqueous solution present in the second compartment or vessel.

In another preferred embodiment, the vaccine composition is a solid formulation, preferably a lyophilised cake which is suitable for immediate dissolution when placed in the mouth. Lyophilised formulations may conveniently be provided in the form of tablets in a pharmaceutical blister pack.

In another aspect the invention provides a rotavirus vaccine in the form of a quick dissolving tablet for oral administration.

In another aspect the invention provides a composition comprising a live attenuated rotavirus strain, in particular a human rotavirus strain, wherein the composition is a lyophilised solid capable of immediate dissolution when placed in the mouth.

Preferably the quick dissolving tablet according to the invention dissolves in the mouth of the subject sufficiently quickly to prevent swallowing of the undissolved tablet. This approach is particularly advantageous for paediatric rotavirus vaccines.

Preferably the virus is a live attenuated human rotavirus which is formulated with an inorganic antacid such as calcium carbonate and a viscous agent such as xanthane gum.

A further aspect of the present invention is to provide a lyophilised formulation wherein the virus component is any rotavirus strain which is formulated with calcium carbonate and xanthane gum.

Vaccines of the invention may be formulated and administered by known techniques, using a suitable amount of live virus to provide effective protection against rotavirus infection without significant adverse side effects in typical vaccinees. A suitable amount of live virus will normally be between 10^4 and 10^7 focus forming units (ffu) per dose. A typical dose of vaccine may comprise 10^5 – 10^6 ffu per dose and may be given in several doses over a period of time, for example in two doses given with a two-month interval. Benefits may however be obtained by having more than 2 doses, for example a 3 or 4 dose regimen, particularly in developing countries. The interval between doses may be more or less than two months long. An optimal amount of live virus for a single dose or for a multiple dose regimen, and optimal timing for the doses, can be ascertained by standard studies involving observation of antibody titres and other responses in subjects.

The vaccine of the invention may also comprise other suitable live viruses for protection against other diseases, for example poliovirus. Alternatively other suitable live virus vaccines for oral administration may be given in a separate dose but on the same occasion as the rotavirus vaccine composition according to the invention.

Sera from twelve 4 to 6 month old infants vaccinated with the P33 material as described in the Vaccine (1998) paper were tested for neutralization of P33, P38, P43 and 89-12C2.

The range of neutralization titers of all the tested sera is similar for P33, P38 and P43. The statistical analysis shows no significant difference in the overall neutralization titers against all three viruses. This suggests that the conformational and non-conformational neutralization epitopes of P33, P38 and P43 are equally well recognized by the anti-P33 sera of P33 vaccinated infants. This observation indirectly suggests that the neutralization epitopes revealed in this in vitro assay were not altered between P33, P38 and P43.

The range of neutralization titers of P89-12C2 however significantly differs from P33, P38 and P43. This observation suggests that the conformational and non-conformational neutralization epitopes of P33, P38 and P43 are not equally well recognized by the anti-P33 sera of P33 vaccinated infants. This observation indirectly suggests that the

neutralization epitopes revealed in this in vitro assay were altered between 89-12 C2 and P33, P38 and P43.

Highly preferred embodiments of the present invention include:

1. The use of an attenuated rotavirus strain from a G1 serotype in the manufacture of a vaccine composition for the induction of an immune response against a rotavirus which is not G1 or G9.

2. The use according to 1 wherein an immune response is additionally induced against rotavirus infection by a G9 serotype and/or G1 serotype.

3. The use according to 1 or 2 wherein the immune response is induced against two or more rotavirus serotypes, these serotypes not being G1 or G9.

4. The use as claimed in any of 1 to 3 wherein the serotype other than G1 or G9-serotype is selected from the list consisting of: G2, G3, G4, G5, G6, G7, G8, G10, G11, G12, G13 and G14.

5. The use according to any of 1 to 4 wherein an immune response is induced against all of the following non-G1 serotypes: G2, G3, G4 and G9.

6. The use according to any of 1 to 5, wherein the G1 rotavirus vaccine composition comprises a single rotavirus variant or substantially a single rotavirus variant, said variant defined by a nucleotide sequence encoding at least one of the major viral proteins designated as VP4 and VP7.

7. The use according to 6 wherein the composition comprises a rotavirus having a VP4 gene comprising, in the nucleotide sequence, at least one of the following: an adenine base (A) at position 788, an adenine base (A) at position 802 and a thymine base (T) at position 501 from the start codon.

8. The use according to 7 wherein the VP4 gene comprises a nucleotide sequence comprising an adenine base (A) at positions 788 and 802 and a thymine base (T) at position 501 from the start codon.

9. The use according to 6 wherein the composition comprises a rotavirus having a VP7 gene comprising, in the nucleotide sequence, at least one of the following: a thymine (T) at position 605, an adenine (A) at position 897 and a guanine (G) at position 897 from the start codon.

10. The use according to 7 wherein the VP7 gene comprises a nucleotide sequence comprising a thymine (T) at position 605 and an adenine (A) or a guanine (G) at position 897 from the start codon.

11. The use according to any of 1 to 10, wherein the composition comprises a rotavirus having a VP4 gene comprising, in the nucleotide sequence, an adenine (A) at positions 788 and 802 and a thymine (T) at position 501 from the start codon; and wherein the VP7 gene comprises, in the nucleotide sequence, a thymine (T) at position 605 and an adenine (A) at position 897 from the start codon.

12. The use according to any of 1 to 11 wherein the composition is capable of protecting against gastro-enteritis and/or diarrhea caused by infection by a rotavirus of a different type to the G1 serotype of the attenuated rotavirus present in the composition.

13. The use according to 12 wherein the composition is up to 83% protective in a population of vaccinated individuals against severe gastro-enteritis caused by infection of rotaviruses of at least two serotypes, these serotypes being different to the G1 serotype of the attenuated rotavirus present in the composition.

14. The use according to 13 wherein the severe gastro-enteritis is caused by infection of a rotavirus of at least four non-G1 serotypes.

15. The use according to 14 wherein the non-G1 serotypes are G2, G3, G4 and G9 serotypes.

16. The use according to any of 1 to 15 wherein the rotavirus strain is ECACC deposit 99081301, or is obtainable or derivable from ECACC deposit 99081301.

17 The use according to any of 1 to 16 wherein the vaccine is used in a 2-dose regime.

The following, non-limiting, examples illustrate the invention.

EXAMPLES

Example 1: Demonstration that strain 89-12 at passage 26 (P26) is a mixture of variants

Sequencing of VP4 and VP7 genes from different passage lots

Sequencing of VP4 and VP7 genes from passage P26 (primary AGMK cells), passage P33 (established (as opposed to primary) AGMK cell line), passage P41 and passage P43 was performed. Total RNA extraction was reverse transcribed and amplified through PCR in one tube/one step.

Primers Rota 5bis and Rota 29bis amplified the entire VP4 gene and primers Rota 1 and Rota 2bis amplified the entire VP7 gene. The PCR material has been sequenced using different primers (see Table 1).

The passage P26 sequence differed from the passage P33 sequence by 3 bases (at positions 501, 788 and 802 bp from the start codon) in VP4 and by three bases in VP7 (108, 605 and 897 bp from the start codon).

The passage P26 sequence scans of VP4 and VP7 show at mutated positions the presence of the passage P33 sequence as a background. Thus it can be seen that passage P26 is a mixture of at least 2 variants.

The passage P33 sequence scans seem homogenous in VP4 and heterogeneous for VP7 (see Table 2).

Passage P38 (derived from passage 33) was passaged 5 times on Vero cells and displayed the same set of VP4 and VP7 sequences as passage P33 (AGMK cell line). Thus there was no major change in populations between P33 and P38.

TABLE 1: Oligonucleotides used for RT-PCR and sequencing

	name	sequence	position
VP7	Rota 1	GGC TTT AAA AGA GAG AAT TTC CGT CTG G	-49 to –22
	Rota 1bis	GGT TAG CTC CTT TTA ATG TAT GGT A	-16 to 10
	Rota 2bis	GGT CAC ATC GAA CAA TTC TAA TCT AAG	1014-988
	Rota 7	CAA GTA CTC AAA TCA ATG ATG G	266-287
	Rota 12	TGT TGA TTT TTC TGT CGA TCC AC	372-394
	Rota 46	GGT TGC TGA GAA TGA GAA ATT AGC TAT AGT GG	651-682
	Rota 18	CCA CTA TAG CTA ATT TCT CAT TCT CAG CAA CC	682-651
VP4	Rota 5	TGG CTT CGC CAT TTT ATA GAC A	2-23
	Rota 6	ATT TCG GAC CAT TTA TAA CC	878-859
	Rota 5bis	TGG CTT CAC TCA TTT ATA GAC A	2-23
	Rota 6bis	ATT TCA GAC CAT TTA TAA CCT AG	878-856
	Rota 25	GGA GTA GTA TAT GAA AGT ACA AAT AAT AG	268-296
	Rota 26	CTA TTA TTT GTA CTT TCA TAT ACT ACT CC	296-268
	Rota 27bis	TCG ATA CAG TAT AAG AGA GCA CAA G	721-745
	Rota 28		
	Rota 31	TTC ATT AAC TTG TGC TCT CTT ATA CTG	753-727
	Rota 32	GTA TAT GTA GAC TAT TGG GAT G	1048-1070
	Rota 45	CAT CCC AAT AGT CTA CAT ATA C	1070-1048
	Rota 53	TGT AAC TCC GGC AAA ATG CAA CG	1205-1227
	Rota 54	CGT TGC ATT TTG CCG GAG TTA CA	1227-1205
	Rota 55	GTA AGA CAA GAT TTA GAG CGC CA	1465-1487
	Rota 40	TGG CGC TCT AAA TCT TGT CTT AC	1487-1465
	Rota 39	CTT GAT GCT GAT GAA GCA GCA TCT G	1703-1727
	Rota 33	CAG ATG CTG CTT CAT CAG CAT CAA G	1727-1703
	Rota 34	CGA TCA TAT CGA ATA TTA AAG GAT G	2008-2032
	Rota 29bis	CAT CCT TTA ATA TTC GAT ATG ATC G	2032-2008
		AGC GTT CAC ACA ATT TAC ATT GTA G	2335-2311

TABLE 2: oligonucleotides used in hybridization

	name	sequence	position
VP7	Rota 41	AGT ATT TTA TAC TAT **A**GT AGA TTA TAT TAA TC	882-913
	Rota 42	AGT ATT TTA TAC TAT **G**GT AGA TTA TAT TAA TC	882-913
VP4	Rota 15	ATC CC**C** ATT ATA CTG CAT T**C**C TTT C	807-783
	Rota 16	ATC CC**T** ATT ATA CTG CAT T**T**C TTT C	807-783
	Rota 35	ATC CC**C** ATT ATA CTG CAT TTC TTT C	807-783
	Rota 36	ATC CC**T** ATT ATA CTG CAT T**C**C TTT C	807-783

The bases shown in bold type in Table 2 are the sites of specific sequence variation in VP4 and VP7.

TABLE 3: sequence variation of VP4 and VP7 genes

3.1

	VP4			VP7		
	501 bp 167 aa	788 bp 263 aa	802 bp 268 aa	108 bp 36 aa	605 bp 202 aa	897 bp 299 aa
P26 (AGMK)	A	G / A	G / A	A	C / T	A
P33 (AGMK)	T	A	A	G / A	T / C	A / G
P38 (VERO)	T	A	A	A / G	T	G / A
P43 (VERO)	T	A	A	A	T	A

N.B. In a second clone from the 3 clones which were developed to the level of production lot, the VP7 897 bp position nucleotide is G, rather than A as in the P43 selected clone. This results in a methionine in place of an isoleucine in the amino acid sequence. Variants corresponding to both the selected P43 clone and the clone in which there is a G in VP7 at 897 bp from the start codon, were excreted in the stools of infants who had been vaccinated with the P33 material.

In Table 3.1, where there are two alternative bases at a particular position, the first of the two represents the base which appears in a major population and the second is the base

which appears in a minor population. Major and minor variant populations are judged by the strength of the signal in sequencing.

3.2

	VP4			VP7		
	501 bp 167 aa	788 bp 263 aa	802 bp 268 aa	108 bp 36 aa	605 bp 202 aa	897 bp 299 aa
P26 (AGMK)	Leu	Gly/Glu	Gly/Arg	Arg	Thr/Met	Ile
P33 (AGMK)	Phe	Glu	Arg	Arg/Arg	Met/Thr	Ile/Met
P38 (VERO)	Phe	Glu	Arg	Arg/Arg	Met	Met/Ile
P43 (VERO)	Phe	Glu	Arg	Arg	Met	Ile

Table 3.2 shows the amino acid changes resulting from the nucleotide differences between the variants.

TABLE 4

	VP4 (788 -802 positions)				VP7 (897 position)	
	G-G	A-A	A-G	G-A	A	G
Probes	Rota 15	Rota 16	Rota 35	Rota 36	Rota 41	Rota 42
Passages						
P26	-	+	+	+	nd	nd
P33	-	+	-	-	+ +	+
P38	-	+	-	-	+	++
P43	-	+	-	-	+	-

Slot blot hybridization

The change in populations between passages P26 to P33 on AGMK cells has been further confirmed by slot blot hybridization. The VP4 and the VP7 gene fragments generated by RT/PCR were hybridized with oligonucleotide probes specific for each variant (see Table 3.1 and 3.2). In contrast to P26 which hybridized with Rota 16, Rota 35 and Rota 36 and not with Rota 15, the VP4 PCR fragment of the P33 material, at positions 788 and 802 hybridized only with Rota 16 and not with either Rota 15 or Rota 35 or Rota 36. These results established the presence of at least 3 variants in P26 (see Table 4).

For the VP7 PCR fragment of the P33 material, position 897 hybridized with Rota 41 and Rota 42. These results established the presence of at least two variants in the P33 material.

Example 2: Isolation and characterization of the P43 clone

To isolate P33 components as a homogeneous virus population, three end-point dilutions of P33/AGMK on Vero cells were performed and the resulting virus was used to infect Vero cells.

Positive wells were selected using two criteria: growth demonstrated by the largest number of foci detected in the wells and the most isolated positive wells on the plates, as is done classically. After 3 end dilution passages in 96 well microtiter plates, 10 positive wells were amplified successively on Vero cells and evaluated for their yield.

Based on yield, three clones were developed to passage level of production lot. Immunorecognition by polyclonal antibodies was shown to be similar both between the three clones and between the clones and P33. Homogeneity of the clones was assessed by slot blot hybridization. The final selection of a single clone was based on yield and sequence.

The selected clone was amplified by successive passages on Vero cells to generate a Master seed, a Working seed and finally production lots.

The selected clone was genetically characterized at different passage levels by sequencing of VP4 and VP7 (identity) and by specific slot blot hybridization of the VP4 and VP7 (homogeneity) of the PCR amplified materials. The sequence of the VP4 and VP7 genes of the P43 material are given in Figures 1 and 2 respectively and are identical to P41.

Homogeneity of the selected clone was assessed by a selective hybridization using oligonucleotide probes discriminating nucleotide changes in VP4 and/or VP7 regions for each variant identified during sequencing of P26/primary AGMK (see Table 4).

The VP4 fragment hybridized with Rota 16 and not with Rota 15, Rota 35 or Rota 36. The VP7 fragment hybridized with Rota 41 and not with Rota 42.

These results confirmed that P43 is a homogeneous population.

Example 3: Removal of potential adventitious virus

Ether was added to P33 (AGMK grown) to a final concentration of 20% for 1 hr. Ether was then bubbled out with N_2 for 35 min. No impact on the titre of P33 seed was observed.

Example 4: Formulation of a live attenuated vaccine

The production lots described above are formulated for oral adminstration to infants by the following method.

1. Lyophilised virus

Standard techniques are used for preparing virus doses. Frozen purified viral bulk is thawed and diluted with appropriate medium composition, in this case Dulbecco's modified eagle Medium, up to a desired standard viral concentration, in this case $10^{6.2}$ ffu/ml. The diluted virus is then further diluted with lyophilisation stabiliser (sucrose 4%, dextran 8%, sorbitol 6%, amino-acid 4%) up to the target viral titre, in this case $10^{5.6}$ ffu/dose. 0.5 ml aliquots of stabilised virus composition are aseptically transferred to 3

ml vials. Each vial is then partially closed with a rubber stopper, the sample is freeze dried under a vacuum, the vial is then fully closed and an aluminium cap is crimped in place around the vial to keep the stopper in place.

For use, the virus is reconstituted using one of the following antacid reconstituents:

(a) Citrate reconstituent

Sodium citrate is dissolved in water, sterilized by filtration and aseptically transferred into reconstituent containers in 1.5 ml amounts at a concentration of 544 mg $Na_3Citrate.2H_2O$ per 1.5ml dose. The reconstituent containers may be for example 3 ml vials, or 4 ml vials, or 2 ml syringes, or soft plastic squeezable capsules for oral administration. As an alternative to maintaining sterile components under sterile conditions, the final container can be autoclaved.

(b) $Al(OH)_3$ reconstituent

An aseptic aluminium hydroxide suspension (Mylanta – trademark) is aseptically diluted in sterile water, aseptically transferred to reconstituent containers (for example 2 ml syringes, or soft plastic squeezable capsules) in 2 ml amounts each containing 48 mg $Al(OH)_3$. An alternative to using sterile components under sterile conditions is to γ irradiate the aluminium hydroxide suspension (preferably at a diluted stage).

Standard ingredients are included to prevent the suspension from settling. Such standard ingredients include for example magnesium stearate, carboxymethylcellulose, hydroxypropylmethylcellulose, microcrystalline cellulose, and silicone polymers. Bacteriostatic agents for example butylparaben, propylparaben or other standard bacteriostatic agents used in food, and flavourings, may also be included.

2. Lyophilised virus with $Al(OH)_3$ in liquid formulation

Standard techniques are used for preparing virus doses. Frozen purified viral bulk is thawed and diluted with appropriate medium composition, in this case Dulbecco's modified eagle Medium, up to a desired standard viral concentration, in this case $10^{6.2}$ ffu/ml. Aluminium hydroxide suspension is added to reach a final quantity of 48 mg/dose and the virus composition is diluted with lyophilisation stabiliser (sucrose 4%, dextran

8%, sorbitol 6%, amino-acid 4%) up to the target viral titre, in this case $10^{5.6}$ ffu/dose. 0.5 ml aliquots of stabilised virus composition are aseptically transferred to 3 ml vials. Lyophilisation and closing of the vials is then carried out as described in part 1.

3. Lyophilised virus with $Al(OH)_3$ for blister presentation

Standard techniques are used for preparing virus doses. Frozen purified viral bulk is thawed and diluted with appropriate medium composition, in this case Dulbecco's modified eagle Medium, up to a desired standard viral concentration, in this case $10^{6.2}$ ffu/ml. Aluminium hydroxide suspension is added to reach a final quantity of 48 mg/dose and the virus composition is diluted with lyophilisation stabiliser which may be sucrose, dextran or amino-acid 4%, or gelatin, or vegetal peptone, or xanthane up to the target viral titre of $10^{5.6}$ ffu/dose. An aseptic filling operation is employed to transfer doses of 0.5 ml or preferably less to blister cavities. The composition is lyophilised, and the blister cavities are sealed by thermic sealing.

Optionally standard ingredients are included to prevent the aluminium hydroxide suspension from settling. Such standard ingredients include for example magnesium stearate, carboxymethylcellulose, hydroxypropylmethylcellulose, microcrystalline cellulose, and silicone polymers. Flavourings may also be included.

Example 5: Rotavirus viral titration for various formulations

5.1:Comparison between lactose and sucrose based formulations:

Table 5

Batch n°	Fomulation composition	Viral titer before lyophilisation	Viral titer after lyophilisation and 1 week at 37°C
98G06/01	Lactose:2%; Dextran:4%; Sorbitol:3%; AminoAcids:2%	$10^{5.22}$	$10^{4.67}$
98G06/03	Sucrose:2%; Dextran:4%; Sorbitol:3%; AminoAcids:2%	$10^{5.28}$	$10^{4.92}$

P43 rotavirus was formulated either with sucrose or with lactose as shown in the table above.
Viral titration before lyophilisation is the viral titre in the completed formulated liquid (containing sucrose dextran sorbitol aminoacids) and without the lyophilisation step.

Good results are those in which a <0.5 log decrease at the lyophilisation step and <0.5 log decrease during the "1 week at 37°C" (accelerated stability test) are achieved.
The precision of the viral titration is around + or – 0.2 log.

The results indicate that sucrose may be used instead of lactose.

5.2: Effect of arginine and replacement of sorbitol by maltitol:

Table 6

Batch n°	Fomulation composition	Viral titer at time = zero after lyophilisation	Viral titer after lyopjhilisation and 1 week at 37°C
98L16/01	Lactose:2%; Dextran:4%; Sorbitol:3%; AminoAcids:2%	$10^{4.8}$	$10^{4.8}$
98L16/02	Lactose:2%; Dextran:4%; Sorbitol:3%; AminoAcids:2% Arginine: 3%	$10^{4.8}$	$10^{4.9}$
98L16/04	Lactose:2%; Dextran:4%; Maltitol:3%; AminoAcids:2% Arginine: 3%	$10^{4.7}$	10^{5}

The results demonstrate that the addition of arginine (which is known to improve the stability of the virus during lyophilisation and also provides a basic medium in order to compensate for the stomach acidity) maintains the viral titer.

Sorbitol tends to decrease the glass transition temperature of the lyophilised cake by too great a degree. This can be overcome by using maltitol instead of sorbitol as shown above and the viral titer is still maintained.

5.3: Various formulation compositions

This experiment demonstrates that a number of formulations are possible.

Table 7

Batch n°	Fomulation composition	Viral titer before lyophilisation	Viral titer after lyophilisation and 1 week at 37°
99C11/01	Sucrose:2%; Dextran:4%; Sorbitol:3%; AminoAcids:2%	$10^{5.24}$	$10^{5.07}$
99C11/02	Sucrose:2%; Dextran:4%; Maltitol:3%; AminoAcids:2%	$10^{5.09}$	$10^{4.92}$
99C11/04	Dextran:4%; Maltitol:3%; AminoAcids:2%	$10^{4.89}$	$10^{5.06}$

Batch n°	Fomulation composition	Viral titer at time = zero after lyophilisation	Viral titer after lyophilisation and 1 week at 37°
99C17/01	Sucrose:2%; Dextran:4%; Sorbitol:3%; AminoAcids:2%	$10^{5.40}$	$10^{5.41}$
99C17/02	Sucrose:2%; Dextran:4%; Sorbitol:1.5%; AminoAcids:2%	$10^{5.30}$	$10^{4.93}$
99C17/03	Sucrose:2%; Dextran:4%; AminoAcids:2%	$10^{5.31}$	$10^{5.24}$
99C17/04	Sucrose:2%; Dextran:4%; Maltitol:3%; AminoAcids:2%	$10^{4.42}$	$10^{4.45}$
99C17/05	Sucrose:2%; Dextran:4%;	$10^{4.39}$	$10^{4.40}$

	Maltitol:1.5%; AminoAcids:2%		
99C17/06	Sucrose:2%; Dextran:4%; Sorbitol:3%;	$10^{5.44}$	$10^{4.97}$
99C17/07	Sucrose:2%; Dextran:4%; Sorbitol:1.5%;	$10^{5.11}$	$10^{4.89}$

5.4: Association between Rotavirus and $Al(OH)_3$ antacid:

Table 8

Rotavirus	$Al(OH)_3$	H_2O	Contact time at room temperature	Centrifugation	Supernatant viral titer in ffu/ml	Pellets viral titer in ffu/ml
$10^{5.6}$ ffu/ml	48 mg in 0.240ml	0.76 ml	30 min	8000rpm, 10 min	$10^{3.66}$	
$10^{5.6}$ ffu/ml	0.48 mg in 0.240ml	0.76 ml	30 min	8000rpm, 10 min	$10^{4.41}$	
$10^{5.6}$ ffu/ml		1 ml	30 min	8000rpm, 10 min	$10^{5.68}$	
Rotavirus in Lyophilised Cake	12 mg in 0.120ml	1.380ml	30 min	8000rpm, 10 min	Below detection	$10^{4.7}$

$Al(OH)_3$ is used as an antacid. This shows that Rotavirus is associated with the insoluble inorganic salt ($Al(OH)_3$) since it centrifuged together with the $Al(OH)_3$ (decrease of viral activity in the supernatant).

5.5: Dissolution of $Al(OH)_3$ antacid by SodiumCitrate before viral titration

Table 9

Viral samples	Dissolution	Conditions	Viral titers ffu/ml
99B10/06 liquid formulation before lyophilisation; $10^{5.43}$	1.5ml $Na_3Citrate$	24h at room temperature	$10^{5.11}$
99B10/06:lyophilized $10^{5.43}$	1.5ml $Na_3Citrate$	24h at room temperature	$10^{4.53}$

When Rotavirus is associated with the $Al(OH)_3$, it is possible to lyophilise everything (including the $Al(OH)_3$). After lyophilisation, it is possible to recover the Rotavirus by dissolving $Al(OH)_3$ in SodiumCitrate. This step does not damage the Rotavirus and retains its activity after this dissolution step.

5.6: Infectivity of Rotavirus after liberation of the $Al(OH)_3$-Rotavirus association:

The mechanism of virus liberation (by dissolution of the carrier) may very well occur *in vivo*. Indeed below pH 6, aluminium hydroxide becomes completely soluble, and thus, Rotavirus will be liberated in the stomach.

$$Al(OH)_3 + 3\ H^+ \dashrightarrow Al^{+++}\ (\text{water soluble}) + 3\ H_2O$$


In the stomach, Al^{+++} ions are not absorbed (J.J. Powell, R.Jugdaohsingh and R.P.H. Thompson, *The regulation of mineral adsorption in the gastrointestinal track*, Proceedings of the Nutrition Society (1999), 58, 147-153).

In the intestine, due to the increase of pH, insoluble forms of aluminium are precipitated ($Al(OH)_3$ or $AlPO_4$), and eliminated by the natural way.

It is unknown whether the newly formed $Al(OH)_3$ (or $AlPO_4$) precipitate will be able to re-associate with free Rotavirus. This raises the question of the infectivity of the $Al(OH)_3$-Rotavirus association itself.

Liberation of Rotavirus from the $Al(OH)_3$-Rotavirus association by other mechanisms is also possible. Lysine, for example, interferes with the viral adsorption on $Al(OH)_3$.

Other anions like borate, sulfate, carbonate and phosphate are known to be specifically adsorbed on aluminium hydroxide, thus, theoretically, it should be possible to displace (by competition for the adsorption site) Rotavirus from the $Al(OH)_3$-Rotavirus association.




Thus, Rotavirus may be liberated from the Rotavirus – $Al(OH)_3$ association and the liberated Rotavirus remains active.

This liberation can be done either by dissolving $Al(OH)_3$ (by HCl in the stomach, or by $Na_3Citrate$ in vitro) or by displacing Rotavirus by a basic amino acid (lysine).

5.7: Infectivity of the $Al(OH)_3$-Rotavirus association

A single dose of lyophilised Rotavirus was reconstituted with water and divided into two parts. The first part, considered as the reference, received an additional volume of water. The second part received 24mg of $Al(OH)_3$ suspended in 0.240 ml of water (Preclinical viral titrations).




When $Al(OH)_3$ is present, Rotavirus is active and the viral titration value is higher compared to the reference sample.

This experiment was repeated without dividing the lyophilised dose, and by adding 12 mg $Al(OH)_3$ or 24 mg $Al(OH)_3$.

Here the reference sample was the one reconstituted with a Citrate-Bicarbonate buffer. Thus, the viral titer is again higher in the presence of $Al(OH)_3$.

DRVC003A46 + 1.5 ml WL buffer	DRVC003A46 + 12 mg Al(OH)3 in 0.120ml + 1.380ml H2O	DRVC003A46 + 24 mg Al(OH)3 in 0.240ml + 1.260ml H2O
5.34	**6.24**	**6.05**
5.32	**5.95**	**6.26**

As in the example above, Rotavirus associates with the $Al(OH)_3$ particles, since the virus can be discarded by centrifugation. DRVC003A46 is a lyophilised formulated Rotavirus (Sucrose:2%; Dextran: 4%,Sorbitol:3%; Amino-acids:2%).



SDSAA= Sucrose 2%, Dextran 4%, Sorbitol 3%, Amino-Acid 2%.

According to the viral titration carried out on the supernatant, the quantity of $Al(OH)_3$ needed to adsorb Rotavirus seems to be low (starting with one lyophilised dose 5.7 log) scaling Up viral titration):

Table 10

Al(OH)3	Adsorption time	Titer in supernatant
12 mg	1 hour RT	2.7
24 mg	1 hour RT	3.4
48 mg	1 hour RT	3.4
72 mg	1 hour RT	2.0
96 mg	1 hour RT	Below detection
12 mg	Overnight	2.7

24 mg	Overnight	Below detection
48 mg	Overnight	2.5
12 mg	Immediate	Below detection
24 mg	Immediate	2.0
48 mg	Immediate	Below detection

Time needed to adsorb Rotavirus on $Al(OH)_3$ seems to be short:

One dose of lyophilised Rotavirus was reconstituted in presence of 24 mg $Al(OH)_3$, and centrifuged after 0, 15, 60 min and 24 hours. The "culot" were resuspended in SDSAA before viral titration:

Table 11

Time	**Culot**	**Supernatant**
0 min	5.26	3.17
15 min	5.34	<1.44
60 min	5.96	<1.44
24 hours	6.13	<1.44

5.8: Using $CaCO_3$ as antacid

In order to avoid aluminium in the vaccine, the antacid $Al(OH)_3$ was replaced by another insoluble inorganic salt: $CaCO_3$ (calcium carbonate).

The phenomena observed with $CaCO_3$ are parallel to those described for $Al(OH)_3$:

- Association of Rotavirus with the inorganic salt;
- Maintainance of Rotavirus activity when associated with the inorganic salt;
- Possibility of liberation of Rotavirus from the association by dissolution of the inorganic base by an acid;
- Possibility of co-lyophilisation of the antacid and the Rotavirus.

$CaCO_3$ and Rotavirus association

In a first trial, lyophilised Rotavirus (viral titer 5.7) was reconstituted with a suspension of $CaCO_3$ in water (50mg in 1.5ml); and then centrifuged, and the viral titer of the supernatant compared to the pellet.

DRVC003A46
+
50mg CaCO3
in
1.5 ml H2O
+
Centrifugation
8000rpm 10
min

Culot
+
1.5 ml
SDSAA

5.83

Supernatant

4.46

DRVC003A46
+
50 mg CaCO3
in
1.5 ml H2O
+
Centrifugation
8000rpm 10
min

Culot
+
1.5 ml
Na
Citrate

5.88

Supernatant

4.33

This indicates that more that 90% of the Rotavirus is associated with $CaCO_3$.

Also, when the virus was associated, it was possible to realise the titration and to recover the original viral quantities.
Also, viral titers are slightly higher that those obtained without $CaCO_3$.

DRVC003A46
+
1.5 ml H2O
+
Centrifugation
8000rpm 10 min

"Culot
"

4.99

Supernatant

5.03

DRVC003A46
+
1.5 ml
W.L Buffer

5.35

Quantity of $CaCO_3$ and Rotavirus association

Lyophilised Rotavirus was reconstituted with a $CaCO_3$ suspension in water (1.5ml):

10 mg

50 mg

100 mg

and then centrifuged, and the viral titer of the supernatant compared to the culot.

Table 12

CaCO3	Extempo + Centri.		1 Hour + Centri	
	Culots	Surpernatant	Culots	Surpernatant
100mg	4.57	3.01	4.79	3.09
50mg	4.17	4.15	4.22	3.86
10mg	3.17	4.77	3.87	4.87

Thus, clearly, more $CaCO_3$ and more virus is associated, and less is found in the supernatant.

However, the full dose is not completely recovered (expected a total of 5.3 at least or even 5.8 as obtained earlier – see above).

$CaCO_3$ Protection of Rotavirus during Baby Rossett-Rice antacid titration

Using 10 doses of lyophilised Rotavirus (DRVC003A46) and 50mg of $CaCO_3$, two types of baby Rossett-Rice titration were carried out:

In a classic Rossett-Rice titration, the antacid is mixed with Rotavirus and HCl is poured into this medium.

In the "inverse" baby Rossett-Rice, the situation is the reverse: antacid is dropped into the HCl pool (as it occurs in vivo).

Table 13

Classical baby Rossett-Rice titration			
Lyophi. Rota	Buffer	Theoretical	Measured

stored at:		Viral Titer	Viral Titer
4°C	60 mg CaCO3	5.3	4.6
-80°C	60 mg CaCO3	5.3	4.6
4°C	24 mg Al(OH)3	5.4	<2.9
-80°C	24 mg Al(OH)3	5.4	<2.9

Inverse baby Rossett-Rice titration			
Lyophi. Rota stored at:	Buffer	Theoretical Viral Titer	Measured Viral Titer
4°C	60 mg CaCO3	5.3	4.6
-80°C	60 mg CaCO3	5.3	4.6
4°C	24 mg Al(OH)3	5.4	<2.9
-80°C	24 mg Al(OH)3	5.4	<2.9

Thus, in this *in vitro* experiment, calcium carbonate is able to protect about 20% of Rotavirus from the presence of HCl, while aluminium hydroxide is not able to.

5.9: Lyophilisation of Rotavirus in presence of $CaCO_3$ antacid:

Table 14

Batch n°	Composition	Viral titer at time = zero after lyophilisation	Viral titer after lyopjhilisation and 1 week at 37°C
99K08/01	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2% $CaCO_3$: 50 mg	$10^{5.28}$	$10^{5.10}$
99K08/02	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2% $CaCO_3$: 60 mg	$10^{5.16}$	$10^{5.15}$
00C24/01	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2% $CaCO_3$: 60 mg Xanthane 0.3%	$10^{5.07}$	$10^{4.69}$
00C24/03	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2%	$10^{5.07}$	$10^{4.85}$

	$CaCO_3$: 60 mg Xanthane 0.3%		
00E09/25	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2% $CaCO_3$: 60 mg Xanthane 0.25%	$10^{5.03}$	$10^{4.91}$
00E09/30	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2% $CaCO_3$: 60 mg Xanthane 0.30%	$10^{5.01}$	$10^{4.87}$
00F26/06	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2% $CaCO_3$: 60 mg Starch: 2%	$10^{4.50}$	$10^{4.70}$

This is the "all in one" - lyophilisation of Rotavirus and antacid (CaCO3) together in the same vial. To prevent sedimentation of $CaCO_3$ during the filling step, viscous agents are needed. Examples of such viscous agents include Xanthane gum and Starch. The Rotavirus activity is maintained even in the presence of Xanthane gum and Starch.

5.10 Lyophilised tablets for quick disintegration when placed in the mouth:

The following formulations demonstrate the "lyoc" concept. That is, quick dissolution of the lyophilised cake in the mouth.

Table 15

Batch n°	Formulation composition	Viral titer before lyophilisation	Viral titer after lyopjhilisation and 1 week at 37°
99B10/06	Sucrose 4% Sodium glutamate 3.7% Al(OH)3 48mg	$10^{5.11}$	$10^{4.53}$
99C11/12	Maltitol 3% Al(OH) 48mg Hydroxypropylmethyl-cellulose: 1%	$10^{4.16}$	$10^{3.79}$

Batch n°	Fomulation composition	Viral titer at time = zero after lyophilisation	Viral titer after lyopjhilisation and 1 week at 37°
00C24/05	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2% $CaCO_3$: 60 mg Xanthane 0.3%	$10^{5.02}$	$10^{4.54}$
00C24/06	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2% $CaCO_3$: 60 mg Xanthane 0.3%	$10^{4.86}$	$10^{4.56}$
00F26/11	Sucrose: 1% Dextran: 2% Sorbitol: 1.5% Am. Acids: 1% $CaCO_3$: 60 mg Starch: 2%	$10^{4.70}$	$10^{4.40}$

In the "lyoc concept" both Xanthane and Starch can be used (maintaining the quick dissolution properties of the lyophilised cake).

Example 6: Use of Calcium Carbonate as the antacid for the Rotavirus vaccine composition

When a suspension of $CaCO_3$ in water is used as the antacid for Rotavirus there is a problem that the calcium carbonate particles sediment rapidly when placed in water since the powder density value approaches 2.6 and the average particle size is 30µm. This sedimentation can be slowed by:

1 increasing the density of the surrounding medium

2 increasing the viscosity of the surrounding medium

3 reducing the particles size

4 keeping particles away from each other

6.1: Increasing density of the surrounding medium:

When the $CaCO_3$-Water suspension (when placed in the syringe) is placed on the lyophilised cake (containing sucrose 2%, dextran 4%; sorbitol 3%; amino-acids 2%) the density of the surrounding medium is increased, but the speed of $CaCO_3$ sedimentation is not very much different from the $CaCO_3$-Water suspension.

6.2 Increasing the viscosity of the surrounding medium:

Pseudoplastic excipents

A pseudoplastic solution is defined as a solution having higher viscosity on standing compared to its viscosity under agitation.

Usual excipients of this type are:

natural polymers for example:

- arabic gum
- adragante gum
- agar-agar
- alginates
- pectines

semi-synthetic polymers for example:

- carboxymethylcellulose (Tyloses C®)
- methylcellulose (Methocels A®, Viscotrans MC®, Tylose MH® and MB®)
- hydroxypropylcellulose (Klucels®)
- hydroxypropylmethylcellulose (Methocels E® and K®, Viscontrans MPHC®)

In general those pseudoplastic excipients are used together with thixotropic agents.

Pseudoplastic excipients with low flowing capacity

Those polymers, at a sufficient concentration, give rise to a structural fluid arrangement resulting in a high viscosity solution having low flowing capacity on standing. A certain quantity of energy needs to be given to the system to allow flowing and transfer. External energies (agitation) are needed to destroy temporarily the structural fluid arrangement in order to obtain a fluid solution.

Examples of such polymers are Carbopols® and Xanthane gum.

Thixotropic excipents

With these excipents, on standing, a gel structure is obtained; while under agitation a fluid solution is obtained.

Examples of thixotropic excipients are: Veegum ®(Magnesium-aluminium silicate) and Avicel RC® (about 89% microcrystalline cellulose and 11% Caboxymethylcellulose Na).

6.3 Reducing the particles size

A reduction in the $CaCO_3$ particle size resulted in a decrease in the antacid capacity of the compound.

6.4 Keeping particles away from each other


This is the case in Veegum® and Avicel® for which insoluble particles smaller (about 1 µm) than the $CaCO_3$ particles, are placed between $CaCO_3$ particles in order to prevent aggregation.

Example 7: Product design

The following schemes demonstrate examples of possible product designs.


7. 1 $CaCO_3$ in the syringe

Having already clinical batches of Rotavirus in lyophilised vials, the antacid can be placed in the reconstituent liquid contained in the syringe.



In this product presentation, sedimentation of $CaCO_3$ must be under control not only during the filling steps, but also during the complete shelf-live of the product (at least 2 years).

7. 2 $CaCO_3$ in the lyophilised vial



7.3. Lyophilisation in a blister

In this case Rotavirus, $CaCO_3$ and Xanthane gum are lyophilised together directly in the blister.



Example 8: Lyophilisation of different strain of Rotavirus

Table 16

Batch n°	Rotavirus strain	Fomulation composition	Viral titer at t = zero after lyophilisation	Viral titer after lyopjhilisation and 1 week at 37°
00F26/01	G1 SB purif n°61 PRO/0232	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2%	$10^{4.6}$	$10^{4.7}$
00F26/02	G2 (DS-1)	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2%	$10^{4.4}$	$10^{4.4}$
00F26/03	G3(P)	Sucrose: 2% Dextran: 4%	$10^{4.6}$	$10^{4.5}$

		Sorbitol: 3% Am. Acids: 2%		
00F26/04	G4 (VA-70)	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2%	$10^{4.8}$	$10^{4.8}$
00F26/05	G9 (W161)	Sucrose: 2% Dextran: 4% Sorbitol: 3% Am. Acids: 2%	$10^{4.6}$	$10^{4.5}$

The strains DS-1, P and VA70 are described as Human rotavirus reference strains for serotype G2, G3 and G4 respectively at page 1361 of "Fields" Raven press 1990, second edition.

In this experiment different Rotavirus strains have been lyophilised.
For all, both the viral titer have been maintained during lyophilisation and accelarated stability (one week at 37°C) has been shown.

Example 9: Phase I safety study in adults of one oral administration of the Rotavirus vaccine.

A Phase I study was carried out to assess the safety and reactogenicity of a single oral dose of $10^{6.0}$ ffu of the P43 vaccine in healthy adults aged 18 to 45 years.

The clinical trial was double blind and randomized. It was placebo-controlled and self-contained. The study was performed in one single centre in Belgium.

Study Population

A total of 33 subjects, 11 in the placebo group and 22 in the vaccine group, were enrolled and all completed the study. All volunteers were Caucasians. Their mean age at the time of vaccination was 35.3 years, with a range of 18 to 44 years. The trial began in January and ran for just over one month.

Material

Vaccine

Clinical lots of P43 vaccine were produced, purified, formulated and lyophilized according to Good Manufacturing Practices. The lots were released by Quality Control and Quality Assurance. Each vial of vaccine contained the following components:

Active ingredient:

P43 strain	Min. $10^{5.8}$ ffu

Excipients, stabilizers:

Sucrose	9 mg
Dextran	18 mg
Sorbitol	13.5 mg
Amino acids	9 mg

Placebo

Vials of placebo were prepared and released. Each vial of placebo contained the following components:

Excipients, stabilizers:

Sucrose	9 mg
Dextran	18 mg
Sorbitol	13.5 mg
Amino acids	9 mg

Diluent

Water for injection was used as diluent to reconstitute vaccine and placebo.

Administration

Approximately 10 to 15 minutes before administration of the vaccine or the placebo, subjects of both groups were given 10 ml of Mylanta® orally. Mylanta® is a registered antacid. The antacid increases the pH of the stomach and prevents inactivation of the rotavirus during its passage through the stomach.

To prepare the vaccine, two vials of lyophilized P43 containing $10^{5.8}$ ffu per vial were reconstituted with 1.5 ml of diluent water for injection. This achieved a calculated viral titer of $10^{6.1}$ ffu per dose. The reconstituted vaccine was administered promptly as a single oral dose.

To prepare the placebo, two vials of lyophilized placebo were reconstituted with 1.5 ml water for injection and administered orally as a single dose.

Safety and Reactogenicity

The following criteria of safety and reactogenicity applied:

Solicited general symptoms were fever, diarrhea, vomiting, nausea, abdominal pain and loss of appetite. They were recorded during eight days post administration.

Unsolicited symptoms were recorded during 30 days post administration.

Serious adverse events were recorded during the entire study period.

Diarrhea samples were to be collected during eight days post administration.

The results were:

No solicited symptoms, no unsolicited and no serious adverse events were reported during the respective observation periods.

No cases of diarrhea were reported.

Conclusions

SB Biologicals P43 vaccine was safe relative to the placebo when administered orally in a double-blind fashion as a single dose at the dose of $10^{6.1}$ ffu to healthy adult volunteers aged 18 to 44.

Example 10 - Efficacy of two doses of a human monovalent Rotavirus vaccine, Rotarix™ in preventing Gastro-enteritis due to G1 and non-G1 (G9) Rotavirus

A randomised, double-blind, placebo-controlled phase II trial was conducted to evaluate the protective efficacy of a vaccine derived from the G1P human strain 89-12 for infant immunisation.

Specifically the vaccine used was named Rotarix™ and comprises as the rotavirus component the attenuated G1 human strain deposited as ECACC deposit 99081301.

493 healthy infants received two doses of Rotarix™ at the viral concentration (10^6 ffu) or placebo (504) at age 2 and 4 months, concomitantly with DTPw-HBV and Hib vaccines, OPV was administered 2 weeks apart. Diarrhoeal samples were tested for the presence of rotavirus (ELISA) and the serotypes determined in positive samples (RT-PCR). Diarrhoeal episodes reported from two weeks after the second dose were considered for the efficacy analysis. Severity was determined using a 20-point scale (Ruuska and Vesikari, 1990). A score ≥ 11 defined severe disease.

Results. An interim analysis of efficacy was performed on the above mentioned group and the isolated serotypes were G1 and G9, almost evenly distributed. The overall attack rate in the placebo group varied from 4.8% for G1 to 3.6% for G9 during the 6 months observation period. Two doses of Rotarix™ at 10^6 ffu protected against all types of diarrhoea caused by G1 with 83% efficacy [95% CI: 50.4-95.7] and 92 % efficacy [95% CI: 47.6-99.8] against severe gastro-enteritis. If the diarrhoea was caused by G9, the protection against all types of diarrhoea was 60% [95% CI: 0.2-86.0] and 81% [95% CI: 33.0-96.4] against severe gastro-enteritis. For each of these efficacy endpoints, there was a statistically significant decrease in diarrhoea episodes in the HRV group as compared to the placebo group ($p < 0.05$, two-sided Fisher's exact test).

Conclusion. These results are highly supportive of the efficacy of 2 doses of a monovalent HRV vaccine, Rotarix™, in protecting young infants against the homologous G1 strain and cross-protect against the G9 strain.

Example 11 - Efficacy of two doses of a human monovalent Rotavirus vaccine, Rotarix™ administered at three different virus concentrations in preventing Gastro-enteritis due to G1 and non-G1 (G2, G3, G4, G9) Rotavirus

A randomised, double-blind, placebo-controlled phase II trial was conducted to evaluate the protective efficacy and efficacy against hospitalization of a vaccine derived from the G1P human strain 89-12 for infant immunisation.

Specifically the vaccine used was named Rotarix™ and comprises as the rotavirus component the attenuated G1 human strain deposited as ECACC deposit 99081301.

Healthy infants received two doses of Rotarix™ at three different virus concentrations (468 received $10^{4.7}$ ffu, 460 received $10^{5.2}$ ffu; 464 received $10^{5.8}$ ffu) or placebo (454) at age 2 and 4 months, concomitantly with DTPw-HBV and Hib vaccines, OPV was administered 2 weeks apart. Diarrhoeal samples were tested for the presence of rotavirus (ELISA) and the serotypes determined in positive samples (RT-PCR). Diarrhoeal episodes reported from two weeks after the second dose until subjects were one year of age were considered for the efficacy analysis. Severity was determined

using a 20-point scale (Ruuska and Vesikari, 1990). A score ≥ 11 defined severe disease.

Results:

Results are illustrated in the tables below. Infants in the vaccine groups had significantly fewer rotavirus gastroenteritis episodes than children in the placebo group ($p < 0.001$, two-sided Fisher's exact test). Depending on the dosage, protective efficacy against severe rotavirus gastroenteritis reached 86% (95%CI: 63%-96%), and 70% (95%CI, 46%-84%) against any rotavirus gastroenteritis. For each of these efficacy endpoints, there was a statistically significant decrease in diarrhoea episodes in the HRV group as compared to the placebo group ($p < 0.001$, two-sided Fisher's exact test). Multiple rotavirus serotypes (G1, G2, G3, G4 and G9) were identified from gastroenteritis stools (ELISA and RT-PCR) allowing to also calculate vaccine efficacy against non-G1 serotypes. As can be seen from table 19 in particular, for non-G1 serotypes (G2, G3, G4 and G9), and depending on the dosage, efficacy against severe rotavirus gastroenteritis reached 83% (95%CI: 40%-97%), providing proof of concept that the monovalent G1-based G1P1A P[8] human rotavirus vaccine elicits cross-protection against heterotypic (i.e. non-G1) strains.

Features of rotavirus gastro-enteritis episodes reported during the study

Table 17

		RIX4414 $10^{4.7}$ ffu	RIX4414 $10^{5.2}$ ffu	RIX4414 $10^{5.8}$ ffu	Placebo
Any rotavirus gastroenteritis		21	22	15	51
no. of episodes (percent) with specific feature among all rotavirus gastroenteritis episodes reported					
Severity scores	<7	4 (19)	8 (36)	2 (13)	5 (10)
	7-10	5 (24)	4 (18)	8 (53)	12 (24)
	≥ 11	12 (57)	10 (45)	5 (33)	34 (67)
Identified rotavirus serotypes					
	wild G1	12 (57)	6 (27)	7 (47)	30 (59)
	G2	0	0	1 (7)	3 (6)
	G3	1 (5)	0	0	2 (4)

G4	0	0	1 (7)	0
G9	8 (38)	14 (64)	7 (47)	15 (29)
Canine	0	0	0	1(2)
Unknown	0	2 (9)	0	0

Protective efficacy of two doses of RIX4414 human rotavirus vaccine against rotavirus gastroenteritis

Table 18

		Any rotavirus gastroenteritis		Severe rotavirus gastroenteritis		Hospitalization for rotavirus gastroenteritis	
	N	**n (%)**	**Efficacy (95% CI)**	**n (%)**	**Efficacy (95% CI)**	**n (%)**	**Efficacy (95% CI)**
Pooled vaccine groups	1392	58 (4)*	61 (42-74)	27 (2)*	74 (56-85)	9 (0.6) *	79 (48-92)
RIX4414 $10^{5.8}$ ffu	464	15 (3)*	70 (46-84)	5 (1)*	86 (63-96)	3 (0.6)‡	79 (25-96)
RIX4414 $10^{5.2}$ ffu	460	22 (5)*	56 (25-75)	10 (2)*	71 (40-87)	1 (0.2)*	93 (54-100)
RIX4414 $10^{4.7}$ ffu	468	21 (4)*	58 (29-76)	12 (3)*	66 (32-84)	5 (1)†	65 (-2-90)
Placebo	454	49 (11)	-	34 (7)	-	14 (3)	-

*p<0.001 for each comparison between the vaccine and placebo groups by two-sided Fisher's exact test (significant level of α=0.05)

†p=0.037 for the comparison between the vaccine and placebo groups by two-sided Fisher's exact test (significant level of α=0.05)

‡p=0.007 for the comparison between the vaccine and placebo groups by two-sided Fisher's exact test (significant level of α=0.05)

N = number of subjects

n/% = number of subjects reporting at least one specified rotavirus gastroenteritis episode

Exact 95% confidence intervals are shown

Protective efficacy of two doses of RIX4414 human rotavirus vaccine against serotype specific severe rotavirus gastroenteritis

Table 19

		Severe rotavirus gastroenteritis		
	N	n(%)	Efficacy (95% CI)	p-value*
G1 wild type rotavirus				
Pooled vaccine groups	1392	13 (0.9)	74 (41- 88)	<0.001
RIX4414 $10^{5.8}$ ffu	464	2 (0.4)	88 (48-99)	<0.001
RIX4414 $10^{5.2}$ ffu	460	4 (0.9)	75 (24-94)	0.006
RIX4414 $10^{4.7}$ ffu	468	7 (1.5)	58 (-9-85)	0.057
Placebo	454	16 (4)	-	-
Non-G1 rotavirus (mainly G9 with G2, G3 and G4 types)				
Pooled vaccine groups	1392	14 (1)	73 (42-88)	<0.001
RIX4414 $10^{5.8}$ ffu	464	3 (0.6)	83 (40-97)	0.001
RIX4414 $10^{5.2}$ ffu	460	6 (1)	65 (7-89)	0.020
RIX4414 $10^{4.7}$ ffu	468	5 (1)	71 (19-92)	0.009
Placebo	454	17 (4)	-	-

*Two-sided Fisher's exact test (significant level of α=0.05) used for each comparison between the vaccine and placebo groups.
N = number of subjects
n/% = number of subjects reporting at least one specified rotavirus gastroenteritis episode
Exact 95% confidence intervals are shown

Conclusion:

These results are highly supportive of the efficacy of 2 doses of a monovalent HRV vaccine, Rotarix™, in protecting young infants against any and severe rotavirus gastroenteritis caused by the homologous G1 strain and broad cross-protection against heterologous strains, namely G2, G3, G4 and G9.

Claims

1. An attenuated rotavirus strain from a G1 serotype for use in the preparation of a composition for inducing an immune response against rotavirus infection from the G1 and at least two of the non-G1 rotavirus serotypes selected from the group consisting of: G2, G3, G4 and G9 serotypes.

2. An attenuated rotavirus strain from a G1 serotype for use according to claim 1, wherein the composition is capable of inducing an immune response against rotavirus infection from at least three other rotavirus strains.

3. . An attenuated rotavirus strain from a G1 serotype for use according to claim 1 or 2 wherein the composition is capable of inducing an immune response to the G1 and to the G2, G3, G4 and G9 serotypes.

4. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 3 wherein the attenuated rotavirus strain is a single rotavirus variant or substantially a single rotavirus variant, said variant defined by a nucleotide sequence encoding at least one of the major viral proteins designated as VP4 and VP7, preferably VP7.

5. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 4 wherein the composition comprises a rotavirus having a VP4 gene comprising, in the nucleotide sequence, at least one of the following: an adenine base (A) at position 788, an adenine base (A) at position 802 and a thymine base (T) at position 501 from the start codon.

6. An attenuated rotavirus strain from a G1 serotype for use according to claim 5 in which the VP4 gene comprises a nucleotide sequence comprising an adenine base (A) at positions 788 and 802 and a thymine base (T) at position 501 from the start codon.

7. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 4 in which the composition comprises a rotavirus having a VP7 gene comprising, in the nucleotide sequence, at least one of the following: a thymine (T) at position 605, an adenine (A) at position 897 and a guanine (G) at position 897 from the start codon.

8. An attenuated rotavirus strain from a G1 serotype for use according to claim 7 in which the VP7 gene comprises a nucleotide sequence comprising a thymine (T) at position 605 and an adenine (A) or a guanine (G) at position 897 from the start codon.

9. An attenuated rotavirus strain from a G1 serotype for use according to claims 6 or 8, wherein the composition comprises a rotavirus having a VP4 gene comprising, in the nucleotide sequence, an adenine (A) at positions 788 and 802 and a thymine (T) at position 501 from the start codon; and wherein the VP7 gene comprises, in the nucleotide sequence, a thymine (T) at position 605 and an adenine (A) at position 897 from the start codon.

10. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 9 wherein the composition is at least 50% protective, in a population of vaccinated individuals, against diarrhea and/or gastroenteritis and/or severe gastroenteritis caused by infection of a rotavirus of at least two non-G1 serotypes.

11. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 9 wherein the composition is up to 60% protective, in a population of vaccinated individuals, against diarrhoea caused by infection of a rotavirus of at least two non-G1 serotypes.

12. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 9 wherein the composition is up to 81% protective against gastro-enteritis caused by infection of a rotavirus of at least two non-G1 serotypes.

13. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 9 wherein the composition is up to 83% protective in a population of vaccinated individuals against severe gastro-enteritis caused by infection of rotaviruses of at least two non-G1 serotypes selected from the group consisting of G2, G3, G4 and G9.

14. An attenuated rotavirus strain from a G1 serotype for use according to claim 10, 12 and 13 wherein the gastro-enteritis is caused by infection of a rotavirus of at least three non-G1 serotypes selected from the group consisting of G2, G3, G4 and G9.

15. An attenuated rotavirus strain from a G1 serotype for use according to claim 14 wherein the gastro-enteritis is caused by infection with G2, G3, G4 and G9 serotypes.

16. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 15 wherein the rotavirus strain is ECACC deposit 99081301, or is obtainable or derivable from ECACC deposit 99081301.

17. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 16 wherein the composition is administered in a 2-dose regime.

18. An attenuated rotavirus strain from a G1 serotype for use according to claims 1 to 17 wherein the attenuated rotavirus strain is formulated with a suitable pharmaceutical carrier or with an antacid buffer or both.

FIG 1 VP4 sequence of P43

```
ATGGCTTCAC TCATTTATAG ACAACTTCTC ACTAATTCAT ATTCAGTAGA   50
TTTACATGAT GAAATAGAGC AAATTGGATC AGAAAAAACT CAGAATGTAA  100
CTATAAATCC GGGTCCATTT GCACAGACTA GATATGCTCC AGTCAATTGG  150
GATCATGGAG AGATAAATGA TTCGACTACA GTAGAACCAA TTTTAGATGG  200
TCCTTATCAG CCAACTACAT TTACTCCACC TAATGATTAT TGGATACTTA  250
TTAATTCAAA TACAAATGGA GTAGTATATG AAAGTACAAA TAATAGTGAC  300
TTTTGGACTG CAGTCGTTGC TATTGAACCG CACGTCAACC CAGTAGATAG  350
ACAATATATG ATATTTGGTG AAAGCAAGCA ATTTAATGTG AGTAACGATT  400
CAAATAAATG GAAGTTTTTA GAAATGTTTA GAAGCAGTAG TCAAAATGAA  450
TTTTATAATA GACGTACATT AACTTCTGAT ACCAGACTTG TAGGAATATT  500
TAAATATGGT GGAAGAGTAT GGACATTTCA TGGTGAAACA CCGAGAGCTA  550
CTACTGACAG TTCAAGTACT GCAAATTTAA ATAATATATC AATTACAATT  600
CATTCAGAAT TTTACATTAT TCCAAGGTCC CAGGAATCTA AATGTAATGA  650
ATATATTAAT AATGGTCTGC CACCAATTCA AAATACTAGA AATGTAGTTC  700
CATTGCCATT ATCATCTAGA TCGATACAGT ATAAGAGAGC ACAAGTTAAT  750
GAAGACATTA TAGTTTCAAA AACTTCATTA TGGAAAGAAA TGCAGTATAA  800
TAGGGATATT ATAATTAGAT TTAAATTTGG TAATAGTATT GTAAAGATGG  850
GAGGACTAGG TTATAAATGG TCTGAAATAT CATATAAGGC AGCAAATTAT  900
CAATATAATT ACTTACGTGA CGGTGAACAA GTAACCGCAC ACACCACTTG  950
TTCAGTAAAT GGAGTGAACA ATTTTAGCTA TAATGGAGGG TTTCTACCCA 1000
CTGATTTTGG TATTTCAAGG TATGAAGTTA TTAAAGAGAA TTCTTATGTA 1050
TATGTAGACT ATTGGGATGA TTCAAAAGCA TTTAGAAATA TGGTATATGT 1100
TAGATCATTA GCAGCTAATT TAAATTCAGT GAAATGTACA GGTGGAAGTT 1150
ATTATTTCAG TATACCAGTA GGTGCATGGC CAGTAATGAA TGGTGGCGCT 1200
GTTTCGTTGC ATTTTGCCGG AGTTACATTA TCCACGCAAT TTACTGATTT 1250
TGTATCATTA AATTCACTAC GATTTAGATT TAGTTTGACA GTTGATGAAC 1300
CACCTTTCTC AATACTGAGA ACACGTACAG TGAATTTGTA TGGATTACCA 1350
GCCGCTAATC CAAATAATGG AAATGAATAC TACGAAATAT CAGGAAGGTT 1400
TTCACTCATT TCTTTAGTTC CAACTAATGA TGATTATCAG ACTCCAATTA 1450
TGAATTCAGT GACGGTAAGA CAAGATTTAG AGCGCCAACT TACTGATTTA 1500
CGAGAAGAAT TTAACTCATT GTCACAAGAA ATAGCTATGG CACAATTGAT 1550
TGATTTAGCA CTGTTGCCTC TAGATATGTT TTCCATGTTT TCAGGAATTA 1600
AAAGTACAAT TGATTTAACT AAATCAATGG CGACTAGTGT AATGAAGAAA 1650
TTTAGAAAAT CAAAATTAGC TACATCAATT TCAGAAATGA CTAATTCATT 1700
GTCAGATGCT GCTTCATCAG CATCAAGAAA CGTTTCTATT AGATCGAATT 1750
TATCTGCGAT TTCAAATTGG ACTAATGTTT CAAATGATGT GTCAAACGTA 1800
ACTAATTCAT TGAACGATAT TTCAACACAA ACATCTACAA TTAGTAAGAA 1850
ACTTAGATTA AAAGAAATGA TTACTCAAAC TGAAGGAATG AGCTTTGACG 1900
ACATTTCAGC AGCTGTACTA AAAACAAAAA TAGATATGTC TACTCAAATT 1950
GGAAAAAATA CTTTACCTGA TATAGTTACA GAAGCATCTG AGAAATTTAT 2000
TCCAAAACGA TCATATCGAA TATTAAAGGA TGATGAAGTA ATGGAAATTA 2050
ATACTGAAGG AAAATTCTTT GCATACAAAA TTAATACATT TGATGAAGTG 2100
CCATTCGATG TAAATAAATT CGCTGAACTA GTAACAGATT CTCCAGTTAT 2150
ATCAGCGATA ATCGATTTTA AGACATTGAA AAATTTAAAT GATAATTATG 2200
GAATCACTCG TACAGAAGCG TTAAATTTAA TTAAATCGAA TCCAAATATG 2250
TTACGTAATT TCATTAATCA AAATAATCCA ATTATAAGGA ATAGAATTGA 2300
ACAGTTAATA CTACAATGTA AATTGTGAGA ACGCTATTGA GGATGTGACC 2350
```

Fig 2 VP7 sequence of P43

```
ATGTATGGTC TTGAATATAC CACAATTCTA ATCTTTCTGA TATCAATTAT  50
TCTACTCAAC TATATATTAA AATCAGTAAC TCGAATAATG GACTACATTA  100
TATATAGATC TTTGTTGATT TATGTAGCAT TATTTGCCTT GACAAGAGCT  150
CAGAATTATG GGCTTAACTT ACCAATAACA GGATCAATGG ACACTGTATA  200
CGCTAACTCT ACTCAAGAAG GAATATTTCT AACATCCACA TTATGTTTGT  250
ATTATCCAAC TGAAGCAAGT ACTCAAATTA ATGATGGTGA ATGGAAAGAC  300
TCATTGTCAC AAATGTTTCT CACAAAAGGT TGGCCAACAG GATCAGTCTA  350
TTTTAAAGAG TATTCAAGTA TTGTTGATTT TTCTGTCGAT CCACAATTAT  400
ATTGTGATTA TAACTTAGTA CTAATGAAAT ATGATCAAAA TCTTGAATTA  450
GATATGTCAG AGTTAGCTGA TTTAATATTG AATGAATGGT TATGTAATCC  500
AATGGATATA ACATTATATT ATTATCAACA ATCGGGAGAA TCAAATAAGT  550
GGATATCAAT GGGATCATCA TGTACTGTGA AAGTGTGTCC ACTGAATACG  600
CAAATGTTAG GAATAGGTTG TCAAACAACA AATGTAGACT CGTTTGAAAT  650
GGTTGCTGAG AATGAGAAAT TAGCTATAGT GGATGTCGTT GATGGGATAA  700
ATCATAAAAT AAATTTGACA ACTACGACAT GTACTATTCG AAATTGTAAG  750
AAGTTAGGTC CAAGAGAGAA TGTAGCTGTA ATACAAGTTG GTGGCTCTAA  800
TGTATTAGAC ATAACAGCAG ATCCAACGAC TAATCCACAA ACTGAGAGAA  850
TGATGAGAGT GAATTGGAAA AAATGGTGGC AAGTATTTTA TACTATAGTA  900
GATTATATTA ACCAAATCGT GCAGGTAATG TCCAAAAGAT CAAGATCATT  950
AAATTCTGCA GCTTTTTATT ATAGAGTATA GATATATCTT AGATTAGATC  1000
GATGTGACC
```