

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 April 2003 (17.04.2003)

PCT

(10) International Publication Number
WO 03/031449 A2

(51) International Patent Classification⁷: **C07D 501/00**,
C07K 9/00, C07D 501/20, A61K 31/545, 38/04, A61P
31/00

TURNER, S., Derek [US/US]; 77 Paloma Avenue, Apt.
37, Pacifica, CA 94044 (US). **AGGEN, James** [US/US];
1105 Bayswater Avenue #1, Burlingame, CA 94010 (US).

(21) International Application Number: PCT/US02/32534

(74) Agents: **BOONE, David, E. et al.**; Theravance, Inc., 901
Gateway Boulevard, South San Francisco, CA 94080 (US).

(22) International Filing Date: 11 October 2002 (11.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/328,889 12 October 2001 (12.10.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **THER-
AVANCE, INC.** [US/US]; 901 Gateway Boulevard, South
San Francisco, CA 94080 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FATHEREE, Paul**
[US/US]; 921 Minnesota Street, San Francisco, CA 94107
(US). **LINSELL, Martin, S.** [GB/US]; 37 Coso Avenue,
San Francisco, CA 94110 (US). **LONG, Daniel, D.**
[GB/US]; 3855 21st Street #3, San Francisco, CA 94114
(US). **MARQUESS, Daniel** [GB/US]; 2037 Touraine
Lane, Half Moon Bay, CA 94019 (US). **MORAN, Ed-
mund, J.** [CA/US]; 131 Chaves, San Francisco, CA 94127
(US). **NODWELL, Matthew, B.** [CA/US]; 1221 Ma-
sonic Avenue, Apt. #12, San Francisco, CA 94117 (US).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 03/031449 A2

(54) Title: CROSS-LINKED GLYCOPEPTIDE-CEPHALOSPORIN ANTIBIOTICS

(57) Abstract: This invention provides cross-linked glycopeptide cephalosporin compounds and pharmaceutically acceptable salts thereof which are useful as antibiotics. This invention also provides pharmaceutical compositions containing such compounds; methods for treating bacterial infections in a mammal using such compounds; and processes and intermediates useful for preparing such compounds.

5

10

This invention is directed to novel cross-linked vancomycin – cephalosporin compounds which are useful as antibiotics. This invention is also directed to pharmaceutical compositions comprising such compounds; methods of using such compounds as antibacterial agents; and processes and intermediates for preparing such compounds.

20

Various classes of antibiotic compounds are known in the art including, for example, β -lactam antibiotics, such as cephalosporins, and glycopeptide antibiotics, such as vancomycin. Cross-linked antibiotic compounds are also known in the art. See, for example, U.S. Patent No. 5,693,791, issued to W. L. Truett and entitled "Antibiotics and Process for Preparation"; and WO 99/64049 A1, published on December 16, 1999, and entitled "Novel Antibacterial Agents."

Despite such compounds, a need exists for new antibiotics having improved properties including, by way of example, increased potency against gram-positive bacteria. In particular, a need exists for new antibiotics which are highly effective against antibiotic-resistant strains of bacteria, such as methicillin-resistant *Staphylococci aureus* (MRSA) and methicillin-resistant *Staphylococci epidermitis* (MRSE).


WO 03/031449

PCT/US02/32534

SUMMARY OF THE INVENTION

Disclosed herein are novel cross-linked glycopeptide – cephalosporin compounds which are useful as antibiotics. Among other properties, compounds of the disclosed embodiments have been found to possess surprising and unexpected potency against gram-positive bacteria including methicillin-resistant *Staphylococci aureus* (MRSA) and methicillin-resistant *Staphylococci epidermitis* (MRSE).

Accordingly, in a first aspect, this invention provides a compound of formula I:



30

or a pharmaceutically-acceptable salt thereof, wherein

WO 03/031449

PCT/US02/32534

X^1 and X^2 are independently selected from the group consisting of hydrogen and chloro;

R^1 is $-Y^a-(W)_n-Y^b-$;

W is selected from the group consisting of $-O-$, $-N(R^d)-$, $-S-$, $-S(O)-$, $-S(O)_2-$, C_{3-6} cycloalkylene, C_{6-10} arylene and C_{2-9} heteroarylene; wherein each arylene, cycloalkylene and heteroarylene group is optionally substituted with 1 to 3 substituents independently selected from R^b ;

Y^a and Y^b are independently C_{1-3} alkylene, or when W is cycloalkylene, arylene or heteroarylene, Y^a and Y^b are independently selected from the group consisting of a covalent bond and C_{1-3} alkylene; wherein each alkylene group is optionally substituted with 1 to 3 substituents independently selected from $-OR^d$, $-NR^dR^e$, $-CO_2R^d$, $-C(O)NR^dR^e$ and $-S(O)_2NR^dR^e$;


R^2 is hydrogen or C_{1-6} alkyl;

each R^3 is independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl, C_{3-6} heterocyclic and R^a ; or two adjacent R^3 groups are joined to form C_{3-6} alkylene or $-O-(C_{1-6}$ alkylene)-O-; wherein each alkyl, alkylene, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^a and R^c ; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^b ;

one of R^4 and R^5 is hydroxy and the other is hydrogen;

R^6 and R^7 are independently hydrogen or methyl;

R^8 is hydrogen or a group of formula (i):



WO 03/031449

PCT/US02/32534

each R^a is independently selected from the group consisting of $-OR^d$, halo, $-SR^d$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)_2NR^dR^e$, $-NR^dR^e$, $-CO_2R^d$, $-OC(O)R^d$, $-C(O)NR^dR^e$, $-NR^dC(O)R^e$, $-OC(O)NR^dR^e$, $-NR^dC(O)OR^e$, $-NR^dC(O)NR^dR^e$, $-CF_3$ and $-OCF_3$;

5 each R^b is independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and R^a ;

each R^c is independently selected from the group consisting of C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; wherein each cycloalkyl, aryl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents

10 independently selected from the group consisting of C_{1-6} alkyl and R^f ;

each R^d and R^e is independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; or R^d and R^e are joined, together with the atoms to which they are attached, to form a C_{3-6} heterocyclic ring having 1 to 3 heteroatoms
15 independently selected from oxygen, nitrogen or sulfur; wherein each alkyl, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^c and R^f ; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-6} alkyl and R^f ;

20 each R^f is independently selected from the group consisting of $-OH$, $-OC_{1-6}$ alkyl, $-SC_{1-6}$ alkyl, $-F$, $-Cl$, $-NH_2$, $-NH(C_{1-6}$ alkyl), $-N(C_{1-6}$ alkyl)₂, $-OC(O)C_{1-6}$ alkyl, $-C(O)OC_{1-6}$ alkyl, $-NHC(O)C_{1-6}$ alkyl, $-C(O)OH$, $-C(O)NH_2$, $-C(O)NHC_{1-6}$ alkyl, $-C(O)N(C_{1-6}$ alkyl)₂, $-CF_3$ and $-OCF_3$;

m is 0, 1, 2 or 3; and


25 n is 0 or 1.

30

WO 03/031449

PCT/US02/32534

Also disclosed herein are intermediates useful for preparing compounds of formula I, and salts thereof. Accordingly, one embodiment provides a compound of formula II:



or a salt thereof; wherein

R¹ and R² are as defined herein;

15 P¹ and P² are independently hydrogen or an amino-protecting group;

P³ is hydrogen or a carboxy-protecting group;

Q is a leaving group or a group of the formula:




where R³ and *m* are as defined herein; and X⁻ is an optionally present anion;

25 which compounds are useful as intermediates for preparing compounds of formula I and/or as antibiotics.

In a second aspect, there is provided a compound of formula II:

30




or a salt thereof; wherein

P¹ and P² are independently hydrogen or an amino-protecting group;

P³ is hydrogen or a carboxy-protecting group;

Q is a leaving group or a group of the formula:



wherein

R¹ is -Y^a-(W)_n-Y^b-;

W is selected from the group consisting of -O-, -N(R^d)-, -S-, -S(O)-, -S(O)₂-, C₃₋₆ cycloalkylene, C₆₋₁₀ arylene and C₂₋₉ heteroarylene; wherein each arylene, cycloalkylene and heteroarylene group is optionally substituted with 1 to 3 substituents independently selected from R^b;

Y^a and Y^b are independently C₁₋₅ alkylene, or when W is cycloalkylene, arylene or heteroarylene, Y^a and Y^b are independently selected from the group consisting of a covalent bond and C₁₋₅ alkylene; wherein each alkylene group is optionally substituted with 1 to 3 substituents independently selected from -OR^d, -NR^dR^e, -CO₂R^d, -C(O)NR^dR^e and -S(O)₂NR^dR^e;

R² is hydrogen or C₁₋₆ alkyl;

each R³ is independently selected from the group consisting of C₁₋₆ alkyl, C₂₋₆

alkenyl, C₂₋₆ alkynyl, C₃₋₆ cycloalkyl, C₆₋₁₀ aryl, C₂₋₉ heteroaryl, C₃₋₆ heterocyclic and R^a; or two adjacent R³ groups are joined to form C₃₋₆ alkylene or -O-(C₁₋₆ alkylene)-O-; wherein each alkyl, alkylene, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^a and R^c; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^b;

each R^a is independently selected from the group consisting of -OR^d, halo, -SR^d, -S(O)R^d, -S(O)₂R^d, -S(O)₂NR^dR^e, -NR^dR^e, -CO₂R^d, -OC(O)R^d, -C(O)NR^dR^e, -NR^dC(O)R^e, -OC(O)NR^dR^e, -NR^dC(O)OR^e, -NR^dC(O)NR^dR^e, -CF₃ and -OCF₃;

each R^b is independently selected from the group consisting of C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl and R^a;

each R^c is independently selected from the group consisting of C₃₋₆ cycloalkyl, C₆₋₁₀ aryl, C₂₋₉ heteroaryl and C₃₋₆ heterocyclic; wherein each cycloalkyl, aryl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C₁₋₆ alkyl and R^f;

each R^d and R^e is independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₆ cycloalkyl, C₆₋₁₀ aryl, C₂₋₉ heteroaryl and C₃₋₆ heterocyclic; or R^d and R^e are joined, together with the atoms to which they are attached, to form a C₃₋₆ heterocyclic ring having 1 to 3 heteroatoms independently selected from oxygen, nitrogen or sulfur; wherein each alkyl, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^c and R^f; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C₁₋₆ alkyl and R^f;

each R^f is independently selected from the group consisting of -OH, -OC₁₋₆ alkyl, -SC₁₋₆ alkyl, -F, -Cl, -NH₂, -NH(C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -OC(O)C₁₋₆ alkyl, -C(O)OC₁₋₆ alkyl, -NHC(O)C₁₋₆ alkyl, -C(O)OH, -C(O)NH₂, -C(O)NHC₁₋₆ alkyl, -C(O)N(C₁₋₆ alkyl)₂, -CF₃ and -OCF₃;

X⁻ is an optionally present anion;


m is 0, 1, 2 or 3; and

n is 0 or 1.

WO 03/031449


PCT/US02/32534

In yet another embodiment, there is provided a compound of formula III:



10 or salts thereof; wherein R^1 , R^2 , P^1 and P^2 are as defined herein, and P^4 is hydrogen or a carboxy-protecting group; which compounds are useful as intermediates for preparing compounds of formula I or II.

15 According to a third aspect, there is provided a compound of formula III:



or salts thereof; wherein

25 P^1 and P^2 are independently hydrogen or an amino-protecting group;

P^4 is hydrogen or a carboxy-protecting group;

R^1 is $-Y^a-(W)_n-Y^b-$;

30 W is selected from the group consisting of $-O-$, $-N(R^d)-$, $-S-$, $-S(O)-$, $-S(O)_2-$, C_{3-6} cycloalkylene, C_{6-10} arylene and C_{2-9} heteroarylene; wherein each arylene, cycloalkylene and heteroarylene group is optionally substituted with 1 to 3 substituents independently selected from R^b ;

Y^a and Y^b are independently C_{1-3} alkylene, or when W is cycloalkylene, arylene or heteroarylene, Y^a and Y^b are independently selected from the group consisting of a covalent bond and C_{1-3} alkylene; wherein each alkylene group is optionally substituted with 1 to 3 substituents independently selected from $-OR^d$, $-NR^dR^e$, $-CO_2R^d$, $-C(O)NR^dR^e$ and $-S(O)_2NR^dR^e$;

R^2 is hydrogen or C_{1-6} alkyl;

each R^a is independently selected from the group consisting of $-OR^d$, halo, $-SR^d$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2NR^dR^e$, $-NR^dR^e$, $-CO_2R^d$, $-OC(O)R^d$, $-C(O)NR^dR^e$, $-NR^dC(O)R^e$, $-OC(O)NR^dR^e$, $-NR^dC(O)OR^e$, $-NR^dC(O)NR^dR^e$, $-CF_3$ and $-OCF_3$;

each R^b is independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and R^2 ;

each R^c is independently selected from the group consisting of C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; wherein each cycloalkyl, aryl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-6} alkyl and R^f ;

each R^d and R^e is independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; or R^d and R^e are joined, together with the atoms to which they are attached, to form a C_{3-6} heterocyclic ring having 1 to 3 heteroatoms independently selected from oxygen, nitrogen or sulfur; wherein each alkyl, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^e and R^f ; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-6} alkyl and R^f ;

each R^f is independently selected from the group consisting of $-OH$, $-OC_{1-6}$ alkyl, $-SC_{1-6}$ alkyl, $-F$, $-Cl$, $-NH_2$, $-NH(C_{1-6}$ alkyl), $-N(C_{1-6}$ alkyl)₂, $-OC(O)C_{1-6}$ alkyl, $-C(O)OC_{1-6}$ alkyl, $-NHC(O)C_{1-6}$ alkyl, $-C(O)OH$, $-C(O)NH_2$, $-C(O)NHC_{1-6}$ alkyl, $-C(O)N(C_{1-6}$ alkyl)₂, $-CF_3$ and $-OCF_3$; and

n is 0 or 1.

In separate and distinct embodiments, there are provided compounds of formulae 2, 5b, 7, 8, 10, 11 and 13 as defined herein, or salts or protected derivatives thereof; which compounds are useful as intermediates for preparing compounds of formula I and/or as antibiotics.

In a fourth aspect, this invention provides a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a therapeutically effective amount of a compound of formula I, or a pharmaceutically-acceptable salt thereof.

While not intending to be limited by theory, the compounds of formula I are believed to inhibit bacterial cell wall biosynthesis thereby inhibiting the growth of the bacteria or causing lysis of the bacteria. Therefore, among other properties, the compounds of formula I are useful as antibiotics.

Accordingly, one embodiment provides a method of treating a bacterial infection in a mammal, the method comprising administering to a mammal a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a therapeutically effective amount of a compound of formula I, or a pharmaceutically-acceptable salt thereof.

In a fifth aspect, there provided a use of a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula I, or a pharmaceutically-acceptable salt thereof, in the manufacture of a medicament for treating a bacterial infection in a mammal in need thereof.

WO 03/031449

PCT/US02/32534

Additionally, in a sixth aspect, this invention provides a method of inhibiting the growth of bacteria, the method comprising contacting bacteria with a growth-inhibiting amount of a compound of formula I, or a pharmaceutically-acceptable salt thereof.

5 In a seventh aspect, this invention provides a method of inhibiting bacterial cell wall biosynthesis, the method comprising contacting bacteria with a cell wall biosynthesis-inhibiting amount of a compound of formula I, or a pharmaceutically-acceptable salt thereof.

10 Disclosed herein are processes for preparing compounds of formula I or a salt thereof. Accordingly, in an eighth aspect, this invention provides a process for preparing a compound of formula I, or a salt thereof, the process comprising:

(a) reacting a glycopeptide of formula 1 as defined herein, with a compound of formula 2 as defined herein; or


15 (b) reacting a compound of formula 10 as defined herein, with a compound of formula 11 as defined herein; or

(c) reacting a compound of formula 9 as defined herein, with a compound of formula 13 as defined herein;


20 to provide a compound of formula I or a salt thereof. In one preferred embodiment, the above process further comprises the step of forming a pharmaceutically-acceptable salt of a compound of formula I.

25 In a ninth aspect, there is provided a process for preparing a compound according to formula I, or a pharmaceutically-acceptable salt thereof; the process comprising reacting a compound of formula 10:

30




or a salt thereof; with a compound of formula **11**:




or a salt thereof; to provide a compound of formula I or a salt thereof.

In a tenth aspect, there is provided a process for preparing a compound according to formula I, or a pharmaceutically-acceptable salt thereof; the process comprising reacting a compound of formula **9**:



or a salt thereof; with a compound of formula **13**:



or a salt thereof; to provide a compound of formula I or a salt thereof.

In an eleventh aspect, there is provided a product prepared by any of these processes.

A twelfth aspect is directed to a compound of formula I, or a pharmaceutically-acceptable salt thereof, for use in therapy.

Additionally, a thirteenth aspect is directed to the use of a compound of formula I, or a pharmaceutically-acceptable salt thereof, for the manufacture of a medicament for the treatment of a bacterial infection in a mammal.

WO 03/031449

PCT/US02/32534

DETAILED DESCRIPTION OF THE INVENTION

This invention provides novel glycopeptide – cephalosporin compounds of formula I or pharmaceutically-acceptable salts thereof. These compounds have multiple chiral centers and, in this regard, the compounds are intended to have the stereochemistry shown. In particular, the glycopeptide portion of the compound is intended to have the stereochemistry of the corresponding naturally-occurring glycopeptide (i.e., vancomycin, chloroorienticin A and the like). The cephalosporin portion of the molecule is intended to have the stereochemistry of known cephalosporin compounds. However, it will be understood by those skilled in the art that minor amounts of isomers having a different stereochemistry from that shown may be present in the compositions of this invention provided that the utility of the composition as a whole is not significantly diminished by the presence of such isomers.

Additionally, the linking portion of the compounds of this invention (i.e., R¹) may contain one or more chiral centers. Typically, this portion of the molecule will be prepared as a racemic mixture. If desired, however, pure stereoisomers (i.e., individual enantiomers or diastereomers) may be used or a stereoisomer-enriched mixture can be employed. All such stereoisomers and enriched mixtures are included within the scope of this invention.

In addition, compounds of this invention contain several acidic groups (i.e., carboxylic acid groups) and several basic groups (i.e., primary and secondary amine groups) and therefore, the compounds of formula I can exist in various salt forms. All such salt forms are included within the scope of this invention. Also, since the compounds of formula I contain a pyridinium ring, an anionic counterion for the pyridinium group may optionally be present including, but not limited to, halides, such as chloride; carboxylates, such as acetate; and the like.

Furthermore, it will be understood by those skilled in the art that labile or chemically unstable compounds which lack any utility due to their instability are not included within the scope of this invention. For example, it is preferred that compounds of formula I contain at least two carbon atoms between any oxygen

WO 03/031449

PCT/US02/32534


(-O-), nitrogen (-N<) or sulfur (-S-) atoms in the -O-R¹-N(R²)- moiety, since when these atoms are separated by a single carbon atom the resulting compound (i.e., containing an acetal, hemiacetal, ketal, hemiketal, aminal, hemiaminal or thioketal group and the like) may be hydrolytically unstable under acidic conditions.

5

Preferred Embodiments

In the compounds of formula I, the following substituents and values are preferred:

10 In a preferred embodiment, the present invention is directed to compounds of formula Ia:



or a pharmaceutically-acceptable salt thereof, wherein R¹, R², R³ and *m* are as

WO 03/031449

PCT/US02/32534

defined herein, including preferred embodiments.

In a preferred embodiment, R^1 is $-Y^a-Y^b-$, i.e., when n is 0. In this embodiment, Y^a and Y^b are independently $C_{1,3}$ alkylene groups wherein each alkylene group is optionally substituted with 1 to 3 substituents independently selected from

5 $-OR^d$, $-NR^dR^e$, $-CO_2R^d$, $-C(O)NR^dR^e$ and $-S(O)_2NR^dR^e$ as defined herein. Preferably, Y^a and Y^b are independently selected from $C_{1,3}$ alkylene; and more preferably, $C_{1,2}$ alkylene. More preferably, Y^a and Y^b are joined together (i.e., R^1) to form a $-(CH_2)_{2,3}-$ group. Still more preferably, Y^a and Y^b are joined together to form

10 a $-(CH_2)_2-$, $-(CH_2)_3-$, $-(CH_2)_4-$, $-(CH_2)_5-$ or $-(CH_2)_6-$ group. In a particularly preferred embodiment, Y^a and Y^b are joined together to form $-(CH_2)_3-$ group.

In another preferred embodiment, R^1 is $-Y^a-W-Y^b-$, i.e., when n is 1. In this embodiment, Y^a and Y^b are independently $C_{1,3}$ alkylene, or when W is cycloalkylene, arylene or heteroarylene, Y^a and Y^b are independently selected from the group consisting of a covalent bond and $C_{1,3}$ alkylene. Each alkylene group in this

15 embodiment is optionally substituted with 1 to 3 substituents selected from $-OR^d$, $-NR^dR^e$, $-CO_2R^d$, $-C(O)NR^dR^e$ and $-S(O)_2NR^dR^e$ as defined herein. When Y^a or Y^b is an alkylene group, the alkylene group is preferably a $C_{1,3}$ alkylene group; more preferably, a $C_{1,2}$ alkylene group; still more preferably, a $-(CH_2)_{1,2}-$ group. In a particularly preferred embodiment, Y^a and Y^b are both $-CH_2-$ and W is C_{6-10} arylene

20 optionally substituted with 1 to 3 substituents independently selected from R^b as defined herein; more preferably, W is phenylene. In another preferred embodiment, Y^a and Y^b are both $-CH_2CH_2-$ and W is $-O-$.

When present, W is preferably C_{6-10} arylene or $-O-$. More preferably, W is phenylene or $-O-$.

25 Preferably, R^2 is hydrogen or $C_{1,3}$ alkyl. More preferably, R^2 is hydrogen.

When present, each R^3 is preferably independently selected from $C_{1,6}$ alkyl, $C_{3,6}$ cycloalkyl, $-OR^d$, $-SR^d$, $-F$ or $-Cl$; or two adjacent R^3 groups are joined to form $C_{3,6}$ alkylene.

In a preferred embodiment, R^4 is hydroxy and R^5 is hydrogen. In another

30 preferred embodiment, R^4 is hydrogen and R^5 is hydroxy.

WO 03/031449

PCT/US02/32534

Preferably, R⁶ is hydrogen and R⁷ is methyl.

Preferably, R⁸ is hydrogen.

Preferably, one of X¹ and X² is chloro and the other is hydrogen; or both are chloro. More preferably, X¹ and X² are both chloro.

5 In a preferred embodiment, R⁴ is hydroxy; R⁵ is hydrogen; R⁶ is hydrogen; R⁷ is methyl; R⁸ is hydrogen; and X¹ and X² are both chloro (i.e., the glycopeptide portion is vancomycin).

In another preferred embodiment, R⁴ is hydrogen; R⁵ is hydroxy; R⁶ is hydrogen; R⁷ is methyl; R⁸ is a group of formula (i); and X¹ and X² are both chloro
10 (i.e., the glycopeptide portion is chloroorienticin or A82846B).

In a preferred embodiment, *m* is 0. In another preferred embodiment, *m* is 1 or 2; more preferably, 1. In still another preferred embodiment, *m* is 2 and the two R³ groups are joined to form a C₃₋₅ alkylene group; more preferably a C₃₋₄ alkylene group.

15 A preferred group of compounds of formula I are those of formula Ia wherein R¹ is -Y^a-(W)_{*n*}-Y^b-, where *n* is 0 and Y^a and Y^b are joined together to form a -(CH₂)₂₋₈- group; R² is hydrogen and *m* is 0; or a pharmaceutically-acceptable salt thereof. In this embodiment, R¹ (i.e., Y^a and Y^b taken together) is preferably a
20 -(CH₂)₂-, -(CH₂)₃-, -(CH₂)₄-, -(CH₂)₅- or -(CH₂)₆- group; more preferably, -(CH₂)₃-.

Another preferred group of compounds of formula I are those of formula Ia wherein R¹, R², R³ and *m* are as defined in Table I, or a pharmaceutically-acceptable salt thereof.

25

30

WO 03/031449

PCT/US02/32534

Table I

Ex. No.	R ¹				R ²	R ³	m	
	Y ^a	W	Y ^b	n				
5	1	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	---	0
	2	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2-CH ₃ -	1
	3	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3-CH ₃ -	1
	4	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-CH ₃ -	1
10	5	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2-CH ₃ O-	1
	6	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3-CH ₃ O-	1
	7	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-CH ₃ O-	1
	8	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2-CH ₃ S-	1
15	9	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3-CH ₃ S-	1
	10	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-CH ₃ S-	1
	11	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2-F-	1
	12	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3-F-	1
20	13	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-F-	1
	14	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2-Cl-	1
	15	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3-Cl-	1
	16	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-Cl-	1
25	17	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2-Ph- ¹	1
	18	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3-Ph-	1
	19	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-Ph-	1
	20	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-cyclopropyl-	1
	21	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-[HOOCCH ₂ S-]	1
	22	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	4-NH ₂ C(O)-	1
	23	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2,3-di-CH ₃ -	2
	24	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3,4-di-CH ₃ -	2

WO 03/031449

PCT/US02/32534

Ex. No.	R ¹				R ²	R ³	m
	Y ^a	W	Y ^b	n			
25	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3,5-di-CH ₃ -	2
26	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3,4-di-CH ₃ O-	2
27	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3-CH ₃ -4-CH ₃ O-	2
28	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	3-CH ₃ O-4-F-	2
29	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2,3-[-(CH ₂) ₄ -]	2
30	-CH ₂ CH ₂ -	---	-CH ₂ -	0	-H	2,3-(-CH ₂ CH ₂ CH ₂ -)	2
31	-(CH ₂) ₃ -	---	-(CH ₂) ₃ -	0	-H	---	0
32	-CH ₂ CH ₂ -	-O-	-CH ₂ CH ₂ -	1	-H	---	0
33	-CH ₂ -	1,4-(-Ph-) ²	-CH ₂ -	1	-H	---	0

10 ¹ Ph = phenyl

² 1,4-(-Ph-) = 1,4-phenylene

In the intermediate of formula II:

Q is preferably halo or the defined pyridinium group.

15 P¹ is preferably hydrogen or *tert*-butoxycarbonyl.

P² is preferably hydrogen or triphenylmethyl.

P³ is preferably hydrogen or *p*-methoxybenzyl.

R¹, R², R³ and *m* are preferably as defined herein including any preferred
embodiments, substituents or values.

20 In the intermediate of formula III:

P¹ is preferably hydrogen or *tert*-butoxycarbonyl.

P² is preferably hydrogen, formyl or triphenylmethyl.

P⁴ is preferably hydrogen, C₁₋₄ alkyl or *p*-methoxybenzyl.

25 R¹ and R² are preferably as defined herein including any preferred
embodiments, substituents or values

WO 03/031449

PCT/US02/32534

Definitions

When describing the compounds, compositions, methods and processes of this invention, the following terms have the following meanings, unless otherwise indicated.

- 5 The term "alkyl" refers to a monovalent saturated hydrocarbon group which may be linear or branched. Unless otherwise defined, such alkyl groups typically contain from 1 to 10 carbon atoms. Representative alkyl groups include, by way of example, methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, isobutyl, *tert*-butyl, *n*-pentyl, *n*-hexyl, *n*-heptyl, *n*-octyl, *n*-nonyl, *n*-decyl and the like.
- 10 The term "alkylene" refers to a divalent saturated hydrocarbon group which may be linear or branched. Unless otherwise defined, such alkylene groups typically contain from 1 to 10 carbon atoms. Representative alkylene groups include, by way of example, methylene, ethane-1,2-diyl ("ethylene"), propane-1,2-diyl, propane-1,3-diyl, butane-1,4-diyl, pentane-1,5-diyl and the like.
- 15 The term "alkenyl" refers to a monovalent unsaturated hydrocarbon group which may be linear or branched and which has at least one, and typically 1, 2 or 3, carbon-carbon double bonds. Unless otherwise defined, such alkenyl groups typically contain from 2 to 10 carbon atoms. Representative alkenyl groups include, by way of example, ethenyl, *n*-propenyl, isopropenyl, *n*-but-2-enyl, *n*-hex-3-enyl and the like.
- 20 The term "alkynyl" refers to a monovalent unsaturated hydrocarbon group which may be linear or branched and which has at least one, and typically 1, 2 or 3, carbon-carbon triple bonds. Unless otherwise defined, such alkynyl groups typically contain from 2 to 10 carbon atoms. Representative alkynyl groups include, by way of example, ethynyl, *n*-propynyl, *n*-but-2-ynyl, *n*-hex-3-ynyl and the like.
- 25 The term "aryl" refers to a monovalent aromatic hydrocarbon having a single ring (i.e., phenyl) or fused rings (i.e., naphthalene). Unless otherwise defined, such aryl groups typically contain from 6 to 10 carbon ring atoms. Representative aryl groups include, by way of example, phenyl and naphthalene-1-yl, naphthalene-2-yl, and the like.
- 30 The term "arylene" refers to a divalent aromatic hydrocarbon having a single

WO 03/031449

PCT/US02/32534

ring (i.e., phenylene) or fused rings (i.e., naphthalenediyl). Unless otherwise defined, such arylene groups typically contain from 6 to 10 carbon ring atoms. Representative arylene groups include, by way of example, 1,2-phenylene, 1,3-phenylene, 1,4-phenylene, naphthalene-1,5-diyl, naphthalene-2,7-diyl, and the like.

5 The term "cycloalkyl" refers to a monovalent saturated carbocyclic hydrocarbon group. Unless otherwise defined, such cycloalkyl groups typically contain from 3 to 10 carbon atoms. Representative cycloalkyl groups include, by way of example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

C The term "cycloalkylene" refers to a divalent saturated carbocyclic
10 hydrocarbon group. Unless otherwise defined, such cycloalkylene groups typically contain from 3 to 10 carbon atoms. Representative cycloalkylene groups include, by way of example, cyclopropane-1,2-diyl, cyclobutyl-1,2-diyl, cyclobutyl-1,3-diyl, cyclopentyl-1,2-diyl, cyclopentyl-1,3-diyl, cyclohexyl-1,2-diyl, cyclohexyl-1,3-diyl, cyclohexyl-1,4-diyl, and the like.

15 The term "halo" refers to fluoro, chloro, bromo and iodo.

 The term "heteroaryl" refers to a monovalent aromatic group having a single ring or two fused rings and containing in the ring at least one heteroatom (typically 1 to 3 heteroatoms) selected from nitrogen, oxygen or sulfur. Unless otherwise defined, such heteroaryl groups typically contain from 5 to 10 total ring atoms. Representative
20 heteroaryl groups include, by way of example, monovalent species of pyrrole, imidazole, thiazole, oxazole, furan, thiophene, triazole, pyrazole, isoxazole, isothiazole, pyridine, pyrazine, pyridazine, pyrimidine, triazine, indole, benzofuran, benzothiophene, benzimidazole, benzthiazole, quinoline, isoquinoline, quinazoline, quinoxaline and the like, where the point of attachment is at any available carbon or
25 nitrogen ring atom.

 The term "heteroarylene" refers to a divalent aromatic group having a single ring or two fused rings and containing at least one heteroatom (typically 1 to 3 heteroatoms) selected from nitrogen, oxygen or sulfur in the ring. Unless otherwise defined, such heteroarylene groups typically contain from 5 to 10 total ring atoms.
30 Representative heteroarylene groups include, by way of example, divalent species of


WO 03/031449

PCT/US02/32534

pyrrole, imidazole, thiazole, oxazole, furan thiophene, triazole, pyrazole, isoxazole, isothiazole, pyridine, pyrazine, pyridazine, pyrimidine, triazine, indole, benzofuran, benzothiophene, benzimidazole, benzthiazole, quinoline, isoquinoline, quinazoline, quinoxaline and the like, where the point of attachment is at any available carbon or
 5 nitrogen ring atom.

The term "heterocyclyl" or "heterocyclic" refers to a monovalent saturated or unsaturated (non-aromatic) group having a single ring or multiple condensed rings and containing in the ring at least one heteroatom (typically 1 to 3 heteroatoms) selected from nitrogen, oxygen or sulfur. Unless otherwise defined, such heterocyclic groups
 10 typically contain from 2 to 9 total ring atoms. Representative heterocyclic groups include, by way of example, monovalent species of pyrrolidine, imidazolidine, pyrazolidine, piperidine, 1,4-dioxane, morpholine, thiomorpholine, piperazine, 3-pyrroline and the like, where the point of attachment is at any available carbon or nitrogen ring atom.

15 The term "cephalosporin" is used herein in its art recognized manner to refer to a β -lactam ring system having the following general formula and numbering system:



25 The term "glycopeptide antibiotic" or "glycopeptide" is used herein in its art recognized manner to refer to the class of antibiotics known as glycopeptides or dalbapheptides. See, for example, R. Nagarajan, "Glycopeptide Antibiotics", Marcel Dekker, Inc. (1994) and references cited therein. Representative glycopeptides include vancomycin, A82846A (eremomycin), A82846B (chloroorienticin A),
 30 A82846C, PA-42867-A (orienticin A), PA-42867-C, PA-42867-D and the like.

WO 03/031449

PCT/US02/32534

The term "vancomycin" is used herein in its art recognized manner to refer to the glycopeptide antibiotic known as vancomycin. In the compounds of the present invention, the point of attachment for the linking moiety is at the "C-terminus" of vancomycin.

5 The term "pharmaceutically-acceptable salt" refers to a salt which is acceptable for administration to a patient, such as a mammal (e.g., salts having acceptable mammalian safety for a given dosage regime). Such salts can be derived from pharmaceutically-acceptable inorganic or organic bases and from pharmaceutically-acceptable inorganic or organic acids. Salts derived from
10 pharmaceutically-acceptable inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium, zinc and the like. Particularly preferred are ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically-acceptable organic bases include salts of primary, secondary and tertiary amines, including substituted
15 amines, cyclic amines, naturally-occurring amines and the like, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperadine,
20 polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like. Salts derived from pharmaceutically-acceptable acids include acetic, ascorbic, benzenesulfonic, benzoic, camphosulfonic, citric, ethanesulfonic, fumaric, gluconic, glucuronic, glutamic, hippuric, hydrobromic, hydrochloric, isethionic, lactic, lactobionic, maleic, malic, mandelic, methanesulfonic,
25 mucic, naphthalenesulfonic, nicotinic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric and tartaric acids.

 The term "salt thereof" refers to a compound formed when the hydrogen of an acid is replaced by a cation, such as a metal cation or an organic cation and the like
30 (e.g., an NH_4^+ cation and the like). Preferably, the salt is a pharmaceutically-

WO 03/031449

PCT/US02/32534

acceptable salt, although this is not required for salts of intermediate compounds which are not intended for administration to a patient.

The term "therapeutically effective amount" refers to an amount sufficient to effect treatment when administered to a patient in need of treatment.

5 The term "treating" or "treatment" as used herein refers to the treating or treatment of a disease or medical condition (such as a bacterial infection) in a patient, such as a mammal (particularly a human or a companion animal) which includes:

- (a) preventing the disease or medical condition from occurring, i.e., prophylactic treatment of a patient;
- 10 (b) ameliorating the disease or medical condition, i.e., eliminating or causing regression of the disease or medical condition in a patient;
- (c) suppressing the disease or medical condition, i.e., slowing or arresting the development of the disease or medical condition in a patient; or
- (d) alleviating the symptoms of the disease or medical condition in a
15 patient.

The term "growth-inhibiting amount" refers to an amount sufficient to inhibit the growth or reproduction of a microorganism or sufficient to cause death or lysis of the microorganism including gram-positive bacteria.

20 The term "cell wall biosynthesis-inhibiting amount" refers to an amount sufficient to inhibit cell wall biosynthesis in a microorganism including gram-positive bacteria.

The term "leaving group" refers to a functional group or atom which can be displaced by another functional group or atom in a substitution reaction, such as a nucleophilic substitution reaction. By way of example, representative leaving groups
25 include chloro, bromo and iodo groups; and sulfonic ester groups, such as mesylate, tosylate, brosylate, nosylate and the like; activated ester groups, such as 7-azabenzotriazole-1-oxy and the like; acyloxy groups, such as acetoxy, trifluoroacetoxy and the like.

30 The term "protected derivatives thereof" refers to a derivative of the specified compound in which one or more functional groups of the compound are protected

WO 03/031449

PCT/US02/32534

from undesired reactions with a protecting or blocking group. Functional groups which may be protected include, by way of example, carboxylic acid groups, amino groups, hydroxyl groups, thiol groups, carbonyl groups and the like. Representative protecting groups for carboxylic acids include esters (such as a *p*-methoxybenzyl ester), amides and hydrazides; for amino groups, carbamates (such as *tert*-butoxycarbonyl) and amides; for hydroxyl groups, ethers and esters; for thiol groups, thioethers and thioesters; for carbonyl groups, acetals and ketals; and the like. Such protecting groups are well-known to those skilled in the art and are described, for example, in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Third Edition, Wiley, New York, 1999, and references cited therein.

The term "amino-protecting group" refers to a protecting group suitable for preventing undesired reactions at an amino group. Representative amino-protecting groups include, but are not limited to, *tert*-butoxycarbonyl (BOC), trityl (Tr), benzyloxycarbonyl (Cbz), 9-fluorenylmethoxycarbonyl (Fmoc), formyl, trimethylsilyl (TMS), *tert*-butyldimethylsilyl (TBS), and the like.

The term "carboxy-protecting group" refers to a protecting group suitable for preventing undesired reactions at a carboxy group. Representative carboxy-protecting groups include, but are not limited to, esters, such as methyl, ethyl, *tert*-butyl, benzyl (Bn), *p*-methoxybenzyl (PMB), 9-fluorenylmethyl (Fm), trimethylsilyl (TMS), *tert*-butyldimethylsilyl (TBS), diphenylmethyl (benzhydryl, DPM) and the like.

General Synthetic Procedures

The cross-linked glycopeptide – cephalosporin compounds of this invention can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be readily determined by one skilled in the art by routine


WO 03/031449

PCT/US02/32534

optimization procedures.

Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary or desired to prevent certain functional groups from undergoing undesired reactions. The choice of a suitable protecting group for a particular functional group as well as suitable conditions for protection and deprotection of such functional groups are well known in the art. Protecting groups other than those illustrated in the procedures described herein may be used, if desired. For example, numerous protecting groups, and their introduction and removal, are described in T. W. Greene and G. M. Wuts, *Protecting Groups in Organic Synthesis*, Third Edition, Wiley, New York, 1999, and references cited therein.

In a preferred method of synthesis, the compounds of formula I are prepared by reacting a glycopeptide of formula 1:




wherein R^4 , R^5 , R^6 , R^7 , R^8 , X^1 and X^2 are as defined herein, or a salt thereof, with a compound of formula 2:

WO 03/031449

PCT/US02/32534

5



10 wherein R^1 , R^2 , R^3 and m are as defined herein, or a salt or protected derivative thereof; to provide a compound of formula I, or a salt or protected derivative thereof.

Typically, this reaction is conducted by coupling glycopeptide 1, or a salt thereof, with about 0.5 to about 1.5 equivalents, preferably about 0.9 to about 1.1 equivalents, of a compound of formula 2 in an inert diluent, such as DMF, using a
 15 conventional carboxylic acid – amine (peptide) coupling reagent. In this reaction, glycopeptide 1, or a salt thereof, is typically first contacted with the coupling reagent in the presence of an excess, preferably about 1.8 to about 2.2 equivalents, of an amine, such as diisopropylethylamine at a temperature ranging from about -20°C to about 25°C , preferably at ambient temperature, for about 0.25 to about 3 hours.
 20 Preferably, excess trifluoroacetic acid (typically about 2 equivalents) is then added to form a TFA salt of any excess diisopropylethylamine. The reaction is then generally cooled to a temperature of about -20°C to about 10°C , preferably to about 0°C , and intermediate 2 is added, followed by excess 2,4,6-collidine. This reaction is typically maintained at about 0°C for about 1 to about 6 hours, or until the reaction is
 25 substantially complete.

A preferred coupling reagent for use in this reaction comprises about 0.5 to about 1.5 equivalents, preferably about 0.9 to about 1.1 equivalents, of benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate (PyBOP) and about 0.5 to about 1.5 equivalents, preferably about 0.9 to about 1.1 equivalents, of 1-
 30 hydroxybenzotriazole (HOBT) or 1-hydroxy-7-azabenzotriazole (HOAT). Other

WO 03/031449

PCT/US02/32534

suitable coupling reagents include *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU); bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOP-Cl); diphenylphosphoryl azide (DPPA); diphenylphosphinic chloride; diphenyl chlorophosphate (DPCP) and HOAT;
5 pentafluorophenyl diphenylphosphinate and the like.

After the coupling reaction is complete, any protecting groups present in the product are then removed using conventional procedures and reagents. Upon completion of this reaction, the reaction product, i.e., a compound of formula I, is isolated and purified using conventional procedures, such as column chromatography,
10 HPLC, recrystallization and the like.

Glycopeptides of formula 1 suitable for use in the above procedure are either commercially available or they can be prepared by fermentation of the appropriate glycopeptide-producing organism, followed by isolation of the glycopeptide from the resulting fermentation broth using art recognized procedures and equipment.

15 The cephalosporin intermediate 2 used in the above procedure is readily prepared from commercially available starting materials and reagents using conventional procedures. By way of example, intermediate 2 can be prepared as shown in Scheme A:

20


25

30

WO 03/031449

PCT/US02/32534

Scheme A



As illustrated in Scheme A, thiazole intermediate **3** (wherein R^9 is an amino-protecting group, such as a trityl group, and R^{10} is a carboxy-protecting group, such as an ethyl group) is first reacted with an ω -functionalized amine of formula **4** (wherein

30

WO 03/031449

PCT/US02/32534

R¹ and R² are as defined herein, R¹¹ is an amino-protecting group, such as a *tert*-butoxycarbonyl (BOC) group, and Z¹ is a leaving group, such as chloro, bromo, iodo, mesylate, tosylate and the like) to provide, after removal of the carboxy-protecting group (i.e., R¹⁰), an intermediate of formula 5a.

5 This reaction is typically conducted by first contacting 3 with about 1.0 to about 1.1 equivalents, preferably with about 1.02 to about 1.06 equivalents, of a compound of formula 4 in an inert diluent, such as DMF, at a temperature ranging from about 0°C to about 50°C, preferably at ambient temperature, for about 0.5 to about 6 hours, or until the reaction is substantially complete. This reaction is typically
10 conducted in the presence of excess, preferably about 1.1 to about 5 equivalents, of a base, such as cesium carbonate. Additionally, when Z¹ is chloro or bromo, a catalytic amount, preferably about 0.2 to about 0.5 equivalents, of an trialkylammonium iodide, such as tetrabutylammonium iodide, is optionally added to facilitate the reaction by generating the iodo derivative of 4 *in situ*.

15 Removal of the carboxy-protecting group (i.e., R¹⁰) then affords intermediate 5a. For example, when the carboxy-protecting group is an alkyl ester, such as an ethyl group, the ester is readily hydrolyzed to the carboxylic acid by contacting the ester with an excess, preferably with about 1.1 to about 2.5 equivalents, of an alkali metal hydroxide, such as sodium hydroxide or potassium hydroxide. This reaction is
20 typically conducted in an inert diluent, such as ethanol, at a temperature ranging from about 0°C to about 100°C for about 0.5 to about 6 hours, or until the reaction is substantially complete, to afford intermediate 5a.

 Thiazole compounds of formula 3 are commercially available from, for example, Aldrich, Milwaukee, WI, or can be prepared from commercially available
25 starting materials and reagents using conventional procedures.

 Similarly, ω -functionalized amines of formula 4 are readily prepared from commercially-available starting materials and reagents using conventional procedures. Preferred compounds of formula 4 include, by way of illustration, *N*-BOC-3-bromopropylamine; *N*-BOC-6-iodohexylamine; *N*-BOC-2-(2-iodoethoxy)-ethylamine;

WO 03/031449

PCT/US02/32534

N-BOC-4-(iodomethyl)benzylamine; and the like. These compounds are readily prepared from commercially available starting using well-known reagents and reaction conditions.

Intermediate 5a is then chlorinated to provide intermediate 5b. This reaction
5 is typically conducted by contacting 5a with about 1.0 to about 1.2 equivalents, of a chlorinating agent, such as *N*-chlorosuccinimide, in an inert diluent, such as chloroform or DMF, at ambient temperature for about 6 to about 24 hours, or until the reaction is substantially complete.

5-Chloro-1,3-thiazole intermediate 5b is then coupled with intermediate 6
10 (wherein R¹² is hydrogen or a suitable carboxyl protecting group, such as a *p*-methoxybenzyl group) to provide intermediate 7. When R¹² is *p*-methoxybenzyl, intermediate 6 is commercially available from Otsuka, Japan. Typically, this reaction is conducted by contacting 5b with about 0.8 to about 1 equivalents of 6 in the presence of a coupling reagent under conventional coupling reaction conditions. A
15 preferred coupling reagent for this reaction is phosphorous oxychloride (typically about 1.1 to about 1.2 equivalents) and an excess amount of an amine, such as 2,4,6-collidine or diisopropylethylamine. The coupling reaction is typically conducted in an inert diluent, such as THF, at a temperature ranging from about -50°C to about 25°C for about 0.5 to about 6 hours, or until the reaction is substantially complete, to afford
20 intermediate 7. To avoid isomerization, this reaction is preferably conducted at -35°C using 2,4,6-collidine as the base.


Intermediate 7 is then reacted with a pyridine or substituted pyridine to afford intermediate 8, where R³ and *n* are as defined herein. This reaction is typically conducted by first exchanging the chloro group in 7 with an iodo group by contacting
25 7 with about one equivalent of sodium iodide in acetone (Finkelstein reaction) or DMF at ambient temperature for about 0.25 to about 2 hours. The resulting iodo intermediate is typically not isolated, but is reacted *in situ* with about 1.1 to about 1.6 equivalents of a pyridine or substituted pyridine to afford 8. Typically, this reaction is conducted at ambient temperature for about 1 to about 12 hours, or until the reaction

WO 03/031449

PCT/US02/32534

is substantially complete. The pyridine or substituted pyridines used in this reaction are either commercially available or can be prepared from commercially available starting materials and reagents using conventional procedures. Representative pyridine derivatives for use in this reaction include pyridine, 2-picoline, 3-picoline, 4-picoline, 2-methoxypyridine, 3-methoxypyridine, 4-methoxypyridine, 2-thiomethoxypyridine, 3-thiomethoxypyridine, 4-thiomethoxypyridine, 4-carboxythiomethoxypyridine, 2-fluoropyridine, 3-fluoropyridine, 4-fluoropyridine, 2-chloropyridine, 3-chloropyridine, 4-chloropyridine, 2-phenylpyridine, 3-phenylpyridine, 4-phenylpyridine, 4-cyclopropylpyridine, nicotinic acid, isonicotinic acid, nicotinamide, isonicotinamide, 2,3-lutidine, 3,4-lutidine, 3,5-lutidine, 3,4-dimethoxypyridine, 4-methoxy-3-methylpyridine, 4-fluoro-3-methoxypyridine, 2,3-cyclopentenopyridine, 2,3-cyclohexenopyridine and the like.

Alternatively, intermediate 5b can be coupled with a compound of formula 9:



20 wherein R^3 , R^{12} and m are as defined herein, to afford intermediate 8. This reaction is typically conducted by contacting 5b with about 0.9 to about 1.1 equivalents of intermediate 9, or a salt thereof, in an inert diluent, such as DMF, in the presence of a coupling reagent, such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC);

25 PyBOP and HOAT or HOBT; HATU; BOP-Cl; DPPA; DPCP and HOAT; and the like. Generally, the coupling reaction is conducted at a temperature ranging from about -40°C to about 25°C for about 1 to about 12 hours, or until the reaction is substantially complete. Compounds of formula 9 are readily prepared from intermediate 6 by reaction of 6 with pyridine or a substituted pyridine under reaction


30 conditions similar to those described above.

WO 03/031449

PCT/US02/32534


Removal of the protecting groups from intermediate **8** using conventional procedures and reagents then affords cephalosporin intermediate **2**. For example, when R^9 is trityl, R^{11} is *tert*-butoxycarbonyl and R^{12} is *para*-methoxybenzyl, the protecting groups are conveniently removed by treating **8** with excess trifluoroacetic acid and excess anisole or triethylsilane in an inert diluent, such as dichloromethane or heptane, at ambient temperature for about 1 to about 12 hours, or until the reaction is complete. The resulting deprotected cephalosporin **2** is typically isolated and purified using conventional procedures, such as precipitation, lyophization and reverse-phase HPLC.

Alternatively, compounds of formula I can be prepared by reacting a glycopeptide derivative of formula **10**:



or a salt thereof, wherein R^1 , R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , X^1 and X^2 are as defined


herein, with a cephalosporin derivative of formula **11**:



10 or a salt or protected derivative thereof (wherein R^3 and m are as defined herein) to afford a compound of formula I, or salt thereof.

This reaction is typically conducted by contacting 10 with about 1 to about 1.5 equivalents of 11 in an inert diluent, such as water, methanol or mixtures thereof, at a pH ranging from about 4 to about 6.5. This reaction is generally conducted at a temperature ranging from about -20°C to about 40°C for about 1 to about 6 hours, or until the reaction is substantially complete.

The vancomycin derivatives of formula **10** employed in this reaction are readily prepared by coupling vancomycin, or a salt thereof, with a phthalimido derivative of the formula:



25 wherein R¹ is as defined herein, under conventional coupling conditions. For example, this reaction is typically conducted by contacting vancomycin with about 1.1 to about 1.2 equivalents of the phthalimido compound in the presence of a coupling reagent, such as PyBOP and HOAT and the like, and a suitable base, such as


30 diisopropylethylamine. Generally, the reaction is conducted in an inert diluent, such

WO 03/031449

PCT/US02/32534

as DMF, at a temperature ranging from about -20°C to about 40°C for about 0.5 to about 6 hours, or until the reaction is substantially complete. The phthalimido compounds employed in this reaction are readily prepared using conventional procedures and reagents.

- 5 Cephalosporin derivatives of formula 11 can be prepared, for example, by coupling a compound of formula 9 above with a thiazole compound of formula 12:



- or a salt thereof, wherein R¹³ is hydrogen or an amino-protecting group (such as a formyl or trityl group). This coupling reaction is typically conducted by contacting 9 with about 0.9 to about 1.1 equivalents of 12 in an inert diluent, such as DMF, in the presence of a coupling reagent, such as EDC and HOAT, and a suitable base, such as 2,4,6-collidine. Generally, this reaction is conducted at a temperature ranging from about -20°C to about 20°C for about 0.5 to about 6 hours, or until the reaction is substantially complete.
- 20


Additionally, compounds of formula I can be prepared by reacting a glycopeptide derivative of formula 13:

25

30

WO 03/031449

PCT/US02/32534



or a salt thereof, wherein R^1 , R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , X^1 and X^2 are as defined herein, with a compound of formula 9 above to afford a compound of formula I, or a salt thereof.

This coupling reaction is typically conducted by contacting 13 with a coupling reagent, such as DIPC and HOAT, and about 0.5 to about 2 equivalents of 9 in an inert diluent, such as DMF, in the presence of a suitable base, such as 2,4,6-collidine. Generally, this reaction is conducted at a temperature ranging from about -20°C to about 40°C for about 1 to about 6 hours, or until the reaction is substantially complete.

The compound of formula 13 used in this reaction is readily prepared by

WO 03/031449

PCT/US02/32534

coupling vancomycin with intermediate 5b above, using conventional coupling procedures described herein.

Further details regarding specific reaction conditions and procedures for preparing representative compounds of this invention or intermediates thereto are
5 described in the Examples set forth below.

Pharmaceutical Formulations

The cross-linked glycopeptide – cephalosporin compounds of this invention are typically administered to a patient in the form of a pharmaceutical composition.
10 Accordingly, in one of its composition aspects, this invention is directed to a pharmaceutical composition comprising a pharmaceutically-acceptable carrier or excipient and a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.

Any conventional carrier or excipient may be used in the pharmaceutical
15 compositions of this invention. The choice of a particular carrier or excipient, or combinations of carriers or excipients, will depend on the mode of administration being used to treat a particular patient or type of bacterial infection. In this regard, the preparation of a suitable pharmaceutical composition for a particular mode of administration, such as oral, topical, inhaled or parenteral administration, is well
20 within the scope of those skilled in the pharmaceutical arts. Additionally, the ingredients for such compositions are commercially-available from, for example, Sigma, P.O. Box 14508, St. Louis, MO 63178. By way of further illustration, conventional formulation techniques are described in *Remington's Pharmaceutical Sciences*, Mace Publishing Co., Philadelphia, PA 17th Ed. (1985) and "*Modern*
25 *Pharmaceutics*," Marcel Dekker, Inc. 3rd Ed. (G.S. Banker & C.T. Rhodes, Eds.).

The pharmaceutical compositions of this invention will typically contain a therapeutically effective amount of a compound of formula I or a pharmaceutically-acceptable salt thereof. Typically, such pharmaceutical compositions will contain
30 from about 0.1 to about 90% by weight of the active agent, and more generally from about 10 to about 30% of the active agent.

WO 03/031449

PCT/US02/32534

Preferred pharmaceutical compositions of this invention are those suitable for parenteral administration, particularly intravenous administration. Such pharmaceutical compositions typically comprise a sterile, physiologically-acceptable aqueous solution containing a therapeutically effective amount of a compound of formula I or a pharmaceutically-acceptable salt thereof.

Physiologically-acceptable aqueous carrier solutions suitable for intravenous administration of active agents are well-known in the art. Such aqueous solutions include, by way of example, 5% dextrose, Ringer's solutions (lactated Ringer's injection, lactated Ringer's plus 5% dextrose injection, acylated Ringer's injection), Normosol-M, Isolyte E, and the like.

Optionally, such aqueous solutions may contain a co-solvent, for example, polyethylene glycol; a chelating agent, for example, ethylenediamine tetracetic acid; a solubilizing agent, for example, a cyclodextrin; an anti-oxidant, for example, sodium metabisulphite; and the like.

If desired, the aqueous pharmaceutical compositions of this invention can be lyophilized and subsequently reconstituted with a suitable carrier prior to administration. In a preferred embodiment, the pharmaceutical composition is a lyophilized composition comprising a pharmaceutically-acceptable carrier and a therapeutically effective amount of a compound of formula I, or a pharmaceutically-acceptable salt thereof. Preferably, the carrier in this composition comprises sucrose, mannitol, dextrose, dextran, lactose or a combination thereof. More preferably, the carrier comprises sucrose, mannitol, or a combination thereof.

In one embodiment, the pharmaceutical compositions of this invention contain a cyclodextrin. When used in the pharmaceutical compositions of this invention, the cyclodextrin is preferably hydroxypropyl- β -cyclodextrin or sulfobutyl ether β -cyclodextrin. In such formulations, the cyclodextrin will comprise about 1 to 25 weight percent; preferably, about 2 to 10 weight percent of the formulation. Additionally, the weight ratio of cyclodextrin to active agent will typically range from about 1:1 to about 10:1.

The pharmaceutical compositions of this invention are preferably packaged in

WO 03/031449

PCT/US02/32534

a unit dosage form. The term "unit dosage form" refers to a physically discrete unit suitable for dosing a patient, i.e., each unit containing a predetermined quantity of active agent calculated to produce the desired therapeutic effect either alone or in combination with one or more additional units. For example, such unit dosage forms
5 may be packaged in sterile, hermetically-sealed ampoules and the like.

The following formulations illustrate representative pharmaceutical compositions of the present invention:

Formulation Example A

10 A frozen solution suitable for preparing an injectable solution is prepared as follows:

Ingredients	Amount
Active Compound	10 to 1000 mg
15 Excipients (e.g., dextrose)	0 to 50 g
Water for Injection Solution	10 to 100 mL

Representative Procedure: The excipients, if any, are dissolved in about 80%
20 of the water for injection and the active compound is added and dissolved. The pH is adjusted with 1 M sodium hydroxide to 3 to 4.5 and the volume is then adjusted to 95% of the final volume with water for injection. The pH is checked and adjusted, if necessary, and the volume is adjusted to the final volume with water for injection. The formulation is then sterile filtered
25 through a 0.22 micron filter and placed into a sterile vial under aseptic conditions. The vial is capped, labeled and stored frozen.

WO 03/031449

PCT/US02/32534

Formulation Example B

A lyophilized powder suitable for preparing an injectable solution is prepared as follows:

5	<u>Ingredients</u>	<u>Amount</u>
	Active Compound	10 to 1000 mg
	Excipients (e.g., mannitol and/or sucrose)	0 to 50 g
	Buffer Agent (e.g., citrate)	0 to 500 mg
10	<u>Water for Injection</u>	<u>10 to 100 mL</u>

Representative Procedure: The excipients and/or buffering agents, if any, are dissolved in about 60% of the water for injection. The active compound is added and dissolved and the pH is adjusted with 1 M sodium hydroxide to 3 to 4.5 and the volume is adjusted to 95% of the final volume with water for injection. The pH is checked and adjusted, if necessary, and the volume is adjusted to the final volume with water for injection. The formulation is then sterile filtered through a 0.22 micron filter and placed into a sterile vial under aseptic conditions. The formulation is then freeze-dried using an appropriate lyophilization cycle. The vial is capped (optionally under partial vacuum or dry nitrogen), labeled and stored under refrigeration.

25

Formulation Example C

An injectable solution for intravenous administration to a patient is prepared from Formulation Example B above as follows:

Representative Procedure: The lyophilized powder of Formulation Example B (e.g., containing 10 to 1000 mg of active compound) is reconstituted with 20 mL of sterile water and the resulting solution is further diluted with 80 mL of sterile saline in a 100 mL infusion bag. The diluted solution is then administered to the patient intravenously over 30 to 120 minutes.

35

WO 03/031449

PCT/US02/32534

Utility

The cross-linked glycopeptide – cephalosporin compounds of the invention are useful as antibiotics. For example, the compounds of this invention are useful for treating or preventing bacterial infections and other bacteria-related medical conditions in mammals, including humans and their companion animals (i.e., dogs, cats, etc.) which are caused by microorganisms susceptible to the compounds of this invention.

Accordingly, in one of its method aspects, this invention provides a method of treating a bacterial infection in a mammal, the method comprising administering to a mammal in need of treatment, a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a therapeutically effective amount of a compound of formula I, or a pharmaceutically-acceptable salt thereof.

By way of illustration, the compounds of this invention are particularly useful for treating or preventing infections caused by Gram-positive bacteria and related microorganisms. For example, the compounds of this invention are effective for treating or preventing infections caused by certain *Enterococcus* spp.; *Staphylococcus* spp., including coagulase negative staphylococci (CNS); *Streptococcus* spp.; *Listeria* spp.; *Clostridium* spp.; *Bacillus* spp.; and the like. Examples of bacterial species effectively treated with the compounds of this invention include, but are not limited to, methicillin-resistant *Staphylococcus aureus* (MRSA); methicillin-susceptible *Staphylococcus aureus* (MSSA); glycopeptide intermediate-susceptible *Staphylococcus aureus* (GISA); methicillin-resistant *Staphylococcus epidermitis* (MRSE); methicillin-sensitive *Staphylococcus epidermitis* (MSSE); vancomycin-sensitive *Enterococcus faecalis* (EFSVS); vancomycin-sensitive *Enterococcus faecium* (EFMVS); penicillin-resistant *Streptococcus pneumoniae* (PRSP); *Streptococcus pyogenes*; and the like. Compounds of this invention are less effective or not effective for treating or preventing infections caused by strains of bacteria which are resistant to both vancomycin and cephalosporins.

Representative types of infections or bacteria-related medical conditions which can be treated or prevented with the compounds of this invention include, but are not

WO 03/031449

PCT/US02/32534

limited to, skin and skin structure infections, urinary tract infections, pneumonia, endocarditis, catheter-related blood stream infections, osteomyelitis, and the like. In treating such conditions, the patient may already be infected with the microorganism to be treated or merely be susceptible to infection in which case the active agent is administered prophylactically.

The compounds of this invention are typically administered in a therapeutically effective amount by any acceptable route of administration. Preferably, the compounds are administered parenterally. The compounds may be administered in a single daily dose or in multiple doses per day. The treatment regimen may require administration over extended periods of time, for example, for several days or for one to six weeks or longer. The amount of active agent administered per dose or the total amount administered will typically be determined by the patient's physician and will depend on such factors as the nature and severity of the infection, the age and general health of the patient, the tolerance of the patient to the active agent, the microorganism(s) causing the infection, the route of administration and the like.

In general, suitable doses will range of from about 0.25 to about 10.0 mg/kg/day of active agent, preferably from about 0.5 to about 2 mg/kg/day. For an average 70 kg human, this would amount to about 15 to about 700 mg per day of active agent, or preferably about 35 to about 150 mg per day.

Additionally, the compounds of this invention are effective for inhibiting the growth of bacteria. In this embodiment, bacteria are contacted either *in vitro* or *in vivo* with a growth-inhibiting amount of a compound of formula I or pharmaceutically-acceptable salt thereof. Typically, a growth-inhibiting amount will range from about 0.008 $\mu\text{g/mL}$ to about 50 $\mu\text{g/mL}$; preferably from about 0.008 $\mu\text{g/mL}$ to about 25 $\mu\text{g/mL}$; and more preferably, from about 0.008 $\mu\text{g/mL}$ to about 10 $\mu\text{g/mL}$. Inhibition of bacterial growth is typically evidenced by a decrease or lack of reproduction by the bacteria and/or by lysis of the bacteria, i.e., by a decrease in colony-forming units in a given volume (i.e., per mL) over a given period of time (i.e., per hour) compared to untreated bacteria.

WO 03/031449

PCT/US02/32534

The compounds of this invention are also effective for inhibiting cell wall biosynthesis in bacteria. In this embodiment, bacterial are contacted either *in vitro* or *in vivo* with a cell wall biosynthesis-inhibiting amount of a compound of formula I or pharmaceutically-acceptable salt thereof. Typically, a cell wall biosynthesis-inhibiting
5 amount will range from about 0.04 $\mu\text{g/mL}$ to about 50 $\mu\text{g/mL}$; preferably from about 0.04 $\mu\text{g/mL}$ to about 25 $\mu\text{g/mL}$; and more preferably, from about 0.04 $\mu\text{g/mL}$ to about 10 $\mu\text{g/mL}$. Inhibition of cell wall biosynthesis in bacteria is typically evidenced by inhibition or lack of growth of the bacteria including lysis of the bacteria.

In addition to surprising and unexpected antibacterial properties, compounds
10 of this invention have also been found to possess acceptable mammalian safety and acceptable aqueous solubility. Additionally, compounds of this invention have been found to have surprising and unexpectedly rapid cidality against certain bacteria, including methicillin-resistant *Staphylococci aureus* (MRSA) and methicillin-resistant *Staphylococci epidermitis* (MRSE). These properties, as well as the antibiotic utility
15 of the compounds of this invention, can be demonstrated using various *in vitro* and *in vivo* assays well-known to those skilled in the art. For example, representative assays are described in further detail in the following Examples.

20

25

30

WO 03/031449

PCT/US02/32534

EXAMPLES

The following synthetic and biological examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of this invention. In the examples below, the following abbreviations have the following meanings unless otherwise indicated. Abbreviations not defined below have their generally accepted meaning.

	BOC	=	<i>tert</i> -butoxycarbonyl
	CFU	=	colony-forming units
10	DCM	=	dichloromethane
	DIPEA	=	diisopropylethylamine
	DMF	=	<i>N,N</i> -dimethylformamide
	DMSO	=	dimethyl sulfoxide
	EtOAc	=	ethyl acetate
15	HOAT	=	1-hydroxy-7-azabenzotriazole
	HPLC	=	high performance liquid chromatography
	MIC	=	minimum inhibitory concentration
	MS	=	mass spectrometry
	PMB	=	<i>p</i> -methoxybenzyl
20	PyBOP	=	benzotriazol-1-yloxytripyrrolidino- phosphonium hexafluorophosphate
	THF	=	tetrahydrofuran
	TLC	=	thin layer chromatography
25	TFA	=	trifluoroacetic acid

All temperatures reported in the following examples are in degrees Celsius (°C) unless otherwise indicated. Also, unless noted otherwise, reagents, starting materials and solvents were purchased from commercial suppliers (such as Aldrich, Fluka, Sigma and the like) and were used without further purification. Vancomycin hydrochloride semi-hydrate was purchased from Alpharma, Inc., Fort Lee, NJ 07024 (Alpharma AS, Oslo, Norway).

Reverse-phase HPLC was typically conducted using a C₁₈ column and (A) 98% water, 2% acetonitrile, 0.1 % TFA, with an excessing gradient (e.g., 0 to about 70%) of (B) 10% water, 90% acetonitrile, 0.1% TFA, unless otherwise stated.

WO 03/031449

PCT/US02/32534

Example A

Synthesis of

(7R)-7-[(Z)-2-(2-Amino-5-chlorothiazol-4-yl)-2-(3-aminopropoxyimino)acetamido]-3-[(1-pyridinio)methyl]-3-cephem-4-carboxylate
bis-Trifluoroacetic Acid Salt

The following synthesis is illustrated, in part, in Scheme A above.

Step 1 – Preparation of *N*-(*tert*-Butoxycarbonyl)-3-bromopropylamine (i.e. Compound 4 where R¹ is -(CH₂)₃-, R² is hydrogen, R¹¹ is BOC, and Z¹ is bromo)

3-Bromopropylamine hydrobromide (100 g, 457 mmol) was suspended in 1.6 L of anhydrous THF. This mixture was cooled to 0°C in an ice/water bath and stirred vigorously while 190 mL of triethylamine was added. To this mixture was added dropwise *tert*-butoxycarbonyl anhydride (112.6 g, 516 mmol) in 200 mL THF. The ice bath was allowed to warm to ambient temperature and the mixture was stirred overnight at which time TLC indicated the reaction was complete. The mixture was then filtered and the filtrate was concentrated under vacuum. The residual oil was diluted with 1500 mL hexane and stored at -20°C for 3 days. The mixture was then decanted and the residual solid was dried under vacuum to give 101 g (94% yield) of the title intermediate as a crystalline white solid.

¹H NMR (DMSO-d₆, 300 MHz): δ 1.35-1.39 (s, 9H), 1.91-1.95 (m, 2H), 2.99-3.04 (t, 2H), 3.43-3.52 (t, 2H), 6.95-6.99 (t, 1H).

Step 2 – Preparation of Ethyl (Z)-2-(2-Triphenylmethylaminothiazol-4-yl)-2-(3-*N*-BOC-aminopropoxyimino)acetate (i.e. ethyl ester of Compound 5a where R¹ is -(CH₂)₃-, R² is hydrogen, R⁹ is triphenylmethyl, R¹¹ is BOC, and A is hydrogen)

Ethyl (Z)-2-(2-triphenylmethylamino)thiazol-4-yl)-2-(hydroxyimino)acetate hydrochloride (100 g, 202.4 mmol) was dissolved in 700 mL of anhydrous DMF. To this stirred mixture was added cesium carbonate (230.8 g, 708.5 mmol) followed by tetrabutylammonium iodide (18.7 g, 50.6 mmol). *N*-BOC-3-bromopropylamine (50.6 g, 212.5 mmol) in DMF (100 mL) was then added dropwise over 30 minutes. The mixture was stirred for two hours at which time HPLC indicated that the reaction was

WO 03/031449

PCT/US02/32534

complete. The mixture was then filtered and the filter cake was washed with 200 mL of DMF. The filtrate was dissolved in 2 L of ethyl acetate and washed with 700 mL of 1N HCl, followed by 700 mL of saturated aqueous sodium bicarbonate, and finally with 500 mL of brine. The organic layer was dried over sodium sulfate, filtered, and concentrated under vacuum. The residual oil was dissolved in 250 mL of boiling ethanol and poured into a beaker. Once the material had completely cooled, the residual clay-like solid was placed in a Büchner funnel and washed with 50 mL of ethanol previously cooled to -20°C (NOTE: the product is moderately soluble in ethanol and use of larger amounts will decrease the overall yield of final product). After air-drying, the residual solid was ground into a fine powder in a mortar and pestle and dried under vacuum to give 117 g (94% yield) of the title intermediate as a fine off-white powder.

¹H NMR (DMSO-d₆, 300 MHz): δ 1.01-1.1 (t, 3H), 1.31 (s, 9H), 1.60-1.70 (t, 2H), 2.94-2.99 (m, 2H), 3.95-4.04 (m, 4H), 6.77-6.81 (t, 1H), 6.95 (s, 1H), 7.16-7.38 (m, 15H), 8.80 (s, 1H).

MS *m/z*: 615.4 [M+H]⁺.

Step 3 – Preparation of (Z)-2-(2-Triphenylmethylaminothiazol-4-yl)-2-(3-N-BOC-aminopropoxyimino)acetate (i.e., Compound 5a where R¹ is -(CH₂)₃-, R² is hydrogen, R³ is triphenylmethyl, R¹¹ is BOC, and A is hydrogen)

The ethyl ester from Step 2 above (84.2 g, 137 mmol) was suspended in 400 mL of anhydrous ethanol and heated in an oil bath to 80°C with stirring. After all material had dissolved, potassium hydroxide (23.1 g, 411 mmol) in 150 mL of ethanol was added dropwise to the solution over 20 minutes. A precipitate began forming 10 minutes after addition of the base was complete, and within another 10 minutes the mixture was solid. The mixture was removed from the oil bath and cooled in an ice bath. Ethyl acetate and water were added to the cooled mixture which was then poured into a separatory funnel. The mixture was washed with 1N phosphoric acid, which caused the formation of a white solid (NOTE: washing the product with a stronger acid, such as 1N HCl causes degradation of the product). Water was added to

WO 03/031449

PCT/US02/32534

the separatory funnel to dissolve this solid, and the organic layer was then washed with saturated aqueous sodium bicarbonate and with brine. The organic layer was dried over sodium sulfate, filtered and concentrated under vacuum to give the title intermediate (80 g, 99% yield) as a dark tan solid.

5

Step 4 – Preparation of (Z)-2-(2-Triphenylmethylamino-5-chlorothiazol-4-yl)-2-(3-N-BOC-aminopropoxyimino)acetate (i.e. Compound 5b where R¹ is -(CH₂)₃-, R² is hydrogen, R⁹ is triphenylmethyl, R¹¹ is BOC, and A is chloro)

The intermediate from Step 3 above (10 g, 17.04 mmol) was dissolved in 70 mL of chloroform and stirred while solid *N*-chlorosuccinimide (2.28 g, 17.04 mmol) was added (NOTE: experiments suggest that excess NCS may produce undesired side products). The mixture was stirred overnight (a minimum of 15 hours) at which time HPLC indicated that the reaction was complete. The mixture was then concentrated under vacuum and the residue was dissolved in a minimal amount of DMF. This mixture was added to vigorously stirred water to form a precipitate which was then collected by filtration. The solid was air-dried to give 9.5 g (90% yield) of the title intermediate as a tan solid. ¹H NMR indicated only a minimal amount of succinimide remaining (NOTE: isolation of chlorinated product is not necessary for successful coupling in next step, but experiments suggest that residual succinimide may interfere with subsequent pyridine displacement). Alternatively, after the chlorination reaction was complete, the reaction mixture was washed with water (3x), brine, and then dried over anhydrous sodium sulfate. This solution was then filtered and concentrated under vacuum to give the title intermediate (90%) as a tan solid.

¹H NMR (DMSO-d₆, 300 MHz): δ 1.37 (s, 9H), 1.63-1.74 (t, 2H), 2.94-2.99 (m, 2H), 3.97-4.05 (t, 2H), 6.80-6.85 (t, 1H), 7.18-7.41 (m, 15H), 8.97 (s, 1H).

MS m/z 621.3 [M+H]⁺.

30

WO 03/031449

PCT/US02/32534

Step 5 – Preparation of (7R)-7-[(Z)-2-(2-Triphenylmethylamino-5-chlorothiazol-4-yl)-2-(3-N-BOC-aminopropoxyimino)acetamido]-3-chloromethyl-3-cephem-4-carboxylate *p*-Methoxybenzyl Ester (i.e. Compound 7 where R¹ is -(CH₂)₃-, R² is hydrogen, R⁹ is triphenylmethyl, R¹¹ is BOC, and R¹² is *p*-methoxybenzyl)

The intermediate from Step 4 (0.62 g, 1 mmol) was dissolved in 6 mL of anhydrous THF, and to this mixture was added 0.34 g (0.83 mmol) of 7-amino-3-chloromethylcephalosporanic acid *p*-methoxybenzyl ester hydrochloride (i.e., compound 6 where R¹² is PMB; obtained from Otsuka, Japan) in 4 mL of anhydrous THF. The resulting mixture was stirred under nitrogen and cooled to -35°C. To this cooled mixture was added diisopropylethylamine (0.52 mL, 3mmol) followed by phosphorous oxychloride (0.11 mL, 1.2 mmol). This mixture was stirred at -20°C for 30 minutes and then quenched with wet THF and diluted with ethyl acetate. This mixture was washed with water, 1N HCl, brine, dried over sodium sulfate, filtered and concentrated to give 0.88 g (100% yield) of the title intermediate as a brown-red solid. ¹H NMR indicated no undesired isomerization and no residual succinimide.

¹H NMR (DMSO-d₆, 300 MHz): δ 1.37 (s, 9H), 1.63-1.74 (t, 2H), 2.94-2.99 (m, 2H), 3.4-3.74 (q, 2H), 3.75 (s, 3H), 3.97-4.05 (t, 2H), 4.40-4.59 (q, 2H), 5.11-5.25 (m, 3H), 5.49-5.54 (m, 1H), 6.75-6.81 (t, 1H), 6.90-6.96 (d, 2H), 7.18-7.41 (m, 17H), 8.97 (s, 1H), 9.41-9.44 (d, 1H).

MS m/z 972.0 [M+H]⁺.

(NOTE: Experiments suggest that DIPEA causes isomerization when the above reaction is carried out on larger scales. A modified procedure which uses 2,4,6-collidine as the base and which maintains the temperature at -35°C for the entire course of the reaction – about 10 minutes – avoids this problem).

WO 03/031449

PCT/US02/32534

Step 6 – Preparation of (7R)-7-[(Z)-2-(2-Triphenylmethylamino-5-chlorothiazol-4-yl)-2-(3-N-BOC-aminopropoxyimino)acetamido]-3-[(1-pyridinio)methyl]-3-cephem-4-carboxylate *p*-Methoxybenzyl Ester (i.e. Compound 8 where R¹ is -(CH₂)₃-, R² is hydrogen, R³ is triphenylmethyl, R¹¹ is BOC, R¹² is *p*-methoxybenzyl and *m* is 0)

The intermediate from Step 5 (500 mg, 0.514 mmol) was dissolved in 2 mL of anhydrous acetone and protected from light using foil. The solution was stirred under a nitrogen atmosphere and 77 mg (0.514 mmol) of sodium iodide was added and the resulting mixture was stirred for 1 hour. Pyridine (63 µL, 0.772 mmol) was added and, after 90 minutes, the mixture was added to 25 mL of ethyl ether. This mixture was centrifuged and the resulting pellet was washed with ethyl ether and centrifuged again. The ether was decanted and the pellet was dried under vacuum to give a quantitative yield of the title intermediate as a tan solid which was used without further purification.

¹H NMR (DMSO-d₆, 300 MHz): δ = 1.37 (s, 9H), 1.63-1.74 (t, 2H), 2.94-2.99 (m, 2H), 3.3-3.50 (q, 2H), 3.4-3.74 (q, 2H), 3.75 (s, 3H), 3.97-4.05 (t, 2H), 5.10-5.12 (d, 1H), 5.21 (s, 2H), 5.50-5.55 (m, 1H), 5.6 (s, 2H), 6.75-6.81 (t, 1H), 6.90-6.96 (d, 2H), 7.18-7.41 (m, 17H), 8.16-8.21 (t, 2H), 8.61-8.70 (t, 1H), 8.96 (s, 1H), 8.98-9.02 (d, 2H), 9.41-9.44 (d, 1H).

MS *m/z* 1014.2 [M+H]⁺.

Step 7 – Preparation of (7R)-7-[(Z)-2-(2-Amino-5-chlorothiazol-4-yl)-2-(3-aminopropoxyimino)acetamido]-3-[(1-pyridinio)methyl]-3-cephem-4-carboxylate Bis-Trifluoroacetic Acid Salt (i.e. Compound 2 where R¹ is -(CH₂)₃-, R² is hydrogen and *m* is 0)

The intermediate from Step 6 (14.4 g) was dissolved in a 1:1 mixture of trifluoroacetic acid and dichloromethane (120 mL). To this stirring mixture was added 6.2 mL of anisole and the resulting mixture was stirred for 3 hours at room temperature. The mixture was then concentrated and the residue dissolved in ethyl acetate and extracted with water. The water layers were lyophilized and the resulting

WO 03/031449

PCT/US02/32534

powder was dissolved in water and purified using reverse-phase prep HPLC. The resulting purified aqueous solution was then lyophilized to give 3.3 g (30% yield) of the title intermediate.

¹H NMR (DMSO-d₆, 300 MHz): δ = 1.80-1.97 (t, 2H), 2.79-2.92 (m, 2H),
5 3.29-3.57 (q, 2H), 4.02-4.15 (t, 2H), 5.15-5.19 (d, 1H), 5.41-5.63 (q, 2H), 5.83-5.92
(m, 1H), 7.39 (s, 2H), 7.77 (s, 3H), 8.17-8.22 (t, 2H), 8.60-8.70 (t, 1H), 9.0-9.08 (d,
2H), 9.59-9.62 (d, 1H).

MS *m/z* 553.1 [M+H]⁺.

(NOTE: The above reaction can also be conducted using triethylsilane in place
10 of the anisole. Additionally, the product can be isolated using ethyl ether trituration).

Example B

Synthesis of
15 **(7R)-7-[(Z)-2-(2-Amino-5-chlorothiazol-4-yl)-2-(3-aminopropoxyimino)acetamido]-3-[(2,3-cyclopenteno-1-pyridinio)methyl]-3-cephem-4-carboxylate bis-Trifluoroacetic Acid Salt**

Using the procedure described in Example A and substituting 2,3-
20 cyclopentenopyridine (obtained from Koei, Japan) for pyridine in Step 6, the title intermediate was obtained.

¹H NMR (DMSO-d₆, 300 MHz): δ = 1.82-1.947 (t, 2H), 2.18-2.29 (m, 2H),
2.40-2.58 (m, 2H), 2.81-2.95 (m, 2H), 3.09-3.17 (t, 2H), 3.21-3.30 (t, 2H), 4.10-4.19
(t, 2H), 5.15-5.19 (d, 1H), 5.40-5.61 (q, 2H), 5.83-5.92 (m, 1H), 7.39 (s, 2H), 7.77 (s,
25 3H), 7.89-7.96 (t, 2H), 8.42-8.48 (d, 1H), 8.62-8.69 (d, 1H), 9.60-9.63 (d, 1H).

MS *m/z* 592.5 [M+H]⁺.

30

Example C

Synthesis of

5 **(7R)-7-[(Z)-2-(2-Amino-5-chlorothiazol-4-yl)-2-(6-aminohexoxyimino)acetamido]-3-[(1-pyridinio)methyl]-3-cephem-4-carboxylate bis-Trifluoroacetic Acid Salt**

Using the procedure described in Example A and substituting *N*-BOC-6-iodohexylamine for *N*-BOC-3-bromopropylamine in Step 2 (and eliminating the
10 tetrabutylammonium iodide), the title intermediate was obtained.

¹H NMR (DMSO-d₆, 300 MHz): δ 1.2 ppm (bs, 4H), 1.3 ppm (m, 2H), 1.5 ppm (m, 2H), 2.7 ppm (m, 2H), 3.3 ppm (dd, 2H), 4.0 ppm (t, 3H), 5.1 ppm (d, 1H), 5.5 ppm (dd, 2H), 5.8 ppm (dd, 1H), 7.25 ppm (bs, 2H), 7.6 ppm⁻ (bs, 3H), 8.2 ppm (dd, 2H), 8.6 ppm (dd, 1H), 9 ppm (dd, 2H), 9.5 ppm (d, 1H).

15 MS *m/z* 594.3(M⁺).

Example D

Synthesis of

20 **(7R)-7-[(Z)-2-(2-Amino-5-chlorothiazol-4-yl)-2-(2-aminoethoxy)ethoxyimino]acetamido]-3-[(1-pyridinio)methyl]-3-cephem-4-carboxylate bis-Trifluoroacetic Acid Salt**

The procedure of Example A was used, except that the following procedure
25 was substituted for Step 2:

Step 2 – Preparation of Ethyl (Z)-2-(2-Triphenylmethyldiaminothiazol-4-yl)-2-[2-(2-*N*-BOC-aminoethyl)ethoxyimino]acetate (i.e. ethyl ester of Compound 5a where R¹ is -(CH₂)₂-O-(CH₂)₂-, R² is hydrogen, R⁹ is triphenylmethyl, R¹¹ is BOC, and A is hydrogen)

30 The intermediate from Step 1 in Example A (42.5 g, 86 mmol) was added to a stirred suspension of *N*-BOC-2-(2-iodoethoxy)-ethylamine (28.5 g, 90 mmol) (prepared in three steps from 2-(2-hydroxyethoxy)ethanol, i.e., (i) BOC₂O, KOH, (ii) MsCl, Et₃N and (iii) NaI) and cesium carbonate (84.1 g, 258 mmol) in DMF (300 mL). The suspension was stirred for 16 h at room temperature at which time HPLC

indicated that the reaction was complete. The reaction mixture was then filtered and the filter cake washed with DMF (100 mL). The filtrate was diluted with ethyl acetate (1 L) and washed with water (300 mL), 1N HCl (200 mL), saturated aqueous sodium bicarbonate (200 mL) and brine (200 mL). The organic layer was dried over magnesium sulfate, filtered, and concentrated *in vacuo*. The residue was purified by flash column chromatography (ethyl acetate:hexane, 1:1) to afford 49.7g (90% yield) of the title intermediate as an off-white solid.

¹H NMR (DMSO-d₆, 300 MHz): δ = 2.96 (s, broad, 2H), 3.20-3.55 (q, 2H), 3.59 (t, 2H), 3.70 (t, 2H), 4.19 (t, 2H), 5.13 (d, 1H), 5.31-5.64 (q, 2H), 5.80 (dd, 1H), 7.40 (s, 2H), 7.87 (s, broad, 3H), 8.20 (t, 2H), 8.64 (t, 1H), 9.23 (d, 2H), 9.55 (d, 1H). MS m/z 503.1 [M-pyridine]⁺.

Example E

15

Synthesis of (7R)-7-[(Z)-2-(2-Amino-5-chlorothiazol-4-yl)-2-(4-aminomethylbenzyloxyimino)acetamido]-3-[(1-pyridinio)methyl]-3-cephem-4-carboxylate bis-Trifluoroacetic Acid Salt

20

Using the procedure described in Example A and Step 2 of Example D and substituting *N*-BOC-4-(iodomethyl)benzylamine (prepared in four step from 4-(aminomethyl)benzoic acid, i.e., (i) BOC₂O, KOH, (ii) LiAlH₄, (iii) MsCl, Et₃N and (iv) NaI) for *N*-BOC-2-(2-iodoethoxy)-ethylamine 3-bromopropylamine hydrobromide in Step 2, the title intermediate was obtained.

25

¹H NMR (DMSO-d₆, 300 MHz): δ = 3.18-3.59 (q, 2H), 4.00 (s, broad, 2H), 5.13 (s, 2H), 5.15 (d, 2H), 5.40-5.64 (q, 2H), 5.85 (dd, 1H), 7.38-7.43 (m, 6H), 8.19-8.23 (m, 4H), 8.64 (t, 1H), 9.17 (d, 2H), 9.71 (d, 1H).

MS m/z 614.1 [M+H]⁺, 535.1 [M-pyridine]⁺.

30

Example F (Comparative)

Synthesis of

(7R)-7-[(Z)-2-(2-Aminothiazol-4-yl)-2-(3-aminopropoxyimino)acetamido]-3-[(1-pyridinio)methyl]-3-cephem-4-carboxylate bis-Trifluoroacetic Acid Salt

5

By eliminating Step 4 in Example A above, the des-chloro derivative of the intermediate of Example A was prepared.

¹H NMR (DMSO-d₆, 300 MHz): δ = 1.75-1.82 (t, 2H), 2.67-2.82 (m, 2H), 3.25-3.61 (q, 2H), 3.98-4.09 (t, 2H), 5.13-5.17 (d, 1H), 5.38-5.58 (q, 2H), 5.79-5.85 (m, 1H), 6.62 (s, 1H) 7.15-7.25 (s, broad, 2H), 7.60-7.75 (s, broad, 3H), 8.16-8.19 (t, 2H), 8.58-8.63 (t, 1H), 8.95-9.01 (d, 2H), 9.57-9.60 (d, 1H).

10

MS m/z 518.6 [M+H]⁺.

Example G

15

Synthesis of

Vancomycin 3-(Aminoxy)propyl Amide

Step 1 – Preparation of N-(3-Aminopropoxy)phthalimide

N-(tert-Butoxycarbonyl)-3-bromopropylamine (from Step 1 in Example A above) (9.58 g, 40.23 mmol) and N-hydroxyphthalimide (6.36 g, 39 mmol) were dissolved in 70 mL of anhydrous DMF. To this solution was added diisopropylethylamine (7.01 mL, 40.23 mmol) resulting in a deep red color. The reaction was stirred at room temperature for 16 hours after which time, the reaction mixture was poured into 500 mL of diethyl ether. The resulting white precipitate was filtered off and discarded. The organic solution was washed with 2x 200 mL saturated sodium bicarbonate and with 2 x 200 mL of water. The organic solution was dried over anhydrous magnesium sulfate, filtered and concentrated to give a white solid. This solid was then dissolved in 50mL of DCM and 50 mL of TFA. After stirring for 1 hour, this solution was poured into 300 mL of diethyl ether. The resulting precipitate was filtered, washed with diethyl ether and dried under vacuum to afford the title intermediate as its trifluoroacetic acid salt.

25

30

¹H NMR (DMSO-d₆, 300 MHz): δ = 1.90 (2H, qn), 2.95 (2H, t), 4.18 (2H, t), 7.79 (4H, s), 7.92 (3H, broad s).

Step 2 – Preparation of Vancomycin 3-(Phthalimidooxy)propyl Amide

5 Vancomycin hydrochloride (10.0 g, 6.74 mmol) and the intermediate from Step 1 (2.70 g, 8.09 mmol) were slurried in 100 mL of anhydrous DMF. Diisopropylethylamine (4.70 mL, 26.98 mmol) was added and the resulting mixture was stirred at room temperature for 10 min. A solution of PyBOP (5.61 g, 10.78mmol) and HOAt (1.65 g, 10.78 mmol) in DMF (20 mL) was then added, and
10 the reaction was stirred at room temperature. After 1 hour, the reaction mixture was added to diethyl ether (500 mL). The resulting precipitate was filtered, washed with diethyl ether and dried under vacuum to yield the title intermediate as an off-white solid.

MS m/z = 1651.8 (M+H)⁺.

15

Step 3 – Preparation of Vancomycin 3-(Aminooxy)propyl Amide

The intermediate from Step 2 (11.2 g, 6.74 mmol) was slurried in 80 mL of anhydrous DMF and hydrazine monohydrate (0.65 mL, 13.48 mmol) was added. The reaction was stirred at room temperature for 4.5 hours and then 1 mL of
20 trifluoroacetic acid was added to the reaction mixture, followed by 300 mL of diethyl ether. After vigorous stirring, the resulting precipitate was filtered, washed with diethyl ether and dried under vacuum. The title compound was purified by reverse phase HPLC using a water/methanol gradient to afford the title intermediate as a lyophilized powder.

25 MS m/z = 1522.9 (M+H)⁺.

30

Example H

Synthesis of

(7R)-7-[2-(2-Amino-5-chlorothiazol-4-yl)-2-oxoacetamido]-3-(1-pyridinio)methyl-3-cephem-4-carboxylate Bis-Trifluoroacetate

5

Step 1 – Preparation of Ethyl 2-(2-Formylamino-5-chlorothiazol-4-yl)-2-oxoacetate

Ethyl 2-(Formylaminothiazol-4-yl)-2-oxoacetate (9.1 g, 39.87 mmol) (obtained from Aldrich, Milwaukee, WI) was slurried in 50 mL of anhydrous DMF. **N**-Chlorosuccinimide (5.6 g, 41.86 mmol) was added as a solid and the suspension was stirred at room temperature. After 18 hours, the reaction mixture was poured into 500 mL of water. The resulting white precipitate was filtered, washed with water and air dried to afford the title intermediate as a white solid.

15

Step 2 – Preparation of 2-(2-Formylamino-5-chlorothiazol-4-yl)-2-oxoacetic Acid

To the intermediate from Step 1 (3.6 g, 13.7 mmol) was added 1M NaOH (30 mL, 30mmol). The resulting suspension was stirred at room temperature for 2 hours (at which time the solution was clear) and 1M HCl (30mL, 30mmol) was then added, followed by 100 mL of water. After vigorous stirring, the resulting precipitate was filtered, washed with a minimum amount of cold water and air dried to afford the title intermediate as an off-white solid.

25

Step 3 – Preparation of (7R)-7-[2-(2-Formylamino-5-chlorothiazol-4-yl)-2-oxoacetamido]-3-chloromethyl-3-cephem-4-carboxylate *p*-Methoxybenzyl Ester

The intermediate from Step 2 (1.03 g, 4.37 mmol), 7-amino-3-chloromethylcephalosporanic acid *p*-methoxybenzyl ester hydrochloride (1.95 g, 4.81 mmol) and HOAt (0.74 g, 4.81 mmol) were slurried in 15 mL of anhydrous DMF. The reaction vessel was purged with nitrogen and then cooled to 0°C with an external

ice bath. 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.92 g, 4.81 mmol) was added to the cold reaction mixture, followed by 2,4,6-collidine (0.64 mL, 4.81 mmol). The reaction was stirred at 0°C for 2 hours and then poured into 200 mL of 0.5M HCl. The resulting precipitate was filtered, washed with water and air dried to afford the title intermediate as a red solid. The compound was used without further purification.

MS m/z = 607 (M+Na)⁺.

Step 4 – Preparation of (7R)-7-[2-(2-Formylamino-5-chlorothiazol-4-yl)-2-oxoacetamido]-3-(1-pyridinio)methyl-3-cephem-4-carboxylate *p*-Methoxybenzyl Ester

The intermediate from Step 3 (2.5 g, 4.27 mmol) and sodium iodide (0.64 g, 4.27 mmol) were dissolved in acetone and shielded from light by foil. The reaction was stirred for 10 minutes and then pyridine (0.42 mL, 5.12 mmol) was added. The reaction was then stirred at room temperature for 1 hour and then 300 mL of water were added. The resulting precipitate was filtered, washed with water and air-dried to provide a red solid. This solid was purified on reverse-phase HPLC and the resulting aqueous solution was lyophilized to afford the title intermediate as a lyophilized powder.

MS m/z = 628.1 (M)⁺.

Step 5 – Preparation of (7R)-7-[2-(2-Amino-5-chlorothiazol-4-yl)-2-oxoacetamido]-3-(1-pyridinio)methyl-3-cephem-4-carboxylate Bis-Trifluoroacetate

The intermediate from Step 4 (0.11 g, 0.18 mmol) was dissolved in 5 mL of methanol and concentrated aqueous hydrochloric acid (0.5 mL) was added. The resulting solution was stirred at room temperature for 1.5 hours. The methanol was removed under vacuum and acetonitrile (10 mL) was added. The solution was then concentrated in vacuum and to the residue as added DCM (2 mL) and TFA (2 mL) and the resulting mixture was stirred at room temperature for 1.5 hours. Diethyl ether (50 mL) was then added and the title intermediate was isolated by centrifugation.

WO 03/031449

PCT/US02/32534

This intermediate was used without further purification.

MS m/z = 479.9 (M)⁺.

Example I

Synthesis of

Compound 13 where R¹ is -(CH₂)₃-, R², R⁵, R⁶, R⁸ are hydrogen, R⁴ is hydroxy, R⁷ is methyl, X¹ and X² are chloro, and *m* is 0

Step 1 – Preparation of (Z)-2-(2-Amino-5-chlorothiazol-4-yl)-2-(3-aminopropoxyimino)acetate

The intermediate from Step 4 in Example A (0.75g, 1.21mmol) was dissolved in 5 mL of DCM and 5 mL of trifluoroacetic acid. After 1 hour stirring at room temperature, 100 mL of diethyl ether was added. The resulting precipitate was filtered, washed with diethyl ether and dried under vacuum to afford the title intermediate as a brown solid

Step 2 – Preparation of Compound 13 where R¹ is -(CH₂)₃-, R², R⁵, R⁶, R⁸ are hydrogen, R⁴ is hydroxy, R⁷ is methyl, X¹ and X² are chloro, and *m* is 0

Vancomycin hydrochloride (1.3 g, 0.88 mmol) and HOAt (0.14 g, .088 mmol) were slurried in 3.5 mL of anhydrous DMSO. A solution of PyBOP (0.46 g, 0.88 mmol) in 3.5 mL of anhydrous DMF was added, followed by DIPEA (154 μ L, 0.88 mmol). After stirring for 20 minutes, a solution of the intermediate from Step 1 (0.22 g, 0.44 mmol) in 1 mL of DMF was added, followed quickly by addition of DIEA (0.54 mL, 3.08 mmol). The reaction mixture was stirred at room temperature for 1 hour, then 0.5 mL of trifluoroacetic acid was added, followed quickly by the addition of 100 mL of Et₂O. The resulting precipitate was filtered, washed with Et₂O and dried in vacuum. The crude product was purified by reverse-phase-HPLC and the resulting aqueous solution was lyophilized to afford the title intermediate as a lyophilized powder.

MS m/z = 1711.0 (M+H)⁺.

Example 1

Synthesis of a
Compound of Formula I where R¹ is -(CH₂)₃-, R², R⁵, R⁶, R⁸ are hydrogen, R⁴ is
5 hydroxy, R⁷ is methyl, X¹ and X² are chloro, and m is 0
(Compound 1 in Table I)

Vancomycin hydrochloride (4.2 g, 2.8 mmol) was dissolved in 40 mL of
DMSO. To this solution was added a solution of PyBOP (1.3 g, 2.6 mmol) and
10 HOAT (0.35 g, 2.6 mmol) in 40 mL of DMF, followed by 0.98 mL (5.68mmol) of
diisopropylethylamine. This mixture was stirred at room temperature for 30 minutes
and then quenched with 0.44 mL (5.7mmol) of trifluoroacetic acid. The mixture was
then cooled to 0°C and a solution of the intermediate from Example A above (1.3 g,
2.6 mmol) in 20 mL of DMF at 0°C was added, followed by 1.5 mL (11.4 mmol) of
15 2,4,6-collidine. The mixture was maintained at 0°C for four hours and then quenched
with 1.1 mL of trifluoroacetic acid. This mixture was then added to ethyl ether to
form a precipitate, centrifuged, washed with ether, decanted and dried under vacuum.
The resulting powder was dissolved in water and purified using prep HPLC. The
fractions containing the desired product were lyophilized to give the tri-trifluoroacetic
20 acid salt of the title compound. The anion of the salt was then exchanged using
Amberlyte resin to afford the tri-hydrochloride salt of the title compound (1.4 g, 27%
yield) as a white powder.

MS m/z 953.3 [[M+H]⁺-pyridine]/2; 992.0 [M+H]⁺/2.

25 Additionally, Compounds 2-30 shown in Table 1 are or were prepared using
the procedures of Example A and Example 1 by using in place of the pyridine in Step
6 of Example A, the following substituted pyridines:

- Example 2 – 2-Picoline
30 Example 3 – 3-Picoline
Example 4 – 4-Picoline
Example 5 – 2-Methoxypyridine

- Example 6 – 3-Methoxypyridine
Example 7 – 4-Methoxypyridine
Example 8 – 2-Thiomethoxypyridine
Example 9 – 3-Thiomethoxypyridine
5 Example 10 – 4-Thiomethoxypyridine
Example 11 – 2-Fluoropyridine
Example 12 – 3-Fluoropyridine
Example 13 – 4-Fluoropyridine
Example 14 – 2-Chloropyridine
10 Example 15 – 3-Chloropyridine
Example 16 – 4-Chloropyridine
Example 17 – 2-Phenylpyridine
Example 18 – 3-Phenylpyridine
Example 19 – 4-Phenylpyridine
15 Example 20 – 4-Cyclopropylpyridine
Example 21 – 4-(Carboxythiomethoxy)pyridine
Example 22 – Isonicotinamide
Example 23 – 2,3-Lutidine
Example 24 – 3,4-Lutidine
20 Example 25 – 3,5-Lutidine
Example 26 – 3,4-Dimethoxypyridine
Example 27 – 4-Methoxy-3-methylpyridine
Example 28 – 4-Fluoro-3-methoxypyridine
Example 29 – 2,3-Cyclohexenopyridine
25 Example 30 – 2,3-Cyclopentenopyridine

The above substituted pyridines are either commercially available or can be prepared by literature procedures.

Example 34 (Comparative)

Synthesis of a
des-Chloro Compound of Formula I where R¹ is -(CH₂)₃-, R², R⁵, R⁶, R⁸ are
5 hydrogen, R⁴ is hydroxy, R⁷ is methyl, X¹ and X² are chloro, and m is 0
(Compound 34)

Using the procedure of Example 1 and substituting the des-chloro
cephalosporin intermediate of Example F for the intermediate of Example A, the title
10 compound was prepared.

MS m/z 935.3 [[M+H]⁺-pyridine]/2; 974.9 [M+H]⁺/2.

Example 35

Determination of Minimal Inhibitory Concentrations (MICs)

15 Minimal inhibitory concentration (MICs) assays were performed using the
broth microdilution method set forth in NCCLS guidelines (see, NCCLS. 2000.
*Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow
Aerobically*; Approved Standard – Fifth Ed., Vol. 20, No. 2). Bacterial strains were
obtained from the American Type Tissue Culture Collection (ATCC), Stanford
20 University Hospital (SU), Kaiser Permanente Regional Laboratory in Berkeley (KPB),
Massachusetts General Hospital (MGH), the Centers for Disease Control (CDC), the
San Francisco Veterans' Administration Hospital (SFVA) or the University of
California San Francisco Hospital (UCSF). Vancomycin-resistant enterococci were
phenotyped as Van A or Van B based on their sensitivity to teicoplanin. Some
25 vancomycin-resistant enterococci that had been genotyped as Van A, Van B, Van C1
or Van C2 were also obtained from the Mayo Clinic.

In this assay, cryopreserved bacterial cultures of reference and clinical strains
were streaked for isolation on appropriate agar medium (i.e., Trypticase Soy Agar,
Trypticase Soy Agar with defibrinated sheep erythrocytes, Brain Heart Infusion Agar,
30 Chocolate Agar). Following incubation to allow formation of colonies, these plates
were sealed with parafilm and stored refrigerated for up to two weeks. For
preparation of assay inocula and to ensure low variability, several colonies from a

bacterial isolate cultured on the agar plates were pricked with an inoculating loop and aseptically transferred to Mueller-Hinton Broth (supplemented with divalent cations to required levels based on manufacturer's certification). The broth culture was grown overnight at 35°C, diluted in fresh prewarmed broth and grown to log phase; this is equivalent to a 0.5 MacFarland standard or 1×10^8 colony forming units per milliliter (CFU/mL). Not all cell suspensions, due to species variability, contained 1×10^8 CFU/mL when turbidity is equivalent to the MacFarland standard, therefore acceptable adjustments (based on NCCLS guidelines) were made in dilutions of different bacterial strains. The inoculum was diluted such that 100 μ L of this culture in Mueller-Hinton Broth, supplemented Mueller-Hinton Broth, or Haemophilus test medium, when over layered onto a 2-fold serially diluted series of antibiotic concentrations also in 100 μ L of corresponding medium, in a 96-well microtiter plate resulted in a starting bacterial concentration of 5×10^5 CFU/mL. The plates were then incubated 18-24 hours at 35°C. The MIC was read visually as the lowest concentration well with no bacterial growth. Bacterial growth is defined as more than three pinpoint colonies, a button of precipitated cells larger than 2 mm in diameter, or obvious turbidity.

Strains routinely tested in the initial screen included methicillin-sensitive *Staphylococcus aureus* (MSSA), methicillin-resistant *Staphylococcus aureus* (MRSA), *Staphylococcus aureus* producing penicillinase, methicillin-sensitive *Staphylococcus epidermidis* (MSSE), methicillin-resistant *Staphylococcus epidermidis* (MRSE), vancomycin-sensitive *Enterococcus faecium* (EFMVS), vancomycin-sensitive *Enterococcus faecalis* (EFSVS), vancomycin-resistant *Enterococcus faecium* also resistant to teicoplanin (EFMVR Van A), vancomycin-resistant *Enterococcus faecium* sensitive to teicoplanin (EFMVR Van B), vancomycin-resistant *Enterococcus faecalis* also resistant to teicoplanin (EFSVR Van A), vancomycin-resistant *Enterococcus faecalis* sensitive to teicoplanin (EFSVR Van B), penicillin-sensitive *Streptococcus pneumoniae* (PSSP) and penicillin-resistant *Streptococcus pneumoniae* (PSRP). Because of the inability of PSSP and PSRP to

grow well in Mueller-Hinton broth, MICs with those strains were determined using either TS broth supplemented with defibrinated blood or Haemophilus test medium.

Test compounds having significant activity against the strains mentioned above were then tested for MIC values in a larger panel of clinical isolates including the species listed above as well as non-specified coagulase negative *Staphylococcus* both sensitive and resistant to methicillin (MS-CNS and MR-CNS). Additionally, these test compounds were also assayed for MICs against gram-negative microorganisms, such as *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Enterobacter cloacae*, *Acinetobacter baumannii*, *Haemophilus influenzae* and *Moraxella catarrhalis*.

Table II shows MIC₉₀ data for a compound of this invention against methicillin-resistant *S. aureus* (MRSA) and methicillin-resistant *S. epidermitis* (MRSE) as compared to the known antibiotic, vancomycin.

Table II
Minimum Inhibitory Concentrations (MICs)

Microorganism	Test Compound	MIC ₉₀ ² (µg/mL)
Methicillin-resistant <i>S. aureus</i> (MRSA) (n = 53) ¹	Compound 1	< 0.1
	Vancomycin	3
Methicillin-resistant <i>S. epidermitis</i> (MRSE) and other coagulase-negative staphylococci (n = 34)	Compound 1	< 0.1
	Vancomycin	4

¹ Number of strains tested.

² Minimum inhibitory concentration for 90% of strains tested.

Additionally, as shown in Table III, compounds of this invention also had surprising and unexpected MICs against various methicillin-resistant *S. aureus* strains when compared to a related des-chloro derivative (i.e., Compound 34).

Table III
Minimum Inhibitory Concentrations

Microorganism	MIC ($\mu\text{g/mL}$)	
	Cmpd. 1	Cmpd. 34
MRSA 33591	≤ 0.1	0.17
MRSA MED-103	≤ 0.1	0.20
MRSA MED-104	0.10	0.58
MRSA MED-107	≤ 0.1	0.34
MRSA MED-110	0.20	0.49
MRSA MED-572	≤ 0.1	< 0.1
MRSA MED-84	≤ 0.1	0.29
MRSA MED-85	≤ 0.1	0.32
MRSA MED-86	≤ 0.1	0.29
MRSA MED-87	≤ 0.1	0.49
MRSA MED-88	≤ 0.1	0.34
MRSA MED-89	≤ 0.1	0.18

Example 36

Time-Kill Assay

This time-kill assay is a method for measuring the rate of bactericidal activity of a test compound. These procedures are similar to those described in V. Lorian, "Antibiotics in Laboratory Medicine", Fourth Edition, Williams and Wilkins (1996), pages 104-105. A rapid time-kill is desirable to quickly prevent bacterial colonization and reduce host tissue damage.

Bacterial inocula were prepared as described in Example 35 for determination of MIC. Bacteria were diluted in prewarmed media in shake flasks and incubated with shaking (200 rpm, 35°C). At 0, 1, 4, and 24 hours samples were withdrawn from

the flasks and bacteria were enumerated by plate counting. Subsequent to the initial sampling, a compound to be assayed was added to the shake flask culture. Plate counts at these intervals previous to and following addition of the compound were expressed graphically in a time-kill curve. Bactericidal activity is defined as a ≥ 3 log decrease (reduction greater than or equal to 99.9%) in bacterial cell numbers by 24 hours.

In this assay, a compound of formula I, i.e., Compound 1, was bactericidal against MSSA 13709 and MRSA 33591 at a concentration of ≤ 1 $\mu\text{g/mL}$ in 4 hours. By comparison, vancomycin was bactericidal against MSSA 13709 and MRSA 33591 at a concentration of 4 $\mu\text{g/mL}$ in 24 hours.

Example 37

In vivo Efficacy Studies in Neutropenic Mice

Animals (male CD-1 mice, 20-30g) were acquired from Charles Rivers Laboratories (Gilroy, CA) and allowed access to food and water *ad libitum*. Neutropenia was induced via 200 mg/kg intraperitoneal (IP) injection of cyclophosphamide given four and two days prior to the inoculation of bacteria.

The organism used was either a susceptible or resistant strain of clinically relevant gram positive pathogens, such as methicillin-susceptible *Staphylococcus aureus* (MSSA 13709) and methicillin-resistant *Staphylococcus aureus* (MRSA 33591). The bacterial inoculum concentration was $\sim 10^6$ CFU/mL. Animals were lightly anesthetized with isoflurane and 50 mL of the bacterial inoculum was injected into the anterior thigh. One hour after the inoculation, animals were dosed intravenously with vehicle or the appropriate dose of the test compound. At 0 hours and 24 hours post-treatment, the animals were euthanized (CO_2 asphyxiation) and the anterior and posterior thigh collected aseptically. The thigh was placed into 10 mL sterile saline and homogenized. Dilutions of the homogenate were plated onto triptic soy agar plates which were incubated overnight. The number of bacterial colonies on a given plate was multiplied by the dilution factor, divided by the thigh weight (in

grams) and expressed as log CFU/g. ED_{50} (dose required to produce 50% of the maximum reduction in thigh titre) was estimated for each test compound.

In this assay using MRSA 33591, a compound of formula I, i.e., Compound 1, had an ED_{50} of < 0.20 mg/kg, iv, compared to an ED_{50} of 9 mg/kg, iv, for vancomycin.

5

Example 38

Determination of Aqueous Solubility

The aqueous solubility of a compound of this invention was determined using the following procedure. A 5 wt. % dextrose buffer solution at pH 2.2 was prepared
10 by adding 1 mL of 1 N hydrochloric acid (Aldrich) to 99 mL of a 5 wt. % aqueous dextrose solution (Baxter).

A 1 mg/mL stock solution for calibration standards was then prepared by dissolving 1 mg of the test compound in 1 mL of DMSO. This solution was vortexed for 30 seconds and then sonicated for 10 minutes. The stock solution was then diluted
15 with water to prepare calibration standards having the following concentrations: 50, 125, 250, 375 and 500 ug/mL.

Each test compound (30 mg) was weighed into a Millipore non-sterile, Ultrafree-MC 0.1 um filter unit (Millipore UFC30VVOO) and a magnetic stir bar was added to each unit. The 5 wt. % dextrose buffer solution (750 uL) was then added to
20 each unit and these mixtures were vortexed for 5 minutes. The filter units were then placed in an Eppendorf tube rack and the tube rack was placed on top of a magnetic stirrer. Each unit was then titrated to pH 3 using 1 N NaOH (VWR) and the resulting solutions centrifuged at 7000 rpms for 5 minutes. Each unit was then diluted 200 fold with 5% dextrose buffer solution and the diluted samples were transferred into auto
25 sampler vials for analysis.

The calibration standards and the test samples were analyzed by reverse-phase HPLC using the following conditions:

30

WO 03/031449

PCT/US02/32534


Column: Luna 150 x 4.6 mm; C18; 5u
Mobile phase: A = 5/95, B = 95/5, both = MeCN/H₂O; 0.1 % TFA
Method: 10m Lido 100 (0-100 % B in 6 min)
Injection volume: 20 uL
5 Wavelength: 214 nm

The solubility of each test sample was calculated by comparing the peak area of the test sample to the calibration curve and multiplying by the dilution factor. Using the above procedure with duplicate sample preparations, Compound 1 was
10 found to have a solubility of >47.9 mg/mL.

While the present invention has been described with reference to specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made
15 to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto. Additionally, all publications, patents, and patent documents cited herein are incorporated by reference herein in their entirety to the same extent as if they had been
20 individually incorporated by reference.

WHAT IS CLAIMED IS:

1. A compound of formula I:



or a pharmaceutically-acceptable salt thereof, wherein

X^1 and X^2 are independently selected from the group consisting of hydrogen and chloro;

R^1 is $-Y^a-(W)_n-Y^b-$;


W is selected from the group consisting of -O-, -N(R^d)-, -S-, -S(O)-, -S(O)₂-, C₃₋₆ cycloalkylene, C₆₋₁₀ arylene and C₂₋₉ heteroarylene; wherein each arylene, cycloalkylene and heteroarylene group is optionally substituted with 1 to 3 substituents independently selected from R^b;

- 5 Y^a and Y^b are independently C₁₋₅ alkylene, or when W is cycloalkylene, arylene or heteroarylene, Y^a and Y^b are independently selected from the group consisting of a covalent bond and C₁₋₅ alkylene; wherein each alkylene group is optionally substituted with 1 to 3 substituents independently selected from -OR^d, -NR^dR^e, -CO₂R^d, -C(O)NR^dR^e and -S(O)₂NR^dR^e;

- 10 R² is hydrogen or C₁₋₆ alkyl;

- each R³ is independently selected from the group consisting of C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₆ cycloalkyl, C₆₋₁₀ aryl, C₂₋₉ heteroaryl, C₃₋₆ heterocyclic and R^a; or two adjacent R³ groups are joined to form C₃₋₆ alkylene or -O-(C₁₋₆ alkylene)-O-; wherein each alkyl, alkylene, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^a and R^c; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^b;

- one of R⁴ and R⁵ is hydroxy and the other is hydrogen;
 20 R⁶ and R⁷ are independently hydrogen or methyl;
 R⁸ is hydrogen or a group of formula (i):



- each R^a is independently selected from the group consisting of -OR^d, halo, -SR^d, -S(O)R^d, -S(O)₂R^d, -S(O)₂OR^d, -S(O)₂NR^dR^e, -NR^dR^e, -CO₂R^d, -OC(O)R^d,
- 30

$-\text{C}(\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^d\text{C}(\text{O})\text{R}^e$, $-\text{OC}(\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^d\text{C}(\text{O})\text{OR}^e$, $-\text{NR}^d\text{C}(\text{O})\text{NR}^d\text{R}^e$, $-\text{CF}_3$
and $-\text{OCF}_3$;

each R^b is independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and R^a ;

5 each R^c is independently selected from the group consisting of C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; wherein each cycloalkyl, aryl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-6} alkyl and R^f ;

each R^d and R^e is independently selected from the group consisting of
10 hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; or R^d and R^e are joined, together with the atoms to which they are attached, to form a C_{3-6} heterocyclic ring having 1 to 3 heteroatoms independently selected from oxygen, nitrogen or sulfur; wherein each alkyl, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently
15 selected from the group consisting of R^c and R^f ; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-6} alkyl and R^f ;

each R^f is independently selected from the group consisting of $-\text{OH}$,
 $-\text{OC}_{1-6}$ alkyl, $-\text{SC}_{1-6}$ alkyl, $-\text{F}$, $-\text{Cl}$, $-\text{NH}_2$, $-\text{NH}(\text{C}_{1-6}$ alkyl), $-\text{N}(\text{C}_{1-6}$ alkyl)₂,
20 $-\text{OC}(\text{O})\text{C}_{1-6}$ alkyl, $-\text{C}(\text{O})\text{OC}_{1-6}$ alkyl, $-\text{NHC}(\text{O})\text{C}_{1-6}$ alkyl, $-\text{C}(\text{O})\text{OH}$, $-\text{C}(\text{O})\text{NH}_2$,
 $-\text{C}(\text{O})\text{NHC}_{1-6}$ alkyl, $-\text{C}(\text{O})\text{N}(\text{C}_{1-6}$ alkyl)₂, $-\text{CF}_3$ and $-\text{OCF}_3$;

m is 0, 1, 2 or 3; and

n is 0 or 1.

25 2. The compound according to Claim 1, wherein n is 0 and Y^a and Y^b are independently C_{1-3} alkylene groups wherein each alkylene group is optionally substituted with 1 to 3 substituents independently selected from $-\text{OR}^d$, $-\text{NR}^d\text{R}^e$, $-\text{CO}_2\text{R}^d$, $-\text{C}(\text{O})\text{NR}^d\text{R}^e$ and $-\text{S}(\text{O})_2\text{NR}^d\text{R}^e$.

30 3. The compound according to Claim 1, wherein n is 0 and

Y^a and Y^b are joined together to form a $-(CH_2)_{2-8}-$ group.

4. The compound according to Claim 3, wherein n is 0 and
5 Y^a and Y^b are joined together to form a $-(CH_2)_2-$, $-(CH_2)_3-$, $-(CH_2)_4-$, $-(CH_2)_5-$ or
 $-(CH_2)_6-$ group.

5. The compound according to Claim 4, wherein n is 0 and
Y^a and Y^b are joined together to form a $-(CH_2)_3-$ group.

10

6. The compound according to Claim 1, wherein n is 1 and Y^a and Y^b are
the same or different and each is selected from the group consisting of a covalent bond
and C₁₋₅ alkylene optionally substituted with 1 to 3 substituents selected from $-OR^d$,
 $-NR^dR^e$, $-CO_2R^d$, $-C(O)NR^dR^e$ and $-S(O)_2NR^dR^e$.

15

7. The compound according to Claim 6, wherein W is C₆₋₁₀ arylene or
 $-O-$.

8. The compound according to Claim 1, wherein n is 1 and Y^a and Y^b are
20 both $-CH_2-$ and W is C₆₋₁₀ arylene optionally substituted with 1 to 3 substituents
independently selected from R^b.


9. The compound according to Claim 8, wherein W is phenylene.

10. The compound according to Claim 1, wherein n is 1 and Y^a and Y^b are
25 both $-CH_2CH_2-$ and W is $-O-$.

11. The compound according to any one of Claims 1 to 10, wherein R² is
hydrogen.

30

12. The compound according to any one of Claims 1 to 11, wherein m is 0.
13. The compound according to any one of Claims 1 to 11, wherein m is 1 or 2 and each R^3 is independently selected from the group consisting of C_{1-6} alkyl, C_{3-6} cycloalkyl, $-OR^d$, $-SR^d$, $-F$ or $-Cl$; or two adjacent R^3 groups are joined to form C_{3-6} alkylene.
14. The compound according to any one of Claims 1 to 13, wherein R^4 is hydroxy; R^5 is hydrogen; R^6 is hydrogen; R^7 is methyl; R^8 is hydrogen; and X^1 and X^2 are both chloro.
15. The compound according to Claim 1, wherein R^1 is $-Y^a-(W)_n-Y^b-$, where n is 0 and Y^a and Y^b are joined together to form a $-(CH_2)_3-$ group; R^2 is hydrogen; R^4 is hydroxy; R^5 is hydrogen; R^6 is hydrogen; R^7 is methyl; R^8 is hydrogen; X^1 and X^2 are both chloro; and m is 0.
16. The compound according to Claim 1, wherein R^4 is hydroxy; R^5 is hydrogen; R^6 is hydrogen; R^7 is methyl; R^8 is hydrogen; X^1 and X^2 are both chloro; and R^1 , R^2 , R^3 and m are as defined in Table I.
17. A compound of formula II:




or a salt thereof; wherein

P^1 and P^2 are independently hydrogen or an amino-protecting group;

P^3 is hydrogen or a carboxy-protecting group;

Q is a leaving group or a group of the formula:



wherein

R^1 is $-Y^a-(W)_n-Y^b-$;

W is selected from the group consisting of $-O-$, $-N(R^d)-$, $-S-$, $-S(O)-$, $-S(O)_2-$, C_{3-6} cycloalkylene, C_{6-10} arylene and C_{2-9} heteroarylene; wherein each arylene, cycloalkylene and heteroarylene group is optionally substituted with 1 to 3 substituents independently selected from R^b ;

Y^a and Y^b are independently C_{1-5} alkylene, or when W is cycloalkylene, arylene or heteroarylene, Y^a and Y^b are independently selected from the group consisting of a covalent bond and C_{1-5} alkylene; wherein each alkylene group is optionally substituted with 1 to 3 substituents independently selected from $-OR^d$, $-NR^dR^e$, $-CO_2R^d$, $-C(O)NR^dR^e$ and $-S(O)_2NR^dR^e$;

R^2 is hydrogen or C_{1-6} alkyl;

each R^3 is independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl, C_{3-6} heterocyclic and R^a ; or two adjacent R^3 groups are joined to form C_{3-6} alkylene or $-O-(C_{1-6} \text{ alkylene})-O-$; wherein each alkyl, alkylene, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^a and R^c ; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^b ;

each R^a is independently selected from the group consisting of $-OR^d$, halo, $-SR^d$, $-S(O)R^d$, $-S(O)_2R^d$, $-S(O)_2OR^d$, $-S(O)_2NR^dR^e$, $-NR^dR^e$, $-CO_2R^d$, $-OC(O)R^d$, $-C(O)NR^dR^e$, $-NR^dC(O)R^e$, $-OC(O)NR^dR^e$, $-NR^dC(O)OR^e$, $-NR^dC(O)NR^dR^e$, $-CF_3$ and $-OCF_3$;

5 each R^b is independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and R^a ;

each R^c is independently selected from the group consisting of C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; wherein each cycloalkyl, aryl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents
10 independently selected from the group consisting of C_{1-6} alkyl and R^f ;

each R^d and R^e is independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; or R^d and R^e are joined, together with the atoms to which they are attached, to form a C_{3-6} heterocyclic ring having 1 to 3 heteroatoms
15 independently selected from oxygen, nitrogen or sulfur; wherein each alkyl, alkenyl and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^c and R^f ; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-6} alkyl and R^f ;

20 each R^f is independently selected from the group consisting of $-OH$, $-OC_{1-6}$ alkyl, $-SC_{1-6}$ alkyl, $-F$, $-Cl$, $-NH_2$, $-NH(C_{1-6}$ alkyl), $-N(C_{1-6}$ alkyl) $_2$, $-OC(O)C_{1-6}$ alkyl, $-C(O)OC_{1-6}$ alkyl, $-NHC(O)C_{1-6}$ alkyl, $-C(O)OH$, $-C(O)NH_2$, $-C(O)NHC_{1-6}$ alkyl, $-C(O)N(C_{1-6}$ alkyl) $_2$, $-CF_3$ and $-OCF_3$;

X^- is an optionally present anion;

25 m is 0, 1, 2 or 3; and


n is 0 or 1.

30

WO 03/031449

PCT/US02/32534

18. A compound of formula III:



or salts thereof; wherein

P¹ and P² are independently hydrogen or an amino-protecting group;

P⁴ is hydrogen or a carboxy-protecting group;

R¹ is -Y^a-(W)_n-Y^b-;

W is selected from the group consisting of -O-, -N(R^d)-, -S-, -S(O)-, -S(O)₂-, C₃₋₆ cycloalkylene, C₆₋₁₀ arylene and C₂₋₉ heteroarylene; wherein each arylene, cycloalkylene and heteroarylene group is optionally substituted with 1 to 3 substituents independently selected from R^b;

Y^a and Y^b are independently C₁₋₃ alkylene, or when W is cycloalkylene, arylene or heteroarylene, Y^a and Y^b are independently selected from the group consisting of a covalent bond and C₁₋₃ alkylene; wherein each alkylene group is optionally substituted with 1 to 3 substituents independently selected from -OR^d, -NR^dR^e, -CO₂R^d, -C(O)NR^dR^e and -S(O)₂NR^dR^e;

R² is hydrogen or C₁₋₆ alkyl;

each R^a is independently selected from the group consisting of -OR^d, halo, -SR^d, -S(O)R^d, -S(O)₂R^d, -S(O)₂OR^d, -S(O)₂NR^dR^e, -NR^dR^e, -CO₂R^d, -OC(O)R^d, -C(O)NR^dR^e, -NR^dC(O)R^e, -OC(O)NR^dR^e, -NR^dC(O)OR^e, -NR^dC(O)NR^dR^e, -CF₃ and -OCF₃;

each R^b is independently selected from the group consisting of C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl and R^a;

each R^e is independently selected from the group consisting of C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; wherein each cycloalkyl, aryl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-6} alkyl and R^f ;

- 5 each R^d and R^e is independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{6-10} aryl, C_{2-9} heteroaryl and C_{3-6} heterocyclic; or R^d and R^e are joined, together with the atoms to which they are attached, to form a C_{3-6} heterocyclic ring having 1 to 3 heteroatoms independently selected from oxygen, nitrogen or sulfur; wherein each alkyl, alkenyl
10 and alkynyl group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of R^e and R^f ; and each aryl, cycloalkyl, heteroaryl and heterocyclic group is optionally substituted with 1 to 3 substituents independently selected from the group consisting of C_{1-6} alkyl and R^f ;


- each R^f is independently selected from the group consisting of -OH,
15 -OC $_{1-6}$ alkyl, -SC $_{1-6}$ alkyl, -F, -Cl, -NH $_2$, -NH(C $_{1-6}$ alkyl), -N(C $_{1-6}$ alkyl) $_2$,
-OC(O)C $_{1-6}$ alkyl, -C(O)OC $_{1-6}$ alkyl, -NHC(O)C $_{1-6}$ alkyl, -C(O)OH, -C(O)NH $_2$,
-C(O)NHC $_{1-6}$ alkyl, -C(O)N(C $_{1-6}$ alkyl) $_2$, -CF $_3$ and -OCF $_3$; and


n is 0 or 1.

- 20 19. A pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a therapeutically effective amount of a compound according to any one of Claims 1 to 16.

- 25 20. A use of a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound according to any one of Claims 1 to 16 in the manufacture of a medicament for treating a bacterial infection in a mammal in need thereof.


23. A process for preparing a compound according to any one of Claims 1 to 16; the process comprising reacting a glycopeptide of formula-1:





or a salt thereof; to provide a compound of formula I, or a salt thereof.


24. A process for preparing a compound according to any one of Claims 1 to 16; the process comprising reacting a compound of formula 10:



WO 03/031449


PCT/US02/32534

or a salt thereof; with a compound of formula 11:



or a salt thereof; to provide a compound of formula I or a salt thereof.


25. A process for preparing a compound according to any one of Claims 1 to 16; the process comprising reacting a compound of formula 9:



or a salt thereof; with a compound of formula 13:

WO 03/031449

PCT/US02/32534



- 20 or a salt thereof; to provide a compound of formula I or a salt thereof.
26. The product prepared by the process of any one of Claims 23 to 25.
27. A compound according to any one of Claims 1 to 16 for use in therapy.
- 25 28. A compound according to any one of Claims 1 to 16 for the manufacture of a medicament for the treatment of a bacterial infection in a mammal.

30