

COMBINATION OF PROANTOCYCIDINS SUCH AS PYCNOGENOL OR GRAPE SEEDS AND CENTELLA
ASIATICA FOR THE TREATMENT OF CARDIOVASCULAR DISORDERS SUCH AS ATHEROSCLEROSIS

5

FIELD OF THE INVENTION

The invention relates to preparations for medical purposes and more specifically to a preparation consisting of the combination of proanthocyanidins such as Pycnogenol or grape seeds and Centella asiatica and / or extracts thereof, for use in the prevention or treatment of atherosclerosis.

10

BACKGROUND OF THE INVENTION

Diseases affecting the heart are commonly classified under the term heart diseases. More particularly, cardiovascular diseases refer to the class of heart diseases that involve the heart and the blood vessels (arteries and veins). One type of cardiovascular diseases is atherosclerosis. Atherosclerosis is a syndrome affecting arterial blood vessels which carry oxygen-rich blood to the heart and other parts of the body. It concerns the process in which deposits of fatty substances, cholesterol, cellular waste products, calcium and other substances build up in the inner lining of arteries and form hard substances called plaque. It usually affects large and medium-sized arteries.

15

20

Studies show that atherosclerosis is a slow, complex disease that typically starts in childhood and often progress when people become older. Over time, plaques can grow large enough to significantly reduce the blood's flow through arteries that produce a diminution of oxygen-rich blood to organs. Most of the damage occurs when plaques become fragile and break (rupture). Plaques that break cause blood clots blocking the blood flow or travelling to another part of the body. When a blood vessel that feeds the heart is blocked, it causes a heart attack. If it blocks a blood vessel that feeds the brain, it causes a stroke. And if blood supply to the arms or legs is reduced, it can cause difficulties in moving and eventually gangrene.

25

30

Certain conditions or habits may increase the events of developing atherosclerosis. These conditions (high blood cholesterol, especially LDL, smoking and exposure to tobacco smoke, high blood pressure, diabetes, obesity, physical inactivity...) are known as risk factors. Changes in lifestyle and/or taking medicines to treat some particular risk factors can often
 5 reduce the genetic influences and prevent from atherosclerosis.

As mentioned above, atherosclerosis is a syndrome which affects people from childhood and progress with age. For centuries, medicine tries to find appropriate treatments to prevent and/or minimize the risks of atherosclerosis. Some attempts have been done with the
 10 compositions containing proanthocyanidins (such as Pycnogenol®).

WO 2007/084648 (Mars, Inc.) discloses compositions containing mixed 4 to 6 procyanidin dimers and certain derivatives thereof, for inducing vasodilation, treating or preventing diseases or disorders of the vascular system (e.g. hypertension, cardiovascular disease,
 15 stroke), and/or treating or preventing NO- responsive conditions or diseases. The disclosed compositions may be used as a pharmaceutical, a food, a food additive, or a dietary supplement. The composition may optionally contain an additional therapeutic or beneficial-to-health agent, or may be administered in combination with another therapeutic or beneficial-to-health agent.

20 Abascal K et al. "Botanicals for chronic venous insufficiency" ALTERNATIVE AND COMPLEMENTARY THERAPIES, Vol. 13, N° 6, Dec. 2007, pp 304-311; also discloses botanicals that are useful as adjunctive therapy for the treatment of chronic venous insufficiency (CVI). Botanicals used include Ruscus aculeatus, horse chestnut seed, Centella
 25 asiatica, Vitis vinifera and oligomeric proanthocyanidins such as Pycnogenol®.

Ramelet A et al. "Veno-active drugs in the management of chronic venous disease. An international consensus statement: Current medical position, prospective views and final resolution" CLINICAL HEMORHEOLOGY AND MICROCIRCULATION, Vol. 33, N° 4,
 30 2005, pp 309-319; discloses veno-active drugs (VAD) having effects on edema and symptoms related to chronic venous disease (CVD) especially venous pain. This review has classified various VAD and the level of Evidence-Based Medicine (EBM) of each drug has been determined. Pycnogenol® and Centelly asiatica are been classified under Grade C as showing less well demonstrated effects on chronic venous disease.

WO 2007/133981 (RATH Matthias et al.) discloses biochemical compositions effective in the prevention and treatment resulting from the inhibition of an atherogenic process, comprising ascorbic acid, lysine, magnesium, cysteine, pyridoxine HCL, riboflavin, folic acid,
 5 cyanocobalamin vitamin B12, S-Adenosyl-L-Methionine, choline bitartrate, copper glycinate, epigallocatechin gallate, quercetin, asiatic acid, and Pycnogenol®. More precisely the disclosed compositions act on the growth of smooth muscle cell and the invasion of extracellular matrix by smooth muscle cell. The compositions may be administered orally, intravenously, or parenterally.

10

Other documents cite some possible associations of plant extract among big lists of compounds. For example EP 1 523 894 (Cognis Iberia S.L.) discloses a preparation containing extracts from the plant *Centella asiatica* or the active component (component a). - Plant extracts contained, include *Camellia sinensis*, *Pinia silvestris*, *Vitis vinifera*, *Litchi chinensis*, *Potentilla erecta* and their mixtures. Components a and b or c are in the mass ratio
 15 90 : 10 to 10 : 90. The preparation is encapsulated.

JP 2002 238497 A (Fankaru KK), XP002578473, discloses food compositions for beautifying purposes, especially the skin, that are prepared by including collagen and *Centella asiatica*, and/or oceanic deep layer water and optionally amino sugars, ceramides and
 20 polyphenols. The agents enhance metabolism of the skin to accelerate the synthesis of collagen and preventing it from deteriorating to keep freshness.

KR 100 846 292 B1 (Intaglio INC), XP002578465, discloses a dispersion type non-aqueous cleansing balm composition for face washing, causes no skin irritation and provides refresh feeling after use. Said dispersion containing at least one polyvalent alcohol (40-75 weight%)
 25 chosen from polyalcohol, polyethylene glycol, propylene glycol, glycerin and butylene glycol as solvent, and cleaning agent (5-30 weight%) chosen from sodium cocoyl isethionate, sodium laurylsulfate, sodium laurylglutamate, hydrogenated castor oil, hydrogenated jojoba oil and hydrogenated olive oil. The dispersion further contains scrubbing agent (0,5-15 weight%) chosen from scrub reagent including powdered jojoba wax, apricot stone, grape
 30 seed and walnut skin, kaolin and porous zeolite, dispersing agent (0,1-5 weight%) chosen from poloxamer and hydroxypropylcellulose, and skin irritation-reducing material (0,01-5 weight%) chosen from carotenoid, astaxanthin, beta-carotene, lycopene, phyloxanthin, proanthocyanidin, flavonoid, riboflavin, vitamin, retinol, ascorbic acid, tocopherol, *Centella asiatica*, bisabolol, chamomile, *Glycyrrhiza uralensis* Fisch and soy isoflavone.

WO 03/080062 (Perrella Segre) discloses compounds for use in sclerotherapy treatment, venous insufficiency, circulation and microcirculation disorders consisting of Folic Acid (vitamin B9), Piridoxine (vitamin B6), Lipoic acid and six different plant extracts: Ginkgo

5 Biloba, Garlic Andrographis, Horse Chestnut, Centella Asiatica, and Grape seeds.

Sclerotherapy is a procedure used to treat blood vessels or blood vessel malformations (in particular veins) and also those of the lymphatic system. A medicine is injected into the vessels (i.e. veins), which makes them shrink. In conventional sclerotherapy, which is an obliterative procedure, a sclerosing solution is injected exclusively into pathological

10 superficial veins. It is used for children and young adults with vascular or lymphatic malformations. In adults, sclerotherapy is often used to treat varicose veins and hemorrhoids. Another procedure, the so called three-dimensional regenerative sclerotherapy, concerns a regenerative, non-obliterative sclerosing solution which is used in the sclerotherapy of ectatic vessels in the lower limbs. This solution comes into contact with the entire network of

15 superficial and perforating veins.

However there is still a need for an effective and safe composition for the treatment or the prevention of atherosclerosis and in particular atherosclerotic plaques, which therefore especially concerns arteries.

20

SUMMARY OF THE INVENTION

Applicants have surprisingly found that the combination of Pycnogenol® and Centella asiatica extract shows an interesting potential in the treatment of atherosclerosis. This safe

25 natural combination is particularly promising in the treatment and prevention of atherosclerotic plaques especially in arteries.

In one aspect of the present invention there is provided a preparation consisting of the combination of proanthocyanidins (preferably procyanidins) and Centella asiatica and / or

30 extracts thereof, in a suitable excipient, for use in the prevention or treatment of atherosclerosis.

In another aspect, the present invention provides for a dietary or food supplement, a food preparation, a beverage, a medicament and a topical preparation comprising the preparation of the present invention.

- 5 In a further aspect, the preparation of the present invention is provided for the treatment or the prevention of atherosclerosis which concerns especially arteries.

In a still further aspect, the present invention provides for a method of treating or preventing atherosclerosis, comprising administering to a subject in need thereof an effective amount of
10 the preparation or the medicament of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

Although methods and materials similar or equivalent to those described herein can be used in
15 the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. The publications and applications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not
20 entitled to antedate such publication by virtue of prior invention. In addition, the materials, methods, and examples are illustrative only and are not intended to be limiting.

In the case of conflict, the present specification, including definitions, will control.
Unless defined otherwise, all technical and scientific terms used herein have the same
25 meaning as is commonly understood by one of skill in art to which the subject matter herein belongs. As used herein, the following definitions are supplied in order to facilitate the understanding of the present invention.

The term "comprise" is generally used in the sense of include, that is to say permitting the
30 presence of one or more features or components.

As used in the specification and claims, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise.

As used herein the term "pine bark extract" refers to a French maritime pine bark extract which is, for example, commercially available as Pycnogenol® (Horphag). The terms "Pycnogenol®", "pine bark extract" and "French maritime pine bark extract" are
5 interchangeable. *Pinus pinaster* (*P. pinaster*) and *Pinus maritima* (*P. maritima*), are understood to refer to the same organism commonly called "French Maritime Pine." Hence, these terms are interchangeable.

The term "extract", as used herein includes any preparation obtained from plants, fruits or
10 vegetables using an extraction method.

The term "food preparation" refers generally to material of either plant or animal origin, or of synthetic sources, that contain essential nutrients such as a carbohydrate, protein, fat, vitamin, mineral, etc. used in the body of an organism to sustain growth, repair, and vital processes
15 and to furnish energy

A "dietary or food supplement" refers to a product that contains substances like vitamins, minerals, foods, botanicals, amino acids and is intended to supplement the usual intake of these substances. Dietary supplements are found in pill, tablet, capsule, powder or liquid form
20 and are meant to be taken by mouth.

The term "nutraceutical" refers to any substance that is a food or a part of a food and provides medical or health benefits, including the prevention and treatment of disease. Such products may range from isolated nutrients, dietary supplements and specific diets to genetically
25 engineered designer foods, herbal products, and processed foods such as cereals, soups and beverages. It also refers to a product isolated or purified from foods, and generally sold in medicinal forms not usually associated with food and demonstrated to have a physiological benefit or provide protection against diseases like chronic diseases for example.

30 The term "beverage" means a liquid for drinking, which may be water, flavored water, soft drinks, alcoholic drink, health drink, or an enriched drink like based on a dairy product (milk) or fruit juice.

“Pharmaceutically acceptable excipients or carriers” are any materials that do not interfere with the pharmacological activity of the active ingredient(s) or degrade the body functions of the subject to which it can be administered but facilitate fabrication of dosage forms or administration of the composition. Examples of pharmaceutically acceptable excipient

5 include but are not limited to maltodextrin, calcium phosphate, and fused silica.

Pharmaceutically acceptable excipients also include flavorants, as well as various additives such as other vitamins and minerals, all solvents, dispersion media, coatings, isotonic and absorption delaying agents, sweeteners and the like, non-toxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, such as for example, sodium
10 acetate, sorbitan monolaurate, triethanolamine oleate, and inert ingredients such as talc and magnesium stearate which are standard excipients in the manufacture of tablets, capsules and other dosage forms.

As used herein the terms "subject" or "patient" are well-recognized in the art, and, are used
15 interchangeably herein to refer to a mammal, including dog, cat, rat, mouse, monkey, cow, horse, goat, sheep, pig, camel, and, most preferably, a human. In some embodiments, the subject is a subject in need of treatment or a subject with a disease or disorder. However, in other embodiments, the subject can be a normal subject. The term does not denote a particular age or sex. Thus, adult and newborn subjects, whether male or female, are intended to be
20 covered.

The term “an effective amount” refers to an amount necessary to obtain a physiological effect. The physiological effect may be achieved by one application dose or by repeated applications. The dosage administered may, of course, vary depending upon known factors, such as the
25 physiological characteristics of the particular composition; the age, health and weight of the subject; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; and the effect desired and can be adjusted by a person skilled in the art.

30 The term “TECA” refers to total triterpenic fraction of *Centella asiatica*.

“Atherosclerosis” is a process of progressive thickening and hardening of the walls of medium-sized and large arteries as a result of fat deposits on their inner lining.

In other words, Atherosclerosis (also known as arteriosclerotic vascular disease or ASVD) is a condition in which an artery wall thickens as the result of a build-up of fatty materials such as cholesterol. It is a syndrome affecting arterial blood vessels, a chronic inflammatory response in the walls of arteries, in large part due to the accumulation of macrophage white blood cells and promoted by low-density lipoproteins (plasma proteins that carry cholesterol and triglycerides) without adequate removal of fats and cholesterol from the macrophages by functional high density lipoproteins (HDL). It is commonly referred to as a hardening or furring of the arteries. It is caused by the formation of multiple plaques within the arteries (see: <http://en.wikipedia.org/wiki/Atherosclerosis>).

10

Applicants have studied specific plants and their potential application in the treatment of atherosclerosis. Surprisingly it was found that the administration of a preparation consisting of the combination of proanthocyanidins (preferably procyanidins) and *Centella asiatica* and / or extracts thereof, in a suitable excipient, considerably reduces the atherosclerosis risks.

15

Indeed, the preparation of the present invention - of completely natural origin - stops the progression of atherosclerosis from subclinical to clinical stages in several patients. The synergy action of a composition comprising proanthocyanidins and a composition comprising *Centella asiatica* and / or extracts thereof seems to be separated by any other pharmacological action (i.e. cholesterol lowering, anti-hypertensive, antiplatelets) (see the Example).

20

Proanthocyanidins designates a group of flavonoids that includes the subgroups procyanidins, prodelphinidins and propelargonidins. Proanthocyanidins are homogeneous or heterogeneous polymers consisting of the monomer units catechin or epicatechin, which are connected either by 4-8 or 4-6 linkages, to the effect that a great number of isomer proanthocyanidins exist.

25

Typically, the proanthocyanidins oligomers have a chain length of 2-12 monomer units. Proanthocyanidins may be synthesized or extracted from a plant material. Non-limiting examples of plant material sources of proanthocyanidins include grape seeds, grape skin, pine barks, ginkgo leaves, peanuts, cocoa beans, tamarind, tomato, peanut, almond, apple, cranberry, blueberry, tea leaves.

30

A well-known product containing proanthocyanidins, which is available in trade as a preparation of a food supplement under the name Pycnogenol®, is an extract of the French maritime pine bark (*Pinus pinaster*), see also US patents 3,436,407 (MASQUELIER

JACQUES); US 5,720,956 (ROHDEWALD, PETER) and US 6,372,266 (SUZUKI NOBUTAKA et al. Horphag Research Ltd.) which are incorporated herein by reference.

Pycnogenol® is a standardized bark extract of the French maritime pine *Pinus pinaster*, Aiton, subspecies *Atlantica* des Villar. The quality of this extract is specified in the United States

- 5 Pharmacopcia (USP 28) (Maritime Pine Extract. In: United States Pharmacopcia. Rockville: United States Pharmacopeial Convention, Inc.; 2005. pp 2115- 2116). The extract consists of a concentrate of polyphenols, which are also contained in fruits and vegetables, but, in low concentrations. The polyphenols are composed from flavonoids, especially procyanidins, and phenolic acids. All these constituents possess the ability to inactivate free radicals.
- 10 Rohdewald P. A review of the French maritime pine bark extract (Pycnogenol®), a herbal medication with a diverse pharmacology. *Int J Clin Pharmacol Ther* 2002;40(4): 158-168. Between 65-75% of Pycnogenol® are procyanidins comprising of catechin and epicatechin subunits with varying chain lengths (Rohdewald P. A review of the French maritime pine bark extract (Pycnogenol®), an herbal medication with a diverse clinical pharmacology. *Int J Clin Pharmacol Ther* 2002; 40: 158-168). Other constituents are polyphenolic monomers,
- 15 phenolic or cinnamic acids and their glycosides (Id.). Pycnogenol® extract is standardized to contain between 65% and 75% procyanidins (70+/- 5% procyanidins) in compliance with USP 28, compounds known for relatively significant antioxidant and anti-inflammatory activity, among other actions (Rohdewald P. "Pycnogenol®, French Maritime Pine Bark extract", *Encyclopedia of Dietary Supplements*, 2005, pp 545-553).
- 20

Centella asiatica is a small herbaceous annual plant used in a number of cosmetics application and health remedies. It is a member of the Mackinlayaceae family (Apiaceae) or Mackinlayoideae subfamily, and is native of Sri Lanka, northern Australia, Indonesia, Iran,

- 25 Malaysia, Melanesia, New Guinea, and other parts of Asia. It is also commonly named TECA, gotu kola, Asiatic Pennywort, Indian Pennywort, Luei Gong Gen, Takip-kohol, Antanan, Pegagan, Pegaga, vallaarai, Kula kud, Bai Bua Bok, Brahmi and rau má. *Centella asiatica* is also currently used in its countries of origin in culinary preparations as accompaniment or as a drink.

30

Centella asiatica extract consists of the leaves and roots of the hydrocotyle *centella asiatica*. The main active ingredients contained in *centella asiatica* are triterpenoids asiatic acid, madecassic acid, asiaticoside (triterpenoid ester glycosides), madecassoside, and brahminoside. A volatile oil comprising p-cymol, b-caryophyllene and farnesene is also found

in the plant. Extracts of *Centella asiatica* are also commercially available. Alternatively, *Centella asiatica* extracts can be prepared from the plant *Centella asiatica* by extraction techniques such as are well known in the art, including extractions in hydrophobic and hydrophilic solvents.

5

The present invention provides for a preparation consisting of the combination of proanthocyanidins (preferably procyanidins) and *Centella asiatica* and / or extracts thereof, in a suitable excipient, for use in the prevention or treatment of atherosclerosis. The preparation of the present invention contains from 20 % to 80 % w/w of proanthocyanidins (preferably

10

procyanidins) and from 20 % to 80 % w/w respectively of *Centella asiatica* and / or extracts thereof and a suitable excipient q.s.p. (quantity sufficient per 100% of the total volume). Preferably, the preparation of the invention comprises about 30 % to 70 % w/w of

15

proanthocyanidins (preferably procyanidins) and from 30 % to 70 % w/w respectively of *Centella asiatica* and / or extracts thereof, more preferably about 40% to 60% w/w of

proanthocyanidins (preferably procyanidins) and respectively 40% to 60% w/w of *Centella asiatica* and even more preferably about 50% w/w of proanthocyanidins (preferably procyanidins) and respectively 50% w/w of *Centella asiatica* and a suitable excipient q.s.p. According to the example, the preparation of the invention comprises the combination of Pycnogenol® and TECA (with or without Aspirin) (tablets, 100 mg for each component).

20

Pycnogenol® extract is standardized to contain between 65% and 75% of proanthocyanidins, preferably procyanidins (70+/- 5% procyanidins). Thus tablets used in the example contain 100 mg of TECA and between 65 mg and 75mg of proanthocyanidins, preferably 65 mg and 75mg of procyanidins.

25

The preparation of the invention comprises proanthocyanidins preferably consisting of procyanidins originated from a plant extract or from a synthesized material (i.e., synthetic proanthocyanidins).

30

The plant extract can be selected from the group consisting of proanthocyanidins containing extracts selected among extracts of pine bark, the cones of cypresses grape seed, apples, peanut skin, walnuts, pomegranates, tomatoes, almonds, tea, hawthorn, cocoa or combination thereof. Proanthocyanidins containing rich extracts are natural and preferably plant extracts having more than 50% by weight (of dried extracts) of proanthocyanidins, more preferably more than 70% by weight and even more preferably more than 75% by weight of

proanthocyanidins (preferably procyanidins). Preferably the plant extract according to the present invention is originated from pine bark and more preferably the plant extract is Pycnogenol®.

- 5 In a preferred embodiment, the preparation comprising proanthocyanidins may be a pine bark extract. The pine bark may be from *P. pinaster*, such as, for example, from Pycnogenol®. In a preferred embodiment, the composition may contain proanthocyanidins at a concentration of 10% to 100% of total weight. For example, a Pycnogenol® composition may be diluted or concentrated to contain 10%, 20%, 30%, 40%, 50%, 60%, 70%, 75%, 80%, 90% or 95%
- 10 proanthocyanidins (preferably procyanidins). Concentration may be performed using known methods such as column chromatography or affinity chromatography.

In a preferred embodiment, *Centella asiatica* extracts, present in the preparation of the invention, are the total triterpenic fraction of *Centella asiatica* (TECA).

15

- Examples of suitable excipients of this invention include, but are not limited to, anti-adherents, binders (e.g., microcrystalline cellulose, gum tragacanth, or gelatin), coatings, disintegrants, fillers, diluents, softeners, emulsifiers, flavoring agents, coloring agents, adjuvants, lubricants, functional agents (e.g., nutrients), viscosity modifiers, bulking agents,
- 20 glidants (e.g., colloidal silicon dioxide) surface active agents, osmotic agents, diluents, or any other non-active ingredient, or combinations thereof.

- For example, the preparation of the present invention may include excipient materials selected from the group consisting of calcium carbonate, coloring agents, whiteners, preservatives, and
- 25 flavors, triacetin, magnesium stearate, sterotes, natural or artificial flavors, essential oils, plant extracts, fruit essences, gelatins, or combinations thereof.

- Optionally the preparation of the present invention may include other artificial or natural sweeteners, bulk sweeteners, or combinations thereof. Bulk sweeteners include both caloric
- 30 and non-caloric compounds. Non-limiting examples of bulk sweeteners include sucrose, dextrose, maltose, dextrin, dried invert sugar, fructose, high fructose corn syrup, levulose, galactose, corn syrup solids, tagatose, polyols (e.g., sorbitol, mannitol, xylitol, lactitol, erythritol, and maltitol), hydrogenated starch hydrolysates, isomalt, trehalose, and combinations thereof.

The suitable excipient can be also a pharmaceutically acceptable excipient.

Optionally the preparation of the present invention can further comprise a non-steroidal anti-inflammatory drug such as acetylsalicylic acid (aspirin).

The present invention further provides for a food preparation, a dietary or food supplement, a nutraceutical, a beverage, a medicament and a topical preparation comprising the preparation of the present invention. As described above, the medicament and the topical preparation may further comprise acetylsalicylic acid (aspirin) as well as a pharmaceutically acceptable excipient.

Preferably, the dietary supplement, the nutraceutical or the medicament of the present invention is administered at a dosage of between 5 mg per day to 2'000 mg per day. Preferably between 50 mg to 1'000 mg per day and even more preferably between 100 mg to 400 mg per day. The dietary supplement, the nutraceutical or the medicament of the present invention contains from 20 % to 80 % w/w of proanthocyanidins (preferably procyanidins) and from 20 % to 80 % w/w respectively of *Centella asiatica* and / or extracts thereof and a suitable excipient q.s.p. It is known that Pycnogenol® extract is standardized to contain between 65% and 75% of proanthocyanidins, preferably procyanidins. Therefore, tablets used in the example do contain 100 mg of TECA and between 65 mg and 75mg of proanthocyanidins, preferably 65 mg and 75mg of procyanidins, more preferably 70 mg of proanthocyanidins or procyanidins,

The present invention further provides the preparation of the invention for use in the treatment or the prevention of atherosclerosis which concerns especially arteries. Preferably said treatment or prevention concerns atherosclerotic plaques.

The present invention also provides for a method of treating or preventing atherosclerosis disorders comprising administering to a subject in need thereof an effective amount of the preparation or of the medicament of the present invention. Said atherosclerosis disorders are preferably atherosclerotic plaques and said subject is a mammal, preferably a human.

The preparation, the dietary supplement, the nutraceutical or the medicament of the present invention can be administered orally, parenterally or topically at a dosage of between 5 mg per day to 2'000 mg per day. Preferably between 50 mg to 1'000 mg per day and more preferably between 100 mg to 400 mg per day. The concentrations of the active ingredients in
5 the preparation are described above.

If intended for oral administration, the medicament of the present invention can be in the form, for example, of a tablet, a caplet, a pill, a hard or soft capsule, a lozenge, a cachet, a dispensable powder, granules, a suspension, an elixir, a dispersion, a liquid, or any other form
10 reasonably adapted for such administration. If intended for parenteral administration, it can be in the form, for example, of a solution for intravenous, intramuscular or subcutaneous injection.

The topical preparations according to the present invention can be, but not limited to, a cream,
15 a patch, a gel, an ointment, a lotion, a tincture, a spray, a mousse, a cleansing composition or a foam. The topical preparations of the present invention can be also in the form of a suspension or dispersion in solvents or fatty substances, or alternatively in the form of an emulsion or micro emulsion, PET-emulsions, multiple emulsions, bickering emulsions, hydrogels, alcoholic gels, lipogels, one or multiphase solutions or a vesicular dispersion and
20 other usual compositions, which can also be applied by pens, as masks or as sprays. The emulsions can also contain anionic, nonionic, cationic or amphoteric surfactant(s).

In a preferred embodiment, the preparation according to the invention comprises proanthocyanidins (preferably procyanidins) and *Centella asiatica* and / or extracts thereof as
25 the sole active ingredients administered to a subject.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications without departing from the spirit
30 or essential characteristics thereof. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations or any two or more of said steps or features. The present disclosure is therefore to be considered as in all aspects illustrated and not restrictive,

the scope of the invention being indicated by the appended Claims, and all changes which come within the meaning and range of equivalency are intended to be embraced therein.

Various references are cited throughout this specification, each of which is incorporated
5 herein by reference in its entirety.

The foregoing description will be more fully understood with reference to the following Examples. Such Examples, are, however, exemplary of methods of practising the present invention and are not intended to limit the scope of the invention.

10

EXAMPLES

**Variation in echogenicity of echolucent carotid and femoral atherosclerotic plaques with combined treatment (Pycnogenol 100 mg + TECA 100 mg, with or without Aspirin 100
15 mg). A prospective, pilot 6-month trial.**
(TECA = total triterpenic fraction of *Centella asiatica*)

This study has evaluated whether a combination of Pycnogenol® and TECA, with or without Aspirin, was effective in reducing plaque progression and cardiovascular risk of events by
20 decreasing platelet aggregation, local arterial inflammation and the intima-media scarring process due to inflammation at the level of carotid and femoral arteries bifurcations.

The reduction in risk of cardiovascular events should have been obtained – by the study hypothesis - by modulating collagen production in plaques mainly by producing a visible
25 increase (ultrasound) in echogenicity in echolucent (soft, black on ultrasound, prone to embolisation and thrombosis causing events) femoral and carotid plaques.

This prospective placebo-controlled trial had the purpose of evaluating the effects of Pycnogenol® and TECA, with or without Aspirin, selectivity on hypoechoic-echolucent
30 plaques.

The sonographic examination of carotid plaques was made with high-resolution ultrasound.

Capturing plaques images, digital image processing, and normalization were standardized. The interobserver, intrascanner, gain-level variability were also standardized using as reference blood (black) for the most echolucent parts of the plaque and the adventitia (white) as the most echogenic part. The technique for plaque characterization and GSM (Grey Scale Median) measurements had been previously described in details.

In previous studies, after the ultrasonic plaque area had been obtained, normalization of echo texture had been obtained and plaque characterization had been used to differentiate echo-texture of plaque associated with cardiovascular events and those that did not cause embolization, thrombosis, or cardiovascular events (i.e. TIAs, strokes, embolisation, thrombosis).

In this study, after identifying plaques at higher risk (black, “soft”, echolucent plaques), patients were treated with the combination Pycnogenol® and TECA (with or without Aspirin) (tablets, 100 mg for each component) once daily for 6 months to evaluate whether this aggregate compound, by its antiplatelet action, anti-inflammatory effects and by modulating local collagen synthesis, could increase the echogenicity and therefore the stability of echolucent plaques. No other treatment was used during the 6 months follow-up.

Asymptomatic patients with carotid/femoral bifurcation, irregular, echolucent plaques (GSM<18) causing >65% stenosis (on duplex) were treated with Pycnogenol® and TECA (with or without Aspirin) or with comparable placebo after informed consent.

All patients (including controls) were treated with antiplatelet agents (in the group of interest the antiplatelet action was obtained by the combined activity of Aspirin and Pycnogenol®). This association may have a combined anti-aggregating effect possibly comparable to 400 mg of Aspirin (which could be not tolerated in most patients). The details of groups are shown in Table 1 and Table 2.

RESULTS: At inclusion the GSM in the hypoechoic plaques treatment group was 15.1 (range, 11-17.4); at 6 months GSM was increased (the increase was significant as GSM raised to 19.5;SD 1.1; P<0.05). There was no change in GSM in the comparable controls.

In conclusion the observed results suggest an important positive action of the combination of Pycnogenol® + TECA and the combination Pycnogenol® + TECA + Aspirin (APT) on the stabilization of hypoechoic, low-density (soft) carotid and femoral plaques which are the most dangerous types of plaques, associated to cardiovascular events due to embolisation of thrombosis.

The preparation of the present invention - of completely natural origin - stops the progression of atherosclerosis from subclinical to clinical stages in several patients. The synergy action of Pycnogenol® and TECA (with or without Aspirin) seems to be separated by any other pharmacological action (i.e. cholesterol lowering, anti-hypertensive, antiplatelets).

TABLE 1

	AGE	Number of patients	Males	Plaques	
				Femoral	Carotid
APT PATIENTS	63.3;7	57	38	66	69
CONTROLS	62.4;5.1	58	34	57	66

TABLE 2:

15 GSM Measurement

Treatment	GSM at baseline	At 6 months*
Pycnogenol 100 mg/day + Centella 100mg/day	15.2; SD 2.3	17/20**
Pycnogenol 100 mg/day + Centella 100 mg/day + Aspirin 100 mg/day	15.1; SD 2.1	19/20**
Pycnogenol® Group 100 mg/day	15.0; SD 2.9	3/18**
Centella Group 100 mg/day	14.9; SD 3.0	8/18**
Aspirin Group	15.3; SD 3.1	2/18

Untreated Group (Control)	15.2; SD 2.8	0/18

** Indicates significant change both from initial value and in comparison with control

* Number of patients with an increase of plaques Grey Scale Median of at least 5 points.

CLAIMS

1. A preparation consisting of the combination of proanthocyanidins containing rich extracts
5 and *Centella asiatica* and / or extracts thereof, in a suitable excipient, wherein said preparation contains from 30 % to 70 % w/w of proanthocyanidins containing rich extracts and from 30 % to 70 % w/w respectively of *Centella asiatica* and / or extracts thereof, for use in the prevention or treatment of atherosclerosis plaques in arteries.
- 10 2. A preparation for use according to claim 1, wherein said preparation contains 50 % w/w of proanthocyanidins containing rich extracts and from 50 % w/w respectively of *Centella asiatica* and / or extracts thereof.
3. The preparation for use according to any of claims 1 to 2, wherein proanthocyanidins
15 containing rich extracts are procyanidins originated from a plant extract or from a synthesized material.
4. The preparation for use according to claim 3, wherein the plant extract is selected from the group consisting of proanthocyanidins containing extracts selected among extracts of pine
20 bark, grape seed, apples, peanut skin, walnuts, pomegranates, tomatoes, almonds, tea, hawthorn, cocoa or combination thereof.
5. The preparation for use according to any of claims 3 to 4, wherein the plant extract is originated from French maritime pine bark (*Pinus pinaster*).
25
6. The preparation for use according to any of claims 1 to 5, wherein *Centella asiatica* extracts are the total triterpenic fraction of *Centella asiatica* (TECA).
7. The preparation for use according to any of claims 1 to 6, wherein said preparation further
30 comprises acetylsalicylic acid.
8. The preparation for use according to any of claims 1 to 7, wherein the suitable excipient is a pharmaceutically acceptable excipient.

9. A food preparation comprising the preparation for use according to any of claims 1 to 8.

10. A dietary supplement comprising the preparation for use according to any of claims 1 to
5 8.

11. A nutraceutical comprising the preparation for use according to any of claims 1 to 8.

12. A beverage comprising the preparation for use according to any of claims 1 to 8.

10

13. A medicament comprising the preparation for use according to any of claims 1 to 8.

14. A topical preparation comprising the preparation for use according to any of claims 1 to
8.

15