

MULTISUBSTITUTED AROMATIC COMPOUNDS AS SERINE PROTEASE INHIBITORS

BACKGROUND OF THE INVENTION

[0001] The present disclosure relates to compounds, e.g., multisubstituted aromatic compounds, which exhibit biological activity, e.g., inhibitory action, against serine proteases, including thrombin and various kallikreins.

[0002] Kallikreins are a subgroup of serine proteases, divided into plasma kallikrein and tissue kallikreins. Plasma kallikrein (KLKB1) liberates kinins (bradykinin and kallidin) from the kininogens, peptides responsible for the regulation of blood pressure and activation of inflammation. In the contact activation pathway of the coagulation cascade, plasma kallikrein assists in the conversion of factor XII to factor XIIa (Keel, M.; Trentz, O. *Injury* 2005, 36, 691–709). Factor XIIa converts FXI into FXIa, which in turn activates FIX, which with its co-factor FVIIIa forms the tenase complex, which finally activates FX to FXa. In the fibrinolysis part of the coagulation cascade, plasma kallikrein serves to convert plasminogen to plasmin. Thus, it has been proposed that plasma kallikrein inhibitors can be useful in the treatment of thrombotic and fibrinolytic diseases and disease conditions (US Patent # 7,625,944; Bird *et al. Thrombosis and Hemostasis* 2012, 107, 1141).

[0003] In rodent models, it has been shown that *activation* of plasma kallikrein in the eye increases retinal vascular permeability; whereas *inhibition* of the kallikrein-kinin system reduces retinal leakage induced by diabetes and hypertension. These findings suggest that intraocular activation of the plasma kallikrein pathway can contribute to excessive retinal vascular permeability that can lead to diabetic macular edema (DME). Thus, evidence suggests that plasma kallikrein inhibitors can provide a new therapeutic opportunity to reduce retinal vascular permeability (Feener, E. P. *Curr Diab Rep* 2010, 10, 270).

[0004] The Kallikrein-kinin system is involved in the regulation of vascular endothelial growth factor (VEGF), endothelial NO synthase, and fibroblast growth factor 2, all of which are involved in angiogenesis (Bader M. 2009, *Arteriosclerosis, Thrombosis, and Vascular Biology*, 29: 617). Tissue kallikrein (KLK1) has been linked to blood vessel growth (Miura S., 2003, *Hypertension*, 41, 1118). Therapies that moderate angiogenesis have been proposed for the treatment of both diabetic macular edema (DME) and age-related macular degeneration (AMD) (Syed, B.A.; Evans, J.B.; Bielory, L., 2012, *Nature Reviews Drug*

33 *Discovery*, 11, 827). Without further wishing to be bound by any theory, it is therefore
34 reasonable to conclude that KLK1 inhibitors can be useful in the treatment of diabetic
35 retinopathy, DME, and AMD.

36 [0005] Studies have shown that inflammation plays an important role in the origin and
37 development of AMD, and treatment often includes anti-inflammatories such as
38 corticosteroid (Telander, D., 2011, *Seminars in Ophthalmology*, 26(3), 192). The connection
39 between the kallikrein-kinin system and inflammation is also well established (Duchene,
40 2011, "Kallikrein-kinin kystem in inflammatory diseases". *Kinins*. De Gruyter. 261). Without
41 further wishing to be bound by any theory, it is reasonable to conclude that the anti-
42 inflammatory nature of kallikrein (e.g. KLK1 and KLKB1) inhibitors can be useful in the
43 treatment of AMD.

44 [0006] Ecallantide (Kalbitor) is a 60-amino acid recombinant protein that acts as a potent
45 reversible inhibitor of plasma kallikrein (Schneider L, *et al.*, *J Allergy Clin Immunol* 2007,
46 120, 416). Ecallantide has been approved by the FDA for the treatment of acute attacks of
47 hereditary angioedema (HAE). Without further wishing to be bound by any theory, it is
48 reasonable to believe that plasma kallikrein inhibition in general can be a useful treatment for
49 HAE, and thus there is strong interest in the development of plasma kallikrein inhibitors as a
50 therapy for HAE.

51 [0007] Tissue kallikreins (KLKs, for example, KLK1) are subdivided into various types,
52 and have been extensively investigated in cancer and inflammation biology. Various
53 kallikrein KLKs have been found to be up- or down-regulated in various cancer types, such
54 as cervical-, testicular-, and non-small-cell lung adenocarcinoma (Caliendo *et al.* *J. Med.*
55 *Chem.*, 2012, 55, 6669). Furthermore, overexpression of various KLKs in the skin has led to
56 the recognition that certain kallikrein inhibitors can be useful for certain dermatological
57 conditions, including atopic dermatitis, psoriasis and rare skin diseases such as Netherton
58 Syndrome (Freitas *et al.* *Bioorganic & Medicinal Chemistry Letters* 2012, 22, 6072–6075). A
59 thorough discussion of tissue kallikrein, plasma kallikreins, their functions and potential roles
60 in various diseases can be found in a variety of references, including the following which are
61 incorporated herein by reference in their entireties and for all purposes: Renné, T.; Gruber, A.
62 *Thromb Haemost* 2012, 107, 1012-3; Sotiropoulou, G.; Pampalakis, G. *Trends in*
63 *Pharmacological Sciences* 2012, 33, 623-634; Pampalakis, G.; Sotiropoulou, G. *Chapter 9*
64 *Pharmacological Targeting of Human Tissue Kallikrein-Related Peptidases. In Proteinases*
65 *as Drug Targets*, Dunn, B., Ed. *The Royal Society of Chemistry*: 2012; pp 199-228; Caliendo,

G.; Santagada, V.; Perissutti, E.; Severino, B.; Fiorino, F.; Frecentese, F.; Juliano, L. *J Med Chem* 2012, 55, 6669-86.


[0008] In mammalian systems, blood vessel injuries result in bleeding events, which are dealt with by the blood coagulation cascade. The cascade includes the extrinsic and intrinsic pathways, involving the activation of at least 13 interconnected factors and a variety of co-factors and other regulatory proteins. Upon vascular injury, plasma factor VII interacts with exposed Tissue Factor (TF), and the resultant TF-fVIIa complex initiates a complex series of events. Factor fXa is produced directly 'downstream' from the TF-fVIIa complex, and amplified manifold via the intrinsic Pathway. FXa then serves as the catalyst for formation of thrombin (fIIa), which in turn is the direct precursor to fibrinolysis. The outcome is a fibrinolytic clot, which stops the bleeding. Fibrinolysis of the polymeric clot into fibrin monomers leads to dissolution and a return of the system to the pre-clot state. The cascade is a complex balance of factors and co-factors and is tightly regulated. In disease states, undesired up- or down-regulation of any factor leads to conditions such as bleeding or thrombosis. Historically, anticoagulants have been used in patients at risk of suffering from thrombotic complications, such as angina, stroke and heart attack. Warfarin has enjoyed dominance as a first-in-line anticoagulant therapeutic. Developed in the 1940s, it is a Vitamin K antagonist and inhibits factors II, VII, IX and X, amongst others. It is administered orally, but its ease of use is tempered by other effects: it has a very long half-life (>2 days) and has serious drug-drug interactions. Importantly, since Vitamin K is a ubiquitous cofactor within the coagulation cascade, antagonism results in the simultaneous inhibition of many clotting factors and thus can lead to significant bleeding complications.

[0009] Much attention has been focused on heparin, the naturally-occurring polysaccharide that activates AT III, the endogenous inhibitor of many of the factors in the coagulation cascade. The need for parenteral administration for the heparin-derived therapeutics, and the inconvenient requirements for close supervision for the orally available warfarin, has resulted in a drive to discover and develop orally available drugs with wide therapeutic windows for safety and efficacy. Indeed, the position of thrombin in the coagulation cascade has made it a popular target for drug discovery. Without wishing to be bound by any theory, it is believed that the ultimate development of direct thrombin inhibitors (DTIs) is usefully based upon the classical D-Phe-Pro-Arg motif, a sequence that mimics fibrinogen, which is a natural substrate of thrombin. Without further wishing to be bound by any theory, it is believed that the use of DTIs is very well precedented, such as with the hirudin-based anticoagulants, and thus there is strong interest in the discovery and development of novel DTIs.

[0010] A thorough discussion of thrombin and its roles in the coagulation process can be found in a variety of references, including the following which are incorporated herein by reference in their entireties and for all purposes: Wieland, H. A., *et al.*, 2003, *Curr Opin Investig Drugs*, **4**:264-71; Gross, P. L. & Weitz, J. I., 2008, *Arterioscler Thromb Vasc Biol*, **28**:380-6; Hirsh, J., *et al.*, 2005, *Blood*, **105**:453-63; Prezelj, A., *et al.*, 2007, *Curr Pharm Des*, **13**:287-312.

BRIEF SUMMARY OF THE INVENTION


[0011] Embodiments of the invention encompass methods for treating or preventing a kallikrein-related disease or disorder in a subject, the methods including administering a compound of Formula (Ia):



or a pharmaceutical composition including the compound and a pharmaceutically acceptable excipient, to a subject in need thereof in an amount effective to treat or prevent said disease or disorder, wherein: Ring A can be substituted or unsubstituted pyrazolyl, or substituted or unsubstituted triazolyl; L^1 , L^2 , and L^4 can be independently absent, a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-; L^3 can be a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-; R^1 , R^2 , and R^4 can be independently absent, hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl, provided that R^1 can be absent when L^1 can be absent, R^2 can be absent when L^2 can be absent, and R^4 can be absent when L^4 can be absent; R^3 can be hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl, provided that R^2 can be absent when L^2 can be absent, R^3 can be

absent when L^3 can be absent, and R^4 can be absent when L^4 can be absent; and R^7 can be hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl. In some embodiments of the methods L^4 and R^4 can be absent.


[0012] In some embodiments, the compound can have the structure of Formula (IIa):



In some embodiments, L^3 can be a bond, and R^3 can be substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl. In some embodiments, the heteroaryl can be pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl. In some embodiments, R^3 can be substituted or unsubstituted phenyl, substituted or unsubstituted morpholinyl, substituted or unsubstituted oxanyl, substituted or unsubstituted oxetanyl, substituted or unsubstituted naphthyl or substituted or unsubstituted benzodioxinyl. In some embodiments, L^3 can be a bond or substituted or unsubstituted alkylene, and R^3 can be substituted or unsubstituted aryl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, L^3 can be $-C(O)O-$, and R^3 can be substituted or unsubstituted alkyl. In some embodiments, L^3 can be $-C(O)NR^5-$, R^5 can be hydrogen or alkyl, and R^3 can be substituted or unsubstituted alkyl, or substituted or unsubstituted aryl. In some embodiments, L^1 can be $-S-$, $-NR^4-$, substituted or unsubstituted alkylene, or substituted or unsubstituted heteroalkylene, and R^1 can be hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, the heteroaryl can be pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl. In some embodiments, R^1 can be chloro-substituted thienyl. In some embodiments, R^1 can be substituted or unsubstituted phenyl, substituted or unsubstituted morpholinyl, substituted or unsubstituted oxanyl, substituted or unsubstituted oxetanyl, substituted or unsubstituted naphthyl or substituted or unsubstituted benzodioxinyl. In some embodiments, L^2 and R^2 can be absent. In some embodiments, L^3 and R^3 can be absent. In some embodiments, L^2 can be substituted or unsubstituted alkylene or $-C(O)-$, and R^2 can be hydrogen, substituted or unsubstituted alkyl, substituted or

unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, the heteroaryl can be pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl. In some embodiments, R^2 can be substituted or unsubstituted phenyl, substituted or unsubstituted morpholinyl, substituted or unsubstituted oxanyl, substituted or unsubstituted oxetanyl, substituted or unsubstituted naphthyl or substituted or unsubstituted benzodioxinyl.


[0013] In some embodiments, the compound can have the structure of Formula (III):



In some embodiments, L^3 can be a bond, or substituted or unsubstituted alkylene, and R^3 can be substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocycloalkyl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, R^3 can be phenyl, naphthyl, pyridyl, pyridazinyl, pyrimidinyl, thienyl, furyl, morpholinyl, oxanyl, oxetanyl, or benzodioxinyl. In some embodiments, L^3 can be $-C(O)O-$, and R^3 can be substituted or unsubstituted alkyl. In some embodiments, L^3 can be $-C(O)NR^7-$, R^7 can be hydrogen or alkyl, and R^3 can be substituted or unsubstituted alkyl, or substituted or unsubstituted aryl. In some embodiments, L^1 can be $-S-$, $-NR^7-$, substituted or unsubstituted alkylene, or substituted or unsubstituted heteroalkylene, R^7 can be a hydrogen or alkyl, and R^1 can be hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, the heteroaryl can be pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl. In some embodiments, R^1 can be chloro-substituted thienyl. In some embodiments, R^1 can be substituted or unsubstituted phenyl, substituted or unsubstituted morpholinyl, substituted or unsubstituted oxanyl, substituted or unsubstituted oxetanyl, substituted or unsubstituted naphthyl or substituted or unsubstituted benzodioxinyl. In some embodiments, L^2 can be a bond and R^2 can be hydrogen. In some embodiments, L^2 can be substituted or unsubstituted alkylene or $-C(O)-$, and R^2 can be hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, the heteroaryl can be pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl. In some embodiments, R^2 can be substituted or unsubstituted phenyl, substituted or

unsubstituted morpholinyl, substituted or unsubstituted oxanyl, substituted or unsubstituted oxetanyl, substituted or unsubstituted naphthyl or substituted or unsubstituted benzodioxinyl.

[0014] In some embodiments, the compound has the following structure of Formula (IV):




In some embodiments, L⁴ can be a bond, and R⁴ can be hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl. In some embodiments, R⁴ can be halogen. In some embodiments, R⁴ can be unsubstituted alkyl. In some embodiments, R⁴ can be phenyl, naphthyl, pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl.

[0015] In some embodiments, the compound can be selected from any of Tables B or C. In some embodiments, the kallikrein-related disorder can be a thrombotic disease, a fibrinolytic disease, a type of cancer, an inflammatory condition, or a dermatological condition. In some embodiments, the kallikrein-related disorder can be an ophthalmic disease. In some embodiments, the ophthalmic disease can be diabetic macular edema, age-related macular degeneration, or diabetic retinopathy. In some embodiments, the type of cancer can be cervical-, testicular-, or non-small-cell lung adenocarcinoma. In some embodiments, the inflammatory condition can be sepsis, inflammatory bowel disease, systemic inflammatory response syndrome, or rheumatoid arthritis. In some embodiments, the dermatological condition can be atopic dermatitis, psoriasis, or Netherton Syndrome. In some embodiments, the compound acts by inhibiting kallikrein. In some embodiments, the compound acts by inhibiting tissue kallikrein. In some embodiments, the compound acts by inhibiting plasma kallikrein. In some embodiments, the compound or pharmaceutical composition can be administered in the form of an ophthalmic composition applied topically to the eye. In some embodiments, the ophthalmic composition can be in the form of eye drops. In some embodiments, the compound or pharmaceutical composition can be administered in the form of an ophthalmic composition via intravitreal injection. Embodiments of the invention also encompass methods for treating or preventing a kallikrein-related disease or disorder in a subject, including administering a compound from Table D or a pharmaceutical composition

including compound and a pharmaceutically acceptable excipient, to a subject in need thereof in an amount effective to treat or prevent the disease or disorder.

[0016] Embodiments of the invention also encompass compounds with structure of Formula (V):



or pharmaceutically acceptable salt, ester, solvate, or prodrug thereof; wherein L^1 can be a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-; L^2 and L^5 can be independently absent, a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-; R^1 can be hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl; R^2 and R^5 can be independently absent, hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl; and R^7 can be hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl. In some embodiments, L^5 and R^5 can be absent. In some embodiments, L^2 and R^2 can be absent. In some embodiments, L^2 can be -C(O)-, and R^2 can be substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl. In some embodiments, R^2 can be substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl. In some embodiments, the heteroaryl can be pyridyl, pyridazinyl,

pyrimidinyl, thienyl, or furyl. In some embodiments, R² can be substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, the heterocycloalkyl can be oxanyl, oxetanyl, or morpholinyl. In some embodiments, the fused ring aryl can be benzodioxinyl or naphthyl. In some embodiments, L¹ can be bond, -S-, -NR⁷-, substituted or unsubstituted alkylene, or substituted or unsubstituted heteroalkylene, and R¹ can be hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, the heteroaryl can be pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl. In some embodiments, R¹ can be chloro-substituted thienyl. In some embodiments, the heterocycloalkyl can be morpholinyl, oxanyl, or oxetanyl. In some embodiments, the fused ring aryl can be benzodioxinyl or naphthyl. In some embodiments, L⁵ can be a bond or substituted or unsubstituted alkylene, and R⁵ can be substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl. In some embodiments, the heteroaryl can be pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl. In some embodiments, the fused ring aryl can be benzodioxinyl or naphthyl. In some embodiments, L⁵ can be substituted or unsubstituted alkylene, and R⁵ can be substituted or unsubstituted heterocycloalkyl. In some embodiments, the heterocycloalkyl can be morpholinyl, oxanyl, or oxetanyl.

[0017] Embodiments of the invention also encompass pharmaceutical compositions including such compounds, or a compound as set forth in Table A, and a pharmaceutically acceptable excipient. Embodiments of the invention also encompass methods for treating a disease or disorder in a subject, including administering such compounds or pharmaceutical compositions to a subject in need thereof in an amount effective to treat or prevent said disease or disorder. In some embodiments, the disease or disorder can be a thrombotic disorder. In some embodiments, the thrombotic disorder can be acute coronary syndrome, venous thromboembolism, arterial thromboembolism, cardiogenic thromboembolism, disseminated intravascular coagulation, or a blood clot thrombus. In some embodiments, the disease or disorder can be fibrosis. In some embodiments, the disease or disorder can be Alzheimer's Disease. In some embodiments, the disease or disorder can be multiple sclerosis. In some embodiments, the disease or disorder can be pain. In some embodiments, the disease or disorder can be cancer. In some embodiments, the compound acts by inhibiting thrombin. In some embodiments, the disease or disorder can be a kallikrein-related disorder. In some embodiments, the kallikrein-related disorder can be a thrombotic disease, a fibrinolytic

disease, a type of cancer, an inflammatory condition, or a dermatological condition. In some embodiments, the kallikrein-related disorder can be an ophthalmic disease. In some embodiments, the ophthalmic disease can be diabetic macular edema, age-related macular degeneration, or diabetic retinopathy. In some embodiments, the type of cancer can be cervical-, testicular-, or non-small-cell lung adenocarcinoma. In some embodiments, the inflammatory condition can be sepsis, inflammatory bowel disease, systemic inflammatory response syndrome, or rheumatoid arthritis. In some embodiments, the dermatological condition can be atopic dermatitis, psoriasis, or Netherton Syndrome. In some embodiments, the compound acts by inhibiting kallikrein. In some embodiments, the compound acts by inhibiting tissue kallikrein. In some embodiments, the compound acts by inhibiting plasma kallikrein. In some embodiments, the compound or pharmaceutical composition can be administered in the form of an ophthalmic composition applied topically to the eye. In some embodiments, the ophthalmic composition can be in the form of eye drops. In some embodiments, the compound or pharmaceutical composition can be administered in the form of an ophthalmic composition via intravitreal injection.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018] Not applicable.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

[0019] The abbreviations used herein have their conventional meaning within the chemical and biological arts. The chemical structures and formulae set forth herein are constructed according to the standard rules of chemical valency known in the chemical arts.

[0020] Where substituent groups are specified by their conventional chemical formulae, written from left to right, they equally encompass the chemically identical substituents that would result from writing the structure from right to left, e.g., $-\text{CH}_2\text{O}-$ is equivalent to $-\text{OCH}_2-$.

[0021] As used herein, the term “attached” signifies a stable covalent bond, certain preferred points of attachment being apparent to those of ordinary skill in the art.

[0022] The terms “halogen” or “halo” include fluorine, chlorine, bromine, and iodine. Additionally, terms such as “haloalkyl” are meant to include monohaloalkyl and

polyhaloalkyl. For example, the term “halo(C₁-C₄)alkyl” includes, but is not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0023] The term “alkyl,” by itself or as part of another substituent, means, unless otherwise stated, a straight (i.e., unbranched) or branched chain, or combination thereof, which can be fully saturated, mono- or polyunsaturated and can include di- and multivalent radicals, having the number of carbon atoms designated (i.e., C₁-C₁₀ means one to ten carbons). Examples of saturated hydrocarbon radicals include, but are not limited to, groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, (cyclohexyl)methyl, homologs and isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated alkyl group is one having one or more double bonds or triple bonds. Examples of unsaturated alkyl groups include, but are not limited to, vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. Accordingly, the term “alkyl” can refer to C₁-C₁₆ straight chain saturated, C₁-C₁₆ branched saturated, C₃-C₈ cyclic saturated and C₁-C₁₆ straight chain or branched saturated aliphatic hydrocarbon groups substituted with C₃-C₈ cyclic saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, this definition shall include but is not limited to methyl (Me), ethyl (Et), propyl (Pr), butyl (Bu), pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, isopropyl (i-Pr), isobutyl (i-Bu), tert-butyl (t-Bu), sec-butyl (s-Bu), isopentyl, neopentyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclopropylmethyl, and the like.

[0024] The term “alkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from an alkyl, as exemplified, but not limited by, -CH₂CH₂CH₂CH₂-. Typically, an alkyl (or alkylene) group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in the compounds disclosed herein. A “lower alkyl” or “lower alkylene” is a shorter chain alkyl or alkylene group, generally having eight or fewer carbon atoms.

[0025] The term “heteroalkyl,” by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or combinations thereof, consisting of at least one carbon atom and at least one heteroatom selected from the group consisting of O, N, P, Si, and S, and wherein the nitrogen and sulfur atoms can optionally be oxidized, and the nitrogen heteroatom can optionally be quaternized. The heteroatom(s) O, N, P, S, and Si can be placed at any interior position of the heteroalkyl group or at the position at which the alkyl group is attached to the remainder of the molecule. Examples include, but are not limited to:

-CH₂-CH₂-O-CH₃, -CH₂-CH₂-NH-CH₃, -CH₂-CH₂-N(CH₃)-CH₃, -CH₂-S-CH₂-CH₃,
-CH₂-CH₂, -S(O)-CH₃, -CH₂-CH₂-S(O)₂-CH₃, -CH=CH-O-CH₃, -Si(CH₃)₃,
-CH₂-CH=N-OCH₃, -CH=CH-N(CH₃)-CH₃, -O-CH₃, -O-CH₂-CH₃, and -CN. Up to two
heteroatoms can be consecutive, such as, for example, -CH₂-NH-OCH₃.

[0026] Similarly, the term “heteroalkylene,” by itself or as part of another substituent, means, unless otherwise stated, a divalent radical derived from heteroalkyl, as exemplified, but not limited by, -CH₂-CH₂-S-CH₂-CH₂- and -CH₂-S-CH₂-CH₂-NH-CH₂-. For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (e.g., alkyleneoxy, alkylenedioxy, alkyleneamino, alkylenediamino, and the like). Still further, for alkylene and heteroalkylene linking groups, no orientation of the linking group is implied by the direction in which the formula of the linking group is written. For example, the formula -C(O)₂R'- represents both -C(O)₂R'- and -R'C(O)₂-. As described above, heteroalkyl groups, as used herein, include those groups that are attached to the remainder of the molecule through a heteroatom, such as -C(O)R', -C(O)NR', -NR'R'', -OR', -SR', and/or -SO₂R'. Where “heteroalkyl” is recited, followed by recitations of specific heteroalkyl groups, such as -NR'R'' or the like, it will be understood that the terms heteroalkyl and -NR'R'' are not redundant or mutually exclusive. Rather, the specific heteroalkyl groups are recited to add clarity. Thus, the term “heteroalkyl” should not be interpreted herein as excluding specific heteroalkyl groups, such as -NR'R'' or the like.

[0027] The terms “cycloalkyl” and “heterocycloalkyl,” by themselves or in combination with other terms, mean, unless otherwise stated, cyclic versions of “alkyl” and “heteroalkyl,” respectively. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include, but are not limited to, 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like. A “cycloalkylene” and a “heterocycloalkylene,” alone or as part of another substituent, means a divalent radical derived from a cycloalkyl and heterocycloalkyl, respectively.

[0028] The term “alkenyl” includes C₂-C₁₆ straight chain unsaturated, C₂-C₁₁ branched unsaturated, C₅-C₈ unsaturated cyclic, and C₂-C₁₆ straight chain or branched unsaturated aliphatic hydrocarbon groups substituted with C₃-C₈ cyclic saturated and unsaturated aliphatic hydrocarbon groups having the specified number of carbon atoms. Double bonds

can occur in any stable point along the chain and the carbon-carbon double bonds can have either the *cis* or *trans* configuration. For example, this definition shall include but is not limited to ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, 1,5-octadienyl, 1,4,7-nonatrienyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, ethylcyclohexenyl, butenylcyclopentyl, 1-pentenyl-3-cyclohexenyl, and the like. Similarly, "heteroalkenyl" refers to heteroalkyl having one or more double bonds.

[0029] The term "alkynyl" refers in the customary sense to alkyl additionally having one or more triple bonds. The term "cycloalkenyl" refers to cycloalkyl additionally having one or more double bonds. The term "heterocycloalkenyl" refers to heterocycloalkyl additionally having one or more double bonds.

[0030] The term "acyl" means, unless otherwise stated, -C(O)R where R is a substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl.

[0031] Each of the above terms (e.g., "alkyl," "heteroalkyl," "aryl," and "heteroaryl") includes both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided herein.

[0032] Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be one or more of a variety of groups selected from, but not limited to, -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -CN, and -NO₂ in a number ranging from zero to (2m'+1), where m' is the total number of carbon atoms in such radical. R', R'', and R''' each preferably independently refer to hydrogen, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl (e.g., aryl substituted with 1-3 halogens), substituted or unsubstituted alkyl, alkoxy, or thioalkoxy groups, or arylalkyl groups. When a compound disclosed herein includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', and R''' group when more than one of these groups is present. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 4-, 5-, 6-, or 7-membered ring. For example, -NR'R'' includes, but is not limited to, 1-pyrrolidiny and 4-

morpholinyl. From the above discussion of substituents, one of skill in the art will understand that the term "alkyl" is meant to include groups including carbon atoms bound to groups other than hydrogen groups, such as haloalkyl (e.g., -CF₃ and -CH₂CF₃) and acyl (e.g., -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the like).

[0033] Similar to the substituents described for the alkyl radical, substituents for the aryl and heteroaryl groups are varied and are selected from, for example: -OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -CN, -NO₂, -R', -N₃, -CH(Ph)₂, fluoro(C₁-C₄)alkoxy, and fluoro(C₁-C₄)alkyl, in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'', and R''' are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl. When a compound disclosed herein includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', and R''' groups when more than one of these groups is present.

[0034] Two or more substituents can optionally be joined to form aryl, heteroaryl, cycloalkyl, or heterocycloalkyl groups. Such so-called ring-forming substituents are typically, though not necessarily, found attached to a cyclic base structure. In one embodiment, the ring-forming substituents are attached to adjacent members of the base structure. For example, two ring-forming substituents attached to adjacent members of a cyclic base structure create a fused ring structure. In another embodiment, the ring-forming substituents are attached to a single member of the base structure. For example, two ring-forming substituents attached to a single member of a cyclic base structure create a spirocyclic structure. In yet another embodiment, the ring-forming substituents are attached to non-adjacent members of the base structure.

[0035] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring can optionally form a ring of the formula -T-C(O)-(CRR')_q-U-, wherein T and U are independently -NR-, -O-, -CRR'-, or a single bond, and q is an integer of from 0 to 3. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring can optionally be replaced with a substituent of the formula -A-(CH₂)_r-B-, wherein A and B are independently -CRR'-, -O-, -NR-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'-, or a single bond, and r is an integer of from 1 to 4. One of the single bonds of the new ring so formed can optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the

aryl or heteroaryl ring can optionally be replaced with a substituent of the formula $-(CRR')_s-X'-(C''R''')_d-$, where s and d are independently integers of from 0 to 3, and X' is $-O-$, $-NR'-$, $-S-$, $-S(O)-$, $-S(O)_2-$, or $-S(O)_2NR'-$. The substituents R , R' , R'' , and R''' are preferably independently selected from hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

[0036] As used herein, the terms "heteroatom" or "ring heteroatom" are meant to include oxygen (O), nitrogen (N), sulfur (S), phosphorus (P), and silicon (Si).

[0037] The term "alkyloxy" (e.g. methoxy, ethoxy, propyloxy, allyloxy, cyclohexyloxy) represents an alkyl group as defined above having the indicated number of carbon atoms attached through an oxygen bridge ($-O-$).

[0038] The term "alkylthio" (e.g. methylthio, ethylthio, propylthio, cyclohexylthio and the like) represents an alkyl group as defined above having the indicated number of carbon atoms attached through a sulfur bridge ($-S-$).

[0039] The term "alkylamino" represents one or two alkyl groups as defined above having the indicated number of carbon atoms attached through an amine bridge. The two alkyl groups can be taken together with the nitrogen to which they are attached forming a cyclic system containing 3 to 8 carbon atoms with or without one C_1 - C_{16} alkyl, aryl C_0 - C_{16} alkyl, or C_0 - C_{16} alkylaryl substituent.

[0040] The term "alkylaminoalkyl" represents an alkylamino group attached through an alkyl group as defined above having the indicated number of carbon atoms.

[0041] The term "alkyloxy(alkyl)amino" (e.g. methoxy(methyl)amine, ethoxy(propyl)amine) represents an alkyloxy group as defined above attached through an amino group, the amino group itself having an alkyl substituent.

[0042] The term "alkylcarbonyl" (e.g. cyclooctylcarbonyl, pentylcarbonyl, 3-hexylcarbonyl) represents an alkyl group as defined above having the indicated number of carbon atoms attached through a carbonyl group.

[0043] The term "alkylcarboxy" (e.g. heptylcarboxy, cyclopropylcarboxy, 3-pentenylcarboxy) represents an alkylcarbonyl group as defined above wherein the carbonyl is in turn attached through an oxygen.

[0044] The term "alkylcarboxyalkyl" represents an alkylcarboxy group attached through an alkyl group as defined above having the indicated number of carbon atoms.

[0045] The term "alkylcarbonylamino" (e.g. hexylcarbonylamino, cyclopentylcarbonylaminomethyl, methylcarbonylaminophenyl) represents an alkylcarbonyl group as defined above wherein the carbonyl is in turn attached through the nitrogen atom of an amino group.

[0046] The nitrogen group can itself be substituted with an alkyl or aryl group.

[0047] The term "aryl" means, unless otherwise stated, a polyunsaturated, aromatic, hydrocarbon substituent, which can be a single ring or multiple rings (preferably from 1 to 3 rings) that are fused together (i.e., a fused ring aryl) or linked covalently. A fused ring aryl refers to multiple rings fused together wherein at least one of the fused rings is an aryl ring. The term "heteroaryl" refers to aryl groups (or rings) that contain from one to four heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. Thus, the term "heteroaryl" includes fused ring heteroaryl groups (i.e., multiple rings fused together wherein at least one of the fused rings is a heteroaromatic ring). A 5,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 5 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. Likewise, a 6,6-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 6 members, and wherein at least one ring is a heteroaryl ring. And a 6,5-fused ring heteroarylene refers to two rings fused together, wherein one ring has 6 members and the other ring has 5 members, and wherein at least one ring is a heteroaryl ring. A heteroaryl group can be attached to the remainder of the molecule through a carbon or heteroatom. Non-limiting examples of aryl and heteroaryl groups include phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxaliny, 5-quinoxaliny, 3-quinolyl, and 6-quinolyl. Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below. An "arylene" and a "heteroarylene," alone or as part of another substituent, mean a divalent radical derived from an aryl and heteroaryl, respectively. Accordingly, the term "aryl" can represent an unsubstituted, mono-, di- or trisubstituted monocyclic, polycyclic, biaryl and heterocyclic aromatic groups covalently attached at any ring position capable of forming a stable covalent bond, certain preferred points of attachment being apparent to those

skilled in the art (e. g. 3-indolyl, 4-imidazolyl). The aryl substituents are independently selected from the group consisting of halo, nitro, cyano, trihalomethyl, C₁₋₁₆alkyl, arylC₁₋₁₆alkyl, C₀₋₁₆alkyloxyC₀₋₁₆alkyl, arylC₀₋₁₆alkyloxyC₀₋₁₆alkyl, C₀₋₁₆alkylthioC₀₋₁₆alkyl, arylC₀₋₁₆alkylthioC₀₋₁₆alkyl, C₀₋₁₆alkylaminoC₀₋₁₆alkyl, arylC₀₋₁₆alkylaminoC₀₋₁₆alkyl, di(arylC₁₋₁₆alkyl)aminoC₀₋₁₆alkyl, C₁₋₁₆alkylcarbonylC₀₋₁₆alkyl, arylC₁₋₁₆alkylcarbonylC₀₋₁₆alkyl, C₁₋₁₆alkylcarboxyC₀₋₁₆alkyl, arylC₁₋₁₆alkylcarboxyC₀₋₁₆alkyl, C₁₋₁₆alkylcarbonylaminoC₀₋₁₆alkyl, arylC₁₋₁₆alkylcarbonylaminoC₀₋₁₆alkyl, -C₀₋₁₆alkylCOOR₄, -C₀₋₁₆alkylCONR₅R₆ wherein R₄, R₅ and R₆ are independently selected from hydrogen, C₁₋₁₆alkyl, arylC₀₋₁₆alkyl, or R₅ and R₆ are taken together with the nitrogen to which they are attached forming a cyclic system containing 3 to 8 carbon atoms with or without one C₁₋₁₆alkyl, arylC₀₋₁₆alkyl, or C₀₋₁₆alkylaryl substituent. Aryl includes but is not limited to pyrazolyl and triazolyl.

[0048] For brevity, the term “aryl” when used in combination with other terms (e.g., aryloxy, arylthioxy, arylalkyl) includes both aryl and heteroaryl rings as defined above. Thus, the terms “arylalkyl,” “aralkyl” and the like are meant to include those radicals in which an aryl group is attached to an alkyl group (e.g., benzyl, phenethyl, pyridylmethyl, and the like) including those alkyl groups in which a carbon atom (e.g., a methylene group) has been replaced by, for example, an oxygen atom (e.g., phenoxymethyl, 2-pyridyloxymethyl, 3-(1-naphthyloxy)propyl, and the like), or a sulfur atom. Accordingly, the terms “arylalkyl” and the like (e.g. (4-hydroxyphenyl)ethyl, (2-aminonaphthyl)hexyl, pyridylcyclopentyl) represents an aryl group as defined above attached through an alkyl group as defined above having the indicated number of carbon atoms.

[0049] The term “oxo,” as used herein, means an oxygen that is double bonded to a carbon atom.

[0050] The term “alkylsulfonyl,” as used herein, means a moiety having the formula -S(O₂)-R', where R' is an alkyl group as defined above. R' can have a specified number of carbons (e.g., “C₁₋₄ alkylsulfonyl”).

[0051] The term “carbonyloxy” represents a carbonyl group attached through an oxygen bridge.

[0052] In the above definitions, the terms “alkyl” and “alkenyl” can be used interchangeably in so far as a stable chemical entity is formed, as would be apparent to those skilled in the art.

[0053] The term “linker” refers to attachment groups interposed between substituents, e.g., R^1 , R^2 , R^3 or R^4 described herein, e.g., Formula (Ia) and generically referred to as R^n , and the group which is substituted, e.g., “ring A” group of e.g., Formula (Ia). In some embodiments, the linker includes amido ($-\text{CONH}-R^n$ or $-\text{NHCO}-R^n$), thioamido ($-\text{CSNH}-R^n$ or $-\text{NHCS}-R^n$), carboxyl ($-\text{CO}_2-R^n$ or $-\text{OCOR}^n$), carbonyl ($-\text{CO}-R^n$), urea ($-\text{NHCONH}-R^n$), thiourea ($-\text{NHCSNH}-R^n$), sulfonamido ($-\text{NHSO}_2-R^n$ or $-\text{SO}_2\text{NH}-R^n$), ether ($-\text{O}-R^n$), sulfonyl ($-\text{SO}_2-R^n$), sulfoxyl ($-\text{SO}-R^n$), carbamoyl ($-\text{NHCO}_2-R^n$ or $-\text{OCONH}-R^n$), or amino ($-\text{NHR}^n$) linking moieties.

[0054] A “substituent group,” as used herein, means a group selected from the following moieties:

(A) $-\text{OH}$, $-\text{NH}_2$, $-\text{SH}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{NO}_2$, oxo, halogen, $-\text{COOH}$, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(B) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, substituted with at least one substituent selected from:

(i) oxo, $-\text{OH}$, $-\text{NH}_2$, $-\text{SH}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{NO}_2$, halogen, $-\text{COOH}$, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(ii) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, substituted with at least one substituent selected from:

(a) oxo, $-\text{OH}$, $-\text{NH}_2$, $-\text{SH}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{NO}_2$, halogen, $-\text{COOH}$, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, unsubstituted heteroaryl, and

(b) alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, substituted with at least one substituent selected from: oxo, $-\text{OH}$, $-\text{NH}_2$, $-\text{SH}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{NO}_2$, halogen, $-\text{COOH}$, unsubstituted alkyl, unsubstituted heteroalkyl, unsubstituted cycloalkyl, unsubstituted heterocycloalkyl, unsubstituted aryl, and unsubstituted heteroaryl.

[0055] A “size-limited substituent” or “size-limited substituent group,” as used herein, means a group selected from all of the substituents described above for a “substituent group,” wherein each substituted or unsubstituted alkyl is a substituted or unsubstituted $\text{C}_1\text{-C}_{20}$ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2-20-membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted $\text{C}_4\text{-}$


C₈ cycloalkyl, and each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 4-8-membered heterocycloalkyl.

[0056] A “lower substituent” or “lower substituent group,” as used herein, means a group selected from all of the substituents described above for a “substituent group,” wherein each substituted or unsubstituted alkyl is a substituted or unsubstituted C₁-C₈ alkyl, each substituted or unsubstituted heteroalkyl is a substituted or unsubstituted 2-8-membered heteroalkyl, each substituted or unsubstituted cycloalkyl is a substituted or unsubstituted C₅-C₇ cycloalkyl, and each substituted or unsubstituted heterocycloalkyl is a substituted or unsubstituted 5-7-membered heterocycloalkyl.

[0057] The term “about” used in the context of a numeric value indicates a range of +/- 10% of the numeric value, unless expressly indicated otherwise.

II. Compounds

[0058] In one aspect, there is provided a compound with structure of Formula (Ia):




or pharmaceutically acceptable salt, ester, solvate, or prodrug thereof. Ring A is substituted or unsubstituted pyrazolyl, or substituted or unsubstituted triazolyl. L¹, L² and L³ are independently absent, a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-. L⁴ is absent, a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-. R¹, R², and R⁴ are independently absent, hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkenyl, or substituted or unsubstituted fused ring aryl. In some embodiments, R¹ can be absent provided L¹ is also absent. In some embodiments, R² can be absent provided L² is also absent. In some embodiments, R⁴ can be absent provided L⁴ is also absent. R³ is hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or


unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl. R^7 is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl.

[0059] In some embodiments, the compound is a pharmaceutically acceptable salt, ester, solvate, or prodrug of a compound of Formula (Ia). In some embodiments, the compound is not an ester, not a solvate, and not a prodrug.

[0060] In some embodiments, L^4 and R^4 are absent, providing a compound with structure of Formula (Ib) following.



[0061] In some embodiments, there is provided a compound according to Formula (Ib) with structure of Formula (IIa) following.



[0062] In some embodiments, the compound has the structure of Formula (IIa), wherein L^3 is a bond, and R^3 is substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heterocycloalkyl, or substituted or unsubstituted heteroaryl. In some embodiments, R^3 is unsubstituted aryl, preferably phenyl. In some embodiments, R^3 is substituted aryl, preferably halogen-substituted phenyl. In some embodiments, R^3 is substituted or unsubstituted phenyl, or substituted or unsubstituted thienyl. In some embodiments, R^3 is unsubstituted thienyl. In some embodiments, R^3 is a chloro-substituted thienyl. In some embodiments, R^3 is substituted or unsubstituted pyridyl, or substituted or

636 unsubstituted pyridazinyl. In some embodiments, R^3 is unsubstituted pyridyl. In some
637 embodiments, R^3 is unsubstituted pyridazinyl. In some embodiments, R^3 is substituted or
638 unsubstituted pyrimidinyl, or substituted or unsubstituted furyl. In some embodiments, R^3 is
639 unsubstituted pyrimidinyl. In some embodiments, R^3 is unsubstituted furyl. In some
640 embodiments, R^3 is substituted or unsubstituted morpholinyl, or substituted or unsubstituted
641 oxanyl, or substituted or unsubstituted oxetanyl. In some embodiments, R^3 is unsubstituted
642 morpholinyl. In some embodiments, R^3 is unsubstituted oxanyl. In some embodiments, R^3 is
643 unsubstituted oxetanyl. In some embodiments, R^3 is substituted or unsubstituted
644 benzodioxinyl, or substituted or unsubstituted naphthyl. In some embodiments, R^3 is
645 unsubstituted benzodioxinyl. In some embodiments, R^3 is unsubstituted naphthyl.

646 **[0063]** In some embodiments, a compound is provided with structure of Formula (IIa),
647 wherein L^3 is a bond, substituted or unsubstituted alkylene, and R^3 is substituted or
648 unsubstituted aryl, or substituted or unsubstituted heterocycloalkyl.

649 **[0064]** In some embodiments, the compound has the structure of Formula (IIa), wherein L^3
650 is $-C(O)O-$, and R^3 is substituted or unsubstituted alkyl, preferably unsubstituted alkyl, more
651 preferably unsubstituted lower alkyl.


652 **[0065]** In some embodiments, the compound has the structure of Formula (IIa), wherein L^3
653 is $-C(O)NR^5-$, R^5 is hydrogen or alkyl, and R^3 is substituted or unsubstituted alkyl, or
654 substituted or unsubstituted aryl.

655 **[0066]** Further to any embodiment above wherein the compound has the structure of
656 Formula (IIa), in some embodiments L^1 is $-S-$, $-NR^7-$, substituted or unsubstituted alkylene, or
657 substituted or unsubstituted heteroalkylene, where R^7 is as described in formula Ia, and R^1 is
658 hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or
659 unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or
660 unsubstituted heterocycloalkyl. In some embodiments, R^1 is substituted or unsubstituted
661 phenyl. In some embodiments R^1 is unsubstituted phenyl. In some embodiments, R^1 is a
662 substituted or unsubstituted pyridyl. In some embodiments, R^1 is a substituted or unsubstituted
663 pyridazinyl. In some embodiments, R^1 is a substituted or unsubstituted pyrimidinyl. In some
664 embodiments, R^1 is a substituted or unsubstituted thienyl. In some embodiments, R^1 is a
665 substituted or unsubstituted furyl. In some embodiments, R^1 is an unsubstituted pyridyl. In
666 some embodiments, R^1 is an unsubstituted pyridazinyl. In some embodiments, R^1 is an
667 unsubstituted pyrimidinyl. In some embodiments, R^1 is an unsubstituted thienyl. In some
668 embodiments, R^1 is a chloro-substituted thienyl. In some embodiments, R^1 is an unsubstituted

furyl. In some embodiments, R¹ is a substituted or unsubstituted morpholinyl. In some
embodiments, R¹ is a substituted or unsubstituted oxanyl. In some embodiments, R¹ is a
substituted or unsubstituted oxetanyl. In some embodiments, R¹ is an unsubstituted
morpholinyl. In some embodiments, R¹ is an unsubstituted oxanyl. In some embodiments, R¹
is an unsubstituted oxetanyl. In some embodiments, R¹ is substituted or unsubstituted
benzodioxinyl. In some embodiments, R¹ is substituted or unsubstituted naphthyl. In some
embodiments, R¹ is unsubstituted benzodioxinyl. In some embodiments, R¹ is unsubstituted
naphthyl. In some embodiments, R³ is substituted or unsubstituted aryl. In some
embodiments, R³ is unsubstituted aryl. In some embodiments, L³ and R³ are absent. In some
embodiments, L² and R² are absent. In some embodiments, L² is a bond. In some
embodiments, L² is a bond and R² is hydrogen.


[0067] Further to any embodiment above wherein the compound has the structure of
Formula (IIa), R² is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted
heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl,
substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl,
substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or
unsubstituted heteroaryl. In some embodiments, R² is substituted or unsubstituted phenyl. In
some embodiments, R² is unsubstituted phenyl. In some embodiments, R² is a substituted
or unsubstituted pyridyl. In some embodiments, R² is a substituted or unsubstituted
pyridazinyl. In some embodiments, R² is a substituted or unsubstituted pyrimidinyl. In some
embodiments, R² is a substituted or unsubstituted thienyl. In some embodiments, R² is a
substituted or unsubstituted furyl. In some embodiments, R² is an unsubstituted pyridyl. In
some embodiments, R² is an unsubstituted pyridazinyl. In some embodiments, R² is an
unsubstituted pyrimidinyl. In some embodiments, R² is an unsubstituted thienyl. In some
embodiments, R² is a chloro-substituted thienyl. In some embodiments, R² is an unsubstituted
furyl. In some embodiments, R² is a substituted or unsubstituted morpholinyl. In some
embodiments, R² is a substituted or unsubstituted oxanyl. In some embodiments, R² is a
substituted or unsubstituted oxetanyl. In some embodiments, R² is an unsubstituted
morpholinyl. In some embodiments, R² is an unsubstituted oxanyl. In some embodiments, R²
is an unsubstituted oxetanyl. In some embodiments, R² is substituted or unsubstituted
benzodioxinyl. In some embodiments, R² is substituted or unsubstituted naphthyl. In some
embodiments, R² is unsubstituted benzodioxinyl. In some embodiments, R² is unsubstituted
naphthyl.

[0068] In some embodiments, the compound of Formula (IIa) has the structure of Formula (IIb) following, wherein L^1 is $-NH-(CH_2)_n-$, n is 0 to 6, preferably 1, and R^1 is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocycloalkyl.




[0069] In some embodiments of the compound of Formula (IIb), L^1 is $-NHCH_2-$ or $-NH(CH_2)_2-$, and R^1 is substituted or unsubstituted aryl. In some embodiments, R^1 is unsubstituted aryl. In some embodiments, R^1 is aryl, preferably phenyl, substituted with halogen, $-CN$ or alkyloxy, preferably methoxy. In some embodiments, R^1 is unsubstituted alkyl, preferably lower alkyl, more preferably methyl or ethyl. In some embodiments, n is 0, and R^1 is hydrogen.

[0070] In some embodiments, the compound of Formula (IIa) has the structure of Formula (IIc) following, wherein L^1 is a bond, and R^1 is unsubstituted alkyl, or substituted or unsubstituted aryl. In some embodiments, R^1 is unsubstituted alkyl, preferably lower alkyl. In some embodiments, R^1 is substituted aryl, preferably halogen-substituted phenyl.



[0071] In some embodiments, there is provided a compound according to Formula (Ib) with structure of Formula (III) following.



[0072] In some embodiments, the compound has the structure of Formula (III). In some embodiments, L^3 is a bond, or substituted or unsubstituted alkylene, and R^3 is substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heterocycloalkyl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, R^3

is substituted or unsubstituted phenyl, or substituted or unsubstituted thienyl. In some embodiments, R^3 is unsubstituted phenyl. In some embodiments, R^3 is unsubstituted thienyl. In some embodiments, R^3 is a chloro-substituted thienyl. In some embodiments, R^3 is substituted or unsubstituted pyridyl, or substituted or unsubstituted pyridazinyl. In some embodiments, R^3 is unsubstituted pyridyl. In some embodiments, R^3 is unsubstituted pyridazinyl. In some embodiments, R^3 is substituted or unsubstituted pyrimidinyl, or substituted or unsubstituted furyl. In some embodiments, R^3 is unsubstituted pyrimidinyl. In some embodiments, R^3 is unsubstituted furyl. In some embodiments, R^3 is substituted or unsubstituted morpholinyl, or substituted or unsubstituted oxanyl, or substituted or unsubstituted oxetanyl. In some embodiments, R^3 is unsubstituted morpholinyl. In some embodiments, R^3 is unsubstituted oxanyl. In some embodiments, R^3 is unsubstituted oxetanyl. In some embodiments, R^3 is substituted or unsubstituted benzodioxinyl, or substituted or unsubstituted naphthyl. In some embodiments, R^3 is unsubstituted benzodioxinyl. In some embodiments, R^3 is unsubstituted naphthyl.


[0073] In some embodiments, the compound has the structure of Formula (III) wherein L^3 is $-C(O)O-$, and R^3 is substituted or unsubstituted alkyl. In some embodiments, the compound has the structure of Formula (III) wherein L^3 is $-C(O)NR^7$, R^7 is hydrogen or alkyl, and R^3 is substituted or unsubstituted alkyl, or substituted or unsubstituted aryl.

[0074] Further to any embodiment above wherein the compound has the structure of Formula (III), in some embodiments, L^1 is $-S-$, $-NR^7-$, substituted or unsubstituted alkylene, or substituted or unsubstituted heteroalkylene, where R^7 is as described in Formula (Ia), and R^1 is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocycloalkyl. In some embodiments, R^1 is substituted or unsubstituted phenyl. In some embodiments, R^1 is unsubstituted phenyl. In some embodiments, R^1 is a substituted or unsubstituted pyridyl. In some embodiments, R^1 is a substituted or unsubstituted pyridazinyl. In some embodiments, R^1 is a substituted or unsubstituted pyrimidinyl. In some embodiments, R^1 is a substituted or unsubstituted thienyl. In some embodiments, R^1 is a substituted or unsubstituted furyl. In some embodiments, R^1 is an unsubstituted pyridyl. In some embodiments, R^1 is an unsubstituted pyridazinyl. In some embodiments, R^1 is an unsubstituted pyrimidinyl. In some embodiments, R^1 is an unsubstituted thienyl. In some embodiments, R^1 is a chloro-substituted thienyl. In some embodiments, R^1 is an unsubstituted furyl. In some embodiments, R^1 is a substituted or unsubstituted morpholinyl. In some embodiments, R^1 is a substituted or unsubstituted oxanyl. In some embodiments, R^1

is a substituted or unsubstituted oxetanyl. In some embodiments, R¹ is an unsubstituted morpholinyl. In some embodiments, R¹ is an unsubstituted oxanyl. In some embodiments, R¹ is an unsubstituted oxetanyl. In some embodiments, R¹ is substituted or unsubstituted benzodioxinyl. In some embodiments, R¹ is substituted or unsubstituted naphthyl. In some embodiments, R¹ is unsubstituted benzodioxinyl. In some embodiments, R¹ is unsubstituted naphthyl.


[0075] Further to any embodiment above wherein the compound has the structure of Formula (III), in some embodiments, L² is a bond. In some embodiments, R² is hydrogen. In some embodiments, L² is substituted or unsubstituted alkylene or -C(O)-, and R² is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl. In some embodiments, R² is substituted or unsubstituted phenyl. In some embodiments, R² is unsubstituted phenyl. In some embodiments, R² is a substituted or unsubstituted pyridyl. In some embodiments, R² is a substituted or unsubstituted pyridazinyl. In some embodiments, R² is a substituted or unsubstituted pyrimidinyl. In some embodiments, R² is a substituted or unsubstituted thienyl. In some embodiments, R² is a substituted or unsubstituted furyl. In some embodiments, R² is an unsubstituted pyridyl. In some embodiments, R² is an unsubstituted pyridazinyl. In some embodiments, R² is an unsubstituted pyrimidinyl. In some embodiments, R² is an unsubstituted thienyl. In some embodiments, R² is a chloro-substituted thienyl. In some embodiments, R² is an unsubstituted furyl. In some embodiments, R² is a substituted or unsubstituted morpholinyl. In some embodiments, R² is a substituted or unsubstituted oxanyl. In some embodiments, R² is a substituted or unsubstituted oxetanyl. In some embodiments, R² is an unsubstituted morpholinyl. In some embodiments, R² is an unsubstituted oxanyl. In some embodiments, R² is an unsubstituted oxetanyl. In some embodiments, R² is substituted or unsubstituted benzodioxinyl. In some embodiments, R² is substituted or unsubstituted naphthyl. In some embodiments, R² is unsubstituted benzodioxinyl. In some embodiments, R² is unsubstituted naphthyl.

[0076] In some embodiments, there is provided a compound according to Formula (Ia) with structure of Formula (IV) following.



[0077] In some embodiments, there is provided a compound according to Formula (IV) wherein L^4 is a bond; and R^4 is hydrogen, halogen, substituted or unsubstituted alkyl, or substituted or unsubstituted heteroalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl. In some embodiments, R^4 is halogen. In some embodiments, R^4 is unsubstituted alkyl. In some embodiments, R^4 is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl. In some embodiments, R^4 is phenyl, naphthyl, pyridyl, pyridazinyl, pyrimidinyl, thienyl, or furyl.

[0078] In some embodiments, there is provided a compound according to Formula (III) with structure of Formula (V) following.



[0079] In some embodiments, the compound is a pharmaceutically acceptable salt, ester, solvate, or prodrug of a compound of Formula (V). In some embodiments, the compound is not an ester, not a solvate, and not a prodrug. In some embodiments, there is provided a compound according to Formula (V) wherein L^1 is a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-. R^1 is hydrogen, a halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl. L^2 is absent, a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-. L^5 is absent, a bond, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene, -S-, -SO-, -SO₂-, -O-, -NHSO₂-, or -NR⁷-. R^2 is absent, hydrogen, a halogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or

815 unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or
816 unsubstituted aryl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted
817 heterocycloalkenyl, substituted or unsubstituted fused ring aryl, or substituted or
818 unsubstituted heteroaryl. In some embodiments, R^2 is absent provided L^2 is also absent. R^5 is
819 absent, hydrogen, a halogen, substituted or unsubstituted alkyl, substituted or unsubstituted
820 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted
821 heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkenyl,
822 substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted fused ring aryl,
823 or substituted or unsubstituted heteroaryl. In some embodiments, R^5 is absent provided L^5 is
824 also absent. R^7 is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted
825 heteroalkyl, substituted or unsubstituted alkylene, substituted or unsubstituted heteroalkylene,
826 substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl,
827 substituted or unsubstituted aryl, substituted or unsubstituted cycloalkenyl, substituted or
828 unsubstituted heterocycloalkenyl, substituted or unsubstituted fused ring aryl, or substituted
829 or unsubstituted heteroaryl.

830 **[0080]** Further to any embodiment above wherein the compound has the structure of
831 Formula (V), in some embodiments, L^2 is a bond. In some embodiments, R^2 is hydrogen. In
832 some embodiments, L^2 is substituted or unsubstituted alkylene or $-C(O)-$, and R^2 is hydrogen,
833 substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or
834 unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or
835 unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or
836 unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted
837 heteroaryl. In some embodiments, R^2 is substituted or unsubstituted phenyl. In some
838 embodiments is an R^2 is unsubstituted phenyl. In some embodiments, R^2 is a substituted or
839 unsubstituted pyridyl. In some embodiments, R^2 is a substituted or unsubstituted pyridazinyl.
840 In some embodiments, R^2 is a substituted or unsubstituted pyrimidinyl. In some embodiments,
841 R^2 is a substituted or unsubstituted thienyl. In some embodiments, R^2 is a substituted or
842 unsubstituted furyl. In some embodiments, R^2 is an unsubstituted pyridyl. In some
843 embodiments, R^2 is an unsubstituted pyridazinyl. In some embodiments, R^2 is an unsubstituted
844 pyrimidinyl. In some embodiments, R^2 is an unsubstituted thienyl. In some embodiments, R^2
845 is a chloro-substituted thienyl. In some embodiments, R^2 is an unsubstituted furyl. In some
846 embodiments, R^2 is a substituted or unsubstituted morpholinyl. In some embodiments, R^2 is a
847 substituted or unsubstituted oxanyl. In some embodiments, R^2 is a substituted or unsubstituted
848 oxetanyl. In some embodiments, R^2 is an unsubstituted morpholinyl. In some embodiments,


849 R^2 is an unsubstituted oxanyl. In some embodiments, R^2 is an unsubstituted oxetanyl. In some
850 embodiments, R^2 is substituted or unsubstituted benzodioxinyl. In some embodiments, R^2 is
851 substituted or unsubstituted naphthyl. In some embodiments, R^2 is unsubstituted
852 benzodioxinyl. In some embodiments, R^2 is unsubstituted naphthyl.

853 **[0081]** Further to any embodiment above wherein the compound has the structure of
854 Formula (V), in some embodiments, L^5 is a bond, or substituted or unsubstituted alkylene,
855 and R^5 is substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl,
856 substituted or unsubstituted heterocycloalkyl, or substituted or unsubstituted
857 heterocycloalkyl. In some embodiments, R^5 is substituted or unsubstituted phenyl, or
858 substituted or unsubstituted thienyl. In some embodiments, R^5 is unsubstituted phenyl. In
859 some embodiments, R^5 is unsubstituted thienyl. In some embodiments, R^5 is a chloro-
860 substituted thienyl. In some embodiments, R^5 is substituted or unsubstituted pyridyl, or
861 substituted or unsubstituted pyridazinyl. In some embodiments, R^5 is unsubstituted pyridyl. In
862 some embodiments, R^5 is unsubstituted pyridazinyl. In some embodiments, R^5 is substituted
863 or unsubstituted pyrimidinyl, or substituted or unsubstituted furyl. In some embodiments, R^5
864 is unsubstituted pyrimidinyl. In some embodiments, R^5 is unsubstituted furyl. In some
865 embodiments, R^5 is substituted or unsubstituted morpholinyl, or substituted or unsubstituted
866 oxanyl, or substituted or unsubstituted oxetanyl. In some embodiments, R^5 is unsubstituted
867 morpholinyl. In some embodiments, R^5 is unsubstituted oxanyl. In some embodiments, R^5 is
868 unsubstituted oxetanyl. In some embodiments, R^5 is substituted or unsubstituted
869 benzodioxinyl, or substituted or unsubstituted naphthyl. In some embodiments, R^5 is
870 unsubstituted benzodioxinyl. In some embodiments, R^5 is unsubstituted naphthyl.

871 **[0082]** Further to any embodiment above wherein the compound has the structure of
872 Formula (V), in some embodiments, L^1 is -S-, $-NR^7$ -, substituted or unsubstituted alkylene, or
873 substituted or unsubstituted heteroalkylene, where R^7 is as described in formula Ia, and R^1 is
874 hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or
875 unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or
876 unsubstituted heterocycloalkyl. In some embodiments, R^1 is substituted or unsubstituted
877 phenyl. In some embodiments, R^1 is unsubstituted phenyl. In some embodiments, R^1 is a
878 substituted or unsubstituted pyridyl. In some embodiments, R^1 is a substituted or unsubstituted
879 pyridazinyl. In some embodiments, R^1 is a substituted or unsubstituted pyrimidinyl. In some
880 embodiments, R^1 is a substituted or unsubstituted thienyl. In some embodiments, R^1 is a
881 substituted or unsubstituted furyl. In some embodiments, R^1 is an unsubstituted pyridyl. In
882 some embodiments, R^1 is an unsubstituted pyridazinyl. In some embodiments, R^1 is an


unsubstituted pyrimidinyl. In some embodiments, R^1 is an unsubstituted thienyl. In some embodiments, R^1 is a chloro-substituted thienyl. In some embodiments, R^1 is an unsubstituted furyl. In some embodiments, R^1 is a substituted or unsubstituted morpholinyl. In some embodiments, R^1 is a substituted or unsubstituted oxanyl. In some embodiments, R^1 is a substituted or unsubstituted oxetanyl. In some embodiments, R^1 is an unsubstituted morpholinyl. In some embodiments, R^1 is an unsubstituted oxanyl. In some embodiments, R^1 is an unsubstituted oxetanyl. In some embodiments, R^1 is substituted or unsubstituted benzodioxinyl. In some embodiments, R^1 is substituted or unsubstituted naphthyl. In some embodiments, R^1 is unsubstituted benzodioxinyl. In some embodiments, R^1 is unsubstituted naphthyl.

[0083] In some embodiments, there is provided a compound according to Formula (V) and its listed embodiments, wherein L^2 and R^2 are absent, providing a compound with structure of Formula (VI) following.



In some embodiments, the compound is a pharmaceutically acceptable salt, ester, solvate, or prodrug of a compound of Formula (VI). In some embodiments, the compound is not an ester, not a solvate, and not a prodrug.

[0084] In some embodiments, there is provided a compound according to Formula (V) and its listed embodiments, wherein L^5 and R^5 are absent, providing a compound with structure of Formula (VII) following.



[0085] In some embodiments, the compound is a pharmaceutically acceptable salt, ester, solvate, or prodrug of a compound of Formula (VII). In some embodiments, the compound is not an ester, not a solvate, and not a prodrug.

[0086] Exemplary compounds, e.g., multisubstituted aromatic compounds, in accordance with the present disclosure are provided herein. In Table A, B, C, and D following,

compound (Cmpd) number, chemical name (i.e., International Union of Pure and Applied Chemistry [IUPAC] name), calculated molecular weight (MW) and biological activity (i.e., inhibition activity in thrombin, KLK1 and KLKB1 assays) are disclosed.

[0087] For Table A following, the disclosed compounds were assayed for inhibition of the protease activity of thrombin as described herein. In Table A, the level of inhibition in the thrombin assay is indicated as follows: a $IC_{50} \leq 0.1 \mu M$; b: $0.1 \mu M < IC_{50} < 1 \mu M$; c: $1 \mu M < IC_{50} < 10 \mu M$; d: $10 \mu M < IC_{50} < 100 \mu M$; e: $IC_{50} \geq 100 \mu M$. Accordingly, in some embodiments, there is provided a compound as expressly set forth in Table A following.

Table A

Cmpd No.	IUPAC name	MW	Thrombin Activity
1	3-(5-amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	176	e
2	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	307	e
3	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	391	a
4	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	401	a
5	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	497	a
6	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	492	a
23	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	415	a
25	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	492	a
26	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	481	a
27	1-[(2-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	516	a
28	1-[(2-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	566	a
29	1-[(2-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	525	a
30	1-[(2-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	431	d
31	1-[(3-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-	516	a

	yl)-1,2-dihydropyridin-2-one		
32	1-[(3-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	566	a
33	1-[(3-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	525	a
34	1-[(3-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	431	c
35	1-[(4-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	516	a
36	1-[(4-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	566	a
37	1-[(4-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	525	a
38	1-[(4-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	431	c
39	1-[(5-chlorothiophen-2-yl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	522	a
40	1-[(5-chlorothiophen-2-yl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	572	a
41	1-[(5-chlorothiophen-2-yl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	531	a
42	1-[(5-chlorothiophen-2-yl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	437	c
43	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	481	a
44	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	511	a
45	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	559	a
46	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	561	a
47	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	531	a
48	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	523	a
49	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	491	a

50	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	491	a
51	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	507	a
52	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	507	a
53	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	586	a
54	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	397	c
55	1-benzyl-3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1,2-dihydropyridin-2-one	535	a
56	2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetonitrile	430	a
57	2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetonitrile	480	a
58	2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetonitrile	440	a
59	2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetic acid	365	d
60	2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetonitrile	346	d
61	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	411	a
62	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	469	a
63	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	491	a
64	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	491	a
65	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	503	a
66	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	502	a
67	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	502	a
68	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	502	e
69	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-	503	a

	2-one		
70	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	507	a
71	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	507	a
72	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-benzyl-1,2-dihydropyridin-2-one	501	a
73	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	425	a
74	3-(5-[(4-fluorophenyl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	470	a
75	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	449	a
76	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-phenylethyl)-1,2-dihydropyridin-2-one	495	a
77	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	471	a
78	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	471	a
79	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(naphthalen-1-ylmethyl)-1,2-dihydropyridin-2-one	531	a
80	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(naphthalen-2-ylmethyl)-1,2-dihydropyridin-2-one	531	a
81	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	483	a
82	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	482	a
83	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	482	a
84	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	482	a
85	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	483	a
86	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	487	a
87	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	487	a
88	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one	518	a

89	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	504	a
90	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	405	a
91	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	407	a
92	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	465	a
93	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	487	a
94	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	498	a
95	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	503	a
96	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	421	a
97	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	479	a
98	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	501	a
99	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	501	a
100	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	513	a
101	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	512	a
102	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	512	a
103	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	512	a
104	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	513	a
105	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	517	a
106	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	517	a

107	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	534	a
108	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	469	a
109	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	527	a
110	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	549	a
111	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	549	a
112	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	561	a
113	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	560	a
114	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	560	a
115	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	560	a
116	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	561	a
117	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	565	a
118	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	565	a
119	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	582	a
120	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	483	a
121	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	471	a
122	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	529	a
123	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	551	a
124	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	563	a

125	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	562	a
126	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	562	a
127	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	562	a
128	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	563	e
129	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	567	a
130	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	567	a
131	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	584	a
132	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	441	a
133	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	499	a
134	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(2-phenylethyl)-1,2-dihydropyridin-2-one	545	a
135	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	521	a
136	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	521	a
137	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(naphthalen-2-ylmethyl)-1,2-dihydropyridin-2-one	581	a
138	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	533	a
139	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	532	a
140	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	532	a
141	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	532	a
142	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	533	a

143	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	537	a
144	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	537	a
145	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one	568	a
146	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	554	a
147	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	455	a
148	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(3-methyloxetan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	419	b
149	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	491	c
150	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	513	a
151	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	513	a
152	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	525	a
153	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	524	a
154	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	524	a
155	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	524	a
156	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	525	a
157	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	529	a
158	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	529	a
159	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	546	e
160	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	459	e

161	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	481	a
162	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	481	a
163	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	493	e
164	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	492	a
165	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	492	a
166	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	492	a
167	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	493	a
168	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	497	a
169	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	497	a
170	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	514	e
171	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	459	e
172	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-phenylethyl)-1,2-dihydropyridin-2-one	505	a
173	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	481	a
174	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(naphthalen-1-ylmethyl)-1,2-dihydropyridin-2-one	541	a
175	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(naphthalen-2-ylmethyl)-1,2-dihydropyridin-2-one	541	a
176	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	493	a
177	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	492	a
178	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	493	a

179	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	497	a
180	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one	528	a
181	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	514	a
182	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	475	a
183	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	497	a
184	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	497	a
185	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	509	e
186	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	508	a
187	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	508	a
188	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	508	a
189	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	509	a
190	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	513	a
191	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	513	a
192	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	530	e
193	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	417	a
194	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	475	a
195	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	497	a
196	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	497	a
197	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-	509	a

	yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one		
198	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	508	a
199	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	508	e
200	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	508	a
201	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	509	a
202	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	513	a
203	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	513	a
204	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	530	a
205	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[2-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	496	c
206	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(2-methoxyethoxy)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	485	a
207	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	496	a
208	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	554	a
209	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	576	a
210	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	576	a
211	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	588	a
212	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	587	a
213	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	587	a
214	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	587	a
215	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	588	a

216	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	592	a
217	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	592	a
218	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	609	a
219	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(1H-1,2,3,4-tetrazol-5-ylmethyl)-1,2-dihydropyridin-2-one	389	d
220	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	365	d
221	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(2-phenylethyl)-1,2-dihydropyridin-2-one	411	d
222	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	387	e
223	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	387	c
224	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(naphthalen-1-ylmethyl)-1,2-dihydropyridin-2-one	447	e
225	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(naphthalen-2-ylmethyl)-1,2-dihydropyridin-2-one	447	e
226	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	399	d
227	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	398	c
228	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	398	c
229	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	398	c
230	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	399	e
231	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	403	c
232	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	403	c
233	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one	434	c
234	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	420	d
235	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-[2-(pyridin-2-yl)ethyl]-1,2-dihydropyridin-2-one	412	d
236	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	321	d
237	3-[5-(benzylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1,2-dihydropyridin-2-one	360	b
238	3-[5-(benzylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	452	a
239	3-[5-(dimethylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1,2-dihydropyridin-2-one	298	e

240	3-[5-(dimethylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	389	c
241	3-1-[(2-aminophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1,2-dihydropyridin-2-one	426	a
242	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	503	a
243	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	525	a
244	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	525	a
245	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	537	a
246	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	536	a
247	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	536	a
248	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	536	a
249	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	537	a
250	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	542	a
251	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	542	a
252	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	558	a
253	3-1-[(4-tert-butylphenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1,2-dihydropyridin-2-one	467	a
254	3-1-[(4-tert-butylphenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-methyl-1,2-dihydropyridin-2-one	481	a
255	3-5-amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl-1,2-dihydropyridin-2-one	270	c
256	3-5-amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	361	c
257	ethyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate	477	a
258	ethyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate	527	a
259	ethyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-	487	a

	dihydropyridin-1-yl]acetate		
260	ethyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate	393	d
261	tert-butyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate	505	a
262	tert-butyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate	555	a
263	tert-butyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate	515	a
264	tert-butyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate	421	e
265	1-[(5-[(5-chlorothiophen-2-yl)methyl]amino-3-(1-methyl-2-oxo-1,2-dihydropyridin-3-yl)-1H-pyrazol-1-yl)carbonyl]cyclopropylmethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate	517	a

918

919 **[0088]** For Table B following, the disclosed compounds were assayed for inhibition of the
 920 protease activity of KLK1 and KLKB1 as described herein. In Table B, C, and D, the level of
 921 inhibition in the KLK1 and KLKB1 assays are indicated as follows: a: $IC_{50} \leq 0.1 \mu M$; b: 0.1
 922 $\mu M < IC_{50} < 1 \mu M$; c: $1 \mu M < IC_{50} < 10 \mu M$; d: $10 \mu M < IC_{50} < 100 \mu M$; e: $IC_{50} \geq 100 \mu M$.
 923 Accordingly, in some embodiments, there is provided a compound as expressly set forth in
 924 Table B following.

925 **Table B**

Cmpd No.	IUPAC name	MW	KLK1 Activity	KLKB1 Activity
4	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	401	d	c
5	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	497		c
19	1-(5-[(4-fluorophenyl)methyl]amino-3-phenyl-1H-pyrazol-1-yl)-2,2-dimethylpropan-1-one	351		e
20	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-2-yl)-1H-pyrazol-1-yl)-2,2-dimethylpropan-1-one	352		e
21	ethyl 3-[(4-fluorophenyl)methyl]amino-1-(pyridin-2-yl)-1H-pyrazole-4-carboxylate	340		e
23	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	415	d	c
24	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-4-fluoro-3-(piperidin-4-yl)-1H-pyrazol-1-yl)-2,2-dimethylpropan-1-one	399		b
25	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	492	d	b
26	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-	481	d	c

	dihydropyridin-2-one			
46	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	561		d
48	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	523		e
49	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	491		b
50	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	491		d
51	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	507		e
52	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	507		b
53	1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	586		e
59	2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetic acid	365	e	c
62	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	469		c
73	3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one	425		e
77	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	471		e
81	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	483	e	c
92	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	465		d
93	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	487		d
94	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	498		c
97	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	479		d
98	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	501		e
100	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	513		c
101	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-	512		d

	methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one			
102	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	512		d
103	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	512		d
104	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	513		c
105	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	517		e
106	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	517		e
109	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	527		c
113	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	560	d	b
116	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	561		c
117	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	565		e
122	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	529		c
123	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	551		d
125	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	562		c
126	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	562	e	c
131	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	584		b
132	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one	441	e	c
138	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one	533	e	b
140	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	532		b
145	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-[2-	568	e	c

	(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one			
149	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	491	e	c
157	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one	529		e
160	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	459		b
179	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	497		d
182	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	475		c
184	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	497		d
194	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	475		b
196	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	497		b
198	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	508		b
201	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one	509		b
203	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one	513		b
208	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	554		c
210	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	576		d
213	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	587	e	b
214	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	587		b
218	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one	609		b
227	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	398		e
229	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	398		e
235	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-	412		d

	3-yl)-1-[2-(pyridin-2-yl)ethyl]-1,2-dihydropyridin-2-one			
237	3-[5-(benzylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1,2-dihydropyridin-2-one	360		d
238	3-[5-(benzylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one	452		c
242	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one	503	e	b
243	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one	525		b
244	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one	525		b
247	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one	536	e	a
248	3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one	536	e	b
265	1-[(5-[(5-chlorothiophen-2-yl)methyl]amino-3-(1-methyl-2-oxo-1,2-dihydropyridin-3-yl)-1H-pyrazol-1-yl)carbonyl]cyclopropylmethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate	517		c
266	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridazin-3-yl)-1H-pyrazol-1-yl)-2,2-dimethylpropan-1-one	353		e
267	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyrimidin-4-yl)-1H-pyrazol-1-yl)-2,2-dimethylpropan-1-one	353		e
268	1-(5-[(4-fluorophenyl)methyl]amino-3-phenyl-1H-pyrazol-1-yl)propan-1-one	323		e
269	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-3-(3-fluoropyridin-2-yl)-1H-pyrazol-1-yl)-2,2-dimethylpropan-1-one	393		e
270	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-3-(furan-2-yl)-1H-pyrazol-1-yl)-2,2-dimethylpropan-1-one	364		e
271	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-3-(pyridin-2-yl)-1H-pyrazol-1-yl)-2,2-dimethylpropan-1-one	375		e
272	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-3-(pyridin-2-yl)-1H-pyrazol-1-yl)-3-methoxy-2,2-dimethylpropan-1-one	405		e
273	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-4-fluoro-3-(oxan-4-yl)-1H-pyrazol-1-yl)-3-hydroxy-2,2-dimethylpropan-1-one	416		d
274	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-4-fluoro-3-(piperidin-4-yl)-1H-pyrazol-1-yl)-3-hydroxy-2,2-dimethylpropan-1-one	415		c
275	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-4-fluoro-3-phenyl-1H-pyrazol-1-yl)-3-hydroxy-2,2-dimethylpropan-1-one	408		e
276	1-[(2-aminophenyl)carbonyl]-N-[(5-chlorothiophen-2-yl)methyl]-3-(3-fluoropyridin-2-yl)-1H-pyrazol-5-amine	428	d	c
277	1-[(2-aminophenyl)carbonyl]-N-[(5-chlorothiophen-2-yl)methyl]-3-(furan-2-yl)-1H-pyrazol-5-amine	399	d	c
278	1-[(2-methoxyphenyl)carbonyl]-3-(pyridin-2-yl)-N-(thiophen-2-ylmethyl)-1H-pyrazol-5-amine	390		e

279	1-[5-(benzylamino)-4-fluoro-3-(pyridin-2-yl)-1H-pyrazol-1-yl]-2,2-dimethylpropan-1-one	352		e
280	1-5-[(furan-2-ylmethyl)amino]-3-(pyridin-2-yl)-1H-pyrazol-1-yl-2,2-dimethylpropan-1-one	324		e
281	2,2-dimethyl-1-[3-(pyridin-2-yl)-5-[(thiophen-2-ylmethyl)amino]-1H-pyrazol-1-yl]propan-1-one	340		e
282	2-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)cyclohexan-1-ol	446	e	c
283	N-(furan-2-ylmethyl)-1-[(2-methoxyphenyl)carbonyl]-3-(pyridin-2-yl)-1H-pyrazol-5-amine	374		e
284	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-(oxan-4-yl)-1H-pyrazol-5-amine	432		e
285	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-(pyridin-2-yl)-1H-pyrazol-5-amine	425	d	c
286	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-phenoxyphenyl)carbonyl]-3-(pyridin-2-yl)-1H-pyrazol-5-amine	487		e
287	N-[(5-chlorothiophen-2-yl)methyl]-1-[(4-methyloxan-4-yl)carbonyl]-3-(pyridin-2-yl)-1H-pyrazol-5-amine	417		d
288	N-[(5-chlorothiophen-2-yl)methyl]-1-[(furan-3-yl)carbonyl]-3-(pyridin-2-yl)-1H-pyrazol-5-amine	385		d
289	N-[(5-chlorothiophen-2-yl)methyl]-1-[(furan-3-yl)carbonyl]-3-(thiophen-2-yl)-1H-pyrazol-5-amine	390		e
290	N-[(5-chlorothiophen-2-yl)methyl]-3-(3-fluoropyridin-2-yl)-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-5-amine	443		e
291	N-[(5-chlorothiophen-2-yl)methyl]-3-(3-fluoropyridin-2-yl)-1-[(furan-3-yl)carbonyl]-1H-pyrazol-5-amine	403		e
292	N-[(5-chlorothiophen-2-yl)methyl]-3-(pyridazin-3-yl)-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-5-amine	402	e	b
293	N-[(5-chlorothiophen-2-yl)methyl]-3-(pyridin-2-yl)-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-5-amine	401	e	b
294	N-[(5-chlorothiophen-2-yl)methyl]-4-fluoro-1-[(furan-3-yl)carbonyl]-3-(oxan-4-yl)-1H-pyrazol-5-amine	410	e	c
295	N-[(5-chlorothiophen-2-yl)methyl]-4-fluoro-1-[(furan-3-yl)carbonyl]-3-phenyl-1H-pyrazol-5-amine	402		e
296	N-benzyl-4-fluoro-1-[(2-methoxyphenyl)carbonyl]-3-phenyl-1H-pyrazol-5-amine	401		e
297	tert-butyl 4-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)piperidine-1-carboxylate	481		e
298	tert-butyl 4-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)piperidine-1-carboxylate	507		e

926

927 [0089] In some embodiments, there is provided a compound as expressly set forth in Table
928 C following.

929 Table C

Cmpd No.	IUPAC name	MW	KLK1 Activity	KLKB1 Activity
7	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)propan-1-one	325		d
8	N-benzyl-1-[(furan-2-yl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	345		c
9	1-[5-(benzylamino)-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl]-3-phenylpropan-1-one	383		e
10	N-[(4-fluorophenyl)methyl]-3-(pyridin-2-yl)-1-[(thiophen-2-yl)carbonyl]-1H-1,2,4-triazol-5-amine	379		c
11	N-[(4-fluorophenyl)methyl]-3-(pyridin-4-yl)-1-[(thiophen-3-yl)carbonyl]-1H-1,2,4-triazol-5-amine	379		c
12	N-[(4-fluorophenyl)methyl]-1-[(morpholin-4-yl)carbonyl]-3-(pyridin-3-yl)-1H-1,2,4-triazol-5-amine	382		e
13	1-[5-(dimethylamino)-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl]propan-1-one	245		e
14	3-(5-[(4-fluorophenyl)methyl]sulfanyl-1-[(2-methoxyphenyl)carbonyl]-1H-1,2,4-triazol-3-yl)pyridine	420		b
15	1-[(2-methoxyphenyl)carbonyl]-5-(methylsulfanyl)-3-(thiophen-2-yl)-1H-1,2,4-triazole	331	c	c
16	N-benzyl-1-[(2-methoxyphenyl)carbonyl]-3-(pyrimidin-4-yl)-1H-1,2,4-triazol-5-amine	386		b
17	N-benzyl-1-[(2-methoxyphenyl)carbonyl]-3-(pyrimidin-5-yl)-1H-1,2,4-triazol-5-amine	386		b
18	N-benzyl-1-[(2-methoxyphenyl)carbonyl]-3-(pyrimidin-2-yl)-1H-1,2,4-triazol-5-amine	386	d	b
22	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	426	d	a
299	1-(1,3-benzothiazol-2-yl)-N-[(4-fluorophenyl)methyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	402		e
300	1-(5-[(4-fluorophenyl)methyl]amino-3-(furan-2-yl)-1H-1,2,4-triazol-1-yl)-2,2-dimethylpropan-1-one	342		d
301	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-2,2-dimethylpropan-1-one	353		d
302	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-2-methylpropan-1-one	339		d
303	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-2-phenylethan-1-one	387		c
304	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-3-methylbutan-1-one	353		d
305	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-3-phenylpropan-1-one	401		d
306	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)butan-1-one	339		d
307	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)propan-1-one	325		d
308	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-2,2-dimethylpropan-1-one	353		d
309	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-2-methoxyethan-1-one	341		e
310	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)propan-1-one	339		d

	1H-1,2,4-triazol-1-yl)-2-methylpropan-1-one			
311	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-2-phenylethan-1-one	387		c
312	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-3-methylbutan-1-one	353		d
313	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-3-phenylpropan-1-one	401		d
314	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)butan-1-one	339		d
315	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl)-2-methylpropan-1-one	339		d
316	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl)-3-methylbutan-1-one	353		e
317	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl)butan-1-one	339		e
318	1-(5-[(4-fluorophenyl)methyl]amino-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl)propan-1-one	325		e
319	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-2,2-dimethylpropan-1-one	365		d
320	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-2-methylpropan-1-one	351		e
321	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-2-phenylethan-1-one	399		d
322	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-3-methylbutan-1-one	365		d
323	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)-3-phenylpropan-1-one	414	d	c
324	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)butan-1-one	351		d
325	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl)propan-1-one	337		d
326	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-2,2-dimethylpropan-1-one	365		e
327	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-2-phenylethan-1-one	399		d
328	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-3-methylbutan-1-one	365		e
329	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)-3-phenylpropan-1-one	414		e
330	1-(5-[(4-methoxyphenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl)butan-1-one	351		d
331	1-(5-[(5-chlorothiophen-2-yl)methyl]amino-3-(oxolan-2-yl)-1H-1,2,4-triazol-1-yl)-2,2-dimethylpropan-1-one	369		c
332	1-[(2-methoxyphenyl)carbonyl]-3-(pyridin-2-yl)-N-(1,3-thiazol-2-ylmethyl)-1H-1,2,4-triazol-5-amine	392	b	d
333	1-[(2-methoxyphenyl)carbonyl]-3-phenyl-N-(1,3-thiazol-2-ylmethyl)-1H-1,2,4-triazol-5-amine	391		e
334	1-[(2-methoxyphenyl)carbonyl]-3-phenyl-N-(thiophen-2-ylmethyl)-1H-1,2,4-triazol-5-amine	390		d
335	1-[(2-methoxyphenyl)carbonyl]-3-phenyl-N-(thiophen-3-ylmethyl)-1H-1,2,4-triazol-5-amine	390		d
336	1-[(furan-2-yl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	375		c
337	1-[(furan-2-yl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3-(pyridin-4-yl)-1H-1,2,4-triazol-5-amine	375		e

338	1-[(furan-3-yl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	375		d
339	1-[(furan-3-yl)carbonyl]-N-[(4-methoxyphenyl)methyl]-3-(pyridin-3-yl)-1H-1,2,4-triazol-5-amine	375		e
340	1-[3-(pyridin-3-yl)-5-[(thiophen-2-ylmethyl)amino]-1H-1,2,4-triazol-1-yl]propan-1-one	313		d
341	1-[4-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-1,2,4-triazol-3-yl)piperidin-1-yl]-2,2-dimethylpropan-1-one	516		b
342	1-[5-(benzylamino)-3-(furan-2-yl)-1H-1,2,4-triazol-1-yl]-2,2-dimethylpropan-1-one	324		e
343	1-[5-(benzylamino)-3-(furan-2-yl)-1H-1,2,4-triazol-1-yl]-2-phenylethan-1-one	358		e
344	1-[5-(benzylamino)-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl]-2,2-dimethylpropan-1-one	335		d
345	1-[5-(benzylamino)-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl]-2-methylpropan-1-one	321		d
346	1-[5-(benzylamino)-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl]-3-methylbutan-1-one	335		d
347	1-[5-(benzylamino)-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl]butan-1-one	321		d
348	1-[5-(benzylamino)-3-(pyridin-2-yl)-1H-1,2,4-triazol-1-yl]propan-1-one	307		d
349	1-[5-(benzylamino)-3-(pyridin-3-yl)-1H-1,2,4-triazol-1-yl]propan-1-one	307		d
350	1-[5-(benzylamino)-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl]-2,2-dimethylpropan-1-one	335		e
351	1-[5-(benzylamino)-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl]-2-methylpropan-1-one	321		c
352	1-[5-(benzylamino)-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl]-2-phenylethan-1-one	369		e
353	1-[5-(benzylamino)-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl]-3-methylbutan-1-one	335		e
354	1-[5-(benzylamino)-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl]-3-phenylpropan-1-one	383		e
355	1-[5-(benzylamino)-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl]butan-1-one	321		e
356	1-[5-(benzylamino)-3-(pyridin-4-yl)-1H-1,2,4-triazol-1-yl]propan-1-one	307		e
357	1-[5-(benzylamino)-3-(thiophen-2-yl)-1H-1,2,4-triazol-1-yl]-2,2-dimethylpropan-1-one	340		e
358	1-benzoyl-N-[(4-fluorophenyl)methyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	373	d	b
359	1-benzoyl-N-[(4-fluorophenyl)methyl]-3-(pyridin-3-yl)-1H-1,2,4-triazol-5-amine	373		b
360	1-benzoyl-N-[(4-methoxyphenyl)methyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	385	c	c
361	1-benzoyl-N-[(4-methoxyphenyl)methyl]-3-(pyridin-3-yl)-1H-1,2,4-triazol-5-amine	385	c	c
362	1-benzoyl-N-benzyl-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	355	d	b
363	1-benzoyl-N-benzyl-3-(pyridin-4-yl)-1H-1,2,4-triazol-5-amine	355		b

364	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-1,2,4-triazol-3-yl)-N,N-dimethylazetidine-1-sulfonamide	511		a
365	3-(pyridin-3-yl)-1-[(pyridin-3-yl)carbonyl]-N-(thiophen-2-ylmethyl)-1H-1,2,4-triazol-5-amine	362		c
366	3-1-[(2-methoxyphenyl)carbonyl]-5-(methylsulfanyl)-1H-1,2,4-triazol-3-ylpyridine	326	c	c
367	N-[(4-fluorophenyl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	403		c
368	N-[(4-fluorophenyl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-(pyridin-3-yl)-1H-1,2,4-triazol-5-amine	403		c
369	N-[(4-fluorophenyl)methyl]-1-[(furan-2-yl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	363		c
370	N-[(4-fluorophenyl)methyl]-1-[(furan-2-yl)carbonyl]-3-(pyridin-3-yl)-1H-1,2,4-triazol-5-amine	363		c
371	N-[(4-fluorophenyl)methyl]-1-[(furan-3-yl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	363		c
372	N-[(4-fluorophenyl)methyl]-1-[(furan-3-yl)carbonyl]-3-(pyridin-3-yl)-1H-1,2,4-triazol-5-amine	363		c
373	N-[(4-fluorophenyl)methyl]-1-[(furan-3-yl)carbonyl]-3-(pyridin-4-yl)-1H-1,2,4-triazol-5-amine	363		e
374	N-[(4-fluorophenyl)methyl]-1-propyl-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	311		e
375	N-[(4-fluorophenyl)methyl]-3-(pyridin-2-yl)-1-[(thiophen-3-yl)carbonyl]-1H-1,2,4-triazol-5-amine	379		b
376	N-[(4-fluorophenyl)methyl]-3-(pyridin-3-yl)-1-[(thiophen-2-yl)carbonyl]-1H-1,2,4-triazol-5-amine	379		c
377	N-[(4-fluorophenyl)methyl]-3-(pyridin-3-yl)-1-[(thiophen-3-yl)carbonyl]-1H-1,2,4-triazol-5-amine	379		b
378	N-[(4-fluorophenyl)methyl]-3-(pyridin-4-yl)-1-[(thiophen-2-yl)carbonyl]-1H-1,2,4-triazol-5-amine	379		e
379	N-[(4-methoxyphenyl)methyl]-3-(pyridin-2-yl)-1-[(thiophen-2-yl)carbonyl]-1H-1,2,4-triazol-5-amine	391		e
380	N-[(4-methoxyphenyl)methyl]-3-(pyridin-2-yl)-1-[(thiophen-3-yl)carbonyl]-1H-1,2,4-triazol-5-amine	391		c
381	N-[(4-methoxyphenyl)methyl]-3-(pyridin-3-yl)-1-[(thiophen-2-yl)carbonyl]-1H-1,2,4-triazol-5-amine	391		e
382	N-[(4-methoxyphenyl)methyl]-3-(pyridin-3-yl)-1-[(thiophen-3-yl)carbonyl]-1H-1,2,4-triazol-5-amine	391		c
383	N-[(5-chlorofuran-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-phenyl-1H-1,2,4-triazol-5-amine	409		e
384	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2,4-dimethoxyphenyl)carbonyl]-3-(oxan-4-yl)-1H-1,2,4-triazol-5-amine	463	d	b
385	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-(oxan-4-yl)-1H-1,2,4-triazol-5-amine	433		b
386	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-(oxolan-2-yl)-1H-1,2,4-triazol-5-amine	419		b
387	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-(piperidin-4-yl)-1H-1,2,4-triazol-5-amine	432	d	a

388	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-[1-(pyrrolidine-1-sulfonyl)azetidin-3-yl]-1H-1,2,4-triazol-5-amine	537		a
389	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-[4-(morpholin-4-yl)phenyl]-1H-1,2,4-triazol-5-amine	510		c
390	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methoxyphenyl)carbonyl]-3-phenyl-1H-1,2,4-triazol-5-amine	425		b
391	N-[(5-chlorothiophen-2-yl)methyl]-1-[(2-methylphenyl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	410	d	b
392	N-[(5-chlorothiophen-2-yl)methyl]-1-[(furan-3-yl)carbonyl]-3-(oxolan-2-yl)-1H-1,2,4-triazol-5-amine	379		b
393	N-[(5-chlorothiophen-2-yl)methyl]-3-[4-(dimethylamino)phenyl]-1-[(2-methoxyphenyl)carbonyl]-1H-1,2,4-triazol-5-amine	468	d	b
394	N-benzyl-1-[(2,6-difluorophenyl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	391		e
395	N-benzyl-1-[(2-chlorophenyl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	390		b
396	N-benzyl-1-[(2-chlorophenyl)carbonyl]-3-(thiophen-2-yl)-1H-1,2,4-triazol-5-amine	395		b
397	N-benzyl-1-[(2-methoxyphenyl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	385		c
398	N-benzyl-1-[(2-methoxyphenyl)carbonyl]-3-(thiophen-2-yl)-1H-1,2,4-triazol-5-amine	390		c
399	N-benzyl-1-[(4-chlorophenyl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	390		b
400	N-benzyl-1-[(furan-2-yl)carbonyl]-3-(pyridin-4-yl)-1H-1,2,4-triazol-5-amine	345		c
401	N-benzyl-1-[(furan-3-yl)carbonyl]-3-(pyridin-2-yl)-1H-1,2,4-triazol-5-amine	345		e
402	N-benzyl-1-[(furan-3-yl)carbonyl]-3-(pyridin-4-yl)-1H-1,2,4-triazol-5-amine	345		d
403	N-benzyl-3-(furan-2-yl)-1-[(2-methoxyphenyl)carbonyl]-1H-1,2,4-triazol-5-amine	374		c
404	N-benzyl-3-(pyridin-2-yl)-1-[(thiophen-2-yl)carbonyl]-1H-1,2,4-triazol-5-amine	361		c
405	N-benzyl-3-(pyridin-2-yl)-1-[(thiophen-3-yl)carbonyl]-1H-1,2,4-triazol-5-amine	361		d
406	N-benzyl-3-(pyridin-4-yl)-1-[(thiophen-2-yl)carbonyl]-1H-1,2,4-triazol-5-amine	361		e
407	N-benzyl-3-(pyridin-4-yl)-1-[(thiophen-3-yl)carbonyl]-1H-1,2,4-triazol-5-amine	361	e	b
408	methyl 5-[(4-methylbenzene)amido]-1H-1,2,4-triazole-3-carboxylate	260		e
409	phenyl 5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazole-1-carboxylate	389		e
410	propan-2-yl 5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazole-1-carboxylate	355		e
411	tert-butyl 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-1,2,4-triazol-3-yl)azetidine-1-carboxylate	504		b
412	tert-butyl 4-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-	532		b

	[(2-methoxyphenyl)carbonyl]-1H-1,2,4-triazol-3-yl)piperidine-1-carboxylate			
413	tert-butyl 5-[(4-fluorophenyl)methyl]amino-3-(pyridin-3-yl)-1H-1,2,4-triazole-1-carboxylate	369		e

930

931

932 [0090] In some embodiments, there is provided a compound as expressly set forth in Table
 933 D following.

934 **Table D**

Cmpd No.	IUPAC name	MW	KLKB1 Activity
414	(3R)-N-[(3-chloro-1H-indol-5-yl)methyl]-1-[(4-chlorophenyl)methyl]-5-oxopyrrolidine-3-carboxamide	416	e
415	(3S)-1-benzyl-N-[(3-chloro-1H-indol-5-yl)methyl]-5-oxopyrrolidine-3-carboxamide	382	e
416	(3S)-N-[(3-chloro-1-methyl-1H-indol-5-yl)methyl]-1-[(4-chlorophenyl)methyl]-5-oxopyrrolidine-3-carboxamide	430	d
417	(3S)-N-[(3-chloro-1H-indol-5-yl)methyl]-1-[(2-chlorophenyl)methyl]-5-oxopyrrolidine-3-carboxamide	416	d
418	(3S)-N-[(3-chloro-1H-indol-5-yl)methyl]-1-[(3-chlorophenyl)methyl]-5-oxopyrrolidine-3-carboxamide	416	e
419	2-N-[(2R)-5-amino-1,2,3,4-tetrahydronaphthalen-2-yl]-4-N-(2-fluorophenyl)-4-N,6-dimethylpyrimidine-2,4-diamine	378	e
420	2-N-[(2R)-5-amino-1,2,3,4-tetrahydronaphthalen-2-yl]-4-N-(3-fluorophenyl)-4-N,6-dimethylpyrimidine-2,4-diamine	378	e
421	2-N-[(2R)-5-amino-1,2,3,4-tetrahydronaphthalen-2-yl]-4-N-(4-fluorophenyl)-4-N,6-dimethylpyrimidine-2,4-diamine	378	e
422	2-N-[(2S)-5-amino-1,2,3,4-tetrahydronaphthalen-2-yl]-4-N,6-dimethyl-4-N-phenylpyrimidine-2,4-diamine	360	e
423	2-N-[(2S)-5-amino-1,2,3,4-tetrahydronaphthalen-2-yl]-4-N-(2-chlorophenyl)-4-N,6-dimethylpyrimidine-2,4-diamine	394	d
424	2-N-[(2S)-5-amino-1,2,3,4-tetrahydronaphthalen-2-yl]-4-N-(3-chlorophenyl)-4-N,6-dimethylpyrimidine-2,4-diamine	394	e
425	2-N-[(2S)-5-amino-1,2,3,4-tetrahydronaphthalen-2-yl]-4-N-(4-chlorophenyl)-4-N,6-dimethylpyrimidine-2,4-diamine	394	e

935

936 [0091] Compounds disclosed herein also include racemic mixtures, stereoisomers and
 937 mixtures of the compounds, including isotopically-labeled and radio-labeled compounds. See
 938 e.g., Goding, 1986, MONOCLONAL ANTIBODIES PRINCIPLES AND PRACTICE; Academic Press,
 939 p. 104. Such isomers can be isolated by standard resolution techniques, including e.g.,

fractional crystallization, chiral chromatography, and the like. See e.g., Eliel, E. L. & Wilen S. H., 1993, STEREOCHEMISTRY IN ORGANIC COMPOUNDS ; John Wiley & Sons, New York.

[0092] In some embodiments, compounds disclosed herein have asymmetric centers and can occur as racemates, racemic mixtures, and as individual enantiomers or diastereoisomers, with all isomeric forms as well as mixtures thereof being contemplated for use in the compounds and methods described herein. The compounds contemplated for use in the compounds and methods described herein do not include those that are known in the art to be too unstable to synthesize and/or isolate.

[0093] The compounds disclosed herein can also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds can be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I), or carbon-14 (^{14}C). All isotopic variations of the compounds disclosed herein, whether radioactive or not, are encompassed within the contemplated scope.

[0094] In some embodiments, metabolites of the compounds disclosed herein are useful for the methods disclosed herein.

[0095] In some embodiments, compounds contemplated herein are provided in the form of a prodrug. The term "prodrug" refers to a compound that can be converted into a compound (e.g., a biologically active compound) described herein *in vivo*. Prodrugs can be useful for a variety of reason known in the art, including e.g., ease of administration due e.g., to enhanced bioavailability in oral administration, and the like. The prodrug can also have improved solubility in pharmaceutical compositions over the biologically active compounds. An example, without limitation, of a prodrug is a compound which is administered as an ester (i.e., the "prodrug") to facilitate transmittal across a cell membrane where water solubility is detrimental to mobility but which then is metabolically hydrolyzed to the carboxylic acid, the active entity, once inside the cell where water solubility is beneficial. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in DESIGN OF PRODRUGS, (ed. H. Bundgaard, Elsevier, 1985), which is hereby incorporated herein by reference for the limited purpose describing procedures and preparation of suitable prodrug derivatives.

[0096] Accordingly, in some embodiments, compounds contemplated herein are provided in the form of a prodrug ester. The term "prodrug ester" refers to derivatives of the compounds disclosed herein formed by the addition of any of a variety of ester-forming groups, e.g., groups known in the art, that are hydrolyzed under physiological conditions.

Examples of prodrug ester groups include pivaloyloxymethyl, acetoxymethyl, phthalidyl, indanyl and methoxymethyl, as well as other such groups known in the art, including a (5-R-2-oxo-1,3-dioxolen-4-yl)methyl group. Other examples of prodrug ester groups can be found in, for example, T. Higuchi and V. Stella, in "Pro-drugs as Novel Delivery Systems", Vol. 14, A.C.S. Symposium Series, American Chemical Society (1975); and BIOREVERSIBLE CARRIERS IN DRUG DESIGN: THEORY AND APPLICATION, edited by E. B. Roche, Pergamon Press: New York, 14-21 (1987) (providing examples of esters useful as prodrugs for compounds containing carboxyl groups). Each of the above-mentioned references is herein incorporated by reference for the limited purpose of disclosing ester-forming groups that can form prodrug esters.

[0097] In some embodiments, prodrugs can be slowly converted to the compounds described herein useful for the methods described herein when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent.

[0098] Certain compounds disclosed herein can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are encompassed within the scope of contemplated compounds. Certain compounds of the present invention can exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the compounds and methods contemplated herein and are intended to be within the scope disclosed herein.

III. Biological Activities

[0099] In some embodiments, compounds described herein exhibit inhibitory activity against thrombin with activities $\geq 1 \mu\text{M}$, e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 μM , or even greater. In some embodiments, the compounds exhibit inhibitory activity against thrombin with activities between 0.1 μM and 1 μM , e.g., about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1.0 μM . In some embodiments, compounds described herein exhibit inhibitory activity against thrombin with activities $\leq 0.1 \mu\text{M}$, e.g., about 1, 2, 5, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, or 100 nM. Ranges of values using a combination of any of the values recited herein as upper and/or lower limits are also contemplated, for example, but not limited to, 1–10 nM, 10–100 nM, 0.1–1 μM , 1–10 μM , 10–100 μM , 100–200 μM , 200–500 μM , or even 500–1000 μM . In some embodiments, the inhibitory activity is in the range of about

1–10 nM, 10–100 nM, 0.1–1 μ M, 1–10 μ M, 10–100 μ M, 100–200 μ M, 200–500 μ M, or even 500–1000 μ M. It is understood that for purposes of quantification, the terms “activity,” “inhibitory activity,” “biological activity,” “thrombin activity and the like in the context of an inhibitory compound disclosed herein can be quantified in a variety of ways known in the art. Unless indicated otherwise, as used herein such terms refer to IC₅₀ in the customary sense (i.e., concentration to achieve half-maximal inhibition).

[0100] Inhibitory activity against thrombin in turn inhibits the blood coagulation process. Accordingly, compounds disclosed herein are indicated in the treatment or management of thrombotic disorders. In some embodiments, a dose or a therapeutically effective dose of a compound disclosed herein will be that which is sufficient to achieve a plasma concentration of the compound or its active metabolite(s) within a range set forth herein, e.g., about 1–10 nM, 10–100 nM, 0.1–1 μ M, 1–10 μ M, 10–100 μ M, 100–200 μ M, 200–500 μ M, or even 500–1000 μ M, preferably about 1–10 nM, 10–100 nM, or 0.1–1 μ M. Without wishing to be bound by any theory, it is believed that such compounds are indicated in the treatment or management of thrombotic disorders.

[0101] In some embodiments, compounds described herein exhibit inhibitory activity against KLK1 and KLKB1 with activities between 1 μ M and 10 μ M, e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 μ M. In some embodiments, compounds described herein exhibit inhibitory activity against KLK1 and KLKB1 with activities $\geq$ 10 μ M, e.g., about 10, 20, 50, 100, 150, 200, 300, 400, 500, 600, 700, 800, 900, 1000 μ M or even greater. In some embodiments, compounds described herein exhibit inhibitory activity against KLK1 and KLKB1 with activities $\leq$ 1 μ M, e.g., about 900, 800, 700, 600, 500, 400, 300, 200, 100, 50 nM or even lower. Ranges of values using a combination of any of the values recited herein as upper and/or lower limits are also contemplated, for example, but not limited to, 1–10 nM, 10–100 nM, 0.1–1 μ M, 1–10 μ M, 10–100 μ M, 100–200 μ M, 200–500 μ M, or even 500–1000 μ M. In some embodiments, the inhibitory activity is in the range of about 1–10 nM, 10–100 nM, 0.1–1 μ M, 1–10 μ M, 10–100 μ M, 100–200 μ M, 200–500 μ M, or even 500–1000 μ M. It is understood that for purposes of quantification, the terms “activity,” “inhibitory activity,” “biological activity,” “KLK1 activity,” “KLKB1 activity” and the like in the context of an inhibitory compound disclosed herein can be quantified in a variety of ways known in the art. Unless indicated otherwise, as used herein such terms refer to IC₅₀ in the customary sense (i.e., concentration to achieve half-maximal inhibition).

[0102] Inhibitory activity against KLKB1 has an effect on the coagulation cascade and the inflammatory response. Thus, it has been proposed that KLKB1 inhibitors can be useful in the treatment of thrombotic and fibrinolytic diseases and disease conditions.

[0103] Accordingly, compounds disclosed herein are indicated in the treatment or management of a variety of diseases or disorders. In some embodiments, a dose or a therapeutically effective dose of a compound disclosed herein will be that which is sufficient to achieve a plasma concentration of the compound or its active metabolite(s) within a range set forth herein, e.g., about 1–10 nM, 10–100 nM, 0.1–1 μ M, 1–10 μ M, 10–100 μ M, 100–200 μ M, 200–500 μ M, or even 500–1000 μ M, preferably about 1–10 nM, 10–100 nM, or 0.1–1 μ M. Without wishing to be bound by any theory, it is believed that such compounds are indicated in the treatment or management of diseases associated with thrombin or kallikrein.

IV. Methods of Treating and Preventing Disease

[0104] Kallikrein-related diseases or disorders are biological conditions associated with or moderated by kallikrein. They include, but are not limited by, those conditions associated with biological pathways that are moderated by tissue and plasma kallikrein. An example of such a pathway is the kallikrein-kinin system (Moreau, M.E. 2005, *Journal of Pharmacological Sciences*, 99, 6). Kallikrein-related diseases or disorders include, but are not limited to, fibrosis, inflammation, thrombosis, hereditary angioedema, skin disorders, cancer, and ophthalmic diseases. Ophthalmic diseases include, but are not limited to, diabetic macular edema, diabetic retinopathy, and age-related macular degeneration.

[0105] **Diabetic Macular Edema.** In rodent models, it has been shown that activation of KLKB1 in the eye increases retinal vascular permeability; whereas inhibition of the kallikrein-kinin system reduces retinal leakage induced by diabetes and hypertension. These findings suggest that intraocular activation of the KLKB1 pathway can contribute to excessive retinal vascular permeability that can lead to diabetic macular edema. Thus, evidence suggests that KLKB1 inhibitors can provide a new therapeutic opportunity to reduce retinal vascular permeability (Feener, E. P. 2010, *Curr Diab Rep* 10, 270).

[0106] **Hereditary Angioedema.** Ecallantide (Kalbitor) is a 60-amino acid recombinant protein that acts as a potent reversible inhibitor of KLKB1 (Schneider L, *et al.* 2007, *J Allergy Clin Immunol*, 120, 416) and has been approved by the FDA for the treatment of acute attacks of hereditary angioedema (HAE). Thus plasma kallikrein inhibition can be a

useful treatment for HAE, and there is strong interest in the development of plasma kallikrein inhibitors as a therapy for HAE.

[0107] Skin. Overexpression of various KLKs in the skin has led to the recognition that certain kallikrein inhibitors can be useful for certain dermatological conditions, including atopic dermatitis, psoriasis and rare skin diseases such as Netherton Syndrome (Freitas *et al. Bioorganic & Medicinal Chemistry Letters* 2012, 22, 6072–6075).

[0108] Thrombosis. Thrombotic diseases are the primary indications for thrombin inhibition, because of thrombin's location in the coagulation cascade and, in turn, the importance of the coagulation cascade in the progression of blood clotting processes. However, without wishing to be bound by any theory, it is believed the coagulation cascade in general, and thrombin in particular, is important in a variety other disease states.

[0109] It has been discovered that compounds described herein, e.g., multisubstituted aromatic compounds, exhibit inhibitory action against thrombin (activated blood-coagulation factor II; EC 3.4.21.5). This, in turn inhibits the blood coagulation process.

[0110] This inhibitory action is useful in the treatment of a variety of thrombotic disorders, such as, but not limited to, acute vascular diseases such as acute coronary syndromes; venous-, arterial- and cardiogenic thromboembolisms; the prevention of other states such as disseminated intravascular coagulation, or other conditions that involve the presence or the potential formation of a blood clot thrombus. Other indications for methods described herein include the following.

[0111] Cancer. Tissue kallikreins (KLKs) are subdivided into various types, and have been extensively investigated in cancer and inflammation biology. Various kallikrein KLKs have been found to be up- or down-regulated in various cancer types, such as cervical-, testicular-, and non-small-cell lung adenocarcinoma (Caliendo *et al. J. Med. Chem.*, 2012, 55, 6669). It has been proposed that KLK1 inhibitors will be useful in certain cancers.

[0112] It has long been recognized that cancer progression is accompanied by venous thrombosis, but it has not been understood how each disease is related. From several clinical trials studying the treatment of VTE, meta-analyses have shown that low molecular weight heparins (LMWHs) improve overall survival in subgroups of cancer patients. *See e.g.*, Zacharski, L. R. & Lee, A. Y., 2008, *Expert Opin Investig Drugs*, 17:1029-1037; Falanga, A. & Piccioli, A., 2005, *Current Opinion in Pulmonary Medicine*, 11:403-407; Smorenburg, S. M., *et al.*, 1999, *Thromb Haemost*, 82:1600-1604; Hettiarachchi, R. J., *et al.*, 1999, *Thromb Haemost*, 82:947-952. This finding was substantiated in later clinical trials that measured

specifically the survival of cancer patients. See e.g., Lee, A. Y.*et al.*, 2005, *J Clin Oncol*, 23:2123-2129; Klerk, C. P.*et al.*, *J Clin Oncol* 2005, 23:2130-2135; Kakkar, A. K., *et al.*, 2004, *J Clin Oncol*, 22:1944-1948; Altinbas, M., *et al.*, 2004, *J Thromb Haemost*, 2:1266-1271.

[0113] More recently, researchers have focused on the specific anticancer effect of DTIs. For example, it was shown that heparin significantly prolonged the survival of patients with limited small cell lung cancer. See e.g., Akl, E. A., *et al.*, 2008, *J Exp Clin Cancer Res*, 27:4. Other investigators found that systemic use of argatroban reduced tumor mass and prolonged survival time in rat glioma models leading to the conclusion that argatroban should be considered as a novel therapeutic for glioma, a notoriously difficult to treat cancer type. See e.g., Hua, Y., *et al.*, 2005, *Acta Neurochir*, Suppl 2005, 95:403-406; Hua, Y., *et al.*, 2005, *J Thromb Haemost*, 3:1917-1923. Very recently, it was demonstrated that dabigatran etexilate, a DTI recently FDA-approved (see e.g., Hughes, B., 2010, *Nat Rev Drug Discov*, 9:903-906) for DVT indications, inhibited both the invasion and metastasis of malignant breast tumors. See e.g., DeFeo, K.*et al.*, 2010, *Thrombosis Research*, 125 (Supplement 2): S188-S188; Defeo, K., *et al.*, 2010, *Cancer Biol Ther*, 10:1001-1008. Thus, dabigatran etexilate treatment led to a 50% reduction in tumor volume at 4 weeks with no weight loss in treated mice. Dabigatran etexilate also reduced tumor cells in the blood and liver micrometastases by 50-60%. These investigators concluded that dabigatran etexilate can be beneficial in not only preventing thrombotic events in cancer patients, but also as adjunct therapy to treat malignant tumors.

[0114] Further, hirudin and the LMWH nadroparin dramatically reduced the number of lung metastases when administered prior to cancer cell inoculation. See e.g., Hu, L., *et al.*, 2004, *Blood*, 104:2746-51.

[0115] The de novo thrombin inhibitor d-Arg-Oic-Pro-d-Ala-Phe(p-Me) has been found to block thrombin-stimulated invasion of prostate cancer cell line PC-3 in a concentration dependent manner. See e.g., Nieman, M. T., *et al.*, 2008, *J Thromb Haemost*, 6:837-845. A reduced rate of tumor growth was observed in mice dosed with the pentapeptide through their drinking water. The mice also showed reduced fold rate in tumor size and reduced overall tumor weight compared to untreated mice. Microscopic examination of treated tumors showed reduced number of large blood vessels thus concluding that the pentapeptide interfered with tumor angiogenesis. Nieman, M. T., *et al.*, *Thromb Haemost*, 104:1044-8.

[0116] In view of these and related studies, it is suggested that anticoagulants affect tumor metastasis; that is, angiogenesis, cancer cell adhesion, migration and invasion processes. See e.g., Van Noorden, C. J., *et al.*, 2010, *Thromb Res*, 125 Suppl 2:S77-79.

[0117] **Fibrosis.** Kallikreins are a subgroup of serine proteases, divided into plasma kallikrein (KLKB1) and tissue kallikreins. KLKB1 liberates kinins (bradykinin and kallidin) from the kininogens, peptides responsible for the regulation of blood pressure and activation of inflammation. In the Contact Activation Pathway of the coagulation cascade, KLKB1 assists in the conversion of factor XII to factor XIIa (Keel, M.; Trentz, O. *Injury* 2005, 36, 691–709). Factor XIIa converts FXI into FXIa, which in turn activates FIX, which with its co-factor FVIIIa forms the tenase complex, which finally activates FX to FXa. In the fibrinolysis part of the coagulation cascade, KLKB1 serves to convert plasminogen to plasmin. Thus, it has been proposed that KLKB1 inhibitors can be useful in the treatment of thrombotic and fibrinolytic diseases and disease conditions (US Patent # 7,625,944; Bird *et al.* *Thrombosis and Hemostasis* 2012, 107, 1141).

[0118] Several studies have shown the utility of anticoagulant therapy in fibrotic disorders. For example, in a rat model of CCl₄-induced chronic liver injury, the DTI SSR182289 decreased liver fibrogenesis significantly after 7 weeks of administration. Similar observations were made in other studies using the LMWHs nadroparin, tinzaparin, enoxaparin, and dalteparin sodium. See e.g., Duplantier, J. G., *et al.*, 2004, *Gut*, 53:1682-1687; Abdel-Salam, O. M., *et al.*, 2005, *Pharmacol Res*, 51:59-67; Assy, N., *et al.*, 2007, *Dig Dis Sci*, 52:1187-1193; Abe, W., *et al.*, 2007, *J Hepatol*, 46:286-294. Thus a thrombin inhibitor as an anticoagulant can be useful in the treatment of fibrinolytic diseases.

[0119] In another example, the DTI melagatran greatly reduced ischemia reperfusion injury in a kidney transplant model in the large white pig. This led to a drastically improved kidney graft survival at 3 months. See e.g., Favreau, F., *et al.*, 2010, *Am J Transplant*, 10:30-39.

[0120] Recent studies have shown that in a bleomycin-induced mouse model of pulmonary fibrosis, dabigatran etexilate treatment reduced important profibrotic events in lung fibroblasts, including the production of collagen and connective tissue growth factor. See e.g., Silver, R. M., *et al.*, 2010, *Am. J. Respir. Crit. Care Med.*, 181:A6780; Bogatkevich, G. S., *et al.*, 2009, *Arthritis Rheum*, 60:3455-3464.

[0121] The above experimental evidence points to a close relationship between thrombin and fibrosis and suggests novel therapeutic opportunities for fibrosis using thrombin inhibitors. See e.g., Calvaruso, V., *et al.*, 2008, *Gut*, 57:1722-1727; Chambers, R. C., 2008,

1167 *Br J Pharmacol*, 153 Suppl 1:S367-378; Chambers, R. C. & Laurent, G. J., 2002, *Biochem*
1168 *Soc Trans*, 30:194-200; Howell, D. C., *et al.*, 2001, *Am J Pathol*, 159:1383-1395.

1169 **[0122] Inflammation.** Kallikrein has long been implicated in inflammation (Clements, J.A.
1170 *The Molecular Biology of the Kallikreins and Their Roles in Inflammation*, Academic Press:
1171 San Diego, CA, 1997; Vol. 5). There is experimental evidence that KLKB1 is associated with
1172 sepsis and inflammatory arthritis (Colman, R.W., 1998, *Clinical Reviews in Allergy and*
1173 *Immunology*, 16: 365). Thus a KLKB1 inhibitor can be useful in the treatment of
1174 inflammatory conditions associated with the kallikrein-kinin system, such as systemic
1175 inflammatory response syndrome, sepsis, rheumatoid arthritis, and inflammatory bowel
1176 disease.

1177 **[0123] Age-Related Macular Degeneration.** KLK1 has been linked to blood vessel
1178 growth moderated by the VEGF pathway (Miura S., 2003, *Hypertension*, 41, 1118). Age-
1179 related macular degeneration (AMD) is associated with the proliferation of abnormal blood
1180 vessels and VEGF expression (Lopez, P.F., 1996, *Investigative Ophthalmology & Visual*
1181 *Science*, 37, 855). Thus, KLK1 inhibitors have been proposed for the treatment of AMD (US
1182 Patent #20120264798; Ferrara, N., 2000, *Current Opinion in Biotechnology*, 11, 617).

1183 **[0124] Alzheimer's Disease.** Very recent experiments confirm higher thrombin levels in
1184 brain endothelial cells of patients with Alzheimer's disease. While 'normal' thrombin levels
1185 are connected to regulatory CNS functions, thrombin accumulation in the brain is toxic. It has
1186 also been found that the neural thrombin inhibitor Protease Nexin 1 (PN-1) is significantly
1187 reduced in the Alzheimer's disease brain, despite the fact that PN-1 mRNA levels are
1188 unchanged. These observations have led some investigators to suggest that reduction of CNS-
1189 resident thrombin will prove useful in Alzheimer's Disease (AD) treatment. See e.g.,
1190 Vaughan, P. J., *et al.*, 1994, *Brain Res*, 668:160-170; Yin, X., *et al.*, 2010, *Am J Pathol*,
1191 176:1600-1606; Akiyama, H., *et al.*, 1992, *Neurosci Lett*, 146:152-154.

1192 **[0125] Multiple Sclerosis.** Investigators found that hirudin treatment in an animal model of
1193 Multiple Sclerosis (MS) showed a dramatic improvement in disease severity. See e.g., Han,
1194 M. H., *et al.*, 2008, *Nature*, 451:1076-1081. Similar results were obtained following treatment
1195 with heparin (a DTI) and dermatan sulfate, another coagulation inhibitor. See e.g.,
1196 Chelmicka-Szorc, E. & Arnason, B. G., 1972, *Arch Neurol*, 27:153-158; Inaba, Y., *et al.*,
1197 1999, *Cell Immunol*, 198:96-102. Other evidence shows that naturally occurring antithrombin
1198 III has anti-inflammatory effects in diseases such as endotoxemia and other sepsis-related
1199 conditions. See e.g., Wiedermann, C. J. & Romisch, J., 2002, *Acta Med Austriaca*, 29:89-92.

Naturally occurring thrombin inhibitors are presumably synthesized *in situ* and have protective roles in CNS inflammation. Therefore, therapeutic thrombin inhibition has been proposed as a potential MS treatment. See e.g., Luo, W., *et al.*, 2009, In: THROMBIN, Maragoudakis, M. E.; Tsopanoglou, N. E., Eds. Springer New York: 2009; pp 133-159.

[0126] Pain. In a rat pain model with partial lesion of the sciatic nerve, intrathecal hirudin prevented the development of neuropathic pain and curbed pain responses for 7 days. The investigators found that following injury, neuropathic pain was mediated by thrombin generation, which in turn activated PAR-1 receptor in the spinal cord. Hirudin inhibited thrombin generation and ultimately led to pain relief. See e.g., Garcia, P. S., *et al.*, 2010, *Thromb Haemost*, 103:1145-1151; Narita, M., *et al.*, 2005, *J Neurosci*, 25:10000-10009. Researchers hypothesize that thrombin and the PARs are involved not just as part of the coagulation cascade, but in inflammation, nociception and neurodevelopment. Development of a DTI to intersect an unexploited pharmacology will lead to pain therapeutics distinct from opioids and NSAIDs, whose shortcomings are well documented. See e.g., Garcia 2010, *Id.*

[0127] Accordingly, in a further aspect, there is provided a method for treating a disease or disorder in a subject in need thereof. The method includes administering a compound of any of Formulae (Ia), (Ib), (IIa), (IIb), (IIc), (III), (IV), (V), (VI) or (VII) as disclosed herein, a compound as set forth in Table A, B, C, or D, pharmaceutically acceptable salt, ester, solvate, or prodrug thereof, or pharmaceutical composition thereof, to a subject in need thereof in an amount effective to treat the disease or disorder. The terms “therapeutically effective amount,” “amount effective to treat,” “amount effective to prevent” and the like refer to that amount of drug or pharmaceutical agent (e.g., compound or pharmaceutical composition disclosed herein) that will elicit the biological or medical response of a tissue, system, animal, or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

[0128] Compounds useful for methods disclosed herein include the compounds set forth for Formulae (Ia), (Ib), (IIa), (IIb), (IIc), (III), (IV), (V), (VI) or (VII) and for the compounds set forth in Table A, B, C, or D above.

[0129] In some embodiments, the diseases or disorders are fibrinolytic diseases. In some embodiments the disease is a fibrotic disorder. In some embodiments, the disease is cancer. In some embodiments, the diseases are inflammatory diseases. In some embodiments the disease is sepsis. In some embodiments the disease is inflammatory arthritis. In some embodiments, the disease is diabetic macular edema. In some embodiments, the disease is

hereditary angioedema. In some embodiments, the disease is diabetic retinopathy. In some embodiments, the disease is age-related macular degeneration. In some embodiments, the diseases are various skin diseases which include but are not limited to atopic dermatitis, psoriasis and rare skin diseases such as Netherton Syndrome. In some embodiments, the diseases or disorder is Alzheimer's disease. In some embodiments, the disease is multiple sclerosis. In some embodiments, the disease is pain.

[0130] In some embodiments, the disease or disorder is cancer. In some embodiments, the cancer is limited small cell lung cancer. In some embodiments, the cancer is a glioma. In some embodiments, the cancer is malignant breast cancer. In some embodiments, the cancer is a micrometastasis. In some embodiments, the micrometastasis is of the blood or liver. In some embodiments, the cancer is a lung metastasis. In some embodiments, the cancer is prostatic cancer.

[0131] In another aspect, there is provided a method for preventing a disease or disorder in a subject. The method includes administering a compound of any of Formulae (Ia), (Ib), (IIa), (IIb), (IIc), (III), (IV), (V), (VI) or (VII) as disclosed herein, compound as set forth in any of Table A, B, C, or D herein, pharmaceutically acceptable salt, ester, solvate, or prodrug thereof, or pharmaceutical composition thereof, to a subject in need thereof in an amount effective to prevent the disease or disorder.

V. Assays

[0132] Compounds described herein can be assayed, by a variety of methods known in the art and described herein, for inhibition of biological activity, e.g., protease activity, of a variety of proteins, e.g., thrombin, KLKB1 and KLK1.

[0133] The KLKB1 kallikrein activity reported herein (e.g., Tables B, C, and D) was obtained as follows. Human KLKB1 protein was obtained from Enzyme Research Labs. The chromogenic substrate S-2302 was obtained from DiaPharma. KLKB1 was assayed in buffer containing 0.05 M Tris (pH 7.4), 0.01 M NaCl and 0.2% w/v PEG-8000. The final concentration of enzyme used was 3 nM KLKB1. The final concentration of substrate used was 250 μ M S-2302 for KLKB1. All assays were performed in 96-well microtiter plates at room temperature (RT). The enzyme and inhibitor were pre-incubated for 10 minutes then substrate was added and read at 405 nm in a SpectraMax Plus Spectrophotometer (Molecular Devices). Inhibitor IC₅₀ values were determined by adding test compound as ten point, three-fold serial dilutions in buffer solution, as known in the art. The plate was read at 10 minutes after substrate addition. The IC₅₀ was calculated by plotting the percent (%) inhibition against

compound concentration and fitting the data to a constrained four parameter sigmoidal curve, as known in the art.

[0134] The KLK1 kallikrein activity reported herein (e.g., Tables B and C) was obtained as follows. Recombinant human tissue kallikrein (KLK1) was obtained from R&D Systems. Pro-Phe-Arg-AMC (I-1295) substrate was obtained from Bachem. KLK1 enzyme is activated by incubating 0.5 mg/ml KLK1 combined with 0.1 µg/ml thermolysin in a buffer of 0.05 M Tris (pH 7.5), 0.15 M NaCl, and 0.01 M CaCl₂ for one hour at 37°C. The thermolysin is then deactivated by the addition of equal parts 20 mM 1, 10 phenanthroline solution in water. The activated KLK1 solution is then added to CHES buffer (0.05 M CHES, 0.15 M NaCl, 0.01 M CaCl₂, pH 10) for a final concentration of 5 nM along with the test article and incubated for 10 minutes. Substrate is then added at a concentration of 2.75 µM. Substrate activation is read 10 minutes after substrate addition using a Synergy H1 multifunction plate reader (Biotek) programmed with a 360 nm excitation wavelength and a 480 nm emission wavelength. Inhibitor response was established by adding test compound as ten point, three-fold serial dilutions, as known in the art. The IC₅₀ was calculated by plotting the percent (%) inhibition against compound concentration and fitting the data to a constrained four parameter sigmoidal curve, as known in the art.

[0135] The thrombin activity reported herein (e.g., Table A) was obtained as follows. Human thrombin was obtained from Haematologic Technologies Inc. The chromogenic substrate S-2238 was obtained from DiaPharma. Thrombin was assayed in buffer containing 0.05 M Tris (pH 7.4), 0.015 M NaCl and 0.01% PEG-8000. The final concentration of enzyme used was 3 nM thrombin. The final concentration of substrate used was 125 µM S-2238 for thrombin. All assays were performed in 96-well microtiter plates at room temperature (RT). The enzyme and inhibitor were pre-incubated for 10 minutes then substrate was added and read at 405 nm in a SpectraMax Plus Spectrophotometer (Molecular Devices). Inhibitor IC₅₀ values were determined by adding test compound as ten point, three-fold serial dilutions in buffer solution, as known in the art. The plate was read at 10 minutes after substrate addition. The IC₅₀ was calculated by plotting the percent (%) inhibition against compound concentration and fitting the data to a constrained four parameter sigmoidal curve, as known in the art.

VI. Pharmaceutical Compositions

[0136] In another aspect, there is provided a pharmaceutical composition comprising a compound disclosed herein and a pharmaceutically acceptable excipient. The compound is a

compound of any of Formulae (Ia), (Ib), (IIa), (IIb), (IIc), (III), (IV), (V), (VI) or (VII) as disclosed herein, a compound as set forth in Table A, B, C, or D herein, or pharmaceutically acceptable salt, ester, solvate, or prodrug thereof. In some embodiments, the compound is set forth in Table A, B, C, or D herein.

[0137] The term “pharmaceutically acceptable salts” is meant to include salts of the active compounds that are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds disclosed herein contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds disclosed herein contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, lactic, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, oxalic, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galacturonic acids and the like (*see*, for example, Berge *et al.*, “Pharmaceutical Salts”, *Journal of Pharmaceutical Science*, **1977**, 66, 1-19). Certain specific compounds disclosed herein contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0138] Compounds disclosed herein can exist as salts, such as with pharmaceutically acceptable acids. Accordingly, the compounds contemplated herein include such salts. Examples of such salts include hydrochlorides, hydrobromides, sulfates, methanesulfonates, nitrates, maleates, acetates, citrates, fumarates, tartrates (e.g., (+)-tartrates, (-)-tartrates, or mixtures thereof including racemic mixtures), succinates, benzoates, and salts with amino acids such as glutamic acid. These salts can be prepared by methods known to those skilled in the art.

[0139] The neutral forms of the compounds are preferably regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents.

[0140] Pharmaceutically acceptable salts of the compounds above, where a basic or acidic group is present in the structure, are also included within the scope of compounds contemplated herein. When an acidic substituent is present, such as $\text{-NHSO}_3\text{H}$, -COOH and -P(O)(OH)_2 , there can be formed the ammonium, sodium, potassium, calcium salt, and the like, for use as the dosage form. Basic groups, such as amino or basic heteroaryl radicals, or pyridyl and acidic salts, such as hydrochloride, hydrobromide, acetate, maleate, palmoate, methanesulfonate, p-toluenesulfonate, and the like, can be used as the dosage form.

[0141] Also, in the embodiments in which R-COOH is present, pharmaceutically acceptable esters can be employed, e. g. , methyl, ethyl, tert-butyl, pivaloyloxymethyl, and the like, and those esters known in the art for modifying solubility or hydrolysis characteristics for use as sustained release or prodrug formulations.

A. Formulations

[0142] The compounds disclosed herein can be prepared and administered in a wide variety of ophthalmic, oral, parenteral, and topical dosage forms. The compounds described herein can be administered by eye drop. Also, compounds described herein can be administered by injection (e.g. intravenously, intramuscularly, intravitreally, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally). As such, compounds described herein can also be administered by intravitreal injection. Also, the compounds described herein can be administered by inhalation, for example, intranasally. Additionally, the compounds disclosed herein can be administered transdermally. It is also envisioned that multiple routes of administration (e.g., intramuscular, oral, ocular) can be used to administer the compounds disclosed herein.

[0143] In some embodiments, the compounds disclosed herein can be prepared in liquid pharmaceutical compositions for ocular administration. The composition for ocular use can contain one or more agents selected from the group of buffering agents, solubilizing agents, coloring agents, viscosity enhancing agents, and preservation agents in order to produce pharmaceutically elegant and convenient preparations.

[0144] In some embodiments, the composition for ocular use can contain preservatives for protection against microbiological contamination, including but not limited to benzalkonium

chloride and/or EDTA. Other possible preservatives include but are not limited to benzyl alcohol, methyl parabens, propyl parabens, and chlorobutanol. Preferably, a preservative, or combination of preservatives, will be employed to impart microbiological protection in addition to protection against oxidation of components.

[0145] In some embodiments, the compounds disclosed herein can be administered orally as tablets, aqueous or oily suspensions, lozenges, troches, powders, granules, emulsions, capsules, syrups or elixirs. The composition for oral use can contain one or more agents selected from the group of sweetening agents, flavoring agents, coloring agents and preserving agents in order to produce pharmaceutically elegant and palatable preparations. Accordingly, there are also provided pharmaceutical compositions comprising a pharmaceutically acceptable carrier or excipient and one or more compounds disclosed herein.

[0146] In some embodiments, tablets contain the acting ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients can be, for example, (1) inert diluents, such as calcium carbonate, lactose, calcium phosphate, carboxymethylcellulose, or sodium phosphate; (2) granulating and disintegrating agents, such as corn starch or alginic acid; (3) binding agents, such as starch, gelatin or acacia; and (4) lubricating agents, such as magnesium stearate, stearic acid or talc. These tablets can be uncoated or coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed.

[0147] For preparing pharmaceutical compositions from the compounds disclosed herein, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substance that can also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

[0148] A compound disclosed herein, in the form of a free compound or a pharmaceutically-acceptable pro-drug, metabolite, analogue, derivative, solvate or salt, can be administered, for in vivo application, parenterally by injection or by gradual perfusion over time. Administration can be intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, or transdermally. For *in vitro* studies the compounds can be

1397 added or dissolved in an appropriate biologically acceptable buffer and added to a cell or
1398 tissue.

1399 **[0149]** In powders, the carrier is a finely divided solid in a mixture with the finely divided
1400 active component. In tablets, the active component is mixed with the carrier having the
1401 necessary binding properties in suitable proportions and compacted in the shape and size
1402 desired.

1403 **[0150]** The powders and tablets preferably contain from 5% to 70% of the active
1404 compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar,
1405 lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium
1406 carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term
1407 “preparation” is intended to include the formulation of the active compound with
1408 encapsulating material as a carrier providing a capsule in which the active component with or
1409 without other carriers, is surrounded by a carrier, which is thus in association with it.
1410 Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and
1411 lozenges can be used as solid dosage forms suitable for oral administration.

1412 **[0151]** For preparing suppositories, a low melting wax, such as a mixture of fatty acid
1413 glycerides or cocoa butter, is first melted and the active component is dispersed
1414 homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into
1415 convenient sized molds, allowed to cool, and thereby to solidify.

1416 **[0152]** Liquid form preparations include solutions, suspensions, and emulsions, for
1417 example, water or water/propylene glycol solutions. For parenteral injection, liquid
1418 preparations can be formulated in solution in aqueous polyethylene glycol solution.

1419 **[0153]** When parenteral application is needed or desired, particularly suitable admixtures
1420 for the compounds disclosed herein are injectable, sterile solutions, preferably oily or
1421 aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. In
1422 particular, carriers for parenteral administration include aqueous solutions of dextrose, saline,
1423 pure water, ethanol, glycerol, propylene glycol, peanut oil, sesame oil, polyoxyethylene-block
1424 polymers, and the like. Ampoules are convenient unit dosages. The compounds disclosed
1425 herein can also be incorporated into liposomes or administered via transdermal pumps or
1426 patches. Pharmaceutical admixtures suitable for use in the pharmaceuticals compositions and
1427 methods disclosed herein include those described, for example, in PHARMACEUTICAL
1428 SCIENCES (17th Ed., Mack Pub. Co., Easton, PA) and WO 96/05309, the teachings of both of
1429 which are hereby incorporated by reference.

[0154] In some embodiments, preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives can also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, growth factors and inert gases and the like.

[0155] Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizers, and thickening agents as desired. Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

[0156] Also included are solid form preparations that are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations can contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

[0157] The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

[0158] The quantity of active component in a unit dose preparation can be varied or adjusted from 0.1 mg to 10000 mg, more typically 1.0 mg to 1000 mg, most typically 10 mg to 500 mg, according to the particular application and the potency of the active component. The composition can, if desired, also contain other compatible therapeutic agents.

[0159] Some compounds can have limited solubility in water and therefore can require a surfactant or other appropriate co-solvent in the composition. Such co-solvents include:

1463 Polysorbate 20, 60, and 80; Pluronic F-68, F-84, and P-103; cyclodextrin; and polyoxyl 35
1464 castor oil. Such co-solvents are typically employed at a level between about 0.01 % and about
1465 2% by weight.

1466 **[0160]** Viscosity greater than that of simple aqueous solutions can be desirable to decrease
1467 variability in dispensing the formulations, to decrease physical separation of components of a
1468 suspension or emulsion of formulation, and/or otherwise to improve the formulation. Such
1469 viscosity building agents include, for example, polyvinyl alcohol, polyvinyl pyrrolidone,
1470 methyl cellulose, hydroxy propyl methylcellulose, hydroxyethyl cellulose, carboxymethyl
1471 cellulose, hydroxy propyl cellulose, chondroitin sulfate and salts thereof, hyaluronic acid and
1472 salts thereof, and combinations of the foregoing. Such agents are typically employed at a
1473 level between about 0.01% and about 2% by weight.

1474 **[0161]** The compositions disclosed herein can additionally include components to provide
1475 sustained release and/or comfort. Such components include high molecular weight, anionic
1476 mucomimetic polymers, gelling polysaccharides, and finely-divided drug carrier substrates.
1477 These components are discussed in greater detail in U.S. Pat. Nos. 4,911,920; 5,403,841;
1478 5,212,162; and 4,861,760. The entire contents of these patents are incorporated herein by
1479 reference in their entirety for all purposes.

1480 **[0162]** By the present, there are provided methods for ameliorating wound healing and for
1481 mediating tissue repair (including but not limited to treatment of peripheral and coronary
1482 vascular disease). According to these methods, a subject having a wound or in need of tissue
1483 repair, is treated at the site of the wound or damaged tissue or treated systemically, with a
1484 compound disclosed herein in the form of a free compound or a pharmaceutically-acceptable
1485 prodrug, metabolite, analogue, derivative, solvate or salt.

1486 **[0163]** Generally, the terms “treating”, “treatment” and the like are used herein to mean
1487 affecting a subject, tissue or cell to obtain a desired pharmacologic and/or physiologic effect.
1488 The effect can be prophylactic in terms of completely or partially preventing a disease or
1489 disorder or sign or symptom thereof, and/or can be therapeutic in terms of a partial or
1490 complete cure for a disorder and/or adverse effect attributable to it. “Treating” as used herein
1491 covers any treatment of, or prevention of a disease or disorder in a vertebrate, a mammal,
1492 particularly a human, and includes: (a) preventing the disease or disorder from occurring in a
1493 subject that can be predisposed to the disease or disorder, but has not yet been diagnosed as
1494 having it; (b) inhibiting the disease or disorder, i.e. , arresting its development; or (c)

relieving or ameliorating the disease or disorder, i.e. , cause regression of the disease or disorder.

[0164] There are provided various pharmaceutical compositions useful for ameliorating certain diseases and disorders. The pharmaceutical compositions according to one embodiment are prepared by formulating a compound disclosed herein in the form of a free compound or a pharmaceutically-acceptable pro-drug, metabolite, analogue, derivative, solvate or salt, either alone or together with other pharmaceutical agents, suitable for administration to a subject using carriers, excipients and additives or auxiliaries. Frequently used carriers or auxiliaries include magnesium carbonate, titanium dioxide, lactose, mannitol and other sugars, talc, milk protein, gelatin, starch, vitamins, cellulose and its derivatives, animal and vegetable oils, polyethylene glycols and solvents, such as sterile water, alcohols, glycerol and polyhydric alcohols. Intravenous vehicles include fluid and nutrient replenishers.

[0165] Preservatives include antimicrobial, anti-oxidants, chelating agents and inert gases. Other pharmaceutically acceptable carriers include aqueous solutions, non-toxic excipients, including salts, preservatives, buffers and the like, as described, for instance, in Remington's Pharmaceutical Sciences, 15th ed. Easton: Mack Publishing Co. , 1405-1412, 1461-1487 (1975) and The National Formulary XIV., 14th ed. Washington: American Pharmaceutical Association (1975), the contents of which are hereby incorporated by reference. The pH and exact concentration of the various components of the pharmaceutical composition are adjusted according to routine skills in the art. *See e.g.*, Goodman and Gilman (eds.), 1990, THE PHARMACOLOGICAL BASIS FOR THERAPEUTICS (7th ed.).

[0166] The pharmaceutical compositions are preferably prepared and administered in dose units. Solid dose units are tablets, capsules and suppositories. For treatment of a subject, depending on activity of the compound, manner of administration, nature and severity of the disease or disorder, age and body weight of the subject, different daily doses can be used.

[0167] Under certain circumstances, however, higher or lower daily doses can be appropriate. The administration of the daily dose can be carried out both by single administration in the form of an individual dose unit or else several smaller dose units and also by multiple administrations of subdivided doses at specific intervals.

[0168] The pharmaceutical compositions contemplated herein can be administered locally or systemically in a therapeutically effective dose. Amounts effective for this use will, of course, depend on the severity of the disease or disorder and the weight and general state of the subject. Typically, dosages used *in vitro* can provide useful guidance in the amounts

useful for in situ administration of the pharmaceutical composition, and animal models can be used to determine effective dosages for treatment of particular disorders.

[0169] Various considerations are described, e. g. , in Langer, 1990, *Science*, **249**: 1527; Goodman and Gilman's (eds.), 1990, *Id.*, each of which is herein incorporated by reference and for all purposes. Dosages for parenteral administration of active pharmaceutical agents can be converted into corresponding dosages for oral administration by multiplying parenteral dosages by appropriate conversion factors. As to general applications, the parenteral dosage in mg/mL times 1.8 = the corresponding oral dosage in milligrams ("mg"). As to oncology applications, the parenteral dosage in mg/mL times 1.6 = the corresponding oral dosage in mg. An average adult weighs about 70 kg. See e.g., Miller-Keane, 1992, *ENCYCLOPEDIA & DICTIONARY OF MEDICINE, NURSING & ALLIED HEALTH*, 5th Ed., (W. B. Saunders Co.), pp. 1708 and 1651.

[0170] The method by which the compound disclosed herein can be administered for oral use would be, for example, in a hard gelatin capsule wherein the active ingredient is mixed with an inert solid diluent, or soft gelatin capsule, wherein the active ingredient is mixed with a co-solvent mixture, such as PEG 400 containing Tween-20. A compound disclosed herein can also be administered in the form of a sterile injectable aqueous or oleaginous solution or suspension. The compound can generally be administered intravenously or as an oral dose of 0.1 µg to 20 mg/kg given, for example, every 3 - 12 hours.

[0171] Formulations for oral use can be in the form of hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin. They can also be in the form of soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, such as peanut oil, liquid paraffin or olive oil.

[0172] Aqueous suspensions normally contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspension. Such excipients can be (1) suspending agent such as sodium carboxymethyl cellulose, methyl cellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; (2) dispersing or wetting agents which can be (a) naturally occurring phosphatide such as lecithin; (b) a condensation product of an alkylene oxide with a fatty acid, for example, polyoxyethylene stearate ; (c) a condensation product of ethylene oxide with a long chain aliphatic alcohol, for example, heptadecaethylenoxycetanol; (d) a condensation product of ethylene oxide with a partial ester derived from a fatty acid and hexitol such as

1561 polyoxyethylene sorbitol monooleate, or (e) a condensation product of ethylene oxide with a
1562 partial ester derived from fatty acids and hexitol anhydrides, for example polyoxyethylene
1563 sorbitan monooleate.

1564 **[0173]** The pharmaceutical compositions can be in the form of a sterile injectable aqueous
1565 or oleagenous suspension. This suspension can be formulated according to known methods
1566 using those suitable dispersing or wetting agents and suspending agents that have been
1567 mentioned above. The sterile injectable preparation can also a sterile injectable solution or
1568 suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a
1569 solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed
1570 are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed
1571 oils are conventionally employed as a solvent or suspending medium. For this purpose, any
1572 bland fixed oil can be employed including synthetic mono-or diglycerides. In addition, fatty
1573 acids such as oleic acid find use in the preparation of injectables.

1574 **[0174]** A compound disclosed herein can also be administered in the form of ophthalmic
1575 compositions applied topically to the eye, preferably in the form of eye drops. A compound
1576 disclosed herein can also be administered in the form of intravitreal injection.

1577 **[0175]** A compound disclosed herein can also be administered in the form of suppositories
1578 for rectal administration of the drug. These compositions can be prepared by mixing the drug
1579 with a suitable non-irritating excipient that is solid at ordinary temperature but liquid at the
1580 rectal temperature and will therefore melt in the rectum to release the drug. Such materials
1581 include cocoa butter and polyethylene glycols.

1582 **[0176]** The compounds disclosed herein as used in the methods disclosed herein can also be
1583 administered in the form of liposome delivery systems, such as small unilamellar vesicles,
1584 large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a
1585 variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

1586 **[0177]** For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing
1587 the compounds disclosed herein, are employed.

1588 **[0178]** In addition, some of the compounds disclosed herein can form solvates with water
1589 or common organic solvents. Such solvates are encompassed within the scope of the methods
1590 contemplated herein.

B. Effective Dosages

[0179] Pharmaceutical compositions provided herein include compositions wherein the active ingredient is contained in a therapeutically effective amount, i.e., in an amount effective to achieve its intended purpose. The actual amount effective for a particular application will depend, *inter alia*, on the condition being treated.

[0180] The dosage and frequency (single or multiple doses) of compound administered can vary depending upon a variety of factors, including route of administration; size, age, sex, health, body weight, body mass index, and diet of the recipient; nature and extent of symptoms of the disease being treated (e.g., the disease responsive to inhibition of thrombin, KLK1, and/or KLKB1); presence of other diseases or other health-related problems; kind of concurrent treatment; and complications from any disease or treatment regimen. Other therapeutic regimens or agents can be used in conjunction with the methods and compounds disclosed herein.

[0181] For any compound described herein, the therapeutically effective amount can be initially determined from a variety of techniques known in the art, e.g., biochemical characterization of inhibition of enzyme (thrombin, KLK1, or KLKB1), cell culture assays, and the like. Target concentrations will be those concentrations of active compound(s) that are capable of decreasing enzymatic activity as measured, for example, using the methods described.

[0182] Therapeutically effective amounts for use in humans can be determined from animal models. For example, a dose for humans can be formulated to achieve a concentration that has been found to be effective in animals. The dosage in humans can be adjusted by monitoring enzymatic inhibition and adjusting the dosage upwards or downwards, as described above.

[0183] Dosages can be varied depending upon the requirements of the patient and the compound being employed. The dose administered to a patient, in the context of the methods disclosed herein, should be sufficient to affect a beneficial therapeutic response in the patient over time. The size of the dose also will be determined by the existence, nature, and extent of any adverse side effects. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under circumstances is reached. In some embodiments of a method disclosed herein, the dosage range is 0.001% to 10% w/v. In some embodiments, the dosage range is 0.1% to 5% w/v.

[0184] Dosage amounts and intervals can be adjusted individually to provide levels of the administered compound effective for the particular clinical indication being treated. This will provide a therapeutic regimen that is commensurate with the severity of the individual's disease state.

[0185] Utilizing the teachings provided herein, an effective prophylactic or therapeutic treatment regimen can be planned that does not cause substantial toxicity and yet is entirely effective to treat the clinical symptoms demonstrated by the particular patient. This planning should involve the careful choice of active compound by considering factors such as compound potency, relative bioavailability, patient body weight, presence and severity of adverse side effects, preferred mode of administration, and the toxicity profile of the selected agent.

[0186] Accordingly, in some embodiments, dosage levels of the compounds disclosed herein as used in the present methods are of the order of e.g., about 0.1 mg to about 1 mg, about 1 mg to about 10 mg, about 0.5 mg to about 20 mg per kilogram body weight, an average adult weighing 70 kilograms, with a preferred dosage range between about 0.1 mg to about 20 mg per kilogram body weight per day (from about 7.0 mg to about 1.4 gm per patient per day). The amount of the compound disclosed herein that can be combined with the carrier materials to produce a single dosage will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for oral administration to humans can contain about 5 μ g to 1 g of a compound disclosed herein with an appropriate and convenient amount of carrier material that can vary from about 5 to 95 percent of the total composition. Dosage unit forms will generally contain between from about 0.1 mg to 500 mg of a compound disclosed herein.

[0187] It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

C. Toxicity


[0188] The ratio between toxicity and therapeutic effect for a particular compound is its therapeutic index and can be expressed as the ratio between LD₅₀ (the amount of compound lethal in 50% of the population) and ED₅₀ (the amount of compound effective in 50% of the population). Compounds that exhibit high therapeutic indices are preferred. Therapeutic

1657 index data obtained from *in vitro* assays, cell culture assays and/or animal studies can be used
1658 in formulating a range of dosages for use in humans. The dosage of such compounds
1659 preferably lies within a range of plasma concentrations that include the ED₅₀ with little or no
1660 toxicity. The dosage can vary within this range depending upon the dosage form employed
1661 and the route of administration utilized. See, *e.g.* Fingl *et al.*, *In: THE PHARMACOLOGICAL*
1662 *BASIS OF THERAPEUTICS*, Ch.1, p.1, 1975. The exact formulation, route of administration, and
1663 dosage can be chosen by the individual practitioner in view of the patient's condition and the
1664 particular method in which the compound is used. For *in vitro* formulations, the exact
1665 formulation and dosage can be chosen by the individual practitioner in view of the patient's
1666 condition and the particular method in which the compound is used.


VII. Examples

[0189] The examples below are meant to illustrate certain embodiments of the invention and not to limit the scope of the invention. Abbreviations used herein have their conventional meaning in the art, unless indicated otherwise. Specific abbreviations include the following: Å = Ångström; Ac₂O = acetic anhydride; AcOH = acetic acid; aq = aqueous; Bt = benzotriazole; BOC = *N*-*tert*-butoxycarbonyl; br = broad; *t*-BuOH = *tert*-butanol; °C = degree Celsius; d = doublet; DABCO = 1,4-diazabicyclo[2.2.2]octane; DCE = 1,2-dichloroethane; DCM = dichloromethane; dd = doublet of doublets; DIEA = diethylisopropylamine; DMAP = 4-dimethylaminopyridine; DMF = *N,N*-dimethylformamide; DMSO = dimethylsulfoxide; δ = chemical shift (given in ppm, unless otherwise indicated); EDCI = 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide; eq = equivalent; Et₂O = diethyl ether; Et₃N = triethylamine; EtOAc = ethyl acetate; EtOH = ethanol; g = gram; h (or hr) = hour; HOBt = hydroxybenzotriazole; HPLC = high performance liquid chromatography; Hz = Hertz; IC₅₀ = inhibitory concentration at 50% inhibition; *J* = coupling constant (given in Hz, unless otherwise indicated); LC = liquid chromatography; LHMDs = lithium hexamethyldisilazide; m = multiplet; M = molar; [M+H]⁺ = parent mass spectrum peak plus H⁺; MS = mass spectrum; ms = molecular sieves; MP = melting point; Me₂NH = dimethylamine; MeOH = methanol; mg = milligram; mL = milliliter; mM = millimolar; mmol = millimole; min = minute; μL = microliter; μM = micromolar; ng = nanogram; nM = nanomolar; NMR = nuclear magnetic resonance; ppm = parts per million; q = quartet; R_f = retention factor; RT = room temperature; s = singlet; t = triplet; TFA = trifluoroacetic acid; THF = tetrahydrofuran; TLC = thin layer chromatography.

[0190] **General Scheme I.** A synthetic scheme useful for synthesis of compounds described herein is disclosed in General Scheme I following, wherein the terms “R^x”, “R^y”, ... and “R^z” are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl, or other groups obvious to those skilled in the art.

1696 **General Scheme I**

1697

1698 *Example 1 - Preparation of Intermediate 1*1699 **[0191]** The synthesis of Intermediate 1 followed General Procedure 1 following.1700 **General Procedure 1**

1701


1702

1703 **[0192]** To a solution of 2-hydroxynicotinic acid (50.0 g, 0.359 moles, 1.0 eq.) in
 1704 dichloromethane (500 mL) at 0°C was added thionyl chloride (133.6 mL, 1.798 moles, 5.0
 1705 eq.) dropwise. After 30 min tetrahydrofuran (500 mL) was added and the reaction stirred for
 1706 14–15 hours at ambient temperature. The reaction mixture was cooled to 0°C, to it was added
 1707 methanol (150 mL) dropwise, and the mixture was stirred for a further 30 min at room
 1708 temperature. The reaction mixture was concentrated under reduced pressure to obtain a solid,
 1709 which was then neutralized with aqueous sodium bicarbonate (pH 7–8), and again
 1710 concentrated to obtain solid product. The solid was dissolved in methanol, filtered, and the
 1711 filtrate concentrated to give desired product 45.0 g, (yield; 81.8 %) m/z 153.99 $[M+H]^+$ ^1H
 1712 NMR (DMSO- d_6 , 400 MHz) δ 8.051-074 (1H, q), 7.661-7.682 (1H, q), 6.259-6.292 (1H, m),
 1713 3.734 (3H, s) ppm.

Example 2 - Preparation of Intermediate 2

[0193] The synthesis of Intermediate 2 followed the procedure of General Procedure 2 following.

General Procedure 2



Intermediate 1


Intermediate 2

[0194] To a cold (-78°C) solution of acetonitrile (8.18 mL, 0.156 moles, 1.2 eq.) in tetrahydrofuran (300 mL) was added $n\text{-BuLi}$ (2.5M in Hexane; 62.68 mL, 0.156 moles, 1.2 eq) dropwise over a period of 60 min. After addition, the reaction was stirred for another 60 min, then to it added methyl 2-oxo-1,2-dihydropyridine-3-carboxylate (**Intermediate 1**, 20.0 g, 130 mmol, 1.0 eq) portionwise to reaction mixture and maintained -78°C for 3 hrs. The reaction was quenched with water and washed with ethyl acetate. The aqueous layer was evaporated to obtain crude product, which was suspended in methanol and stirred for 30 min at room temperature. The solid was filtered through suction and dried over high vacuum to afford **Intermediate 2** (11.5g, 54%).

Example 3 - Preparation of Compound 1

[0195] The synthesis of **Compound 1** followed the procedure of General Procedure 3 following.

General Procedure 3



Intermediate 2

Compound 1

[0196] To a solution of **Intermediate 2** (20.0 g, 0.123 moles, 1.0 eq) in isopropanol (600 mL) and acetic acid (22.2 mL) was added hydrazine monohydrate (7.40 mL, 0.148 moles, 1.2 eq) dropwise and the reaction was heated at 85°C for 4-5 Hrs. After cooling, the reaction mixture was concentrated to give crude product, which was purified by column chromatography using neutral silica gel (60-120 mesh), eluting with 10-25% methanol in dichloromethane as gradient to give the desired product **Compound 1** 13.25 g (yield-61%)


1741 m/z 177.06 [M+H]⁺ 1H NMR (DMSO-d₆, 400 MHz) δ 11.831 (1H, s), 7.857-7.879 (1H, q),
1742 7.383-7.403 (1H, q), 6.303-6.336 (1H, m), 6.048 (1H, s) 4.633 (2H, s) ppm.

1743

1744 *Example 4 - Preparation of Compound 2*

1745 [0197] The synthesis of **Compound 2** followed the procedure of General Procedure 4
1746 following.

1747 **General Procedure 4**



Compound 1


Compound 2

1750 [0198] To a solution of **Compound 1** in dimethylformamide (100 mL) at 10-15°C was
1751 added acetic acid (11.2 mL) dropwise, followed by 5-chlorothiophene-2-carbaldehyde
1752 (9.15 g, 0.0624 moles, 1.1 eq) added portionwise. The reaction was stirred for 30-45 min at
1753 room temperature. Sodium cyanoborohydride (5.35 g, 0.0851 moles, 1.5 eq.) was added
1754 portionwise over a period of 45 min and reaction was stirred for 2 hours. After completion of
1755 reaction, the mixture was poured into ice cold water under stirring and the product was
1756 extracted with ethyl acetate. The organic layer was dried over sodium sulfate and
1757 concentrated under reduced pressure to obtain crude product, which was purified by column
1758 chromatography using neutral silica gel and product was eluted with 10-12% Methanol in
1759 dichloromethane as mobile phase to yield pure desired product **compound 2** (7.3 g, yield:
1760 42.7%) m/z[M+H]⁺ 307.10 1H NMR (DMSO-d₆, 400 MHz) δ 12.034 (1H, s), 11.815 (1H,
1761 s), 7.869-7.882 (1H, q), 7.404-7.415 (1H, d), 6.922-6.931 (1H, d), 6.862-6.871 (1H, d),
1762 6.314-6.331 (1H, d), 6.117 (1H, s), 5.867-5.898 (1H, t), 4.348-4.363 (2H, d) ppm.

1763

1764 *Example 5 - Preparation of Compound 3*


1765 [0199] The synthesis of **Compound 3** followed the procedure of General Procedure 5
1766 following.

1767 **General Procedure 5**1769 **Compound 2**1770 **Compound 3**

1771 **[0200]** To a cooled (0°C) solution of compound **2** in triethylamine (2.98 mL, 0.0215 moles,
 1772 3.0 eq.) and dichloromethane (40 mL) was added pivaloyl chloride (0.776 g, 0.00647 moles,
 1773 0.9 eq) dropwise over a period of 30 minutes. The reaction was stirred for 2-3 hours by
 1774 maintaining the temperature below 10°C. After completion, the reaction was diluted with ice
 1775 cold water under stirring and the product was extracted with dichloromethane. The organic
 1776 phase was dried over sodium sulfate and concentrated under reduced pressure. The resultant
 1777 crude product was purified by column chromatography using neutral silica gel, eluting with
 1778 5-8% methanol in dichloromethane to furnish pure desired product (compound **3**, 0.76 g,
 1779 yield: 43.6%) $m/z[M+H]^+$ 391.24 ¹H NMR (DMSO-d₆, 400 MHz) δ 11.250 (1H, s), 8.086-
 1780 8.109 (1H, q), 7.731-7.761 (1H, t), 7.484 (1H, s), 6.974-6.984 (1H, d), 6.934-6.944 (1H, d),
 1781 6.317-6.350 (1H, t), 6.213 (1H, s), 4.471-4.486 (2H, d), 1.47 (9H, s) ppm.

1783 *Example 6 - Preparation of Compound 4*

1784 **[0201]** The synthesis of Compound **4** followed the procedure of General Procedure 6
 1785 following.

1786 **General Procedure 6**1788 **Compound 2**1789 **Compound 4**


1789 **[0202]** To a solution of furan-3-carboxylic acid (0.338 g, 0.00301 moles, 1.2 eq) in
 1790 dimethylformamide (5.0 mL) was added EDCI.HCl (0.724 g, 0.00337 moles, 1.5 eq), DIEA

(0.811g, 0.00629 moles, 2.5 eq) and finally HOBt (0.074g, 0.00048 moles, 0.5 eq). The reaction mixture was stirred at room temperature for 30 min, followed by the addition of compound **2** (0.770 g, 0.00251 moles, 1.0 eq). The mixture was stirred at 14 hours at room temperature. After checking that the reaction had reached completion by LC-MS, the mixture was poured into ice cold water under stirring. The product was extracted with ethyl acetate. The organic phase was dried over sodium sulfate, concentrated under reduced pressure and purified by column chromatography using neutral silica gel (60-120 mesh), eluting with 15-25% ethyl acetate in *n*-hexane as gradient to give pure desired compound **4** (0.45 g, yield: 45%) m/z [M+H]⁺ 401.84 ¹H NMR (DMSO-*d*₆, 400 MHz) δ 11.923 (1H, s), 9.024-9.029 (1H, q), 8.274-8.297 (1H, q), 7.888-7.893 (1H, d), 7.833-7.884 (1H, q), 7.500-7.512 (1H, d), 7.085-7.091 (1H, q), 6.965-6.990 (2H, q), 6.313-6.347 (2H, t), 5.771 (1H, s), 4.445-4.560 (1H, d) ppm.

Example 7 - Preparation of Compound 5

[0203] The synthesis of Compound 5 followed the procedure of General Procedure 7 following.

General Procedure 7



Compound 4

Compound 5


[0204] To a solution of compound **4** (0.150 g, 0.375 mmol, 1.0 eq) in DMF (5.0 mL) was added anhydrous potassium carbonate (0.129 g, 0.937 mmol, 2.5 eq) and then stirred for 30 minutes at room temperature. 2-(Chloromethyl)thiophene (0.059 g, 0.45 mmol, 1.2 eq) was added to the reaction mixture and the reaction stirred for a further 2-3 hours at room temperature. The mixture was monitored by TLC and LCMS. After completion of reaction, the reaction mixture was poured into ice cold water under stirring and extracted with ethyl acetate. The organic phase was dried over sodium sulfate, concentrated under reduced pressure and purified by column chromatography using neutral silica gel. The product was eluted with 1-5% ethyl acetate as gradient in *n*-hexane to furnish compound **5** (0.036 g, yield: 19.3%) m/z [M+H]⁺ 497.23. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 9.020 (1H, s), 8.274-8.297

(1H, dd), 7.960-7.981 (1H, dd), 7.885-7.893 (1H, t), 7.833-7.864 (1H, t), 7.519-7.539 (1H, dd), 7.430-7.434 (1H, d), 7.117-7.133 (1H, dd), 7.087-7.091 (1H, d), 6.975-6.987 (1H, t), 6.380-6.427 (1H, t), 6.435 (1H, s), 5.189(2H, s), 4.550-4.565(2H, d) ppm.

Example 8 - Preparation of Compound 6

[0205] The synthesis of Compound 6 followed the procedure of General Procedure 8 following.

General Procedure 8



Compound 4

Compound 6


[0206] To a solution of compound **4** (0.150 g, 0.375 mmol, 1.0 eq.) in DMF (5.0 mL) was added cesium carbonate (0.304 g, 0.937 mmol, 2.5 eq.). The reaction mixture was stirred for 30 min at room temperature, followed by the addition of 4-(chloromethyl)pyridine hydrochloride (0.073 g, 0.45 mmol, 1.2 eq.). The reaction was stirred for 3-4 hours at 70°C. The reaction was monitored by TLC and LCMS. After completion of the reaction, the mixture was poured into ice cold water under stirring and extracted into ethyl acetate. The organic phase was dried over sodium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography using neutral silica gel, eluting with 40-55% ethyl acetate as gradient in *n*-hexane to furnish compound **6** (0.032 g, yield: 17.4%) *m/z* [M+H]⁺ 491.95. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 9.030 (1H, s), 8.541-8.526 (2H, d), 8.379-8.356 (1H, dd), 8.020-7.999 (1H, dd), 7.893-7.836 (2H, m), 7.210-7.195 (2H, d), 7.093-7.089 (1H, d), 6.968-6.948 (2H, t), 6.498-6.463 (1H, t), 6.294 (1H, s), 5.255 (2H, s), 4.542-4.526 (2H, d) ppm.

Example 9 - Preparation of Intermediate 3

[0207] General Scheme II. A synthetic scheme useful for synthesis of compounds described herein is disclosed in General Scheme II following, wherein the terms “R^x”, “R^y”, and “R^z” are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted


aryl, or substituted or unsubstituted heteroaryl, or other groups obvious to those skilled in the art.

General Scheme II



[0208] The synthesis of Intermediate 3 followed General Procedure 9 following.


General Procedure 9



Intermediate 3

[0209] A solution of nicotinic acid (9.9 g, 80.9 mmol) in water (30 mL) was added slowly portion-wise to a previously stirred mixture of aminoguanidine sulfate (10 g, 73.5 mmol) in concentrated H_2SO_4 (8.8 mL, 162 mmol), and the reaction mixture was stirred at 140 °C for 72 h. The reaction mixture was diluted with water (50 mL) and neutralized with saturated aqueous K_2CO_3 (30 mL), and the resultant solid was filtered. The residue was washed with water (2×30 mL), Et_2O (2×30 mL) and dried under vacuum to afford Intermediate 3 (4.6 g, 39%) as an off-white solid. ^1H NMR: ($\text{DMSO}-d_6$) δ 12.23 (s, 1H), 9.05 (s, 1H), 8.54 (d, $J = 2.8$ Hz, 1H), 8.17 (d, $J = 7.4$ Hz, 1H), 7.42-7.52 (m, 1H), 6.19 (s, 2H); MS: 162 $[\text{M} + \text{H}]^+$; MP: 234-236 °C; TLC: 20% MeOH/ NH_3 in CHCl_3 : R_f : 0.40.


Example 10 - Preparation of Intermediate 4



Intermediate 4

[0210] General Procedure 9 was followed to obtain Intermediate 4 (8.5 g, 46%). ¹H NMR: (DMSO-d₆) δ 8.60 (d, *J* = 4.4 Hz, 1H), 7.86-7.91 (m, 2H), 7.37 (br s, 1H), 5.79 (br s, 2H); MS: 162 [M + H]⁺; MP: 218-220 °C; TLC: 20% MeOH/NH₃ in CHCl₃; R_f: 0.40.

Example 11 - Preparation of Intermediate 5




Intermediate 5

[0211] General Procedure 9 was followed to obtain Intermediate 5 (12 g, 67%). ¹H NMR: (DMSO-d₆) δ 12.35 (br s, 1H), 8.59 (d, *J* = 5.5 Hz, 2H), 7.76-7.78 (m, 2H), 6.23 (s, 2H); MS: 162 [M + H]⁺; TLC: 20% MeOH/NH₃ in CHCl₃; R_f: 0.40.

Example 12 - Preparation of Intermediate 6

[0212] The synthesis of Intermediate 6 followed the procedure of General Procedure 10 following.

General Procedure 10



Intermediate 3


Intermediate 6

[0213] 4-Fluorobenzaldehyde (3.1 g, 24.8 mmol, 2 eq) and molecular sieves (4Å powder) were added to a solution of Intermediate 3 (2 g, 12.4 mmol) in EtOH (20 mL) at RT and refluxed for 8 h. Then was added a catalytic quantity of AcOH, NaCNBH₃ (1.6 g, 24.8 mmol, 2 eq) at 0 °C and with stirring for 15 h at RT. The solvent was distilled off, and the residue was dissolved in EtOAc (200 mL) and filtered through a Celite® pad to remove inorganic materials. The filtrate was washed with saturated aqueous NaHCO₃ (2 × 20 mL), water (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo*. The resultant compound was purified by column chromatography over silica gel (100-200 mesh) by using a solvent gradient of 0-10% MeOH-CHCl₃ as the eluent to afford Intermediate 6 (1.7 g, 51%). ¹H NMR: (DMSO-d₆) δ 12.50 (s, 1H), 9.06 (d, *J* = 1.4 Hz, 1H), 8.53-8.55 (m, 1H), 8.17-8.20 (m, 1H), 7.33-7.45 (m, 4H), 7.12-7.19 (m, 2H), 4.40 (d, *J* = 6.4 Hz, 2H); MS: 270 [M + H]⁺; MP: 185-186 °C; TLC: 10% MeOH in CHCl₃; R_f: 0.25.

1897

1898 *Example 13 - Preparation of Intermediate 7*

1899




1900

Intermediate 4**Intermediate 7**

1901 **[0214]** General Procedure 10 was followed to obtain Intermediate 7 (2.8 g, 60%). MS: 252
 1902 $[M + H]^+$; MP: 226-228 °C; TLC: 10% MeOH in CHCl₃; R_f: 0.30.

1903 *Example 14 - Preparation of Intermediate 8*

1904




1905

Intermediate 4**Intermediate 8**

1906 **[0215]** General Procedure 10 was followed to obtain Intermediate 8 (1.6 g, 48%). ¹H NMR:
 1907 (DMSO-d₆) δ 13.15 (br s, 1H), 8.60 (d, *J* = 4.0 Hz, 1H), 7.86-7.93 (m, 2H), 7.30-7.42 (m,
 1908 3H), 7.02-7.15 (m, 2H), 6.84 (br s, 1H), 4.37 (d, *J* = 6.2 Hz, 2H); MS: 270 $[M + H]^+$; MP:
 1909 219-220 °C; TLC: 10% MeOH in CHCl₃; R_f: 0.25.

1910 *Example 15 - Preparation of Intermediate 9*

1911



1912

Intermediate 5**Intermediate 9**


1913 **[0216]** General Procedure 10 was followed to obtain Intermediate 9 (1.4 g, 42%). MS: 270
 1914 $[M + H]^+$; TLC: 10% MeOH in CHCl₃; R_f: 0.25.

1915

1916 *Example 16 - Preparation of Compound 7*

1917 **[0217]** The synthesis of Compound 7 followed General Procedure 11 following.

General Procedure 11




Intermediate 6

Compound 7

[0218] Propionyl chloride (39 μ L, 0.44 mmol, 1.2 eq) was added to a solution of Intermediate 6 (100 mg, 0.37 mmol) in triethylamine (3 mL) at RT and stirred for 5 h. The reaction mixture was diluted with water (5 mL) and extracted with EtOAc (20 mL). The organic layer washed with water (2×5 mL), saturated aqueous NaHCO_3 (5 mL), brine (5 mL), dried over Na_2SO_4 , filtered and concentrated *in vacuo*. The crude compound was purified by column chromatography over silica gel (100-200 mesh) by using a gradient mixture of 0-30% EtOAc-hexane as the eluent to afford Compound 7 (40 mg, 33%). ^1H NMR: (DMSO-d_6) δ 9.14 (d, $J = 1.8$ Hz, 1H), 8.66-8.67 (m, 1H), 8.28-8.34 (m, 2H), 7.47-7.53 (m, 3H), 7.13-7.17 (m, 2H), 4.63 (d, $J = 6.2$ Hz, 2H), 3.05 (q, $J = 7.5$ Hz, 2H), 1.16 (t, $J = 7.5$ Hz, 3H); MS: 326 $[\text{M} + \text{H}]^+$; TLC: 50% EtOAc in hexane: R_f : 0.60.


Example 17 - Preparation of Compound 8



Intermediate 7

Compound 8


[0219] General Procedure 11 was followed to obtain Compound 8 (48 mg, 35%). ^1H NMR: (DMSO-d_6) δ 8.71 (d, $J = 4.0$ Hz, 1H), 8.46 (br s, 1H), 8.13-8.23 (m, 3H), 7.92-7.96 (m, 1H), 7.24-7.52 (m, 6H), 6.88-6.89 (m, 1H), 4.74 (d, $J = 6.2$ Hz, 2H); MS: 346 $[\text{M} + \text{H}]^+$; MP: 143-145 $^\circ\text{C}$; TLC: 50% EtOAc in hexane: R_f : 0.60.

1938 *Example 18 - Preparation of Compound 9*

1939

1940 **Intermediate 7****Compound 9**


1941 **[0220]** General Procedure 11 was followed to obtain Compound 9 (25 mg, 16%). ¹H NMR:
 1942 (DMSO-d₆) δ 8.65 (d, *J* = 4.0 Hz, 1H), 8.26 (br s, 1H), 8.03 (d, *J* = 8.1 Hz, 1H), 7.90 (t, *J* =
 1943 7.7 Hz, 1H), 7.19-7.48 (m, 11H), 4.67 (d, *J* = 6.0 Hz, 2H), 3.30-3.41 (m, 2H), 2.99-3.03 (m,
 1944 2H); MS: 384 [M + H]⁺; MP: 118-120 °C; TLC: 50% EtOAc in hexane: R_f: 0.40.

1945 *Example 19 - Preparation of Compound 10*

1946

1947 **Intermediate 8****Compound 10**

1948 **[0221]** General Procedure 11 was followed to obtain Compound 10 (40 mg, 28%). ¹H
 1949 NMR: (DMSO-d₆) δ 8.72 (d, *J* = 4.6 Hz, 1H), 8.47-8.54 (m, 2H), 8.12-8.23 (m, 2H), 7.94-
 1950 7.98 (m, 1H), 7.48-7.52 (m, 3H), 7.34-7.36 (m, 1H), 7.16 (t, *J* = 9.0 Hz, 2H), 4.71 (d, *J* = 6.1
 1951 Hz, 2H); MS: 380 [M + H]⁺; MP: 159-160 °C; TLC: 50% EtOAc in hexane: R_f: 0.60.

1952 *Example 20 - Preparation of Compound 11*

1953


1954 **Intermediate 9****Compound 11**

[0222] General Procedure 11 was followed to obtain Compound 11 (20 mg, 14%). ¹H NMR: (DMSO-d₆) δ 9.19 (d, *J* = 1.3 Hz, 1H), 8.63-8.73 (m, 3H), 8.00 (d, *J* = 5.7 Hz, 2H), 7.72-7.88 (m, 2H), 7.50-7.54 (m, 2H), 7.17 (t, *J* = 8.8 Hz, 2H), 4.70 (d, *J* = 6.2 Hz, 2H); MS: 380 [M + H]⁺; MP: 187-188 °C; TLC: 50% EtOAc in hexane: R_f: 0.60.

Example 21 - Preparation of Compound 12

[0223] The synthesis of Compound 12 followed General Procedure 12 following.

General Procedure 12




Intermediate 6

Compound 12

[0224] A solution of Intermediate 6 (100 mg, 0.37 mmol) in dry DMF (2 mL) was added to a solution of morpholinecarbonyl chloride (86 μL, 0.74 mmol, 2 eq), DABCO (124 mg, 1.11 mmol, 3 eq) in DMF (3 mL) at RT and stirred for 2 h. The reaction mixture was diluted with water (10 mL) and extracted with EtOAc (30 mL). The organic layer washed with water (2 × 5 mL), saturated aqueous NaHCO₃ (2 × 5 mL), brine (10 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo* to get a crude residue. The crude compound was purified by column chromatography over silica gel (100-200 mesh) by using a gradient mixture of 0-50% EtOAc-hexane as the eluent to afford Compound 12 (33 mg, 23%). ¹H NMR: (DMSO-d₆) δ 9.11 (s, 1H), 8.64 (d, *J* = 4.8 Hz, 1H), 8.25 (d, *J* = 7.9 Hz, 1H), 7.90 (s, 1H), 7.46-7.52 (m, 3H), 7.16 (t, *J* = 8.8, 2H), 4.59 (d, *J* = 6.2 Hz, 2H), 3.70-3.99 (m, 8H); MS: 383 [M + H]⁺; TLC: 50% EtOAc in hexane: R_f: 0.40.


Example 22 - Preparation of Compound 13

[0225] A useful scheme for the preparation of compounds of the type of Compound 13 is provided in Scheme 1 following.

1978 **Scheme 1**


1979

1980 **[0226]** A detailed description of the preparation of Intermediates 10-13 and Cmpd 24
 1981 follows.

1982 **Preparation of Intermediate 10**

1983


1984 **[0227]** A solution of cyanogen bromide (1.3 g, 12.6 mmol) in acetone (5 mL) was added
 1985 portion-wise slowly to a mixture of benzotriazole (3 g, 25.2 mmol, 2 eq) in EtOH (50 mL)
 1986 followed by 10% aqueous NaOH (6 mL, 12.6 mmol, 1 eq) at 0 °C. The reaction mixture was
 1987 then stirred at RT for 30 min. Solid formation was observed. The solid was filtered and
 1988 washed with cold EtOH. The resulting material was recrystallized from benzene to afford
 1989 Intermediate 10 (2.2 g, 33%) as a white solid. ¹H NMR: (DMSO-d₆) δ 11.76 (s, 1H), 8.29-
 1990 8.39 (m, 2H), 7.86-8.09 (m, 2H), 7.44-7.72 (m, 4H), MS: 264 [M + H]⁺; TLC: 30% EtOAc in
 1991 hexane: R_f: 0.50.

1992 **Preparation of Intermediate 11**

1993

[0228] Dimethylamine (1.59 mL, 7.60 mmol, 1 eq) was added to Intermediate 10 (2 g, 7.60 mmol) in THF (30 mL) at RT and the resulting mixture was allowed to stir for 24 h. The solvent was evaporated and the residue was dissolved in DCM (100 mL). The organic layer was washed with 10% Na₂CO₃ (3 × 5 mL), brine (10 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford Intermediate 11 (1.2 g, 71%) as a light yellow liquid which was used without additional purification. ¹H NMR: (DMSO-d₆) δ 8.17 (d, *J* = 8.4 Hz, 1H), 7.65-7.80 (m, 3H), 7.49-7.53 (m, 1H), 2.87 (s, 6H); MS: 190 [M + H]⁺; TLC: 30% EtOAc in hexane: R_f: 0.30.

Preparation of Intermediate 12




Intermediate 11

Intermediate 12

[0229] Oxalyl chloride (2 mL, 23.3 mmol, 1.4 eq) was added to a solution of nicotinic acid (2 g, 16.3 mmol) in DCM followed by catalytic amount of DMF (0.5 mL) at 0 °C and stirred for 5 h at RT. The solvent was then evaporated to afford nicotinic acid chloride as a yellow solid. Nicotinic acid chloride (1.1 g, 7.93 mmol, 1.5 eq) was then added to a solution of Intermediate 11 (1 g, 5.29 mmol) in CHCl₃ (30 mL) followed by Et₃N (0.7 mL, 5.29 mmol, 1 eq) at 0 °C. The reaction mixture was allowed to warm to RT for stir for 18 h. The mixture was then diluted with CHCl₃ (20 mL). The organic layer was washed with water (10 mL), brine (10 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo*. The resulting compound was purified by column chromatography over silica gel (100-200 mesh) by using a gradient mixture of 0-50% EtOAc-hexane as the eluent to afford Intermediate 12 (900 mg, 60%) as a white solid. MS: 295 [M + H]⁺; TLC: 50% EtOAc in DCM: R_f: 0.40.

Preparation of Intermediate 13




Intermediate 12

Intermediate 13

[0230] Hydrazine hydrate (5 mL) was added solution of Intermediate 12 (900 mg, 25.2 mmol) in chloroform (20 mL) at RT and the resulting mixture was allowed to stir for 24 h. The mixture was diluted with excess CHCl_3 (20 mL). The organic layer was then washed with water (15 mL), brine (10 mL), dried over Na_2SO_4 , filtered and concentrated *in vacuo*. The crude residue was partially purified by column chromatography over silica gel (100-200 mesh) by using a gradient mixture of 0-50% EtOAc-hexane as the eluent to afford Intermediate 13 (150 mg) as a thick brown mass. MS: 190 $[\text{M} + \text{H}]^+$; TLC: 10% MeOH in CHCl_3 ; R_f : 0.30.

Preparation of Compound 13




Intermediate 13

Compound 13

[0231] General Procedure 11 was followed to obtain Compound 13 (13 mg, 6%). ^1H NMR: ($\text{DMSO}-d_6$) δ 9.15 (s, 1H), 8.68 (d, $J = 3.5$ Hz, 1H), 8.31 (d, $J = 7.9$ Hz, 1H), 7.53 (dd, $J = 7.9, 4.8$ Hz, 1H), 3.04-3.14 (m, 8H), 1.15 (t, $J = 7.3$ Hz, 3H); MS: 246 $[\text{M} + \text{H}]^+$; TLC: 50% EtOAc in DCM; R_f : 0.50.


Example 23 - Preparation of Compound 14

[0232] A general chemical scheme for the formation of compounds of the type of Compound 14 is provided in General Scheme III following, wherein the terms " R^x ", " R^y ", and " R^z " are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl, or other groups obvious to those skilled in the art. **General Scheme III**




[0233] A detailed description of the preparation of Intermediates 14, 15 and Compound 14 follows.

Preparation of Intermediate 14



Intermediate 14


[0234] Oxalyl chloride (5.4 mL, 61.0 mmol, 1.5 eq) and DMF (3 mL) was added sequentially to a solution of nicotinic acid (5 g, 40.7 mmol) in dry DCM (300 mL) at RT. The reaction mixture was allowed to stir at RT for 2 h. The solvent was removed and co-distilled with dry toluene (2 × 50 mL) and to afford 5 g of crude nicotinic acid chloride (5 g, 35.5 mmol). This material was added slowly portion-wise to a solution of thiosemicarbazide (5 g, 54.9 mmol, 1.5 eq) in pyridine (50 mL) at 0 °C over a period of 1 h and then allowed to stir at RT for 14 h. The reaction mixture was neutralized with saturated aqueous NaHCO₃ (30 mL) and extracted with *t*-BuOH (3 × 100 mL) and dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude residue was dissolved in water (20 mL) along with 10% aqueous KOH (50 mL) and the resulting mixture was allowed to stir at 100 °C for 3 h. The reaction mixture was then cooled to 0 °C and neutralized with 10% aqueous AcOH (60 mL), extracted with EtOAc (2 × 150 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford crude Intermediate 14 (1.2 g) as an off-white solid. MS: 179 [M + H]⁺; TLC: 20% MeOH/NH₃ in CHCl₃: R_f: 0.30.

2060 **Preparation of Intermediate 15**

2061

2062 **Intermediate 14****Intermediate 15**

2063 **[0235]** 4-Fluorobenzyl bromide (0.12 mL, 1.01 mmol, 0.6 eq) was added to a solution of
 2064 Intermediate 14 (300 mg, 1.68 mmol) in water (5 mL) and THF (15 mL) at -10 °C and the
 2065 reaction mixture was allowed to stir at -10 °C for 8 h. The solvent was removed and the
 2066 residue was diluted with water (10 mL) and extracted with EtOAc (50 mL). The organic layer
 2067 was washed with water (15 mL), saturated aqueous NaHCO₃ (10 mL), brine, dried over
 2068 Na₂SO₄, filtered and concentrated *in vacuo*. The crude compound was purified by column
 2069 chromatography over silica gel (100-200 mesh) by using a solvent gradient mixture of 0-10%
 2070 MeOH-CHCl₃ as the eluent to afford Intermediate 15 (110 mg, 23%) as an off-white solid.
 2071 MS: 287 [M + H]⁺; TLC: EtOAc: R_f: 0.40.

2072 **Preparation of Compound 14**


2073

2074 **Intermediate 15****Compound 14**

2075 **[0236]** General Procedure 11 was followed to obtain Compound 14 (20 mg, 30%). ¹H
 2076 NMR: (DMSO-d₆) δ 9.13 (s, 1H), 8.71 (d, *J* = 4.0 Hz, 1H), 8.26 (d, *J* = 7.9 Hz, 1H), 7.53-
 2077 7.67 (m, 5H), 7.09-7.25 (m, 4H), 4.64 (s, 2H), 3.75 (s, 3H); MS: 421 [M + H]⁺; MP: 108-112
 2078 °C; TLC: 30% EtOAc in hexane: R_f: 0.40.

2079 *Example 24 - Preparation of Intermediate 16*

2080




2081

2082

Intermediate 16


[0237] 2-Thiophene carboxylic acid chloride (6.5 mL, 60.4 mmol) was added slowly portionwise to a solution of thiosemicarbazide (5 g, 54.9 mmol, 1.1 eq) in pyridine (50 mL) at 0 °C over a period of 1 h and then allowed to stir at RT for 14 h. The reaction mixture was neutralized with saturated aqueous NaHCO₃ (50 mL) and extracted with *t*-BuOH (3 × 100 mL) and dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude residue was dissolved in water (30 mL) along with 10% aqueous KOH (60 mL) and the resulting mixture was allowed to stir at 100 °C for 3 h. The reaction mixture was then cooled to 0 °C and neutralized with 10% aqueous AcOH, extracted with EtOAc (2 × 150 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford crude Intermediate 16 (1.2 g) as an off-white solid. MS: 184 [M + H]⁺; TLC: 10% MeOH/NH₃ in CHCl₃: R_f: 0.60.

Example 25 - Preparation of Intermediate 17



[0238] A solution of methyl iodide (65 μL, 1.04 mmol, 1.6 eq) in EtOH (2 mL) was added to a solution of Intermediate 16 (120 mg, 0.66 mmol) in 1M aqueous NaOH (3 mL) at RT and the resulting mixture was allowed to stir for 3 h. The reaction mixture was then neutralized with 10% aqueous AcOH (5 mL) and extracted with EtOAc (30 mL). The organic layer was washed with water (10 mL), saturated aqueous NaHCO₃ (5 mL), brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude compound was purified by column chromatography over silica gel (100-200 mesh) by using a solvent gradient mixture of 0-10% MeOH-CHCl₃ as the eluent to afford Intermediate 17 (90 mg, 70%) as an off-white solid. ¹H NMR: (DMSO-*d*₆) δ 14.19 (br s, 1H), 7.62-7.67 (m, 2H), 7.16-7.18 (m, 1H), 2.60 (s, 3H); MS: 198 [M + H]⁺; TLC: 50% EtOAc in hexane: R_f: 0.50.

Example 26 - Preparation of Compound 15




[0239] General Procedure 11 was followed to obtain Cmpd 29 (30 mg, 29%). ¹H NMR: (DMSO-d₆) δ 7.72 (d, *J* = 4.8 Hz, 1H), 7.56-7.65 (m, 3H), 7.25 (d, *J* = 8.8 Hz, 1H), 7.09-7.24 (m, 2H), 3.77 (s, 3H), 2.73 (s, 3H); MS: 332 [M + H]⁺; MP: 165-167 °C; TLC: 30% EtOAc in hexane: R_f: 0.40.

Example 27 - Preparation of Compound 16

[0240] **General Scheme IV.** A synthetic scheme useful for synthesis of compounds described herein including Compound 16 is disclosed in General Scheme IV following, wherein the terms “R^x”, “R^y”, and “R^z” are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl, or other groups obvious to those skilled in the art.

General Scheme IV




[0241] A description of the synthesis of Intermediates 19-24 and Compound 16 follows. Synthesis of Intermediate 19 followed General Procedure 13 following.

Preparation of Intermediate 19 [General Procedure 13]

[0242] General Procedure 13 was followed in the preparation of Intermediate 19.


General Procedure 13



Intermediate 19

[0243] Thionyl chloride (3.55 mL, 48.4 mmol, 3 eq) was added dropwise to a solution of pyrimidine-4-carboxylic acid (2 g, 16.1 mmol) in EtOH (15 mL) and the resulting mixture was heated to reflux for 14 h. The mixture was then cooled to RT and made alkaline with saturated aqueous NaHCO₃ to pH 8. The basic solution was then extracted with EtOAc (4 × 50 mL). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford Intermediate 19 (1.7 g, 77%). ¹H NMR: (DMSO-d₆) δ 9.40 (d, *J* = 1.0 Hz, 1H), 9.10 (d, *J* = 5.1 Hz, 1H), 8.05 (dd, *J* = 5.1, 1.3 Hz, 1H), 4.39 (q, *J* = 7.1 Hz, 2H), 1.35 (t, *J* = 7.1 Hz, 3H); MS: 153 [M + H]⁺; TLC: 40% hexane in EtOAc: R_f: 0.40.

Preparation of Intermediate 20




Intermediate 20

[0244] General Procedure 13 was followed to obtain crude Intermediate 20 (950 mg, 86%). ¹H NMR: (DMSO-d₆) δ 9.43 (s, 1H), 9.26 (s, 2H), 4.39 (q, *J* = 7.1 Hz, 2H), 1.35 (t, *J* = 7.1 Hz, 3H); TLC: 40% EtOAc in hexane: R_f: 0.50.

Preparation of Intermediate 21 [General Procedure 14]

[0245] General Procedure 14 was followed in the preparation of Intermediate 21.


General Procedure 14




Intermediate 19

Intermediate 21


[0246] Intermediate 19 (1.6 g, 10.5 mmol) was added dropwise to a vigorously stirring mixture of aminoguanidine sulfate (10.3 g, 42.1 mmol, 4 eq) in freshly prepared NaOMe (using 968 mg, 42.1 mmol of Na in 28 mL of dry MeOH) at 0 °C. The resulting mixture was heated to reflux for 20 h. The mixture was then cooled to RT, carefully poured over ice cold water (20 mL) and concentrated *in vacuo*. The crude residue was purified over neutral alumina using 4-10% MeOH-CHCl₃ as the eluent to give Intermediate 21 (500 mg, 26%). MS: 163 [M + H]⁺; TLC: 20% MeOH in CHCl₃: R_f: 0.20.

2159 **Preparation of Intermediate 22****Intermediate 20****Intermediate 22**

2162 [0247] General Procedure 14 was followed to obtain Intermediate 22 (500 mg, 45%). ¹H
 2163 NMR: (DMSO-d₆) δ 12.44 (br s, 1H), 9.17-9.18 (m, 3H), 6.32 (s, 2H); TLC: 20% MeOH in
 2164 CHCl₃; R_f: 0.20.

2165 **Preparation of Intermediate 23****Intermediate 21****Intermediate 23**


2168 [0248] General Procedure 10 was followed to obtain Intermediate 23 (210 mg, 34%). ¹H
 2169 NMR: (DMSO-d₆) δ 12.80 (s, 1H), 9.18 (s, 1H), 8.83 (s, 1H), 7.92 (d, *J* = 4.4 Hz, 1H), 7.25-
 2170 7.40 (m, 5H), 4.44 (d, *J* = 5.7 Hz, 2H); TLC: EtOAc: R_f: 0.30.

2171 **Preparation of Intermediate 24****Intermediate 22****Intermediate 24**

2174 [0249] General Procedure 10 was followed to obtain Intermediate 24 (160 mg, 20%). MS:
 2175 253 [M + H]⁺; TLC: EtOAc: R_f: 0.30.

2176 **Preparation of Compound 16 [General Procedure 11]**


2177 [0250] General Procedure 11 was followed in the preparation of Compound 16.

2178 **General Procedure 11**

2179

2180 **Intermediate 23****Compound 16**

2181 **[0251]** 2-Methoxybenzoyl chloride (72 μ L, 0.54 mmol, 2 eq) was added to a solution of
 2182 Intermediate 23 (70 mg, 0.27 mmol) in Et₃N (0.18 mL, 1.35 mmol) and DCM (3 mL) at 0 °C.
 2183 The resulting mixture was allowed to stir at RT for 2 h. The reaction mixture was then diluted
 2184 with water (5 mL) and extracted with DCM (3 $\times$ 15 mL). The combined organic layers were
 2185 washed with saturated aqueous NaHCO₃ (10 mL), water (2 $\times$ 5 mL), brine (15 mL), dried
 2186 over Na₂SO₄, filtered and concentrated *in vacuo*. The crude material was purified by column
 2187 chromatography over silica gel (100-200 mesh) using a gradient mixture of 0-70% EtOAc-
 2188 hexane as the eluent to afford Compound 16 (45 mg, 29%). ¹H NMR: (DMSO-d₆) δ 9.21 (s,
 2189 1H), 8.90 (d, *J* = 5.1 Hz, 1H), 8.59 (t, *J* = 6.0 Hz, 1H), 7.93 (d, *J* = 5.1 Hz, 1H), 7.08-7.60 (m,
 2190 10H), 4.72 (d, *J* = 5.7 Hz, 2H), 3.77 (s, 3H); MS: 387 [M + H]⁺; MP: 192-195 °C; TLC: 40%
 2191 hexane in EtOAc: R_f: 0.30.

2192 *Example 28 - Preparation of Compound 17*

2193


2194 **Intermediate 24****Compound 17**

2195 **[0252]** General Procedure 11 was followed by preparative HPLC purification to obtain
 2196 Compound 17 (30 mg, 16%). ¹H NMR: (DMSO-d₆) δ 9.26 (s, 1H), 9.11 (s, 2H), 8.64 (t, *J* =
 2197 6.3 Hz, 1H), 7.07-7.60 (m, 9H), 4.71 (d, *J* = 6.3 Hz, 2H), 3.78 (s, 3H); MS: 387 [M + H]⁺;
 2198 MP: 154-157 °C; TLC: 40% EtOAc in hexane: R_f: 0.20.


2199 *Example 29 - Preparation of Compound 18*

2200 **[0253] General Scheme V.** A synthetic scheme useful for synthesis of compounds
 2201 described herein including Compound 18 is disclosed in General Scheme V following,

wherein the terms “R^x”, “R^y”, and “R^z” are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl, or other groups obvious to those skilled in the art.

General Scheme V


[0254] A description of the synthesis of Intermediates 25, 26 and Compound 18 follows.

Preparation of Intermediate 25**Intermediate 25**

[0255] Oxalyl chloride (2.36 mL, 24.2 mmol, 1.5 eq) and a catalytic quantity of DMF were added to a solution of pyrimidine-2-carboxylic acid (2 g, 16.1 mmol) in dry DCM (30 mL) at 0 °C. The resulting mixture was allowed to warm to RT and stir for 3 h. The volatiles were removed *in vacuo* and the residue was thoroughly dried to afford pyrimidine-2-carboxylic acid chloride (2.1 g, 14.8 mmol) as a black solid. The crude material was added portion-wise to a solution of aminoguanidine sulfate (5.5 g, 22.2 mmol, 1.5 eq) in pyridine (20 mL) at 0 °C. The resulting mixture was allowed to warm to RT and stir for 14 h. The mixture was then neutralized with saturated aqueous NaHCO₃, extracted with *t*-BuOH (5 × 50 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude material was dissolved in water (45 mL) and the resulting solution was heated to 100 °C for 24 h. The reaction mixture was

then cooled to RT, extracted with *t*-BuOH (5 × 30 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo* to afford Intermediate 25 (650 mg, 25 %) as off-white solid. TLC: 30% MeOH in CHCl₃; R_f: 0.20.

Preparation of Intermediate 26




Intermediate 25

Intermediate 26

[0256] General Procedure 10 was followed to obtain Intermediate 26 (120 mg, 17%). MS: 253 [M + H]⁺; TLC: EtOAc: R_f: 0.30.

Preparation of Compound 18





Intermediate 26

Compound 18

[0257] General Procedure 11 was followed to obtain Compound 18 (32 mg, 21%). ¹H NMR: (DMSO-d₆) δ 8.86 (d, *J* = 5.1 Hz, 2H), 8.44 (t, *J* = 6.0 Hz, 1H), 7.08-7.59 (m, 10H), 4.73 (d, *J* = 6.3 Hz, 2H), 3.77 (s, 3H); MS: 387 [M + H]⁺; MP: 203-205 °C; TLC: 40% hexane in EtOAc: R_f: 0.40.

Example 30 - Preparation of Compound 19

[0258] **General Scheme VI.** A synthetic scheme useful for synthesis of compounds described herein including Compound 19 is disclosed in General Scheme VI following, wherein the terms “R^x”, “R^y”, and “R^z” are independently hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl, or other groups obvious to those skilled in the art.

2246 **General Scheme VI**2248 **Preparation of Intermediate 27**2249 **[0259]** General Procedure 10 was followed in the preparation of Intermediate 27.2250 **General Procedure 10**


2251

2252

Intermediate 27

2253 **[0260]** 4-Fluorobenzaldehyde (0.54 mL, 5.03 mmol, 2 eq) and molecular sieves (4 Å
 2254 powder) were added to a solution of 3-amino-5-phenylpyrazole (400 mg, 2.51 mmol) in
 2255 EtOH (20 mL) at RT and the resulting mixture was heated to reflux. After 8 h, the reaction
 2256 mixture was cooled to 0 °C and AcOH (0.4 mL) and NaCNBH₃ (316 mg, 5.03 mmol, 2 eq)
 2257 were added. The mixture was then allowed to warm to RT and stir for 15 h. The solvent was
 2258 evaporated and the residue was dissolved in EtOAc (100 mL) and filtered through a Celite
 2259 pad to remove inorganic materials. The filtrate was then washed with saturated aqueous
 2260 NaHCO₃ (2 × 20 mL), water (20 mL), brine (20 mL), dried over Na₂SO₄, filtered and
 2261 concentrated *in vacuo*. The crude material was purified by column chromatography over
 2262 silica gel (100-200 mesh) by using a solvent gradient of 0-50% EtOAc-petroleum ether as the
 2263 eluent to afford Intermediate 27 (240 mg, 36%) as an off white solid. MS: 268 [M + H]⁺;
 2264 TLC: EtOAc: R_f: 0.60.

2265 **Preparation of Compound 19**2266 **[0261]** General Procedure 15 was followed in the preparation of Compound 19.

2267 **General Procedure 15**


2268

2269 **Intermediate 27****Compound 19**

2270 **[0262]** Pivaloyl chloride (32 μ L, 0.26 mmol, 1.2 eq) was added to a solution of
 2271 Intermediate 27 (60 mg, 0.22 mmol) in triethylamine (3 mL) at RT and stirred for 3 h. The
 2272 reaction mixture was diluted with water (5 mL) and extracted with EtOAc (20 mL). The
 2273 organic layer was washed with water (2×5 mL), saturated aqueous NaHCO₃ (5 mL), brine
 2274 (5 mL), dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude compound was
 2275 purified by column chromatography over silica gel (100-200 mesh) by using a gradient
 2276 mixture of 0-10% EtOAc-hexane as the eluent to afford Compound 19 (23 mg, 29%). ¹H
 2277 NMR: (DMSO-d₆) δ 7.79-7.84 (m, 3H), 7.37-7.49 (m, 5H), 7.17 (t, J = 8.8 Hz, 2H), 5.89 (s,
 2278 1H), 4.38 (d, J = 6.2 Hz, 2H), 1.49 (s, 9H); MS: 352 [M + H]⁺; TLC: 20% EtOAc in hexane:
 2279 R_f: 0.60.

2280 *Example 31 - Preparation of Compound 20*

2281 **[0263] General Scheme VII.** A synthetic scheme useful for synthesis of compounds
 2282 described herein including Compound 20 is disclosed in General Scheme VII following,
 2283 wherein the terms “R^x”, “R^y”, and “R^z” are independently hydrogen, substituted or
 2284 unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted
 2285 cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted
 2286 heterocycloalkenyl, substituted or unsubstituted aryl, or substituted or unsubstituted
 2287 heteroaryl, or other groups obvious to those skilled in the art.

2288 **General Scheme VII**


2289

2290 [0264] A description of the syntheses of Intermediates 28-31 and Compound 20 follows.

2291 **Preparation of Intermediate 28 [General Procedure 16]**

2292 [0265] General Procedure 16 was followed in the preparation of Intermediate 28.

2293 **General Procedure 16**



2294

2295


Intermediate 28

2296 [0266] Thionyl chloride (5.4 mL, 73.2 mmol, 3 eq) was added to a solution of picolinic
2297 acid (3 g, 24.4 mmol) in EtOH (50 mL) at 0°C . The resulting mixture was heated to reflux
2298 and allowed to stir for 2 h. The mixture was then cooled and the solvent was evaporated. The
2299 resulting residue was poured into saturated aqueous NaHCO_3 and extracted with EtOAc ($2 \times$
2300 50 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated *in*
2301 *vacuo*. The crude material was purified by column chromatography over silica gel (100-200
2302 mesh) by using DCM as the eluent to afford Intermediate 28 (3 g, 81%) as a colorless liquid.
2303 MS: 152 $[\text{M} + \text{H}]^+$; TLC: 10% MeOH/ NH_3 in CHCl_3 ; R_f : 0.70.

2304 **Preparation of Intermediate 29 [General Procedure 17]**

2305 [0267] General Procedure 17 was followed in the preparation of Intermediate 29.

2306 **General Procedure 17**



2307

2308

Intermediate 28


Intermediate 29

2309 [0268] A solution of Intermediate 28 (3 g, 19.6 mmol) and CH_3CN (0.8 mL, 19.6 mmol, 1
2310 eq) in dry toluene (10 mL) was slowly added to a mixture of NaH (784 mg, 19.6 mmol, 1 eq,
2311 60% in mineral oil) in toluene (50 mL) at 65°C . The resulting mixture was allowed to stir at
2312 65°C for 16 h. The reaction mixture was then cooled to RT and quenched with ice cold water
2313 (20 mL). The resulting solid was filtered to afford Intermediate 29 (1.5 g, 53%) as a brown
2314 solid. ^1H NMR: (CDCl_3) δ 8.70 (d, $J = 4.8$ Hz, 1H), 8.12 (d, $J = 7.5$ Hz, 1H), 7.90-7.94 (m,
2315 1H), 7.56-7.60 (m, 1H), 4.41 (s, 2H); MS: 147 $[\text{M} + \text{H}]^+$; TLC: EtOAc; R_f : 0.40.

2316 **Preparation of Intermediate 30 [General Procedure 18]**

2317 [0269] General Procedure 18 was followed in the preparation of Intermediate 30.

General Procedure 15




Intermediate 29

Intermediate 30

[0270] Hydrazine hydrate (0.34 mL, 6.8 mmol, 1 eq) was added to a solution of Intermediate 29 (1 g, 6.8 mmol) in EtOH (30 mL) at RT. The mixture was then heated to reflux and allowed to stir for 20 h. The solvent was then evaporated. The resulting crude material was triturated with Et₂O (2 × 20 mL) and dried under vacuum to afford Intermediate 30 (700 mg, 64%) as a brown liquid. ¹H NMR: (DMSO-d₆) δ 8.53 (d, *J* = 4.4 Hz, 1H), 7.78 (d, *J* = 4.4 Hz, 2H), 7.23-7.26 (m, 1H), 5.95 (s, 1H), 4.84 (br s, 2H); MS: 161 [M + H]⁺; TLC: EtOAc: R_f: 0.20.

Preparation of Intermediate 31




Intermediate 30

Intermediate 31

[0271] General Procedure 10 was followed to afford Intermediate 31 (450 mg). MS: 269 [M + H]⁺; TLC: EtOAc: R_f: 0.40.

Preparation of Compound 20



Intermediate 31


Compound 20

[0272] General Procedure 11 was followed to afford Compound 20 (40 mg, 30%). ¹H NMR: (DMSO-d₆) δ 8.58 (d, *J* = 4.4 Hz, 1H), 7.86-7.98 (m, 3H), 7.38-7.46 (m, 3H), 7.18 (t, *J* = 8.8 Hz, 2H), 5.84 (s, 1H), 4.40 (d, *J* = 6.2 Hz, 2H), 1.50 (s, 9H); MS: 353 [M + H]⁺; MP: 102-103 °C; TLC: 20% EtOAc in hexane: R_f: 0.60.


Example 33 - General Scheme VIII

[0273] A synthetic scheme useful for synthesis of compounds described herein is disclosed in General Scheme VIII following, wherein the terms “Ar,” “R¹” and “R²” are as defined in Example 1.

General Scheme VIII




Example 34 - Preparation of Intermediate 32



[0274] A solution of ethyl cyanoacetate (20 g, 176.8 mmol) and triethyl orthoformate (29.4 mL, 176.8 mmol) in acetic anhydride (100 mL) was heated to 140 °C and allowed to stir for 5 h. The solvent was then evaporated to afford crude Intermediate 32 (23 g, 76%) as a low melting solid. MS: 170 [M + H]⁺; TLC: 30% EtOAc in hexane: R_f: 0.40.

2354 *Example 35 - Preparation of Intermediate 33*

2355



2356

Intermediate 32**Intermediate 33**


2357

2358

2359 **[0275]** Sodium acetate (8.2 g, 100 mmol, 2 eq) was added to a solution of Intermediate 32
 2360 (8.45 g, 50.0 mmol) and 2-hydrazinopyridine (5 g, 45.5 mmol, 0.9 eq) in AcOH (100 mL)
 2361 and water (20 mL). The resulting mixture was heated at 110 °C and allowed to stir for 16 h.
 2362 The mixture was then allowed to cool and ice-cold water was added. The precipitate was
 2363 collected by filtration and washed with Et₂O and dried under vacuum to afford Intermediate
 2364 33 (4 g, 38%) as a pale yellow solid. ¹H NMR: (DMSO-d₆) δ 8.48-8.49 (m, 1H), 8.00-8.04
 2365 (m, 1H), 7.87 (d, *J* = 8.3 Hz, 1H), 7.79 (s, 1H), 7.65 (br s, 2H), 7.33-7.36 (m, 1H), 4.22 (q, *J*
 2366 = 7.0 Hz, 2H), 1.28 (t, *J* = 7.0 Hz, 3H); MS: 233 [M + H]⁺; TLC: 15% EtOAc in hexane: R_f:
 2367 0.50.

2368 *Example 36 - Preparation of Compound 21*

2369



2370

Intermediate 33**Compound 21**

2371

2372 **[0276]** Sodium hydride (603 mg, 15.1 mmol, 1 eq, 60% in mineral oil) was added to a
 2373 solution of Intermediate 33 (3.5 g, 15.1 mmol) in DMF (300 mL) at 0°C. After 30 minutes, a
 2374 solution of 4-fluorobenzyl bromide (2.85 g, 15.1 mmol, 1 eq) in DMF (50 mL) was added
 2375 and the resulting mixture was allowed to warm to RT. After 5 h, the reaction mixture was
 2376 diluted with water (100 mL) and extracted with EtOAc (3 × 100 mL). The combined organic
 2377 layers were washed with water (5 × 50 mL), brine (50 mL), dried over Na₂SO₄, filtered and
 2378 concentrated *in vacuo*. The crude material was purified by column chromatography over
 2379 silica gel (100-200 mesh) by using a gradient mixture of 0-5% EtOAc-hexane as the eluent to
 2380 afford a partially pure product. The material was then recrystallized from Et₂O and pentane to

2381 afford Compound 21 (2.8 g, 55%) as a pale yellow solid. ¹H NMR: (DMSO-d₆) δ 9.50 (t, *J* =
 2382 6.6 Hz, 1H), 8.45-8.46 (m, 1H), 8.00-8.05 (m, 1H), 7.82-7.89 (m, 2H), 7.24-7.38 (m, 3H),
 2383 7.11 (t, *J* = 8.8 Hz, 2H), 4.88 (d, *J* = 6.6 Hz, 2H), 4.17 (q, *J* = 7.0 Hz, 2H), 1.24 (t, *J* = 7.0 Hz,
 2384 3H); MS: 341 [M + H]⁺; MP: 99-100 °C; TLC: 15% EtOAc in hexane: R_f: 0.40.

2385 *Example 37 – Trans-corneal permeability of compounds 4 and 22*

2386 [0277] In this example, trans-corneal permeability of compounds are measured in-vitro
 2387 across excised rabbit corneas. Excised rabbit cornea are acquired from Pel-Freez and
 2388 delivered in DMEM culture media on ice overnight. The test apparatus is a curved 9 mm
 2389 Franz chamber (PermeGear) suitable for rabbit eyes. Test compound is prepared in PBS
 2390 buffer. The compound solution is added to the donor chamber of the Franz chamber and the
 2391 entire apparatus placed in an incubator at 37°C for four hours. During incubation and at every
 2392 subsequent hour, a sample is removed from the receiver chamber and analyzed by HPLC
 2393 (Shimadzu Prominence) using a C18 column (Phenomenex 00F-4605-E0) in reverse phase
 2394 with acetonitrile in water. The apparent permeability coefficient P_{app} (cm/s) is calculated as

2395
$$P_{app} = \frac{1}{AC_0} \times \frac{dM}{dt}$$

2396 where dM/dt is the flux (nmol/s) across the cell layers or cornea, A is the exposed surface
 2397 area (cm²) of the insert membrane of rabbit cornea, and C₀ is the initial drug concentration
 2398 (μM) in the donor compartment.

2399 [0278] In Table E following, measured trans-corneal permeability P_{app} in units of cm/s is
 2400 presented for example compounds.

2401 **Table E**

Cmpd No.	P _{app}
4	2.5 × 10 ⁻⁶
22	5.7 × 10 ⁻⁶

2402

2403 *Example 38 – Pharmacokinetics in mice*

2404 [0279] In this example, pharmacokinetics in mice is presented for series of example
 2405 compounds #4, #23, #24, #25, #26. Each compound is administered intravenously (i.v.) as a
 2406 single dose via tail vein or orally (p.o.) as a single dose via gastric gavage to male CD-1 mice
 2407 of nominal weights between 20 g and 26 g. Nominal doses are 1 mg/kg and 5 mg/kg for i.v.
 2408 and p.o., respectively. In some examples (dose type A), both p.o. and i.v. doses are prepared
 2409 by dissolving the test compound in 5% dimethyl acetamide and diluted in tetraethylene glycol

for a final concentration of 0.25 mg/mL. In other examples (dose type B), i.v. doses are prepared by dissolving test compounds in 20% dimethyl acetamide, 40% polyethylene glycol 300 and 40% phosphate buffered saline, and p.o. doses are prepared by dissolving test compounds in carboxymethyl cellulose suspension (1% by weight) in water and 2.5 % dimethyl acetamide.

[0280] Animals are housed in standard holding cages with food and water available ad libitum except for animals used for p.o. dosing which are fasted overnight prior to dosing. Samples are taken in triplicate via cardiac puncture at times prior to dosing and at 0.083 (i.v. only), 0.25, 0.5, 1, 2, 4, 8, and 24 hours after administration. Plasma is obtained by centrifuge and stored frozen until analyzed by LC-MS/MS using a Shimadzu VP System HPLC coupled to a Applied Biosystems MDS SCIEX API 3000 triple quadrapole MS. Assay results are calibrated using reference samples prepared in a range between 1.5 and 5000 ng/mL.

[0281] Pharmacokinetic parameters are calculated from mean concentration values using a non-compartmental analysis as described in the following and as apparent to those of ordinary skill in the art. Half-lives ($t_{1/2}$) and elimination rate constants (λ) are determined by log linear regression using equal weighting on the last three finite sample time points. Concentration at time zero (C_0) for the i.v. data is established by the extrapolation of log linear regression using equal weighting on the first three sample time points. Area under the curve (AUC) values are calculated using linear trapezoidal integration. Systemic clearance (CL) is calculated as the ratio of dosage and AUC. The apparent volume of distribution (V_d) is calculated as the ratio of CL and λ . Percent oral bioavailability (%F) is determined from the ratio of i.v. and p.o. AUC values weighted by dosage.

[0282] In Table F following resulting pharmacokinetic parameters are listed for five example compounds, rounded to the nearest significant digit.

Table F

Cmpd No	4	23	24	25	26
Dose type	A	A	A	B	B
i.v. $t_{1/2}$ (h)	1	0.5	1	0.3	0.3
i.v. C_0 (ng/ml)	300	400	200	100	200
i.v. AUC (h·ng/ml)	100	100	200	40	50

i.v. V_d (ml/kg)	10000	5000	8000	6000	5000
i.v. CL (ml/kg/h)	10000	8000	5000	10000	10000
p.o. $t_{1/2}$ (h)	6	0.6	1	> 0.3	> 0.3
p.o. AUC (h·ng/ml)	300	80	600	10	20
%F	60	10	70	9	8

2435


2436 **[0283]** The contents of all references, patents, and published applications cited herein are
 2437 hereby incorporated by reference in their entirety and for all purposes.

2438 **[0284]** While the invention has been described in detail with reference to certain preferred
 2439 embodiments thereof, it will be understood that modifications and variations are within the
 2440 spirit and scope of that which is described and claimed.

2441

CLAIMS

1. A compound with structure of Formula (V):



or pharmaceutically acceptable salt, ester, solvate, or prodrug thereof;

wherein

L^1 is $-NR^7-$;

L^2 is a bond, or $-C(=O)-$, provided that if L^2 is a bond, R^2 is hydrogen;

L^5 is a bond or substituted or unsubstituted alkylene, provided that if L^5 is a bond, R^5 is hydrogen;

R^1 is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl;

R^2 is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl;

R^5 is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl; and

R^7 is hydrogen, or substituted or unsubstituted alkyl.

2. The compound according to claim 1, wherein L^2 is $-C(=O)-$, and R^2 is substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted

heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl.

3. The compound according to claim 2, wherein R^2 is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl.

4. The compound according to claim 3, wherein R^2 is substituted or unsubstituted heteroaryl selected from the group consisting of substituted or unsubstituted pyridyl, substituted or unsubstituted pyridazinyl, substituted or unsubstituted pyrimidinyl, substituted or unsubstituted thienyl, and substituted or unsubstituted furyl.

5. The compound according to claim 2, wherein R^2 is substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocycloalkyl.

6. The compound according to claim 5, wherein R^2 is substituted or unsubstituted heterocycloalkyl selected from the group consisting of substituted or unsubstituted oxanyl, substituted or unsubstituted oxetanyl, and substituted or unsubstituted morpholinyl.

7. The compound according to claim 2, wherein R^2 is substituted or unsubstituted fused ring aryl selected from the group consisting of substituted or unsubstituted benzodioxinyl and substituted or unsubstituted naphthyl.

8. The compound according to claim 1, wherein R^1 is hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocycloalkyl.

9. The compound according to claim 8, wherein R^1 is substituted alkyl having at least one substituted or unsubstituted heteroaryl substituent selected from the group consisting of substituted or unsubstituted pyridyl, substituted or unsubstituted pyridazinyl, substituted or unsubstituted pyrimidinyl, substituted or unsubstituted thienyl, and substituted or unsubstituted furyl.

10. The compound according to claim 9, wherein R^1 is substituted alkyl having at least one chloro-substituted thienyl substituent.

11. The compound according to claim 8, wherein R¹ is substituted alkyl having at least one substituted or unsubstituted heterocycloalkyl substituent selected from the group consisting of substituted or unsubstituted morpholinyl, substituted or unsubstituted oxanyl, and substituted or unsubstituted oxetanyl.

12. The compound according to claim 8, wherein R¹ is substituted alkyl having at least one substituted or unsubstituted fused ring aryl substituent selected from the group consisting of substituted or unsubstituted benzodioxinyl and substituted or unsubstituted naphthyl.

13. The compound according to any of claims 2 to 12, wherein L⁵ is substituted or unsubstituted alkylene, and R⁵ is substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl.

14. The compound according to claim 13, wherein R⁵ is substituted or unsubstituted heteroaryl selected from the group consisting of substituted or unsubstituted pyridyl, substituted or unsubstituted pyridazinyl, substituted or unsubstituted pyrimidinyl, substituted or unsubstituted thienyl, and substituted or unsubstituted furyl.

15. The compound according to claim 13, wherein R⁵ is substituted or unsubstituted fused ring aryl selected from the group consisting of substituted or unsubstituted benzodioxinyl and substituted or unsubstituted naphthyl.

16. The compound according to claim 2, wherein L⁵ is substituted or unsubstituted alkylene, and R⁵ is substituted or unsubstituted heterocycloalkyl.

17. The compound according to claim 16, wherein R⁵ is substituted or unsubstituted heterocycloalkyl selected from the group consisting of substituted or unsubstituted morpholinyl, substituted or unsubstituted oxanyl, and substituted or unsubstituted oxetanyl.

18. The compound of claim 1, wherein R⁵ is substituted or unsubstituted C₁-C₈ heteroalkyl having O or N as a heteroatom.

19. The compound of claim 1, wherein R⁵ is substituted C₁-C₈ alkyl having one or more substituent group selected from the group consisting of -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NR-C(NR'R''R''')=NR''', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -CN, -NO₂, -CF₃, substituted or unsubstituted alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl, wherein R', R'', R''', and R'''' each independently is hydrogen, substituted or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted alkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted thioalkoxy, or substituted or unsubstituted arylalkyl.

20. The compound of claim 1, wherein R² is substituted or unsubstituted cycloalkyl.

21. The compound of claim 20, wherein R² is substituted or unsubstituted cycloalkyl selected from the group consisting of substituted or unsubstituted cyclopropyl, substituted or unsubstituted cyclobutyl, substituted or unsubstituted cyclopentyl, and substituted or unsubstituted cyclohexyl.

22. The compound of claim 1, wherein R⁷ is substituted or unsubstituted methyl.

23. The compound of claim 1, wherein R⁷ is hydrogen.

24. The compound of any of claims 1-23, as set forth below:

Compound 1, 3-(5-amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 2, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 3, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 4, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 5, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 6, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;;

Compound 23, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one;;

Compound 25, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 26, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 27, 1-[(2-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 28, 1-[(2-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 29, 1-[(2-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 30, 1-[(2-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 31, 1-[(3-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 32, 1-[(3-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 33, 1-[(3-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 34, 1-[(3-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 35, 1-[(4-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 36, 1-[(4-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 37, 1-[(4-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 38, 1-[(4-chlorophenyl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 39, 1-[(5-chlorothiophen-2-yl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 40, 1-[(5-chlorothiophen-2-yl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 41, 1-[(5-chlorothiophen-2-yl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 42, 1-[(5-chlorothiophen-2-yl)methyl]-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 43, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 44, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 45, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 46, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 47, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 48, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 49, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 50, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 51, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 52, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 53, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 54, 1-benzyl-3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 55, 1-benzyl-3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1,2-dihydropyridin-2-one;

Compound 56, 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetonitrile;

Compound 57, 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetonitrile;

Compound 58, 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetonitrile;

Compound 59, 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetic acid;

Compound 60, 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetonitrile;

Compound 61, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 62, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 63, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 64, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 65, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 66, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 67, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 68, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 69, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 70, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 71, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 72, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-benzyl-1,2-dihydropyridin-2-one;

Compound 73, 3-(1-benzoyl-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one;

Compound 74, 3-(5-[(4-fluorophenyl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 75, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 76, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-phenylethyl)-1,2-dihydropyridin-2-one;

Compound 77, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 78, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 79, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(naphthalen-1-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 80, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(naphthalen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 81, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 82, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 83, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 84, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 85, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 86, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 87, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 88, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one;

Compound 89, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 90, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one;

Compound 91, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 92, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 93, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 94, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 95, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-hydroxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 96, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 97, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 98, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 99, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 100, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 101, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 102, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 103, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 104, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 105, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 106, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 107, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(3-methoxy-2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 108, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 109, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 110, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 111, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 112, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 113, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 114, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 115, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 116, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 117, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 118, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 119, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 120, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,3-dihydro-1,4-benzodioxin-5-yl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one;

Compound 121, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 122, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 123, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 124, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 125, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 126, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 127, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 128, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 129, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 130, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 131, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2,4-dimethoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 132, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 133, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 134, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(2-phenylethyl)-1,2-dihydropyridin-2-one;

Compound 135, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound	136,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	137,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(naphthalen-2-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	138,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	139,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	140,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	141,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	142,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	143,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	144,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;
Compound	145,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one;
Compound	146,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;
Compound	147,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one;
Compound	148,	3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(3-methyloxetan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one;

Compound 149, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 150, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 151, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 152, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 153, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 154, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 155, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 156, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 157, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 158, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 159, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(4-methyloxan-4-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 160, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 161, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 162, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 163, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 164, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 165, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 166, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 167, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 168, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 169, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 170, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 171, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 172, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-phenylethyl)-1,2-dihydropyridin-2-one;

Compound 173, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 174, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(naphthalen-1-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 175, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(naphthalen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 176, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 177, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 178, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 179, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 180, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one;

Compound 181, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 182, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 183, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 184, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 185, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 186, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 187, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 188, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 189, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 190, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 191, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 192, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-2-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 193, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 194, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 195, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 196, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 197, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 198, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 199, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 200, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 201, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 202, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 203, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 204, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(thiophen-3-yl)carbonyl]-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 205, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[2-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 206, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(2-methoxyethoxy)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 207, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1,2-dihydropyridin-2-one;

Compound 208, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 209, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 210, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 211, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 212, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 213, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 214, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 215, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 216, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 217, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 218, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[4-(morpholin-4-yl)phenyl]carbonyl-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 219, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(1H-1,2,3,4-tetrazol-5-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 220, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 221, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(2-phenylethyl)-1,2-dihydropyridin-2-one;

Compound 222, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 223, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 224, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(naphthalen-1-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 225, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(naphthalen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 226, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 227, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 228, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 229, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 230, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 231, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 232, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 233, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)-2-oxoethyl]-1,2-dihydropyridin-2-one;

Compound 234, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 235, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-[2-(pyridin-2-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 236, 3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-1-methyl-1,2-dihydropyridin-2-one;

Compound 237, 3-[5-(benzylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1,2-dihydropyridin-2-one;

Compound 238, 3-[5-(benzylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 239, 3-[5-(dimethylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1,2-dihydropyridin-2-one;

Compound 240, 3-[5-(dimethylamino)-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl]-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 241, 3-1-[(2-aminophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1,2-dihydropyridin-2-one;

Compound 242, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(2-methoxyethyl)-1,2-dihydropyridin-2-one;

Compound 243, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(furan-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 244, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(furan-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 245, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridazin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 246, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 247, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridin-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 248, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyridin-4-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 249, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(pyrimidin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 250, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(thiophen-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 251, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-(thiophen-3-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 252, 3-1-[(2-chlorophenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-[2-(morpholin-4-yl)ethyl]-1,2-dihydropyridin-2-one;

Compound 253, 3-1-[(4-tert-butylphenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1,2-dihydropyridin-2-one;

Compound 254, 3-1-[(4-tert-butylphenyl)carbonyl]-5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl-1-methyl-1,2-dihydropyridin-2-one;

Compound 255, 3-5-amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl-1,2-dihydropyridin-2-one;

Compound 256, 3-5-amino-1-[(furan-2-yl)carbonyl]-1H-pyrazol-3-yl-1-(pyridin-2-ylmethyl)-1,2-dihydropyridin-2-one;

Compound 257, ethyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate;

Compound 258, ethyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate;

Compound 259, ethyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate;

Compound 260, ethyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate;

Compound 261, tert-butyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-(2,2-dimethylpropanoyl)-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate;

Compound 262, tert-butyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(2-methoxyphenyl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate;

Compound 263, tert-butyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1-[(furan-3-yl)carbonyl]-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate;

Compound 264, tert-butyl 2-[3-(5-[(5-chlorothiophen-2-yl)methyl]amino-1H-pyrazol-3-yl)-2-oxo-1,2-dihydropyridin-1-yl]acetate;

Compound 265, 1-[(5-[(5-chlorothiophen-2-yl)methyl]amino-3-(1-methyl-2-oxo-1,2-dihydropyridin-3-yl)-1H-pyrazol-1-yl)carbonyl]cyclopropylmethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate.

25. A pharmaceutical composition comprising a pharmaceutically acceptable excipient in combination with a compound according to any of claims 1-24.

26. A compound according to any of claims 1 to 24, or a pharmaceutical composition according to claim 25, for use in therapy.

27. A compound according to any of claims 1 to 24, or a pharmaceutical composition according to claim 25, for use as a medicament.

28. A compound according to any of claims 1 to 24, or a pharmaceutical composition according to claim 25, for use in the treatment of a disease or disorder responsive to inhibition of thrombin.

29. The compound for use according to any of claims 26-28, wherein L^2 is $-C(=O)-$, and R^2 is substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl.

30. The compound for use according to claim 28, wherein said disease or disorder is at least one of a thrombotic disorder, a disease or disorder involving a blood clot thrombus, and a disease or disorder which can lead to formation of a blood clot thrombus.

31. The compound for use according to claim 30, wherein said thrombotic disorder comprises at least one of acute coronary syndrome, thromboembolism, and thrombosis.

32. The compound for use according to claim 31, wherein the thromboembolism comprises at least one of venous thromboembolism, arterial thromboembolism, and cardiogenic thromboembolism.

33. The compound for use according to claim 32, wherein the venous thromboembolism comprises at least one of deep vein thrombosis and pulmonary embolism.

34. The compound for use according to claim 30, wherein said thrombotic disorder involves dysfunctional coagulation or disseminated intravascular coagulation.

35. The compound for use according to claim 28, wherein said disease or disorder comprises at least one of fibrosis, Alzheimer's Disease, multiple sclerosis, pain, cancer, and inflammation.

36. The compound for use according to claim 31, wherein the thrombosis is arterial thrombosis.

37. A compound according to any of claims 1 to 24, or a pharmaceutical composition according to claim 25, for use in the treatment of a disease or disorder responsive to inhibition of kallikrein.

38. The compound for use according to claim 37, wherein said kallikrein-related disorder is an ophthalmic disease.

39. The compound for use of claim 38, wherein said ophthalmic disease comprises at least one of diabetic macular edema, age-related macular degeneration, and diabetic retinopathy.

40. The compound for use according to claim 38, wherein said kallikrein-related disorder is an ophthalmic disease, and wherein said compound or pharmaceutical composition is to be administered orally, in the form of an ophthalmic composition applied topically to the eye, or in the form of an ophthalmic composition via intravitreal injection.

41. The compound for use according to claim 40, wherein said ophthalmic composition is in the form of eye drops.

42. The compound for use according to claim 38, wherein said compound acts by reducing retinal vascular permeability.

43. The compound for use according to claim 37, wherein said kallikrein-related disorder comprises at least one of a thrombotic disease, a fibrinolytic disease, a type of cancer, an inflammatory condition, and a dermatological condition.

44. The compound for use according to claim 43, wherein said type of cancer comprises at least one of cervical-, testicular-, and non-small-cell lung adenocarcinoma.

45. The compound for use according to claim 43, wherein said inflammatory condition comprises at least one of sepsis, inflammatory bowel disease, systemic inflammatory response syndrome, hereditary angioedema, and rheumatoid arthritis.

46. The compound for use according to claim 43, wherein said kallikrein-related disorder is an inflammatory condition and said compound is to be administered orally.

47. The compound for use according to claim 43, wherein said dermatological condition comprises at least one of atopic dermatitis, psoriasis, or Netherton Syndrome.

48. The compound for use according to claim 37, wherein said compound acts by inhibiting kallikrein.

49. The compound for use according to claim 48, wherein said compound acts by inhibiting tissue kallikrein.

50. The compound for use according to claim 48, wherein said compound acts by inhibiting plasma kallikrein.

51. Use of a compound according to any of claims 1 to 24, or a pharmaceutical composition according to claim 25, in the manufacture of a medicament for the treatment of a disease or disorder responsive to inhibition of thrombin.

52. The use according to claim 51, wherein L^2 is $-C(=O)-$, and R^2 is substituted or unsubstituted alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted cycloalkenyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted heterocycloalkyl, substituted or unsubstituted heterocycloalkenyl, substituted or unsubstituted aryl, substituted or unsubstituted fused ring aryl, or substituted or unsubstituted heteroaryl.

53. The use according to claim 51, wherein said disease or disorder is at least one of a thrombotic disorder, a disease or disorder involving a blood clot thrombus, and a disease or disorder which can lead to formation of a blood clot thrombus.

54. The use according to claim 53, wherein said thrombotic disorder comprises at least one of acute coronary syndrome, thromboembolism, and thrombosis.

55. The use according to claim 54, wherein the thromboembolism comprises at least one of venous thromboembolism, arterial thromboembolism, and cardiogenic thromboembolism.

56. The use according to claim 55, wherein the venous thromboembolism comprises at least one of deep vein thrombosis and pulmonary embolism.

57. The use according to claim 53, wherein said thrombotic disorder involves dysfunctional coagulation or disseminated intravascular coagulation.

58. The use according to claim 51, wherein said disease or disorder comprises at least one of fibrosis, Alzheimer's Disease, multiple sclerosis, pain, cancer, and inflammation.

59. The use according to claim 54, wherein the thrombosis is arterial thrombosis.

60. Use of a compound according to any of claims 1 to 24, or a pharmaceutical composition according to claim 25, in the manufacture of a medicament for the treatment of a disease or disorder responsive to inhibition of kallikrein.

61. The use according to claim 60, wherein said kallikrein-related disorder is an ophthalmic disease.

62. The use of claim 61, wherein said ophthalmic disease comprises at least one of diabetic macular edema, age-related macular degeneration, and diabetic retinopathy.

63. The use according to claim 61, wherein said kallikrein-related disorder is an ophthalmic disease, and wherein said compound or pharmaceutical composition is to be administered orally, in the form of an ophthalmic composition applied topically to the eye, or in the form of an ophthalmic composition via intravitreal injection.

64. The use according to claim 63, wherein said ophthalmic composition is in the form of eye drops.

65. The use according to claim 61, wherein said compound acts by reducing retinal vascular permeability.

66. The use according to claim 60, wherein said kallikrein-related disorder comprises at least one of a thrombotic disease, a fibrinolytic disease, a type of cancer, an inflammatory condition, and a dermatological condition.

67. The use according to claim 66, wherein said type of cancer comprises at least one of cervical-, testicular-, and non-small-cell lung adenocarcinoma.

68. The use according to claim 66, wherein said inflammatory condition comprises at least one of sepsis, inflammatory bowel disease, systemic inflammatory response syndrome, hereditary angioedema, and rheumatoid arthritis.

69. The use according to claim 66, wherein said kallikrein-related disorder is an inflammatory condition and said compound is to be administered orally.

70. The use according to claim 66, wherein said dermatological condition comprises at least one of atopic dermatitis, psoriasis, or Netherton Syndrome.