14. The pharmaceutical composition according to any one of claims 10 to 13 wherein the compound is crystalline Form 2 N-[3-[(4aR,7aS)-2-amino-6-(5-fluoropyrimidin-2-yl)-4a,5,6,7-tetrahydropyrrolo[3,4-d][1,3]thiazin-7a(4H)-yl]-4-fluoro-phenyl]-5-methoxy-pyrazine-2-carboxamide.

15. The pharmaceutical composition according to claim 14 wherein the crystalline Form 2 N-[3-[(4aR,7aS)-2-amino-6-(5-fluoropyrimidin-2-yl)-4a,5,6,7-tetrahydropyrrolo[3,4-d][1,3]thiazin-7a(4H)-yl]-4-fluoro-phenyl]-5-methoxy-pyrazine-2-carboxamide is characterized by a substantial peak in the X-ray diffraction spectrum, at diffraction angle 2-theta of 11.8°, with one or more peaks selected from the group consisting of 18.6°, 19.3°, and 26.7°; with a tolerance for the diffraction angles of 0.2 degrees.

16. A compound of the formula:



or a pharmaceutically acceptable salt thereof, in combination with an anti-N3pGlu Abeta monoclonal antibody for use in the treatment of Alzheimer's disease.

17. The compound for use according to claim 16 wherein the anti-N3pGlu Abeta monoclonal antibody is B12L or R17L.

18. The compound for use according to either claim 16 or claim 17 wherein the anti-N3pGlu Abeta monoclonal antibody is B12L.

19. The compound for use according any one of claims 16 to 18 wherein the compound is N-[3-[(4aR,7aS)-2-amino-6-(5-fluoropyrimidin-2-yl)-4a,5,6,7-tetrahydropyrrolo[3,4-d][1,3]thiazin-7a(4H)-yl]-4-fluoro-phenyl]-5-methoxy-pyrazine-2-carboxamide.

20. The compound for use according any one of claims 16 to 19 wherein the compound is crystalline Form 2 N-[3-[(4aR,7aS)-2-amino-6-(5-fluoropyrimidin-2-yl)-

4a,5,6,7-tetrahydropyrrolo[3,4-d][1,3]thiazin-7a(4H)-yl]-4-fluoro-phenyl]-5-methoxy-pyrazine-2-carboxamide.

21. The compound for use according to claim 20 wherein the crystalline Form 2 N-[3-[(4aR,7aS)-2-amino-6-(5-fluoropyrimidin-2-yl)-4a,5,6,7-tetrahydropyrrolo[3,4-d][1,3]thiazin-7a(4H)-yl]-4-fluoro-phenyl]-5-methoxy-pyrazine-2-carboxamide is characterized by a substantial peak in the X-ray diffraction spectrum, at diffraction angle 2-theta of 11.8°, with one or more peaks selected from the group consisting of 18.6°, 19.3°, and 26.7°; with a tolerance for the diffraction angles of 0.2 degrees.