


FIG. 1






FIG. 2B




FIG. 2C




FIG. 2D





FIG. 2E



Treatment	Dose (mg/kg)	Ave Conc. (nM)	Ave Conc. (µg/ml)
ADX-097	30	837	181
ADX-097	10	69.0	14.9
ADX-097	3	4.31	0.93
ADX-097	1	1.36	0.29
Fc-2fH _{1.5}	17	1138	137

FIG. 2F





FIG. 2G



** $P < 0.004$ vs. 3, 10, or 30 mg/kg ADX-097

FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 4A




FIG. 4B




FIG. 4C

Test article	Dose (mg/kg)	uPCR _{AKI} (± SEM)	% change vs. PBS	P value vs. PBS
PBS	-	116 ± 19.5	-	-
ADX-097	3	38.2 ± 9.29	↓ 73 ± 8.7	0.002
	1 (IV)	53.3 ± 10.9	↓ 66 ± 9.6	0.007
	1 (SC)	46.2 ± 10.2	↓ 59 ± 10	0.01
	0.3	81.2 ± 23.2	↓ 32 ± 22	0.24
CVF	100 U/kg (x9)	34.9 ± 11.6	↓ 76 ± 11	0.002
Fc-2fH _{5.5}	1.7	47.3 ± 8.24	↓ 65 ± 7.8	0.006
	0.57	97.3 ± 22.4	↓ 17 ± 21	0.51
	0.17	114 ± 14.4	↓ 1.6 ± 14	0.94

FIG. 4D




FIG. 4E




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 6A

*Spearman correlation matrix of
urinary biomarkers to glomerular C3*





FIG. 6B



FIGs. 7A-7F




FIG. 8C




FIG. 8B




FIG. 8A

