



[Figure 1]

Hydrophobic peptide fraction (IL-6)

Mean $\pm$ SE, n=3 **p<0.01 *p<0.05 vs 0 μ g/mL (Turkey-Kramer)


[Figure 2]

Hydrophobic pyroglutamyl peptide fraction (IL-6)

Mean $\pm$ SE, n=3 **p<0.01 *p<0.05 vs 0 μ g/mL (Turkey-Kramer)


[Figure 3]

Hydrophilic peptide fraction (IL-6)


Mean $\pm$ SE, n=3 **p<0.01 *p<0.05 vs 0 μ g/mL (Turkey-Kramer)

[Figure 4]


Hydrophilic pyroglutamyl peptide fraction (IL-6)

Mean $\pm$ SE, n=3 **p<0.01 *p<0.05 vs 0 μ g/mL (Turkey-Kramer)

[Figure 5]

Hydrophobic peptide fraction (IL-1 β)Mean $\pm$ SE, n=3 **p<0. 01 *p<0. 05 vs 0 μ g/mL (Turkey-Kramer)

[Figure 6]

Hydrophobic pyroglutamyl peptide fraction (IL-1 β)Mean $\pm$ SE, n=3 **p<0. 01 *p<0. 05 vs 0 μ g/mL (Turkey-Kramer)