

1 / 9





FIG. 1

2 / 9



	R_1	R_2	R_3
Iso-mogroside VI (1)	$-\alpha\text{-OH}$	$-\text{Glc}_I \xrightarrow{6} \text{Glc}_{II}$	$-\text{Glc}_{III} \xrightarrow{6} \text{Glc}_{IV}$ 2 Glc_V 4 Glc_{VI}
<i>epi</i> -mogroside V (2)	$-\beta\text{-OH}$	$-\text{Glc}_I \xrightarrow{6} \text{Glc}_{II}$	$-\text{Glc}_{III} \xrightarrow{6} \text{Glc}_{IV}$ 2 Glc_V
neomogroside (3)	$-\alpha\text{-OH}$	$-\text{Glc}_I \xrightarrow{6} \text{Glc}_{II}$ 2 Glc_{VI}	$-\text{Glc}_{III} \xrightarrow{6} \text{Glc}_{IV}$ 2 Glc_V
11-oxo-mogroside V (4)	$=\text{O}$	$-\text{Glc}_I \xrightarrow{6} \text{Glc}_{II}$	$-\text{Glc}_{III} \xrightarrow{6} \text{Glc}_{IV}$ 2 Glc_V
mogroside V (5)	$-\alpha\text{-OH}$	$-\text{Glc}_I \xrightarrow{6} \text{Glc}_{II}$	$-\text{Glc}_{III} \xrightarrow{6} \text{Glc}_{IV}$ 2 Glc_V
mogroside VI (6)	$-\alpha\text{-OH}$	$-\text{Glc}_I \xrightarrow{6} \text{Glc}_{II}$ 2 Glc_{VI}	$-\text{Glc}_{III} \xrightarrow{6} \text{Glc}_{IV}$ 2 Glc_V

FIG. 2




FIG. 3(A)




FIG. 3(B)




FIG. 4




FIG. 4 Cont'd




FIG. 5(A-D)

HSQC-TOCSY with different mixing time*			
10 ms	30 ms	60 ms	100 ms
C-1 	C-1 	C-1 	C-1 
C-2 	C-2 	C-2 	C-2 
	C-3 	C-3 	C-3 
		C-4 	C-4 
			C-5 
			C-6 

FIG. 6

9 / 9




FIG. 7




FIG. 8