960805-5
1 pos copy

THIS SHEET HAS BEEN REPLACED

PC7851AGFF

INDOLE DERIVATIVES

Background of the Invention

The present invention relates to indole derivatives, to processes and intermediates for their preparation, to pharmaceutical compositions containing them and to their medicinal use. The active compounds of the present invention are useful in treating migraine and other disorders.

United States Patents 4,839,377 and 4,855,314 and European Patent Application Publication Number 313397 refer to 5-substituted 3-aminoalkyl indoles. The compounds are said to be useful for the treatment of migraine.


British Patent Application 040279 refers to 3-aminoalkyl-1H-indole-5-thioamides and carboxamides. The compounds are said to be useful in treating hypertension, Raymond's disease and migraine.

European Patent Application Publication Number 303506 refers to 3-poly:hydro-pyridyl-5-substituted-1H-indoles. The compounds are said to have 5HT1-receptor agonist and vasoconstrictor activity and to be useful in treating migraine.

European Patent Application Publication Number 354777 refers to N-piperidinyl:indolyl:ethyl-alkane sulfonamide derivatives. The compounds are said to have 5HT1-receptor agonist and vasoconstrictor activity and to be useful in treating cephalic pain.

Summary of the Invention

The present invention relates to compounds of the formula



I

This Sheet Has Been Replaced

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

wherein n is 0, 1, or 2; R_1 is hydrogen; R_2 is selected from hydrogen, halogen (e.g., fluorine, chlorine, bromine or iodine), cyano, OR_4, $-(CH_2)_m-(C=O)NR_5R_6$, $-(CH_2)_m-SO_2NR_5R_6$, $-(CH_2)_m-NR_7(C=O)R_8$, $-(CH_2)_m-NR_7SO_2R_8$, $-(CH_2)_m-S(O)_xR_8$, $-(CH_2)_m-NR_7(C=O)NR_5R_6$, $-(CH_2)_m-NR_7(C=O)OR_9$, and $-CH=CH(CH_2)_yR_{10}$; R_3 is selected from hydrogen and C_1 to C_6 linear or branched alkyl; R_4 is selected from hydrogen, C_1 to C_6 alkyl, and aryl; R_5 and R_6 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl or R_5 and R_6 taken together to form a 4, 5, or 6 membered ring; R_7 and R_8 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_9 is selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_{10} is selected from $-(C=O)NR_5R_6$ and $-SO_2NR_5R_6$, wherein R_5 and R_6 are defined as above, and $-NR_7(C=O)R_8$, $-NR_7SO_2R_8$, $-NR_7(C=O)NR_5R_6$, $-S(O)_xR_8$ and $-NR_7(C=O)OR_9$, wherein R_7, R_8, and R_9 are as defined above; m is 0, 1, 2, or 3; y is 0, 1, or 2; x is 1 or 2; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen (e.g., fluorine, chlorine, bromine or iodine), hydroxy, cyano, carboxamido, nitro and C_1 to C_4 alkoxy and the pharmaceutically acceptable salts thereof. These compounds are useful in treating migraine and other disorders. Compounds of the formula I wherein R_2 is $-CH=CH-R_{10}$ are also useful as intermediates for preparing other compounds of the formula I.

The compounds of the invention include all optical isomers of formula I (e.g., R and S enantiomers) and their racemic mixtures. The R enantiomers at the designated chiral site in formula I are preferred.

Unless otherwise indicated, the alkyl groups referred to herein, as well as the alkyl moieties of other groups referred to herein (e.g. alkoxy), may be linear or branched,

THIS SHEET HAS BEEN REPLACED

and they may also be cyclic (e.g., cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl) or be linear or branched and contain cyclic moieties.

Preferred compounds of the invention are compounds of the formula I wherein R_1 is hydrogen; R_2 is $-(CH_2)_m-SO_2NHR_5$, $-(CH_2)_m-NHSO_2R_8$, $-(CH_2)_m-SO_2R_8$, $-(CH_2)_m-(C=O)NHR_5$, or $-(CH_2)_m-NH(C=O)R_8$; R_3 is hydrogen or methyl; and m, R_5 and R_8 are as defined above and the pharmaceutically acceptable salts thereof. Of the foregoing preferred compounds, the R enantiomers at the designated chiral site in formula I are more preferred.

The following compounds are particularly preferred:

(R)-5-methoxy-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-bromo-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-ethylsulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylmethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-carboxamido-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylsulfonylethyl)-3-(N-methylpyrrolidin-2-yl-methyl)-1H-indole;

(R)-5-(2-aminosulphonylethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-aminosulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-N,N-dimethylaminosulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-phenylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

(R)-5-(2-phenylsulphonyethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole hemisuccinate;

(R)-5-(2-ethylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole hemisuccinate;

(R)-5-(3-benzenecarbonylaminoprop-1-enyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-(4-methylphenylsulphonyl)ethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(3-methylsulphonylaminoprop-1-enyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-ethylsulphonylethyl)-3-(N-2-propylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-ethylsulphonylethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-(4-methylphenylsulphonyl)ethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylsulfonamidoethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole; and

(R)-5-(2-methylsulfonamidomethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole.

The following are other specific compounds of the present invention:

(R)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-fluoro-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-acetylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;


(R)-5-benzyloxycarbonylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-aminocarbonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-aminocarbonylmethyl-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-methylsulfonamido-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole; and

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

(R)-5-aminosulfonyl-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole.

The present invention also relates to a pharmaceutical composition for treating a condition selected from hypertension, depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain, and chronic paroxysmal hemicrania and headache associated with vascular disorders comprising an amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating such condition and a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition for treating disorders arising from deficient serotonergic neurotransmission (e.g., depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain, and chronic paroxysmal hemicrania and headache associated with vascular disorders) comprising an amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating such condition and a pharmaceutically acceptable carrier.

The present invention also relates to a method for treating a condition selected from hypertension, depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain and chronic paroxysmal hemicrania and headache associated with vascular disorders comprising administering to a mammal (e.g., a human) requiring such treatment an amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating such condition.


The present invention also relates to a method for treating disorders arising from deficient serotonergic neurotransmission (e.g., depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain and chronic paroxysmal hemicrania and headache associated with vascular disorders) comprising administering to a mammal (e.g., a human) requiring such treatment an

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating such condition.


The present invention also relates to a compound of the formula



V


wherein W is $-CO_2R_{11}$ or R_3; Q is CH_2 or C=O; n, R_1, R_2 and R_3 are as defined for formula I; and R_{11} is selected from C_1 to C_6 alkyl, benzyl and aryl, wherein aryl is as defined above. The compounds of formula V are useful as intermediates in preparing compounds of the formula I.

Accordingly, one group of the foregoing intermediates comprises compounds of the formula



II

wherein n, R_1, R_2 and R_{11} are as defined above and a second group of the foregoing intermediates comprises compounds of the formula



III


This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

wherein n, R_1, R_3 and R_{10} are as defined above.

Detailed Description of the Invention


Compounds of formula I are prepared by hydride reduction of a compound of the formula



II

wherein R_1, R_2, n and R_{11} are as defined above with a hydride reducing agent in an inert solvent. Suitable hydride reducing agents include lithium aluminum hydride, diborane, lithium borohydride and sodium borohydride. The preferred reagent is lithium aluminum hydride. Suitable solvents include ethers, such as diethyl ether, tetrahydrofuran, 1,4-dioxane and 1,2-dimethoxyethane. The preferred solvent is tetrahydrofuran. The reaction is conducted at a temperature of about 30°C to about 100°C., preferably about 65°C to about 70°C.

Compounds of formula I are also prepared by catalytic reduction of a compound of the formula



III

wherein R_1, R_3, n and R_{10} are as defined above under an atmosphere of hydrogen, preferably at a pressure of about 1 to about 3 atmospheres, or using a hydrogen source such as ammonium formate or formic acid in an inert solvent. Suitable catalysts include palladium on carbon, Raney nickel, platinum oxide, rhodium, and ruthenium. The


THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

preferred catalyst is palladium on carbon. Suitable solvents include C_1 to C_6 alcohols, N,N-dimethylformamide, ethyl acetate, and acetonitrile. The preferred solvent is ethanol. The reaction is conducted at a temperature of about 0°C to about 60°C, most preferably at about 25°C.

Compounds of formula I are also prepared by alkylation of compounds of formula I where R_3=H and R_2 and R_1, are as defined for formula I with alkyl halides in the presence of a base in an inert solvent. Suitable alkyl halides include alkyl halides R^3 - Halide where the halide is chloride, bromide and iodide. The preferred halide is iodide, or bromide in the presence of a suitable iodide source such as sodium iodide. Suitable bases include tertiary amines and inorganic bases. The preferred base is sodium carbonate. Suitable solvents include N,N-dimethylacetamide, N,N-dimethylformamide, dimethoxyethane, tetrahydrofuran, dichloromethane, acetonitrile. The preferred solvent is N,N-dimethylacetamide. The reaction is conducted at a temperature of about 0°C to about 150°C, preferably at about 120°C.

The compounds of formula II can be prepared by reacting a magnesium salt of an indole derivative of the formula



IV


wherein R_1 and R_2 are defined above, with the acid chloride of an N-CO_2R_{11}-proline, N-CO_2R_{11}-azetidine-2-carboxylic acid, or N-CO_2R_{11}-pipecolinic acid (R, S, or racemate), wherein R_{11} is defined as above. The indole magnesium salt is first prepared from the reaction of an indole of formula IV with an alkyl or aryl magnesium halide, preferably ethylmagnesium bromide. The reaction is generally conducted in an inert solvent at a temperature between about -30°C and about 65°C, preferably at about 25°C. Suitable solvents include diethyl

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

ether, tetrahydrofuran, and other alkyl ethers. The preferred solvent is diethyl ether. The acid chloride of proline, azetidine-2-carboxylic acid, or pipecolinic acid is prepared in a separate reaction vessel by reaction of the N-CO_2R_{11}-proline, N-CO_2R_{11}-azetidine-2-carboxylic acid, or N-CO_2R_{11}-pipecolinic acid (R, S, or racemate), with oxalyl chloride in methylene chloride at about -10°C to about 25°C (Helv. Chim. Acta, 1920 (1976)). Suitable solvents include diethyl ether, tetrahydrofuran, other alkyl ethers, and methylene chloride. The proline, azetidine-2-carboxylic acid, or pipecolinic acid is N-substituted with a protecting group to avoid reaction of the nitrogen with the acid chloride when it is formed. Suitable protecting groups are substituted-aryl or substituted-alkyl carbamates (e.g. benzyloxycarbonyl). Preferably, a solution of the N-CO_2R_{11}-proline acid chloride in an inert solvent (e.g., diethyl ether) is added slowly to the solution of the magnesium salt of an indole of formula IV at a temperature of about -30°C to about 50°C, preferably at about 25°C.

The compounds of formula III can be prepared by reacting a compound of formula




wherein R_1, R_3 and n are defined as above and X is chlorine, bromine or iodine (preferably bromine), with a compound containing a vinyl group (e.g. ethyl vinyl sulfone or N-methylvinylsulfonamide) in the presence of a palladium catalyst, a triarylphosphine and a base in an inert solvent. Suitable catalysts include palladium (II) salts, preferably palladium (II) acetate. Suitable solvents include acetonitrile, N,N-dimethylformamide, and tetrahydrofuran.

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

The preferred solvent is acetonitrile. The preferred triarylphosphine is tri-o-tolylphosphine. Suitable bases include trisubstituted amines. The preferred base is triethylamine. The reaction is conducted at a temperature of about 25°C to 150°C, most preferably at about 80°C.


Compounds of formula I and intermediates to compounds of formula I can be prepared by hydride reduction of a compound of the formula



VI

wherein R_2, n, and R_{11} are as defined above with a hydride reducing agent in an inert solvent. Suitable hydride reducing agents include lithium aluminum hydride, diborane, lithium borohydride, and sodium amide. The preferred reagent is lithium aluminum hydride. Suitable solvents include ethers, such as diethyl ether, tetrahydrofuran, 1,4-dioxane and 1,2-dimethoxyethane. The preferred solvent is tetrahydrofuran. The reduction is conducted at a temperature of about 30°C to about 100°C, preferably about 65°C to about 70°C.

Compounds of formula I and intermediates to compounds of formula I can also be prepared by catalytic reduction of a compound of the formula




VI

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

wherein R_2, n, and R_{11} are as defined above under an atmosphere of hydrogen, preferably at a pressure of about 1 to 3 atmospheres, or using a hydrogen source such as ammonium formate of formic acid in an inert solvent. Suitable catalysts include palladium on carbon, Raney nickel, and platinum oxide. The preferred catalyst is palladium on carbon. Suitable solvents include C_1 to C_6 alcohols, N,N-dimethylformamide, ethyl acetate, and acetonitrile. The preferred solvent is ethanol. The reaction is conducted at a temperature of about 0°C to about 60°C, preferably at about 25°C.

Compounds of formula VI can be prepared by the transition metal catalyzed cyclization of a compound of the formula



VII


wherein R_2, n, and R_{11} are as defined above, and X is chlorine, bromine, or iodine (preferably bromine or iodine), and R_{12} is $-OR_{11}$ as defined above or alkyl, aryl, or trifluoromethyl (preferably trifluoromethyl) in a suitable inert solvent with a phase transfer catalyst and a base. Suitable catalysts include palladium salts such as palladium (II) acetate or palladium (II) chloride (preferably palladium acetate) and rhodium salts, such as tris(triphenyl phosphine)rhodium (I) chloride. Suitable solvents include N,N-dimethylformamide, acetonitrile, and N-methylpyrrolidine. The preferred solvent is N,N-dimethylformamide. Suitable phase transfer catalysts include tetraalkylammonium halides, preferably tetra-n-butylammonium chloride. Suitable bases include tertiary amines, sodium hydrogen carbonate, and sodium carbonate.

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

The preferred base is triethylamine. The reaction is conducted at a temperature of about 80°C to about 180°C, preferably about 150°C to 160°C.


Compounds of formula VI can also be prepared by hydride reduction of a compound of the formula




II

wherein R_2, n, and R_{11} are as defined above with a hydride reducing agent in an inert solvent. Suitable hydride reducing agents include lithium borohydride, sodium borohydride, and sodium cyanoborohydride. The preferred reagent is lithium borohydride. Suitable solvents include ethers, such as diethyl ether, tetrahydrofuran, 1,4-dioxane and 1,2-dimethoxyethane. The preferred solvent is tetrahydrofuran. The reduction is conducted at a temperature of about 30°C to about 100°C, preferably about 65°C to about 70°C.

Compounds of formula VII can be prepared by the Mitsunobu coupling reaction of compounds of formulas



VIII



IX


wherein R_2, n, R_{11}, and R_{12} are as defined above using a phosphine and an azodicarboxylate in a suitable solvent. Suitable phosphines include trialkylphosphines and triarylphosphines, preferably triphenylphosphine. Suitable azodicarboxylates include dialkyl azodicarboxylates,

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

preferably diethyl diazodicarboxylate. Suitable solvents include methylene chloride, ethers, including tetrahydrofuran, diethyl ether, and 1,4-dioxane, N-N-dimethylformamide and acetonitrile. The preferred solvent is tetrahydrofuran. The reaction is conducted at a temperature of about 0°C to about 65°C, most preferably at about 25°C.


Compounds of formula VIII, if not available commercially, can be prepared by reacting a compound of formula X



X

wherein R_2 and X are as defined above with the acid chloride or the symmetrical anhydride of $R_{12}CO_2H$ in a suitable solvent with an suitable base. The preferred acid chloride or anhydride is trifluoroacetic anhydride. Suitable solvents include ethers, including tetrahydrofuran, diethyl ether and 1,4-dioxane, methylene chloride, and chloroform. The preferred solvent is methylene chloride. Suitable bases include triethylamine, pyridine, and sodium hydrogen carbonate. The preferred base is pyridine. The reaction is conducted at a temperature of about 0°C to about 65°C, preferably at about 25°C.

Compounds of formula X, if not available commercially, can be prepared by reacting a compound of formula XI



XI


wherein R_2 is as defined above with either chloride, bromine, or iodine in a suitable solvent with a suitable base. Reaction with bromine is preferred. Suitable solvents

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

include C_1-C_6 alcohols, methylene chloride, chloroform, or carbon tetrachloride. The preferred solvent is methanol. Suitable bases include triethylamine, pyridine, sodium carbonate, and sodium hydrogen carbonate. The preferred base is sodium hydrogen carbonate. The reaction is conducted at a temperature of about 0°C to about 65°C, preferably at about 25°C.

Compounds of the formula IX can be prepared from hydride reduction of a compound of formula XII




XII

wherein R_{11} is defined as above and R_{13} is C_1-C_6 alkyl, aryl, or alkylaryl with a hydride reducing agent in an inert solvent. Suitable hydride reducing agents include lithium aluminum hydride, lithium borohydride, sodium borohydride, and diisobutylaluminum hydride. The preferred reagent is diisobutylaluminum hydride. Suitable solvents include ethers, such as diethyl ether, tetrahydrofuran, 1,4-dioxane and 1,2-dimethoxyethane. The preferred solvent is tetrahydrofuran. The reduction is conducted at a temperature of about -100°C to about 0°C, preferably at about -80°C to about -70°C.


Compounds of the formula XII can be prepared from the Wittig reaction in a suitable solvent involving compounds of the formulas

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced



XIII



XIV

wherein R_{11} and R_{13} are defined as above. Suitable solvents include ethers such as diethyl ether, tetrahydrofuran, and 1,4-dioxane. Tetrahydrofuran is the preferred solvent. The reaction is conducted at a temperature of about -78°C to about 30°C, preferably at about -78°C.

Compounds of the formula XIII can be prepared as outlined in S. Kiyooka, et al., J. Org. Chem., 5409 (1989) and Y. Hamada, et al., Chem. Pharm. Bull., 1921 (1982).

Compounds of the formula XIV are either commercially available or can be prepared as outlined in L. Fieser and M. Fieser, Reagents for Organic Synthesis, John Wiley and Sons, New York, Vol. 1, p. 112 (1967).

Unless indicated otherwise, the pressure of each of the above reactions is not critical. Generally, the reactions will be conducted at a pressure of about one to about three atmospheres, preferably at ambient pressure (about one atmosphere).

The compounds of the formula I which are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate a compound of the formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent, and subsequently convert the free base to a pharmaceutically

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is obtained.

The acids which are used to prepare the pharmaceutically acceptable acid addition salts of the base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate or bisulfate, phosphate or acid phosphate, acetate, lactate, citrate or acid citrate, tartrate or bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts.

Those compounds of the formula I which are also acidic in nature, e.g., where R_2 contains a carboxylate, are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the herein described acidic compounds of formula I. These non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium calcium and magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction of maximum product of yields of the desired final product.

The compounds of the formula I and the pharmaceutically acceptable salts thereof (hereinafter, also referred to as the active compounds of the invention) are useful psychotherapeutics and are potent serotonin (5-HT_1) agonists and may be used in the treatment of depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, chronic paroxysmal hemicrania and headache associated with vascular disorders, pain, and other disorders arising from deficient serotonergic neurotransmission. The compounds can also be used as centrally acting antihypertensives and vasodilators.

The active compounds of the invention are evaluated as anti-migraine agents by testing the extent to which they mimic sumatriptan in contracting the dog isolated saphenous vein strip (P.P.A. Humphrey et al., Br. J. Pharmacol., 94, 1128 (1988)). This effect can be blocked by methiothepin, a known serotonin antagonist. Sumatriptan is known to be useful in the treatment of migraine and produces a selective increase in carotid vascular resistance in the anaesthetized dog. It has been suggested (W. Fenwick et al., Br. J. Pharmacol., 96, 83 (1989)) that this is the basis of its efficacy.

The compositions of the present invention may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers. Thus, the active compounds of the invention may be formulated for oral, buccal, intranasal, parenteral (e.g., intravenous, intramuscular or subcutaneous) or rectal administration or

This Sheet Has Been Replaced

in a form suitable for administration by inhalation or insufflation.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium phosphate); lubricants (e.g. magnesium stearate, talc or silica); disintegrants (e.g. potato starch or sodium starch glycollate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g. sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agents (e.g. lecithin or acacia); non-aqueous vehicles (e.g. almond oil, oily esters or ethyl alcohol); and preservatives (e.g. methyl or propyl p-hydroxybenzoates or sorbic acid).

For buccal administration the composition may take the form of tablets or lozenges formulated in conventional manner.

The active compounds of the invention may be formulated for parenteral administration by injection, including using conventional catheterization techniques or infusion. Formulations for injection may be presented in unit dosage form e.g. in ampules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulating agents such as suspending, stabilizing and/or dispersing agents.

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

Alternatively, the active ingredient may be in powder form for reconstitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

The active compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

For intranasal administration or administration by inhalation, the active compounds of the invention are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped by the patient or as an aerosol spray presentation from a pressurized container or a nebulizer, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

A proposed dose of the active compounds of the invention for oral, parenteral or buccal administration to the average adult human for the treatment of the conditions referred to above (e.g., migraine) is 0.1 to 200 mg of the active ingredient per unit dose which could be administered, for example, 1 to 4 times per day.

Aerosol formulations for treatment of the conditions referred to above (e.g., migraine) in the average adult human are preferably arranged so that each metered dose or "puff" of aerosol contains 20 μg to 1000 μg of the compound of the invention. The overall daily dose with an aerosol will be within the range 100 μg to 10 mg. Administration

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

may be several times daily, for example 2, 3, 4 or 8 times, giving for example, 1, 2 or 3 doses each time.

The following Examples illustrate the preparation of the compounds of the present invention. Melting points are uncorrected. NMR data are reported in parts per million (δ) and are referenced to the deuterium lock signal from the sample solvent. Specific rotations were measured at room temperature using the sodium D line (589 nm).

Commercial reagents were utilized without further purification. Chromatography refers to column chromatography performed using 32-63 μm silica gel and executed under nitrogen pressure (flash chromatography) conditions. Room temperature refers to 20 - 25°C.

EXAMPLE 1

General Procedure for the Reduction of Benzyloxycarbonyl- pyrrolidin-2-ylcarbonyl-1H-indole, N-Benzyloxycarbonyl-azetidin-2-ylcarbonyl-1H-indoles, or N-Benzyloxycarbonyl-piperidin-2-ylcarbonyl-1H-indoles Forming 3-(N-Methyl- pyrrolidin-2-ylmethyl)-1H-indoles, 3-(N-Methyl-azetidin-2-ylmethyl)-1H-indoles, or 3-(N-Methylpiperidin-2-ylmethyl)-1H-indoles, respectively.

To a stirred solution of (R)- or (S)-(N-benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-1H-indole, (R)-, (S), or (R,S)-(N-benzyloxycarbonylazetidin-2-ylcarbonyl)-1H-indole, or (R)-, (S)-, or (R,S)-(N-benzyloxycarbonylpiperidin-2-ylcarbonyl)-1H-indole, (5.00 mmol) in anhydrous tetrahydrofuran (20mL) at room temperature under nitrogen was carefully added lithium aluminum hydride (0.57 g, 15.0 mmol, 3.0 eq) as a powder, and the resulting mixture was stirred at room temperature under nitrogen for 1 hour. The mixture was then heated at reflux (66°C) under nitrogen for 12 hours. The reaction was then quenched with successive additions of water (0.5 mL), aqueous sodium hydroxide (20%, 0.5 mL), and then additional water (1.0 mL), and the resulting mixture filtered through diatomaceous earth (Celite (trademark)).

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

The solids were then washed with copious amounts of ethyl acetate (50 mL). The combined filtrate was then washed with water (20 mL), dried ($MgSO_4$), and evaporated under reduced pressure. The residue was then column chromatographed using silica gel (50 g) and elution with the appropriate solvent system to afford the 3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole, 3-(N-methylazetidin-2-ylmethyl)-1H-indole, or 3-(N-methylpiperidin-2-ylmethyl)-1H-indole. Following this procedure the following compounds were prepared:

A. **(S)-5-Methoxy-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

(S)-(N-Benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-5-methoxy-1H-indole was used. The chromatographic eluent was 8% triethylamine in ethyl acetate to afford the title compound (yields ranged from 22 to 57%) as an oil: IR ($CHCl_3$) 3475, 1625, 1585, 1480, 1455 cm^{-1}; ^1H NMR ($CDCl_3$) δ 8.13 (br s, 1H), 7.23 (d, J=8.8 Hz, 1H), 7.04 (d, J=2.4 Hz, 1H), 6.97 (d, J=2.2 Hz, 1H), 6.84 (dd, J=2.4 and 8.8 Hz, 1H), 3.86 (s, 3H), 3.17-3.10 (m, 2H), 2.58 (dd, J=9.9 and 13.9 Hz, 1H), 2.50-2.40 (m, 1H), 2.47 (s, 3H), 2.26-2.17 (m, 1H), 1.89-1.72 (m, 2H), 1.70-1.52 (m, 2H); ^{13}C NMR ($CDCl_3$) δ 153.8, 131.4, 128.2, 122.7, 113.9, 111.8, 111.7, 101.1, 66.6, 57.5, 56.0, 40.8, 31.5, 30.0, 21.9; LRMS, m/z (relative intensity) 244 (M^+, 7), 160 (20), 145 (16), 117 (21), 84 (100); HRMS: calculated for $C_{15}H_{20}N_2O$: 244.1573; found: 244.1575; $[\alpha]^{25} = -96°$ ($CHCl_3$, c = 1.0).

B. **(R)-5-Methoxy-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

(R)-(N-Benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-5-methoxy-1H-indole was used. The chromatographic eluent was 8% triethylamine in ethyl acetate to afford the title compound (yields ranged from 13 to 61%) as an oil whose spectral and physical properties were identical with the spectral and physical properties of the title compound of Example 1A with the exception of specific rotation of plane

This Sheet Has Been Replaced

polarized light: $[\alpha]^{25}$ = + 100° ($CHCl_3$, c = 1.0). HRMS: calculated for $C_{15}H_{20}N_2O$: 244.1573; found: 244.1547.

C. **(R)-5-Dibenzylamino-3-(N-methylpyrroidin-2-ylmethyl)-1H-indole**

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-5-dibenzylamino-1H-indole was used. Column chromatography using elution with methylene chloride/methanol/ammonium hydroxide [9:1:0.1] afforded the title compound as a pale green foam: ^1H NMR ($CDCl_3$) δ 7.82 (br s, NH), 7.35-7.19 (m, 10 H), 7.20 (d, J=8.6 Hz, 1H), 6.95 (d, J=2.1 Hz, 1H), 6.85 (dd, J=2.3 and 8.7 Hz, 1H), 6.80 (d, J=2.2 Hz, 1H), 4.65 (s, 4H), 3.25-3.02 (m, 2H), 2.52 (dd, J=9.5 and 13.9 Hz, 1H), 2.39-2.15 (m, 2 H), 2.30 (s, 3H), 1.85-1.40 (m, 4H); ^{13}C NMR ($CDCl_3$) δ 143.2, 139.7, 130.5, 128.5, 128.2, 127.3, 126.8, 122.9, 112.5, 112.2, 111.8, 103.4, 67.0, 57.4, 56.4, 40.6, 31.4, 29.7, 21.9. HRMS: calculated for $C_{28}H_{31}N_3$ 409.2520. Found 409.2475.

D. **(R)-5-Methoxy-3-(N-methylpiperid-2-ylmethyl)-1H-indole**

(R)-3-(N-Benzyloxycarbonylpiperid-2-ylcarbonyl)-5-methoxy-1H-indole was used. Column chromatography using elution with 6% triethylamine in ethyl acetate afforded the title compound as a white foam: ^{13}C NMR ($CDCl_3$) δ 153.7, 131.4, 128.3, 123.3, 113.2, 111.7, 111.6, 101.2, 64.4, 57.2, 55.9, 43.4, 31.0, 28.8, 25.9, 24.1; $[\alpha]^{25}$ = +67° ($CDCl_3$, c=1.0); HRMS: calculated for $C_{16}H_{22}N_2O$: 258.1734. Found: 258.1710.

E. **(S)-5-Methoxy-3-(N-methylazetidin-2-ylmethyl)-1H-indole**

(S)-3-(N-Benzyloxycarbonylazetidinyl-2-ylcarbonyl)-5-methoxy-1H-indole was used. The chromatographic eluet was 8% triethylamine in ethyl acetate to afford the title compound as a white solid: mp, 118.0-120.0°C; ^{13}C NMR ($CDCl_3$) δ 153.8, 131.6, 128.0, 122.9, 112.3, 111.9, 111.8, 101.0, 68.5, 56.0, 53.1, 44.7, 32.4, 25.0; $[\alpha]^{25}$ = -44° ($CHCl_3$,

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

c=1.0). Anal. calcd for $C_{14}H_{18}N_2O$: C, 73.01; H, 7.88, N, 12.16. Found: C, 72.65; H, 7.91; N, 12.06.

F. **(R,S)-5-Methoxy-3-(N-methylazetidin-2-ylmethyl)-1H-indole**

(R,S)-3-(N-Benzyloxycarbonylazetidinyl-2-ylcarbonyl)-5-methoxy-1H-indole was used. The chromatographic eluet was 10% triethylamine in ethyl acetate to afford the title compound as a white solid: mp, 116.0-119.0°C; Anal. calcd for $C_{14}H_{18}N_2O$: C, 73.01; H, 7.88; N, 12.16. Found: C, 72.61; H, 7.99; N, 12.10.

EXAMPLE 2

General Method for the Hydrogenation of 5-(2-Sulfonylethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indoles to Form 5-(2-Sulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indoles

A solution of 5-(2-sulfonylethenyl)-3-(N-methylpyrrolidin-2-yl)-1H-indole (0.47 mmol) and 10% Pd/C (0.150 g) in ethanolic hydrogen chloride [prepared from absolute ethanol (10 mL) and acetyl chloride (43 μL)] and N,N-dimethylformamide (7.5 mL) was shaken under a hydrogen atmosphere (15 psi) at room temperature for 20 hours. The resultant reaction mixture was filtered through diatomaceous earth (Celite (trademark)), washed with absolute ethanol, and the combined filtrates were evaporated under reduced pressure. The residue was partitioned between ethyl acetate and water. The organic phase was separated, washed with water (3x), brine (1x), dried (Na_2SO_4), and evaporated under reduced pressure to afford a yellow oil. Column chromatography of this oil using silica gel and elution with methylene chloride/absolute ethanol/ammonia (90:10:1) afforded the appropriate 5-(2-Ethylsulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole. Following this procedure, the following compounds were prepared:

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

A. **(R)-5-(2-Ethylsulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

(R)-5-trans-(2-Ethylsulfonylethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole (Example 4A) was reduced as described above. Chromatography afforded the title compound (0.33 mmol, 70%) as a gum: TLC (CH_2Cl_2:EtOH:NH_3, 90:10:1): R_f = 0.3; $[\alpha]^{25}$ = + 62° (methanol, c = 0.10). Anal. Calcd for $C_{18}H_{26}N_2O_2S \cdot 0.05\ CH_2Cl_2$: C, 63.21; H, 7.67; N, 8.17; found: C, 63.55; H, 7.61; N, 8.41.

B. **(R)-5-(2-Methylaminosulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

(R)-5-trans-(2-Methylaminosulfonylethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole (Example 4B) was reduced as described above. Chromatography afforded the title compound (65%) as a foam. Anal. Calcd for $C_{17}H_{25}N_3O_2S \cdot 0.1\ CH_2Cl_2$: C, 59.71; H, 7.39; N, 12.12; found: C, 59.66; H, 7.14; N, 11.90.

EXAMPLE 3

General Synthesis of 3-(N-Benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-1H-indoles 3-(N-Benzyloxycarbonylazetidin-2-ylcarbonyl)-1H-indoles, or 3-(N-Benzyloxycarbonylpiperidin-2-ylcarbonyl)-1H-indoles.

Two solutions containing the reactants were prepared separately as follows. To a stirred solution of N-carbobenzyloxyproline (D or L, 3.10 g, 12.4 mmol, 1 eq) or N-carbobenzyloxyazetidine-2-carboxylic acid (R or S or racemate, 12.4 mmol) or N-carbobenzyloxypipecolinic acid (R or S or racemate, 12.4 mmol) in anhydrous methylene chloride (7 mL) with one drop dimethylformamide was added oxalyl chloride (1.60 mL, 18.4 mmol, 1.5 eq), and the resulting effervescing solution was stirred at room temperature under nitrogen for 1.5 hours. The solution was then evaporated under reduced pressure, and any remaining solvent was removed from the residual oil using high vacuum to afford the N-benzyloxycarbonylproline acid chloride. At the same time, a solution of ethylmagnesium bromide (3.0 M in ether,

This Sheet Has Been Replaced



This Sheet Has Been REPLACED

4.13 mL, 12.4 mmol, 1 eq) was added to a stirred solution of the indole (12.4 mmol) in anhydrous ether (50 mL), and this cloudy solution was heated at reflux under nitrogen for 1.5 hours to form the indolemagnesium bromide salt. The proline acid chloride was then dissolved in methylene chloride or ethyl ether (3 mL), and this solution was added dropwise to the stirred solution of the indolemagnesium bromide salt at room temperature, and the resultant reaction mixture was stirred at room temperature under nitrogen for 1 hour. A saturated solution of sodium hydrogen carbonate (25 mL) and ethyl acetate (50 mL) was then added to the reaction mixture, and this mixture was vigorously stirred for 15 minutes. The resulting mixture was filtered through diatomaceous earth (Celite (trademark)), the solids washed with copious amounts of ethyl acetate, and the ethyl acetate layer was separated from the aqueous layer which was extracted with ethyl acetate (2 x 25 mL). All ethyl acetate extracts were combined, dried, and evaporated under reduced pressure. The residual oil/solid was flash chromatographed using silica gel (250 g) and eluted with an appropriate solvent system to afford the desired 3-(N-benzyloxycarbonylpyrrolidin-2-ylcarbonyl)indole, 3-(N-benzyloxycarbonylazetidin-2-ylcarbonyl)-1H-indole, or 3-(N-benzyloxycarbonylpiperidin-2-ylcarbonyl)-1H-indole.

A. **(S)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-5-methoxy-1H-indole**

N-Carbobenzyloxy-L-proline was used. Chromatography using 40-60% ethyl acetate gradient in hexanes afforded the title compound (yields ranged from 27 to 43%) as a white powder. Recrystallization in ethyl acetate/hexanes afforded an analytical sample as a white crystalline solid: mp, 164.0-165.0°C; IR (KBr) 3250, 1695, 1660, 1585, 1520, 1485, 1450, 1425 cm^{-1}; ^1H NMR ($CDCl_3$) [Note: the spectrum of the title compound appears as a 1:3 mixture of diastereomers due to slow inversion of the amide nitrogen on an NMR time scale. Therefore, the ^1H NMR will be interpreted for each

This Sheet Has Been Replaced

compound separately with the more abundant conformer quoted first] δ [more abundant conformer] 9.83 (br s, 1H), 7.53 (d, J=3.4 Hz, 1H), 7.42-7.30 (m, 6H), 7.00 (d, J=8.9 Hz, 1H), 6.69 (dd, J=2.4 and 9.0 Hz, 1H), 5.25 (d, J=12.9 Hz, 1H), 5.14 (d, J=12.5 Hz, 1H), 5.07-4.99 (m, 1H), 3.74 (s, 3H), 3.78-3.55 (m, 2H), 2.28-1.84 (m, 4H) and δ [less abundant conformer] 9.28 (br s, 1H), 7.90 (d, J=2.3 Hz, 1H), 7.59 (d, J=3.4 Hz, 1H), 7.24 (d, J=9.0 Hz, 1H), 7.06-6.90 (m, 5H), 6.88 (dd, J=2.7 and 9.0 Hz, 1H), 5.07-4.99 (m, 2H), 4.96-4.88 (m, 1H), 3.86 (s, 3H), 3.78-3.55 (m, 2H), 2.28-1.84 (m, 4H); LRMS, m/z (relative intensity) 379 (8), 378 (M^+,33), 204 (31), 174 (64), 160 (41), 146 (10), 91 (100). Analysis: calculated for $C_{22}H_{22}N_2O_4$: C, 69.83; H, 5.86; N, 7.40; found: C, 69.81; H, 5.67; N, 7.40.

B. **(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-5-methoxy-1H-indole**

N-Carbobenzyloxy-D-proline was used. Chromatography using 40-60% ethyl acetate gradient in hexanes afforded the title compound (yields ranged from 25 to 36%) as a white powder. Recrystallization in ethyl acetate/hexanes afforded an analytical sample as a white crystalline solid: mp, 165.0-166.0°C. The spectral and physical data for the title compound was identical in all respects with the spectral and physical data of its enantiomer (the title compound of Example 3A); HRMS: calculated for $C_{22}H_{22}N_2O_4$: 378.1582; found: 378.1573.

C. **(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-5-dibenzylamino-1H-indole**

N-Carbobenzyloxy-D-proline was used. Trituration of the extraction residue with diethyl ether afforded the title compound as a solid: mp, 176.0-177.0°C; LRMS (m/z, relative intensity) 543 (100, M^+), 453 (10), 407 (7), 339 (40), 307 (10), 247 (10), 154 (38); $[\alpha]^{25}$=+112° (THF, c=1.0); Anal. calcd for $C_{35}H_{33}N_3O_3$: C, 77.32; H, 6.12; N, 7.73. Found: C, 77.35; H, 6.30; N, 7.66.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

D. **(R)-3-(N-Benzyloxycarbonylpiperid-2-ylcarbonyl)-5-methoxy-1H-indole**

N-Carbobenzyloxy-D-pipecolinic acid was used. Column chromatography using elution with 10% ether in methylene chloride afforded the title compound as a tan foam: LRMS (m/z, relative intensity) 392 (90, M^+), 348 (27), 284 (13), 273 (12), 258 (15), 237 (47), 217 (58), 173 (100), HRMS: calculated for $C_{35}H_{33}N_3O_2$: C, 69.22; H, 5.53; N, 7.69. Found: C, 69.35; H, 5.33; N, 7.64.

E. **(S)-3-(N-Benzyloxycarbonylazetidinyl-2-ylcarbonyl)-5-methoxy-1H-indole**

(S)-N-Carbobenzyloxyazetidine-2-carboxylic acid was used. Trituration of the extract residue with absolute methanol afforded the title compound as a white solid: mp, 199.0-200.0°C. Anal. calcd for $C_{21}H_{20}N_2O_4$: C, 69.22; H, 5.53; N, 7.69. Found: C, 69.35; H, 5.33; N, 7.64.

F. **(R,S)-3-(N-Benzyloxycarbonylazetidinyl-2-ylcarbonyl)-5-methoxy-1H-indole**

(R,S)-N-Carbobenzyloxyazetidine-2-carboxylic acid was used. Trituration of the extract residue with absolute methanol afforded the title compound as a white solid: mp, 199.0-200.0°C. Anal. calcd for $C_{21}H_{20}N_2O_4$: C, 69.22; H, 5.53; N, 7.69. Found: C, 68.85; H, 5.47; N, 7.57.

EXAMPLE 4

General Method for the Synthesis of 5-trans-(2-Sulfonylethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indoles

A mixture of the appropriate vinyl sulfone (1.17 mmol, 1.4 eq), tri-o-tolylphosphine (0.075 g, 0.25 mmol, 0.33 eq), palladium (II) acetate (0.013 g), triethylamine (0.25 mL, 1.79 mmol, 2 eq), and (R)-5-bromo-3-(N-methylpyrrolidinyl-methyl)-1H-indole (0.25 g, 0.85 mmol) in anhydrous acetonitrile (3 mL) was heated at reflux under nitrogen for 17 hours. The resultant reaction mixture was evaporated under reduced pressure, and the residue was column chromatographed using silica gel and

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced



elution with methylene chloride/absolute ethanol/ammonia (90:8:1) to afford the title compound.

A. **(R)-5-trans-(2-Ethylsulfonylethenyl)-3-(N-methyl-pyrrolidin-2-ylmethyl)-1H-indole**

Ethyl vinyl sulfone was used, and chromatography afforded the title compound (65%) as a white foam: TLC (CH_2Cl_2/EtOH/NH_3, 90:10:1):R_f = 0.5. Analysis: calculated for $C_{18}H_{24}N_2O_2S \cdot 0.2\ CH_2Cl_2$: C, 62.55; H, 7.04; N, 8.02; found: C, 62.65; H, 6.94; N, 7.92.

B. **(R)-5-trans-(2-Methylaminosulphonylethenyl)-3-(N-methyl-pyrrolidin-2-ylmethyl)-1H-indole**

N-methylvinylsulfonamide was used, and chromatography afforded the title compound (71%) as a white foam. Analysis: calculated for $C_{17}H_{23}N_3O_2S \cdot 0.1\ CH_2Cl_2$: C, 60.06; H, 6.84; N, 12.29; found: C, 59.74; H, 6.77; N, 11.97.

EXAMPLE 5

General Procedure for the Hydride Reduction of 3-(N-Benzyloxycarbonyl-pyrrolidin-2-ylmethyl)-1H-indoles and 3-(N-Benzyloxycarbonylpiperid-2-ylmethyl)-1H-indoles Forming 3-(N-Methylpyrrolidin-2-ylmethyl)-1H-indoles and 3-(N-Methylpiperid-2-ylmethyl)-1H-indoles

To a stirred mixture of lithium aluminum hydride (0.152 g, 4.00 mmol,2 eq) in anhydrous tetrahydrofuran (10 mL) at 0°C was added rapidly a solution of the 3-(N-benzyloxy-carbonylpyrrolidin-2-ylmethyl)-1H-indole or the 3-(N-benzyloxycarbonylpiperid-2-ylmethyl)-1H-indole (2.00 mmol) in anhydrous tetrahydrofuran (5 mL). The resulting mixture is heated at reflux under a nitrogen atmosphere for 3 hours. The reaction mixture is cooled, and water (0.25 mL), 15% aqueous sodium hydroxide (0.25mL), and then more water (0.75 mL) were added sequentially. The resulting mixture was stirred at 25°C for 30 minutes, filtered, and the filtrate was evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 50 g) and elution with a solution methylene chloride: methanol: ammonium hydroxide [9:1:0.1] or other appropriate solvent

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

system to afford the corresponding 3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole or 3-(N-methylpiperid-2-ylmethyl)-1H-indole.

Following this procedure the following compounds were prepared:

A. **(R)-5-(Methylaminosulfonylmethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-5-(methylaminosulfonylmethyl)-1H-indole was used. The reaction residue after aqueous work-up as described above was triturated with absolute methanol to afford the title compound as a white solid: mp, 213.0-214.0°C; ^{1}H NMR (DMSO-d_6) δ 10.9 (br s, indole NH), 7.51 (be d, 1H), 7.31 (d, J=8.3 Hz, 1H), 7.16 (br d, 1H), 7.08 (br dd, J=8.3 Hz, 1H), 6.82 (br q, sulfonamide NH), 4.35 (s, 2H), 3.07-2.95 (m, 2H), 2.54 (d, J=4.7 Hz, 3H), 2.52-2.38 (m, 2H), 2.35 (s, 3H), 2.10 (br, q, J=8.2 Hz, 1H), 1.75-1.40 (m, 4H); $[\alpha]^{25}$=+89° (DMSO-d_6, c=1.0); Anal. calcd for $C_{16}H_{23}N_3SO_2$: C, 59.79; H, 7.21; N, 13.07. Found: C, 59.66; H, 7.29; N, 12.81.

B. **(R)-5-Aminomethyl-3-(N-methylpyrrolidin-2-ylmethyl)-1H-inole**

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-5-cyano-1H-indole was used. Column chromatography using elution with 9:1:0.1 [methylene chloride:methanol:ammonium hydroxide] afforded the title compound as a white foam: ^{13}C NMR δ 135.6, 132.3, 127.5, 123.0, 122.8, 121.4, 117.1, 112.8, 111.5, 66.8, 57.2, 46.4, 40.5, 31.2, 29.2, 21.5; HRMS: calculated for $C_{15}H_{21}N_3$ 243.1737, found 243.1732.

C. **(R,S)-5-(Methylaminosulfonylmethyl)-3-(N-methylpiperid-2-ylmethyl)-1H-indole**

(R,S)-3-(N-Benzyloxycarbonylpiperidin-2-ylmethyl)-5-(methylaminosulfonylmethyl)-1H-indole was used. Column chromatography using elution with 10% triethylamine in ethyl acetate afforded the title compound as a clear, colorless oil: ^{13}C NMR (DMSO-d_6) δ 135.9, 127.7, 124.0, 123.6, 121.0, 119.7, 111.9, 111.1, 63.9, 56.7, 56.3, 43.2, 30.5, 29.0,

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

27.9, 25.5, 23.7; LRMS (m/z, relative intensity) 336 (1, M^+), 241 (5), 143 (31), 142 (13), 99 (34), 98 (100), 70 (16); HRMS calculated for $C_{17}H_{25}N_3O_2S$: 336.1745; found: 336.1756.

EXAMPLE 6

General Procedure for the Catalytic Reduction of 3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indoles and 3-(N-Benzyloxycarbonylpiperid-2-ylmethyl)-1H-indoles Forming 3-(Pyrrolidin-2-ylmethyl)-1H-indoles and 3-(Piperid-2-ylmethyl)-1H-indoles


A mixture of the 3-(N-benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole or the 3-(N-benzyloxycarbonylpiperid-2-ylmethyl)-1H-indcle (2.00 mmol), 10% palladium on carbon (0.20 g), and ammonium formate (1.26 g, 20 mmol, 10 eq) in absolute ethanol (15 mL) was stirred under a nitrogen atmosphere for 4 hours. The resulting reaction mixture was filtered through diatomaceous earth, and the filtrate was evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 50 g) and elution with a solution of methylene chloride: methanol: ammonium hydroxide [8:2:0.2] or other appropriate solvent system to afford the corresponding 3-(pyrrolidin-2-ylmethyl)-1H-indole or 3-(piperid-2-ylmethyl)-1H-indole.

Following this procedure the following compounds were prepared:

A. **(R)-5-(Methylaminosulfonylmethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole**

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-5-(methylaminosulfonylmethyl)-1H-indole was used. Column chromatography as described above afforded the title compound as an off-white gum: ^{13}C NMR (DMSO-d_6) δ 135.9, 127.5, 123.8, 123.7, 120.9, 119.7, 112.4, 111.1, 59.2, 56.6, 45.7, 31.1, 31.0, 29.0, 24.6; $[\alpha]^{25}$ = +4° (DMSO-d_6, c=1.0); $[\alpha]^{25}$=-14° (EtOH/$CHCl_3$ [1:1], c=1.0); HRMS: calculated for [$C_{15}H_{21}N_3O_2S \cdot H^+$]: 308.1433; found: 308.1467.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

B. **(R)-5-Cyano-3-(pyrrolidin-2-ylmethyl)-1H-indole**

(R)-3-(N-Benzyloxycarbonlpyrrolidin-2-ylmethyl)-5-cyano-1H-indole was used. Column chromatography as described above afforded the title compound as an off-white gum: ^{13}C NMR ($CDCl_3/CD_3OD$) δ 138.1, 127.2, 125.0, 124.4, 124.2, 121.0, 113.4, 112.2, 101.5, 59.5, 50.1, 45.7, 31.3, 30.3, 24.7; LRMS (M/z, relative intensity) 225 (M+,3), 179 (3), 155 (10), 70 (100); HRMS: calculated for $C_{14}H_{15}N_3$ 225.1268, found 225.1245.

C. **(R)-3-(Pyrrolidin-2-ylmethyl)-1H-indole**

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole was used. Evaporation of the filtrate residue directly afforded the title compound as a white foam: ^1H NMR ($CDCl_3$) δ 9.05 (br s, indole NH), 7.50 (d, J=8.6 Hz, 1H), 7.23 (d, J=8.6 Hz, 1H), 7.12-6.98 (m, 2H), 6.90 (s, 1H), 4.0 (br s, amine NH), 3.36-3.24 (m, 1H), 2.95-2.75 (m, 3H), 2.70-2.58 (m, 1H), 1.85-1.50 (m, 3H), 1.45-1.29 (m, 1H); $[\alpha]^{25} = +18°$ ($CHCl_3$, c=1.0).

D. **(R)-5-Methoxy-3-(Pyrrolidin-2-ylmethyl)-1H-indole**

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-5-methoxy-1H-indole was used. Evaporation of the filtrate residue directly afforded the title compound as a gum: LRMS (m/z, relative intensity) 231 (100, M+), 161 (10), 155 (17), 135 (11), 119 (32); $[\alpha]^{25}=-12°$ ($CHCl_3$, c=1.0); Anal, calcd for $C_{14}H_{18}N_2O \cdot 0.75\ C_2H_4O_2$ [acetic acid salt]: C, 67.61; H, 7.69; N, 10.17. Found: C, 67.74; H, 7.53; N, 9.90.

E. **(R,S)-5-(Methylaminosulfonylmethyl)-3-(piperid-2-ylmethyl)-1H-indole**

(R,S)-3-(N-Benzyloxycarbonylpiperid-2-ylmethyl)-5-(methylaminosulfonylmethyl)-1H-indole was used. Column chromatography as described above afforded the title compound as a clear, colorless oil: ^{13}C NMR (DMSO-d_6) δ 136.0, 127.5, 124.2, 123.8, 121.0, 119.8, 111.2, 110.9, 56.8, 56.7, 45.8, 31.7, 31.4, 29.0, 25.0, 23.9; LRMS (m/z, relative intensity) 321 (19, M+), 238 (43), 227 (21), 144 (99), 143

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

(100); HRMS: calculated for $C_{16}H_{23}N_3O_2S$: 321.1513; found: 321.1501.

EXAMPLE 7

General Procedure for the Formulation of 3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indoles and 3-(N-Benzyloxycarbonylpiperid-2-ylmethyl)-1H-indoles Via the Palladium Catalyzed Cyclization of 1-(N-Benzyloxycarbonyl-pyrrolidin-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetyl-amino)propenes and 1-(N-Benzyloxycarbonylpiperid-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetylamino)propenes

A mixture of the 1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetylamino)propene or the1-(N-benzyloxycarbonylpiperid-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetylamino)propene (2.00 mmol), tetrabutylammonium chloride (2.00 mmol), and palladium(II) acetate (0.089 g, 0.40 mmol, 0.2 eq) in a solution of triethylamine (8 mL) and anhydrous N,N-dimethylformamide (4 mL) was heated at reflux under nitrogen for 2 hours. The resulting reaction mixture was evaporated under reduced pressure, and the residue was partitioned between ethyl acetate (25 mL) and water (25 mL). The ethyl acetate layer was removed, and the aqueous layer was extracted with additional ethyl acetate (25 mL). The organic extracts were combined, dried ($MgSO_4$), and evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 50 g) amd elution with either a diethyl ether gradient in methylene chloride or an acetone gradient in methylene chloride to afford the corresponding 3-(N-benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole or the 3-(N-benzyloxycarbonylpiperid-2-ylmethyl)-1H-indole.

Following this procedure the following compounds were prepared:

A. **(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole**

(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-iodophenyl)-N-trifluoroacetyl-amino)propene was used.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

Column chromatography afforded the title compound as a clear, pale brown oil: ^{1}H NMR ($CDCl_3$) δ 8.05 (br s, indole NH), 7.49-7.34 (m, 7H), 7.17 (br t, 1H), 7.02 (br s, 1H), 6.95 (br s, 1H), 5.24 (s, 2H), 4.28-4.14 (br m, 1H), 3.52-3.41 (m, 2H), 3.28 (br d, 1H), 2.79-2.63 (m, 1H), 1.90-1.70 (m, 4H); LRMS (m/z, relative intensity) 334 (10, M+), 204 (16), 160 (39), 130 (39), 91 (100).

B. **(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-5-(methylaminosulfonylmethyl)-1H-indole**

(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-methylaminosulfonylmethylphenyl)-N-trifluoroacetylamino)propene was used. Column chromatography afforded the title compound as an off-white foam: IR ($CHCl_3$) 1673, 1410, 1358, 1324, 1118, 1092 cm^{-1}; LRMS (m/z, relative intensity) 441 (9, M+), 237 (29), 204 (77), 160 (97), 143 (73), 91 (100); HRMS: calculated for $C_{13}H_{27}N_3O_4S$: 441.1724; found: 441.1704.

C. **(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl-5-cyano-1H-indole**

(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-cyanophenyl)-N-trifluoroacetylamino)propene was used. Column chromatography afforded the title compound as a white foam: IR (1% solution in $CHCl_3$) 2215, 1687 cm^{-1}; ^{13}C NMR [Note: due to slow nitrogen inversion two conformers of the products are seen by NMR spectroscopy] ($CDCl_3$) δ 155.1, 137.9, 137.0, 128.8, 128.5, 128.4, 128.0, 127.8, 124.9, 124.6, 121.0, 114.0, 113.9, 112.1, 102.3, 67.2, 66.7, 58.5, 57.6, 47.0, 46.7, 30.3, 30.0, 29.6, 28.8, 23.6, 22.7. Anal. calcd for $C_{22}H_{21}N_3O_2 \cdot 0.25\ C_2H_4O_2$ [acetic acid]: C, 72.17; H, 5.92; N, 11.22. Found: C, 72.28; H, 5.76; N, 10.95.

D. **(R,S)-3-(N-Benzyloxycarbonylpiperid-2-ylmethyl)-5-(methylaminosulfonylmethyl-1H-indole**

(R,S)-1-(N-Benzyloxycarbonylpiperid-2-yl)-3-(N-(2-bromo-4-methylaminosulfonyl-methylphenyl)-N-trifluoroacetylamino)propene was used. Column chromatography afforded the title compound as an off-white foam: ^{13}C NMR

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

[Note: due to slow nitrogen inverion two conformers of the products are since by NMR spectroscopy] ($CHCl_3$) δ 162.5, 136.9, 136.2, 128.4, 127.6, 124.5, 123.3, 120.8, 120.3, 111.5, 66.8, 57.4, 39.5, 36.5, 31.4, 29.8, 25.8, 25.5, 18.8; LRMS (m/z, relative intensity) 445 (5, M+), 361 (4), 238 (40), 218 (80), 174 (100), 143 (53); HRMS calculated for $C_{24}H_{29}N_3O_4S$: 455.1880; found: 455.1899.

EXAMPLE 8

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-5-methoxy-1H-indole

To a stirred mixture of lithium borohydride (0.092 g, 4.22 mmol, 2 eq) in anhydrous tetrahydrofuran (5 mL) at 0°C was added a solution of the (R)-3-(N-benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-5-methoxy-1H-indole (0.80 g, 2.11 mmol) in anhydrous tetrahydrofuran (8 mL). The resultant mixture was heated at reflux under nitrogen for 1 hour. The reaction mixture was cooled, and water (1 mL) was added carefully, followed by ethyl acetate (20 mL). The resultant mixture was stirred at room temperature for 30 minutes, dried ($MgSO_4$), filtered through diatomaceous earth, and the filtrate was evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 50 g) and elution with ethyl acetate/hexanes [1:1] afforded (R)-3-(N-benzyloxycarbonylpyrrolidin-2-ylmethyl)-5-methoxy-1H-indole as a colorless gum: ^{13}C NMR [Note: due to slow nitrogen inversion two conformers of the products are seen by NMR spectroscopy] ($CDCl_3$) δ 162.5, 136.9, 1.36.2, 128.4, 127.8, 127.6, 124.5, 123.3, 120.8, 120.3, 111.5, 66.8, 57.4, 39.5, 36.5, 31.4, 29.8, 25.8, 25.5, 18.8; LRMS (m/z, relative intensity) 364 (30, M+), 204 (17), 160 (92), 145 (17), 117 (13), 91 (100). Anal. calcd for $C_{22}H_{24}N_2O_3 \cdot 0.5\ H_2O$: C, 70.76; H, 6.75; N, 7.50. Found: C, 70.70; H, 6.94; N, 7.15.

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

EXAMPLE 9

General Procedure for the Formation of 1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetylamino)propenes and 1-(N-Benzyloxycarbonylpiperid-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetylamino)propenes from the Mitsunobu Coupling of 2-Halo-N-trifluoroacetylanilines with 1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene or 1-(N-Benzyloxycarbonylpiperid-2-yl)-3-hydroxypropene

To a stirred mixture of 1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene or 1-(N-benzyloxycarbonylpiperid-2-yl)-3-hydroxy-propene (R, or S, or racemate 2.00 mmol), the 2-halo-N-trifluoroacetylaniline (2.5 mmol, 1.25 eq), and triphenylphosphine (0.655 g, 2.50 mmol, 1.25 eq) in anhydrous tetrahydrofuran at 0°C under a nitrogen atmosphere was added diethyl azodicarboxylate (0.39 mL, 2.48 mmol, 1.25 eq) dropwise. The reaction solution was slowly warmed to 25°C over the course of 2 hours, and then stirred at 25°C under a nitrogen atmosphere for an additional 12 hours. The resulting reaction solution was evaporated under reduced pressure, and the residue was column chromatographed using silica gel (approximately 50 g) and elution with either a diethyl ether gradient in hexanes or an ethyl acetate gradient in hexanes to afford the corresponding 1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetylamino)propene or 1-(N-benzyloxycarbonylpiperid-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetylamino)propene.

Following this procedure the following compounds were prepared:

A. **(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-iodophenyl)-N-trifluoroacetylamino)propene**

(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene and 2-iodo-N-trifluoro-acetylaniline were used. Column chromatography afforded the title compound as a clear, colorless oil: ^{1}H NMR ($CDCl_3$) δ 7.88 (br d, 1H), 7.43-6.89 (m, 10H), 5.70-5.35 (m, 2H), 5.13 (br s, 2H),

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

5.00-4.75 (m, 1H), 4.40-4.29 (m, 1H), 3.60-3.42 (m, 3H), 2.05-1.45 (m, 4H); LRMS (FAB, m/z, relative intensity) 559 (100, [MH+]), 515 (52), 451 (15), 244 (7).

B. **(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-methylaminosulfonylmethylphenyl)-N-trifluoroacetylamino)propene**

(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene and 2-bromo-4-methylaminosulfonylmethyl-N-trifluoroacetylaniline were used. Column chromatography using elution with 4% acetone in methylene chloride afforded the title compound as a white foam (44%): FAB LRMS (m/z, relative intensity) 620 ([MH+ with ^{81}Br], 618 ([MH+ with ^{79}Br], 98), 576 (50), 574 (63), 512 (17), 484 (33).

C. **(R)-1-(N-Benzyloxycarbonlypyrrolidin-2-yl)-3-(N-(2-bromo-4-cyanophenyl)-N-trifluoroacetylamino)propene**

(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene and 2-bromo-4-cyano-N-trifluoroacetylaniline were used. Column chromatography using elution with a gradient of diethyl ether (5% - 100%) in methylene chloride afforded the title compound as a clear, colorless oil: IR ($CHCl_3$) 2231, 1702, 1157 cm^{-1}; LRMS (m/z, relative intensity) 537 ([MH+ with ^{81}Br], 13), 535 ([MH+ with ^{79}Br], 13), 402 (29), 400 (30), 294 (55), 292 (57), 244 (80), 213 (89), 91 (100); Anal. calcd for $C_{24}BrF_3H_{21}N_3O_3 \cdot 0.2\ H_2O$: C, 53.39; H, 3.99; N, 7.78. Found: C, 53.25; H, 3.95; N, 7.98.

D. **(R,S)-1-(N-Benzyloxycarbonylpiperid-2-yl)-3-(N-(2-bromo-4-methylaminosulfonylmethylphenyl)-N-trifluoroacetylamino)propene**

(R,S)-1-(N-Benzyloxycarbonylpiperid-2-yl)-3-hydroxypropene and 2-bromo-4-methylaminosulfonylmethyl-N-trifluoroacetylaniline were used. Column chromatography using elution with 20% acetonitrile in methylene chloride afforded the title compound as a white foam: FAB LRMS (m/z, relative intensity) 634 ([MH+ with ^{81}Br], 26), 632 ([MH+ with ^{79}Br], 22), 590 (35), 588 (43), 401 (33), 327 (48), 281 (75),

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

207 (90), 147 (100); FAB HRMS: calculated for $C_{26}H_{29}BrF_3N_3O_5S$• [H+] 632.1043, found 632.1047 [for ^{79}Br and ^{32}S].

EXAMPLE 10

General Synthesis of 2-Halo-N-trifluoroacetylanilines from Reaction of 2-Haloanilines and Trifluoroacetic Anhydride

To a stirred solution of the 2-haloaniline (2.00 mmol) and pyridine (0.18 mL, 2.22 mmol, 1.1 eq) in anhydrous methylene chloride (10 mL) at 0°C under a nitrogen atmosphere was added dropwise trifluoroacetic anhydride (0.31 mL, 2.19 mmol, 1.1 eq). The resultant reaction mixture was stirred at 0°C under a nitrogen atmosphere for 3 hours. A saturated solution of sodium hydrogen carbonate was added (15 mL), and this aqueous mixture was extracted with ethyl acetate (3 x 15 mL). The extracts were combined, dried ($MgSO_4$), and evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 50 g) and elution with an ethyl acetate gradient in hexanes to afford the corresponding 2-halo-N-trifluoroacetylaniline.

Following this procedure the following compounds were prepared:


A. **2-Iodo-N-trifluoroacetylaniline**

2-Iodoaniline was used. Evaporation of the ethyl acetate extracts afforded the title compound directly as a white solid: mp, 105.0-106.5°C; FAB LRMS (m/z relative intensity) 316 ([MH+], 8), 155 (80), 135 (26), 119 (100); ^{13}C NMR (acetone-d_6) δ 206.2, 140.4, 130.2, 130.1, 128.2.

B. **2-Bromo-4-methylaminosulfonylmethyl-N-trifluoroacetylaniline**

2-Bromo-4-methylaminosulfonylmethylaniline was used. Evaporation of the ethyl acetate extracts afforded the title compound directly as a white solid: mp, 164.0-166.0°C. Anal. calcd for $C_{10}H_{10}BrF_3N_2O_3S$: C, 32.02; H, 2.69; N, 7.47. Found: C, 32.18; H, 2.67; N, 7.30.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

C. **2-Bromo-4-cyano-N-trifluoroacetylaniline**

2-Bromo-4-aminocarbonylaniline was used. Dehydration of the carboxamide also occurred in this reaction. Column chromatography using ethyl acetate/hexanes afforded the title compound as a white solid: mp, 125-130°C; ^{1}H NMR (DMSO-d_6) δ 11.6 (br s, NH), 8.37 (d, J=1.8 Hz, 1H), 7.96 (dd, J=1.8 and 8.2 Hz, 1H), 7.71 (d, J=8.2 Hz, 1H).

EXAMPLE 11

General Procedure for the Bromination of Anilines to Form 2-Bromoanilines

To a stirred solution of the aniline (2.00 mmol) and sodium hydrogen carbonate (0.21 g, 2.50 mmol, 1.25 eq) in methanol (10 mL) at 0°C was added dropwise bromine (0.113 mL, 2.19 mmol, 1.1 eq). The resulting reaction mixture was then stirred at 25°C for 30 minutes. The reaction mixture was then evaporated under reduced pressure, and the residue was placed in a saturated solution of sodium hydrogen carbonate (10 mL). This aqueous mixture was extracted with ethyl acetate (3 x 15 mL). The extracts were combined, dried ($MgSO_4$), and evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 50 g) and elution with an appropriate solvent system to afford the corresponding 2-bromoaniline.

Following this procedure the following compounds were prepared:

A. **2-Bromo-4-methylaminosulfonylmethylaniline**

4-Methylaminosulfonylmethylaniline (M.D. Dowle, et al. Eur. Pat. Appl. EP225,726) was used. Column chromatography using elution with 40% ethyl acetate in hexanes afforded the title compound as a white solid: mp, 104.0-107.0°C. Anal. calcd for $C_9H_{11}BrN_2O_2S$: C, 34.42; H, 3.97; N, 10.04. Found: C, 34.66; H, 3.96; N, 9.96.

B. **4-Aminocarbonyl-2-bromoaniline**

4-Aminobenzamide was used. Column Chromatography using elution with a ethyl acetate gradient (25-50%) in methylene chloride afforded the title compound as a white solid: mp,

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

144.5-146.0°C; ^{1}H NMR (DMSO-d_6) δ 7.93 (d, J=2.0 Hz, 1H), 7.70 (br s, amide NH), 7.62 (dd, J=2.0 and 8.5 Hz, 1H), 7.05 (br s, amide NH), 6.77 (d, J=8.5 Hz, 1H), 5.85 (s, aniline NH_2).

EXAMPLE 12

1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene or 1-(N-Benzyloxycarbonylpiperid-2-yl)-3-hydroxypropene


To a stirred solution of either ethyl 3-(N-benzyloxycarbonylpyrrolidin-2-yl)-2-propenoate or ethyl-3-(N-benzyloxycarbonylpiperid-2-yl)-2-propenoate (R, or S, or racemate, 10.00 mmol) in anhydrous tetrahydrofuran (75 mL) at -78°C under nitrogen was added dropwise a solution of diisobutylaluminium hydride (1.0 M in hexanes, 12.0 mL, 22.0 mmol, 2.2 eq). The resulting solution was stirred at -78°C under nitrogen for 30 minutes. The reaction solution was then allowed to warmed to room temperature over the course of 2 hours. A saturated solution of sodium hydrogen carbonate (50 mL) was added, and the aqueous mixture was extracted with ethyl acetate (3 x 50 mL). The extracts were combined, dried ($MgSO_4$), and evaporated under reduced pressure. Column chromatography of the residue with diethyl ether/hexanes [1:1] afforded either 1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene or 1-(N-benzyloxycarbonyl-piperid-2-yl)-3-hydroxypropene.

Following the procedure the following compounds were prepared:

A. **(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene**

(R)-Ethyl 3-(N-benzyloxycarbonylpyrrolidin-2-yl)-2-propenoate was used. Chromatography of the extraction residue afforded the title compound as a clear, colorless oil: ^{1}H NMR ($CDCl_3$) δ 7.40-7.25 (m, 5H), 5.75-5.53 (m, 2H), 5.20-5.00 (m, 2H), 4.38 (br m, 1H), 4.06 (br d, J=13.7 Hz, 2H), 3.45 (br t, J=7.0 Hz, 1H), 2.03-1.68 (m, 4H); $[\alpha]^{25}$ = +34° (MeOH, c=1.0); HRMS: calculated for $C_{15}H_{19}NO_3$ 261.1365, found 261.1356.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

B. **(R,S)-1-(N-Benzyloxycarbonylpiperid-2-yl)-3-hydroxypropene**

(R,S)-Ethyl 3-(N-benzyloxycarbonylpiperid-2-yl)-2-propenoate was used. Chromatography of the extraction residue afforded the title compound as a clear, colorless oil: LRMS (m/z, relative intensity) 257 (3), 212 (12), 193 (8), 175 (65), 173 (100), 145 (27), 109 (24), 91 (87); ^{1}H NMR ($CDCl_3$) δ 7.40-7.20 (m, 5H), 5.70-5.61 (m, 2H), 5.14 (d, J=17.6 Hz, 1H), 5.10 (d, J=17.5 Hz, 1H), 4.88 (br m, 1H), 4.14-4.00 (m, 3H), 2.91 (br t, J=12.7 Hz, 1H), 1.78-1.47 (m, 6H). Anal. calcd for $C_{16}H_{21}NO_3 \cdot 0.1\ H_2O$: C, 69.34; H, 7.71; N, 5.05. Found: 69.38; H, 7.84; N, 5.16.

EXAMPLE 13

Synthesis of Ethyl 3-(N-Benzyloxycarbonylpyrrolidin-2-yl)-2-propenoate or Ethyl 3-(N-Benzyloxycarbonylpiperid-2-yl)-2-propenoate

To a stirred solution of N-carbobenzyloxypyrrolidine-2-carboxaldehyde or N-carbobenzyloxypiperidine-2-carboxaldehyde (5.00 mmol) [S. Kiyooka, et al., J. Org. Chem., 5409 (1989) and Y. Hamada, et al., Chem. Pharm. Bull., 1921 (1982)] in anhydrous tetrahydrofuran at -78°C was added (carbethoxymethylene)triphenylphosphorane (2.09 g, 6.00 mmol. 1.2 eq) as a solid portionwise. The resulting reaction mixture was stirred at room temperature under nitrogen for 2 hours, and then heated at reflux under nitrogen for 1 hour. The reaction mixture was evaporated under reduced pressure and the residue was column chromatographed using silica gel (approximately 100 g) and elution with 20% diethyl ether in hexanes to afford either ethyl 3-(N-benzyloxycarbonylpyrrolidin-2-yl)-2-propenoate or ethyl 3-(N-benzyloxycarbonylpiperid-2-yl)-2-propenoate.

A. **(R)-Ethyl 3-(N-Benzyloxycarbonylpyrrolidin-2-yl)-2-propenoate**

(R)-N-Carbobenzyloxypyrrolidine-2-carboxaldehyde was used. Chromatography as described above afforded the title compound as a clear, colorless oil: ^{1}H NMR ($CDCl_3$-d_6) δ 7.34-

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

7.25 (m, 5H), 6.89-6.76 (m, 1H), 5.88-5.74 (m, 1H), 5.18-5.05 (m, 2H), 4.60-4.43 (m, 1H), 4.17 (q, J=7.1 Hz, 2H), 3.55-3.40 (m, 2H), 2.11-2.00 (m, 1H), 1.90-1.75 (m, 3H), 1.28 (t, J=7.1 Hz, 3H); ^{13}C NMR ($CDCl_3$) [Note: due to slow nitrogen inversion two conformers of the products are seen by NMR spectroscopy] δ 166.3, 154.7, 147.9, 147.4, 136.6, 128.4, 127.9, 120.9, 66.9, 65.8, 60.4, 58.1, 57.7, 46.8, 46.4, 31.6, 30.8, 23.6, 22.8, 22.6, 15.3, 14.2.

B. **(R,S)-Ethyl 3-(N-Benzyloxycarbonlpiperid-2-yl)-2-propenoate**

(R,S)-N-Carbobenzyloxypiperidine-2-carboxaldehyde was used. Chromatography as described above afforded the title compound as a clear, colorless oil: ^{1}H NMR ($CDCl_3$-d_6) δ 7.36-7.27 (m, 5H), 6.85 (dd, J=4.4 and 16.3 Hz, 1H), 5.80 (dd, J=2.4 and 16.3 Hz, 1H), 5.11 (s, 2H), 5.01 (br m, 1H), 4.17 (q, J=6.7 Hz, 2H), 4.05 (br d, J=12.6 Hz, 1H), 2.87 (br t, 1H), 1.80-1.35 (m, 6H), 1.27 (t, J=6.6 Hz, 3H); FAB LRMS (m/z, relative intensity) 318 ([MH+], 100), 274 (86), 228 (14), 210 (21), 182 (43), 138 (32).

EXAMPLE 14

(R)-5-Amino-3-(N-methylpyrrolidin-2ylmethyl)indole

A mixture of (R)-5-dibenzylamino-3-(N-methylpyrrolidin-2-ylmethyl)indole (1.08 g, 2.64 mmol) and palladium [II] hydroxide on carbon (0.6 g) in absolute ethanol (25 mL) was shaken under a hydrogen atmoshpere (3 atm) at 40°C for 4 hours. The resulting mixture was filtered through diatmaceous earth, and the filtrate was evaporated under pressure to afford the title compound (0.60 g, 2.62 mmol, 99%) as a white foam: ^{1}H NMR (DMSO-d_6) δ 10.65 (br s, NH), 7.14 (d, J=2.2 Hz, 1H), 7.12 (d, J=8.6 Hz, 1H), 6.85 (d, J=1.6 Hz, 1H), 6.60 (dd, J=2.0 and 8.6 Hz, 1H), 3.63-2.83 (m, 7H), 2.78 (s, 3H), 2.05-1.67 (m, 4H); $[\alpha]^{25}$=+9° (MeOH, c=1.0); HRMS: calculated for $C_{14}H_{19}N_3$: 229.1575; found 229.1593.

THIS SHEET HAS BEEN REPLACED



EXAMPLE 15

General Synthesis of 5-Carbonylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indoles and 5-Sulfonylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indoles

To a stirred solution of (R)-5-amino-3-(N-methylpyrrolidin-2-ylmethyl)indole (0.229 g, 1.00 mmol) and triethylamine (0.21 mL, 1.5 mmol, 1.5 eq) in anhydrous acetonitrile (3 mL) at 0°C under nitrogen was added the appropriate carbonyl chloride or sulfonyl chloride (1.5 mmol, 1.5 eq). The resulting reaction mixture was stirred at room temperature for 12 hours. The reaction mixture was then evaporated under reduced pressure, and the residue was column chromatographed using silica gel (approximately 25 g) and elution with an appropriate solvent system afforded the appropriate 5-carbonylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole or 5-sulfonylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole.

Following this procedure the following compounds were prepared:

A. **(R)-5-Benzyloxycarbonylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

Benzyl chloroformate was used. Column chromatography using elution with triethylamine/acetone/ethyl acetate [2:10:88] afforded the title compound as an off-white foam: ^{13}C NMR ($CDCl_3$) δ 163.3, 136.4, 133.6, 129.8, 128.6, 128.2, 127.9, 126.0, 123.2, 113.8, 111.4, 110.1, 66.8, 66.5, 57.5, 40.8, 31.5, 29.8, 21.8; LRMS (m/z, relative intensity) 363 (M+, 12), 279 (7), 184 (7), 171 (33), 108 (100); HRMS: calculated for $C_{22}H_{25}N_3O_2$ 363.1949, found 363.1926. Anal. calcd for $C_{22}H_{25}N_3O_2 \cdot 0.4\ C_4H_4O_4$ [ethyl acetate]: C, 71.09; H, 7.13; N, 10.54. Found: C, 70.82; H, 7.03; N, 10.58.

B. **(R)-3-(N-Methylpyrrolidin-2-ylmethyl)-5-methylsulfonamido-1H-indole**

Methanesulfonyl chloride was used. Column chromatography using elution with triethylamine/acetone/ethyl acetate [1:3:6] afforded the title compound as a white

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

foam: ^{13}C NMR ($CDCl_3$) δ 134.9, 128.3, 128.2, 123.6, 119.3, 115.0, 113.9, 112.0, 66.7, 57.3, 40.7, 38.7, 31.3, 29.4, 21.7; HRMS: calculated for $C_{15}H_{21}N_3O_2S$ [with ^{32}S] 307.1356, found 307.1323.

C. **(R)-5-Acetylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

Acetyl chloride was used. Column chromatography using elution with triethylamine/acetone/ethyl acetate [1:3:6] afforded the title compound as a white foam: ^{13}C NMR (acetone-d_6) δ 168.3, 134.4, 132.2, 128.7, 124.1, 115.7, 113.8, 111.6, 110.2, 67.3, 58.0, 40.9, 31.9, 30.5, 24.1, 22.5; LRMS (m/z, relative intensity) 271 (M+, 39), 241 (4), 207 (5), 187 (20), 144 (20), 84 (100); HRMS: calculated for $C_{16}H_{21}N_3O$ 271.1686, found 271.1693. Anal. calcd for $C_{16}H_{21}N_3O \cdot 1.15\ H_2O$: C, 65.80; H, 8.04; N, 14.39. Found: C, 65.99; H, 7.90; N, 13.99.

D. **(R)-5-N,N-Dimethylaminocarbonylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

Dimethylcarbamyl chloride was used. Column chromatography using elution with methylene chloride/methanol/ammonium hydroxide [9:1:0.1] afforded the title compound as an off white foam: ^1H NMR ($CDCl_3$) δ 8.95 (br s, 1H), 7.49 (br s, 1H), 7.15-7.06 (m, 2H), 6.82 (d, J=1.9 Hz, 1H), 6.44 (br s, 1H), 3.12-3.05 (m, 2H), 3.00 (s, 6H), 2.58-2.40 (m, 2H), 2.40 (s, 3H), 2.18 (br q, J=8.1 Hz, 1H), 1.83-1.47 (m, 4H); ^{13}C NMR ($CDCl_3$) δ 157.2, 133.8, 130.5, 127.7, 123.2, 117.8, 113.0, 112.0, 111.3, 66.5, 57.4, 40.6, 36.4, 31.4, 29.8, 21.7; LRMS (m/z, relative intensity) 300 (M+, 50), 217 (10), 171 (20), 84 (100); HRMS: calculated for $C_{17}H_{24}N_4O$ 300.1952, found 300.1957.

E. **(R)-5-Trifluoroacetylamino-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

Trifluoroacetic anhydride was used. Column chromatography using elution with methylene chloride/methanol/ammonium hydroxide [9:1:0-.1] afforded the title compound as an off white foam: ^1H NMR ($CDCl_3$) δ 8.99 (br s,

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED


1H), 7.80 (br s, 1H), 7.27-7.19 (m, 2H), 6.95 (d, J=1.4 Hz, 1H0, 3.16-3.08 (m, 2H), 2.58 (dd, J=9.4 and 13.5 Hz, 1H). 2.57-2.43 (m, 1H), 2.43 (m, 1H), 2.43 (s, 3H), 2.22 (dd, J=9.2 and 17.5 Hz, 1H), 1.85-1.46 (m, 4H); ^{13}C NMR ($CDCl_3$) δ 134.5, 127.7, 126.9, 123.8, 116.1, 113.9, 111.9, 111.6, 104.1, 66.6, 57.3, 40.6, 31.3, 29.5, 21.7; HRMS: calculated for $C_{16}H_{18}F_3N_3O$ 325.1403, found 325.1378.

EXAMPLE 16

(R)-3-(N-Methylpyrrolidin-2-ylmethyl)-5-(2-methylsulfonamidomethyl)-1H-indole

To a stirred mixture of (R)-5-aminomethyl-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole (0.113 g, 0.46 mmol) and pyridine (50μL, 0.93 mmol, 2.0 eq) in a solution of dimethylformamide and acetontrile (1:3, respectively, 2 mL total) at 0°C under nitrogen was added methanesulfonyl chloride dropwise (44μL, 0.56 mmol, 1.3 eq). The resulting reaction solution was stirred at room temperature under nitrogen for 1 hour, and then it was evaporated under reduced pressure. The residual oil was column chromatographed using silica gel (6 g) and elution with methylene chloride/methanol/ammonium hydroxide [9:1:0.1] to afford the title compound (0.044 g, 0.14 mmol, 30%) as a white foam: ^1H NMR ($CDCl_3$) δ 8.25 (br s, NH), 7.54 (br s, 1H), 7.35 (d, J=8.4 Hz, 1H), 7.17 (dd, J=1.6 and 8.4 Hz, 1H), 7.06 (d, J=1.8 Hz, 1H), 4.78 (br s, NH), 4.42 (s, 2H), 3.20-3.12 (m, 2H), 2.87 (s, 3H), 2.64 (dd, J=9.4 and 13.9 Hz, 1H). 2.54-2.43 (m, 1H), 2.47 (s, 3H), 2.25 (dd, J=9.3 and 17.3, 1H), 1.86-1.52 (m, 4H); ^{13}C NMR ($CDCl_3$) δ 135.8, 127.8, 127.3, 123.0, 122.0, 118.5, 113.7, 111.6, 66.7, 57.4, 47.9, 40.9, 40.7, 31.3, 29.5, 21.7; LRMS (m/z relative intensity) 321 (28), 320 (M+, 26), 237 (51), 157 (100), 143 (64), 129 (78); HRMS: calculated for $C_{16}H_{22}N_3O_2S$ 320.1435, found 320.1453.

THIS SHEET HAS BEEN REPLACED



EXAMPLE 17

General Synthesis of Allylsulphonamides

A. **Allylsulphonamide**

The title compound was prepared by the method of M. A. Belous and I. Ya. Postouski, Zhur. Obschei. Khim., 1950, 20, 1701.

B. **N-Methylallylsulphonamide**

The title compound was prepared by an analogous procedure to above by using methylamine instead of ammonia. Anal. Calcd for $C_4H_9NO_2S$: C,40.25; H,7.43; N,9.38. Found: C,40.51; H,7.37; N,9.70.

EXAMPLE 18

Preparation of Ethylallylsulphone

The title compound was prepared by the method of R. J. Palmer and C. J. M. Stirling., J. Amer. Chem. Soc., 1980, 102, 7888.

EXAMPLE 19

General Synthesis of Vinyl Sulphonamides

Where the required vinylsulphonamide was not commercially available, they were prepared by the following procedure based on the procedure described in Zhur. Obschei. Khim., 1959, 29, 1494.

A. **N,N-Dimethylvinylsulphonamide**




To a stirred solution of chloroethylsulphonyl chloride (25 g, 153 mmol) in dry diethyl ether (150 mL) at -10°C, was added dropwise a solution of dimethylamine (30.5 mL, 460 mmol) in dry diethyl ether (100 mL) over 5 minutes. After stirring for 90 minutes at -10°C the solution was filtered and evaporated in vacuo. The residue was distilled to give the title compound (9.5 g, 46%): b.p. 120-122°C (20 mm Hg). Anal. Calcd for $C_4H_9NO_2S$: C,35.54; H,6.71; N,10.36%. Found: C,35.36; H,6.37; N,10.19.

B. The following examples were prepared by the general procedure above, using the appropriate amine starting material. Purification was by distillation or column chromatography.

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

R_2NSO_2-CH=CH_2

R_2N	Isolated Form	Analysis % (Theoretical in brackets) C	H	N
MeNH-	Oil b.p. 93-5°C (0.05 mm Hg)	Literature compound U.S. 3,761,473		
	Oil	47.97 (47.97	7.41 7.48	7.81 7.99)
	Oil	44.73 (44.70	6.80 6.88	8.62 8.69)
nPr_2N-	Oil	50.37 (50.23	8.79 8.96	7.68 7.32
nPrNH-	Oil	40.22 (40.24	7.35 7.43	9.1 9.39)
	Oil	40.51 (40.79	5.85 6.16	9.35 9.52)
iPrNH-	Oil	40.42 (40.25	7.33 7.43	9.30 9.39)

EXAMPLE 20

General Synthesis of Vinyl Sulphones

Where the required vinyl sulphone was not commercially available, they were prepared from the corresponding thiols using the procedure described by J. M. Gaillot, Y Gelas-Mialhe and R. Vessiere Can. J. Chem., 1979, 57, 1958. The following examples are representative.

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

$RS\text{-}CH_2CH_2\text{-}OH$

R	Isolated Form	Analysis % (Theoretical in brackets) C	H
nPr	Oil 1/16 EtOAc 1/5 H_2O	48.68 (48.76	9.79 10.06)
nBu	Oil	T.l.c - Rf. 0.26 (SiO_2, Ether/Hexane 1:1)	

$RS\text{-}CH_2CH_2\text{-}Cl$

R	Isolated Form	Analysis % (Theoretical in brackets) C	H
nPr	Oil 1/5 H_2O 1/30 CH_2Cl_2	41.63 (41.65	7.60 7.69)
nBu	Oil 1.0 H_2O	42.31 (42.21	7.84 8.27)

$RSO_2\text{-}CH_2CH_2Cl$

R	Isolated Form	Analysis % (Theoretical in brackets) C	H
nPr	Oil	34.75 (35.19	6.68 6.50)
nBu	Oil 1/15 CH_2Cl_2	38.41 (38.27	7.01 6.95)

$RSO_2\text{-}CH{=}CH_2$

R	Isolated Form	Analysis % (Theoretical in brackets) C	H
nBu	Oil	48.95 (48.62	8.07 8.16)

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced


EXAMPLE 21

General Synthesis of indoles with 5-alkenyl substituents

A. **(R)-5-trans-(2-N,N-Dimethylaminocarbonylethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole**

A mixture of N,N-dimethylacrylamide (134 μL, 1.3 mmol), tri-o-tolylphosphine (91 mg, 0.3 mmol), palladium (II) acetate (15 mg, 0.07 mmol), triethylamine (280 μL, 2 mmol) and (R)-5-bromo-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole was dissolved in anhydrous acetonitrile (5 mL) and refluxed for 24 hours under nitrogen. The reaction was partitioned between ethylacetate and aqueous sodium carbonate. The dried (Na_2SO_4) organic phase was evaporated and the residue purified by column chromatography on silica gel, eluting with CH_2Cl_2: MeOH: NH_4OH 96:3.5:0.5 to afford the title compound as a white foam (145 mg, 47%). Anal. Calcd for $C_{19}H_{25}N_3O \cdot 1/9\ CH_2Cl_2$: C,71.56; H,7.87; N,13.10%. Found: C,71.29; H,8.15; N,13.05.

B. The following examples were prepared using the above procedure with the appropriate alkene starting material (available commercially, or prepared by routes outlined in this patent).




THIS SHEET HAS BEEN REPLACED



This Sheet Has Been Replaced

R^2	Isolated Form	Analysis % (Theoretical in brackets) C	H	N	$[\alpha]^{25}$ D (c=0.1 MeOH)
$MeSO_2CH=CH$	Foam 3/10 CH_2Cl_2	60.45 (60.42	6.43 6.62	8.33 8.15)	
$PhSOCH=CH$	Foam 1/10 CH_2Cl_2	68.04 (68.24	6.27 6.27	6.99 7.20)	
$NH_2SO_2CH=CH$	Foam 1/3 MeOH 1/3 H_2O	58.56 (58.39	6.80 6.85	12.19 12.51)	
$EtSOCH=CH$	Foam 1/20 CH_2Cl_2 1/4 H_2O	66.70 (66.66	7.35 7.62	8.64 8.62)	
	Foam 1/8 CH_2Cl_2 1/2 H_2O	61.74 (61.49	6.93 7.22	10.53 10.69)	
$nBuSO_2CH=CH$	Foam 1/4 CH_2Cl_2 1/10 EtOH	63.56 (63.59	7.77 7.57	7.22 7.25)	
$Me_2NSO_2CH=CH$	Foam 1/3 H_2O	61.14 (61.19	7.06 7.27	11.57 11.89)	

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED


R^2	Isolated Form	Analysis % (Theoretical in brackets) C	H	N	$[\alpha]_D^{25}$ (c=0.1 MeOH)
	Foam 1/2 CH_2Cl_2 1/4 H_2O	59.64 (59.43	6.82 7.07	9.83 9.67)	
$nPr_2NSO_2CH=CH$	Foam 1.0 H_2O 1/10 CH_2Cl_2	61.48 (61.72	7.76 8.25	9.69 9.77)	
$nPrNHSO_2CH=CH$	Foam 1/10 CH_2Cl_2 1/3 H_2O	61.07 (61.01	7.12 7.49	10.91 11.18	
	Foam 1/3 CH_2Cl_2 1 H_2O	56.83 (56.39	6.40 6.87	10.36 10.36)	+34°
$iPrNHSO_2CH=CH$	Foam 1/6 CH_2Cl_2	61.03 (61.27	7.42 7.33	11.17 11.19)	+30°
$PhSO_2CH_2CH=CH$	Foam 1/6 CH_2Cl_2	68.21 (68.27	6.81 6.50	7.15 6.87)	
$Me_2NSO_2CH_2CH=CH$	Foam 1/20 CH_2Cl_2	62.54 (62.55	7.50 7.46	11.21 11.48)	


This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

R^2	Isolated Form	Analysis % (Theoretical in brackets) C	H	N	$[\alpha]_D^{25}$ (c=0.1 MeOH)
$NH_2SO_2CH_2CH=CH$	Foam 0.1 CH_2Cl_2 1.0 MeOH	58.50 (58.13	6.93 7.30	11.22 11.24)	
$EtSO_2CH_2CH=CH$	Foam	65.56 (65.86	7.47 7.56	8.00 8.08)	
$PhCONHCH_2CH=CH$	Foam 0.1 CH_2Cl_2	75.69 (75.78	6.97 7.18	10.76 11.00)	+70°
$MeSO_2NHCH_2CH=CH$	Foam 0.1 CH_2Cl_2	61.05 (61.07	7.31 7.14	11.12 11.80)	

C) The following compounds could be prepared by the procedure a) above but using the corresponding beta-chloroethylsulphone as starting material instead of an alkene. These reactions were preferably carried out in the presence of 3-6 equivalents of triethylamine.



R^2	Isolated Form	Analysis % (Theoretical in brackets) C	H	N	$[\alpha]_D^{25}$ (c=0.1 MeOH)
$nPrSO_2CH=CH$	Foam 1/8 CH_2Cl_2 1/3 H_2O	62.93 (63.25	7.15 7.47	7.71 7.71)	
	Foam 0.15 CH_2Cl_2	62.22 (62.20	5.37 5.49	6.52 6.55)	+48°

This Sheet Has Been Replaced

EXAMPLE 22


General Procedure for Hydrogenation of 5-alkenylindoles

A typical procedure is as follows:

A. (R)-5-(2-Aminosulphonylethyl)-3-(N-methyl-pyrrolidin-2-ylmethyl)-1H-indole


(R)-5-(2-Aminosulphonylethenyl)-3-(N-methyl-pyrrolidin-2-ylmethyl)-1H-indole (157 mg, 0.5 mmol) was dissolved in absolute ethanol (10 mL) and added to a solution of ethanolic hydrogen chloride 25 ml (prepared from acetyl chloride (38 μL, 0.53 mmol) and absolute ethanol (25 mL)). 10% palladium-on-carbon (125 mg) was added. This solution was hydrogenated under a hydrogen atmosphere (15 p.s.i.) at room temperature for 18 hours. The resultant reaction mixture was filtered through diatomaceous earth (Celite trademark or Arbacell-trademark)) washed with absolute ethanol and the combined filtrates evaporated in vacuo. The residue was purified by column chromatography on silica gel, eluting with a gradient solvent mixture up to CH_2Cl_2: MeOH:NH_4OH 93:7:1 to give the title compound as a colourless oil (80 mg, 51%). Anal. Calcd for $C_{16}H_{23}N_3O_2S \cdot 1/4$ MeOH. 1/3 H_2O: C,58.21; H,7.36; N,12.54. Found: C,58.60; H,7.40; N,12.57. $[\alpha]^{25} = +69°$ (c=0.1, MeOH).


B. The following examples were prepared by an analogous procedure to a) above.



THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced


R^2	Isolated Form	Analysis % (Theoretical in brackets) C	H	N	$[\alpha]_D^{25}$ (c=0.1 MeOH)
$Me_2NSO_2CH_2CH_2$	Oil 1/20 CH_2Cl_2	51.52 (51.31	7.40 7.67	11.49 11.89	+48°
$Me_2NCOCH_2CH_2$	Oil 1/10 CH_2Cl_2	70.96 (71.29	8.52 8.46	12.84 13.06)	+78.2°
$MeSO_2CH_2CH_2$	Gum 1/4 H_2O	52.76 (52.83	7.29 7.60	8.41 8.62)	+83°
$EtSOCH_2CH_2$	Oil	51.39 (51.27	7.69 8.03	8.18 7.83)	
	Foam 2/3 H_2O	52.73 (52.81	7.60 8.11	10.64 10.47)	+57°
$PhSO_2CH_2CH_2CH_2$	Oil 2/3 H_2O	57.56 (57.80	7.27 7.23	6.96 6.85)	

R^2	Isolated Form	Analysis % (Theoretical in brackets) C	H	N	$[\alpha]_D^{25}$ (c=0.1 MeOH)
$NH_2SO_2CH_2CH_2CH_2$	Foam 0.65 CH_2Cl_2	56.78 (56.26	7.16 6.80	10.34 10.75)	
	Foam 1/2 H_2O	62.45 (62.47	7.48 7.86	10.74 10.93)	
$Me_2NSO_2CH_2CH_2CH_2$	Oil 0.1 CH_2Cl_2	52.03 (51.66	7.76 7.91	10.41 11.16)	
$^nBuSO_2CH_2CH_2$	Oil 1/3 CH_2Cl_2	52.28 (52.48	7.50 7.91	7.23 7.17)	+48°
$nPrNHSO_2CH_2CH_2$	Foam 1/4 H_2O	52.07 (52.01	7.95 8.08	11.17 11.42)	+57°
$^nPrSO_2CH_2CH_2$	Foam 1/20 CH_2Cl_2 3/4 H_2O	52.80 (52.47	7.72 8.15	7.24 7.65)	+50°
$^nPr_2NSO_2CH_2CH_2$	Gum 1.0 H_2O	52.28 (52.37	8.38 8.80	10.03 9.92)	+40°

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

R^2	Isolated Form	Analysis % (Theoretical in brackets) C	H	N	$[\alpha]_D^{25}$ (c=0.1 MeOH)
$EtSO_2CH_2CH_2CH_2$	Glass 0.5 CH_2Cl_2	59.10 (59.90	7.57 7.47	7.04 7.18)	
	Foam 1/3 CH_2Cl_2	59.07 (59.56	7.10 7.15	10.80 10.78)	+30°
$iPrNHSO_2CH_2CH_2$	Foam 1/8 CH_2Cl_2	61.59 (61.39	7.88 7.88	11.16 11.23)	+58°

EXAMPLE 23

General Synthesis of (R)-5-(2-Ethylsulphonylethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole

A. **(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-yl-methyl)-5-bromo-1H-indole**

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-yl-carbonyl)-5-bromo-1H-indole(0.67 g, 1.57 mmol) was dissolved in anhydrous tetrahydrofuran (20 mL) and at room temperature under nitrogen was added lithium borohydride (2 molar in tetrahydrofuran) (1.2 mL, 2.4 mmol). The reaction mixture was stirred at room temperature for 3 hours and warmed to reflux for 16 hours. After cooling to room temperature, 2NHCl (10 mL) was added dropwise and the reaction mixture partitioned between ethyl acetate and water. The separated organic phase was washed with saturated aqueous sodium hydrogen carbonate (2x), brine (1x), dried (Na_2SO_4), and evaporated in vacuo to give a colourless oil. Purification by column chromatography on silica gel, eluting with dichloromethane gave the title compound as an oil (0.32 g). TLC (SiO_2:CH_2Cl_2): R_f=0.2.

B. **(R)-5-(Ethylsulphonylethenyl)-3-(N-Benzyloxy carbonylpyrrolidin-2-ylmethyl)-1H-indole**

The compound from procedure a) above was coupled with ethyl vinylsulphone under standard conditions described above, to give the title compound as a foam. Anal. Calcd for $C_{25}H_{28}N_2O_4S \cdot 1/8\ CH_2Cl_2$: C,65.15; H,6.15; N,6.05. Found: C,65.16; H,6.17; N,5.97. $[\alpha]^{25}$ = -50° (0.1, MeOH).

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

C. **(R)-5-(2-Ethylsulphonylethyl)-3(pyrrolidin-2-ylmethyl)-1H-indole**

The compound prepared in procedure b) above, was hydrogenated under the standard condition described above, to give the title compound as a foam. Anal. Calcd for $C_{17}H_{24}N_2O_2S \cdot 1/2\ CH_2Cl_2$: C,63.07; H,7.48; N,8.63. Found: C,62.90; H,7.25; N,8.58. $[\alpha]^{25} = -11°$ (c=0.1, MeOH).

EXAMPLE 24

General Synthesis of (R)-3-(N-alkyl-pyrrolidin-2-ylmethyl)indoles


A. **(R)-3-(N-Ethylpyrrolidin-2-ylmethyl)-5-(2-ethylsulphonylethyl)-1H-indole**

To a solution of (R)-3-(pyrrolidin-2-ylmethyl)-5-(2-ethylsulphonylethyl)-1H-indole (0.27 g, 0.8 mmol) in dimethylformamide (dried over 4A sieves) (5 mls), was added sodium carbonate (90 mgs) and ethyl iodide (0.07 mls, 0.88 mmol) at room temperature. The mixture was heated at 120°C under nitrogen for 16 hours. After cooling to room temperature the reaction mixture was partitioned between ethyl acetate and water. The separated organic phase was washed with water (3x), dried (Na_2SO_4) and evaporated in vacuo to give an oil. Purification by column chromatography on silica gel, eluting with CH_2Cl_2: EtOH: NH_4OH (90:10:0.5) gave the title compound as a gum (100 mgs). Anal. Calcd for $C_{19}H_{28}N_2O_2S \cdot 1/4\ CH_2Cl_2$. 1/2 H2O: C,61.04; H,7.85; N,7.40. Found: C,60.80; H,7.69; N,7.48. $[\alpha]^{25}$ + 60° (c=0.1, MeOH).

B. The following examples were prepared using the procedure described in a) above but with the appropriate alkyl halide in place of ethyl iodide. The alkyl halide could be iodide or bromide with the optional presence of sodium iodide. Solvents used were either dimethylformamide or dimethylacetamide.

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced



R^3	Isolated Form	Analysis % (Theoretical in brackets) C	H	N	$[\alpha]_D^{25}$ c=0.1 MeOH
iPr	Gum I/10 CH_2Cl_2 I/4 H_2O	64.18 (64.29	8.17 8.24	7.55 7.46)	+24°
* $CH_3CH(CH_2CH_3)$ (Isomer 1 - R.f. 0.40 SiO_2, CH_2Cl_2: MeOH: NH_3 (90:10:1)	Gum I/2 CH_2Cl_2 I/4 H_2O	60.68 (60.97	7.91 7.97	7.08 6.62)	-3°
* $CH_3CH(CH_2CH_3)$ (Isomer 2 - R.f. 0.38 SiO_2, CH_2Cl_2:MeOH:NH_3 (90:10:1))	Gum I/8 CH_2Cl_2	65.19 (65.53	8.13 8.40	7.45 7.24)	+26°
nPr	Gum I/20 CH_2Cl_2 3/5 H_2O	64.04 (63.77	8.19 8.36	7.52 7.42)	+62°
$(CH_3)_2CHCH_2$	Gum I/2 H_2O	65.32 (65.40	8.49 8.63	6.87 7.26)	+80°
* $CH_3(CH_3CH_2)CHCH_2$ (S-isomer)	Gum 2/3 H_2O	65.72 (65.63	8.82 8.85	7.10 6.96)	+65°

EXAMPLE 25

(R)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole

(R)-5-Bromo-3-(N-methylpyrrolidin-2-yl-methyl)-1H-indole (60 mg, 0.2 mmol) was dissolved in ethanol (1 mL) and hydrogenated over 10% palladium on carbon (45 mg) at 60 p.s.i. of hydrogen pressure at room temperature for 16 hours. The reaction mixture was evaporated to dryness, and the residue partitioned between ethyl acetate and 10% aqueous sodium carbonate. The organic phase was dried (Na_2SO_4), and evaporated in vacuo. The resulting residue was purified by column chromatography on silica gel (eluting

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

with 89:10:1 CH_2Cl_2:MeOH:NH_4OH) to give the title compound (28 mg). Anal. Calcd for $C_{14}H_{18}N_2 \cdot 1/8\ CH_2Cl_2$ C,75.42; H,8.18; N,12.46. Found: C,75.50; H,8.51; N,12.09. $[\alpha]^{25} = +60.2°$ (c=0.088, $CHCl_3$).

EXAMPLE 26

(R)-3-(N-Benzyloxycarbonylpyrrolidin-2-ylcarbonyl)-5-bromo-1H-indole

Two solutions containing the reactants were prepared separately as follows: To a stirred solution of N-benzyloxycarbonyl-D-proline (1.0 g) in anhydrous dichloromethane (2 ml) and N,N-dimethylformamide (1 drop) was added oxalyl chloride (0.5 mL), and the resulting solution was stirred at room temperature for 1.5 hours. The solution was evaporated under reduced pressure, and remaining solvent was removed under high vacuum to give the N-benzyloxycarbonyl-D-proline acid chloride. At the same time, a solution of ethyl magnesium bromide (1.4 mL of a 3M solution in ether) was added dropwise over 5 minutes to a stirred solution of 5-bromoindole (0.75 g) in dry ether (18 mL). The mixture was stirred at room temperature for 10 minutes, heated under reflux for 2 hours, then cooled to -30°C. A solution of the above N-benzyloxycarbonyl-D-proline acid chloride in dry ether (4 mL) was added dropwise with stirring, and stirring was continued for a further 1 hour. Ether (12.5 mL) and saturated aqueous sodium bicarbonate (6.5 mL) were added, and the temperature was allowed to rise to room temperature. Stirring was continued for a further 10 minutes and the mixture was filtered. The solid was washed well with ethyl acetate, and the combined filtrate and washings were washed with water, brine and dried ($MgSO_4$). Evaporation of the solvent gave an oil which was chromatographed on silica gel. Elution with ethyl acetate gave the title compound as a foam (0.82 g):
LRMS, m/z (relative intensity) 428 (M+ with ^{81}Br,5), 426 (M+ with ^{79}Br, 5), 224 (19), 222 (21), 204 (62), 160

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED


(68), 91 (100). Anal Calcd for $C_{21}H_{19}BrN_2O_3$: C,59.02; H,4.48; N,6.56. Found: C,58.85; H,4.51; N,6.38%.

EXAMPLE 27

(R)-5-Bromo-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole

A solution of (R)-3-(N-benzyloxycarbonyl-pyrrolidin-2-ylcarbonyl)-5-bromo-1H-indole (1.04 g) in dry tetrahydrofuran (20 mL) was added dropwise to a stirred suspension of lithium aluminium hydride (0.27 g) in dry tetrahydrofuran (15 mL) at room temperature under an atmosphere of dry nitrogen. The mixture was heated under reflux with stirring for 18 hours and then cooled. Additional lithium aluminium hydride (50 mg) was added and refluxing was continued for an additional 3 hours. The mixture was again cooled, lithium aluminium hydride (40 mg) was added, and refluxing was continued for a further 18 hours. The mixture was cooled and water (0.44 mL) was carefully added with stirring, followed by 20% aqueous sodium hydroxide (0.44 mL), followed by more water (1.33 mL). The mixture was diluted with ethyl acetate and filtered through Celite (trademark) filter aid. The filtrate was washed with water, brine and then dried (Na_2SO_4). Evaporation of the solvent gave an oil which was chromatographed on silica gel. Elution with dichloromethane/ethanol/concentrated aqueous ammonia (90:10:0.5) gave the title compound as a solid (0.51 g), m.p. 137-140°C (from dichloromethane/hexane); IR (KBr) 1620, 1595, 1570, 1480, 1450, 1435 cm^{-1}; ^1H NMR (DMSO-d_6) δ 11.05 (br s, 1H), 7.65 (br d, 1H), 7.31 (d, J=8.6 Hz, 1H), 7.21 (br d, 1H), 7.16 (dd, J=1.8 and 8.6 Hz, 1H), 3.03-2.94 (m, 2H), 2.47 (dd, J=9.2 and 14.0 Hz, 1H), 2.36-2.26 (m, 1H), 2.33 (s, 3H), 2.09 (dd, J=8.7 and 17.3 Hz, 1H), 1.73-1.38 (m,4H); ^{13}C NMR (DMSO-d_6) δ 134.8, 129.5, 124.7, 123.2, 120.7, 113.4, 112.1, 110.9, 66.1, 57.0, 40.5, 30.9, 29.1, 21.6; LRMS, m/z (relative intensity) 294 (M^+ with ^{81}Br, 1), 293 (2), 292 (M^+ with

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

^{79}Br, 1), 210 (14), 208 (15), 154 (8), 129 (42), 128 (19), 101 (26), 85 (57), 84 (100), 83 (30); $[\alpha]^{25}$ = + 62° (methanol, c = 0.10). Anal Calcd for $C_{14}H_{17}N_2Br$. 0.25 H_2O: C, 56.48; H, 5.93; N, 9.41. Found: C, 56.65; H, 5.69; N,9.23.

EXAMPLE 28

(R)-5-(2-Ethylsulphonylethenyl)-3-(N-methyl-pyrrolidin-2-ylmethyl)-1H-indole


A mixture of (R)-5-bromo-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole (0.25 g), ethyl vinyl sulphone (0.14 g), tri-o-tolylphosphine (0.075 g), palladium (II) acetate (0.013 g), triethylamine (0.25 mL) and acetonitrile (3 mL) was heated under reflux for 17 hours in an atmosphere of nitrogen. The mixture was evaporated and the residue was chromatographed on silica gel. Elution with dichloromethane/ethanol/ concentrated aqueous ammonia (90:8:1) gave the title compound as a foam (0.185 g): TLC (dichloromethane/-ethanol/ concentrated aqueous ammonia, 90:10:1): R_f = 0.5. Anal. Calcd for $C_{18}H_{24}N_2O_2S$. 0.2 CH_2Cl_2: C,62.55; H,7.04; N,8.02. Found: C,62.65; H,6.94; N,7.92.

EXAMPLE 29

(R)-5-(2-Ethylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)1H-indole

(R)-5-(2-Ethylsulphonylethenyl)-3-(N-methyl-pyrrolidin-2-ylmethyl)-1H-indole (157 mg) was dissolved in a mixture of ethanolic hydrogen chloride [prepared by addition of acetyl chloride (0.043 mL) to ethanol (10 mL)], N,N-dimethylformamide (7.5 mL) and water (0.1 mL) and the solution was shaken under a hydrogen atmosphere (15 psi) at room temperature for 18 hours in the presence of 10% palladium on carbon (150 mg). The mixture was filtered through Arbacel (trade mark) filter aid and the residue was washed well with ethanol. The combined filtrate and washings were evaporated under reduced pressure and the residual oil was partitioned between

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

ethyl acetate and 2M aqueous sodium carbonate solution. The organic layer was separated, washed three times with water followed by brine and then dried (Na_2SO_4). Evaporation of the solvent gave an oil which was chromatographed on silica gel. Elution with dichloromethane/methanol/ concentrated aqueous ammonia (90:10:1) gave the title compound as a gum (110 mg): TLC ($CH_2Cl_2/C_2H_5OH/NH_3$; 90:10:1): R_f = 0.3; $[\alpha]^{25}$ = +62° (methanol, c = 0.10). Anal. Calcd for $C_{18}H_{26}N_2O_2S$. 0.05 CH_2Cl_2: C,63.21; H,7.67; N,8.17. Found: C,63.55; H,7.61; N,8.41%.

EXAMPLE 30

(R)-5-(2-Ethylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole hemisuccinate


A solution of succinic acid (69 mg) in hot ethanol (3.5 mL) was added slowly with stirring to a solution of (R)-5-(2-ethylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole free base (390 mg) in ethanol (3.5 mL). The solution was evaporated and the residue was triturated first with ether and then with ethyl acetate to give the title compound as a solid (375 mg): mp 59-62°C: $[\alpha]^{25}$ = + 36° (methanol, c = 0.10). Anal. Calcd for $C_{18}H_{26}N_2O_2S$. 0.5 $C_4H_6O_4$. 0.25 $CH_3CO_2C_2H_5$. 0.5 H_2O: C,59.00; H,7.42; H,6.68. Found: C,59.17; H,7.37; N,6.73.

EXAMPLE 31

(R)-5-(2-Benzenesulphonylethenyl)-3-(N-methyl-pyrrolidin-2-ylmethyl)-1H-indole hydrobromide

A mixture of (R)-5-bromo-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole (0.25 g), phenylvinylsulphone (0.19 g), tri-o-tolylphosphine (0.075 g), palladium (II) acetate (0.0125 g), triethylamine (0.25 mL) and acetonitrile (2.5 mL) was heated under reflux for 42 hours in an atmosphere of nitrogen. The solvent was evaporated and the residue was chromatographed on silica gel. Elution with dichloromethane/methanol/

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED

concentrated aqueous ammonia (90:10:1) gave the title compound as a foam (0.24 g): Anal. Calcd for $C_{22}H_{24}N_2O_2S$. HBr. 1/3 CH_2Cl_2: C,54.77; H,5.29; N,5.72. Found: C,55.00; H,4.85; N,5.58.

EXAMPLE 32

(R)-5-(2-Benzenesulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole

A solution of (R)-5-(2-benzenesulphonylethenyl)-3-N-methylpyrrolidin-2-ylmethyl)-1H-indole hydrobromide (0.214 g) and 10% palladium on carbon (0.15 g) in a mixture of absolute ethanol (10 mL), N,N-dimethylformamide (1 mL) and water (2 drops) was shaken under a hydrogen atmosphere (15 psi) at room temperature for 18 hours. The mixture was filtered through Celite (trademark) filter aid and the residue was washed well with ethanol. The combined filtrate and washings were evaporated under reduced pressure and the residue was partitioned between ethyl acetate and 2M aqueous sodium carbonate solution. The organic layer was separated, washed three times with water, followed by brine and dried (Na_2SO_4). Evaporation of the solvent gave a gum which was chromatographed on silica gel. Elution with dichloromethane/methanol/concentrated aqueous ammonia (90:10:0.5) gave the title compound as a foam (0.096 g). Anal. Calcd for $C_{22}H_{26}N_2O_2S$. H_2O: C,65.97; H,7.05; N,7.00. Found: C,65.51; H,6.77; N,7.45.

EXAMPLE 33

(R)-5-(2-Benzenesulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole hemisuccinate

A solution of succinic acid (95 mg) in ethanol (5 mL) was added to a solution of (R)-5-(2-benzenesulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole free base (620 mg) in ethanol (5 mL). The solution was evaporated to give the title compound as a foam (680 mg): $[\alpha]^{25}$ = + 29° (methanol, c = 0.10). Anal.

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

Calcd for $C_{22}H_{26}N_2O_2S$. 0.5 $C_4H_6O_4$. 0.33 C_2H_5OH. 0.5 H_2O; C,63.59; H,6.92; N,6.01. Found: C,63.52; H,6.91; N,6.12.

EXAMPLE 34

(R)-5-[2-(4-Methylphenylsulphonyl)ethenyl]-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole

A mixture of (R)-5-bromo-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole (0.40 g), 4-methylphenylvinylsulphone (0.273 g), tri-o-tolylphosphine (0.085 g), palladium (II) acetate (0.031 g), triethylamine (0.42 g), and acetonitrile (20 mL) was heated under reflux for 16 hours in an atmosphere of nitrogen. The mixture was cooled and partitioned between ethyl acetate and 10% aqueous sodium bicarbonate solution. The organic layer was washed with brine, dried (Na_2SO_4) and evaporated. The residual orange oil was chromatographed on silica gel. Elution was commenced with dichloromethane/methanol (90:10), followed by dichloromethane/methanol/concentrated aqueous ammonia (90:10:0.25), gradually increasing the concentration of concentrated aqueous ammonia to 1%. The later product-containing fractions were evaporated to give the title compound as a foam (226 mg): $[\alpha]^{25}$ = + 71° (methanol, c = 0.10). Anal. Calcd for $C_{23}H_{26}N_2O_2S$. 0.15 CH_2Cl_2: C,68.27; H,6.51; N,6.88. Found: C,68.26; H,6.54; N,6.99.

EXAMPLE 35

(R)-5-[2-(4-Methylphenylsulphonyl)ethyl-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole

A solution of (R)-5-[2-(4-methylphenylsulphonyl)ethenyl]-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole (0.18 g) and 10% palladium on carbon (0.20 g) in ethanolic hydrogen chloride [prepared from absolute ethanol (25 mL) and acetyl chloride (35 µL)] was shaken under a hydrogen atmosphere (15 psi) at room temperature for 16 hours. The reaction mixture was filtered through Celite (trademark) filter aid and the residue was washed

This Sheet Has Been Replaced



THIS SHEET HAS BEEN REPLACED


well with ethanol. The combined filtrate and washings were evaporated under reduced pressure and the residue was partitioned between ethyl acetate and 2M aqueous sodium carbonate solution. The organic layer was saturated, washed three times with water, followed by brine and dried (Na_2SO_4). Evaporation of the solvent gave a gum which was chromatographed on silica gel. Elution with dichloromethane/methanol/concentrated aqueous ammonia (90:10:0.25) gave the title compound as a foam (108 mg): $[\alpha]^{25} = +30°$ (methanol, c = 0.10). Anal. Calcd for $C_{23}H_{28}N_2O_2S$. 0.05 CH_2Cl_2. 0.5 H_2O: C,67.55; H,7.15; N,6.84. Found: C,67.51; H,7.04; N,6.98.

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

CLAIMS

1. A compound of the formula



I

CLMS

wherein n is 0, 1, or 2; R_1 is hydrogen; R_2 is selected from hydrogen, halogen, cyano, OR_4, $-(CH_2)_m-(C=O)NR_5R_6$, $-(CH_2)_m-SO_2NR_5R_6$, $-(CH_2)_m-NR_7(C=O)R_8$, $-(CH_2)_m-NR_7SO_2R_8$, $-(CH_2)_m-S(O)_xR_8$, $-(CH_2)_m-NR_7(C=O)NR_5R_6$, $-(CH_2)_m-NR_7(C=O)OR_9$, and $-CH=CH(CH_2)_yR_{10}$; R_3 is selected from hydrogen and C_1 to C_6 linear or branched alkyl; R_4 is selected from hydrogen, C_1 to C_6 alkyl, and aryl; R_5 and R_6 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl or R_5 and R_6 taken together to form a 4, 5, or 6 membered ring; R_7 and R_8 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_9 is selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_{10} is selected from $-(C=O)NR_5R_6$ and $-SO_2NR_5R_6$, wherein R_5 and R_6 are defined as above, and $-NR_7(C=O)R_8$, $-NR_7SO_2R_8$, $-NR_7(C=O)NR_5R_6$, $-S(O)_xR_8$ and $-NR_7(C=O)OR_9$, wherein R_7, R_8, and R_9 are as defined above; y is 0, 1, or 2; x is 1 or 2; m is 0, 1, 2, or 3; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C_1 to C_4 alkoxy and the pharmaceutically acceptable salts thereof.

2. The R enantiomer of a compound according to claim 1.

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

* C L M S *

3. A compound according to claim 1 wherein R_1 is hydrogen; R_2 is $-(CH_2)_m-SO_2NHR_5$, $-(CH_2)_m-NHSO_2R_8$, $-(CH_2)_m-SO_2R_8$, $-(CH_2)_m-(C=O)NHR_5$, or $-(CH_2)_m-NH(C=O)R_8$; R_3 is hydrogen or methyl; and m, R_5 and R_8 are as defined in claim 1.

4. A compound according to claim 1, said compound being selected from:

(R)-5-methoxy-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-bromo-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-ethylsulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylmethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-carboxamido-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylsulfonylethyl)-3-(N-methylpyrrolidin-2-yl-methyl)-1H-indole;

(R)-5-(2-methylsulfonamidoethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-aminosulphonylethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-aminosulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-N,N-dimethylaminosulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-phenylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole hemisuccinate;

(R)-5-(2-ethylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole hemisuccinate;

This Sheet Has Been Replaced

* C L M S *

THIS SHEET HAS BEEN REPLACED

(R)-5-(2-phenylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(3-benzenecarbonylaminoprop-1-enyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-(4-methylphenylsulphonyl)ethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(3-methylsulphonylaminoprop-1-enyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-ethylsulphonylethyl)-3-(N-2-propylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-ethylsulphonylethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-(4-methylphenylsulphonyl)ethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole; and

(R)-5-(2-methylsulfonamidomethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole.

5. A pharmaceutical composition for treating a condition selected from hypertension, depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain, and chronic paroxysmal hemicrania and headache associated with vascular disorders comprising an amount of a compound according to claim 1 effective in treating such condition and a pharmaceutically acceptable carrier.

6. A pharmaceutical composition for treating disorders arising from deficient serotonergic neurotransmission comprising an amount of a compound according to claim 1 effective in treating such a disorder and a pharmaceutically acceptable carrier.

7. A method for treating a condition selected from hypertension, depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain and chronic paroxysmal hemicrania and headache associated with vascular disorders comprising administering to a mammal requiring such treatment an amount of a compound

This Sheet Has Been Replaced


* C L M S *

THIS SHEET HAS BEEN REPLACED

according to claim 1 effective in treating such condition.

8. A method for treating disorders arising from deficient serotonergic neurotransmission comprising administering to a mammal requiring such treatment an amount of a compound according to claim 1 effective in treating such a disorder.

9. A compound of the formula



V

wherein W is $-CO_2R_{11}$ or R_3; Q is CH_2 or C=O; n is 0, 1 or 2; R_1 is hydrogen; R_2 is selected from hydrogen, halogen, cyano, OR_4, $-(CH_2)_m-(C=O)NR_5R_6$, $-(CH_2)_m-SO_2NR_5R_6$, $-(CH_2)_m-NR_7(C=O)R_8$, $-(CH_2)_m-NR_7SO_2R_8$, $-(CH_2)_m-S(O)_xR_8$, $-(CH_2)_m-NR_7(C=O)NR_5R_6$, $-(CH_2)_m-NR_7(C=O)OR_9$, and $-CH=CH(CH_2)_yR_{10}$; x is 1 or 2; m is 0, 1, 2, or 3; R_3 is selected from hydrogen and C_1 to C_6 linear or branched alkyl; R_4 is selected from hydrogen, C_1 to C_6 alkyl, and aryl, R_5 and R_6 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl or R_5 and R_6 taken together to form a 4, 5, or 6 membered ring; R_7 and R_8 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_9 is selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_{10} is selected from $-(C=O)NR_5R_6$ and $-SO_2NR_5R_6$, wherein R_5 and R_6 are defined as above, and $-NR_7(C=O)R_8$, $-NR_7SO_2R_8$, $-NR_7(C=O)NR_5R_6$, $-S(O)_xR_8$ and $-NR_7(C=O)OR_9$, wherein R_7, R_8, R_9 and x are defined as above; y is 0, 1, or 2; R_{11} is selected from C_1 to C_6 alkyl, benzyl and aryl; and the above aryl groups and the aryl moieties of the above

This Sheet Has Been Replaced


* C L M S *

THIS SHEET HAS BEEN REPLACED

alkyl-aryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C_1 to C_4 alkoxy.

10. The R enantiomer of a compound according to claim 9.

11. A compound according to claim 9, said compound being a compound of the formula



II

wherein n, R_1, R_2 and R_{11} are as defined in claim 9.

12. The R enantiomer of a compound according to claim 11.


13. A compound according to claim 11 wherein R_1 is hydrogen; R_2 is $-(CH_2)_m-SO_2NHR_5$, $-(CH_2)_m-NHSO_2R_6$, $-(CH_2)_m-SO_2R_6$, $-(CH_2)_m-(C=O)NHR_5$ or $-(CH_2)_m-NH(C=O)R_6$; m is 0, 1, 2, or 3; R_5 is hydrogen, C_1 to C_6 alkyl, aryl, or C_1 to C_3 alkyl-aryl; R_{11} is selected from C_1 to C_6 alkyl, benzyl and aryl; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C_1 to C_4 alkoxy.

14. A compound according to claim 9, said compound being a compound of the formula

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

* C L M S *




III

wherein n, R_1, R_3 and R_{10} are as defined in claim 9.

15. The R enantiomer of a compound according to claim 14.

16. A compound according to claim 14 wherein R_1 is hydrogen; R_3 is hydrogen or methyl; and R_{10} is $-SO_2NHR_5$, $NHSO_2R_8$, $-SO_2R_8$, $-(C=O)NHR_5$ or $-NH(C=O)R_8$, wherein R_5 and R_8 are as defined in claim 9.

17. A process for preparing a compound of the formula



I


wherein n is 0, 1, or 2; R_1 is hydrogen; R_2 is selected from hydrogen, halogen, cyano, OR_4, $-(CH_2)_m-(C=O)NR_5R_6$, $-(CH_2)_m-SO_2NR_5R_6$, $-(CH_2)_m-NR_7(C=O)R_8$, $-(CH_2)_m-NR_7SO_2R_8$, $-(CH_2)_m-S(O)_xR_8$, $-(CH_2)_m-NR_7(C=O)NR_5R_6$, $-(CH_2)_m-NR_7(C=O)OR_9$, and $-CH=CH(CH_2)_yR_{10}$; R_3 is selected from hydrogen and C_1 to C_6 linear or branched alkyl; R_4 is selected from hydrogen, C_1 to C_6 alkyl, and aryl; R_5 and R_6 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl or R_5 and R_6 taken together to form a 4, 5, or 6 membered ring; R_7 and R_8 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_9

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

is selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_{10} is selected from $-(C=O)NR_5R_6$ and $-SO_2NR_5R_6$, wherein R_5 and R_6 are defined as above, and $-NR_7(C=O)R_8$, $-NR_7SO_2R_8$, $-NR_7(C=O)NR_5R_6$, $-S(O)_xR_8$ and $-NR_7(C=O)OR_9$, wherein R_7, R_8, and R_9 are as defined above; y is 0, 1, or 2; x is 1 or 2; m is 0, 1, 2, or 3; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C_1 to C_4 alkoxy, comprising

(a) reducing a compound of the formula



V

where Q is CH_2 or C=O, W is $-CO_2R_{11}$ or R_3, and R_{11} is C_1 to C_6 alkyl, benzyl, or aryl, and R_1, R_2, and R_{11} are as defined above;

(b) where R_2 is $-CH=CH(CH_2)_yR_{10}$ and R_{10} is as defined above by reacting a compound of formula I where R_3 is H or C_1 to C_6 linear or branched alkyl and R_2 is halogen and R_1 is defined as above with a compound of the formula $CH_2=CH(CH_2)_yR_{10}$ where R_{10} is as defined above using transition metal catalysis; or

(c) reacting the compound of formula I where R_3 is hydrogen and R_1 and R_2 are as defined above with a compound of the formula R_3-Z where R_3 is C_1 to C_6 linear or branched alkyl and Z is halogen and base; and,


if desired, converting a compound of formula I to a the pharmaceutically acceptable salts thereof.

THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced

is selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_{10} is selected from $-(C=O)NR_5R_6$ and $-SO_2NR_5R_6$, wherein R_5 and R_6 are defined as above, and $-NR_7(C=O)R_8$, $-NR_7SO_2R_8$, $-NR_7(C=O)NR_5R_6$, $-S(O)_xR_8$ and $-NR_7(C=O)OR_9$, wherein R_7, R_8, and R_9 are as defined above; y is 0, 1, or 2; x is 1 or 2; m is 0, 1, 2, or 3; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C_1 to C_4 alkoxy, comprising

(a) reducing a compound of the formula



V

where Q is CH_2 or C=O, W is $-CO_2R_{11}$ or R_3, and R_{11} is C_1 to C_6 alkyl, benzyl, or aryl, and R_1, R_2, and R_{11} are as defined above;

(b) where R_2 is $-CH=CH(CH_2)_yR_{10}$ and R_{10} is as defined above by reacting a compound of formula I where R_3 is H or C_1 to C_6 linear or branched alkyl and R_2 is halogen and R_1 is defined as above with a compound of the formula $CH_2=CH(CH_2)_yR_{10}$ where R_{10} is as defined above using transition metal catalysis; or

(c) reacting the compound of formula I where R_3 is hydrogen and R_1 and R_2 are as defined above with a compound of the formula R_3-Z where R_3 is C_1 to C_6 linear or branched alkyl and Z is halogen and base; and,

if desired, converting a compound of formula I to a the pharmaceutically acceptable salts thereof.

This Sheet Has Been Replaced

THIS SHEET HAS BEEN REPLACED

* C L M S *

18. The process according to claim 17 wherein the compound of formula I is an R enantiomer.

19. A process according to claim 17 wherein for the compound of formula I, R_1 is hydrogen; R_2 is $-(CH_2)_m-SO_2NHR_5$, $-(CH_2)_m-NHSO_2R_8$, $-(CH_2)_m-SO_2R_8$, $-(CH_2)_m-(C=O)NHR_5$, or $-(CH_2)_m-NH(C=O)R_8$; R_3 is hydrogen or methyl; and m, R_5 and R_8 are as defined in claim 17.

20. A process according to claim 17, said compound of formula I being selected from:

(R)-5-methoxy-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-bromo-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-ethylsulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylaminosulfonylmethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-carboxamido-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-methylsulfonylethyl)-3-(N-methylpyrrolidin-2-yl-methyl)-1H-indole;

(R)-5-(2-methylsulfonamidoethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-aminosulphonylethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-aminosulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-N,N-dimethylaminosulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-phenylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole hemisuccinate;

This Sheet Has Been Replaced

* C L M S *

THIS SHEET HAS BEEN REPLACED

(R)-5-(2-ethylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole hemisuccinate;

(R)-5-(2-phenylsulphonylethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(3-benzenecarbonylaminoprop-1-enyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-(4-methylphenylsulphonyl)ethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(3-methylsulphonylaminoprop-1-enyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;


(R)-5-(2-ethylsulphonylethyl)-3-(N-2-propylpyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-ethylsulphonylethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;

(R)-5-(2-(4-methylphenylsulphonyl)ethenyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole; and

(R)-5-(2-methylsulfonamidomethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole.

21. A process for preparing a compound of the formula



V

wherein W is $-CO_2R_{11}$ or R_3; Q is CH_2 or C=O; n is 0, 1 or 2; R_1 is hydrogen; R_2 is selected from hydrogen, halogen, cyano, OR_4, $-(CH_2)_m-(C=O)NR_5R_6$, $-(CH_2)_m-SO_2NR_5R_6$, $-(CH_2)_m-NR_7(C=O)R_8$, $-(CH_2)_m-NR_7SO_2R_8$, $-(CH_2)_m-S(O)_xR_8$, $-(CH_2)_m-NR_7(C=O)NR_5R_6$, $-(CH_2)_m-NR_7(C=O)OR_9$, and $-CH=CH(CH_2)_yR_{10}$; x is 1 or 2; m is 0, 1, 2, or 3; R_3 is selected from hydrogen and C_1 to C_6 linear or branched alkyl; R_4 is selected from hydrogen, C_1 to C_6 alkyl, and aryl, R_5 and R_6 are


This Sheet Has Been Replaced

* C L M S *

THIS SHEET HAS BEEN REPLACED

independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl or R_5 and R_6 taken together to form a 4, 5, or 6 membered ring; R_7 and R_8 are independently selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_9 is selected from hydrogen, C_1 to C_6 alkyl, aryl, and C_1 to C_3 alkyl-aryl; R_{10} is selected from $-(C=O)NR_5R_6$ and $-SO_2NR_5R_6$, wherein R_5 and R_6 are defined as above, and $-NR_7(C=O)R_8$, $-NR_7SO_2R_8$, $-NR_7(C=O)NR_5R_6$, $-S(O)_xR_8$ and $-NR_7(C=O)OR_9$, wherein R_7, R_8, R_9 and x are defined as above; y is 0, 1 or 2; R_{11} is selected from C_1 to C_6 alkyl, benzyl and aryl; and the above aryl groups and the aryl moieties of the above alkyl-aryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C_1 to C_4 alkoxy, comprising

(a) where W is $-CO_2R_{11}$ and R_1, R_2, R_{11}, and Q are as defined above, by reacting a compound of the formula



IV

wherein R_1 and R_2 are as defined above with an acid chloride of the formula N-CO_2R_{11}-proline with base; or

(b) where W is R_3, Q is CH_2, and R_2 is $-CH=CH(CH_2)_yR_{10}$ and R_1, R_3, and R_{10} are as defined above, by reacting the compound of formula V where R_2 is halogen, W is R_3, Q is CH_2 and R_1 and R_3 are as defined above with a compound of the formula $CH_2=CH(CH_2)_yR_{10}$ where R_{10} is as defined above using transition metal catalysis.


22. The process according to claim 21 wherein the compound of formula V is an R enantiomer.

This Sheet Has Been Replaced

* C L M S *

THIS SHEET HAS BEEN REPLACED

23. A process according to claim 21, wherein said compound of formula V is a compound of the formula




II

wherein n, R_1, R_2 and R_{11} are as defined in claim 21.

24. The process according to claim 23 wherein the compound of formula II is an R enantiomer.

25. A process according to claim 23 wherein for the compound of formula II, R_1 is hydrogen; R_2 is $-(CH_2)_m-SO_2NHR_5$, $-(CH_2)_m-NHSO_2R_8$, $-(CH_2)_m-SO_2R_8$, $-(CH_2)_m-(C=O)NHR_5$ or $-(CH_2)_m-NH(C=O)R_8$; m is 0, 1, 2, or 3; R_5 is hydrogen, C_1 to C_6 alkyl, aryl, or C_1 to C_3 alkyl-aryl; R_{11} is selected from C_1 to C_6 alkyl, benzyl and aryl; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C_1 to C_4 alkoxy.

26. A process according to claim 21, wherein said compound of formula V is a compound of the formula



III

wherein n, R_1, R_3 and R_{10} are as defined in claim 21.

This Sheet Has Been Replaced

* C L M S *

THIS SHEET HAS BEEN REPLACED

27. The process according to claim 26 wherein the compound of formula III is an R enantiomer.

28. A process according to claim 26 wherein for the compound of formula III R_1 is hydrogen; R_3 is hydrogen or methyl; and R_{10} is $-SO_2NHR_5$, $NHSO_2R_8$, $-SO_2R_8$, $-(C=O)NHR_5$ or $-NH(C=O)R_8$, wherein R_5 and R_8 are as defined in claim 21.