CLAIMS

1. Use of about 4 g of eicosapentaenoic acid (EPA), or an atrial fibrillation flutter and/or atrial flutter therapeutic agent, in the manufacture of a medicament for treating a patient, wherein~~A method for treating a patient comprising~~:

~~determining or having determined whether~~ the patient has been determined to have~~has~~ a triglyceride level of at least 150 mg/dl and a history of atrial fibrillation and/or atrial flutter by:

~~obtaining or~~ having obtained information which indicates that the patient has the triglyceride level of at least 150 mg/dl and has previously had symptoms of atrial fibrillation and/or atrial flutter and/or was previously diagnosed with atrial fibrillation and/or atrial flutter; and/or

~~monitoring or~~ having monitored the subject for the triglyceride level of at least 150 mg/dl and for symptoms of atrial fibrillation and/or atrial flutter; and

(a) if the patient ~~has~~ had previously had symptoms of atrial fibrillation and/or atrial flutter, ~~was~~ had previously been diagnosed with atrial fibrillation and/or atrial flutter, and/or had symptoms of atrial fibrillation and/or atrial flutter when monitored, then

(i) having determined~~determining~~ whether a risk that the patient will experience symptoms of atrial fibrillation and/or atrial flutter if the patient ~~were~~ had been administered about 4 g of eicosapentaenoic acid (EPA) per day is greater than reducing a risk of one or more cardiovascular events in the patient; or

(ii) having determined ~~determining~~ whether reducing a risk of one or more cardiovascular events if the patient ~~were~~ had been administered about 4 g of EPA per day is greater than a risk that the patient will experience symptoms of atrial fibrillation and/or atrial flutter; and

(b) if the risk is greater that the patient in (i) will experience symptoms of atrial fibrillation and/or atrial flutter, ~~administering~~ an atrial fibrillation flutter and/or atrial flutter

therapeutic agent is to be administered to the patient instead of about 4 g of EPA per day; and

(c) if the risk is greater that the patient in (ii) will have a reduced risk of one or more cardiovascular events, ~~administering~~ about 4 g of EPA per day is to be administered to the patient.

2. The ~~method~~ use of claim 1, wherein the subject in (c) is likely to experience both (1) increased frequency of atrial fibrillation and/or atrial flutter symptoms, and (2) decreased frequency of cardiovascular events if ~~administered~~ the subject is to be administered about 4 g of EPA per day.

3. The ~~method~~ use of claim 1, wherein the subject ~~is being~~had been treated with a statin and regardless if the subject is in (a) or (b), the subject continues to be ~~continuing to~~ treated ~~the subject~~ with the statin.

4. Use of about 4 g of EPA, or atrial fibrillation and/or atrial flutter therapeutic agent, in the manufacture of a medicament ~~A method~~ for treating a subject ~~comprising~~wherein:

~~determining or having determined~~it had been determined whether the subject has a triglyceride level of at least 150 mg/dl and a history of atrial fibrillation and/or atrial flutter by:

(i) ~~obtaining or~~ having obtained information which indicates that the subject has the triglyceride level of at least 150 mg/dl and has previously had symptoms of atrial fibrillation and/or atrial flutter and/or was previously diagnosed with atrial fibrillation and/or atrial flutter; and/or

(ii) ~~monitoring or~~ having monitored the subject for the triglyceride level of at least 150 mg/dl and symptoms of atrial fibrillation and/or atrial flutter; and

(a) if the subject ~~has~~ had previously had symptoms of atrial fibrillation and/or atrial flutter, ~~was~~ had previously been diagnosed with atrial fibrillation and/or atrial flutter, and/or

symptoms of atrial fibrillation and/or atrial flutter ~~were~~ had been determined during the monitoring, then ~~administering~~ atrial fibrillation and/or atrial flutter therapeutic agent is to be administered to the subject, and

(b) if the subject ~~has~~ had not previously had symptoms of atrial fibrillation and/or atrial flutter, ~~was~~ had not been previously diagnosed with atrial fibrillation and/or atrial flutter, and/or symptoms of atrial fibrillation and/or atrial flutter ~~were~~ had not been determined during the monitoring, then ~~administering~~ about 4 g of eicosapentaenoic acid (EPA) per day is to be administered the subject to reduce a risk for a cardiovascular event,

wherein a risk of atrial fibrillation and/or atrial flutter for the subject in (a) is lower following administration of an atrial fibrillation flutter and/or atrial flutter therapeutic agent than it would be if the subject in (a) were to be administered about 4 g of EPA per day.

5. The ~~method~~ use of claim 4, wherein the subject in (a) is likely to experience both (1) increased frequency of atrial fibrillation and/or atrial flutter symptoms, and (2) decreased frequency of cardiovascular events if ~~administered~~ about 4 g of EPA per day is to be administered.

6. The ~~method~~ use of claim 4, wherein the subject ~~is being~~had been treated with a statin and regardless if the subject is in (a) or (b), ~~continuing~~ the subject continues to be treated ~~the subject~~ with the statin.

7. The ~~method~~ use of any one of claims 1 to 6, wherein the atrial fibrillation flutter and/or atrial flutter therapeutic agent is one or more of warfarin, beta blocker, calcium channel blocker.

8. The ~~method~~ use of claim 7, wherein the warfarin is to be administered at a dose of about 25 mg per day.

9. The ~~method~~ use of any one of claims 1 to 8, wherein reducing the risk for the cardiovascular event for the subject further comprises determining that one or more cardiovascular benefits are expected to occur.

10. The ~~method~~ use of any one of claims 1 to 9, wherein the cardiovascular event does not include atrial fibrillation and/or atrial flutter.

11. Use of about 4 g eicosapentaenoic acid (EPA) or a statin in the manufacture of a medicament for treating a subject~~A method of treatment comprising~~:

~~identifying~~ wherein a subject has been identified whose measured triglyceride level is above 150 mg/dl and who has previously experienced symptoms of atrial fibrillation or atrial flutter;

~~determining that~~wherein individuals who had been identified ~~have~~ as having triglyceride levels above 150 mg/dl and who ~~have~~ had previously experienced atrial fibrillation or atrial flutter episodes are, upon treatment with 4 g eicosapentaenoic acid (EPA) per day, likely to experience both (1) increased frequency of atrial fibrillation or atrial flutter episodes, and (2) decreased frequency of cardiovascular events other than atrial fibrillation or atrial flutter; and

are to be treated ~~selecting a treatment~~ according to either (a) or (b) below:

(a) wherein it had been determined~~determining~~ that decreasing the subject's frequency of cardiovascular events other than atrial fibrillation or atrial flutter is worth a risk of increased frequency of atrial fibrillation or atrial flutter in the subject, and so ~~treating~~ the subject is to be treated with 4 g EPA per day; or

(b) ~~determining~~ wherein it had been determined that decreasing the subject's frequency of cardiovascular events other than atrial fibrillation or atrial flutter is not worth a risk of increased frequency of atrial fibrillation or atrial flutter in the subject, and so excluding the

subject from treatment with EPA, and instead ~~treating~~ the subject is to be treated with a statin.

12. Use of 4 g eicosapentaenoic acid (EPA) and/or a statin in the manufacture of a medicament for treating a subject~~A method of treatment comprising~~:

~~identifying~~ wherein a subject has been identified whose measured triglyceride level is above 150 mg/dl, who is being treated with a statin, and who has previously experienced symptoms of atrial fibrillation or atrial flutter;

~~determining that~~wherein individuals who had been identified ~~have~~ as having triglyceride levels above 150 mg/dl, who had been~~are being~~ treated with a statin, and who ~~have~~ had previously experienced atrial fibrillation or atrial flutter episodes are, upon treatment with 4 g eicosapentaenoic acid (EPA) per day, likely to experience both (1) increased frequency of atrial fibrillation or atrial flutter episodes, and (2) decreased frequency of cardiovascular events other than atrial fibrillation or atrial flutter; and

are to be treated ~~selecting a treatment~~ according to either (a) or (b) below:

(a) wherein it had been determined~~determining~~ that decreasing the subject's frequency of cardiovascular events other than atrial fibrillation or atrial flutter is worth a risk of increased frequency of atrial fibrillation or atrial flutter in the subject, and so ~~treating~~ the subject is to be treated with 4 g EPA per day and a statin; or

(b) ~~determining~~ wherein it had been determined that decreasing a subject's frequency of cardiovascular events other than atrial fibrillation or atrial flutter is not worth a risk of increased frequency of atrial fibrillation or atrial flutter in the subject, and so excluding the subject from treatment with EPA, and instead ~~continuing to treat~~ the subject is to continue being treated with a statin.

13. Use of 4 g eicosapentaenoic acid (EPA) or a statin in the manufacture of a medicament for treating a subject~~A method of treatment comprising~~:

Wherein ~~identifying~~ a subject has been identified whose measured triglyceride level is above 150 mg/dl who is being treated with a statin, and who has a previous history of atrial fibrillation or atrial flutter;

~~determining that~~wherein individuals who had been determined to have triglyceride levels above 150 mg/dl, who are being treated with a statin, and who have a previous history of atrial fibrillation or atrial flutter are, upon treatment with 4 g eicosapentaenoic acid (EPA) per day, are likely to experience both (1) increased frequency of atrial fibrillation or atrial flutter episodes, and (2) cardiovascular benefits expected to result from the EPA treatment; and

~~selecting a treatment~~are to be treated according to either (a) or (b) below:

(a) ~~determining~~ wherein it had been determined that the expected cardiovascular benefits are worth a risk of increased frequency of atrial fibrillation or atrial flutter in the subject, and so ~~treating~~ the subject is to be treated with 4 g EPA per day; or

(b) wherein it had been determined ~~determining~~ that the expected cardiovascular benefits are not worth a risk of increased frequency of atrial fibrillation or atrial flutter in the subject, and so excluding the subject from treatment with EPA, and instead ~~treating~~ the subject is to be treated with a statin.

14. Use of at least about 4 g of eicosapentaenoic acid (EPA) in the manufacture of a medicament for~~A method of~~ reducing a risk of a cardiovascular event in a subject having triglyceride levels of at least about 150 mg/dl, ~~the method comprising~~wherein ~~administering to~~ the subject is to be administered at least about 4 g of eicosapentaenoic acid (EPA) per day for a period of time effective to reduce the cardiovascular event, wherein the subject develops atrial fibrillation and/or atrial flutter during the period of time effective to reduce the risk of the cardiovascular event.

15. The ~~method~~ use of claim 14, wherein the subject does not have a history of atrial fibrillation and/or flutter prior to ~~administration of~~ the 4 g of EPA having been administered.

16. Use of at least about 4 g of eicosapentaenoic acid (EPA) in the manufacture of a medicament for~~A method of~~ reducing a risk of a cardiovascular event in a subject having triglyceride levels of at least about 150 mg/dl and less than 65 years of age, ~~the method comprising administering to~~ wherein the subject is to be administered at least about 4 g of eicosapentaenoic acid (EPA) per day for a period of time effective to reduce the cardiovascular event, wherein the subject does not exhibit an increase in symptoms atrial fibrillation and/or atrial flutter.

17. The ~~method~~ use of any one of claims 1 to 16, wherein the subject has established cardiovascular disease.

18. The ~~method~~ use of claim 17, wherein the established cardiovascular disease is determined by the presence of any one of: documented coronary artery disease, documented cerebrovascular disease, documented carotid disease, documented peripheral arterial disease, or combinations thereof.

19. The ~~method~~ use of any one of claims 1 to 16, wherein the subject does not have an established cardiovascular disease but at least two risk factors for cardiovascular disease.

20. The ~~method~~ use of any one of claims 1 to 19, wherein the EPA is to be administered to the subject in 1 to 4 dosage units per day.

21. The ~~method~~ use of any one of claims 1 to 20, wherein the subject has one or more of: a baseline non-HDL-C value of about 200 mg/dL to about 300 mg/dL; a baseline total cholesterol value of about 250 mg/dL to about 300 mg/dL; a baseline VLDL-C value of about 140 mg/dL to about 200 mg/dL; a baseline HDL-C value of about 10 to about 30 mg/dL; and/or a baseline LDL-C value of about 40 to about 100 mg/dL.

22. The ~~method~~ use of any one of claims 1 to 21, wherein the subject is to be administered the EPA for at least about 4 months, at least about 1 year, at least about 2 years, at least about 3 years, at least about 4 years, or at least about 5 years.