Mutated 5-enolpyruvylshikimate-3-phosphate synthase, gene coding for this protein and transformed plants containing this gene

The present invention relates to a new 5-enolpyruvylshikimate-3-phosphate synthase (or EPSPS) which displays increased tolerance with respect to herbicides which are competitive inhibitors with respect to phosphonoenolpyruvate (PEP) of EPSPS activity. This more tolerant EPSP synthase possesses at least one "threonine by isoleucine" substitution. The invention also relates to a gene coding for such a protein, to plant cells transformed by chimeric gene constructions containing this gene, to the plants regenerated from these cells and also to the plants originating from crossing using these transformed plants.

Glyphosate, sulfosate or fosametine are broad-spectrum systemic herbicides of the phosphonomethylglycine family. They act essentially as competitive inhibitors of 5-enolpyruvylshikimate-3-phosphate synthase (EC 2.5.1.19) or EPSPS with respect to the PEP (phosphoenolpyruvate). After their application to the plant, they are translocated in the plant where they accumulate in the rapidly growing parts, in particular the cauline and root apices, causing damage to the point of destruction of sensitive plants.

Plastid EPSPS, the main target of these products, is an enzyme of the pathway of biosynthesis of aromatic amino acids, which is encoded by one or more nuclear genes and synthesized in the form of a cytoplasmic precursor, then imported into the plastids where it accumulates in its mature form.

The tolerance of plants to glyphosate and to products of the family is obtained by stable introduction into their genome of an EPSPS gene, of plant or bacterial origin, which is mutated or otherwise in respect of the characteristics of inhibition by glyphosate of the product of this gene. In view of the mode of action of glyphosate and the degree of tolerance to glyphosate of the product of the genes which are used, it is advantageous to be able to express the product of the translation of this gene so as enable it to be accumulated in substantial amounts in the plastids.

It is known, for example from US Patent 4,535,060, to confer on a plant a tolerance to a herbicide of the above type, especially N-phosphono-methylglycine or glyphosate, by introducing into the genome of plants a gene coding for an EPSPS carrying at least one mutation that makes this enzyme more resistant to its competitive inhibitor (glyphosate) after localization of the enzyme in the plastid compartment. These techniques, however, need to be improved in order to obtain greater reliability in the

use of these plants under agricultural conditions.

In the present description, "plant" is understood to mean any differentiated multicellular organism capable of photosynthesis, and "plant cell" is understood to mean any cell originating from a plant and capable of constituting undifferentiated tissues such as calluses or differentiated tissues such as embryos or plant parts or seeds.

The subject of the present invention is the production of transformed plants having increased tolerance to herbicides of the phosphonomethylglycine family, by regeneration of cells transformed by means of new chimeric genes containing a gene for tolerance to these herbicides.

The subject of the invention is also a chimeric gene for conferring on plants increased tolerance with respect to a herbicide having EPSPS as its target, comprising, in the direction of transcription: a promoter region, optionally a transit peptide region, a sequence of a gene coding for a glyphosate tolerance enzyme and an untranslated polyadenylation signal region at the 3′ end, characterized in that the glyphosate tolerance gene contains, relative to the gene from which it is derived, a "threonine 102 by isoleucine" substitution in the "aroA"(EPSPS) region. Preferably, it comprises, in addition, in the same region, a "proline 106 by serine" substitution. These substitutions can be

introduced or be present in an EPSPS sequence of any origin, in particular of plant, bacterial, algal or fungal origin.

The transit peptides which can be used in the transit peptide region can be, known per se, of plant origin, for example originating from maize, sunflower, pea, tobacco or the like. The first and the second transit peptide can be identical, similar or different. They can, in addition, each comprise one or more transit peptide units according to European Patent Application EP 0 508 909. It is the role of this characteristic region to permit the release of a mature and native protein, and especially the above mutated EPSPS, with maximum efficacy in the plasmid compartment.

The promoter region of the chimeric gene according to the invention may be advantageously composed of at least one gene promoter or promoter fragment which is expressed naturally in plants (tubulin, introns actin, histone).

The untranslated transcription termination signal region at the 3' end of the chimeric gene may be of any origin, for example of bacterial origin, such as that of the nopaline synthase gene, or of plant origin, such as that of the *Arabidopsis thaliana* histone H4A748 gene according to the European Patent Application (European Application 633 317).

The chimeric gene according to the invention

can comprise, in addition to the essential portions above, at least one untranslated intermediate (linker) region, which can be located between the different transcribed regions described above. This intermediate region can be of any origin, for example of bacterial, viral or plant origin.

Isolation of a cDNA coding for a maize EPSPS:

The different steps which led to the obtaining of maize EPSPS cDNA, which served as substrate for the introduction of the two mutations, are described below. All the operations described below are given by way of example, and correspond to a choice made from among the different methods available for arriving at the same result. This choice has no effect on the quality of the result, and consequently any suitable method may be used by a person skilled in the art to arrive at the same result. Most of the methods of engineering of DNA fragments are described in "Current Protocols in Molecular Biology" Volumes 1 and 2, Ausubel F.M. et al., published by Greene Publishing Associates and Wiley-Interscience (1989) (hereinafter, references to protocols described in this work will be designated "ref. CPMB"). The operations relating to DNA which were performed according to the protocols described in this work are especially the following: ligation of DNA fragments, treatment with Klenow DNA polymerase and T4 DNA polymerase, preparation of plasmid and of bacteriophage λ DNA, either as a

minipreparation or as a maxipreparation, and DNA and RNA analyses according to the Southern and Northern techniques, respectively. Other methods described in this work were followed, and only significant modifications or additions to these protocols have been described below.

Example 1:

1. Obtaining of an *Arabidopsis thaliana* EPSPS fragment

a) Two 20-mer oligonucleotides of respective sequences:

5'-GCTCTGCTCATGTCTGCTCC-3'

5'-GCCCGCCCTTGACAAAGAAA-3'

were synthesized from the sequence of an *Arabidopsis thaliana* EPSPS gene (Klee H.J. et al. (1987) Mol. Gen. Genet., 210, 437-442). These two oligonucleotides are at positions 1523 to 1543 and 1737 to 1717, respectively, of the published sequence, and in opposite orientations.

b) *Arabidopsis thaliana (var. columbia)* total DNA was obtained from Clontech (catalogue reference: 6970-1).

c) 50 nanograms (ng) of DNA are mixed with 300 ng of each of the oligonucleotides and subjected to 35 amplification cycles with a Perkin-Elmer 9600 apparatus, under the conditions of standard medium for amplification which are recommended by the supplier.

The resulting 204-bp fragment constitutes the *Arabidopsis thaliana* EPSPS fragment.

2. Construction of a library of a cDNA from a BMS maize cell line

a) 5 g of filtered cells are ground in liquid nitrogen, and the total nucleic acids are extracted according to the method described by Shure et al. with the following modifications:

- the pH of the lysis buffer is adjusted to PH 9.0;
- after precipitation with isopropanol, the pellet is taken up in water and, after dissolution, adjusted to 2.5 M LiCl. After incubation for 12 h at °C, the pellet from centrifugation for 15 min at 30,000 g at 4°C is resolubilized. The LiCl precipitation step is then repeated. The resolubilized pellet constitutes the RNA fraction of the total nucleic acids.

b) The poly(A)$^+$ RNA fraction of the RNA fraction is obtained by chromatography on an oligo(dT)-cellulose column as described in "Current Protocols in Molecular Biology".

c) Synthesis of double-stranded cDNA having a synthetic EcoRI end: this is carried out according to the protocol of the supplier of the different reagents needed for this synthesis penny form of a kit: the

"copy kit" from the company In Vitrogen.

Two single-stranded and partially complementary oligonucleotides of respective sequences:

5'-AATTCCCGGG-3'

5'-CCCGGG-3' (the latter being phosphorylated)

are ligated with the blunt-ended double-stranded cDNAs.

This ligation of the adaptors results in the creation of SmaI sites attached to the double-stranded cDNAs and EcoRI sites in cohesive form at each end of the double-stranded CDNAs.

d) Creation of the library:

The cDNAs possessing the artificial cohesive EcoRI sites at their ends are ligated with bacteriophage λgt10 cDNA which has been cut with EcoRI and dephosphorylated according to the protocol of the supplier New England Biolabs.

An aliquot of the ligation reaction was encapsidated *in vitro* with encapsidation extracts, namely Gigapack Gold, according to the supplier's instructions; this library was titrated using the bacterium *E. coli* C600*hfl*. The library thereby obtained is amplified and stored according to the instructions of the same supplier, and constitutes the BMS maize cell suspension cDNA library.

3. Screening of the BMS maize cell suspension cDNA library with the *Arabidopsis thaliana* EPSP probe

The protocol followed is that of "Current

Protocols in Molecular Biology" Volumes 1 and 2, Ausubel F.M. et al., published by Greene Publishing Associates and Wiley-Interscience (1989)(CPMB). Briefly, approximately 10^6 recombinant phages are plated out on LB dishes at an average density of 100 phages/cm^2. The lytic plaques are replicated in duplicate on Amersham Hybond N membranes.

h) The DNA was fixed to the filters by 1600kJ UV treatment (Stratagene Stratalinker). The filters were prehybridized in 6×SSC/0.1%SDS/0.25 skimmed milk for 2 h at 65°C. The *Arabidopsis thaliana* EPSPS probe was labelled with [^{32}P]dCTP by random priming according to the supplier's instructions (Pharmacia Ready to Go kit). The specific activity obtained is of the order of 10^8 cpm per µg of fragment. After denaturation for 5 min at 100°C, the probe is added to the prehybridization medium and hybridization is continued for 14 hours at 55°C. The filters are fluorographed for 48 h at -80°C with Kodak XAR5 film and Amersham Hyperscreen RPN enhancing screens. Alignment of the positive spots on the filter with the dishes from which they originate enables zones corresponding to the phages displaying a positive hybridization response with the *Arabidopsis thaliana* EPSPS probe to be picked out from the dish. This step of plating out, transfer, hybridization and recovery is repeated until all the spots in the dish of the successively purified phages prove 100% positive in hybridization. A plaque of

independent phage lysis is then picked out in diluent λ medium (Tris-Cl pH 7.5; 10mM $MgSO_4$; 0.1M NaCl; 0.1% gelatin); these phages in solution constitute the EPSP-positive clones of the BMS maize cell suspension.

4. Preparation and analysis of the DNA of the EPSP clones of the BMS maize cell suspension

Approximately 5×10^8 phages are added to 20 ml of C600hfl bacteria at an OD_{600nm} value of 2/ml and incubated for 15 minutes at 37°C. This suspension is then diluted in 200 ml of bacterial growth medium in a 1-l Erlenmeyer and stirred in a rotary stirrer at 250 rpm. Lysis is noted when the medium clarifies, corresponding to the lysis of the turbid bacteria, and takes place after approximately 4 h of stirring. This supernatant is then treated as described in "Current Protocols in Molecular Biology". The DNA obtained corresponds to the EPSP clones of the BMS maize cell suspension.

One to two µg of this DNA are cut with EcoRI and separated on 0.8% LGTA/TBE agarose gel (ref. CPMB). A final verification consists in checking that the purified DNA does indeed display a hybridization signal with the *Arabidopsis thaliana* EPSPS probe. After electrophoresis, the DNA fragments are transferred onto Amersham Hybond N membranes according to the protocol of Southern described in "Current Protocols in Molecular Biology". The filter is hybridized with the *Arabidopsis thaliana* EPSPS probe according to the

conditions described in section 3 above. The clone displaying a hybridization signal with the *Arabidopsis thaliana* EPSPS probe and containing the longest EcoRI fragment has a size estimated on gel as approximately 1.7 kbp.

5. Obtaining of the clone pRPA-ML-711

Ten µg of the phage clone containing the 1.7-kbp insert are digested with EcoRI and separated on 0.8% LGTA/TBE agarose gel (ref. CPMB). The gel fragment containing the 1.7-kbp insert is excised from the gel by BET staining, and the fragment is treated with β-agarse according to the protocol of the supplier, New England Biolabs. The purified DNA of the 1.7-kbp fragment is ligated at 12°C for 14 h with the DNA of plasmid pUC 19 (New England Biolabs) cut with EcoRI according to the ligation protocol described in "Current Protocols in Molecular Biology". Two µl of the above ligation mixture are used for the transformation of an aliquot of electrocompetent E. coli DH10B; transformation is accomplished by electroporation using the following conditions: the mixture of competent bacteria and and ligation medium is introduced into an electroporation cell of thickness 0.2 cm (Biorad) previously cooled to 0°C. The physical conditions of the electroporation using an electroporator made by Biorad are 2500 volts, 25 µF and 200 Ω. Under these conditions, the mean discharge time of the condenser is of the order of 4.2 milliseconds. The bacteria are then

taken up in 1 ml of SOC medium (ref. CPMB) and stirred for 1 hour at 200 rpm on a rotary stirrer in 15-ml Corning tubes. After plating out on LB/agar medium supplemented with 100 µg/ml of carbenicillin, minipreparations of the bacterial clones which have grown after one night at 37°C is produced according to the protocol described in "Current Protocols in Molecular Biology". After digestion of the DNA with EcoRI and separation by electrophoresis on 0.8% LGTA/TBE agarose gel (ref. CPMB), the clones possessing a 1.7-kbp insert are retained. A final verification consists in checking that the purified DNA does indeed display a hybridization signal with the *Arabidopsis thaliana* EPSPS probe. After electrophoresis, the DNA fragments are transferred onto Amersham Hybond N membranes according to the protocol of Southern described in "Current Protocols in Molecular Biology". The filter is hybridized with the *Arabidopsis thaliana* EPSPS probe according to the conditions described in section 3 above. The plasmid clone possessing a 1.7-kbp insert and hybridizing with the *Arabidopsis thaliana* EPSPS probe was prepared on a larger scale, and the DNA resulting from the lysis of the bacteria was purified on a CsCl gradient as described in "Current Protocols in Molecular Biology". The purified DNA was partially sequenced with a Pharmacia kit according to the supplier's instructions and using as primers the M13 direct and reverse universal primers ordered from the

same supplier. The partial sequence produced covers approximately 0.5 kbp. The derived amino acid sequence in the region of the mature protein (approximately 50 amino acid residues) displays 100% identity with the corresponding amino sequence of mature maize EPSPS described in American Patent USP 4,971,908). This clone, corresponding to a 1.7-kbp EcoRI fragment of the EPSP DNA of the BMS maize cell suspension, was designated pRPA-ML-711. The complete sequence of this clone was determined on both strands using the protocol of the Pharmacia kit and synthesizing complementary oligonucleotides and those of the opposite orientation every 250 bp approximately. The complete sequence obtained of this 1713-bp clone is presented in SEQ ID No. 1.

6. Obtaining of the clone pRPA-ML-715

Analysis of the sequence of the clone pRPA-ML-711, and especially comparison of the derived amino acid sequence with that of maize, shows a sequence extension of 92 bp upstream of the GCG codon coding for the NH_2-terminal alanine of the mature portion of maize EPSPS (American Patent USP 4,971,908). Similarly, an extension of 288 bp downstream of the AAT codon coding for the COOH-terminal asparagine of the mature portion of maize EPSPS (American Patent USP 4,971,908) is observed. These two portions could correspond, in the case of the NH_2-terminal extension to a portion of the sequence of a transit peptide for plastid localization,

and, in the case of the COOH-terminal extension, to the untranslated 3′ region of the cDNA.

In order to obtain a cDNA coding for the mature portion of the maize EPSPS cDNA, as described in USP 4,971,908, the following operations were carried out:

a) Removal of the untranslated 3′ region: construction of pRPA-ML-712:

The clone pRPA-ML-711 was cut with the restriction enzyme AseI, and the ends resulting from this cleavage were rendered blunt by treatment with the Klenow fragment of DNA polymerase I according to the protocol described in CPMB. A cleavage with the restriction enzyme SacII was then performed. The DNA resulting from these operations was separated by electrophoresis on 1% LGTA/TBE agarose gel (ref. CPMB).

The gel fragment containing the 0.4-kbp "AseI-blunt ends/SacII" insert was excised from the gel and purified according to the protocol described in section 5 above. The DNA of the clone pRPA-ML-711 was cut with the restriction enzyme HindIII [lacuna] located in the polylinker of the cloning vector pUC19, and the ends resulting from this cleavage were rendered blunt by treatment with the Klenow fragment of DNA polymerase I. A cleavage with the restriction enzyme SacII was then performed. The DNA resulting from these manipulations was separated by electrophoresis on 0.7% LGTA/TBE agarose gel (ref. CPMB).

The gel fragment containing the approximately 3.7-kbp HindIII-blunt ends/SacII insert was excised from the gel and purified according to the protocol described in section 5 above.

The two inserts were ligated, and 2 µl of the ligation mixture were used to transform *E. coli* DH10B as described above in section 5.

The plasmid DNA content of different clones was analysed according to the procedure described for pRPA-ML-711. One of the plasmid clones selected contains an approximately 1.45-kbp EcoRI-HindIII insert. The sequence of the terminal ends of this clone reveals that the 5′ end of the insert corresponds exactly to the corresponding end of pRPA-ML-711, and that the 3′-terminal end possesses the following sequence:

"5′-...AATTAAGCTCTAGAGTCGACCTGCAGGCATGCAAGCTT-3′".

The underlined sequence corresponds to the codon of the COOH-terminal amino acid asparagine, the next codon corresponding to the translation stop codon. The nucleotides downstream correspond to sequence elements of the pUC19 polylinker. This clone comprising the pRPA-ML-711 sequence up to the translation termination site of mature maize EPSPS and followed by sequences of the pUC 19 polylinker up to the HindIII site was designated pRPA-ML-712.

b) Modification of the 5′ end of pRPA-ML-712: construction of pRPA-ML-715:

The clone pRPA-ML-712 was cut with the restriction enzymes PstI and HindIII. The DNA resulting from these manipulations was separated by electrophoresis on 0.8% LGTA/TBE agarose gel (ref. CPMB). The gel fragment containing the 1.3-kbp PstI-EcoRI insert was excised from the gel and purified according to the protocol described in section 5 above. This insert was ligated in the presence of an equimolecular amount of each of the two partially complementary oligonucleotides of sequence:

Oligo 1: 5'-GAGCCGAGCTCCATGGCCGGCGCCGAGGAGATCGTGCTGCA-3'

Oligo 2: 5'-GCACGATCTCCTCGGCGCCGGCCATGGAGCTCGGCTC-3'

as well as in the presence of plasmid pUC19 DNA digested with the restriction enzymes BamHI and HindIII.

Two µl of the ligation mixture were used to transform *E. coli* DH10B as described above in section 5. After analysis of the plasmid DNA content of different clones according to the procedure described above in section 5, one of the clones possessing an approximately 1.3-kbp insert was retained for subsequent analyses. The sequence of the 5'-terminal end of the selected clone reveals that the DNA sequence in this region is the following: sequence of the pUC19 polylinker from the EcoRI to the BamHI sites, followed by the sequence of the oligonucleotides used in the cloning, followed by the remainder of the sequence present in pRPA-ML-712. This clone was designated pRPA-

ML-713. This clone possesses a methionine ATG codon included in an NcoI site upstream of the N-terminal alanine codon of mature EPSP synthase. Furthermore, the alanine and glycine codons of the N-terminal end have been preserved, but modified on the third variable base: initial GCGGGT gives modified GCCGGC.

The clone pRPA-ML-713 was cut with the restriction enzyme HindIII, and the ends of this cleavage were rendered blunt by treatment with the Klenow fragment of DNA polymerase I. A cleavage with restriction enzyme SacI was then performed. The DNA resulting from these manipulations was separated by electrophoresis on 0.8% LGTA/TBE agarose gel (ref. CPMB). The gel fragment containing the 1.3-kbp "HindIII-blunt ends/SacI" insert was excised from the gel and purified according to the protocol described in section 5 above. This insert was ligated in the presence of plasmid pUC19 DNA digested with restriction enzyme XbaI, and the ends of this cleavage were rendered blunt by treatment with the Klenow fragment of DNA polymerase I. A cleavage with the restriction enzyme SacI was then performed. Two µl of the ligation mixture were used to transform *E. coli* DH10B as described above in section 5. After analysis of the plasmid DNA content of different clones according to the procedure described above in section 5, one of the clones possessing an approximately 1.3-kbp insert was retained for subsequent analyses. The sequence of the

terminal ends of the selected clone reveals that the DNA sequence is the following: sequence of the pUC19 polylinker from the EcoRI to SacI sites, followed by the sequence of the oligonucleotides used in the cloning from which the 4 bp GATCC of the oligonucleotide 1 described above have been deleted, followed by the remainder of the sequence present in pRPA-ML-712 up to the HindIII site and sequence of the pUC19 polylinker from XbaI to HindIII. This clone was designated pRPA-ML-715.

7. Obtaining of a cDNA coding for a mutated maize EPSPS

All the mutagenesis steps were carried out with the Pharmacia U.S.E. mutagenesis kit according to the supplier's instructions. The principle of this mutagenesis system is as follows: plasmid DNA is denatured by heat and reassociated in the presence of a molar excess of, on the one hand the mutagenesis oligonucleotide, and on the other hand an oligonucleotide enabling a unique restriction enzyme site present in the polylinker to be eliminated. After the reassociation step, synthesis of the complementary strand is carried out by the action of T4 DNA polymerase in the presence of T4 DNA ligase and gene 32 protein in a suitable buffer which is supplied. The synthesis product is incubated in the presence of the restriction enzyme for which the site is assumed to have disappeared by mutagenesis. The *E. coli* strain

possessing, in particular, the mutS mutation is used as host for the transformation of this DNA. After growth in liquid medium, the total plasmid DNA is prepared and incubated in the presence of the restriction enzyme used before. After these treatments, *E. coli* strain DH10B is used as host for the transformation. The plasmid DNA of the clones isolated is prepared, and the presence of the mutation introduced is verified by sequencing.

A)- modification of sites or sequences without in principle affecting the EPSPS-resistance character of maize to products which are competitive inhibitors of EPSP synthase activity: elimination of an internal NcoI site from pRPA-ML-715.

The pRPA-ML-715 sequence is numbered arbitrarily by placing the first base of the N-terminal alanine codon GCC at position 1. This sequence possesses an NcoI site at position 1217. The site-modification oligonucleotide possesses the sequence:

5'-CCACAGGATGGCGATGGCCTTCTCC-3'.

After sequencing according to the references given above, the sequence read after mutagenesis corresponds to that of the oligonucleotide used. The NcoI site has indeed been eliminated, and the translation into amino acids in this region preserves the initial sequence present in pRPA-ML-715.

This clone was designated pRPA-ML-716.

The 1340-bp sequence of this clone is

presented in SEQ ID No. 2 and SEQ ID No. 3.

B)- sequence modifications enabling the EPSPS-resistance character of maize to products which are competitive inhibitors of EPSP synthase activity to be increased.

The following oligonucleotides were used:

a) mutation Thr 102 → Ile.

5'-GAATGCTGGAATCGCAATGCGGCCATTGACAGC-3'

b) mutation Pro 106 → Ser.

5'-GAATGCTGGAACTGCAATGCGGTCCTTGACAGC-3'

c) mutations Gly 101 → Ala and Thr 102 → Ile.

5'-CTTGGGGAATGCTGCCATCGCAATGCGGCCATTG-3'

d) mutations Thr 102 → Ile and Pro 106 → Ser.

5'-GGGGAATGCTGGAATCGCAATGCGGTCCTTGACAGC-3'

After sequencing, the sequence read after mutagenesis on the three mutated fragments is identical to the parent pRPA-ML-716 DNA sequence, with the exception of the mutagenized region which corresponds to that of the mutagenesis oligonucleotides used. These clones were designated: pRPA-ML-717 for the mutation Thr 102 → Ile, pRPA-ML-718 for the mutation Pro 106 → Ser, pRPA-ML-719 for the mutations Gly 101 → Ala and Thr 102 → Ile and pRPA-ML-720 for the mutations Thr 102 → Ile and Pro 106 → Ser.

The 1340-bp sequence of pRPA-ML-720 is presented in SEQ ID No. 4 and SEQ ID No. 5.

The 1395-bp NcoI-HindIII insert is the basis of all the constructions used for the transformation of plants for the introduction of resistance to herbicides which are competitive inhibitors of EPSPS, and especially glyphosate resistance. This insert will be designated in the remainder of the description "the maize EPSPS double mutant".

Example 2:

Glyphosate tolerance of the different mutants in vitro

2.a: Extraction of EPSP synthase

The different EPSP synthase genes are introduced in the form of an NcoI-HindIII cassette into the plasmid vector pTrc99a (Pharmacia, ref: 27-5007-01) cut with NcoI and HindIII. Recombinant *E. coli* DH10B bacteria overexpressing the different EPSP synthases are sonicated in 40 ml of buffer per 10 g of pelleted cells, and washed with this same buffer (200 mM Tris-HCl pH 7.8, 50 mM mercaptoethanol, 5 mM EDTA and 1 mM PMSF), to which 1 g of polyvinylpyrrolidone is added. The suspension is stirred for 15 minutes at 4°C and then centrifuged for 20 minutes at 27,000 g and 4°C.

Ammonium sulphate is added to the supernatant to bring the solution to 40% saturation with respect to ammonium sulphate. The mixture is centrifuged for 20 minutes at 27,000 g and 4°C. Ammonium sulphate is

added to the new supernatant to bring the solution to 70% saturation with respect to ammonium sulphate. The mixture is centrifuged for 30 minutes at 27,000 g and 4°C. The EPSP synthase present in this protein pellet is taken up in 1 ml of buffer (20 mM Tris-HCl pH 7.8 and 50 mM mercaptoethanol). This solution is dialysed overnight against two litres of this same buffer at 4°C.

2.b: Enzyme activity

The activity of each enzyme, as well as its glyphosate resistance, is measured in vitro over 10 minutes at 37°C in the following reaction mixture: 100 mM maleic acid pH 5.6, 1 mM phosphoenolpyruvate, 3 mM shikimate 3-phosphate (prepared according to Knowles P.F. and Sprinson D.B. 1970. Methods in Enzymol 17A, 351-352 from *Aerobacter aerogenes* strain ATCC 25597) and 10 mM potassium fluoride. The enzyme extract is added at the last moment after the addition of glyphosate, the final concentration of which varies from 0 to 20 mM.

The activity is measured by assaying the phosphate liberated according to the technique of Tausky H.A. and Shorr E. 1953. J. Biol. Chem. 202, 675-685.

Under these conditions, the wild-type (WT) enzyme is already 85% inhibited at a glyphosate concentration of 0.12 mM. At this concentration, the mutant enzyme known as Ser106 is only 50% inhibited,

and the other three mutants, Ile102, Ile102/Ser106 and Ala101/Ile102, show little or no inhibition.

The glyphosate concentration has to be multiplied by ten, that is to say 1.2 mM, in order to produce a 50% inhibition of the mutant enzyme Ile102, the mutants Ile102/Ser106, Ala/Ile and Ala still not being inhibited.

It should be noted that the activity of the mutants Ala/Ile and Ala is not inhibited up to glyphosate concentrations of 10mM, and that that of the mutant Ile102/Ser106 is not reduced even if the glyphosate concentration is multiplied by 2, that is to say 20 mM.

Example 3:

Resistance of transformed tobacco plants

1-1- Transformation

The vector pRPA-RD-173 is introduced into *Agrobacterium tumefaciens* strain EHA101 (Hood et al., 1987) carrying the cosmid pTVK291 (Komari et al., 1986). The transformation technique is based on the procedure of Horsh et al. (1985).

1-2- Regeneration

The regeneration of PBD6 tobacco (source SEITA France) from leaf explants is carried out on a Murashige and Skoog (MS) basal medium comprising 30 g/l of sucrose as well as 200 µg/ml of kanamycin. The leaf explants are removed from plants cultivated in the greenhouse or *in vitro* and transformed according to the

leaf disc technique (Science, 1985, Vol. 227, pp. 1229-1231) in three successive steps: the first comprises the induction of shoots on a medium supplemented with 30 g/l of sucrose containing 0.05 mg/l of naphthylacetic acid (NAA) and 2 mg/l of benzylaminopurine (BAP) for 15 days. The shoots formed during this step are then developed for 10 days by culturing on an MS medium supplemented with 30 g/l of sucrose but not containing any hormone. Shoots which have developed are then removed and cultured on an MS rooting medium having half the content of salts, vitamins and sugar and not containing any hormone. After approximately 15 days, the rooted shoots are transferred to soil.

1-3- Glyphosate resistance

Twenty transformed plants were regenerated and transferred to the greenhouse for the construction of pRPA-RD-173. These plants were treated in the greenhouse at the 5-leaf stage with an aqueous suspension of RoundUp corresponding to 0.8 kg of glyphosate active substance per hectare.

The results correspond to the observation of phytotoxicity indices recorded 3 weeks after treatment. Under these conditions, it is found that the plants transformed with the construction pRPA-RD-173 display very good tolerance, whereas the untransformed control plants are completely destroyed.

These results show clearly the improvement

brought about by the use of a chimeric gene according to the invention for the same gene coding for glyphosate tolerance.

Example 4:

Transformation and selection of maize cells

BMS (Black Mexican Sweet) maize cells in an exponential growth phase are bombarded with the construction pRPA-RD-130 according to the principle and the protocol described by Klein et al. 1987 (Klein T.M., Wolf E.D., Wu R. and Sandford J.C. (1987): High velocity microprojectiles for delivering nucleic acids into living cells, NATURE Vol. 327 pp. 70-73).

Two days after bombardment, the cells are transferred to the same medium containing 2 mM N-(phosphomethyl)glycine.

After 8 weeks of selection on this medium, calluses which develop are selected, then amplified and analysed by PCR, and reveal clearly the presence of the chimeric OTP-EPSPS gene.

Cells not bombarded and grown on the same medium containing 2 mM N-(phosphomethyl)glycine are blocked by the herbicide and do not develop.

The transformed plants according to the invention may be used as parents for obtaining lines and hybrids having the phenotypic character corresponding to the expression of the chimeric gene introduced.

Description of the constructions of the plasmids

pRPA-RD-124: Addition of a "nos" polyadenylation signal to pRPA-ML-720 with creation of a cloning cassette containing the maize double mutant EPSPS gene (Thr 102 → Ile and Pro 106 → Ser). pRPA-ML-720 is digested with HindIII and treated with the Klenow fragment of *E. coli* DNA polymerase I to produce a blunt end. A second digestion is performed with NcoI, and the EPSPS fragment is purified. The EPSPS gene is then ligated with purified pRPA-RD-12 (a cloning cassette containing the polyadenylation signal of nopaline synthase) to give pRPA-RD-124. To obtain the useful purified vector pRPA-RD-12, it was necessary for the latter to be digested beforehand with SalI, treated with Klenow DNA polymerase and then digested a second time with NcoI.

pRPA-RD-125: Addition of an optimized transit peptide (OTP) to pRPA-RD-124 with creation of a cloning cassette containing the EPSPS gene targeted on the plasmids. pRPA-RD-7 (European Patent Application EP 652 286) is digested with SphI, treated with T4 DNA polymerase and then digested with SpeI, and the OTP fragment is purified. This OTP fragment is cloned into pRPA-RD-124 which has previously been digested with NcoI, treated with Klenow DNA polymerase to remove the protruding 3' portion and then digested with SpeI. This clone is then sequenced in order to ensure correct

translational fusion between the OTP and the EPSPS gene. pRPA-RD-125 is then obtained.

pRPA-RD-130: Addition of the H3C4 maize histone promoter and of adh1 intron 1 sequences of pRPA-RD-123 (Patent Application EP 507 698) to pRPA-RD-125 with creation of a cassette for expression in plants for the expression of the double mutant EPSPS gene in the tissues of monocotyledons. pRPA-RD-123 (a cassette containing the H3C4 maize histone promoter fused with the adh1 intron 1) is digested with NcoI and SacI. The DNA fragment containing the promoter derived from pRPA-RD-123 is then purified and ligated with pRPA-RD-125 which has previously been digested with NcoI and SacI.

pRPA-RD-159: Addition of the H4A748 *Arabidopsis* histone double promoter (Patent Application EP 507 698) to pRPA-RD-125 with creation of a cassette for expression in plants for the expression of the "OTP-double mutant EPSPS gene" gene in the tissues of dicotyledons. pRPA-RD-132 (a cassette containing the H4A748 double promoter (Patent Application EP 507 698)) is digested with NcoI and SacI. The purified promoter fragment is then cloned into [lacuna] which has been digested with EcoI and SacI.

pRPA-RD-173: Addition of the "H4A748 promoter-OTP-double mutant EPSPS gene" gene of pRPA-RD-159 to plasmid pRPA-BL-150A (European Patent Application 508 909) with creation of an *Agrobacterium*

tumefaciens transformation vector. pRPA-RD-159 is digested with NotI and treated with Klenow polymerase. This fragment is then cloned into pRPA-BL-150A with SmaI.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Lebrun, Michel
Du Rana, Richard T
Saillard, Alain

(ii) TITLE OF INVENTION: 5-enol pyruvylshikimate-3-phosphate synthase mutee, gene codant pour cette proteine et plantes transformees contenant ce gene

(iii) NUMBER OF SEQUENCES: 5

(iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Francois Chretien
(B) STREET: 1420 rue Pierre Baizet
(C) CITY: Lyon Cedex 09
(E) COUNTRY: France
(F) ZIP: 69263

(v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Chretien, Francois

(ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: (33)72-29-26-46
(B) TELEFAX: (33)72-29-28-43

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1713 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Zea mays
(B) STRAIN: Black Mexican Sweet
(F) TISSUE TYPE: Callus

(vii) IMMEDIATE SOURCE:
(A) LIBRARY: lambda gt10
(B) CLONE: pRPA-ML-711

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AATCAATTTC ACACAGGAAA CAGCTATGAC CATGATTACG AATTCGGGCC CGGGCGCGTG 60

ATCCGGCGGC GGCAGCGGCG GCGGCGGTGC AGCCGGGTGC CGAGGAGATC GTGCTGCAGC 120

CCATCAAGGA GATCTCCGGC ACCGTCAAGC TGCCGGGGTC CAAGTCGCTT TCCAACCGGA 180

TCTGCCTACT CGCCGCCCTG TCCGAGGGGA CAACAGTCGT TGATAACCTG CTGAACAGTG 240

AGGATGTCCA CTACATGCTC GGGGCCTTGA GGACTCTTGG TCTCTCTGTC GAAGCGGACA 300

AAGCTGCCAA AAGAGCTGTA GTTGTTGGCT GTGGTGGAAA GTTCCCAGTT GAGGATGCTA 360

AAGAGGAAGT GCAGCTCTTC TTGGGGAATG CTGGAACTGC AATGCGGCCA TTGACAGCAG 420

CTGTTACTGC TGCTGGTGGA AATGCAACTT ACGTGCTTGA TGGAGTACCA AGAATGAGGG 480

AGATACCCAT TTTTGACTTG GTTTTCGGAT TGAAGCAGCT TGTTGCAGAT GTTGATTGTT 540

TCCTTTTCAC TTACTTCCCA CCTTTTGTTG TCAATGGAAT CGGAGGGCTA CGTGGTGGCA 600

AGTTGAAGCT GTCTTTGTCC ATCATCAGTC AGTACTTGAG TGCCTTTCTG ATGGCTGCTC 660

GTTTTGCTCT TGTTTGATTG GAGATTGAAA TCATTGATAA ATTAATCTCC ATTCCGTACG 720

TCGAAATGAC ATTGAGATTG ATGGAGCGTT TTGGTGTGAA AGCAGAGCAT TCTGATAGCT 780

GGGACAGATT CTACATTAAG GGAGGTCAAA AATACAAGTC CCCTAAAAAT GCCTATGTTG 840

AAGGTGATGC CTCAAGCGCA AGCTATTTCT TGGCTGGTGC TGCAATTACT GGAGGGACTG 900

TGACTGTGGA AGGTTGTGGC ACCACCAGTT TGCAGGGTGA TGTGAAGTTT GCTGAGGTAC 960

TGGAGATGAT GGGAGCGAAG GTTACATGGA CCGAGACTAG CGTAACTGTT ACTGGCCCAC 1020

CGCGGGAGCC ATTTGGGAGG AAACACCTCA AGCCGATTGA TGTCAACATG AACAAGATGC 1080

CTGATGTCGC CATGACTCTT GCTGTGGTTG CCCTCTTTGC CGATGGCCCG ACAGCCATCA 1140

GAGACGTGGC TTCCTGGAGA GTAAAGGAGA CCGAGAGGAT GGTTGCGATC CGGACGGAGC 1200

TAACCAAGCT GGGAGCATCT GTTGAGGAAG GGCCGGACTA CTGCATCATC ACGCCGCCGG 1260

AGAAGCTGAA CGTGACGGCG ATCGACACGT ACGACGACCA CAGGATGGCC ATGGCCTTCT 1320

CCCTTGCCGC CTGTGCCGAG GTCCCCGTCA CCATCCGGGA CCCTGGGTGC ACCCGGAAGA 1380

CCTTCCCCGA CTACTTCGAT GTGCTGAGCA CTTTCGTCAA GAATTAATAA AGCGTGCGAT 1440

ACTACCACGC AGCTTGATTG AAGTGATAGG CTTGTGCTGA GGAAATACAT TTCTTTTGTT 1500

CTGTTTTTCT CTTTCACGGG ATTAAGTTTT GAGTCTGTAA CGTTAGTTGT TTGTAGCAAG 1560

TTTCTATTTC GGATCTTAAG TTTGTGCACT GTAAGCCAAA TTTCATTTCA AGAGTGGTTC 1620

GTTGGAATAA TAAGAATAAT AAATTACGTT TCAGTGAAAA AAAAAAAAA AAAAAAAAA 1680

AAAAAAAAA AAAAAAAAA AACCCGGGAA TTC 1713

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1340 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Zea mays
(B) STRAIN: Black Mexican Sweet

(vii) IMMEDIATE SOURCE:
(B) CLONE: pRPA-ML-716

(ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 6..1337

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CCATG GCC GGC GCC GAG GAG ATC GTG CTG CAG CCC ATC AAG GAG ATC 47
Ala Gly Ala Glu Glu Ile Val Leu Gln Pro Ile Lys Glu Ile
1 5 10

TCC GGC ACC GTC AAG CTG CCG GGG TCC AAG TCG CTT TCC AAC CGG ATC 95
Ser Gly Thr Val Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile
15 20 25 30

CTC CTA CTC GCC GCC CTG TCC GAG GGG ACA ACA GTG GTT GAT AAC CTG 143
Leu Leu Leu Ala Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu
35 40 45

CTG AAC AGT GAG GAT GTC CAC TAC ATG CTC GGG GCC TTG AGG ACT CTT 191
Leu Asn Ser Glu Asp Val His Tyr Met Leu Gly Ala Leu Arg Thr Leu

```
                    50                  55                  60

GGT CTC TCT GTC GAA GCT GAC AAA GCT GCC AAA AGA GCT GTA GTT GTT      239
Gly Leu Ser Val Glu Ala Asp Lys Ala Ala Lys Arg Ala Val Val Val
    65                  70                  75

GGC TGT GGT GGA AAG TTC CCA GTT GAG GAT GCT AAA GAG GAA GTC CAG      287
Gly Cys Gly Gly Lys Phe Pro Val Glu Asp Ala Lys Glu Glu Val Gln
80                  85                  90

CTC TTC TTG GGG AAT GCT GGA ACT GCA ATG CGG CCA TTG ACA GCA GCT      335
Leu Phe Leu Gly Asn Ala Gly Thr Ala Met Arg Pro Leu Thr Ala Ala
95                  100                 105                 110

GTT ACT GCT GCT GGT GGA AAT GCA ACT TAC GTG CTT GAT GGA GTA CCA      383
Val Thr Ala Ala Gly Gly Asn Ala Thr Tyr Val Leu Asp Gly Val Pro
                115                 120                 125

AGA ATG AGG GAG AGA CCC ATT GGC GAC TTG GTT GTC GGA TTG AAG CAG      431
Arg Met Arg Glu Arg Pro Ile Gly Asp Leu Val Val Gly Leu Lys Gln
            130                 135                 140

CTT GGT GCA GAT GTT GAT TGT TTC CTT GGC ACT GAC TGC CCA CCT GTT      479
Leu Gly Ala Asp Val Asp Cys Phe Leu Gly Thr Asp Cys Pro Pro Val
        145                 150                 155

CGT GTC AAT GGA ATC GGA GGG CTA CCT GGT GGC AAG GTC AAG CTG TCT      527
Arg Val Asn Gly Ile Gly Gly Leu Pro Gly Gly Lys Val Lys Leu Ser
    160                 165                 170

GGC TCC ATC AGC AGT CAG TAC TTG AGT GCC TTG CTG ATG GCT GCT CCT      575
Gly Ser Ile Ser Ser Gln Tyr Leu Ser Ala Leu Leu Met Ala Ala Pro
175                 180                 185                 190

TTG GCT CTT GGG GAT GTG GAG ATT GAA ATC ATT GAT AAA TTA ATC TCC      623
Leu Ala Leu Gly Asp Val Glu Ile Glu Ile Ile Asp Lys Leu Ile Ser
                195                 200                 205

ATT CCG TAC GTC GAA ATG ACA TTG AGA TTG ATG GAG CGT TTT GGT GTG      671
Ile Pro Tyr Val Glu Met Thr Leu Arg Leu Met Glu Arg Phe Gly Val
            210                 215                 220

AAA GCA GAG CAT TCT GAT AGC TGG GAC AGA TTC TAC ATT AAG GGA GGT      719
Lys Ala Glu His Ser Asp Ser Trp Asp Arg Phe Tyr Ile Lys Gly Gly
        225                 230                 235

CAA AAA TAC AAG TCC CCT AAA AAT GCC TAT GTT GAA GGT GAT GCC TCA      767
Gln Lys Tyr Lys Ser Pro Lys Asn Ala Tyr Val Glu Gly Asp Ala Ser
    240                 245                 250

AGC GCA AGC TAT TTC TTG GCT GGT GCT GCA ATT ACT GGA GGG ACT GTG      815
Ser Ala Ser Tyr Phe Leu Ala Gly Ala Ala Ile Thr Gly Gly Thr Val
255                 260                 265                 270

ACT GTG GAA GGT TGT GGC ACC ACC AGT TTG CAG GGT GAT GTG AAG TTT      863
Thr Val Glu Gly Cys Gly Thr Thr Ser Leu Gln Gly Asp Val Lys Phe
                275                 280                 285

GCT GAG GTA CTG GAG ATG ATG GGA GCG AAG GTT ACA TGG ACC GAG ACT      911
Ala Glu Val Leu Glu Met Met Gly Ala Lys Val Thr Trp Thr Glu Thr
            290                 295                 300

AGC GTA ACT GTT ACT GGC CCA CCG CGG GAG CCA TTT GGG AGG AAA CAC      959
Ser Val Thr Val Thr Gly Pro Pro Arg Glu Pro Phe Gly Arg Lys His
        305                 310                 315

CTC AAG GCG ATT GAT GTC AAC ATG AAC AAG ATG CCT GAT GTC GCC ATG     1007
Leu Lys Ala Ile Asp Val Asn Met Asn Lys Met Pro Asp Val Ala Met
    320                 325                 330

ACT CTT GCT GTG GTT GCC CTC TTT GCC GAT GGC CCG ACA GCC ATC AGA     1055
Thr Leu Ala Val Val Ala Leu Phe Ala Asp Gly Pro Thr Ala Ile Arg
335                 340                 345                 350

GAC GTG GCT TCC TGG AGA GTA AAG GAG ACC GAG AGG ATG GTT GCG ATC     1103
Asp Val Ala Ser Trp Arg Val Lys Glu Thr Glu Arg Met Val Ala Ile
                355                 360                 365

CGG ACG GAG CTA ACC AAG CTG GGA GCA TCT GTT GAG GAA GGG CCG GAC     1151
Arg Thr Glu Leu Thr Lys Leu Gly Ala Ser Val Glu Glu Gly Pro Asp
```

```
                370                 375                 380

TAC TGC ATC ATC ACC CCT CCT GAG AAG CTG AAC GTG ACT GCT ATC GAC      1199
Tyr Cys Ile Ile Thr Pro Pro Glu Lys Leu Asn Val Thr Ala Ile Asp
        385                 390                 395

ACT TAC GAC GAC CAC AGG ATG GCC ATG GCC TTC TCC CTT GCC GCC TGT      1247
Thr Tyr Asp Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys
    400                 405                 410

GCC GAG GTC CCC GTC ACC ATC CGG GAC CCT GGG TGC ACC CGG AAG ACC      1295
Ala Glu Val Pro Val Thr Ile Arg Asp Pro Gly Cys Thr Arg Lys Thr
415                 420                 425                 430

TTC CCC GAC TAC TTC GAT GTG CTG AGC ACT TTC GTC AAG AAT               1337
Phe Pro Asp Tyr Phe Asp Val Leu Ser Thr Phe Val Lys Asn
                435                 440

TAA                                                                  1340
```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 444 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```
Ala Gly Ala Glu Glu Ile Val Leu Gln Pro Ile Lys Glu Ile Ser Gly
  1               5                  10                  15

Thr Val Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile Leu Leu
             20                  25                  30

Leu Ala Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu Leu Asn
         35                  40                  45

Ser Glu Asp Val His Tyr Met Leu Gly Ala Leu Arg Thr Leu Gly Leu
     50                  55                  60

Ser Val Glu Ala Asp Lys Ala Ala Lys Arg Ala Val Val Val Gly Cys
 65                  70                  75                  80

Gly Gly Lys Phe Pro Val Glu Asp Ala Lys Glu Glu Val Gln Leu Phe
                 85                  90                  95

Leu Gly Asn Ala Gly Thr Ala Met Arg Pro Leu Thr Ala Ala Val Thr
            100                 105                 110

Ala Ala Gly Gly Asn Ala Thr Tyr Val Leu Asp Gly Val Pro Arg Met
        115                 120                 125

Arg Glu Arg Pro Ile Gly Asp Leu Val Val Gly Leu Lys Gln Leu Gly
    130                 135                 140

Ala Asp Val Asp Cys Phe Leu Gly Thr Asp Cys Pro Pro Val Arg Val
145                 150                 155                 160

Asn Gly Ile Gly Gly Leu Pro Gly Gly Lys Val Lys Leu Ser Gly Ser
                165                 170                 175

Ile Ser Ser Gln Tyr Leu Ser Ala Leu Leu Met Ala Ala Pro Leu Ala
            180                 185                 190

Leu Gly Asp Val Glu Ile Glu Ile Ile Asp Lys Leu Ile Ser Ile Pro
        195                 200                 205

Tyr Val Glu Met Thr Leu Arg Leu Met Glu Arg Phe Gly Val Lys Ala
    210                 215                 220

Glu His Ser Asp Ser Trp Asp Arg Phe Tyr Ile Lys Gly Gly Gln Lys
225                 230                 235                 240

Tyr Lys Ser Pro Lys Asn Ala Tyr Val Glu Gly Asp Ala Ser Ser Ala
                245                 250                 255
```

Ser Tyr Phe Leu Ala Gly Ala Ala Ile Thr Gly Gly Thr Val Thr Val
240 245 270

Glu Gly Cys Gly Thr Thr Ser Leu Gln Gly Asp Val Lys Phe Ala Glu
255 260 265

Val Leu Glu Met Met Gly Ala Lys Val Thr Trp Thr Glu Thr Ser Val
270 295 300

Thr Val Thr Gly Pro Pro Arg Glu Pro Phe Gly Arg Lys His Leu Lys
305 310 315 320

Ala Ile Asp Val Asn Met Asn Lys Met Pro Asp Val Ala Met Thr Leu
325 330 335

Ala Val Val Ala Leu Phe Ala Asp Gly Pro Thr Ala Ile Arg Asp Val
340 345 350

Ala Ser Trp Arg Val Lys Glu Thr Glu Arg Met Val Ala Ile Arg Thr
355 360 365

Glu Leu Thr Lys Leu Gly Ala Ser Val Glu Glu Gly Pro Asp Tyr Cys
370 375 380

Ile Ile Thr Pro Pro Glu Lys Leu Asn Val Thr Ala Ile Asp Thr Tyr
385 390 395 400

Asp Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys Ala Glu
405 410 415

Val Pro Val Thr Ile Arg Asp Pro Gly Cys Thr Arg Lys Thr Phe Pro
420 425 430

Asp Tyr Phe Asp Val Leu Ser Thr Phe Val Lys Asn
435 440

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1340 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:
(A) ORGANISM: Zea mays
(B) STRAIN: Black Mexican Sweet

(vii) IMMEDIATE SOURCE:
(B) CLONE: pRPA-ML-720

(ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 6..1337

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCATG GCC GGC GCC GAG GAG ATC GTG CTG CAG CCC ATC AAG GAG ATC 47
Ala Gly Ala Glu Glu Ile Val Leu Gln Pro Ile Lys Glu Ile
1 5 10

TCC GGC ACC GTC AAG CTG CCG GGG TCC AAG TCG CTT TCC AAC CGG ATC 95
Ser Gly Thr Val Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile
15 20 25 30

CTC CTA CTC GCC GCC CTG TCC GAG GGG ACA ACA GTG GTT GAT AAC CTG 143
Leu Leu Leu Ala Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu
35 40 45

CTG AAC AGT GAG GAT GTC CAC TAC ATG CTC GGG GCC TTG AGG ACT CTT 191
Leu Asn Ser Glu Asp Val His Tyr Met Leu Gly Ala Leu Arg Thr Leu
50 55 60

GGT CTC TCT GTC GAA GCG GAC AAA GCT GCC AAA AGA GCT GTA GTT GTT 239
Gly Leu Ser Val Glu Ala Asp Lys Ala Ala Lys Arg Ala Val Val Val
65 70 75

GGC TTT GGT GGA AAG TTC CCA GTT GAG GAT GCT AAA GAG GAA GTG CAG 287
Gly Cys Gly Gly Lys Phe Pro Val Glu Asp Ala Lys Glu Glu Val Gln
80 90

CTC TTC TTG GGG AAT GCT GGA ATC GCA ATG CGC TCC TTG ACA GCA GCT 335
Leu Phe Leu Gly Asn Ala Gly Ile Ala Met Arg Ser Leu Thr Ala Ala
100 105 110

GTT ACT GCT GCT GGT GGA AAT GCA ACT TAC GTG CTT GAT GGA GTA CCA 383
Val Thr Ala Ala Gly Gly Asn Ala Thr Tyr Val Leu Asp Gly Val Pro
115 120 125

AGA ATG AGG GAG AGA CCC ATT GGC GAC TTG GTT GTC GGA TTG AAG CAG 431
Arg Met Arg Glu Arg Pro Ile Gly Asp Leu Val Val Gly Leu Lys Gln
130 135 140

CTT GGT GCA GAT GTT GAT TGT TTC CTT GGC ACT GAC TGC CCA CCT GTT 479
Leu Gly Ala Asp Val Asp Cys Phe Leu Gly Thr Asp Cys Pro Pro Val
145 150 155

CGT GTC AAT GGA ATC GGA GGG CTA CCT GGT GGC AAG GTC AAG CTG TCT 527
Arg Val Asn Gly Ile Gly Gly Leu Pro Gly Gly Lys Val Lys Leu Ser
160 165 170

GGC TCC ATC AGC AGT CAG TAC TTG AGT GCC TTG CTG ATG GCT GCT CCT 575
Gly Ser Ile Ser Ser Gln Tyr Leu Ser Ala Leu Leu Met Ala Ala Pro
175 180 185 190

TTG GCT CTT GGG GAT GTG GAG ATT GAA ATC ATT GAT AAA TTA ATC TCC 623
Leu Ala Leu Gly Asp Val Glu Ile Glu Ile Ile Asp Lys Leu Ile Ser
195 200 205

ATT CCG TAC GTC GAA ATG ACA TTG AGA TTG ATG GAG CGT TTT GGT GTG 671
Ile Pro Tyr Val Glu Met Thr Leu Arg Leu Met Glu Arg Phe Gly Val
210 215 220

AAA GCA GAG CAT TCT GAT AGC TGG GAC AGA TTC TAC ATT AAG GGA GGT 719
Lys Ala Glu His Ser Asp Ser Trp Asp Arg Phe Tyr Ile Lys Gly Gly
225 230 235

CAA AAA TAC AAG TCC CCT AAA AAT GCC TAT GTT GAA GGT GAT GCC TCA 767
Gln Lys Tyr Lys Ser Pro Lys Asn Ala Tyr Val Glu Gly Asp Ala Ser
240 245 250

AGC GCA AGC TAT TTC TTG GCT GGT GCT GCA ATT ACT GGA GGG ACT GTG 815
Ser Ala Ser Tyr Phe Leu Ala Gly Ala Ala Ile Thr Gly Gly Thr Val
255 260 265 270

ACT GTG GAA GGT TGT GGC ACC ACC AGT TTG CAG GGT GAT GTG AAG TTT 863
Thr Val Glu Gly Cys Gly Thr Thr Ser Leu Gln Gly Asp Val Lys Phe
275 280 285

GCT GAG GTA CTG GAG ATG ATG GGA GCG AAG GTT ACA TGG ACC GAG ACT 911
Ala Glu Val Leu Glu Met Met Gly Ala Lys Val Thr Trp Thr Glu Thr
290 295 300

AGC GTA ACT GTT ACT GGC CCA CCG CGG GAG CCA TTT GGG AGG AAA CAC 959
Ser Val Thr Val Thr Gly Pro Pro Arg Glu Pro Phe Gly Arg Lys His
305 310 315

CTC AAG GCG ATT GAT GTC AAC ATG AAC AAG ATG CCT GAT GTC GCC ATG 1007
Leu Lys Ala Ile Asp Val Asn Met Asn Lys Met Pro Asp Val Ala Met
320 325 330

ACT CTT GCT GTG GTT GCC CTC TTT GCC GAT GGC CCG ACA GCC ATC AGA 1055
Thr Leu Ala Val Val Ala Leu Phe Ala Asp Gly Pro Thr Ala Ile Arg
335 340 345 350

GAC GTG GCT TCC TGG AGA GTA AAG GAG ACC GAG AGG ATG GTT GCG ATC 1103
Asp Val Ala Ser Trp Arg Val Lys Glu Thr Glu Arg Met Val Ala Ile
355 360 365

CGG ACG GAG CTA ACC AAG CTG GGA GCA TCT GTT GAG GAA GGG CCG GAC 1151
Arg Thr Glu Leu Thr Lys Leu Gly Ala Ser Val Glu Glu Gly Pro Asp
370 375 380

TAC TGC ATC ATC ACG CCG CCG GAG AAG CTG AAC GTG ACG GCG ATC GAC 1199
Tyr Cys Ile Ile Thr Pro Pro Glu Lys Leu Asn Val Thr Ala Ile Asp
385 390 395

```
ACT TAC GAC GAC CAC AAG ATG GCG ATG GCC TTC TCC CTT GCC GCC TGT      1247
Thr Tyr Asp Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys
    400                 405                 410

GCC GAG GTC CCC GTC ACC ATC CGG GAC CCT GGG TGC ACC CGG AAG ACC      1295
Ala Glu Val Pro Val Thr Ile Arg Asp Pro Gly Cys Thr Arg Lys Thr
415                 420                 425                 430

TTC CCC GAC TAC TTC GAT GTG CTG AGC ACT TTC GTC AAG AAT               1337
Phe Pro Asp Tyr Phe Asp Val Leu Ser Thr Phe Val Lys Asn
                435                 440

TAA                                                                  1340
```

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 444 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```
Ala Gly Ala Glu Glu Ile Val Leu Gln Pro Ile Lys Glu Ile Ser Gly
  1               5                  10                  15

Thr Val Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile Leu Leu
            20                  25                  30

Leu Ala Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu Leu Asn
        35                  40                  45

Ser Glu Asp Val His Tyr Met Leu Gly Ala Leu Arg Thr Leu Gly Leu
    50                  55                  60

Ser Val Glu Ala Asp Lys Ala Ala Lys Arg Ala Val Val Val Gly Cys
65                  70                  75                  80

Gly Gly Lys Phe Pro Val Glu Asp Ala Lys Glu Glu Val Gln Leu Phe
                85                  90                  95

Leu Gly Asn Ala Gly Ile Ala Met Arg Ser Leu Thr Ala Ala Val Thr
            100                 105                 110

Ala Ala Gly Gly Asn Ala Thr Tyr Val Leu Asp Gly Val Pro Arg Met
        115                 120                 125

Arg Glu Arg Pro Ile Gly Asp Leu Val Val Gly Leu Lys Gln Leu Gly
    130                 135                 140

Ala Asp Val Asp Cys Phe Leu Gly Thr Asp Cys Pro Pro Val Arg Val
145                 150                 155                 160

Asn Gly Ile Gly Gly Leu Pro Gly Gly Lys Val Lys Leu Ser Gly Ser
                165                 170                 175

Ile Ser Ser Gln Tyr Leu Ser Ala Leu Leu Met Ala Ala Pro Leu Ala
            180                 185                 190

Leu Gly Asp Val Glu Ile Glu Ile Ile Asp Lys Leu Ile Ser Ile Pro
        195                 200                 205

Tyr Val Glu Met Thr Leu Arg Leu Met Glu Arg Phe Gly Val Lys Ala
    210                 215                 220

Glu His Ser Asp Ser Trp Asp Arg Phe Tyr Ile Lys Gly Gly Gln Lys
225                 230                 235                 240

Tyr Lys Ser Pro Lys Asn Ala Tyr Val Glu Gly Asp Ala Ser Ser Ala
                245                 250                 255

Ser Tyr Phe Leu Ala Gly Ala Ala Ile Thr Gly Gly Thr Val Thr Val
            260                 265                 270

Glu Gly Cys Gly Thr Thr Ser Leu Gln Gly Asp Val Lys Phe Ala Glu
        275                 280                 285
```

Val Leu Glu Met Met Gly Ala Lys Val Thr Trp Thr Glu Thr Ser Val
290 295 300

Thr Val Thr Gly Pro Pro Arg Glu Pro Phe Gly Arg Lys His Leu Lys
305 310 315 320

Ala Ile Asp Val Asn Met Asn Lys Met Pro Asp Val Ala Met Thr Leu
325 330 335

Ala Val Val Ala Leu Phe Ala Asp Gly Pro Thr Ala Ile Arg Asp Val
340 345 350

Ala Ser Trp Arg Val Lys Glu Thr Glu Arg Met Val Ala Ile Arg Thr
355 360 365

Glu Leu Thr Lys Leu Gly Ala Ser Val Glu Glu Gly Pro Asp Tyr Cys
370 375 380

Ile Ile Thr Pro Pro Glu Lys Leu Asn Val Thr Ala Ile Asp Thr Tyr
385 390 395 400

Asp Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys Ala Glu
405 410 415

Val Pro Val Thr Ile Arg Asp Pro Gly Cys Thr Arg Lys Thr Phe Pro
420 425 430

Asp Tyr Phe Asp Val Leu Ser Thr Phe Val Lys Asn
435 440

CLAIMS

1. DNA gene coding for a mutated 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), characterized in that it comprises at least one threonine 102 → isoleucine substitution.

2. DNA gene according to claim 1, characterized in that it comprises, in addition, at least a second mutation in the EPSPS, different from the first mutation.

3. DNA gene according to claim 2, characterized in that it comprises, in addition, a mutation consisting of a substitution of proline 106 by serine.

4. DNA gene according to claim 2, characterized in that it comprises, in addition, a mutation consisting of a substitution of glycine 101 by alanine.

5. DNA gene according to one of claims 1 to 4, characterized in that it is of bacterial origin.

6. DNA gene according to claim 5, characterized in that it originates from a bacterium of the genus *Salmonella typhimurium*.

7. DNA gene according to one of claims 1 to 4, characterized in that it is of plant origin.

8. DNA gene according to claim 7, characterized in that it is of maize origin.

9. Mutated EPSPS protein, characterized in that it comprises at least one substitution of

* C L M S *

threonine 102 by isoleucine.

10. Chimeric gene comprising a coding sequence as well as regulatory elements at positions 5' and 3' which are heterologous and capable of functioning in plants, characterized in that it comprises as coding sequence at least one sequence according to one of claims 1 to 8.

11. Chimeric gene according to claim 9, characterized in that it comprises a plant virus promoter.

12. Chimeric gene according to claim 10, characterized in that it comprises a plant promoter (e.g. α-tubulin, histone, introns, actin, etc.).

13. Vector for the transformation of plants, characterized in that it comprises at least one gene according to one of claims 10 to 12.

14. Plant cell, characterized in that it comprises at least one gene according to one of claims 10 to 12.

15. Plant, characterized in that it is obtained by regeneration from a cell according to claim 14.

16. Method for the production of plants with improved tolerance to a herbicide having EPSP synthase as its target, characterized in that plant cells or protoplates are transformed with a gene according to one of claims 1 to 8, and in that the transformed cells are subjected to a regeneration.

17. Method of treatment of plants with a herbicide having EPSPS as its target, characterized in that the herbicide is applied to plants according to claim 15.

18. Method according to claim 17, characterized in that glyphosate or a glyphosate precursor is applied.

PCT WORLD ORGANISATION FOR INTELLECTUAL PROPERTY
International Office
INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International patent classification[6] : C12N 15/54, 15/82, 9/10, A01H 5/00

A2

(11) International publication number: WO 97/04103

(43) International publication date: 6 February 1997 (06.02.97)

(21) International application number: PCT/FR96/01125

(22) International filing date: 18 July 1996 (18.07.96)

(30) Data relating to the priority:
95/08,979 19 July 1995 (19.07.95) FR

(71) Applicant (for all designated States except US): RHONE-POULENC AGROCHIMIE [FR/FR]; 14-20, rue Pierre-Baizet, F-69009 Lyon (FR).

(72) Inventors; and
(75) Inventors/Applicants (US only): Michel LEBRUN [FR/FR]; 224, rue de Saint-Cyr, F-69009 Lyon (FR). Alain SAILLAND [FR/FR]; 38, rue Ernest-Fabrègue, F-69009 Lyon (FR). Georges FREYSSINET [FR/FR]; 21, rue de Nervieux, F-69450 Saint-Cyr-au-Mont-D'Or (FR).

(74) Representative: François CHRETIEN; Rhône-Poulenc Agrochimie, P.O. Box 9163, F-69263 Lyon Cédex 09 (FR).

(81) Designated States: AL, AU, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO Patent (KE, LS, MW, SD, SZ, UG), Eurasian Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European Patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published
Without the International Search Report and to be re-published once the report has been received.

As printed

(54) Title: **MUTATED 5-ENOL PYRUVYLSHIKIMATE-3-PHOSPHATE SYNTHASE, GENE CODING FOR SAID PROTEIN AND TRANSFORMED PLANTS CONTAINING SAID GENE**

(54) Titre: **5-ENOL PYRUVYLSHIKIMATE-3-PHOSPHATE SYNTHASE MUTEE, GENE CODANT POUR CETTE PROTEINE ET PLANTES TRANSFORMEES CONTENANT CE GENE**

(57) **Abstract**

A mutated glyphosate resistance gene of 5-enol pyruvylshikimate-3-phosphate synthase (EPSPS) including at least one substitution of threonine 102 by isoleucine, and useful for producing glyphosate-resistant transformed plants, is disclosed.

(57) **Abrégé**

Gène de résistance au glyphosate. 1) Gène muté de résistance au glyphosate. 2) Gène EPSPS comprenant au moins une substitution de la Thréonine 102 par l'isoleucine. 3) Il est utilisable pour l'obtention de plantes transformées résistantes au glyphosate.