

METHODS OF GENERATING AN ARRAY

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Patent Application No. 62/723,972,
5 filed August 28, 2018, U.S. Provisional Patent Application No. 62/822,632, filed March 22,
2019, U.S. Provisional Patent Application No. 62/723,950, filed August 28, 2018, U.S.
Provisional Patent Application No. 62/723,957, filed August 28, 2018, U.S. Provisional Patent
Application No. 62/723,960, filed August 28, 2018, U.S. Provisional Patent Application No.
62/723,964, filed August 28, 2018, U.S. Provisional Patent Application No. 62/723,970, filed
10 August 28, 2018, U.S. Provisional Patent Application No. 62/724,483, filed August 29, 2018,
U.S. Provisional Patent Application No. 62/724,487, filed August 29, 2018, U.S. Provisional
Patent Application No. 62/724,489, filed August 29, 2018, U.S. Provisional Patent Application
No. 62/724,561, filed August 29, 2018, U.S. Provisional Patent Application No. 62/788,905,
filed January 6, 2019, U.S. Provisional Patent Application No. 62/788,867, filed January 6, 2019,
15 U.S. Provisional Patent Application No. 62/788,871, filed January 6, 2019, U.S. Provisional
Patent Application No. 62/788,897, filed January 6, 2019, U.S. Provisional Patent Application
No. 62/788,885, filed January 6, 2019, U.S. Provisional Patent Application No. 62/779,342, filed
December 13, 2018, U.S. Provisional Patent Application No. 62/822,565, filed March 22, 2019,
U.S. Provisional Patent Application No. 62/819,496, filed March 15, 2019, U.S. Provisional
20 Patent Application No. 62/819,486, filed March 15, 2019, U.S. Provisional Patent Application
No. 62/819,467, filed March 15, 2019, U.S. Provisional Patent Application No. 62/822,618, filed
March 22, 2019, U.S. Provisional Patent Application No. 62/822,592, filed March 22, 2019, U.S.
Provisional Patent Application No. 62/819,468, filed March 15, 2019, U.S. Provisional Patent
Application No. 62/822,627, filed March 22, 2019, U.S. Provisional Patent Application No.
25 62/819,448, filed March 15, 2019, U.S. Provisional Patent Application No. 62/822,649, filed
March 22, 2019, U.S. Provisional Patent Application No. 62/819,456, filed March 15, 2019, U.S.
Provisional Patent Application No. 62/819,478, filed March 15, 2019, U.S. Provisional Patent
Application No. 62/819,449, filed March 15, 2019, U.S. Provisional Patent Application No.
62/822,554, filed March 22, 2019, U.S. Provisional Patent Application No. 62/822,575, filed
30 March 22, 2019, U.S. Provisional Patent Application No. 62/822,605, filed March 22, 2019, U.S.
Provisional Patent Application No. 62/812,219, filed February 28, 2019, U.S. Provisional Patent

Application No. 62/819,458, filed March 15, 2019, U.S. Provisional Patent Application No. 62/839,223, filed April 26, 2019, U.S. Provisional Patent Application No. 62/839,320, filed April 26, 2019, U.S. Provisional Patent Application No. 62/839,346, filed April 26, 2019, U.S. Provisional Patent Application No. 62/842,463, filed May 2, 2019, U.S. Provisional Patent Application No. 62/860,993, filed June 13, 2019, U.S. Provisional Patent Application No. 62/839,526, filed April 26, 2019 and U.S. Provisional Patent Application No. 62/858,331, filed on June 7, 2019. The contents of each of these applications are incorporated herein by reference in their entireties.

BACKGROUND

Cells within a tissue of a subject have differences in cell morphology and/or function due to varied analyte levels (e.g., gene and/or protein expression) within the different cells. The specific position of a cell within a tissue (e.g., the cell's position relative to neighboring cells or the cell's position relative to the tissue microenvironment) can affect, e.g., the cell's morphology, differentiation, fate, viability, proliferation, behavior, and signaling and cross-talk with other cells in the tissue.

Spatial heterogeneity has been previously studied using techniques that only provide data for a small handful of analytes in the contact of an intact tissue or a portion of a tissue, or provide a lot of analyte data for single cells, but fail to provide information regarding the position of the single cell in a parent biological sample (e.g., tissue sample).

SUMMARY

In one example, provided herein are methods for generating an array, the methods include: providing a first array that includes a plurality of probes on a first substrate, where a probe of the plurality includes a spatial barcode; providing a second array that includes a plurality of primers on a second substrate; generating an oligonucleotide complementary to a probe on the first array; allowing the oligonucleotide to hybridize with a primer on the second array; and extending the primer using the oligonucleotide as a template, thereby generating a probe on the second array. In one aspect, the present disclosure refers to a method for generating an array, the method comprising: providing a first array comprising a plurality of probes immobilized on a first substrate, wherein a probe of the plurality comprises a spatial barcode; providing a second array comprising a plurality of primers immobilized on a second substrate; generating an intermediate oligonucleotide complementary to a probe on the first array; releasing the intermediate oligonucleotide from the probe on the first array; allowing the intermediate oligonucleotide to hybridize with a primer on the second array; and extending the primer using the intermediate oligonucleotide as a template, thereby generating a probe on the second array.

In some examples of any of the methods for generating an array provided herein, the primer on the second array includes a sequence that is identical to a portion of the sequence of the probe on the first array. In some examples, the portion of the sequence is in a functional domain of

the probe. In some examples, the portion of the sequence includes a free 5' end of the probe. In some examples of any of the methods for generating an array provided herein, the plurality of primers on the second array includes a common sequence. In some examples of any of the methods for generating an array provided herein, the probe on the first array is attached to the first substrate via its 3' end. In some examples of any of the methods for generating an array provided herein, the spatial barcode includes 6 to 25 nucleotides. In some examples of any of the methods for generating an array provided herein, the spatial barcode is unique to a particular region on the first array. In some examples, probes associated with a particular region on the first array include a common spatial barcode. In some examples of any of the methods for generating an array provided herein, the plurality of probes on the first array include a common functional domain. In some examples of any of the methods for generating an array provided herein, the methods further include, following generating the oligonucleotide, releasing the oligonucleotide from the probe on the first array. In some examples, releasing the oligonucleotide comprises de-hybridizing the oligonucleotide from the probe using heat. In some examples of any of the methods for generating an array provided herein, the methods further include, following releasing the oligonucleotide from the probe, facilitating migration of the released oligonucleotide to the second array. In some examples, facilitating migration of the released oligonucleotide includes applying an electric field to the first and second arrays, where the electric field increases diffusion of the released oligonucleotide. In some examples of any of the methods for generating an array provided herein, the density of the plurality of probes on the first array is about equal to or higher than the density of the plurality of primers on the second array. In some examples of any of the methods for generating an array provided herein, the number of the plurality of probes on the first array is about equal to or higher than the number of the plurality of primers on the second array. In some examples of any of the methods for generating an array provided herein, the methods further include releasing the oligonucleotide from the probe generated on the second array. In some examples, releasing the oligonucleotide includes de-hybridizing the oligonucleotide from the probe using heat. In some examples of any of the methods for generating an array provided herein, the first substrate includes a plurality of first features, and the probe of the plurality is attached to a feature of the plurality of the first features. In some examples, the spatial barcode is unique to the feature. In some examples of any of the

methods for generating an array provided herein, the second substrate includes a plurality of second features, and the primer on the second array is attached to a feature of the plurality of second features. In some examples, the plurality of first features and/or the plurality of second features are beads, wells, spots, or hydrogel pads.

5 In another example, the present disclosure refers to a method of generating an array, the method comprising: providing a first array comprising a plurality of probes on a first substrate, wherein a probe of the plurality comprises a spatial barcode; providing a primer comprising a sequence complementary to a portion of a probe on the first array; allowing the primer to hybridize with the probe; extending the primer using the probe as a template to generate an
10 oligonucleotide; aligning the first substrate and a second substrate; and attaching the oligonucleotide to the second substrate.

In another example, provided herein are methods of generating an array, the methods include: providing a first array that includes a plurality of probes on a first substrate, where a probe of the plurality includes a spatial barcode; providing a primer that includes a sequence complementary to a portion of a
15 probe on the first array; allowing the primer to hybridize with the probe; extending the primer using the probe as a template to generate an oligonucleotide; aligning the first substrate and a second substrate; and attaching the oligonucleotide to the second substrate. In some examples, the primer includes a linker, and attaching the oligonucleotide to the second substrate includes attaching through the linker. In some
20 examples, the second substrate includes a plurality of features, and attaching the oligonucleotide to the second substrate includes attaching to a feature of the plurality. the plurality of features are beads, wells, spots, or hydrogel pads. In some examples, the oligonucleotide is attached directly to the second substrate. In some examples of any of the methods for generating an array provided herein, the methods further include de-hybridizing the probe on the first array from the oligonucleotide after attaching the
25 oligonucleotide to the second substrate. In some examples of any of the methods for generating an array provided herein, a probe of the plurality on the first array further includes a sequence selected from the group consisting of: a capture domain, a cleavage domain, and a UMI.

In another example, the present disclosure refers to a method of generating an array, the method comprising: providing a first array comprising a plurality of probes on a first substrate, wherein a probe of the plurality comprises a spatial barcode; providing a second array
30 comprising a plurality of primers on a second substrate, wherein a primer of the plurality comprises a sequence complementary to a portion of a probe on the first array; aligning the first array and the second array, thereby allowing the primer to hybridize with the probe; extending the primer using the probe as a template to generate an oligonucleotide; and de-hybridizing the oligonucleotide from the probe, thereby generating an array.

35 In another example, provided herein are methods of generating an array, the methods include: providing a first array that includes a plurality of probes on a first substrate, where a probe of the plurality includes a spatial barcode; providing a second array that includes a plurality of primers on a second substrate, where a primer of the plurality includes a sequence complementary to a portion of a probe on the first array; aligning the first array and the second
40 array, thereby allowing the primer to hybridize with the probe; extending the primer using the probe as a template to generate an oligonucleotide; and de-hybridizing the oligonucleotide from the probe, thereby generating an array. In some examples, the primer includes a linker, and the primer is attached to the second substrate through the linker. In some examples, the primer is attached directly to the second substrate. In some examples of any of the methods

for generating an array described herein, the first substrate includes a plurality of first features, and the probe of the plurality is attached to a feature of the plurality of the first features. In some examples of any of the methods for generating an array described herein, the second substrate includes a plurality of second features, and the primer of the plurality is attached to a feature of the plurality of the second features. In some examples, de-hybridizing the oligonucleotide from the probe includes applying heat.

In one example, provided herein is a method for generating a spatial array that includes:

(a) providing a first array that includes a plurality of features immobilized on a solid substrate, where each feature includes a plurality of identical tagging probes unique to the feature of origin, where each tagging probe includes a barcode and a hybridization domain, where the barcode and the location of the feature are known; (b) providing a second array that includes a plurality of features immobilized on a solid substrate, wherein the plurality of features include a plurality of identical priming oligonucleotides; (c) aligning the first array and the second array, and generating an oligonucleotide that includes a sequence that is complementary to a tagging probe on the first array; (d) releasing the complementary oligonucleotide from the tagging probe on the first array; (e) allowing the complementary oligonucleotide from step (d) to hybridize with a priming oligonucleotide on the second array; and (f) extending the priming oligonucleotide on the second array using the complementary oligonucleotide from step (d) as a template, thereby generating a tagging probe on the second array comprising an oligonucleotide sequence comprising a barcode associated with a known feature from the first array, thereby generating a spatial array. In some examples, the method provided herein further includes (g) releasing the complementary oligonucleotide from step (d) from the second array.

In some examples, the releasing step (d) includes de-hybridizing the complementary oligonucleotide from the tagging probe on the first array using heat. In some examples, step (e) further includes applying an electric field to the first and second array, where the electric field increases diffusion of the complementary oligonucleotide. In some examples, the plurality of tagging probes on the first array further include a functional domain. In some examples, the functional domain of the plurality of tagging probes on the first array is configured to hybridize with the plurality of priming oligonucleotides on the second array. In some examples, the plurality of tagging probes from the first array include a poly-T sequence. In some examples, the density of the plurality of tagging probes on the first array is equal to or

higher than the density of the plurality of priming oligonucleotides on the second array. In some examples, the number of tagging probes on the first array is equal to or higher than the number of the priming oligonucleotides on the second array. In some examples of any of the methods described herein, the plurality of features are spots or beads.

5 All publications, patents, patent applications, and information available on the internet and mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, patent application, or item of information was specifically and individually indicated to be incorporated by reference. To the extent publications, patents, patent applications, and items of information incorporated by reference contradict the disclosure
10 contained in the specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

Where values are described in terms of ranges, it should be understood that the description includes the disclosure of all possible sub-ranges within such ranges, as well as specific numerical values that fall within such ranges irrespective of whether a specific
15 numerical value or specific sub-range is expressly stated.

The term “each,” when used in reference to a collection of items, is intended to identify an individual item in the collection but does not necessarily refer to every item in the collection, unless expressly stated otherwise, or unless the context of the usage clearly indicates otherwise.

Various examples of the features of this disclosure are described herein. However, it
20 should be understood that such examples are provided merely by way of example, and numerous variations, changes, and substitutions can occur to those skilled in the art without departing from the scope of this disclosure. It should also be understood that various alternatives to the specific examples described herein are also within the scope of this disclosure.

DESCRIPTION OF DRAWINGS

25 The following drawings illustrate certain embodiments of the features and advantages of this disclosure. These embodiments are not intended to limit the scope of the appended claims in any manner. Like reference symbols in the drawings indicate like elements.

FIG. 1 shows an exemplary spatial analysis workflow.

FIG. 2 shows an exemplary spatial analysis workflow.

30 **FIG. 3** shows an exemplary spatial analysis workflow.

FIG. 4 shows an exemplary spatial analysis workflow.

FIG. 5 shows an exemplary spatial analysis workflow.

FIG. 6 is a schematic diagram showing an example of a barcoded capture probe, as described herein.

5 **FIG. 7** is a schematic illustrating a cleavable capture probe, wherein the cleaved capture probe can enter into a non-permeabilized cell and bind to target analytes within the sample.

FIG. 8 is a schematic diagram of an exemplary multiplexed spatially-labelled feature.

FIG. 9 is a schematic diagram of an exemplary analyte capture agent.

10 **FIG. 10** is a schematic diagram depicting an exemplary interaction between a feature-immobilized capture probe **1024** and an analyte capture agent **1026**.

FIG. 11A, 11B, and 11C are schematics illustrating how streptavidin cell tags can be utilized in an array-based system to produce a spatially-barcoded cells or cellular contents.

FIG. 12 is a schematic showing the arrangement of barcoded features within an array.

FIG. 13 is a schematic illustrating a side view of a diffusion-resistant medium, e.g., a lid.

15 **FIG. 14A and 14B** are schematics illustrating expanded **FIG. 14A** and side views **FIG. 14B** of an electrophoretic transfer system configured to direct transcript analytes toward a spatially-barcoded capture probe array.

FIG. 15 is a schematic illustrating an exemplary workflow protocol utilizing an electrophoretic transfer system.

20 **FIG. 16** shows an example of a microfluidic channel structure **1600** for partitioning dissociated sample (e.g. biological particles or individual cells from a sample).

FIG. 17A shows an example of a microfluidic channel structure **1700** for delivering spatial barcode carrying beads to droplets.

25 **FIG. 17B** shows a cross-section view of another example of a microfluidic channel structure **1750** with a geometric feature for controlled partitioning.

FIG. 17C shows a workflow schematic.

FIG. 18 is a schematic depicting cell tagging using either covalent conjugation of the analyte binding moiety to the cell surface or non-covalent interactions with cell membrane elements.

30 **FIG. 19** is a schematic depicting cell tagging using either cell-penetrating peptides or delivery systems.

FIG. 20A is a workflow schematic illustrating exemplary, non-limiting, non-exhaustive steps for “pixelating” a sample, wherein the sample is cut, stamped, microdissected, or transferred by hollow-needle or microneedle, moving a small portion of the sample into an individual partition or well.

5 **FIG. 20B** is a schematic depicting multi-needle pixilation, wherein an array of needles punched through a sample on a scaffold and into nanowells containing gel beads and reagents below. Once the needle is in the nanowell, the cell(s) are ejected.

10 **FIG. 21** shows a workflow schematic illustrating exemplary, non-limiting, non-exhaustive steps for dissociating a spatially-barcoded sample for analysis via droplet or flow cell analysis methods.

FIG. 22 is a schematic depicting a master array aligned with a daughter array.

FIG. 23 shows a workflow schematic illustrating exemplary steps of generating a strand complementary to a probe on the master array and using the complementary strand to generate a probe on the daughter array.

15 **FIGS. 24A-24D** show a workflow schematic illustrating exemplary steps of generating probes on a copy-slide using probes on a master-slide as a template. **FIG. 24A** shows a priming step of hybridizing a primer to a probe on a master-slide. **FIG. 24B** shows a step of extending the primer using the probe as a template. **FIG. 24C** shows a step of attaching the extended strands to a copy-slide. **FIG. 24D** shows separation of the master slide from the copy-slide.

20 **FIGS. 25A-25D** show an example of a slide including fiducial markers.

FIG. 26 shows examples of distinct patterns within fiducial marker frames.

FIG. 27A is a schematic diagram showing an example sample handling apparatus that can be used to implement various steps and methods described herein.

25 **FIG. 27B** is a schematic diagram showing an example imaging apparatus that can be used to obtain images of biological samples, analytes, and arrays of features.

FIG. 27C is a schematic diagram of an example of a control unit of the apparatus of **FIGS. 27A** and **27B**.

DETAILED DESCRIPTION

30 I. Introduction

This disclosure describes apparatus, systems, methods, and compositions for spatial analysis of biological samples. This section in particular describes certain general terminology, analytes, sample types, and preparative steps that are referred to in later sections of the disclosure.

5 (a) Spatial Analysis

Tissues and cells can be obtained from any source. For example, tissues and cells can be obtained from single-cell or multicellular organisms (e.g., a mammal). Tissues and cells obtained from a mammal, e.g., a human, often have varied analyte levels (e.g., gene and/or protein expression) which can result in differences in cell morphology and/or function. The position of a cell within a tissue can affect, e.g., the cell's fate, behavior, morphology, and signaling and cross-talk with other cells in the tissue. Information regarding the differences in analyte levels (gene and/or protein expression) within different cells in a tissue of a mammal can also help physicians select or administer a treatment that will be effective in the single-cell or multicellular organisms (e.g., a mammal) based on the detected differences in analyte levels within different cells in the tissue. Differences in analyte levels within different cells in a tissue of a mammal can also provide information on how tissues (e.g., healthy and diseased tissues) function and/or develop. Differences in analyte levels within different cells in a tissue of a mammal can also provide information of different mechanisms of disease pathogenesis in a tissue and mechanism of action of a therapeutic treatment within a tissue. Differences in analyte levels within different cells in a tissue of a mammal can also provide information on drug resistance mechanisms and the development of the same in a tissue of a mammal. Differences in the presence or absence of analytes within different cells in a tissue of a multicellular organism (e.g., a mammal) can provide information on drug resistance mechanisms and the development of the same in a tissue of a multicellular organism.

25 The spatial analysis methodologies provide for the detection of differences in an analyte level (e.g., gene and/or protein expression) within different cells in a tissue of a mammal or within a single cell from a mammal. For example, spatial analysis methodologies can be used to detect the differences in analyte levels (e.g., gene and/or protein expression) within different cells in histological slide samples, the data from which can be reassembled to generate a three-dimensional map of analyte levels (e.g., gene and/or protein expression) of a tissue sample obtained from a mammal, e.g., with a degree of spatial resolution (e.g., single-cell resolution).

Spatial heterogeneity in developing systems has typically been studied via RNA hybridization, immunohistochemistry, fluorescent reporters, or purification or induction of pre-defined subpopulations and subsequent genomic profiling (e.g., RNA-seq). Such approaches, however, rely on a relatively small set of pre-defined markers, therefore introducing selection bias that limits discovery. These prior approaches also rely on *a priori* knowledge. Spatial RNA assays traditionally relied on staining for a limited number of RNA species. In contrast, single-cell RNA-sequencing allows for deep profiling of cellular gene expression (including non-coding RNA), but the established methods separate cells from their native spatial context.

Current spatial analysis methodologies provide a vast amount of analyte level and/or expression data for a variety of multiple analytes within a sample at high spatial resolution, e.g., while retaining the native spatial context. Spatial analysis methods include, e.g., the use of a capture probe including a spatial barcode (e.g., a nucleic acid sequence that provides information as to the position of the capture probe within a cell or a tissue sample (e.g., mammalian cell or a mammalian tissue sample) and a capture domain that is capable of binding to an analyte (e.g., a protein and/or nucleic acid) produced by and/or present in a cell. As described herein, the spatial barcode can be a nucleic acid that has a unique sequence, a unique fluorophore or a unique combination of fluorophores, a unique amino acid sequence, a unique heavy metal or a unique combination of heavy metals, or any other unique detectable agent. The capture domain can be any agent that is capable of binding to an analyte produced by and/or present in a cell (e.g., a nucleic acid that is capable of hybridizing to a nucleic acid from a cell (e.g., an mRNA, genomic DNA, mitochondrial DNA, or miRNA), a substrate or binding partner of an analyte, or an antibody that binds specifically to an analyte). A capture probe can also include a nucleic acid sequence that is complementary to a sequence of a universal forward and/or universal reverse primer. A capture probe can also include a cleavage site (e.g., a cleavage recognition site of a restriction endonuclease), a photolabile bond, a thermosensitive bond, or a chemical-sensitive bond.

The binding of an analyte to a capture probe can be detected using a number of different methods, e.g., nucleic acid sequencing, fluorophore detection, nucleic acid amplification, detection of nucleic acid ligation, and/or detection of nucleic acid cleavage products. In some examples, the detection is used to associate a specific spatial barcode with a specific analyte produced by and/or present in a cell (e.g., a mammalian cell).

Capture probes can be, e.g., attached to a surface, e.g., a solid array, a bead, or a coverslip. In some examples, capture probes are not attached to a surface. In some examples, capture probes can be encapsulated within, embedded within, or layered on a surface of a permeable composition (e.g., any of the substrates described herein). For example, capture probes can be encapsulated or disposed within a permeable bead (e.g., a gel bead). In some examples, capture probes can be encapsulated within, embedded within, or layered on a surface of a substrate (e.g., any of the exemplary substrates described herein, such as a hydrogel or a porous membrane).

In some examples, a cell or a tissue sample including a cell are contacted with capture probes attached to a substrate (e.g., a surface of a substrate), and the cell or tissue sample is permeabilized to allow analytes to be released from the cell and bind to the capture probes attached to the substrate. In some examples, analytes released from a cell can be actively directed to the capture probes attached to a substrate using a variety of methods, e.g., electrophoresis, chemical gradient, pressure gradient, fluid flow, or magnetic field.

In other examples, a capture probe can be directed to interact with a cell or a tissue sample using a variety of methods, e.g., inclusion of a lipid anchoring agent in the capture probe, inclusion of an agent that binds specifically to, or forms a covalent bond with a membrane protein in the capture probe, fluid flow, pressure gradient, chemical gradient, or magnetic field.

Non-limiting aspects of spatial analysis methodologies are described in WO 2011/127099, WO 2014/210233, WO 2014/210225, WO 2016/162309, WO 2018/091676, WO 2012/140224, WO 2014/060483, U.S. Patent No. 10,002,316, U.S. Patent No. 9,727,810, U.S. Patent Application Publication No. 2017/0016053, Rodriques et al., *Science* 363(6434):1463-1467, 2019; WO 2018/045186, Lee et al., *Nat. Protoc.* 10(3):442-458, 2015; WO 2016/007839, WO 2018/045181, WO 2014/163886, Trejo et al., *PLoS ONE* 14(2):e0212031, 2019, U.S. Patent Application Publication No. 2018/0245142, Chen et al., *Science* 348(6233):aaa6090, 2015, Gao et al., *BMC Biol.* 15:50, 2017, WO 2017/144338, WO 2018/107054, WO 2017/222453, WO 2019/068880, WO 2011/094669, U.S. Patent No. 7,709,198, U.S. Patent No. 8,604,182, U.S. Patent No. 8,951,726, U.S. Patent No. 9,783,841, U.S. Patent No. 10,041,949, WO 2016/057552, WO 2017/147483, WO 2018/022809, WO 2016/166128, WO 2017/027367, WO 2017/027368, WO 2018/136856, WO 2019/075091, U.S. Patent No. 10,059,990, WO 2018/057999, WO 2015/161173, and Gupta et al., *Nature*

Biotechnol. 36:1197-1202, 2018, and can be used herein in any combination. Further non-limiting aspects of spatial analysis methodologies are described herein.

(b) General Terminology

Specific terminology is used throughout this disclosure to explain various aspects of the apparatus, systems, methods, and compositions that are described. This sub-section includes explanations of certain terms that appear in later sections of the disclosure. To the extent that the descriptions in this section are in apparent conflict with usage in other sections of this disclosure, the definitions in this section will control.

(i) Barcode

A “barcode” is a label, or identifier, that conveys or is capable of conveying information (e.g., information about an analyte in a sample, a bead, and/or a capture probe). A barcode can be part of an analyte, or independent of an analyte. A barcode can be attached to an analyte. A particular barcode can be unique relative to other barcodes.

Barcodes can have a variety of different formats. For example, barcodes can include polynucleotide barcodes, random nucleic acid and/or amino acid sequences, and synthetic nucleic acid and/or amino acid sequences. A barcode can be attached to an analyte or to another moiety or structure in a reversible or irreversible manner. A barcode can be added to, for example, a fragment of a deoxyribonucleic acid (DNA) or ribonucleic acid (RNA) sample before or during sequencing of the sample. Barcodes can allow for identification and/or quantification of individual sequencing-reads (e.g., a barcode can be or can include a unique molecular identifier or “UMI”).

Barcodes can spatially-resolve molecular components found in biological samples, for example, at single-cell resolution (e.g., a barcode can be or can include a “spatial barcode”). In some embodiments, a barcode includes both a UMI and a spatial barcode. In some embodiments, a barcode includes two or more sub-barcodes that together function as a single barcode. For example, a polynucleotide barcode can include two or more polynucleotide sequences (e.g., sub-barcodes) that are separated by one or more non-barcode sequences.

(ii) Nucleic Acid and Nucleotide

The terms “nucleic acid” and “nucleotide” are intended to be consistent with their use in the art and to include naturally-occurring species or functional analogs thereof. Particularly useful functional analogs of nucleic acids are capable of hybridizing to a nucleic acid in a

sequence-specific fashion (e.g., capable of hybridizing to two nucleic acids such that ligation can occur between the two hybridized nucleic acids) or are capable of being used as a template for replication of a particular nucleotide sequence. Naturally-occurring nucleic acids generally have a backbone containing phosphodiester bonds. An analog structure can have an alternate
5 backbone linkage including any of a variety of those known in the art. Naturally-occurring nucleic acids generally have a deoxyribose sugar (e.g., found in deoxyribonucleic acid (DNA)) or a ribose sugar (e.g. found in ribonucleic acid (RNA)).

A nucleic acid can contain nucleotides having any of a variety of analogs of these sugar moieties that are known in the art. A nucleic acid can include native or non-native nucleotides. In
10 this regard, a native deoxyribonucleic acid can have one or more bases selected from the group consisting of adenine (A), thymine (T), cytosine (C), or guanine (G), and a ribonucleic acid can have one or more bases selected from the group consisting of uracil (U), adenine (A), cytosine (C), or guanine (G). Useful non-native bases that can be included in a nucleic acid or nucleotide are known in the art.

15 (iii) Probe and Target

A “probe” or a “target,” when used in reference to a nucleic acid or sequence of a nucleic acids, is intended as a semantic identifier for the nucleic acid or sequence in the context of a method or composition, and does not limit the structure or function of the nucleic acid or sequence beyond what is expressly indicated.

20 (iv) Oligonucleotide and Polynucleotide

The terms “oligonucleotide” and “polynucleotide” are used interchangeably to refer to a single-stranded multimer of nucleotides from about 2 to about 500 nucleotides in length. Oligonucleotides can be synthetic, made enzymatically (e.g., via polymerization), or using a “split-pool” method. Oligonucleotides can include ribonucleotide monomers (i.e., can be
25 oligoribonucleotides) and/or deoxyribonucleotide monomers (i.e., oligodeoxyribonucleotides). In some examples, oligonucleotides can include a combination of both deoxyribonucleotide monomers and ribonucleotide monomers in the oligonucleotide (e.g., random or ordered combination of deoxyribonucleotide monomers and ribonucleotide monomers). An oligonucleotide can be 4 to 10, 10 to 20, 21 to 30, 31 to 40, 41 to 50, 51 to 60, 61 to 70, 71 to 80,
30 80 to 100, 100 to 150, 150 to 200, 200 to 250, 250 to 300, 300 to 350, 350 to 400, or 400-500 nucleotides in length, for example. Oligonucleotides can include one or more functional moieties

that are attached (e.g., covalently or non-covalently) to the multimer structure. For example, an oligonucleotide can include one or more detectable labels (e.g., a radioisotope or fluorophore).

(v) Subject

A “subject” is an animal, such as a mammal (e.g., human or a non-human simian), or
 5 avian (e.g., bird), or other organism, such as a plant. Examples of subjects include, but are not limited to, a mammal such as a rodent, mouse, rat, rabbit, guinea pig, ungulate, horse, sheep, pig, goat, cow, cat, dog, primate (i.e. human or non-human primate); a plant such as *Arabidopsis thaliana*, corn, sorghum, oat, wheat, rice, canola, or soybean; an algae such as *Chlamydomonas reinhardtii*; a nematode such as *Caenorhabditis elegans*; an insect such as *Drosophila*
 10 *melanogaster*, mosquito, fruit fly, or honey bee; an arachnid such as a spider; a fish such as zebrafish; a reptile; an amphibian such as a frog or *Xenopus laevis*; a *Dictyostelium discoideum*; a fungi such as *Pneumocystis carinii*, *Takifugu rubripes*, yeast, *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe*; or a *Plasmodium falciparum*.

(vi) Genome

15 A “genome” generally refers to genomic information from a subject, which can be, for example, at least a portion of, or the entirety of, the subject’s gene-encoded hereditary information. A genome can include coding regions (e.g., that code for proteins) as well as non-coding regions. A genome can include the sequences of some or all of the subject’s chromosomes. For example, the human genome ordinarily has a total of 46 chromosomes. The
 20 sequences of some or all of these can constitute the genome.

(vii) Adaptor, Adapter, and Tag

An “adaptor,” an “adapter,” and a “tag” are terms that are used interchangeably in this disclosure, and refer to species that can be coupled to a polynucleotide sequence (in a process referred to as “tagging”) using any one of many different techniques including (but not limited
 25 to) ligation, hybridization, and tagmentation. Adaptors can also be nucleic acid sequences that add a function, e.g., spacer sequences, primer sequences/sites, barcode sequences, unique molecular identifier sequences.

(viii) Hybridizing, Hybridize, Annealing, and Anneal

The terms “hybridizing,” “hybridize,” “annealing,” and “anneal” are used
 30 interchangeably in this disclosure, and refer to the pairing of substantially complementary or complementary nucleic acid sequences within two different molecules. Pairing can be achieved

by any process in which a nucleic acid sequence joins with a substantially or fully complementary sequence through base pairing to form a hybridization complex. For purposes of hybridization, two nucleic acid sequences are “substantially complementary” if at least 60% (e.g., at least 70%, at least 80%, or at least 90%) of their individual bases are complementary to one another.

(ix) Primer

A “primer” is a single-stranded nucleic acid sequence having a 3’ end that can be used as a substrate for a nucleic acid polymerase in a nucleic acid extension reaction. RNA primers are formed of RNA nucleotides, and are used in RNA synthesis, while DNA primers are formed of DNA nucleotides and used in DNA synthesis. Primers can also include both RNA nucleotides and DNA nucleotides (e.g., in a random or designed pattern). Primers can also include other natural or synthetic nucleotides described herein that can have additional functionality. In some examples, DNA primers can be used to prime RNA synthesis and vice versa (e.g., RNA primers can be used to prime DNA synthesis). Primers can vary in length. For example, primers can be about 6 bases to about 120 bases. For example, primers can include up to about 25 bases.

(x) Primer Extension

A “primer extension” refers to any method where two nucleic acid sequences (e.g., a constant region from each of two distinct capture probes) become linked (e.g., hybridized) by an overlap of their respective terminal complementary nucleic acid sequences (i.e., for example, 3’ termini). Such linking can be followed by nucleic acid extension (e.g., an enzymatic extension) of one, or both termini using the other nucleic acid sequence as a template for extension. Enzymatic extension can be performed by an enzyme including, but not limited to, a polymerase and/or a reverse transcriptase.

(xi) Proximity Ligation

A “proximity ligation” is a method of ligating two (or more) nucleic acid sequences that are in proximity with each other through enzymatic means (e.g., a ligase). In some embodiments, proximity ligation can include a “gap-filling” step that involves incorporation of one or more nucleic acids by a polymerase, based on the nucleic acid sequence of a template nucleic acid molecule, spanning a distance between the two nucleic acid molecules of interest (see, e.g., U.S. Patent No. 7,264,929, the entire contents of which are incorporated herein by reference).

A wide variety of different methods can be used for proximity ligating nucleic acid molecules, including (but not limited to) “sticky-end” and “blunt-end” ligations. Additionally, single-stranded ligation can be used to perform proximity ligation on a single-stranded nucleic acid molecule. Sticky-end proximity ligations involve the hybridization of complementary single-stranded sequences between the two nucleic acid molecules to be joined, prior to the ligation event itself. Blunt-end proximity ligations generally do not include hybridization of complementary regions from each nucleic acid molecule because both nucleic acid molecules lack a single-stranded overhang at the site of ligation.

(xii) Nucleic Acid Extension

A “nucleic acid extension” generally involves incorporation of one or more nucleic acids (e.g., A, G, C, T, U, nucleotide analogs, or derivatives thereof) into a molecule (such as, but not limited to, a nucleic acid sequence) in a template-dependent manner, such that consecutive nucleic acids are incorporated by an enzyme (such as a polymerase or reverse transcriptase), thereby generating a newly synthesized nucleic acid molecule. For example, a primer that hybridizes to a complementary nucleic acid sequence can be used to synthesize a new nucleic acid molecule by using the complementary nucleic acid sequence as a template for nucleic acid synthesis. Similarly, a 3’ polyadenylated tail of an mRNA transcript that hybridizes to a poly (dT) sequence (e.g., capture domain) can be used as a template for single-strand synthesis of a corresponding cDNA molecule.

(xiii) PCR Amplification

A “PCR amplification” refers to the use of a polymerase chain reaction (PCR) to generate copies of genetic material, including DNA and RNA sequences. Suitable reagents and conditions for implementing PCR are described, for example, in U.S. Patents 4,683,202, 4,683,195, 4,800,159, 4,965,188, and 5,512,462, the entire contents of each of which are incorporated herein by reference. In a typical PCR amplification, the reaction mixture includes the genetic material to be amplified, an enzyme, one or more primers that are employed in a primer extension reaction, and reagents for the reaction. The oligonucleotide primers are of sufficient length to provide for hybridization to complementary genetic material under annealing conditions. The length of the primers generally depends on the length of the amplification domains, but will typically be at least 4 bases, at least 5 bases, at least 6 bases, at least 8 bases, at least 9 bases, at least 10 base pairs (bp), at least 11 bp, at least 12 bp, at least 13 bp, at least 14 bp, at least 15 bp, at least 16 bp,

at least 17 bp, at least 18 bp, at least 19 bp, at least 20 bp, at least 25 bp, at least 30 bp, at least 35 bp, and can be as long as 40 bp or longer, where the length of the primers will generally range from 18 to 50 bp. The genetic material can be contacted with a single primer or a set of two primers (forward and reverse primers), depending upon whether primer extension, linear or
5 exponential amplification of the genetic material is desired.

In some embodiments, the PCR amplification process uses a DNA polymerase enzyme. The DNA polymerase activity can be provided by one or more distinct DNA polymerase enzymes. In certain embodiments, the DNA polymerase enzyme is from a bacterium, e.g., the DNA polymerase enzyme is a bacterial DNA polymerase enzyme. For instance, the DNA
10 polymerase can be from a bacterium of the genus *Escherichia*, *Bacillus*, *Thermophilus*, or *Pyrococcus*.

Suitable examples of DNA polymerases that can be used include, but are not limited to: *E.coli* DNA polymerase I, Bsu DNA polymerase, Bst DNA polymerase, Taq DNA polymerase, VENT™ DNA polymerase, DEEPVENT™ DNA polymerase, LongAmp® Taq DNA
15 polymerase, LongAmp® Hot Start Taq DNA polymerase, Crimson LongAmp® Taq DNA polymerase, Crimson Taq DNA polymerase, OneTaq® DNA polymerase, OneTaq® Quick-Load® DNA polymerase, Hemo KlenTaq® DNA polymerase, REDTaq® DNA polymerase, Phusion® DNA polymerase, Phusion® High-Fidelity DNA polymerase, Platinum Pfx DNA polymerase, AccuPrime Pfx DNA polymerase, Phi29 DNA polymerase, Klenow fragment, Pwo
20 DNA polymerase, Pfu DNA polymerase, T4 DNA polymerase and T7 DNA polymerase enzymes.

The term “DNA polymerase” includes not only naturally-occurring enzymes but also all modified derivatives thereof, including also derivatives of naturally-occurring DNA polymerase enzymes. For instance, in some embodiments, the DNA polymerase can have been modified to
25 remove 5’-3’ exonuclease activity. Sequence-modified derivatives or mutants of DNA polymerase enzymes that can be used include, but are not limited to, mutants that retain at least some of the functional, e.g. DNA polymerase activity of the wild-type sequence. Mutations can affect the activity profile of the enzymes, e.g. enhance or reduce the rate of polymerization, under different reaction conditions, e.g. temperature, template concentration, primer
30 concentration, etc. Mutations or sequence-modifications can also affect the exonuclease activity and/or thermostability of the enzyme.

In some embodiments, PCR amplification can include reactions such as, but not limited to, a strand-displacement amplification reaction, a rolling circle amplification reaction, a ligase chain reaction, a transcription-mediated amplification reaction, an isothermal amplification reaction, and/or a loop-mediated amplification reaction.

5 In some embodiments, PCR amplification uses a single primer that is complementary to the 3' tag of target DNA fragments. In some embodiments, PCR amplification uses a first and a second primer, where at least a 3' end portion of the first primer is complementary to at least a portion of the 3' tag of the target nucleic acid fragments, and where at least a 3' end portion of the second primer exhibits the sequence of at least a portion of the 5' tag of the target nucleic
10 acid fragments. In some embodiments, a 5' end portion of the first primer is non-complementary to the 3' tag of the target nucleic acid fragments, and a 5' end portion of the second primer does not exhibit the sequence of at least a portion of the 5' tag of the target nucleic acid fragments. In some embodiments, the first primer includes a first universal sequence and/or the second primer includes a second universal sequence.

15 In some embodiments (e.g., when the PCR amplification amplifies captured DNA), the PCR amplification products can be ligated to additional sequences using a DNA ligase enzyme. The DNA ligase activity can be provided by one or more distinct DNA ligase enzymes. In some embodiments, the DNA ligase enzyme is from a bacterium, e.g., the DNA ligase enzyme is a bacterial DNA ligase enzyme. In some embodiments, the DNA ligase enzyme is from a virus
20 (e.g., a bacteriophage). For instance, the DNA ligase can be T4 DNA ligase. Other enzymes appropriate for the ligation step include, but are not limited to, Tth DNA ligase, Taq DNA ligase, *Thermococcus sp.* (strain 9oN) DNA ligase (9oNTM DNA ligase, available from New England Biolabs, Ipswich, MA), and Ampligase™ (available from Epicentre Biotechnologies, Madison, WI). Derivatives, e.g. sequence-modified derivatives, and/or mutants thereof, can also be used.

25 In some embodiments, genetic material is amplified by reverse transcription polymerase chain reaction (RT-PCR). The desired reverse transcriptase activity can be provided by one or more distinct reverse transcriptase enzymes, suitable examples of which include, but are not limited to: M-MLV, MuLV, AMV, HIV, ArrayScript™, MultiScribe™, ThermoScript™, and SuperScript® I, II, III, and IV enzymes. "Reverse transcriptase" includes not only naturally
30 occurring enzymes, but all such modified derivatives thereof, including also derivatives of naturally-occurring reverse transcriptase enzymes.

In addition, reverse transcription can be performed using sequence-modified derivatives or mutants of M-MLV, MuLV, AMV, and HIV reverse transcriptase enzymes, including mutants that retain at least some of the functional, e.g. reverse transcriptase, activity of the wild-type sequence. The reverse transcriptase enzyme can be provided as part of a composition that
5 includes other components, e.g. stabilizing components that enhance or improve the activity of the reverse transcriptase enzyme, such as RNase inhibitor(s), inhibitors of DNA-dependent DNA synthesis, e.g. actinomycin D. Many sequence-modified derivative or mutants of reverse transcriptase enzymes, e.g. M-MLV, and compositions including unmodified and modified enzymes are commercially available, e.g. ArrayScript™, MultiScribe™, ThermoScript™, and
10 SuperScript® I, II, III, and IV enzymes.

Certain reverse transcriptase enzymes (e.g. Avian Myeloblastosis Virus (AMV) Reverse Transcriptase and Moloney Murine Leukemia Virus (M-MuLV, MMLV) Reverse Transcriptase) can synthesize a complementary DNA strand using both RNA (cDNA synthesis) and single-stranded DNA (ssDNA) as a template. Thus, in some embodiments, the reverse transcription
15 reaction can use an enzyme (reverse transcriptase) that is capable of using both RNA and ssDNA as the template for an extension reaction, e.g. an AMV or MMLV reverse transcriptase.

In some embodiments, the quantification of RNA and/or DNA is carried out by real-time PCR (also known as quantitative PCR or qPCR), using techniques well known in the art, such as but not limited to “TAQMAN™” or “SYBR®”, or on capillaries (“LightCycler® Capillaries”).

20 In some embodiments, the quantification of genetic material is determined by optical absorbance and with real-time PCR. In some embodiments, the quantification of genetic material is determined by digital PCR. In some embodiments, the genes analyzed can be compared to a reference nucleic acid extract (DNA and RNA) corresponding to the expression (mRNA) and quantity (DNA) in order to compare expression levels of the target nucleic acids.

25 (xiv) Antibody

An “antibody” is a polypeptide molecule that recognizes and binds to a complementary target antigen. Antibodies typically have a molecular structure shape that resembles a Y shape. Naturally-occurring antibodies, referred to as immunoglobulins, belong to one of the immunoglobulin classes IgG, IgM, IgA, IgD, and IgE. Antibodies can also be produced
30 synthetically. For example, recombinant antibodies, which are monoclonal antibodies, can be synthesized using synthetic genes by recovering the antibody genes from source cells,

amplifying into an appropriate vector, and introducing the vector into a host to cause the host to express the recombinant antibody. In general, recombinant antibodies can be cloned from any species of antibody-producing animal using suitable oligonucleotide primers and/or hybridization probes. Recombinant techniques can be used to generate antibodies and antibody fragments, including non-endogenous species.

Synthetic antibodies can be derived from non-immunoglobulin sources. For example, antibodies can be generated from nucleic acids (e.g., aptamers), and from non-immunoglobulin protein scaffolds (such as peptide aptamers) into which hypervariable loops are inserted to form antigen binding sites. Synthetic antibodies based on nucleic acids or peptide structures can be smaller than immunoglobulin-derived antibodies, leading to greater tissue penetration.

Antibodies can also include affimer proteins, which are affinity reagents that typically have a molecular weight of about 12-14 kDa. Affimer proteins generally bind to a target (e.g., a target protein) with both high affinity and specificity. Examples of such targets include, but are not limited to, ubiquitin chains, immunoglobulins, and C-reactive protein. In some embodiments, affimer proteins are derived from cysteine protease inhibitors, and include peptide loops and a variable N-terminal sequence that provides the binding site.

Antibodies can also include single domain antibodies (V_HH domains and VNAR domains), scFvs, and Fab fragments.

(xv) Affinity Group

An “affinity group” is a molecule or molecular moiety which has a high affinity or preference for associating or binding with another specific or particular molecule or moiety. The association or binding with another specific or particular molecule or moiety can be via a non-covalent interaction, such as hydrogen bonding, ionic forces, and van der Waals interactions. An affinity group can, for example, be biotin, which has a high affinity or preference to associate or bind to the protein avidin or streptavidin. An affinity group, for example, can also refer to avidin or streptavidin which has an affinity to biotin. Other examples of an affinity group and specific or particular molecule or moiety to which it binds or associates with include, but are not limited to, antibodies or antibody fragments and their respective antigens, such as digoxigenin and anti-digoxigenin antibodies, lectin, and carbohydrates (e.g., a sugar, a monosaccharide, a disaccharide, or a polysaccharide), and receptors and receptor ligands.

Any pair of affinity group and its specific or particular molecule or moiety to which it binds or associates with can have their roles reversed, for example, such that between a first molecule and a second molecule, in a first instance the first molecule is characterized as an affinity group for the second molecule, and in a second instance the second molecule is characterized as an affinity group for the first molecule.

(xvi) Label, Detectable Label, and Optical Label

The terms “detectable label,” “optical label,” and “label” are used interchangeably herein to refer to a directly or indirectly detectable moiety that is associated with (e.g., conjugated to) a molecule to be detected, e.g., a capture probe or analyte. The detectable label can be directly detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, can be indirectly detectable, e.g., by catalyzing chemical alterations of a substrate compound or composition, which substrate compound or composition is directly detectable. Detectable labels can be suitable for small scale detection and/or suitable for high-throughput screening. As such, suitable detectable labels include, but are not limited to, radioisotopes, fluorophores, chemiluminescent compounds, bioluminescent compounds, and dyes.

The detectable label can be qualitatively detected (e.g., optically or spectrally), or it can be quantified. Qualitative detection generally includes a detection method in which the existence or presence of the detectable label is confirmed, whereas quantifiable detection generally includes a detection method having a quantifiable (e.g., numerically reportable) value such as an intensity, duration, polarization, and/or other properties. In some embodiments, the detectable label is bound to a feature or to a capture probe associated with a feature. For example, detectably labeled features can include a fluorescent, a colorimetric, or a chemiluminescent label attached to a bead (see, for example, Rajeswari et al., *J. Microbiol Methods* 139:22-28, 2017, and Forcucci et al., *J. Biomed Opt.* 10:105010, 2015, the entire contents of each of which are incorporated herein by reference).

In some embodiments, a plurality of detectable labels can be attached to a feature, capture probe, or composition to be detected. For example, detectable labels can be incorporated during nucleic acid polymerization or amplification (e.g., Cy5®-labelled nucleotides, such as Cy5®-dCTP). Any suitable detectable label can be used. In some embodiments, the detectable label is a fluorophore. For example, the fluorophore can be from a group that includes: 7-AAD (7-Aminoactinomycin D), Acridine Orange (+DNA), Acridine Orange (+RNA), Alexa Fluor® 350,

Alexa Fluor® 430, Alexa Fluor® 488, Alexa Fluor® 532, Alexa Fluor® 546, Alexa Fluor® 555,
 Alexa Fluor® 568, Alexa Fluor® 594, Alexa Fluor® 633, Alexa Fluor® 647, Alexa Fluor® 660,
 Alexa Fluor® 680, Alexa Fluor® 700, Alexa Fluor® 750, Allophycocyanin (APC), AMCA /
 AMCA-X, 7-Aminoactinomycin D (7-AAD), 7- Amino-4-methylcoumarin, 6-Aminoquinoline,
 5 Aniline Blue, ANS, APC-Cy7, ATTO-TAG™ CBQCA, ATTO-TAG™ FQ, Auramine O-
 Feulgen, BCECF (high pH), BFP (Blue Fluorescent Protein), BFP / GFP FRET, BOBO™-1 /
 BO-PRO™-1, BOBO™-3 / BO-PRO™-3, BODIPY® FL, BODIPY® TMR, BODIPY® TR-X,
 BODIPY® 530/550, BODIPY® 558/568, BODIPY® 564/570, BODIPY® 581/591, BODIPY®
 630/650-X, BODIPY® 650-665-X, BTC, Calcein, Calcein Blue, Calcium Crimson™, Calcium
 10 Green-1™, Calcium Orange™, Calcofluor® White, 5-Carboxyfluorescein (5-FAM), 5-
 Carboxynaphthofluorescein, 6-Carboxyrhodamine 6G, 5-Carboxytetramethylrhodamine (5-
 TAMRA), Carboxy-X-rhodamine (5-ROX), Cascade Blue®, Cascade Yellow™, CCF2
 (GeneBLAzer™), CFP (Cyan Fluorescent Protein), CFP / YFP FRET, Chromomycin A3, Cl-
 NERF (low pH), CPM, 6-CR 6G, CTC Formazan, Cy2®, Cy3®, Cy3.5®, Cy5®, Cy5.5®,
 15 Cy7®, Cychrome (PE-Cy5), Dansylamine, Dansyl cadaverine, Dansylchloride, DAPI, Dapoxyl,
 DCFH, DHR, DiA (4-Di-16-ASP), DiD (DiI C18(5)), DIDS, DiI (DiI C18(3)), DiO (DiO C18(3)),
 DiR (DiI C18(7)), Di-4 ANEPPS, Di-8 ANEPPS, DM-NERF (4.5-6.5 pH), DsRed (Red
 Fluorescent Protein), EBFP, ECFP, EGFP, ELF® -97 alcohol, Eosin, Erythrosin, Ethidium
 bromide, Ethidium homodimer-1 (EthD-1), Europium (III) Chloride, 5-FAM (5-
 20 Carboxyfluorescein), Fast Blue, Fluorescein-dT phosphoramidite, FITC, Fluo-3, Fluo-4,
 FluorX®, Fluoro-Gold™ (high pH), Fluoro-Gold™ (low pH), Fluoro-Jade, FM® 1-43, Fura-2
 (high calcium), Fura-2 / BCECF, Fura Red™ (high calcium), Fura Red™ / Fluo-3,
 GeneBLAzer™ (CCF2), GFP Red Shifted (rsGFP), GFP Wild Type, GFP / BFP FRET, GFP /
 DsRed FRET, Hoechst 33342 & 33258, 7-Hydroxy-4-methylcoumarin (pH 9), 1,5 IAEDANS,
 25 Indo-1 (high calcium), Indo-1 (low calcium), Indodicarbocyanine, Indotricarbocyanine, JC-1, 6-
 JOE, JOJO™-1 / JO-PRO™-1, LDS 751 (+DNA), LDS 751 (+RNA), LOLO™-1 / LO-PRO™-
 1, Lucifer Yellow, LysoSensor™ Blue (pH 5), LysoSensor™ Green (pH 5), LysoSensor™
 Yellow/Blue (pH 4.2), LysoTracker® Green, LysoTracker® Red, LysoTracker® Yellow, Mag-
 Fura-2, Mag-Indo-1, Magnesium Green™, Marina Blue®, 4-Methylumbelliferone, Mithramycin,
 30 MitoTracker® Green, MitoTracker® Orange, MitoTracker® Red, NBD (amine), Nile Red,
 Oregon Green® 488, Oregon Green® 500, Oregon Green® 514, Pacific Blue, PBF1, PE (R-

phycoerythrin), PE-Cy5, PE-Cy7, PE-Texas Red, PerCP (Peridinin chlorophyll protein), PerCP-Cy5.5 (TruRed), PharRed (APC-Cy7), C-phycoerythrin, R-phycoerythrin (PE), PI (Propidium Iodide), PKH26, PKH67, POPO™-1 / PO-PRO™-1, POPO™-3 / PO-PRO™-3, Propidium Iodide (PI), PyMPO, Pyrene, Pyronin Y, Quantam Red (PE-Cy5), Quinacrine
 5 Mustard, R670 (PE-Cy5), Red 613 (PE-Texas Red), Red Fluorescent Protein (DsRed), Resorufin, RH 414, Rhod-2, Rhodamine B, Rhodamine Green™, Rhodamine Red™, Rhodamine Phalloidin, Rhodamine 110, Rhodamine 123, 5-ROX (carboxy-X-rhodamine), S65A, S65C, S65L, S65T, SBFI, SITS, SNAFL®-1 (high pH), SNAFL®-2, SNARF®-1 (high pH), SNARF®-1 (low pH), Sodium Green™, SpectrumAqua®, SpectrumGreen® #1, SpectrumGreen® #2,
 10 SpectrumOrange®, SpectrumRed®, SYTO® 11, SYTO® 13, SYTO® 17, SYTO® 45, SYTOX® Blue, SYTOX® Green, SYTOX® Orange, 5-TAMRA (5-Carboxytetramethylrhodamine), Tetramethylrhodamine (TRITC), Texas Red® / Texas Red®-X, Texas Red®-X (NHS Ester), Thiadicarbocyanine, Thiazole Orange, TOTO®-1 / TO-PRO®-1, TOTO®-3 / TO-PRO®-3, TO-PRO®-5, Tri-color (PE-Cy5), TRITC (Tetramethylrhodamine),
 15 TruRed (PerCP-Cy5.5), WW 781, X-Rhodamine (XRITC), Y66F, Y66H, Y66W, YFP (Yellow Fluorescent Protein), YOYO®-1 / YO-PRO®-1, YOYO®-3 / YO-PRO®-3, 6-FAM (Fluorescein), 6-FAM (NHS Ester), 6-FAM (Azide), HEX, TAMRA (NHS Ester), Yakima Yellow, MAX, TET, TEX615, ATTO 488, ATTO 532, ATTO 550, ATTO 565, ATTO Rho101, ATTO 590, ATTO 633, ATTO 647N, TYE 563, TYE 665, TYE 705, 5' IRDye® 700, 5'
 20 IRDye® 800, 5' IRDye® 800CW (NHS Ester), WellRED D4 Dye, WellRED D3 Dye, WellRED D2 Dye, Lightcycler® 640 (NHS Ester), and Dy 750 (NHS Ester).

As mentioned above, in some embodiments, a detectable label is or includes a luminescent or chemiluminescent moiety. Common luminescent/chemiluminescent moieties include, but are not limited to, peroxidases such as horseradish peroxidase (HRP), soybean
 25 peroxidase (SP), alkaline phosphatase, and luciferase. These protein moieties can catalyze chemiluminescent reactions given the appropriate substrates (e.g., an oxidizing reagent plus a chemiluminescent compound. A number of compound families are known to provide chemiluminescence under a variety of conditions. Non-limiting examples of chemiluminescent compound families include 2,3-dihydro-1,4-phthalazinedione luminol, 5-amino-6,7,8-trimethoxy-
 30 and the dimethylamino[ca]benz analog. These compounds can luminesce in the presence of alkaline hydrogen peroxide or calcium hypochlorite and base. Other examples of

chemiluminescent compound families include, e.g., 2,4,5-triphenylimidazoles, para-dimethylamino and - methoxy substituents, oxalates such as oxalyl active esters, p-nitrophenyl, N-alkyl acridinium esters, luciferins, lucigenins, or acridinium esters.

(xvii) Template Switching Oligonucleotide

5 A “template switching oligonucleotide” is an oligonucleotide that hybridizes to untemplated nucleotides added by a reverse transcriptase (e.g., enzyme with terminal transferase activity) during reverse transcription. In some embodiments, a template switching oligonucleotide hybridizes to untemplated poly(C) nucleotides added by a reverse transcriptase. In some embodiments, the template switching oligonucleotide adds a common 5' sequence to
10 full-length cDNA that is used for cDNA amplification.

In some embodiments, the template switching oligonucleotide adds a common sequence onto the 5' end of the RNA being reverse transcribed. For example, a template switching oligonucleotide can hybridize to untemplated poly(C) nucleotides added onto the end of a cDNA molecule and provide a template for the reverse transcriptase to continue replication to the 5' end
15 of the template switching oligonucleotide, thereby generating full-length cDNA ready for further amplification. In some embodiments, once a full-length cDNA molecule is generated, the template switching oligonucleotide can serve as a primer in a cDNA amplification reaction.

In some embodiments, a template switching oligonucleotide is added before, contemporaneously with, or after a reverse transcription, or other terminal transferase-based
20 reaction. In some embodiments, a template switching oligonucleotide is included in the capture probe. In certain embodiments, methods of sample analysis using template switching oligonucleotides can involve the generation of nucleic acid products from analytes of the tissue sample, followed by further processing of the nucleic acid products with the template switching oligonucleotide.

25 Template switching oligonucleotides can include a hybridization region and a template region. The hybridization region can include any sequence capable of hybridizing to the target. In some embodiments, the hybridization region can, e.g., include a series of G bases to complement the overhanging C bases at the 3' end of a cDNA molecule. The series of G bases can include 1 G base, 2 G bases, 3 G bases, 4 G bases, 5 G bases, or more than 5 G bases. The
30 template sequence can include any sequence to be incorporated into the cDNA. In other embodiments, the hybridization region can include at least one base in addition to at least one G

base. In other embodiments, the hybridization can include bases that are not a G base. In some embodiments, the template region includes at least 1 (e.g., at least 2, 3, 4, 5 or more) tag sequences and/or functional sequences. In some embodiments, the template region and hybridization region are separated by a spacer.

5 In some embodiments, the template regions include a barcode sequence. The barcode sequence can act as a spatial barcode and/or as a unique molecular identifier. Template switching oligonucleotides can include deoxyribonucleic acids; ribonucleic acids; modified nucleic acids including 2-aminopurine, 2,6-diaminopurine (2-amino-dA), inverted dT, 5-methyl dC, 2'-deoxyInosine, Super T (5-hydroxybutynl-2'-deoxyuridine), Super G (8-aza-7-deazaguanosine),
 10 locked nucleic acids (LNAs), unlocked nucleic acids (UNAs, e.g., UNA-A, UNA-U, UNA-C, UNA-G), Iso-dG, Iso-dC, 2' fluoro bases (e.g., Fluoro C, Fluoro U, Fluoro A, and Fluoro G), or any combination of the foregoing.

In some embodiments, the length of a template switching oligonucleotide can be at least about 1, 2, 10, 20, 50, 75, 100, 150, 200, or 250 nucleotides or longer. In some embodiments, the
 15 length of a template switching oligonucleotide can be at most about 2, 10, 20, 50, 100, 150, 200, or 250 nucleotides or longer.

(xviii) Splint Oligonucleotide

A “splint oligonucleotide” is an oligonucleotide that, when hybridized to other polynucleotides, acts as a “splint” to position the polynucleotides next to one another so that they
 20 can be ligated together. In some embodiments, the splint oligonucleotide is DNA or RNA. The splint oligonucleotide can include a nucleotide sequence that is partially complimentary to nucleotide sequences from two or more different oligonucleotides. In some embodiments, the splint oligonucleotide assists in ligating a “donor” oligonucleotide and an “acceptor” oligonucleotide. In general, an RNA ligase, a DNA ligase, or another other variety of ligase is
 25 used to ligate two nucleotide sequences together

In some embodiments, the splint oligonucleotide is between 10 and 50 oligonucleotides in length, e.g., between 10 and 45, 10 and 40, 10 and 35, 10 and 30, 10 and 25, or 10 and 20 oligonucleotides in length. In some embodiments, the splint oligonucleotide is between 15 and 50, 15 and 45, 15 and 40, 15 and 35, 15 and 30, 15 and 30, or 15 and 25 nucleotides in length.

(c) Analytes

The apparatus, systems, methods, and compositions described in this disclosure can be used to detect and analyze a wide variety of different analytes. For the purpose of this disclosure, an “analyte” can include any biological substance, structure, moiety, or component to be
5 analyzed. The term “target” can similarly refer to an analyte of interest.

Analytes can be broadly classified into one of two groups: nucleic acid analytes, and non-nucleic acid analytes. Examples of non-nucleic acid analytes include, but are not limited to, lipids, carbohydrates, peptides, proteins, glycoproteins (N-linked or O-linked), lipoproteins, phosphoproteins, specific phosphorylated or acetylated variants of proteins, amidation variants
10 of proteins, hydroxylation variants of proteins, methylation variants of proteins, ubiquitylation variants of proteins, sulfation variants of proteins, viral coat proteins, extracellular and intracellular proteins, antibodies, and antigen binding fragments. In some embodiments, the analyte can be an organelle (e.g., nuclei or mitochondria).

Cell surface features corresponding to analytes can include, but are not limited to, a
15 receptor, an antigen, a surface protein, a transmembrane protein, a cluster of differentiation protein, a protein channel, a protein pump, a carrier protein, a phospholipid, a glycoprotein, a glycolipid, a cell-cell interaction protein complex, an antigen-presenting complex, a major histocompatibility complex, an engineered T-cell receptor, a T-cell receptor, a B-cell receptor, a chimeric antigen receptor, an extracellular matrix protein, a posttranslational modification (e.g.,
20 phosphorylation, glycosylation, ubiquitination, nitrosylation, methylation, acetylation or lipidation) state of a cell surface protein, a gap junction, and an adherens junction.

Analytes can be derived from a specific type of cell and/or a specific sub-cellular region. For example, analytes can be derived from cytosol, from cell nuclei, from mitochondria, from microsomes, and more generally, from any other compartment, organelle, or portion of a cell.
25 Permeabilizing agents that specifically target certain cell compartments and organelles can be used to selectively release analytes from cells for analysis.

Examples of nucleic acid analytes include DNA analytes such as genomic DNA, methylated DNA, specific methylated DNA sequences, fragmented DNA, mitochondrial DNA, in situ synthesized PCR products, and RNA/DNA hybrids.

30 Examples of nucleic acid analytes also include RNA analytes such as various types of coding and non-coding RNA. Examples of the different types of RNA analytes include

messenger RNA (mRNA), ribosomal RNA (rRNA), transfer RNA (tRNA), microRNA (miRNA), and viral RNA. The RNA can be a transcript (e.g., present in a tissue section). The RNA can be small (e.g., less than 200 nucleic acid bases in length) or large (e.g., RNA greater than 200 nucleic acid bases in length). Small RNAs mainly include 5.8S ribosomal RNA (rRNA), 5S rRNA, transfer RNA (tRNA), microRNA (miRNA), small interfering RNA (siRNA), small nucleolar RNA (snoRNAs), Piwi-interacting RNA (piRNA), tRNA-derived small RNA (tsRNA), and small rDNA-derived RNA (srRNA). The RNA can be double-stranded RNA or single-stranded RNA. The RNA can be circular RNA. The RNA can be a bacterial rRNA (e.g., 16S rRNA or 23S rRNA).

Additional examples of analytes include mRNA and cell surface features (e.g., using the labelling agents described herein), mRNA and intracellular proteins (e.g., transcription factors), mRNA and cell methylation status, mRNA and accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), mRNA and metabolites (e.g., using the labelling agents described herein), a barcoded labelling agent (e.g., the oligonucleotide tagged antibodies described herein) and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor), mRNA and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein).

Analytes can include a nucleic acid molecule with a nucleic acid sequence encoding at least a portion of a V(D)J sequence of an immune cell receptor (e.g., a TCR or BCR). In some embodiments, the nucleic acid molecule is cDNA first generated from reverse transcription of the corresponding mRNA, using a poly(T) containing primer. The generated cDNA can then be barcoded using a capture probe, featuring a barcode sequence (and optionally, a UMI sequence) that hybridizes with at least a portion of the generated cDNA. In some embodiments, a template switching oligonucleotide hybridizes to a poly(C) tail added to a 3' end of the cDNA by a reverse transcriptase enzyme. The original mRNA template and template switching oligonucleotide can then be denatured from the cDNA and the barcoded capture probe can then hybridize with the cDNA and a complement of the cDNA generated. Additional methods and compositions suitable for barcoding cDNA generated from mRNA transcripts including those encoding V(D)J regions of an immune cell receptor and/or barcoding methods and composition including a template switch oligonucleotide are described in PCT Patent Application PCT/US2017/057269, filed October 18, 2017, and U.S. Patent Application Serial No. 15/825,740, filed November 29, 2017,

both of which are incorporated herein by reference in their entireties. V(D)J analysis can also be completed with the use of one or more labelling agents that bind to particular surface features of immune cells and associated with barcode sequences. The one or more labelling agents can include an MHC or MHC multimer.

As described above, the analyte can include a nucleic acid capable of functioning as a component of a gene editing reaction, such as, for example, clustered regularly interspaced short palindromic repeats (CRISPR)-based gene editing. Accordingly, the capture probe can include a nucleic acid sequence that is complementary to the analyte (e.g., a sequence that can hybridize to the CRISPR RNA (crRNA), single guide RNA (sgRNA), or an adapter sequence engineered into a crRNA or sgRNA).

In certain embodiments, an analyte can be extracted from a live cell. Processing conditions can be adjusted to ensure that a biological sample remains live during analysis, and analytes are extracted from (or released from) live cells of the sample. Live cell-derived analytes can be obtained only once from the sample, or can be obtained at intervals from a sample that continues to remain in viable condition.

In general, the systems, apparatus, methods, and compositions can be used to analyze any number of analytes. For example, the number of analytes that are analyzed can be at least about 2, at least about 3, at least about 4, at least about 5, at least about 6, at least about 7, at least about 8, at least about 9, at least about 10, at least about 11, at least about 12, at least about 13, at least about 14, at least about 15, at least about 20, at least about 25, at least about 30, at least about 40, at least about 50, at least about 100, at least about 1,000, at least about 10,000, at least about 100,000 or more different analytes present in a region of the sample or within an individual feature of the substrate. Methods for performing multiplexed assays to analyze two or more different analytes will be discussed in a subsequent section of this disclosure.

(d) Biological Samples

(i) Types of Biological Samples

A “biological sample” is obtained from the subject for analysis using any of a variety of techniques including, but not limited to, biopsy, surgery, and laser capture microscopy (LCM), and generally includes cells and/or other biological material from the subject. In addition to the subjects described above, a biological sample can also be obtained from a prokaryote such as a bacterium, e.g., *Escherichia coli*, *Staphylococci* or *Mycoplasma pneumoniae*; an archaea; a virus

such as Hepatitis C virus or human immunodeficiency virus; or a viroid. A biological sample can be obtained from non-mammalian organisms (e.g., a plants, an insect, an archnid, a nematode, a fungi, or an amphibian). A biological sample can also be obtained from a eukaryote, such as a patient derived organoid (PDO) or patient derived xenograft (PDX). Subjects from which
5 biological samples can be obtained can be healthy or asymptomatic individuals, individuals that have or are suspected of having a disease (e.g., a patient with a disease such as cancer) or a pre-disposition to a disease, and/or individuals that are in need of therapy or suspected of needing therapy.

The biological sample can include any number of macromolecules, for example, cellular
10 macromolecules and organelles (e.g., mitochondria and nuclei). The biological sample can be a nucleic acid sample and/or protein sample. The biological sample can be a carbohydrate sample or a lipid sample. The biological sample can be obtained as a tissue sample, such as a tissue section, biopsy, a core biopsy, needle aspirate, or fine needle aspirate. The sample can be a fluid sample, such as a blood sample, urine sample, or saliva sample. The sample can be a skin
15 sample, a colon sample, a cheek swab, a histology sample, a histopathology sample, a plasma or serum sample, a tumor sample, living cells, cultured cells, a clinical sample such as, for example, whole blood or blood-derived products, blood cells, or cultured tissues or cells, including cell suspensions.

Cell-free biological samples can include extracellular polynucleotides. Extracellular
20 polynucleotides can be isolated from a bodily sample, e.g., blood, plasma, serum, urine, saliva, mucosal excretions, sputum, stool, and tears.

Biological samples can be derived from a homogeneous culture or population of the subjects or organisms mentioned herein or alternatively from a collection of several different organisms, for example, in a community or ecosystem.

25 Biological samples can include one or more diseased cells. A diseased cell can have altered metabolic properties, gene expression, protein expression, and/or morphologic features. Examples of diseases include inflammatory disorders, metabolic disorders, nervous system disorders, and cancer. Cancer cells can be derived from solid tumors, hematological malignancies, cell lines, or obtained as circulating tumor cells.

30 Biological samples can also include fetal cells. For example, a procedure such as amniocentesis can be performed to obtain a fetal cell sample from maternal circulation.

Sequencing of fetal cells can be used to identify any of a number of genetic disorders, including, e.g., aneuploidy such as Down's syndrome, Edwards syndrome, and Patau syndrome. Further, cell surface features of fetal cells can be used to identify any of a number of disorders or diseases.

5 Biological samples can also include immune cells. Sequence analysis of the immune repertoire of such cells, including genomic, proteomic, and cell surface features, can provide a wealth of information to facilitate an understanding the status and function of the immune system. By way of example, determining the status (e.g., negative or positive) of minimal residue disease (MRD) in a multiple myeloma (MM) patient following autologous stem cell
10 transplantation is considered a predictor of MRD in the MM patient (see, e.g., U.S. Patent Application Publication No. 2018/0156784, the entire contents of which are incorporated herein by reference).

 Examples of immune cells in a biological sample include, but are not limited to, B cells, T cells (e.g., cytotoxic T cells, natural killer T cells, regulatory T cells, and T helper cells),
15 natural killer cells, cytokine induced killer (CIK) cells, myeloid cells, such as granulocytes (basophil granulocytes, eosinophil granulocytes, neutrophil granulocytes/hypersegmented neutrophils), monocytes/macrophages, mast cells, thrombocytes/megakaryocytes, and dendritic cells.

 As discussed above, a biological sample can include a single analyte of interest, or more
20 than one analyte of interest. Methods for performing multiplexed assays to analyze two or more different analytes in a single biological sample will be discussed in a subsequent section of this disclosure.

(ii) Preparation of Biological Samples

 A variety of steps can be performed to prepare a biological sample for analysis. Except
25 where indicated otherwise, the preparative steps described below can generally be combined in any manner to appropriately prepare a particular sample for analysis.

(1) Tissue Sectioning

 A biological sample can be harvested from a subject (e.g., via surgical biopsy, whole
subject sectioning) or grown in vitro on a growth substrate or culture dish as a population of
30 cells, and prepared for analysis as a tissue slice or tissue section. Grown samples may be sufficiently thin for analysis without further processing steps. Alternatively, grown samples, and

samples obtained via biopsy or sectioning, can be prepared as thin tissue sections using a mechanical cutting apparatus such as a vibrating blade microtome. As another alternative, in some embodiments, a thin tissue section can be prepared by applying a touch imprint of a biological sample to a suitable substrate material.

5 The thickness of the tissue section can be a fraction of (e.g., less than 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2, or 0.1) the maximum cross-sectional dimension of a cell. However, tissue sections having a thickness that is larger than the maximum cross-section cell dimension can also be used. For example, cryostat sections can be used, which can be, e.g., 10-20 micrometers thick.

10 More generally, the thickness of a tissue section typically depends on the method used to prepare the section and the physical characteristics of the tissue, and therefore sections having a wide variety of different thicknesses can be prepared and used. For example, the thickness of the tissue section can be at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.7, 1.0, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 20, 30, 40, or 50 micrometers. Thicker sections can also be used if desired or convenient, e.g., at least 70, 80, 90, or 100 micrometers or more. Typically, the thickness of a tissue section
15 is between 1-100 micrometers, 1-50 micrometers, 1-30 micrometers, 1-25 micrometers, 1-20 micrometers, 1-15 micrometers, 1-10 micrometers, 2-8 micrometers, 3-7 micrometers, or 4-6 micrometers, but as mentioned above, sections with thicknesses larger or smaller than these ranges can also be analysed.

20 Multiple sections can also be obtained from a single biological sample. For example, multiple tissue sections can be obtained from a surgical biopsy sample by performing serial sectioning of the biopsy sample using a sectioning blade. Spatial information among the serial sections can be preserved in this manner, and the sections can be analysed successively to obtain three-dimensional information about the biological sample.

(2) Freezing

25 In some embodiments, the biological sample (e.g., a tissue section as described above) can be prepared by deep freezing at a temperature suitable to maintain or preserve the integrity (e.g., the physical characteristics) of the tissue structure. Such a temperature can be, e.g., less than -20°C, or less than -25°C, -30°C, -40°C, -50°C, -60°C, -70°C, -80°C -90°C, -100°C, -110°C, -120°C, -130°C, -140°C, -150°C, -160°C, -170°C, -180°C, -190°C, or -200°C. The
30 frozen tissue sample can be sectioned, e.g., thinly sliced, onto a substrate surface using any number of suitable methods. For example, a tissue sample can be prepared using a chilled

microtome (e.g., a cryostat) set at a temperature suitable to maintain both the structural integrity of the tissue sample and the chemical properties of the nucleic acids in the sample. Such a temperature can be, e.g., less than -15°C, less than -20°C, or less than -25°C.

(3) Formalin Fixation and Paraffin Embedding

5 In some embodiments, the biological sample can be prepared using formalin-fixation and paraffin-embedding (FFPE), which are established methods. In some embodiments, cell suspensions and other non-tissue samples can be prepared using formalin-fixation and paraffin-embedding. Following fixation of the sample and embedding in a paraffin or resin block, the sample can be sectioned as described above. Prior to analysis, the paraffin-embedding material
10 can be removed from the tissue section (e.g., deparaffinization) by incubating the tissue section in an appropriate solvent (e.g., xylene) followed by a rinse (e.g., 99.5% ethanol for 2 minutes, 96% ethanol for 2 minutes, and 70% ethanol for 2 minutes).

(4) Fixation

As an alternative to formalin fixation described above, a biological sample can be fixed
15 in any of a variety of other fixatives to preserve the biological structure of the sample prior to analysis. For example, a sample can be fixed via immersion in ethanol, methanol, acetone, paraformaldehyde-Triton, and combinations thereof.

In some embodiments, acetone fixation is used with fresh frozen samples, which can include, but are not limited to, cortex tissue, mouse olfactory bulb, human brain tumor, human
20 post-mortem brain, and breast cancer samples. When acetone fixation is performed, pre-permeabilization steps (described below) may not be performed. Alternatively, acetone fixation can be performed in conjunction with permeabilization steps.

(5) Embedding

As an alternative to paraffin embedding described above, a biological sample can be
25 embedded in any of a variety of other embedding materials to provide structural substrate to the sample prior to sectioning and other handling steps. In general, the embedding material is removed prior to analysis of tissue sections obtained from the sample. Suitable embedding materials include, but are not limited to, waxes, resins (e.g., methacrylate resins), epoxies, and agar.

30

(6) Staining

To facilitate visualization, biological samples can be stained using a wide variety of stains and staining techniques. In some embodiments, for example, a sample can be stained using any number of stains, including but not limited to, acridine orange, Bismarck brown, carmine, 5 coomassie blue, cresyl violet, DAPI, eosin, ethidium bromide, acid fuchsin, haematoxylin, Hoechst stains, iodine, methyl green, methylene blue, neutral red, Nile blue, Nile red, osmium tetroxide, propidium iodide, rhodamine, or safranin.

The sample can be stained using hematoxylin and eosin (H&E) staining techniques, using Papanicolaou staining techniques, Masson's trichrome staining techniques, silver staining 10 techniques, Sudan staining techniques, and/or using Periodic Acid Schiff (PAS) staining techniques. PAS staining is typically performed after formalin or acetone fixation. In some embodiments, the sample can be stained using Romanowsky stain, including Wright's stain, Jenner's stain, Can-Grunwald stain, Leishman stain, and Giemsa stain.

In some embodiments, biological samples can be destained. Methods of destaining or 15 discolored a biological sample are known in the art, and generally depend on the nature of the stain(s) applied to the sample. For example, in some embodiments, one or more immunofluorescent stains are applied to the sample via antibody coupling. Such stains can be removed using techniques such as cleavage of disulfide linkages via treatment with a reducing agent and detergent washing, chaotropic salt treatment, treatment with antigen retrieval solution, 20 and treatment with an acidic glycine buffer. Methods for multiplexed staining and destaining are described, for example, in Bolognesi et al., *J. Histochem. Cytochem.* 2017; 65(8): 431-444, Lin et al., *Nat Commun.* 2015; 6:8390, Pirici et al., *J. Histochem. Cytochem.* 2009; 57:567-75, and Glass et al., *J. Histochem. Cytochem.* 2009; 57:899-905, the entire contents of each of which are incorporated herein by reference.

(7) Hydrogel Embedding

In some embodiments, the biological sample can be embedded in a hydrogel matrix. Embedding the sample in this manner typically involves contacting the biological sample with a hydrogel such that the biological sample becomes surrounded by the hydrogel. For example, the sample can be embedded by contacting the sample with a suitable polymer material, and 30 activating the polymer material to form a hydrogel. In some embodiments, the hydrogel is formed such that the hydrogel is internalized within the biological sample.

In some embodiments, the biological sample is immobilized in the hydrogel via cross-linking of the polymer material that forms the hydrogel. Cross-linking can be performed chemically and/or photochemically, or alternatively by any other hydrogel-formation method known in the art.

5 The composition and application of the hydrogel-matrix to a biological sample typically depends on the nature and preparation of the biological sample (e.g., sectioned, non-sectioned, type of fixation). As one example, where the biological sample is a tissue section, the hydrogel-matrix can include a monomer solution and an ammonium persulfate (APS) initiator/tetramethylethylenediamine (TEMED) accelerator solution. As another example, where
10 the biological sample consists of cells (e.g., cultured cells or cells disassociated from a tissue sample), the cells can be incubated with the monomer solution and APS/TEMED solutions. For cells, hydrogel-matrix gels are formed in compartments, including but not limited to devices used to culture, maintain, or transport the cells. For example, hydrogel-matrices can be formed with monomer solution plus APS/TEMED added to the compartment to a depth ranging from
15 about 0.1 μm to about 2 mm.

Additional methods and aspects of hydrogel embedding of biological samples are described for example in Chen et al., *Science* 347(6221):543–548, 2015, the entire contents of which are incorporated herein by reference.

(8) Isometric Expansion

20 In some embodiments, a biological sample embedded in a hydrogel can be isometrically expanded. Isometric expansion methods that can be used include hydration, a preparative step in expansion microscopy, as described in Chen et al., *Science* 347(6221):543–548, 2015.

Isometric expansion can be performed by anchoring one or more components of a biological sample to a gel, followed by gel formation, proteolysis, and swelling. Isometric
25 expansion of the biological sample can occur prior to immobilization of the biological sample on a substrate, or after the biological sample is immobilized to a substrate. In some embodiments, the isometrically expanded biological sample can be removed from the substrate prior to contacting the substrate with capture probes, as will be discussed in greater detail in a subsequent section.

30 In general, the steps used to perform isometric expansion of the biological sample can depend on the characteristics of the sample (e.g., thickness of tissue section, fixation, cross-

linking), and/or the analyte of interest (e.g., different conditions to anchor RNA, DNA, and protein to a gel).

In some embodiments, proteins in the biological sample are anchored to a swellable gel such as a polyelectrolyte gel. An antibody can be directed to the protein before, after, or in conjunction with being anchored to the swellable gel. DNA and/or RNA in a biological sample can also be anchored to the swellable gel via a suitable linker. Examples of such linkers include, but are not limited to, 6-((Acryloyl)amino) hexanoic acid (Acryloyl-X SE) (available from ThermoFisher, Waltham, MA), Label-IT Amine (available from MirusBio, Madison, WI) and Label X (described for example in Chen et al., *Nat. Methods* 13:679-684, 2016, the entire contents of which are incorporated herein by reference).

Isometric expansion of the sample can increase the spatial resolution of the subsequent analysis of the sample. The increased resolution in spatial profiling can be determined by comparison of an isometrically expanded sample with a sample that has not been isometrically expanded.

In some embodiments, a biological sample is isometrically expanded to a size at least 2x, 2.1x, 2.2x, 2.3x, 2.4x, 2.5x, 2.6x, 2.7x, 2.8x, 2.9x, 3x, 3.1x, 3.2x, 3.3x, 3.4x, 3.5x, 3.6x, 3.7x, 3.8x, 3.9x, 4x, 4.1x, 4.2x, 4.3x, 4.4x, 4.5x, 4.6x, 4.7x, 4.8x, or 4.9x its non-expanded size. In some embodiments, the sample is isometrically expanded to at least 2x and less than 20x of its non-expanded size.

(9) Substrate Attachment

In some embodiments, the biological sample can be attached to a substrate. Examples of substrates suitable for this purpose are described in detail below. Attachment of the biological sample can be irreversible or reversible, depending upon the nature of the sample and subsequent steps in the analytical method.

In certain embodiments, the sample can be attached to the substrate reversibly by applying a suitable polymer coating to the substrate, and contacting the sample to the polymer coating. The sample can then be detached from the substrate using an organic solvent that at least partially dissolves the polymer coating. Hydrogels are examples of polymers that are suitable for this purpose.

More generally, in some embodiments, the substrate can be coated or functionalized with one or more substances to facilitate attachment of the sample to the substrate. Suitable

substances that can be used to coat or functionalize the substrate include, but are not limited to, lectins, poly-lysine, antibodies, and polysaccharides.

(10) Disaggregation of Cells

In some embodiments, the biological sample corresponds to cells (e.g., derived from a cell culture or a tissue sample). In a cell sample with a plurality of cells, individual cells can be naturally unaggregated. For example, the cells can be derived from a suspension of cells and/or disassociated or disaggregated cells from a tissue or tissue section.

Alternatively, the cells in the sample may be aggregated, and may be disaggregated into individual cells using, for example, enzymatic or mechanical techniques. Examples of enzymes used in enzymatic disaggregation include, but are not limited to, dispase, collagenase, trypsin, and combinations thereof. Mechanical disaggregation can be performed, for example, using a tissue homogenizer.

(11) Suspended and Adherent Cells

In some embodiments, the biological sample can be derived from a cell culture grown in vitro. Samples derived from a cell culture can include one or more suspension cells which are anchorage-independent within the cell culture. Examples of such cells include, but are not limited to, cell lines derived from hematopoietic cells, and from the following cell lines: Colo205, CCRF-CEM, HL-60, K562, MOLT-4, RPMI-8226, SR, HOP-92, NCI-H322M, and MALME-3M.

Samples derived from a cell culture can include one or more adherent cells which grow on the surface of the vessel that contains the culture medium. Non-limiting examples of adherent cells include DU145 (prostate cancer) cells, H295R (adrenocortical cancer) cells, HeLa (cervical cancer) cells, KBM-7 (chronic myelogenous leukemia) cells, LNCaP (prostate cancer) cells, MCF-7 (breast cancer) cells, MDA-MB-468 (breast cancer) cells, PC3 (prostate cancer) cells, SaOS-2 (bone cancer) cells, SH-SY5Y (neuroblastoma, cloned from a myeloma) cells, T-47D (breast cancer) cells, THP-1 (acute myeloid leukemia) cells, U87 (glioblastoma) cells, National Cancer Institute's 60 cancer cell line panel (NCI60), vero (African green monkey *Chlorocebus* kidney epithelial cell line) cells, MC3T3 (embryonic calvarium) cells, GH3 (pituitary tumor) cells, PC12 (pheochromocytoma) cells, dog MDCK kidney epithelial cells, *Xenopus* A6 kidney epithelial cells, zebrafish AB9 cells, and Sf9 insect epithelial cells.

Additional examples of adherent cells are shown in Table 1 and catalogued, for example, in “A Catalog of in Vitro Cell Lines, Transplantable Animal and Human Tumors and Yeast,” The Division of Cancer Treatment and Diagnosis (DCTD), National Cancer Institute (2013), and in Abaan et al., “The exomes of the NCI-60 panel: a genomic resource for cancer biology and systems pharmacology,” *Cancer Research* 73(14):4372-82, 2013, the entire contents of each of which are incorporated by reference herein.

Table 1: Examples of adherent cells

Cell Line	Species	Organ of Origin	Disease
BT549	Human	Breast	Ductal Carcinoma
HS 578T	Human	Breast	Carcinoma
MCF7	Human	Breast	Adenocarcinoma
MDA-MB-231	Human	Breast	Adenocarcinoma
MDA-MB-468	Human	Breast	Adenocarcinoma
T-47D	Human	Breast	Ductal Carcinoma
SF268	Human	CNS	Anaplastic Astrocytoma
SF295	Human	CNS	Glioblastoma-Multiforme
SF539	Human	CNS	Glioblastoma
SNB-19	Human	CNS	Glioblastoma
SNB-75	Human	CNS	Astrocytoma
U251	Human	CNS	Glioblastoma
Colo205	Human	Colon	Dukes' type D, Colorectal adenocarcinoma
HCC 2998	Human	Colon	Carcinoma
HCT-116	Human	Colon	Carcinoma
HCT-15	Human	Colon	Dukes' type C, Colorectal adenocarcinoma
HT29	Human	Colon	Colorectal adenocarcinoma

KM12	Human	Colon	Adenocarcinoma, Grade III
SW620	Human	Colon	Adenocarcinoma
786-O	Human	Kidney	renal cell adenocarcinoma
A498	Human	Kidney	Adenocarcinoma
ACHN	Human	Kidney	renal cell adenocarcinoma
CAKI	Human	Kidney	clear cell carcinoma
RXF 393	Human	Kidney	Poorly Differentiated Hypernephroma
SN12C	Human	Kidney	Carcinoma
TK-10	Human	Kidney	Spindle Cell carcinoma
UO-31	Human	Kidney	Carcinoma
A549	Human	Lung	Adenocarcinoma
EKVX	Human	Lung	Adenocarcinoma
HOP-62	Human	Lung	Adenocarcinoma
HOP-92	Human	Lung	Large Cell, Undifferentiated
NCI-H226	Human	Lung	squamous cell carcinoma; mesothelioma
NCI-H23	Human	Lung	adenocarcinoma; non-small cell lung cancer
NCI-H460	Human	Lung	carcinoma; large cell lung cancer
NCI-H522	Human	Lung	adenocarcinoma; non-small cell lung cancer
LOX IMVI	Human	Melanoma	Malignant Amelanotic melanoma
M14	Human	Melanoma	malignant melanoma
MALME-3M	Human	Melanoma	malignant melanoma
MDA-MB-435	Human	Melanoma	Adenocarcinoma
SK-MEL-2	Human	Melanoma	malignant melanoma

SK-MEL-28	Human	Melanoma	malignant melanoma
SK-MEL-5	Human	Melanoma	malignant melanoma
UACC-257	Human	Melanoma	malignant melanoma
UACC-62	Human	Melanoma	malignant melanoma
IGROV1	Human	Ovary	Cystadenocarcinoma
OVCAR-3	Human	Ovary	Adenocarcinoma
OVCAR-4	Human	Ovary	Adenocarcinoma
OVCAR-5	Human	Ovary	Adenocarcinoma
OVCAR-8	Human	Ovary	Adenocarcinoma
SK-OV-3	Human	Ovary	Adenocarcinoma
NCI-ADR-RES	Human	Ovary	Adenocarcinoma
DU145	Human	Prostate	Carcinoma
PC-3	Human	Prostate	grade IV, adenocarcinoma

In some embodiments, the adherent cells are cells that correspond to one or more of the following cell lines: BT549, HS 578T, MCF7, MDA-MB-231, MDA-MB-468, T-47D, SF268, SF295, SF539, SNB-19, SNB-75, U251, Colo205, HCC 2998, HCT-116, HCT-15, HT29, KM12, SW620, 786-O, A498, ACHN, CAKI, RXF 393, SN12C, TK-10, UO-31, A549, EKVX, HOP-62, HOP-92, NCI-H226, NCI-H23, NCI-H460, NCI-H522, LOX IMVI, M14, MALME-3M, MDA-MB-435, SK-, EL-2, SK-MEL-28, SK-MEL-5, UACC-257, UACC-62, IGROV1, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, SK-OV-3, NCI-ADR-RES, DU145, PC-3, DU145, H295R, HeLa, KBM-7, LNCaP, MCF-7, MDA-MB-468, PC3, SaOS-2, SH-SY5Y, T-47D, THP-1, U87, vero, MC3T3, GH3, PC12, dog MDCK kidney epithelial, Xenopus A6 kidney epithelial, zebrafish AB9, and Sf9 insect epithelial cell lines.

(12) Tissue Permeabilization

In some embodiments, a biological sample can be permeabilized to facilitate transfer of analytes out of the sample, and/or to facilitate transfer of species (such as capture probes) into the sample. If a sample is not permeabilized sufficiently, the amount of analyte captured from the sample may be too low to enable adequate analysis. Conversely, if the tissue sample is too permeable, the relative spatial relationship of the analytes within the tissue sample can be lost.

Hence, a balance between permeabilizing the tissue sample enough to obtain good signal intensity while still maintaining the spatial resolution of the analyte distribution in the sample is desirable.

In general, a biological sample can be permeabilized by exposing the sample to one or
5 more permeabilizing agents. Suitable agents for this purpose include, but are not limited to, organic solvents (e.g., acetone, ethanol, and methanol), cross-linking agents (e.g., paraformaldehyde), detergents (e.g., saponin, Triton X-100™ or Tween-20™), and enzymes (e.g., trypsin, proteases). In some embodiments, the biological sample can be incubated with a cellular permeabilizing agent to facilitate permeabilization of the sample. Additional methods for
10 sample permeabilization are described, for example, in Jamur et al., *Method Mol. Biol.* 588:63-66, 2010, the entire contents of which are incorporated herein by reference. Any suitable method for sample permeabilization can generally be used in connection with the samples described herein.

In some embodiments, where a diffusion-resistant medium is used to limit migration of
15 analytes or other species during the analytical procedure, the diffusion-resistant medium can include at least one permeabilization reagent. For example, the diffusion-resistant medium can include wells (e.g., micro-, nano-, or picowells) containing a permeabilization buffer or reagents. In some embodiments, where the diffusion-resistant medium is a hydrogel, the hydrogel can include a permeabilization buffer. In some embodiments, the hydrogel is soaked in
20 permeabilization buffer prior to contacting the hydrogel with a sample. In some embodiments, the hydrogel or other diffusion-resistant medium can contain dried reagents or monomers to deliver permeabilization reagents when the diffusion-resistant medium is applied to a biological sample. In some embodiments, the diffusion-resistant medium, (i.e. hydrogel) is covalently attached to a solid substrate (i.e. an acrylated glass slide). In some embodiments, the hydrogel
25 can be modified to both contain capture probes and deliver permeabilization reagents. For example, a hydrogel film can be modified to include spatially-barcoded capture probes. The spatially-barcoded hydrogel film is then soaked in permeabilization buffer before contacting the spatially-barcoded hydrogel film to the sample. The spatially-barcoded hydrogel film thus delivers permeabilization reagents to a sample surface in contact with the spatially-barcoded
30 hydrogel, enhancing analyte migration and capture. In some embodiments, the spatially-barcoded hydrogel is applied to a sample and placed in a permeabilization bulk solution. In some

embodiments, the hydrogel film soaked in permeabilization reagents is sandwiched between a sample and a spatially-barcoded array. In some embodiments, target analytes are able to diffuse through the permeabilizing reagent soaked hydrogel and hybridize or bind the capture probes on the other side of the hydrogel. In some embodiments, the thickness of the hydrogel is proportional to the resolution loss. In some embodiments, wells (e.g., micro-, nano-, or picowells) can contain spatially-barcoded capture probes and permeabilization reagents and/or buffer. In some embodiments, spatially-barcoded capture probes and permeabilization reagents are held between spacers. In some embodiments, the sample is punch, cut, or transferred into the well, wherein a target analyte diffuses through the permeabilization reagent/buffer and to the spatially-barcoded capture probes. In some embodiments, resolution loss may be proportional to gap thickness (e.g. the amount of permeabilization buffer between the sample and the capture probes). In some embodiments, the diffusion-resistant medium (e.g. hydrogel) is between approximately 50-500 micrometers thick including 500, 450, 400, 350, 300, 250, 200, 150, 100, or 50 micrometers thick, or any thickness within 50 and 500 micrometers.

In some embodiments, permeabilization solution can be delivered to a sample through a porous membrane. In some embodiments, a porous membrane is used to limit diffusive analyte losses, while allowing permeabilization reagents to reach a sample. Membrane chemistry and pore size can be manipulated to minimize analyte loss. In some embodiments, the porous membrane may be made of glass, silicon, paper, hydrogel, polymer monoliths, or other material. In some embodiments, the material may be naturally porous. In some embodiments, the material may have pores or wells etched into solid material. In some embodiments, the permeabilization reagents are flowed through a microfluidic chamber or channel over the porous membrane. In some embodiments, the flow controls the sample's access to the permeabilization reagents. In some embodiments, a porous membrane is sandwiched between a spatially-barcoded array and the sample, wherein permeabilization solution is applied over the porous membrane. The permeabilization reagents diffuse through the pores of the membrane and into the tissue.

In some embodiments, the biological sample can be permeabilized by adding one or more lysis reagents to the sample. Examples of suitable lysis agents include, but are not limited to, bioactive reagents such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, such as lysozymes, achromopeptidase,

lysostaphin, labiase, kitalase, lyticase, and a variety of other commercially available lysis enzymes.

Other lysis agents can additionally or alternatively be added to the biological sample to facilitate permeabilization. For example, surfactant-based lysis solutions can be used to lyse sample cells. Lysis solutions can include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). More generally, chemical lysis agents can include, without limitation, organic solvents, chelating agents, detergents, surfactants, and chaotropic agents.

In some embodiments, the biological sample can be permeabilized by non-chemical permeabilization methods. Non-chemical permeabilization methods are known in the art. For example, non-chemical permeabilization methods that can be used include, but are not limited to, physical lysis techniques such as electroporation, mechanical permeabilization methods (e.g., bead beating using a homogenizer and grinding balls to mechanically disrupt sample tissue structures), acoustic permeabilization (e.g., sonication), and thermal lysis techniques such as heating to induce thermal permeabilization of the sample.

(13) Selective Enrichment of RNA Species

In some embodiments, where RNA is the analyte, one or more RNA analyte species of interest can be selectively enriched. For example, one or more species of RNA of interest can be selected by addition of one or more oligonucleotides to the sample. In some embodiments, the additional oligonucleotide is a sequence used for priming a reaction by a polymerase. For example, one or more primer sequences with sequence complementarity to one or more RNAs of interest can be used to amplify the one or more RNAs of interest, thereby selectively enriching these RNAs. In some embodiments, an oligonucleotide with sequence complementarity to the complementary strand of captured RNA (e.g., cDNA) can bind to the cDNA. For example, biotinylated oligonucleotides with sequence complementary to one or more cDNA of interest binds to the cDNA and can be selected using biotinylation-streptavidin affinity using any of a variety of methods known to the field (e.g., streptavidin beads).

Alternatively, one or more species of RNA can be down-selected (e.g., removed) using any of a variety of methods. For example, probes can be administered to a sample that selectively hybridize to ribosomal RNA (rRNA), thereby reducing the pool and concentration of rRNA in the sample. Subsequent application of the capture probes to the sample can result in improved capture of other types of RNA due to the reduction in non-specific RNA present in the sample.

Additionally and alternatively, duplex-specific nuclease (DSN) treatment can remove rRNA (see, e.g., Archer, et al, Selective and flexible depletion of problematic sequencesw from RNA-seq libraries at the cDNA stage, *BMC Genomics*, 15 401, (2014), the entire contents of which are incorporated herein by reference). Furthermore, hydroxyapatite chromatography can remove
5 abundant species (e.g., rRNA) (see, e.g., Vandernoot, V.A., cDNA normalization by hydroxyapatite chromatography to enrich transcriptome diversity in RNA-seq applications, *Biotechniques*, 53(6) 373-80, (2012), the entire contents of which are incorporated herein by reference).

(14) Other Reagents

10 Additional reagents can be added to a biological sample to perform various functions prior to analysis of the sample. In some embodiments, DNase and RNase inactivating agents or inhibitors such as proteinase K, and/or chelating agents such as EDTA, can be added to the sample.

In some embodiments, the sample can be treated with one or more enzymes. For
15 example, one or more endonucleases to fragment DNA, DNA polymerase enzymes, and dNTPs used to amplify nucleic acids can be added. Other enzymes that can also be added to the sample include, but are not limited to, polymerase, transposase, ligase, and DNase, and RNase.

In some embodiments, reverse transcriptase enzymes can be added to the sample, including enzymes with terminal transferase activity, primers, and switch oligonucleotides.

20 Template switching can be used to increase the length of a cDNA, e.g., by appending a predefined nucleic acid sequence to the cDNA.

(15) Pre-processing for Capture Probe Interaction

In some embodiments, analytes in a biological sample can be pre-processed prior to interaction with a capture probe. For example, prior to interaction with capture probes,
25 polymerization reactions catalyzed by a polymerase (e.g., DNA polymerase or reverse transcriptase) are performed in the biological sample. In some embodiments, a primer for the polymerization reaction includes a functional group that enhances hybridization with the capture probe. The capture probes can include appropriate capture domains to capture biological analytes of interest (e.g., poly(dT) sequence to capture poly(A) mRNA).

30 In some embodiments, biological analytes are pre-processed for library generation via next generation sequencing. For example, analytes can be pre-processed by addition of a

modification (e.g., ligation of sequences that allow interaction with capture probes). In some embodiments, analytes (e.g., DNA or RNA) are fragmented using fragmentation techniques (e.g., using transposases and/or fragmentation buffers).

Fragmentation can be followed by a modification of the analyte. For example, a
5 modification can be the addition through ligation of an adapter sequence that allows hybridization with the capture probe. In some embodiments, where the analyte of interest is RNA, poly(A) tailing is performed. Addition of a poly(A) tail to RNA that does not contain a poly(A) tail can facilitate hybridization with a capture probe that includes a capture domain with a functional amount of poly(dT) sequence.

10 In some embodiments, prior to interaction with capture probes, ligation reactions catalyzed by a ligase are performed in the biological sample. In some embodiments, ligation can be performed by chemical ligation. In some embodiments, the ligation can be performed using click chemistry as further below. In some embodiments, the capture domain includes a DNA sequence that has complementarity to a RNA molecule, where the RNA molecule has
15 complementarity to a second DNA sequence, and where the RNA-DNA sequence complementarity is used to ligate the second DNA sequence to the DNA sequence in the capture domain. In these embodiments, direct detection of RNA molecules is possible.

In some embodiments, prior to interaction with capture probes, target-specific reactions are performed in the biological sample. Examples of target specific reactions include, but are not
20 limited to, ligation of target specific adaptors, probes and/or other oligonucleotides, target specific amplification using primers specific to one or more analytes, and target-specific detection using in situ hybridization, DNA microscopy, and/or antibody detection. In some embodiments, a capture probe includes capture domains targeted to target-specific products (e.g., amplification or ligation).

25 II. General Spatial Array-Based Analytical Methodology

This section of the disclosure describes methods, apparatus, systems, and compositions for spatial array-based analysis of biological samples.

(a) Spatial Analysis Methods

30 Array-based spatial analysis methods involve the transfer of one or more analytes from a biological sample to an array of features on a substrate, each of which is associated with a unique

spatial location on the array. Subsequent analysis of the transferred analytes includes determining the identity of the analytes and the spatial location of each analyte within the sample. The spatial location of each analyte within the sample is determined based on the feature to which each analyte is bound in the array, and the feature's relative spatial location within the array.

There are at least two general methods to associate a spatial barcode with one or more neighboring cells, such that the spatial barcode identifies the one or more cells, and/or contents of the one or more cells, as associated with a particular spatial location. One general method is to drive target analytes out of a cell and towards the spatially-barcode array. **FIG. 1** depicts an exemplary embodiment of this general method. In **FIG. 1**, the spatially-barcode array populated with capture probes (as described further herein) is contacted with a sample **101**, and sample is permeabilized, allowing the target analyte to migrate away from the sample and toward the array. The target analyte interacts with a capture probe on the spatially-barcode array **102**. Once the target analyte hybridizes/is bound to the capture probe, the sample is optionally removed from the array and the capture probes are analyzed in order to obtain spatially-resolved analyte information **103**.

Another general method is to cleave the spatially-barcode capture probes from an array, and drive the spatially-barcode capture probes towards and/or into or onto the sample. **FIG. 2** depicts an exemplary embodiment of this general method, the spatially-barcode array populated with capture probes (as described further herein) can be contacted with a sample **201**. The spatially-barcode capture probes are cleaved and then interact with cells within the provided sample **202**. The interaction can be a covalent or non-covalent cell-surface interaction. The interaction can be an intracellular interaction facilitated by a delivery system or a cell penetration peptide. Once the spatially-barcode capture probe is associated with a particular cell, the sample can be optionally removed for analysis. The sample can be optionally dissociated before analysis. Once the tagged cell is associated with the spatially-barcode capture probe, the capture probes can be analyzed to obtain spatially-resolved information about the tagged cell **203**.

FIG. 3 shows an exemplary workflow that includes preparing a sample on a spatially-barcode array **301**. Sample preparation may include placing the sample on a slide, fixing the sample, and/or staining the sample for imaging. The stained sample is then imaged on the array **302** using both brightfield (to image the sample hematoxylin and eosin stain) and fluorescence

(to image features) modalities. In some embodiments, target analytes are then released from the sample and capture probes forming the spatially-barcoded array hybridize or bind the released target analytes **303**. The sample is then removed from the array **304** and the capture probes cleaved from the array **305**. The sample and array are then optionally imaged a second time in both modalities **305B** while the analytes are reverse transcribed into cDNA, and an amplicon library is prepared **306** and sequenced **307**. The two sets of images are then spatially-overlaid in order to correlate spatially-identified sample information **308**. When the sample and array are not imaged a second time, **305B**, a spot coordinate file is supplied by the manufacturer instead. The spot coordinate file replaces the second imaging step **305B**. Further, amplicon library preparation **306** can be performed with a unique PCR adapter and sequenced **307**.

FIG. 4 shows another exemplary workflow that utilizes a spatially-labelled array on a substrate, where capture probes labelled with spatial barcodes are clustered at areas called features. The spatially-labelled capture probes can include a cleavage domain, one or more functional sequences, a spatial barcode, a unique molecular identifier, and a capture domain. The spatially-labelled capture probes can also include a 5' end modification for reversible attachment to the substrate. The spatially-barcoded array is contacted with a sample **401**, and the sample is permeabilized through application of permeabilization reagents **402**. Permeabilization reagents may be administered by placing the array/sample assembly within a bulk solution. Alternatively, permeabilization reagents may be administered to the sample via a diffusion-resistant medium and/or a physical barrier such as a lid, wherein the sample is sandwiched between the diffusion-resistant medium and/or barrier and the array-containing substrate. The analytes are migrated toward the spatially-barcoded capture array using any number of techniques disclosed herein. For example, analyte migration can occur using a diffusion-resistant medium lid and passive migration. As another example, analyte migration can be active migration, using an electrophoretic transfer system, for example. Once the analytes are in close proximity to the spatially-barcoded capture probes, the capture probes can hybridize or otherwise bind a target analyte **403**. The sample can be optionally removed from the array **404**.

The capture probes can be optionally cleaved from the array **405**, and the captured analytes can be spatially-tagged by performing a reverse transcriptase first strand cDNA reaction. A first strand cDNA reaction can be optionally performed using template switching oligonucleotides. For example, a template switching oligonucleotide can hybridize to a poly(C)

tail added to a 3' end of the cDNA by a reverse transcriptase enzyme. The original mRNA template and template switching oligonucleotide can then be denatured from the cDNA and the barcoded capture probe can then hybridize with the cDNA and a complement of the cDNA can be generated. The first strand cDNA can then be purified and collected for downstream
5 amplification steps. The first strand cDNA can be amplified using PCR **406**, wherein the forward and reverse primers flank the spatial barcode and target analyte regions of interest, generating a library associated with a particular spatial barcode. In some embodiments, the cDNA comprises a sequencing by synthesis (SBS) primer sequence. The library amplicons are sequenced and analyzed to decode spatial information **407**.

FIG. 5 depicts an exemplary workflow where the sample is removed from the spatially-
barcoded array and the spatially-barcoded capture probes are removed from the array for
barcoded analyte amplification and library preparation. Another embodiment includes
performing first strand synthesis using template switching oligonucleotides on the spatially-
barcoded array without cleaving the capture probes. In this embodiment, sample preparation **501**
15 and permeabilization **502** are performed as described elsewhere herein. Once the capture probes
capture the target analyte(s), first strand cDNA created by template switching and reverse
transcriptase **503** is then denatured and the second strand is then extended **504**. The second
strand cDNA is then denatured from the first strand cDNA, neutralized, and transferred to a tube
505. cDNA quantification and amplification can be performed using standard techniques
20 discussed herein. The cDNA can then be subjected to library preparation **506** and indexing **507**,
including fragmentation, end-repair, and a-tailing, and indexing PCR steps.

In some non-limiting examples of the workflow above, the sample can be immersed in
100% chilled methanol and incubated for 30 minutes at -20 °C. After 20 minutes, the sample can
be removed and rinsed in ultrapure water. After rinsing the sample, fresh eosin solution is
25 prepared, and the sample can be covered in isopropanol. After incubating the sample in
isopropanol for 1 minute, the reagent can be removed by holding the slide at an angle, where the
bottom edge of the slide can be in contact with a laboratory wipe and air dried. The sample can
be uniformly covered in hematoxylin solution and incubated for 7 minutes at room temperature.
After incubating the sample in hematoxylin for 7 minutes, the reagent can be removed by
30 holding the slide at an angle, where the bottom edge of the slide can be in contact with a
laboratory wipe. The slide containing the sample can be immersed in water and the excess liquid

can be removed. After that, the sample can be covered with blueing buffer and can be incubated for 2 minutes at room temperature. The slide containing the sample can again be immersed in water, and uniformly covered with eosin solution and incubated for 1 minute at room temperature. The slide can be air-dried and incubated for 5 minutes at 37 °C. The sample can be
5 imaged using the methods disclosed herein.

The following are non-limiting, exemplary steps for sample permeabilization and cDNA generation. The sample can be exposed to a permeabilization enzyme and incubated for 6 minutes at 37 °C. Other permeabilization methods are described herein. The permeabilization enzyme can be removed and the sample prepared for analyte capture by adding SSC buffer. The
10 sample can then subjected to a pre-equilibration thermocycling protocol and the SSC buffer can be removed. A Master Mix, containing nuclease-free water, a reverse transcriptase reagent, a template switch oligo, a reducing agent, and a reverse transcriptase enzyme can be added, and the sample with the Master Mix can be subjected to a thermocycling protocol. The reagents can be removed from the sample and NaOH can be applied and incubated for 5 minutes at room
15 temperature. The NaOH can be removed and elution buffer can be added and removed from the sample. A Second Strand Mix, including a second strand reagent, a second strand primer, and a second strand enzyme, can be added to the sample and the sample can be sealed and incubated. At the end of the incubation, the reagents can be removed and elution buffer can be added and removed from the sample, and NaOH can be added again to the sample and the sample can be
20 incubated for 10 minutes at room temperature. Tris-HCl can be added and the reagents can be mixed.

The following steps are non-limiting, exemplary steps for cDNA amplification and quality control. A qPCR Mix, including nuclease-free water, qPCR Master Mix, and cDNA primers, can be prepared and the NaOH/Tris-HCl mix can be mixed with the qPCR Mix and the
25 sample, and thermocycled according to a predetermined thermocycling protocol. After completing the thermocycling, a cDNA amplification mix can be prepared and combined with the sample and mixed. The sample can then be incubated and thermocycled. The sample can then be resuspended in SPRIselect Reagent and pipetted to ensure proper mixing. The sample can then be incubated at 5 minutes at room temperature, and cleared by placing the sample on a
30 magnet (e.g., the magnet is in the high position). The supernatant can be removed and 80% ethanol can be added to the pellet, and incubated for 30 seconds. The ethanol can be removed

and the pellet can be washed again. The sample can then be centrifuged and placed on a magnet (e.g., the magnet is on the low position). Any remaining ethanol can be removed and the sample can be air dried. The magnet can be removed and elution buffer can be added to the sample, mixed, and incubated for 2 minutes at room temperature. The sample can then be placed on the magnet (e.g., on high position) until the solution clears. A portion of the sample can be run on an Agilent Bioanalyzer High Sensitivity chip, where a region can be selected and the cDNA concentration can be measured to calculate the total cDNA yield. Alternatively, the quantification can be determined by Agilent Bioanalyzer or Agilent TapeStation.

The following steps are non-limiting, exemplary steps for spatial gene expression library construction. A Fragmentation Mix, including a fragmentation buffer and fragmentation enzyme, can be prepared on ice. Elution buffer and fragmentation mix can be added to each sample, mixed, and centrifuged. The sample mix can then be placed in a thermocycler and cycled according to a predetermined protocol. The SPRIselect Reagent can be added to the sample and incubated at 5 minutes at room temperature. The sample can be placed on a magnet (e.g., in the high position) until the solution clears, and the supernatant can be transferred to a new tube strip. SPRIselect Reagent can be added to the sample, mixed, and incubated for 5 minutes at room temperature. The sample can be placed on a magnet (e.g., in the high position) until the solution clears. The supernatant can be removed and 80% ethanol can be added to the pellet, the pellet can be incubated for 30 seconds, and the ethanol can be removed. The ethanol wash can be repeated and the sample placed on a magnet (e.g., in the low position) until the solution clears. The remaining ethanol can be removed and elution buffer can be added to the sample, mixed, and incubated for 2 minutes at room temperature. The sample can be placed on a magnet (e.g., in the high position) until the solution clears, and a portion of the sample can be moved to a new tube strip. An Adaptor Ligation Mix, including ligation buffer, DNA ligase, and adaptor oligos, can be prepared and centrifuged. The Adaptor Ligation Mix can be added to the sample, pipette-mixed, and centrifuged briefly. The sample can then be thermocycled according to a predetermined protocol. The SPRIselect Reagent can be added to the sample, incubated for 5 minutes at room temperature, and placed on a magnet (e.g., in the high position) until the solution clears. The supernatant can be removed and the pellet can be washed with 80% ethanol, incubated for 30 seconds, and the ethanol can be removed. The ethanol wash can be repeated, and the sample can be centrifuged briefly before placing the sample on a magnet (e.g., in the low

position). Any remaining ethanol can be removed and the sample can be air dried. Elution buffer can be added to the sample, the sample can be removed from the magnet, and the sample can be pipette-mixed, incubated for 2 minutes at room temperature, and placed on a magnet (e.g., in the low position) until the solution clears. A portion of the sample can be transferred to a new tube strip. A Sample Index PCR Mix, including amplification mix and SI primer, can be prepared and combined with the sample. The sample / Sample Index PCR Mix can be loaded into an individual Chromium i7 Sample Index well and a thermocycling protocol can be used.

SPRIsselect Reagent can be added to each sample, mixed, and incubated for 5 minutes at room temperature. The sample can be placed on a magnet (e.g., in the high position) until the solution clears, and the supernatant can be transferred to a new tube strip. The SPRIsselect Reagent can be added to each sample, pipette-mixed, and incubated for 5 minutes at room temperature. The sample can then placed on a magnet (e.g., in the high position) until the solution clears. The supernatant can be removed, and the pellet can be washed with 80% ethanol, incubated for 30 seconds, and then the ethanol can be removed. The ethanol wash can be repeated, the sample centrifuged, and placed on a magnet (e.g., in the low position) to remove any remaining ethanol. The sample can be removed from the magnet and Elution Buffer can be added to the sample, pipette-mixed, and incubated at 2 minutes at room temperature. The sample can be placed on a magnet (e.g., in the low position) until the solution clears and a portion of the sample can be transferred to a new tube strip. The average fragment size can be determined using a Bioanalyzer trace or an Agilent TapeStation.

In some embodiments, performing correlative analysis of data produced by this workflow, and other workflows described herein, can yield over 95% correlation of genes expressed across two capture areas (e.g. 95% or greater, 96% or greater, 97% or greater, 98% or greater, or 99% or greater). When performing the described workflows using single cell RNA sequencing of nuclei, in some embodiments, correlative analysis of the data can yield over 90% (e.g. over 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) correlation of genes expressed across two capture areas.

(b) Capture Probes

A “capture probe” refers to any molecule capable of capturing (directly or indirectly) and/or labelling an analyte of interest in a biological sample. In some embodiments, the capture probe is a nucleic acid or a polypeptide. In some embodiments, the capture probe is a conjugate

(e.g., an oligonucleotide-antibody conjugate). In some embodiments, the capture probe includes a barcode (e.g., a spatial barcode and/or a unique molecular identifier (UMI)) and a capture domain.

FIG. 6 is a schematic diagram showing an example of a capture probe, as described herein. As shown, the capture probe **602** is optionally coupled to a feature **601** by a cleavage domain **603**, such as a disulfide linker. The capture probe can include functional sequences that are useful for subsequent processing, such as functional sequence **604**, which can include a sequencer specific flow cell attachment sequence, e.g., a P5 sequence, as well as functional sequence **606**, which can include sequencing primer sequences, e.g., a R1 primer binding site. In some embodiments, sequence **604** is a P7 sequence and sequence **606** is a R2 primer binding site. A spatial barcode **605** can be included within the capture probe for use in barcoding the target analyte. The functional sequences can generally be selected for compatibility with any of a variety of different sequencing systems, e.g., 454 Sequencing, Ion Torrent Proton or PGM, Illumina X10, PacBio, Nanopore, etc., and the requirements thereof. In some embodiments, functional sequences can be selected for compatibility with non-commercialized sequencing systems. Examples of such sequencing systems and techniques, for which suitable functional sequences can be used, include (but are not limited to) Roche 454 sequencing, Ion Torrent Proton or PGM sequencing, Illumina X10 sequencing, PacBio SMRT sequencing, and Oxford Nanopore sequencing. Further, in some embodiments, functional sequences can be selected for compatibility with other sequencing systems, including non-commercialized sequencing systems.

In some embodiments, the spatial barcode **605**, functional sequences **604** (e.g., flow cell attachment sequence) and **606** (e.g., sequencing primer sequences) can be common to all of the probes attached to a given feature. The spatial barcode can also include a capture domain **607** to facilitate capture of a target analyte.

Capture Domain

As discussed above, each capture probe includes at least one capture domain. The “capture domain” is an oligonucleotide, a polypeptide, a small molecule, or any combination thereof, that binds specifically to a desired analyte. In some embodiments, a capture domain can be used to capture or detect a desired analyte.

In some embodiments, the capture domain is a functional nucleic acid sequence configured to interact with one or more analytes, such as one or more different types of nucleic

acids (e.g., RNA molecules and DNA molecules). In some embodiments, the functional nucleic acid sequence can include an N-mer sequence (e.g., a random N-mer sequence), which N-mer sequences are configured to interact with a plurality of DNA molecules. In some embodiments, the functional sequence can include a poly(T) sequence, which poly(T) sequences are configured to interact with messenger RNA (mRNA) molecules via the poly(A) tail of an mRNA transcript. In some embodiments, the functional nucleic acid sequence is the binding target of a protein (e.g., a transcription factor, a DNA binding protein, or a RNA binding protein), where the analyte of interest is a protein.

Capture probes can include ribonucleotides and/or deoxyribonucleotides as well as synthetic nucleotide residues that are capable of participating in Watson-Crick type or analogous base pair interactions. In some embodiments, the capture domain is capable of priming a reverse transcription reaction to generate cDNA that is complementary to the captured RNA molecules. In some embodiments, the capture domain of the capture probe can prime a DNA extension (polymerase) reaction to generate DNA that is complementary to the captured DNA molecules. In some embodiments, the capture domain can template a ligation reaction between the captured DNA molecules and a surface probe that is directly or indirectly immobilized on the substrate. In some embodiments, the capture domain can be ligated to one strand of the captured DNA molecules. For example, SplintR ligase along with RNA or DNA sequences (e.g., degenerate RNA) can be used to ligate a single-stranded DNA or RNA to the capture domain. In some embodiments, ligases with RNA-templated ligase activity, e.g., SplintR ligase, T4 RNA ligase 2 or KOD ligase, can be used to ligate a single-stranded DNA or RNA to the capture domain. In some embodiments, a capture domain includes a splint oligonucleotide. In some embodiments, a capture domain captures a splint oligonucleotide.

In some embodiments, the capture domain is located at the 3' end of the capture probe and includes a free 3' end that can be extended, e.g. by template dependent polymerization, to form an extended capture probe as described herein. In some embodiments, the capture domain includes a nucleotide sequence that is capable of hybridizing to nucleic acid, e.g. RNA or other analyte, present in the cells of the tissue sample contacted with the array. In some embodiments, the capture domain can be selected or designed to bind selectively or specifically to a target nucleic acid. For example, the capture domain can be selected or designed to capture mRNA by way of hybridization to the mRNA poly(A) tail. Thus, in some embodiments, the capture domain

includes a poly(T) DNA oligonucleotide, i.e., a series of consecutive deoxythymidine residues linked by phosphodiester bonds, which is capable of hybridizing to the poly(A) tail of mRNA. In some embodiments, the capture domain can include nucleotides that are functionally or structurally analogous to a poly(T) tail. For example, a poly(U) oligonucleotide or an oligonucleotide included of deoxythymidine analogues. In some embodiments, the capture domain includes at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides. In some embodiments, the capture domain includes at least 25, 30, or 35 nucleotides.

In some embodiments, random sequences, e.g., random hexamers or similar sequences, can be used to form all or a part of the capture domain. For example, random sequences can be used in conjunction with poly(T) (or poly(T) analogue) sequences. Thus, where a capture domain includes a poly(T) (or a “poly(T)-like”) oligonucleotide, it can also include a random oligonucleotide sequence (e.g., “poly(T)-random sequence” probe). This can, for example, be located 5’ or 3’ of the poly(T) sequence, e.g. at the 3’ end of the capture domain. The poly(T)-random sequence probe can facilitate the capture of the mRNA poly(A) tail. In some embodiments, the capture domain can be an entirely random sequence. In some embodiments, degenerate capture domains can be used.

In some embodiments, a pool of two or more capture probes form a mixture, where the capture domain of one or more capture probes includes a poly(T) sequence and the capture domain of one or more capture probes includes random sequences. In some embodiments, a pool of two or more capture probes form a mixture where the capture domain of one or more capture probes includes poly(T)-like sequence and the capture domain of one or more capture probes includes random sequences. In some embodiments, a pool of two or more capture probes form a mixture where the capture domain of one or more capture probes includes a poly(T)-random sequences and the capture domain of one or more capture probes includes random sequences. In some embodiments, probes with degenerate capture domains can be added to any of the preceding combinations listed herein. In some embodiments, probes with degenerate capture domains can be substituted for one of the probes in each of the pairs described herein.

The capture domain can be based on a particular gene sequence or particular motif sequence or common/conserved sequence, that it is designed to capture (i.e., a sequence-specific capture domain). Thus, in some embodiments, the capture domain is capable of binding selectively to a desired sub-type or subset of nucleic acid, for example a particular type of RNA,

such as mRNA, rRNA, tRNA, SRP RNA, tmRNA, snRNA, snoRNA, SmY RNA, scaRNA, gRNA, RNase P, RNase MRP, TERC, SL RNA, aRNA, cis-NAT, crRNA, lncRNA, miRNA, piRNA, siRNA, shRNA, tasiRNA, rasiRNA, 7SK, eRNA, ncRNA or other types of RNA. In a non-limiting example, the capture domain can be capable of binding selectively to a desired subset of ribonucleic acids, for example, microbiome RNA, such as 16S rRNA.

In some embodiments, a capture domain includes an “anchor” or “anchoring sequence”, which is a sequence of nucleotides that is designed to ensure that the capture domain hybridizes to the intended biological analyte. In some embodiments, an anchor sequence includes a sequence of nucleotides, including a 1-mer, 2-mer, 3-mer or longer sequence. In some embodiments, the short sequence is random. For example, a capture domain including a poly(T) sequence can be designed to capture an mRNA. In such embodiments, an anchoring sequence can include a random 3-mer (e.g., GGG) that helps ensure that the poly(T) capture domain hybridizes to an mRNA. In some embodiments, an anchoring sequence can be VN, N, or NN. Alternatively, the sequence can be designed using a specific sequence of nucleotides. In some embodiments, the anchor sequence is at the 3' end of the capture domain. In some embodiments, the anchor sequence is at the 5' end of the capture domain.

In some embodiments, capture domains of capture probes are blocked prior to contacting the biological sample with the array, and blocking probes are used when the nucleic acid in the biological sample is modified prior to its capture on the array. In some embodiments, the blocking probe is used to block or modify the free 3' end of the capture domain. In some embodiments, blocking probes can be hybridized to the capture probes to mask the free 3' end of the capture domain, e.g., hairpin probes or partially double stranded probes. In some embodiments, the free 3' end of the capture domain can be blocked by chemical modification, e.g., addition of an azidomethyl group as a chemically reversible capping moiety such that the capture probes do not include a free 3' end. Blocking or modifying the capture probes, particularly at the free 3' end of the capture domain, prior to contacting the biological sample with the array, prevents modification of the capture probes, e.g., prevents the addition of a poly(A) tail to the free 3' end of the capture probes.

Non-limiting examples of 3' modifications include dideoxy C-3' (3'-ddC), 3' inverted dT, 3' C3 spacer, 3' Amino, and 3' phosphorylation. In some embodiments, the nucleic acid in the biological sample can be modified such that it can be captured by the capture domain. For

example, an adaptor sequence (including a binding domain capable of binding to the capture domain of the capture probe) can be added to the end of the nucleic acid, e.g., fragmented genomic DNA. In some embodiments, this is achieved by ligation of the adaptor sequence or extension of the nucleic acid. In some embodiments, an enzyme is used to incorporate additional nucleotides at the end of the nucleic acid sequence, e.g., a poly(A) tail. In some embodiments, the capture probes can be reversibly masked or modified such that the capture domain of the capture probe does not include a free 3' end. In some embodiments, the 3' end is removed, modified, or made inaccessible so that the capture domain is not susceptible to the process used to modify the nucleic acid of the biological sample, e.g., ligation or extension.

In some embodiments, the capture domain of the capture probe is modified to allow the removal of any modifications of the capture probe that occur during modification of the nucleic acid molecules of the biological sample. In some embodiments, the capture probes can include an additional sequence downstream of the capture domain, i.e., 3' to the capture domain, namely a blocking domain.

In some embodiments, the capture domain of the capture probe can be a non-nucleic acid domain. Examples of suitable capture domains that are not exclusively nucleic-acid based include, but are not limited to, proteins, peptides, aptamers, antigens, antibodies, and molecular analogs that mimic the functionality of any of the capture domains described herein.

Cleavage Domain

Each capture probe can optionally include at least one cleavage domain. The cleavage domain represents the portion of the probe that is used to reversibly attach the probe to an array feature, as will be described further below. Further, one or more segments or regions of the capture probe can optionally be released from the array feature by cleavage of the cleavage domain. As an example spatial barcodes and/or universal molecular identifiers (UMIs) can be released by cleavage of the cleavage domain.

FIG. 7 is a schematic illustrating a cleavable capture probe, wherein the cleaved capture probe can enter into a non-permeabilized cell and bind to target analytes within the sample. The capture probe **701** contains a cleavage domain **702**, a cell penetrating peptide **703**, a reporter molecule **704**, and a disulfide bond (-S-S-). **705** represents all other parts of a capture probe, for example a spatial barcode and a capture domain.

In some embodiments, the cleavage domain linking the capture probe to a feature is a disulfide bond. A reducing agent can be added to break the disulfide bonds, resulting in release of the capture probe from the feature. As another example, heating can also result in degradation of the cleavage domain and release of the attached capture probe from the array feature. In some
5 embodiments, laser radiation is used to heat and degrade cleavage domains of capture probes at specific locations. In some embodiments, the cleavage domain is a photo-sensitive chemical bond (i.e., a chemical bond that dissociates when exposed to light such as ultraviolet light).

Other examples of cleavage domains include labile chemical bonds such as, but not limited to, ester linkages (e.g., cleavable with an acid, a base, or hydroxylamine), a vicinal diol
10 linkage (e.g., cleavable via sodium periodate), a Diels-Alder linkage (e.g., cleavable via heat), a sulfone linkage (e.g., cleavable via a base), a silyl ether linkage (e.g., cleavable via an acid), a glycosidic linkage (e.g., cleavable via an amylase), a peptide linkage (e.g., cleavable via a protease), or a phosphodiester linkage (e.g., cleavable via a nuclease (e.g., DNAase)).

In some embodiments, the cleavage domain includes a sequence that is recognized by one
15 or more enzymes capable of cleaving a nucleic acid molecule, e.g., capable of breaking the phosphodiester linkage between two or more nucleotides. A bond can be cleavable via other nucleic acid molecule targeting enzymes, such as restriction enzymes (e.g., restriction endonucleases). For example, the cleavage domain can include a restriction endonuclease (restriction enzyme) recognition sequence. Restriction enzymes cut double-stranded or single
20 stranded DNA at specific recognition nucleotide sequences known as restriction sites. In some embodiments, a rare-cutting restriction enzyme, i.e., enzymes with a long recognition site (at least 8 base pairs in length), is used to reduce the possibility of cleaving elsewhere in the capture probe.

In some embodiments, the cleavage domain includes a poly(U) sequence which can be
25 cleaved by a mixture of Uracil DNA glycosylase (UDG) and the DNA glycosylase-lyase Endonuclease VIII, commercially known as the USER™ enzyme. Releasable capture probes can be available for reaction once released. Thus, for example, an activatable capture probe can be activated by releasing the capture probes from a feature.

In some embodiments, where the capture probe is attached indirectly to a substrate, e.g.,
30 via a surface probe, the cleavage domain includes one or more mismatch nucleotides, so that the complementary parts of the surface probe and the capture probe are not 100% complementary

(for example, the number of mismatched base pairs can one, two, or three base pairs). Such a mismatch is recognized, e.g., by the MutY and T7 endonuclease I enzymes, which results in cleavage of the nucleic acid molecule at the position of the mismatch.

In some embodiments, where the capture probe is attached to a feature indirectly, e.g., via
5 a surface probe, the cleavage domain includes a nickase recognition site or sequence. Nickases are endonucleases which cleave only a single strand of a DNA duplex. Thus, the cleavage domain can include a nickase recognition site close to the 5' end of the surface probe (and/or the 5' end of the capture probe) such that cleavage of the surface probe or capture probe destabilizes the duplex between the surface probe and capture probe thereby releasing the capture probe)
10 from the feature.

Nickase enzymes can also be used in some embodiments where the capture probe is attached to the feature directly. For example, the substrate can be contacted with a nucleic acid molecule that hybridizes to the cleavage domain of the capture probe to provide or reconstitute a nickase recognition site, e.g., a cleavage helper probe. Thus, contact with a nickase enzyme will
15 result in cleavage of the cleavage domain thereby releasing the capture probe from the feature. Such cleavage helper probes can also be used to provide or reconstitute cleavage recognition sites for other cleavage enzymes, e.g., restriction enzymes.

Some nickases introduce single-stranded nicks only at particular sites on a DNA molecule, by binding to and recognizing a particular nucleotide recognition sequence. A number
20 of naturally-occurring nickases have been discovered, of which at present the sequence recognition properties have been determined for at least four. Nickases are described in U.S. Patent No. 6,867,028, which is incorporated herein by reference in its entirety. In general, any suitable nickase can be used to bind to a complementary nickase recognition site of a cleavage domain. Following use, the nickase enzyme can be removed from the assay or inactivated
25 following release of the capture probes to prevent unwanted cleavage of the capture probes.

Examples of suitable capture domains that are not exclusively nucleic-acid based include, but are not limited to, proteins, peptides, aptamers, antigens, antibodies, and molecular analogs that mimic the functionality of any of the capture domains described herein.

In some embodiments, a cleavage domain is absent from the capture probe. Examples of
30 substrates with attached capture probes lacking a cleavage domain are described for example in

Macosko et al., (2015) Cell 161, 1202–1214, the entire contents of which are incorporated herein by reference.

In some embodiments, the region of the capture probe corresponding to the cleavage domain can be used for some other function. For example, an additional region for nucleic acid extension or amplification can be included where the cleavage domain would normally be positioned. In such embodiments, the region can supplement the functional domain or even exist as an additional functional domain. In some embodiments, the cleavage domain is present but its use is optional.

Functional Domain

Each capture probe can optionally include at least one functional domain. Each functional domain typically includes a functional nucleotide sequence for a downstream analytical step in the overall analysis procedure.

In some embodiments, the capture probe can include a functional domain for attachment to a sequencing flow cell, such as, for example, a P5 sequence for Illumina® sequencing. In some embodiments, the capture probe or derivative thereof can include another functional domain, such as, for example, a P7 sequence for attachment to a sequencing flow cell for Illumina® sequencing. The functional domains can be selected for compatibility with a variety of different sequencing systems, e.g., 454 Sequencing, Ion Torrent Proton or PGM, Illumina X10, etc., and the requirements thereof.

In some embodiments, the functional domain includes a primer. The primer can include an R1 primer sequence for Illumina® sequencing, and in some embodiments, an R2 primer sequence for Illumina® sequencing. Examples of such capture probes and uses thereof are described in U.S. Patent Publication Nos. 2014/0378345 and 2015/0376609, the entire contents of each of which are incorporated herein by reference.

Spatial Barcode

As discussed above, the capture probe can include one or more spatial barcodes (e.g., two or more, three or more, four or more, five or more) spatial barcodes. A “spatial barcode” is a contiguous nucleic acid segment or two or more non-contiguous nucleic acid segments that function as a label or identifier that conveys or is capable of conveying spatial information. In some embodiments, a capture probe includes a spatial barcode that possesses a spatial aspect,

where the barcode is associated with a particular location within an array or a particular location on a substrate.

A spatial barcode can be part of an analyte, or independent from an analyte (i.e., part of the capture probe). A spatial barcode can be a tag attached to an analyte (e.g., a nucleic acid molecule) or a combination of a tag in addition to an endogenous characteristic of the analyte (e.g., size of the analyte or end sequence(s)). A spatial barcode can be unique. In some embodiments where the spatial barcode is unique, the spatial barcode functions both as a spatial barcode and as a unique molecular identifier (UMI), associated with one particular capture probe.

Spatial barcodes can have a variety of different formats. For example, spatial barcodes can include polynucleotide spatial barcodes; random nucleic acid and/or amino acid sequences; and synthetic nucleic acid and/or amino acid sequences. In some embodiments, a spatial barcode is attached to an analyte in a reversible or irreversible manner. In some embodiments, a spatial barcode is added to, for example, a fragment of a DNA or RNA sample before, during, and/or after sequencing of the sample. In some embodiments, a spatial barcode allows for identification and/or quantification of individual sequencing-reads. In some embodiments, a spatial barcode is used as a fluorescent barcode for which fluorescently labeled oligonucleotide probes hybridize to the spatial barcode.

In some embodiments, the spatial barcode is a nucleic acid sequence that does not substantially hybridize to analyte nucleic acid molecules in a biological sample. In some embodiments, the spatial barcode has less than 80% sequence identity (e.g., less than 70%, 60%, 50%, or less than 40% sequence identity) to the nucleic acid sequences across a substantial part (e.g., 80% or more) of the nucleic acid molecules in the biological sample.

The spatial barcode sequences can include from about 6 to about 20 or more nucleotides within the sequence of the capture probes. In some embodiments, the length of a spatial barcode sequence can be about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some embodiments, the length of a spatial barcode sequence can be at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some embodiments, the length of a spatial barcode sequence is at most about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or shorter.

These nucleotides can be completely contiguous, i.e., in a single stretch of adjacent nucleotides, or they can be separated into two or more separate subsequences that are separated by 1 or more nucleotides. Separated spatial barcode subsequences can be from about 4 to about 16 nucleotides in length. In some embodiments, the spatial barcode subsequence can be about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some embodiments, the spatial barcode subsequence can be at least about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some embodiments, the spatial barcode subsequence can be at most about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or shorter.

For multiple capture probes that are attached to a common array feature, the one or more spatial barcode sequences of the multiple capture probes can include sequences that are the same for all capture probes coupled to the feature, and/or sequences that are different across all capture probes coupled to the feature.

FIG. 8 is a schematic diagram of an exemplary multiplexed spatially-labelled feature. In **FIG. 8**, the feature **801** can be coupled to spatially-barcoded capture probes, wherein the spatially-barcoded probes of a particular feature can possess the same spatial barcode, but have different capture domains designed to associate the spatial barcode of the feature with more than one target analyte. For example, a feature may be coupled to four different types of spatially-barcoded capture probes, each type of spatially-barcoded capture probe possessing the spatial barcode **802**. One type of capture probe associated with the feature includes the spatial barcode **802** in combination with a poly(T) capture domain **803**, designed to capture mRNA target analytes. A second type of capture probe associated with the feature includes the spatial barcode **802** in combination with a random N-mer capture domain **804** for gDNA analysis. A third type of capture probe associated with the feature includes the spatial barcode **802** in combination with a capture domain complementary to the capture domain on an analyte capture agent capture agent barcode domain **805**. A fourth type of capture probe associated with the feature includes the spatial barcode **802** in combination with a capture probe that can specifically bind a nucleic acid molecule **806** that can function in a CRISPR assay (e.g., CRISPR/Cas9). While only four different capture probe-barcoded constructs are shown in **FIG. 8**, capture-probe barcoded constructs can be tailored for analyses of any given analyte associated with a nucleic acid and capable of binding with such a construct. For example, the schemes shown in **FIG. 8** can also be used for concurrent analysis of other analytes disclosed herein, including, but not limited to: (a)

mRNA, a lineage tracing construct, cell surface or intracellular proteins and metabolites, and gDNA; (b) mRNA, accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq) cell surface or intracellular proteins and metabolites, and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein); (c) mRNA, cell surface or intracellular proteins and/or metabolites, a barcoded labelling agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor).

Capture probes attached to a single array feature can include identical (or common) spatial barcode sequences, different spatial barcode sequences, or a combination of both. Capture probes attached to a feature can include multiple sets of capture probes. Capture probes of a given set can include identical spatial barcode sequences. The identical spatial barcode sequences can be different from spatial barcode sequences of capture probes of another set.

The plurality of capture probes can include spatial barcode sequences (e.g., nucleic acid barcode sequences) that are associated with specific locations on a spatial array. For example, a first plurality of capture probes can be associated with a first region, based on a spatial barcode sequence common to the capture probes within the first region, and a second plurality of capture probes can be associated with a second region, based on a spatial barcode sequence common to the capture probes within the second region. The second region may or may not be associated with the first region. Additional pluralities of capture probes can be associated with spatial barcode sequences common to the capture probes within other regions. In some embodiments, the spatial barcode sequences can be the same across a plurality of capture probe molecules.

In some embodiments, multiple different spatial barcodes are incorporated into a single arrayed capture probe. For example, a mixed but known set of spatial barcode sequences can provide a stronger address or attribution of the spatial barcodes to a given spot or location, by providing duplicate or independent confirmation of the identity of the location. In some embodiments, the multiple spatial barcodes represent increasing specificity of the location of the particular array point.

Unique Molecular Identifier

The capture probe can include one or more (e.g., two or more, three or more, four or more, five or more) Unique Molecular Identifiers (UMIs). A unique molecular identifier is a contiguous nucleic acid segment or two or more non-contiguous nucleic acid segments that

function as a label or identifier for a particular analyte, or for a capture probe that binds a particular analyte (e.g., via the capture domain).

A UMI can be unique. A UMI can include one or more specific polynucleotides sequences, one or more random nucleic acid and/or amino acid sequences, and/or one or more
5 synthetic nucleic acid and/or amino acid sequences.

In some embodiments, the UMI is a nucleic acid sequence that does not substantially hybridize to analyte nucleic acid molecules in a biological sample. In some embodiments, the UMI has less than 80% sequence identity (e.g., less than 70%, 60%, 50%, or less than 40% sequence identity) to the nucleic acid sequences across a substantial part (e.g., 80% or more) of
10 the nucleic acid molecules in the biological sample.

The UMI can include from about 6 to about 20 or more nucleotides within the sequence of the capture probes. In some embodiments, the length of a UMI sequence can be about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or longer. In some embodiments, the length of a UMI sequence can be at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20
15 nucleotides or longer. In some embodiments, the length of a UMI sequence is at most about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 nucleotides or shorter.

These nucleotides can be completely contiguous, i.e., in a single stretch of adjacent nucleotides, or they can be separated into two or more separate subsequences that are separated by 1 or more nucleotides. Separated UMI subsequences can be from about 4 to about 16
20 nucleotides in length. In some embodiments, the UMI subsequence can be about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some embodiments, the UMI subsequence can be at least about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or longer. In some embodiments, the UMI subsequence can be at most about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 nucleotides or shorter.

In some embodiments, a UMI is attached to an analyte in a reversible or irreversible manner. In some embodiments, a UMI is added to, for example, a fragment of a DNA or RNA sample before, during, and/or after sequencing of the analyte. In some embodiments, a UMI allows for identification and/or quantification of individual sequencing-reads. In some
25 embodiments, a UMI is used as a fluorescent barcode for which fluorescently labeled
30 oligonucleotide probes hybridize to the UMI.

Other Aspects of Capture Probes

For capture probes that are attached to an array feature, an individual array feature can include one or more capture probes. In some embodiments, an individual array feature includes hundreds or thousands of capture probes. In some embodiments, the capture probes are associated with a particular individual feature, where the individual feature contains a capture
5 probe including a spatial barcode unique to a defined region or location on the array.

In some embodiments, a particular feature can contain capture probes including more than one spatial barcode (e.g., one capture probe at a particular feature can include a spatial barcode that is different than the spatial barcode included in another capture probe at the same particular feature, while both capture probes include a second, common spatial barcode), where
10 each spatial barcode corresponds to a particular defined region or location on the array. For example, multiple spatial barcode sequences associated with one particular feature on an array can provide a stronger address or attribution to a given location by providing duplicate or independent confirmation of the location. In some embodiments, the multiple spatial barcodes represent increasing specificity of the location of the particular array point. In a non-limiting
15 example, a particular array point can be coded with two different spatial barcodes, where each spatial barcode identifies a particular defined region within the array, and an array point possessing both spatial barcodes identifies the sub-region where two defined regions overlap, e.g., such as the overlapping portion of a Venn diagram.

In another non-limiting example, a particular array point can be coded with three
20 different spatial barcodes, where the first spatial barcode identifies a first region within the array, the second spatial barcode identifies a second region, where the second region is a subregion entirely within the first region, and the third spatial barcode identifies a third region, where the third region is a subregion entirely within the first and second subregions.

In some embodiments, capture probes attached to array features are released from the
25 array features for sequencing. Alternatively, in some embodiments, capture probes remain attached to the array features, and the probes are sequenced while remaining attached to the array features (e.g., via in-situ sequencing). Further aspects of the sequencing of capture probes are described in subsequent sections of this disclosure.

In some embodiments, an array feature can include different types of capture probes
30 attached to the feature. For example, the array feature can include a first type of capture probe with a capture domain designed to bind to one type of analyte, and a second type of capture

probe with a capture domain designed to bind to a second type of analyte. In general, array features can include one or more (e.g., two or more, three or more, four or more, five or more, six or more, eight or more, ten or more, 12 or more, 15 or more, 20 or more, 30 or more, 50 or more) different types of capture probes attached to a single array feature.

5 In some embodiments, the capture probe is nucleic acid. In some embodiments, the capture probe is attached to the array feature via its 5' end. In some embodiments, the capture probe includes from the 5' to 3' end: one or more barcodes (e.g., a spatial barcode and/or a UMI) and one or more capture domains. In some embodiments, the capture probe includes from the 5' to 3' end: one barcode (e.g., a spatial barcode or a UMI) and one capture domain. In some
10 embodiments, the capture probe includes from the 5' to 3' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), and a capture domain. In some embodiments, the capture probe includes from the 5' to 3' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), a second functional domain, and a capture domain. In some embodiments, the capture probe includes from
15 the 5' to 3' end: a cleavage domain, a functional domain, a spatial barcode, a UMI, and a capture domain. In some embodiments, the capture probe does not include a spatial barcode. In some embodiments, the capture probe does not include a UMI. In some embodiments, the capture probe includes a sequence for initiating a sequencing reaction.

 In some embodiments, the capture probe is immobilized on a feature via its 3' end. In
20 some embodiments, the capture probe includes from the 3' to 5' end: one or more barcodes (e.g., a spatial barcode and/or a UMI) and one or more capture domains. In some embodiments, the capture probe includes from the 3' to 5' end: one barcode (e.g., a spatial barcode or a UMI) and one capture domain. In some embodiments, the capture probe includes from the 3' to 5' end: a
25 cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), and a capture domain. In some embodiments, the capture probe includes from the 3' to 5' end: a cleavage domain, a functional domain, a spatial barcode, a UMI, and a capture domain.

 In some embodiments, a capture probe includes an in situ synthesized oligonucleotide. In some embodiments, the in situ synthesized oligonucleotide includes one or more constant sequences, one or more of which serves as a priming sequence (e.g., a primer for amplifying
30 target nucleic acids). In some embodiments, a constant sequence is a cleavable sequence. In some embodiments, the in situ synthesized oligonucleotide includes a barcode sequence, e.g., a

variable barcode sequence. In some embodiments, the in situ synthesized oligonucleotide is attached to a feature of an array.

In some embodiments, a capture probe is a product of two or more oligonucleotide sequences, e.g., two or more oligonucleotide sequences that are ligated together. In some
5 embodiments, one of the oligonucleotide sequences is an in situ synthesized oligonucleotide.

In some embodiments, the capture probe includes a splint oligonucleotide. Two or more oligonucleotides can be ligated together using a splint oligonucleotide and any variety of ligases known in the art or described herein (e.g., SplintR ligase).

In some embodiments, one of the oligonucleotides includes: a constant sequence (e.g., a
10 sequence complementary to a portion of a splint oligonucleotide), a degenerate sequence, and a capture domain (e.g., as described herein). In some embodiments, the capture probe is generated by having an enzyme add polynucleotides at the end of an oligonucleotide sequence. The capture probe can include a degenerate sequence, which can function as a unique molecular identifier.

A capture probe can include a degenerate sequence, which is a sequence in which some
15 positions of a nucleotide sequence contain a number of possible bases. A degenerate sequence can be a degenerate nucleotide sequence including about or at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, or 50 nucleotides. In some embodiments, a nucleotide sequence contains 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, or more degenerate positions within the nucleotide sequence. In some embodiments, the degenerate sequence is used
20 as a UMI.

In some embodiments, a capture probe includes a restriction endonuclease recognition sequence or a sequence of nucleotides cleavable by specific enzyme activities. For example, uracil sequences can be cleaved by specific enzyme activity. As another example, other modified bases (e.g., modified by methylation) can be recognized and cleaved by specific endonucleases.
25 The capture probes can be subjected to an enzymatic cleavage, which removes the blocking domain and any of the additional nucleotides that are added to the 3' end of the capture probe during the modification process. The removal of the blocking domain reveals and/or restores the free 3' end of the capture domain of the capture probe. In some embodiments, additional nucleotides can be removed to reveal and/or restore the 3' end of the capture domain of the
30 capture probe.

In some embodiments, a blocking domain can be incorporated into the capture probe when it is synthesized, or after its synthesis. The terminal nucleotide of the capture domain is a reversible terminator nucleotide (e.g., 3'-O-blocked reversible terminator and 3'-unblocked reversible terminator), and can be included in the capture probe during or after probe synthesis.

Extended Capture Probes

An "extended capture probe" is a capture probe with an enlarged nucleic acid sequence. For example, where the capture probe includes nucleic acid, an "extended 3' end" indicates that further nucleotides were added to the most 3' nucleotide of the capture probe to extend the length of the capture probe, for example, by standard polymerization reactions utilized to extend nucleic acid molecules including templated polymerization catalyzed by a polymerase (e.g., a DNA polymerase or reverse transcriptase).

In some embodiments, extending the capture probe includes generating cDNA from the captured (hybridized) RNA. This process involves synthesis of a complementary strand of the hybridized nucleic acid, e.g., generating cDNA based on the captured RNA template (the RNA hybridized to the capture domain of the capture probe). Thus, in an initial step of extending the capture probe, e.g., the cDNA generation, the captured (hybridized) nucleic acid, e.g., RNA, acts as a template for the extension, e.g., reverse transcription, step.

In some embodiments, the capture probe is extended using reverse transcription. For example, reverse transcription includes synthesizing cDNA (complementary or copy DNA) from RNA, e.g., (messenger RNA), using a reverse transcriptase. In some embodiments, reverse transcription is performed while the tissue is still in place, generating an analyte library, where the analyte library includes the spatial barcodes from the adjacent capture probes. In some embodiments, the capture probe is extended using one or more DNA polymerases.

In some embodiments, the capture domain of the capture probe includes a primer for producing the complementary strand of the nucleic acid hybridized to the capture probe, e.g., a primer for DNA polymerase and/or reverse transcription. The nucleic acid, e.g., DNA and/or cDNA, molecules generated by the extension reaction incorporate the sequence of the capture probe. The extension of the capture probe, e.g., a DNA polymerase and/or reverse transcription reaction, can be performed using a variety of suitable enzymes and protocols.

In some embodiments, a full-length DNA, e.g. cDNA, molecule is generated. In some embodiments, a "full-length" DNA molecule refers to the whole of the captured nucleic acid

molecule. However, if the nucleic acid, e.g. RNA, was partially degraded in the tissue sample, then the captured nucleic acid molecules will not be the same length as the initial RNA in the tissue sample. In some embodiments, the 3' end of the extended probes, e.g., first strand cDNA molecules, is modified. For example, a linker or adaptor can be ligated to the 3' end of the extended probes. This can be achieved using single stranded ligation enzymes such as T4 RNA
5 ligase or CircligaseTM (available from Epicentre Biotechnologies, Madison, WI). In some embodiments, template switching oligonucleotides are used to extend cDNA in order to generate a full-length cDNA (or as close to a full-length cDNA as possible). In some embodiments, a second strand synthesis helper probe (a partially double stranded DNA molecule capable of
10 hybridizing to the 3' end of the extended capture probe), can be ligated to the 3' end of the extended probe, e.g., first strand cDNA, molecule using a double stranded ligation enzyme such as T4 DNA ligase. Other enzymes appropriate for the ligation step are known in the art and include, e.g., Tth DNA ligase, Taq DNA ligase, *Thermococcus* sp. (strain 9^oN) DNA ligase (9^oNTM DNA ligase, New England Biolabs), AmpligaseTM (available from Epicentre
15 Biotechnologies, Madison, WI), and SplintR (available from New England Biolabs, Ipswich, MA). In some embodiments, a polynucleotide tail, e.g., a poly(A) tail, is incorporated at the 3' end of the extended probe molecules. In some embodiments, the polynucleotide tail is incorporated using a terminal transferase active enzyme.

In some embodiments, double-stranded extended capture probes are treated to remove
20 any unextended capture probes prior to amplification and/or analysis, e.g. sequence analysis. This can be achieved by a variety of methods, e.g., using an enzyme to degrade the unextended probes, such as an exonuclease enzyme, or purification columns.

In some embodiments, extended capture probes are amplified to yield quantities that are sufficient for analysis, e.g., via DNA sequencing. In some embodiments, the first strand of the
25 extended capture probes (e.g., DNA and/or cDNA molecules) acts as a template for the amplification reaction (e.g., a polymerase chain reaction).

In some embodiments, the amplification reaction incorporates an affinity group onto the extended capture probe (e.g., RNA-cDNA hybrid) using a primer including the affinity group. In some embodiments, the primer includes an affinity group and the extended capture probes
30 includes the affinity group. The affinity group can correspond to any of the affinity groups described previously.

In some embodiments, the extended capture probes including the affinity group can be coupled to an array feature specific for the affinity group. In some embodiments, the substrate can include an antibody or antibody fragment. In some embodiments, the array feature includes avidin or streptavidin and the affinity group includes biotin. In some embodiments, the array feature includes maltose and the affinity group includes maltose-binding protein. In some embodiments, the array feature includes maltose-binding protein and the affinity group includes maltose. In some embodiments, amplifying the extended capture probes can function to release the extended probes from the array feature, insofar as copies of the extended probes are not attached to the array feature.

In some embodiments, the extended capture probe or complement or amplicon thereof is released from an array feature. The step of releasing the extended capture probe or complement or amplicon thereof from an array feature can be achieved in a number of ways. In some embodiments, an extended capture probe or a complement thereof is released from the feature by nucleic acid cleavage and/or by denaturation (e.g. by heating to denature a double-stranded molecule).

In some embodiments, the extended capture probe or complement or amplicon thereof is released from the array feature by physical means. For example, methods for inducing physical release include denaturing double stranded nucleic acid molecules. Another method for releasing the extended capture probes is to use a solution that interferes with the hydrogen bonds of the double stranded molecules. In some embodiments, the extended capture probe is released by applying heated water such as water or buffer of at least 85°C, e.g., at least 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99°C. In some embodiments, a solution including salts, surfactants, etc. that can further destabilize the interaction between the nucleic acid molecules is added to release the extended capture probe from the array feature. In some embodiments, a formamide solution can be used to destabilize the interaction between nucleic acid molecules to release the extended capture probe from the array feature.

Analyte Capture Agents

This disclosure also provides methods and materials for using analyte capture agents for spatial profiling of biological analytes (e.g., mRNA, genomic DNA, accessible chromatin, and cell surface or intracellular proteins and/or metabolites). As used herein, an “analyte capture agent” (also referred to previously at times as a “cell labelling” agent”) refers to an agent that

interacts with an analyte (e.g., an analyte in a sample) and with a capture probe (e.g., a capture probe attached to a substrate) to identify the analyte. In some embodiments, the analyte capture agent includes an analyte binding moiety and a capture agent barcode domain.

FIG. 9 is a schematic diagram of an exemplary analyte capture agent **902** comprised of an analyte binding moiety **904** and a capture agent barcode domain **908**. An analyte binding moiety **904** is a molecule capable of binding to an analyte **906** and interacting with a spatially-barcoded capture probe. The analyte binding moiety can bind to the analyte **906** with high affinity and/or with high specificity. The analyte capture agent can include a capture agent barcode domain **908**, a nucleotide sequence (e.g., an oligonucleotide), which can hybridize to at least a portion or an entirety of a capture domain of a capture probe. The analyte binding moiety **904** can include a polypeptide and/or an aptamer (e.g., an oligonucleotide or peptide molecule that binds to a specific target analyte). The analyte binding moiety **904** can include an antibody or antibody fragment (e.g., an antigen-binding fragment).

As used herein, the term “analyte binding moiety” refers to a molecule or moiety capable of binding to a macromolecular constituent (e.g., an analyte, e.g., a biological analyte). In some embodiments of any of the spatial profiling methods described herein, the analyte binding moiety of the analyte capture agent that binds to a biological analyte can include, but is not limited to, an antibody, or an epitope binding fragment thereof, a cell surface receptor binding molecule, a receptor ligand, a small molecule, a bi-specific antibody, a bi-specific T-cell engager, a T-cell receptor engager, a B-cell receptor engager, a pro-body, an aptamer, a monobody, an affimer, a darpin, and a protein scaffold, or any combination thereof. The analyte binding moiety can bind to the macromolecular constituent (e.g., analyte) with high affinity and/or with high specificity. The analyte binding moiety can include a nucleotide sequence (e.g., an oligonucleotide), which can correspond to at least a portion or an entirety of the analyte binding moiety. The analyte binding moiety can include a polypeptide and/or an aptamer (e.g., a polypeptide and/or an aptamer that binds to a specific target molecule, e.g., an analyte). The analyte binding moiety can include an antibody or antibody fragment (e.g., an antigen-binding fragment) that binds to a specific analyte (e.g., a polypeptide).

In some embodiments, analyte capture agents are capable of binding to analytes present inside a cell. In some embodiments, analyte capture agents are capable of binding to cell surface analytes that can include, without limitation, a receptor, an antigen, a surface protein, a

transmembrane protein, a cluster of differentiation protein, a protein channel, a protein pump, a carrier protein, a phospholipid, a glycoprotein, a glycolipid, a cell-cell interaction protein complex, an antigen-presenting complex, a major histocompatibility complex, an engineered T-cell receptor, a T-cell receptor, a B-cell receptor, a chimeric antigen receptor, an extracellular
5 matrix protein, a posttranslational modification (e.g., phosphorylation, glycosylation, ubiquitination, nitrosylation, methylation, acetylation or lipidation) state of a cell surface protein, a gap junction, and an adherens junction. In some embodiments, the analyte capture agents are capable of binding to cell surface analytes that are post-translationally modified. In such
10 embodiments, analyte capture agents can be specific for cell surface analytes based on a given state of posttranslational modification (e.g., phosphorylation, glycosylation, ubiquitination, nitrosylation, methylation, acetylation or lipidation), such that a cell surface analyte profile can include posttranslational modification information of one or more analytes.

In some embodiments, the analyte capture agent includes a capture agent barcode domain that is conjugated or otherwise attached to the analyte binding moiety. In some embodiments, the
15 capture agent barcode domain is covalently-linked to the analyte binding moiety. In some embodiments, a capture agent barcode domain is a nucleic acid sequence. In some embodiments, a capture agent barcode domain includes an analyte binding moiety barcode and an analyte capture sequence.

As used herein, the term “analyte binding moiety barcode” refers to a barcode that is
20 associated with or otherwise identifies the analyte binding moiety. In some embodiments, by identifying an analyte binding moiety by identifying its associated analyte binding moiety barcode, the analyte to which the analyte binding moiety binds can also be identified. An analyte binding moiety barcode can be a nucleic acid sequence of a given length and/or sequence that is associated with the analyte binding moiety. An analyte binding moiety barcode can generally
25 include any of the variety of aspects of barcodes described herein. For example, an analyte capture agent that is specific to one type of analyte can have coupled thereto a first capture agent barcode domain (e.g., that includes a first analyte binding moiety barcode), while an analyte capture agent that is specific to a different analyte can have a different capture agent barcode domain (e.g., that includes a second barcode analyte binding moiety barcode) coupled thereto. In
30 some aspects, such a capture agent barcode domain can include an analyte binding moiety barcode that permits identification of the analyte binding moiety to which the capture agent

barcode domain is coupled. The selection of the capture agent barcode domain can allow significant diversity in terms of sequence, while also being readily attachable to most analyte binding moieties (e.g., antibodies) as well as being readily detected, (e.g., using sequencing or array technologies). In some embodiments, the analyte capture agents can include analyte
5 binding moieties with capture agent barcode domains attached to them. For example, an analyte capture agent can include a first analyte binding moiety (e.g., an antibody that binds to an analyte, e.g., a first cell surface feature) having associated with it a capture agent barcode domain that includes a first analyte binding moiety barcode.

In some embodiments, the capture agent barcode domain of an analyte capture agent
10 includes an analyte capture sequence. As used herein, the term “analyte capture sequence” refers to region or moiety of configured to hybridize to, bind to, couple to, or otherwise interact with a capture domain of a capture probe. In some embodiments, an analyte capture sequence includes a nucleic acid sequence that is complementary to or substantially complementary to the capture domain of a capture probe such that the analyte capture sequence hybridizes to the capture
15 domain of the capture probe. In some embodiments, an analyte capture sequence comprises a poly(A) nucleic acid sequence that hybridizes to a capture domain that comprises a poly(T) nucleic acid sequence. In some embodiments, an analyte capture sequence comprises a poly(T) nucleic acid sequence that hybridizes to a capture domain that comprises a poly(A) nucleic acid sequence. In some embodiments, an analyte capture sequence comprises a non-homopolymeric
20 nucleic acid sequence that hybridizes to a capture domain that comprises a non-homopolymeric nucleic acid sequence that is complementary (or substantially complementary) to the non-homopolymeric nucleic acid sequence of the analyte capture region.

In some embodiments of any of the spatial analysis methods described herein that employ an analyte capture agent, the capture agent barcode domain can be directly coupled to the analyte
25 binding moiety, or they can be attached to a bead, molecular lattice, e.g., a linear, globular, cross-linked, or other polymer, or other framework that is attached or otherwise associated with the analyte binding moiety, which allows attachment of multiple capture agent barcode domains to a single analyte binding moiety. Attachment (coupling) of the capture agent barcode domains to the analyte binding moieties can be achieved through any of a variety of direct or indirect,
30 covalent or non-covalent associations or attachments. For example, in the case of a capture agent barcode domain coupled to an analyte binding moiety that includes an antibody or antigen-

binding fragment, such capture agent barcode domains can be covalently attached to a portion of the antibody or antigen-binding fragment using chemical conjugation techniques (e.g., Lightning-Link® antibody labelling kits available from Innova Biosciences). In some embodiments, a capture agent barcode domain can be coupled to an antibody or antigen-binding
5 fragment using non-covalent attachment mechanisms (e.g., using biotinylated antibodies and oligonucleotides or beads that include one or more biotinylated linker, coupled to oligonucleotides with an avidin or streptavidin linker.) Antibody and oligonucleotide biotinylation techniques can be used, and are described for example in Fang et al., *Nucleic Acids Res.* (2003), 31(2): 708-715, the entire contents of which are incorporated by reference herein.

10 Likewise, protein and peptide biotinylation techniques have been developed and can be used, and are described for example in U.S. Patent No. 6,265,552, the entire contents of which are incorporated by reference herein. Furthermore, click reaction chemistry such as a methyltetrazine-PEG5-NHS ester reaction, a TCO-PEG4-NHS ester reaction, or the like, can be used to couple capture agent barcode domains to analyte binding moieties. The reactive moiety
15 on the analyte binding moiety can also include amine for targeting aldehydes, amine for targeting maleimide (e.g., free thiols), azide for targeting click chemistry compounds (e.g., alkynes), biotin for targeting streptavidin, phosphates for targeting EDC, which in turn targets active ester (e.g., NH₂). The reactive moiety on the analyte binding moiety can be a chemical compound or group that binds to the reactive moiety on the analyte binding moiety. Exemplary strategies to
20 conjugate the analyte binding moiety to the capture agent barcode domain include the use of commercial kits (e.g., Solulink, Thunder link), conjugation of mild reduction of hinge region and maleimide labelling, stain-promoted click chemistry reaction to labeled amides (e.g., copper-free), and conjugation of periodate oxidation of sugar chain and amine conjugation. In the cases where the analyte binding moiety is an antibody, the antibody can be modified prior to or
25 contemporaneously with conjugation of the oligonucleotide. For example, the antibody can be glycosylated with a substrate-permissive mutant of β -1,4-galactosyltransferase, GalT (Y289L) and azide-bearing uridine diphosphate-N-acetylgalactosamine analog uridine diphosphate - GalNAz. The modified antibody can be conjugated to an oligonucleotide with a dibenzocyclooctyne-PEG4-NHS group. In some embodiments, certain steps (e.g., COOH
30 activation (e.g., EDC) and homobifunctional cross linkers) can be avoided to prevent the analyte binding moieties from conjugating to themselves. In some embodiments of any of the spatial

profiling methods described herein, the analyte capture agent (e.g., analyte binding moiety coupled to an oligonucleotide) can be delivered into the cell, e.g., by transfection (e.g., using transfectamine, cationic polymers, calcium phosphate or electroporation), by transduction (e.g., using a bacteriophage or recombinant viral vector), by mechanical delivery (e.g., magnetic
5 beads), by lipid (e.g., 1,2-Dioleoyl-sn-glycero-3-phosphocholine (DOPC)), or by transporter proteins. An analyte capture agent can be delivered into a cell using exosomes. For example, a first cell can be generated that releases exosomes comprising an analyte capture agent. An analyte capture agent can be attached to an exosome membrane. An analyte capture agent can be contained within the cytosol of an exosome. Released exosomes can be harvested and provided
10 to a second cell, thereby delivering the analyte capture agent into the second cell. An analyte capture agent can be releasable from an exosome membrane before, during, or after delivery into a cell. In some embodiments, the cell is permeabilized to allow the analyte capture agent to couple with intracellular cellular constituents (such as, without limitation, intracellular proteins, metabolites and nuclear membrane proteins). Following intracellular delivery, analyte capture
15 agents can be used to analyze intracellular constituents as described herein.

In some embodiments of any of the spatial profiling methods described herein, the capture agent barcode domain coupled to an analyte capture agent can include modifications that render it non-extendable by a polymerase. In some embodiments, when binding to a capture domain of a capture probe or nucleic acid in a sample for a primer extension reaction, the capture
20 agent barcode domain can serve as a template, not a primer. When the capture agent barcode domain also includes a barcode (e.g., an analyte binding moiety barcode), such a design can increase the efficiency of molecular barcoding by increasing the affinity between the capture agent barcode domain and unbarcoded sample nucleic acids, and eliminate the potential formation of adaptor artifacts. In some embodiments, the capture agent barcode domain can
25 include a random N-mer sequence that is capped with modifications that render it non-extendable by a polymerase. In some cases, the composition of the random N-mer sequence can be designed to maximize the binding efficiency to free, unbarcoded ssDNA molecules. The design can include a random sequence composition with a higher GC content, a partial random sequence with fixed G or C at specific positions, the use of guanosines, the use of locked nucleic
30 acids, or any combination thereof.

A modification for blocking primer extension by a polymerase can be a carbon spacer group of different lengths or a dideoxynucleotide. In some embodiments, the modification can be an abasic site that has an apurine or apyrimidine structure, a base analog, or an analogue of a phosphate backbone, such as a backbone of N-(2-aminoethyl)-glycine linked by amide bonds, tetrahydrofuran, or 1', 2'-Dideoxyribose. The modification can also be a uracil base, 2'OMe modified RNA, C3-18 spacers (e.g., structures with 3-18 consecutive carbon atoms, such as C3 spacer), ethylene glycol multimer spacers (e.g., spacer 18 (hexa-ethyleneglycol spacer), biotin, di-deoxynucleotide triphosphate, ethylene glycol, amine, or phosphate.

In some embodiments of any of the spatial profiling methods described herein, the capture agent barcode domain coupled to the analyte binding moiety includes a cleavable domain. For example, after the analyte capture agent binds to an analyte (e.g., a cell surface analyte), the capture agent barcode domain can be cleaved and collected for downstream analysis according to the methods as described herein. In some embodiments, the cleavable domain of the capture agent barcode domain includes a U-excising element that allows the species to release from the bead. In some embodiments, the U-excising element can include a single-stranded DNA (ssDNA) sequence that contains at least one uracil. The species can be attached to a bead via the ssDNA sequence. The species can be released by a combination of uracil-DNA glycosylase (e.g., to remove the uracil) and an endonuclease (e.g., to induce an ssDNA break). If the endonuclease generates a 5' phosphate group from the cleavage, then additional enzyme treatment can be included in downstream processing to eliminate the phosphate group, e.g., prior to ligation of additional sequencing handle elements, e.g., Illumina full P5 sequence, partial P5 sequence, full R1 sequence, and/or partial R1 sequence.

In some embodiments, an analyte binding moiety of an analyte capture agent includes one or more antibodies or antigen binding fragments thereof. The antibodies or antigen binding fragments including the analyte binding moiety can specifically bind to a target analyte. In some embodiments, the analyte is a protein (e.g., a protein on a surface of the biological sample (e.g., a cell) or an intracellular protein). In some embodiments, a plurality of analyte capture agents comprising a plurality of analyte binding moieties bind a plurality of analytes present in a biological sample. In some embodiments, the plurality of analytes includes a single species of analyte (e.g., a single species of polypeptide). In some embodiments in which the plurality of analytes includes a single species of analyte, the analyte binding moieties of the plurality of

analyte capture agents are the same. In some embodiments in which the plurality of analytes includes a single species of analyte, the analyte binding moieties of the plurality of analyte capture agents are the different (e.g., members of the plurality of analyte capture agents can have two or more species of analyte binding moieties, wherein each of the two or more species of analyte binding moieties binds a single species of analyte, e.g., at different binding sites). In some embodiments, the plurality of analytes includes multiple different species of analyte (e.g., multiple different species of polypeptides).

In some embodiments, multiple different species of analytes (e.g., polypeptides) from the biological sample can be subsequently associated with the one or more physical properties of the biological sample. For example, the multiple different species of analytes can be associated with locations of the analytes in the biological sample. Such information (e.g., proteomic information when the analyte binding moiety(ies) recognizes a polypeptide(s)) can be used in association with other spatial information (e.g., genetic information from the biological sample, such as DNA sequence information, transcriptome information (i.e., sequences of transcripts), or both).

For example, a cell surface protein of a cell can be associated with one or more physical properties of the cell (e.g., a shape, size, activity, or a type of the cell). The one or more physical properties can be characterized by imaging the cell. The cell can be bound by an analyte capture agent comprising an analyte binding moiety that binds to the cell surface protein and an analyte binding moiety barcode that identifies that analyte binding moiety, and the cell can be subjected to spatial analysis (e.g., any of the variety of spatial analysis methods described herein). For example, the analyte capture agent bound to the cell surface protein can be bound to a capture probe (e.g., a capture probe on an array), which capture probe includes a capture domain that interacts with an analyte capture sequence present on the capture agent barcode domain of the analyte capture agent. All or part of the capture agent barcode domain (including the analyte binding moiety barcode) can be copied with a polymerase using a 3' end of the capture domain as a priming site, generating an extended capture probe that includes the all or part of the capture probe (including a spatial barcode present on the capture probe) and a copy of the analyte binding moiety barcode. In some embodiments, the spatial array with the extended capture probe(s) can be contacted with a sample, where the analyte capture agent(s) associated with the spatial array capture the target analyte(s). The analyte capture agent(s) containing the extended capture probe(s), which includes the spatial barcode(s) of the capture probe(s) and the analyte

binding moiety barcode(s), can then be denatured from the capture probe(s) of the spatial array. This allows the spatial array to be reused. The sample can be dissociated into non-aggregated cells (e.g. single cells) and analyzed by the single cell / droplet methods described herein. The extended capture probe can be sequenced to obtain a nucleic acid sequence, in which the spatial
 5 barcode of the capture probe is associated with the analyte binding moiety barcode of the analyte capture agent. The nucleic acid sequence of the extended capture probe can thus be associated with the analyte (e.g., cell surface protein), and in turn, with the one or more physical properties of the cell (e.g., a shape or cell type). In some embodiments, the nucleic acid sequence of the extended capture probe can be associated with an intracellular analyte of a nearby cell, where the
 10 intracellular analyte was released using any of the cell permeabilization or analyte migration techniques described herein.

In some embodiments of any of the spatial profiling methods described herein, the capture agent barcode domains released from the analyte capture agents can then be subjected to sequence analysis to identify which analyte capture agents were bound to analytes. Based upon
 15 the capture agent barcode domains that are associated with a feature (e.g., a feature at a particular location) on a spatial array and the presence of the analyte binding moiety barcode sequence, an analyte profile can be created for a biological sample. Profiles of individual cells or populations of cells can be compared to profiles from other cells, e.g., ‘normal’ cells, to identify variations in analytes, which can provide diagnostically relevant information. In some embodiments, these
 20 profiles can be useful in the diagnosis of a variety of disorders that are characterized by variations in cell surface receptors, such as cancer and other disorders.

FIG. 10 is a schematic diagram depicting an exemplary interaction between a feature-immobilized capture probe **1024** and an analyte capture agent **1026**. The feature-immobilized capture probe **1024** can include a spatial barcode **1008** as well as one or more functional
 25 sequences **1006** and **1010**, as described elsewhere herein. The capture probe can also include a capture domain **1012** that is capable of binding to an analyte capture agent **1026**. The analyte capture agent **1026** can include a functional sequence **1018**, capture agent barcode domain **1016**, and an analyte capture sequence **1014** that is capable of binding to the capture domain **1012** of the capture probe **1024**. The analyte capture agent can also include a linker **1020** that allows the
 30 capture agent barcode domain **1016** to couple to the analyte binding moiety **1022**.

In some embodiments of any of the spatial profiling methods described herein, the methods are used to identify immune cell profiles. Immune cells express various adaptive immunological receptors relating to immune function, such as T cell receptors (TCRs) and B cell receptors (BCRs). T cell receptors and B cell receptors play a part in the immune response by specifically recognizing and binding to antigens and aiding in their destruction.

The T cell receptor, or TCR, is a molecule found on the surface of T cells that is generally responsible for recognizing fragments of antigen as peptides bound to major histocompatibility complex (MHC) molecules. The TCR is generally a heterodimer of two chains, each of which is a member of the immunoglobulin superfamily, possessing an N-terminal variable (V) domain, and a C terminal constant domain. In humans, in 95% of T cells, the TCR consists of an alpha (α) and beta (β) chain, whereas in 5% of T cells, the TCR consists of gamma and delta (γ/δ) chains. This ratio can change during ontogeny and in diseased states as well as in different species. When the TCR engages with antigenic peptide and MHC (peptide/MHC or pMHC), the T lymphocyte is activated through signal transduction.

Each of the two chains of a TCR contains multiple copies of gene segments – a variable ‘V’ gene segment, a diversity ‘D’ gene segment, and a joining ‘J’ gene segment. The TCR alpha chain (TCRa) is generated by recombination of V and J segments, while the beta chain (TCRb) is generated by recombination of V, D, and J segments. Similarly, generation of the TCR gamma chain involves recombination of V and J gene segments, while generation of the TCR delta chain occurs by recombination of V, D, and J gene segments. The intersection of these specific regions (V and J for the alpha or gamma chain, or V, D and J for the beta or delta chain) corresponds to the CDR3 region that is important for antigen-MHC recognition. Complementarity determining regions (*e.g.*, CDR1, CDR2, and CDR3), or hypervariable regions, are sequences in the variable domains of antigen receptors (*e.g.*, T cell receptor and immunoglobulin) that can complement an antigen. Most of the diversity of CDRs is found in CDR3, with the diversity being generated by somatic recombination events during the development of T lymphocytes. A unique nucleotide sequence that arises during the gene arrangement process can be referred to as a clonotype.

The B cell receptor, or BCR, is a molecule found on the surface of B cells. The antigen binding portion of a BCR is composed of a membrane-bound antibody that, like most antibodies (*e.g.*, immunoglobulins), has a unique and randomly determined antigen-binding site. The antigen binding portion of a BCR includes membrane-bound immunoglobulin molecule of one

isotype (*e.g.*, IgD, IgM, IgA, IgG, or IgE). When a B cell is activated by its first encounter with a cognate antigen, the cell proliferates and differentiates to generate a population of antibody-secreting plasma B cells and memory B cells. The various immunoglobulin isotypes differ in their biological features, structure, target specificity and distribution. A variety of molecular mechanisms exist to generate initial diversity, including genetic recombination at multiple sites.

The BCR is composed of two genes IgH and IgK (or IgL) coding for antibody heavy and light chains. Immunoglobulins are formed by recombination among gene segments, sequence diversification at the junctions of these segments, and point mutations throughout the gene. Each heavy chain gene contains multiple copies of three different gene segments – a variable ‘V’ gene segment, a diversity ‘D’ gene segment, and a joining ‘J’ gene segment. Each light chain gene contains multiple copies of two different gene segments for the variable region of the protein – a variable ‘V’ gene segment and a joining ‘J’ gene segment.

The recombination can generate a molecule with one of each of the V, D, and J segments. Furthermore, several bases can be deleted and others added (called N and P nucleotides) at each of the two junctions, thereby generating further diversity. After B cell activation, a process of affinity maturation through somatic hypermutation occurs. In this process, progeny cells of the activated B cells accumulate distinct somatic mutations throughout the gene with higher mutation concentration in the CDR regions leading to the generation of antibodies with higher affinity to the antigens.

In addition to somatic hypermutation, activated B cells undergo the process of isotype switching. Antibodies with the same variable segments can have different forms (isotypes) depending on the constant segment. Whereas all naïve B cells express IgM (or IgD), activated B cells mostly express IgG but also IgM, IgA and IgE. This expression switching from IgM (and/or IgD) to IgG, IgA, or IgE occurs through a recombination event causing one cell to specialize in producing a specific isotype. A unique nucleotide sequence that arises during the gene arrangement process can similarly be referred to as a clonotype.

Certain methods described herein are utilized to analyze the various sequences of TCRs and BCRs from immune cells, for example, various clonotypes. In some embodiments, the methods are used to analyze the sequence of a TCR alpha chain, a TCR beta chain, a TCR delta chain, a TCR gamma chain, or any fragment thereof (*e.g.*, variable regions including V(D)J or VJ regions, constant regions, transmembrane regions, fragments thereof, combinations thereof,

and combinations of fragments thereof). In some embodiments, the methods described herein can be used to analyze the sequence of a B cell receptor heavy chain, B cell receptor light chain, or any fragment thereof (e.g., variable regions including V(D)J or VJ regions, constant regions, transmembrane regions, fragments thereof, combinations thereof, and combinations of fragments thereof).

Where immune cells are to be analyzed, primer sequences useful in any of the various operations for attaching barcode sequences and/or amplification reactions can include gene specific sequences which target genes or regions of genes of immune cell proteins, for example immune receptors. Such gene sequences include, but are not limited to, sequences of various T cell receptor alpha variable genes (TRAV genes), T cell receptor alpha joining genes (TRAJ genes), T cell receptor alpha constant genes (TRAC genes), T cell receptor beta variable genes (TRBV genes), T cell receptor beta diversity genes (TRBD genes), T cell receptor beta joining genes (TRBJ genes), T cell receptor beta constant genes (TRBC genes), T cell receptor gamma variable genes (TRGV genes), T cell receptor gamma joining genes (TRGJ genes), T cell receptor gamma constant genes (TRGC genes), T cell receptor delta variable genes (TRDV genes), T cell receptor delta diversity genes (TRDD genes), T cell receptor delta joining genes (TRDJ genes), and T cell receptor delta constant genes (TRDC genes).

In some embodiments, the analyte binding moiety is based on the Major Histocompatibility Complex (MHC) class I or class II. In some embodiments, the analyte binding moiety is an MHC multimer including, without limitation, MHC dextramers, MHC tetramers, and MHC pentamers (see, for example, U.S. Patent Application Publication Nos. US 2018/0180601 and US 2017/0343545, the entire contents of each of which are incorporated herein by reference. MHCs (e.g., a soluble MHC monomer molecule), including full or partial MHC-peptides, can be used as analyte binding moieties of analyte capture agents that are coupled to capture agent barcode domains that include an analyte binding moiety barcode that identifies its associated MHC (and, thus, for example, the MHC's TCR binding partner). In some embodiments, MHCs are used to analyze one or more cell-surface features of a T-cell, such as a TCR. In some cases, multiple MHCs are associated together in a larger complex (MHC multimer) to improve binding affinity of MHCs to TCRs via multiple ligand binding synergies.

FIG. 11A, 11B, and 11C are schematics illustrating how streptavidin cell tags can be utilized in an array-based system to produce a spatially-barcoded cell or cellular contents. For

example, as shown in **FIG. 11**, peptide-bound major histocompatibility complex (pMHCs) can be individually associated with biotin and bound to a streptavidin moiety such that the streptavidin moiety comprises multiple pMHC moieties. Each of these moieties can bind to a TCR such that the streptavidin binds to a target T-cell via multiple MCH/TCR binding interactions. Multiple interactions synergize and can substantially improve binding affinity. Such improved affinity can improve labelling of T-cells and also reduce the likelihood that labels will dissociate from T-cell surfaces. As shown in **FIG. 11B**, a capture agent barcode domain **1101** can be modified with streptavidin **1102** and contacted with multiple molecules of biotinylated MHC **1103** (such as a pMHC) such that the biotinylated MHC **1103** molecules are coupled with the streptavidin conjugated capture agent barcode domain **1101**. The result is a barcoded MHC multimer complex **1105**. As shown in **FIG. 11B**, the capture agent barcode domain sequence **1101** can identify the MHC as its associated label and also includes optional functional sequences such as sequences for hybridization with other oligonucleotides. As shown in **FIG. 11C**, one example oligonucleotide is capture probe **1106** that comprises a complementary sequence (e.g., rGrGrG corresponding to C C C), a barcode sequence and other functional sequences, such as, for example, a UMI, an adapter sequence (e.g., comprising a sequencing primer sequence (e.g., R1 or a partial R1 ("pR1")), a flow cell attachment sequence (e.g., P5 or P7 or partial sequences thereof)), etc. In some cases, capture probe **1106** may at first be associated with a feature (e.g., a gel bead) and released from the feature. In other embodiments, capture probe **1106** can hybridize with a capture agent barcode domain **1101** of the MHC-oligonucleotide complex **1105**. The hybridized oligonucleotides (Spacer C C C and Spacer rGrGrG) can then be extended in primer extension reactions such that constructs comprising sequences that correspond to each of the two spatial barcode sequences (the spatial barcode associated with the capture probe, and the barcode associated with the MHC-oligonucleotide complex) are generated. In some cases, one or both of these corresponding sequences may be a complement of the original sequence in capture probe **1106** or capture agent barcode domain **1101**. In other embodiments, the capture probe and the capture agent barcode domain are ligated together. The resulting constructs can be optionally further processed (e.g., to add any additional sequences and/or for clean-up) and subjected to sequencing. As described elsewhere herein, a sequence derived from the capture probe **1106** spatial barcode sequence may be used to identify a feature and the sequence derived from spatial barcode sequence on the capture agent barcode

domain **1101** may be used to identify the particular peptide MHC complex **1104** bound on the surface of the cell (e.g., when using MHC-peptide libraries for screening immune cells or immune cell populations).

(c) Substrate

5 For the spatial array-based analytical methods described in this section, the substrate functions as a support for direct or indirect attachment of capture probes to features of the array. In addition, in some embodiments, a substrate (e.g., the same substrate or a different substrate) can be used to provide support to a biological sample, particularly, for example, a thin tissue section. Accordingly, a “substrate” is a support that is insoluble in aqueous liquid and which
10 allows for positioning of biological samples, analytes, features, and/or capture probes on the substrate.

A wide variety of different substrates can be used for the foregoing purposes. In general, a substrate can be any suitable support material. Exemplary substrates include, but are not limited to, glass, modified and/or functionalized glass, hydrogels, films, membranes, plastics
15 (including e.g., acrylics, polystyrene, copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, Teflon™, cyclic olefins, polyimides etc.), nylon, ceramics, resins, Zeonor, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, optical fiber bundles, and polymers, such as polystyrene, cyclic olefin copolymers (COCs), cyclic olefin polymers (COPs), polypropylene, polyethylene
20 and polycarbonate.

The substrate can also correspond to a flow cell. Flow cells can be formed of any of the foregoing materials, and can include channels that permit reagents, solvents, features, and molecules to pass through the cell.

Among the examples of substrate materials discussed above, polystyrene is a
25 hydrophobic material suitable for binding negatively charged macromolecules because it normally contains few hydrophilic groups. For nucleic acids immobilized on glass slides, by increasing the hydrophobicity of the glass surface the nucleic acid immobilization can be increased. Such an enhancement can permit a relatively more densely packed formation (e.g., provide improved specificity and resolution).

In some embodiments, a substrate is coated with a surface treatment such as poly(L)-lysine. Additionally or alternatively, the substrate can be treated by silanation, e.g. with epoxy-silane, amino-silane, and/or by a treatment with polyacrylamide.

The substrate can generally have any suitable form or format. For example, the substrate
5 can be flat, curved, e.g. convexly or concavely curved towards the area where the interaction between a biological sample, e.g. tissue sample, and the substrate takes place. In some embodiments, the substrate is a flat, e.g., planar, chip or slide. The substrate can contain one or more patterned surfaces within the substrate (e.g., channels, wells, projections, ridges, divots, etc.).

10 A substrate can be of any desired shape. For example, a substrate can be typically a thin, flat shape (e.g., a square or a rectangle). In some embodiments, a substrate structure has rounded corners (e.g., for increased safety or robustness). In some embodiments, a substrate structure has one or more cut-off corners (e.g., for use with a slide clamp or cross-table). In some embodiments, where a substrate structure is flat, the substrate structure can be any appropriate
15 type of support having a flat surface (e.g., a chip or a slide such as a microscope slide).

Substrates can optionally include various structures such as, but not limited to, projections, ridges, and channels. A substrate can be micropatterned to limit lateral diffusion (e.g., to prevent overlap of spatial barcodes). A substrate modified with such structures can be modified to allow association of analytes, features (e.g., beads), or probes at individual sites. For
20 example, the sites where a substrate is modified with various structures can be contiguous or non-contiguous with other sites.

In some embodiments, the surface of a substrate can be modified so that discrete sites are formed that can only have or accommodate a single feature. In some embodiments, the surface of a substrate can be modified so that features adhere to random sites.

25 In some embodiments, the surface of a substrate is modified to contain one or more wells, using techniques such as (but not limited to) stamping techniques, microetching techniques, and molding techniques. In some embodiments in which a substrate includes one or more wells, the substrate can be a concavity slide or cavity slide. For example, wells can be formed by one or more shallow depressions on the surface of the substrate. In some embodiments, where a
30 substrate includes one or more wells, the wells can be formed by attaching a cassette (e.g., a cassette containing one or more chambers) to a surface of the substrate structure.

In some embodiments, the structures of a substrate (e.g., wells) can each bear a different capture probe. Different capture probes attached to each structure can be identified according to the locations of the structures in or on the surface of the substrate. Exemplary substrates include arrays in which separate structures are located on the substrate including, for example, those
5 having wells that accommodate features.

In some embodiments, a substrate includes one or more markings on a surface of the substrate, e.g., to provide guidance for correlating spatial information with the characterization of the analyte of interest. For example, a substrate can be marked with a grid of lines (e.g., to allow the size of objects seen under magnification to be easily estimated and/or to provide reference
10 areas for counting objects). In some embodiments, fiducial markers can be included on the substrate. Such markings can be made using techniques including, but not limited to, printing, sand-blasting, and depositing on the surface.

In some embodiments where the substrate is modified to contain one or more structures, including but not limited to wells, projections, ridges, or markings, the structures can include
15 physically altered sites. For example, a substrate modified with various structures can include physical properties, including, but not limited to, physical configurations, magnetic or compressive forces, chemically functionalized sites, chemically altered sites, and/or electrostatically altered sites.

In some embodiments where the substrate is modified to contain various structures, including but not limited to wells, projections, ridges, or markings, the structures are applied in a
20 pattern. Alternatively, the structures can be randomly distributed.

In some embodiments, a substrate is treated in order to minimize or reduce non-specific analyte hybridization within or between features. For example, treatment can include coating the substrate with a hydrogel, film, and/or membrane that creates a physical barrier to non-specific
25 hybridization. Any suitable hydrogel can be used. For example, hydrogel matrices prepared according to the methods set forth in U.S. Patent Nos. 6,391,937, 9,512,422, and 9,889,422, and U.S. Patent Application Publication Nos. U.S. 2017/0253918 and U.S. 2018/0052081, can be used. The entire contents of each of the foregoing documents are incorporated herein by reference.

Treatment can include adding a functional group that is reactive or capable of being
30 activated such that it becomes reactive after receiving a stimulus (e.g., photoreactive). Treatment

can include treating with polymers having one or more physical properties (e.g., mechanical, electrical, magnetic, and/or thermal) that minimize non-specific binding (e.g., that activate a substrate at certain locations to allow analyte hybridization at those locations).

5 The substrate (e.g., a bead or a feature on an array) can include tens to hundreds of thousands or millions of individual oligonucleotide molecules (e.g., at least about 10,000, 50,000, 100,000, 500,000, 1,000,000, 10,000,000, 100,000,000, 1,000,000,000, or 10,000,000,000 oligonucleotide molecules).

10 In some embodiments, the surface of the substrate is coated with a cell-permissive coating to allow adherence of live cells. A “cell-permissive coating” is a coating that allows or helps cells to maintain cell viability (e.g., remain viable) on the substrate. For example, a cell-permissive coating can enhance cell attachment, cell growth, and/or cell differentiation, e.g., a cell-permissive coating can provide nutrients to the live cells. A cell-permissive coating can include a biological material and/or a synthetic material. Non-limiting examples of a cell-permissive coating include coatings that feature one or more extracellular matrix (ECM) components (e.g., proteoglycans and fibrous proteins such as collagen, elastin, fibronectin and laminin), poly-lysine, poly(L)-ornithine, and/or a biocompatible silicone (e.g., CYTOSOFT®). For example, a cell-permissive coating that includes one or more extracellular matrix components can include collagen Type I, collagen Type II, collagen Type IV, elastin, fibronectin, laminin, and/or vitronectin. In some embodiments, the cell-permissive coating includes a solubilized basement membrane preparation extracted from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma (e.g., MATRIGEL®). In some embodiments, the cell-permissive coating includes collagen. A cell-permissive coating can be used to culture adherent cells on a spatially-barcoded array, or to maintain cell viability of a tissue sample or section while in contact with a spatially-barcoded array.

25 Where the substrate includes a gel (e.g., a hydrogel or gel matrix), oligonucleotides within the gel can attach to the substrate. The terms “hydrogel” and “hydrogel matrix” are used interchangeably herein to refer to a macromolecular polymer gel including a network. Within the network, some polymer chains can optionally be cross-linked, although cross-linking does not always occur.

30 In some embodiments, a hydrogel can include hydrogel subunits. A “hydrogel subunit” is a hydrophilic monomer, a molecular precursor, or a polymer that can be polymerized (e.g., cross-

linked) to form a three-dimensional (3D) hydrogel network. The hydrogel subunits can include any convenient hydrogel subunits, such as, but not limited to, acrylamide, bis-acrylamide, polyacrylamide and derivatives thereof, poly(ethylene glycol) and derivatives thereof (e.g. PEG-acrylate (PEG-DA), PEG-RGD), gelatin-methacryloyl (GelMA), methacrylated hyaluronic acid (MeHA),
5 polyaliphatic polyurethanes, polyether polyurethanes, polyester polyurethanes, polyethylene copolymers, polyamides, polyvinyl alcohols, polypropylene glycol, polytetramethylene oxide, polyvinyl pyrrolidone, polyacrylamide, poly(hydroxyethyl acrylate), and poly(hydroxyethyl methacrylate), collagen, hyaluronic acid, chitosan, dextran, agarose, gelatin, alginate, protein polymers, methylcellulose, and the like, and combinations thereof.

10 In some embodiments, a hydrogel includes a hybrid material, e.g., the hydrogel material includes elements of both synthetic and natural polymers. Examples of suitable hydrogels are described, for example, in U.S. Patent Nos. 6,391,937, 9,512,422, and 9,889,422, and in U.S. Patent Application Publication Nos. 2017/0253918, 2018/0052081 and 2010/0055733, the entire contents of each of which are incorporated herein by reference.

15 In some embodiments, cross-linkers and/or initiators are added to hydrogel subunits. Examples of cross-linkers include, without limitation, bis-acrylamide and diazirine. Examples of initiators include, without limitation, azobisisobutyronitrile (AIBN), riboflavin, and L-arginine. Inclusion of cross-linkers and/or initiators can lead to increased covalent bonding between interacting biological macromolecules in later polymerization steps.

20 In some embodiments, hydrogels can have a colloidal structure, such as agarose, or a polymer mesh structure, such as gelatin.

In some embodiments, some hydrogel subunits are polymerized (e.g., undergo “formation”) covalently or physically cross-linked, to form a hydrogel network. For example, hydrogel subunits can be polymerized by any method including, but not limited to, thermal
25 crosslinking, chemical crosslinking, physical crosslinking, ionic crosslinking, photo-crosslinking, irradiative crosslinking (e.g., x-ray, electron beam), and combinations thereof. Techniques such as lithographic photopolymerization can also be used to form hydrogels.

Polymerization methods for hydrogel subunits can be selected to form hydrogels with different properties (e.g., pore size, swelling properties, biodegradability, conduction,
30 transparency, and/or permeability of the hydrogel). For example, a hydrogel can include pores of sufficient size to allow the passage of macromolecules, (e.g., nucleic acids, proteins, chromatin,

metabolites, gRNA, antibodies, carbohydrates, peptides, metabolites, and/or small molecules) into the sample (e.g., tissue section). It is known that pore size generally decreases with increasing concentration of hydrogel subunits and generally increases with an increasing ratio of hydrogel subunits to crosslinker. Therefore, a fixative/hydrogel composition can be prepared that includes a concentration of hydrogel subunits that allows the passage of such biological macromolecules.

In some embodiments, the hydrogel can form the substrate. In some embodiments, the substrate includes a hydrogel and one or more second materials. In some embodiments, the hydrogel is placed on top of one or more second materials. For example, the hydrogel can be pre-formed and then placed on top of, underneath, or in any other configuration with one or more second materials. In some embodiments, hydrogel formation occurs after contacting one or more second materials during formation of the substrate. Hydrogel formation can also occur within a structure (e.g., wells, ridges, projections, and/or markings) located on a substrate.

In some embodiments, hydrogel formation on a substrate occurs before, contemporaneously with, or after features (e.g., beads) are attached to the substrate. For example, when a capture probe is attached (e.g., directly or indirectly) to a substrate, hydrogel formation can be performed on the substrate already containing the capture probes.

In some embodiments, hydrogel formation occurs within a biological sample. In some embodiments, a biological sample (e.g., tissue section) is embedded in a hydrogel. In some embodiments, hydrogel subunits are infused into the biological sample, and polymerization of the hydrogel is initiated by an external or internal stimulus.

In embodiments in which a hydrogel is formed within a biological sample, functionalization chemistry can be used. In some embodiments, functionalization chemistry includes hydrogel-tissue chemistry (HTC). Any hydrogel-tissue backbone (e.g., synthetic or native) suitable for HTC can be used for anchoring biological macromolecules and modulating functionalization. Non-limiting examples of methods using HTC backbone variants include CLARITY, PACT, ExM, SWITCH and ePACT. In some embodiments, hydrogel formation within a biological sample is permanent. For example, biological macromolecules can permanently adhere to the hydrogel allowing multiple rounds of interrogation. In some embodiments, hydrogel formation within a biological sample is reversible.

In some embodiments, additional reagents are added to the hydrogel subunits before, contemporaneously with, and/or after polymerization. For example, additional reagents can include but are not limited to oligonucleotides (e.g., capture probes), endonucleases to fragment DNA, fragmentation buffer for DNA, DNA polymerase enzymes, dNTPs used to amplify the nucleic acid and to attach the barcode to the amplified fragments. Other enzymes can be used, including without limitation, RNA polymerase, transposase, ligase, proteinase K, and DNase. Additional reagents can also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers, and switch oligonucleotides. In some embodiments, optical labels are added to the hydrogel subunits before, contemporaneously with, and/or after polymerization.

In some embodiments, HTC reagents are added to the hydrogel before, contemporaneously with, and/or after polymerization. In some embodiments, a cell labelling agent is added to the hydrogel before, contemporaneously with, and/or after polymerization. In some embodiments, a cell-penetrating agent is added to the hydrogel before, contemporaneously with, and/or after polymerization.

Hydrogels embedded within biological samples can be cleared using any suitable method. For example, electrophoretic tissue clearing methods can be used to remove biological macromolecules from the hydrogel-embedded sample. In some embodiments, a hydrogel-embedded sample is stored before or after clearing of hydrogel, in a medium (e.g., a mounting medium, methylcellulose, or other semi-solid mediums).

A “conditionally removable coating” is a coating that can be removed from the surface of a substrate upon application of a releasing agent. In some embodiments, a conditionally removable coating includes a hydrogel as described herein, e.g., a hydrogel including a polypeptide-based material. Non-limiting examples of a hydrogel featuring a polypeptide-based material include a synthetic peptide-based material featuring a combination of spider silk and a trans-membrane segment of human muscle L-type calcium channel (e.g., PEPGEL®), an amphiphilic 16 residue peptide containing a repeating arginine-alanine-aspartate-alanine sequence (RADARADARADARADA) (e.g., PURAMATRIX®), EAK16 (AEAEAKAKAEAEAKAK), KLD12 (KLDLKLDLKLDL), and PGMATRIX™.

In some embodiments, the hydrogel in the conditionally removable coating is a stimulus-responsive hydrogel. A stimulus-responsive hydrogel can undergo a gel-to-solution and/or gel-

to-solid transition upon application of one or more external triggers (e.g., a releasing agent). See, e.g., Willner, *Acc. Chem. Res.* 50:657–658, 2017, which is incorporated herein by reference in its entirety. Non-limiting examples of a stimulus-responsive hydrogel include a thermoresponsive hydrogel, a pH-responsive hydrogel, a light-responsive hydrogel, a redox-responsive hydrogel, an analyte-responsive hydrogel, or a combination thereof. In some embodiments, a stimulus-responsive hydrogel can be a multi-stimuli-responsive hydrogel.

A “releasing agent” or “external trigger” is an agent that allows for the removal of a conditionally removable coating from a substrate when the releasing agent is applied to the conditionally removable coating. An external trigger or releasing agent can include physical triggers such as thermal, magnetic, ultrasonic, electrochemical, and/or light stimuli as well as chemical triggers such as pH, redox reactions, supramolecular complexes, and/or biocatalytically driven reactions. See e.g., Echeverria, et al., *Gels* (2018), 4, 54; doi:10.3390/gels4020054, which is incorporated herein by reference in its entirety. The type of “releasing agent” or “external trigger” can depend on the type of conditionally removable coating. For example, a conditionally removable coating featuring a redox-responsive hydrogel can be removed upon application of a releasing agent that includes a reducing agent such as dithiothreitol (DTT). As another example, a pH-responsive hydrogel can be removed upon the application of a releasing agent that changes the pH.

(d) Arrays

In many of the methods described herein, features (as described further below) are collectively positioned on a substrate. An “array” is a specific arrangement of a plurality of features that is either irregular or forms a regular pattern. Individual features in the array differ from one another based on their relative spatial locations. In general, at least two of the plurality of features in the array include a distinct capture probe (e.g., any of the examples of capture probes described herein).

Arrays can be used to measure large numbers of analytes simultaneously. In some embodiments, oligonucleotides are used, at least in part, to create an array. For example, one or more copies of a single species of oligonucleotide (e.g., capture probe) can correspond to or be directly or indirectly attached to a given feature in the array. In some embodiments, a given feature in the array includes two or more species of oligonucleotides (e.g., capture probes). In some embodiments, the two or more species of oligonucleotides (e.g., capture probes) attached

directly or indirectly to a given feature on the array include a common (e.g., identical) spatial barcode.

A “feature” is an entity that acts as a support or repository for various molecular entities used in sample analysis. Examples of features include, but are not limited to, a bead, a spot of
5 any two- or three-dimensional geometry (e.g., an ink jet spot, a masked spot, a square on a grid), a well, and a hydrogel pad. In some embodiments, features are directly or indirectly attached or fixed to a substrate. In some embodiments, the features are not directly or indirectly attached or fixed to a substrate, but instead, for example, are disposed within an enclosed or partially enclosed three dimensional space (e.g., wells or divots).

10 In addition to those above, a wide variety of other features can be used to form the arrays described herein. For example, in some embodiments, features that are formed from polymers and/or biopolymers that are jet printed, screen printed, or electrostatically deposited on a substrate can be used to form arrays. Jet printing of biopolymers is described, for example, in PCT Patent Application Publication No. WO 2014/085725. Jet printing of polymers is described,
15 for example, in de Gans et al., *Adv Mater.* 16(3): 203-213 (2004). Methods for electrostatic deposition of polymers and biopolymers are described, for example, in Hoyer et al., *Anal. Chem.* 68(21): 3840-3844 (1996). The entire contents of each of the foregoing references are incorporated herein by reference.

20 As another example, in some embodiments, features are formed by metallic micro- or nanoparticles. Suitable methods for depositing such particles to form arrays are described, for example, in Lee et al., *Beilstein J. Nanotechnol.* 8: 1049-1055 (2017), the entire contents of which are incorporated herein by reference.

25 As a further example, in some embodiments, features are formed by magnetic particles that are assembled on a substrate. Examples of such particles and methods for assembling arrays are described in Ye et al., *Scientific Reports* 6: 23145 (2016), the entire contents of which are incorporated herein by reference.

30 As another example, in some embodiments, features correspond to regions of a substrate in which one or more optical labels have been incorporated, and/or which have been altered by a process such as permanent photobleaching. Suitable substrates to implement features in this manner include a wide variety of polymers, for example. Methods for forming such features are

described, for example, in Moshrefzadeh et al., *Appl. Phys. Lett.* 62: 16 (1993), the entire contents of which are incorporated herein by reference.

As yet another example, in some embodiments, features can correspond to colloidal particles assembled (e.g., via self-assembly) to form an array. Suitable colloidal particles are described for example in Sharma, *Resonance* 23(3): 263-275 (2018), the entire contents of which are incorporated herein by reference.

As a further example, in some embodiments, features can be formed via spot-array photopolymerization of a monomer solution on a substrate. In particular, two-photon and three-photon polymerization can be used to fabricate features of relatively small (e.g., sub-micron) dimensions. Suitable methods for preparing features on a substrate in this manner are described for example in Nguyen et al., *Materials Today* 20(6): 314-322 (2017), the entire contents of which are incorporated herein by reference.

In some embodiments, features are directly or indirectly attached or fixed to a substrate that is liquid permeable. In some embodiments, features are directly or indirectly attached or fixed to a substrate that is biocompatible. In some embodiments, features are directly or indirectly attached or fixed to a substrate that is a hydrogel.

FIG. 12 depicts an exemplary arrangement of barcoded features within an array. From left to right, **FIG. 12** shows (L) a slide including six spatially-barcoded arrays, (C) an enlarged schematic of one of the six spatially-barcoded arrays, showing a grid of barcoded features in relation to a biological sample, and (R) an enlarged schematic of one section of an array, showing the specific identification of multiple features within the array (labelled as ID578, ID579, ID560, etc.).

As used herein, the term “bead array” refers to an array that includes a plurality of beads as the features in the array. In some embodiments, the beads are attached to a substrate. For example, the beads can optionally attach to a substrate such as a microscope slide and in proximity to a biological sample (e.g., a tissue section that includes cells). The beads can also be suspended in a solution and deposited on a surface (e.g., a membrane, a tissue section, or a substrate (e.g., a microscope slide)).

Examples of arrays of beads on or within a substrate include beads located in wells such as the BeadChip array (available from Illumina Inc., San Diego, CA), arrays used in sequencing platforms from 454 LifeSciences (a subsidiary of Roche, Basel, Switzerland), and array used in

sequencing platforms from Ion Torrent (a subsidiary of Life Technologies, Carlsbad, CA).

Examples of bead arrays are described in, e.g., U.S. Patent Nos. 6,266,459; 6,355,431; 6,770,441; 6,859,570; 6,210,891; 6,258,568; and 6,274,320; U.S. Pat. Application Publication Nos. 2009/0026082; 2009/0127589; 2010/0137143; and 2010/0282617; and PCT Patent

5 Application Publication Nos. WO 00/063437 and WO 2016/162309, the entire contents of each of which is incorporated herein by reference.

In some embodiments, the bead array includes a plurality of beads. For example, the bead array can include at least 10,000 beads (e.g., at least 100,000 beads, at least 1,000,000 beads, at least 5,000,000 beads, at least 10,000,000 beads). In some embodiments, the plurality of beads
10 includes a single type of beads (e.g., substantially uniform in size, shape, and other physical properties, such as translucence). In some embodiments, the plurality of beads includes two or more types of different beads.

In some embodiments, a bead array is formed when beads are embedded in a hydrogel layer where the hydrogel polymerizes and secures the relative bead positions. The bead-arrays
15 can be pre-equilibrated and combined with reaction buffers and enzymes (e.g., reverse-transcription mix). In some embodiments, the bead arrays are frozen.

A “flexible array” includes a plurality of spatially-barcoded features attached to, or embedded in, a flexible substrate (e.g., a membrane or tape) placed onto a biological sample. In some embodiments, a flexible array includes a plurality of spatially-barcoded features embedded
20 within a hydrogel matrix. To form such an array, features of a microarray are copied into a hydrogel, and the size of the hydrogel is reduced by removing water. These steps can be performed multiple times. For example, in some embodiments, a method for preparing a high-density spatially barcoded array can include copying a plurality of features from a microarray into a first hydrogel, where the first hydrogel is in contact with the microarray; reducing the size
25 of the first hydrogel including the copied features by removing water, forming a first shrunken hydrogel including the copied features; copying the features in the first shrunken hydrogel into a second hydrogel, where the second hydrogel is in contact with the first hydrogel; and reducing the size of the second hydrogel including the copied features by removing water, forming a second shrunken hydrogel including the copied features, thus generating a high-density spatially
30 barcoded array. The result is a high-density flexible array including spatially-barcoded features.

In some embodiments, spatially-barcoded beads can be loaded onto a substrate (e.g., a hydrogel) to produce a high-density self-assembled bead array.

Flexible arrays can be pre-equilibrated, combined with reaction buffers and enzymes at functional concentrations (e.g., a reverse-transcription mix). In some embodiments, the flexible
5 bead-arrays can be stored for extended periods (e.g., days) or frozen until ready for use. In some embodiments, permeabilization of biological samples (e.g., a tissue section) can be performed with the addition of enzymes/detergents prior to contact with the flexible array. The flexible array can be placed directly on the sample, or placed in indirect contact with the biological sample (e.g., with an intervening layer or substance between the biological sample and the
10 flexible bead-array). In some embodiments, once a flexible array is applied to the sample, reverse transcription and targeted capture of analytes can be performed on solid microspheres, or circular beads of a first size and circular beads of a second size.

A “microcapillary array” is an arrayed series of features that are partitioned by microcapillaries. A “microcapillary channel” is an individual partition created by the
15 microcapillaries. For example, microcapillary channels can be fluidically isolated from other microcapillary channels, such that fluid or other contents in one microcapillary channel in the array are separated from fluid or other contents in a neighboring microcapillary channel in the array. The density and order of the microcapillaries can be any suitable density or order of discrete sites.

20 In some embodiments, microcapillary arrays are treated to generate conditions that facilitate loading. An example is the use of a corona wand (BD-20AC, Electro Technic Products) to generate a hydrophilic surface. In some embodiments, a feature (e.g., a bead with capture probe attached) is loaded onto a microcapillary array such that the exact position of the feature within the array is known. For example, a capture probe containing a spatial barcode can be
25 placed into a microcapillary channel so that the spatial barcode can enable identification of the location from which the barcode sequence of the barcoded nucleic acid molecule was derived.

In some embodiments, when random distribution is used to distribute features, empirical testing can be performed to generate loading/distribution conditions that facilitate a single feature per microcapillary. In some embodiments, it can be desirable to achieve distribution conditions
30 that facilitate only a single feature (e.g., bead) per microcapillary channel. In some embodiments,

it can be desirable to achieve distribution conditions that facilitate more than one feature (e.g., bead) per microcapillary channel, by flowing the features through the microcapillary channel.

In some embodiments, the microcapillary array is placed in contact with a sample (e.g., on top or below) so that microcapillaries containing a feature (e.g., a bead, which can include a capture probe) are in contact with the biological sample. In some embodiments, a biological sample is placed onto an exposed side of a microcapillary array and mechanical compression is applied, moving the biological sample into the microcapillary channel to create a fluidically isolated reaction chamber containing the biological sample.

In some embodiments, a biological sample is partitioned by contacting a microcapillary array to the biological sample, thereby creating microcapillary channels including a bead and a portion of the biological sample. In some embodiments, a portion of a biological sample contained in a microcapillary channel is one or more cells. In some embodiments, a feature is introduced into a microcapillary array by flow after one or more cells are added to a microcapillary channel.

In some embodiments, reagents are added to the microcapillary array. The added reagents can include enzymatic reagents, and reagent mixtures for performing amplification of a nucleic acid. In some embodiments, the reagents include a reverse transcriptase, a ligase, one or more nucleotides, and any combinations thereof. One or more microcapillary channels can be sealed after reagents are added to the microcapillary channels, e.g. using silicone oil, mineral oil, a non-porous material, or lid.

In some embodiments, a reagent solution is removed from each microcapillary channel following an incubation for an amount of time and at a certain temperature or range of temperatures, e.g., following a hybridization or an amplification reaction. Reagent solutions can be processed individually for sequencing, or pooled for sequencing analysis.

In some embodiments, some or all features in an array include a capture probe. In some embodiments, an array can include a capture probe attached directly or indirectly to the substrate.

The capture probe includes a capture domain (e.g., a nucleotide sequence) that can specifically bind (e.g., hybridize) to a target analyte (e.g., mRNA, DNA, or protein) within a sample. In some embodiments, the binding of the capture probe to the target (e.g., hybridization) can be detected and quantified by detection of a visual signal, e.g. a fluorophore, a heavy metal (e.g., silver ion), or chemiluminescent label, which has been incorporated into the target. In some

embodiments, the intensity of the visual signal correlates with the relative abundance of each analyte in the biological sample. Since an array can contain thousands or millions of capture probes (or more), an array of features with capture probes can interrogate many analytes in parallel.

5 In some embodiments, a substrate includes one or more capture probes that are designed to capture analytes from one or more organisms. In a non-limiting example, a substrate can contain one or more capture probes designed to capture mRNA from one organism (e.g., a human) and one or more capture probes designed to capture DNA from a second organism (e.g., a bacterium).

10 The capture probes can be attached to a substrate or feature using a variety of techniques. In some embodiments, the capture probe is directly attached to a feature that is fixed on an array. In some embodiments, the capture probes are immobilized to a substrate by chemical immobilization. For example, a chemical immobilization can take place between functional groups on the substrate and corresponding functional elements on the capture probes. Exemplary
15 corresponding functional elements in the capture probes can either be an inherent chemical group of the capture probe, e.g. a hydroxyl group, or a functional element can be introduced on to the capture probe. An example of a functional group on the substrate is an amine group. In some embodiments, the capture probe to be immobilized includes a functional amine group or is chemically modified in order to include a functional amine group. Means and methods for such a
20 chemical modification are well known in the art.

In some embodiments, the capture probe is a nucleic acid. In some embodiments, the capture probe is immobilized on the feature or the substrate via its 5' end. In some embodiments, the capture probe is immobilized on a feature or a substrate via its 5' end and includes from the 5' to 3' end: one or more barcodes (e.g., a spatial barcode and/or a UMI) and one or more
25 capture domains. In some embodiments, the capture probe is immobilized on a feature via its 5' end and includes from the 5' to 3' end: one barcode (e.g., a spatial barcode or a UMI) and one capture domain. In some embodiments, the capture probe is immobilized on a feature or a substrate via its 5' end and includes from the 5' to 3' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), and a capture domain.

30 In some embodiments, the capture probe is immobilized on a feature or a substrate via its 5' end and includes from the 5' to 3' end: a cleavage domain, a functional domain, one or more

barcodes (e.g., a spatial barcode and/or a UMI), a second functional domain, and a capture domain. In some embodiments, the capture probe is immobilized on a feature or a substrate via its 5' end and includes from the 5' to 3' end: a cleavage domain, a functional domain, a spatial barcode, a UMI, and a capture domain. In some embodiments, the capture probe is immobilized on a feature or a substrate via its 5' end and does not include a spatial barcode. In some
5 embodiments, the capture probe is immobilized on a feature or a substrate via its 5' end and does not include a UMI. In some embodiments, the capture probe includes a sequence for initiating a sequencing reaction.

 In some embodiments, the capture probe is immobilized on a feature or a substrate via its
10 3' end. In some embodiments, the capture probe is immobilized on a feature or a substrate via its 3' end and includes from the 3' to 5' end: one or more barcodes (e.g., a spatial barcode and/or a UMI) and one or more capture domains. In some embodiments, the capture probe is immobilized on a feature or a substrate via its 3' end and includes from the 3' to 5' end: one barcode (e.g., a spatial barcode or a UMI) and one capture domain. In some embodiments, the capture probe is
15 immobilized on a feature or a substrate via its 3' end and includes from the 3' to 5' end: a cleavage domain, a functional domain, one or more barcodes (e.g., a spatial barcode and/or a UMI), and a capture domain. In some embodiments, the capture probe is immobilized on a feature or a substrate via its 3' end and includes from the 3' to 5' end: a cleavage domain, a functional domain, a spatial barcode, a UMI, and a capture domain.

20 The localization of the functional group within the capture probe to be immobilized can be used to control and shape the binding behavior and/or orientation of the capture probe, e.g. the functional group can be placed at the 5' or 3' end of the capture probe or within the sequence of the capture probe. In some embodiments, a capture probe can further include a substrate (e.g., a support attached to the capture probe, a support attached to the feature, or a support attached to
25 the substrate). A typical substrate for a capture probe to be immobilized includes moieties which are capable of binding to such capture probes, e.g., to amine-functionalized nucleic acids. Examples of such substrates are carboxy, aldehyde, or epoxy supports.

 In some embodiments, the substrates on which capture probes can be immobilized can be chemically activated, e.g. by the activation of functional groups, available on the substrate. The
30 term "activated substrate" relates to a material in which interacting or reactive chemical functional groups are established or enabled by chemical modification procedures. For example,

a substrate including carboxyl groups can be activated before use. Furthermore, certain substrates contain functional groups that can react with specific moieties already present in the capture probes.

5 In some embodiments, a covalent linkage is used to directly couple a capture probe to a substrate. In some embodiments a capture probe is indirectly coupled to a substrate through a linker separating the “first” nucleotide of the capture probe from the substrate, i.e., a chemical linker. In some embodiments, a capture probe does not bind directly to the array, but interacts indirectly, for example by binding to a molecule which itself binds directly or indirectly to the array. In some embodiments, the capture probe is indirectly attached to a substrate (e.g., via a
10 solution including a polymer).

In some embodiments where the capture probe is immobilized on the feature of the array indirectly, e.g. via hybridization to a surface probe capable of binding the capture probe, the capture probe can further include an upstream sequence (5' to the sequence that hybridizes to the nucleic acid, e.g. RNA of the tissue sample) that is capable of hybridizing to 5' end of the
15 surface probe. Alone, the capture domain of the capture probe can be seen as a capture domain oligonucleotide, which can be used in the synthesis of the capture probe in embodiments where the capture probe is immobilized on the array indirectly.

In some embodiments, a substrate is comprised of an inert material or matrix (e.g., glass slides) that has been functionalization by, for example, treatment with a material comprising
20 reactive groups which enable immobilization of capture probes. See, for example, WO 2017/019456, the entire contents of which are herein incorporated by reference. Non-limiting examples include polyacrylamide hydrogels supported on an inert substrate (e.g., glass slide; see WO 2005/065814 and U.S. Patent Application No. 2008/0280773, the entire contents of which are incorporated herein by reference).

25 In some embodiments, functionalized biomolecules (e.g., capture probes) are immobilized on a functionalized substrate using covalent methods. Methods for covalent attachment include, for example, condensation of amines and activated carboxylic esters (e.g., N-hydroxysuccinimide esters); condensation of amine and aldehydes under reductive amination conditions; and cycloaddition reactions such as the Diels–Alder [4+2] reaction, 1,3-dipolar
30 cycloaddition reactions, and [2+2] cycloaddition reactions. Methods for covalent attachment also include, for example, click chemistry reactions, including [3+2] cycloaddition reactions (e.g.,

Huisgen 1,3-dipolar cycloaddition reaction and copper(I)-catalyzed azide-alkyne cycloaddition (CuAAC)); thiol-ene reactions; the Diels–Alder reaction and inverse electron demand Diels–Alder reaction; [4+1] cycloaddition of isonitriles and tetrazines; and nucleophilic ring-opening of small carbocycles (e.g., epoxide opening with amino oligonucleotides). Methods for covalent attachment also include, for example, maleimides and thiols; and *para*-nitrophenyl ester-functionalized oligonucleotides and polylysine-functionalized substrate. Methods for covalent attachment also include, for example, disulfide reactions; radical reactions (see, e.g., U.S. Patent No. 5,919,626, the entire contents of which are herein incorporated by reference); and hydrazide-functionalized substrate (e.g., wherein the hydrazide functional group is directly or indirectly attached to the substrate) and aldehyde-functionalized oligonucleotides (see, e.g., Yershov *et al.* (1996) Proc. Natl. Acad. Sci. USA 93, 4913–4918, the entire contents of which are herein incorporated by reference).

In some embodiments, functionalized biomolecules (e.g., capture probes) are immobilized on a functionalized substrate using photochemical covalent methods. Methods for photochemical covalent attachment include, for example, immobilization of anthraquinone-conjugated oligonucleotides (see, e.g., Koch *et al.* (2000) Bioconjugate Chem. 11, 474–483, the entire contents of which are herein incorporated by reference).

In some embodiments, functionalized biomolecules (e.g., capture probes) are immobilized on a functionalized substrate using non-covalent methods. Methods for non-covalent attachment include, for example, biotin-functionalized oligonucleotides and streptavidin-treated substrates (see, e.g., Holmstrøm *et al.* (1993) Analytical Biochemistry 209, 278–283 and Gilles *et al.* (1999) Nature Biotechnology 17, 365–370, the entire contents of which are herein incorporated by reference).

In some embodiments, an oligonucleotide (e.g., a capture probe) can be attached to a substrate or feature according to the methods set forth in U.S. Patent Nos. 6,737,236, 7,259,258, 7,375,234, 7,427,678, 5,610,287, 5,807,522, 5,837,860, and 5,472,881; U.S. Patent Application Publication Nos. 2008/0280773 and 2011/0059865; Shalon *et al.* (1996) Genome Research, 639–645; Rogers *et al.* (1999) Analytical Biochemistry 266, 23–30; Stimpson *et al.* (1995) Proc. Natl. Acad. Sci. USA 92, 6379–6383; Beattie *et al.* (1995) Clin. Chem. 45, 700–706; Lamture *et al.* (1994) Nucleic Acids Research 22, 2121–2125; Beier *et al.* (1999) Nucleic Acids Research 27, 1970–1977; Joos *et al.* (1997) Analytical Biochemistry 247, 96–101; Nikiforov *et al.* (1995)

Analytical Biochemistry 227, 201–209; Timofeev *et al.* (1996) Nucleic Acids Research 24, 3142–3148; Chrisey *et al.* (1996) Nucleic Acids Research 24, 3031–3039; Guo *et al.* (1994) Nucleic Acids Research 22, 5456–5465; Running and Urdea (1990) BioTechniques 8, 276–279; Fahy *et al.* (1993) Nucleic Acids Research 21, 1819–1826; Zhang *et al.* (1991) 19, 3929–3933; and Rogers *et al.* (1997) Gene Therapy 4, 1387–1392. The entire contents of each of the foregoing documents are incorporated herein by reference.

Arrays can be prepared by a variety of methods. In some embodiments, arrays are prepared through the synthesis (e.g., in-situ synthesis) of oligonucleotides on the array, or by jet printing or lithography. For example, light-directed synthesis of high-density DNA oligonucleotides can be achieved by photolithography or solid-phase DNA synthesis. To implement photolithographic synthesis, synthetic linkers modified with photochemical protecting groups can be attached to a substrate and the photochemical protecting groups can be modified using a photolithographic mask (applied to specific areas of the substrate) and light, thereby producing an array having localized photo-deprotection. Many of these methods are known in the art, and are described e.g., in Miller et al., “Basic concepts of microarrays and potential applications in clinical microbiology.” Clinical microbiology reviews 22.4 (2009): 611–633; US201314111482A; US9593365B2; US2019203275; and WO2018091676, which are incorporated herein by reference in the entirety.

In some embodiments, the arrays are “spotted” or “printed” with oligonucleotides and these oligonucleotides (e.g., capture probes) are then attached to the substrate. The oligonucleotides can be applied by either noncontact or contact printing. A noncontact printer can use the same method as computer printers (e.g., bubble jet or inkjet) to expel small droplets of probe solution onto the substrate. The specialized inkjet-like printer can expel nanoliter to picoliter volume droplets of oligonucleotide solution, instead of ink, onto the substrate. In contact printing, each print pin directly applies the oligonucleotide solution onto a specific location on the surface. The oligonucleotides can be attached to the substrate surface by the electrostatic interaction of the negative charge of the phosphate backbone of the DNA with a positively charged coating of the substrate surface or by UV-cross-linked covalent bonds between the thymidine bases in the DNA and amine groups on the treated substrate surface. In some embodiments, the substrate is a glass slide. In some embodiments, the oligonucleotides

(e.g., capture probes) are attached to the substrate by a covalent bond to a chemical matrix, e.g. epoxy-silane, amino-silane, lysine, polyacrylamide, etc.

The arrays can also be prepared by in situ-synthesis. In some embodiments, these arrays can be prepared using photolithography. The method typically relies on UV masking and light-directed combinatorial chemical synthesis on a substrate to selectively synthesize probes directly on the surface of the array, one nucleotide at a time per spot, for many spots simultaneously. In some embodiments, a substrate contains covalent linker molecules that have a protecting group on the free end that can be removed by light. UV light is directed through a photolithographic mask to deprotect and activate selected sites with hydroxyl groups that initiate coupling with incoming protected nucleotides that attach to the activated sites. The mask is designed in such a way that the exposure sites can be selected, and thus specify the coordinates on the array where each nucleotide can be attached. The process can be repeated, a new mask is applied activating different sets of sites and coupling different bases, allowing arbitrary oligonucleotides to be constructed at each site. This process can be used to synthesize hundreds of thousands of different oligonucleotides. In some embodiments, maskless array synthesizer technology can be used. It uses an array of programmable micromirrors to create digital masks that reflect the desired pattern of UV light to deprotect the features.

In some embodiments, the inkjet spotting process can also be used for in-situ oligonucleotide synthesis. The different nucleotide precursors plus catalyst can be printed on the substrate, and are then combined with coupling and deprotection steps. This method relies on printing picoliter volumes of nucleotides on the array surface in repeated rounds of base-by-base printing that extends the length of the oligonucleotide probes on the array.

Arrays can also be prepared by active hybridization via electric fields to control nucleic acid transport. Negatively charged nucleic acids can be transported to specific sites, or features, when a positive current is applied to one or more test sites on the array. The surface of the array can contain a binding molecule, e.g., streptavidin, which allows for the formation of bonds (e.g., streptavidin-biotin bonds) once electronically addressed biotinylated probes reach their targeted location. The positive current is then removed from the active features, and new test sites can be activated by the targeted application of a positive current. The process are repeated until all sites on the array are covered.

An array for spatial analysis can be generated by various methods as described herein. In some embodiments, the array has a plurality of capture probes comprising spatial barcodes. These spatial barcodes and their relationship to the locations on the array can be determined. In some cases, such information is readily available, because the oligonucleotides are spotted,
5 printed, or synthesized on the array with a pre-determined pattern. In some cases, the spatial barcode can be decoded by methods described herein, e.g., by in-situ sequencing, by various labels associated with the spatial barcodes etc. In some embodiments, an array can be used as a template to generate a daughter array. Thus, the spatial barcode can be transferred to the daughter array with a known pattern.

10 In some embodiments, an array comprising barcoded probes can be generated through ligation of a plurality of oligonucleotides. In some instances, an oligonucleotide of the plurality contains a portion of a barcode, and the complete barcode is generated upon ligation of the plurality of oligonucleotides. For example, a first oligonucleotide containing a first portion of a barcode can be attached to a substrate (e.g., using any of the methods of attaching an
15 oligonucleotide to a substrate described herein), and a second oligonucleotide containing a second portion of the barcode can then be ligated onto the first oligonucleotide to generate a complete barcode. Different combinations of the first, second and any additional portions of a barcode can be used to increase the diversity of the barcodes. In instances where the second oligonucleotide is also attached to the substrate prior to ligation, the first and/or the second
20 oligonucleotide can be attached to the substrate via a surface linker which contains a cleavage site. Upon ligation, the ligated oligonucleotide is linearized by cleaving at the cleavage site.

To increase the diversity of the barcodes, a plurality of second oligonucleotides comprising two or more different barcode sequences can be ligated onto a plurality of first oligonucleotides that comprise the same barcode sequence, thereby generating two or more
25 different species of barcodes. To achieve selective ligation, a first oligonucleotide attached to a substrate containing a first portion of a barcode can initially be protected with a protective group (e.g., a photocleavable protective group), and the protective group can be removed prior to ligation between the first and second oligonucleotide. In instances where the barcoded probes on an array are generated through ligation of two or more oligonucleotides, a concentration gradient
30 of the oligonucleotides can be applied to a substrate such that different combinations of the

oligonucleotides are incorporated into a barcoded probe depending on its location on the substrate.

Barcoded probes on an array can also be generated by adding single nucleotides to existing oligonucleotides on an array, for example, using polymerases that function in a
5 template-independent manner. Single nucleotides can be added to existing oligonucleotides in a concentration gradient, thereby generating probes with varying length, depending on the location of the probes on the array.

Arrays can also be prepared by modifying existing arrays, for example, by modifying the oligonucleotides attached to the arrays. For instance, probes can be generated on an array that
10 comprises oligonucleotides that are attached to the array at the 3' end and have a free 5' end. The oligonucleotides can be in situ synthesized oligonucleotides, and can include a barcode. The length of the oligonucleotides can be less than 50 nucleotides (nts) (e.g., less than 45, 40, 35, 30, 25, 20, 15, or 10 nts). To generate probes using these oligonucleotides, a primer complementary to a portion of an oligonucleotide (e.g., a constant sequence shared by the oligonucleotides) can
15 be used to hybridize with the oligonucleotide and extend (using the oligonucleotide as a template) to form a duplex and to create a 3' overhang. The 3' overhang thus allows additional nucleotides or oligonucleotides to be added on to the duplex. A capture probe can be generated by, for instance, adding one or more oligonucleotides to the end of the 3' overhang (e.g., via splint oligonucleotide mediated ligation), where the added oligonucleotides can include the
20 sequence or a portion of the sequence of a capture domain.

In instances where the oligonucleotides on an existing array include a recognition sequence that can hybridize with a splint oligonucleotide, probes can also be generated by directly ligating additional oligonucleotides onto the existing oligonucleotides via the splint oligonucleotide. The recognition sequence can at the free 5' end or the free 3' end of an
25 oligonucleotide on the existing array. Recognition sequences useful for the methods of the present disclosure may not contain restriction enzyme recognition sites or secondary structures (e.g., hairpins), and may include high contents of Guanine and Cytosine nucleotides and thus have high stability.

Bead arrays can be generated by attaching beads (e.g., barcoded beads) to a substrate in a
30 regular pattern, or an irregular arrangement. Beads can be attached to selective regions on a substrate by, e.g., selectively activating regions on the substrate to allow for attachment of the

beads. Activating selective regions on the substrate can include activating a coating (e.g., a photocleavable coating) or a polymer that is applied on the substrate. Beads can be attached iteratively, e.g., a subset of the beads can be attached at one time, and the same process can be repeated to attach the remaining beads. Alternatively, beads can be attached to the substrate all in one step.

Barcoded beads, or beads comprising a plurality of barcoded probes, can be generated by first preparing a plurality of barcoded probes on a substrate, depositing a plurality of beads on the substrate, and generating probes attached to the beads using the probes on the substrate as a template.

Large scale commercial manufacturing methods allow for millions of oligonucleotides to be attached to an array. Commercially available arrays include those from Roche NimbleGen, Inc., (Wisconsin) and Affymetrix (ThermoFisher Scientific).

In some embodiments, arrays can be prepared according to the methods set forth in WO 2012/140224, WO 2014/060483, WO 2016/162309, WO 2017/019456, WO 2018/091676, and WO 2012/140224, and U.S. Patent Application No. 2018/0245142. The entire contents of the foregoing documents are herein incorporated by reference.

In some embodiments, a feature on the array includes a bead. In some embodiments, two or more beads are dispersed onto a substrate to create an array, where each bead is a feature on the array. Beads can optionally be dispersed into wells on a substrate, e.g., such that only a single bead is accommodated per well.

A “bead” is a particle. A bead can be porous, non-porous, solid, semi-solid, and/or a combination thereof. In some embodiments, a bead can be dissolvable, disruptable, and/or degradable, whereas in certain embodiments, a bead is not degradable.

A bead can generally be of any suitable shape. Examples of bead shapes include, but are not limited to, spherical, non-spherical, oval, oblong, amorphous, circular, cylindrical, and variations thereof. A cross section (e.g., a first cross-section) can correspond to a diameter or maximum cross-sectional dimension of the bead. In some embodiments, the bead can be approximately spherical. In such embodiments, the first cross-section can correspond to the diameter of the bead. In some embodiments, the bead can be approximately cylindrical. In such embodiments, the first cross-section can correspond to a diameter, length, or width along the approximately cylindrical bead.

Beads can be of uniform size or heterogeneous size. “Polydispersity” generally refers to heterogeneity of sizes of molecules or particles. The polydispersity index (PDI) of a bead can be calculated using the equation $PDI = M_w/M_n$, where M_w is the weight-average molar mass and M_n is the number-average molar mass. In certain embodiments, beads can be provided as a population or plurality of beads having a relatively monodisperse size distribution. Where it can be desirable to provide relatively consistent amounts of reagents, maintaining relatively consistent bead characteristics, such as size, can contribute to the overall consistency.

In some embodiments, the beads provided herein can have size distributions that have a coefficient of variation in their cross-sectional dimensions of less than 50%, less than 40%, less than 30%, less than 20%, less than 15%, less than 10%, less than 5%, or lower. In some embodiments, a plurality of beads provided herein has a polydispersity index of less than 50%, less than 45%, less than 40%, less than 35%, less than 30%, less than 25%, less than 20%, less than 15%, less than 10%, less than 5%, or lower.

In some embodiments, the bead can have a diameter or maximum dimension no larger than 100 μm (e.g., no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , 10 μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm .)

In some embodiments, a plurality of beads has an average diameter no larger than 100 μm . In some embodiments, a plurality of beads has an average diameter or maximum dimension no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , 10 μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm .

In some embodiments, the volume of the bead can be at least about 1 μm^3 , e.g., at least 1 μm^3 , 2 μm^3 , 3 μm^3 , 4 μm^3 , 5 μm^3 , 6 μm^3 , 7 μm^3 , 8 μm^3 , 9 μm^3 , 10 μm^3 , 12 μm^3 , 14 μm^3 , 16 μm^3 , 18 μm^3 , 20 μm^3 , 25 μm^3 , 30 μm^3 , 35 μm^3 , 40 μm^3 , 45 μm^3 , 50 μm^3 , 55 μm^3 , 60 μm^3 , 65 μm^3 , 70 μm^3 , 75 μm^3 , 80 μm^3 , 85 μm^3 , 90 μm^3 , 95 μm^3 , 100 μm^3 , 125 μm^3 , 150 μm^3 , 175 μm^3 , 200 μm^3 , 250 μm^3 , 300 μm^3 , 350 μm^3 , 400 μm^3 , 450 μm^3 , 500 μm^3 , 550 μm^3 , 600 μm^3 , 650 μm^3 , 700 μm^3 , 750 μm^3 , 800 μm^3 , 850 μm^3 , 900 μm^3 , 950 μm^3 , 1000 μm^3 , 1200 μm^3 , 1400 μm^3 , 1600 μm^3 , 1800 μm^3 , 2000 μm^3 , 2200 μm^3 , 2400 μm^3 , 2600 μm^3 , 2800 μm^3 , 3000 μm^3 , or greater.

In some embodiments, the bead can have a volume of between about $1\ \mu\text{m}^3$ and $100\ \mu\text{m}^3$, such as between about $1\ \mu\text{m}^3$ and $10\ \mu\text{m}^3$, between about $10\ \mu\text{m}^3$ and $50\ \mu\text{m}^3$, or between about $50\ \mu\text{m}^3$ and $100\ \mu\text{m}^3$. In some embodiments, the bead can include a volume of between about $100\ \mu\text{m}^3$ and $1000\ \mu\text{m}^3$, such as between about $100\ \mu\text{m}^3$ and $500\ \mu\text{m}^3$ or between about $500\ \mu\text{m}^3$ and $1000\ \mu\text{m}^3$. In some embodiments, the bead can include a volume between about $1000\ \mu\text{m}^3$ and $3000\ \mu\text{m}^3$, such as between about $1000\ \mu\text{m}^3$ and $2000\ \mu\text{m}^3$ or between about $2000\ \mu\text{m}^3$ and $3000\ \mu\text{m}^3$. In some embodiments, the bead can include a volume between about $1\ \mu\text{m}^3$ and $3000\ \mu\text{m}^3$, such as between about $1\ \mu\text{m}^3$ and $2000\ \mu\text{m}^3$, between about $1\ \mu\text{m}^3$ and $1000\ \mu\text{m}^3$, between about $1\ \mu\text{m}^3$ and $500\ \mu\text{m}^3$, or between about $1\ \mu\text{m}^3$ and $250\ \mu\text{m}^3$.

The bead can include one or more cross-sections that can be the same or different. In some embodiments, the bead can have a first cross-section that is different from a second cross-section. The bead can have a first cross-section that is at least about 0.0001 micrometer, 0.001 micrometer, 0.01 micrometer, 0.1 micrometer, or 1 micrometer. In some embodiments, the bead can include a cross-section (e.g., a first cross-section) of at least about 1 micrometer (μm), 2 μm , 3 μm , 4 μm , 5 μm , 6 μm , 7 μm , 8 μm , 9 μm , 10 μm , 11 μm , 12 μm , 13 μm , 14 μm , 15 μm , 16 μm , 17 μm , 18 μm , 19 μm , 20 μm , 25 μm , 30 μm , 35 μm , 40 μm , 45 μm , 50 μm , 55 μm , 60 μm , 65 μm , 70 μm , 75 μm , 80 μm , 85 μm , 90 μm , 100 μm , 120 μm , 140 μm , 160 μm , 180 μm , 200 μm , 250 μm , 300 μm , 350 μm , 400 μm , 450 μm , 500 μm , 550 μm , 600 μm , 650 μm , 700 μm , 750 μm , 800 μm , 850 μm , 900 μm , 950 μm , 1 millimeter (mm), or greater. In some embodiments, the bead can include a cross-section (e.g., a first cross-section) of between about 1 μm and 500 μm , such as between about 1 μm and 100 μm , between about 100 μm and 200 μm , between about 200 μm and 300 μm , between about 300 μm and 400 μm , or between about 400 μm and 500 μm . For example, the bead can include a cross-section (e.g., a first cross-section) of between about 1 μm and 100 μm . In some embodiments, the bead can have a second cross-section that is at least about 1 μm . For example, the bead can include a second cross-section of at least about 1 micrometer (μm), 2 μm , 3 μm , 4 μm , 5 μm , 6 μm , 7 μm , 8 μm , 9 μm , 10 μm , 11 μm , 12 μm , 13 μm , 14 μm , 15 μm , 16 μm , 17 μm , 18 μm , 19 μm , 20 μm , 25 μm , 30 μm , 35 μm , 40 μm , 45 μm , 50 μm , 55 μm , 60 μm , 65 μm , 70 μm , 75 μm , 80 μm , 85 μm , 90 μm , 100 μm , 120 μm , 140 μm , 160 μm , 180 μm , 200 μm , 250 μm , 300 μm , 350 μm , 400 μm , 450 μm , 500 μm , 550 μm , 600 μm , 650 μm , 700 μm , 750 μm , 800 μm , 850 μm , 900 μm , 950 μm , 1 millimeter (mm), or greater. In some embodiments, the bead can include a second cross-section of between about 1 μm and 500

μm, such as between about 1 μm and 100 μm, between about 100 μm and 200 μm, between about 200 μm and 300 μm, between about 300 μm and 400 μm, or between about 400 μm and 500 μm. For example, the bead can include a second cross-section of between about 1 μm and 100 μm.

5 In some embodiments, beads can be of a nanometer scale (e.g., beads can have a diameter or maximum cross-sectional dimension of about 100 nanometers (nm) to about 900 nanometers (nm) (e.g., 850 nm or less, 800 nm or less, 750 nm or less, 700 nm or less, 650 nm or less, 600 nm or less, 550 nm or less, 500 nm or less, 450 nm or less, 400 nm or less, 350 nm or less, 300 nm or less, 250 nm or less, 200 nm or less, 150 nm or less). A plurality of beads can have an
10 average diameter or average maximum cross-sectional dimension of about 100 nanometers (nm) to about 900 nanometers (nm) (e.g., 850 nm or less, 800 nm or less, 750 nm or less, 700 nm or less, 650 nm or less, 600 nm or less, 550 nm or less, 500 nm or less, 450 nm or less, 400 nm or less, 350 nm or less, 300 nm or less, 250 nm or less, 200 nm or less, 150 nm or less). In some
15 embodiments, a bead has a diameter or size that is about the size of a single cell (e.g., a single cell under evaluation).

 In some embodiments, the bead can be a gel bead. A “gel” is a semi-rigid material permeable to liquids and gases. Exemplary gels include, but are not limited to, those having a colloidal structure, such as agarose; polymer mesh structures, such as gelatin; hydrogels; and cross-linked polymer structures, such as polyacrylamide, SFA (see, for example, U.S. Patent
20 Application Publication No. 2011/0059865, which is incorporated herein by reference in its entirety) and PAZAM (see, for example, U.S. Patent Application Publication No. 2014/0079923, which is incorporated herein by reference in its entirety).

 A gel can be formulated into various shapes and dimensions depending on the context of intended use. In some embodiments, a gel is prepared and formulated as a gel bead (e.g., a gel
25 bead including capture probes attached or associated with the gel bead). A gel bead can be a hydrogel bead. A hydrogel bead can be formed from molecular precursors, such as a polymeric or monomeric species.

 In some embodiments, a hydrogel bead can include a polymer matrix (e.g., a matrix formed by polymerization or cross-linking). A polymer matrix can include one or more polymers

(e.g., polymers having different functional groups or repeat units). Cross-linking can be via covalent, ionic, and/or inductive interactions, and/or physical entanglement.

A semi-solid bead can be a liposomal bead.

Solid beads can include metals including, without limitation, iron oxide, gold, and silver.

- 5 In some embodiments, the bead can be a silica bead. In some embodiments, the bead can be rigid. In some embodiments, the bead can be flexible and/or compressible.

The bead can be a macromolecule. The bead can be formed of nucleic acid molecules bound together. The bead can be formed via covalent or non-covalent assembly of molecules (e.g., macromolecules), such as monomers or polymers. Polymers or monomers can be natural or
10 synthetic. Polymers or monomers can be or include, for example, nucleic acid molecules (e.g., DNA or RNA).

A bead can be rigid, or flexible and/or compressible. A bead can include a coating including one or more polymers. Such a coating can be disruptable or dissolvable. In some
15 embodiments, a bead includes a spectral or optical label (e.g., dye) attached directly or indirectly (e.g., through a linker) to the bead. For example, a bead can be prepared as a colored preparation (e.g., a bead exhibiting a distinct color within the visible spectrum) that can change color (e.g., colorimetric beads) upon application of a desired stimulus (e.g., heat and/or chemical reaction) to form differently colored beads (e.g., opaque and/or clear beads).

A bead can include natural and/or synthetic materials. For example, a bead can include a
20 natural polymer, a synthetic polymer or both natural and synthetic polymers. Examples of natural polymers include, without limitation, proteins, sugars such as deoxyribonucleic acid, rubber, cellulose, starch (e.g., amylose, amylopectin), enzymes, polysaccharides, silks, polyhydroxyalkanoates, chitosan, dextran, collagen, carrageenan, ispaghula, acacia, agar, gelatin, shellac, sterculia gum, xanthan gum, corn sugar gum, guar gum, gum karaya, agarose, alginic
25 acid, alginate, or natural polymers thereof. Examples of synthetic polymers include, without limitation, acrylics, nylons, silicones, spandex, viscose rayon, polycarboxylic acids, polyvinyl acetate, polyacrylamide, polyacrylate, polyethylene glycol, polyurethanes, polylactic acid, silica, polystyrene, polyacrylonitrile, polybutadiene, polycarbonate, polyethylene, polyethylene terephthalate, poly(chlorotrifluoroethylene), poly(ethylene oxide), poly(ethylene terephthalate),
30 polyethylene, polyisobutylene, poly(methyl methacrylate), poly(oxymethylene), polyformaldehyde, polypropylene, polystyrene, poly(tetrafluoroethylene), poly(vinyl acetate),

poly(vinyl alcohol), poly(vinyl chloride), poly(vinylidene dichloride), poly(vinylidene difluoride), poly(vinyl fluoride) and/or combinations (e.g., co-polymers) thereof. Beads can also be formed from materials other than polymers, including for example, lipids, micelles, ceramics, glass-ceramics, material composites, metals, and/or other inorganic materials.

5 In some embodiments, a bead is a degradable bead. A degradable bead can include one or more species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) with a labile bond such that, when the bead/species is exposed to the appropriate stimuli, the labile bond is broken and the bead degrades. The labile bond can be a chemical bond (e.g., covalent bond, ionic bond) or can be another type of physical interaction (e.g., van der Waals interactions, dipole-dipole
10 interactions, etc.). In some embodiments, a crosslinker used to generate a bead can include a labile bond. Upon exposure to the appropriate conditions, the labile bond can be broken and the bead degraded. For example, upon exposure of a polyacrylamide gel bead including cystamine crosslinkers to a reducing agent, the disulfide bonds of the cystamine can be broken and the bead degraded.

15 Degradation can refer to the disassociation of a bound or entrained species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) from a bead, both with and without structurally degrading the physical bead itself. For example, entrained species can be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore sizes due to osmotic pressure differences can generally
20 occur without structural degradation of the bead itself. In some embodiments, an increase in pore size due to osmotic swelling of a bead can permit the release of entrained species within the bead. In some embodiments, osmotic shrinking of a bead can cause a bead to better retain an entrained species due to pore size contraction.

 Any suitable agent that can degrade beads can be used. In some embodiments, changes in
25 temperature or pH can be used to degrade thermo-sensitive or pH-sensitive bonds within beads. In some embodiments, chemical degrading agents can be used to degrade chemical bonds within beads by oxidation, reduction or other chemical changes. For example, a chemical degrading agent can be a reducing agent, such as DTT, where DTT can degrade the disulfide bonds formed between a crosslinker and gel precursors, thus degrading the bead. In some embodiments, a
30 reducing agent can be added to degrade the bead, which can cause the bead to release its contents. Examples of reducing agents can include, without limitation, dithiothreitol (DTT), β -

mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris(2-carboxyethyl) phosphine (TCEP), or combinations thereof.

Any of a variety of chemical agents can be used to trigger the degradation of beads. Examples of chemical agents include, but are not limited to, pH-mediated changes to the integrity of a component within the bead, degradation of a component of a bead via cleavage of cross-linked bonds, and depolymerization of a component of a bead.

In some embodiments, a bead can be formed from materials that include degradable chemical crosslinkers, such as N,N'-bis-(acryloyl)cystamine (BAC) or cystamine. Degradation of such degradable crosslinkers can be accomplished through any variety of mechanisms. In some examples, a bead can be contacted with a chemical degrading agent that can induce oxidation, reduction or other chemical changes. For example, a chemical degrading agent can be a reducing agent, such as dithiothreitol (DTT). Additional examples of reducing agents can include β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris(2-carboxyethyl) phosphine (TCEP), or combinations thereof.

In some embodiments, exposure to an aqueous solution, such as water, can trigger hydrolytic degradation, and thus degradation of the bead. Beads can also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety of changes to a bead. For example, heat can cause a solid bead to liquefy. A change in heat can cause melting of a bead such that a portion of the bead degrades. In some embodiments, heat can increase the internal pressure of the bead components such that the bead ruptures or explodes. Heat can also act upon heat-sensitive polymers used as materials to construct beads.

Where degradable beads are used, it can be beneficial to avoid exposing such beads to the stimulus or stimuli that cause such degradation prior to a given time, in order to, for example, avoid premature bead degradation and issues that arise from such degradation, including for example poor flow characteristics and aggregation. By way of example, where beads include reducible cross-linking groups, such as disulfide groups, it will be desirable to avoid contacting such beads with reducing agents, e.g., DTT or other disulfide cleaving reagents. In such embodiments, treatment of the beads described herein will, in some embodiments be provided free of reducing agents, such as DTT. Because reducing agents are often provided in commercial enzyme preparations, it can be desirable to provide reducing agent free (or DTT free) enzyme preparations in treating the beads described herein. Examples of such enzymes include, e.g.,

polymerase enzyme preparations, reverse transcriptase enzyme preparations, ligase enzyme preparations, as well as many other enzyme preparations that can be used to treat the beads described herein. The terms “reducing agent free” or “DTT free” preparations refer to a preparation having less than about 1/10th, less than about 1/50th, or less than about 1/100th of the lower ranges for such materials used in degrading the beads. For example, for DTT, the
5 reducing agent free preparation can have less than about 0.01 millimolar (mM), 0.005 mM, 0.001 mM DTT, 0.0005 mM DTT, or less than about 0.0001 mM DTT. In some embodiments, the amount of DTT can be undetectable.

A degradable bead can be useful to more quickly release an attached capture probe (e.g., a nucleic acid molecule, a spatial barcode sequence, and/or a primer) from the bead when the
10 appropriate stimulus is applied to the bead as compared to a bead that does not degrade. For example, for a species bound to an inner surface of a porous bead or in the case of an encapsulated species, the species can have greater mobility and accessibility to other species in solution upon degradation of the bead. In some embodiments, a species can also be attached to a degradable bead via a degradable linker (e.g., disulfide linker). The degradable linker can
15 respond to the same stimuli as the degradable bead or the two degradable species can respond to different stimuli. For example, a capture probe having one or more spatial barcodes can be attached, via a disulfide bond, to a polyacrylamide bead including cystamine. Upon exposure of the spatially barcoded bead to a reducing agent, the bead degrades and the capture probe having
20 the one or more spatial barcode sequences is released upon breakage of both the disulfide linkage between the capture probe and the bead and the disulfide linkages of the cystamine in the bead.

The addition of multiple types of labile bonds to a bead can result in the generation of a bead capable of responding to varied stimuli. Each type of labile bond can be sensitive to an associated stimulus (e.g., chemical stimulus, light, temperature, pH, enzymes, etc.) such that
25 release of reagents attached to a bead via each labile bond can be controlled by the application of the appropriate stimulus. Some non-limiting examples of labile bonds that can be coupled to a precursor or bead include an ester linkage (e.g., cleavable with an acid, a base, or hydroxylamine), a vicinal diol linkage (e.g., cleavable via sodium periodate), a Diels-Alder linkage (e.g., cleavable via heat), a sulfone linkage (e.g., cleavable via a base), a silyl ether
30 linkage (e.g., cleavable via an acid), a glycosidic linkage (e.g., cleavable via an amylase), a peptide linkage (e.g., cleavable via a protease), or a phosphodiester linkage (e.g., cleavable via a

nuclease (e.g., DNAase)). A bond can be cleavable via other nucleic acid molecule targeting enzymes, such as restriction enzymes (e.g., restriction endonucleases). Such functionality can be useful in controlled release of reagents from a bead. In some embodiments, another reagent including a labile bond can be linked to a bead after gel bead formation via, for example, an activated functional group of the bead as described above. In some embodiments, a gel bead including a labile bond is reversible. In some embodiments, a gel bead with a reversible labile bond is used to capture one or more regions of interest of a biological sample. For example, without limitation, a bead including a thermolabile bond can be heated by a light source (e.g., a laser) that causes a change in the gel bead that facilitates capture of a biological sample in contact with the gel bead. Capture probes having one or more spatial barcodes that are releasably, cleavably, or reversibly attached to the beads described herein include capture probes that are released or releasable through cleavage of a linkage between the capture probe and the bead, or that are released through degradation of the underlying bead itself, allowing the capture probes having the one or more spatial barcodes to be accessed or become accessible by other reagents, or both.

Beads can have different physical properties. Physical properties of beads can be used to characterize the beads. Non-limiting examples of physical properties of beads that can differ include size, shape, circularity, density, symmetry, and hardness. For example, beads can be of different sizes. Different sizes of beads can be obtained by using microfluidic channel networks configured to provide specific sized beads (e.g., based on channel sizes, flow rates, etc.). In some embodiments, beads have different hardness values that can be obtained by varying the concentration of polymer used to generate the beads. In some embodiments, a spatial barcode attached to a bead can be made optically detectable using a physical property of the capture probe. For example, a nucleic acid origami, such as a deoxyribonucleic acid (DNA) origami, can be used to generate an optically detectable spatial barcode. To do so, a nucleic acid molecule, or a plurality of nucleic acid molecules, can be folded to create two-and/or three-dimensional geometric shapes. The different geometric shapes can be optically detected.

In some embodiments, special types of nanoparticles with more than one distinct physical property can be used to make the beads physically distinguishable. For example, Janus particles with both hydrophilic and hydrophobic surfaces can be used to provide unique physical properties.

In some embodiments, a bead is able to identify multiple analytes (e.g., nucleic acids, proteins, chromatin, metabolites, drugs, gRNA, and lipids) from a single cell. In some embodiments, a bead is able to identify a single analyte from a single cell (e.g., mRNA).

A bead can have a tunable pore size. The pore size can be chosen to, for instance, retain
5 denatured nucleic acids. The pore size can be chosen to maintain diffusive permeability to exogenous chemicals such as sodium hydroxide (NaOH) and/or endogenous chemicals such as inhibitors. A bead can be formed of a biocompatible and/or biochemically compatible material, and/or a material that maintains or enhances cell viability. A bead can be formed from a material that can be depolymerized thermally, chemically, enzymatically, and/or optically.

10 In some embodiments, beads can be non-covalently loaded with one or more reagents. The beads can be non-covalently loaded by, for instance, subjecting the beads to conditions sufficient to swell the beads, allowing sufficient time for the reagents to diffuse into the interiors of the beads, and subjecting the beads to conditions sufficient to de-swell the beads. Swelling of the beads can be accomplished, for instance, by placing the beads in a thermodynamically
15 favorable solvent, subjecting the beads to a higher or lower temperature, subjecting the beads to a higher or lower ion concentration, and/or subjecting the beads to an electric field.

The swelling of the beads can be accomplished by various swelling methods. In some embodiments, swelling is reversible (e.g., by subjecting beads to conditions that promote de-swelling). In some embodiments, the de-swelling of the beads is accomplished, for instance, by
20 transferring the beads in a thermodynamically unfavorable solvent, subjecting the beads to lower or higher temperatures, subjecting the beads to a lower or higher ion concentration, and/or adding or removing an electric field. The de-swelling of the beads can be accomplished by various de-swelling methods. In some embodiments, de-swelling is reversible (e.g., subject beads to conditions that promote swelling). In some embodiments, the de-swelling of beads can include
25 transferring the beads to cause pores in the bead to shrink. The shrinking can then hinder reagents within the beads from diffusing out of the interiors of the beads. The hindrance created can be due to steric interactions between the reagents and the interiors of the beads. The transfer can be accomplished microfluidically. For instance, the transfer can be achieved by moving the beads from one co-flowing solvent stream to a different co-flowing solvent stream. The
30 swellability and/or pore size of the beads can be adjusted by changing the polymer composition of the bead.

A bead can include a polymer that is responsive to temperature so that when the bead is heated or cooled, the characteristics or dimensions of the bead can change. For example, a polymer can include poly(N-isopropylacrylamide). A gel bead can include poly(N-isopropylacrylamide) and when heated the gel bead can decrease in one or more dimensions (e.g., a cross-sectional diameter, multiple cross-sectional diameters). A temperature sufficient for changing one or more characteristics of the gel bead can be, for example, at least about 0 degrees Celsius (°C), 1°C, 2°C, 3°C, 4°C, 5°C, 10°C, or higher. For example, the temperature can be about 4°C. In some embodiments, a temperature sufficient for changing one or more characteristics of the gel bead can be, for example, at least about 25°C, 30°C, 35°C, 37°C, 40°C, 45°C, 50°C, or higher. For example, the temperature can be about 37°C.

Functionalization of beads for attachment of capture probes can be achieved through a wide range of different approaches, including, without limitation, activation of chemical groups within a polymer, incorporation of active or activatable functional groups in the polymer structure, or attachment at the pre-polymer or monomer stage in bead production. The bead can be functionalized to bind to targeted analytes, such as nucleic acids, proteins, carbohydrates, lipids, metabolites, peptides, or other analytes.

In some embodiments, a bead can contain molecular precursors (e.g., monomers or polymers), which can form a polymer network via polymerization of the molecular precursors. In some embodiments, a precursor can be an already polymerized species capable of undergoing further polymerization via, for example, a chemical cross-linkage. In some embodiments, a precursor can include one or more of an acrylamide or a methacrylamide monomer, oligomer, or polymer. In some embodiments, the bead can include prepolymers, which are oligomers capable of further polymerization. For example, polyurethane beads can be prepared using prepolymers. In some embodiments, a bead can contain individual polymers that can be further polymerized together (e.g., to form a co-polymer). In some embodiments, a bead can be generated via polymerization of different precursors, such that they include mixed polymers, co-polymers, and/or block co-polymers. In some embodiments, a bead can include covalent or ionic bonds between polymeric precursors (e.g., monomers, oligomers, and linear polymers), nucleic acid molecules (e.g., oligonucleotides), primers, and other entities. In some embodiments, covalent bonds can be carbon-carbon bonds or thioether bonds.

Cross-linking of polymers can be permanent or reversible, depending upon the particular cross-linker used. Reversible cross-linking can allow the polymer to linearize or dissociate under appropriate conditions. In some embodiments, reversible cross-linking can also allow for reversible attachment of a material bound to the surface of a bead. In some embodiments, a cross-linker can form a disulfide linkage. In some embodiments, a chemical cross-linker forming a disulfide linkage can be cystamine or a modified cystamine.

For example, where the polymer precursor material includes a linear polymer material, such as a linear polyacrylamide, PEG, or other linear polymeric material, the activation agent can include a cross-linking agent, or a chemical that activates a cross-linking agent within formed droplets. Likewise, for polymer precursors that include polymerizable monomers, the activation agent can include a polymerization initiator. For example, in certain embodiments, where the polymer precursor includes a mixture of acrylamide monomer with a N,N'-bis-(acryloyl)cystamine (BAC) comonomer, an agent such as tetraethylmethylenediamine (TEMED) can be provided, which can initiate the copolymerization of the acrylamide and BAC into a cross-linked polymer network, or other conditions sufficient to polymerize or gel the precursors. The conditions sufficient to polymerize or gel the precursors can include exposure to heating, cooling, electromagnetic radiation, and/or light.

Following polymerization or gelling, a polymer or gel can be formed. The polymer or gel can be diffusively permeable to chemical or biochemical reagents. The polymer or gel can be diffusively impermeable to macromolecular constituents. The polymer or gel can include one or more of disulfide cross-linked polyacrylamide, agarose, alginate, polyvinyl alcohol, polyethylene glycol (PEG)-diacrylate, PEG-acrylate, PEG-thiol, PEG-azide, PEG-alkyne, other acrylates, chitosan, hyaluronic acid, collagen, fibrin, gelatin, or elastin. The polymer or gel can include any other polymer or gel.

In some embodiments, disulfide linkages can be formed between molecular precursor units (e.g., monomers, oligomers, or linear polymers) or precursors incorporated into a bead and nucleic acid molecules (e.g., oligonucleotides, capture probes). Cystamine (including modified cystamines), for example, is an organic agent including a disulfide bond that can be used as a crosslinker agent between individual monomeric or polymeric precursors of a bead.

Polyacrylamide can be polymerized in the presence of cystamine or a species including cystamine (e.g., a modified cystamine) to generate polyacrylamide gel beads including disulfide

linkages (e.g., chemically degradable beads including chemically-reducible cross-linkers). The disulfide linkages can permit the bead to be degraded (or dissolved) upon exposure of the bead to a reducing agent.

5 In some embodiments, chitosan, a linear polysaccharide polymer, can be cross-linked with glutaraldehyde via hydrophilic chains to form a bead. Crosslinking of chitosan polymers can be achieved by chemical reactions that are initiated by heat, pressure, change in pH, and/or radiation.

10 In some embodiments, a bead can include an acrydite moiety, which in certain aspects can be used to attach one or more capture probes to the bead. In some embodiments, an acrydite moiety can refer to an acrydite analogue generated from the reaction of acrydite with one or more species (e.g., disulfide linkers, primers, other oligonucleotides, etc.), such as, without limitation, the reaction of acrydite with other monomers and cross-linkers during a polymerization reaction. Acrydite moieties can be modified to form chemical bonds with a species to be attached, such as a capture probe. Acrydite moieties can be modified with thiol
15 groups capable of forming a disulfide bond or can be modified with groups already including a disulfide bond. The thiol or disulfide (via disulfide exchange) can be used as an anchor point for a species to be attached or another part of the acrydite moiety can be used for attachment. In some embodiments, attachment can be reversible, such that when the disulfide bond is broken (e.g., in the presence of a reducing agent), the attached species is released from the bead. In some
20 embodiments, an acrydite moiety can include a reactive hydroxyl group that can be used for attachment of species.

In some embodiments, precursors (e.g., monomers or cross-linkers) that are polymerized to form a bead can include acrydite moieties, such that when a bead is generated, the bead also includes acrydite moieties. The acrydite moieties can be attached to a nucleic acid molecule (e.g.,
25 an oligonucleotide), which can include a priming sequence (e.g., a primer for amplifying target nucleic acids, random primer, primer sequence for messenger RNA) and/or one or more capture probes. The one or more capture probes can include sequences that are the same for all capture probes coupled to a given bead and/or sequences that are different across all capture probes coupled to the given bead. The capture probe can be incorporated into the bead. In some
30 embodiments, the capture probe can be incorporated or attached to the bead such that the capture probe retains a free 3' end. In some embodiments, the capture probe can be incorporated or

attached to the bead such that the capture probe retains a free 5' end. In some embodiments, beads can be functionalized such that each bead contains a plurality of different capture probes. For example, a bead can include a plurality of capture probes e.g., Capture Probe 1, Capture Probe 2, and Capture Probe 3, and each of Capture Probes 1, Capture Probes 2, and Capture
5 Probes 3 contain a distinct capture domain (e.g., capture domain of Capture Probe 1 includes a poly(dT) capture domain, capture domain of Capture Probe 2 includes a gene-specific capture domain, and capture domain of Capture Probe 3 includes a CRISPR-specific capture domain). By functionalizing beads to contain a plurality of different capture domains per bead, the level of multiplex capability for analyte detection can be improved.

10 In some embodiments, precursors (e.g., monomers or cross-linkers) that are polymerized to form a bead can include a functional group that is reactive or capable of being activated such that when it becomes reactive it can be polymerized with other precursors to generate beads including the activated or activatable functional group. The functional group can then be used to attach additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) to the
15 beads. For example, some precursors including a carboxylic acid (COOH) group can co-polymerize with other precursors to form a bead that also includes a COOH functional group. In some embodiments, acrylic acid (a species including free COOH groups), acrylamide, and bis(acryloyl)cystamine can be co-polymerized together to generate a bead including free COOH groups. The COOH groups of the bead can be activated (e.g., *via* 1-Ethyl-3-(3-
20 dimethylaminopropyl)carbodiimide (EDC) and N-Hydroxysuccinimide (NHS) or 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)) such that they are reactive (e.g., reactive to amine functional groups where EDC/NHS or DMTMM are used for activation). The activated COOH groups can then react with an appropriate species (e.g., a species including an amine functional group where the carboxylic acid groups are activated to be
25 reactive with an amine functional group) as a functional group on a moiety to be linked to the bead.

Beads including disulfide linkages in their polymeric network can be functionalized with additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) via reduction of some of the disulfide linkages to free thiols. The disulfide linkages can be reduced via, for
30 example, the action of a reducing agent (e.g., DTT, TCEP, etc.) to generate free thiol groups, without dissolution of the bead. Free thiols of the beads can then react with free thiols of a

species or a species including another disulfide bond (e.g., via thiol-disulfide exchange) such that the species can be linked to the beads (e.g., via a generated disulfide bond). In some embodiments, free thiols of the beads can react with any other suitable group. For example, free thiols of the beads can react with species including an acrydite moiety. The free thiol groups of the beads can react with the acrydite via Michael addition chemistry, such that the species including the acrydite is linked to the bead. In some embodiments, uncontrolled reactions can be prevented by inclusion of a thiol capping agent such as N-ethylmaleimide or iodoacetate.

Activation of disulfide linkages within a bead can be controlled such that only a small number of disulfide linkages are activated. Control can be exerted, for example, by controlling the concentration of a reducing agent used to generate free thiol groups and/or concentration of reagents used to form disulfide bonds in bead polymerization. In some embodiments, a low concentration of reducing agent (e.g., molecules of reducing agent:gel bead ratios) of less than or equal to about 1:100,000,000,000, less than or equal to about 1:10,000,000,000, less than or equal to about 1:1,000,000,000, less than or equal to about 1:100,000,000, less than or equal to about 1:10,000,000, less than or equal to about 1:1,000,000, less than or equal to about 1:100,000, or less than or equal to about 1:10,000) can be used for reduction. Controlling the number of disulfide linkages that are reduced to free thiols can be useful in ensuring bead structural integrity during functionalization. In some embodiments, optically-active agents, such as fluorescent dyes can be coupled to beads via free thiol groups of the beads and used to quantify the number of free thiols present in a bead and/or track a bead.

In some embodiments, addition of moieties to a bead after bead formation can be advantageous. For example, addition of a capture probe after bead formation can avoid loss of the species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) during chain transfer termination that can occur during polymerization. In some embodiments, smaller precursors (e.g., monomers or cross linkers that do not include side chain groups and linked moieties) can be used for polymerization and can be minimally hindered from growing chain ends due to viscous effects. In some embodiments, functionalization after bead synthesis can minimize exposure of species (e.g., oligonucleotides) to be loaded with potentially damaging agents (e.g., free radicals) and/or chemical environments. In some embodiments, the generated hydrogel can possess an upper critical solution temperature (UCST) that can permit temperature driven swelling and collapse of a bead. Such functionality can aid in oligonucleotide (e.g., a primer)

infiltration into the bead during subsequent functionalization of the bead with the oligonucleotide. Post-production functionalization can also be useful in controlling loading ratios of species in beads, such that, for example, the variability in loading ratio is minimized. Species loading can also be performed in a batch process such that a plurality of beads can be

5 functionalized with the species in a single batch.

Reagents can be encapsulated in beads during bead generation (e.g., during polymerization of precursors). Such reagents can or cannot participate in polymerization. Such reagents can be entered into polymerization reaction mixtures such that generated beads include the reagents upon bead formation. In some embodiments, such reagents can be added to the

10 beads after formation. Such reagents can include, for example, capture probes (e.g., oligonucleotides), reagents for a nucleic acid amplification reaction (e.g., primers, polymerases, dNTPs, co-factors (e.g., ionic co-factors), buffers) including those described herein, reagents for enzymatic reactions (e.g., enzymes, co-factors, substrates, buffers), reagents for nucleic acid modification reactions such as polymerization, ligation, or digestion, and/or reagents for template

15 preparation (e.g., tagmentation) for one or more sequencing platforms (e.g., Nextera® for Illumina®). Such reagents can include one or more enzymes described herein, including without limitation, polymerase, reverse transcriptase, restriction enzymes (e.g., endonuclease), transposase, ligase, proteinase K, DNase, etc. Such reagents can also or alternatively include one or more reagents such as lysis agents, inhibitors, inactivating agents, chelating agents, stimulus

20 agents. Trapping of such reagents can be controlled by the polymer network density generated during polymerization of precursors, control of ionic charge within the bead (e.g., via ionic species linked to polymerized species), or by the release of other species. Encapsulated reagents can be released from a bead upon bead degradation and/or by application of a stimulus capable of releasing the reagents from the bead.

25 In some embodiments, the beads can also include (e.g., encapsulate or have attached thereto) a plurality of capture probes that include spatial barcodes, and the optical properties of the spatial barcodes can be used for optical detection of the beads. For example, the absorbance of light by the spatial barcodes can be used to distinguish the beads from one another. In some embodiments, a detectable label can directly or indirectly attach to a spatial barcode and provide

30 optical detection of the bead. In some embodiments, each bead in a group of one or more beads

has a unique detectable label, and detection of the unique detectable label determines the location of the spatial barcode sequence associated with the bead.

Optical properties giving rise to optical detection of beads can be due to optical properties of the bead surface (e.g., a detectable label attached to the bead or the size of the bead), or optical properties from the bulk region of the bead (e.g., a detectable label incorporated during bead formation or an optical property of the bead itself). In some embodiments, a detectable label can be associated with a bead or one or more moieties coupled to the bead.

In some embodiments, the beads include a plurality of detectable labels. For example, a fluorescent dye can be attached to the surface of the beads and/or can be incorporated into the beads. Different intensities of the different fluorescent dyes can be used to increase the number of optical combinations that can be used to differentiate between beads. For example, if N is the number of fluorescent dyes (e.g., between 2 and 10 fluorescent dyes, such as 4 fluorescent dyes) and M is the possible intensities for the dyes (e.g., between 2 and 50 intensities, such as 20 intensities), then M^N are the possible distinct optical combinations. In one example, 4 fluorescent dyes with 20 possible intensities can be used to generate 160,000 distinct optical combinations.

One or more optical properties of the beads or biological contents, such as cells or nuclei, can be used to distinguish the individual beads or biological contents from other beads or biological contents. In some embodiments, the beads are made optically detectable by including a detectable label having optical properties to distinguish the beads from one another.

In some embodiments, optical properties of the beads can be used for optical detection of the beads. For example, without limitation, optical properties can include absorbance, birefringence, color, fluorescence, luminosity, photosensitivity, reflectivity, refractive index, scattering, or transmittance. For example, beads can have different birefringence values based on degree of polymerization, chain length, or monomer chemistry.

In some embodiments, nanobeads, such as quantum dots or Janus beads, can be used as optical labels or components thereof. For example, a quantum dot can be attached to a spatial barcode of a bead.

Optical labels of beads can provide enhanced spectral resolution to distinguish between beads with unique spatial barcodes (e.g., beads including unique spatial barcode sequences). In some embodiments, a first bead includes a first optical label and spatial barcodes each having a first spatial barcode sequence. A second bead includes a second optical label and spatial

barcodes each having a second spatial barcode sequence. The first optical label and second optical label can be different (e.g., provided by two different fluorescent dyes or the same fluorescent dye at two different intensities). The first and second spatial barcode sequences can be different nucleic acid sequences. In some embodiments, the beads can be imaged to identify the first and second optical labels, and the first and second optical barcodes can then be used to associate the first and second optical labels with the first and second spatial barcode sequences, respectively.

Optical labels can be included while generating the beads. For example, optical labels can be included in the polymer structure of a gel bead, or attached at the pre-polymer or monomer stage in bead production. In some embodiments, the beads include moieties that attach to one or more optical labels (e.g., at a surface of a bead and/or within a bead). In some embodiments, optical labels can be loaded into the beads with one or more reagents. For example, reagents and optical labels can be loaded into the beads by diffusion of the reagents (e.g., a solution of reagents including the optical barcodes). In some embodiments, optical labels can be included while preparing spatial barcodes. For example, spatial barcodes can be prepared by synthesizing molecules including barcode sequences (e.g., using a split pool or combinatorial approach). Optical labels can be attached to spatial barcodes prior to attaching the spatial barcodes to a bead. In some embodiments, optical labels can be included after attaching spatial barcodes to a bead. For example, optical labels can be attached to spatial barcodes coupled to the bead. In some embodiments, spatial barcodes or sequences thereof can be releasably or cleavably attached to the bead. Optical labels can be releasably or non-releasably attached to the bead. In some embodiments, a first bead (e.g., a bead including a plurality of spatial barcodes) can be coupled to a second bead including one or more optical labels. For example, the first bead can be covalently coupled to the second bead via a chemical bond. In some embodiments, the first bead can be non-covalently associated with the second bead.

The first and/or second bead can include a plurality of spatial barcodes. The plurality of spatial barcodes coupled to a given bead can include the same barcode sequences. Where both the first and second beads include spatial barcodes, the first and second beads can include spatial barcodes including the same barcode sequences or different barcode sequences.

Bead arrays containing captured analytes can be processed in bulk or partitioned into droplet emulsions for preparing sequencing libraries. In some embodiments, next generation

sequencing reads are clustered and correlated to the spatial position of the spatial barcode on the bead array. For example, the information can be computationally superimposed over a high-resolution image of the tissue section to identify the location(s), where the analytes were detected.

5 In some embodiments, de-cross linking can be performed to account for de-crosslinking chemistries that may be incompatible with certain barcoding/library prep biochemistry (e.g., presence of proteases). For example, a two-step process is possible. In the first step, beads can be provided in droplets such that DNA binds to the beads after the conventional de-crosslinking chemistry is performed. In the second step, the emulsion is broken and beads collected and then
10 re-encapsulated after washing for further processing.

In some embodiments, beads can be affixed or attached to a substrate using photochemical methods. For example, a bead can be functionalized with perfluorophenylazide silane (PFPA silane), contacted with a substrate, and then exposed to irradiation (see, e.g., Liu et al. (2006) *Journal of the American Chemical Society* 128, 14067–14072). For example,
15 immobilization of anthraquinone-functionalized substrates (see, e.g., Koch et al. (2000) *Bioconjugate Chem.* 11, 474–483, the entire contents of which are herein incorporated by reference).

The arrays can also be prepared by bead self-assembly. Each bead can be covered with hundreds of thousands of copies of a specific oligonucleotide. In some embodiments, each bead
20 can be covered with about 1,000 to about 1,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 1,000,000 to about 10,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 2,000,000 to about 3,000,000, about 3,000,000 to about 4,000,000, about 4,000,000 to about 5,000,000, about 5,000,000 to about 6,000,000, about 6,000,000 to about 7,000,000, about 7,000,000 to about 8,000,000, about 8,000,000 to
25 about 9,000,000, or about 9,000,000 to about 10,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 10,000,000 to about 100,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 100,000,000 to about 1,000,000,000 oligonucleotides. In some embodiments, each bead can be covered with about 1,000,000,000 to about 10,000,000,000 oligonucleotides. The beads can be irregularly
30 distributed across etched substrates during the array production process. During this process, the beads can be self-assembled into arrays (e.g., on a fiber-optic bundle substrate or a silica slide

substrate). In some embodiments, the beads irregularly arrive at their final location on the array. Thus, the bead location may need to be mapped or the oligonucleotides may need to be synthesized based on a predetermined pattern.

Beads can be affixed or attached to a substrate covalently, non-covalently, with adhesive, or a combination thereof. The attached beads can be, for example, layered in a monolayer, a bilayer, a trilayer, or as a cluster. As defined herein, a “monolayer” generally refers to an arrayed series of probes, beads, spots, dots, features, micro-locations, or islands that are affixed or attached to a substrate, such that the beads are arranged as one layer of single beads. In some embodiments, the beads are closely packed.

As defined herein, the phrase “substantial monolayer” or “substantially form(s) a monolayer” generally refers to (the formation of) an arrayed series of probes, beads, microspheres, spots, dots, features, micro-locations, or islands that are affixed or attached to a substrate, such that about 50% to about 99% (e.g., about 50% to about 98%) of the beads are arranged as one layer of single beads. This arrangement can be determined using a variety of methods, including microscopic imaging.

In some embodiments, the monolayer of beads is located in a predefined area on the substrate. For example, the predefined area can be partitioned with physical barriers, a photomask, divots in the substrate, or wells in the substrate.

As used herein, the term “reactive element” generally refers to a molecule or molecular moiety that can react with another molecule or molecular moiety to form a covalent bond. Reactive elements include, for example, amines, aldehydes, alkynes, azides, thiols, haloacetyls, pyridyl disulfides, hydrazides, carboxylic acids, alkoxyamines, sulfhydryls, maleimides, Michael acceptors, hydroxyls, and active esters. Some reactive elements, for example, carboxylic acids, can be treated with one or more activating agents (e.g., acylating agents, isourea-forming agents) to increase susceptibility of the reactive element to nucleophilic attack. Non-limiting examples of activating agents include *N*-hydroxysuccinimide, *N*-hydroxysulfosuccinimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide, dicyclohexylcarbodiimide, diisopropylcarbodiimide, 1-hydroxybenzotriazole, (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate, (benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate, 4-(*N,N*-dimethylamino)pyridine, and carbonyldiimidazole.

In some embodiments, the reactive element is bound directly to a bead. For example, hydrogel beads can be treated with an acrylic acid monomer to form acrylic acid-functionalized hydrogel beads. In some cases, the reactive element is bound indirectly to the bead via one or more linkers. As used herein, a “linker” generally refers to a multifunctional (e.g., bifunctional, trifunctional) reagent used for conjugating two or more chemical moieties. A linker can be a cleavable linker that can undergo induced dissociation. For example, the dissociation can be induced by a solvent (e.g., hydrolysis and solvolysis); by irradiation (e.g., photolysis); by an enzyme (e.g., enzymolysis); or by treatment with a solution of specific pH (e.g., pH 4, 5, 6, 7, or 8).

In some embodiments, the reactive element is bound directly to a substrate. For example, a glass slide can be coated with (3-aminopropyl)triethoxysilane. In some embodiments, the reactive element is bound indirectly to a substrate via one or more linkers.

Methods for Covalently Bonding Beads to a Substrate

Provided herein are methods for the covalent bonding of beads (e.g., optically labeled beads, hydrogel beads, microsphere beads) to a substrate.

In some embodiments, the beads are coupled to a substrate via a covalent bond between a first reactive element and a second reactive element. In some embodiments, the covalently-bound beads substantially form a monolayer of beads (e.g., hydrogel beads, microsphere beads) on the substrate.

In some embodiments, the beads are functionalized with a first reactive element, which is directly bound to the beads. In some embodiments, the beads are functionalized with a first reactive element, which is indirectly bound to the beads via a linker. In some embodiments, the linker is a benzophenone. In some embodiments, the linker is an amino methacrylamide. For example, the linker can be 3-aminopropyl methacrylamide. In some embodiments, the linker is a PEG linker. In some embodiments, the linker is a cleavable linker.


In some embodiments, the substrate is functionalized with a second reactive element, which is directly bound to the substrate. In some embodiments, the substrate is functionalized with a second reactive element, which is indirectly bound to the beads via a linker. In some embodiments, the linker is a benzophenone. For example, the linker can be benzophenone. In some embodiments, the linker is an amino methacrylamide. For example, the linker can be 3-

aminopropyl methacrylamide. In some embodiments, the linker is a PEG linker. In some embodiments, the linker is a cleavable linker.

In some embodiments, the substrate is a glass slide. In some embodiments, the substrate is a pre-functionalized glass slide.

5 In some embodiments, about 99% of the covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 50% to about 98% form a monolayer of beads on the substrate. For example, about 50% to about 95%, about 50% to about 90%, about 50% to about 85%, about 50% to about 80%, about 50% to about 75%, about 50% to about 70%, about 50% to about 65%, about 50% to about 60%, or about 50% to about 55% of the
10 covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 98%, about 60% to about 98%, about 65% to about 98%, about 70% to about 98%, about 75% to about 98%, about 80% to about 98%, about 85% to about 98%, about 90% to about 95%, or about 95% to about 98% of the covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 95%, about 60% to about
15 90%, about 65% to about 95%, about 70% to about 95%, about 75% to about 90%, about 75% to about 95%, about 80% to about 90%, about 80% to about 95%, about 85% to about 90%, or about 85% to about 95% of the covalently-bound beads for a monolayer of beads on the substrate.

In some embodiments, at least one of the first reactive element and the second reactive
20 element is selected from the group consisting of:



wherein

R¹ is selected from H, C₁-C₆ alkyl, or -SO₃;

R² is C₁-C₆ alkyl; and


X is a halo moiety.

In some embodiments, at least one of the first reactive element or the second reactive

element comprises  or  ;

wherein the  indicates the point of attachment of the first reactive element or the second reactive element to the bead (e.g., hydrogel bead or microsphere bead) or to the substrate.

5 In some embodiments, at least one of the first reactive element or the second reactive element is selected from the group consisting of:




wherein

R^1 is selected from H, C_1 – C_6 alkyl, or $-SO_3$;


10 R^2 is C_1 – C_6 alkyl; and

X is a halo moiety.



In some embodiments, at least one of the first reactive element or the second reactive


element comprises  , wherein R^1 is selected from H, C_1 – C_6 alkyl, or $-SO_3$. In some
embodiments, R^1 is H. In some embodiments, R^1 is C_1 – C_6 alkyl. In some embodiments, R^1 is –
15 SO_3 .


In some embodiments, at least one of the first reactive element or the second reactive

element comprises  , wherein R^2 is C_1 – C_6 alkyl. In some embodiments, R^2 is methyl.

In some embodiments, at least one of the first reactive element or the second reactive


element comprises  . In some embodiments,  can be reacted with an activating




agent to form an active ester. In some embodiments, the active ester is . In some


- embodiments, the activating agent is an acylating agent (e.g., *N*-hydroxysuccinimide and *N*-hydroxysulfosuccinimide). In some embodiments, the activating agent is an *O*-acylisourea-forming agent (e.g., 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC),
- 5 dicyclohexylcarbodiimide, and diisopropylcarbodiimide). In some embodiments, the activating agent is a combination of at least one acylating agent and at least one *O*-isourea-forming agents (e.g., *N*-hydroxysuccinimide (NHS), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC), *N*-hydroxysulfosuccinimide (sulfo-NHS), and a combination thereof).


In some embodiments, at least one of the first reactive element or the second reactive




- 10 element comprises .


In some embodiments, at least one of the first reactive element or the second reactive




element comprises , wherein X is a halo moiety. For example, X is chloro, bromo, or iodo.


In some embodiments, at least one of the first reactive element or the second reactive




- 15 element comprises .


In some embodiments, at least one of the first reactive element or the second reactive




element comprises .

In some embodiments, at least one of the first reactive element or the second reactive



element comprises .

In some embodiments, at least one of the first reactive element or the second reactive element is selected from the group consisting of:



wherein


5 R^3 is H or C₁–C₆ alkyl; and

R^4 is H or trimethylsilyl.

In some embodiments, at least one of the first reactive element or the second reactive

element comprises $\begin{array}{c} \text{---} \\ \backslash \\ \text{C} \equiv \text{C} \end{array} \text{---R}^4$, wherein R^4 is H or trimethylsilyl. In some embodiments, R^4 is H.

10 In some embodiments, at least one of the first reactive element or the second reactive element is selected from the group consisting of:



wherein R^3 is H or C₁–C₆ alkyl. In some embodiments, R^3 is H. In some embodiments, R^3 is C₁–C₆ alkyl.

In some embodiments, at least one of the first reactive element or the second reactive


15 element comprises $\begin{array}{c} \text{R}^3 \\ | \\ \text{---NH} \end{array}$, wherein R^3 is H or C₁–C₆ alkyl. In some embodiments, R^3 is H. In some embodiments, R^3 is C₁–C₆ alkyl.

In some embodiments, at least one of the first reactive elements or the second reactive elements comprises ---SH .

In some embodiments, at least one of the first reactive elements or the second reactive

20 elements comprises $\begin{array}{c} \text{---CH}_2\text{---} \\ | \\ \text{H} \end{array} \text{C(=O)---}$.

In some embodiments, one of the first reactive elements or the second reactive elements is selected from the group consisting of:




wherein

R^1 is selected from H, C_1 – C_6 alkyl, or $-SO_3$;

R^2 is C_1 – C_6 alkyl;

5 X is a halo moiety;

and the other of the first reactive element or the second reactive element is selected from the group consisting of:




wherein

10 R^3 is H or C_1 – C_6 alkyl; and

R^4 is H or trimethylsilyl.


In some embodiments, one of the first reactive elements or the second reactive elements

is selected from the group consisting of  and , wherein R^3 is H or C_1 – C_6 alkyl;

and the other of the first reactive element or the second reactive element is , wherein

15 R^4 is H or trimethylsilyl. In some embodiments, R^3 is H. In some embodiments, R^3 is C_1 – C_6 alkyl. In some embodiments, R^4 is H. In some embodiments, R^4 is trimethylsilyl.

In some embodiments, one of the first reactive element or the second reactive element is selected from the group consisting of:




wherein

R^1 is selected from H, C_1 – C_6 alkyl, or $-SO_3$;

R^2 is C_1 – C_6 alkyl;


5 X is a halo moiety;

and the other of the first reactive element or the second reactive element is selected from the group consisting of:



10 wherein R^3 is H or C_1 – C_6 alkyl. In some embodiments, R^1 is H. In some embodiments, R^1 is C_1 – C_6 alkyl. In some embodiments, R^1 is $-SO_3$. In some embodiments, R^2 is methyl. In some embodiments, X is iodo. In some embodiments, R^3 is H. In some embodiments, R^3 is C_1 – C_6 alkyl.

In some embodiments, one of the first reactive elements or the second reactive elements is selected from the group consisting of:




15


wherein

R^1 is selected from H, C_1 – C_6 alkyl, or $-SO_3$;

R^2 is C_1 – C_6 alkyl;

and the other of the first reactive elements or the second reactive elements comprises , wherein R³ is H or C₁–C₆ alkyl. In some embodiments, R¹ is H. In some embodiments, R¹ is C₁–C₆ alkyl. In some embodiments, R¹ is –SO₃. In some embodiments, R² is methyl. In some embodiments, R³ is H. In some embodiments, R³ is C₁–C₆ alkyl.



5 In some embodiments, one of the first reactive element or the second reactive element is selected from the group consisting of:




wherein X is a halo moiety;

and the other of the first reactive element or the second reactive element comprises . In some embodiments, X is bromo. In some embodiments, X is iodo.

In some embodiments, one of the first reactive element or the second reactive element is

selected from the group consisting of  and .

and the other of the first reactive element or the second reactive element comprises .

The term “halo” refers to fluoro (F), chloro (Cl), bromo (Br), or iodo (I).

15 The term “alkyl” refers to a hydrocarbon chain that may be a straight chain or branched chain, containing the indicated number of carbon atoms. For example, C₁₋₁₀ indicates that the group may have from 1 to 10 (inclusive) carbon atoms in it. Non-limiting examples include methyl, ethyl, *iso*-propyl, *tert*-butyl, *n*-hexyl.

The term “haloalkyl” refers to an alkyl, in which one or more hydrogen atoms is/are replaced with an independently selected halo.

The term “alkoxy” refers to an -O-alkyl radical (e.g., -OCH₃).

The term “alkylene” refers to a divalent alkyl (e.g., -CH₂-).

The term “alkenyl” refers to a hydrocarbon chain that may be a straight chain or branched chain having one or more carbon-carbon double bonds. The alkenyl moiety contains the indicated

number of carbon atoms. For example, C₂₋₆ indicates that the group may have from 2 to 6 (inclusive) carbon atoms in it.

The term “alkynyl” refers to a hydrocarbon chain that may be a straight chain or branched chain having one or more carbon-carbon triple bonds. The alkynyl moiety contains the indicated
5 number of carbon atoms. For example, C₂₋₆ indicates that the group may have from 2 to 6 (inclusive) carbon atoms in it.

The term “aryl” refers to a 6-20 carbon mono-, bi-, tri- or polycyclic group wherein at least one ring in the system is aromatic (e.g., 6-carbon monocyclic, 10-carbon bicyclic, or 14-carbon
10 tricyclic aromatic ring system); and wherein 0, 1, 2, 3, or 4 atoms of each ring may be substituted by a substituent. Examples of aryl groups include phenyl, naphthyl, tetrahydronaphthyl, and the like.

Methods for Non-Covalently Bonding Beads to a Substrate

Provided herein are methods for the non-covalent bonding of beads (e.g., optically-
15 labeled beads, hydrogel beads, or microsphere beads) to a substrate.

In some embodiments, beads are coupled to a substrate via a non-covalent bond between a first affinity group and a second affinity group. In some embodiments, the non-covalently-bound beads substantially form a monolayer of beads (e.g., hydrogel beads, microsphere beads) on the substrate.

In some embodiments, the beads are functionalized with a first affinity group, which is
20 directly bound to the beads. In some embodiments, the beads are functionalized with a first affinity group, which is indirectly bound to the beads via a linker. In some embodiments, the linker is a benzophenone. In some embodiments, the linker is an amino methacrylamide. For example, the linker can be 3-aminopropyl methacrylamide. In some embodiments, the linker is a
25 PEG linker. In some embodiments, the linker is a cleavable linker.

In some embodiments, the substrate is functionalized with a second affinity group, which is directly bound to the substrate. In some embodiments, the substrate is functionalized with a second affinity group, which is indirectly bound to the beads via a linker. In some embodiments, the linker is a benzophenone. In some embodiments, the linker is an amino methacrylamide. For
30 example, the linker can be 3-aminopropyl methacrylamide. In some embodiments, the linker is a PEG linker. In some embodiments, the linker is a cleavable linker.

In some embodiments the first affinity group or the second affinity group is biotin, and the other of the first affinity group or the second affinity group is streptavidin.

In some embodiments, about 99% of the non-covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 50% to about 98% form a monolayer of beads on the substrate. For example, about 50% to about 95%, about 50% to about 90%, about 50% to about 85%, about 50% to about 80%, about 50% to about 75%, about 50% to about 70%, about 50% to about 65%, about 50% to about 60%, or about 50% to about 55% of the non-covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 98%, about 60% to about 98%, about 65% to about 98%, about 70% to about 98%, about 75% to about 98%, about 80% to about 98%, about 85% to about 98%, about 90% to about 95%, or about 95% to about 98% of the non-covalently-bound beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 95%, about 60% to about 90%, about 65% to about 95%, about 70% to about 95%, about 75% to about 90%, about 75% to about 95%, about 80% to about 90%, about 80% to about 95%, about 85% to about 90%, or about 85% to about 95% of the non-covalently-bound beads form a monolayer of beads on the substrate.

In some embodiments, the monolayer of beads is formed in a predefined area on the substrate. In some embodiments, the predefined area is partitioned with physical barriers. For example, divots or wells in the substrate. In some embodiments, the predefined area is partitioned using a photomask. For example, the substrate is coated with a photo-activated solution, dried, then irradiated under a photomask. In some embodiments, the photo-activated solution is UV-activated.

As used herein, an “adhesive” generally refers to a substance used for sticking objects or materials together. Adhesives include, for example, glues, pastes, liquid tapes, epoxy, bioadhesives, gels, and mucilage. In some embodiments, an adhesive is liquid tape. In some embodiments, the adhesive is glue.

In some embodiments, beads are adhered to a substrate using an adhesive (e.g., liquid tape, glue, paste). In some embodiments, the adhered beads substantially form a monolayer of beads on the substrate (e.g., a glass slide). In some embodiments, the beads are hydrogel beads. In some embodiments, the beads are microsphere beads. In some embodiments, the beads are coated with the adhesive, and then the beads are contacted with the substrate. In some

embodiments, the substrate is coated with the adhesive, and then the substrate is contacted with the beads. In some embodiments, both the substrate is coated with the adhesive and the beads are coated with the adhesive, and then the beads and substrate are contacted with one another.

In some embodiments, about 99% of the adhered beads form a monolayer of beads on the substrate. In some embodiments, about 50% to about 98% form a monolayer of beads on the substrate. For example, about 50% to about 95%, about 50% to about 90%, about 50% to about 85%, about 50% to about 80%, about 50% to about 75%, about 50% to about 70%, about 50% to about 65%, about 50% to about 60%, or about 50% to about 55% of the adhered beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 98%, about 60% to about 98%, about 65% to about 98%, about 70% to about 98%, about 75% to about 98%, about 80% to about 98%, about 85% to about 98%, about 90% to about 95%, or about 95% to about 98% of the adhered beads form a monolayer of beads on the substrate. In some embodiments, about 55% to about 95%, about 60% to about 90%, about 65% to about 95%, about 70% to about 95%, about 75% to about 90%, about 75% to about 95%, about 80% to about 90%, about 80% to about 95%, about 85% to about 90%, or about 85% to about 95% of the adhered beads form a monolayer of beads on the substrate.

In some embodiments, beads can be deposited onto a biological sample such that the deposited beads form a monolayer of beads on the biological sample (e.g., over or under the biological sample). In some embodiments, beads deposited on the substrate can self-assemble into a monolayer of beads that saturate the intended surface area of the biological sample under investigation. In this approach, bead arrays can be designed, formulated, and prepared to evaluate a plurality of analytes from a biological sample of any size or dimension. In some embodiments, the concentration or density of beads (e.g., gel beads) applied to the biological sample is such that the area as a whole, or one or more regions of interest in the biological sample, is saturated with a monolayer of beads. In some embodiments, the beads are contacted with the biological sample by pouring, pipetting, spraying, and the like, onto the biological sample. Any suitable form of bead deposition can be used.

In some embodiments, the biological sample can be confined to a specific region or area of the array. For example, a biological sample can be affixed to a glass slide and a chamber, gasket, or cage positioned over the biological sample to act as a containment region or frame within which the beads are deposited. As will be apparent, the density or concentration of beads

needed to saturate an area or biological sample can be readily determined by one of ordinary skill in the art (e.g., through microscopic visualization of the beads on the biological sample). In some embodiments, the bead array contains microfluidic channels to direct reagents to the spots or beads of the array.

5

Feature Geometric Attributes

Features on an array can have a variety of sizes. In some embodiments, a feature of an array can have a diameter or maximum dimension between 1 μm to 100 μm . For example, between 1 μm to 10 μm , 1 μm to 20 μm , 1 μm to 30 μm , 1 μm to 40 μm , 1 μm to 50 μm , 1 μm to 60 μm , 1 μm to 70 μm , 1 μm to 80 μm , 1 μm to 90 μm , 90 μm to 100 μm , 80 μm to 100 μm , 70 μm to 100 μm , 60 μm to 100 μm , 50 μm to 100 μm , 40 μm to 100 μm , 30 μm to 100 μm , 20 μm to 100 μm , or 10 μm to 100 μm . In some embodiments, the feature has a diameter or maximum dimension between 30 μm to 100 μm , 40 μm to 90 μm , 50 μm to 80 μm , 60 μm to 70 μm , or any range within the disclosed sub-ranges. In some embodiments, the feature has a diameter or maximum dimension no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , 10 μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm . In some embodiments, the feature has a diameter or maximum dimension of approximately 65 μm .

In some embodiments, the size and/or shape of a plurality of features of an array are approximately uniform. In some embodiments, the size and/or shape of a plurality of features of an array is not uniform. For example, in some embodiments, features in an array can have an average cross-sectional dimension, and a distribution of cross-sectional dimensions among the features can have a full-width and half-maximum value of 0% or more (e.g., 5% or more, 10% or more, 20% or more, 30% or more, 40% or more, 50% or more, 70% or more, or 100% or more) of the average cross-sectional dimension for the distribution.

In certain embodiments, features in an array can have an average cross-sectional dimension of between about 1 μm and about 10 μm . This range in average feature cross-sectional dimension corresponds to the approximate diameter of a single mammalian cell. Thus, an array of such features can be used to detect analytes at, or below, mammalian single-cell resolution.

30

In some embodiments, a plurality of features has a mean diameter or mean maximum dimension of about 0.1 μm to about 100 μm (e.g., about 0.1 μm to about 5 μm , about 1 μm to about 10 μm , about 1 μm to about 20 μm , about 1 μm to about 30 μm , about 1 μm to about 40 μm , about 1 μm to about 50 μm , about 1 μm to about 60 μm , about 1 μm to about 70 μm , about 1 μm to about 80 μm , about 1 μm to about 90 μm , about 90 μm to about 100 μm , about 80 μm to about 100 μm , about 70 μm to about 100 μm , about 60 μm to about 100 μm , about 50 μm to about 100 μm , about 40 μm to about 100 μm , about 30 μm to about 100 μm , about 20 μm to about 100 μm , or about 10 μm to about 100 μm). In some embodiments, the plurality of features has a mean diameter or mean maximum dimension between 30 μm to 100 μm , 40 μm to 90 μm , 50 μm to 80 μm , 60 μm to 70 μm , or any range within the disclosed sub-ranges. In some embodiments, the plurality of features has a mean diameter or a mean maximum dimension no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , 10 μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm . In some embodiments, the plurality of features has a mean average diameter or a mean maximum dimension of approximately 65 μm .

In some embodiments, where the feature is a bead, the bead can have a diameter or maximum dimension no larger than 100 μm (e.g., no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , 10 μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm).

In some embodiments, a plurality of beads has an average diameter no larger than 100 μm . In some embodiments, a plurality of beads has an average diameter or maximum dimension no larger than 95 μm , 90 μm , 85 μm , 80 μm , 75 μm , 70 μm , 65 μm , 60 μm , 55 μm , 50 μm , 45 μm , 40 μm , 35 μm , 30 μm , 25 μm , 20 μm , 15 μm , 14 μm , 13 μm , 12 μm , 11 μm , 10 μm , 9 μm , 8 μm , 7 μm , 6 μm , 5 μm , 4 μm , 3 μm , 2 μm , or 1 μm .

In some embodiments, the volume of the bead can be at least about 1 μm^3 , e.g., at least 1 μm^3 , 2 μm^3 , 3 μm^3 , 4 μm^3 , 5 μm^3 , 6 μm^3 , 7 μm^3 , 8 μm^3 , 9 μm^3 , 10 μm^3 , 12 μm^3 , 14 μm^3 , 16 μm^3 , 18 μm^3 , 20 μm^3 , 25 μm^3 , 30 μm^3 , 35 μm^3 , 40 μm^3 , 45 μm^3 , 50 μm^3 , 55 μm^3 , 60 μm^3 , 65 μm^3 , 70 μm^3 , 75 μm^3 , 80 μm^3 , 85 μm^3 , 90 μm^3 , 95 μm^3 , 100 μm^3 , 125 μm^3 , 150 μm^3 , 175 μm^3 , 200 μm^3 , 250 μm^3 , 300 μm^3 , 350 μm^3 , 400 μm^3 , 450 μm^3 , 500 μm^3 , 550 μm^3 , 600 μm^3 , 650 μm^3 , 700 μm^3 , 750 μm^3 , 800 μm^3 , 850 μm^3 , 900 μm^3 , 950 μm^3 , 1000 μm^3 , 1200 μm^3 , 1400 μm^3 .

μm^3 , 1600 μm^3 , 1800 μm^3 , 2000 μm^3 , 2200 μm^3 , 2400 μm^3 , 2600 μm^3 , 2800 μm^3 , 3000 μm^3 , or greater.

In some embodiments, the bead can have a volume of between about 1 μm^3 and 100 μm^3 , such as between about 1 μm^3 and 10 μm^3 , between about 10 μm^3 and 50 μm^3 , or between about 50 μm^3 and 100 μm^3 . In some embodiments, the bead can include a volume of between about 100 μm^3 and 1000 μm^3 , such as between about 100 μm^3 and 500 μm^3 or between about 500 μm^3 and 1000 μm^3 . In some embodiments, the bead can include a volume between about 1000 μm^3 and 3000 μm^3 , such as between about 1000 μm^3 and 2000 μm^3 or between about 2000 μm^3 and 3000 μm^3 . In some embodiments, the bead can include a volume between about 1 μm^3 and 3000 μm^3 , such as between about 1 μm^3 and 2000 μm^3 , between about 1 μm^3 and 1000 μm^3 , between about 1 μm^3 and 500 μm^3 , or between about 1 μm^3 and 250 μm^3 .

The bead can include one or more cross-sections that can be the same or different. In some embodiments, the bead can have a first cross-section that is different from a second cross-section. The bead can have a first cross-section that is at least about 0.0001 micrometer, 0.001 micrometer, 0.01 micrometer, 0.1 micrometer, or 1 micrometer. In some embodiments, the bead can include a cross-section (e.g., a first cross-section) of at least about 1 micrometer (μm), 2 μm , 3 μm , 4 μm , 5 μm , 6 μm , 7 μm , 8 μm , 9 μm , 10 μm , 11 μm , 12 μm , 13 μm , 14 μm , 15 μm , 16 μm , 17 μm , 18 μm , 19 μm , 20 μm , 25 μm , 30 μm , 35 μm , 40 μm , 45 μm , 50 μm , 55 μm , 60 μm , 65 μm , 70 μm , 75 μm , 80 μm , 85 μm , 90 μm , 100 μm , 120 μm , 140 μm , 160 μm , 180 μm , 200 μm , 250 μm , 300 μm , 350 μm , 400 μm , 450 μm , 500 μm , 550 μm , 600 μm , 650 μm , 700 μm , 750 μm , 800 μm , 850 μm , 900 μm , 950 μm , 1 millimeter (mm), or greater. In some embodiments, the bead can include a cross-section (e.g., a first cross-section) of between about 1 μm and 500 μm , such as between about 1 μm and 100 μm , between about 100 μm and 200 μm , between about 200 μm and 300 μm , between about 300 μm and 400 μm , or between about 400 μm and 500 μm . For example, the bead can include a cross-section (e.g., a first cross-section) of between about 1 μm and 100 μm . In some embodiments, the bead can have a second cross-section that is at least about 1 μm . For example, the bead can include a second cross-section of at least about 1 micrometer (μm), 2 μm , 3 μm , 4 μm , 5 μm , 6 μm , 7 μm , 8 μm , 9 μm , 10 μm , 11 μm , 12 μm , 13 μm , 14 μm , 15 μm , 16 μm , 17 μm , 18 μm , 19 μm , 20 μm , 25 μm , 30 μm , 35 μm , 40 μm , 45 μm , 50 μm , 55 μm , 60 μm , 65 μm , 70 μm , 75 μm , 80 μm , 85 μm , 90 μm , 100 μm , 120 μm , 140 μm , 160 μm , 180 μm , 200 μm , 250 μm , 300 μm , 350 μm , 400 μm , 450 μm , 500 μm , 550 μm , 600 μm ,

650 μm , 700 μm , 750 μm , 800 μm , 850 μm , 900 μm , 950 μm , 1 millimeter (mm), or greater. In some embodiments, the bead can include a second cross-section of between about 1 μm and 500 μm , such as between about 1 μm and 100 μm , between about 100 μm and 200 μm , between about 200 μm and 300 μm , between about 300 μm and 400 μm , or between about 400 μm and 500 μm . For example, the bead can include a second cross-section of between about 1 μm and 100 μm .

In some embodiments, beads can be of a nanometer scale (e.g., beads can have a diameter or maximum cross-sectional dimension of about 100 nanometers (nm) to about 900 nanometers (nm) (e.g., 850 nm or less, 800 nm or less, 750 nm or less, 700 nm or less, 650 nm or less, 600 nm or less, 550 nm or less, 500 nm or less, 450 nm or less, 400 nm or less, 350 nm or less, 300 nm or less, 250 nm or less, 200 nm or less, 150 nm or less). A plurality of beads can have an average diameter or average maximum cross-sectional dimension of about 100 nanometers (nm) to about 900 nanometers (nm) (e.g., 850 nm or less, 800 nm or less, 750 nm or less, 700 nm or less, 650 nm or less, 600 nm or less, 550 nm or less, 500 nm or less, 450 nm or less, 400 nm or less, 350 nm or less, 300 nm or less, 250 nm or less, 200 nm or less, 150 nm or less). In some embodiments, a bead has a diameter or size that is about the size of a single cell (e.g., a single cell under evaluation).

Beads can be of uniform size or heterogeneous size. "Polydispersity" generally refers to heterogeneity of sizes of molecules or particles. The polydispersity (PDI) can be calculated using the equation $\text{PDI} = \text{Mw}/\text{Mn}$, where Mw is the weight-average molar mass and Mn is the number-average molar mass. In certain embodiments, beads can be provided as a population or plurality of beads having a relatively monodisperse size distribution. Where it can be desirable to provide relatively consistent amounts of reagents, maintaining relatively consistent bead characteristics, such as size, can contribute to the overall consistency.

In some embodiments, the beads provided herein can have size distributions that have a coefficient of variation in their cross-sectional dimensions of less than 50%, less than 40%, less than 30%, less than 20%, less than 15%, less than 10%, less than 5%, or lower. In some embodiments, a plurality of beads provided herein has a polydispersity index of less than 50%, less than 45%, less than 40%, less than 35%, less than 30%, less than 25%, less than 20%, less than 15%, less than 10%, less than 5%, or lower.

Array Geometric Attributes

In some embodiments, an array includes a plurality of features. For example, an array includes between 4,000 and 10,000 features, or any range within 4,000 to 6000 features. For example, an array includes between 4,000 to 4,400 features, 4,000 to 4,800 features, 4,000 to 5,200 features, 4,000 to 5,600 features, 5,600 to 6,000 features, 5,200 to 6,000 features, 4,800 to 6,000 features, or 4,400 to 6,000 features. In some embodiments, the array includes between 4,100 and 5,900 features, between 4,200 and 5,800 features, between 4,300 and 5,700 features, between 4,400 and 5,600 features, between 4,500 and 5,500 features, between 4,600 and 5,400 features, between 4,700 and 5,300 features, between 4,800 and 5,200 features, between 4,900 and 5,100 features, or any range within the disclosed sub-ranges. For example, the array can include about 4,000 features, about 4,200 features, about 4,400 features, about 4,800 features, about 5,000 features, about 5,200 features, about 5,400 features, about 5,600 features, or about 6,000 features. In some embodiments, the array comprises at least 4,000 features. In some embodiments, the array includes approximately 5,000 features.

In some embodiments, features within an array have an irregular arrangement or relationship to one another, such that no discernable pattern or regularity is evident in the geometrical spacing relationships among the features. For example, features within an array may be positioned randomly with respect to one another. Alternatively, features within an array may be positioned irregularly, but the spacings may be selected deterministically to ensure that the resulting arrangement of features is irregular.

In some embodiments, features within an array are positioned regularly with respect to one another to form a pattern. A wide variety of different patterns of features can be implemented in arrays. Examples of such patterns include, but are not limited to, square arrays of features, rectangular arrays of features, hexagonal arrays of features (including hexagonal close-packed arrays), radial arrays of features, spiral arrays of features, triangular arrays of features, and more generally, any array in which adjacent features in the array are reached from one another by regular increments in linear and/or angular coordinate dimensions.

In some embodiments, features within an array are positioned with a degree of regularity with respect to one another such that the array of features is neither perfectly regular nor perfectly irregular (i.e., the array is “partially regular”). For example, in some embodiments, adjacent features in an array can be separated by a displacement in one or more linear and/or angular coordinate dimensions that is 10% or more (e.g., 20% or more, 30% or more, 40% or

more, 50% or more, 60% or more, 70% or more, 80% or more, 90% or more, 100% or more, 110% or more, 120% or more, 130% or more, 140% or more, 150% or more, 160% or more, 170% or more, 180% or more, 190% or more, 200% or more) of an average displacement or a nominal displacement between adjacent features in the array. In certain embodiments, the distribution of displacements (linear and/or angular) between adjacent features in an array has a full-width at half-maximum of between 0% and 200% (e.g., between 0% and 100%, between 0% and 75%, between 0% and 50%, between 0% and 25%, between 0% and 15%, between 0% and 10%) of an average displacement or nominal displacement between adjacent features in the array.

In some embodiments, arrays of features can have a variable geometry. For example, a first subset of features in an array can be arranged according to a first geometrical pattern, and a second subset of features in the array can be arranged according to a second geometrical pattern that is different from the first pattern. Any of the patterns described above can correspond to the first and/or second geometrical patterns, for example.

In general, arrays of different feature densities can be prepared by adjusting the spacing between adjacent features in the array. In some embodiments, the geometric center-to-center spacing between adjacent features in an array is between 100 nm and 100 μm . For example, the center-to-center spacing can be between 20 μm to 40 μm , 20 μm to 60 μm , 20 μm to 80 μm , 80 μm to 100 μm , 60 μm to 100 μm , or 40 μm to 100 μm . In some embodiments, the center-to-center spacing between adjacent array features is between 30 μm and 100 μm , 40 μm and 90 μm , 50 μm and 80 μm , 60 μm and 70 μm , 80 μm and 120 μm , or any range within the disclosed sub-ranges. In some embodiments, the center-to-center spacing between adjacent array features of a feature of an array is approximately 65 μm .

In some embodiments, an array of features can have a spatially varying resolution. In general, an array with a spatially varying resolution is an array in which the center-to-center spacing (along linear, angular, or both linear and angular coordinate dimensions) between adjacent features in the array varies. Such arrays can be useful in a variety of applications. For example, in some embodiments, depending upon the spatial resolution at which the sample is to be investigated, the sample can be selectively associated with the portion of the array that corresponds approximately to the desired spatial resolution of the measurement.

Arrays of spatially varying resolution can be implemented in a variety of ways. In some embodiments, for example, the center-to-center spacing between adjacent features in the array varies continuously along one or more linear and/or angular coordinate directions. Thus, for a rectangular array, the spacing between successive rows of features, between successive columns of features, or between both successive rows and successive columns of features, can vary continuously.

In certain embodiments, arrays of spatially varying resolution can include discrete domains with populations of features. Within each domain, adjacent features can have regular center-to-center spacings. Thus, for example, an array can include a first domain within which adjacent features are spaced from one another along linear and/or angular coordinate dimensions by a first set of uniform coordinate displacements, and a second domain within which adjacent features are spaced from one another along linear and/or angular coordinate dimensions by a second set of uniform coordinate displacements. The first and second sets of displacements differ in at least one coordinate displacement, such that adjacent features in the two domains are spaced differently, and the resolution of the array in the first domain is therefore different from the resolution of the array in the second domain.

In some embodiments, the center-to-center spacing of array features can be sufficiently small such that array features are effectively positioned continuously or nearly continuously along one or more array dimensions, with little or no displacement between array features along those dimensions. For example, in a feature array where the features correspond to regions of a substrate (i.e., oligonucleotides are directly bound to the substrate), the displacement between adjacent oligonucleotides can be very small – effectively, the molecular width of a single oligonucleotide. In such embodiments, each oligonucleotide can include a distinct spatial barcode such that the spatial location of each oligonucleotide in the array can be determined during sample analysis. Arrays of this type can have very high spatial resolution, but may only include a single oligonucleotide corresponding to each distinct spatial location in a sample.

In general, the size of the array (which corresponds to the maximum dimension of the smallest boundary that encloses all features in the array along one coordinate direction) can be selected as desired, based on criteria such as the size of the sample, the feature sizes, and the density of capture probes within each feature. For example, in some embodiments, the array can be a rectangular or square array for which the maximum array dimension along each coordinate

direction is 10 mm or less (e.g., 9 mm or less, 8 mm or less, 7 mm or less, 6 mm or less, 5 mm or less, 4 mm or less, 3 mm or less). Thus, for example, a square array of features can have dimensions of 8 mm by 8 mm, 7 mm by 7 mm, 5 mm by 5 mm, or be smaller than 5 mm by 5 mm.

5

(e) Analyte Capture

In this section, general aspects of methods and systems for capturing analytes are described. Individual method steps and system features can be present in combination in many different embodiments; the specific combinations described herein do not in any way limit other combinations of steps and features.

Generally, analytes can be captured when contacting a biological sample with, e.g., a substrate comprising capture probes (e.g., substrate with capture probes embedded, spotted, printed on the substrate or a substrate with features (e.g., beads, wells) comprising capture probes).

As used herein, “contact,” “contacted,” and/ or “contacting,” a biological sample with a substrate comprising features refers to any contact (e.g., direct or indirect) such that capture probes can interact (e.g., capture) with analytes from the biological sample. For example, the substrate may be near or adjacent to the biological sample without direct physical contact, yet capable of capturing analytes from the biological sample. In some embodiments the biological sample is in direct physical contact with the substrate. In some embodiments, the biological sample is in indirect physical contact with the substrate. For example, a liquid layer may be between the biological sample and the substrate. In some embodiments, the analytes diffuse through the liquid layer. In some embodiments the capture probes diffuse through the liquid layer. In some embodiments reagents may be delivered via the liquid layer between the biological sample and the substrate. In some embodiments, indirect physical contact may be the presence of a second substrate (e.g., a hydrogel, a film, a porous membrane) between the biological sample and the first substrate comprising features with capture probes. In some embodiments, reagents may be delivered by the second substrate to the biological sample.

Diffusion-Resistant Media / Lids

To increase efficiency by encouraging analyte diffusion toward the spatially-labelled capture probes, a diffusion-resistant medium can be used. In general, molecular diffusion of

biological analytes occurs in all directions, including toward the capture probes (i.e. toward the spatially-barcoded array), and away from the capture probes (i.e. into the bulk solution). Increasing diffusion toward the spatially-barcoded array reduces analyte diffusion away from the spatially-barcoded array and increases the capturing efficiency of the capture probes.

5 In some embodiments, a biological sample is placed on the top of a spatially-barcoded substrate and a diffusion-resistant medium is placed on top of the biological sample. For example, the diffusion-resistant medium can be placed onto an array that has been placed in contact with a biological sample. In some embodiments, the diffusion-resistant medium and spatially-labelled array are the same component. For example, the diffusion-resistant medium
10 can contain spatially-labelled capture probes within or on the diffusion-resistant medium (e.g., coverslip, slide, hydrogel, or membrane). In some embodiments, a sample is placed on a substrate and a diffusion-resistant medium is placed on top of the biological sample. Additionally, a spatially-barcoded capture probe array can be placed in close proximity over the diffusion-resistant medium. For example, a diffusion-resistant medium may be sandwiched
15 between a spatially-labelled array and a sample on a substrate. In some embodiments, the diffusion-resistant medium is disposed or spotted onto the sample. In other embodiments, the diffusion-resistant medium is placed in close proximity to the sample.

 In general, the diffusion-resistant medium can be any material known to limit diffusivity of biological analytes. For example, the diffusion-resistant medium can be a solid lid (e.g.,
20 coverslip or glass slide). In some embodiments, the diffusion-resistant medium may be made of glass, silicon, paper, hydrogel polymer monoliths, or other material. In some embodiments, the glass side can be an acrylated glass slide. In some embodiments, the diffusion-resistant medium is a porous membrane. In some embodiments, the material may be naturally porous. In some
25 embodiments, the material may have pores or wells etched into solid material. In some embodiments, the pore size can be manipulated to minimize loss of target analytes. In some embodiments, the membrane chemistry can be manipulated to minimize loss of target analytes. In some embodiments, the diffusion-resistant medium (i.e. hydrogel) is covalently attached to a substrate (i.e. glass slide). In some embodiments, the diffusion-resistant medium can be any material known to limit diffusivity of poly(A) transcripts. In some embodiments, the diffusion-
30 resistant medium can be any material known to limit the diffusivity of proteins. In some

embodiments, the diffusion-resistant medium can be any material known to limit the diffusivity of macromolecular constituents.

In some embodiments, a diffusion-resistant medium includes one or more diffusion-resistant media. For example, one or more diffusion-resistant media can be combined in a variety of ways prior to placing the media in contact with a biological sample including, without

limitation, coating, layering, or spotting. As another example, a hydrogel can be placed onto a biological sample followed by placement of a lid (e.g., glass slide) on top of the hydrogel. In some embodiments, a force (e.g., hydrodynamic pressure, ultrasonic vibration, solute contrasts, microwave radiation, vascular circulation, or other electrical, mechanical, magnetic, centrifugal, and/or thermal forces) is applied to control diffusion and enhance analyte capture. In some embodiments, one or more forces and one or more diffusion-resistant media are used to control diffusion and enhance capture. For example, a centrifugal force and a glass slide can be used contemporaneously. Any of a variety of combinations of a force and a diffusion-resistant medium can be used to control or mitigate diffusion and enhance analyte capture.

In some embodiments, the diffusion-resistant medium, along with the spatially-barcoded array and sample, is submerged in a bulk solution. In some embodiments, the bulk solution includes permeabilization reagents. In some embodiments, the diffusion-resistant medium includes at least one permeabilization reagent. In some embodiments, the diffusion-resistant medium (i.e. hydrogel) is soaked in permeabilization reagents before contacting the diffusion-resistant medium to the sample. In some embodiments, the diffusion-resistant medium can include wells (e.g., micro-, nano-, or picowells) containing a permeabilization buffer or reagents. In some embodiments, the diffusion-resistant medium can include permeabilization reagents. In some embodiments, the diffusion-resistant medium can contain dried reagents or monomers to deliver permeabilization reagents when the diffusion-resistant medium is applied to a biological sample. In some embodiments, the diffusion-resistant medium is added to the spatially-barcoded array and sample assembly before the assembly is submerged in a bulk solution. In some embodiments, the diffusion-resistant medium is added to the spatially-barcoded array and sample assembly after the sample has been exposed to permeabilization reagents. In some embodiments, the permeabilization reagents are flowed through a microfluidic chamber or channel over the diffusion-resistant medium. In some embodiments, the flow controls the sample's access to the permeabilization reagents. In some embodiments, the target analytes diffuse out of the sample

and toward a bulk solution and get embedded in a spatially-labelled capture probe-embedded diffusion-resistant medium. In some embodiments, a free solution is sandwiched between the biological sample and a diffusion-resistant medium.

FIG. 13 is an illustration of an exemplary use of a diffusion-resistant medium. A diffusion-resistant medium **1302** can be contacted with a sample **1303**. In **FIG. 13**, a glass slide **1304** is populated with spatially-barcoded capture probes **1306**, and the sample **1303**, **1305** is contacted with the array **1304**, **1306**. A diffusion-resistant medium **1302** can be applied to the sample **1303**, wherein the sample **1303** is sandwiched between a diffusion-resistant medium **1302** and a capture probe coated slide **1304**. When a permeabilization solution **1301** is applied to the sample, using the diffusion-resistant medium/lid **1302** directs migration of the analytes **1305** toward the capture probes **1306** by reducing diffusion of the analytes out into the medium. Alternatively, the lid may contain permeabilization reagents.

Conditions for capture

Capture probes on the substrate (or on a feature on the substrate) interact with released analytes through a capture domain, described elsewhere, to capture analytes. In some embodiments, certain steps are performed to enhance the transfer or capture of analytes by the capture probes of the array. Examples of such modifications include, but are not limited to, adjusting conditions for contacting the substrate with a biological sample (e.g., time, temperature, orientation, pH levels, pre-treating of biological samples, etc.), using force to transport analytes (e.g., electrophoretic, centrifugal, mechanical, etc.), performing amplification reactions to increase the amount of biological analytes (e.g., PCR amplification, in situ amplification, clonal amplification), and/or using labeled probes for detecting of amplicons and barcodes.

In some embodiments, capture of analytes is facilitated by treating the biological sample with permeabilization reagents. If a biological sample is not permeabilized sufficiently, the amount of analyte captured on the substrate can be too low to enable adequate analysis. Conversely, if the biological sample is too permeable, the analyte can diffuse away from its origin in the biological sample, such that the relative spatial relationship of the analytes within the biological sample is lost. Hence, a balance between permeabilizing the biological sample enough to obtain good signal intensity while still maintaining the spatial resolution of the analyte distribution in the biological sample is desired. Methods of preparing biological samples to

facilitation are known in the art and can be modified depending on the biological sample and how the biological sample is prepared (e.g., fresh frozen, FFPE, etc.).

Passive capture methods

In some embodiments, analytes can be migrated from a sample to a substrate. Methods
5 for facilitating migration can be passive (e.g., diffusion) and/or active (e.g., electrophoretic migration of nucleic acids). Non-limiting examples of passive migration can include simple diffusion and osmotic pressure created by the rehydration of dehydrated objects.

Passive migration by diffusion uses concentration gradients. Diffusion is movement of
10 untethered objects toward equilibrium. Therefore, when there is a region of high object concentration and a region of low object concentration, the object (capture probe, the analyte, etc.) moves to an area of lower concentration. In some embodiments, untethered analytes move down a concentration gradient.

In some embodiments, different reagents may be added to the biological sample, such
15 that the biological sample is rehydrated while improving capture of analytes. In some embodiments, the biological sample can be rehydrated with permeabilization reagents. In some embodiments, the biological sample can be rehydrated with a staining solution (e.g., hematoxylin and eosin stain).

Active capture methods

In some examples of any of the methods described herein, an analyte in a cell or a
20 biological sample can be transported (e.g., passively or actively) to a capture probe (e.g., a capture probe affixed to a solid surface).

For example, analytes in a cell or a biological sample can be transported to a capture
25 probe (e.g., an immobilized capture probe) using an electric field (e.g., using electrophoresis), a pressure gradient, fluid flow, a chemical concentration gradient, a temperature gradient, and/or a magnetic field. For example, analytes can be transported through, e.g., a gel (e.g., hydrogel matrix), a fluid, or a permeabilized cell, to a capture probe (e.g., an immobilized capture probe).

In some examples, an electrophoretic field can be applied to analytes to facilitate
migration of the analytes towards a capture probe. In some examples, a sample contacts a
substrate and capture probes fixed on a substrate (e.g., a slide, cover slip, or bead), and an
30 electric current is applied to promote the directional migration of charged analytes towards the capture probes fixed on the substrate. An electrophoresis assembly, where a cell or a biological

sample is in contact with a cathode and capture probes (e.g., capture probes fixed on a substrate), and where the capture probes (e.g., capture probes fixed on a substrate) is in contact with the cell or biological sample and an anode, can be used to apply the current.

Electrophoretic transfer of analytes can be performed while retaining the relative spatial alignment of the analytes in the sample. As such, an analyte captured by the capture probes (e.g., capture probes fixed on a substrate) retains the spatial information of the cell or the biological sample. Applying an electrophoretic field to analytes can also result in an increase in temperature (e.g., heat). In some embodiments, the increased temperature (e.g., heat) can facilitate the migration of the analytes towards a capture probe.

In some examples, a spatially-addressable microelectrode array is used for spatially-constrained capture of at least one charged analyte of interest by a capture probe. The microelectrode array can be configured to include a high density of discrete sites having a small area for applying an electric field to promote the migration of charged analyte(s) of interest. For example, electrophoretic capture can be performed on a region of interest using a spatially-addressable microelectrode array.

A high density of discrete sites on a microelectrode array can be used for small device. The surface can include any suitable density of discrete sites (e.g., a density suitable for processing the sample on the conductive substrate in a given amount of time). In an embodiment, the surface has a density of discrete sites greater than or equal to about 500 sites per 1 mm². In some embodiments, the surface has a density of discrete sites of about 100, about 200, about 300, about 400, about 500, about 600, about 700, about 800, about 900, about 1,000, about 2,000, about 3,000, about 4,000, about 5,000, about 6,000, about 7,000, about 8,000, about 9,000, about 10,000, about 20,000, about 40,000, about 60,000, about 80,000, about 100,000, or about 500,000 sites per 1 mm². In some embodiments, the surface has a density of discrete sites of at least about 200, at least about 300, at least about 400, at least about 500, at least about 600, at least about 700, at least about 800, at least about 900, at least about 1,000, at least about 2,000, at least about 3,000, at least about 4,000, at least about 5,000, at least about 6,000, at least about 7,000, at least about 8,000, at least about 9,000, at least about 10,000, at least about 20,000, at least about 40,000, at least about 60,000, at least about 80,000, at least about 100,000, or at least about 500,000 sites per 1 mm².

Schematics illustrating an electrophoretic transfer system configured to direct transcript analytes toward a spatially-barcoded capture probe array are shown in **FIG. 14A** and **FIG. 14B**.

In this exemplary configuration of an electrophoretic system, a sample **1402** is sandwiched between the cathode **1401** and the spatially-barcoded capture probe array **1404, 1405**, and the spatially-barcoded capture probe array **1404, 1405** is sandwiched between the sample **1402** and the anode **1403**, such that the sample **1402, 1406** is in contact with the spatially-barcoded capture probes **1407**. When an electric field is applied to the electrophoretic transfer system, negatively charged mRNA analytes **1406** will be pulled toward the positively charged anode **1403** and into the spatially-barcoded array **1404, 1405** containing the spatially-barcoded capture probes **1407**.

The spatially-barcoded capture probes **1407** then interact with/hybridize with/immobilize the mRNA target analytes **1406**, making the analyte capture more efficient. The electrophoretic system set-up may change depending on the target analyte. For example, proteins may be positive, negative, neutral, or polar depending on the protein as well as other factors (e.g. isoelectric point, solubility, etc.). The skilled practitioner has the knowledge and experience to arrange the electrophoretic transfer system to facilitate capture of a particular target analyte.

FIG. 15 is an illustration showing an exemplary workflow protocol utilizing an electrophoretic transfer system. In the example, **Panel A** depicts a flexible spatially-barcoded feature array being contacted with a sample. The sample can be a flexible array, wherein the array is immobilized on a hydrogel, membrane, or other flexible substrate. **Panel B** depicts contact of the array with the sample and imaging of the array-sample assembly. The image of the sample/array assembly can be used to verify sample placement, choose a region of interest, or any other reason for imaging a sample on an array as described herein. **Panel C** depicts application of an electric field using an electrophoretic transfer system to aid in efficient capture of a target analyte. Here, negatively charged mRNA target analytes migrate toward the positively charged anode. **Panel D** depicts application of reverse transcription reagents and first strand cDNA synthesis of the captured target analytes. **Panel E** depicts array removal and preparation for library construction (**Panel F**) and next-generation sequencing (**Panel G**).

Region of Interest

A biological sample can have regions that show morphological feature(s) that may indicate the presence of disease or the development of a disease phenotype. For example, morphological features at a specific site within a tumor biopsy sample can indicate the

aggressiveness, therapeutic resistance, metastatic potential, migration, stage, diagnosis, and/or prognosis of cancer in a subject. A change in the morphological features at a specific site within a tumor biopsy sample often correlate with a change in the level or expression of an analyte in a cell within the specific site, which can, in turn, be used to provide information regarding the aggressiveness, therapeutic resistance, metastatic potential, migration, stage, diagnosis, and/or prognosis of cancer in a subject. A region or area within a biological sample that is selected for specific analysis (e.g., a region in a biological sample that has morphological features of interest) is often described as “a region of interest.”

A region of interest in a biological sample can be used to analyze a specific area of interest within a biological sample, and thereby, focus experimentation and data gathering to a specific region of a biological sample (rather than an entire biological sample). This results in increased time efficiency of the analysis of a biological sample.

A region of interest can be identified in a biological sample using a variety of different techniques, e.g., expansion microscopy, bright field microscopy, dark field microscopy, phase contrast microscopy, electron microscopy, fluorescence microscopy, reflection microscopy, interference microscopy, confocal microscopy, and visual identification (e.g., by eye), and combinations thereof. For example, the staining and imaging of a biological sample can be performed to identify a region of interest. In some examples, the region of interest can correspond to a specific structure of cytoarchitecture. In some embodiments, a biological sample can be stained prior to visualization to provide contrast between the different regions of the biological sample. The type of stain can be chosen depending on the type of biological sample and the region of the cells to be stained. In some embodiments, more than one stain can be used to visualize different aspects of the biological sample, e.g., different regions of the sample, specific cell structures (e.g. organelles), or different cell types. In other embodiments, the biological sample can be visualized or imaged without staining the biological sample.

In some embodiments, imaging can be performed using one or more fiducial markers, i.e., objects placed in the field of view of an imaging system which appear in the image produced. Fiducial markers are typically used as a point of reference or measurement scale. Fiducial markers can include, but are not limited to, detectable labels such as fluorescent, radioactive, chemiluminescent, and colorimetric labels. The use of fiducial markers to stabilize and orient biological samples is described, for example, in Carter et al., *Applied Optics* 46:421-

427, 2007), the entire contents of which are incorporated herein by reference. In some embodiments, a fiducial marker can be a physical particle (e.g., a nanoparticle, a microsphere, a nanosphere, a bead, or any of the other exemplary physical particles described herein or known in the art).

5 In some embodiments, a fiducial marker can be present on a substrate to provide orientation of the biological sample. In some embodiments, a microsphere can be coupled to a substrate to aid in orientation of the biological sample. In some examples, a microsphere coupled to a substrate can produce an optical signal (e.g., fluorescence). In another example, a microsphere can be attached to a portion (e.g., corner) of an array in a specific pattern or design
10 (e.g., hexagonal design) to aid in orientation of a biological sample on an array of features on the substrate. In some embodiments, a quantum dot can be coupled to the substrate to aid in the orientation of the biological sample. In some examples, a quantum dot coupled to a substrate can produce an optical signal.

In some embodiments, a fiducial marker can be an immobilized molecule with which a
15 detectable signal molecule can interact to generate a signal. For example, a marker nucleic acid can be linked or coupled to a chemical moiety capable of fluorescing when subjected to light of a specific wavelength (or range of wavelengths). Such a marker nucleic acid molecule can be contacted with an array before, contemporaneously with, or after the tissue sample is stained to visualize or image the tissue section. Although not required, it can be advantageous to use a
20 marker that can be detected using the same conditions (e.g., imaging conditions) used to detect a labelled cDNA.

In some embodiments, fiducial markers are included to facilitate the orientation of a tissue sample or an image thereof in relation to an immobilized capture probes on a substrate. Any number of methods for marking an array can be used such that a marker is detectable only
25 when a tissue section is imaged. For instance, a molecule, e.g. a fluorescent molecule that generates a signal, can be immobilized directly or indirectly on the surface of a substrate. Markers can be provided on a substrate in a pattern (e.g., an edge, one or more rows, one or more lines, etc.).

In some embodiments, a fiducial marker can be randomly placed in the field of view. For
30 example, an oligonucleotide containing a fluorophore can be randomly printed, stamped, synthesized, or attached to a substrate (e.g., a glass slide) at a random position on the substrate. A

tissue section can be contacted with the substrate such that the oligonucleotide containing the fluorophore contacts, or is in proximity to, a cell from the tissue section or a component of the cell (e.g., an mRNA or DNA molecule). An image of the substrate and the tissue section can be obtained, and the position of the fluorophore within the tissue section image can be determined (e.g., by reviewing an optical image of the tissue section overlaid with the fluorophore detection). In some embodiments, fiducial markers can be precisely placed in the field of view (e.g., at known locations on a substrate). In this instance, a fiducial marker can be stamped, attached, or synthesized on the substrate and contacted with a biological sample. Typically, an image of the sample and the fiducial marker is taken, and the position of the fiducial marker on the substrate can be confirmed by viewing the image.

In some embodiments, a fiducial marker can be an immobilized molecule (e.g., a physical particle) attached to the substrate. For example, a fiducial marker can be a nanoparticle, e.g., a nanorod, a nanowire, a nanocube, a nanopyramid, or a spherical nanoparticle. In some examples, the nanoparticle can be made of a heavy metal (e.g., gold). In some embodiments, the nanoparticle can be made from diamond. In some embodiments, the fiducial marker can be visible by eye.

As noted herein, any of the fiducial markers described herein (e.g., microspheres, beads, or any of the other physical particles described herein) can be located at a portion (e.g., corner) of an array in a specific pattern or design (e.g., hexagonal design) to aid in orientation of a biological sample on an array of features on the substrate. In some embodiments, the fiducial markers located at a portion (e.g., corner) of an array (e.g., an array on a substrate) can be patterned or designed in at least 1, at least 2, at least 3, or at least 4 unique patterns. In some examples, the fiducial markers located at the corners of the array (e.g., an array on a substrate) can have four unique patterns of fiducial markers.

In some examples, fiducial markers can surround the array. In some embodiments the fiducial markers allow for detection of, e.g., mirroring. In some embodiments, the fiducial markers may completely surround the array. In some embodiments, the fiducial markers may not completely surround the array. In some embodiments, the fiducial markers identify the corners of the array. In some embodiments, one or more fiducial markers identify the center of the array. In some embodiments, the fiducial markers comprise patterned spots, wherein the diameter of one or more patterned spot fiducial markers is approximately 100 micrometers. The diameter of the

fiducial markers can be any useful diameter including, but not limited to, 50 micrometers to 500 micrometers in diameter. The fiducial markers may be arranged in such a way that the center of one fiducial marker is between 100 micrometers and 200 micrometers from the center of one or more other fiducial markers surrounding the array. In some embodiments, the array with the surrounding fiducial markers is approximately 8 mm by 8 mm. In some embodiments, the array without the surrounding fiducial markers is smaller than 8 mm by 50 mm.

In some embodiments, an array can be enclosed within a frame. Put another way, the perimeter of an array can have fiducial markers such that the array is enclosed, or substantially enclosed. In some embodiments, the perimeter of an array can be fiducial markers (e.g., any fiducial marker described herein). In some embodiments, the perimeter of an array can be uniform. For example, the fiducial markings can connect, or substantially connect, consecutive corners of an array in such a fashion that the non-corner portion of the array perimeter is the same on all sides (e.g., four sides) of the array. In some embodiments, the fiducial markers attached to the non-corner portions of the perimeter can be patterned or designed to aid in the orientation of the biological sample on the array. In some embodiments, the particles attached to the non-corner portions of the perimeter can be patterned or designed in at least 1, at least 2, at least 3, or at least 4 patterns. In some embodiments, the patterns can have at least 2, at least 3, or at least 4 unique patterns of fiducial markings on the non-corner portion of the array perimeter.

In some embodiments, an array can include at least two fiducial markers (e.g., at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 12, at least 15, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90, at least 100 fiducial markers or more (e.g., several hundred, several thousand, or tens of thousands of fiducial markers)) in distinct positions on the surface of a substrate. Fiducial markers can be provided on a substrate in a pattern (e.g., an edge, one or more rows, one or more lines, etc.).

In some embodiments, staining and imaging a biological sample prior to contacting the biological sample with a spatial array is performed to select samples for spatial analysis. In some embodiments, the staining includes applying a fiducial marker as described above, including fluorescent, radioactive, chemiluminescent, or colorimetric detectable markers. In some embodiments, the staining and imaging of biological samples allows the user to identify the specific sample (or region of interest) the user wishes to assess.

In some embodiments, a lookup table (LUT) can be used to associate one property with another property of a feature. These properties include, e.g., locations, barcodes (e.g., nucleic acid barcode molecules), spatial barcodes, optical labels, molecular tags, and other properties.

5 In some embodiments, a lookup table can associate a nucleic acid barcode molecule with a feature. In some embodiments, an optical label of a feature can permit associating the feature with a biological particle (e.g., cell or nuclei). The association of a feature with a biological particle can further permit associating a nucleic acid sequence of a nucleic acid molecule of the biological particle to one or more physical properties of the biological particle (e.g., a type of a cell or a location of the cell). For example, based on the relationship between the barcode and the
10 optical label, the optical label can be used to determine the location of a feature, thus associating the location of the feature with the barcode sequence of the feature. Subsequent analysis (e.g., sequencing) can associate the barcode sequence and the analyte from the sample. Accordingly, based on the relationship between the location and the barcode sequence, the location of the biological analyte can be determined (e.g., in a specific type of cell or in a cell at a specific
15 location of the biological sample).

In some embodiments, a feature can have a plurality of nucleic acid barcode molecules attached thereto. The plurality of nucleic acid barcode molecules can include barcode sequences. The plurality of nucleic acid molecules attached to a given feature can have the same barcode sequences, or two or more different barcode sequences. Different barcode sequences can be used
20 to provide improved spatial location accuracy.

In some embodiments, a substrate is treated in order to minimize or reduce non-specific analyte hybridization within or between features. For example, treatment can include coating the substrate with a hydrogel, film, and/or membrane that creates a physical barrier to non-specific hybridization. Any suitable hydrogel can be used. For example, hydrogel matrices prepared
25 according to the methods set forth in U.S. Patent Nos. 6,391,937, 9,512,422, and 9,889,422, and U.S. Patent Application Publication Nos. U.S. 2017/0253918 and U.S. 2018/0052081, can be used. The entire contents of each of the foregoing documents are incorporated herein by reference.

Treatment can include adding a functional group that is reactive or capable of being
30 activated such that it becomes reactive after receiving a stimulus (e.g., photoreactive). Treatment can include treating with polymers having one or more physical properties (e.g., mechanical,

electrical, magnetic, and/or thermal) that minimize non-specific binding (e.g., that activate a substrate at certain locations to allow analyte hybridization at those locations).

In some examples, an array (e.g., any of the exemplary arrays described herein) can be contacted with only a portion of a biological sample (e.g., a cell, a feature, or a region of interest). In some examples, a biological sample is contacted with only a portion of an array (e.g., any of the exemplary arrays described herein). In some examples, a portion of the array can be deactivated such that it does not interact with the analytes in the biological sample (e.g., optical deactivation, chemical deactivation, heat deactivation, or blocking of the capture probes in the array (e.g., using blocking probes)). In some examples, a region of interest can be removed from a biological sample and then the region of interest can be contacted to the array (e.g., any of the arrays described herein). A region of interest can be removed from a biological sample using microsurgery, laser capture microdissection, chunking, a microtome, dicing, trypsinization, labelling, and/or fluorescence-assisted cell sorting.

(f) Partitioning

As discussed above, in some embodiments, the sample can optionally be separated into single cells, cell groups, or other fragments/pieces that are smaller than the original, unfragmented sample. Each of these smaller portions of the sample can be analyzed to obtain spatially-resolved analyte information for the sample.

For samples that have been separated into smaller fragments – and particularly, for samples that have been disaggregated, dissociated, or otherwise separated into individual cells – one method for analyzing the fragments involves partitioning the fragments into individual partitions (e.g., fluid droplets), and then analyzing the contents of the partitions. In general, each partition maintains separation of its own contents from the contents of other partitions. The partition can be a droplet in an emulsion, for example.

In addition to analytes, a partition can include additional components, and in particular, one or more beads. A partition can include a single gel bead, a single cell bead, or both a single cell bead and single gel bead.

A partition can also include one or more reagents. Unique identifiers, such as barcodes, can be injected into the droplets previous to, subsequent to, or concurrently with droplet generation, such as via a microcapsule (e.g., bead). Microfluidic channel networks (e.g., on a chip) can be utilized to generate partitions. Alternative mechanisms can also be employed in the

partitioning of individual biological particles, including porous membranes through which aqueous mixtures of cells are extruded into non-aqueous fluids.

The partitions can be flowable within fluid streams. The partitions can include, for example, micro-vesicles that have an outer barrier surrounding an inner fluid center or core. In some cases, the partitions can include a porous matrix that is capable of entraining and/or retaining materials within its matrix. The partitions can be droplets of a first phase within a second phase, wherein the first and second phases are immiscible. For example, the partitions can be droplets of aqueous fluid within a non-aqueous continuous phase (e.g., oil phase). In another example, the partitions can be droplets of a non-aqueous fluid within an aqueous phase. In some examples, the partitions can be provided in a water-in-oil emulsion or oil-in-water emulsion. A variety of different vessels are described in, for example, U.S. Patent Application Publication No. 2014/0155295, the entire contents of which are incorporated herein by reference. Emulsion systems for creating stable droplets in non-aqueous or oil continuous phases are described, for example, in U.S. Patent Application Publication No. 2010/0105112, the entire contents of which are incorporated herein by reference.

For droplets in an emulsion, allocating individual particles to discrete partitions can be accomplished, for example, by introducing a flowing stream of particles in an aqueous fluid into a flowing stream of a non-aqueous fluid, such that droplets are generated at the junction of the two streams. Fluid properties (e.g., fluid flow rates, fluid viscosities, etc.), particle properties (e.g., volume fraction, particle size, particle concentration, etc.), microfluidic architectures (e.g., channel geometry, etc.), and other parameters can be adjusted to control the occupancy of the resulting partitions (e.g., number of analytes per partition, number of beads per partition, etc.). For example, partition occupancy can be controlled by providing the aqueous stream at a certain concentration and/or flow rate of analytes.

To generate single analyte partitions, the relative flow rates of the immiscible fluids can be selected such that, on average, the partitions can contain less than one analyte per partition to ensure that those partitions that are occupied are primarily singly occupied. In some cases, partitions among a plurality of partitions can contain at most one analyte. In some embodiments, the various parameters (e.g., fluid properties, particle properties, microfluidic architectures, etc.) can be selected or adjusted such that a majority of partitions are occupied, for example, allowing for only a small percentage of unoccupied partitions. The flows and channel architectures can be

controlled as to ensure a given number of singly occupied partitions, less than a certain level of unoccupied partitions and/or less than a certain level of multiply occupied partitions.

The channel segments described herein can be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure can have a variety of geometries. For example, a microfluidic channel structure can have one or more than one channel junction. As another example, a microfluidic channel structure can have 2, 3, 4, or 5 channel segments each carrying particles that meet at a channel junction. Fluid can be directed to flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can include compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid can also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary, and/or gravity flow.

A partition can include one or more unique identifiers, such as barcodes. Barcodes can be previously, subsequently, or concurrently delivered to the partitions that hold the compartmentalized or partitioned biological particle. For example, barcodes can be injected into droplets previous to, subsequent to, or concurrently with droplet generation. The delivery of the barcodes to a particular partition allows for the later attribution of the characteristics of the individual biological particle to the particular partition. Barcodes can be delivered, for example on a nucleic acid molecule (e.g., an oligonucleotide), to a partition via any suitable mechanism. Barcoded nucleic acid molecules can be delivered to a partition via a microcapsule. A microcapsule, in some instances, can include a bead.

In some embodiments, barcoded nucleic acid molecules can be initially associated with the microcapsule and then released from the microcapsule. Release of the barcoded nucleic acid molecules can be passive (e.g., by diffusion out of the microcapsule). In addition or alternatively, release from the microcapsule can be upon application of a stimulus which allows the barcoded nucleic acid molecules to dissociate or to be released from the microcapsule. Such stimulus can disrupt the microcapsule, an interaction that couples the barcoded nucleic acid molecules to or within the microcapsule, or both. Such stimulus can include, for example, a thermal stimulus, photo-stimulus, chemical stimulus (e.g., change in pH or use of a reducing

agent(s)), a mechanical stimulus, a radiation stimulus; a biological stimulus (e.g., enzyme), or any combination thereof.

In some embodiments, one more barcodes (e.g., spatial barcodes, UMIs, or a combination thereof) can be introduced into a partition as part of the analyte. As described previously, barcodes can be bound to the analyte directly, or can form part of a capture probe or analyte capture agent that is hybridized to, conjugated to, or otherwise associated with an analyte, such that when the analyte is introduced into the partition, the barcode(s) are introduced as well. As described above, **FIG. 16** shows an example of a microfluidical channel structure for partitioning individual analytes (e.g., cells) into discrete partitions.

FIG. 16 shows an example of a microfluidic channel structure for partitioning individual analytes (e.g., cells) into discrete partitions. The channel structure can include channel segments **1601**, **1602**, **1603**, and **1604** communicating at a channel junction **1605**. In operation, a first aqueous fluid **1606** that includes suspended biological particles (or cells) **1607** may be transported along channel segment **1601** into junction **1605**, while a second fluid **1608** that is immiscible with the aqueous fluid **1606** is delivered to the junction **1605** from each of channel segments **1602** and **1603** to create discrete droplets **1609**, **1610** of the first aqueous fluid **1606** flowing into channel segment **1604**, and flowing away from junction **1605**. The channel segment **1604** may be fluidically coupled to an outlet reservoir where the discrete droplets can be stored and/or harvested. A discrete droplet generated may include an individual biological particle **1607** (such as droplets **1609**). A discrete droplet generated may include more than one individual biological particle **1607**. A discrete droplet may contain no biological particle **1607** (such as droplet **1610**). Each discrete partition may maintain separation of its own contents (e.g., individual biological particle **1607**) from the contents of other partitions.

FIG. 17A shows another example of a microfluidic channel structure **1700** for delivering beads to droplets. The channel structure includes channel segments **1701**, **1702**, **1703**, **1704** and **1705** communicating at a channel junction **1706**. During operation, the channel segment **1701** can transport an aqueous fluid **1707** that includes a plurality of beads **1708** along the channel segment **1701** into junction **1706**. The plurality of beads **1708** can be sourced from a suspension of beads. For example, the channel segment **1701** can be connected to a reservoir that includes an aqueous suspension of beads **1708**. The channel segment **1702** can transport the aqueous fluid **1707** that includes a plurality of particles **1709** (e.g., cells) along the channel segment **1702** into

junction **1706**. In some embodiments, the aqueous fluid **1707** in either the first channel segment **1701** or the second channel segment **1702**, or in both segments, can include one or more reagents, as further described below.

A second fluid **1710** that is immiscible with the aqueous fluid **1707** (e.g., oil) can be delivered to the junction **1706** from each of channel segments **1703** and **1704**. Upon meeting of the aqueous fluid **1707** from each of channel segments **1701** and **1702** and the second fluid **1710** from each of channel segments **1703** and **1704** at the channel junction **1706**, the aqueous fluid **1707** can be partitioned as discrete droplets **1711** in the second fluid **1710** and flow away from the junction **1706** along channel segment **1705**. The channel segment **1705** can deliver the discrete droplets to an outlet reservoir fluidly coupled to the channel segment **1705**, where they can be harvested.

As an alternative, the channel segments **1701** and **1702** can meet at another junction upstream of the junction **1706**. At such junction, beads and biological particles can form a mixture that is directed along another channel to the junction **1706** to yield droplets **1711**. The mixture can provide the beads and biological particles in an alternating fashion, such that, for example, a droplet includes a single bead and a single biological particle.

The second fluid **1710** can include an oil, such as a fluorinated oil, that includes a fluorosurfactant for stabilizing the resulting droplets, for example, inhibiting subsequent coalescence of the resulting droplets **1711**.

The partitions described herein can include small volumes, for example, less than about 10 microliters (μL), 5 μL , 1 μL , 900 picoliters (pL), 800 pL, 700 pL, 600 pL, 500 pL, 400pL, 300 pL, 200 pL, 100pL, 50 pL, 20 pL, 10 pL, 1 pL, 500 nanoliters (nL), 100 nL, 50 nL, or less. In the foregoing discussion, droplets with beads were formed at the junction of different fluid streams. In some embodiments, droplets can be formed by gravity-based partitioning methods.

FIG. 17B shows a cross-section view of another example of a microfluidic channel structure **1750** with a geometric feature for controlled partitioning. A channel structure **1750** can include a channel segment **1752** communicating at a channel junction **1758** (or intersection) with a reservoir **1754**. In some instances, the channel structure **1750** and one or more of its components can correspond to the channel structure **1700** and one or more of its components.

An aqueous fluid **1760** comprising a plurality of particles **1756** may be transported along the channel segment **1752** into the junction **1758** to meet a second fluid **1762** (e.g., oil, etc.) that

is immiscible with the aqueous fluid **1760** in the reservoir **1754** to create droplets **1764** of the aqueous fluid **1760** flowing into the reservoir **1754**. At the junction **1758** where the aqueous fluid **1760** and the second fluid **1762** meet, droplets can form based on factors such as the hydrodynamic forces at the junction **1758**, relative flow rates of the two fluids **1760**, **1762**, fluid properties, and certain geometric parameters (e.g., Δh , etc.) of the channel structure **1750**. A plurality of droplets can be collected in the reservoir **1754** by continuously injecting the aqueous fluid **1760** from the channel segment **1752** at the junction **1758**.

A discrete droplet generated may comprise one or more particles of the plurality of particles **1756**. As described elsewhere herein, a particle may be any particle, such as a bead, cell bead, gel bead, biological particle, macromolecular constituents of biological particle, or other particles. Alternatively, a discrete droplet generated may not include any particles.

In some instances, the aqueous fluid **1760** can have a substantially uniform concentration or frequency of particles **1756**. As described elsewhere herein, the particles **1756** (e.g., beads) can be introduced into the channel segment **1752** from a separate channel (not shown in **FIG. 17**). The frequency of particles **1756** in the channel segment **1752** may be controlled by controlling the frequency in which the particles **1756** are introduced into the channel segment **1752** and/or the relative flow rates of the fluids in the channel segment **1752** and the separate channel. In some instances, the particles **1756** can be introduced into the channel segment **1752** from a plurality of different channels, and the frequency controlled accordingly. In some instances, different particles may be introduced via separate channels. For example, a first separate channel can introduce beads and a second separate channel can introduce biological particles into the channel segment **1752**. The first separate channel introducing the beads may be upstream or downstream of the second separate channel introducing the biological particles.

In some instances, the second fluid **1762** may not be subjected to and/or directed to any flow in or out of the reservoir **1754**. For example, the second fluid **1762** may be substantially stationary in the reservoir **1754**. In some instances, the second fluid **1762** may be subjected to flow within the reservoir **1754**, but not in or out of the reservoir **1754**, such as via application of pressure to the reservoir **1754** and/or as affected by the incoming flow of the aqueous fluid **1760** at the junction **1758**. Alternatively, the second fluid **1762** may be subjected and/or directed to flow in or out of the reservoir **1754**. For example, the reservoir **1754** can be a channel directing the second fluid **1762** from upstream to downstream, transporting the generated droplets.

The channel structure **1750** at or near the junction **1758** may have certain geometric features that at least partly determine the sizes and/or shapes of the droplets formed by the channel structure **1750**. The channel segment **1752** can have a first cross-section height, h_1 , and the reservoir **1754** can have a second cross-section height, h_2 . The first cross-section height, h_1 , and the second cross-section height, h_2 , may be different, such that at the junction **1758**, there is a height difference of Δh . The second cross-section height, h_2 , may be greater than the first cross-section height, h_1 . In some instances, the reservoir may thereafter gradually increase in cross-section height, for example, the more distant it is from the junction **1758**. In some instances, the cross-section height of the reservoir may increase in accordance with expansion angle, β , at or near the junction **1758**. The height difference, Δh , and/or expansion angle, β , can allow the tongue (portion of the aqueous fluid **1760** leaving channel segment **1752** at junction **1758** and entering the reservoir **1754** before droplet formation) to increase in depth and facilitate decrease in curvature of the intermediately formed droplet. For example, droplet size may decrease with increasing height difference and/or increasing expansion angle.

The height difference, Δh , can be at least about 1 μm . Alternatively, the height difference can be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500 μm or more. Alternatively, the height difference can be at most about 500, 400, 300, 200, 100, 90, 80, 70, 60, 50, 45, 40, 35, 30, 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 μm or less. In some instances, the expansion angle, β , may be between a range of from about 0.5° to about 4° , from about 0.1° to about 10° , or from about 0° to about 90° . For example, the expansion angle can be at least about 0.01° , 0.1° , 0.2° , 0.3° , 0.4° , 0.5° , 0.6° , 0.7° , 0.8° , 0.9° , 1° , 2° , 3° , 4° , 5° , 6° , 7° , 8° , 9° , 10° , 15° , 20° , 25° , 30° , 35° , 40° , 45° , 50° , 55° , 60° , 65° , 70° , 75° , 80° , 85° , or higher. In some instances, the expansion angle can be at most about 89° , 88° , 87° , 86° , 85° , 84° , 83° , 82° , 81° , 80° , 75° , 70° , 65° , 60° , 55° , 50° , 45° , 40° , 35° , 30° , 25° , 20° , 15° , 10° , 9° , 8° , 7° , 6° , 5° , 4° , 3° , 2° , 1° , 0.1° , 0.01° , or less.

In some instances, the flow rate of the aqueous fluid **1760** entering the junction **1758** can be between about 0.04 microliters (μL)/minute (min) and about 40 $\mu\text{L}/\text{min}$. In some instances, the flow rate of the aqueous fluid **1760** entering the junction **1758** can be between about 0.01 microliters (μL)/minute (min) and about 100 $\mu\text{L}/\text{min}$. Alternatively, the flow rate of the aqueous fluid **1760** entering the junction **1758** can be less than about 0.01 $\mu\text{L}/\text{min}$. alternatively, the flow

rate of the aqueous fluid **1760** entering the junction **1758** can be greater than about 40 $\mu\text{L}/\text{min}$, such as 45 $\mu\text{L}/\text{min}$, 50 $\mu\text{L}/\text{min}$, 55 $\mu\text{L}/\text{min}$, 60 $\mu\text{L}/\text{min}$, 65 $\mu\text{L}/\text{min}$, 70 $\mu\text{L}/\text{min}$, 75 $\mu\text{L}/\text{min}$, 80 $\mu\text{L}/\text{min}$, 85 $\mu\text{L}/\text{min}$, 90 $\mu\text{L}/\text{min}$, 95 $\mu\text{L}/\text{min}$, 100 $\mu\text{L}/\text{min}$, 110 $\mu\text{L}/\text{min}$, 120 $\mu\text{L}/\text{min}$, 130 $\mu\text{L}/\text{min}$, 140 $\mu\text{L}/\text{min}$, 150 $\mu\text{L}/\text{min}$, or greater. At lower flow rates, such as flow rates of about less than or equal to 10 microliters/minute, the droplet radius may not be dependent on the flow rate of the aqueous fluid **1760** entering the junction **1758**. The second fluid **1762** may be stationary, or substantially stationary, in the reservoir **1754**. Alternatively, the second fluid **1762** may be flowing, such as at the above flow rates described for the aqueous fluid **1760**.

While **FIG. 17B** illustrates the height difference, Δh , being abrupt at the junction **1758** (e.g., a step increase), the height difference may increase gradually (e.g., from about 0 μm to a maximum height difference). Alternatively, the height difference may decrease gradually (e.g., taper) from a maximum height difference. A gradual increase or decrease in height difference, as used herein, may refer to a continuous incremental increase or decrease in height difference, wherein an angle between any one differential segment of a height profile and an immediately adjacent differential segment of the height profile is greater than 90° . For example, at the junction **1758**, a bottom wall of the channel and a bottom wall of the reservoir can meet at an angle greater than 90° . Alternatively or in addition, a top wall (e.g., ceiling) of the channel and a top wall (e.g., ceiling) of the reservoir can meet an angle greater than 90° . A gradual increase or decrease may be linear or non-linear (e.g., exponential, sinusoidal, etc.). Alternatively or in addition, the height difference may variably increase and/or decrease linearly or non-linearly. While **FIG. 17B** illustrates the expanding reservoir cross-section height as linear (e.g., constant expansion angle, β), the cross-section height may expand non-linearly. For example, the reservoir may be defined at least partially by a dome-like (e.g., hemispherical) shape having variable expansion angles. The cross-section height may expand in any shape.

A variety of different beads can be incorporated into partitions as described above. In some embodiments, for example, non-barcoded beads can be incorporated into the partitions. For example, where the biological particle (e.g., a cell) that is incorporated into the partitions carries one or more barcodes (e.g., spatial barcode(s), UMI(s), and combinations thereof), the bead can be a non-barcoded bead.

In some embodiments, a barcode carrying bead can be incorporated into partitions. For example, a nucleic acid molecule, such as an oligonucleotide, can be coupled to a bead by a

releasable linkage, such as, for example, a disulfide linker. The same bead can be coupled (e.g., via releasable linkage) to one or more other nucleic acid molecules. The nucleic acid molecule can be or include a barcode. As noted elsewhere herein, the structure of the barcode can include a number of sequence elements.

5 The nucleic acid molecule can include a functional domain that can be used in subsequent processing. For example, the functional domain can include one or more of a sequencer specific flow cell attachment sequence (e.g., a P5 sequence for Illumina® sequencing systems) and a sequencing primer sequence (e.g., a R1 primer for Illumina® sequencing systems). The nucleic acid molecule can include a barcode sequence for use in barcoding the sample (e.g., DNA, RNA,
10 protein, etc.). In some cases, the barcode sequence can be bead-specific such that the barcode sequence is common to all nucleic acid molecules coupled to the same bead. Alternatively or in addition, the barcode sequence can be partition-specific such that the barcode sequence is common to all nucleic acid molecules coupled to one or more beads that are partitioned into the same partition. The nucleic acid molecule can include a specific priming sequence, such as an
15 mRNA specific priming sequence (e.g., poly(T) sequence), a targeted priming sequence, and/or a random priming sequence. The nucleic acid molecule can include an anchoring sequence to ensure that the specific priming sequence hybridizes at the sequence end (e.g., of the mRNA). For example, the anchoring sequence can include a random short sequence of nucleotides, such as a 1-mer, 2-mer, 3-mer or longer sequence, which can ensure that a poly(T) segment is more
20 likely to hybridize at the sequence end of the poly(A) tail of the mRNA.

 The nucleic acid molecule can include a unique molecular identifying sequence (e.g., unique molecular identifier (UMI)). In some embodiments, the unique molecular identifying sequence can include from about 5 to about 8 nucleotides. Alternatively, the unique molecular identifying sequence can include less than about 5 or more than about 8 nucleotides. The unique
25 molecular identifying sequence can be a unique sequence that varies across individual nucleic acid molecules coupled to a single bead.

 In some embodiments, the unique molecular identifying sequence can be a random sequence (e.g., such as a random N-mer sequence). For example, the UMI can provide a unique identifier of the starting mRNA molecule that was captured, in order to allow quantitation of the
30 number of original expressed RNA.

In general, an individual bead can be coupled to any number of individual nucleic acid molecules, for example, from one to tens to hundreds of thousands or even millions of individual nucleic acid molecules. The respective barcodes for the individual nucleic acid molecules can include both common sequence segments or relatively common sequence segments and variable or unique sequence segments between different individual nucleic acid molecules coupled to the same bead.

FIG. 17C depicts a workflow wherein cells are partitioned into droplets along with barcode-bearing beads **1770**. See **FIG. 17A**. The droplet forms an isolated reaction chamber wherein the cells can be lysed **1771** and target analytes within the cells can then be captured **1772** and amplified **1773**, **1774** according to previously described methods. After sequence library preparation clean-up **1775**, the material is sequenced and/or quantified **1776** according to methods described herein.

It should be noted that while the example workflow in **FIG. 17C** includes steps specifically for the analysis of mRNA, analogous workflows can be implemented for a wide variety of other analytes, including any of the analytes described previously.

By way of example, in the context of analyzing sample RNA as shown in **FIG. 17C**, the poly(T) segment of one of the released nucleic acid molecules (e.g., from the bead) can hybridize to the poly(A) tail of a mRNA molecule. Reverse transcription can result in a cDNA transcript of the mRNA, which transcript includes each of the sequence segments of the nucleic acid molecule. If the nucleic acid molecule includes an anchoring sequence, it will more likely hybridize to and prime reverse transcription at the sequence end of the poly(A) tail of the mRNA.

Within any given partition, all of the cDNA transcripts of the individual mRNA molecules can include a common barcode sequence segment. However, the transcripts made from the different mRNA molecules within a given partition can vary at the unique molecular identifying sequence segment (e.g., UMI segment). Beneficially, even following any subsequent amplification of the contents of a given partition, the number of different UMIs can be indicative of the quantity of mRNA originating from a given partition. As noted above, the transcripts can be amplified, cleaned up and sequenced to identify the sequence of the cDNA transcript of the mRNA, as well as to sequence the barcode segment and the UMI segment. While a poly(T) primer sequence is described, other targeted or random priming sequences can also be used in

priming the reverse transcription reaction. Likewise, although described as releasing the barcoded oligonucleotides into the partition, in some cases, the nucleic acid molecules bound to the bead can be used to hybridize and capture the mRNA on the solid phase of the bead, for example, in order to facilitate the separation of the RNA from other cell contents.

5 In some embodiments, precursors that include a functional group that is reactive or capable of being activated such that it becomes reactive can be polymerized with other precursors to generate gel beads that include the activated or activatable functional group. The functional group can then be used to attach additional species (e.g., disulfide linkers, primers, other oligonucleotides, etc.) to the gel beads. For example, some precursors featuring a
10 carboxylic acid (COOH) group can co-polymerize with other precursors to form a bead that also includes a COOH functional group. In some cases, acrylic acid (a species comprising free COOH groups), acrylamide, and bis(acryloyl)cystamine can be co-polymerized together to generate a bead with free COOH groups. The COOH groups of the bead can be activated (e.g., via 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and N-Hydroxysuccinimide (NHS) or 4-(4,6-
15 Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)) such that they are reactive (e.g., reactive to amine functional groups where EDC/NHS or DMTMM are used for activation). The activated COOH groups can then react with an appropriate species (e.g., a species comprising an amine functional group where the carboxylic acid groups are activated to be reactive with an amine functional group) comprising a moiety to be linked to the bead.

20 In some embodiments, a degradable bead can be introduced into a partition, such that the bead degrades within the partition and any associated species (e.g., oligonucleotides) are released within the droplet when the appropriate stimulus is applied. The free species (e.g., oligonucleotides, nucleic acid molecules) can interact with other reagents contained in the partition. For example, a polyacrylamide bead featuring cystamine and linked, via a disulfide
25 bond, to a barcode sequence, can be combined with a reducing agent within a droplet of a water-in-oil emulsion. Within the droplet, the reducing agent can break the various disulfide bonds, resulting in bead degradation and release of the barcode sequence into the aqueous, inner environment of the droplet. In another example, heating of a droplet with a bead-bound barcode sequence in basic solution can also result in bead degradation and release of the attached barcode
30 sequence into the aqueous, inner environment of the droplet.

Any suitable number of species (e.g., primer, barcoded oligonucleotide) can be associated with a bead such that, upon release from the bead, the species (e.g., primer, e.g., barcoded oligonucleotide) are present in the partition at a pre-defined concentration. Such pre-defined concentration can be selected to facilitate certain reactions for generating a sequencing library, e.g., amplification, within the partition. In some cases, the pre-defined concentration of the primer can be limited by the process of producing nucleic acid molecule (e.g., oligonucleotide) bearing beads.

A degradable bead can include one or more species with a labile bond such that, when the bead/species is exposed to the appropriate stimulus, the bond is broken and the bead degrades.

The labile bond can be a chemical bond (e.g., covalent bond, ionic bond) or can be another type of physical interaction (e.g., van der Waals interactions, dipole-dipole interactions, etc.) In some embodiments, a crosslinker used to generate a bead can include a labile bond. Upon exposure to the appropriate conditions, the labile bond can be broken and the bead degraded. For example, upon exposure of a polyacrylamide gel bead that includes cystamine crosslinkers to a reducing agent, the disulfide bonds of the cystamine can be broken and the bead degraded.

A degradable bead can be useful in more quickly releasing an attached species (e.g., a nucleic acid molecule, a barcode sequence, a primer, etc.) from the bead when the appropriate stimulus is applied to the bead as compared to a bead that does not degrade. For example, for a species bound to an inner surface of a porous bead or in the case of an encapsulated species, the species can have greater mobility and accessibility to other species in solution upon degradation of the bead. In some embodiments, a species can also be attached to a degradable bead via a degradable linker (e.g., disulfide linker). The degradable linker can respond to the same stimuli as the degradable bead or the two degradable species can respond to different stimuli. For example, a barcode sequence can be attached, via a disulfide bond, to a polyacrylamide bead comprising cystamine. Upon exposure of the barcoded-bead to a reducing agent, the bead degrades and the barcode sequence is released upon breakage of both the disulfide linkage between the barcode sequence and the bead and the disulfide linkages of the cystamine in the bead.

As will be appreciated from the above description, while referred to as degradation of a bead, in many embodiments, degradation can refer to the disassociation of a bound or entrained species from a bead, both with and without structurally degrading the physical bead itself. For example, entrained species can be released from beads through osmotic pressure differences due

to, for example, changing chemical environments. By way of example, alteration of bead pore sizes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some cases, an increase in pore size due to osmotic swelling of a bead can permit the release of entrained species within the bead. In some embodiments, osmotic shrinking
5 of a bead can cause a bead to better retain an entrained species due to pore size contraction. Numerous chemical triggers can be used to trigger the degradation of beads within partitions. Examples of these chemical changes can include, but are not limited to pH-mediated changes to the integrity of a component within the bead, degradation of a component of a bead via cleavage of cross-linked bonds, and depolymerization of a component of a bead.

10 In some embodiments, a bead can be formed from materials that include degradable chemical cross-linkers, such as BAC or cystamine. Degradation of such degradable cross-linkers can be accomplished through a number of mechanisms. In some examples, a bead can be contacted with a chemical degrading agent that can induce oxidation, reduction or other chemical changes. For example, a chemical degrading agent can be a reducing agent, such as dithiothreitol
15 (DTT). Additional examples of reducing agents can include β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris(2-carboxyethyl) phosphine (TCEP), or combinations thereof. A reducing agent can degrade the disulfide bonds formed between gel precursors forming the bead, and thus, degrade the bead.

20 In certain embodiments, a change in pH of a solution, such as an increase in pH, can trigger degradation of a bead. In other embodiments, exposure to an aqueous solution, such as water, can trigger hydrolytic degradation, and thus degradation of the bead. In some cases, any combination of stimuli can trigger degradation of a bead. For example, a change in pH can enable a chemical agent (e.g., DTT) to become an effective reducing agent.

25 Beads can also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety of changes to a bead. For example, heat can cause a solid bead to liquefy. A change in heat can cause melting of a bead such that a portion of the bead degrades. In other cases, heat can increase the internal pressure of the bead components such that the bead ruptures or explodes. Heat can also act upon heat-sensitive polymers used as materials to construct beads.

30 In addition to beads and analytes, partitions that are formed can include a variety of different reagents and species. For example, when lysis reagents are present within the partitions,

the lysis reagents can facilitate the release of analytes within the partition. Examples of lysis agents include bioactive reagents, such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, etc., such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other lysis enzymes
5 available from, e.g., Sigma-Aldrich, Inc. (St. Louis, MO), as well as other commercially available lysis enzymes. Other lysis agents can additionally or alternatively be co-partitioned to cause the release analytes into the partitions. For example, in some cases, surfactant-based lysis solutions can be used to lyse cells, although these can be less desirable for emulsion based systems where the surfactants can interfere with stable emulsions. In some embodiments, lysis
10 solutions can include non-ionic surfactants such as, for example, TritonX-100 and Tween 20. In some embodiments, lysis solutions can include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). Electroporation, thermal, acoustic or mechanical cellular disruption can also be used in certain embodiments, e.g., non-emulsion based partitioning such as encapsulation of analytes that can be in addition to or in place of droplet partitioning, where
15 any pore size of the encapsulate is sufficiently small to retain nucleic acid fragments of a given size, following cellular disruption.

Examples of other species that can be co-partitioned with analytes in the partitions include, but are not limited to, DNase and RNase inactivating agents or inhibitors, such as proteinase K, chelating agents, such as EDTA, and other reagents employed in removing or
20 otherwise reducing negative activity or impact of different cell lysate components on subsequent processing of nucleic acids. Additional reagents can also be co-partitioned, including endonucleases to fragment DNA, DNA polymerase enzymes and dNTPs used to amplify nucleic acid fragments and to attach the barcode molecular tags to the amplified fragments.

Additional reagents can also include reverse transcriptase enzymes, including enzymes
25 with terminal transferase activity, primers and oligonucleotides, and switch oligonucleotides (also referred to herein as “switch oligos” or “template switching oligonucleotides”) which can be used for template switching. In some embodiments, template switching can be used to increase the length of a cDNA. Template switching can be used to append a predefined nucleic acid sequence to the cDNA. In an example of template switching, cDNA can be generated from
30 reverse transcription of a template, e.g., cellular mRNA, where a reverse transcriptase with terminal transferase activity can add additional nucleotides, e.g., poly(C), to the cDNA in a

template independent manner. Switch oligos can include sequences complementary to the additional nucleotides, e.g., poly(G). The additional nucleotides (e.g., poly(C)) on the cDNA can hybridize to the additional nucleotides (e.g., poly(G)) on the switch oligo, whereby the switch oligo can be used by the reverse transcriptase as template to further extend the cDNA.

- 5 Template switching oligonucleotides can include a hybridization region and a template region. The hybridization region can include any sequence capable of hybridizing to the target. In some cases, the hybridization region includes a series of G bases to complement the overhanging C bases at the 3' end of a cDNA molecule. The series of G bases can include 1 G base, 2 G bases, 3 G bases, 4 G bases, 5 G bases or more than 5 G bases. The template sequence can include any
- 10 sequence to be incorporated into the cDNA. In some cases, the template region includes at least 1 (e.g., at least 2, 3, 4, 5 or more) tag sequences and/or functional sequences. Switch oligos can include deoxyribonucleic acids; ribonucleic acids; bridged nucleic acids, modified nucleic acids including 2-Aminopurine, 2,6-Diaminopurine (2-Amino-dA), inverted dT, 5-Methyl dC, 2'-deoxyInosine, Super T (5-hydroxybutynl-2'-deoxyuridine), Super G (8-aza-7-deazaguanosine),
- 15 locked nucleic acids (LNAs), unlocked nucleic acids (UNAs, e.g., UNA-A, UNA-U, UNA-C, UNA-G), Iso-dG, Iso-dC, 2' Fluoro bases (e.g., Fluoro C, Fluoro U, Fluoro A, and Fluoro G), and combinations of the foregoing.

In some embodiments, beads that are partitioned with the analyte can include different types of oligonucleotides bound to the bead, where the different types of oligonucleotides bind to

20 different types of analytes. For example, a bead can include one or more first oligonucleotides (which can be capture probes, for example) that can bind or hybridize to a first type of analyte, such as mRNA for example, and one or more second oligonucleotides (which can be capture probes, for example) that can bind or hybridize to a second type of analyte, such as gDNA for example. Partitions can also include lysis agents that aid in releasing nucleic acids from the co-

25 partitioned cell, and can also include an agent (e.g., a reducing agent) that can degrade the bead and/or break covalent linkages between the oligonucleotides and the bead, releasing the oligonucleotides into the partition. The released barcoded oligonucleotides (which can also be barcoded) can hybridize with mRNA released from the cell and also with gDNA released from the cell.

30 Barcoded constructs thus formed from hybridization can include a first type of construct that includes a sequence corresponding to an original barcode sequence from the bead and a

sequence corresponding to a transcript from the cell, and a second type of construct that includes a sequence corresponding to the original barcode sequence from the bead and a sequence corresponding to genomic DNA from the cell. The barcoded constructs can then be released/removed from the partition and, in some embodiments, further processed to add any additional sequences. The resulting constructs can then be sequenced, the sequencing data processed, and the results used to spatially characterize the mRNA and the gDNA from the cell.

In another example, a partition includes a bead that includes a first type of oligonucleotide (e.g., a first capture probe) with a first barcode sequence, a poly(T) priming sequence that can hybridize with the poly(A) tail of an mRNA transcript, and a UMI barcode sequence that can uniquely identify a given transcript. The bead also includes a second type of oligonucleotide (e.g., a second capture probe) with a second barcode sequence, a targeted priming sequence that is capable of specifically hybridizing with a third barcoded oligonucleotide (e.g., an analyte capture agent) coupled to an antibody that is bound to the surface of the partitioned cell. The third barcoded oligonucleotide includes a UMI barcode sequence that uniquely identifies the antibody (and thus, the particular cell surface feature to which it is bound).

In this example, the first and second barcoded oligonucleotides include the same spatial barcode sequence (e.g., the first and second barcode sequences are the same), which permits downstream association of barcoded nucleic acids with the partition. In some embodiments, however, the first and second barcode sequences are different.

The partition also includes lysis agents that aid in releasing nucleic acids from the cell and can also include an agent (e.g., a reducing agent) that can degrade the bead and/or break a covalent linkage between the barcoded oligonucleotides and the bead, releasing them into the partition. The first type of released barcoded oligonucleotide can hybridize with mRNA released from the cell and the second type of released barcoded oligonucleotide can hybridize with the third type of barcoded oligonucleotide, forming barcoded constructs.

The first type of barcoded construct includes a spatial barcode sequence corresponding to the first barcode sequence from the bead and a sequence corresponding to the UMI barcode sequence from the first type of oligonucleotide, which identifies cell transcripts. The second type of barcoded construct includes a spatial barcode sequence corresponding to the second barcode sequence from the second type of oligonucleotide, and a UMI barcode sequence corresponding

to the third type of oligonucleotide (e.g., the analyte capture agent) and used to identify the cell surface feature. The barcoded constructs can then be released/removed from the partition and, in some embodiments, further processed to add any additional sequences. The resulting constructs are then sequenced, sequencing data processed, and the results used to characterize the mRNA and cell surface feature of the cell.

The foregoing discussion involves two specific examples of beads with oligonucleotides for analyzing two different analytes within a partition. More generally, beads that are partitioned can have any of the structures described previously, and can include any of the described combinations of oligonucleotides for analysis of two or more (e.g., three or more, four or more, five or more, six or more, eight or more, ten or more, 12 or more, 15 or more, 20 or more, 25 or more, 30 or more, 40 or more, 50 or more) different types of analytes within a partition. Examples of beads with combinations of different types of oligonucleotides (e.g., capture probes) for concurrently analyzing different combinations of analytes within partitions include, but are not limited to: (a) genomic DNA and cell surface features (e.g., using the analyte capture agents described herein); (b) mRNA and a lineage tracing construct; (c) mRNA and cell methylation status; (d) mRNA and accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq); (e) mRNA and cell surface or intracellular proteins and/or metabolites; (f) a barcoded analyte capture agent (e.g., the MHC multimers described herein) and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); and (g) mRNA and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein).

Additionally, in some embodiments, the unaggregated cell or disaggregated cells introduced and processed within partitions or droplets as described herein, can be removed from the partition, contacted with a spatial array, and spatially barcoded according to methods described herein. For example, single cells of an unaggregated cell sample can be partitioned into partitions or droplets as described herein. The partitions or droplets can include reagents to permeabilize a cell, barcode targeted cellular analyte(s) with a cellular barcode, and amplify the barcoded analytes. The partitions or droplets can be contacted with any of the spatial arrays described herein. In some embodiments, the partition can be dissolved, such that the contents of the partition are placed in contact with the capture probes of the spatial array. The capture probes

of the spatial array can then capture target analytes from the ruptured partitions or the droplets, and processed by the spatial workflows described herein.

(g) Analysis of Captured Analytes

5 Removal of sample from array

In some embodiments, after contacting a biological sample with a substrate that includes capture probes, a removal step can optionally be performed to remove all or a portion of the biological sample from the substrate. In some embodiments, the removal step includes enzymatic and/or chemical degradation of cells of the biological sample. For example, the removal step can
10 include treating the biological sample with an enzyme (e.g., a proteinase, e.g., proteinase K) to remove at least a portion of the biological sample from the substrate. In some embodiments, the removal step can include ablation of the tissue (e.g., laser ablation).

In some embodiments, provided herein are methods for spatially detecting an analyte (e.g., detecting the location of an analyte, e.g., a biological analyte) from a biological sample
15 (e.g., present in a biological sample), the method comprising: (a) optionally staining and/or imaging a biological sample on a substrate; (b) permeabilizing (e.g., providing a solution comprising a permeabilization reagent to) the biological sample on the substrate; (c) contacting the biological sample with an array comprising a plurality of capture probes, wherein a capture probe of the plurality captures the biological analyte; and (d) analyzing the captured biological
20 analyte, thereby spatially detecting the biological analyte; wherein the biological sample is fully or partially removed from the substrate.

In some embodiments, a biological sample is not removed from the substrate. For example, the biological sample is not removed from the substrate prior to releasing a capture probe (e.g., a capture probe bound to an analyte) from the substrate. In some embodiments, such
25 releasing comprises cleavage of the capture probe from the substrate (e.g., via a cleavage domain). In some embodiments, such releasing does not comprise releasing the capture probe from the substrate (e.g., a copy of the capture probe bound to an analyte can be made and the copy can be released from the substrate, e.g., via denaturation). In some embodiments, the biological sample is not removed from the substrate prior to analysis of an analyte bound to a
30 capture probe after it is released from the substrate. In some embodiments, the biological sample remains on the substrate during removal of a capture probe from the substrate and/or analysis of

an analyte bound to the capture probe after it is released from the substrate. In some embodiments, analysis of an analyte bound to capture probe from the substrate can be performed without subjecting the biological sample to enzymatic and/or chemical degradation of the cells (e.g., permeabilized cells) or ablation of the tissue (e.g., laser ablation).

5 In some embodiments, at least a portion of the biological sample is not removed from the substrate. For example, a portion of the biological sample can remain on the substrate prior to releasing a capture probe (e.g., a capture probe bound to an analyte) from the substrate and/or analyzing an analyte bound to a capture probe released from the substrate. In some embodiments, at least a portion of the biological sample is not subjected to enzymatic and/or chemical
10 degradation of the cells (e.g., permeabilized cells) or ablation of the tissue (e.g., laser ablation) prior to analysis of an analyte bound to a capture probe from the substrate.

 In some embodiments, provided herein are methods for spatially detecting an analyte (e.g., detecting the location of an analyte, e.g., a biological analyte) from a biological sample (e.g., present in a biological sample) that include: (a) optionally staining and/or imaging a
15 biological sample on a substrate; (b) permeabilizing (e.g., providing a solution comprising a permeabilization reagent to) the biological sample on the substrate; (c) contacting the biological sample with an array comprising a plurality of capture probes, wherein a capture probe of the plurality captures the biological analyte; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte; where the biological sample is not removed
20 from the substrate.

 In some embodiments, provided herein are methods for spatially detecting a biological analyte of interest from a biological sample that include: (a) staining and imaging a biological sample on a substrate; (b) providing a solution comprising a permeabilization reagent to the biological sample on the substrate; (c) contacting the biological sample with an array on a
25 substrate, wherein the array comprises one or more capture probe pluralities thereby allowing the one or more pluralities of capture probes to capture the biological analyte of interest; and (d) analyzing the captured biological analyte, thereby spatially detecting the biological analyte of interest; where the biological sample is not removed from the substrate.

 In some embodiments, the method further includes selecting a region of interest in the
30 biological sample to subject to spatial transcriptomic analysis. In some embodiments, one or more of the one or more capture probes include a capture domain. In some embodiments, one or

more of the one or more capture probe pluralities comprise a unique molecular identifier (UMI). In some embodiments, one or more of the one or more capture probe pluralities comprise a cleavage domain. In some embodiments, the cleavage domain comprises a sequence recognized and cleaved by a uracil-DNA glycosylase, apurinic/apyrimidinic (AP) endonuclease (APE1), U
5 uracil-specific excision reagent (USER), and/or an endonuclease VIII. In some embodiments, one or more capture probes do not comprise a cleavage domain and is not cleaved from the array.

A set of experiments performed determined methods that did not remove the biological sample from the substrate yielded higher quality sequencing data, higher median genes per cell, and higher median UMI counts per cell compared to a similar methods where the biological
10 sample was removed from the substrate (data not shown).

In some embodiments, a capture probe can be extended. For example, extending a capture probe can includes generating cDNA from a captured (hybridized) RNA. This process involves synthesis of a complementary strand of the hybridized nucleic acid, e.g., generating cDNA based on the captured RNA template (the RNA hybridized to the capture domain of the
15 capture probe). Thus, in an initial step of extending a capture probe, e.g., the cDNA generation, the captured (hybridized) nucleic acid, e.g., RNA, acts as a template for the extension, e.g., reverse transcription, step.

In some embodiments, the capture probe is extended using reverse transcription. For example, reverse transcription includes synthesizing cDNA (complementary or copy DNA) from
20 RNA, e.g., (messenger RNA), using a reverse transcriptase. In some embodiments, reverse transcription is performed while the tissue is still in place, generating an analyte library, where the analyte library includes the spatial barcodes from the adjacent capture probes. In some embodiments, the capture probe is extended using one or more DNA polymerases.

In some embodiments, a capture domain of a capture probe includes a primer for
25 producing the complementary strand of a nucleic acid hybridized to the capture probe, e.g., a primer for DNA polymerase and/or reverse transcription. The nucleic acid, e.g., DNA and/or cDNA, molecules generated by the extension reaction incorporate the sequence of the capture probe. The extension of the capture probe, e.g., a DNA polymerase and/or reverse transcription reaction, can be performed using a variety of suitable enzymes and protocols.

30 In some embodiments, a full-length DNA, e.g. cDNA, molecule is generated. In some embodiments, a “full-length” DNA molecule refers to the whole of the captured nucleic acid

molecule. However, if the nucleic acid, e.g. RNA, was partially degraded in the tissue sample, then the captured nucleic acid molecules will not be the same length as the initial RNA in the tissue sample. In some embodiments, the 3' end of the extended probes, e.g., first strand cDNA molecules, is modified. For example, a linker or adaptor can be ligated to the 3' end of the extended probes. This can be achieved using single stranded ligation enzymes such as T4 RNA ligase or CircligaseTM (available from Epicentre Biotechnologies, Madison, WI). In some embodiments, template switching oligonucleotides are used to extend cDNA in order to generate a full-length cDNA (or as close to a full-length cDNA as possible). In some embodiments, a second strand synthesis helper probe (a partially double stranded DNA molecule capable of hybridizing to the 3' end of the extended capture probe), can be ligated to the 3' end of the extended probe, e.g., first strand cDNA, molecule using a double stranded ligation enzyme such as T4 DNA ligase. Other enzymes appropriate for the ligation step are known in the art and include, e.g., Tth DNA ligase, Taq DNA ligase, *Thermococcus* sp. (strain 9^oN) DNA ligase (9^oNTM DNA ligase, New England Biolabs), AmpligaseTM (available from Epicentre Biotechnologies, Madison, WI), and SplintR (available from New England Biolabs, Ipswich, MA). In some embodiments, a polynucleotide tail, e.g., a poly(A) tail, is incorporated at the 3' end of the extended probe molecules. In some embodiments, the polynucleotide tail is incorporated using a terminal transferase active enzyme.

In some embodiments, double-stranded extended capture probes are treated to remove any unextended capture probes prior to amplification and/or analysis, e.g. sequence analysis. This can be achieved by a variety of methods, e.g., using an enzyme to degrade the unextended probes, such as an exonuclease enzyme, or purification columns.

In some embodiments, extended capture probes are amplified to yield quantities that are sufficient for analysis, e.g., via DNA sequencing. In some embodiments, the first strand of the extended capture probes (e.g., DNA and/or cDNA molecules) acts as a template for the amplification reaction (e.g., a polymerase chain reaction).

In some embodiments, the amplification reaction incorporates an affinity group onto the extended capture probe (e.g., RNA-cDNA hybrid) using a primer including the affinity group. In some embodiments, the primer includes an affinity group and the extended capture probes includes the affinity group. The affinity group can correspond to any of the affinity groups described previously.

In some embodiments, the extended capture probes including the affinity group can be coupled to a substrate specific for the affinity group. In some embodiments, the substrate can include an antibody or antibody fragment. In some embodiments, the substrate includes avidin or streptavidin and the affinity group includes biotin. In some embodiments, the substrate includes maltose and the affinity group includes maltose-binding protein. In some embodiments, the substrate includes maltose-binding protein and the affinity group includes maltose. In some embodiments, amplifying the extended capture probes can function to release the extended probes from the surface of the substrate, insofar as copies of the extended probes are not immobilized on the substrate.

In some embodiments, the extended capture probe or complement or amplicon thereof is released. The step of releasing the extended capture probe or complement or amplicon thereof from the surface of the substrate can be achieved in a number of ways. In some embodiments, an extended capture probe or a complement thereof is released from the array by nucleic acid cleavage and/or by denaturation (e.g. by heating to denature a double-stranded molecule).

In some embodiments, the extended capture probe or complement or amplicon thereof is released from the surface of the substrate (e.g., array) by physical means. For example, where the extended capture probe is indirectly immobilized on the array substrate, e.g. via hybridization to a surface probe, it can be sufficient to disrupt the interaction between the extended capture probe and the surface probe. Methods for disrupting the interaction between nucleic acid molecules include denaturing double stranded nucleic acid molecules art. A straightforward method for releasing the DNA molecules (i.e., of stripping the array of the extended probes) is to use a solution that interferes with the hydrogen bonds of the double stranded molecules. In some embodiments, the extended capture probe is released by applying heated water such as water or buffer of at least 85°C, e.g., at least 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99°C. In some embodiments, a solution including salts, surfactants, etc. that can further destabilize the interaction between the nucleic acid molecules is added to release the extended capture probe from the substrate.

In some embodiments, where the extended capture probe includes a cleavage domain, the extended capture probe is released from the surface of the substrate by cleavage. For example, the cleavage domain of the extended capture probe can be cleaved by any of the methods described herein. In some embodiments, the extended capture probe is released from the surface

of the substrate, e.g., via cleavage of a cleavage domain in the extended capture probe, prior to the step of amplifying the extended capture probe.

Capture probes can optionally include a “cleavage domain,” where one or more segments or regions of the capture probe (e.g., spatial barcodes and/or UMIs) can be releasably, cleavably, or reversibly attached to a feature, or some other substrate, so that spatial barcodes and/or UMIs can be released or be releasable through cleavage of a linkage between the capture probe and the feature, or released through degradation of the underlying support, allowing the spatial barcode(s) and/or UMI(s) of the cleaved capture probe to be accessed or be accessible by other reagents, or both.

In some embodiments, the capture probe is linked, via a disulfide bond, to a feature. In some embodiments, the capture probe is linked to a feature via a propylene group (e.g., Spacer C3). A reducing agent can be added to break the various disulfide bonds, resulting in release of the capture probe including the spatial barcode sequence. In another example, heating can also result in degradation and release of the attached capture probe. In some embodiments, the heating is done by laser (e.g., laser ablation) and features at specific locations can be degraded. In addition to thermally cleavable bonds, disulfide bonds, photo-sensitive bonds, and UV sensitive bonds, other non-limiting examples of labile bonds that can be coupled to a capture probe (i.e., spatial barcode) include an ester linkage (e.g., cleavable with an acid, a base, or hydroxylamine), a vicinal diol linkage (e.g., cleavable via sodium periodate), a Diels-Alder linkage (e.g., cleavable via heat), a sulfone linkage (e.g., cleavable via a base), a silyl ether linkage (e.g., cleavable via an acid), a glycosidic linkage (e.g., cleavable via an amylase), a peptide linkage (e.g., cleavable via a protease), or a phosphodiester linkage (e.g., cleavable via a nuclease (e.g., DNAase)).

In some embodiments, the cleavage domain includes a sequence that is recognized by one or more enzymes capable of cleaving a nucleic acid molecule, e.g., capable of breaking the phosphodiester linkage between two or more nucleotides. A bond can be cleavable via other nucleic acid molecule targeting enzymes, such as restriction enzymes (e.g., restriction endonucleases). For example, the cleavage domain can include a restriction endonuclease (restriction enzyme) recognition sequence. Restriction enzymes cut double-stranded or single stranded DNA at specific recognition nucleotide sequences known as restriction sites. In some embodiments, a rare-cutting restriction enzyme, i.e., enzymes with a long recognition site (at

least 8 base pairs in length), is used to reduce the possibility of cleaving elsewhere in the capture probe.

In some embodiments, the cleavage domain includes a poly(U) sequence which can be cleaved by a mixture of Uracil DNA glycosylase (UDG) and the DNA glycosylase-lyase
5 Endonuclease VIII, commercially known as the USER™ enzyme. In some embodiments, the cleavage domain can be a single U. In some embodiments, the cleavage domain can be an abasic site that can be cleaved with an abasic site-specific endonuclease (e.g., Endonuclease IV or Endonuclease VIII). Releasable capture probes can be available for reaction once released. Thus, for example, an activatable capture probe can be activated by releasing the capture probes from a
10 feature.

In some embodiments, the cleavage domain of the capture probe is a nucleotide sequence within the capture probe that is cleaved specifically, e.g., physically by light or heat, chemically or enzymatically. The location of the cleavage domain within the capture probe will depend on whether or not the capture probe is immobilized on the substrate such that it has a free 3' end
15 capable of functioning as an extension primer (e.g. by its 5' or 3' end). For example, if the capture probe is immobilized by its 5' end, the cleavage domain will be located 5' to the spatial barcode and/or UMI, and cleavage of said domain results in the release of part of the capture probe including the spatial barcode and/or UMI and the sequence 3' to the spatial barcode, and optionally part of the cleavage domain, from a feature. Alternatively, if the capture probe is
20 immobilized by its 3' end, the cleavage domain will be located 3' to the capture domain (and spatial barcode) and cleavage of said domain results in the release of part of the capture probe including the spatial barcode and the sequence 3' to the spatial barcode from a feature. In some embodiments, cleavage results in partial removal of the cleavage domain. In some embodiments, cleavage results in complete removal of the cleavage domain, particularly when the capture
25 probes are immobilized via their 3' end as the presence of a part of the cleavage domain can interfere with the hybridization of the capture domain and the target nucleic acid and/or its subsequent extension.

In some embodiments, where the capture probe is immobilized to the substrate indirectly, e.g., via a surface probe defined below, the cleavage domain includes one or more mismatch
30 nucleotides, so that the complementary parts of the surface probe and the capture probe are not 100% complementary (for example, the number of mismatched base pairs can one, two, or three

base pairs). Such a mismatch is recognized, e.g., by the MutY and T7 endonuclease I enzymes, which results in cleavage of the nucleic acid molecule at the position of the mismatch.

In some embodiments, where the capture probe is immobilized to the feature indirectly, e.g., via a surface probe, the cleavage domain includes a nickase recognition site or sequence. In this respect, nickase enzymes cleave only one strand in a nucleic acid duplex. Nickases are endonucleases which cleave only a single strand of a DNA duplex. Thus, the cleavage domain can include a nickase recognition site close to the 5' end of the surface probe (and/or the 5' end of the capture probe) such that cleavage of the surface probe or capture probe destabilizes the duplex between the surface probe and capture probe thereby releasing the capture probe) from the feature.

Nickase enzymes can also be used in some embodiments where the capture probe is immobilized to the feature directly. For example, the substrate can be contacted with a nucleic acid molecule that hybridizes to the cleavage domain of the capture probe to provide or reconstitute a nickase recognition site, e.g., a cleavage helper probe. Thus, contact with a nickase enzyme will result in cleavage of the cleavage domain thereby releasing the capture probe from the feature. Such cleavage helper probes can also be used to provide or reconstitute cleavage recognition sites for other cleavage enzymes, e.g., restriction enzymes.

Some nickases introduce single-stranded nicks only at particular sites on a DNA molecule, by binding to and recognizing a particular nucleotide recognition sequence. A number of naturally-occurring nickases have been discovered, of which at present the sequence recognition properties have been determined for at least four. Nickases are described in U.S. Patent No. 6,867,028, which is herein incorporated by reference in its entirety. In general, any suitable nickase can be used to bind to a complementary nickase recognition site of a cleavage domain. Following use, the nickase enzyme can be removed from the assay or inactivated following release of the capture probes to prevent unwanted cleavage of the capture probes.

In some embodiments, a cleavage domain for separating spatial barcodes from a feature is absent from the capture probe. For example, a substrate having a capture probe lacking a cleavage domain can be used for spatial analysis (see, e.g., corresponding substrates and probes described Macosko et al., (2015) Cell 161, 1202–1214, the entire contents of which are incorporated herein by reference.

In some embodiments, the region of the capture probe corresponding to the cleavage domain can be used for some other function. For example, an additional region for nucleic acid extension or amplification can be included where the cleavage domain would normally be positioned. In such embodiments, the region can supplement the functional domain or even exist
5 as an additional functional domain. In some embodiments, the cleavage domain is present but its use is optional.

After analytes from the sample have hybridized or otherwise been associated with capture probes, analyte capture agents, or other barcoded oligonucleotide sequences according to any of the methods described above in connection with the general spatial cell-based analytical
10 methodology, the barcoded constructs that result from hybridization/association are analyzed via sequencing to identify the analytes.

In some embodiments, where a sample is barcoded directly via hybridization with capture probes or analyte capture agents hybridized, bound, or associated with either the cell surface, or introduced into the cell, as described above, sequencing can be performed on the intact sample.
15 Alternatively, if the barcoded sample has been separated into fragments, cell groups, or individual cells, as described above, sequencing can be performed on individual fragments, cell groups, or cells. For analytes that have been barcoded via partitioning with beads, as described above, individual analytes (e.g., cells, or cellular contents following lysis of cells) can be extracted from the partitions by breaking the partitions, and then analyzed by sequencing to
20 identify the analytes.

A wide variety of different sequencing methods can be used to analyze barcoded analyte constructs. In general, sequenced polynucleotides can be, for example, nucleic acid molecules such as deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), including variants or derivatives thereof (e.g., single stranded DNA or DNA/RNA hybrids, and nucleic acid molecules
25 with a nucleotide analog).

Sequencing of polynucleotides can be performed by various commercial systems. More generally, sequencing can be performed using nucleic acid amplification, polymerase chain reaction (PCR) (e.g., digital PCR and droplet digital PCR (ddPCR), quantitative PCR, real time PCR, multiplex PCR, PCR-based singleplex methods, emulsion PCR), and/or isothermal
30 amplification.

Other examples of methods for sequencing genetic material include, but are not limited to, DNA hybridization methods (e.g., Southern blotting), restriction enzyme digestion methods, Sanger sequencing methods, next-generation sequencing methods (e.g., single-molecule real-time sequencing, nanopore sequencing, and Polony sequencing), ligation methods, and
5 microarray methods. Additional examples of sequencing methods that can be used include targeted sequencing, single molecule real-time sequencing, exon sequencing, electron microscopy-based sequencing, panel sequencing, transistor-mediated sequencing, direct sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel
10 electrophoresis, duplex sequencing, cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, co-amplification at lower denaturation temperature-PCR (COLD-PCR), sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-
15 synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, 454 sequencing, Solexa Genome Analyzer sequencing, SOLiD™ sequencing, MS-PET sequencing, and any combinations thereof.

Sequence analysis of the nucleic acid molecules (including barcoded nucleic acid molecules or derivatives thereof) can be direct or indirect. Thus, the sequence analysis substrate
20 (which can be viewed as the molecule which is subjected to the sequence analysis step or process) can directly be the barcoded nucleic acid molecule or it can be a molecule which is derived therefrom (e.g., a complement thereof). Thus, for example, in the sequence analysis step of a sequencing reaction, the sequencing template can be the barcoded nucleic acid molecule or it can be a molecule derived therefrom. For example, a first and/or second strand DNA molecule
25 can be directly subjected to sequence analysis (e.g. sequencing), i.e., can directly take part in the sequence analysis reaction or process (e.g. the sequencing reaction or sequencing process, or be the molecule which is sequenced or otherwise identified). Alternatively, the barcoded nucleic acid molecule can be subjected to a step of second strand synthesis or amplification before sequence analysis (e.g. sequencing or identification by another technique). The sequence analysis
30 substrate (e.g., template) can thus be an amplicon or a second strand of a barcoded nucleic acid molecule.

In some embodiments, both strands of a double stranded molecule can be subjected to sequence analysis (e.g., sequenced). In some embodiments, single stranded molecules (e.g. barcoded nucleic acid molecules) can be analyzed (e.g. sequenced). To perform single molecule sequencing, the nucleic acid strand can be modified at the 3' end.

5 Massively parallel sequencing techniques can be used for sequencing nucleic acids, as described above. In one embodiment, a massively parallel sequencing technique can be based on reversible dye-terminators. As an example, DNA molecules are first attached to primers on, e.g., a glass or silicon substrate, and amplified so that local clonal colonies are formed (bridge amplification). Four types of ddNTPs are added, and non-incorporated nucleotides are washed
10 away. Unlike pyrosequencing, the DNA is only extended one nucleotide at a time due to a blocking group (e.g., 3' blocking group present on the sugar moiety of the ddNTP). A detector acquires images of the fluorescently labelled nucleotides, and then the dye along with the terminal 3' blocking group is chemically removed from the DNA, as a precursor to a subsequent cycle. This process can be repeated until the required sequence data is obtained.

15 As another example, massively parallel pyrosequencing techniques can also be used for sequencing nucleic acids. In pyrosequencing, the nucleic acid is amplified inside water droplets in an oil solution (emulsion PCR), with each droplet containing a single nucleic acid template attached to a single primer-coated bead that then forms a clonal colony. The sequencing system contains many picolitre-volume wells each containing a single bead and sequencing enzymes.
20 Pyrosequencing uses luciferase to generate light for detection of the individual nucleotides added to the nascent nucleic acid and the combined data are used to generate sequence reads.

 As another example application of pyrosequencing, released PPi can be detected by being immediately converted to adenosine triphosphate (ATP) by ATP sulfurylase, and the level of ATP generated can be detected via luciferase-produced photons, such as described in Ronaghi, et
25 al., *Anal. Biochem.* 242(1), 84-9 (1996); Ronaghi, *Genome Res.* 11(1), 3-11 (2001); Ronaghi et al. *Science* 281 (5375), 363 (1998); and U.S. Pat. Nos. 6,210,891, 6,258,568, and 6,274,320, the entire contents of each of which are incorporated herein by reference.

 In some embodiments, sequencing is performed by detection of hydrogen ions that are released during the polymerization of DNA. A microwell containing a template DNA strand to
30 be sequenced can be flooded with a single type of nucleotide. If the introduced nucleotide is complementary to the leading template nucleotide, it is incorporated into the growing

complementary strand. This causes the release of a hydrogen ion that triggers a hypersensitive ion sensor, which indicates that a reaction has occurred. If homopolymer repeats are present in the template sequence, multiple nucleotides will be incorporated in a single cycle. This leads to a corresponding number of released hydrogen ions and a proportionally higher electronic signal.

5 In some embodiments, sequencing can be performed in-situ. In-situ sequencing methods are particularly useful, for example, when the biological sample remains intact after analytes on the sample surface (e.g., cell surface analytes) or within the sample (e.g., intracellular analytes) have been barcoded. In-situ sequencing typically involves incorporation of a labeled nucleotide (e.g., fluorescently labeled mononucleotides or dinucleotides) in a sequential, template-
10 dependent manner or hybridization of a labeled primer (e.g., a labeled random hexamer) to a nucleic acid template such that the identities (i.e., nucleotide sequence) of the incorporated nucleotides or labeled primer extension products can be determined, and consequently, the nucleotide sequence of the corresponding template nucleic acid. Aspects of in-situ sequencing are described, for example, in Mitra et al., (2003) Anal. Biochem. 320, 55-65, and Lee et al.,
15 (2014) Science, 343(6177), 1360-1363, the entire contents of each of which are incorporated herein by reference.

In addition, examples of methods and systems for performing in-situ sequencing are described in PCT Patent Application Publication Nos. WO2014/163886, WO2018/045181, WO2018/045186, and in U.S. Patent Nos. 10,138,509 and 10,179,932, the entire contents of each
20 of which are incorporated herein by reference. Example techniques for in-situ sequencing include, but are not limited to, STARmap (described for example in Wang et al., (2018) Science, 361(6499) 5691), MERFISH (described for example in Moffitt, (2016) Methods in Enzymology, 572, 1-49), and FISSEQ (described for example in U.S. Patent Application Publication No. 2019/0032121). The entire contents of each of the foregoing references are incorporated herein
25 by reference.

For analytes that have been barcoded via partitioning, barcoded nucleic acid molecules or derivatives thereof (e.g., barcoded nucleic acid molecules to which one or more functional sequences have been added, or from which one or more features have been removed) can be pooled and processed together for subsequent analysis such as sequencing on high throughput
30 sequencers. Processing with pooling can be implemented using barcode sequences. For example, barcoded nucleic acid molecules of a given partition can have the same barcode, which is

different from barcodes of other spatial partitions. Alternatively, barcoded nucleic acid molecules of different partitions can be processed separately for subsequent analysis (e.g., sequencing).

In some embodiments, where capture probes do not contain a spatial barcode, the spatial
5 barcode can be added after the capture probe captures analytes from a biological sample and before analysis of the analytes. When a spatial barcode is added after an analyte is captured, the barcode can be added after amplification of the analyte (e.g., reverse transcription and polymerase amplification of RNA). In some embodiments, analyte analysis uses direct sequencing of one or more captured analytes, such as direct sequencing of hybridized RNA. In
10 some embodiments, direct sequencing is performed after reverse transcription of hybridized RNA. In some embodiments direct sequencing is performed after amplification of reverse transcription of hybridized RNA.

In some embodiments, direct sequencing of captured RNA is performed by sequencing-by-synthesis (SBS). In some embodiments, a sequencing primer is complementary to a sequence
15 in one or more of the domains of a capture probe (e.g., functional domain). In such embodiments, sequencing-by-synthesis can include reverse transcription and/or amplification in order to generate a template sequence (e.g., functional domain) from which a primer sequence can bind.

SBS can involve hybridizing an appropriate primer, sometimes referred to as a
20 sequencing primer, with the nucleic acid template to be sequenced, extending the primer, and detecting the nucleotides used to extend the primer. Preferably, the nucleic acid used to extend the primer is detected before a further nucleotide is added to the growing nucleic acid chain, thus allowing base-by-base in situ nucleic acid sequencing. The detection of incorporated nucleotides is facilitated by including one or more labelled nucleotides in the primer extension reaction. To
25 allow the hybridization of an appropriate sequencing primer to the nucleic acid template to be sequenced, the nucleic acid template should normally be in a single stranded form. If the nucleic acid templates making up the nucleic acid spots are present in a double stranded form these can be processed to provide single stranded nucleic acid templates using methods well known in the art, for example by denaturation, cleavage etc. The sequencing primers which are hybridized to
30 the nucleic acid template and used for primer extension are preferably short oligonucleotides, for example, 15 to 25 nucleotides in length. The sequencing primers can be provided in solution or

in an immobilized form. Once the sequencing primer has been annealed to the nucleic acid template to be sequenced by subjecting the nucleic acid template and sequencing primer to appropriate conditions, primer extension is carried out, for example using a nucleic acid polymerase and a supply of nucleotides, at least some of which are provided in a labelled form, and conditions suitable for primer extension if a suitable nucleotide is provided.

Preferably after each primer extension step, a washing step is included in order to remove unincorporated nucleotides which can interfere with subsequent steps. Once the primer extension step has been carried out, the nucleic acid colony is monitored to determine whether a labelled nucleotide has been incorporated into an extended primer. The primer extension step can then be repeated to determine the next and subsequent nucleotides incorporated into an extended primer. If the sequence being determined is unknown, the nucleotides applied to a given colony are usually applied in a chosen order which is then repeated throughout the analysis, for example dATP, dTTP, dCTP, dGTP.

SBS techniques which can be used are described for example, but not limited to, those in U.S. Patent App. Pub. No. 2007/0166705, U.S. Patent App. Pub. No. 2006/0188901, U.S. Patent 7,057,026, U.S. Patent App. Pub. No. 2006/0240439, U.S. Patent App. Pub. No. 2006/0281109, PCT Patent App. Pub. No. WO 05/065814, U.S. Patent App. Pub. No. 2005/0100900, PCT Patent App. Pub. No. WO 06/064199, PCT Patent App. Pub. No. WO07/010,251, U.S. Patent App. Pub. No. 2012/0270305, U.S. Patent App. Pub. No. 2013/0260372, and U.S. Patent App. Pub. No. 2013/0079232, the entire contents of each of which are incorporated herein by reference.

In some embodiments, direct sequencing of captured RNA is performed by sequential fluorescence hybridization (e.g., sequencing by hybridization). In some embodiments, a hybridization reaction where RNA is hybridized to a capture probe is performed in situ. In some embodiments, captured RNA is not amplified prior to hybridization with a sequencing probe. In some embodiments, RNA is amplified prior to hybridization with sequencing probes (e.g., reverse transcription to cDNA and amplification of cDNA). In some embodiments, amplification is performed using single-molecule hybridization chain reaction. In some embodiments, amplification is performed using rolling chain amplification.

Sequential fluorescence hybridization can involve sequential hybridization of probes including degenerate primer sequences and a detectable label. A degenerate primer sequence is a

short oligonucleotide sequence which is capable of hybridizing to any nucleic acid fragment independent of the sequence of said nucleic acid fragment. For example, such a method could include the steps of: (a) providing a mixture including four probes, each of which includes either A, C, G, or T at the 5'-terminus, further including degenerate nucleotide sequence of 5 to 11 nucleotides in length, and further including a functional domain (e.g., fluorescent molecule) that is distinct for probes with A, C, G, or T at the 5'-terminus; (b) associating the probes of step (a) to the target polynucleotide sequences, whose sequence needs will be determined by this method; (c) measuring the activities of the four functional domains and recording the relative spatial location of the activities; (d) removing the reagents from steps (a)-(b) from the target polynucleotide sequences; and repeating steps (a)-(d) for n cycles, until the nucleotide sequence of the spatial domain for each bead is determined, with modification that the oligonucleotides used in step (a) are complementary to part of the target polynucleotide sequences and the positions 1 through n flanking the part of the sequences. Because the barcode sequences are different, in some embodiments, these additional flanking sequences are degenerate sequences. The fluorescent signal from each spot on the array for cycles 1 through n can be used to determine the sequence of the target polynucleotide sequences.

In some embodiments, direct sequencing of captured RNA using sequential fluorescence hybridization is performed in vitro. In some embodiments, captured RNA is amplified prior to hybridization with a sequencing probe (e.g., reverse transcription to cDNA and amplification of cDNA). In some embodiments, a capture probe containing captured RNA is exposed to the sequencing probe targeting coding regions of RNA. In some embodiments, one or more sequencing probes are targeted to each coding region. In some embodiments, the sequencing probe is designed to hybridize with sequencing reagents (e.g., a dye-labeled readout oligonucleotides). A sequencing probe can then hybridize with sequencing reagents. In some embodiments, output from the sequencing reaction is imaged. In some embodiments, a specific sequence of cDNA is resolved from an image of a sequencing reaction. In some embodiments, reverse transcription of captured RNA is performed prior to hybridization to the sequencing probe. In some embodiments, the sequencing probe is designed to target complementary sequences of the coding regions of RNA (e.g., targeting cDNA).

In some embodiments, a captured RNA is directly sequenced using a nanopore-based method. In some embodiments, direct sequencing is performed using nanopore direct RNA

sequencing in which captured RNA is translocated through a nanopore. A nanopore current can be recorded and converted into a base sequence. In some embodiments, captured RNA remains attached to a substrate during nanopore sequencing. In some embodiments, captured RNA is released from the substrate prior to nanopore sequencing. In some embodiments, where the
5 analyte of interest is a protein, direct sequencing of the protein can be performed using nanopore-based methods. Examples of nanopore-based sequencing methods that can be used are described in Deamer et al., Trends Biotechnol. 18, 147-151 (2000); Deamer et al., Acc. Chem. Res. 35:817-825 (2002); Li et al., Nat. Mater. 2:611-615 (2003); Soni et al., Clin. Chem. 53, 1996-2001 (2007); Healy et al., Nanomed. 2, 459-481 (2007); Cockroft et al., J. Am. Chem. Soc. 130,
10 818-820 (2008); and in U.S. Patent 7,001,792. The entire contents of each of the foregoing references are incorporated herein by reference.

In some embodiments, direct sequencing of captured RNA is performed using single molecule sequencing by ligation. Such techniques utilize DNA ligase to incorporate oligonucleotides and identify the incorporation of such oligonucleotides. The oligonucleotides
15 typically have different labels that are correlated with the identity of a particular nucleotide in a sequence to which the oligonucleotides hybridize. Aspects and features involved in sequencing by ligation are described, for example, in Shendure et al. Science (2005), 309: 1728-1732, and in U.S. Patent Nos. 5,599,675; 5,750,341; 6,969,488; 6,172,218; and 6,306,597, the entire contents of each of which are incorporated herein by reference.

In some embodiments, nucleic acid hybridization can be used for sequencing. These
20 methods utilize labeled nucleic acid decoder probes that are complementary to at least a portion of a barcode sequence. Multiplex decoding can be performed with pools of many different probes with distinguishable labels. Non-limiting examples of nucleic acid hybridization sequencing are described for example in U.S. Pat. No. 8,460,865, and in Gunderson et al.,
25 Genome Research 14:870-877 (2004), the entire contents of each of which are incorporated herein by reference.

In some embodiments, commercial high-throughput digital sequencing techniques can be used to analyze barcode sequences, in which DNA templates are prepared for sequencing not one at a time, but in a bulk process, and where many sequences are read out preferably in parallel, or
30 alternatively using an ultra-high throughput serial process that itself may be parallelized. Examples of such techniques include Illumina[®] sequencing (e.g., flow cell-based sequencing

techniques), sequencing by synthesis using modified nucleotides (such as commercialized in TruSeq™ and HiSeq™ technology by Illumina, Inc., San Diego, CA), HeliScope™ by Helicos Biosciences Corporation, Cambridge, MA, and PacBio RS by Pacific Biosciences of California, Inc., Menlo Park, CA), sequencing by ion detection technologies (Ion Torrent, Inc., South San Francisco, CA), and sequencing of DNA nanoballs (Complete Genomics, Inc., Mountain View, CA).

In some embodiments, detection of a proton released upon incorporation of a nucleotide into an extension product can be used in the methods described herein. For example, the sequencing methods and systems described in U.S. Patent Application Publication Nos. 2009/0026082, 2009/0127589, 2010/0137143, and 2010/0282617, can be used to directly sequence barcodes.

In some embodiments, real-time monitoring of DNA polymerase activity can be used during sequencing. For example, nucleotide incorporations can be detected through fluorescence resonance energy transfer (FRET), as described for example in Levene et al., Science (2003), 299, 682-686, Lundquist et al., Opt. Lett. (2008), 33, 1026-1028, and Korlach et al., Proc. Natl. Acad. Sci. USA (2008), 105, 1176-1181. The entire contents of each of the foregoing references are incorporated herein by reference herein.

In some embodiments, the methods described herein can be used to assess analyte levels and/or expression in a cell or a biological sample over time (e.g., before or after treatment with an agent or different stages of differentiation). In some examples, the methods described herein can be performed on multiple similar biological samples or cells obtained from the subject at a different time points (e.g., before or after treatment with an agent, different stages of differentiation, different stages of disease progression, different ages of the subject, or before or after development of resistance to an agent).

(h) Spatially Resolving Analyte Information

In some embodiments, a lookup table (LUT) can be used to associate one property with another property of a feature. These properties include, e.g., locations, barcodes (e.g., nucleic acid barcode molecules), spatial barcodes, optical labels, molecular tags, and other properties.

In some embodiments, a lookup table can associate the plurality of nucleic acid barcode molecules with the features. In some embodiments, the optical label of a feature can permit associating the feature with the biological particle (e.g., cell or nuclei). The association of the

feature with the biological particle can further permit associating a nucleic acid sequence of a nucleic acid molecule of the biological particle to one or more physical properties of the biological particle (e.g., a type of a cell or a location of the cell). For example, based on the relationship between the barcode and the optical label, the optical label can be used to determine the location of a feature, thus associating the location of the feature with the barcode sequence of the feature. Subsequent analysis (e.g., sequencing) can associate the barcode sequence and the analyte from the sample. Accordingly, based on the relationship between the location and the barcode sequence, the location of the biological analyte can be determined (e.g., in a specific type of cell, in a cell at a specific location of the biological sample).

In some embodiments, the feature can have a plurality of nucleic acid barcode molecules attached thereto. The plurality of nucleic acid barcode molecules can include barcode sequences. The plurality of nucleic acid molecules attached to a given feature can have the same barcode sequences, or two or more different barcode sequences. Different barcode sequences can be used to provide improved spatial location accuracy.

As discussed above, analytes obtained from a sample, such as RNA, DNA, peptides, lipids, and proteins, can be further processed. In particular, the contents of individual cells from the sample can be provided with unique spatial barcode sequences such that, upon characterization of the analytes, the analytes can be attributed as having been derived from the same cell. More generally, spatial barcodes can be used to attribute analytes to corresponding spatial locations in the sample. For example, hierarchical spatial positioning of multiple pluralities of spatial barcodes can be used to identify and characterize analytes over a particular spatial region of the sample. In some embodiments, the spatial region corresponds to a particular spatial region of interest previously identified, e.g., a particular structure of cytoarchitecture previously identified. In some embodiments, the spatial region corresponds to a small structure or group of cells that cannot be seen with the naked eye. In some embodiments, a unique molecular identifier can be used to identify and characterize analytes at a single cell level.

The analyte can include a nucleic acid molecule, which can be barcoded with a barcode sequence of a nucleic acid barcode molecule. In some embodiments, the barcoded analyte can be sequenced to obtain a nucleic acid sequence. In some embodiments, the nucleic acid sequence can include genetic information associate with the sample. The nucleic acid sequence can include the barcode sequence, or a complement thereof. The barcode sequence, or a complement thereof,

of the nucleic acid sequence can be electronically associated with the property (e.g., color and/or intensity) of the analyte using the LUT to identify the associated feature in an array.

In some embodiments, two- or three-dimensional spatial profiling of one or more analytes present in a biological sample can be performed using a proximity capture reaction, which is a reaction that detects two analytes that are spatially close to each other and/or interacting with each other. For example, a proximity capture reaction can be used to detect sequences of DNA that are close in space to each other, e.g., the DNA sequences can be within the same chromosome, but separated by about 700 bp or less. As another example, a proximity capture reaction can be used to detect protein associations, e.g., two proteins that interact with each other. A proximity capture reaction can be performed in situ to detect two analytes that are spatially close to each other and/or interacting with each other inside a cell. Non-limiting examples of proximity capture reactions include DNA nanoscopy, DNA microscopy, and chromosome conformation capture methods. Chromosome conformation capture (3C) and derivative experimental procedures can be used to estimate the spatial proximity between different genomic elements. Non-limiting examples of chromatin capture methods include chromosome conformation capture (3-C), conformation capture-on-chip (4-C), 5-C, ChIA-PET, Hi-C, targeted chromatin capture (T2C). Examples of such methods are described, for example, in Miele et al., *Methods Mol Biol.* (2009), 464, Simonis et al., *Nat. Genet.* (2006), 38(11): 1348-54, Raab et al., *Embo. J.* (2012), 31(2): 330–350, and Eagen et al., *Trends Biochem. Sci.* (2018) 43(6): 469-478, the entire contents of each of which is incorporated herein by reference.

In some embodiments, the proximity capture reaction includes proximity ligation. In some embodiments, proximity ligation can include using antibodies with attached DNA strands that can participate in ligation, replication, and sequence decoding reactions. For example, a proximity ligation reaction can include oligonucleotides attached to pairs of antibodies that can be joined by ligation if the antibodies have been brought in proximity to each oligonucleotide, e.g., by binding the same target protein (complex), and the DNA ligation products that form are then used to template PCR amplification, as described for example in Söderberg et al., *Methods.* (2008), 45(3): 227-32, the entire contents of which are incorporated herein by reference. In some embodiments, proximity ligation can include chromosome conformation capture methods. In some embodiments, the proximity capture reaction is performed on analytes within about 400 nm distance (e.g., about 300 nm, about 200 nm, about 150 nm, about 100 nm, about 50 nm,

about 25 nm, about 10 nm, or about 5 nm) from each other. In general, proximity capture reactions can be reversible or irreversible.

III. General Spatial Cell-Based Analytical Methodology

5 (a) Barcoding Biological Sample

In some embodiments, provided herein are methods and materials for attaching and/or introducing a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a biological sample (e.g., to a cell in a biological sample) for use in spatial analysis. In some embodiments, a plurality of molecules (e.g., a plurality of nucleic acid molecules) having a
10 plurality of barcodes (e.g., a plurality of spatial barcodes) are introduced to a biological sample (e.g., to a plurality of cells in a biological sample) for use in spatial analysis.

FIG. 18 is a schematic diagram depicting cell tagging using either covalent conjugation of the analyte binding moiety to the cell surface or non-covalent interactions with cell membrane elements. **FIG. 18** lists non-exhaustive examples of a covalent analyte binding moiety/cell
15 surface interactions, including protein targeting, amine conjugation using NHS chemistry, cyanuric chloride, thiol conjugation via maleimide addition, as well as targeting glycoproteins/glycolipids expressed on the cell surface via click chemistry. Non-exhaustive examples of non-covalent interactions with cell membrane elements include lipid modified oligos, biocompatible anchor for cell membrane (oleyl-PEG-NHS), lipid modified positive
20 neutral polymer, and antibody to membrane proteins. The cell tag can be used in combination with an analyte capture agent and cleavable or non-cleavable spatially-barcoded capture probes for spatial and multiplexing applications.

In some embodiments, a plurality of molecules (e.g., a plurality of nucleic acid molecules) having a plurality of barcodes (e.g., a plurality of spatial barcodes) are introduced to a
25 biological sample (e.g., to a plurality of cells in a biological sample) for use in spatial analysis, wherein the plurality of molecules are introduced to the biological sample in an arrayed format. In some embodiments, a plurality of molecules (e.g., a plurality of nucleic acid molecules) having a plurality of barcodes are provided on a substrate (e.g., any of the variety of substrates described herein) in any of the variety of arrayed formats described herein, and the biological
30 sample is contacted with the molecules on the substrate such that the molecules are introduced to the biological sample. In some embodiments, the molecules that are introduced to the biological

sample are cleavably attached to the substrate, and are cleaved from the substrate and released to the biological sample when contacted with the biological sample. In some embodiments, the molecules that are introduced to the biological sample are attached to the substrate covalently prior to cleavage. In some embodiments, the molecules that are introduced to the biological sample are non-covalently attached to the substrate (e.g., via hybridization), and are released from the substrate to the biological sample when contacted with the biological sample.

In some embodiments, a plurality of molecules (e.g., a plurality of nucleic acid molecules) having a plurality of barcodes (e.g., a plurality of spatial barcodes) are migrated or transferred from a substrate to cells of a biological sample. In some embodiments, migrating a plurality of molecules from a substrate to cells of a biological sample includes applying a force (e.g., mechanical, centrifugal, or electrophoretic) to the substrate and/or the biological sample to facilitate migration of the plurality of molecules from the substrate to the biological sample.

In some embodiments of any of the spatial analysis methods described herein, physical force is used to facilitate attachment to or introduction of a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) into a biological sample (e.g., a cell present in a biological sample). As used herein, “physical force” refers to the use of a physical force to counteract the cell membrane barrier in facilitating intracellular delivery of molecules. Examples of physical force instruments and methods that can be used in accordance with materials and methods described herein include the use of a needle, ballistic DNA, electroporation, sonoporation, photoporation, magnetofection, hydroporation, and combinations thereof.

In some embodiments, biological samples (e.g., cells in a biological sample) can be labelled using cell-tagging agents where the cell-tagging agents facilitate the introduction of the molecules (e.g., nucleic acid molecules) having barcodes (e.g., spatial barcodes) into the biological sample (e.g., into cells in a biological sample). As used herein, the term “cell-tagging agent” refers to a molecule having a moiety that is capable of attaching to the surface of a cell (e.g., thus attaching the barcode to the surface of the cell) and/or penetrating and passing through the cell membrane (e.g., thus introducing the barcode to the interior of the cell). In some embodiments, a cell-tagging agent includes a barcode (e.g., a spatial barcode). The barcode of a barcoded cell-tagging agent can be any of the variety of barcodes described herein. In some embodiments, the barcode of a barcoded cell-tagging agent is a spatial barcode. In some embodiments, a cell-tagging agent comprises a nucleic acid molecule that includes the barcode

(e.g., the spatial barcode). In some embodiments, the barcode of a barcoded cell-tagging agent identifies the associated molecule, where each spatial barcode is associated with a particular molecule. In some embodiments, one or more molecules are applied to a sample. In some embodiments, a nucleic acid molecule that includes the barcode is covalently attached to the cell-tagging agent. In some embodiments, a nucleic acid molecule that includes the barcode is non-covalently attached to the cell-tagging agent. A non-limiting example of non-covalent attachment includes hybridizing the nucleic acid molecule that includes the barcode to a nucleic acid molecule on the cell-tagging agent (which nucleic acid molecule on the cell-tagging agent can be bound to the cell-tagging agent covalently or non-covalently). In some embodiments, a nucleic acid molecule that is attached to a cell-tagging agent that includes a barcode (e.g., a spatial barcode) also includes one or more additional domains. Such additional domains include, without limitation, a PCR handle, a sequencing priming site, a domain for hybridizing to another nucleic acid molecule, and combinations thereof.

In some embodiments, a cell-tagging agent attaches to the surface of a cell. When the cell-tagging agent includes a barcode (e.g., a nucleic acid that includes a spatial barcode), the barcode is also attached to the surface of the cell. In some embodiments of any of the spatial analysis methods described herein, a cell-tagging agent attaches covalently to the cell surface to facilitate introduction of the spatial profiling reagents. In some embodiments of any of the spatial analysis methods described herein, a cell-tagging agent attaches non-covalently to the cell surface to facilitate introduction of the spatial profiling reagents.

In some embodiments, once a cell or cells in a biological sample is spatially tagged with a cell-tagging agent(s), spatial analysis of analytes present in the biological sample is performed. In some embodiments, such spatial analysis includes dissociating the spatially-tagged cells of the biological sample (or a subset of the spatially-tagged cells of the biological sample) and analyzing analytes present in those cells on a cell-by-cell basis. Any of a variety of methods for analyzing analytes present in cells on a cell-by-cell basis can be used. Non-limiting examples include any of the variety of methods described herein and methods described in PCT Application Publication No. WO 2019/113533A1, the content of which is incorporated herein by reference in its entirety. For example, the spatially-tagged cells can be encapsulated with beads comprising one or more nucleic acid molecules having a barcode (e.g., a cellular barcode) (e.g., an emulsion). The nucleic acid present on the bead can have a domain that hybridizes to a

domain on a nucleic acid present on the tagged cell (e.g., a domain on a nucleic acid that is attached to a cell-tagging agent), thus linking the spatial barcode of the cell to the cellular barcode of the bead. Once the spatial barcode of the cell and the cellular barcode of the bead are linked, analytes present in the cell can be analyzed using capture probes (e.g., capture probes present on the bead). This allows the nucleic acids produced (using these methods) from specific cells to be amplified and sequenced separately (e.g. within separate partitions or droplets).

In some embodiments, once a cell or cells in a biological sample is spatially tagged with a cell-tagging agent(s), spatial analysis of analytes present in the biological sample is performed in which the cells of the biological sample are not dissociated into single cells. In such embodiments, various methods of spatial analysis such as any of those provided herein can be employed. For example, once a cell or cells in a biological sample is spatially tagged with a cell-tagging agent(s), analytes in the cells can be captured and assayed. In some embodiments, cell-tagging agents include both a spatial barcode and a capture domain that can be used to capture analytes present in a cell. For example, cell-tagging agents that include both a spatial barcode and a capture domain can be introduced to cells of the biological sample in a way such that locations of the cell-tagging agents are known (or can be determined after introducing them to the cells). One non-limiting example of introducing cell-tagging agents to a biological sample is to provide the cell-tagging agents in an arrayed format (e.g., arrayed on a substrate such as any of the variety of substrates and arrays provided herein), where the positions of the cell-tagging agents on the array are known at the time of introduction (or can be determined after introduction). The cells can be permeabilized as necessary (e.g., using permeabilization agents and methods described herein), reagents for analyte analysis can be provided to the cells (e.g., a reverse transcriptase, a polymerase, nucleotides, etc., in the case where the analyte is a nucleic acid that binds to the capture probe), and the analytes can be assayed. In some embodiments, the assayed analytes (and/or copies thereof) can be released from the substrate and analyzed. In some embodiments, the assayed analytes (and/or copies thereof) are assayed in situ.

Introducing a Cell-Tagging Agent to the Surface of a Cell

Non-limiting examples of cell-tagging agents and systems that attach to the surface of a cell (e.g., thus introducing the cell-tagging agent and any barcode attached thereto to the exterior of the cell) that can be used in accordance with materials and methods provided herein for

spatially profiling an analyte or analytes in a biological sample include: lipid tagged primers / lipophilic-tagged moieties, positive or neutral oligo-conjugated polymers, antibody-tagged primers, streptavidin-conjugated oligonucleotides, dye-tagged oligonucleotides, click-chemistry, receptor-ligand systems, covalent binding systems via amine or thiol functionalities, and combinations thereof.

Lipid tagged primers / Lipophilic-tagged moieties

In some embodiments of any of the spatial profiling methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is coupled to a lipophilic molecule. In some embodiments, the lipophilic molecule enables the delivery of the molecule to the cell membrane or the nuclear membrane. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) coupled to a lipophilic molecule can associate with and/or insert into lipid membranes such as cell membranes and nuclear membranes. In some cases, the insertion can be reversible. In some cases, the association between the lipophilic molecule and the cell may be such that the cell retains the lipophilic molecule (e.g., and associated components, such as nucleic acid barcode molecules) during subsequent processing (e.g., partitioning, cell permeabilization, amplification, pooling, etc.). In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) coupled to a lipophilic molecule may enter into the intracellular space and/or a cell nucleus.

Non-limiting examples of lipophilic molecules that can be used in embodiments described herein include sterol lipids such as cholesterol, tocopherol, steryl, palmitate, lignoceric acid, and derivatives thereof. In some embodiments, the lipophilic molecules are neutral lipids that are conjugated to hydrophobic moieties (e.g., cholesterol, squalene, or fatty acids) (See Raouane et al. Bioconjugate Chem., 23(6):1091-1104 (2012) which is herein incorporated by reference in its entirety). In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) may be attached to the lipophilic moiety via a linker, such as a tetra-ethylene glycol (TEG) linker. Other exemplary linkers include, but are not limited to, Amino Linker C6, Amino Linker C12, Spacer C3, Spacer C6, Spacer C12, Spacer 9, and Spacer 18. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is indirectly coupled (e.g., via hybridization or ligand-ligand interactions, such as biotin-streptavidin) to a lipophilic molecule. Other lipophilic molecules that may be used

in accordance with methods provided herein include amphiphilic molecules wherein the headgroup (e.g., charge, aliphatic content, and/or aromatic content) and/or fatty acid chain length (e.g., C12, C14, C16, or C18) can be varied. For instance, fatty acid side chains (e.g., C12, C14, C16, or C18) can be coupled to glycerol or glycerol derivatives (e.g., 3-t-

5 butyldiphenylsilylglycerol), which can also comprise, e.g., a cationic head group. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) disclosed herein can then be coupled (either directly or indirectly) to these amphiphilic molecules. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) coupled to an amphiphilic molecule may associate with and/or insert into
10 a membrane (e.g., a cell, cell bead, or nuclear membrane). In some cases, an amphiphilic or lipophilic moiety may cross a cell membrane and provide a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to an internal region of a cell and/or cell bead.

In some embodiments, wherein the molecule (e.g., with a nucleic acid sequence) has an amino group within the molecule, the molecule (e.g., a nucleic acid molecule) having a barcode
15 (e.g., a spatial barcode) and an amino group can be coupled to an amine-reactive lipophilic molecule. For example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and an amino group can be conjugated to DSPE-PEG(2000)-cyanuric chloride (1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[cyanur(polyethylene glycol)-2000]).

20 In some embodiments, a cell tagging agent can attach to a surface of a cell through a combination of lipophilic and covalent attachment. For example, a cell tagging agent can include an oligonucleotide attached to a lipid to target the oligonucleotide to a cell membrane, and an amine group that can be covalently linked to a cell surface protein(s) via any number of chemistries described herein. In these embodiments, the lipid can increase the surface
25 concentration of the oligonucleotide and can promote the covalent reaction.

Positive or neutral oligo-conjugated polymers

In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a
30 glycol chitosan derivative. The glycol chitosan derivative (e.g., glycol chitosan-cholesterol) can serve as a hydrophobic anchor (see Wang et al. J. Mater. Chem. B., 30:6165 (2015), which is

herein incorporated by reference in its entirety). Non-limiting examples of chitosan derivatives that can be coupled to a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be found in Cheung et al., *Marine Drugs*, 13(8): 5156-5186 (2015), which is herein incorporated by reference in its entirety.

5 Antibody-tagged primers

In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an antibody or antigen binding fragment thereof in a manner that facilitates attachment of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to the surface
10 of a cell. In some embodiments, facilitating attachment to the cell surface facilitates introduction of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) into the cell. In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an antibody that is directed to an antigen that is present on the surface of a cell. In some embodiments, the molecule (e.g., a nucleic acid molecule) having a
15 barcode (e.g., a spatial barcode) can be coupled to an antibody that is directed to an antigen that is present on the surface of a plurality of cells (e.g., a plurality of cells in a biological sample). In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an antibody that is directed to an antigen that is present on the surface of all or substantially all the cells present in a biological sample. Any of the exemplary methods
20 described herein of attaching a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to another molecule (e.g., a cell-tagging agent) can be used.

Streptavidin-conjugated oligonucleotides

In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can attach to the surface
25 of a cell using biotin-streptavidin. In some embodiments, primary amines in the side chain of lysine residues of cell surface polypeptides are labelled with NHS-activated biotin reagents. For example, the N-terminus of a polypeptide can react with NHS-activated biotin reagents to form stable amide bonds. In some embodiments, cell-tagging agents include molecules (e.g., a nucleic acid molecule) having barcodes (e.g., a spatial barcode) conjugated to streptavidin. In some
30 cases, streptavidin can be conjugated to the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) using click chemistry (e.g., maleimide modification) as

described herein. In some embodiments, a cell containing NHS-activated biotin incorporated into lysine side chains of a cell surface protein forms a non-covalent bond with the streptavidin conjugated to the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) conjugated to streptavidin is itself part of a cell-tagging agent.

Dye-tagged oligonucleotides

In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is directly linked to a fluorescent tag. In some embodiments, the physical properties of the fluorescent tags (e.g., hydrophobic properties) can overcome the hydrophilic nature of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). For example, in some embodiments, wherein the molecule is a nucleic acid molecule, a fluorescent tag (e.g., BODIPY, Cy3, Atto 647N, and Rhodamine Red C2) can be coupled to a 5' end of the nucleic acid molecule having a barcode (e.g., a spatial barcode). In some embodiments, wherein the molecule is a nucleic acid molecule, any fluorescent tag having hydrophobic properties can be coupled to the nucleic acid molecule having a barcode (e.g., a spatial barcode) in a manner that overcomes the hydrophilic nature of the nucleic acid molecule. Non-limiting examples of fluorescent tags with hydrophobic properties include BODIPY, Cy3, Atto 647N, and Rhodamine Red C2.

Click-chemistry

In some embodiments of any of the spatial analysis methods described herein, molecules (e.g., a nucleic acid molecule) having barcodes (e.g., a spatial barcode) are coupled to click-chemistry moieties. As used herein, the term "click chemistry," generally refers to reactions that are modular, wide in scope, give high yields, generate only inoffensive byproducts, such as those that can be removed by nonchromatographic methods, and are stereospecific (but not necessarily enantioselective) (see, e.g., Angew. Chem. Int. Ed., 2001, 40(11):2004-2021, which is incorporated herein by reference in its entirety). In some cases, click chemistry can describe pairs of functional groups that can selectively react with each other in mild, aqueous conditions.

An example of a click chemistry reaction is the Huisgen 1,3-dipolar cycloaddition of an azide and an alkyne, i.e., copper-catalysed reaction of an azide with an alkyne to form the 5-membered heteroatom ring 1,2,3-triazole. The reaction is also known as a Cu(I)-Catalyzed Azide-Alkyne Cycloaddition (CuAAC), a Cu(I) click chemistry or a Cu⁺ click chemistry.

Catalysts for the click chemistry include, but are not limited to, Cu(I) salts, or Cu(I) salts made in situ by reducing Cu(II) reagents to Cu(I) reagents with a reducing reagent (Pharm Res. 2008, 25(10): 2216–2230, which is incorporated herein by reference in its entirety). Known Cu(II) reagents for the click chemistry can include, but are not limited to, the Cu(II)-(TBTA) complex and the Cu(II) (THPTA) complex. TBTA, which is tris-[(1-benzyl-1H-1,2,3-triazol-4-yl)methyl]amine, also known as tris-(benzyltriazolylmethyl)amine, can be a stabilizing ligand for Cu(I) salts. THPTA, which is tris-(hydroxypropyltriazolylmethyl)amine, is another example of a stabilizing agent for Cu(I). Other conditions can also be used to construct the 1,2,3-triazole ring from an azide and an alkyne using copper-free click chemistry, such as the Strain-promoted Azide- Alkyne Click chemistry reaction (SPAAC) (see, e.g., Chem. Commun., 2011, 47:6257-6259 and Nature, 2015, 519(7544):486-90, each of which is incorporated herein by reference in its entirety).

Receptor-ligand systems

In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a ligand, wherein the ligand is part of a receptor-ligand interaction on the surface of a cell. For example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a ligand that interacts selectively with a cell surface receptor thereby targeting the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a specific cell. Non-limiting examples of receptor-ligand systems that can be used include integrin receptor-ligand interactions, GPCR receptor-ligand interactions, RTK receptor-ligand interactions, and TLR-ligand interactions (see Juliano, Nucleic Acids Res., 44(14): 6518-6548 (2016), which is incorporated herein by reference in its entirety). Any of the methods described herein for attaching a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a ligand (e.g., any of the methods described herein relating to attaching a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to an antibody) can be used.

Covalent binding systems via amine or thiol functionalities

In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can incorporate reactive functional groups at sites within the molecule (e.g., with a nucleic acid sequence). In such cases,

the reactive functional groups can facilitate conjugation to ligands and/or surfaces. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can include thiol modifiers that are designed to react with a broad array of activated accepting groups (e.g., maleimide and gold microspheres). For example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) having thiol modifiers can interact with a maleimide-conjugated peptide thereby resulting in labelling of the peptide. In some embodiments, maleimide-conjugated peptides are present on the surface of a cell whereupon interaction with the thiol-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode), the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is coupled to the surface of the cell. Non-limiting examples of thiol modifiers include: 5' thiol modifier C6 S-S, 3' thiol modifier C3 S-S, dithiol, 3' thiol modifier oxa 6-S-S, and dithiol serinol.

In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can include amine modifiers, e.g., amine modifiers that are designed to attach to another molecule in the presence of an acylating agent. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can include amine modifiers that are designed to attach to a broad array of linkage groups (e.g., carbonyl amide, thiourea, sulfonamide, and carboxamide). For example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and an amine modifier can interact with a sulfonamide-conjugated peptide thereby resulting in labelling of the peptide. In some embodiments, sulfonamide-conjugated peptides are present on the surface of a cell whereupon interaction with the amine-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode), the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is coupled to the surface of the cell. Non-limiting example of amine modifiers include: DMS(O)MT-Amino-Modifier-C6, Amino-Modifier-C3-TFA, Amino-Modifier-C12, Amino-Modifier-C6-TFA, Amino-dT, Amino-Modifier-5, Amino-Modifier-C2-dT, Amino-Modifier-C6-dT, and 3'-Amino-Modifier-C7.

As another example, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can incorporate reactive functional groups at sites within the molecule (e.g., with a nucleic acid sequence) such as N-hydroxysuccinimide (NHS). In some embodiments, amines (e.g., amine-containing peptides) are present on the surface of a cell whereupon interaction with

the NHS-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode), the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is coupled to the surface of the cell. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) is reacted with a bifunctional NHS linker to form an NHS-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode).

In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a biocompatible anchor for cell membrane (BAM). For example, a BAM can include molecules that comprise an oleyl group and PEG. The oleyl group can facilitate anchoring the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a cell, and the PEG can increase water solubility. In some embodiments, oleyl-PEG-NHS can be coupled to a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) using NHS chemistry.

Azide-based systems

In some embodiments, wherein the molecule (e.g., with a nucleic acid sequence) incorporates reactive functional groups at sites within the molecule, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to an azide group on a cell surface. In some embodiments, the reactive functional group is an alkynyl group. In some embodiments, click chemistry as described herein can be used to attach the alkynyl-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to an azide group on the cell surface. An azide group can be attached to the cell surface through a variety of methods. For example, NHS chemistry can be used to attach an azide group to the cell surface. In some embodiments, N-azidoacetylmannosamine-tetraacylated (Ac4ManNAz), which contains an azide group, can react with sialic acid on the surface of a cell to attach azide to the cell surface. In some embodiments, azide is attached to the cell surface by bio-orthogonal expression of azide.

Lectin-based systems

In some embodiments of any of the spatial analysis methods described herein, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) can be coupled to a lectin that facilitates attachment of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a cell surface. Lectin can bind to glycans, e.g., glycans on the surface of cells. In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode

(e.g., a spatial barcode) has an incorporated reactive functional group such as an azide group. In some embodiments, the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and an azide group is reacted with a modified lectin, e.g., a lectin modified using NHS chemistry to introduce an azide reactive group. In some embodiments, a live cell is labelled with a lectin-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). In some embodiments, a fixed cell is labelled with a lectin-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). In some embodiments, a permeabilized cell is labelled with a lectin-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode). In some embodiments, organelles in the secretory pathway can be labelled with a lectin-modified molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode).

(b) Introducing a Cell-Tagging Agent to the Interior of a Cell

Non-limiting examples of cell-tagging agents and systems that penetrate and/or pass through the cell membrane (e.g., thus introducing the cell-tagging agent and any barcode attached thereto to the interior of the cell) that can be used in accordance with materials and methods provided herein for spatially profiling an analyte or analytes in a biological sample include: a cell-penetrating agent (e.g., a cell-penetrating peptide), a nanoparticle, a liposome, a polymersome, a peptide-based chemical vector, electroporation, sonoporation, lentiviral vectors, retroviral vectors, and combinations thereof.

FIG. 19 is a schematic showing an exemplary cell tagging method. Non-exhaustive examples of oligo delivery vehicles may include a cell penetrating peptide or a nanoparticle. Non-exhaustive examples of the delivery systems can include lipid-based polymeric and metallic nanoparticles or oligos that can be conjugated or encapsulated within the delivery system. The cell tag can be used in combination with a capture agent barcode domain and a cleavable or non-cleavable spatially barcoded capture probes for spatial and multiplexing applications.

Cell-Penetrating Agent

In some embodiments of any of the spatial profiling methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a cell-penetrating agent. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is coupled to a cell-penetrating agent, and the cell-penetrating agent allows the molecule

to interact with an analyte inside the cell. A “cell-penetrating agent” as used herein refers to an agent capable of facilitating the introduction of a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain into a cell of a biological sample (see, e.g., Lovatt et al. Nat Methods. 2014 Feb;11(2):190-6, which is incorporated herein by reference in its entirety). In some embodiments, a cell-penetrating agent is a cell-penetrating peptide. A “cell-penetrating peptide” as used herein refers to a peptide (e.g., a short peptide, e.g., a peptide not usually exceeding 30 residues) that has the capacity to cross cellular membranes.

In some embodiments of any of the spatial profiling methods described herein, a cell-penetrating peptide coupled to a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain can cross a cellular membrane using an energy dependent or an energy independent mechanism. For example, a cell-penetrating peptide can cross a cellular membrane through direct translocation through physical perturbation of the plasma membrane, endocytosis, adaptive translocation, pore-formation, electroporation-like permeabilization, and/or entry at microdomain boundaries. Non-limiting examples of a cell-penetrating peptide include: penetratin, tat peptide, pVEC, transportan, MPG, Pep-1, a polyarginine peptide, MAP, R6W3, (D-Arg)⁹, Cys(Npys)-(D-Arg)⁹, Anti-BetaGamma (MPS - Phosducin - like protein C terminus), Cys(Npys) antennapedia, Cys(Npys)-(Arg)⁹, Cys(Npys)-TAT (47-57), HIV-1 Tat (48-60), KALA, mastoparan, penetratin-Arg, pep-1-cysteamine, TAT(47-57)GGG-Cys(Npys), Tat-NR2Bct, transdermal peptide, SynB1, SynB3, PTD-4, PTD-5, FHV Coat-(35-49), BMV Gag-(7-25), HTLV-II Rex-(4-16), R9-tat, SBP, FBP, MPG, MPG(ΔNLS), Pep-2, MTS, plsl, and a polylysine peptide (see, e.g., Bechara et al. FEBS Lett. 2013 Jun 19;587(12):1693-702, which is incorporated by reference herein in its entirety).

Nanoparticles

In some embodiments of any of the spatial profiling methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by an inorganic particle (e.g., a nanoparticle). In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is coupled to an inorganic particle (e.g., a nanoparticle), and the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain uses the nanoparticle to get access to analytes inside the cell. Non-limiting examples of nanoparticles that can be used in embodiments herein to deliver a molecule (e.g., a nucleic acid

molecule) having a barcode (e.g., a spatial barcode) and a capture domain into a cell and/or cell bead include inorganic nanoparticles prepared from metals, (e.g., iron, gold, and silver), inorganic salts, and ceramics (e.g., phosphate or carbonate salts of calcium, magnesium, or silicon). The surface of a nanoparticle can be coated to facilitate binding of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain, or the surface can be chemically modified to facilitate attachment of the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain. Magnetic nanoparticles (e.g., supermagnetic iron oxide), fullerenes (e.g., soluble carbon molecules), carbon nanotubes (e.g., cylindrical fullerenes), quantum dots and supramolecular systems can also be used.

Liposomes

In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a liposome. Various types of lipids, including cationic lipids, can be used in liposome delivery. In some cases, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is delivered to a cell via a lipid nano-emulsion. A lipid emulsion refers to a dispersion of one immiscible liquid in another stabilized by emulsifying agent. Labeling cells can comprise use of a solid lipid nanoparticle.

Polymersomes

In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a polymersome. In some embodiments, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is contained in the polymersome, and the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain uses the polymersome to get access to analytes inside the cell. A “polymersome” as referred to herein is an artificial vesicle. For example, a polymersome can be a vesicle similar to a liposome, but the membrane comprises amphiphilic synthetic block copolymers (see, e.g., Rideau et al. Chem. Soc. Rev., 2018, 47, 8572-8610, which is incorporated by reference herein in its entirety). In some embodiments, polymersomes comprise di-(AB) or tri-block copolymers (e.g., ABA or ABC), where A and C

are a hydrophilic block and B is a hydrophobic block. In some embodiments, a polymersome comprises poly(butadiene)-b-poly(ethylene oxide), poly(ethyl ethylene)-b-poly(ethylene oxide), polystyrene-b-poly(ethylene oxide), poly(2-vinylpyridine)-b-poly(ethylene oxide), polydimethylsiloxane-b-poly(ethylene oxide), polydimethylsiloxane-g-poly(ethylene oxide),
 5 polycaprolactone-b-poly(ethylene oxide), polyisobutylene-b-poly(ethylene oxide), polystyrene-b-polyacrylic acid, polydimethylsiloxane-b-poly-2-methyl-2-oxazoline, or a combination thereof (wherein b=block and g=grafted).

Peptide-Based Chemical Vectors

In some embodiments of any of the spatial analysis methods described herein, capture of
 10 a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a peptide-based chemical vector, e.g., a cationic peptide-based chemical vector. Cationic peptides can be rich in basic residues like lysine and/or arginine. In some embodiments of any of the spatial analysis methods described herein, capture
 15 of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by a polymer-based chemical vector. Cationic polymers, when mixed with a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain, can form nanosized complexes called polyplexes. Polymer based vectors can comprise natural proteins, peptides and/or polysaccharides. Polymer based vectors can comprise synthetic polymers. In some embodiments, a polymer-based vector
 20 comprises polyethylenimine (PEI). PEI can condense DNA into positively-charged particles, which bind to anionic cell surface residues and are brought into the cell via endocytosis. In some embodiments, a polymer-based chemical vector comprises poly(L)-lysine (PLL), poly (DL-lactic acid) (PLA), poly (DL-lactide-co-glycoside) (PLGA), polyornithine, polyarginine, histones, protamines, or a combination thereof. Polymer-based vectors can comprise a mixture of
 25 polymers, for example, PEG and PLL. Other non-limiting examples of polymers include dendrimers, chitosans, synthetic amino derivatives of dextran, and cationic acrylic polymers.

Electroporation

In some embodiments of any of the spatial analysis methods described herein, capture of
 a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial
 30 barcode) and a capture domain is facilitated by electroporation. With electroporation, a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial

barcode) and a capture domain can enter a cell through one or more pores in the cellular membrane formed by applied electricity. The pore of the membrane can be reversible based on the applied field strength and pulse duration.

Sonoporation

5 In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by sonoporation. Cell membranes can be temporarily permeabilized using sound waves, allowing cellular uptake of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain.

Lentiviral vectors and retroviral vectors

10 In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by vectors. For example, a vector as described herein can be an expression vector where the expression vector includes a promoter sequence
15 operably linked to the sequence encoding the molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain. Non-limiting examples of vectors include plasmids, transposons, cosmids, and viral vectors (e.g., any adenoviral vectors (e.g., pSV or pCMV vectors), adeno-associated virus (AAV) vectors, lentivirus vectors, and retroviral
20 vectors), and any Gateway® vectors. A vector can, for example, include sufficient cis-acting elements for expression where other elements for expression can be supplied by the host mammalian cell or in an in vitro expression system. Skilled practitioners will be capable of selecting suitable vectors and mammalian cells for introducing any of spatial profiling reagents described herein.

Other Methods and Cell-Tagging Agents for Intracellular Introduction of a Molecule

25 In some embodiments of any of the spatial analysis methods described herein, capture of a biological analyte by a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain is facilitated by the use of a needle, for example for injection (e.g., microinjection), particle bombardment, photoporation, magnetofection, and/or
30 hydroporation. For example, with particle bombardment, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain can be coated with

heavy metal particles and delivered to a cell at a high speed. In photoporation, a transient pore in a cell membrane can be generated using a laser pulse, allowing cellular uptake of a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain. In magnetofection, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial
5 barcode) and a capture domain can be coupled to a magnetic particle (e.g., magnetic nanoparticle, nanowires, etc.) and localized to a target cell via an applied magnetic field. In hydroporation, a molecule (e.g., a nucleic acid molecule) having a barcode (e.g., a spatial barcode) and a capture domain can be delivered to a cell and/or cell bead via hydrodynamic pressure.

10 (c) Methods for separating sample into single cells or cell groups

Some embodiments of any of the methods described herein can include separating a biological sample into single cells, cell groups, types of cells, or a region or regions of interest. For example, a biological sample can be separated into single cells, cell groups, types of cells, or a region or regions of interest before being contacted with one or more capture probes. In other
15 examples, a biological sample is first contacted with one or more capture probes, and then separated into single cells, cell groups, types of cells, or a region or regions of interest.

In some embodiments, a biological sample can be separated into chunks using pixelation. Pixelation can include the steps of providing a biological sample, and punching out one or more portions of the biological sample. The punched out portions of the biological sample can then be
20 used to perform any of the methods described herein. In some embodiments, the punched-out portions of the biological sample can be in a random pattern or a designed pattern. In some embodiments, the punched-out portions of the biological sample can be focused on a region of interest or a subcellular structure in the biological sample.

FIG. 20A is a workflow schematic illustrating exemplary, non-limiting, non-exhaustive steps for “pixelating” a sample, wherein the sample is cut, stamped, microdissected, or
25 transferred by hollow-needle or microneedle, moving a small portion of the sample into an individual partition or well.

FIG. 20B is a schematic depicting multi-needle pixelation, wherein an array of needles punched through a sample on a scaffold and into nanowells containing gel beads and reagents
30 below. Once the needle is in the nanowell, the cell(s) are ejected.

In some embodiments, a biological sample is divided into chunks before performance of any of the spatial analysis methods described herein. In some embodiments, the methods can include spatial barcoding of FFPE “chunks” via barcodes applied in spatially well-defined pattern (like in DNA microarray printing). The DNA barcode is either long so that it will not
5 diffuse out in subsequent steps or is covalently applied to the FFPE sample. To enable barcodes to get embedded into an FFPE slide, the wax can be heated, barcodes can be added to the slide before cooling, and then the chunks can be cut. The cutting can be done in various ways such as using laser microdissection, or via mechanical or acoustic means. Other alternates are to embed some fluorophores/Qdots, etc. to preserve spatial information into the sample. The barcoding at
10 this step enables massively parallel random encapsulation of chunks while retaining local spatial information (e.g., tumor vs normal cells).

In some embodiments, a biological sample can be divided or portioned using laser capture microdissection (e.g., highly-multiplexed laser capture microdissection).

(d) Release and Amplification of Analytes

15 In some embodiments, lysis reagents can be added to the sample to facilitate the release of analyte(s) from a sample. Examples of lysis agents include, but are not limited to, bioactive reagents such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other commercially available lysis enzymes.

20 Other lysis agents can additionally or alternatively be co-partitioned with the biological sample to cause the release of the sample’s contents into the partitions. In some embodiments, surfactant-based lysis solutions can be used to lyse cells, although these can be less desirable for emulsion-based systems where the surfactants can interfere with stable emulsions. Lysis solutions can include ionic surfactants such as, for example, sarcosyl and sodium dodecyl sulfate (SDS). Electroporation, thermal, acoustic or mechanical cellular disruption can also be used in
25 certain embodiments, e.g., non-emulsion based partitioning such as encapsulation of biological materials that can be in addition to or in place of droplet partitioning, where any pore size of the encapsulate is sufficiently small to retain nucleic acid fragments of a given size, following cellular disruption.

30 In addition to the permeabilization agents, other reagents can also be added to interact with the biological sample, including, for example, DNase and RNase inactivating agents or

inhibitors, such as proteinase K, chelating agents, such as EDTA, and other reagents to allow for subsequent processing of analytes from the sample.

Further reagents that can be added to a sample, include, for example, endonucleases to fragment DNA, DNA polymerase enzymes, and dNTPs used to amplify nucleic acids. Other
5 enzymes that can also be added to the sample include, but are not limited to, polymerase, transposase, ligase, proteinase K, and DNase, etc. Additional reagents can also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers, and switch oligonucleotides. In some embodiments, template switching can be used to increase the length of a cDNA, e.g., by appending a predefined nucleic acid sequence to the cDNA.

10 If a tissue sample is not permeabilized sufficiently, the amount of analyte captured on the substrate can be too low to enable adequate analysis. Conversely, if the tissue sample is too permeable, the analyte can diffuse away from its origin in the tissue sample, such that the relative spatial relationship of the analytes within the tissue sample is lost. Hence, a balance between permeabilizing the tissue sample enough to obtain good signal intensity while still maintaining
15 the spatial resolution of the analyte distribution in the tissue sample is desired.

In some embodiments, where the biological sample includes live cells, permeabilization conditions can be modified so that the live cells experience only brief permeabilization (e.g., through short repetitive bursts of electric field application), thereby allowing one or more analytes to migrate from the live cells to the substrate while retaining cellular viability.

20 In some embodiments, after contacting a biological sample with a substrate that include capture probes, a removal step is performed to remove all or a portion of the biological sample from the substrate. In some embodiments, the removal step includes enzymatic or chemical degradation of the permeabilized cells of the biological sample. For example, the removal step can include treating the biological samples with an enzyme (e.g., proteinase K) to remove at least a portion
25 of the biological sample from the first substrates. In some embodiments, the removal step can include ablation of the tissue (e.g., laser ablation).

In some embodiments, where RNA is captured from cells in a sample, one or more RNA species of interest can be selectively enriched. For example, one or more species of RNA of interest can be selected by addition of one or more oligonucleotides. One or more species of
30 RNA can be selectively down-selected (e.g., removed) using any of a variety of methods. For example, probes can be administered to a sample that selectively hybridize to ribosomal RNA

(rRNA), thereby reducing the pool and concentration of rRNA in the sample. Subsequent application of the capture probes to the sample can result in improved RNA capture due to the reduction in non-specific RNA present in the sample. In some embodiments, the additional oligonucleotide is a sequence used for priming a reaction by a polymerase. For example, one or more primer sequences with sequence complementarity to one or more RNAs of interest, can be used to amplify the one or more RNAs of interest, thereby selectively enriching these RNAs. In some embodiments, an oligonucleotide with sequence complementarity to the complementary strand of captured RNA (e.g., cDNA) can bind to the cDNA. In one non-limiting example, biotinylated oligonucleotides with sequence complementary to one or more cDNA of interest binds to the cDNA and can be selected using biotinylation-streptavidin affinity in any number of methods known to the field (e.g., streptavidin beads).

Nucleic acid analytes can be amplified using a polymerase chain reaction (e.g., digital PCR, quantitative PCR, or real time PCR), or isothermal amplification, or any of the nucleic acid amplification or extension reactions described herein.

(e) Partitioning

As discussed above, in some embodiments, the sample can optionally be separated into single cells, cell groups, or other fragments/pieces that are smaller than the original, unfragmented sample. Each of these smaller portions of the sample can be analyzed to obtain spatially-resolved analyte information from the sample. Non-limiting partitioning methods are described herein.

For samples that have been separated into smaller fragments – and particularly, for samples that have been disaggregated, dissociated, or otherwise separated into individual cells – one method for analyzing the fragments involves partitioning the fragments into individual partitions (e.g., fluid droplets), and then analyzing the contents of the partitions. In general, each partition maintains separation of its own contents from the contents of other partitions. For example, the partition can be a droplet in an emulsion.

In addition to analytes, a partition can include additional components, and in particular, one or more beads. A partition can include a single gel bead, a single cell bead, or both a single cell bead and single gel bead.

A partition can also include one or more reagents. Unique identifiers, such as barcodes, can be injected into the droplets previous to, subsequent to, or concurrently with droplet generation, such as via a microcapsule (e.g., bead). Microfluidic channel networks (e.g., on a chip) can be utilized to generate partitions. Alternative mechanisms can also be employed in the partitioning of individual biological particles, including porous membranes through which aqueous mixtures of cells are extruded into non-aqueous fluids.

The partitions can be flowable within fluid streams. The partitions can include, for example, micro-vesicles that have an outer barrier surrounding an inner fluid center or core. In some cases, the partitions can include a porous matrix that is capable of entraining and/or retaining materials within its matrix. The partitions can be droplets of a first phase within a second phase, wherein the first and second phases are immiscible. For example, the partitions can be droplets of aqueous fluid within a non-aqueous continuous phase (e.g., oil phase). In another example, the partitions can be droplets of a non-aqueous fluid within an aqueous phase. In some examples, the partitions can be provided in a water-in-oil emulsion or oil-in-water emulsion. A variety of different vessels are described in, for example, U.S. Patent Application Publication No. 2014/0155295, the entire contents of which are incorporated herein by reference. Emulsion systems for creating stable droplets in non-aqueous or oil continuous phases are described, for example, in U.S. Patent Application Publication No. 2010/0105112, the entire contents of which are incorporated herein by reference.

For droplets in an emulsion, allocating individual particles to discrete partitions can be accomplished, for example, by introducing a flowing stream of particles in an aqueous fluid into a flowing stream of a non-aqueous fluid, such that droplets are generated at the junction of the two streams. Fluid properties (e.g., fluid flow rates, fluid viscosities, etc.), particle properties (e.g., volume fraction, particle size, particle concentration, etc.), microfluidic architectures (e.g., channel geometry, etc.), and other parameters can be adjusted to control the occupancy of the resulting partitions (e.g., number of analytes per partition, number of beads per partition, etc.) For example, partition occupancy can be controlled by providing the aqueous stream at a certain concentration and/or flow rate of analytes.

To generate single analyte partitions, the relative flow rates of the immiscible fluids can be selected such that, on average, the partitions can contain less than one analyte per partition to ensure that those partitions that are occupied are primarily singly occupied. In some cases,

partitions among a plurality of partitions can contain at most one analyte. In some embodiments, the various parameters (e.g., fluid properties, particle properties, microfluidic architectures, etc.) can be selected or adjusted such that a majority of partitions are occupied, for example, allowing for only a small percentage of unoccupied partitions. The flows and channel architectures can be controlled as to ensure a given number of singly occupied partitions, less than a certain level of unoccupied partitions and/or less than a certain level of multiply occupied partitions.

The channel segments described herein can be coupled to any of a variety of different fluid sources or receiving components, including reservoirs, tubing, manifolds, or fluidic components of other systems. As will be appreciated, the microfluidic channel structure can have a variety of geometries. For example, a microfluidic channel structure can have one or more than one channel junction. As another example, a microfluidic channel structure can have 2, 3, 4, or 5 channel segments each carrying particles that meet at a channel junction. Fluid can be directed to flow along one or more channels or reservoirs via one or more fluid flow units. A fluid flow unit can include compressors (e.g., providing positive pressure), pumps (e.g., providing negative pressure), actuators, and the like to control flow of the fluid. Fluid can also or otherwise be controlled via applied pressure differentials, centrifugal force, electrokinetic pumping, vacuum, capillary, and/or gravity flow.

A partition can include one or more unique identifiers, such as barcodes. Barcodes can be previously, subsequently, or concurrently delivered to the partitions that hold the compartmentalized or partitioned biological particle. For example, barcodes can be injected into droplets previous to, subsequent to, or concurrently with droplet generation. The delivery of the barcodes to a particular partition allows for the later attribution of the characteristics of the individual biological particle to the particular partition. Barcodes can be delivered, for example on a nucleic acid molecule (e.g., an oligonucleotide), to a partition via any suitable mechanism. Barcoded nucleic acid molecules can be delivered to a partition via a microcapsule. A microcapsule, in some instances, can include a bead.

In some embodiments, barcoded nucleic acid molecules can be initially associated with the microcapsule and then released from the microcapsule. Release of the barcoded nucleic acid molecules can be passive (e.g., by diffusion out of the microcapsule). In addition or alternatively, release from the microcapsule can be upon application of a stimulus which allows the barcoded nucleic acid nucleic acid molecules to dissociate or to be released from the microcapsule. Such

stimulus can disrupt the microcapsule, an interaction that couples the barcoded nucleic acid molecules to or within the microcapsule, or both. Such stimulus can include, for example, a thermal stimulus, photo-stimulus, chemical stimulus (e.g., change in pH or use of a reducing agent(s)), a mechanical stimulus, a radiation stimulus; a biological stimulus (e.g., enzyme), or
5 any combination thereof.

In some embodiments, one more barcodes (e.g., spatial barcodes, UMIs, or a combination thereof) can be introduced into a partition as part of the analyte. As described previously, barcodes can be bound to the analyte directly, or can form part of a capture probe or analyte capture agent that is hybridized to, conjugated to, or otherwise associated with an analyte, such
10 that when the analyte is introduced into the partition, the barcode(s) are introduced as well.

FIG. 21 depicts an exemplary workflow, where a sample is contacted with a spatially-barcoded capture probe array and the sample is fixed, stained, and imaged **2101**, as described elsewhere herein. The capture probes can be cleaved from the array **2102** using any method as described herein. The capture probes can diffuse toward the cells by either passive or active
15 migration as described elsewhere herein. The capture probes may then be introduced to the sample **2103** as described elsewhere herein, wherein the capture probe is able to gain entry into the cell in the absence of cell permeabilization, using one of the cell penetrating peptides or lipid delivery systems described herein. The sample can then be optionally imaged in order to confirm probe uptake, via a reporter molecule incorporated within the capture probe **2104**. The sample
20 can then be separated from the array and undergo dissociation **2105**, wherein the sample is separated into single cells or small groups of cells. Once the sample is dissociated, the single cells can be introduced to an oil-in water droplet **2106**, wherein a single cell is combined with reagents within the droplet and processed so that the spatial barcode that penetrated the cell labels the contents of that cell within the droplet. Other cells undergo separately partitioned
25 reactions concurrently. The contents of the droplet is then sequenced **2107** in order to associate a particular cell or cells with a particular spatial location within the sample **2108**.

As described above, **FIG. 16** shows an example of a microfluidic channel structure for partitioning individual analytes (e.g., cells) into discrete partitions. **FIGs. 17A** and **17C** also show other examples of microfluidic channel structures that can be used for delivering beads to
30 droplets.

A variety of different beads can be incorporated into partitions as described above. In some embodiments, for example, non-barcoded beads can be incorporated into the partitions. For example, where the biological particle (e.g., a cell) that is incorporated into the partitions carries one or more barcodes (e.g., spatial barcode(s), UMI(s), and combinations thereof), the bead can
5 be a non-barcoded bead.

In some embodiments, a barcode carrying bead can be incorporated into partitions. For example, a nucleic acid molecule, such as an oligonucleotide, can be coupled to a bead by a releasable linkage, such as, for example, a disulfide linker. The same bead can be coupled (e.g., via releasable linkage) to one or more other nucleic acid molecules. The nucleic acid molecule
10 can be or include a barcode. As noted elsewhere herein, the structure of the barcode can include a number of sequence elements.

The nucleic acid molecule can include a functional domain that can be used in subsequent processing. For example, the functional domain can include one or more of a sequencer specific flow cell attachment sequence (e.g., a P5 sequence for Illumina® sequencing systems) and a
15 sequencing primer sequence (e.g., a R1 primer for Illumina® sequencing systems). The nucleic acid molecule can include a barcode sequence for use in barcoding the sample (e.g., DNA, RNA, protein, etc.). In some cases, the barcode sequence can be bead-specific such that the barcode sequence is common to all nucleic acid molecules coupled to the same bead. Alternatively or in
20 addition, the barcode sequence can be partition-specific such that the barcode sequence is common to all nucleic acid molecules coupled to one or more beads that are partitioned into the same partition. The nucleic acid molecule can include a specific priming sequence, such as an mRNA specific priming sequence (e.g., poly (T) sequence), a targeted priming sequence, and/or a random priming sequence. The nucleic acid molecule can include an anchoring sequence to ensure that the specific priming sequence hybridizes at the sequence end (e.g., of the mRNA).
25 For example, the anchoring sequence can include a random short sequence of nucleotides, such as a 1-mer, 2-mer, 3-mer or longer sequence, which can ensure that a poly(T) segment is more likely to hybridize at the sequence end of the poly(A) tail of the mRNA.

The nucleic acid molecule can include a unique molecular identifying sequence (e.g., unique molecular identifier (UMI)). In some embodiments, the unique molecular identifying
30 sequence can include from about 5 to about 8 nucleotides. Alternatively, the unique molecular identifying sequence can include less than about 5 or more than about 8 nucleotides. The unique

molecular identifying sequence can be a unique sequence that varies across individual nucleic acid molecules coupled to a single bead.

In some embodiments, the unique molecular identifying sequence can be a random sequence (e.g., such as a random N-mer sequence). For example, the UMI can provide a unique
5 identifier of the starting mRNA molecule that was captured, in order to allow quantitation of the number of original expressed RNA.

In general, an individual bead can be coupled to any number of individual nucleic acid molecules, for example, from one to tens to hundreds of thousands or even millions of individual nucleic acid molecules. The respective barcodes for the individual nucleic acid molecules can
10 include both common sequence segments or relatively common sequence segments and variable or unique sequence segments between different individual nucleic acid molecules coupled to the same bead.

Within any given partition, all of the cDNA transcripts of the individual mRNA molecules can include a common barcode sequence segment. However, the transcripts made
15 from the different mRNA molecules within a given partition can vary at the unique molecular identifying sequence segment (e.g., UMI segment). Beneficially, even following any subsequent amplification of the contents of a given partition, the number of different UMIs can be indicative of the quantity of mRNA originating from a given partition. As noted above, the transcripts can be amplified, cleaned up and sequenced to identify the sequence of the cDNA transcript of the
20 mRNA, as well as to sequence the barcode segment and the UMI segment. While a poly(T) primer sequence is described, other targeted or random priming sequences can also be used in priming the reverse transcription reaction. Likewise, although described as releasing the barcoded oligonucleotides into the partition, in some cases, the nucleic acid molecules bound to the bead can be used to hybridize and capture the mRNA on the solid phase of the bead, for
25 example, in order to facilitate the separation of the RNA from other cell contents.

In some embodiments, precursors that include a functional group that is reactive or capable of being activated such that it becomes reactive can be polymerized with other precursors to generate gel beads that include the activated or activatable functional group. The functional group can then be used to attach additional species (e.g., disulfide linkers, primers,
30 other oligonucleotides, etc.) to the gel beads. For example, some precursors featuring a carboxylic acid (COOH) group can co-polymerize with other precursors to form a bead that also

includes a COOH functional group. In some cases, acrylic acid (a species comprising free COOH groups), acrylamide, and bis(acryloyl)cystamine can be co-polymerized together to generate a bead with free COOH groups. The COOH groups of the bead can be activated (e.g., via 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and N-Hydroxysuccinimide (NHS) or 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)) such that they are reactive (e.g., reactive to amine functional groups where EDC/NHS or DMTMM are used for activation). The activated COOH groups can then react with an appropriate species (e.g., a species comprising an amine functional group where the carboxylic acid groups are activated to be reactive with an amine functional group) comprising a moiety to be linked to the bead.

In some embodiments, a degradable bead can be introduced into a partition, such that the bead degrades within the partition and any associated species (e.g., oligonucleotides) are released within the droplet when the appropriate stimulus is applied. The free species (e.g., oligonucleotides, nucleic acid molecules) can interact with other reagents contained in the partition. For example, a polyacrylamide bead featuring cystamine and linked, via a disulfide bond, to a barcode sequence, can be combined with a reducing agent within a droplet of a water-in-oil emulsion. Within the droplet, the reducing agent can break the various disulfide bonds, resulting in bead degradation and release of the barcode sequence into the aqueous, inner environment of the droplet. In another example, heating of a droplet with a bead-bound barcode sequence in basic solution can also result in bead degradation and release of the attached barcode sequence into the aqueous, inner environment of the droplet.

Any suitable number of species (e.g., primer, barcoded oligonucleotide) can be associated with a bead such that, upon release from the bead, the species (e.g., primer, e.g., barcoded oligonucleotide) are present in the partition at a pre-defined concentration. Such pre-defined concentration can be selected to facilitate certain reactions for generating a sequencing library, e.g., amplification, within the partition. In some cases, the pre-defined concentration of the primer can be limited by the process of producing nucleic acid molecule (e.g., oligonucleotide) bearing beads.

A degradable bead can include one or more species with a labile bond such that, when the bead/species is exposed to the appropriate stimulus, the bond is broken and the bead degrades.

The labile bond can be a chemical bond (e.g., covalent bond, ionic bond) or can be another type of physical interaction (e.g., van der Waals interactions, dipole-dipole interactions, etc.). In some

embodiments, a crosslinker used to generate a bead can include a labile bond. Upon exposure to the appropriate conditions, the labile bond can be broken and the bead degraded. For example, upon exposure of a polyacrylamide gel bead that includes cystamine crosslinkers to a reducing agent, the disulfide bonds of the cystamine can be broken and the bead degraded.

- 5 A degradable bead can be useful in more quickly releasing an attached species (e.g., a nucleic acid molecule, a barcode sequence, a primer, etc.) from the bead when the appropriate stimulus is applied to the bead as compared to a bead that does not degrade. For example, for a species bound to an inner surface of a porous bead or in the case of an encapsulated species, the species can have greater mobility and accessibility to other species in solution upon degradation of the
- 10 bead. In some embodiments, a species can also be attached to a degradable bead via a degradable linker (e.g., disulfide linker). The degradable linker can respond to the same stimuli as the degradable bead or the two degradable species can respond to different stimuli. For example, a barcode sequence can be attached, via a disulfide bond, to a polyacrylamide bead comprising cystamine. Upon exposure of the barcoded-bead to a reducing agent, the bead degrades and the
- 15 barcode sequence is released upon breakage of both the disulfide linkage between the barcode sequence and the bead and the disulfide linkages of the cystamine in the bead.

As will be appreciated from the above description, while referred to as degradation of a bead, in many embodiments, degradation can refer to the disassociation of a bound or entrained species from a bead, both with and without structurally degrading the physical bead itself. For example,

20 entrained species can be released from beads through osmotic pressure differences due to, for example, changing chemical environments. By way of example, alteration of bead pore sizes due to osmotic pressure differences can generally occur without structural degradation of the bead itself. In some cases, an increase in pore size due to osmotic swelling of a bead can permit the release of entrained species within the bead. In some embodiments, osmotic shrinking of a bead

25 can cause a bead to better retain an entrained species due to pore size contraction.

Numerous chemical triggers can be used to trigger the degradation of beads within partitions. Examples of these chemical changes can include, but are not limited to pH-mediated changes to the integrity of a component within the bead, degradation of a component of a bead via cleavage of cross-linked bonds, and depolymerization of a component of a bead.

- 30 In some embodiments, a bead can be formed from materials that include degradable chemical cross-linkers, such as BAC or cystamine. Degradation of such degradable cross-linkers can be

accomplished through a number of mechanisms. In some examples, a bead can be contacted with a chemical degrading agent that can induce oxidation, reduction or other chemical changes. For example, a chemical degrading agent can be a reducing agent, such as dithiothreitol (DTT). Additional examples of reducing agents can include β -mercaptoethanol, (2S)-2-amino-1,4-dimercaptobutane (dithiobutylamine or DTBA), tris(2-carboxyethyl) phosphine (TCEP), or combinations thereof. A reducing agent can degrade the disulfide bonds formed between gel precursors forming the bead, and thus, degrade the bead.

In certain embodiments, a change in pH of a solution, such as an increase in pH, can trigger degradation of a bead. In other embodiments, exposure to an aqueous solution, such as water, can trigger hydrolytic degradation, and thus degradation of the bead. In some cases, any combination of stimuli can trigger degradation of a bead. For example, a change in pH can enable a chemical agent (e.g., DTT) to become an effective reducing agent.

Beads can also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety of changes to a bead. For example, heat can cause a solid bead to liquefy. A change in heat can cause melting of a bead such that a portion of the bead degrades. In other cases, heat can increase the internal pressure of the bead components such that the bead ruptures or explodes. Heat can also act upon heat-sensitive polymers used as materials to construct beads.

In addition to beads and analytes, partitions that are formed can include a variety of different reagents and species. For example, when lysis reagents are present within the partitions, the lysis reagents can facilitate the release of analytes within the partition. Examples of lysis agents include bioactive reagents, such as lysis enzymes that are used for lysis of different cell types, e.g., gram positive or negative bacteria, plants, yeast, mammalian, etc., such as lysozymes, achromopeptidase, lysostaphin, labiase, kitalase, lyticase, and a variety of other lysis enzymes available from, e.g., Sigma-Aldrich, Inc. (St. Louis, MO), as well as other commercially available lysis enzymes. Other lysis agents can additionally or alternatively be co-partitioned to cause the release analytes into the partitions. For example, in some cases, surfactant-based lysis solutions can be used to lyse cells, although these can be less desirable for emulsion based systems where the surfactants can interfere with stable emulsions. In some embodiments, lysis solutions can include non-ionic surfactants such as, for example, TritonX-100 and Tween 20. In some embodiments, lysis solutions can include ionic surfactants such as, for example, sarcosyl

and sodium dodecyl sulfate (SDS). Electroporation, thermal, acoustic or mechanical cellular disruption can also be used in certain embodiments, e.g., non-emulsion based partitioning such as encapsulation of analytes that can be in addition to or in place of droplet partitioning, where any pore size of the encapsulate is sufficiently small to retain nucleic acid fragments of a given size, following cellular disruption.

Examples of other species that can be co-partitioned with analytes in the partitions include, but are not limited to, DNase and RNase inactivating agents or inhibitors, such as proteinase K, chelating agents, such as EDTA, and other reagents employed in removing or otherwise reducing negative activity or impact of different cell lysate components on subsequent processing of nucleic acids. Additional reagents can also be co-partitioned, including endonucleases to fragment DNA, DNA polymerase enzymes and dNTPs used to amplify nucleic acid fragments and to attach the barcode molecular tags to the amplified fragments.

Additional reagents can also include reverse transcriptase enzymes, including enzymes with terminal transferase activity, primers and oligonucleotides, and switch oligonucleotides (also referred to herein as “switch oligos” or “template switching oligonucleotides”) which can be used for template switching. In some embodiments, template switching can be used to increase the length of a cDNA. Template switching can be used to append a predefined nucleic acid sequence to the cDNA. In an example of template switching, cDNA can be generated from reverse transcription of a template, e.g., cellular mRNA, where a reverse transcriptase with terminal transferase activity can add additional nucleotides, e.g., poly(C), to the cDNA in a template independent manner. Switch oligos can include sequences complementary to the additional nucleotides, e.g., poly(G). The additional nucleotides (e.g., poly(C)) on the cDNA can hybridize to the additional nucleotides (e.g., poly(G)) on the switch oligo, whereby the switch oligo can be used by the reverse transcriptase as template to further extend the cDNA.

Template switching oligonucleotides can include a hybridization region and a template region. The hybridization region can include any sequence capable of hybridizing to the target. In some cases, the hybridization region includes a series of G bases to complement the overhanging C bases at the 3' end of a cDNA molecule. The series of G bases can include 1 G base, 2 G bases, 3 G bases, 4 G bases, 5 G bases or more than 5 G bases. The template sequence can include any sequence to be incorporated into the cDNA. In some cases, the template region includes at least 1 (e.g., at least 2, 3, 4, 5 or more) tag sequences and/or functional sequences. Switch oligos can

include deoxyribonucleic acids; ribonucleic acids; modified nucleic acids including 2-Aminopurine, 2,6-Diaminopurine (2-Amino-dA), inverted dT, 5-Methyl dC, 2'-deoxyInosine, Super T (5-hydroxybutynl-2'-deoxyuridine), Super G (8-aza-7-deazaguanosine), locked nucleic acids (LNAs), unlocked nucleic acids (UNAs, e.g., UNA-A, UNA-U, UNA-C, UNA-G), Iso-dG, Iso-dC, 2' Fluoro bases (e.g., Fluoro C, Fluoro U, Fluoro A, and Fluoro G), and combinations of the foregoing.

In some embodiments, beads that are partitioned with the analyte can include different types of oligonucleotides bound to the bead, where the different types of oligonucleotides bind to different types of analytes. For example, a bead can include one or more first oligonucleotides (which can be capture probes, for example) that can bind or hybridize to a first type of analyte, such as mRNA for example, and one or more second oligonucleotides (which can be capture probes, for example) that can bind or hybridize to a second type of analyte, such as gDNA for example. Partitions can also include lysis agents that aid in releasing nucleic acids from the co-partitioned cell, and can also include an agent (e.g., a reducing agent) that can degrade the bead and/or break covalent linkages between the oligonucleotides and the bead, releasing the oligonucleotides into the partition. The released barcoded oligonucleotides (which can also be barcoded) can hybridize with mRNA released from the cell and also with gDNA released from the cell.

Barcoded constructs thus formed from hybridization can include a first type of construct that includes a sequence corresponding to an original barcode sequence from the bead and a sequence corresponding to a transcript from the cell, and a second type of construct that includes a sequence corresponding to the original barcode sequence from the bead and a sequence corresponding to genomic DNA from the cell. The barcoded constructs can then be released/removed from the partition and, in some embodiments, further processed to add any additional sequences. The resulting constructs can then be sequenced, the sequencing data processed, and the results used to spatially characterize the mRNA and the gDNA from the cell.

In another example, a partition includes a bead that includes a first type of oligonucleotide (e.g., a first capture probe) with a first barcode sequence, a poly(T) priming sequence that can hybridize with the poly(A) tail of an mRNA transcript, and a UMI barcode sequence that can uniquely identify a given transcript. The bead also includes a second type of oligonucleotide (e.g., a second capture probe) with a second barcode sequence, a targeted

priming sequence that is capable of specifically hybridizing with a third barcoded oligonucleotide (e.g., an analyte capture agent) coupled to an antibody that is bound to the surface of the partitioned cell. The third barcoded oligonucleotide includes a UMI barcode sequence that uniquely identifies the antibody (and thus, the particular cell surface feature to which it is bound).

In this example, the first and second barcoded oligonucleotides include the same spatial barcode sequence (e.g., the first and second barcode sequences are the same), which permits downstream association of barcoded nucleic acids with the partition. In some embodiments, however, the first and second barcode sequences are different.

The partition also includes lysis agents that aid in releasing nucleic acids from the cell and can also include an agent (e.g., a reducing agent) that can degrade the bead and/or break a covalent linkage between the barcoded oligonucleotides and the bead, releasing them into the partition. The first type of released barcoded oligonucleotide can hybridize with mRNA released from the cell and the second type of released barcoded oligonucleotide can hybridize with the third type of barcoded oligonucleotide, forming barcoded constructs.

The first type of barcoded construct includes a spatial barcode sequence corresponding to the first barcode sequence from the bead and a sequence corresponding to the UMI barcode sequence from the first type of oligonucleotide, which identifies cell transcripts. The second type of barcoded construct includes a spatial barcode sequence corresponding to the second barcode sequence from the second type of oligonucleotide, and a UMI barcode sequence corresponding to the third type of oligonucleotide (e.g., the analyte capture agent) and used to identify the cell surface feature. The barcoded constructs can then be released/removed from the partition and, in some embodiments, further processed to add any additional sequences. The resulting constructs are then sequenced, sequencing data processed, and the results used to characterize the mRNA and cell surface feature of the cell.

The foregoing discussion involves two specific examples of beads with oligonucleotides for analyzing two different analytes within a partition. More generally, beads that are partitioned can have any of the structures described previously, and can include any of the described combinations of oligonucleotides for analysis of two or more (e.g., three or more, four or more, five or more, six or more, eight or more, ten or more, 12 or more, 15 or more, 20 or more, 25 or more, 30 or more, 40 or more, 50 or more) different types of analytes within a partition.

Examples of beads with combinations of different types of oligonucleotides (e.g., capture probes) for concurrently analyzing different combinations of analytes within partitions include, but are not limited to: (a) genomic DNA and cell surface features (e.g., using the analyte capture agents described herein); (b) mRNA and a lineage tracing construct; (c) mRNA and cell
5 methylation status; (d) mRNA and accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq); (e) mRNA and cell surface or intracellular proteins and/or metabolites; (f) a barcoded analyte capture agent (e.g., the MHC multimers described herein) and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); and (g) mRNA and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense
10 oligonucleotide as described herein).

(f) Sequencing Analysis

After analytes from the sample have hybridized or otherwise been associated with capture probes, analyte capture agents, or other barcoded oligonucleotide sequences according to any of the methods described above in connection with the general spatial cell-based analytical
15 methodology, the barcoded constructs that result from hybridization/association are analyzed via sequencing to identify the analytes.

In some embodiments, where a sample is barcoded directly via hybridization with capture probes or analyte capture agents hybridized, bound, or associated with either the cell surface, or introduced into the cell, as described above, sequencing can be performed on the intact sample.

20 Alternatively, if the barcoded sample has been separated into fragments, cell groups, or individual cells, as described above, sequencing can be performed on individual fragments, cell groups, or cells. For analytes that have been barcoded via partitioning with beads, as described above, individual analytes (e.g., cells, or cellular contents following lysis of cells) can be extracted from the partitions by breaking the partitions, and then analyzed by sequencing to
25 identify the analytes.

A wide variety of different sequencing methods can be used to analyze barcoded analyte constructs. In general, sequenced polynucleotides can be, for example, nucleic acid molecules such as deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), including variants or derivatives thereof (e.g., single stranded DNA or DNA/RNA hybrids, and nucleic acid molecules
30 with a nucleotide analog).

Sequencing of polynucleotides can be performed by various commercial systems. More generally, sequencing can be performed using nucleic acid amplification, polymerase chain reaction (PCR) (e.g., digital PCR and droplet digital PCR (ddPCR), quantitative PCR, real time PCR, multiplex PCR, PCR-based singleplex methods, emulsion PCR), and/or isothermal
5 amplification.

Other examples of methods for sequencing genetic material include, but are not limited to, DNA hybridization methods (e.g., Southern blotting), restriction enzyme digestion methods, Sanger sequencing methods, next-generation sequencing methods (e.g., single-molecule real-time sequencing, nanopore sequencing, and Polony sequencing), ligation methods, and
10 microarray methods. Additional examples of sequencing methods that can be used include targeted sequencing, single molecule real-time sequencing, exon sequencing, electron microscopy-based sequencing, panel sequencing, transistor-mediated sequencing, direct sequencing, random shotgun sequencing, Sanger dideoxy termination sequencing, whole-genome sequencing, sequencing by hybridization, pyrosequencing, capillary electrophoresis, gel
15 electrophoresis, duplex sequencing, cycle sequencing, single-base extension sequencing, solid-phase sequencing, high-throughput sequencing, massively parallel signature sequencing, co-amplification at lower denaturation temperature-PCR (COLD-PCR), sequencing by reversible dye terminator, paired-end sequencing, near-term sequencing, exonuclease sequencing, sequencing by ligation, short-read sequencing, single-molecule sequencing, sequencing-by-
20 synthesis, real-time sequencing, reverse-terminator sequencing, nanopore sequencing, 454 sequencing, Solexa Genome Analyzer sequencing, SOLiD™ sequencing, MS-PET sequencing, and any combinations thereof.

Sequence analysis of the nucleic acid molecules (including barcoded nucleic acid molecules or derivatives thereof) can be direct or indirect. Thus, the sequence analysis substrate
25 (which can be viewed as the molecule which is subjected to the sequence analysis step or process) can directly be the barcoded nucleic acid molecule or it can be a molecule which is derived therefrom (e.g., a complement thereof). Thus, for example, in the sequence analysis step of a sequencing reaction, the sequencing template can be the barcoded nucleic acid molecule or it can be a molecule derived therefrom. For example, a first and/or second strand DNA molecule
30 can be directly subjected to sequence analysis (e.g. sequencing), i.e., can directly take part in the sequence analysis reaction or process (e.g. the sequencing reaction or sequencing process, or be

the molecule which is sequenced or otherwise identified). Alternatively, the barcoded nucleic acid molecule can be subjected to a step of second strand synthesis or amplification before sequence analysis (e.g. sequencing or identification by another technique). The sequence analysis substrate (e.g., template) can thus be an amplicon or a second strand of a barcoded nucleic acid molecule.

In some embodiments, both strands of a double stranded molecule can be subjected to sequence analysis (e.g., sequenced). In some embodiments, single stranded molecules (e.g. barcoded nucleic acid molecules) can be analyzed (e.g. sequenced). To perform single molecule sequencing, the nucleic acid strand can be modified at the 3' end.

Massively parallel sequencing techniques can be used for sequencing nucleic acids, as described above. In one embodiment, a massively parallel sequencing technique can be based on reversible dye-terminators. As an example, DNA molecules are first attached to primers on, e.g., a glass or silicon substrate, and amplified so that local clonal colonies are formed (bridge amplification). Four types of ddNTPs are added, and non-incorporated nucleotides are washed away. Unlike pyrosequencing, the DNA is only extended one nucleotide at a time due to a blocking group (e.g., 3' blocking group present on the sugar moiety of the ddNTP). A detector acquires images of the fluorescently labelled nucleotides, and then the dye along with the terminal 3' blocking group is chemically removed from the DNA, as a precursor to a subsequent cycle. This process can be repeated until the required sequence data is obtained.

As another example, massively parallel pyrosequencing techniques can also be used for sequencing nucleic acids. In pyrosequencing, the nucleic acid is amplified inside water droplets in an oil solution (emulsion PCR), with each droplet containing a single nucleic acid template attached to a single primer-coated bead that then forms a clonal colony. The sequencing system contains many picolitre-volume wells each containing a single bead and sequencing enzymes. Pyrosequencing uses luciferase to generate light for detection of the individual nucleotides added to the nascent nucleic acid and the combined data are used to generate sequence reads.

As another example application of pyrosequencing, released PPi can be detected by being immediately converted to adenosine triphosphate (ATP) by ATP sulfurylase, and the level of ATP generated can be detected via luciferase-produced photons, such as described in Ronaghi, et al., *Anal. Biochem.* 242(1), 84-9 (1996); Ronaghi, *Genome Res.* 11(1), 3-11 (2001); Ronaghi et

al. Science 281 (5375), 363 (1998); and U.S. Pat. Nos. 6,210,891, 6,258,568, and 6,274,320, the entire contents of each of which are incorporated herein by reference.

In some embodiments, sequencing is performed by detection of hydrogen ions that are released during the polymerization of DNA. A microwell containing a template DNA strand to be sequenced can be flooded with a single type of nucleotide. If the introduced nucleotide is complementary to the leading template nucleotide, it is incorporated into the growing complementary strand. This causes the release of a hydrogen ion that triggers a hypersensitive ion sensor, which indicates that a reaction has occurred. If homopolymer repeats are present in the template sequence, multiple nucleotides will be incorporated in a single cycle. This leads to a corresponding number of released hydrogen ions and a proportionally higher electronic signal.

In some embodiments, sequencing can be performed in-situ. In-situ sequencing methods are particularly useful, for example, when the biological sample remains intact after analytes on the sample surface (e.g., cell surface analytes) or within the sample (e.g., intracellular analytes) have been barcoded. In-situ sequencing typically involves incorporation of a labeled nucleotide (e.g., fluorescently labeled mononucleotides or dinucleotides) in a sequential, template-dependent manner or hybridization of a labeled primer (e.g., a labeled random hexamer) to a nucleic acid template such that the identities (i.e., nucleotide sequence) of the incorporated nucleotides or labeled primer extension products can be determined, and consequently, the nucleotide sequence of the corresponding template nucleic acid. Aspects of in-situ sequencing are described, for example, in Mitra et al., (2003) Anal. Biochem., 320, 55-65, and Lee et al., (2014) Science, 343(6177), 1360-1363, the entire contents of each of which are incorporated herein by reference.

In addition, examples of methods and systems for performing in-situ sequencing are described in PCT Patent Application Publication Nos. WO2014/163886, WO2018/045181, WO2018/045186, and in U.S. Patent Nos. 10,138,509 and 10,179,932, the entire contents of each of which are incorporated herein by reference. Example techniques for in-situ sequencing include, but are not limited to, STARmap (described for example in Wang et al., (2018) Science, 361(6499) 5691), MERFISH (described for example in Moffitt, (2016) Methods in Enzymology, 572, 1-49), and FISSEQ (described for example in U.S. Patent Application Publication No. 2019/0032121). The entire contents of each of the foregoing references are incorporated herein by reference.

For analytes that have been barcoded via partitioning, barcoded nucleic acid molecules or derivatives thereof (e.g., barcoded nucleic acid molecules to which one or more functional sequences have been added, or from which one or more features have been removed) can be pooled and processed together for subsequent analysis such as sequencing on high throughput
5 sequencers. Processing with pooling can be implemented using barcode sequences. For example, barcoded nucleic acid molecules of a given partition can have the same barcode, which is different from barcodes of other spatial partitions. Alternatively, barcoded nucleic acid molecules of different partitions can be processed separately for subsequent analysis (e.g., sequencing).

10 In some embodiments, where capture probes do not contain a spatial barcode, the spatial barcode can be added after the capture probe captures analytes from a biological sample and before analysis of the analytes. When a spatial barcode is added after an analyte is captured, the barcode can be added after amplification of the analyte (e.g., reverse transcription and polymerase amplification of RNA). In some embodiments, analyte analysis uses direct
15 sequencing of one or more captured analytes, such as direct sequencing of hybridized RNA. In some embodiments, direct sequencing is performed after reverse transcription of hybridized RNA. In some embodiments direct sequencing is performed after amplification of reverse transcription of hybridized RNA.

In some embodiments, direct sequencing of captured RNA is performed by sequencing-
20 by-synthesis (SBS). In some embodiments, a sequencing primer is complementary to a sequence in one or more of the domains of a capture probe (e.g., functional domain). In such embodiments, sequencing-by-synthesis can include reverse transcription and/or amplification in order to generate a template sequence (e.g., functional domain) from which a primer sequence can bind.

25 SBS can involve hybridizing an appropriate primer, sometimes referred to as a sequencing primer, with the nucleic acid template to be sequenced, extending the primer, and detecting the nucleotides used to extend the primer. Preferably, the nucleic acid used to extend the primer is detected before a further nucleotide is added to the growing nucleic acid chain, thus allowing base-by-base in situ nucleic acid sequencing. The detection of incorporated nucleotides
30 is facilitated by including one or more labelled nucleotides in the primer extension reaction. To allow the hybridization of an appropriate sequencing primer to the nucleic acid template to be

sequenced, the nucleic acid template should normally be in a single stranded form. If the nucleic acid templates making up the nucleic acid spots are present in a double stranded form these can be processed to provide single stranded nucleic acid templates using methods well known in the art, for example by denaturation, cleavage etc. The sequencing primers which are hybridized to the nucleic acid template and used for primer extension are preferably short oligonucleotides, for example, 15 to 25 nucleotides in length. The sequencing primers can be greater than 25 nucleotides in length as well. For example, sequencing primers can be about 20 to about 60 nucleotides in length, or more than 60 nucleotides in length. The sequencing primers can be provided in solution or in an immobilized form. Once the sequencing primer has been annealed to the nucleic acid template to be sequenced by subjecting the nucleic acid template and sequencing primer to appropriate conditions, primer extension is carried out, for example using a nucleic acid polymerase and a supply of nucleotides, at least some of which are provided in a labelled form, and conditions suitable for primer extension if a suitable nucleotide is provided.

Preferably after each primer extension step, a washing step is included in order to remove unincorporated nucleotides which can interfere with subsequent steps. Once the primer extension step has been carried out, the nucleic acid colony is monitored to determine whether a labelled nucleotide has been incorporated into an extended primer. The primer extension step can then be repeated to determine the next and subsequent nucleotides incorporated into an extended primer. If the sequence being determined is unknown, the nucleotides applied to a given colony are usually applied in a chosen order which is then repeated throughout the analysis, for example dATP, dTTP, dCTP, dGTP.

SBS techniques which can be used are described for example, but not limited to, those in U.S. Patent App. Pub. No. 2007/0166705, U.S. Patent App. Pub. No. 2006/0188901, U.S. Patent 7,057,026, U.S. Patent App. Pub. No. 2006/0240439, U.S. Patent App. Pub. No. 2006/0281109, PCT Patent App. Pub. No. WO 05/065814, U.S. Patent App. Pub. No. 2005/0100900, PCT Patent App. Pub. No. WO 06/064199, PCT Patent App. Pub. No. WO07/010,251, U.S. Patent App. Pub. No. 2012/0270305, U.S. Patent App. Pub. No. 2013/0260372, and U.S. Patent App. Pub. No. 2013/0079232, the entire contents of each of which are incorporated herein by reference.

In some embodiments, direct sequencing of captured RNA is performed by sequential fluorescence hybridization (e.g., sequencing by hybridization). In some embodiments, a

hybridization reaction where RNA is hybridized to a capture probe is performed in situ. In some embodiments, captured RNA is not amplified prior to hybridization with a sequencing probe. In some embodiments, RNA is amplified prior to hybridization with sequencing probes (e.g., reverse transcription to cDNA and amplification of cDNA). In some embodiments, amplification is performed using single-molecule hybridization chain reaction. In some embodiments, amplification is performed using rolling chain amplification.

Sequential fluorescence hybridization can involve sequential hybridization of probes including degenerate primer sequences and a detectable label. A degenerate primer sequence is a short oligonucleotide sequence which is capable of hybridizing to any nucleic acid fragment independent of the sequence of said nucleic acid fragment. For example, such a method could include the steps of: (a) providing a mixture including four probes, each of which includes either A, C, G, or T at the 5'-terminus, further including degenerate nucleotide sequence of 5 to 11 nucleotides in length, and further including a functional domain (e.g., fluorescent molecule) that is distinct for probes with A, C, G, or T at the 5'-terminus; (b) associating the probes of step (a) to the target polynucleotide sequences, whose sequence needs will be determined by this method; (c) measuring the activities of the four functional domains and recording the relative spatial location of the activities; (d) removing the reagents from steps (a)-(b) from the target polynucleotide sequences; and repeating steps (a)-(d) for n cycles, until the nucleotide sequence of the spatial domain for each bead is determined, with modification that the oligonucleotides used in step (a) are complementary to part of the target polynucleotide sequences and the positions 1 through n flanking the part of the sequences. Because the barcode sequences are different, in some embodiments, these additional flanking sequences are degenerate sequences. The fluorescent signal from each spot on the array for cycles 1 through n can be used to determine the sequence of the target polynucleotide sequences.

In some embodiments, direct sequencing of captured RNA using sequential fluorescence hybridization is performed in vitro. In some embodiments, captured RNA is amplified prior to hybridization with a sequencing probe (e.g., reverse transcription to cDNA and amplification of cDNA). In some embodiments, a capture probe containing captured RNA is exposed to the sequencing probe targeting coding regions of RNA. In some embodiments, one or more sequencing probes are targeted to each coding region. In some embodiments, the sequencing probe is designed to hybridize with sequencing reagents (e.g., a dye-labeled readout

oligonucleotides). A sequencing probe can then hybridize with sequencing reagents. In some embodiments, output from the sequencing reaction is imaged. In some embodiments, a specific sequence of cDNA is resolved from an image of a sequencing reaction. In some embodiments, reverse transcription of captured RNA is performed prior to hybridization to the sequencing probe. In some embodiments, the sequencing probe is designed to target complementary sequences of the coding regions of RNA (e.g., targeting cDNA).

In some embodiments, a captured RNA is directly sequenced using a nanopore-based method. In some embodiments, direct sequencing is performed using nanopore direct RNA sequencing in which captured RNA is translocated through a nanopore. A nanopore current can be recorded and converted into a base sequence. In some embodiments, captured RNA remains attached to a substrate during nanopore sequencing. In some embodiments, captured RNA is released from the substrate prior to nanopore sequencing. In some embodiments, where the analyte of interest is a protein, direct sequencing of the protein can be performed using nanopore-based methods. Examples of nanopore-based sequencing methods that can be used are described in Deamer et al., Trends Biotechnol. 18, 147-151 (2000); Deamer et al., Acc. Chem. Res. 35:817-825 (2002); Li et al., Nat. Mater. 2:611-615 (2003); Soni et al., Clin. Chem. 53, 1996-2001 (2007); Healy et al., Nanomed. 2, 459-481 (2007); Cockroft et al., J. Am. Chem. Soc. 130, 818-820 (2008); and in U.S. Patent 7,001,792. The entire contents of each of the foregoing references are incorporated herein by reference.

In some embodiments, direct sequencing of captured RNA is performed using single molecule sequencing by ligation. Such techniques utilize DNA ligase to incorporate oligonucleotides and identify the incorporation of such oligonucleotides. The oligonucleotides typically have different labels that are correlated with the identity of a particular nucleotide in a sequence to which the oligonucleotides hybridize. Aspects and features involved in sequencing by ligation are described, for example, in Shendure et al. Science (2005), 309: 1728-1732, and in U.S. Patent Nos. 5,599,675; 5,750,341; 6,969,488; 6,172,218; and 6,306,597, the entire contents of each of which are incorporated herein by reference.

In some embodiments, nucleic acid hybridization can be used for sequencing. These methods utilize labeled nucleic acid decoder probes that are complementary to at least a portion of a barcode sequence. Multiplex decoding can be performed with pools of many different probes with distinguishable labels. Non-limiting examples of nucleic acid hybridization

sequencing are described for example in U.S. Pat. No. 8,460,865, and in Gunderson et al., Genome Research 14:870-877 (2004), the entire contents of each of which are incorporated herein by reference.

In some embodiments, commercial high-throughput digital sequencing techniques can be used to analyze barcode sequences, in which DNA templates are prepared for sequencing not one at a time, but in a bulk process, and where many sequences are read out preferably in parallel, or alternatively using an ultra-high throughput serial process that itself may be parallelized.

Examples of such techniques include Illumina[®] sequencing (e.g., flow cell-based sequencing techniques), sequencing by synthesis using modified nucleotides (such as commercialized in TruSeq[™] and HiSeq[™] technology by Illumina, Inc., San Diego, CA), HeliScope[™] by Helicos Biosciences Corporation, Cambridge, MA, and PacBio RS by Pacific Biosciences of California, Inc., Menlo Park, CA), sequencing by ion detection technologies (Ion Torrent, Inc., South San Francisco, CA), and sequencing of DNA nanoballs (Complete Genomics, Inc., Mountain View, CA).

In some embodiments, detection of a proton released upon incorporation of a nucleotide into an extension product can be used in the methods described herein. For example, the sequencing methods and systems described in U.S. Patent Application Publication Nos. 2009/0026082, 2009/0127589, 2010/0137143, and 2010/0282617, can be used to directly sequence barcodes. The entire contents of each of the foregoing references are incorporated herein by reference.

In some embodiments, real-time monitoring of DNA polymerase activity can be used during sequencing. For example, nucleotide incorporations can be detected through fluorescence resonance energy transfer (FRET), as described for example in Levene et al., Science (2003), 299, 682-686, Lundquist et al., Opt. Lett. (2008), 33, 1026-1028, and Korlach et al., Proc. Natl. Acad. Sci. USA (2008), 105, 1176-1181. The entire contents of each of the foregoing references are herein incorporated by reference.

IV. Multiplexing

(a) Multiplexing Generally

In various embodiments of spatial analysis as described herein, features can include different types of capture probes for analyzing both intrinsic and extrinsic information for

individual cells. For example, a feature can include one or more of the following: 1) a capture probe featuring a capture domain that binds to one or more endogenous nucleic acids in the cell; 2) a capture probe featuring a capture domain that binds to one or more exogenous nucleic acids in the cell (e.g., nucleic acids from a microorganism (e.g., a virus, a bacterium)) that infects the cell, nucleic acids introduced into the cell (e.g., such as plasmids or nucleic acid derived therefrom), nucleic acids for gene editing (e.g., CRISPR-related RNA such as crRNA, guide RNA); 3) a capture probe featuring a capture domain that binds to a analyte capture agent (e.g., an antibody coupled to a oligonucleotide that includes a capture agent barcode domain having an analyte capture sequence that binds the capture domain), and 4) a capture moiety featuring a domain that binds to a protein (e.g., an exogenous protein expressed in the cell, a protein from a microorganism (e.g., a virus, a bacterium)) that infects the cell, or a binding partner for a protein of the cell (e.g., an antigen for an immune cell receptor).

In some embodiments of any of the spatial analysis methods as described herein, spatial profiling includes concurrent analysis of two different types of analytes. A feature can be a gel bead, which is coupled (e.g., reversibly coupled) to one or more capture probes. The capture probes can include a spatial barcode sequence and a poly (T) priming sequence that can hybridize with the poly (A) tail of an mRNA transcript. The capture probe can also include a UMI sequence that can uniquely identify a given transcript. The capture probe can also include a spatial barcode sequence and a random N-mer priming sequence that is capable of randomly hybridizing with gDNA. In this configuration, capture probes can include the same spatial barcode sequence, which permits association of downstream sequencing reads with the feature.

In some embodiments of any of the spatial analysis methods as described herein, a feature can be a gel bead, which is coupled (e.g., reversibly coupled) to capture probes. The Capture probe can include a spatial barcode sequence and a poly(T) priming sequence that can hybridize with the poly(A) tail of an mRNA transcript. The capture probe can also include a UMI sequence that can uniquely identify a given transcript. The capture probe can include a spatial barcode sequence and a capture domain that is capable of specifically hybridizing with an analyte capture agent. The analyte capture agent can include an oligonucleotide that includes an analyte capture sequence that interacts with the capture domain coupled to the feature. The oligonucleotide of the analyte capture agent can be coupled to an antibody that is bound to the surface of a cell. The oligonucleotide includes a barcode sequence (e.g., an analyte binding

moiety barcode) that uniquely identifies the antibody (and thus, the particular cell surface feature to which it is bound). In this configuration, the capture probes include the same spatial barcode sequence, which permit downstream association of barcoded nucleic acids with the location on the spatial array. In some embodiments of any of the spatial profiling methods described herein, the analyte capture agents can be produced by any suitable route, including via example coupling schemes described elsewhere herein.

In some embodiments of any of the spatial analysis methods described herein, other combinations of two or more biological analytes that can be concurrently measured include, without limitation: (a) genomic DNA and cell surface features (e.g., via analyte capture agents that bind to a cell surface feature), (b) mRNA and a lineage tracing construct, (c) mRNA and cell methylation status, (d) mRNA and accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), (e) mRNA and cell surface or intracellular proteins and/or metabolites, (f) mRNA and chromatin (spatial organization of chromatin in a cell), (g) an analyte capture agent (e.g., any of the MHC multimers described herein) and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor), (h) mRNA and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein), (i) genomic DNA and a perturbation agent, (j) an analyte capture agent and a perturbation reagents, (k) accessible chromatin and a perturbation reagent, (l) chromatin (e.g., spatial organization of chromatin in a cell) and a perturbation reagent, and (m) cell surface or intracellular proteins and/or metabolites and a perturbation reagent, or any combination thereof.

In some embodiments of any of the spatial analysis methods described herein, the first analyte can include a nucleic acid molecule with a nucleic acid sequence (e.g., mRNA, complementary DNA derived from reverse transcription of mRNA) encoding at least a portion of a V(D)J sequence of an immune cell receptor (e.g., a TCR or BCR). In some embodiments, the nucleic acid molecule with a nucleic acid sequence encoding at least a portion of a V(D)J sequence of an immune cell receptor is cDNA first generated from reverse transcription of the corresponding mRNA, using a poly(T) containing primer. The cDNA that is generated can then be barcoded using a primer, featuring a spatial barcode sequence (and optionally, a UMI sequence) that hybridizes with at least a portion of the cDNA that is generated. In some embodiments, a template switching oligonucleotide in conjunction a terminal transferase or a reverse transcriptase having terminal transferase activity can be employed to generate a priming

region on the cDNA to which a barcoded primer can hybridize during cDNA generation.

Terminal transferase activity can, for example, add a poly(C) tail to a 3' end of the cDNA such that the template switching oligonucleotide can bind via a poly(G) priming sequence and the 3' end of the cDNA can be further extended. The original mRNA template and template switching
5 oligonucleotide can then be denatured from the cDNA and the barcoded primer comprising a sequence complementary to at least a portion of the generated priming region on the cDNA can then hybridize with the cDNA and a barcoded construct comprising the barcode sequence (and any optional UMI sequence) and a complement of the cDNA generated. Additional methods and compositions suitable for barcoding cDNA generated from mRNA transcripts including those
10 encoding V(D)J regions of an immune cell receptor and/or barcoding methods and composition including a template switch oligonucleotide are described, for example, in PCT Patent Application Publication No. WO 2018/075693, and in U.S. Patent Application Publication No. 2018/0105808, the entire contents of each of which are incorporated herein by reference.

In some embodiments, V(D)J analysis can be performed using methods similar to those
15 described herein. For example, V(D)J analysis can be completed with the use of one or more analyte capture agents that bind to particular surface features of immune cells and are associated with barcode sequences (e.g., analyte binding moiety barcodes). The one or more analyte capture agents can include an MHC or MHC multimer. A barcoded oligonucleotide coupled to a bead that can be used for V(D)J analysis. The oligonucleotide is coupled to a bead by a releasable
20 linkage, such as a disulfide linker. The oligonucleotide can include functional sequences that are useful for subsequent processing, such as functional sequence, which can include a sequencer specific flow cell attachment sequence, e.g., a P5 sequence, as well as functional sequence, which can include sequencing primer sequences, e.g., a R1 primer binding site. In some
25 embodiments, the sequence can include a P7 sequence and a R2 primer binding site. A barcode sequence can be included within the structure for use in barcoding the template polynucleotide. The functional sequences can be selected for compatibility with a variety of different sequencing systems, e.g., 454 Sequencing, Ion Torrent Proton or PGM, Illumina X10, etc., and the requirements thereof. In some embodiments, the barcode sequence, functional sequences (e.g.,
flow cell attachment sequence) and additional sequences (e.g., sequencing primer sequences) can
30 be common to all of the oligonucleotides attached to a given bead. The barcoded oligonucleotide can also include a sequence to facilitate template switching (e.g., a poly(G) sequence). In some

embodiments, the additional sequence provides a unique molecular identifier (UMI) sequence segment, as described elsewhere herein.

In an exemplary method of cellular polynucleotide analysis using a barcode oligonucleotide, a cell is co-partitioned along with a bead bearing a barcoded oligonucleotide and additional reagents such as a reverse transcriptase, primers, oligonucleotides (e.g., template switching oligonucleotides), dNTPs, and a reducing agent into a partition (e.g., a droplet in an emulsion). Within the partition, the cell can be lysed to yield a plurality of template polynucleotides (e.g., DNA such as genomic DNA, RNA such as mRNA, etc.).

A reaction mixture featuring a template polynucleotide from a cell and (i) the primer having a sequence towards a 3' end that hybridizes to the template polynucleotide (e.g., poly(T)) and (ii) a template switching oligonucleotide that includes a first oligonucleotide towards a 5' end can be subjected to an amplification reaction to yield a first amplification product. In some embodiments, the template polynucleotide is an mRNA with a poly(A) tail and the primer that hybridizes to the template polynucleotide includes a poly(T) sequence towards a 3' end, which is complementary to the poly(A) segment. The first oligonucleotide can include at least one of an adaptor sequence, a barcode sequence, a unique molecular identifier (UMI) sequence, a primer binding site, and a sequencing primer binding site or any combination thereof. In some cases, a first oligonucleotide is a sequence that can be common to all partitions of a plurality of partitions. For example, the first oligonucleotide can include a flow cell attachment sequence, an amplification primer binding site, or a sequencing primer binding site and the first amplification reaction facilitates the attachment the oligonucleotide to the template polynucleotide from the cell. In some embodiments, the first oligonucleotide includes a primer binding site. In some embodiments, the first oligonucleotide includes a sequencing primer binding site.

The sequence towards a 3' end (e.g., poly(T)) of the primer hybridizes to the template polynucleotide. In a first amplification reaction, extension reaction reagents, e.g., reverse transcriptase, nucleoside triphosphates, co-factors (e.g., Mg^{2+} or Mn^{2+}), that are also co-partitioned, can extend the primer sequence using the cell's nucleic acid as a template, to produce a transcript, e.g., cDNA, having a fragment complementary to the strand of the cell's nucleic acid to which the primer annealed. In some embodiments, the reverse transcriptase has terminal transferase activity and the reverse transcriptase adds additional nucleotides, e.g., poly(C), to the cDNA in a template independent manner.

The template switching oligonucleotide, for example a template switching oligonucleotide which includes a poly(G) sequence, can hybridize to the cDNA and facilitate template switching in the first amplification reaction. The transcript, therefore, can include the sequence of the primer, a sequence complementary to the template polynucleotide from the cell, and a sequence complementary to the template switching oligonucleotide.

In some embodiments of any of the spatial analysis methods described herein, subsequent to the first amplification reaction, the first amplification product or transcript can be subjected to a second amplification reaction to generate a second amplification product. In some embodiments, additional sequences (e.g., functional sequences such as flow cell attachment sequence, sequencing primer binding sequences, barcode sequences, etc.) are attached. The first and second amplification reactions can be performed in the same volume, such as for example in a droplet. In some embodiments, the first amplification product is subjected to a second amplification reaction in the presence of a barcoded oligonucleotide to generate a second amplification product having a barcode sequence. The barcode sequence can be unique to a partition, that is, each partition can have a unique barcode sequence. The barcoded oligonucleotide can include a sequence of at least a segment of the template switching oligonucleotide and at least a second oligonucleotide. The segment of the template switching oligonucleotide on the barcoded oligonucleotide can facilitate hybridization of the barcoded oligonucleotide to the transcript, e.g., cDNA, to facilitate the generation of a second amplification product. In addition to a barcode sequence, the barcoded oligonucleotide can include a second oligonucleotide such as at least one of an adaptor sequence, a unique molecular identifier (UMI) sequence, a primer binding site, and a sequencing primer binding site, or any combination thereof.

In some embodiments of any of the spatial analysis methods described herein, the second amplification reaction uses the first amplification product as a template and the barcoded oligonucleotide as a primer. In some embodiments, the segment of the template switching oligonucleotide on the barcoded oligonucleotide can hybridize to the portion of the cDNA or complementary fragment having a sequence complementary to the template switching oligonucleotide or that which was copied from the template switching oligonucleotide. In the second amplification reaction, extension reaction reagents, e.g., polymerase, nucleoside triphosphates, co-factors (e.g., Mg^{2+} or Mn^{2+}), that are also co-partitioned, can extend the primer

sequence using the first amplification product as template. The second amplification product can include a second oligonucleotide, a sequence of a segment of the template polynucleotide (e.g., mRNA), and a sequence complementary to the primer.

In some embodiments of any of the spatial analysis methods described herein, the second
5 amplification product uses the barcoded oligonucleotide as a template and at least a portion of the first amplification product as a primer. The segment of the first amplification product (e.g., cDNA) having a sequence complementary to the template switching oligonucleotide can hybridize to the segment of the barcoded oligonucleotide comprising a sequence of at least a segment of the template switching oligonucleotide. In the second amplification reaction,
10 extension reaction reagents, e.g., polymerase, nucleoside triphosphates, co-factors (e.g., Mg^{2+} or Mn^{2+}), that are also co-partitioned, can extend the primer sequence (e.g., first amplification product) using the barcoded oligonucleotide as template. The second amplification product can include the sequence of the primer, a sequence which is complementary to the sequence of the template polynucleotide (e.g., mRNA), and a sequence complementary to the second
15 oligonucleotide.

In some embodiments of any of the spatial analysis methods described herein, three or more classes of biological analytes can be concurrently measured. For example, a feature can include capture probes that can participate in an assay of at least three different types of analytes via three different capture domains. A bead can be coupled to a barcoded oligonucleotide that
20 includes a capture domain that includes a poly(T) priming sequence for mRNA analysis; a barcoded oligonucleotide that includes a capture domain that includes a random N-mer priming sequence for gDNA analysis; and a barcoded oligonucleotide that includes a capture domain that can specifically bind a an analyte capture agent (e.g., an antibody with a spatial barcode), via its analyte capture sequence.

In some embodiments of any of the spatial analysis methods described herein, other
25 combinations of three or more biological analytes that can be concurrently measured include, without limitation: (a) mRNA, a lineage tracing construct, and cell surface and/or intracellular proteins and/or metabolites; (b) mRNA, accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), and cell surface and/or intracellular proteins and/or metabolites; (c) mRNA,
30 genomic DNA, and a perturbation reagent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein); (d) mRNA, accessible

chromatin, and a perturbation reagent; (e) mRNA, an analyte capture agent (e.g., any of the MHC multimers described herein), and a perturbation reagent; (f) mRNA, cell surface and/or intracellular proteins and/or metabolites, and a perturbation reagent; (g) mRNA, a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor), and a perturbation reagent; (h) mRNA, an analyte capture agent, and a V(D)J sequence of an immune cell receptor; (i) cell surface and/or intracellular proteins and/or metabolites, a an analyte capture agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor; (j) methylation status, mRNA, and cell surface and/or intracellular proteins and/or metabolites; (k) mRNA, chromatin (e.g., spatial organization of chromatin in a cell), and a perturbation reagent; (l) a V(D)J sequence of an immune cell receptor, chromatin (e.g., spatial organization of chromatin in a cell); and a perturbation reagent; and (m) mRNA, a V(D)J sequence of an immune cell receptor, and chromatin (e.g., spatial organization of chromatin in a cell), or any combination thereof.

In some embodiments of any of the spatial analysis methods described herein, four or more classes biological analytes can be concurrently measured. A feature can be a bead that is coupled to barcoded primers that can each participate in an assay of a different type of analyte. The feature is coupled (e.g., reversibly coupled) to a capture probe that includes a capture domain that includes a poly(T) priming sequence for mRNA analysis and is also coupled (e.g., reversibly coupled) to capture probe that includes a capture domain that includes a random N-mer priming sequence for gDNA analysis. Moreover, the feature is also coupled (e.g., reversibly coupled) to a capture probe that binds an analyte capture sequence of an analyte capture agent via its capture domain. The feature can also be coupled (e.g., reversibly coupled) to a capture probe that can specifically bind a nucleic acid molecule that can function as a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein), via its capture domain.

In some embodiments of any of the spatial analysis methods described herein, each of the various spatially barcoded capture probes present at a given feature or on a given bead include the same spatial barcode sequence. In some embodiments, each barcoded capture probe can be released from the feature in a manner suitable for analysis of its respective analyte. For example, barcoded constructs A, B, C and D can be generated as described elsewhere herein and analyzed. Barcoded construct A can include a sequence corresponding to the barcode sequence from the bead (e.g., a spatial barcode) and a DNA sequence corresponding to a target mRNA. Barcoded

construct B can include a sequence corresponding to the barcode sequence from the bead (e.g., a spatial barcode) and a sequence corresponding to genomic DNA. Barcoded construct C can include a sequence corresponding to the barcode sequence from the bead (e.g., a spatial barcode) and a sequence corresponding to barcode sequence associated with an analyte capture agent (e.g., an analyte binding moiety barcode). Barcoded construct D can include a sequence corresponding to the barcode sequence from the bead (e.g., a spatial barcode) and a sequence corresponding to a CRISPR nucleic acid (which, in some embodiments, also includes a barcode sequence). Each construct can be analyzed (e.g., via any of a variety of sequencing methods) and the results can be associated with the given cell from which the various analytes originated. Barcoded (or even non-barcoded) constructs can be tailored for analyses of any given analyte associated with a nucleic acid and capable of binding with such a construct.

In some embodiments of any of the spatial analysis methods described herein, other combinations of four or more biological analytes that can be concurrently measured include, without limitation: (a) mRNA, a lineage tracing construct, cell surface and/or intracellular proteins and/or metabolites, and gDNA; (b) mRNA, accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq), cell surface and/or intracellular proteins and/or metabolites, and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein); (c) mRNA, cell surface and/or intracellular proteins and/or metabolites, an analyte capture agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); (d) mRNA, genomic DNA, a perturbation reagent, and accessible chromatin; (e) mRNA, cell surface and/or intracellular proteins and/or metabolites, an analyte capture agent (e.g., the MHC multimers described herein), and a perturbation reagent; (f) mRNA, cell surface and/or intracellular proteins and/or metabolites, a perturbation reagent, and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); (g) mRNA, a perturbation reagent, an analyte capture agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor); (h) mRNA, chromatin (e.g., spatial organization of chromatin in a cell), and a perturbation reagent; (i) a V(D)J sequence of an immune cell receptor, chromatin (e.g., spatial organization of chromatin in a cell); and a perturbation reagent; (j) mRNA, a V(D)J sequence of an immune cell receptor, chromatin (e.g., spatial organization of chromatin in a cell), and genomic DNA; (k) mRNA, a V(D)J sequence of an immune cell receptor, chromatin (e.g.,

spatial organization of chromatin in a cell), and a perturbation reagent, or any combination thereof.

(b) Construction of spatial arrays for multi-analyte analysis

This disclosure also provides methods and materials for constructing a spatial array
5 capable of multi-analyte analysis. In some embodiments, a spatial array includes a plurality of features on a substrate where one or more members of the plurality of features include a plurality of oligonucleotides having a first type functional sequence and oligonucleotides having a second, different type of functional sequence. In some embodiments, a feature can include oligonucleotides with two types of functional sequences. A feature can be coupled to
10 oligonucleotides comprising a TruSeq functional sequence and also to oligonucleotides comprising a Nextera functional sequence. In some embodiments, one or more members of the plurality of features comprises both types of functional sequences. In some embodiments, one or more members of the plurality features includes a first type of functional sequence. In some
15 embodiments, one or more members of the plurality of features includes a second type of functional sequence. In some embodiments, an additional oligonucleotide can be added to the functional sequence to generate a full oligonucleotide where the full oligonucleotide includes a spatial barcode sequence, an optional UMI sequence, a priming sequence, and a capture domain. Attachment of these sequences can be via ligation (including via splint ligation as is described in U.S. Patent Application Publication No. 20140378345, the entire contents of which are
20 incorporated herein by reference), or any other suitable route. As discussed herein, oligonucleotides can be hybridized with splint sequences that can be helpful in constructing complete full oligonucleotides (e.g., oligonucleotides that are capable of spatial analysis).

In some embodiments, the oligonucleotides that hybridize to the functional sequences (e.g., TruSeq and Nextera) located on the features include capture domains capable of capturing
25 different types of analytes (e.g., mRNA, genomic DNA, cell surface proteins, or accessible chromatin). In some examples, oligonucleotides that can bind to the TruSeq functional sequences can include capture domains that include poly(T) capture sequences. In addition to the poly(T) capture sequences, the oligonucleotides that can bind the TruSeq functional groups can also include a capture domain that includes a random N-mer sequence for capturing genomic DNA
30 (e.g., or any other sequence or domain as described herein capable of capturing any of the biological analytes described herein). In such cases, the spatial arrays can be constructed by

applying ratios of TruSeq-poly(T) and TruSeq-N-mer oligonucleotides to the features comprising the functional TruSeq sequences. This can produce spatial arrays where a portion of the oligonucleotides can capture mRNA and a different portion of oligonucleotides can capture genomic DNA. In some embodiments, one or more members of a plurality of features include both TruSeq and Nextera functional sequences. In such cases, a feature including both types of functional sequences is capable of binding oligonucleotides specific to each functional sequence. For example, an oligonucleotide capable of binding to a TruSeq functional sequence could be used to deliver an oligonucleotide including a poly(T) capture domain and an oligonucleotide capable of binding to a Nextera functional sequence could be used to deliver an oligonucleotide including an N-mer capture domain for capturing genomic DNA. It will be appreciated by a person of ordinary skill in the art that any combination of capture domains (e.g., capture domains having any of the variety of capture sequences described herein capable of binding to any of the different types of analytes as described herein) could be combined with oligonucleotides capable of binding to TruSeq and Nextera functional sequences to construct a spatial array.

In some embodiments, an oligonucleotide that includes a capture domain (e.g., an oligonucleotide capable of coupling to an analyte) or an analyte capture agent can include an oligonucleotide sequence that is capable of binding or ligating to an assay primer. The adapter can allow the capture probe or the analyte capture agent to be attached to any suitable assay primers and used in any suitable assays. The assay primer can include a priming region and a sequence that is capable of binding or ligating to the adapter. In some embodiments, the adapter can be a non-specific primer (e.g., a 5' overhang) and the assay primer can include a 3' overhang that can be ligated to the 5' overhang. The priming region on the assay primer can be any primer described herein, e.g., a poly (T) primer, a random N-mer primer, a target-specific primer, or an analyte capture agent capture sequence.

In some examples, an oligonucleotide can includes an adapter, e.g., a 5' overhang with 10 nucleotides. The adapter can be ligated to assay primers, each of which includes a 3' overhang with 10 nucleotides that complementary to the 5' overhang of the adapter. The capture probe can be used in any assay by attaching to the assay primer designed for that assay.

Adapters and assay primers can be used to allow the capture probe or the analyte capture agent to be attached to any suitable assay primers and used in any suitable assays. A capture probe that includes a spatial barcode can be attached to a bead that includes a poly(dT) sequence.

A capture probe including a spatial barcode and a poly(T) sequence can be used to assay multiple biological analytes as generally described herein (e.g., the biological analyte includes a poly(A) sequence or is coupled to or otherwise is associated with an analyte capture agent comprising a poly(A) sequence as the analyte capture sequence).

5 A splint oligonucleotide with a poly(A) sequence can be used to facilitate coupling to a capture probe that includes a spatial barcode and a second sequence that facilitates coupling with an assay primer. Assay primers include a sequence complementary to the splint oligo second sequence and an assay-specific sequence that determines assay primer functionality (e.g., a poly(T) primer, a random N-mer primer, a target-specific primer, or an analyte capture agent
10 capture sequence as described herein).

In some embodiments of any of the spatial profiling methods described herein, a feature can include a capture probe that includes a spatial barcode comprising a switch oligonucleotide, e.g., with a 3' end 3rG. For example, a feature (e.g., a gel bead) with a spatial barcode functionalized with a 3rG sequence can be used that enables template switching (e.g., reverse
15 transcriptase template switching), but is not specific for any particular assay. In some embodiments, the assay primers added to the reaction can determine which type of analytes are analyzed. For example, the assay primers can include binding domains capable of binding to target biological analytes (e.g., poly (T) for mRNA, N-mer for genomic DNA, etc.). A capture probe (e.g., an oligonucleotide capable of spatial profiling) can be generated by using a reverse
20 transcriptase enzyme/polymerase to extend, which is followed by template switching onto the barcoded adapter oligonucleotide to incorporate the barcode and other functional sequences. In some embodiments, the assay primers include capture domains capable of binding to a poly(T) sequence for mRNA analysis, random primers for genomic DNA analysis, or a capture sequence that can bind a nucleic acid molecule coupled to an analyte binding moiety (e.g., a an analyte
25 capture sequence of an analyte capture agent) or a nucleic acid molecule that can function in as a perturbation reagent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein).

V. Systems for Sample Analysis

30 The methods described above for analyzing biological samples can be implemented using a variety of hardware components. In this section, examples of such components are described.

However, it should be understood that in general, the various steps and techniques discussed herein can be performed using a variety of different devices and system components, not all of which are expressly set forth.

FIG. 27A is a schematic diagram showing an example sample handling apparatus **2700**.

5 Sample handling apparatus **2700** includes a sample chamber **2702** that, when closed or sealed, is fluid-tight. Within chamber **2702**, a first holder **2704** holds a first substrate **2706** on which a sample **2708** is positioned. Sample chamber **2702** also includes a second holder **2710** that holds a second substrate **2712** with an array of features **2714**, as described above.

10 A fluid reservoir **2716** is connected to the interior volume of sample chamber **2702** via a fluid inlet **2718**. Fluid outlet **2720** is also connected to the interior volume of sample chamber **2702**, and to valve **2722**. In turn, valve **2722** is connected to waste reservoir **2724** and, optionally, to analysis apparatus **2726**. A control unit **2728** is electrically connected to second holder **2710**, to valve **2722**, to waste reservoir **2724**, and to fluid reservoir **2716**.

15 During operation of apparatus **2700**, any of the reagents, solutions, and other biochemical components described above can be delivered into sample chamber **2702** from fluid reservoir **2716** via fluid inlet **2718**. Control unit **2728**, connected to fluid reservoir **2716**, can control the delivery of reagents, solutions, and components, and adjust the volumes and flow rates according to programmed analytical protocols for various sample types and analysis procedures. In some embodiments, fluid reservoir **2716** includes a pump, which can be controlled by control unit
20 **2728**, to facilitate delivery of substances into sample chamber **2702**.

In certain embodiments, fluid reservoir **2716** includes a plurality of chambers, each of which is connected to fluid inlet **2718** via a manifold (not shown). Control unit **2728** can selectively deliver substances from any one or more of the multiple chambers into sample chamber **2702** by adjusting the manifold to ensure that the selected chambers are fluidically
25 connected to fluid inlet **2718**.

In general, control unit **2728** can be configured to introduce substances from fluid reservoir **2716** into sample chamber **2702** before, after, or both before and after, sample **2708** on first substrate **2706** has interacted with the array of features **2714** on first substrate **2712**. Many examples of such substances have been described previously. Examples of such substances
30 include, but are not limited to, permeabilizing agents, buffers, fixatives, staining solutions,

washing solutions, and solutions of various biological reagents (e.g., enzymes, peptides, oligonucleotides, primers).

To initiate interaction between sample **2708** and feature array **2714**, the sample and array are brought into spatial proximity. To facilitate this step, second holder **2710** – under the control
5 of control unit **2728** – can translate second substrate **2712** in any of the x-, y-, and z-coordinate directions. In particular, control unit **2728** can direct second holder **2710** to translate second substrate **2712** in the z-direction so that sample **2708** contacts, or nearly contacts, feature array **2714**.

In some embodiments, apparatus **2700** can optionally include an alignment sub-system
10 **2730**, which can be electrically connected to control unit **2728**. Alignment sub-system **2730** functions to ensure that sample **2708** and feature array **2714** are aligned in the x-y plane prior to translating second substrate **2712** in the z-direction so that sample **2708** contacts, or nearly contacts, feature array **2714**.

Alignment sub-system **2730** can be implemented in a variety of ways. In some
15 embodiments, for example, alignment sub-system **2730** includes an imaging unit that obtains one or more images showing fiducial markings on first substrate **2706** and/or second substrate **2712**. Control unit **2718** analyzes the image(s) to determine appropriate translations of second substrate **2712** in the x- and/or y-coordinate directions to ensure that sample **2708** and feature array **2714** are aligned prior to translation in the z-coordinate direction.

In certain embodiments, control unit **2728** can optionally regulate the removal of
20 substances from sample chamber **2702**. For example, control unit **2728** can selectively adjust valve **2722** so that substances introduced into sample chamber **2702** from fluid reservoir **2716** are directed into waste reservoir **2724**. In some embodiments, waste reservoir **2724** can include a reduced-pressure source (not shown) electrically connected to control unit **2728**. Control unit
25 **2728** can adjust the fluid pressure in fluid outlet **2720** to control the rate at which fluids are removed from sample chamber **2702** into waste reservoir **2724**.

In some embodiments, analytes from sample **2708** or from feature array **2714** can be selectively delivered to analysis apparatus **2726** via suitable adjustment of valve **2722** by control unit **2728**. As described above, in some embodiments, analysis apparatus **2726** includes a
30 reduced-pressure source (not shown) electrically connected to control unit **2728**, so that control unit **2728** can adjust the rate at which analytes are delivered to analysis apparatus **2726**. As such,

fluid outlet **2720** effectively functions as an analyte collector, while analysis of the analytes is performed by analysis apparatus **2726**. It should be noted that not all of the workflows and methods described herein are implemented via analysis apparatus **2726**. For example, in some embodiments, analytes that are captured by feature array **2714** remain bound to the array (i.e.,
5 are not cleaved from the array), and feature array **2714** is directly analyzed to identify specifically-bound sample components.

In addition to the components described above, apparatus **2700** can optionally include other features as well. In some embodiments, for example, sample chamber **2702** includes a heating sub-system **2732** electrically connected to control unit **2728**. Control unit **2728** can
10 activate heating sub-system **2732** to heat sample **2708** and/or feature array **2714**, which can help to facilitate certain steps of the methods described herein.

In certain embodiments, sample chamber **2702** includes an electrode **2734** electrically connected to control unit **2728**. Control unit **2728** can optionally activate electrode **2734**, thereby establishing an electric field between the first and second substrates. Such fields can be
15 used, for example, to facilitate migration of analytes from sample **2708** toward feature array **2714**.

In some of the methods described herein, one or more images of a sample and/or a feature array are acquired. Imaging apparatus that is used to obtain such images can generally be implemented in a variety of ways. **FIG. 27B** shows one example of an imaging apparatus **2750**.
20 Imaging apparatus **2750** includes a light source **2752**, light conditioning optics **2754**, light delivery optics **2756**, light collection optics **2760**, light adjusting optics **2762**, and a detection sub-system **2764**. Each of the foregoing components can optionally be connected to control unit **2728**, or alternatively, to another control unit. For purposes of explanation below, it will be assumed that control unit **2728** is connected to the components of imaging apparatus **2750**.

25 During operation of imaging apparatus **2750**, light source **2752** generates light. In general, the light generated by source **2752** can include light in any one or more of the ultraviolet, visible, and/or infrared regions of the electromagnetic spectrum. A variety of different light source elements can be used to generate the light, including (but not limited to) light emitting diodes, laser diodes, laser sources, fluorescent sources, incandescent sources, and
30 glow-discharge sources.

The light generated by light source **2752** is received by light conditioning optics **2754**. In general, light conditioning optics **2754** modify the light generated by light source **2752** for specific imaging applications. For example, in some embodiments, light conditioning optics **2754** modify the spectral properties of the light, e.g., by filtering out certain wavelengths of the light. For this purpose, light conditioning optics **2754** can include a variety of spectral optical elements, such as optical filters, gratings, prisms, and chromatic beam splitters.

In certain embodiments, light conditioning optics **2754** modify the spatial properties of the light generated by light source **2752**. Examples of components that can be used for this purpose include (but are not limited to) apertures, phase masks, apodizing elements, and diffusers.

After modification by light conditioning optics **2754**, the light is received by light delivery optics **2756** and directed onto sample **2708** or feature array **2714**, either of which is positioned on a mount **2758**. Light conditioning optics **2754** generally function to collect and direct light onto the surface of the sample or array. A variety of different optical elements can be used for this purpose, and examples of such elements include, but are not limited to, lenses, mirrors, beam splitters, and various other elements having non-zero optical power.

Light emerging from sample **2708** or feature array **2714** is collected by light collection optics **2760**. In general, light collection optics **2760** can include elements similar to any of those described above in connection with light delivery optics **2756**. The collected light can then optionally be modified by light adjusting optics **2762**, which can generally include any of the elements described above in connection with light conditioning optics **2754**.

The light is then detected by detection sub-system **2764**. Generally, detection sub-system **2764** functions to generate one or more images of sample **2708** or feature array **2714** by detecting light from the sample or feature array. A variety of different imaging elements can be used in detection sub-system **2764**, including CCD detectors and other image capture devices.

Each of the foregoing components can optionally be connected to control unit **2728** as shown in **FIG. 27B**, so that control unit **2728** can adjust various properties of the imaging apparatus. For example, control unit **2728** can adjust the position of sample **2708** or feature array **2714** relative to the position of the incident light, and also with respect to the focal plane of the incident light (if the incident light is focused). Control unit **2728** can also selectively filter both the incident light and the light emerging from the sample.

Imaging apparatus **2750** can typically obtain images in a variety of different imaging modalities. In some embodiments, for example, the images are transmitted light images, as shown in **FIG. 27B**. In certain embodiments, apparatus **2750** is configured to obtain reflection images. In some embodiments, apparatus **2750** can be configured to obtain birefringence images, fluorescence images, phosphorescence images, multiphoton absorption images, and more generally, any known image type.

In general, control unit **2728** can perform any of the method steps described herein that do not expressly require user intervention by transmitting suitable control signals to the components of sample handling apparatus **2700** and/or imaging apparatus **2750**. To perform such steps, control unit **2728** generally includes software instructions that, when executed, cause control unit **2728** to undertake specific steps. In some embodiments, control unit **2728** includes an electronic processor and software instructions that are readable by the electronic processor, and cause the processor to carry out the steps describe herein. In certain embodiments, control unit **2728** includes one or more application-specific integrated circuits having circuit configurations that effectively function as software instructions.

Control unit **2728** can be implemented in a variety of ways. **FIG. 27C** is a schematic diagram showing one example of control unit **2728**, including an electronic processor **2780**, a memory unit **2782**, a storage device **2784**, and an input/output interface **2786**. Processor **2780** is capable of processing instructions stored in memory unit **2782** or in storage device **2784**, and to display information on input/output interface **2786**.

Memory unit **2782** stores information. In some embodiments, memory unit **2782** is a computer-readable medium. Memory unit **2782** can include volatile memory and/or non-volatile memory. Storage device **2784** is capable of providing mass storage, and in some embodiments, is a computer-readable medium. In certain embodiments, storage device **2784** may be a floppy disk device, a hard disk device, an optical disk device, a tape device, a solid state device, or another type of writeable medium.

The input/output interface **2786** implements input/output operations. In some embodiments, the input/output interface **2786** includes a keyboard and/or pointing device. In some embodiments, the input/output interface **2786** includes a display unit for displaying graphical user interfaces and/or display information.

Instructions that are executed and cause control unit **2728** to perform any of the steps or procedures described herein can be implemented in digital electronic circuitry, or in computer hardware, firmware, or in combinations of these. The instructions can be implemented in a computer program product tangibly embodied in an information carrier, e.g., in a machine-readable storage device, for execution by a programmable processor (e.g., processor **2780**). The computer program can be written in any form of programming language, including compiled or interpreted languages, and it can be deployed in any form, including as a stand-alone program or as a module, component, subroutine, or other unit suitable for use in a computing environment. Storage devices suitable for tangibly embodying computer program instructions and data include all forms of non-volatile memory, including by way of example semiconductor memory devices, such as EPROM, EEPROM, and flash memory devices; magnetic disks such as internal hard disks and removable disks; magneto-optical disks; and CD-ROM and DVD-ROM disks. The processor and the memory can be supplemented by, or incorporated in, ASICs (application-specific integrated circuits).

Processor **2780** can include any one or more of a variety of suitable processors. Suitable processors for the execution of a program of instructions include, by way of example, both general and special purpose microprocessors, and the sole processor or one of multiple processors of any kind of computer or computing device.

VI. Methods of preparing an array for spatial analysis

A variety of methods are useful for generating an array of the present disclosure. Barcoded probes on an array can be generated through ligation of a plurality of oligonucleotides. In some instances, an oligonucleotide of the plurality contains a portion of a barcode, and the complete barcode is generated upon ligation of the plurality of oligonucleotides. For example, a first oligonucleotide containing a first portion of a barcode can be attached to a substrate (e.g., using any of the methods of attaching an oligonucleotide to a substrate described herein), and a second oligonucleotide containing a second portion of the barcode can then be ligated onto the first oligonucleotide to generate a complete barcode. The first oligonucleotide and the second oligonucleotide can be delivered to the array with a pre-determined spatial pattern. Additional oligonucleotides can be delivered. Different combinations of the first, second and any additional portions of a barcode can be used to increase the diversity of the barcodes. In instances where the

second oligonucleotide is also attached to the substrate prior to ligation, the first and/or the second oligonucleotide can be attached to the substrate via a surface linker.

To increase the diversity of the barcodes, a plurality of second oligonucleotides comprising two or more different barcode sequences can be ligated onto a plurality of first
5 oligonucleotides that comprise the same barcode sequence, thereby generating two or more different species of barcodes. To achieve selective ligation, a first oligonucleotide attached to a substrate containing a first portion of a barcode can initially be protected with a protective group (e.g., a photocleavable protective group), and the protective group can be removed prior to ligation between the first and second oligonucleotide.

10 In instances where the barcoded probes on an array are generated through ligation of two or more oligonucleotides, a concentration gradient of the oligonucleotides can be applied to a substrate such that different combinations of the oligonucleotides are incorporated into a probe depending on its location on the substrate. Barcoded probes on an array can also be generated by adding single nucleotides to existing oligonucleotides on an array, for example, using
15 polymerases that function in a template-independent manner. Single nucleotides can be added to existing oligonucleotides in a concentration gradient, thereby generating probes with varying length, depending on the location of the probes on the array.

A daughter array can be generated using a master array or a template array, where probes on the master array can be used as templates to synthesize probes for the daughter array. The
20 daughter probes can be synthesized on a substrate, or can be synthesized before attaching to a substrate. The master array can be any of the arrays provided herein. The probes generated on the daughter array can be the same or different from their corresponding template probes on the master array. For instance, a probe on the daughter array can share the same sequence as or be complementary to a corresponding probe on the master array. In some embodiments, the master
25 array can be used to generate multiple daughter arrays.

Arrays can also be prepared by modifying existing arrays, for example, by modifying the oligonucleotides attached to the arrays, including arrays obtained commercially. For instance, probes can be generated on an array that comprises oligonucleotides that are attached to the array at the 3' end and have a free 5' end. The oligonucleotides can be in situ synthesized
30 oligonucleotides, and can include a barcode. The length of the oligonucleotides can be less than 50 nucleotides (e.g., less than 45, 40, 35, 30, 25, 20, 15, or 10 nucleotides). To generate probes

using these oligonucleotides, a primer complementary to a portion of an oligonucleotide (e.g., a constant sequence shared by the oligonucleotides) can be used to hybridize with the oligonucleotide and extend (using the oligonucleotide as a template) to form a duplex and to create a 3' overhang. The 3' overhang thus allows additional nucleotides or oligonucleotides to be added onto the duplex. A capture probe can be generated by, for instance, adding one or more oligonucleotides to the end of the 3' overhang (e.g., via splint oligonucleotide mediated ligation), where the added oligonucleotides can include the sequence or a portion of the sequence of a capture domain.

In instances where the oligonucleotides on an existing array include a recognition sequence that can hybridize with a splint oligonucleotide, probes can also be generated by directly ligating additional oligonucleotides on to the existing oligonucleotides via the splint oligonucleotide. The recognition sequence can be located at the free 5' end or the free 3' end of an oligonucleotide on the existing array. Recognition sequences useful for the methods of the present disclosure can include high contents of Guanine and Cytosine nucleotides and thus have high stability.

Bead arrays can be generated by attaching beads (e.g., barcoded beads) to a substrate in a random or non-random pattern. Beads can be attached to selective regions on a substrate by, e.g., selectively activating regions on the substrate to allow for attachment of the beads. Activating selective regions on the substrate can include activating a coating (e.g., a photocleavable coating) or a polymer that is applied on the substrate. Beads can be attached iteratively, e.g., a subset of the beads can be attached at one time, and the same process can be repeated to attach the remaining beads. Alternatively, beads can be attached to the substrate all in one step.

Barcoded beads, or beads comprising a plurality of barcoded probes, can be generated by first preparing a plurality of barcoded probes on a substrate, depositing a plurality of beads on the substrate, and generating probes attached to the beads using the probes on the substrate as a template.

A probe on an array generated using any of the methods provided herein can be further modified. For example, oligonucleotides or some other functional groups (e.g., linkers, labels) can be added to the probe. In some embodiments, cleavage domain, functional domain, spatial barcode, UMI, and/or capture domain are added.

Also provided herein are methods of using an array generated as described herein for various methods as described in the present disclosure. For example, in some embodiments, an array generated by a method described herein can be used for profiling at least one biological analyte present in a biological sample. In some embodiments, the method involves contacting a biological sample with an array produced by any of the methods described herein; releasing a biological analyte from the biological sample, wherein the biological analyte is bound by a capture probe at a distinct spatial position of the substrate; detecting the bound biological analyte; and correlating the biological analyte with a barcode of the capture probe at the distinct spatial position of the substrate, thus profiling at least one biological analyte as having been present in the biological sample at a distinct spatial position.

VII. Methods of generating a daughter array using a master array

The present disclosure provides methods of generating a second array (daughter array) using a first array (master array) as a template. Upon generation of a daughter array, the master array can be re-used for generating further arrays. Accordingly, the master array can be used as a template to generate one or more daughter arrays.

The first array can be any of the arrays described herein, or generated by any of the methods described herein. An exemplary first array can include a first substrate comprising a plurality of probes. The first substrate can be any of the substrates described herein. A probe of the plurality can include, for example, a spatial barcode, a functional domain, a capture domain, a cleavage domain, and/or a UMI. The spatial barcode can correspond to a particular region on the array. In instances where two or more probes are associated with the same region, the two or more probes can share a common (e.g., identical) spatial barcode, or can have different spatial barcodes, or a combination of both. The sequence of a spatial barcode can include about 6 to about 25 or more nucleotides (e.g., at least or about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides). The length of a spatial barcode sequence can be, e.g., at least or about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides. In some instances, the length of a spatial barcode sequence can be, e.g., at most about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides. A probe of the plurality can be attached to the first substrate via its 3' end or 5' end. For example, a probe of the plurality can be attached to the first substrate via its 3' end and includes, from 3' to

5', a capture domain (e.g., a poly(T) sequence), and a spatial barcode, and an additional sequence at the 5' end.

The first array can also include a plurality of first features on the substrate, where the plurality of probes are attached to the features. A first feature can have a diameter or maximum dimension of between 100 nm and 100 μ m (e.g., between 100 nm and 1 μ m, 100 nm and 10 μ m, 100 nm and 20 μ m, 100 nm and 30 μ m, 100 nm and 40 μ m, 100 nm and 50 μ m, 100 nm and 60 μ m, 100 nm and 70 μ m, 100 nm and 80 μ m, 100 nm and 90 μ m, 90 μ m and 100 μ m, 80 μ m and 100 μ m, 70 μ m and 100 μ m, 60 μ m and 100 μ m, 50 μ m and 100 μ m, 40 μ m and 100 μ m, 30 μ m and 100 μ m, 20 μ m and 100 μ m, 10 μ m and 100 μ m, 1 μ m and 100 μ m, 900 nm and 100 μ m, 800 nm and 100 μ m, 700 nm and 100 μ m, 600 nm and 100 μ m, 500 nm and 100 μ m, 400 nm and 100 μ m, 300 nm and 100 μ m, or 200 nm and 100 μ m). The plurality of first features can be beads that have a diameter or maximum dimension of no larger than 100 μ m (e.g., no larger than 95 μ m, 90 μ m, 85 μ m, 80 μ m, 75 μ m, 70 μ m, 65 μ m, 60 μ m, 55 μ m, 50 μ m, 45 μ m, 40 μ m, 35 μ m, 30 μ m, 25 μ m, 20 μ m, 15 μ m, 14 μ m, 13 μ m, 12 μ m, 11 μ m, 10 μ m, 9 μ m, 8 μ m, 7 μ m, 6 μ m, 5 μ m, 4 μ m, 3 μ m, 2 μ m, or 1 μ m). The first array can include between 4000 and 10,000 features, or any range within 4000 to 6000 features. Alternatively, the first array can include at least 10,000 features (e.g., at least 100,000 features, at least 1,000,000 features, at least 5,000,000 features, or at least 10,000,000 features). The features of the first array can be arranged in a pattern. In some instances, the center of a feature on the first array is between 1 μ m and 100 μ m (e.g., between 20 μ m to 40 μ m, 20 μ m to 60 μ m, 20 μ m to 80 μ m, 80 μ m to 100 μ m, 60 μ m to 100 μ m, or 40 μ m to 100 μ m) from the center of another feature of the first array.

The first array can be a "high resolution" array, and can include features having an average diameter of about 1 μ m to about 10 μ m. The first array can additionally or alternatively have a high-density pattern, produced by compacting or compressing features together in one or more dimensions.

The spatial barcode in a probe on the first array can correspond to the location of the corresponding feature on the array. In instances where a first feature includes two or more probes, the two or more probes can include a common (e.g., identical) spatial barcode, or can have different spatial barcodes, or a combination of both. Probes attached to a feature can include multiple sets of probes. Probes of a given set can include identical spatial barcodes. The identical spatial barcodes can be different from spatial barcodes in the probes of another set.

A second array generated by the presently disclosed methods can include a second substrate comprising probes that are generated using the probes on the first array as a template. The second substrate can be any of the substrates described herein. The probes on the second array can have substantially the same relative positions as the corresponding template probes on the first array. The density of the probes on the second array can be about equal, higher than, or lower than the density of the probes on the first array. A probe of the second array can include a sequence that is at least 80% (e.g., at least 82%, 84%, 86%, 88%, 90%, 92%, 94%, 96%, or 98%) identical to a sequence in a corresponding template probe on the first array. A probe of the second array can also include a sequence that is complementary to a sequence in a corresponding template probe on the first array. A probe of the second array can include a spatial barcode that is the same as or is complementary to the spatial barcode in a corresponding template probe on the first array.

A second array generated by the presently disclosed methods can also include a plurality of second features on the substrate, where the probes are attached to the features. In instances where the first array includes a plurality of first features, and the second array includes a plurality of second features, the density of the first features and second features can be about equal or different. For example, the density of the second features on the second substrate can be lower than the density of the first features on the first array.

Probes of the second array can be synthesized on the second substrate, e.g., through the extension of primers that are attached, directly or indirectly, to the second substrate (or to the features on the substrate). Alternatively, the probes of the second array can be synthesized prior to being attached, directly or indirectly, to the second substrate (or to the features on the substrate).

In synthesizing a probe on the second substrate, a primer attached to the substrate (or to a feature on the substrate) can be used as an extension primer to generate an oligonucleotide that includes a common sequence as a corresponding template probe on the first array. In some instances, the corresponding template probe on the first array can be attached to the first substrate (or to a first feature on the first substrate) at the 3' end, and includes from 3' to 5', a capture domain (e.g., a poly(T) sequence), a spatial barcode, and a sequence shared with a primer on the second substrate. The second substrate can include a plurality of primers, and the density of the primers can be about equal, higher, or lower than that of the probes on the first array. The

primers can have a length of about 3 to about 35 nucleotides (e.g., about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, or 34 nucleotides). In some instances, the plurality of primers can include a common sequence (e.g., a primer sequence). To replicate the sequence of the corresponding probe, a primer can e.g., share
5 a portion of the sequence of the corresponding probe on the first array, and thus can hybridize with a complementary strand of the corresponding probe. The primers can be of sufficient length to provide for hybridization to the complementary strand. The portion of the sequence can have a length of about 5 to about 25 nucleotides (e.g., about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 nucleotides). The portion of the sequence can be, can include, or can be
10 part of a functional domain of the probe. In instances where the probe has a free 5' end, the portion of the sequence can be a sequence at the 5' end of the probe.

The second array can also include a plurality of second features on the substrate, where the plurality of probes (e.g., primers) are attached to the features. A second feature can have a diameter or maximum dimension of between 100 nm and 100 μ m (e.g., between 100 nm and 1
15 μ m, 100 nm and 10 μ m, 100 nm and 20 μ m, 100 nm and 30 μ m, 100 nm and 40 μ m, 100 nm and 50 μ m, 100 nm and 60 μ m, 100 nm and 70 μ m, 100 nm and 80 μ m, 100 nm and 90 μ m, 90 μ m and 100 μ m, 80 μ m and 100 μ m, 70 μ m and 100 μ m, 60 μ m and 100 μ m, 50 μ m and 100 μ m, 40 μ m and 100 μ m, 30 μ m and 100 μ m, 20 μ m and 100 μ m, 10 μ m and 100 μ m, 1 μ m and 100 μ m, 900 nm and 100 μ m, 800 nm and 100 μ m, 700 nm and 100 μ m, 600 nm and 100 μ m, 500 nm and
20 100 μ m, 400 nm and 100 μ m, 300 nm and 100 μ m, or 200 nm and 100 μ m). The plurality of second features can be beads that have a diameter or maximum dimension of no larger than 100 μ m (e.g., no larger than 95 μ m, 90 μ m, 85 μ m, 80 μ m, 75 μ m, 70 μ m, 65 μ m, 60 μ m, 55 μ m, 50 μ m, 45 μ m, 40 μ m, 35 μ m, 30 μ m, 25 μ m, 20 μ m, 15 μ m, 14 μ m, 13 μ m, 12 μ m, 11 μ m, 10 μ m, 9 μ m, 8 μ m, 7 μ m, 6 μ m, 5 μ m, 4 μ m, 3 μ m, 2 μ m, or 1 μ m). The second array can include
25 between 4000 and 10,000 features, or any range within 4000 to 6000 features. Alternatively, the second array can include at least 10,000 features (e.g., at least 100,000 features, at least 1,000,000 features, at least 5,000,000 features, or at least 10,000,000 features). The features of the second array can be arranged in a pattern. In some instances, the center of a feature on the second array is between 1 μ m and 100 μ m (e.g., between 20 μ m to 40 μ m, 20 μ m to 60 μ m, 20
30 μ m to 80 μ m, 80 μ m to 100 μ m, 60 μ m to 100 μ m, or 40 μ m to 100 μ m) from the center of another feature of the second array.

A strand complementary to the probe on the first array can be generated and released to allow hybridization with a primer on the second substrate (e.g., via the portion of sequence shared by the primer and the probe). The complementary strand can be released by applying heat to facilitate de-hybridization between the complementary strand and the probe. The first array and the second substrate can be aligned such that hybridization between the complementary strand and the primer can occur. To facilitate the migration of the released complementary strand, an electric field can be applied to the first array and the substrate. Upon hybridization between the primer and the complementary strand, primer extension can be carried out using the complementary strand as a template, thereby generating a probe on the second array. Following primer extension, the complementary strand can optionally be released from the newly synthesized probe (e.g., using heat).

In instances where the first array includes a plurality of first features, and the second substrate includes a plurality of second features, the first features of the first array and the second features of the second substrate can be uniformly positioned or non-uniformly positioned. The relative positions of the second features can be substantially the same or different from the relative positions of the first features. The second features can be arranged on the second substrate such that a second feature is in sufficient proximity to at least one first feature when the first array and the second substrate are aligned, such that at least one primer on the second feature interacts with a complementary released from the first feature for a sufficient period of time to allow hybridization.

To synthesize a probe of the second array prior to attaching the probe to a substrate, a primer is provided that can hybridize with a portion of the sequence of a template probe on the first array. The primer can include any sequence that is capable of hybridizing with the template probe on the first array. The portion of the sequence can be, can include, or can be a part of, a functional domain of the probe. The primer can be of sufficient length to provide for hybridization to the template probe. The portion of the sequence can have a length of about 3 to about 35 nucleotides (e.g., about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, or 34 nucleotides). Depending on the master array used, the primer can hybridize with the probe at the free 3' end, the free 5' end, or at a portion of the probe that is between the 3' end and the 5' end. Upon hybridization, primer extension can be carried out to generate a complementary strand of the probe. Following primer extension, the

second substrate can be provided and the newly synthesized strand can be attached to the second substrate using any of the methods described herein or known in the art suitable for attaching an oligonucleotide to a substrate. The density and/or number of the probes on the second array generated using the presently disclosed methods can be about equal to, higher, or lower than those of the first array.

FIGS. 22 and 23 show exemplary schematics of some of the methods described herein. As shown in **FIG. 22**, a daughter array that includes a plurality of primers on a substrate is provided. The primers can be attached to a plurality of features (e.g., immobilized beads). A master array that includes a plurality of probes on a substrate is also provided. In some instances, the probes can be attached to a plurality of features (e.g., immobilized beads). The probes on the master array can have known spatial barcodes. The relationship between the location of the probes on the master array and the spatial barcodes is readily known or can be determined by the methods described herein. As shown in **FIG. 23**, the probes on the master array can further include a primer sequence at the 5' end and a poly-T sequence at the 3' end, with the 3' end of the probes being attached to the substrate. The daughter array is aligned with the master array, and an intermediate oligonucleotide complementary to the probe on the master array is generated, where the intermediate oligonucleotide contains a poly-A sequence at the 5' end and a sequence at the 3' end that can hybridize with the primer on the daughter array. Upon hybridization, the primer on the daughter array is then extended to generate a probe that contains the known spatial barcode of the probe from the master array. The intermediate oligonucleotide can optionally be released from the newly generated probe.

Figures 24A-24D show further exemplary schematics of some of the methods described herein. A plurality of master oligonucleotides can be spotted to the surface of a substrate, and/or can be attached to the substrate using a surface linker. An exemplary master oligonucleotide can be attached to the substrate at the 5' end, and have a free 3' end. In the priming step, a number of primers are provided, wherein the primers have surface linkers at one end and are complementary to a portion of a master oligonucleotide (e.g., to a sequence located at the 3' end of the master oligonucleotide). Upon hybridization between the primer and the master oligonucleotide, the primer is used for nucleic acid extension to generate a complementary strand of the master oligonucleotide. Following extension, a copy-surface (or copy-slide) is provided and the extended complementary strand can be attached to the copy-surface. For example, the 5'

end of the extended complementary strand can be attached to the copy-surface. The result is a daughter array based on the master array. Daughter arrays generated by the presently disclosed methods can have a relative spatial alignment between the probes that corresponds to a relative spatial alignment of the probes on the master array. This process can be repeated multiple times to make multiple daughter arrays using the master array.

VIII. Fiducial Markers

In some embodiments, imaging can be performed using one or more fiducial markers.

Fiducial markers can be any one or more objects placed in the field of view of an imaging system and that are captured in an image of a sample, an array, a substrate, or any of the other aspects described herein. Fiducial markers are typically used as a point of reference or measurement scale for alignment (e.g., to align a sample and an array, to align two substrates, to determine a location of a sample or array on a substrate relative to a marker) and/or for quantitative measurements of sizes and/or distances.

Fiducial markers can be formed from materials that include, but are not limited to, metallic, reflective, fluorescent, radioactive, chemiluminescent, and colorimetric labels. In some embodiments, a fiducial marker can be present on a substrate to provide orientation of the biological sample. The use of fiducial markers to stabilize and orient biological samples is described, for example, in U.S. Pat. No. 7,848,553, and in U.S. Pat. Pub. 2003/0153850, both of which are herein incorporated by reference in their entireties.

FIGS. 25A-25D show an example substrate **2500** having a surface **2501** that includes four arrays. Specifically, substrate **2500** includes a first array **2504a**, a second array **2504b**, a third array **2504c**, and a fourth array **2504d**, wherein each array is configured to receive a biological sample **2528** within a defined perimeter of the array. That is, the first array **2504a**, the second array **2504b**, the third array **2504c**, and the fourth array **2504d** are arrays of features as described above.

Fiducial markers **2532**, arranged on the surface **2501** of the substrate **2500** in the form of a frame **2534**, outline the outer perimeter of each array **2504a-2504d**, so that each array is defined by a plurality of fiducial markers **2532** as shown in **FIGS. 25B-D**. The frame **2534**

provides a visual guide that allows for proper placement of the biological sample **2528** on one of the arrays (or, allows for proper placement of samples on multiple arrays).

In general, substrate **2500** can be any of the substrates described herein, and fiducial markers **2532** can be arranged on any surface of substrate **2500**. Further, biological sample **2528** can generally be any of the different types of biological samples described herein, including (but not limited to) tissue sections.

Fiducial markers **2532** can be fabricated using a variety of techniques. Such techniques include, but are limited to, jet printing, screen printing, sand-blasting, etching, vapor deposition, photomasking, and various other deposition processes.

In some embodiments, substrate **2500** can optionally include a label **2502** that provides identification information such as, but not limited to, sample name and/or sample number. Label **2502** can be implemented as a bar code label, a laser-scribed identifier, an electronic identification tag, and other types of labels and/or combinations thereof.

FIG. 25B shows an enlarged view of first array **2504a**. First array **2504a** includes a plurality of fiducial markers **2532** arranged in multiple patterns to form frame **2534**. Frame **2534** provides a visual indicator for placing the biological sample **2528** in a suitable position such that the sample will be positioned on the array **2504a** within the perimeter defined by frame **2534**.

FIGS. 25C-D show enlarged views of frame **2534** or portions thereof (for illustrative purposes, remaining portions of array **2504a** have been excluded). Frame **2534** includes a first side **2520**, a second side **2522**, a third side **2524**, and a fourth side **2526**. Each of the sides of frame **2534** have a distinct fiducial marker pattern. For example, the first side **2520** includes fiducial markers arranged in three staggered rows **2506** (depicted in dotted sections in **FIG. 25D**) that form a two-dimensional lattice structure. In various embodiments, each row may be positioned relative to a neighboring row in either a stacked or staggered arrangement.

In some embodiments, a row **2506** of fiducial markers **2532** can include adjacent markers that are spaced apart by a distance e , as shown in **FIG. 25D**. In some embodiments, the marker-to-marker distance is the distance between the geometric centers of two adjacent markers. In some embodiments, a portion of or all of the markers **2532** of row **2506** are equidistant from one or more adjacent markers. In some embodiments, at least one or more markers **2532** can be positioned at different distances from one or more adjacent markers.

In some embodiments, some or all fiducial markers **2532** of a row can be arranged in a linear geometrical pattern. In some embodiments, some or all fiducial markers **2532** of a row can be arranged in a non-linear geometrical pattern, for example, an alternating staggered configuration where one fiducial marker is displaced from a successive fiducial marker in two orthogonal coordinate directions (i.e., the x- and y-directions), as shown in fourth side **2526**.

Still referring to **FIGS. 25C-D**, first side **2520** has three rows of fiducial markers **2532** arranged in an intercalated pattern. Similarly, the second side **2522**, third side **2524**, and fourth side **2526** include fiducial marker arranged in geometrical patterns that differ from the pattern of first side **2520**. In some embodiments, the first array **2504a**, the second array **2504b**, the third array **2504c**, and the fourth array **2504d** (shown in **FIG. 25A**) each have a different frame formed of fiducial markers (note that distinct features of the frames for the second, third, and fourth arrays are not shown in **FIG. 25A**) that enables each array to be differentiated based on the pattern geometry of its corresponding fiducial marker frame.

In some embodiments, each side of a single frame contains a unique pattern of fiducial markers that allows a particular side to be automatically differentiated from the other sides in the frame to facilitate determination of an array orientation. Individual sides can be manually identified or differentiated from one another by a human user, or alternatively, specific sides can be automatically identified or differentiated from one another by capturing an image of the frame, and analyzing the fiducial markings in the image to correlate specific patterns of fiducial markings with particular frame sides using an electronic processor executing software-based instructions. In some embodiments, the pattern of fiducial markers **2532** includes multiple repeated pattern segments to make software-based pattern recognition easier, even when some of the fiducial markers are obscured by the biological sample.

The pattern of fiducial markers forming the frame can further include glyphs in the corners of the frame, as shown in **FIGS. 25B-C**. The first glyph **2512** is shown to be rectangular in shape and can be positioned in the corner between the first side **2520** and the fourth side **2526**. Furthermore, the second glyph **2514** is shown as having a filled hexagonal shape and can be positioned in the corner between the first side **2520** and the second side **2522**. Similarly, third glyph **2516** is shown to be triangular in shape and can be positioned in the corner between the fourth side **2526** and the third side **2524**. The fourth glyph **2518** is shown as having an empty hexagonal shape and can be positioned in the corner between the second side **2522** and the third

side **2524**. While the glyphs illustrated in **FIGS. 25B-C** have specific patterns for illustrative purposes, it should be understood that in general, glyphs corresponding to any specific patterns or shapes of fiducial markers can be positioned at any location on a substrate. The examples shown in the figures are in no way intended to imply limitations.

5 In general, a “glyph” is a distinct pattern of fiducial markers that forms a geometric shape. Examples of such shapes include, but are not limited to, a circle, an hourglass, a hexagon, a square, a rectangle, a triangle, a pentagon, a heptagon, an octagon, a nonagon, a decagon, an ellipse, a regular polygon, and/or any other regular or irregular shape. Glyphs can also correspond to combinations of any of the above shapes. Glyphs can further correspond to
10 “empty” or “filled” shapes.

 In some embodiments, glyphs can be observed in an image (e.g., a bright-field microscope image) and identified by the a human user in a manual analysis process. In some embodiments, fiducial markers forming glyphs and frames can include a colorant or label (including any of the labels described above) that allows the pattern of fiducial markers defining
15 the glyph or frame to be visible to the naked eye, with or without the aid of magnification (e.g., in an optical microscope). In certain embodiments, fiducial markers can include one or more fluorescent species so that glyphs and frames can be excited by incident light and fluorescence emission from the fiducial markers can be observed and imaged.

 In some embodiments, the one or more fiducial markers **2532** can be printed or deposited
20 on a surface of a substrate. For example, in some embodiments, the one or more fiducial markers **2532** can be printed using contact or non-contact printing. In another example, the one or more fiducial markers **2532** can be fabricated lithographically. In some embodiments, the substrate is manufactured with carboxyl groups exposed on the surface and the fiducial marker includes a reactive chemical group (e.g., an amine group) that forms a covalent bond to a desired functional
25 group on the substrate surface.

 In some examples, one or more fiducial markers **2532** includes metallic nanoparticles. Suitable metallic nanoparticles include, but are not limited to, gold nanoparticles. In some embodiments, the metallic nanoparticles have an average diameter of between 5 nm and 1 micron (e.g., between 10 nm and 1 micron, between 20 nm and 1 micron, between 20 nm and
30 800 nm, between 20 nm and 600 nm, between 20 nm and 400 nm, between 30 nm and 100 nm).

In some embodiments, the metallic nanoparticles are coated with a coating material. Suitable coating materials include, for example, a variety of polymer materials. In some embodiments, metallic nanoparticles can be linked to polymer ligands. For example, in **FIG. 25C**, gold nanoparticle **2530** is linked to a polymer ligand having a length of approximately 50 nm.

Polymer ligands linked to metallic nanoparticles, and the metallic nanoparticles themselves, can be functionalized in various ways. In some embodiments, for example, metallic nanoparticles are linked to polymer ligands having an amine group at a terminus. In some embodiments, a metallic nanoparticle (such as gold nanoparticle **2530**) can be functionalized with a chemical compound having a carboxylic acid group on its terminus that covalently attaches to the amine group associated with the coated gold nanoparticle **2530**. Moreover, in some embodiments, metallic nanoparticles are fluorescent.

In some embodiments, the fiducial markers are formed of a metal material such as chromium and can be applied by using chemical deposition. For example, a pattern of discrete markers or a thin film/layer composed of a metal such as chromium can be electroplated onto the substrate to create a visible frame for each of the arrays on the substrate. In some embodiments, the fiducial markers are laser etched onto the surface of the substrate. In certain embodiments, the fiducial markers are formed from TubeWriter™ 360 ink, permanent marker ink, pen ink, inkjet printer ink, or combinations thereof. In some embodiments, fiducial markers are formed from colored and/or fluorescent microspheres or microbeads.

In some embodiments, fiducial markers are formed from amine-functionalized iron oxide nanoparticles. In some embodiments, fiducial markers are formed from amine-functionalized thulium doped upconversion nanophosphors. In some embodiments, fiducial markers are formed from amine-functionalized quantum dots, such as but not limited to, cadmium-based quantum dots. In some embodiments, fiducial markers include colorimetric labels such as fluorescent oligonucleotides (e.g., oligonucleotides with fluorescent moieties).

In general, fiducial markers can have a variety of two-dimensional shapes when viewed in a plane parallel to the substrate. For example, in some embodiments, the fiducial markers **2532** have the shape of a circular spot, as shown in **FIGS. 25A-D**. In some embodiments, the fiducial markers **2532** are provided in the shape of a hexagon, a square, a rectangle, a triangle, a pentagon, a heptagon, an octagon, a nonagon, a decagon, an ellipse, a regular polygon, or any

other regular or irregular geometric shape. In some embodiments, some or all of the fiducial markers of a frame have an identical shape. In some embodiments, some of the fiducial markers of a frame have a first shape, and some of the fiducial markers have a second shape different from the first shape.

5 In some embodiments, fiducial markers can surround all or a portion of an array. In some embodiments, the fiducial markers can completely surround the array. Alternatively, in some embodiments, the fiducial markers may not completely surround the array (e.g., may only surround a portion of the array). In some embodiments, the fiducial markers identify the corners or other specific portions of an array, such as the center of the array.

10 In some embodiments, the average maximum cross-sectional dimension of fiducial markers that form a frame is between 50 microns and 500 microns (e.g., between 50 microns and 400 microns, between 50 microns and 300 microns, between 100 microns and 500 microns, between 100 microns and 300 microns).

In some embodiments, fiducial markers are arranged in such a way that the center of one
15 fiducial marker is between 100 microns and 200 microns (e.g., between 110 microns and 190 microns, between 120 microns and 170 microns, between 120 microns and 150 microns) from the center of a closest next fiducial marker surrounding an array.

In some embodiments, the array with the surrounding fiducial markers has a maximum dimension on the substrate of 10 mm or less (e.g., 8 mm or less, 6 mm or less, 5 mm or less, 4
20 mm or less, 3 mm or less). In some embodiments, the array is a square or rectangular array with a side length of about 8 mm.

In some embodiments, a fiducial marker can be present on a substrate to provide orientation of the biological sample. In some embodiments, a microsphere can be coupled to a substrate to aid in orientation of the biological sample. In some examples, a microsphere
25 coupled to a substrate can produce an optical signal (e.g., fluorescence). In another example, a microsphere can be attached to a portion (e.g., corner) of an array in a specific pattern or design (e.g., hexagonal pattern) to aid in orientation of a biological sample on the array of features on a substrate.

In some embodiments, a fiducial marker can be an immobilized molecule with which a
30 detectable signal molecule can interact to generate a signal. For example, a marker nucleic acid can be linked or coupled to a chemical moiety capable of fluorescing when subjected to light of a

specific wavelength (or range of wavelengths). Such a marker nucleic acid molecule can be contacted with an array before, contemporaneously with, or after the tissue sample is stained to visualize or image the tissue section. Although not required, it can be advantageous to use a marker that can be detected using the same conditions (e.g., imaging conditions) used to detect a labelled cDNA.

In some embodiments, fiducial markers are included to facilitate the orientation of a tissue sample or an image thereof in relation to an immobilized capture probe on a substrate. Any number of methods for marking an array can be used such that a fiducial marker is detectable only when a tissue section is imaged. For instance, a molecule, e.g., a fluorescent molecule that generates a signal, can be immobilized directly or indirectly on the surface of a substrate.

In some embodiments, a fiducial marker can be randomly located on a substrate surface. For example, an oligonucleotide containing a fluorophore can be randomly printed, stamped, synthesized, or attached to a substrate (e.g., a glass slide) at a random position on the substrate. A tissue section can be contacted with the substrate such that the oligonucleotide containing the fluorophore contacts, or is in proximity to, a cell from the tissue section or a component of the cell (e.g., an mRNA or DNA molecule). An image of the substrate and the tissue section can be obtained, and the position of the fluorophore within the tissue section image can be determined (e.g., by reviewing an optical image of the tissue section overlaid with the fluorophore detection).

In some embodiments, fiducial markers can be applied to known, selected locations on a substrate. For example, fiducial markers can be stamped, attached, or synthesized on the substrate. An image of the substrate that includes the fiducial marker is obtained, and the position of the fiducial marker on the substrate can be confirmed qualitatively by viewing the image, and/or quantitatively by measuring the location of the fiducial marker in the image.

As described above, a single substrate (e.g., a glass slide) can include multiple framed arrays, wherein at least one frame has a unique pattern that allows a particular array to be differentiated from the other arrays on the substrate. In particular, the pattern of fiducial markers of at least one frame can contain a slight variation as compared to the other frames. For example, **FIG. 26** shows a first array **2602** surrounded by a first frame **2610**, a second array **2604** surrounded by a second frame **2612**, a third array **2606** surrounded by a third frame **2616**, and a

fourth array **2608** surrounded by a fourth frame **2614**. In some embodiments, a specific motif in the pattern of fiducial markers can be varied from frame to frame so as to differentiate individual arrays within a same slide. As shown in **FIG. 26**, the region corresponding to a first top motif **2618a** in frame **2610** can be varied by altering the number of fiducial markers in counterpart
5 regions of frames **2612**, **2614**, and **2616**. By doing so, the second top motif **2618b**, the third top motif **2618c**, and the fourth top motif **2618d** include unique, distinguishable patterns of fiducial markers that can be used to differentiate each array.

In some embodiments, one or more additional regions within any one or more sides of the frames and/or glyphs can be altered to generate unique patterns of fiducial markers within each
10 array. For example, a first lateral motif **2620a**, a second lateral motif **2620b**, a third lateral motif **2620c**, and a fourth lateral motif **2620d** can include unique patterns of fiducial markers in addition to, or as an alternative to, the unique patterns of fiducial markers of the top motifs of each frame. Such variation in the patterning of the fiducial markers can permit a user and/or analysis software to determine which of the four arrays is being scanned and/or evaluated. In
15 some embodiments, the end user and/or analysis software can also determine whether the array has “shifted” in orientation (e.g., rotated about an axis orthogonal to, or parallel to, the plane of the substrate), allowing for correction of the array rotation before sample analysis occurs.

IX. Additional embodiments

20 It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

Preparation of solid supports

25 The solid support typically is a glass, plastic or silicon chip or slide. Attachment of the capture molecule is done using known techniques in the art. See, for example, Seurnynck-Servoss et al., Anal Biochem. (2007) 371(1): 105–115 and Gerdts et al., Microarrays (2016) 5:16, for methods of immobilizing proteins to a solid support. Methods of immobilizing oligonucleotides
30 to a support are known in the art (beads: Dressman et al. (2003) Proc. Natl. Acad. Sci. USA 100:8817, Brenner et al. (2000) Nat. Biotech. 18:630, Albretsen et al. (1990) Anal. Biochem.

189:40, and Lang et al. Nucleic Acids Res. (1988) 16: 10861; nitrocellulose: Ranki et al. (1983) Gene 21:77; cellulose: Goldkom (1986) Nucleic Acids Res. 14:9171; polystyrene: Ruth et al. (1987) Conference of Therapeutic and Diagnostic Applications of Synthetic Nucleic Acids, Cambridge U.K.; teflon-acrylamide: Duncan et al. (1988) Anal. Biochem. 169:104;
5 polypropylene: Polsky-Cynkin et al. (1985) Clin. Chem. 31:1438; nylon: Van Ness et al. (1991) Nucleic Acids Res. 19:3345; agarose: Polsky-Cynkin et al., Clin. Chem. (1985) 31:1438; and sephacryl: Langdale et al. (1985) Gene 36:201; latex: Wolf et al. (1987) Nucleic Acids Res. 15:2911).

In some embodiments, the solid support is first coated with a polymer to enhance
10 attachment of the capture molecule. The solid support may be of any size required to mount a tissue section of the sample. For example, a glass microscope slide measures about 75 mm by 25 mm (3" by 1") and is about 1 mm thick. A range of other sizes are available for various special purposes, such as, e.g., 75 x 50 mm, 46 x 27 mm, and 48 x 28 mm, or may be produced for the specific assay. Any size solid support may be used. Functionalized solid supports may be
15 commercially available and can be adapted for the methods disclosed herein.

The solid support may be derivatized or coated to enhance attachment of the capture molecule. Methods for derivatizing or coating solid supports are known in the art and may depend on the nature of the capture molecule. For example, if the solid support is a glass slide then it may be derivatized with amines to facilitate the attachment of a protein, e.g., an antibody,
20 or a nucleic acid capture molecule.

In some embodiments, the solid support comprises surface linkers to attach the capture molecule to the solid support, i.e., an indirect attachment of the capture molecule to the solid support.

The solid support may comprise markings to provide guidance for correlating spatial
25 information with the characterization of the analyte of interest. In some embodiments, the solid support may be a microarray. In some embodiments, the markings comprise a grid. In some embodiments, the solid support comprises a visual grid. In some embodiments, the solid support may be physically separated into regions by, for example, canals, grooves and the like. In some embodiments, the solid support may be physically separated into regions by, for example,
30 chemical means such as a hydrophobic coating. Spatially defined regions may be "addressable"

in that its location and the identity of the immobilized probe at that location are known or determinable.

High Resolution Arrays

5 High density nucleic acid arrays or microarrays may be used in the methods described herein. A microarray is a multiplex technology used in molecular biology. A typical microarray consists of an arrayed series of microscopic spots (features or sites) of capture molecules (millions of spots can be incorporated on a single array and several arrays can be presented on a single solid support). The distinct (discrete) position of each capture molecule spot (each species
10 of capture molecule) is known as a "feature" or "site" (and hence in the methods provided herein each species of capture molecule may be viewed as a specific feature of the array; each feature occupies a distinct (discrete) position on the array), and typically each separate feature contains in the region of picomoles (10-12 moles) of a specific, e.g., DNA sequence (a "species"), which are known as "probes" (or "reporters"). Typically, these can be a short section of a gene or other
15 nucleic acid element to which an oligonucleotide from the sample (or "target") can hybridize under high-stringency hybridization conditions.

For creating an accurate "map" of the analyte of interest in the sample, it is desirable to have a high resolution array, i.e., a high density of capture molecules per unit area (e.g., μm^2). In some embodiments, there is a plurality of capture molecules. In some embodiments, there is a
20 plurality of more than 1, 2, 3, 4, 5 or more species of capture molecules.

In certain embodiments, capture molecules are immobilized via one or more of the cleavable linkers described herein.

In some embodiments, the capture molecules, e.g., oligonucleotides, may be synthesized directly on the solid substrate. Suitable methods for such an approach are known to the person
25 skilled in the art. For instance, features (i.e., clusters of capture molecules) may be made by a solid phase amplification method known as bridge amplification. Other examples including manufacture techniques developed by Agilent Inc., Affymetrix Inc., Roche Nimblegen Inc. or Flexgen BV. Typically, lasers and a set of mirrors that specifically activate the spots where nucleotide additions are to take place are used. Such an approach may provide, for example,
30 spot sizes (i.e., features) of around $30\mu\text{m}$ or larger.

Master-Slide Strategy

In one aspect, the present invention provides methods for making a manufacturing master-surface (or master-slide) on a solid support for spatial analysis. The master-slide can be used as a template for creating printed copy consumable slides, e.g., copy-slides. The master-surface includes a number of spatial analysis barcodes characterized by high barcode diversity. The method of making the master-surface include, without limitation, spotting and printing of oligonucleotides onto the surface. FIG. 24A-D depict an example of the process for manufacturing a master-surface. A plurality of master oligonucleotides may be spotted to the surface of a substrate using a surface linker to bind each master oligonucleotide to the solid support or surface. In the priming step, a number of copy oligonucleotides are provided, wherein the copy oligonucleotides have surface linkers at one end and a priming region at the other end, wherein the priming region is complementary (at least in part) to a region of one or more master oligonucleotides. The priming regions of the copy oligonucleotides are permitted to prime to the master oligonucleotides. The primed configuration is used for nucleic acid extension to make complements of the master oligonucleotides, thereby providing duplex nucleic acid molecules from the original master oligonucleotides and the newly extended complements. Following extension, a copy-surface (or copy-slide) is provided and the surface linkers on the copy oligonucleotides are used to link the extended complements to the copy-surface. The result is a copy-slide based on the master-slide. This process can be repeated multiple times to make multiple copy-slides of the master-slide. The synthesized or extended nucleic acid sequences based on the master-slide are localized to the copy-slide. Alternately, the copy-slide could have bound universal oligonucleotides that can be contacted with the master-slide for priming and extension while the two slides are in contact (FIG. 24D).

Method of using a master array to build a daughter array

In some embodiments, the disclosure relates to methods of using a master array to build a daughter array. In some embodiments, provided herein is a master array that includes a plurality of features immobilized on a solid substrate (e.g. any of the solid support or solid substrate described herein), where each feature includes a plurality of tagging probes. In some embodiments, each tagging probe comprises a barcode (e.g. any of the spatial barcode described herein). In some embodiments, the plurality of tagging probes on each feature are identical (e.g.,

the same barcode sequence). In some embodiments, the tagging probes include one or more hybridization domains (e.g. a poly-T sequence or a poly-A sequence). In some embodiments, the tagging probes include a functional domain (e.g. any of the functional domains described herein). In some embodiments, the barcode of the tagging probes and the location of the plurality of features on the master array are known.

In some embodiments, provided herein is a daughter array that includes a plurality of features immobilized on a solid substrate (e.g. any of the solid supports or solid substrates described herein), where the plurality of features include a plurality of priming oligonucleotides (e.g. any of the priming oligonucleotides described herein). In some embodiments, the priming oligonucleotides are identical (e.g. the same nucleotide sequence).

In some embodiments, the methods described herein include aligning the master and daughter array, and generating an oligonucleotide that includes a sequence that is complementary to a tagging probe on the master array. In some embodiments, the methods described herein include releasing the complementary oligonucleotide from the tagging probe on the master array.

In some embodiments, the complementary oligonucleotide is released (e.g. de-hybridized) from the tagging probe on the master array using heat. In some embodiments, the released complementary oligonucleotide is allowed to diffuse from the master array to reach the daughter array. In some embodiments, the methods provided herein further include applying an electric field to the master and daughter arrays, where the electric field increases diffusion of the complementary oligonucleotide. In some embodiments, the methods provided herein include allowing the complementary oligonucleotide to hybridize with a priming oligonucleotide (e.g. any of the priming oligonucleotides described herein) on the daughter array. In some embodiments, the complementary oligonucleotide hybridizes with the priming oligonucleotide through a functional domain (e.g. any of the functional domains described herein). In some embodiments, the methods provided herein include extending the priming oligonucleotide on the daughter array using the complementary oligonucleotide as a template, thereby generating a tagging probe on the daughter array that includes an oligonucleotide sequence comprising a barcode. In some embodiments, the barcode is associated with a known feature (e.g. feature with a known location) on the master array.

In some embodiments, the density of the plurality of tagging probes on the master array is substantially the same as (e.g. equal to) the density of the plurality of tagging probes on the

daughter array. In some embodiments, the density of the plurality of tagging probes on the master array is higher than the density of the plurality of tagging probes on the daughter array. In some embodiments, the number of the plurality of tagging probes on the master array is substantially the same as (e.g. equal to) the number of the plurality of tagging probes on the daughter array. In some embodiments, the number of the plurality of tagging probes on the master array is higher than the number of the plurality of tagging probes on the daughter array.

Priming oligonucleotide

As used herein, the term “priming oligonucleotide” typically refers to a nucleic acid molecule (e.g. oligonucleotide) which includes a priming sequence (e.g., a primer for generating a nucleic acid molecule using a template). In some embodiments, the priming oligonucleotides are configured to hybridize with one or more functional domains on a probe. In some embodiments, the priming oligonucleotides can be attached to a feature (e.g. any of the features described herein) or a solid support (e.g. any of the solid supports or substrates described herein) at the 5' end or at the 3' end. In some embodiments, the priming oligonucleotides on two or more features (e.g. any of the features described herein) are identical.

Method for building a spatial array

To build a daughter array using a master array, a daughter array that includes a plurality of beads immobilized on a solid substrate is provided, which also includes a plurality of priming oligonucleotides immobilized on the beads. A master array is provided that includes a plurality of beads immobilized on a solid substrate, and further includes a plurality of tagging probes with known spatial barcodes immobilized on the beads. In addition, the location of the tagging probes on the master array is known. The tagging probes on the master array further contain a priming sequence at the 5' end and a poly-T sequence at the 3' end, with the 3' end of the tagging probes being attached to the beads.

The daughter array is aligned with the master array. An oligonucleotide complementary to a tagging probe on the master array is generated through PCR, where the oligonucleotide contains a poly-A sequence at the 5' end and is able to hybridize with the priming oligonucleotides on the daughter array through the priming sequence. The newly generated oligonucleotide is released and allowed to hybridize with a priming oligonucleotide on the

daughter array, which is then extended to generate a tagging probe that contains the known barcode of the tagging probe from the master array. The strand that is not immobilized on the daughter array can optionally be released from the newly generated tagging probe.

WHAT IS CLAIMED IS:

1. A method for generating an array, the method comprising:
 - providing a first array comprising a plurality of probes immobilized on a first substrate, wherein a probe of the plurality comprises a spatial barcode;
 - providing a second array comprising a plurality of primers immobilized on a second substrate;
 - generating an intermediate oligonucleotide complementary to a probe on the first array;
 - releasing the intermediate oligonucleotide from the probe on the first array;
 - allowing the intermediate oligonucleotide to hybridize with a primer on the second array; and
 - extending the primer using the intermediate oligonucleotide as a template, thereby generating a probe on the second array.
2. The method of claim 1, wherein the primer on the second array comprises a sequence that is identical to a portion of the sequence of the probe on the first array.
3. The method of claim 2, wherein the portion of the sequence is in a functional domain of the probe.
4. The method of claim 2, wherein the probe has a free 5' end and the portion of the sequence comprises a sequence at the 5' end of the probe.
5. The method of any one of the preceding claims, wherein the plurality of primers on the second array comprise a common sequence.
6. The method of any one of the preceding claims, wherein the probe on the first array is attached to the first substrate via its 3' end.

7. The method of any one of the preceding claims, wherein the spatial barcode comprises 6 to 25 nucleotides.
- 5 8. The method of any one of the preceding claims, wherein the spatial barcode is unique to a particular region on the first array.
9. The method of claim 8, wherein probes associated with a particular region on the first array comprise a common spatial barcode.
- 10 10. The method of any one of the preceding claims, wherein the plurality of probes on the first array comprise a common functional domain.
11. The method of any one of the preceding claims, wherein releasing the intermediate oligonucleotide comprises de-hybridizing the intermediate oligonucleotide from the probe using heat.
- 15 12. The method of any one of the preceding claims, further comprising, following releasing the intermediate oligonucleotide from the probe, facilitating migration of the released intermediate oligonucleotide to the second array.
- 20 13. The method of claim 12, wherein facilitating migration of the released intermediate oligonucleotide comprises applying an electric field to the first and second arrays, wherein the electric field increases diffusion of the released intermediate oligonucleotide.
- 25 14. The method of any one of the preceding claims, wherein the density of the plurality of probes on the first array is about equal to or higher than the density of the plurality of primers on the second array.
- 30 15. The method of any one of the preceding claims, wherein the number of the plurality of probes on the first array is about equal to or higher than the number of the plurality of primers on the second array.

16. The method of any one of the preceding claims, further comprising releasing the intermediate oligonucleotide from the probe generated on the second array.

5 17. The method of claim 16, wherein releasing the intermediate oligonucleotide comprises de-hybridizing the intermediate oligonucleotide from the probe using heat.

18. The method of any one of the preceding claims, wherein the first substrate comprises a plurality of first features, and wherein the probe of the plurality is attached to a feature of
10 the plurality of the first features.

19. The method of claim 18, wherein the spatial barcode is unique to the feature.

20. The method of any one of the preceding claims, wherein the second substrate comprises a plurality of second features, and wherein the primer on the second array is attached to a
15 feature of the plurality of second features.

21. The method of any of claims 18-20, wherein the plurality of first features and/or the plurality of second features are beads, wells, spots, or hydrogel pads.
20