

CLAIMS

1. An antibody or antigen-binding fragment thereof, comprising the aCD25-a-686-HCDR3 amino acid sequence (SEQ ID NO: 4) as variable heavy chain complementarity determining region 3.
2. The antibody or antigen-binding fragment thereof of claim 1, further comprising
- a) aCD25-a-686-HCDR1 amino acid sequence (SEQ ID NO: 2) as variable heavy chain complementarity determining region 1; and
- b) aCD25-a-686-HCDR2 amino acid sequence (SEQ ID NO: 3) as variable heavy chain complementarity determining region 2.
3. The antibody or antigen-binding fragment thereof of claim 1 or claim 2, further comprising:
- a) aCD25-a-686-LCDR1 amino acid sequence (SEQ ID NO: 6) as variable light chain complementarity determining region 1;
- b) aCD25-a-686-LCDR2 amino acid sequence (SEQ ID NO: 7) as variable light chain complementarity determining region 2; and
- c) aCD25-a-686-LCDR3 amino acid sequence (SEQ ID NO: 8) as variable light chain complementarity determining region 3.
4. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment thereof comprises a variable heavy chain comprising aCD25-a-686-HCDR123 amino acid sequence (SEQ ID NO:5).
5. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment thereof comprises a variable light chain comprising aCD25-a-686-LCDR123 amino acid (SEQ ID NO: 9).
6. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment thereof is a monoclonal antibody, a domain antibody, a single chain antibody, a Fab fragment, a F(ab')₂ fragment, a single chain variable fragment (scFv), a scFv-Fc fragment, a single chain antibody (scAb), or a single domain antibody.

- 5 7. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment thereof is a rabbit, mouse, chimeric, humanized or fully human antigen-binding antibody.
8. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody is selected from the group consisting of IgG1, IgG2, IgG3, and IgG4 isotype antibodies.
- 10 9. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment thereof is comprised in a bispecific antibody, a multispecific antibody, or an immunoconjugate further comprising a therapeutic or diagnostic agent.
- 15 10. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment thereof binds the extracellular domain of human CD25.
- 20 11. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment thereof binds cells expressing human CD25 on their surface and is an anti-CD25 Antibody Agent.
- 25 12. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment does not inhibit the binding of Interleukin-2 (IL-2) to CD25.
- 30 13. The antibody or antigen-binding fragment thereof of any preceding claim, wherein the antibody or antigen-binding fragment does not inhibit the signalling of Interleukin-2 (IL-2) via CD25.
- 35 14. The antibody or an antigen-binding fragment thereof of any preceding claim wherein the antibody or antigen-binding fragment thereof is afucosylated.
15. An antibody or antigen-binding fragment that specifically binds to an epitope of human CD25, wherein the epitope comprises one or more amino acid residues comprised in amino acids 70-84 of SEQ ID NO:1.

16. The antibody or antigen-binding fragment of claim 15, wherein the epitope comprises amino acids 70-84 of SEQ ID NO:1.
- 5 17. An affinity matured variant of an antibody or antigen-binding fragment thereof of any one of claims 1 to 16.
- 10 18. An antibody or antigen-binding fragment of claim 17, wherein the antibody or antigen-binding fragment thereof comprises an HCDR2 region selected from the group consisting of AIIPVFGTASYAQKFQG (SEQ ID NO: 13), GIIPIFGDASYAQKFQG (SEQ ID NO: 14), GIIPIFGDANYAQKLQG (SEQ ID NO: 15), GIIPLFGRANYAQKFQG (SEQ ID NO: 16) and GIIPVFGQANYAQKFQG (SEQ ID NO: 17).
- 15 19. An antibody or antigen-binding fragment of claim 17, wherein the antibody or antigen-binding fragment thereof comprises an HCDR1 region selected from the group consisting of GTFSSLAIT (SEQ ID NO: 10), GTFSSLAIS (SEQ ID NO: 2), GTFSSALAIS (SEQ ID NO: 11) and GTFSSLAIF (SEQ ID NO: 12).
- 20 20. An antibody or antigen-binding fragment of any of claims 17 to 19, wherein the antibody or antigen-binding fragment thereof comprises a variable heavy chain comprising a sequence selected from aCD25-a-686-m1-HCDR123 amino acid sequence (SEQ ID NO: 18), aCD25-a-686-m2-HCDR123 amino acid sequence (SEQ ID NO: 19), aCD25-a-686-m3-HCDR123 amino acid sequence (SEQ ID NO: 20), aCD25-a-686-m4-HCDR123 amino acid sequence (SEQ ID NO: 21), and aCD25-a-686-m5-HCDR123 amino acid sequence (SEQ ID NO: 22).
- 25 21. An antibody or antigen-binding fragment of any of claims 17 to 20, wherein the antibody or antigen-binding fragment thereof comprises a variable light chain comprising a sequence selected from aCD25-a-686-m1-LCDR123 amino acid sequence (SEQ ID NO: 9), aCD25-a-686-m2-LCDR123 amino acid sequence (SEQ ID NO: 9), aCD25-a-686-m3-LCDR123 amino acid sequence (SEQ ID NO: 9), aCD25-a-686-m4-LCDR123 amino acid sequence (SEQ ID NO: 9), and aCD25-a-686-m5-LCDR123 amino acid sequence (SEQ ID NO: 9).
- 30
- 35

22. An antibody or antigen-binding fragment of any of claims 17 to 21, wherein the antibody or antigen-binding fragment thereof comprises:
- a) a variable heavy chain comprising a aCD25-a-686-m1-HCDR123 amino acid sequence (SEQ ID NO: 18) and a variable light chain comprising a aCD25-a-686-m1-LCDR123 amino acid sequence (SEQ ID NO: 9);
 - b) a variable heavy chain comprising a aCD25-a-686-m2-HCDR123 amino acid sequence (SEQ ID NO: 19) and a variable light chain comprising a aCD25-a-686-m2-LCDR123 amino acid sequence (SEQ ID NO: 9);
 - c) a variable heavy chain comprising a aCD25-a-686-m3-HCDR123 amino acid sequence (SEQ ID NO: 20) and a variable light chain comprising a aCD25-a-686-m3-LCDR123 amino acid sequence (SEQ ID NO: 9);
 - d) a variable heavy chain comprising a aCD25-a-686-m4-HCDR123 amino acid sequence (SEQ ID NO: 21) and a variable light chain comprising a aCD25-a-686-m4-LCDR123 amino acid sequence (SEQ ID NO: 9); or
 - e) a variable heavy chain comprising a aCD25-a-686-m5-HCDR123 amino acid sequence (SEQ ID NO: 22) and a variable light chain comprising a aCD25-a-686-m5-LCDR123 amino acid sequence (SEQ ID NO: 9).
23. A nucleic acid molecule encoding the antibody or antigen-binding fragment thereof of any of the Claims 1 to 22.
24. A nucleic acid vector comprising the nucleic acid molecule of claim 23.
25. A host cell comprising the nucleic acid vector of claim 24.
26. A method for producing an antibody or antigen-binding fragment thereof of any one of claims 1 to 22 comprising culturing a host cell of claim 25.
27. A composition comprising an antibody or antigen-binding fragment thereof of any one of claims 1 to 22.
28. The composition of claim 27 that further comprises a pharmaceutically acceptable carrier or excipient.
29. The pharmaceutical composition of claim 27 or claim 28, wherein said composition is for use in the treatment of cancer.

- 5 30. Use of an antibody or antigen-binding fragment thereof of any one of claims 1 to 22, or of a composition of claim 27 or 28 in the manufacture of a medicament for treating cancer.
31. Use of an antibody or antigen-binding fragment thereof that competes with an antibody according to anyone of claims 1 to 22 for the binding of CD25 in the manufacture of a medicament for treating cancer.
- 10 32. A method of treating cancer in a subject, comprising administering to the subject an effective amount of the composition of claim 27 or claim 28.
- 15 33. A method of treating cancer in a subject, comprising administering to the subject an effective amount of an antibody or antigen-binding fragment thereof that competes with an antibody according to anyone of claims 1 to 22 for the binding of CD25.
- 20 34. The method of claim 32 or 33, further comprising administering, simultaneously or sequentially in any order, a second agent to the subject.
35. The method of claim 34 wherein the second agent is an immune checkpoint inhibitor or a cancer vaccine.
- 25 36. The method of claim 35 wherein the second agent is an immune checkpoint inhibitor, wherein the immune checkpoint inhibitor is a PD-1 antagonist.
37. The method of claim 36 wherein the PD-1 antagonist is an anti-PD-1 antibody or an anti-PD-L1 antibody.
- 30 38. The method of any one of claims 32 to 37 wherein the subject has a solid tumor.
- 35 39. The method of any one of claims 32 to 37 wherein the subject has a haematological cancer tumour.

- 5
40. A method of depleting regulatory T cells in a tumour in a subject comprising the step of administering an effective amount of the composition of claim 27 or claim 28 or an antibody or antigen-binding fragment that competes with an antibody according to anyone of claims 1 to 22 for the binding of CD25.
41. A method according to claim 40, wherein the subject has a solid tumor.
42. A method according to claim 40, wherein the subject has a haematological cancer tumor.
- 10
43. An antibody or antigen-binding fragment that competes with an antibody according to anyone of claims 1 to 22 for the binding of CD25 for use in the treatment of cancer.
- 15
44. A kit comprising the composition of claim 27 or claim 28 in a container.
45. A method of preparing an anti-CD25 antibody comprising providing an antibody according to any one of claims 1 to 22, and subjecting the antibody to affinity maturation, wherein the anti-CD25 antibody produced has a greater affinity to CD25 than the parental antibody,
- 20
46. A method of preparing a pharmaceutical composition comprising providing an antibody prepared according to a method of claim 45 and co-formulating the antibody with at least one of more pharmaceutically acceptable excipient.
- 25