

SPECIFICATION

Title of the Invention: Pyrimidine derivative

Technical Field

[0001]

The present invention relates to a novel pyrimidine derivative. More specifically, the present invention relates to a pyrimidine derivative having an mPGES-1 inhibitory action, and useful as an active ingredient of a medicament for prophylactic and/or therapeutic treatment of such diseases as inflammation, pain, and rheumatism.

Background Art

[0002]

Prostaglandin E2 (PGE2) is involved in inflammation, pain, pyrexia, and the like by means of PGE receptors, and can suppress the PGE2 production to suppress inflammation. Non-steroidal anti-inflammatory drugs (NSAIDs) inhibit cyclooxygenase (COX) in the upstream of the prostaglandin biosynthesis pathway, and thereby exhibit anti-inflammatory activity. However, they totally suppress the prostaglandin biosynthesis pathway downstream from the prostanoïd production in which COX is involved, and therefore they cause gastric mucosal injury as side effects due to suppression of secretion of gastric mucus or blood flow in gastric mucosa.

[0003]

There are two types of isozymes of COX, COX-1 and COX-2. Among them, COX-2 is expressed and induced in inflammatory tissues by various inflammation-promoting stimuli (for example, those of cytokines such as interleukin-1 β). Medicaments that selectively inhibit this COX-2 suppress the production of PGI2, which has vasodilatation and platelet aggregation actions; however, since they do not inhibit the production of thromboxane A2 (TXA2) catalyzed by COX-1 (TXA2 causes vasoconstriction and platelet coagulation), they are considered to increase risk of thrombosis, and increase cardiovascular events, either.

[0004]

In the downstream of the biosynthesis pathway of PGE2, PGE2 is biosynthesized from PGH2 by the prostaglandin E synthase (PGE synthase, PGES). As PGES, there are three kinds of enzymes, mPGES-1 (microsomal prostaglandin E2 synthase-1), mPGES-2 (microsomal prostaglandin E2 synthase-2), and cPGES (cytosolic PGE synthase). Among them, mPGES-1 is an inducible trimer enzyme, of

which expression is increased by inflammatory stimuli (Proc. Natl. Acad. Sci. USA, 96, pp.7220-7225, 1999), and it is known to participate in cancer, inflammation, pain, pyrexia, tissue repair, and the like.

[0005]

Since mPGES-1 inhibitors can selectively inhibit the final step of the PGE2 biosynthesis pathway in inflammation lesions (Pharmacol. Rev., 59, pp.207-224, 2007; J. Biol. Chem., 279, pp.33684-33695, 2004), they are expected as anti-inflammatory agents that do not cause gastric mucosal injuries, unlike the non-steroidal anti-inflammatory agents. There are also expected efficacies of mPGES-1 inhibitors for prophylactic and/or therapeutic treatment of pain, rheumatism, osteoarthritis, pyrexia, Alzheimer's disease, multiple sclerosis, arteriosclerosis, ocular hypertension such as glaucoma, ischemic retinopathy, systemic scleroderma, malignant tumors such as large intestine tumor, and diseases for which suppression of the PGE2 production exhibits efficacy (refer to International Patent Publication WO2015/125842 for PGE2, PGES, and mPGES-1, as well as uses of mPGES-1 inhibitors, and the like). In addition, it is also known that mPGES-1 inhibitors increase productions of other prostanoids in connection with the suppression of the PGE2 production (J. Biol. Chem., 280, pp.16579-16585, 2005).

[0006]

As such mPGES-1 inhibitors, there are known the heterocyclic derivatives disclosed in Japanese Patent No. 5601422, the substituted pyrimidine compounds disclosed in International Patent Publication WO2015/59618, the triazine compounds disclosed in International Patent Publication WO2015/125842, and the like. International Patent Publication WO2015/59618 discloses a pyrimidine compound substituted with p-trifluoromethylphenyl group and 2-chloro-5-isobutyramidobenzyl group (Example 2), and International Patent Publication WO2015/125842 discloses triazine compounds substituted with p-trifluoromethylphenyl group and 2-chloro-5-isobutyramidobenzyl group (Examples 1 to 28).

Prior art references

Patent documents

[0007]

Patent document 1: Japanese Patent No. 5601422

Patent document 2: International Patent Publication WO2015/59618

Patent document 3: International Patent Publication WO2015/125842

Non-patent documents

[0008]

Non-patent document 1: Proc. Natl. Acad. Sci. USA, 96, pp.7220-7225, 1999

Non-patent document 2: Pharmacol. Rev., 59, pp.207-224, 2007

Non-patent document 3: J. Biol. Chem., 279, pp.33684-33695, 2004

Non-patent document 4: J. Biol. Chem., 280, pp.16579-16585, 2005

Summary of the Invention

Object to be Achieved by the Invention

[0009]

An object of the present invention is to provide a novel compound having an mPGES-1 inhibitory action, and useful as an active ingredient of a medicament for prophylactic and/or therapeutic treatment of such diseases as inflammation, pain, and rheumatism.

Means for Achieving the Object


[0010]

The inventors of the present invention conducted various researches in order to achieve the aforementioned object. As a result, they found that pyrimidine derivatives represented by the following general formula (1) have a potent mPGES-1 inhibitory action, and are useful as active ingredients of medicaments for prophylactic and/or therapeutic treatment of such diseases as inflammation, pain, and rheumatism, and accomplished the present invention.

[0011]

The present invention thus provides a compound represented by the following general formula (1):

[Formula 1]



(in the formula, X represents carbonyl group, or sulfonyl group; R¹ represents hydrogen atom, a halogen atom, an alkyl group, an alkanoyl group, cyano group, or carboxyl group; R² represents an alkyl group, a cyclic carbon group which may have a

substituent, or a heterocyclic group which may have a substituent; R³ represents hydrogen atom, or 1 to 3 substituents substituting on the benzene ring (these substituents are selected from the group consisting of a halogen atom, an alkyl group (this alkyl group may be substituted with a halogen atom), and an alkoxy group (this alkoxy group may be substituted with a halogen atom)); R⁴ and R⁵ independently represent hydrogen atom, a halogen atom, or an alkyl group; and R⁶ represents an alkyl group (this alkyl group may be substituted with hydroxy group, a halogen atom, or an alkoxy group), or an alkoxy group, or a salt thereof.

[0012]

According to preferred embodiments of the aforementioned invention, there are provided the compound represented by the aforementioned general formula (1), or a salt thereof, wherein X is carbonyl group; the compound represented by the aforementioned general formula (1), or a salt thereof, wherein R⁶ is a branched C₁₋₆ alkyl group (this alkyl group may be substituted with a C₁₋₆ alkoxy group); the compound represented by the aforementioned general formula (1), or a salt thereof, wherein both R⁴ and R⁵ are hydrogen atoms; the compound represented by the aforementioned general formula (1), or a salt thereof, wherein R³ consists of one halogen atom; the compound represented by the aforementioned general formula (1), or a salt thereof, wherein R¹ is hydrogen atom, an alkyl group, or cyano group; and the compound represented by the aforementioned general formula (1), or a salt thereof, wherein R² is a saturated or partially saturated 3- to 7-membered monocyclic hydrocarbon group which may have a substituent, a phenyl group which may have a substituent, a saturated or partially saturated 3- to 7-membered monocyclic heterocyclic group which may have a substituent (this heterocyclic group contains 1 to 3 ring-constituting heteroatoms), or a monocyclic aromatic heterocyclic group which may have a substituent (this heterocyclic group contains 1 to 3 ring-constituting heteroatoms).

[0013]

As other aspects, the present invention provides an mPGES-1 inhibitor containing a compound represented by the aforementioned general formula (1), or a salt thereof; and a PGE2 biosynthesis inhibitor containing a compound represented by the aforementioned general formula (1), or a salt thereof.

[0014]

As still another aspect, the present invention provides a medicament containing a compound represented by the aforementioned general formula (1) or a physiologically acceptable salt thereof as an active ingredient. This medicament can

be used for prophylactic and/or therapeutic treatment of, for example, inflammation, pain, rheumatism, osteoarthritis, pyrexia, Alzheimer's disease, multiple sclerosis, arteriosclerosis, ocular hypertension such as glaucoma, ischemic retinopathy, systemic scleroderma, malignant tumors such as large intestine tumor, and diseases for which suppression of the PGE2 production exhibits efficacy.

[0015]

The present invention also provides use of a compound represented by the aforementioned general formula (1) or a salt thereof for manufacture of the aforementioned mPGES-1 inhibitor, the aforementioned PGE2 biosynthesis inhibitor, or the aforementioned medicament; a method for inhibiting mPGES-1 in a living body of a mammal including human, which comprises the step of administering an effective amount of a compound represented by the aforementioned general formula (1) or a physiologically acceptable salt thereof to the mammal including human; a method for inhibiting biosynthesis of PGE2 in a living body of a mammal including human, which comprises the step of administering an effective amount of a compound represented by the aforementioned general formula (1) or a physiologically acceptable salt thereof to the mammal including human; and a method for promoting production of a prostanoid other than PGE2 by inhibiting biosynthesis of PGE2 in a living body of a mammal including human, which comprises the step of administering an effective amount of a compound represented by the aforementioned general formula (1) or a physiologically acceptable salt thereof to the mammal including human.

Effect of the Invention

[0016]

The compounds represented by the aforementioned general formula (1) and salts thereof provided by the present invention can exhibit a potent inhibitory action against mPGES-1 to inhibit the biosynthesis of PGE2. Therefore, they are useful as an active ingredient of a medicament for prophylactic and/or therapeutic treatment of, for example, pain, rheumatism, osteoarthritis, pyrexia, Alzheimer's disease, multiple sclerosis, arteriosclerosis, ocular hypertension such as glaucoma, ischemic retinopathy, systemic scleroderma, malignant tumors such as large intestine tumor, and diseases for which suppression of the PGE2 production exhibits efficacy.

Modes for Carrying out the Invention

[0017]

In the aforementioned general formula (1), X represents carbonyl group, or

sulfonyl group. It is preferred that X is carbonyl group.

R¹ represents hydrogen atom, a halogen atom, an alkyl group, an alkanoyl group, cyano group, or carboxyl group. In this specification, the term halogen atom may mean fluorine atom, chlorine atom, bromine atom, or iodine atom. As the halogen atom, fluorine atom or chlorine atom is preferred.

[0018]

In this specification, the term alkyl group may mean a straight, branched, or cyclic alkyl group, or an alkyl group consisting of a combination of the foregoing alkyl groups. Although carbon number of the alkyl group is not particularly limited, it is, for example, 1 to 12, preferably 1 to 6, particularly preferably 1 to 4. The same shall apply to alkyl moieties of other substituents having an alkyl moiety (for example, alkoxy group). As R¹, hydrogen atom, an alkyl group, or cyano group is preferred, hydrogen atom, methyl group, or cyano group is more preferred, and hydrogen atom is particularly preferred.

[0019]

R² represents an alkyl group, a cyclic carbon group which may have a substituent, or a heterocyclic group which may have a substituent. As the cyclic carbon group, for example, an aromatic hydrocarbon group, or a saturated or partially saturated cyclic hydrocarbon group can be used. As the aromatic hydrocarbon group, for example, phenyl group, naphthyl group, and the like can be used, and as the saturated or partially saturated cyclic hydrocarbon group, for example, a saturated, or partially saturated 3- to 12-membered monocyclic or bicyclic cyclic hydrocarbon group can be used. As the aromatic hydrocarbon group, phenyl group can be preferably used. As the cyclic hydrocarbon group, a saturated 3- to 7-membered monocyclic cyclic hydrocarbon group can be preferably used, and cyclopropyl group, cyclobutyl group, cyclopentyl group, cyclohexyl group, cycloheptyl group, and the like are more preferred.

[0020]

As the heterocyclic group, a saturated or partially saturated heterocyclic group having 1 or 2 or more ring-constituting heteroatoms, or an aromatic heterocyclic group having 1 or 2 or more ring-constituting heteroatoms can be used. As the ring-constituting heteroatoms, nitrogen atom, oxygen atom, sulfur atom, and the like can be used. As the saturated or partially saturated heterocyclic group having 1 or 2 or more ring-constituting heteroatoms, for example, a saturated or partially saturated 3- to 7-membered monocyclic heterocyclic group, or 8- to 12-membered bicyclic heterocyclic group can be used. Examples of the saturated or partially saturated 3- to 7-membered monocyclic heterocyclic group include, for example, 1-aziridinyl group, 1-azetidiny

group, 1-pyrrolidinyl group, 2-pyrrolidinyl group, 3-pyrrolidinyl group, 2-tetrahydrofuryl group, 3-tetrahydrofuryl group, thiolanyl group, 1-imidazolidinyl group, 2-imidazolidinyl group, 4-imidazolidinyl group, 1-pyrazolidinyl group, 3-pyrazolidinyl group, 4-pyrazolidinyl group, 1-(2-pyrrolinyl) group, 1-(2-imidazoliny) group, 2-(2-imidazoliny) group, 1-(2-pyrazoliny) group, 3-(2-pyrazoliny) group, piperidino group, 2-piperidinyl group, 3-piperidinyl group, 4-piperidinyl group, 1-homopiperidinyl group, 2-tetrahydropyranyl group, morpholino group, (thiomorpholin)-4-yl group, 1-piperazinyl group, 1-homopiperazinyl group, and the like, and examples of the saturated or partially saturated 8- to 12-membered bicyclic heterocyclic group include, for example, 2-quinuclidinyl group, 2-cromanyl group, 3-cromanyl group, 4-cromanyl group, 5-cromanyl group, 6-cromanyl group, 7-cromanyl group, 8-cromanyl group, 1-isocromanyl group, 3-isocromanyl group, 4-isocromanyl group, 5-isocromanyl group, 6-isocromanyl group, 7-isocromanyl group, 8-isocromanyl group, 2-thiocromanyl group, 3-thiocromanyl group, 4-thiocromanyl group, 5-thiocromanyl group, 6-thiocromanyl group, 7-thiocromanyl group, 8-thiocromanyl group, 1-isothiocromanyl group, 3-isothiocromanyl group, 4-isothiocromanyl group, 5-isothiocromanyl group, 6-isothiocromanyl group, 7-isothiocromanyl group, 8-isothiocromanyl group, 1-indolinyl group, 2-indolinyl group, 3-indolinyl group, 4-indolinyl group, 5-indolinyl group, 6-indolinyl group, 7-indolinyl group, 1-isoindolinyl group, 2-isoindolinyl group, 4-isoindolinyl group, 5-isoindolinyl group, 2-(4H-chromenyl) group, 3-(4H-chromenyl) group, 4-(4H-chromenyl) group, 5-(4H-chromenyl) group, 6-(4H-chromenyl) group, 7-(4H-chromenyl) group, 8-(4H-chromenyl) group, 1-isochromenyl group, 3-isochromenyl group, 4-isochromenyl group, 5-isochromenyl group, 6-isochromenyl group, 7-isochromenyl group, 8-isochromenyl group, 1-(1H-pyrrolidinyl) group, 2-(1H-pyrrolidinyl) group, 3-(1H-pyrrolidinyl) group, 5-(1H-pyrrolidinyl) group, 6-(1H-pyrrolidinyl) group, 7-(1H-pyrrolidinyl) group, and the like, but the examples are not limited to these.

[0021]

As for the aromatic heterocyclic group having 1 or 2 or more ring-constituting heteroatoms, examples of monocyclic aromatic heterocyclic group include, for example, 2-furyl group, 3-furyl group, 2-thienyl group, 3-thienyl group, 1-pyrrolyl group, 2-pyrrolyl group, 3-pyrrolyl group, 2-oxazolyl group, 4-oxazolyl group, 5-oxazolyl group, 3-isoxazolyl group, 4-isoxazolyl group, 5-isoxazolyl group, 2-thiazolyl group, 4-thiazolyl group, 5-thiazolyl group, 3-isothiazolyl group, 4-isothiazolyl group, 5-isothiazolyl group, 1-imidazolyl group, 2-imidazolyl group, 4-imidazolyl group, 5-imidazolyl group, 1-pyrazolyl group, 3-pyrazolyl group, 4-pyrazolyl group, 5-pyrazolyl group, (1,2,3-

oxadiazol)-4-yl group, (1,2,3-oxadiazol)-5-yl group, (1,2,4-oxadiazol)-3-yl group, (1,2,4-oxadiazol)-5-yl group, (1,2,5-oxadiazol)-3-yl group, (1,2,5-oxadiazol)-4-yl group, (1,3,4-oxadiazol)-2-yl group, (1,3,4-oxadiazol)-5-yl group, furazanyl group, (1,2,3-thiadiazol)-4-yl group, (1,2,3-thiadiazol)-5-yl group, (1,2,4-thiadiazol)-3-yl group, (1,2,4-thiadiazol)-5-yl group, (1,2,5-thiadiazol)-3-yl group, (1,2,5-thiadiazol)-4-yl group, (1,3,4-thiadiazolyl)-2-yl group, (1,3,4-thiadiazolyl)-5-yl group, (1H-1,2,3-triazol)-1-yl group, (1H-1,2,3-triazol)-4-yl group, (1H-1,2,3-triazol)-5-yl group, (2H-1,2,3-triazol)-2-yl group, (2H-1,2,3-triazol)-4-yl group, (1H-1,2,4-triazol)-1-yl group, (1H-1,2,4-triazol)-3-yl group, (1H-1,2,4-triazol)-5-yl group, (4H-1,2,4-triazol)-3-yl group, (4H-1,2,4-triazol)-4-yl group, (1H-tetrazol)-1-yl group, (1H-tetrazol)-5-yl group, (2H-tetrazol)-2-yl group, (2H-tetrazol)-5-yl group, 2-pyridyl group, 3-pyridyl group, 4-pyridyl group, 3-pyridazinyl group, 4-pyridazinyl group, 2-pyrimidinyl group, 4-pyrimidinyl group, 5-pyrimidinyl group, 2-pyrazinyl group, (1,2,3-triazin)-4-yl group, (1,2,3-triazin)-5-yl group, (1,2,4-triazin)-3-yl group, (1,2,4-triazin)-5-yl group, (1,2,4-triazin)-6-yl group, (1,3,5-triazin)-2-yl group, 1-azepinyl group, 2-azepinyl group, 3-azepinyl group, 4-azepinyl group, (1,4-oxazepin)-2-yl group, (1,4-oxazepin)-3-yl group, (1,4-oxazepin)-5-yl group, (1,4-oxazepin)-6-yl group, (1,4-oxazepin)-7-yl group, (1,4-thiazepin)-2-yl group, (1,4-thiazepin)-3-yl group, (1,4-thiazepin)-5-yl group, (1,4-thiazepin)-6-yl group, (1,4-thiazepin)-7-yl group, and the like, but the examples are not limited to these.

[0022]

Examples of condensed polycyclic aromatic heterocyclic group include, for example, 8- to 14-membered condensed polycyclic aromatic heterocyclic groups such as 2-benzofuranyl group, 3-benzofuranyl group, 4-benzofuranyl group, 5-benzofuranyl group, 6-benzofuranyl group, 7-benzofuranyl group, 1-isobenzofuranyl group, 4-isobenzofuranyl group, 5-isobenzofuranyl group, 2-benzo[b]thienyl group, 3-benzo[b]thienyl group, 4-benzo[b]thienyl group, 5-benzo[b]thienyl group, 6-benzo[b]thienyl group, 7-benzo[b]thienyl group, 1-benzo[c]thienyl group, 4-benzo[c]thienyl group, 5-benzo[c]thienyl group, 1-indolyl group, 2-indolyl group, 3-indolyl group, 4-indolyl group, 5-indolyl group, 6-indolyl group, 7-indolyl group, (2H-isoindol)-1-yl group, (2H-isoindol)-2-yl group, (2H-isoindol)-4-yl group, (2H-isoindol)-5-yl group, (1H-indazol)-1-yl group, (1H-indazol)-3-yl group, (1H-indazol)-4-yl group, (1H-indazol)-5-yl group, (1H-indazol)-6-yl group, (1H-indazol)-7-yl group, (2H-indazol)-1-yl group, (2H-indazol)-2-yl group, (2H-indazol)-4-yl group, (2H-indazol)-5-yl group, 2-benzoxazolyl group, 4-benzoxazolyl group, 5-benzoxazolyl group, 6-benzoxazolyl group, 7-benzoxazolyl group, (1,2-benzisoxazol)-3-yl group, (1,2-benzisoxazol)-4-yl group, (1,2-benzisoxazol)-5-yl group, (1,2-benzisoxazol)-6-yl group, (1,2-benzisoxazol)-7-yl group,

(2,1-benzisoxazol)-3-yl group, (2,1-benzisoxazol)-4-yl group, (2,1-benzisoxazol)-5-yl group, (2,1-benzisoxazol)-6-yl group, (2,1-benzisoxazol)-7-yl group, 2-benzothiazolyl group, 4-benzothiazolyl group, 5-benzothiazolyl group, 6-benzothiazolyl group, 7-benzothiazolyl group, (1,2-benzisothiazol)-3-yl group, (1,2-benzisothiazol)-4-yl group, (1,2-benzisothiazol)-5-yl group, (1,2-benzisothiazol)-6-yl group, (1,2-benzisothiazol)-7-yl group, (2,1-benzisothiazol)-3-yl group, (2,1-benzisothiazol)-4-yl group, (2,1-benzisothiazol)-5-yl group, (2,1-benzisothiazol)-6-yl group, (2,1-benzisothiazol)-7-yl group, (1,2,3-benzoxadiazol)-4-yl group, (1,2,3-benzoxadiazol)-5-yl group, (1,2,3-benzoxadiazol)-6-yl group, (1,2,3-benzoxadiazol)-7-yl group, (2,1,3-benzoxadiazol)-4-yl group, (2,1,3-benzoxadiazol)-5-yl group, (1,2,3-benzothiadiazol)-4-yl group, (1,2,3-benzothiadiazol)-5-yl group, (1,2,3-benzothiadiazol)-6-yl group, (1,2,3-benzothiadiazol)-7-yl group, (2,1,3-benzothiadiazol)-4-yl group, (2,1,3-benzothiadiazol)-5-yl group, (1H-benzotriazol)-1-yl group, (1H-benzotriazol)-4-yl group, (1H-benzotriazol)-5-yl group, (1H-benzotriazol)-6-yl group, (1H-benzotriazol)-7-yl group, (2H-benzotriazol)-2-yl group, (2H-benzotriazol)-4-yl group, (2H-benzotriazol)-5-yl group, 2-quinolyl group, 3-quinolyl group, 4-quinolyl group, 5-quinolyl group, 6-quinolyl group, 7-quinolyl group, 8-quinolyl group, 1-isoquinolyl group, 3-isoquinolyl group, 4-isoquinolyl group, 5-isoquinolyl group, 6-isoquinolyl group, 7-isoquinolyl group, 8-isoquinolyl group, 3-cinnolinyl group, 4-cinnolinyl group, 5-cinnolinyl group, 6-cinnolinyl group, 7-cinnolinyl group, 8-cinnolinyl group, 2-quinazolinyl group, 4-quinazolinyl group, 5-quinazolinyl group, 6-quinazolinyl group, 7-quinazolinyl group, 8-quinazolinyl group, 2-quinoxaliny group, 5-quinoxaliny group, 6-quinoxaliny group, 1-phthalazinyl group, 5-phthalazinyl group, 6-phthalazinyl group, 2-naphthyridinyl group, 3-naphthyridinyl group, 4-naphthyridinyl group, 2-puriny group, 6-puriny group, 7-puriny group, 8-puriny group, 2-pteridinyl group, 4-pteridinyl group, 6-pteridinyl group, 7-pteridinyl group, 1-carbazolyl group, 2-carbazolyl group, 3-carbazolyl group, 4-carbazolyl group, 9-carbazolyl group, 2-(α -carboliny) group, 3-(α -carboliny) group, 4-(α -carboliny) group, 5-(α -carboliny) group, 6-(α -carboliny) group, 7-(α -carboliny) group, 8-(α -carboliny) group, 9-(α -carboliny) group, 1-(β -carbonylyl) group, 3-(β -carbonylyl) group, 4-(β -carbonylyl) group, 5-(β -carbonylyl) group, 6-(β -carbonylyl) group, 7-(β -carbonylyl) group, 8-(β -carbonylyl) group, 9-(β -carbonylyl) group, 1-(γ -carboliny) group, 2-(γ -carboliny) group, 4-(γ -carboliny) group, 5-(γ -carboliny) group, 6-(γ -carboliny) group, 7-(γ -carboliny) group, 8-(γ -carboliny) group, 9-(γ -carboliny) group, 1-acridinyl group, 2-acridinyl group, 3-acridinyl group, 4-acridinyl group, 9-acridinyl group, 1-phenoxazinyl group, 2-phenoxazinyl group, 3-phenoxazinyl group, 4-phenoxazinyl group, 10-phenoxazinyl group, 1-phenothiazinyl group, 2-phenothiazinyl group, 3-phenothiazinyl group, 4-

phenothiazinyl group, 10-phenothiazinyl group, 1-phenazinyl group, 2-phenazinyl group, 1-phenanthridinyl group, 2-phenanthridinyl group, 3-phenanthridinyl group, 4-phenanthridinyl group, 6-phenanthridinyl group, 7-phenanthridinyl group, 8-phenanthridinyl group, 9-phenanthridinyl group, 10-phenanthridinyl group, 2-phenanthrolinyl group, 3-phenanthrolinyl group, 4-phenanthrolinyl group, 5-phenanthrolinyl group, 6-phenanthrolinyl group, 7-phenanthrolinyl group, 8-phenanthrolinyl group, 9-phenanthrolinyl group, 10-phenanthrolinyl group, 1-thianthrenyl group, 2-thianthrenyl group, 1-indolizinyl group, 2-indolizinyl group, 3-indolizinyl group, 5-indolizinyl group, 6-indolizinyl group, 7-indolizinyl group, 8-indolizinyl group, 1-phenoxathiinyl group, 2-phenoxathiinyl group, 3-phenoxathiinyl group, 4-phenoxathiinyl group, thieno[2,3-b]furyl group, pyrrolo[1,2-b]pyridazinyl group, pyrazolo[1,5-a]pyridyl group, imidazo[11,2-a]pyridyl group, imidazo[1,5-a]pyridyl group, imidazo[1,2-b]pyridazinyl group, imidazo[1,2-a]pyrimidinyl group, 1,2,4-triazolo[4,3-a]pyridyl group, and 1,2,4-triazolo[4,3-a]pyridazinyl group, but the examples are not limited to these.

[0023]

When the expression "which may have a substituent" is used for a certain functional group in this specification, it means that the functional group is unsubstituted, or the functional group has one or two or more substituents at chemically substitutable positions, unless otherwise indicated. Type, number, and substitution position of substituent existing on a functional group are not particularly limited, and when there are two or more substituents, they may be the same or different. Examples of the substituent existing on a functional group include, for example, an alkyl group, a halogen atom, oxo group, thioxo group, nitro group, nitroso group, cyano group, isocyano group, cyanato group, tiocyanato group, isocyanato group, isotiocyanato group, hydroxy group, sulfanyl group, carboxy group, sulfanylcarbonyl group, oxalo group, mesoxalo group, thiocarboxy group, dithiocarboxy group, carbamoyl group, thiocarbamoyl group, sulfo group, sulfamoyl group, sulfinio group, sulfinamoyl group, sulfeno group, sulfenamoyl group, phosphono group, hydroxyphosphonyl group, a hydrocarbon group, a heterocyclic group, a hydrocarbon-oxy group, a (heterocyclic ring)-oxy group, a hydrocarbon-sulfanyl group, a (heterocyclic ring)-sulfanyl group, an acyl group, amino group, hydrazino group, hydrazono group, diazenyl group, ureido group, thioureido group, guanidino group, carbamoimidoyl group (amidino group), azido group, imino group, hydroxyamino group, hydroxyimino group, aminoxy group, diazo group, semicarbazino group, semicarbazono group, allophanyl group, hydantoyl group, phosphano group, phosphoroso group, phospho group, boryl group, silyl group,

stanyl group, selanyl group, oxido group, and the like, but the examples are not limited to these examples.

[0024]

The substituent in the aforementioned definitions may be substituted with another substituent at a chemically substitutable position on the substituent. Type, number, and substitution position of the substituent are not particularly limited, and when the substituent is substituted with two or more substituents, they may be the same or different. Examples of such a substituent include, for example, a halogenated alkyl group (for example, trifluoromethyl group and the like), a hydroxyalkyl group (for example, hydroxymethyl group and the like), a halogenated alkyl-carbonyl group (for example, trifluoroacetyl and the like), a halogenated alkyl-sulfonyl group (for example, trifluoromethanesulfonyl and the like), an acyl-oxy group, an acyl-sulfanyl group, an N-hydrocarbon-amino group, an N,N-di(hydrocarbon)-amino group, an N-(heterocyclic ring)-amino group, an N-hydrocarbon-N-(heterocyclic ring)-amino group, an acyl-amino group, a di(acyl)-amino group, and the like, but the examples are not limited to these examples.

[0025]

As R², a saturated or partially saturated 3- to 7-membered monocyclic cyclic hydrocarbon group which may have a substituent, a phenyl group which may have a substituent, a saturated or partially saturated 3- to 7-membered monocyclic heterocyclic group which may have a substituent (this heterocyclic group contains 1 to 3 ring-constituting heteroatoms), or a aromatic monocyclic heterocyclic group which may have a substituent (this heterocyclic group contains 1 to 3 ring-constituting heteroatoms) is preferred. As R², a saturated cyclic hydrocarbon group such as cyclopentyl group, and cyclohexyl group, phenyl group, thienyl group, pyridyl group, pyrimidyl group, thiazolyl group, pyrazolyl group, oxodihydropyridyl group, benzothiophene group, and the like are preferred, and examples of substituents substituting on the rings of these cyclic groups include, for example, one or two or more substituents selected from methyl group, isobutyl group, chlorine atom, fluorine atom, cyano group, trifluoromethyl group, difluoromethyl group, methoxy group, difluoromethoxy group, trifluoromethoxy group, ethoxy group, methoxyethoxy group, ethoxyethoxy group, trifluoroethoxy group, isopropoxy group, methoxypropoxy group, methylsulfanyl group, cyclopropyl group, cyclopropylethynyl group, propynyl group, tetrahydropyran-4-ylmethoxy group, and the like, but the examples are not limited to these.

[0026]

R³ represents hydrogen atom, or 1 to 3 substituents substituting on the benzene ring, and these substituents are selected from the group consisting of a halogen atom, an alkyl group (this alkyl group may be substituted with a halogen atom), and an alkoxy group (this alkoxy group may be substituted with a halogen atom). R³ preferably represents one or two substituents substituting on the benzene ring, and it is preferred that these substituents are chlorine atom, fluorine atom, trifluoromethyl group, or difluoromethoxy group. When there are two substituents, chlorine atom, fluorine atom, or trifluoromethyl group is preferred. Although substitution position of R³ is not particularly limited, when R³ consists of one substituent substituting on the benzene ring, it is preferred that it substitutes at the ortho-position of the pyrimidinyl group substituting on the benzene ring, and it is more preferred that this substituent substitutes at the ortho-position of the pyrimidinyl group substituting on the benzene ring, and the para-position of the aminomethyl group substituting on the benzene ring. Although substitution position of R³ is not particularly limited, when R³ consists of two substituents substituting on the benzene ring, it is preferred that they substitute at the ortho-positions of the pyrimidinyl group substituting on the benzene ring, and it is more preferred that these substituents substitute at the ortho-positions of the pyrimidinyl group substituting on the benzene ring, and the ortho-position and the para-position of the aminomethyl group substituting on the benzene ring.

[0027]

R⁴ and R⁵ independently represent hydrogen atom, a halogen atom, or an alkyl group. It is preferred that both R⁴ and R⁵ are hydrogen atoms.

R⁶ represents an alkyl group (this alkyl group may be substituted with hydroxy group, a halogen atom, or an alkoxy group), or an alkoxy group. R⁶ is preferably a straight, branched, or cyclic C₁₋₆ alkyl group, and the branched C₁₋₆ alkyl group may be substituted with a C₁₋₆ alkoxy group. As R⁶, ethyl group, propyl group, isopropyl group, 1-methyl-1-methoxyethyl group, cyclopropyl group, t-butyl group, and the like can be preferably used, and ethyl group, propyl group, and isopropyl group can be particularly preferably used.

[0028]

According to the present invention, the compounds represented by the formula (1) can be prepared by, for example, one of the methods of (a) to (d) mentioned below.


[0029]

Method (a):

A pyrimidine derivative represented by the formula (1) wherein R¹ is cyano


group can be prepared by reacting an amidine derivative represented by the following formula (2):

[Formula 2]




(in the formula, R³, R⁴, R⁵, R⁶, and X have the same meanings as those defined above), an ester represented by the following formula (3)

[Formula 3]



(in the formula, R¹ is cyano group), and an aldehyde represented by the following formula (4)

[Formula 4]



(in the formula, R² has the same meaning as that defined above).

[0030]

Examples of base used for this reaction include organic bases (amines, for example, mono- to trialkylamines such as methylamine, ethylamine, diethylamine, triethylamine, propylamine, isopropylamine, and diisopropylethylamine; alkanolamines such as ethanolamine; alkylene polyamines such as ethylenediamine, and diethylenetriamine, and the like), inorganic bases [metal hydroxides (alkali metal or alkaline earth metal hydroxides, and the like) such as sodium hydroxide, potassium hydroxide, calcium hydroxide, iron hydroxide, and aluminum hydroxide; alkali metal carbonates such as sodium carbonate, sodium hydrogencarbonate, potassium carbonate,

rubidium carbonate, and cesium carbonate; ammonia, and the like], and the like. Preferred examples include alkali metal bases and organic bases, and particularly preferred examples include potassium carbonate, sodium ethoxide, and diisopropylethylamine. A salt may be formed with one or two or more kinds of these bases.

[0031]


Reaction time, reaction temperature, and the like may be selected from conventionally used ranges. Reaction temperature is preferably 0 to 140°C, particularly preferably 20 to 80°C. Reaction time is preferably 0.25 to 48 hours, particularly preferably 0.5 to 24 hours.

Solvent used for this reaction may be selected from conventionally used solvents. Water, N,N-dimethylacetamide, methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 1-pentanol, and the like are preferred.


[0032]

The ester represented by the formula (3) used in this reaction may be an ester represented by the following formula (3)-1 or (3)-2.

[Formula 5]



(3)-1



(3)-2


If the pyrimidine derivative is prepared by using an ester represented by the formula (3)-1 or (3)-2, the compound represented by the formula (1) wherein R¹ is hydrogen atom can be prepared.

[0033]

Method (b):

A pyrimidine derivative represented by the formula (1) can be prepared by reacting an amidine derivative represented by the formula (2) (in the formula, R³, R⁴, R⁵, R⁶, and X have the same meanings as those defined above), and a β-ketoester represented by the following formula (5):

[Formula 6]



Formula (5)

(in the formula, R¹ and R² have the same meanings as those defined above).

[0034]

Solvent used for this reaction may be selected from conventionally used solvents. Methanol, ethanol, 1-propanol, and 2-propanol are preferred.

Reaction time, reaction temperature, and the like may be selected from conventionally used ranges. Reaction temperature is preferably 25 to 120°C, particularly preferably 60 to 80°C. Reaction time is preferably 2 to 48 hours, particularly preferably 10 to 24 hours.


The ester represented by the formula (3) used in this reaction may be an ester represented by the aforementioned formula (3)-1.

[0035]

Method (c):

A pyrimidine derivative represented by the formula (1) can be prepared by reacting a pyrimidine derivative represented by the following formula (1):


[Formula 7]



Formula (1)

(in the formula, R¹, R², R³, R⁴, R⁵, R⁶, and X have the same meanings as those defined above) with a deprotecting agent, and reacting the resultant with a compound represented by the following formula (7):

[Formula 8]



Formula (7)

(in the formula, R⁶ and X have the same meanings as those defined above).

[0036]

Examples of the deprotecting agent include, for example, inorganic acids (hydrochloric acid, sulfuric acid, nitric acid, and the like), organic acids [sulfonic acids (aliphatic sulfonic acids such as methanesulfonic acid, halogenated aliphatic sulfonic acids such as trifluoromethanesulfonic acid, aromatic sulfonic acid such as benzenesulfonic acid, and toluenesulfonic acid, and the like), carboxylic acids (halogenated carboxylic acids such as trifluoroacetic acid, and mono-, di-, or trichloroacetic acid), and the like]. Among these strong acids, inorganic acids such as hydrochloric acid, carboxylic acids such as trifluoroacetic acid, and sulfonic acids such as trifluoromethanesulfonic acid are preferred.

[0037]

Solvent used for the deprotection reaction may be selected from conventionally used solvents. Dichloromethane, and the like are preferred, and it is also preferable to perform the reaction without solvent.

Reaction time, reaction temperature, and the like of the deprotection reaction may be selected from conventionally used ranges. Reaction temperature is preferably 0 to 50°C, particularly preferably 10 to 30°C. Reaction time is preferably 1 to 48 hours, particularly preferably 3 to 24 hours.

[0038]

Solvent used for the amidation reaction may be selected from conventionally used solvents. N,N-Dimethylformamide, dichloromethane, and the like are preferred.


Reaction time, reaction temperature, and the like of the amidation reaction may be selected from conventionally used ranges. Reaction temperature is preferably -10 to 60°C, particularly preferably 0 to 30°C. Reaction time is preferably 1 to 48 hours, particularly preferably 3 to 24 hours.

[0039]

Method (d):

By reacting an amidine derivative represented by the following formula (7):

[Formula 9]




Formula (7)

(in the formula, R³ has the same meaning as that defined above),


a compound represented by the following formula (3)-1:

[Formula 10]



and an aldehyde represented by the following formula (4):


[Formula 11]



Formula (4)

(in the formula, R² has the same meaning as that defined above), a pyrimidine derivative represented by the formula (8) can be prepared.

[Formula 12]



Formula (8)

[0040]

Examples of base used for this reaction include organic bases (amines, for example, mono- to trialkylamines such as methylamine, ethylamine, diethylamine, triethylamine, propylamine, isopropylamine, and diisopropylethylamine; alkanolamines such as ethanolamine; alkylene polyamines such as ethylenediamine, and diethylenetriamine, and the like), inorganic bases [metal hydroxides (alkali metal or alkaline earth metal hydroxides, and the like) such as sodium hydroxide, potassium hydroxide, calcium hydroxide, iron hydroxide, and aluminum hydroxide; alkali metal carbonates such as sodium carbonate, sodium hydrogencarbonate, potassium carbonate, rubidium carbonate, and cesium carbonate; ammonia, and the like], and the like.

Preferred examples include alkali metal bases and organic bases, and particularly preferred examples include potassium carbonate, sodium ethoxide, and diisopropylethylamine. A salt may be formed with one or two or more kinds of these bases.

[0041]

Reaction time, reaction temperature, and the like may be selected from conventionally used ranges. Reaction temperature is preferably 0 to 140°C, particularly preferably 20 to 80°C. Reaction time is preferably 0.25 to 48 hours, particularly preferably 0.5 to 24 hours.


[0042]

Solvent used for this reaction may be selected from conventionally used solvents. Water, N,N-dimethylacetamide, methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 1-pentanol, and the like are preferred.

[0043]

Then, a pyrimidine derivative represented by the formula (1) wherein R¹, R⁴, and R⁵ are hydrogen atoms, X is C=O, and R⁶ is isopropyl group can be prepared by reacting the pyrimidine derivative represented by the formula (8) with a compound represented by the following formula:

[Formula 13]



[0044]

Examples of base used for this reaction include inorganic bases [alkali metal carbonates such as sodium carbonate, sodium hydrogencarbonate, potassium carbonate, rubidium carbonate, and cesium carbonate, alkali metal phosphates such as tripotassium phosphate], organic bases (amines, for example, trialkylamines such as triethylamine, and diisopropylethylamine), and the like. Preferred examples include sodium carbonate, potassium carbonate, and cesium carbonate.

[0045]

Examples of metal catalyst used for this reaction include, $\text{Pd}(\text{OAc})_2$, $\text{Pd}(\text{dba})_2$, $\text{Pd}_2(\text{dba})_3$, $\text{Pd}(\text{PPh}_3)_4$, $\text{PdCl}_2(\text{dppf})\cdot\text{CH}_2\text{Cl}_2$, and the like. Preferred examples include $\text{Pd}(\text{OAc})_2$, and $\text{Pd}(\text{dba})_2$.

Examples of ligand used for this reaction include XPhos, SPhos, RuPhos, and the like.

[0046]

Reaction time, reaction temperature, and the like may be selected from conventionally used ranges. Reaction temperature is preferably 30 to 180°C, particularly preferably 60 to 140°C. Reaction time is preferably 0.5 to 48 hours, particularly preferably 1 to 24 hours.

Solvent used for this reaction may be selected from conventionally used solvents. 1,4-Dioxane/water, CPME/water, toluene/water, THF/water, i-PrOH/water, EtOH/water, MeOH, THF, CPME, DME, 1,4-dioxane, and the like are preferred.

[0047]

Examples of the compounds of the present invention falling within the scope of the general formula (1) include, for example;

N-{4-chloro-3-[4-(4-chloro-2-fluorophenyl)-5-cyano-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}-2,2-dimethylpropionamide;

N-[4-chloro-3-(5-cyano-6-oxo-4-thiophen-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]-2,2-dimethylpropionamide;

N-[4-chloro-3-(5-cyano-6-oxo-4-thiophen-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;

N-[4-chloro-3-(6-oxo-4-thiophen-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;

N-{4-chloro-3-[4-(4-chloro-2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{4-chloro-3-[4-(4-chlorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{4-chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-[4-chloro-3-(6-oxo-4-phenyl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;

N-{4-chloro-3-[6-oxo-4-(6-trifluoromethylpyridin-3-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

[0048]

N-{4-chloro-3-[4-(4-chloro-3-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-[4-chloro-3-(6-oxo-4-thiophen-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]-2-methoxy-2-methylpropionamide;

N-{4-chloro-3-[6-oxo-4-(5-trifluoromethylpyridin-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{4-chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}-2-methoxy-2-methylpropionamide;

N-{4-chloro-3-[4-(3-chloro-2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(2-chloro-4-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(4-difluoromethoxy-2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(2-fluoro-4-trifluoromethoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(2-fluoro-4-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(4-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(4-difluoromethoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[6-oxo-4-(4-trifluoromethylphenyl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 [0049]
 N-{4-chloro-3-[4-(3-fluorothiophen-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-[4-chloro-3-(4-cyclopentyl-6-oxo-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
 N-{4-chloro-3-[6-oxo-4-(4-trifluoromethoxyphenyl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(2,4-difluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(2-chlorothiophen-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-[4-chloro-3-(4-cyclohexyl-6-oxo-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
 N-{4-chloro-3-[4-(2-fluoro-5-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(5-chloro-2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[6-oxo-4-(6-trifluoromethylpyridin-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 [0050]

N-{3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-[4-chloro-3-(6-oxo-4-thiophen-2-yl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
 N-[4-chloro-3-(5-methyl-6-oxo-4-phenyl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
 N-{4-chloro-3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(6-ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(3-fluoro-4-methylphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-chloro-3-{4-[4-(2-methoxyethoxy)phenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(5-methylpyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)cyclopropanecarboxamide;
 [0051]
 N-(4-chloro-3-{4-[4-(difluoromethyl)phenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[5-(trifluoromethyl)thiophen-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[2-(trifluoromethyl)thiazol-5-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(2-fluoro-4-methylphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{4-[4-(2-ethoxyethoxy)-3-fluorophenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{4-[4-(2-ethoxyethoxy)-2-fluorophenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(6-isopropoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{3-[4-(1-tert-butyl-1H-pyrazol-4-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-

chlorobenzyl}isobutyramide;

N-{4-chloro-3-[4-(1-isobutyl-1H-pyrazol-4-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{4-chloro-3-[4-(2-methoxythiazol-5-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-(4-chloro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide;

N-{4-chloro-3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}-3,3,3-trifluoro-2,2-dimethylpropionamide;

[0052]

N-(4-chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide;

N-(4-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)-2,2-dimethylpropionamide;

N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)-2,2-dimethylpropionamide;

N-(4-difluoromethoxy-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-(4-chloro-3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-(4-chloro-3-{4-[6-(3-methoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-(4-chloro-3-{4-[6-(2-methoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-{4-chloro-3-[4-(3-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

[0053]

N-(4-chloro-3-{6-oxo-4-[4-(pyridin-2-ylethynyl)phenyl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-[4-chloro-3-(5-fluoro-6-oxo-4-phenyl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;

N-{4-chloro-3-[5-fluoro-4-(4-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{4-chloro-3-[5-fluoro-6-oxo-4-(thiophen-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{4-chloro-3-[5-fluoro-6-oxo-4-(thiophen-3-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{3-[4-(benzo[b]thiophen-7-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-chloro-
 benzyl}isobutyramide;
 N-{4-chloro-3-[4-(4-cyanophenyl)-6-oxo-1,6-dihydropyrimidin-2-
 yl]benzyl}isobutyramide;
 N-(4-chloro-3-{4-[4-(methylsulfanyl)phenyl]-6-oxo-1,6-dihydropyrimidin-2-
 yl}benzyl)isobutyramide;
 N-{4-chloro-3-[6-oxo-4-(p-toluy)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-chloro-3-{4-[4-(diethylsulfamoyl)phenyl]-6-oxo-1,6-dihydropyrimidin-2-
 yl}benzyl)isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[4-(piperidine-1-sulfonyl)phenyl]-1,6-dihydropyrimidin-2-
 yl}benzyl)isobutyramide;
 [0054]
 N-{4-chloro-3-[4-(4-cyclopropylphenyl)-6-oxo-1,6-dihydropyrimidin-2-
 yl]benzyl}isobutyramide
 N-(3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-
 (trifluoromethyl)benzyl)isobutyramide;
 N-{3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-
 (trifluoromethyl)benzyl}isobutyramide;
 N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-
 (trifluoromethyl)benzyl)isobutyramide;
 N-{3-[4-(6-ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-
 (trifluoromethyl)benzyl}isobutyramide;
 N-(4-methyl-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-
 yl}benzyl)isobutyramide;
 N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-
 methylbenzyl)isobutyramide;
 N-{3-[4-(6-ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-
 methylbenzyl}isobutyramide;
 N-[4-(difluoromethyl)-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-
 2-yl}benzyl]isobutyramide;
 N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-
 (difluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[2-(trifluoromethyl)pyridin-4-yl]-1,6-dihydropyrimidin-2-
 yl}benzyl)isobutyramide;
 [0055]
 N-(4-chloro-3-{4-[2-(difluoromethoxy)pyridin-4-yl]-6-oxo-1,6-dihydropyrimidin-2-

yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(3-fluoro-4-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 4-{2-[2-chloro-5-(isobutyrylaminomethyl)phenyl]-6-oxo-1,6-dihydropyrimidin-4-yl}-N,N-dimethylbenzamide;
 N-{4-chloro-3-[4-(4-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[4-(6-methoxypyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(6-cyclopropylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{3-[4-(6-ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-fluorobenzyl}isobutyramide;
 N-[4-chloro-3-(2'-methoxy-6-oxo-1,6-dihydro[4,5']bipyrimidinyl-2-yl)benzyl]isobutyramide;
 N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-[4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl)isobutyramide;
 [0056]
 N-(4-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)-1-(trifluoromethyl)cyclopropane-1-carboxamide;
 methyl 4-{2-[2-chloro-5-(isobutyrylaminomethyl)phenyl]-6-oxo-1,6-dihydropyrimidin-4-yl}benzoate;
 N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)-1-(trifluoromethyl)cyclopropane-1-carboxamide;
 N-(2,4-difluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{4-[6-(2-butoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-chlorobenzyl)isobutyramide;

N-{4-chloro-3-[6-oxo-2'-(trifluoromethyl)-1,6-dihydro[4,5']-bipyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-chloro-3-{4-[6-(3-ethoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-[3-(4-{6-[2-(2-butoxyethoxy)ethoxy]pyridin-3-yl}-6-oxo-1,6-dihydropyrimidin-2-yl)-4-chlorobenzyl]isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[6-(pyridin-2-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 [0057]
 N-(3-{6-oxo-4-[6-(pyridin-2-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-3-{4-[6-(6-methylpyridin-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[6-oxo-4-(2-oxo-1-pentyl-1,2-dihydropyridin-4-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{3-[4-(1-butyl-2-oxo-1,2-dihydropyridin-4-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-chlorobenzyl}isobutyramide;
 N-(4-chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide;
 N-(4-chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide;
 N-(4-chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)butyramide;
 N-(4-chloro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)butyramide;
 N-{4-chloro-3-[4-(5-fluoropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)propionamide;
 N-(4-chloro-3-{4-[5-(difluoromethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)butyramide;
 N-{3-[4-(5-chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
 N-{3-[4-(5-fluoropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-

(trifluoromethyl)benzyl}isobutyramide;
N-(4-chloro-3-{4-[5-(difluoromethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide;
[0058]
N-{4-chloro-3-[4-(5-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-{3-[4-(5-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
N-(4-chloro-3-{6-oxo-4-[6-propoxypyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)propionamide;
N-(4-chloro-3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide;
N-{4-chloro-3-[4-(5-methylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-{3-[4-(5-methylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
N-{4-chloro-2-fluoro-3-[4-(5-methylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-{4-chloro-3-[4-(5-methoxypyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-{4-chloro-3-[4-(1-isopropyl-1H-pyrazol-4-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-{4-chloro-3-[6-oxo-4-(1-propyl-1H-pyrazol-4-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-(4-difluoromethoxy-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-[4-chloro-3-(4-{6-[2-(2-methoxyethoxy)ethoxy]pyridin-3-yl}-6-oxo-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
[0059]
N-{4-chloro-3-[4-(6-{2-[2-(2-ethoxyethoxy)ethoxy]ethoxy}pyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-(4-chloro-3-{4-[6-(6-methylpyridin-2-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-(4-chloro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)propionamide;
N-(4-chloro-3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-

yl}benzyl)butyramide;
N-(4-chloro-3-{4-[6-(2-fluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)butyramide;
N-(4-chloro-3-{4-[6-(3-fluoropropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide;
N-(3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
N-(3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
N-{3-[4-(6-cyclopropylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
N-(3-{4-[6-(2-butoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
[0060]
N-(3-{4-[6-(3-methoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
N-[4-chloro-3-(6-oxo-4-pyridin-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
N-{3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-methylbenzyl}isobutyramide;
N-(4-chloro-3-{6-oxo-4-[2-(trifluoromethyl)pyridin-4-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-[4-(difluoromethyl)-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl]isobutyramide;
N-(3-{4-[2-(cyclopropylmethoxy)pyridin-4-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
N-(3-{4-[6-(cyclopropylmethoxy)pyridazin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
4-(2-{2-chloro-5-[(3,3,3-trifluoro-2,2-dimethylpropionylamino)methyl]phenyl}-6-oxo-1,6-dihydropyrimidin-4-yl)-N,N-dimethylbenzamide;
N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide;
N-[3-(2'-methoxy-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide;
N-{4-chloro-3-[4-(5-chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-{4-chloro-3-[4-(6-cyclopropylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-

fluorobenzyl}isobutyramide;
 N-[4-chloro-3-(2'-cyclopropyl-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl)benzyl]isobutyramide;
 N-(4-chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 [0061]
 N-(4-chloro-3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-(3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(6-ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide;
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide;
 N-(3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide;
 N-(2-chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;
 N-(2-chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(5-chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide;
 N-(2-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(2-chloro-4-fluoro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide;
 [0062]
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(3,3,3-trifluoropropoxy)pyridin-3-yl]-1,6-

dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(2-chloro-4-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[5-(2-propoxyethoxy)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-(2-chloro-3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;
 N-(4-chloro-3-{4-[5-(difluoromethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-(4-chloro-3-{4-[5-(2-ethoxyethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-{3-[4-(5-chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-fluorobenzyl}isobutyramide;
 N-(2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-3-{4-[5-(cyclopropylmethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-{4-chloro-2-fluoro-3-[4-(5-fluoropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(2-chloro-3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;
 N-(2-chloro-3-{4-[5-(cyclopropylmethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;
 N-(2-chloro-4-fluoro-3-{6-oxo-4-[6-(3,3,3-trifluoropropoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 [0063]
 N-(4-chloro-3-{6-oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{6-oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

N-(4-chloro-3-{6-oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)butyramide;
 N-{4-chloro-2-fluoro-3-[6-oxo-4-(4-trifluoromethylthiazol-2-yl)-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{4-chloro-3-[4-(5-ethynylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[5-(1-propynyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{4-[6-(3-methyloxetan-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-{3-[4-(5-ethynylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
 N-(4-chloro-3-{4-[6-(3-methyloxetan-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{6-oxo-4-[5-(1-propynyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{4-[6-(3-ethoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(3-{6-oxo-4-[6-(3,3,3-trifluoropropoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 [0064]
 N-(3-{4-[6-(2-fluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(3-{4-[6-(3-fluoropropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(3-{4-[2-(difluoromethoxy)pyridin-4-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-

(trifluoromethyl)benzyl)isobutyramide;
N-{3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
N-(3-{6-oxo-4-[5-(2-propoxyethoxy)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
N-(3-{4-[5-(difluoromethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
N-(4-chloro-3-{4-[5-(cyclopropylmethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-(3-{4-[5-(cyclopropylmethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)butyramide;
N-(4-chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)butyramide;
N-[3-(2'-cyclopropylmethoxy-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide;
N-{3-[4-(6-butoxypyridazin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
N-{4-chloro-3-[6-oxo-2'-(trifluoromethyl)-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]benzyl}isobutyramide;
[0065]
N-(4-chloro-3-{4-[(2-cyclopropylmethoxy)pyridin-4-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-{4-chloro-2-fluoro-3-[4-(5-fluoropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
N-{2-fluoro-3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
N-(4-chloro-3-{4-[6-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
N-{2-chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-fluorobenzyl}isobutyramide;
N-{4-chloro-3-[4-(4-cyanophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide;
N-{4-chloro-2-fluoro-3-[6-oxo-4-(6-phenylpyridin-3-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-(2,4-difluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-{3-[4-(5-chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2,4-difluorobenzyl}isobutyramide;
 N-(4-chloro-3-{4-[6-(difluoromethyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-(2-chloro-3-{4-[6-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;
 N-(4-chloro-3-{4-[5-(difluoromethyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 [0066]
 N-{3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2,4-difluorobenzyl}isobutyramide;
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(2-chloro-4-fluoro-3-{6-oxo-4-[4-(trifluoromethyl)thiazol-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{4-[2-(cyclopropylethynyl)thiazol-5-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-(2-chloro-3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;
 N-(4-chloro-3-{4-[6-(3-ethyloxetan-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(3-{4-[6-(3-ethyloxetan-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-{4-chloro-2-fluoro-3-[4-(4-methylthiazol-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[4-(1-propynyl)thiazol-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(4-chloro-3-{4-[5-(cyclopropylethynyl)thiazol-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-(4-chloro-3-{4-[4-(cyclopropylethynyl)thiazol-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;
 N-(3-{4-[5-(cyclopropylethynyl)thiazol-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(3-{4-[6-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-

(trifluoromethyl)benzyl)isobutyramide;

[0067]

N-(4-chloro-3-{4-[5-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-(3-{4-[5-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

N-(3-{4-[6-(3-morpholin-4-ylpropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

N-(3-{6-oxo-4-[5-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

N-(4-chloro-2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-[3-(2'-butyl-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide;

N-(3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide;

N-(4-chloro-2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)butyramide;

N-{4-chloro-2-fluoro-3-[6-oxo-4-(6-propylpyridin-3-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{3-[6-oxo-4-(6-propylpyridin-3-yl)-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;

N-{3-[2'-(3,3-dimethyl-1-butynyl)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;

[0068]

N-{4-chloro-3-[2'-(3,3-dimethyl-1-butynyl)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-2-fluorobenzyl}isobutyramide;

N-(4-chloro-3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide;

N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(1-propynyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-(2,4-difluoro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-{4-chloro-2-fluoro-3-[4-(6-methoxypyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;

N-{2-fluoro-3-[4-(5-fluoropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-

(trifluoromethyl)benzyl}isobutyramide;
N-[4-chloro-2-fluoro-3-(4-{6-[(1-hydroxycyclohexyl)ethynyl]pyridin-3-yl}-6-oxo-1,6-dihydropyrimidin-2-yl)benzyl}isobutyramide;
N-(4-chloro-2-fluoro-3-{6-oxo-4-[5-(1-propynyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-{3-[4-(6-ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluoro-4-(trifluoromethyl)benzyl}isobutyramide;
N-{3-[4-(benzothiazol-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-chloro-2-fluorobenzyl}isobutyramide;
N-(3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
[0069]
N-(4-chloro-2-fluoro-3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-(2,4-difluoro-3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-(2-chloro-4-fluoro-3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-(3-{6-oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide;
N-(3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide;
N-(3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide;
N-(3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide;
N-{3-[4-(benzothiazol-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2,4-difluorobenzyl}isobutyramide;
N-{3-[4-(benzothiazol-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
N-(2-chloro-4-fluoro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
N-(3-{6-oxo-4-[4-(trifluoromethyl)thiazol-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
[0070]
N-{4-chloro-3-[4-(5-ethynylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-

fluorobenzyl}isobutyramide;
 N-[3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide;
 N-[4-chloro-2-fluoro-3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
 N-[2,4-difluoro-3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
 N-[2-chloro-4-fluoro-3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide;
 N-(3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide;
 N-(4-chloro-3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)butyramide;
 N-(3-{4-[5-(3,3-dimethyl-1-butynyl)pyrazin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(3-{4-[5-(cyclopropylethynyl)pyrazin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-{3-[2'-(4-methyl-1-pentynyl)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
 N-{3-[6-oxo-4-(5-propylpyridin-2-yl)-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide;
 [0071]
 N-{4-chloro-2-fluoro-3-[6-oxo-4-(5-propylpyridin-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide;
 N-{4-chloro-3-[2'-(cyclopropylmethoxy)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-2-fluorobenzyl}isobutyramide;
 N-{3-[2'-(cyclopropylmethoxy)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-4-(trifluoromethyl)benzyl}butyramide;
 N-{4-chloro-3-[2'-(cyclopropylmethoxy)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-2-fluorobenzyl}butyramide;
 N-(3-{4-[5-(cyclopropylmethoxy)pyrazin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;
 N-(4-chloro-2-fluoro-3-{6-oxo-4-[2-(1-propynyl)thiazol-5-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;
 N-(2-chloro-3-{4-[2-(cyclopropylethynyl)thiazol-5-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide;

N-(3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide;

N-(3-{4-[6-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide;

N-(2-fluoro-3-{4-[5-(isobutyrylaminomethyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

N-(2-fluoro-3-{6-oxo-4-[5-(1-propynyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

[0072]

N-(3-{4-[6-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide;

N-(3-{4-[6-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

N-(2-fluoro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

N-(2-fluoro-3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide;

N-(3-{4-[2-(cyclopropylethynyl)thiazol-5-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide;

N-[2-fluoro-3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide;

N-(4-chloro-2-fluoro-3-{6-oxo-4-[5-(2,2,2-trifluoroethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide;

N-(3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide, and the like, but the examples are not limited to these.

[0073]

The compounds represented by the general formula (1) may be in the form of salt. The salt is not particularly limited, and appropriately selected depending on the purpose. Examples include, for example, salts with alkali metals such as sodium and potassium; salts with alkaline earth metals such as calcium and magnesium; salts with organic amines such as methylamine, ethylamine, and diethanolamine, mineral acid salts such as hydrochlorides, sulfates, and nitrates, organic acid salts such as p-toluenesulfonates, maleates, and tartrates, and the like.

[0074]

The compounds represented by the general formula (1) and salts thereof may

exist in the form of hydrate or solvate. Type of solvent that forms the solvate is not particularly limited, and examples include, for example, ethanol, ethyl acetate, acetone, and the like. The compounds represented by the general formula (1) may exist as an enantiomer, diastereoisomer, or geometrical isomer depending on type of substituent, and besides arbitrary isomers in a pure form, mixtures of arbitrary isomers also fall within the scope of the present invention.

[0075]

The compounds represented by the general formula (1) and salts thereof can be easily synthesized by performing common chemical reactions widely used by those skilled in the art with starting compounds easily obtainable for those skilled in the art. Specific preparation methods of the compounds of the present invention are shown in the examples mentioned in this specification. By referring to those synthesis methods, those skilled in the art can easily prepare the compounds of the present invention falling within the scope of the general formula (1).

[0076]

The compounds of the present invention represented by the general formula (1) have an mPGES-1 inhibitory action, and can inhibit the PGE2 biosynthesis on the basis of the inhibitory action. Therefore, on the basis of the mPGES-1 inhibitory action, the medicament of the present invention containing a compound represented by the general formula (1) or a physiologically acceptable salt thereof of the present invention as an active ingredient can be used for prophylactic and/or therapeutic treatment of, for example, inflammation, pain, rheumatism, osteoarthritis, pyrexia, Alzheimer's disease, multiple sclerosis, arteriosclerosis, ocular hypertension such as glaucoma, ischemic retinopathy, systemic scleroderma, malignant tumors such as large intestine tumor, and diseases for which suppression of the PGE2 production exhibits efficacy.

[0077]

More specifically, the medicament of the present invention can be used as a medicament for prophylactic and/or therapeutic treatment of, for example, inflammatory colitis, irritable bowel syndrome, migraine, headache, low back pain, lumbar spinal canal stenosis, intervertebral disc herniation, temporomandibular arthrosis, neck-shoulder-arm syndrome, cervical spondylosis, endometriosis, adenomyosis uteri, premature delivery, threatened premature delivery, dysmenorrhea, overactive bladder, bladder outlet obstruction associated with benign prostatic hyperplasia, nocturia, urinary incontinence, neurogenic bladder, interstitial cystitis, bladder pain syndrome, urinary calculus, benign prostatic hyperplasia, chronic

prostatitis, intrapelvic pain syndrome, erectile dysfunction, cognitive disorder, neurodegenerative disease, Alzheimer's disease, pulmonary hypertension, psoriasis, rheumatoid arthritis, rheumatic fever, fibromyalgia, neuralgia, complex regional pain syndrome, fascia dyscrasia, ischemic heart disease, hypertension, angina pectoris, viral infectious disorders, bacterial infection, fungal infectious disorders, burn, inflammation and pain after operation, trauma, or extraction of a tooth, malignant tumor, myocardial infarction, atherosclerosis, thrombosis, embolism, type I diabetes mellitus, type II diabetes mellitus, cerebral apoplexy, gout, arthritis, osteoarthritis, juvenile arthritis, ankylosing spondilitis, tenosynovitis, ligamentum osteosis, systemic erythematodes, vasculitis, pancreatitis, nephritis, conjunctivitis, iritis, scleritis, uveitis, wound treatment, dermatitis, eczema, osteoporosis, asthma, chronic obstructive pulmonary disease, fibroid lung, allergic conditions, familial adenomatous polyposis, pachydermia, bursitis, hystero myoma, or pain in cancer. As for the relation of mPGES-1 inhibitory action and use as medicament, for example, International Patent Publication WO2015/125842 can be referred to. The entire disclosures of this international patent publication and all the references cited therein are incorporated into the disclosure of this specification by reference.

[0078]

Although a compound represented by the aforementioned general formula (1) or a physiologically acceptable salt thereof as the active ingredient of the medicament of the present invention may be administered as the medicament of the present invention, a pharmaceutical composition for oral or parenteral administration can be preferably prepared by a method well known to those skilled in the art, and administered. Examples of pharmaceutical composition suitable for oral administration include, for example, tablets, powders, capsules, subtilized granules, solutions, granules, syrups, and the like, and pharmaceutical composition suitable for parenteral administration include, for example, injections such as injections for intravenous injection and intramuscular injection, fusion drips, inhalants, eye drops, nose drops, suppositories, transdermal preparations, transmucosal preparations, and the like, but the pharmaceutical composition is not limited to these.

[0079]

The aforementioned pharmaceutical composition can be produced by a method well known to those skilled in the art using pharmaceutical additives commonly used for preparation of pharmaceutical compositions in this industry. Such pharmaceutical additives are not particularly limited, and can be appropriately chosen depending on form of the pharmaceutical composition, purpose thereof such as impartation of

properties for sustained release, and the like. Examples of the pharmaceutical additives include, for example, excipients, binders, fillers, disintegrating agents, surfactants, lubricants, dispersing agents, buffering agents, preservatives, corrigents, perfumes, coating agents, diluents, and the like, but the pharmaceutical additives are not limited to these.

[0080]

Dose of the medicament of the present invention is not particularly limited, and can be appropriately chosen depending on type of disease to be prevented or treated, purpose of administration such as prevention or treatment, type of active ingredient, weight, age, conditions of patient, administration route, and the like. In the case of oral administration, for example, it can be used at a dose in the range of about 0.01 to 500 mg in terms of weight of the active ingredient as the daily dose for adults. However, the dose can be appropriately chosen by those skilled in the art, and is not limited to the aforementioned range.

Examples


[0081]

Hereafter, the present invention will be explained in more detail with reference to examples. However, the present invention is not limited by these examples. The chemical formulas of the compounds of the examples are shown in Tables 1-1 to 1-18.

[0082]

Reference Example 1: N-(4-Chloro-3-cyanobenzyl)-2,2-dimethylpropionamide

[Formula 14]



[0083]

To a solution of 5-aminomethyl-2-chlorobenzonitrile (2.44 g) in methylene chloride (50 mL), N,N-diisopropylethylamine (11 mL) was added. Under ice cooling, pivaloyl chloride was added to the resulting mixture. The mixture was returned to room temperature, and stirred for 3 hours. The reaction mixture was poured into water, chloroform was added to the mixture, and the organic layer was separated. The organic layer was washed with saturated brine. The organic layer was dried over sodium sulfate, and then the solvent was evaporated. Silica gel column

chromatography was performed to obtain the title compound (3.28 g).


$^1\text{H-NMR}$ (CDCl_3 , δ): 1.24 (9H, s), 4.43 (2H, d, $J=5.9\text{Hz}$), 6.11 (1H, brs), 7.4-7.6 (3H, m)

MS (m/z): 250 (M^+)

[0084]

Reference Example 2: N-(3-Carbamimidoyl-4-chlorobenzyl)-2,2-dimethylpropanoic acid amide hydrochloride

[Formula 15]




[0085]

A suspension of ammonium chloride (2.09 g) in toluene (49 mL) was cooled on ice under nitrogen substitution, and a solution of trimethylaluminum in toluene (1.8 M, 22 mL) was added to the suspension. The resulting mixture was returned to room temperature, and stirred for 1 hour. N-(4-Chloro-3-cyanobenzyl)-2,2-dimethylpropionamide (3.26 g) was added to the mixture, and the residues were washed off into the mixture with toluene (5 mL). At an external temperature of 80°C, the mixture was stirred overnight. The reaction mixture was cooled on ice, and methanol (105 mL) was added to the mixture. The reaction mixture was warmed to 80°C, and stirred for 30 minutes. The reaction mixture was returned to room temperature, stirred for 18 minutes, and then stirred for 27 minutes under ice cooling. The insoluble matter was removed by filtration, and the solvent of the filtrate was evaporated to obtain the title compound (3.91 g). The obtained compound was used for the following reaction without purification.

[0086]

Reference Example 3: N-(4-Chloro-3-cyanobenzyl)isobutyramide

[Formula 16]



[0087]

By performing operations similar to those of Reference Example 1 using isobutyryl chloride, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 1.19 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.43 (2H, d, J=6.4Hz), 5.96 (1H, brs), 7.4-7.6 (3H, m)

MS (m/z): 236 (M⁺)

[0088]

Reference Example 4: N-(3-Carbamimidoyl-4-chlorobenzyl)isobutyramide hydrochloride

[Formula 17]




[0089]

By performing operations similar to those of Reference Example 2 using N-(4-chloro-3-cyanobenzyl)isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0090]

Reference Example 5: N-(4-Chloro-3-cyanobenzyl)-2-methoxy-2-methylpropionamide

[Formula 18]



[0091]


A mixture of 5-aminomethyl-2-chlorobenzonitrile (1.53 g), 2-methoxy-2-methylpropionic acid (876 mg), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (2.13 g), 1-hydroxybenzotriazol monohydrate (1.50 g), N,N-diisopropylethylamine (3.9 mL), and acetonitrile (18 mL) was stirred overnight. The reaction mixture was poured into saturated aqueous sodium hydrogencarbonate, and extracted with ethyl acetate. The organic layer was washed with saturated brine. The organic layer was dried over sodium sulfate, and then the solvent was evaporated. Silica gel column chromatography was performed to obtain the title compound (1.20 g). ¹H-NMR (CDCl₃, δ): 1.41 (6H, s), 3.29 (3H, s), 4.43 (2H, d, J=6.4Hz), 7.18 (1H, brs), 7.4-7.7 (3H, m)

MS (m/z): 266 (M⁺)

[0092]

Reference Example 6: N-(3-Carbamimidoyl-4-chlorobenzyl)-2-methoxy-2-methylpropionamide hydrochloride

[Formula 19]




[0093]

By performing operations similar to those of Reference Example 2 using N-(4-chloro-3-cyanobenzyl)-2-methoxy-2-methylpropionamide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0094]

Reference Example 7: tert-Butyl {3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}carbamate

[Formula 20]



[0095]

By performing operations similar to those of Example 4 using 5-chloropyridine-2-carbaldehyde, and tert-butyl (3-carbamimidoylbenzyl)carbamate, the title compound was obtained.


¹H-NMR (DMSO-d₆, δ): 1.41 (9H, m), 4.24 (2H, d, J=6.4Hz), 7.17 (1H, brs), 7.4-7.5 (4H, m), 8.0-8.1 (2H, m), 8.19 (1H, brs), 8.48 (1H, d, J=8.8Hz), 8.78 (1H, d, J=2.0Hz)

MS (m/z): 412 (M⁺)

[0096]

Reference Example 8: 2-(3-Aminomethylphenyl)-6-(5-chloropyridin-2-yl)-3H-pyrimidin-4-one

[Formula 21]



[0097]

To tert-butyl {3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}carbamate (125 mg), trifluoroacetic acid (0.5 mL) was added, and the resulting mixture was stirred at room temperature for 3 hours. The reaction mixture was neutralized with a saturated sodium hydrogencarbonate solution, and then the produced solid was taken by filtration using a filter, and dried under reduced pressure to obtain the title compound (92 mg). The obtained compound was used for the following reaction without purification.

[0098]

Example 1: N-{4-Chloro-3-[4-(4-chloro-2-fluorophenyl)-5-cyano-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}-2,2-dimethylpropionamide

To N-(3-carbamimidoyl-4-chlorobenzyl)-2,2-dimethylpropanoic acid amide hydrochloride (807 mg), potassium carbonate (611 mg), and water (15 mL) were added, and the resulting mixture was stirred for 10 minutes. Ethyl cyanoacetate (0.47 mL), and 4-chloro-2-fluorobenzaldehyde (350 mg) were added to the reaction mixture, and the resulting mixture was heated to 100°C for 40 minutes by applying microwaves. After the reaction, the reaction mixture was extracted with ethyl acetate, and the solvent was evaporated. Silica gel column chromatography was performed to obtain the title compound (48 mg).

¹H-NMR (DMSO-d₆, δ): 1.12 (9H, s), 4.28 (2H, d, J=5.8Hz), 7.3-7.8 (6H, m), 8.0-8.2 (1H, m)

MS (m/z): 472 (M⁺)

[0099]

Example 2: N-[4-Chloro-3-(5-cyano-6-oxo-4-thiophen-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]-2,2-dimethylpropionamide

By performing operations similar to those of Example 1 using thiophene-3-carbaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.13 (9H, s), 4.30 (2H, d, J=5.8Hz), 7.3-7.9 (5H, m), 8.0-8.2 (1H, m), 8.5-8.6 (1H, m), 13.78 (1H, brs)

MS (m/z): 426 (M⁺)

[0100]

Example 3: N-[4-Chloro-3-(5-cyano-6-oxo-4-thiophen-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 1 using thiophene-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.31 (2H, d, J=5.9Hz), 7.3-7.9 (5H, m), 8.3-8.4 (1H, m), 8.5-8.6 (1H, m)

MS (m/z): 412 (M⁺)

[0101]

Example 4: N-[4-Chloro-3-(6-oxo-4-thiophen-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

To N-(3-carbamimidoyl-4-chlorobenzyl)isobutyramide hydrochloride (1.039 g), potassium carbonate (990 mg), and water (15 mL) were added, and the resulting mixture was stirred for 10 minutes. Ethyl phenylsulfonylacetate (1.634 g), and thiophene-3-carbaldehyde (0.31 mL) were added to the mixture, and the resulting mixture was heated to 100°C for 40 minutes by applying microwaves. After the reaction, the reaction mixture was extracted with ethyl acetate, and the solvent was evaporated. Purification was performed by silica gel column chromatography to obtain the title compound (89 mg).

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.31 (2H, d, J=5.9Hz), 6.84 (1H, s), 7.3-7.8 (5H, m), 8.2-8.4 (2H, m), 12.45 (1H, brs)

MS (m/z): 387 (M⁺)

[0102]

Example 5: N-{4-Chloro-3-[4-(4-chloro-2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-chloro-2-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.30 (2H, d, J=5.9Hz), 6.78 (1H, s), 7.3-7.7 (5H, m), 7.9-8.1 (1H, m), 8.2-8.4 (1H, m), 12.95 (1H, brs)

MS (m/z): 433 (M⁺)

[0103]

Example 6: N-{4-Chloro-3-[4-(4-chlorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-chlorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.31 (2H, d, J=5.8Hz),

6.98 (1H, s), 7.3-7.6 (5H, m), 8.0-8.4 (3H, m), 12.89 (1H, brs)

MS (m/z): 415 (M⁺)

[0104]

Example 7: N-{4-Chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-2-carbaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.04 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.32 (2H, d, J=5.9Hz), 7.21 (1H, s), 7.4-7.7 (3H, m), 8.0-8.4 (3H, m), 8.78 (1H, d, J=1.9Hz), 13.04 (1H, brs)

MS (m/z): 416 (M⁺)

[0105]

Example 8: N-[4-Chloro-3-(6-oxo-4-phenyl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using benzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.31 (2H, d, J=5.9Hz), 6.94 (1H, s), 7.3-7.6 (6H, m), 8.0-8.2 (2H, m), 8.3-8.4 (1H, m), 12.85 (1H, brs)

MS (m/z): 381 (M⁺)

[0106]

Example 9: N-{4-Chloro-3-[6-oxo-4-(6-trifluoromethylpyridin-3-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-trifluoromethylpyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.32 (2H, d, J=6.4Hz), 7.25 (1H, s), 7.4-7.7 (3H, m), 8.03 (1H, d, J=8.3Hz), 8.2-8.4 (1H, m), 8.69 (1H, dd, J=1.4Hz, 8.3Hz), 9.40 (1H, s), 13.13 (1H, brs)

MS (m/z): 450 (M⁺)

[0107]

Example 10: N-{4-Chloro-3-[4-(4-chloro-3-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-chloro-3-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.04 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.31 (2H, d, J=6.3Hz), 7.08 (1H, s), 7.3-7.8 (4H, m), 7.97 (1H, dd, J=1.4Hz, 8.3Hz), 8.09 (1H, dd, J=1.9Hz, 9.3Hz), 8.2-8.4 (1H, m), 12.90 (1H, brs)

MS (m/z): 433 (M⁺)

[0108]

Example 11: N-[4-Chloro-3-(6-oxo-4-thiophen-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]-2-methoxy-2-methylpropionamide

By performing operations similar to those of Example 4 using thiophene-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)-2-methoxy-2-methylpropionamide hydrochloride, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.27 (6H, s), 3.16 (3H, s), 4.32 (2H, d, J=6.3Hz), 6.83 (1H, s), 7.3-7.8 (5H, m), 8.2-8.5 (2H, m), 12.75 (1H, brs)

MS (m/z): 417 (M⁺)

[0109]

Example 12: N-{4-Chloro-3-[6-oxo-4-(5-trifluoromethylpyridin-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-trifluoromethylpyridine-2-carbaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.04 (6H, d, J=7.3Hz), 2.3-2.5 (1H, m), 4.32 (2H, d, J=5.8Hz), 7.31 (1H, s), 7.4-7.7 (3H, m), 8.2-8.5 (3H, m), 9.12 (1H, d, J=0.9Hz), 13.14 (1H, brs)

MS (m/z): 450 (M⁺)

[0110]

Example 13: N-{4-Chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}-2-methoxy-2-methylpropionamide

By performing operations similar to those of Example 4 using 5-chloropyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)-2-methoxy-2-methylpropionamide hydrochloride, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.28 (6H, s), 3.17 (3H, s), 4.33 (2H, d, J=5.8Hz), 7.21 (1H, s), 7.4-7.7 (3H, m), 8.0-8.5 (3H, m), 8.78 (1H, d, J=1.9Hz), 13.06 (1H, brs)

MS (m/z): 446 (M⁺)

[0111]

Example 14: N-{4-Chloro-3-[4-(3-chloro-2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 3-chloro-2-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=7.2Hz), 2.43 (1H, sept, J=6.8Hz), 4.30 (2H, d, J=6.4Hz), 6.78 (1H, brs), 7.35 (1H, t, J=7.2Hz), 7.42 (1H, dd, J=2.0Hz, 8.0Hz), 7.5-7.6 (2H, m), 7.7-7.8 (1H, m), 7.91 (1H, t, J=6.8Hz), 8.33 (1H, t, J=6.0Hz), 13.08 (1H, s)

MS (m/z): 433 (M⁺)

[0112]

Example 15: N-{4-Chloro-3-[4-(2-chloro-4-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-chloro-4-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.02 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.29 (2H, d, J=5.6Hz), 6.62 (1H, brs), 7.3-7.5 (2H, m), 7.51 (1H, d, J=2.0Hz), 7.5-7.6 (2H, m), 7.72 (1H, dd, J=6.0Hz, 8.8Hz), 8.2-8.3 (1H, m), 13.03 (1H, brs)

MS (m/z): 433 (M⁺)

[0113]

Example 16: N-{4-Chloro-3-[4-(4-difluoromethoxy-2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-difluoromethoxy-2-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (1H, m), 4.30 (2H, d, J=5.6Hz), 6.74 (1H, brs), 7.16 (1H, dd, J=2.8Hz, 8.8Hz), 7.29 (1H, dd, J=2.4Hz, 12.8Hz), 7.40 (1H, t, J=73.2Hz), 7.41 (1H, dd, J=2.4Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.07 (1H, t, J=8.8Hz), 8.33 (1H, t, J=6.0Hz), 13.01 (1H, brs)

MS (m/z): 465 (M⁺)

[0114]

Example 17: N-{4-Chloro-3-[4-(2-fluoro-4-trifluoromethoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-fluoro-4-trifluoromethoxybenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (1H, m), 4.30 (2H, d, J=6.0Hz), 6.79 (1H, brs), 7.37 (1H, d, J=8.8Hz), 7.41 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (3H, m), 8.12 (1H, t, J=8.8Hz), 8.33 (1H, t, J=6.0Hz), 13.04 (1H, brs)

MS (m/z): 483 (M⁺)

[0115]

Example 18: N-{4-Chloro-3-[4-(2-fluoro-4-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-fluoro-4-methoxybenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.84 (3H, s), 4.31 (2H, d, J=6.0Hz), 6.68 (1H, brs), 6.90 (1H, dd, J=2.4Hz, 8.8Hz), 6.98 (1H, dd, J=2.4Hz, 13.6Hz), 7.41 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.00 (1H, t, J=8.8Hz), 8.33 (1H, t, J=6.4Hz), 12.89 (1H, brs)

MS (m/z): 429 (M⁺)

[0116]

Example 19: N-{4-Chloro-3-[4-(4-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-methoxybenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 3.82 (3H, s), 4.31 (2H, d, J=6.0Hz), 6.84 (1H, brs), 7.0-7.1 (2H, m), 7.41 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.04 (2H, d, J=8.8Hz), 8.34 (1H, t, J=6.0Hz), 12.75 (1H, brs)

MS (m/z): 411 (M⁺)

[0117]

Example 20: N-{4-Chloro-3-[4-(4-difluoromethoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-difluoromethoxybenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.5 (1H, m), 4.31 (2H, d, J=5.6Hz), 6.95 (1H, brs), 7.27 (2H, d, J=8.8Hz), 7.34 (1H, t, J=73.6Hz), 7.42 (1H, d, J=8.4Hz), 7.5-7.6 (2H, m), 8.14 (2H, d, J=8.4Hz), 8.3-8.4 (1H, m), 12.90 (1H, brs)

MS (m/z): 447 (M⁺)

[0118]

Example 21: N-{4-Chloro-3-[6-oxo-4-(4-trifluoromethylphenyl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-trifluoromethylbenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=5.6Hz), 7.08 (1H, brs), 7.42 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 7.84 (2H, d, J=8.0Hz), 8.2-8.4 (3H, m), 13.04 (1H, brs)

MS (m/z): 449 (M⁺)

[0119]

Example 22: N-{4-Chloro-3-[4-(3-fluorothiophen-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 3-fluorothiophene-2-carbaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.04 (6H, d, J=7.2Hz), 2.3-2.6 (1H, m), 4.31 (2H, d, J=6.0Hz), 6.61 (1H, brs), 7.18 (1H, d, J=5.6Hz), 7.42 (1H, d, J=8.4Hz), 7.49 (1H, s), 7.57 (1H, d, J=8.0Hz), 7.81 (1H, t, J=5.6Hz), 8.34 (1H, t, J=6.0Hz), 12.91 (1H, brs)

MS (m/z): 405 (M⁺)

[0120]

Example 23: N-[4-Chloro-3-(4-cyclopentyl-6-oxo-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

Under a nitrogen atmosphere, N-(3-carbamimidoyl-4-chlorobenzyl)isobutyramide hydrochloride (580 mg) was dissolved in ethanol (8 mL), and ethyl 3-cyclopentyl-3-oxopropanoate (405 mg) was added to the solution. Under ice cooling, a 20% solution of sodium ethoxide in ethanol (1.5 mL) was slowly added dropwise to the solution, and the resulting mixture was refluxed overnight by heating. Ethanol was removed under reduced pressure, the residue was dissolved in ethyl acetate, and the solution was successively washed with 2 N hydrochloric acid, water, and saturated brine. The solvent was removed under reduced pressure, and then the obtained residue was purified by silica gel column chromatography, and dried under reduced pressure to obtain the title compound (88 mg).

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 1.5-2.0 (8H, m), 2.42 (1H, sept, J=6.8Hz), 2.8-3.0 (1H, m), 4.29 (2H, d, J=5.6Hz), 6.19 (1H, brs), 7.38 (1H, dd, J=2.0Hz, 8.4Hz), 7.41 (1H, d, J=1.6Hz), 7.53 (1H, d, J=8.4Hz), 8.3-8.4 (1H, m), 12.62 (1H, brs)

MS (m/z): 373 (M⁺)

[0121]

Example 24: N-{4-Chloro-3-[6-oxo-4-(4-trifluoromethoxyphenyl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-trifluoromethoxybenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.4Hz), 7.01 (1H, brs), 7.4-7.6 (5H, m), 8.21 (2H, d, J=8.8Hz), 8.34 (1H, t, J=6.4Hz), 12.96 (1H, brs)

MS (m/z): 465 (M⁺)

[0122]

Example 25: N-{4-Chloro-3-[4-(2,4-difluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2,4-difluorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.30 (2H, d, J=6.0Hz), 6.73 (1H, brs), 7.2-7.3 (1H, m), 7.4-7.6 (4H, m), 8.07 (1H, ddd, J=2.4Hz, 6.8Hz, 8.8Hz), 8.32 (1H, t, J=6.4Hz), 12.96 (1H, brs)

MS (m/z): 417 (M⁺)

[0123]

Example 26: N-{4-Chloro-3-[4-(2-chlorothiophen-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-chlorothiophene-3-carbaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.42 (1H, sept, J=6.8Hz), 4.30 (2H, d, J=5.6Hz), 6.83 (1H, brs), 7.41 (1H, dd, J=2.0Hz, 8.0Hz), 7.45 (1H, d, J=6.0Hz), 7.5-7.6 (3H, m), 8.3-8.4 (1H, m), 12.96 (1H, brs)

MS (m/z): 422 (M⁺)

[0124]

Example 27: N-[4-Chloro-3-(4-cyclohexyl-6-oxo-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 23 using ethyl 3-cyclohexyl-3-oxopropanoate, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 1.2-1.4 (5H, m), 1.7-1.9 (5H, m), 2.3-2.6 (2H, m), 4.29 (2H, d, J=5.6Hz), 6.13 (1H, brs), 7.4-7.5 (2H, m), 7.52 (1H, d, J=8.0Hz), 8.3-8.4 (1H, m), 12.63 (1H, brs)

MS (m/z): 387 (M⁺)

[0125]

Example 28: N-{4-Chloro-3-[4-(2-fluoro-5-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-fluoro-5-methoxybenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (1H, m), 3.77 (3H, s), 4.31 (2H, d, J=6.0Hz), 6.76 (1H, brs), 7.0-7.1 (1H, m), 7.2-7.3 (1H, m), 7.41 (1H, dd, J=2.0Hz, 8.4Hz), 7.4-7.5 (1H, m), 7.5-7.6 (2H, m), 8.33 (1H, t, J=6.0Hz), 13.00 (1H, brs)

MS (m/z): 429 (M⁺)

[0126]

Example 29: N-{4-Chloro-3-[4-(5-chloro-2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloro-2-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.04 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.0Hz), 6.79 (1H, brs), 7.4-7.6 (5H, m), 7.98 (1H, dd, J=2.8Hz, 6.8Hz), 8.34 (1H, t, J=6.4Hz), 13.06 (1H, brs)

MS (m/z): 433 (M⁺)

[0127]

Example 30: N-{4-Chloro-3-[6-oxo-4-(6-trifluoromethylpyridin-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-trifluoromethylpyridine-2-carbaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.4Hz), 7.18 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.03 (1H, dd, J=0.8Hz, 8.0Hz), 8.25 (1H, t, J=8.0Hz), 8.34 (1H, t, J=6.4Hz), 8.49 (1H, d, J=8.0Hz), 13.13 (1H, brs)

MS (m/z): 450 (M⁺)

[0128]

Example 31: N-{4-Chloro-3-[4-(2-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.30 (2H, d, J=6.0Hz), 6.75 (1H, brs), 7.3-7.4 (3H, m), 7.5-7.6 (3H, m), 8.00 (1H, dt, J=2.0Hz, 8.0Hz), 8.33 (1H, t, J=6.0Hz), 12.98 (1H, brs)

MS (m/z): 399 (M⁺)

[0129]

Example 32: N-{3-[4-(5-Chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

To a solution of 2-(3-aminomethylphenyl)-6-(5-chloropyridin-2-yl)-3H-pyrimidin-4-one (92 mg) in N,N-dimethylformamide (4 mL), diisopropylethylamine (178 μL) was added. Under ice cooling, isobutyryl chloride (46 μL) was added dropwise to the mixture, and the resulting mixture was stirred overnight at room temperature. Ethyl acetate was added to the reaction mixture, the resulting mixture was washed twice with water, and the organic layer was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography to obtain the title compound (10 mg).

¹H-NMR (DMSO-d₆, δ): 1.07 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.37 (2H, d, J=5.6Hz), 7.19 (1H, brs), 7.4-7.6 (2H, m), 8.0-8.3 (4H, m), 8.47 (1H, d, J=8.4Hz), 8.7-8.8 (1H, m), 12.94 (1H, brs)

MS (m/z): 382 (M⁺)

[0130]

Example 33: N-[4-Chloro-3-(6-oxo-4-thiophen-2-yl)-1,6-dihydropyrimidin-2-

yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using thiophene-2-carbaldehyde, the title compound was obtained.

¹H-NMR (DMSO-d₆, δ): 1.04 (6H, d, J=7.2Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=5.6Hz), 6.86 (1H, brs), 7.1-7.2 (1H, m), 7.42 (1H, dd, J=2.0Hz, 8.4Hz), 7.50 (1H, d, J=2.0Hz), 7.57 (1H, d, J=8.4Hz), 7.77 (1H, dd, J=0.8Hz, 4.8Hz), 7.92 (1H, d, J=3.2Hz), 8.34 (1H, t, J=5.6Hz), 12.82 (1H, brs)

MS (m/z): 387 (M⁺)

[0131]

Example 34: N-[4-Chloro-3-(5-methyl-6-oxo-4-phenyl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 23 using ethyl 2-methyl-3-oxo-3-phenylpropanoate, the title compound was obtained.


¹H-NMR (DMSO-d₆, δ): 1.02 (6H, d, J=6.8Hz), 2.07 (3H, s), 2.3-2.6 (1H, m), 4.28 (2H, d, J=6.0Hz), 7.18 (1H, brs), 7.3-7.6 (7H, m), 8.3-8.4 (1H, m), 12.38 (1H, brs)

MS (m/z): 395 (M⁺)

[0132]

Reference Example 9: N-(4-Chloro-3-cyanobenzyl)cyclopropanecarboxamide

[Formula 22]



[0133]

By performing operations similar to those of Reference Example 1 using cyclopropanecarbonyl chloride, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 0.7-1.1 (4H, m), 1.3-1.5 (1H, m), 4.44 (2H, d, J=5.8Hz), 6.16 (1H, brs), 7.4-7.5 (2H, m), 7.58 (1H, brs)

MS (m/z): 234 (M⁺)

[0134]

Reference Example 10: N-(3-Carbamimidoyl-4-chlorobenzyl)cyclopropanecarboxamide hydrochloride

[Formula 23]




[0135]

By performing operations similar to those of Reference Example 2 using N-(4-chloro-3-cyanobenzyl)cyclopropanecarboxamide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0136]

Reference Example 11: 4-(2-Ethoxyethoxy)-2-fluorobenzaldehyde

[Formula 24]



[0137]

2-Fluoro-4-hydroxybenzaldehyde (560 mg), 1-bromo-2-ethoxyethane (1.84 g), potassium carbonate (1.11 g), and acetonitrile (10 mL) were mixed, and the mixture was heated to 85°C under a nitrogen atmosphere. After 16 hours, the reaction mixture was left to cool. The reaction mixture was poured into brine, the resulting mixture was extracted with ethyl acetate, and then the solvent was evaporated. Silica gel column chromatography was performed to obtain the title compound (670 mg).


¹H-NMR (CDCl₃, δ): 1.25 (3H, t, J=7.1Hz), 3.60 (2H, q, J=7.0Hz), 3.7-3.9 (2H, m), 4.1-4.3 (2H, m), 6.68 (1H, dd, J=2.4Hz, 12.7Hz), 6.81 (1H, dd, J=2.4Hz, 8.8Hz), 7.81 (1H, t, J=8.3Hz), 10.21 (1H, s)

MS (m/z): 212 (M⁺)

[0138]

Reference Example 12: N-(4-Chloro-3-cyanobenzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide

[Formula 25]



[0139]

By performing operations similar to those of Reference Example 5 using 3,3,3-

trifluoro-2,2-dimethylpropionic acid, the title compound was obtained.


$^1\text{H-NMR}$ (CDCl_3 , δ): 1.44 (6H, s), 4.48 (2H, d, $J=6.4\text{Hz}$), 6.33 (1H, brs), 7.43 (1H, dd, $J=1.9\text{Hz}$, 8.3Hz), 7.48 (1H, d, $J=8.3\text{Hz}$), 7.54 (1H, d, $J=2.5\text{Hz}$)

MS (m/z): 304 (M^+)

[0140]

Reference Example 13: N-(3-Carbamimidoyl-4-chlorobenzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide hydrochloride

[Formula 26]




[0141]

By performing operations similar to those of Reference Example 2 using N-(4-chloro-3-cyanobenzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0142]

Reference Example 14: N-(3-Cyano-4-fluorobenzyl)-2,2-dimethylpropionamide

[Formula 27]



[0143]

By performing operations similar to those of Reference Example 1 using 5-aminomethyl-2-fluorobenzonitrile, the title compound was obtained.


$^1\text{H-NMR}$ (CDCl_3 , δ): 1.23 (9H, s), 4.42 (2H, d, $J=6.4\text{Hz}$), 6.10 (1H, brs), 7.18 (1H, t, $J=8.6\text{Hz}$), 8.4-8.6 (2H, m)

MS (m/z): 234 (M^+)

[0144]

Reference Example 15: N-(3-Carbamimidoyl-4-fluorobenzyl)-2,2-dimethylpropionamide hydrochloride

[Formula 28]




[0145]

By performing operations similar to those of Reference Example 2 using N-(3-cyano-4-fluorobenzyl)-2,2-dimethylpropionamide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0146]

Reference Example 16: [3-Cyano-4-(difluoromethoxy)benzyl]carbamic acid tert-butyl ester

[Formula 29]




[0147]

5-Bromo-2-(difluoromethoxy)benzonitrile (4.48 g), potassium N-BOC-aminomethyltrifluoroborate (4.50 g), palladium acetate (205 mg), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (745 mg), potassium carbonate (7.51 g), toluene (60 mL), and water (15 mL) were mixed, and the mixture was heated to 85°C under a nitrogen atmosphere. After 17 hours, the reaction mixture was left to cool. The reaction mixture was poured into water, the resulting mixture was extracted with ethyl acetate, and then the organic layer was dried over sodium sulfate. The solvent was evaporated, and then silica gel column chromatography was performed to obtain the title compound (6.38 g).

[0148]

Reference Example 17: 5-Aminomethyl-2-(difluoromethoxy)benzonitrile

[Formula 30]



[0149]

To [3-cyano-4-(difluoromethoxy)benzyl]carbamic acid tert-butyl ester (6.38 g),

dichloromethane (25 mL), and trifluoroacetic acid (12.4 mL) were added, and the resulting mixture was stirred at room temperature for 1.2 hours. The reaction mixture was poured into water, and the aqueous layer was washed twice with chloroform. To the aqueous layer, 2 N aqueous sodium hydroxide (100 mL) was added, and the resulting mixture was extracted twice with chloroform. The organic layer was dried over sodium sulfate, and then the solvent was evaporated to obtain the title compound (2.73 g).


$^1\text{H-NMR}$ (CDCl_3 , δ): 1.41 (2H, brs), 3.92 (2H, s), 6.63 (1H, t, $J=71.8\text{Hz}$), 7.2-7.4 (1H, m), 7.58 (1H, dd, $J=1.9\text{Hz}$, 8.3Hz), 7.68 (1H, d, $J=1.9\text{Hz}$)

MS (m/z): 197 (M^+-1)

[0150]

Reference Example 18: N-[3-Cyano-4-(difluoromethoxy)benzyl]isobutyramide

[Formula 31]



[0151]

By performing operations similar to those of Reference Example 1 using 5-aminomethyl-2-(difluoromethoxy)benzonitrile, and isobutyryl chloride, the title compound was obtained.


$^1\text{H-NMR}$ (CDCl_3 , δ): 1.19 (6H, d, $J=6.8\text{Hz}$), 2.42 (1H, sept, $J=6.8\text{Hz}$), 4.44 (2H, d, $J=6.3\text{Hz}$), 5.97 (1H, brs), 6.63 (1H, t, $J=71.6\text{Hz}$), 7.2-7.4 (1H, m), 7.53 (1H, dd, $J=2.4\text{Hz}$, 8.8Hz), 7.57 (1H, d, $J=2.0\text{Hz}$)

MS (m/z): 268 (M^+)

[0152]

Reference Example 19: N-[3-Carbamimidoyl-4-(difluoromethoxy)benzyl]isobutyramide hydrochloride

[Formula 32]




[0153]

By performing operations similar to those of Reference Example 2 using N-[3-cyano-4-(difluoromethoxy)benzyl]isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0154]

Reference Example 20: 6-(3-Methoxypropoxy)pyridine-3-carbaldehyde

[Formula 33]



[0155]

To 3-methoxypropan-1-ol (25 g), 60% sodium hydride (1.12 g) was added portionwise under ice cooling. Then, 6-chloropyridine-3-carbaldehyde (2.48 g) was added portionwise to the mixture. The resulting mixture was returned to room temperature, and stirred for 14 hours under a nitrogen atmosphere. The reaction mixture was poured into saturated aqueous ammonium chloride, and the resulting mixture was extracted with t-butyl methyl ether. The organic layer was washed with saturated brine, and the solvent was evaporated. Silica gel column chromatography was performed to obtain the title compound (866 mg).


¹H-NMR (CDCl₃, δ): 2.0-2.2 (2H, m), 3.36 (3H, s), 3.55 (2H, t, J=6.4Hz), 4.50 (2H, t, J=6.6Hz), 6.83 (1H, d, J=8.8Hz), 8.06 (1H, dd, J=2.5Hz, 8.8Hz), 8.62 (1H, d, J=1.9Hz), 9.95 (1H, s)

MS (m/z): 196 (M⁺⁺1)

[0156]

Reference Example 21: 4-(Pyridin-2-ylethynyl)benzaldehyde

[Formula 34]



[0157]

To a solution of 4-ethynylbenzaldehyde (1.24 g) in triethylamine (20 mL), 2-bromopyridine (975 μL), bis(triphenylphosphine)palladium chloride (133 mg), and copper iodide (46 mg) were added under a nitrogen atmosphere, and the resulting

mixture was refluxed by heating for 2 hours. After cooling, triethylamine was removed under reduced pressure, chloroform was added to the obtained residue, and the resulting mixture was filtered through a Celite layer. The filtrate was washed successively with 15% potassium carbonate, water, and saturated brine, dried over magnesium sulfate, and filtered, and the solvent was evaporated under reduced pressure. To the obtained residue, ethanol was added, and the resulting mixture was heated, cooled, filtered, and dried to obtain the title compound (953 mg).


$^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$, δ): 7.4-7.5 (1H, m), 7.7-7.8 (1H, m), 7.8-7.9 (3H, m), 7.9-8.0 (2H, m), 8.6-8.7 (1H, m), 10.06 (1H, s)

MS (m/z): 207 (M^+)

[0158]

Reference Example 22: Ethyl 2-fluoro-3-oxo-3-phenylpropanoate

[Formula 35]



[0159]

To a solution of ethyl 3-oxo-3-phenylpropanoate (961 mg) in acetonitrile (20 mL), Selectfluor (registered trade mark, 1.95 g) was added under a nitrogen atmosphere, and the resulting mixture was stirred at 80°C for 6 hours. After cooling, acetonitrile was removed under reduced pressure. To the obtained residue, ethyl acetate was added, and the organic layer was washed successively with water, and saturated brine, and dried over magnesium sulfate. The organic layer was filtered, and the solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography to obtain the title compound (858 mg).


$^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$, δ): 1.26 (3H, t, $J=7.2\text{Hz}$), 4.2-4.3 (2H, m), 5.86 (1H, d, $J=48.8\text{Hz}$), 7.4-7.5 (2H, m), 7.6-7.7 (1H, m), 8.0-8.1 (2H, m)

MS (m/z): 210 (M^+)

[0160]

Reference Example 23: Ethyl 2-fluoro-3-(4-methoxyphenyl)-3-oxopropanoate

[Formula 36]



[0161]

By performing operations similar to those of Reference Example 22 using ethyl 4-methoxybenzoate, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 1.27 (3H, t, J=7.2Hz), 3.89 (3H, s), 4.2-4.3 (2H, m), 5.81 (1H, d, J=48.8Hz), 6.9-7.0 (2H, m), 8.0-8.1 (2H, m)

MS (m/z): 240 (M⁺)

[0162]

Reference Example 24: Ethyl 2-fluoro-3-oxo-3-(thiophen-2-yl)propanoate

[Formula 37]



[0163]

By performing operations similar to those of Reference Example 22 using ethyl 3-oxo-3-(thiophen-2-yl)propanoate, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.16 (3H, t, J=6.8Hz), 4.22 (2H, q, J=6.8Hz), 6.53 (1H, d, J=46.8Hz), 7.35 (1H, dd, J=4.8Hz), 8.1-8.2 (1H, m), 8.22 (1H, dd, J=0.8Hz, 4.8Hz)

MS (m/z): 216 (M⁺)

[0164]

Reference Example 25: Ethyl 3-oxo-3-(thiophen-3-yl)propanoate

[Formula 38]



[0165]

Under a nitrogen atmosphere, a solution of diethyl carbonate (2.3 g) in toluene (10 mL) was warmed to 65°C, *t*-butoxy potassium (875 mg) was added to the solution, and the resulting mixture was stirred at 65°C for 30 minutes. The reaction mixture was warmed to 80°C, and then a solution of 3-acetylthiophene (618 mg) dissolved in toluene (5 mL) was added dropwise, and the resulting mixture was stirred at 80°C for 1.5 hours. The reaction mixture was left to cool to room temperature, then ethyl acetate was added to the reaction mixture, and the organic layer was washed successively with water, and saturated brine, and dried over magnesium sulfate. The obtained residue was purified by silica gel column chromatography to obtain the title compound (765 mg).


¹H-NMR (CDCl₃, δ): 1.26 (3H, t, J=7.2Hz), 3.89 (2H, s), 4.21 (2H, q, J=7.2Hz), 7.3-7.4 (1H, m), 7.56 (1H, dd, J=1.6Hz, 5.6Hz), 8.11 (1H, dd, J=1.6Hz, 2.8Hz)

MS (m/z): 198 (M⁺)

[0166]

Reference Example 26: Ethyl 2-fluoro-3-oxo-3-(thiophen-3-yl)propanoate

[Formula 39]



[0167]

By performing operations similar to those of Reference Example 22 using ethyl 3-oxo-3-(thiophen-3-yl)propanoate, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 1.28 (3H, t, J=6.8Hz), 4.2-4.4 (2H, m), 5.65 (1H, d, J=49.2Hz), 7.36 (1H, dd, J=2.8Hz, 4.8Hz), 7.65 (1H, dd, J=0.8Hz, 6.4Hz), 8.40 (1H, quin, J=1.6Hz)

MS (m/z): 216 (M⁺)

[0168]

Reference Example 27: N,N-Diethyl-4-formylbenzenesulfonamide

[Formula 40]



[0169]

To a solution of 4-formylbenzenesulfonyl chloride (1.0 g) in chloroform (10 mL), diethylamine (731 mg), and then saturated aqueous sodium hydrogencarbonate (11 mL) were added, and the resulting mixture was vigorously stirred at room temperature for 2 hours. The chloroform layer was separated, washed with saturated brine, dried over magnesium sulfate, and filtered, and the solvent was evaporated under reduced pressure. To the obtained residue, a mixture of hexane and ethyl acetate (3:1) was added, and the resulting mixture was heated, cooled, filtered, and dried to obtain the title compound (946 mg).


¹H-NMR (d₆-DMSO, δ): 1.05 (6H, t, J=7.2Hz), 3.21 (4H, q, J=7.2Hz), 7.9-8.0 (2H, m), 8.0-8.1 (2H, m), 10.11 (1H, s)

MS (m/z): 241 (M⁺)

[0170]

Reference Example 28: 4-(Piperidine-1-sulfonyl)benzaldehyde

[Formula 41]



[0171]

To a solution of 4-formylbenzenesulfonyl chloride (818 mg) in chloroform (20 mL), diisopropylethylamine (1.4 mL), and then piperidine (394 μ L) were added, and the resulting mixture was stirred overnight at room temperature. Water was added to the reaction mixture, and the chloroform layer was washed twice with water. The chloroform layer was further washed with saturated brine, dried over magnesium sulfate, and filtered, and the solvent was evaporated under reduced pressure to obtain the title compound (988 mg).


$^1\text{H-NMR}$ (d_6 -DMSO, δ): 1.4-1.5 (2H, m), 1.65 (4H, quin, $J=5.6\text{Hz}$), 3.04 (4H, t, $J=5.6\text{Hz}$), 7.93 (2H, dd, $J=2.0\text{Hz}$, 6.8Hz), 8.04 (2H, dd, $J=2.0\text{Hz}$, 6.4Hz), 10.12 (1H, s)

MS (m/z): 253 (M^+)

[0172]

Reference Example 29: N-[3-Cyano-4-(trifluoromethyl)benzyl]isobutyramide

[Formula 42]



[0173]

By performing operations similar to those of Reference Example 1 using 5-aminomethyl-2-(trifluoromethyl)benzonitrile, and isobutyryl chloride, the title compound was obtained.


$^1\text{H-NMR}$ (d_6 -DMSO, δ): 1.05 (6H, d, $J=7.6\text{Hz}$), 2.45 (1H, sept, $J=6.8\text{Hz}$), 4.38 (2H, d, $J=6.0\text{Hz}$), 7.76 (1H, d, $J=8.8\text{Hz}$), 7.9-8.0 (2H, m), 8.40 (1H, t, $J=6.0\text{Hz}$)

MS (m/z): 270 (M^+)

[0174]

Reference Example 30: N-[3-Carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride

[Formula 43]




[0175]

By performing operations similar to those of Reference Example 2 using N-[3-cyano-4-(trifluoromethyl)benzyl]isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

Reference Example 31: N-(3-Bromo-4-methylbenzyl)isobutyramide

[Formula 44]



[0176]

By performing operations similar to those of Reference Example 1 using 4-aminomethyl-2-bromo-1-methylbenzene, and isobutyryl chloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.02 (6H, d, J=6.8Hz), 2.2-2.5 (4H, m), 4.20 (2H, d, J=6.0Hz), 7.14 (1H, dd, J=2.0Hz, 7.6Hz), 7.29 (1H, d, J=7.6Hz), 7.42 (1H, d, J=1.2Hz), 8.23 (1H, t, J=5.2Hz)

MS (m/z): 269 (M⁺-1)

[0177]

Reference Example 32: N-(3-Cyano-4-methylbenzyl)isobutyramide

[Formula 45]



[0178]

To a solution of N-(3-bromo-4-methylbenzyl)isobutyramide (2.1 g) in N-methylpyrrolidinone (15 mL), potassium ferrocyanide trihydrate (1.7 g), palladium acetate (37 mg), diphenylphosphinoferrocene (172 mg), and potassium carbonate (1.6 g) were added under a nitrogen atmosphere, and the resulting mixture was stirred


overnight at 160°C. The reaction mixture was left to cool to room temperature, and then filtered through a Celite layer, ethyl acetate and water were added to the filtrate, and the organic layer was washed with a saturated sodium hydrogencarbonate solution, and saturated brine, and dried over magnesium sulfate. The organic layer was filtered, the solvent was evaporated under reduced pressure, and the obtained residue was purified by silica gel column chromatography to obtain the title compound (1.1 g). ¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (4H, m), 4.24 (2H, d, J=5.6Hz), 7.4-7.5 (2H, m), 7.58 (1H, s), 8.30 (1H, t, J=6.0Hz)

MS (m/z): 216 (M⁺)

[0179]

Reference Example 33: N-(3-Carbamimidoyl-4-methylbenzyl)isobutyramide hydrochloride

[Formula 46]




[0180]

By performing operations similar to those of Reference Example 2 using N-(3-cyano-4-methylbenzyl)isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0181]

Reference Example 34: N-[3-Cyano-4-(difluoromethyl)benzyl]isobutyramide

[Formula 47]



[0182]

By performing operations similar to those of Reference Example 1 using 5-aminomethyl-2-(difluoromethyl)benzonitrile and isobutyryl chloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.44 (1H, sept, J=6.8Hz), 4.34 (2H, d, J=5.6Hz), 7.23 (1H, t, J=5.4Hz), 7.69 (1H, d, J=8.0Hz), 7.78 (1H, d, J=8.0Hz), 7.85 (1H,


s), 8.36 (1H, t, J=5.6Hz)

MS (m/z): 252 (M⁺)

[0183]

Reference Example 35: N-[3-Carbamimidoyl-4-(difluoromethyl)benzyl]isobutyramide hydrochloride

[Formula 48]




[0184]

By performing operations similar to those of Reference Example 2 using N-[3-cyano-4-(difluoromethyl)benzyl]isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0185]

Reference Example 36: N-(3-Cyano-4-fluorobenzyl)isobutyramide

[Formula 49]



[0186]

By performing operations similar to those of Reference Example 1 using 5-aminomethyl-2-fluorobenzonitrile, and isobutyryl chloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=7.2Hz), 2.42 (1H, sept, J=6.8Hz), 4.26 (2H, d, J=5.6Hz), 7.49 (1H, t, J=8.8Hz), 7.6-7.7 (1H, m), 7.7-7.8 (1H, m), 8.30 (1H, t, J=5.6Hz)

MS (m/z): 220 (M⁺)

[0187]

Reference Example 37: N-(3-Carbamimidoyl-4-fluorobenzyl)isobutyramide hydrochloride

[Formula 50]




[0188]

By performing operations similar to those of Reference Example 2 using N-(3-cyano-4-fluorobenzyl)isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0189]

Reference Example 38: 3-Bromomethyl-6-chloro-2-fluorobenzonitrile

[Formula 51]



[0190]

Under a nitrogen atmosphere, 6-chloro-2-fluoro-3-methylbenzonitrile (2.29 g) was dissolved in acetonitrile (54 mL), 1,3-dibromo-5,5-dimethylhydantoin (1.93 g), and 2,2'-azobis(isobutyronitrile) (222 mg) were added to the solution, and the resulting mixture was refluxed for 3 hours by heating. 1 N Hydrochloric acid was added to the reaction mixture, the resulting mixture was extracted with ethyl acetate, and the solvent was removed under reduced pressure. The obtained residue was purified by silica gel column chromatography, and dried under reduced pressure to obtain the title compound (2.40 g).


¹H-NMR (d₆-DMSO, δ): 4.74 (2H, d, J=1.2Hz), 7.66 (1H, dd, J=1.2Hz, 8.8Hz), 7.95 (1H, t, J=8.8Hz)

MS (m/z): 249 (M⁺⁺¹)

[0191]

Reference Example 39: 3-Aminomethyl-6-chloro-2-fluorobenzonitrile

[Formula 52]



[0192]


3-Bromomethyl-6-chloro-2-fluorobenzonitrile (2.34 g) was dissolved in N,N-dimethylformamide (24 mL), sodium azide (613 mg) was added to the solution under ice cooling, and the resulting mixture was stirred for 35 minutes. Saturated brine was added to the reaction mixture, the resulting mixture was extracted with ethyl acetate, and then the organic layer was washed with saturated brine. The solvent was removed under reduced pressure, and the obtained compound was used for the following reaction without purification.

The obtained compound was dissolved in tetrahydrofuran (24 mL), and water (1 mL), triphenylphosphine (2.97 g) was added to the solution under ice cooling, and the resulting mixture was stirred overnight at room temperature. 1 N Hydrochloric acid was added to the reaction mixture, and the resulting mixture was washed with diethyl ether. The aqueous layer was made alkaline by adding 1 N sodium hydroxide. The aqueous layer was extracted with dichloromethane, then the organic layer was dried over magnesium sulfate, and the obtained organic layer was used as it was for the following reaction.

[0193]

Reference Example 40: N-(4-Chloro-3-cyano-2-fluorobenzyl)isobutyramide

[Formula 53]



[0194]

By performing operations similar to those of Reference Example 1 using 3-aminomethyl-6-chloro-2-fluorobenzonitrile, and isobutyryl chloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.02 (6H, d, J=6.8Hz), 2.41 (1H, sept, J=6.8Hz), 4.29 (2H, d, J=5.6Hz), 7.5-7.7 (2H, m), 8.33 (1H, t, J=5.6Hz)

MS (m/z): 254 (M⁺)

[0195]

Reference Example 41: N-(3-Carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride

[Formula 54]




[0196]

By performing operations similar to those of Reference Example 2 using N-(4-chloro-3-cyano-2-fluorobenzyl)isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0197]

Reference Example 42: N-(3-Cyano-4-fluorobenzyl)-1-(trifluoromethyl)cyclopropane-1-carboxamide

[Formula 55]



[0198]

By performing operations similar to those of Reference Example 1 using 5-aminomethyl-2-fluorobenzonitrile, and 1-(trifluoromethyl)cyclopropane-1-carbonylchloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.2-1.4 (4H, m), 4.29 (2H, d, J=6.0Hz), 7.49 (1H, t, J=8.8Hz), 7.5-7.7 (1H, m), 7.7-7.8 (1H, m), 8.41 (1H, t, J=5.6Hz)

MS (m/z): 286 (M⁺)

[0199]

Reference Example 43: N-(3-Carbamimidoyl-4-fluorobenzyl)-1-(trifluoromethyl)cyclopropane-1-carboxamide hydrochloride

[Formula 56]



[0200]


By performing operations similar to those of Reference Example 2 using N-(3-cyano-4-fluorobenzyl)-1-(trifluoromethyl)cyclopropane-1-carboxamide, the title

compound was obtained. The obtained compound was used for the following reaction without purification.

[0201]

Reference Example 44: N-(3-Cyano-2,4-difluorobenzyl)isobutyramide

[Formula 57]



[0202]

By performing operations similar to those of Reference Example 1 using 3-aminomethyl-2,6-difluorobenzonitrile, and isobutyryl chloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.02 (6H, d, J=6.8Hz), 2.41 (1H, sept, J=6.8Hz), 4.28 (2H, d, J=6.0Hz), 7.3-7.5 (1H, m), 7.6-7.8 (1H, m), 8.32 (1H, t, J=5.6Hz)

MS (m/z): 238 (M⁺)

[0203]

Reference Example 45: N-(3-Carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride

[Formula 58]




[0204]

By performing operations similar to those of Reference Example 2 using N-(3-cyano-2,4-difluorobenzyl)isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0205]

Reference Example 46: 6-(2-Propoxyethoxy)pyridine-3-carbaldehyde

[Formula 59]



[0206]

By performing operations similar to those of Reference Example 20 using 2-propoxyethanol, the title compound was obtained.

¹H-NMR (CDCl₃, δ): 0.93 (3H, t, J=7.3Hz), 1.5-1.8 (2H, m), 3.49 (2H, t, J=6.8Hz), 3.7-3.9 (2H, m), 4.5-4.7 (2H, m), 6.90 (1H, d, J=8.3Hz), 8.06 (1H, dd, J=2.5Hz, 8.8Hz), 8.61 (1H, d, J=1.9Hz), 9.95 (1H, s)

MS (m/z): 210 (M⁺⁺¹)

[0207]

Example 35: N-{4-Chloro-3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-methoxypyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.92 (3H, s), 4.31 (2H, d, J=5.8Hz), 6.91 (1H, d, J=8.3Hz), 6.96 (1H, brs), 7.3-7.6 (3H, m), 8.2-8.4 (2H, m), 8.89 (1H, d, J=2.0Hz), 12.79 (1H, brs)

MS (m/z): 412 (M⁺)

[0208]

Example 36: N-{4-Chloro-3-[4-(6-ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-ethoxypyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.34 (3H, t, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=5.8Hz), 4.37 (2H, q, J=7.0Hz), 6.88 (1H, d, J=8.8Hz), 6.95 (1H, brs), 7.3-7.6 (3H, m), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.81 (1H, brs)

MS (m/z): 426 (M⁺)

[0209]

Example 37: N-{4-Chloro-3-[4-(3-fluoro-4-methylphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 3-fluoro-4-methylbenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.29 (3H, d, J=1.0Hz), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.4Hz), 6.98 (1H, brs), 7.3-7.9 (6H, m), 8.2-8.4 (1H, m), 12.91 (1H, brs)

MS (m/z): 413 (M⁺)

[0210]

Example 38: N-(4-Chloro-3-{4-[4-(2-methoxyethoxy)phenyl]-6-oxo-1,6-

dihydropyrimidin-2-yl}benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-(2-methoxyethoxy)benzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.32 (3H, s), 3.6-3.8 (2H, m), 4.1-4.2 (2H, m), 4.31 (2H, d, J=6.3Hz), 6.84 (1H, brs), 7.03 (2H, d, J=9.3Hz), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.53 (1H, d, J=2.0Hz), 7.56 (1H, d, J=8.3Hz), 8.03 (2H, d, J=8.8Hz), 8.2-8.4 (1H, m), 12.77 (1H, brs)

MS (m/z): 455 (M⁺)

[0211]

Example 39: N-{4-Chloro-3-[4-(5-methylpyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-methylpyridine-2-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.37 (3H, s), 2.3-2.5 (1H, m), 4.32 (2H, d, J=5.9Hz), 7.20 (1H, brs), 7.42 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 7.76 (1H, dd, J=2.4Hz, 8.3Hz), 8.14 (1H, d, J=8.3Hz), 8.3-8.4 (1H, m), 8.57 (1H, d, J=2.4Hz)

MS (m/z): 396 (M⁺)

[0212]

Example 40: N-(4-Chloro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)cyclopropanecarboxamide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)cyclopropanecarboxamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.6-0.8 (4H, m), 1.5-1.7 (1H, m), 4.35 (2H, d, J=6.4Hz), 7.25 (1H, brs), 7.4-7.7 (3H, m), 8.03 (1H, d, J=8.3Hz), 8.5-8.8 (2H, m), 9.41 (1H, s), 13.13 (1H, brs)

MS (m/z): 448 (M⁺)

[0213]

Example 41: N-(4-Chloro-3-{4-[4-(difluoromethyl)phenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-(difluoromethyl)benzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.4Hz), 7.03 (1H, brs), 7.10 (1H, t, J=56.0Hz), 7.42 (1H, dd, J=2.4Hz, 8.3Hz), 7.56 (1H, d, J=2.0Hz), 7.58 (1H, d, J=8.3Hz), 7.68 (2H, d, J=8.3Hz), 8.21 (2H, d, J=8.3Hz), 8.3-8.4 (1H, m), 12.96 (1H, brs)

MS (m/z): 431 (M⁺)

[0214]

Example 42: N-(4-Chloro-3-{6-oxo-4-[5-(trifluoromethyl)thiophen-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)thiophene-2-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.44 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.3Hz), 7.12 (1H, brs), 7.43 (1H, dd, J=2.4Hz, 8.3Hz), 7.52 (1H, d, J=2.0Hz), 7.58 (1H, d, J=8.3Hz), 7.68 (1H, dd, J=1.2Hz, 4.2Hz), 8.04 (1H, dd, J=1.5Hz, 3.9Hz), 8.3-8.4 (1H, m), 13.01 (1H, brs)

MS (m/z): 455 (M⁺)

[0215]

Example 43: N-(4-Chloro-3-{6-oxo-4-[2-(trifluoromethyl)thiazol-5-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(trifluoromethyl)thiazole-5-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=5.9Hz), 7.24 (1H, brs), 7.44 (1H, dd, J=1.9Hz, 8.4Hz), 7.52 (1H, d, J=1.9Hz), 7.58 (1H, d, J=8.3Hz), 8.2-8.4 (1H, m), 8.91 (1Hs), 13.15 (1H, brs)

MS (m/z): 456 (M⁺)

[0216]

Example 44: N-{4-Chloro-3-[4-(2-fluoro-4-methylphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-fluoro-4-methylbenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.37 (3H, s), 2.3-2.5 (1H, m), 4.31 (2H, d, J=5.9Hz), 6.74 (1H, brs), 7.14 (1H, d, J=7.8Hz), 7.19 (1H, d, J=12.7Hz), 7.41 (1H, dd, J=2.4Hz, 8.3Hz), 7.54 (1H, d, J=1.9Hz), 7.57 (1H, d, J=8.3Hz), 7.92 (1H, t, J=8.1Hz), 8.2-8.4 (1H, m), 12.93 (1H, brs)

MS (m/z): 413 (M⁺)

[0217]

Example 45: N-(4-Chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=5.8Hz), 7.08 (1H, brs), 7.20 (1H, d, J=8.8Hz), 7.42 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6

(2H, m), 7.77 (1H, t, J=73.0Hz), 8.2-8.4 (1H, m), 8.53 (1H, dd, J=2.5Hz, 8.3Hz), 8.95 (1H, d, J=2.0Hz), 12.97 (1H, brs)

MS (m/z): 448 (M⁺)

[0218]

Example 46: N-(4-Chloro-3-{4-[4-(2-ethoxyethoxy)-3-fluorophenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-(2-ethoxyethoxy)-3-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 1.13 (3H, t, J=7.1Hz), 2.44 (1H, sept, J=6.8Hz), 3.52 (2H, q, J=7.0Hz), 3.7-3.8 (2H, m), 4.2-4.3 (2H, m), 4.32 (2H, d, J=6.4Hz), 6.94 (1H, brs), 7.27 (1H, t, J=8.8Hz), 7.42 (1H, dd, J=2.0Hz, 8.3Hz), 7.54 (1H, d, J=2.0Hz), 7.57 (1H, d, J=8.3Hz), 7.8-8.0 (2H, m), 8.2-8.4 (1H, m), 12.80 (1H, brs)

MS (m/z): 487 (M⁺)

[0219]

Example 47: N-(4-Chloro-3-{4-[4-(2-ethoxyethoxy)-2-fluorophenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-(2-ethoxyethoxy)-2-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.13 (3H, t, J=7.1Hz), 2.43 (1H, sept, J=6.8Hz), 3.50 (2H, q, J=7.0Hz), 3.6-3.8 (2H, m), 4.1-4.3 (2H, m), 4.31 (2H, d, J=5.9Hz), 6.70 (1H, brs), 6.91 (1H, dd, J=2.5Hz, 8.8Hz), 6.99 (1H, dd, J=2.4Hz, 14.2Hz), 7.41 (1H, dd, J=2.0Hz, 8.3Hz), 7.54 (1H, d, J=2.4Hz), 7.56 (1H, d, J=8.3Hz), 8.00 (1H, t, J=9.3Hz), 8.2-8.4 (1H, m), 12.84 (1H, brs)

MS (m/z): 487 (M⁺)

[0220]

Example 48: N-{4-Chloro-3-[4-(6-isopropoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-isopropoxypyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.31 (6H, d, J=5.9Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=5.8Hz), 5.32 (1H, quin, J=6.4Hz), 6.83 (1H, d, J=8.3Hz), 6.94 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.55 (1H, d, J=1.9Hz), 7.57 (1H, d, J=8.3Hz), 8.30 (1H, dd, J=2.5Hz, 8.8Hz), 8.34 (1H, d, J=6.3Hz), 8.86 (1H, d, J=2.5Hz), 12.84 (1H, brs)

MS (m/z): 440 (M⁺)

[0221]

Example 49: N-(4-Chloro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2,2,2-trifluoroethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.3Hz), 5.07 (2H, q, J=9.1Hz), 7.02 (1H, brs), 7.09 (1H, d, J=8.8Hz), 7.42 (1H, dd, J=2.5Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (1H, m), 8.44 (1H, dd, J=2.4Hz, 8.3Hz), 8.91 (1H, d, J=2.5Hz), 12.91 (1H, brs)

MS (m/z): 480 (M⁺)

[0222]

Example 50: N-{3-[4-(1-tert-Butyl-1H-pyrazol-4-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-chlorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 1-tert-butyl-1H-pyrazole-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.54 (9H, s), 2.43 (1H, sept, J=6.8Hz), 4.30 (2H, d, J=5.8Hz), 6.66 (1H, brs), 7.39 (1H, dd, J=2.0Hz, 8.3Hz), 7.47 (1H, d, J=2.0Hz), 7.54 (1H, d, J=8.3Hz), 8.01 (1H, s), 8.3-8.4 (2H, m), 12.44 (1H, brs)

MS (m/z): 427 (M⁺)

[0223]

Example 51: N-{4-Chloro-3-[4-(1-isobutyl-1H-pyrazol-4-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 1-isobutyl-1H-pyrazole-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.84 (6H, d, J=6.8Hz), 1.03 (6H, d, J=6.8Hz), 2.12 (1H, sept, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.93 (2H, d, J=7.3Hz), 4.30 (2H, d, J=5.9Hz), 6.62 (1H, brs), 7.39 (1H, dd, J=1.9Hz, 8.3Hz), 7.48 (1H, d, J=2.0Hz), 7.54 (1H, d, J=8.3Hz), 8.02 (1H, s), 8.28 (1H, s), 8.3-8.4 (1H, m), 12.61 (1H, brs)

MS (m/z): 427 (M⁺)

[0224]

Example 52: N-{4-Chloro-3-[4-(2-methoxythiazol-5-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-methoxythiazole-5-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.06 (3H, s), 4.30 (2H, d, J=5.9Hz), 6.83 (1H, brs), 7.41 (1H, dd, J=2.0Hz, 8.3Hz), 7.47 (1H, d, J=1.9Hz), 7.57 (1H, d, J=8.3Hz), 8.05 (1H, s), 8.3-8.4 (1H, m), 12.85 (1H, brs)

MS (m/z): 418 (M⁺)

[0225]

Example 53: N-(4-Chloro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.38 (6H, s), 4.37 (2H, d, J=5.9Hz), 7.25 (1H, s), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.57 (1H, d, J=1.9Hz), 7.60 (1H, d, J=8.3Hz), 8.03 (1H, d, J=8.3Hz), 8.5-8.7 (1H, m), 8.69 (1H, dd, J=1.9Hz, 8.3Hz), 9.41 (1H, d, J=1.5Hz), 13.11 (1H, brs)

MS (m/z): 518 (M⁺)

[0226]

Example 54: N-{4-Chloro-3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}-3,3,3-trifluoro-2,2-dimethylpropionamide

By performing operations similar to those of Example 4 using 6-methoxypyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.38 (6H, s), 3.92 (3H, s), 4.37 (2H, d, J=5.9Hz), 6.91 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.40 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.6 (1H, m), 7.58 (1H, d, J=8.3Hz), 8.34 (1H, dd, J=2.0Hz, 8.8Hz), 8.5-8.7 (1H, m), 8.89 (1H, d, J=1.9Hz), 12.87 (1H, brs)

MS (m/z): 480 (M⁺)

[0227]

Example 55: N-(4-Chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.38 (6H, s), 4.37 (2H, d, J=6.4Hz), 7.08 (1H, brs), 7.20 (1H, d, J=8.8Hz), 7.40 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.7 (2H, m), 7.77 (1H, t, J=72.5Hz), 8.54 (1H, dd, J=2.5Hz, 8.8Hz), 8.5-8.7 (1H, m), 8.95 (1H, d, J=2.0Hz), 12.95 (1H, brs)

MS (m/z): 516 (M⁺)

[0228]

Example 56: N-(4-Fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-

dihydropyrimidin-2-yl}benzyl)-2,2-dimethylpropionamide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-fluorobenzyl)-2,2-dimethylpropionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.14 (9H, s), 4.32 (2H, d, J=6.3Hz), 7.23 (1H, s), 7.3-7.5 (2H, m), 7.6-7.8 (1H, m), 8.04 (1H, d, J=8.3Hz), 8.1-8.2 (1H, m), 8.71 (1H, dd, J=1.4Hz, 7.8Hz), 9.43 (1H, d, J=1.5Hz), 12.95 (1H, brs)

MS (m/z): 448 (M⁺)

[0229]

Example 57: N-(3-{4-[6-(Difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)-2,2-dimethylpropionamide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-fluorobenzyl)-2,2-dimethylpropionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.14 (9H, s), 4.32 (2H, d, J=5.9Hz), 7.08 (1H, brs), 7.22 (1H, d, J=9.3Hz), 7.3-7.5 (2H, m), 7.72 (1H, d, J=5.3Hz), 7.79 (1H, t, J=72.6Hz), 8.1-8.3 (1H, m), 8.57 (1H, dd, J=2.4Hz, 8.8Hz), 8.99 (1H, d, J=2.5Hz), 12.85 (1H, brs)

MS (m/z): 446 (M⁺)

[0230]

Example 58: N-(4-Difluoromethoxy-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(difluoromethoxy)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.3Hz), 7.20 (1H, t, J=73.5Hz), 7.21 (1H, brs), 7.30 (1H, d, J=8.3Hz), 7.49 (1H, dd, J=2.5Hz, 8.8Hz), 7.66 (1H, d, J=1.9Hz), 8.03 (1H, d, J=8.3Hz), 8.2-8.4 (1H, m), 8.70 (1H, dd, J=1.9Hz, 8.3Hz), 9.40 (1H, d, J=1.4Hz), 12.92 (1H, brs)

MS (m/z): 482 (M⁺)

[0231]

Example 59: N-(4-Chloro-3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-ethoxyethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.12 (3H, t, J=7.1Hz), 2.43 (1H, sept, J=6.8Hz), 3.49 (2H, q, J=7.0Hz), 3.6-3.8 (2H, m), 4.31 (2H, d, J=5.8Hz), 4.4-4.5 (2H, m),

6.93 (1H, d, J=8.7Hz), 6.96 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.55 (1H, d, J=1.9Hz), 7.57 (1H, d, J=8.3Hz), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.86 (1H, brs)
MS (m/z): 470 (M⁺)

[0232]

Example 60: N-(4-Chloro-3-{4-[6-(3-methoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-methoxypropoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.97 (2H, quin, J=6.5Hz), 2.43 (1H, sept, J=6.8Hz), 3.24 (3H, s), 3.47 (2H, q, J=6.4Hz), 4.31 (2H, d, J=5.8Hz), 4.37 (2H, t, J=6.6Hz), 6.89 (1H, d, J=8.7Hz), 6.95 (1H, brs), 7.41 (1H, dd, J=2.4Hz, 8.3Hz), 7.55 (1H, d, J=2.4Hz), 7.57 (1H, d, J=8.3Hz), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.81 (1H, brs)

MS (m/z): 470 (M⁺)

[0233]

Example 61: N-(4-Chloro-3-{4-[6-(2-methoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-methoxyethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.30 (3H, s), 3.6-3.8 (2H, m), 4.31 (2H, d, J=6.3Hz), 4.4-4.5 (2H, m), 6.92 (1H, d, J=8.3Hz), 6.96 (1H, brs), 7.41 (1H, dd, J=2.4Hz, 8.3Hz), 7.55 (1H, d, J=1.9Hz), 7.57 (1H, d, J=8.3Hz), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.85 (1H, brs)

MS (m/z): 456 (M⁺)

[0234]

Example 62: N-{4-Chloro-3-[4-(3-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 3-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=7.2Hz), 2.43 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.03 (1H, s), 7.3-7.4 (1H, m), 7.4-7.5 (1H, m), 7.5-7.6 (3H, m), 7.88 (1H, d, J=10.0Hz), 7.94 (1H, d, J=8.0Hz), 8.3-8.4 (1H, m), 12.96 (1H, s)

MS (m/z): 399 (M⁺)

[0235]

Example 63: N-(4-Chloro-3-{6-oxo-4-[4-(pyridin-2-ylethynyl)phenyl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-(pyridin-2-ylethynyl)benzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.03 (1H, s), 7.4-7.5 (2H, m), 7.5-7.6 (2H, m), 7.6-7.8 (3H, m), 7.87 (1H, ddd, J=1.6Hz, 7.6Hz, 7.6Hz), 8.17 (2H, d, J=8.0Hz), 8.35 (1H, t, J=6.0Hz), 8.6-8.7 (1H, m), 12.96 (1H, s)

MS (m/z): 482 (M⁺)

[0236]

Example 64: N-[4-Chloro-3-(5-fluoro-6-oxo-4-phenyl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 23 using ethyl 2-fluoro-3-oxo-3-phenylpropanoate, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.30 (2H, d, J=6.0Hz), 7.42 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (5H, m), 7.9-8.0 (2H, m), 8.34 (1H, t, J=6.0Hz), 13.47 (1H, s)

MS (m/z): 399 (M⁺)

[0237]

Example 65: N-{4-Chloro-3-[5-fluoro-4-(4-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 23 using ethyl 2-fluoro-3-(4-methoxyphenyl)-3-oxopropanoate, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (1H, m), 3.83 (3H, s), 4.30 (2H, d, J=5.6Hz), 7.0-7.1 (2H, m), 7.41 (1H, dd, J=2.4Hz, 8.4Hz), 7.5-7.6 (2H, m), 7.95 (2H, d, J=8.8Hz), 8.33 (1H, t, J=6.0Hz), 13.35 (1H, s)

MS (m/z): 429 (M⁺)

[0238]

Example 66: N-{4-Chloro-3-[5-fluoro-6-oxo-4-(thiophen-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 23 using ethyl 2-fluoro-3-oxo-3-(thiophen-2-yl)propanoate, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.4Hz), 7.28 (1H, dd, J=4.0Hz, 4.8Hz), 7.43 (1H, dd, J=2.0Hz, 8.4Hz), 7.51 (1H, d, J=2.0Hz), 7.57 (1H, d, J=8.4Hz), 7.83 (1H, d, J=4.0Hz), 7.91 (1H, dd, J=0.8Hz, 4.8Hz), 8.34 (1H, t, J=6.0Hz), 13.38 (1H, s)

MS (m/z): 405 (M⁺)

[0239]

Example 67: N-{4-Chloro-3-[5-fluoro-6-oxo-4-(thiophen-3-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 23 using ethyl 2-fluoro-3-oxo-3-(thiophen-3-yl)propanoate, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.30 (2H, d, J=5.6Hz), 7.41 (1H, dd, J=2.4Hz, 8.4Hz), 7.53 (1H, d, J=2.0Hz), 7.57 (1H, d, J=8.4Hz), 7.65 (1H, d, J=4.8Hz), 7.72 (1H, dd, J=2.8Hz, 4.8Hz), 8.25 (1H, d, J=2.0Hz), 8.34 (1H, t, J=6.0Hz), 13.36 (1H, s)

MS (m/z): 405 (M⁺)

[0240]

Example 68: N-{3-[4-(Benzo[b]thiophen-7-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-chloro-benzyl}isobutyramide

By performing operations similar to those of Example 4 using benzo[b]thiophene-7-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 4.35 (2H, d, J=6.4Hz), 7.11 (1H, s), 7.45 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 7.60 (1H, d, J=8.4Hz), 7.65 (1H, d, J=2.0Hz), 7.74 (1H, d, J=5.6Hz), 8.0-8.1 (2H, m), 8.36 (1H, t, J=6.0Hz), 13.03 (1H, s)

MS (m/z): 437 (M⁺)

[0241]

Example 69: N-{4-Chloro-3-[4-(4-cyanophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-cyanobenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.0Hz), 7.11 (1H, s), 7.4-7.5 (1H, m), 7.5-7.6 (2H, m), 7.95 (2H, d, J=8.0Hz), 8.30 (2H, d, J=8.4Hz), 8.34 (1H, t, J=6.0Hz), 13.05 (1H, s)

MS (m/z): 406 (M⁺)

[0242]

Example 70: N-(4-Chloro-3-{4-[4-(methylsulfanyl)phenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-(methylsulfanyl)benzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 2.52 (3H, s), 4.31 (2H, d, J=6.0Hz), 6.91 (1H, s), 7.34 (2H, d, J=8.8Hz), 7.42 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.02 (2H, d, J=8.4Hz), 8.34 (1H, t, J=6.0Hz), 12.84 (1H, s)

MS (m/z): 427 (M⁺)

[0243]

Example 71: N-{4-Chloro-3-[6-oxo-4-(p-toluy)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-methylbenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.36 (3H, s), 2.3-2.6 (1H, m), 4.31 (2H, d, J=6.0Hz), 6.88 (1H, s), 7.29 (2H, d, J=8.0Hz), 7.41 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 7.97 (2H, d, J=8.4Hz), 8.34 (1H, t, J=6.0Hz), 12.83 (1H, s)

MS (m/z): 395 (M⁺)

[0244]

Example 72: N-(4-Chloro-3-{4-[4-(diethylsulfamoyl)phenyl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using N,N-diethyl-4-formylbenzenesulfonamide, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.0-1.1 (12H, m), 2.3-2.6 (1H, m), 3.19 (4H, q, J=7.2Hz), 4.31 (2H, d, J=6.4Hz), 7.07 (1H, s), 7.42 (1H, dd, J=2.4Hz, 8.4Hz), 7.5-7.6 (2H, m), 7.88 (2H, d, J=8.0Hz), 8.27 (2H, d, J=8.8Hz), 8.34 (1H, t, J=6.0Hz), 13.02 (1H, s)

MS (m/z): 516 (M⁺)

[0245]

Example 73: N-(4-Chloro-3-{6-oxo-4-[4-(piperidine-1-sulfonyl)phenyl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-(piperidin-1-ylsulfonyl)benzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=7.2Hz), 1.3-1.4 (2H, m), 1.5-1.6 (4H, m), 2.3-2.6 (1H, m), 2.9-3.0 (4H, m), 4.31 (2H, d, J=5.6Hz), 7.09 (1H, s), 7.43 (1H, dd, J=2.8Hz, 8.4Hz), 7.5-7.6 (2H, m), 7.81 (2H, d, J=8.8Hz), 8.3-8.4 (3H, m), 13.05 (1H, s)

MS (m/z): 529 (M⁺⁺¹)

[0246]

Example 74: N-{4-Chloro-3-[4-(4-cyclopropylphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-cyclopropylbenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-0.8 (2H, m), 1.0-1.1 (8H, m), 1.9-2.0 (1H, m), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=5.6Hz), 6.87 (1H, s), 7.17 (2H, d, J=8.4Hz), 7.41 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 7.95 (2H, d, J=8.4Hz), 8.33 (1H, t, J=6.0Hz), 12.80

(1H, s)

MS (m/z): 421 (M⁺)

[0247]

Example 75: N-(3-{6-Oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained. ¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.0Hz), 7.26 (1H, s), 7.6-7.7 (2H, m), 7.90 (1H, d, J=8.4Hz), 8.03 (1H, d, J=8.0Hz), 8.40 (1H, t, J=6.0Hz), 8.66 (1H, d, J=8.0Hz), 9.39 (1H, s), 13.23 (1H, s)

MS (m/z): 484 (M⁺)

[0248]

Example 76: N-{3-[4-(6-Methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-methoxypyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained. ¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 3.91 (3H, s), 4.40 (2H, d, J=6.0Hz), 6.91 (1H, d, J=8.8Hz), 6.98 (1H, s), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.31 (1H, dd, J=2.0Hz, 8.4Hz), 8.39 (1H, t, J=6.0Hz), 8.87 (1H, s), 12.97 (1H, s)

MS (m/z): 446 (M⁺)

[0249]

Example 77: N-(3-{4-[6-(Difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained. ¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.0Hz), 7.10 (1H, s), 7.20 (1H, d, J=8.8Hz), 7.61 (2H, dd, J=4.4Hz, 12.8Hz), 7.77 (1H, t, J=72.4Hz), 7.89 (1H, d, J=8.4Hz), 8.39 (1H, t, J=6.0Hz), 8.51 (1H, d, J=8.4Hz), 8.93 (1H, s), 13.09 (1H, s)

MS (m/z): 482 (M⁺)

[0250]

Example 78: N-{3-[4-(6-Ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-ethoxypyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 1.33 (3H, t, J=7.6Hz), 2.4-2.6 (1H, m), 4.3-4.4 (4H, m), 6.88 (1H, d, J=8.8Hz), 6.97 (1H, s), 7.6-7.7 (2H, m), 7.88 (1H, d, J=7.6Hz), 8.30 (1H, dd, J=2.4Hz, 8.8Hz), 8.39 (1H, t, J=6.0Hz), 8.85 (1H, d, J=2.4Hz), 12.93 (1H, s)

MS (m/z): 460 (M⁺)

[0251]

Example 79: N-(4-Methyl-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-methylbenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (4H, m), 4.29 (2H, d, J=6.0Hz), 7.19 (1H, s), 7.32 (2H, s), 7.44 (1H, s), 8.03 (1H, d, J=8.4Hz), 8.24 (1H, t, J=6.0Hz), 8.70 (1H, dd, J=1.2Hz, 8.4Hz), 9.41 (1H, s), 12.90 (1H, s)

MS (m/z): 430 (M⁺)

[0252]

Example 80: N-(3-{4-[6-(Difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-methylbenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-methylbenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (4H, m), 4.29 (2H, d, J=5.6Hz), 7.02 (1H, s), 7.20 (1H, d, J=8.8Hz), 7.31 (2H, s), 7.43 (1H, s), 7.77 (1H, t, J=72.8Hz), 8.24 (1H, t, J=5.6Hz), 8.55 (1H, dd, J=2.4Hz, 8.8Hz), 8.95 (1H, d, J=2.0Hz), 12.77 (1H, s)

MS (m/z): 428 (M⁺)

[0253]

Example 81: N-{3-[4-(6-Ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-methylbenzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-ethoxypyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-methylbenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.34 (3H, t, J=6.8Hz), 2.3-2.6 (4H, m),

4.28 (2H, d, J=6.0Hz), 4.37 (2H, q, J=6.8Hz), 6.87 (1H, s), 6.89 (1H, s), 7.30 (2H, s), 7.42 (1H, s), 8.24 (1H, t, J=6.0Hz), 8.33 (1H, dd, J=2.4Hz, 8.8Hz), 8.87 (1H, d, J=2.4Hz), 12.64 (1H, s)

MS (m/z): 406 (M⁺)

[0254]

Example 82: N-[4-(Difluoromethyl)-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(difluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.39 (2H, d, J=6.0Hz), 7.26 (1H, s), 7.4-7.7 (2H, m), 7.80 (1H, d, J=8.0Hz), 8.05 (1H, d, J=8.4Hz), 8.34 (1H, t, J=5.6Hz), 8.72 (1H, d, J=7.2Hz), 9.44 (1H, s), 13.17 (1H, s)

MS (m/z): 466 (M⁺)

[0255]

Example 83: N-(3-{4-[6-(Difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(difluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(difluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.38 (2H, d, J=6.0Hz), 7.09 (1H, s), 7.22 (2H, d, J=8.0Hz), 7.4-8.0 (4H, m), 8.33 (1H, t, J=6.0Hz), 8.58 (1H, d, J=8.4Hz), 8.99 (1H, s), 13.04 (1H, s)

MS (m/z): 464 (M⁺)

[0256]

Example 84: N-(4-Chloro-3-{6-oxo-4-[2-(trifluoromethyl)pyridin-4-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(trifluoromethyl)pyridine-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.36 (1H, s), 7.44 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.3-8.4 (2H, m), 8.45 (1H, s), 8.91 (1H, d, J=5.6Hz), 13.19 (1H, s)

MS (m/z): 450 (M⁺)

[0257]

Example 85: N-(4-Chloro-3-{4-[2-(difluoromethoxy)pyridin-4-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(difluoromethoxy)pyridine-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=7.6Hz), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.22 (1H, s), 7.43 (1H, dd, J=1.2Hz, 8.4Hz), 7.58 (2H, d, J=8.8Hz), 7.68 (1H, s), 7.75 (1H, t, J=72.8Hz), 7.90 (1H, d, J=5.6Hz), 8.34 (1H, t, J=6.0Hz), 8.39 (1H, d, J=5.2Hz), 13.14 (1H, s)

MS (m/z): 448 (M⁺)

[0258]

Example 86: N-{4-Chloro-3-[4-(3-fluoro-4-methoxyphenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 3-fluoro-4-methoxybenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=7.2Hz), 2.43 (1H, sept, J=6.8Hz), 3.91 (3H, s), 4.31 (2H, d, J=6.0Hz), 6.93 (1H, s), 7.26 (1H, t, J=8.8Hz), 7.42 (1H, dd, J=2.0Hz, 8.0Hz), 7.54 (1H, d, J=1.6Hz), 7.57 (1H, d, J=8.0Hz), 7.9-8.0 (2H, m), 8.34 (1H, t, J=6.0Hz), 12.84 (1H, s)

MS (m/z): 429 (M⁺)

[0259]

Example 87: 4-{2-[2-Chloro-5-(isobutyrylaminomethyl)phenyl]-6-oxo-1,6-dihydropyrimidin-4-yl}-N,N-dimethylbenzamide

By performing operations similar to those of Example 4 using 4-formyl-N,N-dimethylbenzamide, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 2.92 (3H, s), 3.00 (3H, s), 4.32 (2H, d, J=5.6Hz), 6.99 (1H, s), 7.42 (1H, dd, J=1.6Hz, 8.0Hz), 7.49 (2H, d, J=8.4Hz), 7.55 (1H, d, J=1.6Hz), 7.57 (1H, d, J=8.4Hz), 8.13 (2H, d, J=8.4Hz), 8.34 (1H, t, J=5.6Hz), 12.95 (1H, s)

MS (m/z): 452 (M⁺)

[0260]

Example 88: N-{4-Chloro-3-[4-(4-fluorophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-fluorobenzaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.0Hz), 6.95 (1H, s), 7.2-7.4 (2H, m), 7.42 (1H, dd, J=2.0Hz, 8.0Hz), 7.54 (1H, d, J=1.6Hz), 7.57 (1H, d, J=8.4Hz), 8.1-8.2 (2H, m), 8.34 (1H, t, J=6.0Hz), 12.91 (1H, s)

MS (m/z): 399 (M⁺)

[0261]

Example 89: N-{4-Chloro-3-[4-(6-methoxypyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-methoxypyridine-2-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=7.2Hz), 2.44 (1H, sept, J=6.8Hz), 3.98 (3H, s), 4.32 (2H, d, J=6.4Hz), 6.9-7.0 (1H, m), 7.24 (1H, s), 7.42 (1H, dd, J=2.0Hz, 8.4Hz), 7.55 (1H, d, J=2.0Hz), 7.58 (1H, d, J=8.0Hz), 7.8-7.9 (2H, m), 8.34 (1H, t, J=6.4Hz), 12.96 (1H, s)

MS (m/z): 412 (M⁺)

[0262]

Example 90: N-(4-Fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.6Hz), 7.23 (1H, s), 7.3-7.4 (1H, m), 7.4-7.5 (1H, m), 7.74 (1H, d, J=5.6Hz), 8.04 (1H, d, J=8.0Hz), 8.34 (1H, t, J=6.0Hz), 8.72 (1H, dd, J=1.2Hz, 8.4Hz), 9.44 (1H, d, J=1.6Hz), 12.97 (1H, s)

MS (m/z): 434 (M⁺)

[0263]

Example 91: N-{4-Chloro-3-[4-(6-cyclopropylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-cyclopropylpyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.9-1.1 (4H, m), 1.03 (6H, d, J=6.8Hz), 2.1-2.2 (1H, m), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.0Hz), 6.99 (1H, s), 7.39 (1H, d, J=8.4Hz), 7.43 (1H, d, J=2.0Hz), 7.55 (1H, d, J=1.2Hz), 7.57 (1H, d, J=8.4Hz), 8.25 (1H, dd, J=2.0Hz, 8.4Hz), 8.33 (1H, t, J=6.0Hz), 9.06 (1H, d, J=1.6Hz), 12.91 (1H, s)

MS (m/z): 422 (M⁺)

[0264]

Example 92: N-{3-[4-(6-Ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-ethoxypyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-fluorobenzyl)isobutyramide

hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=7.6Hz), 1.34 (3H, t, J=7.2Hz), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 4.38 (2H, q, J=7.2Hz), 6.89 (1H, d, J=8.8Hz), 6.94 (1H, s), 7.3-7.4 (1H, m), 7.4-7.5 (1H, m), 7.73 (1H, d, J=5.6Hz), 8.3-8.4 (2H, m), 8.92 (1H, d, J=2.0Hz), 12.72 (1H, s)

MS (m/z): 410 (M⁺)

[0265]

Example 93: N-[4-Chloro-3-(2'-methoxy-6-oxo-1,6-dihydro[4,5']bipyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 2-methoxypyrimidine-5-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.99 (3H, s), 4.31 (2H, d, J=6.0Hz), 7.07 (1H, s), 7.42 (1H, dd, J=1.6Hz, 8.0Hz), 7.5-7.6 (2H, m), 8.33 (1H, t, J=6.0Hz), 9.22 (2H, s), 12.98 (1H, s)

MS (m/z): 413 (M⁺)

[0266]

Example 94: N-(3-{4-[6-(Difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.07 (1H, s), 7.22 (1H, d, J=9.6Hz), 7.3-7.4 (1H, m), 7.4-7.5 (1H, m), 7.74 (1H, d, J=5.6Hz), 7.78 (1H, t, J=72.0Hz), 8.34 (1H, t, J=6.0Hz), 8.57 (1H, dd, J=2.4Hz, 8.8Hz), 8.99 (1H, d, J=2.4Hz), 12.84 (1H, s)

MS (m/z): 433 (M⁺⁺¹)

[0267]

Example 95: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=6.0Hz), 7.30 (1H, s), 7.4-7.6 (2H, m), 8.02 (1H, d, J=8.4Hz), 8.36 (1H, t, J=5.6Hz), 8.67 (1H, d, J=8.4Hz), 9.39 (1H, s), 13.37 (1H, s)

MS (m/z): 468 (M⁺)

[0268]

Example 96: N-(4-Chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.13 (1H, s), 7.19 (1H, d, J=8.4Hz), 7.4-7.6 (2H, m), 7.77 (1H, t, J=72.4Hz), 8.36 (1H, t, J=6.0Hz), 8.51 (1H, dd, J=2.0Hz, 8.8Hz), 8.93 (1H, s), 13.22 (1H, s)

MS (m/z): 466 (M⁺)

[0269]

Example 97: N-(4-Fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)-1-(trifluoromethyl)cyclopropane-1-carboxamide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-fluorobenzyl)-1-(trifluoromethyl)cyclopropane-1-carboxamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.2-1.4 (4H, m), 4.35 (2H, d, J=5.6Hz), 7.23 (1H, s), 7.3-7.4 (1H, m), 7.4-7.5 (1H, m), 7.75 (1H, d, J=5.6Hz), 8.04 (1H, d, J=8.0Hz), 8.46 (1H, t, J=6.0Hz), 8.72 (1H, d, J=8.4Hz), 9.44 (1H, s), 12.99 (1H, s)

MS (m/z): 500 (M⁺)

[0270]

Example 98: Methyl 4-{2-[2-chloro-5-(isobutyrylaminomethyl)phenyl]-6-oxo-1,6-dihydropyrimidin-4-yl}benzoate

By performing operations similar to those of Example 4 using methyl 4-formylbenzoate, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.88 (3H, s), 4.31 (2H, d, J=6.4Hz), 7.06 (1H, s), 7.42 (1H, dd, J=2.0Hz, 8.4Hz), 7.55 (1H, d, J=2.0Hz), 7.58 (1H, d, J=8.4Hz), 8.04 (2H, d, J=8.4Hz), 8.20 (2H, d, J=8.4Hz), 8.34 (1H, t, J=6.0Hz), 13.00 (1H, s)

MS (m/z): 439 (M⁺)

[0271]

Example 99: N-(3-{4-[6-(Difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)-1-(trifluoromethyl)cyclopropane-1-carboxamide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-fluorobenzyl)-1-

(trifluoromethyl)cyclopropane-1-carboxamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.2-1.4 (4H, m), 4.35 (2H, d, J=6.0Hz), 7.08 (1H, s), 7.22 (1H, d, J=8.8Hz), 7.3-7.4 (1H, m), 7.4-7.5 (1H, m), 7.75 (1H, d, J=6.4Hz), 7.78 (1H, t, J=72.0Hz), 8.46 (1H, t, J=6.0Hz), 8.58 (1H, dd, J=2.4Hz, 8.8Hz), 9.00 (1H, d, J=2.4Hz), 12.84 (1H, s)

MS (m/z): 498 (M⁺)

[0272]

Example 100: N-(2,4-Difluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.2-7.4 (2H, m), 7.4-7.6 (1H, m), 8.03 (1H, d, J=8.4Hz), 8.33 (1H, t, J=6.0Hz), 8.67 (1H, d, J=8.4Hz), 9.39 (1H, s), 13.37 (1H, s)

MS (m/z): 452 (M⁺)

[0273]

Example 101: N-(4-Chloro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.3Hz), 1.03 (6H, d, J=6.8Hz), 1.4-1.6 (2H, m), 2.43 (1H, sept, J=6.8Hz), 3.40 (2H, t, J=6.6Hz), 3.6-3.8 (2H, m), 4.31 (2H, d, J=5.8Hz), 4.4-4.5 (2H, m), 6.92 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.7 (2H, m), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.84 (1H, brs)

MS (m/z): 484 (M⁺)

[0274]

Example 102: N-(3-{4-[6-(2-Butoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-chlorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-butoxyethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.3Hz), 1.03 (6H, d, J=6.8Hz), 1.2-1.4 (2H, m), 1.4-1.6 (2H, m), 2.43 (1H, sept, J=6.8Hz), 3.44 (2H, t, J=6.6Hz), 3.6-3.8 (2H, m), 4.31 (2H, d, J=5.8Hz), 4.4-4.5 (2H, m), 6.92 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.7 (2H, m), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.0Hz), 12.87 (1H, brs)

MS (m/z): 498 (M⁺)

[0275]

Example 103: N-{4-Chloro-3-[6-oxo-2'-(trifluoromethyl)-1,6-dihydro[4,5']-bipyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-(trifluoromethyl)pyrimidine-5-carboxaldehyde, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.34 (1H, s), 7.44 (1H, dd, J=2.0Hz, 8.0Hz), 7.5-7.6 (2H, m), 8.34 (1H, t, J=6.0Hz), 9.26 (2H, s), 13.21 (1H, s)

MS (m/z): 451 (M⁺)

[0276]

Reference Example 47: 6-(3-Ethoxypropoxy)pyridine-3-carbaldehyde

[Formula 60]



[0277]

By performing operations similar to those of Reference Example 20 using 3-ethoxypropan-1-ol, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 1.1-1.3 (3H, m), 2.0-2.2 (2H, m), 3.4-3.7 (4H, m), 4.5-4.7 (2H, m), 6.83 (1H, d, J=8.3Hz), 8.06 (1H, dd, J=2.4Hz, 8.8Hz), 8.62 (1H, d, J=1.4Hz), 9.95 (1H, s)

MS (m/z): 209 (M⁺)

[0278]

Reference Example 48: 6-(Cyclopropylmethoxy)pyridine-3-carbaldehyde

[Formula 61]




[0279]

By performing operations similar to those of Reference Example 20 using cyclopropylmethanol, the title compound was obtained.

¹H-NMR (CDCl₃, δ): 0.3-0.7 (4H, m), 1.2-1.4 (1H, m), 4.24 (2H, d, J=7.3Hz), 6.86 (1H, d, J=8.3Hz), 8.06 (1H, dd, J=2.5Hz, 8.8Hz), 8.60 (1H, d, J=2.5Hz), 9.95 (1H, s)

Reference Example 49: 6-[2-(2-Butoxyethoxy)ethoxy]pyridine-3-carbaldehyde

[Formula 62]



[0280]

By performing operations similar to those of Reference Example 20 using 2-(2-butoxyethoxy)ethanol, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 0.91 (3H, t, J=7.3Hz), 1.3-1.7 (4H, m), 3.4-4.0 (8H, m), 4.5-4.7 (2H, m), 6.88 (1H, d, J=8.8Hz), 8.06 (1H, dd, J=2.0Hz, 8.3Hz), 8.60 (1H, d, J=2.4Hz), 9.95 (1H, s)

MS (m/z): 267 (M⁺)

[0281]

Reference Example 50: 6-(Pyridin-2-ylmethoxy)pyridine-3-carbaldehyde

[Formula 63]



[0282]

By performing operations similar to those of Reference Example 20 using pyridin-2-ylmethanol, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 5.61 (2H, s), 6.98 (1H, d, J=8.3Hz), 7.2-7.3 (1H, m), 7.45 (1H, d, J=7.8Hz), 7.71 (1H, dt, J=2.0Hz, 7.8Hz), 8.11 (1H, dd, J=2.4Hz, 8.8Hz), 8.6-8.7 (2H, m), 9.97 (1H, s)

MS (m/z): 214 (M⁺)

[0283]

Reference Example 51: 6-(6-Methylpyridin-3-ylmethoxy)pyridine-3-carbaldehyde

[Formula 64]



[0284]

By performing operations similar to those of Reference Example 20 using (6-methylpyridin-3-yl)methanol, the title compound was obtained.

¹H-NMR (CDCl₃, δ): 2.57 (3H, s), 5.47 (2H, s), 6.88 (1H, d, J=8.8Hz), 7.18 (1H, d, J=8.3Hz), 7.69 (1H, dd, J=2.5Hz, 7.9Hz), 8.09 (1H, dd, J=2.4Hz, 8.3Hz), 8.5-8.7 (2H, m).


9.97 (1H, s)

MS (m/z): 228 (M⁺)

[0285]

Reference Example 52: 2-Oxo-1-pentyl-1,2-dihydropyridine-4-carbaldehyde

[Formula 65]



[0286]

To a solution of 2-oxo-1,2-dihydropyridine-4-carbaldehyde (2.46 g) in DMF (50 mL), 60% sodium hydride (960 mg) was added under ice cooling. The resulting mixture was stirred at room temperature for 30 minutes. 1-Bromopentane (3.0 mL) was added to the reaction mixture, and the resulting mixture was stirred at 50°C for 5 hours. The reaction mixture was poured into saturated aqueous ammonium chloride, and the resulting mixture was extracted with t-butyl methyl ether. The organic layer was washed with water, and dried over sodium sulfate. The organic layer was filtered, and then the solvent was evaporated. Silica gel column chromatography was performed to obtain the title compound (606 mg).


¹H-NMR (CDCl₃, δ): 0.91 (3H, t, J=7.3Hz), 1.2-1.5 (4H, m), 1.6-1.9 (2H, m), 3.95 (2H, t, J=7.8Hz), 6.56 (1H, dd, J=1.5Hz, 6.8Hz), 6.99 (1H, d, J=2.0Hz), 7.37 (1H, d, J=6.8Hz), 9.88 (1H, d, J=0.9Hz)

MS (m/z): 193 (M⁺)

[0287]

Reference Example 53: 1-Butyl-2-oxo-1,2-dihydropyridine-4-carbaldehyde

[Formula 66]



[0288]

By performing operations similar to those of Reference Example 52 using 1-bromobutane, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 0.97 (3H, t, J=7.3Hz), 1.3-1.5 (2H, m), 1.6-1.9 (2H, m), 3.96 (2H, t, J=7.3Hz), 6.56 (1H, dd, J=2.0Hz, 7.3Hz), 7.00 (1H, d, J=1.5Hz), 7.38 (1H, d, J=7.3Hz), 9.87 (1H, s)

MS (m/z): 179 (M⁺)

[0289]

Reference Example 54: N-(4-Chloro-3-cyanobenzyl)propionamide

[Formula 67]



[0290]

By performing operations similar to those of Reference Example 1 using propionyl chloride, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 1.19 (3H, t, J=7.3Hz), 2.29 (2H, q, J=7.3Hz), 4.44 (2H, d, J=5.8Hz), 5.95 (1H, brs), 7.4-7.6 (2H, m), 7.57 (1H, s)

MS (m/z): 222 (M⁺)

[0291]

Reference Example 55: N-(3-Carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride

[Formula 68]




[0292]

By performing operations similar to those of Reference Example 2 using N-(4-chloro-3-cyanobenzyl)propionamide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0293]

Reference Example 56: N-(4-Chloro-3-cyanobenzyl)butyramide

[Formula 69]



[0294]

By performing operations similar to those of Reference Example 1 using

butyryl chloride, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 0.96 (3H, t, J=7.3Hz), 1.6-1.8 (2H, m), 2.23 (2H, t, J=7.8Hz), 4.44 (2H, d, J=5.8Hz), 5.95 (1H, brs), 7.4-7.6 (2H, m), 7.57 (1H, s)

MS (m/z): 236 (M⁺)

[0295]

Reference Example 57: N-(3-Carbamimidoyl-4-chlorobenzyl)butyramide hydrochloride
[Formula 70]

H-Cl




[0296]

By performing operations similar to those of Reference Example 2 using N-(4-chloro-3-cyanobenzyl)butyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0297]

Reference Example 58: 6-{2-[2-(2-Ethoxyethoxy)ethoxy]ethoxy}pyridine-3-carbaldehyde
[Formula 71]



[0298]


By performing operations similar to those of Reference Example 20 using 2-[2-(2-ethoxyethoxy)ethoxy]ethanol, the title compound was obtained.

¹H-NMR (CDCl₃, δ): 1.21 (3H, t, J=7.3Hz), 3.52 (2H, q, J=6.8Hz), 3.5-4.0 (10H, m), 4.5-4.7 (2H, m), 6.88 (1H, d, J=8.8Hz), 8.06 (1H, dd, J=2.5Hz, 8.8Hz), 8.60 (1H, d, J=1.9Hz), 9.95 (1H, s)

MS (m/z): 283 (M⁺)

[0299]

Reference Example 59: 6-(6-Methylpyridin-2-ylmethoxy)pyridine-3-carbaldehyde
[Formula 72]



[0300]

By performing operations similar to those of Reference Example 20 using (6-methylpyridin-2-yl)methanol, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 2.59 (3H, s), 5.57 (2H, s), 6.97 (1H, d, J=8.8Hz), 7.10 (1H, d, J=7.8Hz), 7.23 (1H, d, J=7.3Hz), 7.59 (1H, t, J=7.9Hz), 8.10 (1H, dd, J=2.4Hz, 8.8Hz), 8.63 (1H, d, J=2.4Hz), 9.96 (1H, s)

MS (m/z): 228 (M⁺)

[0301]

Reference Example 60: 6-(3-Fluoropropoxy)pyridine-3-carbaldehyde

[Formula 73]



[0302]

By performing operations similar to those of Reference Example 20 using 3-fluoropropan-1-ol, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 1.1-1.3 (2H, m), 4.4-4.7 (3H, m), 4.70 (1H, t, J=5.4Hz), 6.84 (1H, d, J=8.8Hz), 8.08 (1H, dd, J=2.5Hz, 8.8Hz), 8.62 (1H, d, J=2.0Hz), 9.96 (1H, s)

MS (m/z): 183 (M⁺)

[0303]

Reference Example 61: 3-(Bromomethyl)-2-chloro-6-fluorobenzonitrile

[Formula 74]




[0304]

Under a nitrogen atmosphere, 2-chloro-6-fluoro-3-methylbenzonitrile (5.18 g) was dissolved in acetonitrile (150 mL), 1,3-dibromo-5,5-dimethylhydantoin (4.37 g), and 2,2'-azobis(isobutyronitrile) (501 mg) were added to the solution, and the resulting mixture was refluxed for 3 hours by heating. 1 N Hydrochloric acid was added to the reaction mixture, the resulting mixture was extracted with ethyl acetate, and then the obtained organic layer was washed with 1 N hydrochloric acid, and saturated brine. The solvent was removed under reduced pressure, and the obtained compound was used for the following reaction without purification.

[0305]

Reference Example 62: 3-(Aminomethyl)-2-chloro-6-fluorobenzonitrile

[Formula 75]



[0306]


3-(Bromomethyl)-2-chloro-6-fluorobenzonitrile was dissolved in N,N-dimethylformamide (75 mL), sodium azide (1.99 g) was added to the solution under ice cooling, and the resulting mixture was stirred for 30 minutes. Saturated brine was added to the reaction mixture, the resulting mixture was extracted with ethyl acetate, and then the organic layer was washed with saturated brine. The solvent was removed under reduced pressure, and the obtained compound was used for the following reaction without purification.

The obtained compound was dissolved in tetrahydrofuran (75 mL), and water (3 mL), triphenylphosphine (9.62 g) was added to the solution under ice cooling, and the resulting mixture was stirred overnight at room temperature. 1 N Hydrochloric acid was added to the reaction mixture, and the resulting mixture was washed with diethyl ether. 1 N Aqueous sodium hydroxide was added, the resulting mixture was extracted with dichloromethane, then the organic layer was dried over magnesium sulfate, and the obtained the organic layer was used as it was for the following reaction.

[0307]

Reference Example 63: N-(2-Chloro-3-cyano-4-fluorobenzyl)isobutyramide

[Formula 76]



[0308]

By performing operations similar to those of Reference Example 1 using isobutyryl chloride, the title compound was obtained.

¹H-NMR (CDCl₃, δ): 1.17 (6H, d, J=6.8Hz), 2.40 (1H, sept, J=6.8Hz), 4.50 (2H, d, J=6.4Hz), 5.98 (1H, s), 7.13 (1H, t, J=8.8Hz), 7.6-7.7 (1H, m)


MS (m/z): 254 (M⁺)

[0309]

Reference Example 64: N-(3-Carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide

hydrochloride

[Formula 77]




[0310]

By performing operations similar to those of Reference Example 2 using N-(2-chloro-3-cyano-4-fluorobenzyl)isobutyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0311]

Reference Example 65: 3-Chloro-2-fluoro-6-(trifluoromethyl)benzamidine hydrochloride

[Formula 78]




[0312]

By performing operations similar to those of Reference Example 2 using 3-chloro-2-fluoro-6-(trifluoromethyl)benzonitrile, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0313]

Reference Example 66: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-4(3H)-one

[Formula 79]



[0314]

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-

(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 7.38 (1H, s), 7.87 (1H, d, J=8.4Hz), 8.04 (1H, d, J=8.4Hz), 8.10 (1H, t, J=7.6Hz), 8.65 (1H, d, J=7.6Hz), 9.38 (1H, s), 13.44 (1H, brs)

MS (m/z): 437 (M⁺)

[0315]

Reference Example 67: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(difluoromethoxy)pyridin-3-yl]pyrimidin-4(3H)-one

[Formula 80]



[0316]

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 7.2-7.3 (2H, m), 7.77 (1H, t, J=72.8Hz), 7.86 (1H, d, J=8.8Hz), 8.09 (1H, t, J=8.0Hz), 8.50 (1H, dd, J=2.4Hz, 8.8Hz), 8.92 (1H, d, J=2.0Hz), 13.30 (1H, brs)

MS (m/z): 435 (M⁺)

[0317]

Reference Example 68: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(2-ethoxyethoxy)pyridin-3-yl]pyrimidin-4(3H)-one

[Formula 81]




[0318]

By performing operations similar to those of Example 4 using 6-(2-ethoxyethoxy)pyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.

$^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$, δ): 1.12 (3H, t, $J=6.8\text{Hz}$), 3.49 (2H, q, $J=6.8\text{Hz}$), 3.71 (2H, t, $J=5.2\text{Hz}$), 4.43 (2H, t, $J=5.2\text{Hz}$), 6.94 (1H, d, $J=8.8\text{Hz}$), 7.06 (1H, s), 7.85 (1H, d, $J=8.0\text{Hz}$), 8.08 (1H, t, $J=8.0\text{Hz}$), 8.30 (1H, d, $J=7.2\text{Hz}$), 8.84 (1H, s), 13.20 (1H, brs)
MS (m/z): 457 (M^+)

[0319]

Reference Example 69: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[5-(trifluoromethyl)pyridin-2-yl]pyrimidin-4(3H)-one
[Formula 82]




[0320]

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-2-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.
 $^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$, δ): 7.41 (1H, s), 7.88 (1H, d, $J=8.8\text{Hz}$), 8.11 (1H, t, $J=8.4\text{Hz}$), 8.34-8.40 (2H, m), 9.13 (1H, s), 13.45 (1H, brs)
MS (m/z): 437 (M^+)

[0321]

Reference Example 70: 6-(Tetrahydropyran-4-ylmethoxy)pyridine-3-carbaldehyde
[Formula 83]



[0322]

To a solution of (tetrahydropyran-4-yl)methanol (8.36 g) in DMF (150 mL), potassium tert-butoxide (5.84 g) was added, and the resulting mixture was stirred for 30 minutes under a nitrogen atmosphere. 6-Chloropyridine-3-carbaldehyde (4.24 g) was added to the reaction mixture, and the resulting mixture was stirred for 14 hours under a nitrogen atmosphere. The reaction mixture was poured into water, and the resulting mixture was extracted with t-butyl methyl ether. The solvent was evaporated, and then the residue was purified by silica gel column chromatography to

obtain the title compound (0.80 g).


¹H-NMR (CDCl₃, δ): 1.3-1.8 (4H, m), 2.0-2.2 (1H, m), 3.3-3.6 (2H, m), 3.9-4.1 (2H, m), 4.27 (2H, d, J=6.9Hz), 6.83 (1H, d, J=8.3Hz), 8.07 (1H, dd, J=2.4Hz, 8.8Hz), 8.61 (1H, d, J=2.5Hz), 9.96 (1H, s)

MS (m/z): 221 (M⁺)

[0323]

Reference Example 71: 5-(1-Propynyl)pyridine-3-carbaldehyde

[Formula 84]



[0324]

5-Bromopyridine-3-carbaldehyde (595 mg), dichlorobis(triphenylphosphine)palladium(II) (225 mg), copper(I) iodide (120 mg), N,N-diisopropylethylamine (2.2 mL), tetrahydrofuran (10 mL), a 1 M solution of tetrabutylammonium fluoride in tetrahydrofuran (3.4 mL), and 1-(trimethylsilyl)-1-propyne (1.43 mL) were mixed in order. Microwaves were irradiated on the mixture at 50°C for 30 minutes under a nitrogen atmosphere. The reaction mixture was filtered, then ethyl acetate was added to the filtrate, and the resulting mixture was washed with water. The organic layer was separated, and the solvent was evaporated. The residue was purified by silica gel column chromatography to obtain the title compound (322 mg).


¹H-NMR (CDCl₃, δ): 2.11 (3H, s), 8.12 (1H, s), 8.83 (1H, s), 8.94 (1H, s), 10.08 (1H, s)

MS (m/z): 145 (M⁺)

[0325]

Reference Example 72: 6-(3-Methyloxetan-3-ylmethoxy)pyridine-3-carbaldehyde

[Formula 85]



[0326]

By performing operations similar to those of Reference Example 70 using 3-methyl-3-oxetanemethanol, the title compound was obtained.

¹H-NMR (CDCl₃, δ): 1.44 (3H, s), 4.4-4.8 (6H, m), 6.87 (1H, d, J=8.8Hz), 8.10 (1H, dd,


J=2.5Hz, 8.8Hz), 8.63 (1H, d, J=2.0Hz), 9.97 (1H, s)

MS (m/z): 207 (M⁺)

[0327]

Reference Example 73: 6-Butoxypyridazine-3-carbaldehyde

[Formula 86]



[0328]

To a solution of (6-butoxypyridazin-3-yl)methanol (1.51 g) in chloroform (50 mL), 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3-(1H)-one (4.22 g) was slowly added with ice cooling under a nitrogen atmosphere. The resulting mixture was stirred at room temperature for 3 hours, and then saturated aqueous sodium hydrogencarbonate, and aqueous sodium thiosulfate were added to the mixture. The organic layer was separated, dried over magnesium sulfate, and filtered. The solvent was evaporated under reduced pressure, and the obtained residue was purified by silica gel column chromatography to obtain the title compound (790 mg).


¹H-NMR (CDCl₃, δ): 0.98 (3H, t, J=7.6Hz), 1.42 (2H, sext, J=7.6Hz), 1.85 (2H, quin, J=7.6Hz), 4.26 (2H, t, J=7.6Hz), 6.96 (1H, d, J=9.2Hz), 7.74 (1H, d, J=9.6Hz), 9.74 (1H, d, J=1.2Hz)

MS (m/z): 180 (M⁺)

[0329]

Reference Example 74: N-(4-Chloro-3-cyano-2-fluorobenzyl)butyramide

[Formula 87]



[0330]

By performing operations similar to those of Reference Example 1 using 3-aminomethyl-6-chloro-2-fluorobenzonitrile, and butyryl chloride, the title compound was obtained.

¹H-NMR (CDCl₃, δ): 0.94 (3H, t, J=7.6Hz), 1.67 (2H, sext, J=7.6Hz), 2.19 (2H, t, J=7.6Hz), 4.46 (2H, t, J=6.4Hz), 7.29 (1H, dd, J=1.2Hz, 8.4Hz), 7.61 (1H, t, J=8.4Hz)


MS (m/z): 254 (M⁺)

[0331]

Reference Example 75: N-(3-Carbamimidoyl-4-chloro-2-fluorobenzyl)butyramide hydrochloride

[Formula 88]

H-Cl




[0332]

By performing operations similar to those of Reference Example 2 using N-(4-chloro-3-cyano-2-fluorobenzyl)butyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0333]

Reference Example 76: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-(6-methoxypyridin-3-yl)pyrimidin-4(3H)-one

[Formula 89]



[0334]

By performing operations similar to those of Example 4 using 6-methoxypyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 3.92 (3H, s), 6.92 (1H, d, J=8.8Hz), 7.07 (1H, s), 7.85 (1H, d, J=8.4Hz), 8.08 (1H, t, J=8.4Hz), 8.30 (1H, dd, J=2.4Hz, 8.8Hz), 8.86 (1H, d, J=2.0Hz), 13.17 (1H, brs)

MS (m/z): 399 (M⁺)

[0335]

Reference Example 77: 6-(Cyclopropylethynyl)pyridine-3-carbaldehyde

[Formula 90]



[0336]

To a solution of 6-bromopyridine-3-carbaldehyde (930 mg) in tetrahydrofuran (25 mL), cyclopropylacetylene (0.64 mL), dichlorobis(triphenylphosphine)palladium(II) (70 mg), copper iodide (38 mg), and triethylamine (2.1 mL) were added under a nitrogen atmosphere, and the resulting mixture was stirred overnight at 40°C. After the reaction, the reaction mixture was filtered through a Celite layer, the solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (595 mg).


¹H-NMR (CDCl₃, δ): 0.9-1.0 (4H, m), 1.5-1.6 (1H, m), 7.48 (1H, d, J=7.6Hz), 8.08 (1H, dd, J=2.0Hz, 8.0Hz), 8.97 (1H, d, J=2.0Hz), 10.07 (1H, s)

MS (m/z): 171 (M⁺)

[0337]

Reference Example 78: 2-(Cyclopropylethynyl)thiazole-5-carbaldehyde

[Formula 91]



[0338]

To a solution of 2-bromothiazole-5-carbaldehyde (1.92 g) in tetrahydrofuran (40 mL), cyclopropylacetylene (1.27 mL), dichlorobis(triphenylphosphine)palladium(II) (140 mg), copper iodide (76 mg), and triethylamine (4.2 mL) were added under a nitrogen atmosphere, and the resulting mixture was stirred at 40°C for 3 hours. After the reaction, the reaction mixture was filtered through a Celite layer, the solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (1.42 g).


¹H-NMR (CDCl₃, δ): 0.9-1.1 (4H, m), 1.5-1.6 (1H, m), 8.32 (1H, s), 9.98 (1H, s)

MS (m/z): 177 (M⁺)

[0339]

Reference Example 79: 5-(Cyclopropylethynyl)pyridine-2-carbaldehyde

[Formula 92]



[0340]


To a solution of 5-bromopyridine-2-carbaldehyde (1.86 g) in tetrahydrofuran (40 mL), cyclopropylacetylene (1.27 mL), dichlorobis(triphenylphosphine)palladium(II) (140 mg), copper iodide (76 mg), and triethylamine (4.2 mL) were added under a nitrogen atmosphere, and the resulting mixture was stirred overnight at 40°C. After the reaction, the reaction mixture was filtered through a Celite layer, the solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (1.29 g).

¹H-NMR (CDCl₃, δ): 0.8-1.0 (4H, m), 1.4-1.6 (1H, m), 7.79 (1H, dd, J=2.0Hz, 8.0Hz), 7.87 (1H, d, J=8.0Hz), 8.72 (1H, d, J=2.0Hz), 10.05 (1H, s)

MS (m/z): 171 (M⁺)

[0341]

Reference Example 80: 6-(3-Ethylloxetan-3-ylmethoxy)pyridine-3-carbaldehyde
[Formula 93]



[0342]


Microwaves were irradiated on a mixture of 3-ethyl-3-oxetanemethanol (4.65 g), 6-chloropyridine-3-carbaldehyde (2.83 g), potassium carbonate (5.53 g), and dimethyl sulfoxide (13 mL) at 110°C for 2 hours, and at 130°C for further 2 hours. The reaction mixture was poured into water, and the resulting mixture was extracted with t-butyl methyl ether. The insoluble matter was removed by filtration. The solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (0.86 g).

¹H-NMR (CDCl₃, δ): 0.96 (3H, t, J=7.3Hz), 1.87 (2H, q, J=7.3Hz), 4.4-4.7 (6H, m), 6.89 (1H, d, J=8.3Hz), 8.09 (1H, dd, J=2.5Hz, 8.8Hz), 8.63 (1H, d, J=2.5Hz), 9.97 (1H, s)

MS (m/z): 221 (M⁺)

[0343]

Reference Example 81: 4-(1-Propynyl)thiazole-2-carbaldehyde
[Formula 94]



[0344]

By performing operations similar to those of Reference Example 71 using 4-bromothiazole-2-carbaldehyde, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 2.10 (3H, s), 7.69 (1H, d, J=1.0Hz), 9.96 (1H, d, J=1.5Hz)

MS (m/z): 151 (M⁺)

[0345]

Reference Example 82: 5-(Cyclopropylethynyl)thiazole-2-carbaldehyde

[Formula 95]



[0346]

By performing operations similar to those of Reference Example 77 using 5-bromothiazole-2-carbaldehyde, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 0.8-1.1 (4H, m), 1.4-1.6 (1H, m), 7.99 (1H, s), 9.90 (1H, s)

MS (m/z): 177 (M⁺)

[0347]

Reference Example 83: 4-(Cyclopropylethynyl)thiazole-2-carbaldehyde

[Formula 96]



[0348]

By performing operations similar to those of Reference Example 77 using 4-bromothiazole-2-carbaldehyde, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 0.8-1.0 (4H, m), 1.4-1.6 (1H, m), 7.67 (1H, d, J=1.5Hz), 9.95 (1H, d, J=1.0Hz)

MS (m/z): 177 (M⁺)

[0349]

Reference Example 84: 6-(Cyclopropylethynyl)pyridine-2-carbaldehyde

[Formula 97]



[0350]

By performing operations similar to those of Reference Example 77 using 6-bromopyridine-2-carbaldehyde, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 0.8-1.1 (4H, m), 1.4-1.7 (1H, m), 7.56 (1H, dd, J=1.4Hz, 7.8Hz), 7.7-7.9 (2H, m), 10.05 (1H, s)

MS (m/z): 171 (M⁺)

[0351]

Reference Example 85: 5-(Cyclopropylethynyl)pyridine-3-carbaldehyde

[Formula 98]



[0352]

By performing operations similar to those of Reference Example 77 using 5-bromopyridine-3-carbaldehyde, the title compound was obtained.


¹H-NMR (CDCl₃, δ): 0.8-1.0 (4H, m), 1.4-1.6 (1H, m), 8.10 (1H, t, J=1.7Hz), 8.79 (1H, d, J=2.4Hz), 8.92 (1H, d, J=1.9Hz), 10.08 (1H, s)

MS (m/z): 171 (M⁺)

[0353]

Reference Example 86: N-[3-Cyano-4-(trifluoromethyl)benzyl]butyramide

[Formula 99]



[0354]

By performing operations similar to those of Reference Example 1 using 5-aminomethyl-2-(trifluoromethyl)benzonitrile, and butyryl chloride, the title compound was obtained.

$^1\text{H-NMR}$ (CDCl_3 , δ): 0.98 (3H, t, $J=7.2\text{Hz}$), 1.72 (2H, quin, $J=7.6\text{Hz}$), 2.26 (2H, t, $J=7.2\text{Hz}$), 4.53 (2H, d, $J=5.6\text{Hz}$), 6.06 (1H, s), 7.4-7.8 (3H, m)


MS (m/z): 270 (M^+)

[0355]

Reference Example 87: N-[3-Carbamimidoyl-4-(trifluoromethyl)benzyl]butyramide hydrochloride

[Formula 100]

H-Cl




[0356]

By performing operations similar to those of Reference Example 2 using N-[3-cyano-4-(trifluoromethyl)benzyl]butyramide, the title compound was obtained. The obtained compound was used for the following reaction without purification.

[0357]

Reference Example 88: 6-(3-Morpholin-4-ylpropoxy)pyridine-3-carbaldehyde

[Formula 101]



[0358]

By performing operations similar to those of Reference Example 70 using 4-(3-hydroxypropyl)morpholine, the title compound was obtained.


$^1\text{H-NMR}$ (CDCl_3 , δ): 2.0 (2H, sext, $J=6.8\text{Hz}$), 2.4-2.6 (6H, m), 3.73 (4H, t, $J=4.8\text{Hz}$), 4.47 (2H, t, $J=6.4\text{Hz}$), 6.83 (1H, d, $J=8.8\text{Hz}$), 8.06 (1H, dd, $J=2.4\text{Hz}$, 8.8Hz), 8.62 (1H, d, $J=2.4\text{Hz}$), 9.95 (1H, s)

MS (m/z): 250 (M^+)

[0359]

Reference Example 89: 6-(1-Propynyl)pyridine-3-carbaldehyde

[Formula 102]



[0360]

6-Bromopyridine-3-carbaldehyde (595 mg), dichlorobis(triphenylphosphine)palladium(II) (225 mg), copper(I) iodide (122 mg), N,N-diisopropylethylamine (2.2 mL), tetrahydrofuran (10 mL), a 1 M solution of tetrabutylammonium fluoride in tetrahydrofuran (3.4 mL), and 1-(trimethylsilyl)-1-propyne (1.42 mL) were mixed in order. Microwaves were irradiated on the mixture at 50°C for 30 minutes under a nitrogen atmosphere. After the reaction, the reaction mixture was filtered through a Celite layer, the solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (370 mg).


¹H-NMR (CDCl₃, δ): 2.14 (3H, s), 7.51 (1H, d, J=8.4Hz), 8.10 (1H, dd, J=2.0Hz, 8.4Hz), 8.99 (1H, d, J=1.6Hz), 10.09 (1H, s)

MS (m/z): 145 (M⁺)

[0361]

Reference Example 90: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-(5-fluoropyridin-3-yl)pyrimidin-4(3H)-one

[Formula 103]



[0362]

By performing operations similar to those of Example 4 using 5-fluoropyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 7.31 (1H, s), 7.87 (1H, d, J=8.8Hz), 8.10 (1H, t, J=8.4Hz), 8.2-8.3 (1H, m), 8.72 (1H, d, J=2.4Hz), 9.11 (1H, s), 13.37 (1H, brs)

MS (m/z): 387 (M⁺)

[0363]

Reference Example 91: 6-[(1-Hydroxycyclohexyl)ethynyl]pyridine-3-carbaldehyde

[Formula 104]



[0364]

To a solution of 6-bromopyridine-3-carbaldehyde (1.86 g) in tetrahydrofuran (40 mL), 1-ethynyl-1-cyclohexanol (1.86 g), dichlorobis(triphenylphosphine)palladium(II) (140 mg), copper(I) iodide (76 mg), and triethylamine (4.2 mL) were added under a nitrogen atmosphere, and the resulting mixture was stirred at 50°C for 3 hours. After the reaction, the reaction mixture was filtered through a Celite layer, the solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (2.22 g).


¹H-NMR (CDCl₃, δ): 1.2-1.4 (1H, m), 1.5-1.8(7H, m), 2.0-2.1 (2H, m), 2.29 (1H, s), 7.58 (1H, d, J=8.0Hz), 8.13 (1H, dd, J=1.6Hz, 8.0Hz), 9.03 (1H, d, J=1.6Hz), 10.11 (1H, s)

MS (m/z): 229 (M⁺)

[0365]

Reference Example 92: 5-(1-Propynyl)pyridine-2-carbaldehyde

[Formula 105]



[0366]

5-Bromopyridine-2-carbaldehyde (930 mg), dichlorobis(triphenylphosphine)palladium(II) (351 mg), copper(I) iodide (190 mg), N,N-diisopropylethylamine (3.4 mL), tetrahydrofuran (15 mL), a 1 M solution of tetrabutylammonium fluoride in tetrahydrofuran (5.3 mL), and 1-(trimethylsilyl)-1-propyne (2.22 mL) were mixed in order, and the resulting mixture was stirred at 50°C for 1 hours under a nitrogen atmosphere. After the reaction, the reaction mixture was filtered through a Celite layer, the solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (454 mg).

¹H-NMR (CDCl₃, δ): 2.13 (3H, s), 7.8-7.9 (2H, m), 8.75 (1H, d, J=1.2Hz), 10.06 (1H, s)


MS (m/z): 145 (M⁺)

[0367]

Reference Example 93: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-(6-

ethoxypyridin-3-yl)pyrimidin-4(3H)-one

[Formula 106]



[0368]

By performing operations similar to those of Example 4 using 6-ethoxypyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.33 (3H, t, J=6.8Hz), 4.37 (2H, q, J=6.8Hz), 6.89 (1H, d, J=8.8Hz), 7.05 (1H, s), 7.85 (1H, d, J=8.8Hz), 8.08 (1H, t, J=7.6Hz), 8.29 (1H, dd, J=1.6Hz, 8.8Hz), 8.83 (1H, s), 13.19 (1H, brs)

MS (m/z): 413 (M⁺)

[0369]

Example 104: N-(4-Chloro-3-{4-[6-(3-ethoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-ethoxypropoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.10 (3H, t, J=6.8Hz), 1.96 (2H, quint, J=6.3Hz), 2.43 (1H, sept, J=6.8Hz), 3.42 (2H, q, J=6.8Hz), 3.50 (2H, t, J=5.9Hz), 4.31 (2H, d, J=6.3Hz), 4.37 (2H, t, J=6.4Hz), 6.89 (1H, d, J=8.8Hz), 6.95 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.80 (1H, brs)

MS (m/z): 484 (M⁺)

[0370]

Example 105: N-(4-Chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylmethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.03 (6H, d, J=6.8Hz), 1.1-1.4 (1H, m), 2.43 (1H, sept, J=6.8Hz), 4.16 (2H, d, J=6.8Hz), 4.31 (2H, d, J=5.8Hz), 6.8-7.0 (2H, m), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.85 (1H, d, J=1.9Hz), 12.86 (1H, brs)

MS (m/z): 452 (M⁺)

[0371]

Example 106: N-[3-(4-{6-[2-(2-Butoxyethoxy)ethoxy]pyridin-3-yl}-6-oxo-1,6-dihydropyrimidin-2-yl)-4-chlorobenzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-[2-(2-butoxyethoxy)ethoxy]pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.3Hz), 1.03 (6H, d, J=6.8Hz), 1.2-1.5 (4H, m), 2.43 (1H, sept, J=7.8Hz), 3.36 (2H, t, J=6.4Hz), 3.4-3.6 (4H, m), 3.7-3.8 (2H, m), 4.31 (2H, d, J=5.8Hz), 4.4-4.5 (2H, m), 6.92 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.77 (1H, brs)

MS (m/z): 542 (M⁺)

[0372]

Example 107: N-(4-Chloro-3-{6-oxo-4-[6-(pyridin-2-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(pyridin-2-ylmethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=5.8Hz), 5.49 (2H, d, J=6.8Hz), 6.97 (1H, brs), 7.05 (1H, d, J=8.8Hz), 7.3-7.4 (1H, m), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.46 (1H, d, J=7.8Hz), 7.5-7.6 (2H, m), 7.81 (1H, dt, J=1.5Hz, 7.3Hz), 8.33 (1H, t, J=5.8Hz), 8.39 (1H, dd, J=2.4Hz, 8.8Hz), 8.5-8.6 (1H, m), 8.87 (1H, d, J=2.4Hz), 12.87 (1H, brs)

MS (m/z): 489 (M⁺)

[0373]

Example 108: N-(3-{6-Oxo-4-[6-(pyridin-2-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(pyridin-2-ylmethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=5.8Hz), 5.49 (2H, s), 6.98 (1H, brs), 7.04 (1H, d, J=8.8Hz), 7.2-7.4 (1H, m), 7.46 (1H, d, J=7.8Hz), 7.5-7.7 (2H, m), 7.80 (1H, dt, J=1.5Hz, 7.9Hz), 7.88 (1H, d, J=7.9Hz), 8.3-8.6 (3H, m), 8.8-8.9 (1H, m), 12.95 (1H, brs)

MS (m/z): 523 (M⁺)

[0374]

Example 109: N-(4-Chloro-3-{4-[6-(6-methylpyridin-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(6-methylpyridin-3-ylmethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 2.46 (3H, s), 4.31 (2H, d, J=5.8Hz), 5.42 (2H, s), 6.9-7.1 (2H, m), 7.26 (1H, d, J=1.3Hz), 7.42 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.6 (2H, m), 7.77 (1H, dd, J=2.4Hz, 7.8Hz), 8.2-8.4 (2H, m), 8.55 (1H, d, J=2.0Hz), 8.90 (1H, d, J=2.5Hz)

MS (m/z): 503 (M⁺)

[0375]

Example 110: N-{4-Chloro-3-[6-oxo-4-(2-oxo-1-pentyl-1,2-dihydropyridin-4-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-oxo-1-pentyl-1,2-dihydropyridine-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.3Hz), 1.03 (6H, d, J=6.8Hz), 1.1-1.4 (4H, m), 1.5-1.7 (2H, m), 2.43 (1H, sept, J=6.8Hz), 3.89 (2H, t, J=6.9Hz), 4.31 (2H, d, J=5.8Hz), 6.80 (1H, dd, J=1.9Hz, 6.8Hz), 7.01 (1H, brs), 7.06 (1H, d, J=1.5Hz), 7.42 (1H, dd, J=2.0Hz, 8.3Hz), 7.53 (1H, d, J=2.0Hz), 7.57 (1H, d, J=8.3Hz), 7.77 (1H, d, J=7.3Hz), 8.34 (1H, t, J=6.1Hz), 13.04 (1H, brs)

MS (m/z): 468 (M⁺)

[0376]

Example 111: N-{3-[4-(1-Butyl-2-oxo-1,2-dihydropyridin-4-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-chlorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 1-butyl-2-oxo-1,2-dihydropyridine-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.90 (3H, t, J=7.3Hz), 1.03 (6H, d, J=6.8Hz), 1.2-1.4 (2H, m), 1.5-1.7 (2H, m), 2.43 (1H, sept, J=6.8Hz), 3.90 (2H, t, J=7.0Hz), 4.31 (2H, d, J=5.8Hz), 6.80 (1H, dd, J=1.4Hz, 6.8Hz), 7.01 (1H, brs), 7.06 (1H, d, J=1.9Hz), 7.42 (1H, dd, J=2.5Hz, 8.3Hz), 7.53 (1H, d, J=2.0Hz), 7.57 (1H, d, J=8.3Hz), 7.77 (1H, d, J=7.3Hz), 8.34 (1H, t, J=5.9Hz), 13.03 (1H, brs)

MS (m/z): 454 (M⁺)

[0377]

Example 112: N-(4-Chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.02 (3H, t, J=7.8Hz), 2.16 (2H, q, J=7.8Hz), 4.31 (2H, d,

J=6.4Hz), 7.08 (1H, brs), 7.20 (1H, d, J=8.8Hz), 7.43 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.7 (2H, m), 7.77 (1H, t, J=72.7Hz), 8.3-8.4 (1H, m), 8.54 (1H, dd, J=2.5Hz, 8.8Hz), 8.95 (1H, d, J=2.0Hz), 12.95 (1H, brs)

MS (m/z): 434 (M⁺)

[0378]

Example 113: N-(4-Chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide

By performing operations similar to those of Example 4 using 6-(cyclopropylmethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.02 (3H, t, J=7.3Hz), 1.2-1.4 (1H, m), 2.16 (2H, q, J=7.8Hz), 4.16 (2H, d, J=7.3Hz), 4.32 (2H, d, J=5.9Hz), 6.91 (1H, d, J=8.8Hz), 6.95 (1H, brs), 7.43 (1H, dd, J=2.0Hz, 8.8Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.85 (1H, d, J=2.0Hz), 12.80 (1H, brs)

MS (m/z): 438 (M⁺)

[0379]

Example 114: N-(4-Chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.3Hz), 1.4-1.6 (2H, m), 2.13 (2H, t, J=7.3Hz), 4.32 (2H, d, J=5.8Hz), 7.08 (1H, brs), 7.20 (1H, d, J=8.7Hz), 7.43 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.7 (2H, m), 7.77 (1H, t, J=72.2Hz), 8.3-8.5 (1H, m), 8.54 (1H, dd, J=2.5Hz, 8.8Hz), 8.95 (1H, d, J=2.5Hz), 12.98 (1H, brs)

MS (m/z): 448 (M⁺)

[0380]

Example 115: N-(4-Chloro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-0.9 (6H, m), 1.4-1.6 (4H, m), 2.13 (2H, t, J=7.3Hz), 3.40 (2H, t, J=6.8Hz), 3.6-3.8 (2H, m), 4.32 (2H, d, J=6.3Hz), 4.4-4.5 (2H, m), 6.92 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.3-7.7 (3H, m), 8.2-8.5 (2H, m), 8.87 (1H, d, J=2.0Hz), 12.87 (1H, brs)

MS (m/z): 484 (M⁺)

[0381]

Example 116: N-{4-Chloro-3-[4-(5-fluoropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-fluoropyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.4Hz), 7.19 (1H, brs), 7.43 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.7 (2H, m), 8.2-8.4 (2H, m), 8.70 (1H, d, J=2.9Hz), 9.14 (1H, s), 13.05 (1H, brs)

MS (m/z): 400 (M⁺)

[0382]

Example 117: N-(4-Chloro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)propionamide

By performing operations similar to those of Example 4 using 6-(2,2,2-trifluoroethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.02 (3H, t, J=7.3Hz), 2.16 (2H, q, J=7.8Hz), 4.32 (2H, d, J=5.9Hz), 5.07 (2H, q, J=8.8Hz), 7.02 (1H, brs), 7.10 (1H, d, J=8.8Hz), 7.43 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.3-8.4 (1H, m), 8.44 (1H, dd, J=2.4Hz, 8.8Hz), 8.91 (1H, d, J=2.5Hz), 12.83 (1H, brs)

MS (m/z): 466 (M⁺)

[0383]

Example 118: N-(4-Chloro-3-{4-[5-(difluoromethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)butyramide

By performing operations similar to those of Example 4 using 5-(difluoromethoxy)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.8Hz), 1.4-1.6 (2H, m), 2.13 (2H, t, J=7.3Hz), 4.33 (2H, d, J=5.9Hz), 7.18 (1H, brs), 7.41 (1H, t, J=7.8Hz), 7.44 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.7 (2H, m), 7.80 (1H, dd, J=2.4Hz, 8.3Hz), 8.30 (1H, d, J=8.8Hz), 8.38 (1H, t, J=5.9Hz), 8.63 (1H, d, J=2.9Hz), 13.04 (1H, brs)

MS (m/z): 448 (M⁺)

[0384]

Example 119: N-{3-[4-(5-Chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-

chloropyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.9Hz), 4.40 (2H, d, J=5.8Hz), 7.23 (1H, brs), 7.5-7.7 (2H, m), 7.90 (1H, d, J=8.3Hz), 8.3-8.5 (1H, m), 8.48 (1H, s), 8.74 (1H, d, J=1.9Hz), 9.19 (1H, s), 13.17 (1H, brs)
MS (m/z): 450 (M⁺)
[0385]

Example 120: N-{3-[4-(5-Fluoropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-fluoropyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 4.41 (2H, d, J=5.8Hz), 7.21 (1H, brs), 7.5-7.7 (2H, m), 7.90 (1H, d, J=8.3Hz), 8.28 (1H, d, J=9.8Hz), 8.3-8.5 (1H, m), 8.70 (1H, d, J=2.9Hz), 9.13 (1H, s), 13.16 (1H, brs)
MS (m/z): 434 (M⁺)
[0386]

Example 121: N-(4-Chloro-3-{4-[5-(difluoromethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide

By performing operations similar to those of Example 4 using 5-(difluoromethoxy)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.03 (3H, t, J=7.8Hz), 2.16 (2H, q, J=7.3Hz), 4.32 (2H, d, J=5.8Hz), 7.19 (1H, brs), 7.42 (1H, t, J=73.3Hz), 7.44 (1H, dd, J=2.5Hz, 8.8Hz), 7.5-7.7 (2H, m), 7.80 (1H, dd, J=2.4Hz, 8.3Hz), 8.31 (1H, d, J=8.8Hz), 8.35 (1H, t, J=5.8Hz), 8.63 (1H, d, J=2.4Hz), 12.96 (1H, brs)
MS (m/z): 434 (M⁺)
[0387]

Example 122: N-{4-Chloro-3-[4-(5-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-methoxypyridine-3-carbaldehyde, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.90 (3H, s), 4.32 (2H, d, J=5.9Hz), 7.14 (1H, brs), 7.42 (1H, dd, J=2.4Hz, 8.3Hz), 7.5-7.7 (2H, m), 7.94 (1H, d, J=2.2Hz), 8.34 (1H, t, J=5.9Hz), 8.40 (1H, d, J=3.0Hz), 8.85 (1H, d, J=1.9Hz), 12.98 (1H, brs)

MS (m/z): 412 (M⁺)

[0388]

Example 123: N-{3-[4-(5-Methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-methoxypyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 3.88 (3H, s), 4.40 (2H, d, J=5.8Hz), 7.17 (1H, brs), 7.5-7.7 (2H, m), 7.8-8.0 (2H, m), 8.3-8.5 (2H, m), 8.83 (1H, s), 13.04 (1H, brs)

MS (m/z): 446 (M⁺)

[0389]

Example 124: N-(4-Chloro-3-{6-oxo-4-[6-propoxypyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)propionamide

By performing operations similar to those of Example 4 using 6-propoxypyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.9-1.1 (6H, m), 1.6-1.8 (2H, m), 2.16 (2H, q, J=7.8Hz), 4.2-4.4 (4H, m), 6.89 (1H, d, J=8.8Hz), 6.95 (1H, brs), 7.43 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.86 (1H, d, J=2.5Hz), 12.85 (1H, brs)

MS (m/z): 426 (M⁺)

[0390]

Example 125: N-(4-Chloro-3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide

By performing operations similar to those of Example 4 using 6-(2,2-difluoroethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.02 (3H, t, J=7.8Hz), 2.16 (2H, q, J=7.3Hz), 4.32 (2H, d, J=5.9Hz), 4.64 (2H, dt, J=3.4Hz, 15.1Hz), 6.41 (1H, tt, J=3.6Hz, 54.7Hz), 7.00 (1H, brs), 7.03 (1H, d, J=8.8Hz), 7.43 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.35 (1H, t, J=5.9Hz), 8.40 (1H, dd, J=2.5Hz, 8.8Hz), 8.90 (1H, d, J=2.0Hz), 12.89 (1H, brs)

MS (m/z): 448 (M⁺)

[0391]

Example 126: N-{4-Chloro-3-[4-(5-methylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-

methylpyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.36 (3H, s), 2.3-2.5 (1H, m), 4.32 (2H, d, J=5.9Hz), 7.05 (1H, brs), 7.42 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.23 (1H, s), 8.35 (1H, t, J=5.8Hz), 8.52 (1H, s), 9.03 (1H, d, J=1.4Hz), 12.95 (1H, brs)

MS (m/z): 396 (M⁺)

[0392]

Example 127: N-{3-[4-(5-Methylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-methylpyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.35 (3H, s), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.4Hz), 7.08 (1H, brs), 7.5-7.7 (2H, m), 7.89 (1H, d, J=7.8Hz), 8.21 (1H, s), 8.40 (1H, t, J=5.9Hz), 8.51 (1H, s), 9.02 (1H, s), 13.01 (1H, brs)

MS (m/z): 430 (M⁺)

[0393]

Example 128: N-{4-Chloro-2-fluoro-3-[4-(5-methylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-methylpyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.36 (3H, s), 2.3-2.6 (1H, m), 4.32 (2H, d, J=5.9Hz), 7.11 (1H, brs), 7.4-7.6 (2H, m), 8.21 (1H, s), 8.36 (1H, t, J=5.4Hz), 8.52 (1H, s), 9.01 (1H, d, J=1.2Hz), 13.17 (1H, brs)

MS (m/z): 414 (M⁺)

[0394]

Example 129: N-{4-Chloro-3-[4-(5-methoxypyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-methoxypyridine-2-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.44 (1H, sept, J=6.8Hz), 3.90 (3H, s), 4.32 (2H, d, J=5.9Hz), 7.12 (1H, brs), 7.42 (1H, dd, J=2.4Hz, 8.3Hz), 7.50 (1H, dd, J=3.0Hz, 8.8Hz), 7.5-7.6 (2H, m), 8.21 (1H, d, J=8.8Hz), 8.34 (1H, t, J=5.9Hz), 8.43 (1H, d, J=3.0Hz), 12.84 (1H, brs)

MS (m/z): 412 (M⁺)

[0395]

Example 130: N-{4-Chloro-3-[4-(1-isopropyl-1H-pyrazol-4-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 1-isopropyl-1H-pyrazole-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.42 (6H, d, J=6.4Hz), 2.43 (1H, sept, J=6.8Hz), 4.30 (2H, d, J=5.8Hz), 4.52 (1H, sept, J=6.3Hz), 6.62 (1H, brs), 7.39 (1H, dd, J=1.9Hz, 8.3Hz), 7.47 (1H, d, J=1.9Hz), 7.54 (1H, d, J=8.3Hz), 8.00 (1H, s), 8.2-8.4 (2H, m), 12.39 (1H, brs)

MS (m/z): 413 (M⁺)

[0396]

Example 131: N-{4-Chloro-3-[6-oxo-4-(1-propyl-1H-pyrazol-4-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 1-propyl-1H-pyrazole-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.82 (3H, t, J=7.6Hz), 1.03 (6H, d, J=6.8Hz), 1.7-1.9 (2H, m), 2.43 (1H, sept, J=6.8Hz), 4.08 (2H, t, J=6.8Hz), 4.30 (2H, d, J=6.3Hz), 6.61 (1H, brs), 7.39 (1H, dd, J=1.9Hz, 8.3Hz), 7.48 (1H, d, J=2.5Hz), 7.54 (1H, d, J=7.8Hz), 8.01 (1H, s), 8.2-8.4 (2H, m), 12.13 (1H, brs)

MS (m/z): 413 (M⁺)

[0397]

Example 132: N-(4-Difluoromethoxy-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(difluoromethoxy)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.4Hz), 6.9-8.0 (7H, m), 8.34 (1H, t, J=5.9Hz), 8.56 (1H, dd, J=2.4Hz, 8.8Hz), 8.97 (1H, d, J=2.0Hz), 12.77 (1H, brs)

MS (m/z): 480 (M⁺)

[0398]

Example 133: N-[4-Chloro-3-(4-{6-[2-(2-methoxyethoxy)ethoxy]pyridin-3-yl}-6-oxo-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-[2-(2-methoxyethoxy)ethoxy]pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.24 (3H, s), 3.4-3.6 (4H, m), 3.7-3.8 (2H, m), 4.31 (2H, d, J=5.8Hz), 4.4-4.5 (2H, m), 6.92 (1H, d,

J=8.8Hz), 6.96 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.87 (1H, d, J=1.9Hz), 12.86 (1H, brs)

MS (m/z): 500 (M⁺)

[0399]

Example 134: N-{4-Chloro-3-[4-(6-{2-[2-(2-ethoxyethoxy)ethoxy]ethoxy}pyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-{2-[2-(2-ethoxyethoxy)ethoxy]ethoxy}pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.08 (3H, t, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 3.3-3.8 (12H, m), 4.31 (2H, d, J=5.8Hz), 4.4-4.5 (2H, m), 6.92 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.42 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.84 (1H, brs)

MS (m/z): 558 (M⁺)

[0400]

Example 135: N-(4-Chloro-3-{4-[6-(6-methylpyridin-2-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(6-methylpyridin-2-ylmethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 2.47 (3H, s), 4.31 (2H, d, J=5.8Hz), 5.44 (2H, s), 6.97 (1H, brs), 7.04 (1H, d, J=8.8Hz), 7.18 (1H, d, J=7.8Hz), 7.24 (1H, d, J=7.3Hz), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 7.68 (1H, t, J=7.8Hz), 8.33 (1H, t, J=5.8Hz), 8.38 (1H, dd, J=2.4Hz, 8.8Hz), 8.87 (1H, d, J=2.4Hz), 12.71 (1H, brs)

MS (m/z): 503 (M⁺)

[0401]

Example 136: N-(4-Chloro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)propionamide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.3Hz), 1.02 (3H, t, J=7.6Hz), 1.4-1.6 (2H, m), 2.16 (2H, q, J=7.8Hz), 3.40 (2H, t, J=6.8Hz), 3.6-3.8 (2H, m), 4.32 (2H, d, J=5.9Hz), 4.4-4.5 (2H, m), 6.93 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.43 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.3-8.4 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.74 (1H, brs)

MS (m/z): 470 (M⁺)

[0402]

Example 137: N-(4-Chloro-3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(2-ethoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=8.1Hz), 1.12 (3H, t, J=7.8Hz), 1.4-1.6 (2H, m), 2.13 (2H, t, J=7.3Hz), 3.49 (2H, q, J=7.3Hz), 3.6-3.8 (2H, m), 4.32 (2H, d, J=5.8Hz), 4.3-4.5 (2H, m), 6.93 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.43 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.7 (2H, m), 8.2-8.5 (2H, m), 8.87 (1H, d, J=2.4Hz), 12.87 (1H, brs)

MS (m/z): 470 (M⁺)

[0403]

Example 138: N-(4-Chloro-3-{4-[6-(2-fluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(2-fluoroethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.3Hz), 1.4-1.6 (2H, m), 2.13 (2H, t, J=7.1Hz), 4.32 (2H, d, J=5.8Hz), 4.5-4.9 (4H, m), 6.97 (1H, d, J=8.3Hz), 6.97 (1H, brs), 7.43 (1H, dd, J=2.5Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.3-8.5 (2H, m), 8.88 (1H, d, J=2.5Hz), 12.87 (1H, brs)

MS (m/z): 444 (M⁺)

[0404]

Example 139: N-(4-Chloro-3-{4-[6-(3-fluoropropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)propionamide

By performing operations similar to those of Example 4 using 6-(3-fluoropropoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)propionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.02 (3H, t, J=7.8Hz), 2.0-2.3 (4H, m), 4.32 (2H, d, J=6.4Hz), 4.43 (2H, t, J=6.3Hz), 4.55 (1H, t, J=6.4Hz), 4.66 (1H, t, J=5.9Hz), 6.92 (1H, d, J=8.8Hz), 6.96 (1H, brs), 7.43 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.3-8.4 (2H, m), 8.88 (1H, d, J=2.5Hz), 12.87 (1H, brs)

MS (m/z): 444 (M⁺)

[0405]

Example 140: N-(3-{4-[6-(2-Ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-

ethoxyethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 1.12 (3H, t, J=7.2Hz), 2.46 (1H, sept, J=7.2Hz), 3.49 (2H, q, J=7.2Hz), 3.7-3.8 (2H, m), 4.3-4.4 (4H, m), 6.92 (1H, d, J=8.8Hz), 6.96 (1H, s), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.0Hz), 8.31 (1H, d, J=8.8Hz), 8.39 (1H, t, J=6.0Hz), 8.84 (1H, s), 12.97 (1H, s)

MS (m/z): 504 (M⁺)

[0406]

Example 141: N-(3-{6-Oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.2Hz), 1.05 (6H, d, J=7.2Hz), 1.5-1.6 (2H, m), 2.46 (1H, sept, J=6.8Hz), 3.40 (2H, t, J=6.8Hz), 3.6-3.7 (2H, m), 4.4-4.5 (4H, m), 6.92 (1H, d, J=8.4Hz), 6.97 (1H, s), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.0Hz), 8.31 (1H, d, J=8.0Hz), 8.39 (1H, t, J=6.0Hz), 8.85 (1H, s), 12.97 (1H, s)

MS (m/z): 518 (M⁺)

[0407]

Example 142: N-{3-[4-(6-Cyclopropylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-cyclopropylpyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.0-1.1(10H, m), 2.1-2.2 (1H, m), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.0Hz), 7.01 (1H, s), 7.39 (1H, d, J=8.0Hz), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.0Hz), 8.23 (1H, d, J=6.8Hz), 8.39 (1H, t, J=6.0Hz), 9.04 (1H, s), 13.00 (1H, s)

MS (m/z): 456 (M⁺)

[0408]

Example 143: N-(3-{4-[6-(2-Butoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-butoxyethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.2Hz), 1.05 (6H, d, J=6.8Hz), 1.2-1.3 (2H, m), 1.4-1.5 (2H, m), 2.4-2.6 (1H, m), 3.43 (2H, t, J=6.8Hz), 3.6-3.7 (2H, m), 4.4-4.5 (4H, m),

6.91 (1H, d, J=8.8Hz), 6.96 (1H, s), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.0Hz), 8.31 (1H, d, J=8.8Hz), 8.39 (1H, t, J=6.0Hz), 8.84 (1H, s), 12.96 (1H, s)

MS (m/z): 532 (M⁺)

[0409]

Example 144: N-(3-{4-[6-(3-Methoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-methoxypropoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 1.96 (2H, quin, J=6.8Hz), 2.4-2.6 (1H, m), 3.24 (3H, s), 3.46 (2H, t, J=6.4Hz), 4.36 (2H, t, J=6.8Hz), 4.40 (2H, d, J=5.6Hz), 6.89 (1H, d, J=8.8Hz), 6.96 (1H, s), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.30 (1H, d, J=8.4Hz), 8.39 (1H, t, J=6.4Hz), 8.84 (1H, s), 12.97 (1H, s)

MS (m/z): 504 (M⁺)

[0410]

Example 145: N-[4-Chloro-3-(6-oxo-4-pyridin-3-yl-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.32 (2H, d, J=6.0Hz), 7.09 (1H, s), 7.42 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (3H, m), 8.34 (1H, t, J=6.0Hz), 8.4-8.5 (1H, m), 8.68 (1H, dd, J=1.6Hz, 4.8Hz), 9.24 (1H, d, J=2.0Hz), 12.99 (1H, s)

MS (m/z): 382 (M⁺)

[0411]

Example 146: N-{3-[4-(6-Methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-methylbenzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-methoxypyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-methylbenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (4H, m), 3.92 (3H, s), 4.28 (2H, d, J=6.0Hz), 6.9-7.0 (2H, m), 7.2-7.3 (2H, m), 7.42 (1H, s), 8.24 (1H, t, J=6.0Hz), 8.34 (1H, dd, J=2.4Hz, 8.8Hz), 8.89 (1H, d, J=2.0Hz), 12.65 (1H, s)

MS (m/z): 392 (M⁺)

[0412]

Example 147: N-(4-Chloro-3-{6-oxo-4-[2-(trifluoromethyl)pyridin-4-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(trifluoromethyl)pyridine-4-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.36 (1H, s), 7.44 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.3-8.4 (2H, m), 8.44 (1H, s), 8.91 (1H, d, J=5.6Hz), 13.19 (1H, s)

MS (m/z): 450 (M⁺)

[0413]

Example 148: N-[4-(Difluoromethyl)-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(difluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.39 (2H, d, J=6.0Hz), 7.26 (1H, s), 7.43 (1H, m), 7.58 (1H, d, J=8.0Hz), 7.68 (1H, t, J=98.4Hz), 7.80 (1H, d, J=8.0Hz), 8.05 (1H, d, J=8.4Hz), 8.34 (1H, t, J=5.6Hz), 8.72 (1H, d, J=7.2Hz), 9.44 (1H, s), 13.17 (1H, s)

MS (m/z): 466 (M⁺)

[0414]

Example 149: N-(3-{4-[2-(Cyclopropylmethoxy)pyridin-4-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(cyclopropylmethoxy)pyridine-4-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.05 (6H, d, J=6.8Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 4.13 (2H, d, J=6.8Hz), 4.40 (2H, d, J=6.0Hz), 7.14 (1H, s), 7.40 (1H, s), 7.5-7.7 (3H, m), 7.89 (1H, d, J=8.0Hz), 8.23 (1H, d, J=5.6Hz), 8.39 (1H, t, J=6.4Hz), 13.13 (1H, s)

MS (m/z): 486 (M⁺)

[0415]

Example 150: N-(3-{4-[6-(Cyclopropylmethoxy)pyridazin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylmethoxy)pyridazine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.4-0.6 (4H, m), 1.05 (6H, d, J=7.2Hz), 1.3-1.4 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.03 (2H, d, J=7.6Hz), 4.40 (2H, d, J=5.6Hz), 7.0-7.1 (2H, m), 7.6-7.7

(2H, m), 7.88 (1H, d, J=8.4Hz), 8.0-8.1 (1H, m), 8.40 (1H, t, J=6.4Hz), 13.10 (1H, s)

MS (m/z): 487 (M⁺)

[0416]

Example 151: 4-(2-(2-Chloro-5-[(3,3,3-trifluoro-2,2-dimethylpropionylamino)methyl]phenyl)-6-oxo-1,6-dihydropyrimidin-4-yl)-N,N-dimethylbenzamide

By performing operations similar to those of Example 4 using 4-formyl-N,N-dimethylbenzamide, and N-(3-carbamimidoyl-4-chlorobenzyl)-3,3,3-trifluoro-2,2-dimethylpropionamide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.38 (6H, s), 2.92 (3H, s), 3.00 (3H, s), 4.37 (2H, d, J=6.0Hz), 7.00 (1H, s), 7.40 (1H, dd, J=2.0Hz, 8.4Hz), 7.49 (2H, d, J=8.4Hz), 7.55 (1H, d, J=2.0Hz), 7.58 (1H, d, J=8.4Hz), 8.13 (2H, d, J=8.0Hz), 8.61 (1H, t, J=6.0Hz), 12.89 (1H, brs)

MS (m/z): 520 (M⁺)

[0417]

Example 152: N-(3-{4-[6-(Difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.32 (2H, d, J=5.6Hz), 7.12 (1H, s), 7.20 (1H, d, J=8.8Hz), 7.29 (1H, t, J=8.8Hz), 7.4-7.6 (1H, m), 7.77 (1H, t, J=72.8Hz), 8.33 (1H, t, J=5.6Hz), 8.52 (1H, dd, J=2.0Hz, 8.8Hz), 8.93 (1H, d, J=1.6Hz), 13.23 (1H, brs)

MS (m/z): 450 (M⁺)

[0418]

Example 153: N-[3-(2'-Methoxy-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 2-methoxypyrimidine-5-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 3.99 (3H, s), 4.40 (2H, d, J=6.0Hz), 7.10 (1H, s), 7.6-7.7 (2H, m), 7.89 (1H, d, J=8.4Hz), 8.39 (1H, t, J=6.0Hz), 9.19 (2H, s), 13.05 (1H, brs)

MS (m/z): 447 (M⁺)

[0419]

Example 154: N-{4-Chloro-3-[4-(5-chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-

yl}benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=7.2Hz), 2.4-2.6 (1H, m), 4.32 (2H, d, J=6.0Hz), 7.19 (1H, s), 7.43 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.34 (1H, t, J=5.6Hz), 8.50 (1H, s), 8.75 (1H, d, J=2.0Hz), 9.21 (1H, s), 13.08 (1H, brs)

MS (m/z): 416 (M⁺)

[0420]

Example 155: N-{4-Chloro-3-[4-(6-cyclopropylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-cyclopropylpyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.9-1.1 (4H, m), 1.04 (6H, d, J=6.8Hz), 2.1-2.2 (1H, m), 2.4-2.6 (1H, m), 4.32 (2H, d, J=5.6Hz), 7.04 (1H, s), 7.39 (1H, d, J=8.4Hz), 7.4-7.5 (2H, m), 8.23 (1H, dd, J=2.0Hz, 8.4Hz), 8.35 (1H, t, J=5.6Hz), 9.03 (1H, s), 13.13 (1H, brs)

MS (m/z): 440 (M⁺)

[0421]

Example 156: N-[4-Chloro-3-(2'-cyclopropyl-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-cyclopropylpyrimidine-5-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=7.2Hz), 1.0-1.2 (4H, m), 2.2-2.3 (1H, m), 2.43 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.4Hz), 7.10 (1H, s), 7.42 (1H, dd, J=2.0Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.33 (1H, t, J=6.0Hz), 9.21 (2H, s), 13.01 (1H, brs)

MS (m/z): 423 (M⁺)

[0422]

Example 157: N-(4-Chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylmethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.04 (6H, d, J=6.8Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 4.16 (2H, d, J=7.2Hz), 4.32 (2H, d, J=5.6Hz), 6.90 (1H, d, J=8.4Hz), 7.00 (1H, s), 7.4-7.5 (2H, m), 8.30 (1H, dd, J=2.4Hz, 8.8Hz), 8.35 (1H, t, J=5.6Hz), 8.82 (1H, d, J=2.0Hz), 13.06 (1H, brs)

MS (m/z): 470 (M⁺)

[0423]

Example 158: N-(4-Chloro-3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-ethoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 1.10 (3H, t, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 3.49 (2H, q, J=6.8Hz), 3.71 (2H, t, J=4.8Hz), 4.32 (2H, d, J=5.6Hz), 4.43 (2H, t, J=4.4Hz), 6.92 (1H, d, J=8.8Hz), 7.00 (1H, s), 7.4-7.5 (2H, m), 8.3-8.4 (2H, m), 8.84 (1H, d, J=1.6Hz), 13.10 (1H, brs)

MS (m/z): 488 (M⁺)

[0424]

Example 159: N-(3-{4-[6-(Cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylmethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.05 (6H, d, J=7.2Hz), 1.2-1.3 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.16 (2H, d, J=6.8Hz), 4.40 (2H, d, J=5.6Hz), 6.90 (1H, d, J=8.8Hz), 6.95 (1H, s), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.30 (1H, d, J=7.2Hz), 8.39 (1H, t, J=5.6Hz), 8.83 (1H, s), 12.95 (1H, brs)

MS (m/z): 486 (M⁺)

[0425]

Example 160: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.2Hz), 1.04 (6H, d, J=6.8Hz), 1.51 (2H, sext, J=7.2Hz), 2.45 (1H, sept, J=6.8Hz), 3.40 (2H, t, J=6.8Hz), 3.71 (2H, t, J=4.8Hz), 4.32 (2H, d, J=6.0Hz), 4.44 (2H, t, J=4.8Hz), 6.92 (1H, d, J=8.8Hz), 7.02 (1H, s), 7.4-7.5 (2H, m), 8.31 (1H, dd, J=2.8Hz, 8.8Hz), 8.36 (1H, t, J=6.0Hz), 8.84 (1H, d, J=2.4Hz), 13.00 (1H, brs)

MS (m/z): 502 (M⁺)

[0426]

Example 161: N-(4-Chloro-3-[4-(6-ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-ethoxypyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 1.33 (3H, t, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.2Hz), 4.37 (2H, q, J=6.8Hz), 6.88 (1H, d, J=8.8Hz), 7.00 (1H, s), 7.4-7.5 (2H, m), 8.30 (1H, dd, J=2.4Hz, 8.8Hz), 8.36 (1H, t, J=5.6Hz), 8.84 (1H, d, J=2.0Hz), 13.08 (1H, brs)

MS (m/z): 444 (M⁺)

[0427]

Example 162: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.33 (2H, d, J=5.6Hz), 7.34 (1H, s), 7.4-7.6 (2H, m), 8.3-8.5 (3H, m), 9.13 (1H, s), 13.42 (1H, brs)

MS (m/z): 468 (M⁺)

[0428]

Example 163: N-(3-{4-[6-(2-Ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-ethoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 1.12 (3H, t, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 3.49 (2H, q, J=6.8Hz), 3.71 (2H, t, J=4.8Hz), 4.32 (2H, d, J=5.6Hz), 4.44 (2H, t, J=4.8Hz), 6.93 (1H, d, J=8.8Hz), 7.00 (1H, s), 7.28 (1H, t, J=8.8Hz), 7.4-7.6 (1H, m), 8.3-8.4 (2H, m), 8.85 (1H, s), 13.11 (1H, brs)

MS (m/z): 472 (M⁺)

[0429]

Example 164: N-(3-{4-[6-(Cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylmethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.04 (6H, d, J=6.8Hz), 1.2-1.3 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.16 (2H, d, J=7.6Hz), 4.32 (2H, d, J=5.6Hz), 6.91 (1H, d, J=8.8Hz), 6.98 (1H, s), 7.28 (1H, t, J=8.8Hz), 7.4-7.6 (1H, m), 8.3-8.4 (2H, m), 8.83 (1H, d, J=1.6Hz), 13.10 (1H, brs)

MS (m/z): 454 (M⁺)

[0430]

Example 165: N-(2-Chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.34 (2H, d, J=5.2Hz), 7.13 (1H, s), 7.18 (1H, d, J=8.8Hz), 7.4-7.6 (2H, m), 7.76 (1H, t, J=72.8Hz), 8.37 (1H, t, J=6.0Hz), 8.51 (1H, dd, J=2.4Hz, 8.4Hz), 8.92 (1H, d, J=2.4Hz), 13.22 (1H, brs)

MS (m/z): 466 (M⁺)

[0431]

Example 166: N-(2-Chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylmethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.07 (6H, d, J=7.2Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 4.16 (2H, d, J=6.8Hz), 4.34 (2H, d, J=5.6Hz), 6.90 (1H, d, J=8.8Hz), 7.00 (1H, s), 7.4-7.6 (2H, m), 8.30 (1H, dd, J=2.4Hz, 8.8Hz), 8.37 (1H, t, J=5.6Hz), 8.82 (1H, d, J=2.4Hz), 12.95 (1H, brs)

MS (m/z): 470 (M⁺)

[0432]

Example 167: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2,2,2-trifluoroethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.32 (2H, d, J=6.0Hz), 5.07 (2H, q, J=8.8Hz), 7.0-7.1 (2H, m), 7.4-7.6 (2H, m), 8.36 (1H, t, J=6.0Hz), 8.41 (1H, d, J=8.0Hz), 8.88 (1H, s), 13.16 (1H, brs)

MS (m/z): 498 (M⁺)

[0433]

Example 168: N-{4-Chloro-3-[4-(5-chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=6.0Hz), 7.25 (1H, s), 7.4-7.6 (2H, m), 8.36 (1H, t, J=6.0Hz), 8.4-8.5 (1H, m), 8.75 (1H, d, J=2.4Hz), 9.18 (1H, d, J=2.0Hz), 13.29 (1H, brs)

MS (m/z): 434 (M⁺)

[0434]

Example 169: N-(2-Fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(trifluoromethyl)pyridin-3-yl]pyrimidin-4(3H)-one (109 mg), potassium N-[(trifluoroborate)methyl]isobutyramide (104 mg), bis(dibenzylideneacetone)palladium(0) (7.2 mg), 2-dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl (12 mg), sodium carbonate (53 mg), 1,4-dioxane (4 mL), and water (0.4 mL) were mixed, and reacted at 130°C for 100 minutes by applying microwaves. After the reaction, the reaction mixture was extracted with ethyl acetate, and the organic layer was dried over magnesium sulfate. The solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (35 mg).

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 2.48 (1H, sept, J=6.8Hz), 4.41 (2H, d, J=5.6Hz), 7.31 (1H, s), 7.67 (1H, t, J=8.0Hz), 7.79 (1H, d, J=8.4Hz), 8.01 (1H, d, J=8.4Hz), 8.40 (1H, t, J=6.0Hz), 8.64 (1H, d, J=7.6Hz), 9.36 (1H, s), 13.38 (1H, brs)

MS (m/z): 502 (M⁺)

[0435]

Example 170: N-(2-Chloro-4-fluoro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2,2,2-trifluoroethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=7.2Hz), 2.4-2.6 (1H, m), 4.34 (2H, d, J=6.0Hz), 5.06 (2H, q, J=9.2Hz), 7.0-7.1 (2H, m), 7.4-7.6 (2H, m), 8.38 (1H, t, J=6.0Hz), 8.41 (1H, dd, J=2.0Hz, 8.8Hz), 8.88 (1H, d, J=2.0Hz), 13.15 (1H, brs)

MS (m/z): 498 (M⁺)

[0436]

Example 171: N-{4-Chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=5.6Hz), 7.24 (1H, s), 7.4-7.6 (2H, m), 8.06 (1H, dd, J=2.4Hz, 8.8Hz), 8.22 (1H, d, J=8.8Hz), 8.37 (1H, t, J=5.6Hz), 8.79 (1H, d, J=2.4Hz), 13.31 (1H, brs)

MS (m/z): 434 (M⁺)

[0437]

Example 172: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[6-(3,3,3-trifluoropropoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3,3,3-trifluoropropoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 2.7-2.9 (2H, m), 4.32 (2H, d, J=6.0Hz), 4.56 (2H, t, J=5.6Hz), 6.93 (1H, d, J=8.8Hz), 7.02 (1H, s), 7.4-7.5 (2H, m), 8.3-8.4 (2H, m), 8.87 (1H, d, J=2.0Hz), 13.12 (1H, brs)

MS (m/z): 512 (M⁺)

[0438]

Example 173: N-(2-Chloro-4-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.35 (2H, d, J=5.6Hz), 7.35 (1H, s), 7.4-7.6 (2H, m), 8.34 (1H, dd, J=2.0Hz, 8.8Hz), 8.3-8.5 (2H, m), 9.13 (1H, s), 13.38 (1H, brs)

MS (m/z): 468 (M⁺)

[0439]

Example 174: N-(3-{4-[6-(Difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(difluoromethoxy)pyridin-3-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 2.48 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=5.6Hz), 7.14 (1H, s), 7.20 (1H, d, J=8.8Hz), 7.66 (1H, t, J=7.6Hz), 7.77 (1H, t, J=72.0Hz), 7.79 (1H, d, J=8.0Hz), 8.44 (1H, t, J=5.6Hz), 8.49 (1H, d, J=7.2Hz), 8.91 (1H, s), 13.29 (1H, brs)

MS (m/z): 500 (M⁺)

[0440]

Example 175: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[5-(2-propoxyethoxy)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.6Hz), 1.05 (6H, d, J=7.2Hz), 1.4-1.6 (2H, m), 2.46 (1H, sept, J=6.8Hz), 3.42 (2H, t, J=6.8Hz), 3.74 (2H, t, J=4.4Hz), 4.26 (2H, t, J=4.4Hz), 4.33 (2H, d, J=6.0Hz), 7.13 (1H, s), 7.4-7.6 (3H, m), 8.15 (1H, d, J=8.8Hz), 8.36 (1H, t, J=6.0Hz), 8.44 (1H, d, J=2.8Hz), 13.13 (1H, brs)

MS (m/z): 502 (M⁺)

[0441]

Example 176: N-(3-{4-[6-(2-Ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(2-ethoxyethoxy)pyridin-3-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 1.12 (3H, t, J=7.2Hz), 2.48 (1H, sept, J=6.8Hz), 3.49 (2H, q, J=6.8Hz), 3.70 (2H, t, J=4.8Hz), 4.4-4.5 (4H, m), 6.92 (1H, d, J=8.8Hz), 7.01 (1H, s), 7.65 (1H, t, J=7.6Hz), 7.78 (1H, d, J=7.6Hz), 8.29 (1H, d, J=8.8Hz), 8.43 (1H, t, J=6.0Hz), 8.82 (1H, s), 13.17 (1H, brs)

MS (m/z): 522 (M⁺)

[0442]

Example 177: N-(4-Chloro-3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2,2-difluoroethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 4.60-4.68 (2H, m), 6.2-6.6 (1H, m), 7.02 (1H, d, J=8.0Hz), 7.05 (1H, s), 7.4-7.5 (2H, m), 8.3-8.4 (2H, m), 8.87 (1H, d, J=2.0Hz), 13.12 (1H, brs)

MS (m/z): 480 (M⁺)

[0443]

Example 178: N-(2-Chloro-3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2,2-difluoroethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=7.2Hz), 2.4-2.6 (1H, m), 4.34 (2H, d, J=6.0Hz), 4.6-4.7 (2H, m), 6.2-6.6 (1H, m), 7.02 (1H, d, J=8.8Hz), 7.04 (1H, s), 7.4-7.6 (2H, m), 8.3-8.4 (2H, m), 8.86 (1H, d, J=2.0Hz), 13.14 (1H, brs)

MS (m/z): 480 (M⁺)

[0444]

Example 179: N-(4-Chloro-3-{4-[5-(difluoromethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(difluoromethoxy)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.46 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=6.0Hz), 7.21 (1H, s), 7.42 (1H, t, J=7.2Hz), 7.4-7.6 (2H, m), 7.78 (1H, dd, J=2.8Hz, 8.4Hz), 8.27 (1H, d, J=8.4Hz), 8.37 (1H, t, J=6.0Hz), 8.63 (1H, d, J=2.8Hz), 13.27 (1H, brs)

MS (m/z): 466 (M⁺)

[0445]

Example 180: N-(4-Chloro-3-{4-[5-(2-ethoxyethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(2-ethoxyethoxy)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 1.13 (3H, t, J=7.2Hz), 2.46 (1H, sept, J=7.2Hz), 3.51 (2H, q, J=7.2Hz), 3.73 (2H, t, J=4.4Hz), 4.25 (2H, t, J=4.4Hz), 4.33 (2H, d, J=5.6Hz), 7.14 (1H, s), 7.4-7.6 (3H, m), 8.15 (1H, d, J=8.8Hz), 8.36 (1H, t, J=6.0Hz), 8.44 (1H, d, J=2.8Hz), 13.12 (1H, brs)

MS (m/z): 488 (M⁺)

[0446]

Example 181: N-{3-[4-(5-Chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.44 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=5.6Hz), 7.19 (1H, s), 7.3-7.4 (1H, m), 7.4-7.6 (1H, m), 7.74 (1H, d, J=6.4Hz), 8.35 (1H, t, J=5.6Hz), 8.54 (1H, t, J=2.0Hz), 8.76 (1H, d, J=2.0Hz), 9.24 (1H, d, J=1.6Hz), 12.93 (1H, brs)

MS (m/z): 400 (M⁺)

[0447]

Example 182: N-(2-Fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[5-(trifluoromethyl)pyridin-2-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.41 (2H, d, J=5.6Hz), 7.35 (1H, s), 7.67 (1H, t, J=7.2Hz), 7.80 (1H, d, J=8.4Hz), 8.36 (2H, s), 8.45 (1H, t, J=5.6Hz), 9.13 (1H, s), 13.49 (1H, brs)

MS (m/z): 502 (M⁺)

[0448]

Example 183: N-(4-Chloro-3-{4-[5-(cyclopropylmethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylmethoxy)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.7 (2H, m), 1.05 (6H, d, J=7.6Hz), 1.1-1.4 (1H, m), 2.46 (1H, sept, J=7.2Hz), 3.97 (2H, d, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.13 (1H, s), 7.45-7.50 (3H, m), 8.14 (1H, d, J=8.4Hz), 8.36 (1H, t, J=5.6Hz), 8.42 (1H, d, J=3.2Hz), 13.12 (1H, brs)

MS (m/z): 470 (M⁺)

[0449]

Example 184: N-{4-Chloro-2-fluoro-3-[4-(5-fluoropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-fluoropyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d,

J=5.6Hz), 7.24 (1H, s), 7.4-7.6 (2H, m), 8.29 (1H, d, J=9.6Hz), 8.36 (1H, t, J=5.6Hz), 8.71 (1H, d, J=2.8Hz), 9.11 (1H, s), 13.30 (1H, brs)

MS (m/z): 418 (M⁺)

[0450]

Example 185: N-(2-Chloro-3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-ethoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=7.2Hz), 1.12 (3H, t, J=6.8Hz), 2.4-2.6 (1H, m), 3.49 (2H, q, J=6.8Hz), 3.71 (2H, t, J=4.4Hz), 4.34 (2H, d, J=6.0Hz), 4.43 (2H, t, J=4.4Hz), 6.92 (1H, d, J=8.8Hz), 7.01 (1H, s), 7.4-7.6 (2H, m), 8.31 (1H, dd, J=2.4Hz, 8.8Hz), 8.37 (1H, t, J=6.0Hz), 8.83 (1H, d, J=2.0Hz), 13.09 (1H, brs)

MS (m/z): 488 (M⁺)

[0451]

Example 186: N-(2-Chloro-3-{4-[5-(cyclopropylmethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylmethoxy)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.7 (2H, m), 1.07 (6H, d, J=6.8Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 3.97 (2H, d, J=7.6Hz), 4.35 (2H, d, J=6.0Hz), 7.14 (1H, s), 7.41-7.51 (3H, m), 8.14 (1H, d, J=8.8Hz), 8.38 (1H, t, J=5.6Hz), 8.42 (1H, d, J=2.8Hz), 13.11 (1H, brs)

MS (m/z): 470 (M⁺)

[0452]

Example 187: N-(2-Chloro-4-fluoro-3-{6-oxo-4-[6-(3,3,3-trifluoropropoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3,3,3-trifluoropropoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 2.7-2.9 (2H, m), 4.34 (2H, d, J=5.6Hz), 4.56 (2H, t, J=6.0Hz), 6.92 (1H, d, J=8.8Hz), 7.02 (1H, s), 7.4-7.6 (2H, m), 8.34 (1H, dd, J=2.8Hz, 8.8Hz), 8.37 (1H, t, J=6.0Hz), 8.86 (1H, d, J=1.6Hz), 13.11 (1H, brs)

MS (m/z): 512 (M⁺)

[0453]

Example 188: N-(4-Chloro-3-{6-oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(tetrahydropyran-4-ylmethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.33 (2H, dq, J=4.4Hz, 12.2Hz), 1.66 (2H, dd, J=2.0Hz, 12.7Hz), 1.9-2.1 (1H, m), 2.43 (1H, sept, J=6.8Hz), 3.2-3.4 (2H, m), 3.87 (2H, dd, J=3.0Hz, 11.8Hz), 4.19 (2H, d, J=6.4Hz), 4.31 (2H, t, J=5.8Hz), 6.90 (1H, d, J=8.8Hz), 6.95 (1H, brs), 7.41 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.86 (1H, d, J=2.5Hz), 12.83 (1H, brs)

MS (m/z): 496 (M⁺)

[0454]

Example 189: N-(3-{6-Oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(tetrahydropyran-4-ylmethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 1.32 (2H, dq, J=4.4Hz, 12.2Hz), 1.6-1.7 (2H, m), 1.9-2.1 (1H, m), 2.46 (1H, sept, J=6.8Hz), 3.2-3.4 (2H, m), 3.87 (2H, dd, J=3.4Hz, 11.2Hz), 4.18 (2H, d, J=6.3Hz), 4.40 (2H, d, J=6.3Hz), 6.90 (1H, d, J=8.8Hz), 6.97 (1H, brs), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.3Hz), 8.31 (1H, d, J=8.8Hz), 8.39 (1H, t, J=5.9Hz), 8.84 (1H, d, J=2.5Hz), 12.94 (1H, brs)

MS (m/z): 530 (M⁺)

[0455]

Example 190: N-(4-Chloro-3-{6-oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(tetrahydropyran-4-ylmethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chlorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.3Hz), 1.33 (2H, dq, J=4.4Hz, 12.2Hz), 1.4-1.7 (4H, m), 1.9-2.1 (1H, m), 2.13 (2H, t, J=7.3Hz), 3.2-3.4 (2H, m), 3.87 (2H, dd, J=3.0Hz, 11.3Hz), 4.19 (2H, d, J=6.9Hz), 4.32 (2H, t, J=5.8Hz), 6.90 (1H, d, J=8.8Hz), 6.95 (1H, brs), 7.43 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.33 (1H, dd, J=2.4Hz, 8.8Hz), 8.38 (1H, t, J=5.8Hz), 8.86 (1H, d, J=2.5Hz), 12.87 (1H, brs)

MS (m/z): 496 (M⁺)

[0456]

Example 191: N-{4-Chloro-2-fluoro-3-[6-oxo-4-(4-trifluoromethylthiazol-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-trifluoromethylthiazole-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=5.8Hz), 7.05 (1H, brs), 7.4-7.6 (2H, m), 8.37 (1H, t, J=5.4Hz), 8.76 (1H, s), 13.51 (1H, brs)

MS (m/z): 474 (M⁺)

[0457]

Example 192: N-{4-Chloro-3-[4-(5-ethynylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-ethynylpyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.8Hz), 4.52 (1H, s), 7.17 (1H, brs), 7.43 (1H, dd, J=1.9Hz, 8.3Hz), 7.5-7.7 (2H, m), 8.34 (1H, t, J=5.8Hz), 8.49 (1H, s), 8.78 (1H, d, J=1.9Hz), 9.24 (1H, d, J=2.0Hz), 13.05 (1H, brs)

MS (m/z): 406 (M⁺)

[0458]

Example 193: N-(4-Chloro-3-{6-oxo-4-[5-(1-propynyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(1-propynyl)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.10 (3H, s), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.8Hz), 7.14 (1H, brs), 7.43 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.7 (2H, m), 8.35 (1H, t, J=5.8Hz), 8.39 (1H, s), 8.68 (1H, d, J=2.0Hz), 9.17 (1H, d, J=1.5Hz), 13.02 (1H, brs)

MS (m/z): 420 (M⁺)

[0459]

Example 194: N-(3-{4-[6-(3-Methyloxetan-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-methyloxetan-3-ylmethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 1.35 (3H, s), 2.46 (1H, sept, J=6.9Hz), 4.30 (2H, d, J=5.8Hz), 4.3-4.5 (4H, m), 4.50 (2H, d, J=5.9Hz), 6.95 (1H, d, J=9.0Hz), 6.99 (1H, brs), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.3Hz), 8.33 (1H, dd, J=1.9Hz, 8.8Hz), 8.39 (1H, t, J=5.4Hz), 8.86 (1H, d, J=1.5Hz), 12.90 (1H, brs)

ESI(+)-MS: 517 (M⁺+1)

[0460]

Example 195: N-{3-[4-(5-Ethynylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-ethynylpyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.9Hz), 4.41 (2H, d, J=5.8Hz), 4.51 (1H, s), 7.20 (1H, brs), 7.5-7.7 (2H, m), 7.89 (1H, d, J=8.3Hz), 8.40 (1H, t, J=6.3Hz), 8.47 (1H, s), 8.77 (1H, d, J=2.0Hz), 9.22 (1H, d, J=2.0Hz), 13.12 (1H, brs)

MS (m/z): 440 (M⁺)

[0461]

Example 196: N-(4-Chloro-3-{4-[6-(3-methyloxetan-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-ylbenzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-(3-methyloxetan-3-ylmethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.36 (3H, s), 2.43 (1H, sept, J=6.8Hz), 4.2-4.4 (4H, m), 4.43 (2H, s), 4.50 (2H, d, J=5.8Hz), 6.9-7.1 (2H, m), 7.42 (1H, dd, J=2.5Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.88 (1H, d, J=2.4Hz), 12.86 (1H, brs)

ESI(+)-MS: 483 (M⁺+1)

[0462]

Example 197: N-(3-{6-Oxo-4-[5-(1-propynyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(1-propynyl)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.10 (3H, s), 2.46 (1H, sept, J=6.8Hz), 4.41 (2H, d, J=5.8Hz), 7.17 (1H, brs), 7.5-7.7 (2H, m), 7.89 (1H, d, J=7.8Hz), 8.3-8.5 (2H, m), 8.67 (1H, d, J=1.9Hz), 9.15 (1H, d, J=2.0Hz), 13.10 (1H, brs)

MS (m/z): 454 (M⁺)

[0463]

Example 198: N-(4-Chloro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-3-yl]-1,6-

dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.3-2.6 (1H, m), 4.32 (2H, d, J=6.4Hz), 7.31 (1H, s), 7.44 (1H, dd, J=2.4Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.34 (1H, t, J=6.0Hz), 8.73 (1H, s), 9.10 (1H, d, J=1.2Hz), 9.54 (1H, d, J=1.2Hz), 13.11 (1H, s)

MS (m/z): 450 (M⁺)

[0464]

Example 199: N-(3-{4-[6-(3-Ethoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-ethoxypropoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.6Hz), 1.10 (3H, t, J=6.8Hz), 1.95 (2H, quin, J=6.4Hz), 2.4-2.6 (1H, m), 3.42 (2H, q, J=6.8Hz), 3.50 (2H, t, J=6.4Hz), 4.36 (2H, t, J=6.8Hz), 4.40 (2H, d, J=6.0Hz), 6.89 (1H, d, J=8.4Hz), 6.96 (1H, s), 7.6-7.7 (2H, m), 7.88 (1H, d, J=8.0Hz), 8.30 (1H, d, J=8.8Hz), 8.39 (1H, t, J=6.0Hz), 8.85 (1H, s), 12.97 (1H, s)

MS (m/z): 518 (M⁺)

[0465]

Example 200: N-(4-Chloro-3-{4-[6-(2,2-difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2,2-difluoroethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.31 (2H, d, J=6.0Hz), 4.63 (2H, dt, J=3.6Hz, 15.2Hz), 6.41 (1H, tt, J=3.6Hz, 54.4Hz), 6.96 (1H, s), 7.02 (1H, d, J=8.8Hz), 7.40 (1H, dd, J=2.0Hz, 8.0Hz), 7.5-7.6 (2H, m), 8.34 (1H, t, J=6.0Hz), 8.39 (1H, dd, J=2.4Hz, 8.8Hz), 8.89 (1H, d, J=2.0Hz), 12.86 (1H, s)

MS (m/z): 462 (M⁺)

[0466]

Example 201: N-(3-{4-[6-(2,2-Difluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2,2-difluoroethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.6Hz), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.0Hz),

4.64 (2H, dt, J=3.2Hz, 15.2Hz), 6.41 (1H, tt, J=3.2Hz, 54.8Hz), 7.02 (2H, d, J=8.8Hz), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.3-8.4 (2H, m), 8.87 (1H, d, J=2.0Hz), 12.96 (1H, s)

MS (m/z): 496 (M⁺)

[0467]

Example 202: N-(3-{6-Oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2,2,2-trifluoroethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.4Hz), 5.06 (2H, q, J=8.8Hz), 7.04 (1H, s), 7.09 (1H, d, J=8.8Hz), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.3-8.4 (2H, m), 8.89 (1H, d, J=1.2Hz), 13.01 (1H, s)

MS (m/z): 514 (M⁺)

[0468]

Example 203: N-(3-{6-Oxo-4-[6-(3,3,3-trifluoropropoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3,3,3-trifluoropropoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 2.8-2.9 (2H, m), 4.40 (2H, d, J=6.0Hz), 4.56 (2H, t, J=6.0Hz), 6.93 (1H, d, J=8.8Hz), 6.98 (1H, s), 7.6-7.7 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.34 (1H, d, J=7.6Hz), 8.39 (1H, t, J=6.0Hz), 8.87 (1H, s), 12.99 (1H, s)

MS (m/z): 528 (M⁺)

[0469]

Example 204: N-(3-{4-[6-(2-Fluoroethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-fluoroethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.0Hz), 4.5-4.9 (4H, m), 6.9-7.0 (2H, m), 7.5-7.6 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.34 (1H, d, J=8.8Hz), 8.39 (1H, t, J=6.0Hz), 8.86 (1H, s), 12.98 (1H, s)

MS (m/z): 478 (M⁺)

[0470]

Example 205: N-(3-{4-[6-(3-Fluoropropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-fluoropropoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.0-2.2 (2H, m), 2.4-2.6 (1H, m), 4.3-4.7 (6H, m), 6.91 (1H, d, J=8.8Hz), 6.98 (1H, s), 7.6-7.7 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.32 (1H, dd, J=2.0Hz, 8.8Hz), 8.39 (1H, t, J=6.0Hz), 8.86 (1H, d, J=2.0Hz), 12.96 (1H, s)
MS (m/z): 492 (M⁺)
[0471]

Example 206: N-(3-{4-[2-(Difluoromethoxy)pyridin-4-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(difluoromethoxy)pyridine-4-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=5.6Hz), 7.25 (1H, s), 7.6-7.7 (3H, m), 7.73 (1H, t, J=72.8Hz), 7.8-7.9 (2H, m), 8.3-8.4 (2H, m), 13.23 (1H, s)
MS (m/z): 482 (M⁺)
[0472]

Example 207: N-(3-{6-Oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.41 (2H, d, J=5.6Hz), 7.33 (1H, s), 7.6-7.7 (2H, m), 7.90 (1H, d, J=8.4Hz), 8.3-8.4 (3H, m), 9.12 (1H, s), 13.23 (1H, s)
MS (m/z): 484 (M⁺)
[0473]

Example 208: N-(3-[4-(5-Chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.0Hz),

7.21 (1H, s), 7.6-7.7 (2H, m), 7.89 (1H, d, J=7.6Hz), 8.08 (1H, dd, J=2.8Hz, 8.8Hz), 8.19 (1H, d, J=8.0Hz), 8.40 (1H, t, J=6.0Hz), 8.78 (1H, d, J=2.4Hz), 13.17 (1H, s)

MS (m/z): 450 (M⁺)

[0474]

Example 209: N-(3-{6-Oxo-4-[5-(2-propoxyethoxy)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(propoxyethoxy)pyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.2Hz), 1.05 (6H, d, J=7.2Hz), 1.52 (2H, sext, J=7.2Hz), 2.4-2.6 (1H, m), 3.41 (2H, t, J=7.2Hz), 3.7-3.8 (2H, m), 4.2-4.3 (2H, m), 4.40 (2H, d, J=6.4Hz), 7.11 (1H, s), 7.52 (1H, dd, J=3.2Hz, 8.8Hz), 7.6-7.7 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.14 (1H, d, J=8.8Hz), 8.39 (1H, t, J=6.0Hz), 8.43 (1H, d, J=3.2Hz), 12.98 (1H, s)

MS (m/z): 518 (M⁺)

[0475]

Example 210: N-(3-{4-[5-(Difluoromethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(difluoromethoxy)pyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.0Hz), 7.18 (1H, s), 7.41 (1H, t, J=73.2Hz), 7.6-7.7 (2H, m), 7.80 (1H, dd, J=3.2Hz, 8.8Hz), 7.89 (1H, d, J=8.4Hz), 8.24 (1H, d, J=8.4Hz), 8.40 (1H, t, J=6.0Hz), 8.63 (1H, d, J=2.4Hz), 13.14 (1H, s)

MS (m/z): 482 (M⁺)

[0476]

Example 211: N-(4-Chloro-3-{4-[5-(cyclopropylmethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylmethoxy)pyridine-2-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.04 (6H, d, J=7.2Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 3.98 (2H, d, J=7.2Hz), 4.31 (2H, d, J=5.6Hz), 7.10 (1H, s), 7.41 (1H, dd, J=2.0Hz, 8.4Hz), 7.48 (1H, dd, J=2.8Hz, 8.4Hz), 7.5-7.6 (2H, m), 8.18 (1H, d, J=8.8Hz), 8.34 (1H, t, J=5.6Hz), 8.42 (1H, d, J=2.8Hz), 12.86 (1H, s)

MS (m/z): 452 (M⁺)

[0477]

Example 212: N-(3-{4-[5-(Cyclopropylmethoxy)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylmethoxy)pyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.05 (6H, d, J=7.2Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 3.97 (2H, d, J=6.8Hz), 4.40 (2H, d, J=6.0Hz), 7.11 (1H, s), 7.48 (1H, dd, J=2.8Hz, 8.4Hz), 7.6-7.7 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.13 (1H, d, J=8.4Hz), 8.3-8.4 (2H, m), 12.97 (1H, s)

MS (m/z): 486 (M⁺)

[0478]

Example 213: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.87 (3H, t, J=7.2Hz), 1.55 (2H, sext, J=7.2Hz), 2.14 (2H, t, J=7.2Hz), 4.33 (2H, d, J=6.0Hz), 7.31 (1H, s), 7.4-7.5 (2H, m), 8.02 (1H, d, J=8.4Hz), 8.40 (1H, t, J=6.0Hz), 8.67 (1H, d, J=8.0Hz), 9.39 (1H, s), 13.34 (1H, s)

MS (m/z): 468 (M⁺)

[0479]

Example 214: N-(4-Chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)butyramide

By performing operations similar to those of Example 4 using 6-(difluoromethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.87 (3H, t, J=7.6Hz), 1.55 (2H, sext, J=7.6Hz), 2.14 (2H, t, J=7.6Hz), 4.33 (2H, d, J=5.2Hz), 7.13 (1H, s), 7.19 (1H, d, J=8.4Hz), 7.4-7.5 (2H, m), 7.77 (1H, t, J=7.4Hz), 8.40 (1H, t, J=5.2Hz), 8.51 (1H, dd, J=0.8Hz, 8.8Hz), 8.92 (1H, s), 13.19 (1H, s)

MS (m/z): 466 (M⁺)

[0480]

Example 215: N-[3-(2'-Cyclopropylmethoxy-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 2-

(cyclopropylmethoxy)pyrimidine-5-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.05 (6H, d, J=6.8Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 4.22 (2H, d, J=6.8Hz), 4.40 (2H, d, J=6.0Hz), 7.09 (1H, s), 7.5-7.6 (2H, m), 7.89 (1H, d, J=8.4Hz), 8.39 (1H, t, J=6.0Hz), 9.1-9.2 (2H, m), 13.07 (1H, s)
MS (m/z): 487 (M⁺)

[0481]

Example 216: N-{3-[4-(6-Butoxypyridazin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-butoxypyridazine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.93 (3H, t, J=7.6Hz), 1.05 (6H, d, J=7.2Hz), 1.34 (2H, sext, J=7.6Hz), 1.77 (2H, quin, J=7.2Hz), 2.4-2.6 (1H, m), 4.17 (2H, t, J=7.6Hz), 4.39 (2H, d, J=6.0Hz), 6.96 (1H, s), 7.03 (1H, d, J=10.0Hz), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.4Hz), 7.99 (1H, d, J=9.6Hz), 8.39 (1H, t, J=6.0Hz), 13.17 (1H, s)
MS (m/z): 489 (M⁺)

[0482]

Example 217: N-{4-Chloro-3-[6-oxo-2'-(trifluoromethyl)-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-(trifluoromethyl)pyrimidine-5-carbaldehyde, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.34 (1H, s), 7.44 (1H, dd, J=2.0Hz, 8.0Hz), 7.5-7.6 (2H, m), 8.34 (1H, t, J=6.0Hz), 9.63 (2H, s), 13.21 (1H, s)
MS (m/z): 451 (M⁺)

[0483]

Example 218: N-(4-Chloro-3-{4-[(2-cyclopropylmethoxy)pyridin-4-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(cyclopropylmethoxy)pyridine-4-carbaldehyde, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.04 (6H, d, J=6.8Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 4.14 (2H, d, J=7.2Hz), 4.32 (2H, d, J=6.4Hz), 7.11 (1H, s), 7.4-7.5 (2H, m), 7.5-7.6 (3H, m), 8.24 (1H, d, J=5.6Hz), 8.34 (1H, t, J=5.6Hz), 13.05 (1H, s)
MS (m/z): 452 (M⁺)

[0484]

Example 219: N-{4-Chloro-2-fluoro-3-[4-(5-fluoropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-fluoropyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=6.0Hz), 7.21 (1H, s), 7.4-7.6 (2H, m), 7.84 (1H, dt, J=2.8Hz, 8.8Hz), 8.2-8.3 (1H, m), 8.37 (1H, t, J=6.0Hz), 8.74 (1H, d, J=3.2Hz), 13.28 (1H, brs)

MS (m/z): 418 (M⁺)

[0485]

Example 220: N-{2-Fluoro-3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-(6-methoxypyridin-3-yl)pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 2.48 (1H, sept, J=6.8Hz), 3.91 (3H, s), 4.40 (2H, d, J=6.0Hz), 6.91 (1H, d, J=8.8Hz), 7.01 (1H, s), 7.65 (1H, t, J=8.0Hz), 7.78 (1H, d, J=7.6Hz), 8.30 (1H, d, J=8.4Hz), 8.44 (1H, t, J=6.0Hz), 8.85 (1H, s), 13.17 (1H, brs)

MS (m/z): 464 (M⁺)

[0486]

Example 221: N-(4-Chloro-3-{4-[6-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylethynyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.0 (4H, m), 1.04 (6H, d, J=6.8Hz), 1.6-1.7 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.6Hz), 7.14 (1H, s), 7.4-7.5 (3H, m), 8.3-8.4 (2H, m), 9.13 (1H, d, J=1.6Hz), 13.22 (1H, brs)

MS (m/z): 464 (M⁺)

[0487]

Example 222: N-{2-Chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-2-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 4.35 (2H, d, J=6.0Hz), 7.25 (1H, s), 7.4-7.5 (2H, m), 8.05 (1H, dd, J=2.4Hz, 8.4Hz), 8.21 (1H, d, J=8.4Hz), 8.38 (1H, t, J=6.0Hz), 8.79 (1H, d, J=2.4Hz), 13.29 (1H, brs)

MS (m/z): 434 (M⁺)

[0488]

Example 223: N-{4-Chloro-3-[4-(4-cyanophenyl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-formylbenzonitrile, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.17 (1H, s), 7.4-7.5 (2H, m), 7.95 (2H, d, J=8.4Hz), 8.24 (2H, d, J=8.4Hz), 8.36 (1H, t, J=6.0Hz), 13.27 (1H, brs)

MS (m/z): 424 (M⁺)

[0489]

Example 224: N-{4-Chloro-2-fluoro-3-[6-oxo-4-(6-phenylpyridin-3-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-phenylpyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 4.34 (2H, d, J=5.6Hz), 7.20 (1H, s), 7.4-7.5 (5H, m), 8.09 (1H, d, J=8.4Hz), 8.1-8.2 (2H, m), 8.37 (1H, t, J=5.6Hz), 8.48 (1H, dd, J=2.0Hz, 8.0Hz), 9.31 (1H, d, J=1.6Hz), 13.17 (1H, brs)

MS (m/z): 476 (M⁺)

[0490]

Example 225: N-(2,4-Difluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=6.0Hz), 7.30 (1H, t, J=8.8Hz), 7.35 (1H, s), 7.53 (1H, q, J=8.0Hz), 8.3-8.4 (3H, m), 9.13 (1H, s), 13.40 (1H, brs)

MS (m/z): 452 (M⁺)

[0491]

Example 226: N-{3-[4-(5-Chloropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2,4-

difluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.24 (1H, s), 7.29 (1H, t, J=8.8Hz), 7.4-7.5 (1H, m), 8.33 (1H, t, J=5.6Hz), 8.49 (1H, s), 8.75 (1H, d, J=2.4Hz), 9.18 (1H, d, J=1.6Hz), 13.31 (1H, brs)

MS (m/z): 418 (M⁺)

[0492]

Example 227: N-(4-Chloro-3-{4-[6-(difluoromethyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(difluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=6.0Hz), 7.04 (1H, t, J=54.8Hz), 7.25 (1H, s), 7.4-7.5 (2H, m), 7.81 (1H, d, J=8.4Hz), 8.39 (1H, t, J=6.0Hz), 8.60 (1H, d, J=8.0Hz), 9.31 (1H, s), 13.35 (1H, brs)

MS (m/z): 450 (M⁺)

[0493]

Example 228: N-(2-Chloro-3-{4-[6-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylethynyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.0 (4H, m), 1.07 (6H, d, J=6.8Hz), 1.6-1.7 (1H, m), 2.4-2.6 (1H, m), 4.34 (2H, d, J=6.0Hz), 7.14 (1H, s), 7.4-7.5 (3H, m), 8.3-8.4 (2H, m), 9.13 (1H, d, J=1.6Hz), 13.21 (1H, brs)

MS (m/z): 464 (M⁺)

[0494]

Example 229: N-(4-Chloro-3-{4-[5-(difluoromethyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(difluoromethyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.46 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=6.0Hz), 7.23 (1H, t, J=55.2Hz), 7.32 (1H, s), 7.4-7.6 (2H, m), 8.15 (1H, d, J=7.6Hz),

8.3-8.4 (2H, m), 8.93 (1H, s), 13.33 (1H, brs)

MS (m/z): 450 (M⁺)

[0495]

Example 230: N-{3-[4-(5-Chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2,4-difluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-chloropyridine-2-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.23 (1H, s), 7.29 (1H, t, J=8.8Hz), 7.4-7.6 (1H, m), 8.08 (1H, dd, J=2.4Hz, 8.4Hz), 8.22 (1H, d, J=8.4Hz), 8.34 (1H, t, J=6.0Hz), 8.78 (1H, d, J=2.4Hz), 13.32 (1H, brs)

MS (m/z): 418 (M⁺)

[0496]

Example 231: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[6-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(trifluoromethyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.46 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=5.6Hz), 7.26 (1H, s), 7.4-7.5 (2H, m), 8.05 (1H, d, J=7.6Hz), 8.25 (1H, t, J=8.0Hz), 8.37 (1H, t, J=5.6Hz), 8.47 (1H, d, J=8.0Hz), 13.40 (1H, brs)

MS (m/z): 468 (M⁺)

[0497]

Example 232: N-(2-Chloro-4-fluoro-3-{6-oxo-4-[4-(trifluoromethyl)thiazol-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-(trifluoromethyl)thiazole-2-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=7.2Hz), 2.4-2.6 (1H, m), 4.35 (2H, d, J=6.0Hz), 7.05 (1H, s), 7.4-7.6 (2H, m), 8.39 (1H, t, J=6.0Hz), 8.76 (1H, s), 13.51 (1H, brs)

MS (m/z): 474 (M⁺)

[0498]

Example 233: N-(4-Chloro-3-{4-[2-(cyclopropylethynyl)thiazol-5-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-

(cyclopropylethynyl)thiazole-5-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.0 (4H, m), 1.04 (6H, d, J=6.8Hz), 1.6-1.7 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.0Hz), 7.07 (1H, s), 7.4-7.5 (2H, m), 8.36 (1H, t, J=5.6Hz), 8.59 (1H, s), 13.22 (1H, brs)

MS (m/z): 470 (M⁺)

[0499]

Example 234: N-(2-Chloro-3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.0 (4H, m), 1.07 (6H, d, J=7.2Hz), 1.6-1.7 (1H, m), 2.4-2.6 (1H, m), 4.34 (2H, d, J=6.0Hz), 7.23 (1H, s), 7.4-7.5 (2H, m), 7.88 (1H, dd, J=2.0Hz, 8.4Hz), 8.15 (1H, d, J=8.4Hz), 8.38 (1H, t, J=6.0Hz), 8.69 (1H, d, J=2.0Hz), 13.26 (1H, brs)

MS (m/z): 464 (M⁺)

[0500]

Example 235: N-(4-Chloro-3-{4-[6-(3-ethyloxetan-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-ethyloxetan-3-ylmethoxy)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.90 (3H, t, J=7.8Hz), 1.04 (6H, d, J=6.8Hz), 1.77 (2H, q, J=7.3Hz), 2.43 (1H, sept, J=6.8Hz), 4.2-4.6 (8H, m), 6.9-7.0 (2H, m), 7.42 (1H, dd, J=2.0Hz, 8.3Hz), 7.5-7.6 (2H, m), 8.2-8.4 (2H, m), 8.89 (1H, d, J=2.2Hz), 12.85 (1H, brs)

ESI(+)-MS: 497 (M⁺+1)

[0501]

Example 236: N-(3-{4-[6-(3-Ethyloxetan-3-ylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-ethyloxetan-3-ylmethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.90 (3H, t, J=7.3Hz), 1.05 (6H, d, J=6.8Hz), 1.77 (2H, q, J=7.3Hz), 2.46 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=5.9Hz), 4.40 (2H, d, J=5.9Hz), 4.45 (2H, d, J=5.9Hz), 4.48 (2H, s), 6.9-7.0 (2H, m), 7.5-7.7 (2H, m), 7.87 (1H, d, J=7.8Hz), 8.32 (1H, dd, J=2.4Hz, 8.7Hz), 8.40 (1H, t, J=5.9Hz), 8.86 (1H, d, J=2.0Hz), 12.92 (1H,

brs)

ESI(+)-MS: 531 (M⁺+1)

[0502]

Example 237: N-{4-Chloro-2-fluoro-3-[4-(4-methylthiazol-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 4-methylthiazole-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 2.47 (3H, s), 4.33 (2H, d, J=5.9Hz), 6.96 (1H, brs), 7.4-7.6 (2H, m), 7.59 (1H, s), 8.37 (1H, t, J=6.1Hz), 13.31 (1H, brs)

MS (m/z): 420 (M⁺)

[0503]

Example 238: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[4-(1-propynyl)thiazol-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-(1-propynyl)thiazole-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.09 (3H, s), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=5.9Hz), 6.99 (1H, brs), 7.4-7.6 (2H, m), 8.10 (1H, s), 8.36 (1H, t, J=5.8z), 13.39 (1H, brs)

MS (m/z): 444 (M⁺)

[0504]

Example 239: N-(4-Chloro-3-{4-[5-(cyclopropylethynyl)thiazol-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)thiazole-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.05 (6H, d, J=6.8Hz), 1.6-1.7 (1H, s), 2.45 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=5.9Hz), 6.99 (1H, brs), 7.4-7.6 (2H, m), 8.13 (1H, s), 8.37 (1H, t, J=6.1Hz), 13.38 (1H, brs)

MS (m/z): 470 (M⁺)

[0505]

Example 240: N-(4-Chloro-3-{4-[4-(cyclopropylethynyl)thiazol-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-

(cyclopropylethynyl)thiazole-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.04 (6H, d, J=6.8Hz), 1.5-1.7 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.9Hz), 6.98 (1H, brs), 7.4-7.6 (2H, m), 8.08 (1H, s), 8.36 (1H, t, J=5.4Hz), 13.39 (1H, brs)

MS (m/z): 470 (M⁺)

[0506]

Example 241: N-(3-{4-[5-(Cyclopropylethynyl)thiazol-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)thiazole-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.05 (6H, d, J=6.8Hz), 1.5-1.7 (1H, m), 2.46 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=5.9Hz), 6.94 (1H, brs), 7.5-7.7 (2H, m), 7.89 (1H, d, J=8.3Hz), 8.11 (1H, s), 8.40 (1H, t, J=5.8Hz), 13.23 (1H, brs)

MS (m/z): 486 (M⁺)

[0507]

Example 242: N-(3-{4-[6-(Cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylethynyl)pyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.0 (4H, m), 1.05 (6H, d, J=6.8Hz), 1.5-1.7 (1H, m), 2.46 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=6.4Hz), 7.18 (1H, brs), 7.54 (1H, dd, J=0.9Hz, 7.8Hz), 7.5-7.7 (2H, m), 7.8-8.0 (2H, m), 8.10 (1H, d, J=7.4Hz), 8.40 (1H, t, J=5.8Hz), 13.11 (1H, brs)

MS (m/z): 480 (M⁺)

[0508]

Example 243: N-(4-Chloro-3-{4-[5-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyridine-3-carbaldehyde, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.04 (6H, d, J=6.8Hz), 1.5-1.7 (1H, m), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.9Hz), 7.13 (1H, brs), 7.43 (1H, dd, J=2.5Hz, 8.3Hz), 7.5-7.7 (2H, m), 8.3-8.4 (2H, m), 8.65 (1H, d, J=1.9Hz), 9.15 (1H, d, J=2.0Hz), 13.02 (1H, brs)

MS (m/z): 446 (M⁺)

[0509]

Example 244: N-(3-{4-[5-(Cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained. ¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.05 (6H, d, J=6.8Hz), 1.5-1.7 (1H, m), 2.46 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=5.9Hz), 7.16 (1H, brs), 7.5-7.7 (2H, m), 7.89 (1H, d, J=8.3Hz), 8.34 (1H, s), 8.40 (1H, t, J=5.8Hz), 8.64 (1H, d, J=2.0Hz), 9.13 (1H, d, J=1.5Hz), 13.11 (1H, brs)

MS (m/z): 480 (M⁺)

[0510]

Example 245: N-(3-{4-[6-(3-Morpholin-4-ylpropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(3-morpholin-4-ylpropoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained. ¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 1.89 (2H, sext, J=6.8Hz), 2.3-2.6 (7H, m), 3.56 (4H, t, J=4.8Hz), 4.35 (2H, t, J=6.8Hz), 4.40 (2H, d, J=5.6Hz), 6.88 (1H, d, J=8.8Hz), 6.98 (1H, s), 7.5-7.6 (2H, m), 7.88 (1H, d, J=7.6Hz), 8.30 (1H, dd, J=2.4Hz, 8.8Hz), 8.39 (1H, t, J=6.0Hz), 8.85 (1H, d, J=2.4Hz), 12.90 (1H, s)

MS (m/z): 559 (M⁺)

[0511]

Example 246: N-(3-{6-Oxo-4-[5-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained. ¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 4.41 (2H, d, J=5.6Hz), 7.35 (1H, s), 7.6-7.7 (2H, m), 7.91 (1H, d, J=8.0Hz), 8.42 (1H, t, J=6.0Hz), 8.73 (1H, s), 9.10 (1H, s), 9.54 (1H, s), 13.23 (1H, s)

MS (m/z): 484 (M⁺)

[0512]

Example 247: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.43 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=6.0Hz), 7.36 (1H, s), 7.4-7.5 (2H, m), 8.36 (1H, t, J=6.0Hz), 8.70 (1H, s), 9.10 (1H, d, J=1.2Hz), 9.51 (1H, d, J=1.6Hz), 13.33 (1H, s)

MS (m/z): 468 (M⁺)

[0513]

Example 248: N-[3-(2'-Butyl-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 2-butylpyrimidine-5-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.90 (3H, t, J=7.2Hz), 1.05 (6H, d, J=6.8Hz), 1.34 (2H, sext, J=7.6Hz), 1.76 (2H, quin, J=7.6Hz), 2.46 (1H, sept, J=6.8Hz), 2.93 (2H, t, J=7.2Hz), 4.40 (2H, d, J=6.0Hz), 7.17 (1H, s), 7.6-7.7 (2H, m), 7.89 (1H, d, J=8.4Hz), 8.39 (1H, t, J=6.4Hz), 9.28 (2H, s), 13.15 (1H, s)

MS (m/z): 473 (M⁺)

[0514]

Example 249: N-(3-{6-Oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.2Hz), 1.55 (2H, sext, J=7.2Hz), 2.15 (2H, t, J=7.2Hz), 4.42 (2H, d, J=6.0Hz), 7.31 (1H, s), 7.6-7.7 (2H, m), 7.90 (1H, d, J=7.6Hz), 8.38 (2H, s), 8.44 (1H, t, J=6.0Hz), 9.12 (1H, s), 13.26 (1H, s)

MS (m/z): 484 (M⁺)

[0515]

Example 250: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)butyramide

By performing operations similar to those of Example 4 using 5-(trifluoromethyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.87 (3H, t, J=7.2Hz), 1.55 (2H, sext, J=7.2Hz), 2.14 (2H, t, J=7.2Hz), 4.34 (2H, d, J=6.0Hz), 7.36 (1H, s), 7.5-7.6 (2H, m), 8.3-8.5 (3H, m), 9.13 (1H,

d, J=1.2Hz), 13.38 (1H, s)

MS (m/z): 468 (M⁺)

[0516]

Example 251: N-{4-Chloro-2-fluoro-3-[6-oxo-4-(6-propylpyridin-3-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-propylpyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.91 (3H, t, J=7.2Hz), 1.04 (6H, d, J=6.8Hz), 1.07 (2H, sext, J=7.2Hz), 2.45 (1H, sept, J=6.8Hz), 2.76 (2H, t, J=7.6Hz), 4.32 (2H, d, J=6.0Hz), 7.09 (1H, s), 7.36 (1H, d, J=7.6Hz), 7.4-7.5 (2H, m), 8.29 (1H, dd, J=2.4Hz, 8.0Hz), 8.36 (1H, t, J=6.0Hz), 9.11 (1H, d, J=2.0Hz), 13.10 (1H, s)

MS (m/z): 442 (M⁺)

[0517]

Example 252: N-{3-[6-Oxo-4-(6-propylpyridin-3-yl)-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-propylpyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.91 (3H, t, J=7.2Hz), 1.05 (6H, d, J=7.2Hz), 1.71 (2H, sext, J=7.2Hz), 2.4-2.6 (1H, m), 2.76 (2H, t, J=7.2Hz), 4.40 (2H, d, J=6.0Hz), 7.05 (1H, s), 7.35 (1H, d, J=8.4Hz), 7.6-7.7 (2H, m), 7.88 (1H, d, J=8.0Hz), 8.29 (1H, dd, J=2.4Hz, 8.4Hz), 8.39 (1H, t, J=6.0Hz), 9.12 (1H, d, J=1.6Hz), 13.00 (1H, s)

MS (m/z): 458 (M⁺)

[0518]

Example 253: N-{3-[2'-(3,3-Dimethyl-1-butynyl)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-(3,3-dimethyl-1-butynyl)pyrimidine-5-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 1.33 (9H, s), 2.4-2.6 (1H, m), 4.40 (2H, d, J=6.4Hz), 7.24 (1H, s), 7.6-7.7 (2H, m), 7.89 (1H, d, J=7.6Hz), 8.40 (1H, t, J=6.0Hz), 9.32 (2H, s), 13.19 (1H, s)

MS (m/z): 497 (M⁺)

[0519]

Example 254: N-{4-Chloro-3-[2'-(3,3-dimethyl-1-butynyl)-6-oxo-1,6-dihydro-[4,5'-

bipyrimidin-2-yl]-2-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-(3,3-dimethyl-1-butynyl)pyrimidine-5-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 1.33 (9H, s), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.6Hz), 7.28 (1H, s), 7.4-7.5 (2H, m), 8.36 (1H, t, J=6.0Hz), 9.31 (2H, s), 13.31 (1H, s)

MS (m/z): 481 (M⁺)

[0520]

Example 255: N-(4-Chloro-3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.0 (4H, m), 1.04 (6H, d, J=6.8Hz), 1.6-1.7 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.23 (1H, s), 7.4-7.5 (2H, m), 7.89 (1H, dd, J=2.0Hz, 8.4Hz), 8.15 (1H, d, J=8.0Hz), 8.36 (1H, t, J=5.6Hz), 8.69 (1H, d, J=2.0Hz), 13.26 (1H, brs)

MS (m/z): 464 (M⁺)

[0521]

Example 256: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[6-(1-propynyl)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(1-propynyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.11 (3H, s), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.14 (1H, s), 7.4-7.6 (3H, m), 8.3-8.4 (2H, m), 9.15 (1H, s), 13.23 (1H, brs)

MS (m/z): 438 (M⁺)

[0522]

Example 257: N-(2,4-Difluoro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.2Hz), 1.04 (6H, d, J=6.8Hz), 1.51 (2H, sext,

J=7.2Hz), 2.45 (1H, sept, J=6.8Hz), 3.40 (2H, t, J=6.8Hz), 3.71 (2H, t, J=4.4Hz), 4.32 (2H, d, J=6.0Hz), 4.44 (2H, t, J=4.4Hz), 6.92 (1H, d, J=8.8Hz), 7.00 (1H, s), 7.28 (1H, t, J=8.8Hz), 7.4-7.6 (1H, m), 8.3-8.4 (2H, m), 8.85 (1H, s), 13.11 (1H, brs)

MS (m/z): 486 (M⁺)

[0523]

Example 258: N-{4-Chloro-2-fluoro-3-[4-(6-methoxypyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 6-methoxypyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.45 (1H, sept, J=6.8Hz), 3.98 (3H, s), 4.33 (2H, d, J=6.0Hz), 6.9-7.0 (1H, m), 7.29 (1H, s), 7.4-7.5 (2H, m), 7.8-7.9 (2H, m), 8.37 (1H, t, J=6.0Hz), 13.15 (1H, brs)

MS (m/z): 430 (M⁺)

[0524]

Example 259: N-{2-Fluoro-3-[4-(5-fluoropyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-(5-fluoropyridin-3-yl)pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 2.48 (1H, sept, J=6.8Hz), 4.41 (2H, d, J=5.6Hz), 7.25 (1H, s), 7.66 (1H, t, J=7.2Hz), 7.80 (1H, d, J=8.0Hz), 8.27 (1H, d, J=10.4Hz), 8.44 (1H, t, J=5.6Hz), 8.71 (1H, d, J=2.4Hz), 9.10 (1H, s), 13.38 (1H, brs)

MS (m/z): 452 (M⁺)

[0525]

Example 260: N-[4-Chloro-2-fluoro-3-(4-{6-[(1-hydroxycyclohexyl)ethynyl]pyridin-3-yl}-6-oxo-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-[(1-hydroxycyclohexyl)ethynyl]pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 1.2-1.3 (1H, m), 1.4-1.7 (7H, m), 1.8-1.9 (2H, m), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 5.58 (1H, s), 7.16 (1H, s), 7.4-7.5 (2H, m), 7.57 (1H, d, J=8.0Hz), 8.3-8.4 (2H, m), 9.19 (1H, s), 13.25 (1H, brs)

MS (m/z): 522 (M⁺)

[0526]

Example 261: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[5-(1-propynyl)pyridin-2-yl]-1,6-

dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(1-propynyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.12 (3H, s), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=6.0Hz), 7.24 (1H, s), 7.4-7.5 (2H, m), 7.92 (1H, dd, J=2.0Hz, 8.4Hz), 8.17 (1H, d, J=8.4Hz), 8.36 (1H, t, J=5.6Hz), 8.72 (1H, d, J=2.0Hz), 13.27 (1H, brs)

MS (m/z): 438 (M⁺)

[0527]

Example 262: N-{3-[4-(6-Ethoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluoro-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-(6-ethoxypyridin-3-yl)pyrimidin-4(3H)-one, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 1.33 (3H, t, J=6.8Hz), 2.48 (1H, sept, J=6.8Hz), 4.3-4.5 (4H, m), 6.88 (1H, d, J=8.8Hz), 7.00 (1H, s), 7.65 (1H, t, J=8.0Hz), 7.78 (1H, d, J=8.4Hz), 8.28 (1H, d, J=8.4Hz), 8.43 (1H, t, J=5.6Hz), 8.83 (1H, s), 13.16 (1H, brs)

MS (m/z): 478 (M⁺)

[0528]

Reference Example 94: 6-(2-Isopropoxyethoxy)pyridine-3-carbaldehyde

[Formula 107]




A mixture of 6-chloropyridine-3-carbaldehyde (2.83 g), 2-isopropoxyethanol (10 mL), and potassium carbonate (3.32 g) was stirred at 150°C for 3 hours. The reaction mixture was left to cool, and then poured into saturated aqueous ammonium chloride. The resulting mixture was extracted with t-butyl methyl ether, and the organic layer was washed with brine. The solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (2.27 g).

¹H-NMR (CDCl₃, δ): 1.20 (6H, d, J=5.8Hz), 3.49 (1H, sept, J=6.3Hz), 3.7-3.9 (2H, m), 4.5-4.7 (2H, m), 6.89 (1H, d, J=8.8Hz), 8.06 (1H, dd, J=2.5Hz, 8.8Hz), 8.61 (1H, d, J=1.9Hz), 9.95 (1H, s)

MS (m/z): 210 (M⁺⁺¹)

[0529]

Reference Example 95: 6-[2-(2,2,2-Trifluoroethoxy)ethoxy]pyridine-3-carbaldehyde
[Formula 108]




By performing operations similar to those of Reference Example 94 using 2-(2,2,2-trifluoroethoxy)ethanol, the title compound was obtained.

¹H-NMR (CDCl₃, δ): 3.94 (2H, q, J=8.3Hz), 4.01 (2H, t, J=4.4Hz), 4.5-4.7 (2H, m), 6.90 (1H, d, J=8.3Hz), 8.09 (1H, dd, J=1.9Hz, 8.3Hz), 8.61 (1H, d, J=2.0Hz), 9.97 (1H, s)

MS (m/z): 249 (M⁺)

[0530]

Reference Example 96: Methyl 5-(3,3-dimethyl-1-butynyl)pyrazine-2-carboxylate
[Formula 109]




To a solution of methyl 5-chloropyrazine-2-carboxylate (1.5 g) in triethylamine (25 mL), 3,3-dimethyl-1-butyne (854 mg), dichlorobis(triphenylphosphine)palladium(II) (309 mg), and copper(I) iodide (42 mg) were added, and the resulting mixture was stirred at 60°C for 3 hours. After cooling, the solvent was evaporated under reduced pressure. To the obtained residue, ethyl acetate was added, the resulting mixture was washed with 2 N hydrochloric acid, saturated aqueous sodium hydrogencarbonate, and saturated brine, and the organic layer was dried over magnesium sulfate. The organic layer was filtered, and the solvent was evaporated under reduced pressure. Then, the obtained residue was purified by silica gel column chromatography, and dried under reduced pressure to obtain the title compound (1.54 g).

¹H-NMR (CDCl₃, δ): 1.38 (9H, s), 4.04 (3H, s), 8.67 (1H, d, J=1.2Hz), 9.21 (1H, d, J=1.2Hz)

MS (m/z): 218 (M⁺)

[0531]

Reference Example 97: [5-(3,3-Dimethyl-1-butynyl)pyrazin-2-yl]methanol
[Formula 110]




To a solution of lithium aluminum hydride (268 mg) in tetrahydrofuran (25 mL), a solution of methyl 5-(3,3-dimethyl-1-butynyl)pyrazine-2-carboxylate (1.54 g) in tetrahydrofuran (10 mL) was slowly added dropwise under a nitrogen atmosphere with ice cooling. The resulting mixture was warmed to room temperature, and stirred for 2 hours. Under ice cooling, saturated aqueous sodium hydrogencarbonate was slowly added to the reaction mixture, and ethyl acetate was added to the resulting mixture. The reaction mixture was filtered through a Celite layer, and then the organic layer was washed with saturated aqueous sodium hydrogencarbonate, and saturated brine, and dried over magnesium sulfate. The organic layer was filtered, the solvent was evaporated under reduced pressure, and the residue was dried under reduced pressure to obtain the title compound (1.21 g).

$^1\text{H-NMR}$ (CDCl_3 , δ): 1.17 (9H, s), 3.7-3.8 (1H, m), 4.82 (2H, d, $J=8.4\text{Hz}$), 8.4-8.6 (2H, m)
 MS (m/z): 190 (M^+)

[0532]

Reference Example 98: 5-(3,3-Dimethyl-1-butynyl)pyrazine-2-carbaldehyde

[Formula 111]



By performing operations similar to those of Reference Example 73 using [5-(3,3-dimethyl-1-butynyl)pyrazine-2-yl]methanol, the title compound was obtained.


$^1\text{H-NMR}$ (CDCl_3 , δ): 1.19 (9H, s), 8.64 (1H, d, $J=0.8\text{Hz}$), 9.06 (1H, d, $J=1.2\text{Hz}$), 10.11 (1H, s)

MS (m/z): 188 (M^+)

[0533]

Reference Example 99: Methyl 5-(cyclopropylethynyl)pyrazine-2-carboxylate

[Formula 112]



By performing operations similar to those of Reference Example 96 using ethynylcyclopropane, the title compound was obtained.


$^1\text{H-NMR}$ (CDCl_3 , δ): 0.9-1.1 (4H, m), 1.5-1.6 (1H, m), 4.03 (3H, s), 8.65 (1H, d, $J=1.6\text{Hz}$), 9.19 (1H, d, $J=1.6\text{Hz}$)

MS (m/z): 202 (M^+)

[0534]

Reference Example 100: [5-(Cyclopropylethynyl)pyrazin-2-yl]methanol

[Formula 113]



By performing operations similar to those of Reference Example 97 using methyl 5-(cyclopropylethynyl)pyrazine-2-carboxylate, the title compound was obtained.


$^1\text{H-NMR}$ (CDCl_3 , δ): 0.5-0.7 (2H, m), 0.9-1.0 (2H, m), 1.6-1.7 (1H, m), 3.7-3.8 (1H, m), 4.7-4.8 (2H, m), 8.3-8.6 (2H, m)

MS (m/z): 174 (M^+)

[0535]

Reference Example 101: 5-(Cyclopropylethynyl)pyrazine-2-carbaldehyde

[Formula 114]




By performing operations similar to those of Reference Example 73 using [5-(cyclopropylethynyl)pyrazin-2-yl]methanol, the title compound was obtained.

$^1\text{H-NMR}$ (CDCl_3 , δ): 0.7-0.8 (2H, m), 0.9-1.1 (2H, m), 1.7-1.8 (1H, m), 8.53 (1H, d, $J=1.6\text{Hz}$), 9.01 (1H, d, $J=1.2\text{Hz}$), 10.08 (1H, s)

MS (m/z): 172 (M^+)

[0536]

Reference Example 102: 2-(4-Methyl-1-pentynyl)pyrimidine-5-carbaldehyde
[Formula 115]




To a solution of 2-mercaptopyrimidine-5-carbaldehyde (259 mg) in acetonitrile (12 mL), 4-methyl-1-pentyne (347 μ L), palladium acetate (4.5 mg), triphenylphosphine (5.0 mg), and copper(I) 2-thiophenecarboxylate (706 mg) were added, and microwaves were irradiated on the resulting mixture at 110°C for 23 minutes. The reaction was cooled, then filtered through a Celite layer, and washed with ethyl acetate. The organic layer was washed with water, and saturated brine, and dried over magnesium sulfate. The organic layer was filtered, and the solvent was evaporated under reduced pressure. Then, the obtained residue was purified by silica gel column chromatography, and dried under reduced pressure to obtain the title compound (258 mg).

$^1\text{H-NMR}$ (CDCl_3 , δ): 1.09 (6H, d, $J=6.0\text{Hz}$), 2.03 (1H, sept, $J=6.8\text{Hz}$), 2.43 (2H, d, $J=6.4\text{Hz}$), 9.12 (2H, s), 10.13 (1H, s)

MS (m/z): 188 (M^+)

[0537]

Reference Example 103: 2-(1-Propynyl)thiazole-5-carbaldehyde
[Formula 116]




By performing operations similar to those of Reference Example 92 using 2-bromothiazole-5-carbaldehyde, the title compound was obtained.

$^1\text{H-NMR}$ (CDCl_3 , δ): 2.18 (3H, s), 8.34 (1H, s), 10.00 (1H, s)

MS (m/z): 151 (M^+)

[0538]

Reference Example 104: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-(5-chloropyridin-2-yl)pyrimidin-4(3H)-one
[Formula 117]



By performing operations similar to those of Example 4 using 5-chloropyridine-2-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.


$^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$, δ): 7.29 (1H, s), 7.87 (1H, d, $J=8.8\text{Hz}$), 8.0-8.2 (3H, m), 8.79 (1H, d, $J=2.4\text{Hz}$), 13.40 (1H, brs)

MS (m/z): 403 (M^+)

[0539]

Reference Example 105: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[5-(1-propynyl)pyridin-2-yl]pyrimidin-4(3H)-one

[Formula 118]



By performing operations similar to those of Example 4 using 5-(1-propynyl)pyridine-2-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.


$^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$, δ): 2.12 (3H, s), 7.29 (1H, s), 7.86 (1H, d, $J=8.4\text{Hz}$), 7.94 (1H, dd, $J=2.0\text{Hz}$, 8.0Hz), 8.0-8.2 (2H, m), 8.73 (1H, d, $J=2.0\text{Hz}$), 13.37 (1H, brs)

MS (m/z): 407 (M^+)

[0540]

Reference Example 106: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(cyclopropylethynyl)pyridin-3-yl]pyrimidin-4(3H)-one

[Formula 119]




By performing operations similar to those of Example 4 using 6-(cyclopropylethynyl)pyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.5-1.7 (1H, m), 7.19 (1H, s), 7.52 (1H, d, J=8.4Hz), 7.86 (1H, d, J=8.8Hz), 8.09 (1H, t, J=7.2Hz), 8.33 (1H, d, J=8.4Hz), 9.13 (1H, s), 13.32 (1H, brs)

MS (m/z): 433 (M⁺)

[0541]

Reference Example 107: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(2-propoxyethoxy)pyridin-3-yl]pyrimidin-4(3H)-one

[Formula 120]




By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.2Hz), 1.51 (2H, sext, J=7.2Hz), 3.40 (2H, t, J=6.8Hz), 3.71 (2H, t, J=4.8Hz), 4.44 (2H, t, J=4.8Hz), 6.93 (1H, d, J=8.8Hz), 7.06 (1H, s), 7.85 (1H, d, J=8.8Hz), 8.08 (1H, t, J=8.4Hz), 8.30 (1H, dd, J=2.0Hz, 8.8Hz), 8.84 (1H, s), 13.19 (1H, brs)

MS (m/z): 471 (M⁺)

[0542]

Reference Example 108: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(2-isopropoxyethoxy)pyridin-3-yl]pyrimidin-4(3H)-one

[Formula 121]



By performing operations similar to those of Example 4 using 6-(2-isopropoxyethoxy)pyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 1.09 (6H, d, J=6.0Hz), 3.61 (1H, sept, J=6.0Hz), 3.70 (2H, t, J=4.8Hz), 4.41 (2H, t, J=4.8Hz), 6.93 (1H, d, J=8.8Hz), 7.07 (1H, s), 7.85 (1H, d, J=8.4Hz), 8.08 (1H, t, J=8.4Hz), 8.30 (1H, dd, J=2.0Hz, 8.8Hz), 8.84 (1H, d, J=1.6Hz), 13.18 (1H, brs)

MS (m/z): 471 (M⁺)

[0543]

Reference Example 109: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[2-(cyclopropylethynyl)thiazol-5-yl]pyrimidin-4(3H)-one

[Formula 122]



By performing operations similar to those of Example 4 using 2-(cyclopropylethynyl)thiazole-5-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.


¹H-NMR (d₆-DMSO, δ): 0.8-1.1 (4H, m), 1.6-1.8 (1H, m), 7.13 (1H, s), 7.85 (1H, d, J=8.0Hz), 8.09 (1H, t, J=8.0Hz), 8.60 (1H, s), 13.31 (1H, brs)

MS (m/z): 439 (M⁺)

[0544]

Reference Example 110: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}pyrimidin-4(3H)-one

[Formula 123]



By performing operations similar to those of Example 4 using 6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridine-3-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzimidine hydrochloride, the title compound was obtained.


$^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$, δ): 3.95 (2H, t, $J=4.0\text{Hz}$), 4.14 (2H, q, $J=9.2\text{Hz}$), 4.48 (2H, t, $J=4.4\text{Hz}$), 6.95 (1H, d, $J=8.8\text{Hz}$), 7.08 (1H, s), 7.85 (1H, d, $J=8.8\text{Hz}$), 8.08 (1H, t, $J=8.4\text{Hz}$), 8.31 (1H, dd, $J=2.4\text{Hz}$, 8.8Hz), 8.84 (1H, d, $J=2.0\text{Hz}$), 13.15 (1H, brs)

MS (m/z): 511 (M^+)

[0545]

Reference Example 111: 2,4,6-Trichlorophenyl 5-(2,2,2-trifluoroethyl)pyridine-2-carboxylate

[Formula 124]



To a solution of 2-chloro-5-(2,2,2-trifluoroethyl)pyridine (1.93 g), 2,4,6-trichlorophenyl formate (2.67 g), palladium(II) acetate (67 mg), and 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (343 mg) in toluene (12 mL), a solution of triethylamine (2.8 mL) in toluene (20 mL) was added dropwise at 100°C under a nitrogen atmosphere, and the resulting mixture was stirred at 100°C for 1.5 hours. After the reaction, the reaction mixture was filtered through a Celite layer, the solvent was evaporated, and then the residue was purified by silica gel column chromatography to obtain the title compound (1.63 g).


$^1\text{H-NMR}$ (CDCl_3 , δ): 3.55 (2H, q, 10.8Hz), 7.44 (2H, s), 7.92 (1H, dd, $J=1.6\text{Hz}$, 8.0Hz), 8.33 (1H, d, $J=8.4\text{Hz}$), 8.81 (1H, d, $J=2.0\text{Hz}$)

MS (m/z): 383 (M^+)

[0546]

Reference Example 112: N-Methoxy-N-methyl-5-(2,2,2-trifluoroethyl)pyridine-2-carboxamide

[Formula 125]



A solution of 2,4,6-trichlorophenyl 5-(2,2,2-trifluoroethyl)pyridine-2-carboxylate (1.61 g), N,O-dimethylhydroxylamine hydrochloride (1.02 g), and tripotassium phosphate (4.45 g) in acetonitrile (40 mL) was refluxed overnight by heating under a nitrogen atmosphere. After the reaction, water was added to the reaction mixture, and the resulting mixture was extracted with ethyl acetate. Then, the organic layer was dried over sodium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography to obtain the title compound (471 mg).


$^1\text{H-NMR}$ (CDCl_3 , δ): 3.42 (3H, s), 3.45 (2H, q, 10.8Hz), 3.76 (3H, s), 7.6-7.8 (2H, m), 8.55 (1H, s)

MS (m/z): 217 (M^+-31)

[0547]

Reference Example 113: 5-(2,2,2-Trifluoroethyl)pyridine-2-carbaldehyde

[Formula 126]



To a solution of bis(cyclopentadienyl)zirconium(IV) chloride hydride (585 mg) in tetrahydrofuran (12 mL), a solution of N-methoxy-N-methyl-5-(2,2,2-trifluoroethyl)pyridine-2-carboxamide (469 mg) in tetrahydrofuran (5 mL) was added, and the resulting mixture was stirred for 30 minutes under a nitrogen atmosphere. After the reaction, silica gel was added to the reaction mixture, the resulting mixture was filtered through a Celite layer, and then purification was performed by silica gel column chromatography to obtain the title compound (142 mg).


$^1\text{H-NMR}$ (CDCl_3 , δ): 3.51 (2H, q, 10.8Hz), 7.85 (1H, d, $J=7.2\text{Hz}$), 7.99 (1H, d, $J=8.0\text{Hz}$), 8.73 (1H, d, $J=1.2\text{Hz}$), 10.10 (1H, s)

MS (m/z): 189 (M^+)

[0548]

Reference Example 114: 2-[3-Chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[5-(cyclopropylethynyl)pyridin-2-yl]pyrimidin-4(3H)-one

[Formula 127]



By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyridine-2-carbaldehyde, and 3-chloro-2-fluoro-6-(trifluoromethyl)benzamidinium hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.5-1.7 (1H, m), 7.28 (1H, s), 7.86 (1H, d, J=8.8Hz), 7.91 (1H, dd, J=2.0Hz, 8.4Hz), 8.0-8.2 (2H, m), 8.69 (1H, d, J=1.2Hz), 13.35 (1H, brs)

MS (m/z): 433 (M⁺)

[0549]

Example 263: N-{3-[4-(Benzothiazol-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-chloro-2-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using benzothiazole-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.48 (1H, sept, J=6.8Hz), 4.35 (2H, d, J=5.9Hz), 7.23 (1H, brs), 7.4-7.7 (4H, m), 8.1-8.3 (2H, m), 8.38 (1H, t, J=5.4Hz), 13.50 (1H, brs)

MS (m/z): 456 (M⁺)

[0550]

Example 264: N-(3-{4-[6-(2-Isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-isopropoxyethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 1.09 (6H, d, J=5.9Hz), 2.46 (1H, sept, J=6.8Hz), 3.61 (1H, sept, J=6.4Hz), 3.6-3.8 (2H, m), 4.3-4.5 (4H, m), 6.92 (1H, d, J=8.8Hz), 6.98 (1H, brs), 7.5-7.7 (2H, m), 7.88 (1H, d, J=7.8Hz), 8.31 (1H, dd, J=1.9Hz, 8.8Hz), 8.39 (1H, t, J=5.8Hz), 8.85 (1H, d, J=1.5Hz), 12.95 (1H, brs)

MS (m/z): 518 (M⁺)

[0551]

Example 265: N-(4-Chloro-2-fluoro-3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-

dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-isopropoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 1.10 (6H, d, J=5.9Hz), 2.45 (1H, sept, J=6.8Hz), 3.61 (1H, sept, J=6.3Hz), 3.70 (2H, t, J=4.9Hz), 4.32 (2H, d, J=5.8Hz), 4.41 (2H, t, J=4.9Hz), 6.92 (1H, d, J=8.8Hz), 7.01 (1H, brs), 7.4-7.6 (2H, m), 8.31 (1H, dd, J=2.4Hz, 8.7Hz), 8.36 (1H, t, J=5.8Hz), 8.84 (1H, d, J=2.0Hz), 13.08 (1H, brs)

MS (m/z): 502 (M⁺)

[0552]

Example 266: N-(2,4-Difluoro-3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-isopropoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=7.3Hz), 1.10 (6H, d, J=6.4Hz), 2.45 (1H, sept, J=6.8Hz), 3.62 (1H, sept, J=6.3Hz), 3.70 (2H, t, J=4.9Hz), 4.32 (2H, d, J=5.9Hz), 4.41 (2H, t, J=5.4Hz), 6.92 (1H, d, J=8.8Hz), 7.01 (1H, brs), 7.28 (1H, t, J=8.8Hz), 7.50 (1H, dd, J=7.9Hz), 8.2-8.4 (2H, m), 8.85 (1H, d, J=1.9Hz), 13.02 (1H, brs)

MS (m/z): 486 (M⁺)

[0553]

Example 267: N-(2-Chloro-4-fluoro-3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-isopropoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=6.8Hz), 1.09 (6H, d, J=6.4Hz), 2.4-2.6 (1H, m), 3.61 (1H, sept, J=5.9Hz), 3.70 (2H, t, J=4.9Hz), 4.34 (2H, d, J=5.8Hz), 4.41 (2H, t, J=5.4Hz), 6.91 (1H, d, J=8.8Hz), 7.01 (1H, brs), 7.3-7.6 (2H, m), 8.31 (1H, dd, J=2.4Hz, 8.8Hz), 8.37 (1H, t, J=5.9Hz), 8.84 (1H, d, J=2.0Hz), 13.06 (1H, brs)

MS (m/z): 502 (M⁺)

[0554]

Example 268: N-(3-{6-Oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(tetrahydropyran-4-ylmethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-

(trifluoromethyl)benzyl]butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.3Hz), 1.2-1.4 (2H, m), 1.55 (2H, sext, J=7.3Hz), 1.65 (2H, d, J=2.7Hz), 1.9-2.1 (1H, m), 2.15 (2H, t, J=7.3Hz), 3.2-3.5 (2H, m), 3.87 (2H, dd, J=2.9Hz, 11.2Hz), 4.18 (2H, d, J=6.3Hz), 4.41 (2H, d, J=5.4Hz), 6.90 (1H, d, J=8.8Hz), 6.98 (1H, s), 7.5-7.7 (2H, m), 7.87 (1H, d, J=8.3Hz), 8.30 (1H, dd, J=2.0Hz, 8.3Hz), 8.43 (1H, t, J=5.8Hz), 8.84 (1H, d, J=2.0Hz), 12.54 (1H, brs)

MS (m/z): 530 (M⁺)

[0555]

Example 269: N-(3-{4-[6-(2-Isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(2-isopropoxyethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.3Hz), 1.09 (6H, d, J=5.9Hz), 1.55 (2H, sext, J=7.3Hz), 2.15 (2H, t, J=7.3Hz), 3.61 (1H, sept, J=5.9Hz), 3.6-3.8 (2H, m), 4.3-4.5 (4H, m), 6.92 (1H, d, J=8.8Hz), 6.98 (1H, brs), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.8Hz), 8.31 (1H, dd, J=2.4Hz, 8.8Hz), 8.43 (1H, t, J=5.8Hz), 8.85 (1H, d, J=2.7Hz), 12.81 (1H, brs)

MS (m/z): 518 (M⁺)

[0556]

Example 270: N-(3-{4-[6-(2-Ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(2-ethoxyethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.86 (3H, t, J=7.3Hz), 1.12 (3H, t, J=7.3Hz), 1.55 (2H, sext, J=7.3Hz), 2.15 (2H, t, J=7.6Hz), 3.49 (2H, q, J=6.9Hz), 3.6-3.8 (2H, m), 4.3-4.5 (4H, m), 6.92 (1H, d, J=8.8Hz), 6.98 (1H, brs), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.3Hz), 8.2-8.4 (1H, m), 8.43 (1H, t, J=5.8Hz), 8.85 (1H, brs), 12.95 (1H, brs)

MS (m/z): 504 (M⁺)

[0557]

Example 271: N-(3-{6-Oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-0.9 (6H, m), 1.4-1.7 (4H, m), 2.15 (2H, t, J=7.6Hz), 3.40 (2H,

t, J=6.9Hz), 3.6-3.8 (2H, m), 4.3-4.5 (4H, m), 6.92 (1H, d, J=8.3Hz), 6.98 (1H, brs), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.4Hz), 8.31 (1H, dd, J=1.9Hz, 8.8Hz), 8.43 (1H, t, J=5.8Hz), 8.85 (1H, d, J=1.5Hz), 12.93 (1H, brs)

MS (m/z): 518 (M⁺)

[0558]

Example 272: N-{3-[4-(Benzothiazol-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2,4-difluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using benzothiazole-2-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 4.34 (2H, d, J=5.3Hz), 7.23 (1H, brs), 7.32 (1H, t, J=8.8Hz), 7.5-7.7 (3H, m), 8.1-8.3 (2H, m), 8.35 (1H, t, J=5.6Hz), 13.48 (1H, brs)

MS (m/z): 440 (M⁺)

[0559]

Example 273: N-{3-[4-(Benzothiazol-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using benzothiazole-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 2.47 (1H, sept, J=6.8Hz), 4.42 (2H, d, J=5.9Hz), 7.20 (1H, brs), 7.5-7.8 (4H, m), 7.92 (1H, d, J=8.3Hz), 8.1-8.3 (2H, m), 8.41 (1H, t, J=5.9Hz), 13.35 (1H, brs)

MS (m/z): 472 (M⁺)

[0560]

Example 274: N-(2-Chloro-4-fluoro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(2-propoxyethoxy)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.3Hz), 1.07 (6H, d, J=6.8Hz), 1.51 (2H, sext, J=7.3Hz), 2.4-2.6 (1H, m), 3.40 (2H, t, J=6.3Hz), 3.6-3.8 (2H, m), 4.35 (2H, d, J=5.9Hz), 4.4-4.5 (2H, m), 6.91 (1H, d, J=8.8Hz), 7.01 (1H, brs), 7.3-7.6 (2H, m), 8.31 (1H, dd, J=2.4Hz, 8.8Hz), 8.37 (1H, t, J=5.9Hz), 8.84 (1H, d, J=2.4Hz), 13.10 (1H, brs)

MS (m/z): 502 (M⁺)

[0561]

Example 275: N-(3-{6-Oxo-4-[4-(trifluoromethyl)thiazol-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 4-(trifluoromethyl)thiazole-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 4.41 (2H, d, J=6.8Hz), 7.01 (1H, brs), 7.5-7.7 (2H, m), 7.91 (1H, d, J=8.3Hz), 8.41 (1H, t, J=5.8Hz), 8.73 (1H, s), 13.36 (1H, brs)

MS (m/z): 490 (M⁺)

[0562]

Example 276: N-{4-Chloro-3-[4-(5-ethynylpyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-ethynylpyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 4.33 (2H, d, J=5.9Hz), 4.52 (1H, s), 7.22 (1H, brs), 7.4-7.6 (2H, m), 8.36 (1H, t, J=5.8Hz), 8.4-8.5 (1H, m), 8.78 (1H, d, J=2.0Hz), 9.22 (1H, d, J=2.0Hz), 13.23 (1H, brs)

MS (m/z): 424 (M⁺)

[0563]

Example 277: N-[3-(6-Oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 2.46 (1H, sept, J=6.8Hz), 3.95 (2H, t, J=4.7Hz), 4.14 (2H, q, J=9.3Hz), 4.40 (2H, d, J=5.9Hz), 4.4-4.6 (2H, m), 6.94 (1H, d, J=8.8Hz), 6.99 (1H, brs), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.3Hz), 8.33 (1H, dd, J=2.0Hz, 8.3Hz), 8.39 (1H, t, J=5.9Hz), 8.86 (1H, d, J=2.0Hz), 12.95 (1H, brs)

MS (m/z): 558 (M⁺)

[0564]

Example 278: N-[4-Chloro-2-fluoro-3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.3Hz), 2.45 (1H, sept, J=6.8Hz), 3.9-4.0 (2H, m), 4.14 (2H, q, J=9.7Hz), 4.32 (2H, d, J=5.8Hz), 4.4-4.6 (2H, m), 6.94 (1H, d, J=8.8Hz), 7.02 (1H, brs), 7.4-7.6 (2H, m), 8.2-8.4 (2H, m), 8.85 (1H, d, J=1.9Hz), 13.05 (1H, brs)
MS (m/z): 542 (M⁺)

[0565]

Example 279: N-[2,4-Difluoro-3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.45 (1H, sept, J=6.8Hz), 3.9-4.0 (2H, m), 4.14 (2H, q, J=9.8Hz), 4.32 (2H, d, J=5.9Hz), 4.4-4.6 (2H, m), 6.95 (1H, d, J=8.8Hz), 7.01 (1H, brs), 7.28 (1H, t, J=8.8Hz), 7.51 (1H, q, J=8.4Hz), 8.2-8.4 (2H, m), 8.86 (1H, d, J=2.0Hz), 13.12 (1H, brs)

MS (m/z): 526 (M⁺)

[0566]

Example 280: N-[2-Chloro-4-fluoro-3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)benzyl]isobutyramide

By performing operations similar to those of Example 4 using 6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.07 (6H, d, J=6.8Hz), 2.4-2.6 (1H, m), 3.9-4.0 (2H, m), 4.14 (2H, q, J=9.7Hz), 4.34 (2H, d, J=5.8Hz), 4.4-4.6 (2H, m), 6.93 (1H, d, J=8.8Hz), 7.01 (1H, brs), 7.3-7.6 (2H, m), 8.32 (1H, dd, J=2.4Hz, 8.8Hz), 8.37 (1H, t, J=5.9Hz), 8.84 (1H, d, J=1.9Hz), 13.10 (1H, brs)

MS (m/z): 542 (M⁺)

[0567]

Example 281: N-(3-{4-[5-(Cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)butyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-0.9 (2H, m), 0.86 (3H, t, J=7.2Hz), 0.9-1.0 (2H, m), 1.5-1.7 (3H, m), 2.15 (2H, t, J=7.2Hz), 4.41 (2H, d, J=5.6Hz), 7.21 (1H, s), 7.6-7.7 (2H, m), 7.8-8.0 (2H, m), 8.13 (1H, d, J=8.4Hz), 8.43 (1H, t, J=5.6Hz), 8.68 (1H, d, J=1.6Hz), 13.06 (1H, s)

MS (m/z): 480 (M⁺)

[0568]

Example 282: N-(4-Chloro-3-{4-[5-(cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)butyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-0.9 (2H, m), 0.87 (3H, t, J=7.2Hz), 0.9-1.0 (2H, m), 1.4-1.7 (3H, m), 2.14 (2H, t, J=6.8Hz), 4.33 (2H, d, J=5.6Hz), 7.24 (1H, s), 7.4-7.6 (2H, m), 7.89 (1H, dd, J=2.8Hz, 8.4Hz), 8.15 (1H, d, J=8.4Hz), 8.40 (1H, t, J=6.0Hz), 8.69 (1H, d, J=2.0Hz), 13.20 (1H, s)

MS (m/z): 464 (M⁺)

[0569]

Example 283: N-(3-{4-[5-(3,3-Dimethyl-1-butynyl)pyrazin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(3,3-dimethyl-1-butynyl)pyrazine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=6.8Hz), 1.14 (9H, s), 2.4-2.6 (1H, m), 4.41 (2H, d, J=5.6Hz), 7.17 (1H, s), 7.6-7.7 (2H, m), 7.91 (1H, d, J=8.0Hz), 8.40 (1H, t, J=5.6Hz), 8.81 (1H, s), 9.19 (1H, s), 13.18 (1H, s)

MS (m/z): 497 (M⁺)

[0570]

Example 284: N-(3-{4-[5-(Cyclopropylethynyl)pyrazin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyrazine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.6-0.7 (2H, m), 0.8-1.0 (2H, m), 1.05 (6H, d, J=6.8Hz), 1.6-1.8 (1H, m), 2.4-2.6 (1H, m), 4.41 (2H, d, J=6.0Hz), 7.15 (1H, s), 7.6-7.7 (2H, m), 7.90 (1H, d, J=8.0Hz), 8.40 (1H, t, J=6.0Hz), 8.73 (1H, s), 9.14 (1H, s), 13.17 (1H, s)

MS (m/z): 481 (M⁺)

[0571]

Example 285: N-(3-[2'-(4-Methyl-1-pentynyl)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(4-methyl-1-

pentynyl)pyrimidine-5-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 1.02 (6H, d, J=6.8Hz), 1.05 (6H, d, J=7.2Hz), 1.91 (1H, sept, J=6.8Hz), 2.4-2.6 (3H, m), 4.40 (2H, d, J=6.0Hz), 7.23 (1H, s), 7.6-7.7 (2H, m), 7.89 (1H, d, J=8.0Hz), 8.40 (1H, t, J=6.0Hz), 9.32 (2H, s), 13.19 (1H, s)
MS (m/z): 497 (M⁺)
[0572]

Example 286: N-{3-[6-Oxo-4-(5-propylpyridin-2-yl)-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-propylpyridine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.91 (3H, t, J=7.2Hz), 1.05 (6H, d, J=7.2Hz), 1.71 (2H, sext, J=7.2Hz), 2.46 (1H, sept, J=6.8Hz), 2.76 (2H, t, J=7.2Hz), 4.40 (2H, d, J=6.0Hz), 7.05 (1H, s), 7.35 (1H, d, J=8.4Hz), 7.5-7.7 (2H, m), 7.88 (1H, d, J=8.0Hz), 8.29 (1H, dd, J=2.0Hz, 8.0Hz), 8.39 (1H, t, J=6.0Hz), 9.12 (1H, d, J=1.2Hz), 12.98 (1H, s)
MS (m/z): 458 (M⁺)
[0573]

Example 287: N-{4-Chloro-2-fluoro-3-[6-oxo-4-(5-propylpyridin-2-yl)-1,6-dihydropyrimidin-2-yl]benzyl}isobutyramide

By performing operations similar to those of Example 4 using 5-propylpyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.91 (3H, t, J=7.2Hz), 1.05 (6H, d, J=6.8Hz), 1.71 (2H, m), 2.3-2.6 (1H, m), 2.76 (2H, t, J=7.6Hz), 4.32 (2H, d, J=4.8Hz), 7.09 (1H, s), 7.36 (1H, d, J=8.0Hz), 7.4-7.6 (2H, m), 8.2-8.5 (2H, m), 9.11 (1H, s), 12.99 (1H, s)
MS (m/z): 442 (M⁺)
[0574]

Example 288: N-{4-Chloro-3-[2'-(cyclopropylmethoxy)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-2-fluorobenzyl}isobutyramide

By performing operations similar to those of Example 4 using 2-(cyclopropylmethoxy)pyrimidine-5-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.7 (2H, m), 1.04 (6H, d, J=6.4Hz), 1.2-1.4 (1H, m), 2.3-2.6 (1H, m), 4.22 (2H, d, J=7.2Hz), 4.32 (2H, d, J=4.8Hz), 7.13 (1H, s), 7.4-7.6 (2H, m), 8.3-8.4 (1H, m), 9.1-9.2 (2H, m), 13.01 (1H, s)

MS (m/z): 471 (M⁺)

[0575]

Example 289: N-{3-[2'-(Cyclopropylmethoxy)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-4-(trifluoromethyl)benzyl}butyramide

By performing operations similar to those of Example 4 using 2-(cyclopropylmethoxy)pyrimidine-5-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 0.86 (3H, t, J=7.2Hz), 1.2-1.4 (1H, m), 1.55 (2H, sext, J=7.2Hz), 2.15 (2H, t, J=7.2Hz), 4.22 (2H, d, J=7.2Hz), 4.41 (2H, d, J=5.6Hz), 7.10 (1H, s), 7.6-7.7 (2H, m), 7.88 (1H, d, J=7.6Hz), 8.43 (1H, t, J=5.6Hz), 9.1-9.2 (2H, m), 12.74 (1H, s)

MS (m/z): 487 (M⁺)

[0576]

Example 290: N-{4-Chloro-3-[2'-(cyclopropylmethoxy)-6-oxo-1,6-dihydro-[4,5'-bipyrimidin]-2-yl]-2-fluorobenzyl}butyramide

By performing operations similar to those of Example 4 using 2-(cyclopropylmethoxy)pyrimidine-5-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)butyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 0.87 (3H, t, J=7.2Hz), 1.2-1.4 (1H, m), 1.55 (2H, sext, J=7.2Hz), 2.14 (2H, t, J=7.2Hz), 4.22 (2H, d, J=7.2Hz), 4.33 (2H, d, J=5.6Hz), 7.12 (1H, s), 7.4-7.6 (2H, m), 8.39 (1H, t, J=4.8Hz), 9.17 (2H, s), 12.43 (1H, s)

MS (m/z): 471 (M⁺)

[0577]

Example 291: N-(3-{4-[5-(Cyclopropylmethoxy)pyrazin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylmethoxy)pyrazine-2-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.3-0.4 (2H, m), 0.5-0.6 (2H, m), 1.05 (6H, d, J=6.8Hz), 1.2-1.3 (1H, m), 2.4-2.6 (1H, m), 4.21 (2H, d, J=7.6Hz), 4.40 (2H, d, J=5.6Hz), 7.06 (1H, s), 7.5-7.7 (2H, m), 7.90 (1H, d, J=8.0Hz), 8.3-8.5 (2H, m), 8.8-8.9 (1H, m), 13.05 (1H, s)

MS (m/z): 487 (M⁺)

[0578]

Example 292: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[2-(1-propynyl)thiazol-5-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(1-propynyl)thiazole-5-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.04 (6H, d, J=6.8Hz), 2.17 (3H, s), 2.45 (1H, sept, J=6.8Hz), 4.31 (2H, d, J=6.0Hz), 7.08 (1H, s), 7.4-7.5 (2H, m), 8.36 (1H, t, J=6.0Hz), 8.61 (1H, s), 13.23 (1H, brs)

MS (m/z): 444 (M⁺)

[0579]

Example 293: N-(2-Chloro-3-{4-[2-(cyclopropylethynyl)thiazol-5-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-fluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 2-(cyclopropylethynyl)thiazole-5-carbaldehyde, and N-(3-carbamimidoyl-2-chloro-4-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.1 (4H, m), 1.07 (6H, d, J=6.8Hz), 1.6-1.8 (1H, m), 2.4-2.6 (1H, m), 4.33 (2H, d, J=6.0Hz), 7.07 (1H, s), 7.4-7.6 (2H, m), 8.37 (1H, t, J=6.0Hz), 8.59 (1H, s), 13.22 (1H, brs)

MS (m/z): 470 (M⁺)

[0580]

Example 294: N-(3-{4-[5-(Cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(cyclopropylethynyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.04 (6H, d, J=6.8Hz), 1.5-1.7 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.6Hz), 7.23 (1H, s), 7.28 (1H, t, J=8.8Hz), 7.4-7.6 (1H, m), 7.91 (1H, dd, J=2.4Hz, 8.4Hz), 8.17 (1H, d, J=8.0Hz), 8.33 (1H, t, J=5.6Hz), 8.69 (1H, d, J=1.2Hz), 13.26 (1H, brs)

MS (m/z): 448 (M⁺)

[0581]

Example 295: N-(3-{4-[6-(Cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylethynyl)pyridine-3-carbaldehyde, and N-(3-carbamimidoyl-2,4-difluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.0 (4H, m), 1.04 (6H, d, J=6.8Hz), 1.5-1.7 (1H, m), 2.44 (1H, sept, J=6.8Hz), 4.32 (2H, d, J=5.6Hz), 7.14 (1H, s), 7.28 (1H, t, J=8.8Hz), 7.4-7.6 (2H,

m), 8.3-8.4 (2H, m), 9.14 (1H, d, J=2.0Hz), 13.22 (1H, brs)

MS (m/z): 448 (M⁺)

[0582]

Example 296: N-(2-Fluoro-3-{4-[5-(isobutyrylaminomethyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-(5-chloropyridin-2-yl)pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.03 (6H, d, J=6.8Hz), 1.06 (6H, d, J=6.8Hz), 2.3-2.6 (2H, m), 4.34 (2H, d, J=5.6Hz), 4.41 (2H, d, J=5.6Hz), 7.24 (1H, s), 7.66 (1H, t, J=8.0Hz), 7.7-7.9 (2H, m), 8.11 (1H, d, J=7.6Hz), 8.35 (1H, t, J=5.6Hz), 8.45 (1H, t, J=6.0Hz), 8.60 (1H, d, J=1.6Hz), 13.30 (1H, brs)

MS (m/z): 533 (M⁺)

[0583]

Example 297: N-(2-Fluoro-3-{6-oxo-4-[5-(1-propynyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[5-(1-propynyl)pyridin-2-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 2.12 (3H, s), 2.48 (1H, sept, J=6.8Hz), 4.41 (2H, d, J=5.6Hz), 7.24 (1H, s), 7.66 (1H, t, J=7.6Hz), 7.79 (1H, d, J=7.6Hz), 7.93 (1H, dd, J=2.0Hz, 8.0Hz), 8.12 (1H, d, J=8.0Hz), 8.44 (1H, t, J=5.6Hz), 8.72 (1H, d, J=2.0Hz), 13.34 (1H, brs)

MS (m/z): 472 (M⁺)

[0584]

Example 298: N-(3-{4-[6-(Cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(cyclopropylethynyl)pyridin-3-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.06 (6H, d, J=7.2Hz), 1.5-1.7 (1H, m), 2.48 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=6.0Hz), 7.15 (1H, s), 7.51 (1H, d, J=8.4Hz), 7.65 (1H, t, J=8.0Hz), 7.78 (1H, d, J=8.4Hz), 8.33 (1H, d, J=8.4Hz), 8.43 (1H, t, J=6.0Hz), 9.12 (1H, s), 13.29 (1H, brs)

MS (m/z): 498 (M⁺)

[0585]

Example 299: N-(3-{4-[6-(Cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 4 using 6-(cyclopropylethynyl)pyridine-3-carbaldehyde, and N-[3-carbamimidoyl-4-(trifluoromethyl)benzyl]isobutyramide hydrochloride, the title compound was obtained.
¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.05 (6H, d, J=7.2Hz), 1.5-1.7 (1H, m), 2.45 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=6.0Hz), 7.11 (1H, s), 7.5-7.7 (3H, m), 7.88 (1H, d, J=8.4Hz), 8.34 (1H, dd, J=2.0Hz, 8.4Hz), 8.39 (1H, t, J=6.0Hz), 9.14 (1H, s), 13.06 (1H, brs)

MS (m/z): 480 (M⁺)

[0586]

Example 300: N-(2-Fluoro-3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(2-propoxyethoxy)pyridin-3-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.85 (3H, t, J=7.6Hz), 1.06 (6H, d, J=6.8Hz), 1.51 (2H, sext, J=7.6Hz), 2.48 (1H, sept, J=6.8Hz), 3.39 (2H, t, J=6.8Hz), 3.71 (2H, t, J=4.8Hz), 4.40 (2H, d, J=6.0Hz), 4.43 (2H, t, J=4.4Hz), 6.92 (1H, d, J=8.8Hz), 7.02 (1H, s), 7.65 (1H, t, J=7.6Hz), 7.79 (1H, d, J=8.0Hz), 8.30 (1H, d, J=8.4Hz), 8.46 (1H, t, J=6.0Hz), 8.83 (1H, s), 13.20 (1H, brs)

MS (m/z): 536 (M⁺)

[0587]

Example 301: N-(2-Fluoro-3-{4-[6-(2-isopropoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[6-(2-isopropoxyethoxy)pyridin-3-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.0-1.2 (12H, m), 2.48 (1H, sept, J=6.8Hz), 3.61 (1H, sept, J=6.0Hz), 3.70 (2H, t, J=4.8Hz), 4.3-4.5 (4H, m), 6.92 (1H, d, J=8.8Hz), 7.01 (1H, s), 7.65 (1H, t, J=8.0Hz), 7.78 (1H, d, J=8.0Hz), 8.29 (1H, d, J=8.4Hz), 8.43 (1H, t, J=6.0Hz), 8.83 (1H, s), 13.17 (1H, brs)

MS (m/z): 536 (M⁺)

[0588]

Example 302: N-(3-{4-[2-(Cyclopropylethynyl)thiazol-5-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[2-(cyclopropylethynyl)thiazol-5-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.8-1.1 (4H, m), 1.06 (6H, d, J=6.8Hz), 1.6-1.8 (1H, m), 2.47 (1H, sept, J=6.8Hz), 4.40 (2H, d, J=5.6Hz), 7.08 (1H, s), 7.65 (1H, t, J=7.6Hz), 7.77 (1H, d, J=8.0Hz), 8.43 (1H, t, J=5.6Hz), 8.59 (1H, s), 13.29 (1H, brs)

MS (m/z): 504 (M⁺)

[0589]

Example 303: N-[2-Fluoro-3-(6-oxo-4-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}-1,6-dihydropyrimidin-2-yl)-4-(trifluoromethyl)benzyl]isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-{6-[2-(2,2,2-trifluoroethoxy)ethoxy]pyridin-3-yl}pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.06 (6H, d, J=6.8Hz), 2.48 (1H, sept, J=6.8Hz), 3.95 (2H, t, J=4.4Hz), 4.14 (2H, q, J=9.2Hz), 4.40 (2H, d, J=5.2Hz), 4.48 (2H, t, J=4.4Hz), 6.94 (1H, d, J=8.8Hz), 7.02 (1H, s), 7.65 (1H, t, J=7.6Hz), 7.78 (1H, d, J=8.0Hz), 8.31 (1H, d, J=7.6Hz), 8.43 (1H, t, J=5.6Hz), 8.83 (1H, s), 13.18 (1H, brs)

MS (m/z): 576 (M⁺)

[0590]

Example 304: N-(4-Chloro-2-fluoro-3-{6-oxo-4-[5-(2,2,2-trifluoroethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide

By performing operations similar to those of Example 4 using 5-(2,2,2-trifluoroethyl)pyridine-2-carbaldehyde, and N-(3-carbamimidoyl-4-chloro-2-fluorobenzyl)isobutyramide hydrochloride, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 1.05 (6H, d, J=7.2Hz), 2.46 (1H, sept, J=6.8Hz), 3.84 (2H, q, J=11.2Hz), 4.33 (2H, d, J=6.0Hz), 7.28 (1H, s), 7.4-7.6 (2H, m), 7.95 (1H, dd, J=2.0Hz, 8.0Hz), 8.23 (1H, d, J=7.6Hz), 8.37 (1H, t, J=6.0Hz), 8.72 (1H, d, J=1.6Hz), 13.28 (1H, brs)

MS (m/z): 482 (M⁺)

[0591]

Example 305: N-(3-{4-[5-(Cyclopropylethynyl)pyridin-2-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide

By performing operations similar to those of Example 169 using 2-[3-chloro-2-fluoro-6-(trifluoromethyl)phenyl]-6-[5-(cyclopropylethynyl)pyridin-2-yl]pyrimidin-4(3H)-one, the title compound was obtained.

¹H-NMR (d₆-DMSO, δ): 0.7-1.0 (4H, m), 1.06 (6H, d, J=6.8Hz), 1.5-1.7 (1H, m), 2.48 (1H,

sept, J=6.8Hz), 4.40 (2H, d, J=6.0Hz), 7.23 (1H, s), 7.66 (1H, t, J=8.0Hz), 7.79 (1H, d, J=8.4Hz), 7.89 (1H, dd, J=2.0Hz, 8.0Hz), 8.10 (1H, d, J=8.0Hz), 8.44 (1H, t, J=6.0Hz), 8.69 (1H, d, J=1.2Hz), 13.33 (1H, brs)

MS (m/z): 498 (M⁺)

[0592]

Test Example 1: Test for mPGES-1 inhibitory activity

Microsomes were prepared from COS-1 cells transiently transfected with a plasmid containing human mPGES-1 cDNA, and used as mPGES-1 enzyme. The mPGES-1 enzyme was diluted with a sodium phosphate buffer (pH 7.2) containing reduced glutathione (2.5 mM) and EDTA (1 mM), DMSO or a DMSO solution of a test compound (final concentration of DMSO was 1%) was added to the enzyme, and the mixture was preincubated at 4°C for 15 minutes. Then, PGH2 as the substrate was added at a final concentration of 1 μM to start the enzymatic reaction, and after incubation at 4°C for 4 minutes, a solution of ferric chloride (25 mM) and citric acid (50 mM) was added to terminate the enzymatic reaction. Generated PGE2 was measured by using Prostaglandin E2 Express EIA Kit (Cayman Chemical). IC₅₀ values were determined by using a standard method. The results are shown in Tables 2-1 to 2-5 mentioned below.

















[0593]

[Table 1-1]

Example No.	Structural formula	Example No.	Structural formula
1		10	
2		11	
3		12	
4		13	
5		14	
6		15	
7		16	
8		17	
9		18	



















[0594]

[Table 1-2]

Example No.	Structural formula	Example No.	Structural formula
19		28	
20		29	
21		30	
22		31	
23		32	
24		33	
25		34	
26			
27			

[0595]

[Table 1-3]

Example No.	Structural formula	Example No.	Structural formula
35		44	
36		45	
37		46	
38		47	
39		48	
40		49	
41		50	
42		51	
43		52	

[0596]

[Table 1-4]

Example No.	Structural formula	Example No.	Structural formula
53		62	
54		63	
55		64	
56		65	
57		66	
58		67	
59		68	
60		69	
61		70	



















[0597]

[Table 1-5]

Example No.	Structural formula	Example No.	Structural formula
71		80	
72		81	
73		82	
74		83	
75		84	
76		85	
77		86	
78		87	
79		88	



















[0598]

[Table 1-6]

Example No.	Structural formula	Example No.	Structural formula
89		98	
90		99	
91		100	
92		101	
93		102	
94		103	
95		104	
96		105	
97		106	

[0599]

[Table 1-7]

Example No.	Structural formula	Example No.	Structural formula
107		116	
108		117	
109		118	
110		119	
111		120	
112		121	
113		122	
114		123	
115		124	

[0600]

[Table 1-8]

Example No.	Structural formula	Example No.	Structural formula
125		134	
126		135	
127		136	
128		137	
129		138	
130		139	
131		140	
132		141	
133		142	



















[0601]

[Table 1-9]

Example No.	Structural formula	Example No.	Structural formula
143		152	
144		153	
145		154	
146		155	
147		156	
148		157	
149		158	
150		159	
151		160	

[0602]

[Table 1-10]

Example No.	Structural formula	Example No.	Structural formula
161		170	
162		171	
163		172	
164		173	
165		174	
166		175	
167		176	
168		177	
169		178	

[Table 1-11]

188

[0604]

[Table 1-12]

Example No.	Structural formula	Example No.	Structural formula
197		206	
198		207	
199		208	
200		209	
201		210	
202		211	
203		212	
204		213	
205		214	

[0605]

[Table 1-13]

Example No.	Structural formula	Example No.	Structural formula
215		224	
216		225	
217		226	
218		227	
219		228	
220		229	
221		230	
222		231	
223		232	

[0606]

[Table 1-14]

Example No.	Structural formula	Example No.	Structural formula
233		242	
234		243	
235		244	
236		245	
237		246	
238		247	
239		248	
240		249	
241		250	

[0607]

[Table 1-15]

Example No.	Structural formula	Example No.	Structural formula
251		260	
252		261	
253		262	
254		263	
255		264	
256		265	
257		266	
258		267	
259		268	

[0608]

[Table 1-16]

Example No.	Structural formula	Example No.	Structural formula
269		278	
270		279	
271		280	
272		281	
273		282	
274		283	
275		284	
276		285	
277		286	


[0609]

[Table 1-17]

Example No.	Structural formula	Example No.	Structural formula
287		296	
288		297	
289		298	
290		299	
291		300	
292		301	
293		302	
294		303	
295		304	

[0610]

[Table 1-18]

Example No.	Structural formula
305	

[0611]

[Table 2-1]

Example No.	human mPGES1 IC ₅₀ (nM)	Example No.	human mPGES1 IC ₅₀ (nM)
1	0.02	17	0.86
2	0.01	18	3.7
3	0.48	19	8.5
4	12	21	0.88
5	2.8	22	0.11
6	1.6	23	8.1
7	0.48	24	0.31
8	28	25	1.4
9	13	26	9
10	62	27	1.1
11	7.9	28	57
12	0.18	29	100
13	7.4	30	90
14	7.2	31	11
16	2.3	33	7.6

[0612]

[Table 2-2]

Example No.	human mPGES1 IC ₅₀ (nM)	Example No.	human mPGES1 IC ₅₀ (nM)
20	0.06	69	1.1
35	25	70	3.5
36	24	71	1.6
37	1	72	1.3
38	59	74	0.11
39	5.2	75	1.4
40	5.4	76	17
41	2.5	77	16
42	1.5	78	<1.0
43	0.13	79	19
44	19	80	3.9
45	0.76	81	41
46	8.8	82	7.3
47	7.4	83	25
48	2.3	84	18
49	2.7	85	13
50	7.3	86	3.3
51	5.4	87	29
52	7.1	88	10
53	2.7	89	2.3
54	3.2	90	4.1
55	1.3	91	6.4
56	19	92	26
57	11	93	16
58	33	94	9.1
59	1.4	95	7.8
60	2.7	96	1.5
61	41	97	38
62	27	98	13
63	1.0	99	25
64	1.6	100	28
65	1.1	101	12
66	1.8	102	4.1
67	47	103	37
68	25		

[0613]

[Table 2-3]

Example No.	human mPGES1 IC ₅₀ (nM)	Example No.	human mPGES1 IC ₅₀ (nM)
104	9.4	141	4.1
105	<1	142	110
106	3.7	143	8.7
107	76	144	7.3
108	2.8	145	3.9
109	6.2	147	18
110	3.4	148	7.3
111	1.4	149	7.3
112	55	150	12
113	23	151	130
114	13	152	27
115	23	153	30
116	4.5	154	6.4
117	15	155	52
118	10	156	75
119	6.4	157	0.78
120	15	158	4.7
122	7.5	159	2.1
123	5.2	160	2.5
124	15	161	23
125	100	162	<1
126	4.5	163	6.9
127	<1	164	2.4
128	2.1	165	18
129	31	166	6.4
130	47	167	3.1
131	7.0	168	7.7
132	18	169	1.1
133	87	170	1.7
134	89	171	2.0
135	8.8	172	6.1
136	66	173	38
137	92	174	6.0
138	15	175	18
139	23	176	1.3
140	3.3	177	33

[0614]

[Table 2-4]

Example No.	human mPGES1 IC ₅₀ (nM)	Example No.	human mPGES1 IC ₅₀ (nM)
178	<1	214	4.1
179	32	215	<1
180	67	216	22
181	69	217	37
182	1.2	218	6.3
183	6.2	219	7.1
184	20	220	1.1
185	52	221	<1
186	32	222	30
187	5.9	223	<1
188	1.9	224	8.0
189	4.3	225	1.5
190	32	226	19
191	<1	227	1.7
192	<1	228	3.1
193	<1	229	5.8
194	3.1	230	29
195	2.1	231	25
196	8.3	232	1.6
197	4.2	233	<1
198	3.6	234	12
199	6.5	235	1.0
200	4.6	236	2.2
201	3.2	237	<1
202	<1	238	<1
203	<1	239	1.5
204	3.7	240	<1
205	9.9	241	15
206	8.9	242	45
207	1.9	243	2.9
208	4.8	244	3.7
209	7.5	245	68
210	1.4	246	1.4
211	31	247	2.0
212	37	248	2.3
213	4.7	249	5.2

[0615]

[Table 2-5]


Example No.	human mPGES1 IC ₅₀ (nM)	Example No.	human mPGES1 IC ₅₀ (nM)
250	5.8	286	3.4
251	39	287	1.2
252	26	288	<1
253	3.6	289	2.5
254	3.9	290	<1
255	<1	291	4.8
256	1.0	292	<1
257	11	293	2.4
258	15	294	<1
259	4.7	295	<1
260	4.5	296	62
261	3.4	297	2.8
262	2.2	298	2.6
263	<1	299	3.7
264	1.2	300	<1
265	1.5	301	1.2
266	<1	302	<1
267	6.2	303	<1
268	13	304	4.7
269	23	305	5.0
270	27		
271	23		
272	2.9		
273	2.4		
274	36		
275	1.9		
276	3.2		
277	2.0		
278	<1		
279	1.6		
280	8.7		
281	<1		
282	<1		
283	1.9		
284	3.1		
285	<1		

What is claimed is:

[Claim 1]

A compound represented by the following general formula (1):

[Formula 1]



(in the formula, X represents carbonyl group, or sulfonyl group; R¹ represents hydrogen atom, a halogen atom, an alkyl group, an alkanoyl group, cyano group, or carboxyl group; R² represents an alkyl group, a cyclic carbon group which may have a substituent, or a heterocyclic group which may have a substituent; R³ represents hydrogen atom, or 1 to 3 substituents substituting on the benzene ring (these substituents are selected from the group consisting of a halogen atom, an alkyl group (this alkyl group may be substituted with a halogen atom), and an alkoxy group (this alkoxy group may be substituted with a halogen atom)); R⁴ and R⁵ independently represent hydrogen atom, a halogen atom, or an alkyl group; and R⁶ represents an alkyl group (this alkyl group may be substituted with hydroxy group, a halogen atom, or an alkoxy group), or an alkoxy group), or a salt thereof.

[Claim 2]

The compound represented by the general formula (1), or a salt thereof according to claim 1, wherein X is carbonyl group.

[Claim 3]

The compound represented by the general formula (1), or a salt thereof according to claim 1 or 2, wherein R⁶ is a branched C₁₋₆ alkyl group (this alkyl group may be substituted with a C₁₋₆ alkoxy group).

[Claim 4]

The compound represented by the general formula (1), or a salt thereof according to any one of claims 1 to 3, wherein both R⁴ and R⁵ are hydrogen atoms.

[Claim 5]

The compound represented by the general formula (1), or a salt thereof according to any one of claims 1 to 4, wherein R³ consists of one or two halogen atoms,

or alkyl groups which may be substituted with a halogen atom.

[Claim 6]

The compound represented by the general formula (1), or a salt thereof according to any one of claims 1 to 5, wherein R¹ is hydrogen atom, an alkyl group, or cyano group.

[Claim 7]

The compound represented by the general formula (1), or a salt thereof according to any one of claims 1 to 6, wherein R² is a saturated or partially saturated 3- to 7-membered monocyclic cyclic hydrocarbon group which may have a substituent, a phenyl group which may have a substituent, a saturated or partially saturated 3- to 7-membered monocyclic heterocyclic group which may have a substituent (this heterocyclic group contains 1 to 3 ring-constituting heteroatoms), or a monocyclic aromatic heterocyclic group which may have a substituent (this heterocyclic group contains 1 to 3 ring-constituting heteroatoms).

[Claim 8]

The compound represented by the general formula (1), or a salt thereof according to any one of claims 1 to 7, wherein R² is a phenyl group which may have a substituent, or a pyridyl group which may have a substituent.

[Claim 9]

The compound represented by the general formula (1), or a salt thereof according to claim 1, which is:

N-(4-chloro-3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide,

N-(3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide,

N-(3-{6-oxo-4-[6-(2-propoxyethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide,

N-(4-chloro-3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide,

N-(4-chloro-2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide,

N-(3-{4-[6-(cyclopropylmethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2,4-difluorobenzyl)isobutyramide,

N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide,

N-{4-chloro-3-[4-(5-chloropyridin-2-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-2-

fluorobenzyl}isobutyramide,
N-(4-chloro-2-fluoro-3-{6-oxo-4-[6-(3,3,3-trifluoropropoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide,
N-(2-chloro-4-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}benzyl)isobutyramide,
N-(3-{4-[6-(difluoromethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide,
N-(3-{4-[6-(2-ethoxyethoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluoro-4-(trifluoromethyl)benzyl)isobutyramide,
N-(2-fluoro-3-{6-oxo-4-[5-(trifluoromethyl)pyridin-2-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide,
N-(3-{6-oxo-4-[6-(tetrahydropyran-4-ylmethoxy)pyridin-3-yl]-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide,
N-(3-{4-[6-(3-ethoxypropoxy)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-4-(trifluoromethyl)benzyl)isobutyramide,
N-{2-fluoro-3-[4-(6-methoxypyridin-3-yl)-6-oxo-1,6-dihydropyrimidin-2-yl]-4-(trifluoromethyl)benzyl}isobutyramide, or
N-(4-chloro-3-{4-[6-(cyclopropylethynyl)pyridin-3-yl]-6-oxo-1,6-dihydropyrimidin-2-yl}-2-fluorobenzyl)isobutyramide.

[Claim 10]

An mPGES-1 inhibitor containing the compound represented by the aforementioned general formula (1), or a salt thereof according to any one of claims 1 to 9.

[Claim 11]

A PGE2 biosynthesis inhibitor containing the compound represented by the aforementioned general formula (1), or a salt thereof according to any one of claims 1 to 9.

[Claim 12]

A medicament containing the compound represented by the aforementioned general formula (1), or a physiologically acceptable salt thereof according to any one of claims 1 to 9.

[Claim 13]

The medicament according to claim 12, which is for use in prophylactic and/or therapeutic treatment of inflammation, pain, rheumatism, osteoarthritis, pyrexia, Alzheimer's disease, multiple sclerosis, arteriosclerosis, ocular hypertension, ischemic retinopathy, systemic scleroderma, malignant tumor, overactive bladder, bladder outlet

obstruction associated with benign prostatic hyperplasia, nocturia, urinary incontinence, neurogenic bladder, interstitial cystitis, bladder pain syndrome, urinary calculus, benign prostatic hyperplasia, or a disease for which suppression of the PGE₂ production exhibits efficacy.