

Process for manufacturing vaccines

The present invention relates to improved methods of conducting carbodiimide condensation reactions. In particular, it relates to the conjugation of saccharides and proteins using carbodiimide condensation. It also relates to immunogenic compositions that may be made comprising the saccharide-protein conjugates of the invention.

The use of bacterial capsular polysaccharides has been widely used in immunology for many years for the prevention of bacterial disease. A problem with such a use, however, is the T-independent nature of the immune response. These antigens are thus poorly immunogenic in young children. This problem has been overcome through conjugating the polysaccharide antigens to a protein carrier (a source of T-helper epitopes) which may then be used to elicit a T-dependent immune response, even in the first year of life.

Various conjugation techniques are known in the art. Conjugates can be prepared by direct reductive amination methods as described in US 4365170 (Jennings) and US 4673574 (Anderson). Other methods are described in EP-0-161-188, EP-208375 and EP-0-477508. The conjugation method may alternatively rely on activation of hydroxyl groups of the saccharide with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated saccharide may thus be coupled directly or via a spacer (linker) group to an amino group on the carrier protein. For example, the cyanate ester can be coupled with hexane diamine or adipic acid dihydrazide (ADH or AH) and the amino-derivatised saccharide is conjugated to the carrier protein using carbodiimide (e.g. EDAC or EDC) chemistry via a carboxyl group on the protein carrier. Such conjugates are described in PCT published application WO 93/15760 Uniformed Services University and WO 95/08348 and WO 96/29094. See also Chu C. et al Infect. Immunity, 1983 245 256.

In general the following types of chemical groups on a protein carrier can be used for coupling / conjugation:

- A) Carboxyl (for instance via aspartic acid or glutamic acid) which may be conjugated to natural or derivatised amino groups on saccharide moieties using carbodiimide chemistry;
- B) Amino group (for instance via lysine) which may be conjugated to natural or derivatised carboxyl groups on saccharide moieties using carbodiimide chemistry;
- C) Sulphydryl (for instance via cysteine);
- D) Hydroxyl group (for instance via tyrosine);

- E) Imidazolyl group (for instance via histidine);
- F) Guanidyl group (for instance via arginine); and
- G) Indolyl group (for instance via tryptophan).

- 5 On a saccharide, in general the following groups can be used for a coupling: OH, COOH or NH₂. Aldehyde groups can be generated after different treatments known in the art such as: periodate, acid hydrolysis, hydrogen peroxide, etc.

Direct coupling approaches:

10

Saccharide-OH + CNBr or CDAP -----> cyanate ester + NH₂-Prot -----> conjugate
 Saccharide-aldehyde + NH₂-Prot -----> Schiff base + NaCNBH₃ -----> conjugate
 Saccharide-COOH + NH₂-Prot + EDAC -----> conjugate
 Saccharide-NH₂ + COOH-Prot + EDAC -----> conjugate

15

Indirect coupling via spacer (linker) approaches:

Saccharide-OH + CNBr or CDAP ---> cyanate ester + NH₂----NH₂ -----> saccharide----NH₂ + COOH-Prot + EDAC -----> conjugate

20

Saccharide-OH + CNBr or CDAP -----> cyanate ester + NH₂-----SH -----> saccharide----SH + SH-Prot (native Protein with an exposed cysteine or obtained after modification of amino groups of the protein by SPDP for instance) -----> saccharide-S-S-Prot

25

Saccharide-OH + CNBr or CDAP ---> cyanate ester + NH₂----SH -----> saccharide----SH + maleimide-Prot (modification of amino groups) -----> conjugate

Saccharide-COOH + EDAC + NH₂-----NH₂ ---> saccharide-----NH₂ + EDAC + COOH-Prot -----> conjugate

30

Saccharide-COOH + EDAC+ NH₂----SH -----> saccharide----SH + SH-Prot (native Protein with an exposed cysteine or obtained after modification of amino groups of the protein by SPDP for instance) -----> saccharide-S-S-Prot

35

Saccharide-COOH + EDAC+ NH₂----SH -----> saccharide----SH + maleimide-Prot (modification of amino groups) -----> conjugate

Saccharide-Aldehyde + NH₂-----NH₂ ----> saccharide---NH₂ + EDAC + COOH-Prot ----> conjugate

- 5 As can be observed carbodiimide chemistry (e.g. using EDAC) is very convenient for conjugation reactions as it makes use of groups on the saccharide and/or protein which may be naturally present or easily inserted by derivatisation. It also conveniently links moieties through a peptide bond.
- 10 Carbodiimides (RN=C=NR') are unsaturated compounds with an allene structure (Nakajima and Ikada 1995 Bioconjugate Chem. 6 :123-130; Hoare and Koshland 1967 JBC 242:2447-2453). The chemical is relatively unstable at its reaction pH (4.5-6.5), and therefore all components of the saccharide/protein/carbodiimide conjugation reaction tend to be added together in the art.

15

- The present inventors have found that depending on the nature of the saccharide and protein to be conjugated, better characteristics of the final conjugate for vaccine use may be achieved by adding a certain component of the reaction slowly to the mix. In so doing one or more benefits/improvements may be realised such as: saccharide yield in the
- 20 conjugate, sterile filterability of the conjugate, better control of the conjugation, easier reproducibility, and/or prevention of intra-moiety cross-links.

Accordingly, in one embodiment there is provided a method of conjugating a saccharide to a protein carrier using carbodiimide condensation chemistry, wherein the saccharide

25 comprises (for instance as part of its repeating unit), or has been derivatised to comprise, amino and/or carboxyl groups, and wherein the protein carrier comprises, or has been derivatised to comprise, amino and/or carboxyl groups, comprising the steps of:

- I) - if the protein carrier comprises both amino and carboxyl groups and the saccharide comprises either amino or carboxyl groups:
- 30 a) mixing the saccharide and aliquot of carbodiimide required to perform the conjugation, and
- b) adding the aliquot of protein carrier required over a period of 35 seconds to 6 hours;
- II) - if the saccharide comprises both amino and carboxyl groups and the protein carrier comprises either amino or carboxyl groups:
- 35 a) mixing the protein carrier and aliquot of carbodiimide required to perform the conjugation, and

b) adding the aliquot of saccharide required over a period of 35 seconds to 6 hours;

III) - if the saccharide comprises both amino and carboxyl groups and the protein carrier comprises both amino and carboxyl groups:

a) mixing the protein carrier and saccharide, and

b) adding the aliquot of carbodiimide required to perform the conjugation over a period of 35 seconds to 6 hours.

10 Detailed description

Any suitable carbodiimide may be used as long as it is capable of conjugating saccharides and proteins in an aqueous medium. In one embodiment the carbodiimide may be EDAC (1-ethyl-3-(3-dimethyl-aminopropyl) carbodiimide) [also known as EDC] or it may be a carbodiimide other than EDAC.

The term "saccharide" throughout this specification may indicate polysaccharide or oligosaccharide and includes both. It may indicate lipopolysaccharide (LPS) or lipooligosaccharide (LOS). Before use Polysaccharides (such as bacterial polysaccharides) may be isolated from a source strain (e.g. of bacteria) or isolated from the source strain and sized to some degree by known methods (see for example EP497524 and EP497525; Shousun Chen Szu *et al.* – Carbohydrate Research Vol 152 p7-20 (1986)) for instance by microfluidisation. Polysaccharides can be sized in order to reduce viscosity in polysaccharide samples and/or to improve filterability for conjugated products. Oligosaccharides have a low number of repeat units (typically 5-30 repeat units) and are typically hydrolysed polysaccharides.

The term "protein carrier" is intended to cover both small peptides and large polypeptides (>10 kDa). Clearly large polypeptides are more likely to contain both reactive amino and carboxyl groups without any modification.

For the purposes of the invention, "native polysaccharide" refers to a saccharide that has not been subjected to a process, the purpose of which is to reduce the size of the saccharide. A polysaccharide can become slightly reduced in size during normal purification procedures. Such a saccharide is still native. Only if the polysaccharide has been subjected to sizing techniques would the polysaccharide not be considered native.

For the purposes of the invention, "sized by a factor up to x^2 " means that the saccharide is subject to a process intended to reduce the size of the saccharide but to retain a size more than half the size of the native polysaccharide. X^3 , x^4 etc. are to be interpreted in the same way i.e. the saccharide is subject to a process intended to reduce the size of the polysaccharide but to retain a size more than a third, a quarter etc. the size of the native polysaccharide.

The 35 second to 6 hour time period in step b) of the method for the addition of the full aliquot of the final component can be 50 seconds to 5 hours, 1 minute to 4 hours, 2 minutes to 3 hours, 3 minutes to 2 hours, 4 to 60 minutes, 5 to 50 minutes, 6 to 40 minutes, 7 to 30 minutes or 8 to 20 minutes. It may be 1 minute to 5 hours, 10 minutes to 4 hours, 20 minutes to 3 hours, 30 minutes to 2 hours, 40 to 90 minutes, or 50 to 70 minutes. This time can be adjusted according to the precise saccharide and protein being conjugated.

In one embodiment the aliquot of the final component (e.g. of carbodiimide, saccharide or protein) is added to the reaction mixture at a constant rate during the time period (this is conveniently achieved using a pump operating at a constant rate). Alternatively it may be added in stages over the time period. Although this may be done in many ways, in general parts of the aliquot should be added throughout the period. For instance at least one quarter of the aliquot may be added over the first half of the period, and at least one quarter of the aliquot over the second half of the period. The total amount of the aliquot 'a' measured, for instance, in mL or mg may be added in 4-100 stages ('s') throughout the period. In one embodiment the stages are arranged such that an even amount (a/s) is introduced at all the stages. In one embodiment the stages are evenly spaced throughout the period 'p' (in seconds). Thus if one stage takes place at time zero of the period 'p', then each subsequent stage could take place at a time which is $p/(s-1)$. The volume of the aliquot of the final component added in step b) may be adjusted in terms of ease of addition of the aliquot to the reaction within the desired time period. The carbodiimide may be added as an aqueous solution (typically buffered at pH 7.5 before being added to the reaction) or as solid powder (EDAC for instance is highly soluble in aqueous media). Of course if the carbodiimide is the last component added to the reaction (situation III step b)), a slow dissolving carbodiimide may be used such that the entire aliquot of powder is added to the reaction all at once but it dissolves at a rate consistent with the desired period over which the aliquot is to be made available to the reaction.

If the protein and/or saccharide has no amino or carboxyl groups (or only has one of these), it may be derivatised to give it one (or to give it the other it does not already have). For instance for a saccharide only comprising reactive hydroxyl groups (e.g. meningococcal serogroup A capsular saccharide), such a group should be used for derivatising on amino or carboxyl groups so that EDAC condensation may be carried out.

This may take place within a repeat subunit, or may be a group only present at the end of the saccharide molecule.

It should be noted that where derivatisation takes place, it can be beneficial to only partially derivatise the moiety. For saccharides with repeating subunits, the target epitope may be present in each repeat. Therefore if partial derivatisation takes place (for this it is meant 0.5-20, 1-15, 3-12, or 5-10% of the targeted reactive group is actually derivatised) this can have the benefit of conserving the majority of the epitopes, and preventing too much cross-linking.

If a saccharide or protein already has amino or carboxyl groups only (e.g. Vi saccharide from *Salmonella typhi* which naturally has carboxyl but not amino groups), derivatisation can take place to give it the other type of group (i.e. amino groups for Vi). It should be noted, however, that as derivatisation can be partial this action can change the preferred reaction of the invention from a type I to a type III. For instance if Vi saccharide is conjugated to a protein carrier comprising both amino and carboxyl groups situation I adds the aliquot of protein slowly in step b). If the Vi saccharide carboxyl group is partially derivatised with amino groups it will have both carboxyl and amino groups, thus situation III adding the aliquot of carbodiimide slowly in step b) becomes most relevant.

Derivatisation may occur through the addition of a hetero- or homo-bifunctional linker. It may take place with similar chemistry as described above for saccharide-protein conjugation step (e.g. CDAP or carbodiimide chemistry). The linker may have between 4 and 20, 4 and 12, or 5 and 10 carbon atoms. It may have two reactive amino groups, two reactive carboxyl groups, or one of each (e.g. hexane diamine, 6-aminocaproic acid, or adipic acid dihydrazide). Typically derivatization takes place through reacting a large excess of the linker with the saccharide and/or protein carrier to be derivatised. This allows derivatization to take place with minimal intra-moiety cross-linking (which otherwise might be possible if for instance a carboxyl group on a saccharide was being derivatised with amino groups using carbodiimide condensation). Excess linker is readily removed using techniques such as diafiltration.

In one embodiment the saccharide comprises a reactive hydroxyl group as part of its repeating unit which is partially derivatised via an amino group on the linker (e.g. with CDAP chemistry). In another embodiment the saccharide comprises a reactive amino group as part of its repeating unit which is partially derivatised via a carboxyl group on the linker (e.g. with carbodiimide chemistry). In a further embodiment the saccharide comprises a reactive carboxyl group as part of its repeating unit which is partially derivatised via an amino group on the linker (e.g. with carbodiimide chemistry).

The aliquot of carbodiimide required to perform the conjugation (whether present in step a) or b) of the reaction of the invention) is 0.01 to 3, 0.05 to 2, or 0.09 to 1 mg

carbodiimide/mg saccharide. Although these numbers are calculated in respect of EDAC being the carbodiimide, these numbers may be adjusted if any other carbodiimide is used by multiplying the numbers in the range by: (molecular weight of other carbodiimide)/(molecular weight of EDAC).

5

In general, the saccharide may be present in the methods of the invention at a final concentration of 0.5-50 mg/ml in step b). This will depend on the size and nature of the saccharide, and the extent of any derivatisation. For instance for oligosaccharides a larger concentration will be required, but for large polysaccharides a much smaller concentration will be more appropriate. If it is towards the high end of partially derivatised with amino or carboxyl groups a smaller concentration may be appropriate to reduce the possibility of any cross-linking. The protein carrier may be present at a final concentration of 1-50 mg/ml in step b).

10

15 The initial ratio of protein carrier to saccharide in the methods of the invention can be 5:1 to 1:5, 4:1 to 1:1, or 3:1 to 2:1 (w/w). Again this will depend on the size and nature of the saccharide, and the extent of any derivatisation.

20 Salt conditions (e.g. NaCl) may also be varied according to the nature of the saccharide/protein. Usually around 0.2M NaCl may be present in step b) of the methods of the invention, but may be 0-2, 0.1-1 or 0.2-0.5 M.

25 In terms of pH in step b) of the methods of the invention, the reaction pH may be any pH where the carbodiimide is activated – for instance pH 4.5-6.5, 4.7-6.0, or 5-5.5. This pH is typically maintained throughout the reaction by addition of acid/base as required. EDAC is usually stable at pH 7.5, though if the conjugation requires to be done at higher pH compounds which are known to keep the reaction intermediate stable (such as N-hydroxysuccinimide) may also be present in the reaction in step b), in which case the reaction pH in step b) may be maintained at pH 4.5-7.5.

30

The reaction temperature during step b) of the methods of the invention can be 4-37, 10-32, 17-30, or 22-27 °C, and is typically maintained throughout the reaction.

35 In the methods of the invention, once the entire aliquot has been added in step b) the reaction is typically maintained for a further 10 minutes to 72 hours, 20 minutes to 48 hours, 30 minutes to 24 hours, 40 minutes to 12 hours, 50 minutes to 6 hours, or 1-3 hours. Once the reaction is completed the pH is adjusted to 7.5-9 (towards the higher end of this if N-hydroxysuccinimide is present) to go back to the stable pH range of carbodiimide.

40

Once conjugated, the saccharide-protein conjugate may be purified from: unreacted components, free saccharide, etc by injecting it on a size exclusion chromatography

column (for instance Sephacryl S400HR, Pharmacia). This is typically carried out at 2-8 °C. The conjugate may be sterile filtered then stored. Ultimately an effective dose (for instance 1-20, 2-15, or 3-10 µg saccharide /dose) of the saccharide-protein conjugate can be formulated with a pharmaceutically acceptable excipient (for instance a salt or adjuvant) to manufacture an immunogenic composition or vaccine.

In terms of the saccharides of the invention, any saccharide of viral, fungal, bacterial or eukaryotic source may be conjugated using the methods of the invention. It may be the Vi saccharide from *Salmonella typhi*, or a saccharide other than Vi. It may be the capsular saccharide Hib from *H. influenzae* type b, or may be a saccharide other than Hib. In one embodiment the saccharide is a bacterial capsular saccharide, for instance derived from a bacterium selected from a list consisting of: *N. meningitidis* serogroup A (MenA), B (MenB), C (MenC), W135 (MenW) or Y (MenY), *Streptococcus pneumoniae* serotypes 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F or 33F, Group B Streptococcus group Ia, Ib, II, III, IV, V, VI, or VII, *Staphylococcus aureus* type 5, *Staphylococcus aureus* type 8, *Salmonella typhi* (Vi saccharide), *Vibrio cholerae*, or *H. influenzae* type b.

The weight-average molecular weight of the saccharide may be 1000-2000000, 5000-1000000, 10000-500000, 50000-400000, 75000-300000, or 100000-200000. The molecular weight or average molecular weight of a saccharide herein refers to the weight-average molecular weight (Mw) of the saccharide measured prior to conjugation and is measured by MALLS. The MALLS technique is well known in the art and is typically carried out as described in example 2. For MALLS analysis of saccharides, two columns (TSKG6000 and 5000PWxl) may be used in combination and the saccharides are eluted in water. Saccharides are detected using a light scattering detector (for instance Wyatt Dawn DSP equipped with a 10mW argon laser at 488nm) and an interferometric refractometer (for instance Wyatt Otilab DSP equipped with a P100 cell and a red filter at 498nm). In an embodiment, the polydispersity of the saccharide is 1-1.5, 1-1.3, 1-1.2, 1-1.1 or 1-1.05 and after conjugation to a carrier protein, the polydispersity of the conjugate is 1.0-2.5, 1.0-2.0, 1.0-1.5, 1.0-1.2, 1.5-2.5, 1.7-2.2 or 1.5-2.0. All polydispersity measurements are by MALLS.

The saccharide may be either a native polysaccharide or may have been sized by a factor of no more than 2, 4, 6, 8, 10 or 20 fold (for instance by microfluidization [e.g. by Emulsiflex C-50 apparatus] or other known technique [for instance heat, chemical, oxidation, sonication methods]). Oligosaccharides may have been sized substantially further [for instance by known heat, chemical, or oxidation methods].

The structures of most of these saccharides are known (and therefore whether they naturally have any amino or carboxyl groups for carbodiimide chemistry, or any other reactive group which may be derivatised with amino or carboxyl groups (see table below).

	Natural NH ₂ group	Natural COOH group	Other reactive group
<i>S. aureus</i>			
PS5	No	Yes	OH
PS8	No	Yes	OH
<i>N. meningitidis</i>			
MenA	No	No	OH
MenC	No	Yes	OH
MenW135	No	Yes	OH
MenY	No	Yes	OH
MenB	No (can be generated if de-N-acetylated)	Yes	OH / N-propyl
Gp. B Streptococcus			
Ia, Ib	No	Yes	OH
II	No	Yes	OH
III	No	Yes	OH
IV	No	Yes	OH
V	No	Yes	OH
VI	No	Yes	OH
VII	No	Yes	OH
<i>S. typhi</i>			
Vi	No	Yes	No
<i>S. pneumoniae</i>			
PS1	Yes	Yes	OH
PS3, 4, 5, 8, 9, 12F	No	Yes	OH
<i>Vibrio cholerea</i>			
Capsular saccharide	yes	No	OH
<i>H.influenzae B Hib</i>	No	No	OH
LOS			
Nmen/ Mcat/ Hi	Yes on PEA	Yes on KDO	OH

- 5 The saccharide may be a bacterial lipooligosaccharide or lipopolysaccharide (see above table), for instance derived from a bacterium selected from a list consisting of: *N. meningitidis*, *H. influenzae*, *E. coli*, *Salmonella* or *M. catarrhalis*. The LOS may be meningococcal immunotype L2, L3 or L10. It may be detoxified by alkaline treatment of its Lipid A moiety.

- 10 In an embodiment, the MenA capsular saccharide, is at least partially O-acetylated such that at least 50%, 60%, 70%, 80%, 90%, 95% or 98% of the repeat units are O-acetylated at at least one position. O-acetylation is for example present at least at the O-3 position of at least 50%, 60%, 70%, 80%, 90%, 95% or 98% of the repeat units. In an embodiment, the MenC capsular saccharide, is at least partially O-acetylated such that at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 98% of ($\alpha 2 \rightarrow 9$)-linked NeuNAc repeat units

are O-acetylated at at least one or two positions. O-acetylation is for example present at the O-7 and/or O-8 position of at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 98% of the repeat units. In an embodiment, the MenW capsular saccharide, is at least partially O-acetylated such that at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or

98% of the repeat units are O-acetylated at at least one or two positions. O-acetylation is for example present at the O-7 and/or O-9 position of at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 98% of the repeat units. In an embodiment, the MenY capsular saccharide, is at least partially O-acetylated such that at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 98% of the repeat units are O-acetylated at at least one or two positions. O-acetylation is present at the 7 and/or 9 position of at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 98% of the repeat units. The percentage of O-acetylation refers to the percentage of the repeat units containing O-acetylation. This may be measured in the saccharide prior to conjugate and/or after conjugation.

The protein carrier may be any peptide or protein. It may comprise one or more T-helper epitopes. In one embodiment of the invention the protein carrier is selected from the group consisting of: TT, DT, CRM197, fragment C of TT, protein D of *H. influenzae*, pneumococcal PhtD, and pneumococcal Pneumolysin. The carrier protein may be tetanus toxoid (TT), tetanus toxoid fragment C, non-toxic mutants of tetanus toxin [note all such variants of TT are considered to be the same type of carrier protein for the purposes of this invention], diphtheria toxoid (DT), CRM197, other non-toxic mutants of diphtheria toxin [such as CRM176, CRM 197, CRM228, CRM 45 (Uchida et al *J. Biol. Chem.* 218; 3838-3844, 1973); CRM 9, CRM 45, CRM102, CRM 103 and CRM107 and other mutations described by Nicholls and Youle in *Genetically Engineered Toxins*, Ed: Frankel, Maecel Dekker Inc, 1992; deletion or mutation of Glu-148 to Asp, Gln or Ser and/or Ala 158 to Gly and other mutations disclosed in US 4709017 or US 4950740; mutation of at least one or more residues Lys 516, Lys 526, Phe 530 and/or Lys 534 and other mutations disclosed in US 5917017 or US 6455673; or fragment disclosed in US 5843711] (note all such variants of DT are considered to be the same type of carrier protein for the purposes of this invention), pneumococcal pneumolysin (Kuo et al (1995) *Infect Immun* 63; 2706-13), OMPC (meningococcal outer membrane protein – usually extracted from *N. meningitidis* serogroup B – EP0372501), synthetic peptides (EP0378881, EP0427347), heat shock proteins (WO 93/17712, WO 94/03208), pertussis proteins (WO 98/58668, EP0471177), cytokines, lymphokines, growth factors or hormones (WO 91/01146), artificial proteins comprising multiple human CD4+ T cell epitopes from various pathogen derived antigens (Falugi et al (2001) *Eur J Immunol* 31; 3816-3824) such as N19 protein (Baraldoi et al (2004) *Infect Immun* 72; 4884-7) pneumococcal surface protein PspA (WO 02/091998), iron uptake proteins (WO 01/72337), toxin A or B of *C. difficile* (WO 00/61761), *H. influenzae* Protein D (EP594610 and WO 00/56360), pneumococcal PhtA (WO 98/18930, also referred to Sp36),

pneumococcal PhtD (disclosed in WO 00/37105, and is also referred to Sp036D), pneumococcal PhtB (disclosed in WO 00/37105, and is also referred to Sp036B), or PhtE (disclosed in WO00/30299 and is referred to as BVH-3).

- 5 In a further aspect of the invention there is provided a saccharide-protein carrier conjugate (or an immunogenic composition or vaccine) obtainable or obtained by the method of the invention.

- 10 A use of the immunogenic composition or vaccine of the invention in the manufacture of a medicament for the prevention or treatment of disease, and a method of preventing or treating disease comprising the step of administering an effective dose of the immunogenic composition or vaccine of the invention to a patient in need thereof is further provided. The use or method may be such that the disease is caused by a bacterium selected from a list consisting of: *N. meningitidis*, *Streptococcus pneumoniae*, *M.*
15 *catarrhalis*, Group B Streptococcus, *Staphylococcus aureus*, *Salmonella typhi*, *Vibrio cholerae*, *E. coli*, and *H. influenzae*.

- The immunogenic compositions of the invention may also comprise a DTPa or DTPw vaccine (for instance one containing DT, TT, and either a whole cell pertussis (Pw)
20 vaccine or an acellular pertussis (Pa) vaccine (comprising for instance pertussis toxoid, FHA, pertactin, and, optionally agglutinogens 2 and 3). Such combinations may also comprise a vaccine against hepatitis B (for instance it may comprise hepatitis B surface antigen [HepB], optionally adsorbed onto aluminium phosphate). In one embodiment the immunogenic composition of the invention comprises Hib, MenA and MenC saccharide
25 conjugates, or Hib and MenC saccharide conjugates, or Hib, MenC and MenY saccharide conjugates, or MenA, MenC, MenW and MenY saccharide conjugates, wherein at least one, two or all the saccharide conjugates are made according the method of the invention.

- Immunogenic compositions of the invention optionally comprise additional viral antigens
30 conferring protection against disease caused by measles and/or mumps and/or rubella and/or varicella. For example, immunogenic composition of the invention contains antigens from measles, mumps and rubella (MMR) or measles, mumps, rubella and varicella (MMRV). In an embodiment, these viral antigens are optionally present in the same container as the meningococcal and/or Hib saccharide conjugate(s) present in the
35 composition. In an embodiment, these viral antigens are lyophilised.

In an embodiment, the immunogenic composition of the invention further comprises an antigen from *N. meningitidis* serogroup B. The antigen is optionally an outer membrane

vesicle preparation from *N. meningitidis* serogroup B as described in EP301992, WO 01/09350, WO 04/14417, WO 04/14418 and WO 04/14419.

In general, the immunogenic composition of the invention may comprise a dose of each
5 saccharide conjugate between 0.1 and 20 μ g, 2 and 10 μ g, 2 and 6 μ g or 4 and 7 μ g of saccharide.

“Around” or “approximately” are defined as within 10% more or less of the given figure for the purposes of the invention.

10

In an embodiment, the immunogenic composition of the invention is adjusted to or buffered at, or adjusted to between pH 7.0 and 8.0, pH 7.2 and 7.6 or around or exactly pH 7.4.

15 The immunogenic composition or vaccines of the invention are optionally lyophilised in the presence of a stabilising agent for example a polyol such as sucrose or trehalose.

Optionally, the immunogenic composition or vaccine of the invention contains an amount of an adjuvant sufficient to enhance the immune response to the immunogen. Suitable
20 adjuvants include, but are not limited to, aluminium salts (aluminium phosphate or aluminium hydroxide), squalene mixtures (SAF-1), muramyl peptide, saponin derivatives, mycobacterium cell wall preparations, monophosphoryl lipid A, mycolic acid derivatives, non-ionic block copolymer surfactants, Quil A, cholera toxin B subunit, polyphosphazene and derivatives, and immunostimulating complexes (ISCOMs) such as those described by
25 Takahashi et al. (1990) Nature 344:873-875.

For *N. meningitidis* or HibMen combinations, it may be advantageous not to use any aluminium salt adjuvant or any adjuvant at all.

30 As with all immunogenic compositions or vaccines, the immunologically effective amounts of the immunogens must be determined empirically. Factors to be considered include the immunogenicity, whether or not the immunogen will be complexed with or covalently attached to an adjuvant or carrier protein or other carrier, route of administrations and the number of immunising dosages to be administered.

35

The active agent can be present in varying concentrations in the pharmaceutical composition or vaccine of the invention. Typically, the minimum concentration of the substance is an amount necessary to achieve its intended use, while the maximum concentration is the maximum amount that will remain in solution or homogeneously suspended within the initial mixture. For instance, the minimum amount of a therapeutic agent is optionally one which will provide a single therapeutically effective dosage. For bioactive substances, the minimum concentration is an amount necessary for bioactivity upon reconstitution and the maximum concentration is at the point at which a homogeneous suspension cannot be maintained. In the case of single-dosed units, the amount is that of a single therapeutic application. Generally, it is expected that each dose will comprise 1-100 μ g of protein antigen, optionally 5-50 μ g or 5-25 μ g. For example, doses of bacterial saccharides are 10-20 μ g, 5-10 μ g, 2.5-5 μ g or 1-2.5 μ g of saccharide in the conjugate.

The vaccine preparations of the present invention may be used to protect or treat a mammal (for example a human patient) susceptible to infection, by means of administering said vaccine via systemic or mucosal route. A human patient is optionally an infant (under 12 months), a toddler (12-24, 12-16 or 12-14 months), a child (2-10, 3-8 or 3-5 years) an adolescent (12-21, 14-20 or 15-19 years) or an adult. These administrations may include injection *via* the intramuscular, intraperitoneal, intradermal or subcutaneous routes; or *via* mucosal administration to the oral/alimentary, respiratory, genitourinary tracts. Intranasal administration of vaccines for the treatment of pneumonia or otitis media is preferred (as nasopharyngeal carriage of pneumococci can be more effectively prevented, thus attenuating infection at its earliest stage). Although the vaccine of the invention may be administered as a single dose, components thereof may also be co-administered together at the same time or at different times (for instance if saccharides are present in a vaccine these could be administered separately at the same time or 1-2 weeks after the administration of a bacterial protein vaccine for optimal coordination of the immune responses with respect to each other). In addition to a single route of administration, 2 different routes of administration may be used. For example, viral antigens may be administered ID (intradermal), whilst bacterial proteins may be administered IM (intramuscular) or IN (intranasal). If saccharides are present, they may be administered IM (or ID) and bacterial proteins may be administered IN (or ID). In addition, the vaccines of the invention may be administered IM for priming doses and IN for booster doses.

Vaccine preparation is generally described in Vaccine Design ("The subunit and adjuvant approach" (eds Powell M.F. & Newman M.J.) (1995) Plenum Press New York). Encapsulation within liposomes is described by Fullerton, US Patent 4,235,877.

- 5 A further aspect of the invention is a process for making the immunogenic composition or vaccine of the invention, comprising the step of mixing the MenA and MenC saccharides of the invention made by the method of the invention, with MenW and MenY that have not been made according to the invention, and with a pharmaceutically acceptable excipient.
- 10 The terms "comprising", "comprise" and "comprises" herein are intended by the inventors to be optionally substitutable with the terms "consisting of", "consist of" and "consists of", respectively, in every instance.

- All references or patent applications cited within this patent specification are incorporated
- 15 by reference herein.

- The invention is illustrated in the accompanying examples. The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative , but
- 20 do not limit the invention.

Examples

Example 1 – preparation of polysaccharide conjugates

5 Example 1a – preparation of meningococcal MenA and MenC capsular polysaccharide conjugate according to the invention

MenC –TT conjugates were produced using native polysaccharides (of over 150kDa as measured by MALLS) or were slightly microfluidised. MenA-TT conjugates were produced
10 using either native polysaccharide or slightly microfluidised polysaccharide of over 60kDa as measured by the MALLS method of example 2. Sizing was by microfluidisation using a homogenizer Emulsiflex C-50 apparatus. The polysaccharides were then filtered through a 0.2µm filter.

15 In order to conjugate MenA capsular polysaccharide to tetanus toxoid via a spacer, the following method was used. The covalent binding of the polysaccharide and the spacer (ADH) is carried out by a coupling chemistry by which the polysaccharide is activated under controlled conditions by a cyanylating agent, 1-cyano-4-dimethylamino-pyridinium tetrafluoroborate (CDAP). The spacer reacts with the cyanylated PS through its hydrazino
20 groups, to form a stable isourea link between the spacer and the polysaccharide.

A 10mg/ml solution of MenA (pH 6.0) [3.5 g] was treated with a freshly prepared 100mg/ml solution of CDAP in acetonitrile/water (50/50 (v/v)) to obtain a CDAP/MenA ratio of 0.75 (w/w). After 1.5 minutes, the pH was raised to pH 10.0. Three minutes later,
25 ADH was added to obtain an ADH/MenA ratio of 8.9. The pH of the solution was decreased to 8.75 and the reaction proceeded for 2 hours maintaining this pH (with temperature kept at 25 °C).

The PSA_{AH} solution was concentrated to a quarter of its initial volume and then diafiltered with 30 volumes of 0.2M NaCl using a Filtron Omega membrane with a cut-off of 10kDa,
30 and the retentate was filtered.

Prior to the conjugation (carbodiimide condensation) reaction, the purified TT solution and the PSA_{AH} solution were diluted to reach a concentration of 10 mg/ml for PSA_{AH} and 10mg/ml for TT.

EDAC (1-ethyl-3-(3-dimethyl-aminopropyl) carbodiimide) was added to the PSA_{AH} solution
35 (2g saccharide) in order to reach a final ratio of 0.9 mg EDAC/mg PSA_{AH}. The pH was adjusted to 5.0. The purified tetanus toxoid was added with a peristaltic pump (in 60 minutes) to reach 2 mg TT/mg PSA_{AH}. The resulting solution was left 60 min at +25°C under stirring to obtain a final coupling time of 120 min. The solution was neutralised by addition of 1M Tris-Hcl pH 7.5 (1/10 of the final volume) and left 30 minutes at +25°C then
40 overnight at +2°C to +8°C.

The conjugate was clarified using a 10µm filter and was purified using a Sephacryl S400HR column (Pharmacia, Sweden). The column was equilibrated in 10 mM Tris-HCl (pH 7.0), 0.075 M NaCl and the conjugate (approx. 660mL) was loaded on the column (+2°C to +8°C). The elution pool was selected as a function of optical density at 280 nm.

- 5 Collection started when absorbance increased to 0.05. Harvest continued until the Kd reached 0.30. The conjugate was filter sterilised at +20°C, then stored at +2°C to +8°C. The resultant conjugate had a polysaccharide:protein ratio of 1:2-1:4 (w/w).

- 10 In order to conjugate MenC capsular polysaccharide to tetanus toxoid via a spacer, the following method was used. The covalent binding of the polysaccharide and the spacer (ADH) is carried out by a coupling chemistry by which the polysaccharide is activated under controlled conditions by a cyanylating agent, 1-cyano-4-dimethylamino-pyridinium tetrafluoroborate (CDAP). The spacer reacts with the cyanylated PS through its hydrazino groups, to form a stable isourea link between the spacer and the polysaccharide.

- 15 A 20mg/ml solution of MenC (pH6.0) (3.5 g) was treated with a freshly prepared 100mg/ml solution of CDAP in acetonitrile/water (50/50 (v/v)) to obtain a CDAP/MenC ratio of 1.5 (w/w). After 1.5 minutes, the pH was raised to pH 10.0. At activation pH 5M NaCl was added to achieve a final concentration of 2M NaCl. Three minutes later, ADH was added to obtain an ADH/MenC ratio of 8.9. The pH of the solution was decreased to 8.75 and the
20 reaction proceeded for 2 hours (retained at 25 °C).

The PSC_{AH} solution was concentrated to a minimum of 150 mL and then diafiltered with 30 volumes of 0.2M NaCl using a Filtron Omega membrane with a cut-off of 10kDa, and the retentate was filtered.

- 25 Prior to the conjugation reaction, the purified TT solution and the PSC_{AH} solution (2g scale) were diluted in 0.2M NaCl to reach a concentration of 15 mg/ml for PSC_{AH} and 20mg/ml for TT.

- The purified tetanus toxoid was added to the PSC_{AH} solution in order to reach 2 mg TT/mg PSC_{AH}. The pH was adjusted to 5.0. EDAC (16.7 mg/ml in Tris 0.1M pH 7.5) was added with a peristaltic pump (in 10 minutes) to reach a final ratio of 0.5 mg EDAC/mg PSC_{AH}.
30 The resulting solution was left 110 min at +25°C under stirring and pH regulation to obtain a final coupling time of 120 min. The solution was then neutralized by addition of 1M Tris-Hcl pH 9.0 (1/10 of final volume) and left 30 minutes at +25°C then overnight at +2°C to +8°C.

- The conjugate was clarified using a 10µm filter and was purified using a Sephacryl S400HR column (Pharmacia, Sweden). The column was equilibrated in 10 mM Tris-HCl (pH 7.0), 0.075 M NaCl and the conjugate (approx. 460mL) was loaded on the column (+2°C to +8°C). The elution pool was selected as a function of optical density at 280 nm. Collection started when absorbance increased to 0.05. Harvest continued until the Kd reached 0.20. The conjugate was filter sterilised at +20°C, then stored at +2°C to +8°C.
40 The resultant conjugate had a polysaccharide:protein ratio of 1:2-1:4 (w/w).

Various experiments adding EDAC over 10-45 minutes were carried out – in each case good quality MenC conjugates resulted. If, however the TT carrier was added last slowly to the MenC-ADH + EDAC mix this led to a gel – a conjugate that could not be purified.

Experiments were also carried out adding the EDAC all at once into the reaction but the final TT/PS ratio (2.7/1) (w/w) of the conjugate was lower than for the conjugate obtained via the reaction where EDAC was added over 10 minutes (3.3/1); furthermore the α TT and α PS antigenicity were both lower than that measured in respect of the conjugate made by the reaction where EDAC was added over 10 minutes.

Note on approximate % Derivatisation of the polysaccharides

MenCAH: after CDAP treatment with ADH about 3.47% of hydroxyl groups were derivatized with ADH (with an estimation of two available hydroxyl groups per repeat subunit). For MenA: about 11.5% of hydroxyl groups derivatized with ADH (considering there is only one available hydroxyl group per repeat unit).

Example 1b – preparation of pneumococcal capsular PS 3 polysaccharide conjugate

1) PS03-TT_{AH} process : PS03-TT_{AH}208

Sizing by Emulsiflex

PS was weighed on the basis of 10% theoretical moisture content. The native PS was dissolved overnight in 2M NaCl at an initial concentration of 3 mg/ml. Before sizing, the solution of native PS was clarified on 5 μ m cut-off filter.

A homogenizer EMULSIFLEX C-50 apparatus was used to reduce the molecular weight and the viscosity of the polysaccharide before the activation step. The efficiency of the sizing depends on the circuit pressure, the plunger alimentation pressure and on the total cycles number. In order to improve the efficiency of sizing (and consequently reduce the total number of cycles), the homogenizing cell of Emulsiflex was replaced with a cell with a fixed geometry (Microfluidics F20Y-0.75 μ m interaction chamber). The aim of the sizing was to reduce the molecular weight and the viscosity of the PS without a critical decrease of its antigenicity.

The size reduction was done at 6000 ± 500 psi and followed *in-process* by a measure of viscosity. The sizing was stopped when the target of 2.0 ± 0.2 cp was reached.

Filtration of sized PS on 0.22 μ m

Sized PS was filtered on a Millipak 40 membrane (cut-off 0.22 μ m) at a flow-rate of 10 ml/min.

TT derivatization

The derivatization step was performed at 25°C under continuous stirring in a T° controlled waterbath. TT was diluted in NaCl 0.2M to obtain a final TT concentration of 25mg/ml.

- 5 ADH was added in solid form to the TT solution to reach a 0.2M final concentration. After complete ADH dissolution, the solution was set at pH 6.2+/- 0.1 with HCl. EDAC was then added to the TT/ADH solution to reach a final 0.02M concentration. The pH was set at 6.2+/-0.1 with HCl and was kept under pH regulation during 1 hour. After the derivatization step, the pH was raised up to pH9.5 with NaOH to stop the reaction. The solution was left during 2 hours under pH regulation before the diafiltration step.

Diafiltration

- 15 TT_{AH} derivative was diafiltered in order to remove unreacted ADH and EDAC by-products. The diafiltration was performed on a centramate membrane (0.09 m², 10 kDa cut-off). The solution was dialysed against 20 volumes of 0.2M NaCl. The follow-up of the diafiltration step was performed by a quantification of ADH (TNBS assay) in the permeate after 5, 10, 15 and 20 volumes of diafiltration.

20

Filtration on 0.22 µm

TT_{AH} was finally filtered on 0.22 µm cut-off membrane (Millipack 40) at a flow-rate of 10 ml/min. The filtered TT_{AH} was then stored at -70°C

25

PS3-TT_{AH} conjugate

The conditions of process were the following:

- 30 An initial PS3 concentration of 2 mg/ml in 2 M NaCl, an initial TT_{AH}/PS3 ratio of 1.5/1 (w/w), an EDAC concentration of 0.5 mg/mg PS, and a TT concentration of 10 mg/ml in 0.15M NaCl.

- 35 50mg of PS3 were diluted in 2M NaCl to obtain a final PS concentration of 2 mg/ml. The purified TT_{AH} solution was diluted in 0.2M NaCl to reach a concentration of 10 mg/ml.

The PS3 solution was adjusted to pH5 with HCl.

- 40 EDAC was added in solid form to the PS3 solution in order to reach a final concentration of 0.5 mg EDAC/ mg PS. The pH was adjusted to 5.0 ± 0.05 with HCl and TT_{AH} was manually added in 11 minutes (aliquots/min). The resulting solution was incubated 109 min at + 25°C with stirring and pH regulation to obtain a final coupling time of 120 min. Then the solution was neutralized by addition of 1M Tris-HCl pH 7.5 and left 30 min at +25°C. The conjugate was finally clarified on a 5µm membrane and injected on a Sephacryl S400HR column.

2) PS03-TT_{AH} process : PS03_{AH}-TT215**Sizing by Emulsiflex**

5

As above.

Filtration of sized PS on 0.22 µm

10

As above.

PS3 derivatization

15

The derivatization step was performed at 25°C under continuous stirring in a T° controlled waterbath. PS3 was diluted in NaCl 2M to obtain a final PS concentration of 3mg/ml. The PS solution was set at pH6.0 before the addition of CDAP (0.25mg/mg PS, dissolution at 100mg/ml in a mix of acetonitrile/ WFI). The pH was increased to pH9.5 with NaOH before the addition of ADH (8.9mg ADH/mg PS, dissolution at 100mg/ml in 0.2M NaCl). The pH was kept at 9.5 and regulated during 60 minutes. The percentage of derivatization corresponded to 2.4% (2.4 mg ADH/ 100 mg PS). This can be measured with known techniques: TNBS for the estimating ADH; and DMAB or resorcinol (Monsigny et al (1988) Anal. Biochem. 175, 525-530) for the PS quantification. In this case, TNBS dosage was 228 µg/ml and PS dosage: 5250 µg/ml.

20

25

Given the Mw of ADH is 174.2, and the Mw of the repeat unit of PS3 is 338.27 (having 1 COOH and 4 OH groups), there is 1.3 µmoles of ADH / 15.52 µmole of repeat unit, or 1.3 µmoles of ADH / 62.08 µmole of reactive hydroxyl group. 2.09% of PS3 hydroxyl groups were ADH modified hydroxyl groups.

30

Diafiltration

PS3_{AH} derivative was diafiltered in order to remove unreacted ADH and CDAP by-products. The diafiltration was performed on a UFP-30-C-H24LA membrane (42 cm², 30 kDa cut-off). The solution was dialysed against 20 volumes of 0.2M NaCl.

35

The follow-up of the diafiltration step was performed by a quantification of ADH (TNBS assay) in the permeate after 5, 10, 15 and 20 volumes of diafiltration.

Filtration on 0.22 µm

40

PS_{AH} was finally filtered on 0.22 µm cut-off membrane (Millipack 40) at a flow-rate of 10 ml/min. The filtered PS3_{AH} was then stored at 4°C.

PS3_{AH}-TT conjugate

The conditions of the process were the following:

- 5 An initial PS3 concentration of 2 mg/ml in 2 M NaCl, an initial TT/PS3_{AH} ratio of 1.5/1 (w/w), an EDAC concentration of 0.5 mg/mg PS, and a TT concentration of 10 mg/ml in 0.15M NaCl.

50mg of PS3_{AH} was diluted in 0.2M NaCl to obtain a final PS concentration of 2 mg/ml.

- 10 The purified TT solution was diluted in 0.2M NaCl to reach a concentration of 10 mg/ml.

The PS3_{AH} solution was adjusted to pH5 with HCl.

EDAC was added in solid form to the PS3_{AH} solution in order to reach a final concentration of 0.5 mg EDAC/ mg PS. The pH was adjusted to 5.0 ± 0.05 with HCl and

- 15 TT was added in 10 minutes using a peristaltic pump. The resulting solution was incubated 110 min at + 25°C with stirring and pH regulation to obtain a final coupling time of 120 min. Then the solution was neutralized by addition of 1M Tris-HCl pH 7.5 and left 30 min at +25°C. The conjugate was finally clarified on a 5µm membrane and injected on a Sephacryl S400HR column.

- 20 **3) PS03_{AH}-TT process : PS3_{AH}-TT217**

Sizing by Emulsiflex

As above.

25

Filtration of sized PS on 0.22 µm

As above.

- 30 **PS3 derivatization**

As for 215 conjugate.

Diafiltration

35

As for 215 conjugate.

Filtration on 0.22 µm

- 40 As for 215 conjugate.

PS3_{AH}-TT conjugate

The conditions of the process were the following:

- 5 An initial PS3 concentration of 2 mg/ml in 2 M NaCl, an initial TT/PS3_{AH} ratio of 1.5/1 (w/w), an EDAC concentration of 0.5 mg/mg PS, and a TT concentration of 10 mg/ml in 0.15M NaCl.

- 50mg of PS3_{AH} was diluted in 0.2M NaCl to obtain a final PS concentration of 2 mg/ml.
10 The purified TT solution was diluted in 0.2M NaCl to reach a concentration of 10 mg/ml.
The PS3_{AH} and TT solutions were mixed and adjusted to pH5 with HCl.
EDAC was dissolved in a Tris 1M pH7.5 buffer. 40µl of EDAC were added each minute (10 minutes to reach the EDAC/PS ratio (0.5mg EDAC/ mg PS)). The resulting solution was incubated 110 min at + 25°C under stirring and pH regulation to obtain a final
15 coupling time of 120 min. Then the solution was neutralized by addition of 1M Tris-HCl pH 7.5 and left 30 min at +25°C. The conjugate was finally clarified on a 5µm membrane and injected on a Sephacryl S400HR column.

4) PS03_{AH}-TT process : PS3_{AH}-TT218

20

Sizing by Emulsiflex

As above.

- 25 **Filtration of sized PS on 0.22 µm**

As above.

PS3 derivatization

30

- The derivatization step was performed at 25°C with continuous stirring in a T° controlled waterbath. PS3 was diluted in NaCl 2M to obtain a final PS concentration of 3mg/ml. EDAC was added in solid form to reach an EDAC/ PS ratio of 0.1mg/mg PS. After complete dissolution, the pH of the solution was set at 5. ADH (8.9mg ADH/mg PS,
35 dissolution at 100mg/ml in 0.2M NaCl) was then added using a peristaltic pump in 44 minutes (though as such an excess of ADH was present, direct addition would also have been OK). The pH was kept at 5.0+/-0.1 and regulated during 120 minutes (44' + 76'). The reaction was stopped by increasing the pH to 7.5 using sodium hydroxide. The percentage of derivatization corresponded to 3.7% (mg ADH/ mg PS). TNBS dosage was
40 220 µg/ml and PS dosage was 5902 µg/ml, thus there is 1.26 µmoles of ADH / 17.44 µmole of repeat unit (=µmole of reactive COOH group). Thus, 7.22% of PS3 carboxyl groups were ADH modified COOH groups.

Diafiltration

PS_{3AH} derivative was diafiltered in order to remove unreacted ADH and EDAC by-products. The diafiltration was performed on a UFP-30-C-H24LA membrane (42 cm², 30 kDa cut-off). The solution was dialysed against 23 volumes of 0.2M NaCl.

The follow-up of the diafiltration step was performed by a quantification of ADH (TNBS assay) in the permeate after 5, 10, 15 and 20 volumes of diafiltration

Filtration on 0.22 µm

PS_{3AH} was finally filtered on 0.22 µm cut-off membrane (Millipack 40) at a flow-rate of 10 ml/min. The filtered PS_{3AH} was then stored at 4°C.

PS_{3AH}-TT conjugate

The conditions of the process were the following:

An initial PS_{3AH} concentration of 2 mg/ml in 2 M NaCl, an initial TT/PS_{3AH} ratio of 1.5/1 (w/w), an EDAC concentration of 0.5 mg/mg PS, and a TT concentration of 10 mg/ml in 0.15M NaCl.

50mg of PS_{3AH} was diluted in 0.2M NaCl to obtain a final PS concentration of 2 mg/ml. The purified TT solution was diluted in 0.2M NaCl to reach a concentration of 10 mg/ml.

The PS_{3AH} and TT solutions were mixed together.

The pH was adjusted to 5.0 ± 0.05 with HCl and EDAC was manually added in 10 minutes (equal part-aliquots added each minute). The resulting solution was incubated 110 min at + 25°C with stirring and pH regulation to obtain a final coupling time of 120 min. Then the solution was neutralized by addition of 1M Tris-HCl pH 7.5 and left 30 min at +25°C. The conjugate was finally clarified on a 5µm membrane and injected on a Sephacryl S400HR column.

Conclusions:

Different conjugates were made using carbodiimide chemistry in the conjugation step. The last component added in the reaction solution can be either the TT protein or the EDAC reagent. The time of addition can have an effect on the resulting conjugates.

PS_{3AH}TT215 & 217 conjugates:

The same components and conditions were used to prepare both conjugates. The way in which the last component was added was different. PS_{3AH}TT217 conjugate led to a product which met *in-vitro* criteria. This one was made by adding EDAC in 10 minutes.

PS_{3AH}TT215 conjugate, however, could not be filtered on sterile membrane. For this one, the last component added in the reaction medium was the TT (in 10 minutes).

Final TT/PS ratios were highly different for both conjugates. (0.98/1 vs 0.50/1). If EDAC is added first to the PS_{3AH} (having both reactive amino and carboxyl groups) this can lead to intra cross-linking of hydrazine and carboxylic groups present on the polysaccharide, and thus could lead to a more cross-linked conjugate with a weaker final ratio after the addition of TT in 10 minutes.

This effect is not observed for the PS_{3AH}TT217 conjugate. The TT incorporation worked better by the addition of EDAC in 10 minutes, perhaps due to lower intra cross-linking, and better inter cross-linking between hydrazine groups of the PS_{3AH} and carboxylic groups of the protein.

In the case of the 218 conjugate, as the PS3 EDAC derivatisation only partially derivatises the polysaccharide (to keep the majority of the polysaccharides epitopes intact), again both reactive amino and carboxyl groups are present, hence why slow addition of EDAC in a final conjugation step is also beneficial.

Slow TT addition in the final conjugation step was beneficial (however) for the 208 conjugate where the TT was ADH derivatised (and comprises amino and carboxyl groups), whereas the PS3 was left with its native reactive –OH and –COOH groups as part of its repeating subunit. The addition of EDAC to PS 3 did not have the above cross-linking effect, and the slow addition of the derivatised TT yielded conjugate with good in vitro characteristics – see below.

In-vitro characterization:

Conj.	Derivatization/Chemistry	Conjugation/ Chemistry	Final component addition
208	TT/ADH → EDAC	PS-TT _{AH} → EDAC	TT _{AH} added in 11 minutes
215	PS3/ ADH → CDAP	PS _{AH} -TT → EDAC	TT added in 10 minutes
217	PS3/ ADH → CDAP	PS _{AH} -TT → EDAC	EDAC added in 10 minutes
218	PS3/ ADH → EDAC	PS _{AH} -TT → EDAC	EDAC added in 10 minutes

Conj.	PS	[PS] (mg/ml)	[TT] (mg/ml)	In.TT/PS ratio (w/w)	[EDAC] (mg/mg PS)	Coupl.time (min)
208	C6E02	2.0	10 (TT _{AH}), pump	1.5/1	0.5/1	120
215	3 _{AH} 001 (CDAP)	2.0	10 pump	1.5/1	0.5/1	120

217	3 _{AH} 001 (CDAP)	2.0	10	1.5/1	0.5/1 (Fractions)	120
218	3 _{AH} 002 (EDAC)	2.0	10	1.5/1	0.5/1 (Fractions)	120

Conj.	F.TT/PS ratio (w/w)	Yield PS rec (%)	Filtr.yield rec (%)	Free PS (%)	αPS/ αPS (%) Antigenicity	αTT/ αPS (%) Antigenicity
208	1.84/1	69	95	10.2	99	103 100*
215	0.50/1	17	27	-	-	-
217	0.98/1	66	100	0.7	17	103 100*
218	0.88/1	74	101	11.0	34	222 216*

*relative to the 208 conjugate

5 Example 1c – preparation of S. typhi Vi polysaccharide conjugate of the invention

Sizing by Emulsiflex

PS was weighed on the basis of 15% theoretical moisture content. The native PS is dissolved overnight in WFI at an initial concentration of 7 mg/ml. Before the sizing, the solution of native PS is clarified on 10 μm cut-off filter at a flow-rate of 50 ml/min. A homogenizer EMULSIFLEX C-50 apparatus was used to reduce the molecular weight and the viscosity of the polysaccharide before the activation step. The efficiency of the sizing depends on the circuit pressure, the plunger alimentation pressure and on the total cycles number. In order to improve the efficiency of sizing (and consequently reduce the total number of cycles), the homogenizing cell of Emulsiflex was replaced by a cell with a fixed geometry (Microfluidics F20Y-0.75μm interaction chamber). The aim of the sizing is to reduce the molecular weight and the viscosity of the PS without a critical decrease of its antigenicity.

The size reduction was realized at 15000 ± 500 psi and followed *in-process* by a measure of viscosity. The sizing is stopped when the target of 5.0 ± 0.3 cp is reached.

Filtration of sized PS on 0.22 μm

Sized PS is filtered on a Millipak 40 membrane (cut-off 0.22 μm) at a flow-rate of 10 ml/min. The filtered sized PS is stored at -20°C.

Derivatization of polysaccharide Vi

1.5 g of sized Vi PS was dissolved at 25°C in EPI under agitation (5 mg/ml). 13.35 g of ADH (8.9 mg ADH/mg PS) is added to the PS solution. After complete dissolution pH was adjusted at pH 5.0 ± 0.05 with 1N HCl. EDAC (0.1 mg/mg PS) was added in a solid form. The solution was left 60 min at 25°C. Then the solution was neutralized by addition of 1M Tris-HCl pH 7.5 and left at least 30 min at 25°C (maximum 2 hours). The level of derivatization was estimated to be 4.55% using the TNBS dosage (mg ADH/ 100 mg PS). TNBS dosage was 200 µg/ml and PS dosage was 4034 µg/ml; thus 0.0697 µmoles of ADH / 16.46 µmole of repeat unit (Mw 245). 1.3 µmoles of ADH / 16.46 µmole of reactive COOH group on Vi, thus 7% of Vi COOH groups were ADH modified COOH groups.

Diafiltration

PSVi_{ADH} derivative was diafiltered in order to remove unreacted ADH and EDAC by-products. The diafiltration was performed on a centramate membrane (0.09 m², 10 kDa cut-off). The solution was dialysed against 20 volumes of 0.2M NaCl. The follow-up of the diafiltration step was performed by a quantification of ADH (TNBS assay) in the permeate after 3, 5, 10 and 20 volumes of diafiltration

Filtration on 0.22 µm

PSVi_{ADH} was finally filtered on 0.22 µm cut-off membrane (Millipack 40) at a flow-rate of 10 ml/min. The filtered PSVi_{ADH} was stored at +2/+8°C for a maximum of 4 days.

PSVi_{ADH}-TT conjugates

The conditions of process were the following:

An initial PSVi_{ADH} concentration of 2 mg/ml in 0.2 M NaCl, an initial TT/PSVi_{ADH} ratio of 2.5/1 (w/w), an EDAC concentration of 0.25 mg/mg PS and a TT concentration of 10 mg/ml in 0.2M NaCl.

1g of PSVi_{ADH} was diluted in 0.2M NaCl to obtain a final PS concentration of 2 mg/ml (uronic acid dosage). The purified TT solution was diluted in 0.2M NaCl to reach a concentration of 10 mg/ml.

TT was added to the PSVi_{ADH} solution in order to reach a final ratio of 2.5 mg TT/ mg PS. The pH is adjusted to 5.0 ± 0.05 with 1N HCl. The EDAC solution (7.5 mg/ml in 0.1M Tris pH 7.5) was then added (in 10 minutes with a peristaltic pump) to reach 0.25 mg EDAC/ mg PSVi_{ADH}. The resulting solution was incubated 50 min at + 25°C with stirring and pH regulation to obtain a final coupling time of 60 min. Then the solution was neutralized by addition of 1M Tris-HCl pH 7.5 and left 30 min at +25°C. The conjugate was transferred at 4°C and is left overnight under continuous slow stirring before the chromatography step.

Purification

Prior to the elution on Sephacryl S400HR, the conjugate was clarified using a 10 μ m Kleenpak filter. The flow rate was fixed at 100 ml/min. The conjugate was then injected on
5 Sephacryl S400HR and the collection pool was based on a Kd value. The following criterion was used for the pool collection: from OD= 0.05 at 280 nm harvesting started, and finished when Kd= 0.22.

Sterilizing filtration

10 Before filtration, the bulk was brought back to room temperature. Then the conjugate was filtered on an Opticap 4" sterilizing membrane. The flow rate was fixed at 30 ml/min.

Analytical

15 The resulting conjugate had a final TT/PS ratio (w/w) of 2.44/1, a free PS content of 3.7% and a α PS/ α PS antigenicity of 58%.

Example 1d – preparation of other polysaccharide conjugates

20 The covalent binding of *Haemophilus influenzae* (Hib) PRP polysaccharide to TT was carried out by a coupling chemistry developed by Chu et al (Infection and Immunity 1983, 40 (1); 245-256). Hib PRP polysaccharide was activated by adding CNBr and incubating at pH10.5 for 6 minutes. The pH was lowered to pH8.75 and adipic acid dihydrazide
25 (ADH) was added and incubation continued for a further 90 minutes. The activated PRP was coupled to purified tetanus toxoid via carbodiimide condensation using 1-ethyl-3-(3-dimethyl-aminopropyl)carbodiimide (EDAC). EDAC was added to the activated PRP to reach a final ratio of 0.6mg EDAC/mg activated PRP. The pH was adjusted to 5.0 and purified tetanus toxoid was added to reach 2mg TT/mg activated PRP. The resulting
30 solution was left for three days with mild stirring. After filtration through a 0.45 μ m membrane, the conjugate was purified on a sephacryl S500HR (Pharmacia, Sweden) column equilibrated in 0.2M NaCl.

35 MenC –TT conjugates were produced using native polysaccharides (of over 150kDa as measured by MALLS) or were slightly microfluidised. MenA-TT conjugates were produced using either native polysaccharide or slightly microfluidised polysaccharide of over 60kDa as measured by the MALLS method of example 2. MenW and MenY-TT conjugates were produced using sized polysaccharides of around 100-200kDa as measured by MALLS (see example 2). Sizing was by microfluidisation using a homogenizer Emulsiflex
40 C-50 apparatus. The polysaccharides were then filtered through a 0.2 μ m filter.

Activation and direct coupling were performed as described in WO96/29094 and WO 00/56360. Briefly, the polysaccharide at a concentration of 10-20mg/ml in 2M NaCl pH

5.5-6.0 was mixed with CDAP solution (100mg/ml freshly prepared in acetonitrile/WFI, 50/50) to a final CDAP/polysaccharide ratio of 0.75/1 or 1.5/1. After 1.5 minutes, the pH was raised with sodium hydroxide to pH10.0. After three minutes tetanus toxoid was added to reach a protein/polysaccharide ratio of 1.5/1 for MenW, 1.2/1 for MenY, 1.5/1 for MenA or 1.5/1 for MenC. The reaction continued for one to two hours.

After the coupling step, glycine was added to a final ratio of glycine/PS (w/w) of 7.5/1 and the pH was adjusted to pH9.0. The mixture was left for 30 minutes. The conjugate was clarified using a 10µm Kleenpak filter and was then loaded onto a Sephacryl S400HR column using an elution buffer of 150mM NaCl, 10mM or 5mM Tris pH7.5. Clinical lots were filtered on an Opticap 4 sterilizing membrane. The resultant conjugates had an average polysaccharide:protein ratio of 1:1-1:5 (w/w).

Example 2 – determination of molecular weight using MALLS

Detectors were coupled to a HPLC size exclusion column from which the samples were eluted. On one hand, the laser light scattering detector measured the light intensities scattered at 16 angles by the macromolecular solution and on the other hand, an interferometric refractometer placed on-line allowed the determination of the quantity of sample eluted. From these intensities, the size and shape of the macromolecules in solution can be determined.

The mean molecular weight in weight (M_w) is defined as the sum of the weights of all the species multiplied by their respective molecular weight and divided by the sum of weights of all the species.

a) Weight-average molecular weight: - M_w -

$$M_w = \frac{\sum W_i \cdot M_i}{\sum W_i} = \frac{m_2}{m_1}$$

b) Number-average molecular weight: - M_n -

$$M_n = \frac{\sum N_i \cdot M_i}{\sum N_i} = \frac{m_1}{m_0}$$

c) Root mean square radius: - R_w - and R^2_w is the square radius defined by:

$$R^2_w \text{ or } (r^2)_w = \frac{\sum m_i \cdot r_i^2}{\sum m_i}$$

(-m_i- is the mass of a scattering centre i and -r_i- is the distance between the

5 scattering centre i and the center of gravity of the macromolecule).

d) The polydispersity is defined as the ratio -M_w / M_n-.

10 Meningococcal polysaccharides were analysed by MALLS by loading onto two HPLC columns (TSKG6000 and 5000PWxl) used in combination. 25µl of the polysaccharide were loaded onto the column and was eluted with 0.75ml of filtered water. The polysaccharides are detected using a light scattering detector (Wyatt Dawn DSP equipped with a 10mW argon laser at 488nm) and an interferometric refractometer (Wyatt Otilab DSP
15 equipped with a P100 cell and a red filter at 498nm).

The molecular weight polydispersities and recoveries of all samples were calculated by the Debye method using a polynomial fit order of 1 in the Astra 4.72 software.

20

Example 3 – Clinical trial assessing the effect of a linker in MenA in a MenACWY conjugate vaccine

25 A single dose of different formulations of MenACWY vaccine was administered to teenagers of 15-19 years in 5 groups of 25 subjects in a 1:1:1:1:1 randomised trial. The formulations tested were:

30 F1 – MenACWY conjugated to tetanus toxoid with the MenA conjugate containing an AH (ADH) spacer (made according to example 1) – 5/5/5/5µg

F2 – MenACWY conjugated to tetanus toxoid with the MenA conjugate containing an AH spacer (made according to example 1) – 2.5/5/2.5/2.5µg

F3 – MenACWY conjugated to tetanus toxoid with the MenA conjugate containing an AH spacer (made according to example 1) – 5/5/2.5/2.5µg

35 F4 – MenACWY conjugated to tetanus toxoid with no spacer in any conjugate – 5/5/5/5µg
Control group – Mencevax™ ACWY

On day 30 after inoculation, a blood sample was taken from the patients.

The blood samples were used to assess the percentage of SBA-MenA, SBA-MenC, SBA-MenW135 and SBA-MenY responders one month after the vaccine dose. A vaccine response was defined as 1) for initially seronegative subjects – a post-vaccination antibody titre $\geq 1/32$ at 1 month or 2) for initially seropositive subjects – antibody titre of ≥ 4 fold the pre-vaccination antibody titre.

Results

As shown in the Table below, the use of a spacer in the MenA conjugate led to an increased immune response against MenA. The percentage of responders rose from 66% to 90-95% when the AH spacer was added. This was reflected in an increase in SBA GMT from 4335 to 10000 and an increase in GMC from 5 to 20-40. Surprisingly, the use of a AH spacer also led to an increased immune response against MenC as seen by an increase in the percentage of responders and an increase in the SBA GMT. An increase could also be seen in the SBA-GMT against MenY (6742-7122) and against MenW (4621-5418) when a spacer was introduced.

Formulation	% SBA MenA responders	SBA-MenA GMT	Anti-PSA GMC $\mu\text{g/ml}$ ELISA
F 1 5AH/5/5/5	90.9	9805	20.38
F2 2.5AH/5/2.5/2.5	75	8517	29.5
F3 5AH/5/2.5/2.5	95.5	10290	47.83
F4 5/5/5/5	66.7	4335	5.46
Mencevax™	85.7	8022	27.39
Formulation	% SBA MenC responders	SBA-MenC GMT	Anti-PSC GMC $\mu\text{g/ml}$ ELISA
F 1 5AH/5/5/5	69.6	3989	12.11
F2 2.5AH/5/2.5/2.5	81.8	3524	12.78
F3 5AH/5/2.5/2.5	81.8	3608	8.4
F4 5/5/5/5	73.9	2391	8.84
Mencevax™	90.0	5447	38.71
Formulation	% SBA MenW responders	SBA-MenW GMT	Anti-PSW GMC $\mu\text{g/ml}$ ELISA
F 1 5AH/5/5/5	95	5418	9.65
F2 2.5AH/5/2.5/2.5	85	4469	14.55
F3 5AH/5/2.5/2.5	95.5	4257	6.39
F4 5/5/5/5	95.5	4621	10.7
Mencevax™	86.4	2714	13.57
Formulation	% SBY MenY responders	SBA-MenY GMT	Anti-PSY GMC $\mu\text{g/ml}$ ELISA

F 1 5AH/5/5/5	91.3	7122	16.3
F2 2.5AH/5/2.5/2.5	87.5	5755	12.52
F3 5AH/5/2.5/2.5	80	5928	8.88
F4 5/5/5/5	91.3	6742	13.88
Mencevax™	91.7	4854	21.02

Example 4 – Clinical trial assessing the effect of a linker in MenA and MenC conjugates in a MenACWY conjugate vaccine

- 5 A single dose of different formulations of MenACWY vaccine was administered to teenagers of 15-19 years in 5 groups of 25 subjects in a 1:1:1:1:1 randomised trial. The formulations tested were:

- 10 F1 – MenACWY conjugated to tetanus toxoid with the MenA and MenC conjugates containing an AH spacer (made according to example 1) – 2.5/2.5/2.5/2.5µg
 F2 – MenACWY conjugated to tetanus toxoid with the MenA and MenC conjugates containing an AH spacer (made according to example 1) – 5/5/2.5/2.5µg
 F3 – MenACWY conjugated to tetanus toxoid with the MenA and MenC conjugates containing an AH spacer (made according to example 1) – 5/5/5/5µg
 15 F4 – MenACWY conjugated to tetanus toxoid with the MenA conjugate containing an AH spacer (made according to example 1) – 5/5/5/5µg
 Control group – Mencevax™ ACWY

- 20 On day 30 after inoculation, a blood sample was taken from the patients.

- The blood samples were used to assess the percentage of SBA-MenA, SBA-MenC, SBA-MenW135 and SBA-MenY responders one month after the vaccine dose. A vaccine response was defined as 1) for initially seronegative subjects – a post-vaccination antibody titre $\geq 1/32$ at 1 month or 2) for initially seropositive subjects – antibody titre of ≥ 4 fold the pre-vaccination antibody titre.

Results

- 30 The introduction of an AH spacer into the MenC conjugate led to an increase in the immune response against MenC as shown in the Table below. This is demonstrated by an increase in SBA GMT from 1943 to 4329 and an increase in anti-PSC GMC from 7.65 to 13.13. Good immune responses against MenA, MenW and MenY were maintained.

Formulation	% SBA MenA responders	SBA-MenA GMT	Anti-PSA GMC µg/ml ELISA
F 12.5AH/2.5AH/2.5/2.5	75	8417	20.23
F2 5AH/5AH/2.5/2.5	72	6299	16.07
F3 5AH/5AH/5/5	87	9264	27.26
F4 5AH/5/5/5	77.3	9632	20.39

Mencevax™	78.3	8284	12.93
Formulation	% SBA MenC responders	SBA-MenC GMT	Anti-PSC GMC μg/ml ELISA
F 12.5AH/2.5AH/2.5/2.5	88	3619	12.82
F2 5AH/5AH/2.5/2.5	88	2833	13.32
F3 5AH/5AH/5/5	95.8	4329	13.13
F4 5AH/5/5/5	95.8	1943	7.65
Mencevax™	91.7	1567	16.55
Formulation	% SBA MenW responders	SBA-MenW GMT	Anti-PSW GMC μg/ml ELISA
F 12.5AH/2.5AH/2.5/2.5	100	5656	7
F2 5AH/5AH/2.5/2.5	96	4679	5.4
F3 5AH/5AH/5/5	91.3	4422	4.45
F4 5AH/5/5/5	88	4947	7.67
Mencevax™	96	3486	11.93
Formulation	% SBY MenY responders	SBA-MenY GMT	Anti-PSY GMC μg/ml ELISA
F 1 2.5AH/2.5AH/2.5/2.5	75	3891	17.81
F2 5AH/5AH/2.5/2.5	92	3968	11.96
F3 5AH/5AH/5/5	79.2	2756	9.51
F4 5AH/5/5/5	80	3914	16.76
Mencevax™	88	3056	21.41

CLAIMS

1. A method of conjugating a saccharide to a protein carrier using carbodiimide condensation chemistry, wherein the saccharide comprises (for instance as part of its repeating unit), or has been derivatised to comprise, amino and/or carboxyl groups, and wherein the protein carrier comprises, or has been derivatised to comprise, amino and/or carboxyl groups, comprising the steps of:
- 5 I) - if the protein carrier comprises both amino and carboxyl groups and the saccharide comprises either amino or carboxyl groups:
- 10 a) mixing the saccharide and aliquot of carbodiimide required to perform the conjugation, and
- b) adding the aliquot of protein carrier required over a period of 35 seconds to 6 hours;
- 15 II) - if the saccharide comprises both amino and carboxyl groups and the protein carrier comprises either amino or carboxyl groups:
- a) mixing the protein carrier and aliquot of carbodiimide required to perform the conjugation, and
- 20 b) adding the aliquot of saccharide required over a period of 35 seconds to 6 hours;
- 25 III) - if the saccharide comprises both amino and carboxyl groups and the protein carrier comprises both amino and carboxyl groups:
- a) mixing the protein carrier and saccharide, and
- b) adding the aliquot of carbodiimide required to perform the conjugation over a period of 35 seconds to 6 hours.
2. The method of claim 1, wherein in step b) the period is 50 seconds to 5 hours, 1 minute to 4 hours, 2 minutes to 3 hours, 3 minutes to 2 hours, 4 to 60 minutes, 5 to 50 minutes, 6 to 40 minutes, 7 to 30 minutes or 8 to 20 minutes.
3. The method of claim 1, wherein in step b) the period is 1 minute to 5 hours, 10 minutes to 4 hours, 20 minutes to 3 hours, 30 minutes to 2 hours, 40 to 90 minutes, or 50 to 70 minutes.
4. The method of claims 1-3, wherein the carbodiimide is EDAC (1-ethyl-3-(3-dimethyl-aminopropyl) carbodiimide) or a carbodiimide other than EDAC.
5. The method of claims 1-4, wherein the aliquot of carbodiimide required to perform the conjugation is 0.01 to 3, 0.05 to 2 or 0.09 to 1 mg/mg saccharide.
6. The method of claims 1-5, wherein the saccharide and/or protein carrier has been derivatised to comprise amino or carboxyl groups.

7. The method of claim 6, wherein the derivatisation is through the addition of a hetero- or homo-bifunctional linker.
8. The method of claim 7, wherein the linker has between 4 and 12 carbon atoms.
9. The method of claim 7 or 8, wherein the linker has two reactive amino groups.
10. The immunogenic composition of claims 7-9 wherein the linker is ADH.
11. The method of claim 7 or 8, wherein the linker has two reactive carboxylic acid groups.
12. The method of claim 7 or 8, wherein the linker has a reactive amino group at one end and a reactive carboxylic acid group at the other end.
13. The method of claims 7-12, wherein the derivatization takes place through reacting a large excess of linker with the saccharide and/or protein carrier to be derivatised.
14. The method of claims 7-13, wherein the saccharide comprises a reactive hydroxyl group as part of its repeating unit which is partially derivatised via an amino group on the linker.
15. The method of claim 14, wherein the saccharide is partially derivatised with CDAP chemistry.
16. The method of claims 7-13, wherein the saccharide comprises a reactive amino group as part of its repeating unit which is partially derivatised via a carboxyl group on the linker.
17. The method of claim 16, wherein the saccharide is partially derivatised with carbodiimide condensation chemistry.
18. The method of claims 7-13, wherein the saccharide comprises a reactive carboxyl group as part of its repeating unit which is partially derivatised via an amino group on the linker.
19. The method of claim 18, wherein the saccharide is partially derivatised with carbodiimide chemistry.
20. The method of claims 1-19, wherein in step b) the aliquot of carbodiimide, saccharide or protein carrier is added at a constant rate using a pump.

21. The method of claims 1-19, wherein in step b) the aliquot of carbodiimide, saccharide or protein carrier is added in stages over the period.
- 5 22. The method of claim 21, wherein at least one quarter of the aliquot is added over the first half of the period, and at least one quarter of the aliquot over the second half of the period.
- 10 23. The method of claim 21 or 22, wherein the aliquot 'a' is added in 4-100 stages 's'.
24. The method of claim 23, wherein a/s of the aliquot is added at each stage.
- 15 25. The method of claim 23 or 24, wherein if one stage takes place at time zero of the period 'p', each subsequent stage takes place at a time which is $p/(s-1)$.
26. The method of claims 1-25, wherein the saccharide is present at a final concentration of 0.5-50 mg/ml in step b).
- 20 27. The method of claims 1-26, wherein the initial ratio of protein carrier to saccharide is 5:1 to 1:5, 4:1 to 1:1, or 3:1 to 2:1 (w/w).
28. The method of claims 1-27, wherein the concentration of salt, for instance NaCl, present in step b) is 0-2, 0.1-1 or 0.2-0.5 M.
- 25 29. The method of claims 1-28, wherein the protein carrier is present at a final concentration of 1-50 mg/ml in step b).
- 30 30. The method of claims 1-29, wherein the reaction pH in step b) is maintained at pH 4.5-6.5, 4.7-6.0, or 5-5.5.
31. The method of claims 1-29, wherein N-hydroxysuccinimide is also present in the reaction in step b), and the reaction pH in step b) is maintained at pH 4.5-7.5.
- 35 32. The method of claims 1-31, wherein the temperature of the reaction in step b) is maintained at 4-37, 10-32, 17-30, or 22-27 °C.
- 40 33. The method of claims 1-32, wherein after the aliquot has all been added in step b) the reaction is maintained for a further 10 minutes to 72 hours, 20 minutes to 48 hours, 30 minutes to 24 hours, 40 minutes to 12 hours, 50 minutes to 6 hours, or 1-3 hours.

34. The method of claims 1-33, wherein once the reaction is completed the pH is adjusted to 7.5-9.
- 5 35. The method of claims 1-34, comprising a subsequent step c), wherein the saccharide-protein conjugate is purified on a size exclusion chromatography column.
- 10 36. The method of claims 1-35, comprising a subsequent step d), wherein the saccharide-protein conjugate is sterile filtered.
- 15 37. The method of claims 1-36, comprising a subsequent step e), wherein an effective dose of the saccharide-protein conjugate is formulated with a pharmaceutically acceptable excipient to manufacture an immunogenic composition or vaccine.
- 20 38. The method of claims 1-37, wherein the saccharide is a bacterial capsular saccharide, for instance derived from a bacterium selected from a list consisting of: *N. meningitidis* serogroup A, B, C, W135 or Y, *Streptococcus pneumoniae* serotypes 1, 2, 3, 4, 5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F or 33F, Group B Streptococcus group Ia, Ib, II, III, IV, V, VI, or VII, *Staphylococcus aureus* type 5, *Staphylococcus aureus* type 8, *Salmonella typhi* (Vi saccharide), *Vibrio cholerae*, or *H. influenzae* type b.
- 25 39. The method of claims 1-38, wherein the weight-average molecular weight of the saccharide is 1000-2000000, 5000-1000000, 10000-500000, 50000-400000, 75000-300000, or 100000-200000.
- 30 40. The method of claims 1-39, wherein the saccharide is either a native polysaccharide or is sized by a factor of no more than x10 (for instance by microfluidization).
- 35 41. The method of claims 1-37, wherein the saccharide is a bacterial lipooligosaccharide or lipopolysaccharide, for instance derived from a bacterium selected from a list consisting of: *N. meningitidis*, *H. influenzae*, *E. coli*, *Salmonella* or *M. catarrhalis*.
42. The method of claims 1-41, wherein the protein carrier comprises one or more T-helper epitopes.
- 40 43. The method of claims 1-42, wherein the protein carrier is selected from the group consisting of: TT, DT, CRM197, fragment C of TT, protein D of *H. influenzae*, pneumococcal PhtD, and pneumococcal Pneumolysin.

44. A saccharide-protein carrier conjugate obtainable by the method of claims 1-43.

45. An immunogenic composition or vaccine obtainable by the method of claims 1-43.

46. Use of the immunogenic composition or vaccine of claim 45 in the manufacture of a medicament for the prevention or treatment of disease.

47. Method of preventing or treating disease comprising the step of administering an effective dose of the immunogenic composition or vaccine of claim 45 to a patient in need thereof.

48. The use or method of claim 46 or 47, wherein the disease is caused by a bacterium selected from a list consisting of: *N. meningitidis*, *Streptococcus pneumoniae*, *M. catarrhalis*, Group B Streptococcus, *Staphylococcus aureus*, *Salmonella typhi*, *Vibrio cholerae*, *E. coli*, and *H. influenzae*.