

LIPIDS FOR LIPID NANOPARTICLE DELIVERY OF ACTIVE AGENTS

BACKGROUND

Technical Field

The present disclosure generally relates to novel cationic lipids that can
5 be used in combination with other lipid components, such as neutral lipids, cholesterol
and polymer conjugated lipids, to form lipid nanoparticles with oligonucleotides, to
facilitate the intracellular delivery of therapeutic nucleic acids (*e.g.*, oligonucleotides,
messenger RNA) both *in vitro* and *in vivo*.

Description of the Related Art

10 There are many challenges associated with the delivery of nucleic acids
to affect a desired response in a biological system. Nucleic acid based therapeutics
have enormous potential but there remains a need for more effective delivery of nucleic
acids to appropriate sites within a cell or organism in order to realize this potential.
Therapeutic nucleic acids include, *e.g.*, messenger RNA (mRNA), antisense
15 oligonucleotides, ribozymes, DNAzymes, plasmids, immune stimulating nucleic acids,
antagomir, antimir, mimic, supermir, and aptamers. Some nucleic acids, such as
mRNA or plasmids, can be used to effect expression of specific cellular products as
would be useful in the treatment of, for example, diseases related to a deficiency of a
protein or enzyme. The therapeutic applications of translatable nucleotide delivery are
20 extremely broad as constructs can be synthesized to produce any chosen protein
sequence, whether or not indigenous to the system. The expression products of the
nucleic acid can augment existing levels of protein, replace missing or non-functional
versions of a protein, or introduce new protein and associated functionality in a cell or
organism.

25 Some nucleic acids, such as miRNA inhibitors, can be used to effect
expression of specific cellular products that are regulated by miRNA as would be useful
in the treatment of, for example, diseases related to deficiency of protein or enzyme.
The therapeutic applications of miRNA inhibition are extremely broad as constructs can

be synthesized to inhibit one or more miRNA that would in turn regulate the expression of mRNA products. The inhibition of endogenous miRNA can augment its downstream target endogenous protein expression and restore proper function in a cell or organism as a means to treat disease associated to a specific miRNA or a group of miRNA.

5 Other nucleic acids can down-regulate intracellular levels of specific mRNA and, as a result, down-regulate the synthesis of the corresponding proteins through processes such as RNA interference (RNAi) or complementary binding of antisense RNA. The therapeutic applications of antisense oligonucleotide and RNAi are also extremely broad, since oligonucleotide constructs can be synthesized with any
10 nucleotide sequence directed against a target mRNA. Targets may include mRNAs from normal cells, mRNAs associated with disease-states, such as cancer, and mRNAs of infectious agents, such as viruses. To date, antisense oligonucleotide constructs have shown the ability to specifically down-regulate target proteins through degradation of the cognate mRNA in both *in vitro* and *in vivo* models. In addition, antisense
15 oligonucleotide constructs are currently being evaluated in clinical studies.


 However, two problems currently face the use of oligonucleotides in therapeutic contexts. First, free RNAs are susceptible to nuclease digestion in plasma. Second, free RNAs have limited ability to gain access to the intracellular compartment where the relevant translation machinery resides. Lipid nanoparticles formed from
20 cationic lipids with other lipid components, such as neutral lipids, cholesterol, PEG, PEGylated lipids, and oligonucleotides have been used to block degradation of the RNAs in plasma and facilitate the cellular uptake of the oligonucleotides.

 There remains a need for improved cationic lipids and lipid nanoparticles for the delivery of oligonucleotides. Preferably, these lipid nanoparticles would provide
25 optimal drug:lipid ratios, protect the nucleic acid from degradation and clearance in serum, be suitable for systemic or local delivery, and provide intracellular delivery of the nucleic acid. In addition, these lipid-nucleic acid particles should be well-tolerated and provide an adequate therapeutic index, such that patient treatment at an effective dose of the nucleic acid is not associated with unacceptable toxicity and/or risk to the
30 patient. The present disclosure provides these and related advantages.

BRIEF SUMMARY

In brief, the present disclosure provides lipid compounds, including stereoisomers, pharmaceutically acceptable salts or tautomers thereof, which can be used alone or in combination with other lipid components such as neutral lipids, charged lipids, steroids (including for example, all sterols) and/or their analogs, and/or polymer conjugated lipids to form lipid nanoparticles for the delivery of therapeutic agents. In some instances, the lipid nanoparticles are used to deliver nucleic acids such as antisense and/or messenger RNA. Methods for use of such lipid nanoparticles for treatment of various diseases or conditions, such as those caused by infectious entities and/or insufficiency of a protein, are also provided.

In one embodiment, compounds having the following structure (I) are provided:



(I)

or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein R^1 , R^2 , R^3 , R^4 , R^5 , L^1 , L^2 , L^3 , G^1 , G^2 , and G^3 are as defined herein.

Pharmaceutical compositions comprising one or more of the foregoing compounds of structure (I) and a therapeutic agent are also provided. In some embodiments, the pharmaceutical compositions further comprise one or more components selected from neutral lipids, charged lipids, steroids, and polymer conjugated lipids. Such compositions are useful for formation of lipid nanoparticles for the delivery of the therapeutic agent.

In other embodiments, the present disclosure provides a method for administering a therapeutic agent to a patient in need thereof, the method comprising preparing or providing a composition of lipid nanoparticles comprising the compound

of structure (I) and a therapeutic agent and delivering or administering the composition to the patient.

These and other aspects of the disclosure will be apparent upon reference to the following detailed description.

5 DETAILED DESCRIPTION

In the following description, certain specific details are set forth in order to provide a thorough understanding of various embodiments of the disclosure. However, one skilled in the art will understand that the disclosure may be practiced without these details.

10 The present disclosure is based, in part, upon the discovery of novel cationic (amino) lipids that provide advantages when used in lipid nanoparticles for the *in vivo* delivery of an active or therapeutic agent such as a nucleic acid into a cell of a mammal. In particular, embodiments of the present disclosure provide nucleic acid-lipid nanoparticle compositions comprising one or more of the novel cationic lipids
15 described herein that provide increased activity of the nucleic acid and improved tolerability of the compositions *in vivo*, resulting in a significant increase in the therapeutic index as compared to nucleic acid-lipid nanoparticle compositions previously described. In other embodiments, the disclosed lipids, and lipid nanoparticles comprising the same, have increased safety and/or tolerability when used
20 for delivery of active agents, such as nucleic acids.

In particular embodiments, the present disclosure provides novel cationic lipids that enable the formulation of improved compositions for the *in vitro* and *in vivo* delivery of mRNA and/or other oligonucleotides. In some embodiments, these improved lipid nanoparticle compositions are useful for expression of protein encoded
25 by mRNA. In other embodiments, these improved lipid nanoparticles compositions are useful for upregulation of endogenous protein expression by delivering miRNA inhibitors targeting one specific miRNA or a group of miRNA regulating one target mRNA or several mRNA. In other embodiments, these improved lipid nanoparticle compositions are useful for down-regulating (*e.g.*, silencing) the protein levels and/or
30 mRNA levels of target genes. In some other embodiments, the lipid nanoparticles are

also useful for delivery of mRNA and plasmids for expression of transgenes. In yet other embodiments, the lipid nanoparticle compositions are useful for inducing a pharmacological effect resulting from expression of a protein, *e.g.*, increased production of red blood cells through the delivery of a suitable erythropoietin mRNA, or protection
5 against infection through delivery of mRNA encoding for a suitable antigen or antibody.

The lipid nanoparticles and compositions of embodiments of the present disclosure may be used for a variety of purposes, including the delivery of encapsulated or associated (*e.g.*, complexed) therapeutic agents such as nucleic acids to cells, both *in vitro* and *in vivo*. Accordingly, embodiments of the present disclosure provide methods
10 of treating or preventing diseases or disorders in a subject in need thereof by contacting the subject with a lipid nanoparticle that encapsulates or is associated with a suitable therapeutic agent, wherein the lipid nanoparticle comprises one or more of the novel cationic lipids described herein.

As described herein, embodiments of the lipid nanoparticles of the present disclosure are particularly useful for the delivery of nucleic acids, including, *e.g.*, mRNA, antisense oligonucleotide, plasmid DNA, microRNA (miRNA), miRNA inhibitors (antagomirs/antimirs), messenger-RNA-interfering complementary RNA (micRNA), DNA, multivalent RNA, dicer substrate RNA, complementary DNA
15 (cDNA), etc. Therefore, the lipid nanoparticles and compositions of certain embodiments of the present disclosure may be used to induce expression of a desired protein both *in vitro* and *in vivo* by contacting cells with a lipid nanoparticle comprising one or more novel cationic lipids described herein, wherein the lipid nanoparticle encapsulates or is associated with a nucleic acid that is expressed to produce the desired
20 protein (*e.g.*, a messenger RNA or plasmid encoding the desired protein) or inhibit processes that terminate expression of mRNA (*e.g.*, miRNA inhibitors). Alternatively, the lipid nanoparticles and compositions of embodiments of the present disclosure may be used to decrease the expression of target genes and proteins both *in vitro* and *in vivo* by contacting cells with a lipid nanoparticle comprising one or more novel cationic
25 lipids described herein, wherein the lipid nanoparticle encapsulates or is associated with a nucleic acid that reduces target gene expression (*e.g.*, an antisense oligonucleotide or

small interfering RNA (siRNA)). The lipid nanoparticles and compositions of embodiments of the present disclosure may also be used for co-delivery of different nucleic acids (*e.g.* mRNA and plasmid DNA) separately or in combination, such as may be useful to provide an effect requiring co-localization of different nucleic acids (*e.g.* mRNA encoding for a suitable gene modifying enzyme and DNA segment(s) for incorporation into the host genome).

Nucleic acids for use with embodiments of this disclosure may be prepared according to any available technique. For mRNA, the primary methodology of preparation is, but not limited to, enzymatic synthesis (also termed *in vitro* transcription) which currently represents the most efficient method to produce long sequence-specific mRNA. *In vitro* transcription describes a process of template-directed synthesis of RNA molecules from an engineered DNA template comprised of an upstream bacteriophage promoter sequence (*e.g.*, including but not limited to that from the T7, T3 and SP6 coliphage) linked to a downstream sequence encoding the gene of interest. Template DNA can be prepared for *in vitro* transcription from a number of sources with appropriate techniques which are well known in the art including, but not limited to, plasmid DNA and polymerase chain reaction amplification (see Linpinsel, J.L and Conn, G.L., General protocols for preparation of plasmid DNA template and Bowman, J.C., Azizi, B., Lenz, T.K., Ray, P., and Williams, L.D. in RNA *in vitro* transcription and RNA purification by denaturing PAGE in Recombinant and *in vitro* RNA syntheses Methods v. 941 Conn G.L. (ed), New York, N.Y. Humana Press, 2012).

Transcription of the RNA occurs *in vitro* using the linearized DNA template in the presence of the corresponding RNA polymerase and adenosine, guanosine, uridine, and cytidine ribonucleoside triphosphates (rNTPs) under conditions that support polymerase activity while minimizing potential degradation of the resultant mRNA transcripts. *In vitro* transcription can be performed using a variety of commercially available kits including, but not limited to RiboMax Large Scale RNA Production System (Promega), MegaScript Transcription kits (Life Technologies), as well as with commercially available reagents including RNA polymerases and rNTPs. The methodology for *in vitro* transcription of mRNA is well known in the art. (see, *e.g.*

Losick, R., 1972, In vitro transcription, Ann Rev Biochem v.41 409-46; Kamakaka, R. T. and Kraus, W. L. 2001. In Vitro Transcription. Current Protocols in Cell Biology. 2:11.6:11.6.1–11.6.17; Beckert, B. And Masquida, B.,(2010) Synthesis of RNA by In Vitro Transcription in RNA in Methods in Molecular Biology v. 703 (Neilson, H. Ed), New York, N.Y. Humana Press, 2010; Brunelle, J.L. and Green, R., 2013, Chapter Five – In vitro transcription from plasmid or PCR-amplified DNA, Methods in Enzymology v. 530, 101-114; all of which are incorporated herein by reference).

The desired *in vitro* transcribed mRNA is then purified from the undesired components of the transcription or associated reactions (including unincorporated rNTPs, protein enzyme, salts, short RNA oligos, etc.). Techniques for the isolation of the mRNA transcripts are well known in the art. Well known procedures include phenol/chloroform extraction or precipitation with either alcohol (ethanol, isopropanol) in the presence of monovalent cations or lithium chloride. Additional, non-limiting examples of purification procedures which can be used include size exclusion chromatography (Lukavsky, P.J. and Puglisi, J.D., 2004, Large-scale preparation and purification of polyacrylamide-free RNA oligonucleotides, RNA v.10, 889-893), silica-based affinity chromatography and polyacrylamide gel electrophoresis (Bowman, J.C., Azizi, B., Lenz, T.K., Ray, P., and Williams, L.D. in RNA in vitro transcription and RNA purification by denaturing PAGE in Recombinant and in vitro RNA syntheses Methods v. 941 Conn G.L. (ed), New York, N.Y. Humana Press, 2012). Purification can be performed using a variety of commercially available kits including, but not limited to SV Total Isolation System (Promega) and In Vitro Transcription Cleanup and Concentration Kit (Norgen Biotek).

Furthermore, while reverse transcription can yield large quantities of mRNA, the products can contain a number of aberrant RNA impurities associated with undesired polymerase activity which may need to be removed from the full-length mRNA preparation. These include short RNAs that result from abortive transcription initiation as well as double-stranded RNA (dsRNA) generated by RNA-dependent RNA polymerase activity, RNA-primed transcription from RNA templates and self-complementary 3' extension. It has been demonstrated that these contaminants with dsRNA structures can lead to undesired immunostimulatory activity through interaction

with various innate immune sensors in eukaryotic cells that function to recognize specific nucleic acid structures and induce potent immune responses. This in turn, can dramatically reduce mRNA translation since protein synthesis is reduced during the innate cellular immune response. Therefore, additional techniques to remove these dsRNA contaminants have been developed and are known in the art including but not limited to scaleable HPLC purification (see, *e.g.*, Kariko, K., Muramatsu, H., Ludwig, J. And Weissman, D., 2011, Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA, Nucl Acid Res, v. 39 e142; Weissman, D., Pardi, N., Muramatsu, H., and Kariko, K., HPLC Purification of in vitro transcribed long RNA in Synthetic Messenger RNA and Cell Metabolism Modulation in Methods in Molecular Biology v.969 (Rabinovich, P.H. Ed), 2013). HPLC purified mRNA has been reported to be translated at much greater levels, particularly in primary cells and *in vivo*.

A significant variety of modifications have been described in the art which are used to alter specific properties of *in vitro* transcribed mRNA, and improve its utility. These include, but are not limited to modifications to the 5' and 3' termini of the mRNA. Endogenous eukaryotic mRNA typically contain a cap structure on the 5'-end of a mature molecule which plays an important role in mediating binding of the mRNA Cap Binding Protein (CBP), which is in turn responsible for enhancing mRNA stability in the cell and efficiency of mRNA translation. Therefore, highest levels of protein expression are achieved with capped mRNA transcripts. The 5'-cap contains a 5'-5'-triphosphate linkage between the 5'-most nucleotide and guanine nucleotide. The conjugated guanine nucleotide is methylated at the N7 position. Additional modifications include methylation of the ultimate and penultimate most 5'-nucleotides on the 2'-hydroxyl group.

Multiple distinct cap structures can be used to generate the 5'-cap of *in vitro* transcribed synthetic mRNA. 5'-capping of synthetic mRNA can be performed co-transcriptionally with chemical cap analogs (*i.e.*, capping during *in vitro* transcription). For example, the Anti-Reverse Cap Analog (ARCA) cap contains a 5'-5'-triphosphate guanine-guanine linkage where one guanine contains an N7 methyl group as well as a

3'-O-methyl group. However, up to 20% of transcripts remain uncapped during this co-transcriptional process and the synthetic cap analog is not identical to the 5'-cap structure of an authentic cellular mRNA, potentially reducing translatability and cellular stability. Alternatively, synthetic mRNA molecules may also be enzymatically capped post-transcriptionally. These may generate a more authentic 5'-cap structure that more closely mimics, either structurally or functionally, the endogenous 5'-cap which have enhanced binding of cap binding proteins, increased half-life and reduced susceptibility to 5' endonucleases and/or reduced 5' decapping. Numerous synthetic 5'-cap analogs have been developed and are known in the art to enhance mRNA stability and translatability (see, *e.g.*, Grudzien-Nogalska, E., Kowalska, J., Su, W., Kuhn, A.N., Slepnev, S.V., Darynkiewicz, E., Sahin, U., Jemielity, J., and Rhoads, R.E., Synthetic mRNAs with superior translation and stability properties in Synthetic Messenger RNA and Cell Metabolism Modulation in Methods in Molecular Biology v.969 (Rabinovich, P.H. Ed), 2013).

On the 3'-terminus, a long chain of adenine nucleotides (poly-A tail) is normally added to mRNA molecules during RNA processing. Immediately after transcription, the 3' end of the transcript is cleaved to free a 3' hydroxyl to which poly-A polymerase adds a chain of adenine nucleotides to the RNA in a process called polyadenylation. The poly-A tail has been extensively shown to enhance both translational efficiency and stability of mRNA (see Bernstein, P. and Ross, J., 1989, Poly (A), poly (A) binding protein and the regulation of mRNA stability, Trends Bio Sci v. 14 373-377; Guhaniyogi, J. And Brewer, G., 2001, Regulation of mRNA stability in mammalian cells, Gene, v. 265, 11-23; Dreyfus, M. And Regnier, P., 2002, The poly (A) tail of mRNAs: Bodyguard in eukaryotes, scavenger in bacteria, Cell, v.111, 611-613).

Poly (A) tailing of *in vitro* transcribed mRNA can be achieved using various approaches including, but not limited to, cloning of a poly (T) tract into the DNA template or by post-transcriptional addition using Poly (A) polymerase. The first case allows *in vitro* transcription of mRNA with poly (A) tails of defined length, depending on the size of the poly (T) tract, but requires additional manipulation of the template. The latter case involves the enzymatic addition of a poly (A) tail to *in vitro*

transcribed mRNA using poly (A) polymerase which catalyzes the incorporation of adenine residues onto the 3' termini of RNA, requiring no additional manipulation of the DNA template, but results in mRNA with poly(A) tails of heterogeneous length. 5'-capping and 3'-poly (A) tailing can be performed using a variety of commercially available kits including, but not limited to Poly (A) Polymerase Tailing kit (EpiCenter), mMESSAGE mMACHINE T7 Ultra kit and Poly (A) Tailing kit (Life Technologies) as well as with commercially available reagents, various ARCA caps, Poly (A) polymerase, etc.

In addition to 5' cap and 3' poly adenylation, other modifications of the *in vitro* transcripts have been reported to provide benefits as related to efficiency of translation and stability. It is well known in the art that pathogenic DNA and RNA can be recognized by a variety of sensors within eukaryotes and trigger potent innate immune responses. The ability to discriminate between pathogenic and self DNA and RNA has been shown to be based, at least in part, on structure and nucleoside modifications since most nucleic acids from natural sources contain modified nucleosides. In contrast, *in vitro* synthesized RNA lacks these modifications, thus rendering it immunostimulatory which in turn can inhibit effective mRNA translation as outlined above. The introduction of modified nucleosides into *in vitro* transcribed mRNA can be used to prevent recognition and activation of RNA sensors, thus mitigating this undesired immunostimulatory activity and enhancing translation capacity (see, e.g., Kariko, K. And Weissman, D. 2007, Naturally occurring nucleoside modifications suppress the immunostimulatory activity of RNA: implication for therapeutic RNA development, Curr Opin Drug Discov Devel, v.10 523-532; Pardi, N., Muramatsu, H., Weissman, D., Kariko, K., In vitro transcription of long RNA containing modified nucleosides in Synthetic Messenger RNA and Cell Metabolism Modulation in Methods in Molecular Biology v.969 (Rabinovich, P.H. Ed), 2013; Kariko, K., Muramatsu, H., Welsh, F.A., Ludwig, J., Kato, H., Akira, S., Weissman, D., 2008, Incorporation of Pseudouridine Into mRNA Yields Superior Nonimmunogenic Vector With Increased Translational Capacity and Biological Stability, Mol Ther v.16, 1833-1840). The modified nucleosides and nucleotides used in the synthesis of modified RNAs can be prepared monitored and utilized using general methods and

procedures known in the art. A large variety of nucleoside modifications are available that may be incorporated alone or in combination with other modified nucleosides to some extent into the *in vitro* transcribed mRNA (see, *e.g.*, US 2012/0251618). *In vitro* synthesis of nucleoside-modified mRNA has been reported to have reduced ability to
5 activate immune sensors with a concomitant enhanced translational capacity.

Other components of mRNA which can be modified to provide benefit in terms of translatability and stability include the 5' and 3' untranslated regions (UTR). Optimization of the UTRs (favorable 5' and 3' UTRs can be obtained from cellular or viral RNAs), either both or independently, have been shown to increase mRNA stability
10 and translational efficiency of *in vitro* transcribed mRNA (see, *e.g.*, Pardi, N., Muramatsu, H., Weissman, D., Kariko, K., *In vitro* transcription of long RNA containing modified nucleosides in Synthetic Messenger RNA and Cell Metabolism Modulation in Methods in Molecular Biology v.969 (Rabinovich, P.H. Ed), 2013).

In addition to mRNA, other nucleic acid payloads may be used for this
15 disclosure. For oligonucleotides, methods of preparation include but are not limited to chemical synthesis and enzymatic, chemical cleavage of a longer precursor, *in vitro* transcription as described above, etc. Methods of synthesizing DNA and RNA nucleotides are widely used and well known in the art (see, *e.g.*, Gait, M. J. (ed.) *Oligonucleotide synthesis: a practical approach*, Oxford [Oxfordshire], Washington,
20 D.C.: IRL Press, 1984; and Herdewijn, P. (ed.) *Oligonucleotide synthesis: methods and applications*, Methods in Molecular Biology, v. 288 (Clifton, N.J.) Totowa, N.J.: Humana Press, 2005; both of which are incorporated herein by reference).

For plasmid DNA, preparation for use with embodiments of this disclosure commonly utilizes, but is not limited to, expansion and isolation of the
25 plasmid DNA *in vitro* in a liquid culture of bacteria containing the plasmid of interest. The presence of a gene in the plasmid of interest that encodes resistance to a particular antibiotic (penicillin, kanamycin, etc.) allows those bacteria containing the plasmid of interest to selectively grow in antibiotic-containing cultures. Methods of isolating plasmid DNA are widely used and well known in the art (see, *e.g.*, Heilig, J., Elbing, K.
30 L. and Brent, R., (2001), *Large-Scale Preparation of Plasmid DNA*, Current Protocols in Molecular Biology, 41:II:1.7:1.7.1–1.7.16; Rozkov, A., Larsson, B., Gillström, S.,

Björnstedt, R. and Schmidt, S. R., (2008), Large-scale production of endotoxin-free plasmids for transient expression in mammalian cell culture, *Biotechnol. Bioeng.*, 99: 557–566; and US 6,197,553 B1). Plasmid isolation can be performed using a variety of commercially available kits including, but not limited to Plasmid Plus (Qiagen),
5 GenJET plasmid MaxiPrep (Thermo), and PureYield MaxiPrep (Promega) kits as well as with commercially available reagents.

Various exemplary embodiments of the cationic lipids of the present disclosure, lipid nanoparticles and compositions comprising the same, and their use to deliver active (*e.g.*, therapeutic agents), such as nucleic acids, to modulate gene and
10 protein expression, are described in further detail below.

As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

Unless the context requires otherwise, throughout the present specification and claims, the word "comprise" and variations thereof, such as,
15 "comprises" and "comprising" are to be construed in an open and inclusive sense, that is, as "including, but not limited to".

Reference throughout this specification to "one embodiment" or "an embodiment" means that a particular feature, structure or characteristic described in connection with the embodiment is included in at least one embodiment of the present
20 disclosure. Thus, the appearances of the phrases "in one embodiment" or "in an embodiment" in various places throughout this specification are not necessarily all referring to the same embodiment. Furthermore, the particular features, structures, or characteristics may be combined in any suitable manner in one or more embodiments.

Unless defined otherwise, all technical and scientific terms used herein
25 have the same meaning as is commonly understood by one of skill in the art to which this disclosure belongs. As used in the specification and claims, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise.

The phrase "induce expression of a desired protein" refers to the ability of a nucleic acid to increase expression of the desired protein. To examine the extent of
30 protein expression, a test sample (*e.g.*, a sample of cells in culture expressing the desired protein) or a test mammal (*e.g.*, a mammal such as a human or an animal)

model such as a rodent (*e.g.*, mouse) or a non-human primate (*e.g.*, monkey) model is contacted with a nucleic acid (*e.g.*, nucleic acid in combination with a lipid of the present disclosure). Expression of the desired protein in the test sample or test animal is compared to expression of the desired protein in a control sample (*e.g.*, a sample of
5 cells in culture expressing the desired protein) or a control mammal (*e.g.*, a mammal such as a human or an animal) model such as a rodent (*e.g.*, mouse) or non-human primate (*e.g.*, monkey) model that is not contacted with or administered the nucleic acid. When the desired protein is present in a control sample or a control mammal, the expression of a desired protein in a control sample or a control mammal may be
10 assigned a value of 1.0. In particular embodiments, inducing expression of a desired protein is achieved when the ratio of desired protein expression in the test sample or the test mammal to the level of desired protein expression in the control sample or the control mammal is greater than 1, for example, about 1.1, 1.5, 2.0, 5.0 or 10.0. When a desired protein is not present in a control sample or a control mammal, inducing
15 expression of a desired protein is achieved when any measurable level of the desired protein in the test sample or the test mammal is detected. One of ordinary skill in the art will understand appropriate assays to determine the level of protein expression in a sample, for example dot blots, northern blots, in situ hybridization, ELISA, immunoprecipitation, enzyme function, and phenotypic assays, or assays based on
20 reporter proteins that can produce fluorescence or luminescence under appropriate conditions.

The phrase "inhibiting expression of a target gene" refers to the ability of a nucleic acid to silence, reduce, or inhibit the expression of a target gene. To examine the extent of gene silencing, a test sample (*e.g.*, a sample of cells in culture expressing
25 the target gene) or a test mammal (*e.g.*, a mammal such as a human or an animal) model such as a rodent (*e.g.*, mouse) or a non-human primate (*e.g.*, monkey) model is contacted with a nucleic acid that silences, reduces, or inhibits expression of the target gene. Expression of the target gene in the test sample or test animal is compared to expression of the target gene in a control sample (*e.g.*, a sample of cells in culture
30 expressing the target gene) or a control mammal (*e.g.*, a mammal such as a human or an animal) model such as a rodent (*e.g.*, mouse) or non-human primate (*e.g.*, monkey)

model that is not contacted with or administered the nucleic acid. The expression of the target gene in a control sample or a control mammal may be assigned a value of 100%. In particular embodiments, silencing, inhibition, or reduction of expression of a target gene is achieved when the level of target gene expression in the test sample or the test mammal relative to the level of target gene expression in the control sample or the control mammal is about 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5%, or 0%. In other words, the nucleic acids are capable of silencing, reducing, or inhibiting the expression of a target gene by at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% in a test sample or a test mammal relative to the level of target gene expression in a control sample or a control mammal not contacted with or administered the nucleic acid. Suitable assays for determining the level of target gene expression include, without limitation, examination of protein or mRNA levels using techniques known to those of skill in the art, such as, *e.g.*, dot blots, northern blots, *in situ* hybridization, ELISA, immunoprecipitation, enzyme function, as well as phenotypic assays known to those of skill in the art.

An "effective amount" or "therapeutically effective amount" of an active agent or therapeutic agent such as a therapeutic nucleic acid is an amount sufficient to produce the desired effect, *e.g.*, an increase or inhibition of expression of a target sequence in comparison to the normal expression level detected in the absence of the nucleic acid. An increase in expression of a target sequence is achieved when any measurable level is detected in the case of an expression product that is not present in the absence of the nucleic acid. In the case where the expression product is present at some level prior to contact with the nucleic acid, an increase in expression is achieved when the fold increase in value obtained with a nucleic acid such as mRNA relative to control is about 1.05, 1.1, 1.2, 1.3, 1.4, 1.5, 1.75, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 75, 100, 250, 500, 750, 1000, 5000, 10000, or greater. Inhibition of expression of a target gene or target sequence is achieved when the value obtained with a nucleic acid such as antisense oligonucleotide relative to the control is about 95%, 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5%, or 0%. Suitable assays for measuring expression of a target gene

or target sequence include, *e.g.*, examination of protein or RNA levels using techniques known to those of skill in the art such as dot blots, northern blots, *in situ* hybridization, ELISA, immunoprecipitation, enzyme function, fluorescence, or luminescence of suitable reporter proteins, as well as phenotypic assays known to those of skill in the art.

The term "nucleic acid" as used herein refers to a polymer containing at least two deoxyribonucleotides or ribonucleotides in either single- or double-stranded form and includes DNA, RNA, and hybrids thereof. DNA may be in the form of antisense molecules, plasmid DNA, cDNA, PCR products, or vectors. RNA may be in the form of small hairpin RNA (shRNA), messenger RNA (mRNA), antisense RNA, miRNA, micRNA, multivalent RNA, dicer substrate RNA or viral RNA (vRNA), and combinations thereof. Nucleic acids include nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, and which have similar binding properties as the reference nucleic acid. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2'-O-methyl ribonucleotides, and peptide-nucleic acids (PNAs). Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides that have similar binding properties as the reference nucleic acid.

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions), alleles, orthologs, single nucleotide polymorphisms, and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer et al., *Nucleic Acid Res.*, 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.*, 260:2605-2608 (1985); Rossolini et al., *Mol. Cell. Probes*, 8:91-98 (1994)).

"Nucleotides" contain a sugar deoxyribose (DNA) or ribose (RNA), a base, and a phosphate group. Nucleotides are linked together through the phosphate groups.

"Bases" include purines and pyrimidines, which further include natural compounds adenine, thymine, guanine, cytosine, uracil, inosine, and natural analogs, and synthetic


derivatives of purines and pyrimidines, which include, but are not limited to, modifications which place new reactive groups such as, but not limited to, amines, alcohols, thiols, carboxylates, and alkylhalides.

The term "gene" refers to a nucleic acid (*e.g.*, DNA or RNA) sequence
5 that comprises partial length or entire length coding sequences necessary for the production of a polypeptide or precursor polypeptide.

"Gene product," as used herein, refers to a product of a gene such as an RNA transcript or a polypeptide.

The term "lipid" refers to a group of organic compounds that include, but
10 are not limited to, esters of fatty acids and are generally characterized by being poorly soluble in water, but soluble in many organic solvents. They are usually divided into at least three classes: (1) "simple lipids," which include fats and oils as well as waxes; (2) "compound lipids," which include phospholipids and glycolipids; and (3) "derived lipids" such as steroids.

15 A "steroid" is a compound comprising the following carbon skeleton:



Non-limiting examples of steroids include cholesterol, and the like.

A "cationic lipid" refers to a lipid capable of being positively charged. Exemplary cationic lipids include one or more amine group(s) which bear the positive
20 charge. Preferred cationic lipids are ionizable such that they can exist in a positively charged or neutral form depending on pH. The ionization of the cationic lipid affects the surface charge of the lipid nanoparticle under different pH conditions. This charge state can influence plasma protein absorption, blood clearance, and tissue distribution (Semple, S.C., et al., Adv. Drug Deliv Rev 32:3-17 (1998)) as well as the ability to
25 form endosomolytic non-bilayer structures (Hafez, I.M., et al., Gene Ther 8:1188-1196 (2001)) critical to the intracellular delivery of nucleic acids.

The term "polymer conjugated lipid" refers to a molecule comprising both a lipid portion and a polymer portion. An example of a polymer conjugated lipid

is a pegylated lipid. The term "pegylated lipid" refers to a molecule comprising both a lipid portion and a polyethylene glycol portion. Pegylated lipids are known in the art and include 1-(monomethoxy-polyethyleneglycol)-2,3-dimyristoylglycerol (PEG-DMG) and the like.

5 The term "neutral lipid" refers to any of a number of lipid species that exist either in an uncharged or neutral zwitterionic form at a selected pH. At physiological pH, such lipids include, but are not limited to, phosphatidylcholines such as 1,2-Distearoyl-*sn*-glycero-3-phosphocholine (DSPC), 1,2-Dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC), 1,2-Dimyristoyl-*sn*-glycero-3-phosphocholine (DMPC), 1-
10 Palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC), 1,2-dioleoyl-*sn*-glycero-3-phosphocholine (DOPC), phosphatidylethanolamines such as 1,2-Dioleoyl-*sn*-glycero-3-phosphoethanolamine (DOPE), sphingomyelins (SM), ceramides, steroids such as sterols and their derivatives. Neutral lipids may be synthetic or naturally derived.

 The term "charged lipid" refers to any of a number of lipid species that
15 exist in either a positively charged or negatively charged form independent of the pH within a useful physiological range, *e.g.*, pH ~3 to pH ~9. Charged lipids may be synthetic or naturally derived. Examples of charged lipids include phosphatidylserines, phosphatidic acids, phosphatidylglycerols, phosphatidylinositols, sterol hemisuccinates, dialkyl trimethylammonium-propanes, (*e.g.*, DOTAP, DOTMA), dialkyl
20 dimethylaminopropanes, ethyl phosphocholines, dimethylaminoethane carbamoyl sterols (*e.g.*, DC-Chol).

 The term "lipid nanoparticle" refers to particles having at least one dimension on the order of nanometers (*e.g.*, 1-1,000 nm) which include one or more of the compounds of structure (I) or other specified cationic lipids. In some embodiments,
25 lipid nanoparticles comprising the disclosed cationic lipids (*e.g.*, compounds of structure (I)) are included in a formulation that can be used to deliver an active agent or therapeutic agent, such as a nucleic acid (*e.g.*, mRNA) to a target site of interest (*e.g.*, cell, tissue, organ, tumor, and the like). In some embodiments, the lipid nanoparticles comprise a compound of structure (I) and a nucleic acid. Such lipid nanoparticles
30 typically comprise a compound of structure (I) and one or more excipient selected from neutral lipids, charged lipids, steroids, and polymer conjugated lipids. In some

embodiments, the active agent or therapeutic agent, such as a nucleic acid, may be encapsulated in the lipid portion of the lipid nanoparticle or an aqueous space enveloped by some or all of the lipid portion of the lipid nanoparticle, thereby protecting it from enzymatic degradation or other undesirable effects induced by the mechanisms of the host organism or cells, *e.g.*, an adverse immune response.

In various embodiments, the lipid nanoparticles have a mean diameter of from about 30 nm to about 150 nm, from about 40 nm to about 150 nm, from about 50 nm to about 150 nm, from about 60 nm to about 130 nm, from about 70 nm to about 110 nm, from about 70 nm to about 100 nm, from about 80 nm to about 100 nm, from about 90 nm to about 100 nm, from about 70 to about 90 nm, from about 80 nm to about 90 nm, from about 70 nm to about 80 nm, or about 30 nm, 35 nm, 40 nm, 45 nm, 50 nm, 55 nm, 60 nm, 65 nm, 70 nm, 75 nm, 80 nm, 85 nm, 90 nm, 95 nm, 100 nm, 105 nm, 110 nm, 115 nm, 120 nm, 125 nm, 130 nm, 135 nm, 140 nm, 145 nm, or 150 nm. In some embodiments, the lipid nanoparticles are substantially non-toxic. In certain embodiments, nucleic acids, when present in the lipid nanoparticles, are resistant in aqueous solution to degradation with a nuclease. Lipid nanoparticles comprising nucleic acids and their method of preparation are disclosed in, *e.g.*, U.S. Patent Pub. Nos. 2004/0142025, 2007/0042031 and PCT Pub. Nos. WO 2013/016058 and WO 2013/086373, the full disclosures of which are herein incorporated by reference in their entirety for all purposes.

As used herein, "lipid encapsulated" refers to a lipid nanoparticle that provides an active agent or therapeutic agent, such as a nucleic acid (*e.g.*, mRNA), with full encapsulation, partial encapsulation, or both. In an embodiment, the nucleic acid (*e.g.*, mRNA) is fully encapsulated in the lipid nanoparticle.

As used herein, the term "aqueous solution" refers to a composition comprising water.

"Serum-stable" in relation to nucleic acid-lipid nanoparticles means that the nucleotide is not significantly degraded after exposure to a serum or nuclease assay that would significantly degrade free DNA or RNA. Suitable assays include, for example, a standard serum assay, a DNase assay, or an RNase assay.

"Systemic delivery," as used herein, refers to delivery of a therapeutic product that can result in a broad exposure of an active agent within an organism.

Some techniques of administration can lead to the systemic delivery of certain agents, but not others. Systemic delivery means that a useful, preferably therapeutic, amount of an agent is exposed to most parts of the body. Systemic delivery of lipid nanoparticles can be by any means known in the art including, for example, intravenous, intraarterial, subcutaneous, and intraperitoneal delivery. In some embodiments, systemic delivery of lipid nanoparticles is by intravenous delivery.

"Local delivery," as used herein, refers to delivery of an active agent directly to a target site within an organism. For example, an agent can be locally delivered by direct injection into a disease site such as a tumor, other target site such as a site of inflammation, or a target organ such as the liver, heart, pancreas, kidney, and the like. Local delivery can also include topical applications or localized injection techniques such as intramuscular, subcutaneous, or intradermal injection. Local delivery does not preclude a systemic pharmacological effect.

"Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, which is saturated or unsaturated (*i.e.*, contains one or more double (alkenyl) and/or triple bonds (alkynyl)), having, for example, from one to twenty-four carbon atoms (C₁-C₂₄ alkyl), four to twenty carbon atoms (C₄-C₂₀ alkyl), six to sixteen carbon atoms (C₆-C₁₆ alkyl), six to nine carbon atoms (C₆-C₉ alkyl), one to fifteen carbon atoms (C₁-C₁₅ alkyl), one to twelve carbon atoms (C₁-C₁₂ alkyl), one to eight carbon atoms (C₁-C₈ alkyl) or one to six carbon atoms (C₁-C₆ alkyl) and which is attached to the rest of the molecule by a single bond, *e.g.*, methyl, ethyl, *n*-propyl, 1-methylethyl (iso propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl (*t*-butyl), 3-methylhexyl, 2-methylhexyl, ethenyl, prop-1-enyl, but-1-enyl, pent-1-enyl, penta-1,4-dienyl, ethynyl, propynyl, butynyl, pentynyl, hexynyl, and the like. Unless stated otherwise specifically in the specification, an alkyl group is optionally substituted.

"Alkylene" or "alkylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, which is saturated or unsaturated (*i.e.*, contains one or more

double (alkenylene) and/or triple bonds (alkynylene)), and having, for example, from one to twenty-four carbon atoms (C₁-C₂₄ alkylene), one to fifteen carbon atoms (C₁-C₁₅ alkylene), one to twelve carbon atoms (C₁-C₁₂ alkylene), one to eight carbon atoms (C₁-C₈ alkylene), one to six carbon atoms (C₁-C₆ alkylene), two to four carbon atoms (C₂-C₄ alkylene), one to two carbon atoms (C₁-C₂ alkylene), *e.g.*, methylene, ethylene, propylene, *n*-butylene, ethenylene, propenylene, *n*-butenylene, propynylene, *n*-butynylene, and the like. The alkylene chain is attached to the rest of the molecule through a single or double bond and to the radical group through a single or double bond. The points of attachment of the alkylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkylene chain may be optionally substituted.

"Cycloalkyl" or "carbocyclic ring" refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which may include fused or bridged ring systems, having from three to fifteen carbon atoms, preferably having from three to ten carbon atoms, and which is saturated or unsaturated and attached to the rest of the molecule by a single bond. Monocyclic radicals include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic radicals include, for example, adamantyl, norbornyl, decalyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like. Unless otherwise stated specifically in the specification, a cycloalkyl group may be optionally substituted.

"Cycloalkylene" is a divalent cycloalkyl group. Unless otherwise stated specifically in the specification, a cycloalkylene group may be optionally substituted.

"Heterocyclyl" or "heterocyclic ring" refers to a stable 3- to 18-membered (*e.g.*, 5, 6 or 7-membered) non-aromatic ring radical having one to twelve ring carbon atoms (*e.g.*, two to twelve) and from one to six ring heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. Examples of such heterocyclyl radicals include, but are not limited to, azetidyl, pyrrolidyl, piperidyl, imidazolidyl, tetrahydropyrimidyl, and the like. Unless stated otherwise specifically in the specification, a heterocyclyl group may be optionally substituted.

"Heteroaryl" refers to a 5- to 14-membered ring system radical comprising hydrogen atoms, one to thirteen ring carbon atoms, one to six ring heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, and at least one aromatic ring. Examples include, but are not limited to, pyrrolyl, imidazolyl, triazolyl, thiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, and the like. Unless stated otherwise specifically in the specification, a heteroaryl group may be optionally substituted.

The term "substituted" used herein means any of the above groups (*e.g.*, alkyl, alkylene, cycloalkyl or cycloalkylene) wherein at least one hydrogen atom is replaced by a bond to a non-hydrogen atom such as, but not limited to: a halogen atom such as F, Cl, Br, or I; oxo groups (=O); hydroxyl groups (-OH); C₁-C₁₂ alkyl groups; cycloalkyl groups; -(C=O)OR'; -O(C=O)R'; -C(=O)R'; -OR'; -S(O)_xR'; -S-SR'; C(=O)SR'; -SC(=O)R'; -NR'R'; -NR'C(=O)R'; -C(=O)NR'R'; -NR'C(=O)NR'R'; -OC(=O)NR'R'; -NR'C(=O)OR'; -NR'S(O)_xNR'R'; -NR'S(O)_xR'; and -S(O)_xNR'R', wherein: R' is, at each occurrence, independently H, C₁-C₁₅ alkyl, or cycloalkyl, and x is 0, 1 or 2. In some embodiments the substituent is a C₁-C₁₂ alkyl group. In other embodiments, the substituent is a cycloalkyl group. In other embodiments, the substituent is a halo group, such as fluoro. In other embodiments, the substituent is an oxo group. In other embodiments, the substituent is a hydroxyl group. In other embodiments, the substituent is an alkoxy group (-OR'). In other embodiments, the substituent is a carboxyl group. In other embodiments, the substituent is an amine group (-NR'R').

"Optional" or "optionally" (*e.g.*, optionally substituted) means that the subsequently described event or circumstances may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted alkyl" means that the alkyl radical may or may not be substituted and that the description includes both substituted alkyl radicals and alkyl radicals having no substitution.

The disclosure disclosed herein is also meant to encompass all pharmaceutically acceptable compounds of the compound of structure (I) being isotopically-labelled by having one or more atoms replaced by an atom having a different atomic mass or mass number. Examples of isotopes that can be incorporated

into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, chlorine, and iodine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I , and ^{125}I , respectively. These radiolabeled compounds could be useful to help determine or measure the effectiveness of the compounds, by characterizing, for example, the site or mode of action, or binding affinity to pharmacologically important site of action. Certain isotopically-labelled compounds of structure (I), (IA) or (IB), for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, *i.e.*, ^3H , and carbon-14, *i.e.*, ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

Substitution with heavier isotopes such as deuterium, *i.e.*, ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances.

Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled compounds of structure (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the Preparations and Examples as set out below using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

Embodiments disclosed herein are also meant to encompass the *in vivo* metabolic products of the disclosed compounds. Such products may result from, for example, the oxidation, reduction, hydrolysis, amidation, esterification, and the like of the administered compound, primarily due to enzymatic processes. Accordingly, embodiments of the disclosure include compounds produced by a process comprising administering a compound of this disclosure to a mammal for a period of time sufficient to yield a metabolic product thereof. Such products are typically identified by administering a radiolabeled compound of the disclosure in a detectable dose to an animal, such as rat, mouse, guinea pig, monkey, or to human, allowing sufficient time

for metabolism to occur, and isolating its conversion products from the urine, blood, or other biological samples.

"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity
5 from a reaction mixture, and formulation into an efficacious therapeutic agent.

"Mammal" includes humans and both domestic animals such as laboratory animals and household pets (*e.g.*, cats, dogs, swine, cattle, sheep, goats, horses, rabbits), and non-domestic animals such as wildlife and the like.

"Pharmaceutically acceptable carrier, diluent or excipient" includes
10 without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals.

15 "Pharmaceutically acceptable salt" includes both acid and base addition salts.

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such
20 as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid,
25 cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid,
30 malic acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic

acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, *p*-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

- 5 "Pharmaceutically acceptable base addition salt" refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, 10 magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, 15 isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, 20 piperazine, piperidine, *N*-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

- Often crystallizations produce a solvate of a compound of the disclosure (*i.e.*, a compound of structure (I)). As used herein, the term "solvate" refers to an 25 aggregate that comprises one or more molecules of a compound of the disclosure with one or more molecules of solvent. The solvent may be water, in which case the solvate may be a hydrate. Alternatively, the solvent may be an organic solvent. Thus, the compounds of the present disclosure may exist as a hydrate, including a monohydrate, dihydrate, hemihydrate, sesquihydrate, trihydrate, tetrahydrate and the like, as well as 30 the corresponding solvated forms. Solvates of compound of the disclosure may be true

solvates, while in other cases the compound of the disclosure may merely retain adventitious water or be a mixture of water plus some adventitious solvent.

A "pharmaceutical composition" refers to a formulation of a compound of the disclosure and a medium generally accepted in the art for the delivery of the
5 biologically active compound to mammals, *e.g.*, humans. Such a medium includes all pharmaceutically acceptable carriers, diluents or excipients therefor.

"Effective amount" or "therapeutically effective amount" refers to that amount of a compound of the disclosure which, when administered to a mammal, preferably a human, is sufficient to effect treatment in the mammal, preferably a human.
10 The amount of a lipid nanoparticle of the disclosure which constitutes a "therapeutically effective amount" will vary depending on the compound, the condition and its severity, the manner of administration, and the age of the mammal to be treated, but can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

15 "Treating" or "treatment" as used herein covers the treatment of the disease or condition of interest in a mammal, preferably a human, having the disease or condition of interest, and includes:

(i) preventing the disease or condition from occurring in a mammal, in particular, when such mammal is predisposed to the condition but has not yet been
20 diagnosed as having it;

(ii) inhibiting the disease or condition, *i.e.*, arresting its development;

(iii) relieving the disease or condition, *i.e.*, causing regression of the disease or condition; or

(iv) relieving the symptoms resulting from the disease or condition,
25 *i.e.*, relieving pain without addressing the underlying disease or condition. As used herein, the terms "disease" and "condition" may be used interchangeably or may be different in that the particular malady or condition may not have a known causative agent (so that etiology has not yet been worked out) and it is therefore not yet recognized as a disease but only as an undesirable condition or syndrome, wherein a
30 more or less specific set of symptoms have been identified by clinicians.

The compounds of the disclosure, or their pharmaceutically acceptable salts may contain one or more stereocenters and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or, as (*D*)- or (*L*)- for amino acids. The present disclosure is meant to include all such possible isomers, as well as their racemic and optically pure forms. Optically active (+) and (-), (*R*)- and (*S*)-, or (*D*)- and (*L*)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, for example, chromatography and fractional crystallization. Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of the racemate (or the racemate of a salt or derivative) using, for example, chiral high pressure liquid chromatography (HPLC). When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both *E* and *Z* geometric isomers. Likewise, all tautomeric forms are also intended to be included.


A "stereoisomer" refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not interchangeable. The present disclosure contemplates various stereoisomers and mixtures thereof and includes "enantiomers", which refers to two stereoisomers whose molecules are non-superimposable mirror images of one another.

A "tautomer" refers to a proton shift from one atom of a molecule to another atom of the same molecule. The present disclosure includes tautomers of any said compounds.

Compounds

In an aspect, the disclosure provides novel lipid compounds which are capable of combining with other lipid components such as neutral lipids, charged lipids, steroids and/or polymer conjugated-lipids to form lipid nanoparticles with oligonucleotides. Without wishing to be bound by theory, it is thought that these lipid nanoparticles shield oligonucleotides from degradation in the serum and provide for effective delivery of oligonucleotides to cells *in vitro* and *in vivo*.

In one embodiment, the compounds have the following structure (I):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein:

5 R^1 is optionally substituted C_1 - C_{24} alkyl or optionally substituted C_2 - C_{24} alkenyl;

R^2 and R^3 are each independently optionally substituted C_1 - C_{36} alkyl;

R^4 and R^5 are each independently optionally substituted C_1 - C_6 alkyl, or
 10 R^4 and R^5 join, along with the N to which they are attached, to form a heterocyclyl or heteroaryl;


L^1 , L^2 , and L^3 are each independently optionally substituted C_1 - C_{18} alkylene;

G^1 is a direct bond, $-(CH_2)_nO(C=O)-$, $-(CH_2)_n(C=O)O-$, or $-(C=O)-$;


G^2 and G^3 are each independently $-(C=O)O-$ or $-O(C=O)-$; and

15 n is an integer greater than 0.

In some embodiments, the compound has the following structure (IA):



In some embodiments, the compound has the following structure (IB):




(IB)

In some embodiments, R¹ is optionally substituted C₆-C₁₈ alkyl or C₁₄-C₁₈ alkenyl. In certain embodiments, R¹ is C₈ alkyl, C₉ alkyl, C₁₀ alkyl, C₁₂ alkyl, C₁₄ alkyl, or C₁₆ alkyl. In some more specific embodiments, R¹ is C₁₆ alkenyl. In certain more specific embodiments, R¹ is unbranched. In some embodiments, R¹ is branched. In certain embodiments, R¹ is unsubstituted.

In some embodiments, G¹ is a direct bond, -(CH₂)_nO(C=O)-, or -(CH₂)_n(C=O)O-. In certain embodiments, G¹ is a direct bond. In some more specific embodiments, G¹ is -(CH₂)_n(C=O)O- and n is greater than 1. In some embodiments, n is 1-20. In some embodiments n is 1-10. In some embodiments n is 5-11. In some embodiments, n is 6-10. In certain more specific embodiments, n is 5, 6, 7, 8, 9, or 10. In some embodiments, n is 5. In some embodiments, n is 6. In some embodiments, n is 7. In certain embodiments, n is 8. In some embodiments, n is 9. In some embodiments, n is 10.

In some embodiments, L¹ is C₁-C₆ alkylene. In certain embodiments, L¹ is C₂ alkylene, C₃ alkylene, or C₄ alkylene. In some more specific embodiments, L¹ is unbranched. In certain more specific embodiments, L¹ is unsubstituted.


In some embodiments, R² is C₈-C₂₄ alkyl. In some embodiments, R³ is C₈-C₂₄ alkyl. In some more specific embodiments, R² and R³ are both C₈-C₂₄ alkyl. In some embodiments, R² and R³ are each independently C₁₁ alkyl, C₁₂ alkyl, C₁₃ alkyl, C₁₄ alkyl, C₁₅ alkyl, C₁₆ alkyl, C₁₈ alkyl, or C₂₀ alkyl. In certain embodiments, R² is branched. In more specific embodiments, R³ is branched. In some more specific embodiments, R² and R³ each independently have one of the following structures:



wherein:

R^6 and R^7 are each independently C_2 - C_{12} alkyl.


In some embodiments, R^2 and R^3 each independently have one of the following structures:



In some embodiments, L^2 and L^3 are each independently C_4 - C_{10} alkylene. In certain embodiments, L^2 and L^3 are both C_5 alkylene. In some more specific embodiments, L^2 and L^3 are both C_6 alkylene. In certain embodiments, L^2 and L^3 are both C_8 alkylene. In some more specific embodiments, L^2 and L^3 are both C_9 alkylene. In some embodiments, L^2 is unbranched. In some embodiments, L^3 is unbranched. In more specific embodiments, L^2 is unsubstituted. In some embodiments, L^2 is unsubstituted.






In some embodiments, R^4 and R^5 are each independently C_1 - C_6 alkyl. In more specific embodiments, R^4 and R^5 are both methyl. In certain embodiments, R^4 and R^5 are both ethyl. In certain embodiments, R^4 is methyl and R^5 is n-butyl. In some embodiments, R^4 and R^5 are both n-butyl. In different embodiments, R^4 is methyl and R^5 is n-hexyl.








In some embodiments, R^4 and R^5 join, along with the N to which they are attached, to form a heterocyclyl. In certain embodiments, the heterocyclyl is a 5-membered heterocyclyl. In some embodiments, the heterocyclyl has the following structure:










In various different embodiments, the compound has one of the structures set forth in Table 1 below.







Table 1. Compounds included in embodiments of compounds of structure (I)








No.	Structure	pKa
I-1		-
I-2		-
I-3		-
I-4		-
I-5		-


No.	Structure	pKa
I-6		-
I-7		6.74
I-8		6.68
I-9		6.83
I-10		-
I-11		-
I-12		-

No.	Structure	pKa
I-13		-
I-14		-
I-15		-
I-16		6.77
I-17		-
I-18		6.47
I-19		-

No.	Structure	pKa
I-20		6.84
I-21		-
I-22		-
I-23		-
I-24		-
I-25		6.20
I-26		-

No.	Structure	pKa
I-27		-
I-28		-
I-29		6.81
I-30		6.47
I-31		5.05
I-32		6.41

No.	Structure	pKa
I-33		6.19
I-34		-
I-35		-
I-36		-
I-37		-
I-38		-
I-39		-

No.	Structure	pKa
I-40		-

It is understood that any embodiment of the compounds of structure (I), as set forth above, and any specific substituent and/or variable in the compound structure (I), as set forth above, may be independently combined with other embodiments and/or substituents and/or variables of compounds of structure (I) to form

5 embodiments of the disclosures not specifically set forth above. In addition, in the event that a list of substituents and/or variables is listed for any particular R group, G group, L group or variable n, in a particular embodiment and/or claim, it is understood that each individual substituent and/or variable may be deleted from the particular embodiment and/or claim and that the remaining list of substituents and/or variables

10 will be considered to be within the scope of the disclosure.

It is understood that in the present description, combinations of substituents and/or variables of the depicted formulae are permissible only if such contributions result in stable compounds.

In some embodiments, lipid nanoparticles comprising a compound of structure (I) are provided. The lipid nanoparticles optionally include excipients selected

15 from a neutral lipid, a steroid and a polymer conjugated lipid.

In some embodiments, compositions comprising any one or more of the compounds of structure (I) and a therapeutic agent are provided. For example, in some embodiments, the compositions comprise any of the compounds of structure (I) and a

20 therapeutic agent and one or more excipient selected from neutral lipids, steroids and polymer conjugated lipids. Other pharmaceutically acceptable excipients and/or carriers are also included in various embodiments of the compositions.

In some embodiments, the neutral lipid is selected from DSPC, DPPC, DMPC, DOPC, POPC, DOPE and SM. In some embodiments, the neutral lipid is


DSPC. In various embodiments, the molar ratio of the compound to the neutral lipid ranges from about 2:1 to about 8:1.

In various embodiments, the compositions further comprise a steroid or steroid analogue. In certain embodiments, the steroid or steroid analogue is cholesterol.

- 5 In some of these embodiments, the molar ratio of the compound to cholesterol ranges from about 5:1 to 1:1.

- In various embodiments, the polymer conjugated lipid is a pegylated lipid. For example, some embodiments include a pegylated diacylglycerol (PEG-DAG) such as 1-(monomethoxy-polyethyleneglycol)-2,3-dimyristoylglycerol (PEG-DMG), a
 10 pegylated phosphatidylethanolamine (PEG-PE), a PEG succinate diacylglycerol (PEG-S-DAG) such as 4-O-(2',3'-di(tetradecanoyloxy)propyl-1-O-(ω-methoxy(polyethoxy)ethyl)butanedioate (PEG-S-DMG), a pegylated ceramide (PEG-cer), or a PEG dialkoxypentylcarbamate such as ω-methoxy(polyethoxy)ethyl-N-(2,3-di(tetradecanoyloxy)propyl)carbamate or 2,3-di(tetradecanoyloxy)propyl-N-(ω-methoxy(polyethoxy)ethyl)carbamate. In various embodiments, the molar ratio of the
 15 compound to the pegylated lipid ranges from about 100:1 to about 20:1.

In some embodiments, the composition comprises a pegylated lipid having the following structure (II):



20

(II)

or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein:

R^8 and R^9 are each independently a straight or branched, saturated or unsaturated alkyl chain containing from 10 to 30 carbon atoms, wherein the alkyl chain is optionally interrupted by one or more ester bonds; and

25

w has a mean value ranging from 30 to 60.

In some embodiments, R^8 and R^9 are each independently straight, saturated alkyl chains containing from 12 to 16 carbon atoms. In other embodiments, the average w ranges from about 42 to 55, for example about 49.

In some embodiments of the foregoing composition, the therapeutic agent comprises a nucleic acid. For example, in some embodiments, the nucleic acid is selected from antisense and messenger RNA. In some of the foregoing embodiments, the composition comprises a lipid nanoparticle.


5 Some related embodiments provide a lipid nanoparticle comprising the compound of any one of the foregoing embodiments (*e.g.*, a compound of structure (I)). In certain embodiments, the lipid nanoparticle further comprises a therapeutic agent (*e.g.*, a nucleic acid such as antisense and messenger RNA).

10 In some embodiments, the lipid nanoparticle further comprises one or more excipient selected from neutral lipids, steroids and polymer conjugated lipids. In some embodiments, the neutral lipids are selected from DSPC, DPPC, DMPC, DOPC, POPC, DOPE and SM. In more specific embodiments, the neutral lipid is DSPC.

15 In some more specific embodiments, the molar ratio of the compound to the neutral lipid ranges from about 2:1 to about 8:1. In some embodiments, the steroid is cholesterol. In some embodiments, the molar ratio of the compound to cholesterol ranges from 5:1 to 1:1.

 In certain embodiments, the polymer conjugated lipid is pegylated lipid. In certain more specific embodiments, the molar ratio of the compound to pegylated lipid ranges from about 100:1 to about 20:1.

20 In some embodiments, the pegylated lipid is PEG-DAG, PEG-PE, PEG-S-DAG, PEG-cer or a PEG dialkoxypentylcarbamate. In other embodiments, the pegylated lipid has the following structure (II):



(II)

25 or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein:

R^8 and R^9 are each independently a straight or branched, saturated or unsaturated alkyl chain containing from 10 to 30 carbon atoms, wherein the alkyl chain is optionally interrupted by one or more ester bonds; and

w has a mean value ranging from 30 to 60.

In some more specific embodiments of structure (II), R⁸ and R⁹ are each independently straight, saturated alkyl chains containing from 12 to 16 carbon atoms. In more specific embodiments, the average w is about 49.

In other different embodiments, the disclosure is directed to a method for
5 administering a therapeutic agent to a patient in need thereof, the method comprising preparing or providing any of the foregoing compositions and administering the composition to the patient

For the purposes of administration, embodiments of the compounds of the present disclosure (typically in the form of lipid nanoparticles in combination with a
10 therapeutic agent) may be administered as a raw chemical or may be formulated as pharmaceutical compositions. Pharmaceutical compositions of embodiments of the present disclosure comprise a compound of structure (I) and one or more pharmaceutically acceptable carrier, diluent or excipient. In some embodiments, the compound of structure (I) is present in the composition in an amount which is effective
15 to form a lipid nanoparticle and deliver the therapeutic agent, *e.g.*, for treating a particular disease or condition of interest. Appropriate concentrations and dosages can be readily determined by one skilled in the art.

Administration of the compositions of embodiments of the disclosure can be carried out via any of the accepted modes of administration of agents for serving
20 similar utilities. The pharmaceutical compositions of embodiments of the disclosure may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suspensions, suppositories, injections, inhalants, gels, microspheres, and aerosols. Typical routes of administering such pharmaceutical compositions include, without limitation, oral, topical,
25 transdermal, inhalation, parenteral, sublingual, buccal, rectal, vaginal, and intranasal. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intradermal, intrasternal injection or infusion techniques. Pharmaceutical compositions of embodiments of the disclosure are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of
30 the composition to a patient. Compositions that will be administered to a subject or patient in some embodiments take the form of one or more dosage units, where for

example, a tablet may be a single dosage unit, and a container of a compound of an
embodiments of the disclosure in aerosol form may hold a plurality of dosage units.
Actual methods of preparing such dosage forms are known, or will be apparent, to those
skilled in this art; for example, see *Remington: The Science and Practice of Pharmacy*,
5 20th Edition (Philadelphia College of Pharmacy and Science, 2000). In some
embodiments, the composition to be administered will, in any event, contain a
therapeutically effective amount of a compound of the disclosure, or a pharmaceutically
acceptable salt thereof, for treatment of a disease or condition of interest in accordance
with the teachings of this disclosure.

10 A pharmaceutical composition of embodiments of the disclosure may be
in the form of a solid or liquid. In one aspect, the carrier(s) are particulate, so that the
compositions are, for example, in tablet or powder form. The carrier(s) may be liquid,
with the compositions being, for example, oral syrup, injectable liquid or an aerosol,
which is useful in, for example, inhalatory administration.

15 When intended for oral administration, the pharmaceutical composition
of certain embodiments is preferably in either solid or liquid form, where semi-solid,
semi-liquid, suspension and gel forms are included within the forms considered herein
as either solid or liquid.

 As a solid composition for oral administration, the pharmaceutical
20 composition of some embodiments may be formulated into a powder, granule,
compressed tablet, pill, capsule, chewing gum, wafer or the like form. Such a solid
composition will typically contain one or more inert diluents or edible carriers. In
addition, one or more of the following may be present: binders such as
carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or
25 gelatin; excipients such as starch, lactose or dextrans, disintegrating agents such as
alginic acid, sodium alginate, Primogel, corn starch and the like; lubricants such as
magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening
agents such as sucrose or saccharin; a flavoring agent such as peppermint, methyl
salicylate or orange flavoring; and a coloring agent.

When the pharmaceutical composition of some embodiments is in the form of a capsule, for example, a gelatin capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or oil.

The pharmaceutical composition of some embodiments may be in the form of a liquid, for example, an elixir, syrup, solution, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred composition contain, in addition to a compound of structure (I), one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

The liquid pharmaceutical compositions of embodiments of the disclosure, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose; agents to act as cryoprotectants such as sucrose or trehalose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

A liquid pharmaceutical composition of embodiments of the disclosure intended for either parenteral or oral administration should contain an amount of a compound of the disclosure such that a suitable dosage will be obtained.

The pharmaceutical composition of embodiments of the disclosure may be intended for topical administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil,

diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or iontophoresis device.

5 The pharmaceutical composition of embodiments of the disclosure may be intended for rectal administration, in the form, for example, of a suppository, which will melt in the rectum and release the drug. A composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter and polyethylene glycol.

10 The pharmaceutical composition of embodiments of the disclosure may include various materials, which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials that form the coating shell are typically inert, and may be selected from, for example, sugar, shellac, and other enteric coating
15 agents. Alternatively, the active ingredients may be encased in a gelatin capsule.

 The pharmaceutical composition of embodiments of the disclosure in solid or liquid form may include an agent that binds to the compound of the disclosure and thereby assists in the delivery of the compound. Suitable agents that may act in this capacity include a monoclonal or polyclonal antibody, or a protein.

20 The pharmaceutical composition of embodiments of the disclosure may consist of dosage units that can be administered as an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or by a suitable pump system that dispenses the active ingredients. Aerosols of
25 compounds of embodiments of the disclosure may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, activators, valves, sub-containers, and the like, which together may form a kit. One skilled in the art, without undue experimentation, may determine preferred aerosols.

30 The pharmaceutical compositions of embodiments of the disclosure may be prepared by methodology well known in the pharmaceutical art. For example, a

pharmaceutical composition intended to be administered by injection can be prepared by combining the lipid nanoparticles of the disclosure with sterile, distilled water or other carrier so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that
5 non-covalently interact with the compound of the disclosure so as to facilitate dissolution or homogeneous suspension of the compound in the aqueous delivery system.

The compositions of embodiments of the disclosure, or their pharmaceutically acceptable salts, are administered in a therapeutically effective
10 amount, which will vary depending upon a variety of factors including the activity of the specific therapeutic agent employed; the metabolic stability and length of action of the therapeutic agent; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disorder or condition; and the subject undergoing therapy.

15 Compositions of embodiments of the disclosure may also be administered simultaneously with, prior to, or after administration of one or more other therapeutic agents. Such combination therapy includes administration of a single pharmaceutical dosage formulation of a composition of embodiments of the disclosure and one or more additional active agents, as well as administration of the composition
20 of embodiments of the disclosure and each active agent in its own separate pharmaceutical dosage formulation. For example, a composition of embodiments of the disclosure and the other active agent can be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations. Where separate dosage formulations are used, the
25 compounds of embodiments of the disclosure and one or more additional active agents can be administered at essentially the same time, *i.e.*, concurrently, or at separately staggered times, *i.e.*, sequentially; combination therapy is understood to include all these regimens.


Preparation methods for the above compounds and compositions are
30 described herein below and/or known in the art.

It will be appreciated by those skilled in the art that in the process described herein the functional groups of intermediate compounds may need to be protected by suitable protecting groups. Such functional groups include hydroxy, amino, mercapto and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl or diarylalkylsilyl (for example, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl, and the like. Suitable protecting groups for amino, amidino and guanidino include *t*-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for mercapto include -C(O)-R" (where R" is alkyl, aryl or arylalkyl), *p*-methoxybenzyl, trityl and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters. Protecting groups may be added or removed in accordance with standard techniques, which are known to one skilled in the art and as described herein. The use of protecting groups is described in detail in Green, T.W. and P.G.M. Wutz, *Protective Groups in Organic Synthesis* (1999), 3rd Ed., Wiley. As one of skill in the art would appreciate, the protecting group may also be a polymer resin such as a Wang resin, Rink resin or 2-chlorotrityl-chloride resin.

It will also be appreciated by those skilled in the art, although such protected derivatives of compounds of this disclosure may not possess pharmacological activity as such, they may be administered to a mammal and thereafter metabolized in the body to form compounds of the disclosure which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". All prodrugs of compounds of this disclosure are included within the scope of the disclosure.

Furthermore, compounds of embodiments of the disclosure which exist in free base or acid form can be converted to their pharmaceutically acceptable salts by treatment with the appropriate inorganic or organic base or acid by methods known to one skilled in the art. Salts of compounds of embodiments of the disclosure can be converted to their free base or acid form by standard techniques.


The following General Reaction Scheme 1 illustrates exemplary methods to make compounds of this disclosure, *i.e.*, compounds of structure (I):



(I)

or a pharmaceutically acceptable salt, tautomer or stereoisomer thereof, wherein R^1 , R^2 , R^3 , R^4 , R^5 , L^1 , L^2 , L^3 , G^1 , G^2 , and G^3 are as defined herein. It is understood that one skilled in the art may be able to make these compounds by similar methods or by combining other methods known to one skilled in the art. It is also understood that one skilled in the art would be able to make, in a similar manner as described below, other compounds of structure (I) not specifically illustrated below by using the appropriate starting components and modifying the parameters of the synthesis as needed. In general, starting components may be obtained from sources such as Sigma Aldrich, Lancaster Synthesis, Inc., Maybridge, Matrix Scientific, TCI, and Fluorochem USA, etc. or synthesized according to sources known to those skilled in the art (see, for example, Advanced Organic Chemistry: Reactions, Mechanisms, and Structure, 5th edition (Wiley, December 2000)) or prepared as described in this disclosure.

15

GENERAL REACTION SCHEME 1

General Reaction Scheme I provides an exemplary method for preparation of compounds of structure (I). R^1 , R^2 , R^3 , R^4 , R^5 , L^1 , L^2 , L^3 , G^1 , G^2 , and G^3 in General reaction Scheme 1 are as defined herein. X^1 and X^2 are reactive moieties selected to facilitate the desired reaction (*e.g.*, halo). Compounds of structure A1 are purchased or prepared according to methods known in the art. Reaction of A1 under appropriate reducing conditions (*e.g.*, sodium triacetoxyborohydride) yields the product of the reductive amination between A1 and A2, A3. A3 is then reacted with A4 under suitable basic conditions (*e.g.*, using triethylamine and DMAP) to afford compound A5. A5 is then reacted with amine A6 using appropriate conditions (*e.g.*, heat) to yield a compound of structure (I) as shown.

It should be noted that various alternative strategies for preparation of compounds of structure (I) are available to those of ordinary skill in the art. For example, other compounds of structure (I) wherein can be prepared according to analogous methods using the appropriate starting material. The use of protecting groups as needed and other modification to the above General Reaction Scheme will be readily apparent to one of ordinary skill in the art.

The following examples are provided for purpose of illustration and not limitation.

EXAMPLE 1

LUCIFERASE MRNA IN VIVO EVALUATION USING THE LIPID NANOPARTICLE COMPOSITIONS

Lipid nanoparticles were prepared and tested according to the general procedures described in PCT Pub. Nos. WO 2015/199952 and WO 2017/004143, the full disclosures of which are incorporated herein by reference. Briefly, cationic lipid, DSPC, cholesterol and PEG-lipid were solubilized in ethanol at a molar ratio of about 50:10:38.5:1.5 or about 47.5:10:40.8:1.7. Lipid nanoparticles (LNP) were prepared at a total lipid to mRNA weight ratio of approximately 10:1 to 30:1. The mRNA is diluted to 0.2 mg/mL in 10 to 50 mM citrate or acetate buffer, pH 4. Syringe pumps were used to mix the ethanolic lipid solution with the mRNA aqueous solution at a ratio of about 1:5 to 1:3 (vol/vol) with total flow rates above 15 mL/min. The ethanol was then

removed and the external buffer replaced with PBS by dialysis. Finally, the lipid nanoparticles were filtered through a 0.2 µm pore sterile filter. Lipid nanoparticle particle size was approximately 55-95 nm diameter, and in some instances approximately 70-90 nm diameter as determined by quasi-elastic light scattering using a
5 Malvern Zetasizer Nano ZS (Malvern, UK).

Studies were performed in 6-8 week old female C57BL/6 mice (Charles River) or 8-10 week old CD-1 (Harlan) mice (Charles River) according to guidelines established by an institutional animal care committee (ACC) and the Canadian Council on Animal Care (CCAC). Varying doses of mRNA-lipid nanoparticle were
10 systemically administered by tail vein injection and animals euthanized at a specific time point (e.g., 4 hours) post-administration. Liver and spleen were collected in pre-weighed tubes, weights determined, immediately snap frozen in liquid nitrogen and stored at -80°C until processing for analysis.

For liver, approximately 50 mg was dissected for analyses in a 2 mL
15 FastPrep tubes (MP Biomedicals, Solon OH). ¼" ceramic sphere (MP Biomedicals) is added to each tube and 500 µL of Glo Lysis Buffer – GLB (Promega, Madison WI) equilibrated to room temperature is added to liver tissue. Liver tissues were homogenized with the FastPrep24 instrument (MP Biomedicals) at 2×6.0 m/s for 15 seconds. Homogenate was incubated at room temperature for 5 minutes prior to a 1:4
20 dilution in GLB and assessed using SteadyGlo Luciferase assay system (Promega). Specifically, 50 µL of diluted tissue homogenate was reacted with 50 µL of SteadyGlo substrate, shaken for 10 seconds followed by 5 minute incubation and then quantitated using a CentroXS³ LB 960 luminometer (Berthold Technologies, Germany). The amount of protein assayed was determined by using the BCA protein assay kit (Pierce,
25 Rockford IL). Relative luminescence units (RLU) were then normalized to total µg protein assayed. To convert RLU to ng luciferase a standard curve was generated with QuantiLum Recombinant Luciferase (Promega).

The FLuc mRNA (L-6107 or L-7202) from Trilink Biotechnologies will express a luciferase protein, originally isolated from the firefly, *photinus pyralis*. FLuc
30 is commonly used in mammalian cell culture to measure both gene expression and cell viability. It emits bioluminescence in the presence of the substrate, luciferin. This

capped and poly-adenylated mRNA was fully substituted with respect to uridine and/or cytidine nucleosides.

EXAMPLE 2

DETERMINATION OF pK_a OF FORMULATED LIPIDS

5 As described elsewhere, the pK_a of formulated cationic lipids is correlated with the effectiveness of LNPs for delivery of nucleic acids (see Jayaraman et al, *Angewandte Chemie*, International Edition (2012), 51(34), 8529-8533; Semple et al, *Nature Biotechnology* 28, 172–176 (2010)). The preferred range of pK_a is ~5 to ~7. The pK_a of each cationic lipid was determined in lipid nanoparticles using an assay
10 based on fluorescence of 2-(p-toluidino)-6-naphthalene sulfonic acid (TNS). Lipid nanoparticles comprising cationic lipid/DSPC/cholesterol/PEG-lipid (50/10/38.5/1.5 mol%) in PBS at a concentration of 0.4 mM total lipid were prepared using the in-line process as described in Example 1. TNS was prepared as a 100 μ M stock solution in distilled water. Vesicles were diluted to 24 μ M lipid in 2 mL of buffered solutions
15 containing, 10 mM HEPES, 10 mM MES, 10 mM ammonium acetate, 130 mM NaCl, where the pH ranged from 2.5 to 11. An aliquot of the TNS solution was added to give a final concentration of 1 μ M and following vortex mixing fluorescence intensity was measured at room temperature in a SLM Aminco Series 2 Luminescence Spectrophotometer using excitation and emission wavelengths of 321 nm and 445 nm.
20 A sigmoidal best fit analysis was applied to the fluorescence data and the pK_a was measured as the pH giving rise to half-maximal fluorescence intensity.

EXAMPLE 3

DETERMINATION OF EFFICACY OF LIPID NANOPARTICLE FORMULATIONS

CONTAINING VARIOUS CATIONIC LIPIDS USING AN IN VIVO






25 LUCIFERASE mRNA EXPRESSION RODENT MODEL




The cationic lipids shown in Table 2 have previously been tested with nucleic acids. For comparative purposes, these lipids were also used to formulate lipid nanoparticles containing the FLuc mRNA (L-6107) using an in line mixing method, as

described in Example 1 and in PCT/US10/22614, which is hereby incorporated by reference in its entirety. Lipid nanoparticles were formulated using the following molar ratio: 50% Cationic lipid / 10% distearoylphosphatidylcholine (DSPC) / 38.5% Cholesterol / 1.5% PEG lipid ("PEG-DMG", *i.e.*,

- 5 1-(monomethoxy-polyethyleneglycol)-2,3-dimyristoylglycerol, with an average PEG molecular weight of 2000). In alternate embodiments, cationic lipid, DSPC, cholesterol, and PEG-lipid are formulated at a molar ratio of approximately 47.5:10:40.8:1.7. Relative activity was determined by measuring luciferase expression in the liver 4 hours following administration via tail vein injection as described in
- 10 Example 1. The activity was compared at a dose of 0.3 and 1.0 mg mRNA/kg and expressed as ng luciferase/g liver measured 4 hours after administration, as described in Example 1.






Table 2. Comparator Lipids showing activity with mRNA





Compound	Liver Luc @ 0.3 mg/kg dose	Liver Luc @ 1.0 mg/kg dose	Structure
MC2	4 ± 1	N/D	
DLinDMA	13 ± 3	67 ± 20	
MC4	41 ± 10	N/D	
XTC2	80 ± 28	237 ± 99	
MC3	198 ± 126	757 ± 528	

Compound	Liver Luc @ 0.3 mg/kg dose	Liver Luc @ 1.0 mg/kg dose	Structure
319 (2% PEG)	258 ± 67	681 ± 203	
137	281 ± 203	588 ± 303	
A	77 ± 40	203 ± 122	

Representative compounds of the disclosure shown in Table 3 were formulated using the following molar ratio: 50% cationic lipid / 10% distearoylphosphatidylcholine (DSPC) / 38.5% Cholesterol / 1.5% PEG lipid ("PEG-DMA" 2-[2-(ω-methoxy(polyethyleneglycol₂₀₀₀)ethoxy]-N,N-ditetradecylacetamide) or 47.5% cationic lipid / 10% DSPC / 40.7% Cholesterol / 1.8% PEG lipid. Relative activity was determined by measuring luciferase expression in the liver 4 hours following administration via tail vein injection as described in Example 1. The activity was compared at a dose of 0.5 mg mRNA/kg and expressed as ng luciferase/g liver measured 4 hours after administration, as described in Example 1. Compound numbers in Table 3 refer to the compound numbers of Table 1.


Table 3. Novel Cationic Lipids and Associated Activity

Cmp . No.	pK _a	Liver Luc @ 0.5 mg/kg (ng luc/g liver)	Structure
I-7	6.7 4	2420 ± 1079	
I-8	6.6 8	7384 ± 1916	
I-9	6.8 3	2223 ± 777	
I-16	6.7 7	4678 ± 1243	
I-18	6.4 7	1330 1 ± 8071	

Cmp . No.	pK _a	Liver Luc @ 0.5 mg/kg (ng luc/g liver)	Structure
I-33	6.1 9	1304 ± 278	
I-35	6.3 9	1431 2 ± 4174	
I-36	6.5 5	1037 7 ± 1435	
I-37	6.3 0	1523 4 ± 5248	

EXAMPLE 4

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-DECYL-5-(DIMETHYLAMINO)PENTANAMIDO)NONADECANEDIOATE (COMPOUND I-7)



5 Synthesis of Compound 4-2

A solution of the ketone **4-1** (1.10 g, 1.62 mmol) and 1-decylamine (2.43 mmol, 382 mg, 0.486 mL) in DCE (10 mL) was stirred at room temperature for 15 minutes, followed by addition of sodium triacetoxyborohydride (2.43 mmol, 515 mg) and acetic acid (2.43 mmol, 146 mg; 0.138 mL). After the mixture was stirred at room temperature for 2 days, the reaction mixture was concentrated. The residue was diluted with a mixture of hexanes and washed with dilute NaOH, saturated NaHCO₃ and brine. The organic phase was separated, dried over sodium sulfate and concentrated (colorless oil, 1.41 g). The crude product was purified by column chromatography on silica gel (hexane/EtOAc/Et₃N, 95:5:0 to 80:20:1). The desired product was obtained as colorless oil (863 mg colorless oil, 1.05 mmol, 65% yield). ¹H NMR (400 MHz, CDCl₃) δ: 3.98 (d, 5.8 Hz, 4H), 2.54 (t, 7.1 Hz, 2H), 2.43 (quintet, 5.5 Hz, 1H), 2.30 (t, 7.5 Hz, 4H), 1.68-1.57 (m, 6H), 1.50-1.41 (m, 2H), 1.41-1.08 (70H), 0.92-0.86 (m, 15H), 0.86-0.77 (br. 1H).

15 **Synthesis of Compound 4-3**

To a stirred solution of 5-bromovaleric acid (1.12 mmol, 204 mg) in CH₂Cl₂ (1 mL) at room temperature was added a solution of thionyl chloride (3.36 mmol, 400 mg, 0.25 mL) in CH₂Cl₂ (5 mL) in a period of 1 minute, followed by addition of DMF (ca 16 mg). The mixture was then heated to reflux for 2 hours. The reaction mixture was then concentrated *in vacuo*. The acid chloride was directly used for the next step.

A solution of the above 5-bromopentanoyl chloride in benzene (5 mL) was added dropwise to a solution of **4-2** (230 mg, 0.28 mmol) and triethylamine (5.6 mmol, 565 mg, 0.780 mL) and DMAP (5 mg) in benzene (5 mL) at room temperature in 2 minutes. After addition, the reaction was stirred at room temperature for 1 hour. Methanol (1 mL) was added and the mixture was stirred for 2 hour. The reaction mixture was concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel (hexane/EtOAc, 98:2 to 85:15). The desired product was obtained as colorless oil (250 mg, 0.25 mmol, 91%, colorless oil).


30

Synthesis of Compound I-7

To **4-2** (250 mg, 0.25 mmol) was added dimethyl amine (2M in THF, 10 mL). The solution was stirred at 64 °C overnight. The reaction mixture was concentrated. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a pad of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a brownish oil (ca 233 mg). The crude product (233 mg) was purified by flash dry column chromatography on silica gel (0 to 6% methanol in chloroform). The desired product was obtained as (194 mg, colorless oil, 0.20 mmol, 82%). ¹H NMR (400 MHz, CDCl₃) δ: 4.60-4.20 (br, estimated 0.3H, due to slow isomerization about amide bond), 3.97, 3.96 (2 sets of doublets, 5.8 Hz, 4H), 3.60 (quintet-like, 7.0 Hz, 0.7H), 3.06-2.99 (m, 2H), 2.34-2.24 (m, 8H), 2.21 (singlet, 6H), 1.72-1.56 (m, 8H), 1.56-1.37 (m, 8H), 1.37-1.10 (66H), 0.91-0.85 (m, 15H).

EXAMPLE 5

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-DECYL-4-(DIMETHYLAMINO)BUTANAMIDO)NONADECANEDIOATE (COMPOUND I-18)



5 Synthesis of Compound I-18, (Method A)

To a stirred solution of 4-(dimethylamino)butyric acid hydrochloride (1.12 mmol, 188 mg) and DMF (10-20 μ L) in CH_2Cl_2 (10 mL) at room temperature was added oxalyl chloride (5.6 mmol, 722 mg, 0.496 mL). The resulting mixture was stirred at room temperature overnight. The reaction mixture (deep red) was concentrated under reduced pressure. The resulting acid chloride (brick red solid) was

10 concentrated under reduced pressure. The resulting acid chloride (brick red solid) was directly used for the next step.

A solution of the above acyl chloride in CH_2Cl_2 (10 mL) was added dropwise to a solution of **4-2** (230 mg, 0.28 mmol) and triethylamine (5.6 mmol, 565 mg, 0.780 μ L) and DMAP (5 mg) in CH_2Cl_2 (5 mL) at room temperature. The resulting

15 mixture was stirred at room temperature for overnight and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel (hexane/EtOAc/ Et_3N , from 80:20:0.1 to 75:25:1) and further was purified by flash dry


column chromatography on silica gel (0 to 5% methanol in chloroform). The desired product was obtained as (92 mg, colorless oil, 0.10 mmol, 35%). ¹H NMR (400 MHz, CDCl₃) δ: 4.57-4.29 (br. 0.4H), 3.97, 3.96 (2 sets of doublets, 5.8 Hz, 5.8 Hz, 4H), 3.63 (quintet-like, 6.8 Hz, 0.6H), 3.06-3.00 (m, 2H), 2.35-2.26 (m, 8H), 2.213, 2.211 (2 sets of singlet, 6H), 1.82 (sextet-like, 7.6 Hz, 2H), 1.65-1.56 (m, 6H), 1.54-1.48 (m, 2H), 1.47-1.37 (m, 4H), 1.36-1.06 (66H), 0.91-0.86 (m, 15H).

Synthesis of Compound I-18, (Method B)

To a solution of 4-dimethylaminobutyric acid hydrochloride (2 equiv, 3.04 mmol, 510 mg) and 4-dimethylaminopyridine (3 equiv, DMAP, 4.56 mmol, 557 mg) in acetonitrile (30 mL) was added DCC (2.2 equiv, 3.34 mmol, 690 mg) and the mixture stirred at room temperature for 45 min. A solution of **4-2** (1.25 g, 1.52 mmol) in CH₂Cl₂ (6 mL) was added and the resulting mixture was stirred overnight. On the next day, more DCC (450 mg) was added and the mixture was stirred for another day. The mixture was then concentrated under reduce pressure. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a colorless oil. The product was further purified by flash dry column chromatography on silica gel (0 to 5% methanol in chloroform with a trace of Et₃N). The desired product was obtained as colorless oil (1.14 g, 81%).

EXAMPLE 6


SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-DECYL-5-(DIETHYLAMINO)PENTANAMIDO)NONADECANEDIOATE (COMPOUND I-8)



5 Synthesis of Compound I-8

A mixture of **4-3** (179 mg, 0.18 mmol), diethylamine (0.90 mmol, 66 mg, 0.093 mL), N,N-diisopropylethylamine (0.36 mmol, 46 mg, 0.063 mL) in acetonitrile (6 mL) was sealed and heated at 83 °C for 24 hours. The reaction mixture was concentrated. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a pad of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a brownish oil (153 mg). The crude product (233 mg) was purified by flash dry column chromatography on silica gel (0 to 6% MeOH in chloroform). The desired product was obtained as (136 mg, colorless oil, 0.14 mmol, 77%).

EXAMPLE 7


SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-DECYL-5-(PYRROLIDIN-1-
YL)PENTANAMIDO)NONADECANEDIOATE (COMPOUND I-9)

5 Synthesis of Compound I-9

A mixture of **4-3** (200 mg, 0.20 mmol), pyrrolidine (50 eq. 0.83 mL, 10 mmol) in THF (10 mL) was sealed and heated at 64 °C for 24 hours. The reaction mixture was concentrated. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a pad of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a brownish oil. The crude product (233 mg) was purified by flash dry column chromatography on silica gel (0 to 6% MeOH in chloroform). The desired product was obtained as (158 mg, colorless oil, 0.16 mmol, 80%).

EXAMPLE 8

SYNTHESIS OF BIS(2-HEXYLDECYL) 7-(N-DECYL-4-(DIMETHYLAMINO)BUTANAMIDO)TRIDECANEDIOATE (COMPOUND I-16)

5 Synthesis of **8-2**

A solution of **8-1** (1 eq., 1.15 g, 1.62 mmol) and 1-decylamine (1.5 eq., 2.43 mmol, 382 mg, 0.486 mL) in DCE (10 mL) was stirred at RT for about 15 min. To the solution was added sodium triacetoxyborohydride (1.5 eq., 2.43 mmol, 515 mg) and AcOH (1.5 eq, 2.43 mmol, 146 mg, 0.14 mL). The mixture was stirred at RT for 3 days.

- 10 The reaction mixture was then concentrated. The residue was diluted with hexanes/EtOAc (99:1) and washed with dilute NaOH solution, sat NaHCO₃ and brine. The organic extract was dried over sodium sulfate and was poured over a short column of silica gel. The column was eluted with a mixture of hexane, EtOAc and Et₃N (95:5:0 to 80:20:1). The fractions containing the pure product were combined and concentrated.

The desired product was obtained as colorless oil (1.28 g, 1.51 mmol, 93%). ¹HNMR (400 MHz, CDCl₃) δ: 3.97 (d, 5.8 Hz, 4H), 2.53 (t, 7.2 Hz, 2H), 2.43 (quintet-like, 5.5 Hz, 1H), 2.30 (t, 7.5 Hz, 4H), 1.68-1.57 (m, 6H), 1.49-1.40 (m, 2H), 1.40-1.08 (74H), 0.91-0.85 (m, 15H), 0.83-0.74 (br. 1H).

5

Synthesis Compound I-16


To a stirred solution of 4-(dimethylamino)butyric acid hydrochloride (2.1 mmol, 352 mg) and DMF (ca 13 mg) in CH₂Cl₂ (15 mL) at room temperature was added oxalyl chloride (3 eq, 6.3 mmol, 800 mg, 0.55 mL) under Ar. The resulting mixture was stirred at RT overnight. The reaction mixture (light orange solution) was concentrated in vacuo. The resulting acid chloride (**8-3**, light brown solid) was directly used for the following reaction.

A solution of the above acyl chloride in CH₂Cl₂ (10 mL) was added to a solution of **8-2** (300 mg, 0.35 mmol) and triethylamine (10.5 mmol, 1.06 g, 1.5 mL) and DMAP (5 mg) in CH₂Cl₂ (5 mL) at RT. After addition, the reaction mixture was stirred at RT for overnight. After concentration of the mixture, the product was isolated by column chromatography on silica gel (hexane, EtOAc and Et₃N, from 80:20:0.1 to 70:30:1) and the product was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform). The desired product was obtained as slightly yellow oil (110 mg, 0.11 mmol, 32%). ¹HNMR (400 MHz, CDCl₃ at 7.26 ppm) δ: 4.57-4.34 (br. 0.4H), 3.98-3.94 (m, 4H), 3.64 (quintet-like, 6.8 Hz, 0.6H), 3.06-3.00 (m, 2H), 2.35-2.25 (m, 8H), 2.213, 2.210 (2 sets of singlet, 6H), 1.82 (sextet-like, 7.4 Hz, 2H), 1.65-1.56 (m, 6H), 1.54-1.48 (m, 2H), 1.48-1.37 (m, 4H), 1.37-1.06 (70H), 0.91-0.86 (m, 15H).

25

EXAMPLE 9

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(4-(DIMETHYLAMINO)-N-(2-ETHYLHEXYL)BUTANAMIDO)NONADECANEDIOATE (COMPOUND I-20)



Synthesis of 9-2

A solution of **9-1** (1 eq., 0.82 g, 1.21 mmol) and 2-ethyl-1-hexylamine (1.5 eq, 1.81 mmol, 234 mg) in DCE (8 mL) was stirred at RT for about 15 min. To the solution was added sodium triacetoxyborohydride (1.5 eq., 1.81 mmol, 384 mg) and AcOH (1.5 eq, 1.81 mmol, 109 mg). The mixture was stirred at RT for 2 days. The reaction mixture was then concentrated. The residue was diluted with hexanes and EtOAc (ca 99:5) and washed with dilute NaOH, sat NaHCO₃ and brine. The extract was filtered through a short column of silica gel. The column was washed with a mixture of hexane and EtOAc (95:5) and then a mixture of hexanes, EtOAc and Et₃N (80:20:0.5). The filtrate from the latter washing was concentrated to dryness. This gave the pure product as colorless oil (888 mg, 1.12 mmol, 93%). ¹HNMR (400 MHz, CDCl₃) δ: 3.98 (d, 5.8 Hz, 4H), 2.45 (d, 5.1 Hz, 2H), 2.39 (quintet-like, 5.5 Hz, 1H), 2.30 (t, 7.5 Hz, 4H), 1.68-1.57 (m, 6H), 1.41-1.08 (65H), 0.92-0.85 (m, 18H), 0.84-0.78 (br. 1H).

15

Synthesis of 9-4

To a stirred solution of 5-bromovaleric acid (2.24 mmol, 405 mg) in CH₂Cl₂ (2 mL) at room temperature was slowly added a solution of thionyl chloride (3 eq. 6.72 mmol, 800 mg, 0.49 mL) in CH₂Cl₂ (5 mL) in a period of 1 min. DMF (two tiny drops, ca 16 mg) was added to the reaction mixture. The mixture was then heated to reflux for 2 h. The reaction mixture was concentrated in vacuo. The resulting acid chloride, **9-3**, was directly used for the next step.


A solution of the above 5-bromopentanoyl chloride in benzene (8 mL) was added to a solution of **9-2** (444 mg, 0.56 mmol) and triethylamine (1.56 mL) and DMAP (5 mg) in benzene (5 mL) at RT in 2 min. After addition, the mixture was stirred at RT overnight. After evaporation of solvents in vacuo, the product was isolated by column chromatography on silica gel (hexane/EtOAc, 99:1 to 90:10). The desired product was pure enough for the next step (colorless oil, 527 mg, 0.55 mmol, 98%).

Synthesis of Compound I-20

To a pressure flask containing **9-4** (260 mg, 0.27 mmol) was added dimethyl amine (2M in THF, 10 mL). The solution was stirred at 64 °C (oil bath temperature) overnight. Excess amine and solvent was evaporated. The residue was
5 taken up in a mixture of ethyl acetate and hexane (95:5) and filtered through a pad of silica gel. The pad was washed with a mixture of hexane and EtOAc (95:5) and then a mixture of hexanes, EtOAc and Et₃N (80:20:1). The filtrate from the latter washing was concentrated to dryness. This gave the crude product as brown oil (233 mg). The crude product was further purified by flash dry column chromatography on silica gel (0
10 to 5% MeOH in chloroform). The desired product was obtained as colorless oil (204 mg, 0.22 mmol, 82%). ¹HNMR (400 MHz, CDCl₃ at 7.27 ppm) δ: 3.97 (d, 5.8 Hz, 4H), 3.66-3.57 (m, ca 1H), 3.19-2.99 (2 sets of peaks, 2H), 2.38-2.24 (m, 8H), 2.22 (singlet, 6H), 1.72-1.37 (m, 15H), 1.37-1.10 (60H), 0.91-0.85 (m, 18H).

EXAMPLE 10

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(3-(DIMETHYLAMINO)-N-NONYLPROPANAMIDO)NONADECANEDIOATE (COMPOUND I-25)



5 Synthesis of 10-1

To a solution of 3-bromopropionic acid (2.02 mmol, 311 mg) in CH₂Cl₂ (5 mL) and DMF (0.01 mL) was added oxalyl chloride (5.05 mmol, 641 mg, 0.44 mL) at RT. The resulting mixture was stirred at RT overnight. The mixture was then concentrated in vacuo. The residual liquid/solid (yellow) was dissolved in 10 mL of CH₂Cl₂ and added to a solution of **4-2** (833 mg, 1.02 mmol) and triethylamine (5.05


mmol, 0.7 mL) and DMAP (5 mg) in CH₂Cl₂ (10 mL) at RT in 4 min. After addition, the mixture was stirred at RT for 2 h. After evaporation of solvents in vacuo, the product was isolated by column chromatography on silica gel (hexane/EtOAc, 99:1 to 90:10). The desired product was obtained as a colorless oil (794 mg, 0.83 mmol, 81%).

5

Synthesis of Compound I-25

A mixture of **10-1** (283 mg, 0.30 mmol) and dimethylamine (2M in THF, 12 mL) in a pressure flask was stirred at 68 °C (oil bath temperature) overnight. The reaction mixture was concentrated. The residue was taken up in a mixture of
10 hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a pad of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a brownish oil (302 mg). The crude product (302 mg) was purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product was obtained as a colorless oil (169 mg, 0.18 mmol, 61%). ¹HNMR
15 (400 MHz, CDCl₃ at 7.26) δ: 4.50-4.31 (br, estimated 0.3H, due to slow isomerization about amide bond), 3.97 (shouldered doublet, 5.8 Hz, 4H), 3.60 (quintet-like, 7.0 Hz, 0.7H), 3.07-3.00 (m, 2H), 2.65 (q-like, 7.6 Hz, 2H), 2.48 (q-like, 7.6 Hz, 2H), 2.29 (shouldered triplet, 7.6 Hz, 4H), 2.26, 2.25 (2 sets of singlet, 6H), 1.66-1.56 (m, 6H), 1.56-1.48 (m, 2H), 1.48-1.37 (m, 4H), 1.37-1.10 (66H), 0.91-0.85 (m, 15H).

EXAMPLE 11

SYNTHESIS OF BIS(2-HEXYLDECYL) 7-(N-DECYL-4-(PYRROLIDIN-1-
YL)BUTANAMIDO)TRIDECANEDIOATE (COMPOUND I-29)

5 Synthesis of 11-1

To a solution of 4-bromobutyric acid (0.97 mmol, 161mg) in CH_2Cl_2 (3 mL) and DMF (0.01 mL) was added oxalyl chloride (3 eq, 2.91 mmol, 370 mg, 0.25 mL) at RT. The mixture was stirred at RT overnight. The reaction mixture was then concentrated in vacuo. The residual liquid/solid (slightly yellow) was dissolved in 5 mL of CH_2Cl_2 and was added a solution of **8-2** (410 mg, 0.48 mmol), triethylamine (0.4 mL) and DMAP (2 mg) in CH_2Cl_2 (20 mL) at RT in 2 min. After addition, the resulting mixture was stirred at RT for 2.5 h. TLC (Hexane/Ethyl acetate = 9:1) showed two


major spots. The reaction mixture was then concentrated at RT under reduced pressure. The residue was used for the next reaction without any purification.

Synthesis of Compound I-29

5 The above residue containing **11-1** was taken up in a mixture of pyrrolidine (2.10 mL, 25 mmol) and THF (15 mL). The mixture was transferred into a pressure flask and heated at 68 °C overnight. The mixture was cooled and concentrated under reduce pressure. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and filtered through a short column of silica gel and washed with
10 the same solvent mixture. Concentration of the filtrate gave yellow oil/solid. The crude product (300 mg) was purified by column chromatography on silica gel (0% to 10% MeOH and 0% to 0.5% Et₃N in CH₂Cl₂). The desired product was obtained as yellow oil (215 mg). The product (215 mg) was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform). The desired product was
15 obtained as colorless oil (162 mg, 0.16 mmol, 34%). ¹HNMR (400 MHz, CDCl₃ at 7.26 ppm) δ: 4.57-4.34 (br. 0.4H), 3.98-3.94 (m, 4H), 3.64 (quintet-like, 6.8 Hz, 0.6H), 3.06-3.00 (m, 2H), 2.51-2.44 (m, 6H), 2.37-2.21(m, 6H), 1.86 (sextet-like, 7.6 Hz, 2H), 1.80-1.71 (m, 4H), 1.65-1.48 (m, 8H), 1.48-1.37 (m, 4H), 1.37-1.06 (70H), 0.91-0.86 (m, 15H).

EXAMPLE 12

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(4-(DIMETHYLAMINO)-N-(2-ETHYLHEXYL)BUTANAMIDO)NONADECANEDIOATE (COMPOUND I-30)



5 Synthesis of Compound I-30


To a solution of 4-dimethylaminobutyric acid hydrochloride (0.50 mmol, 85 mg) and 4-dimethylaminopyridine (3 equiv, DMAP, 0.75 mmol, 92 mg) in acetonitrile (5 mL) was added DCC (1.1 mmol x 2, 226, mg) and the mixture stirred at room temperature for 45 min. A solution of **9-2** (160 mg, 0.20 mmol) in CH₂Cl₂ (1 mL) was added to the reaction mixture and the resulting mixture was stirred over the weekend. More DCC (135 mg) was added and stirred for another day. No progress was observed based on TLC analysis. The reaction mixture was concentrated under reduced pressure. The residue was taken up in a mixture of hexanes and EtOAc (ca 99:5) and was filtered through a short column of silica gel. The column was washed with a mixture of hexane and EtOAc (95:5) and then a mixture of hexanes, EtOAc and Et₃N (80:20:1). The filtrate from the latter washing was concentrated to dryness (133 mg). The crude product (133 mg) was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product

was obtained as (48 mg, colorless oil, 0.053 mmol, 27%). ^1H NMR (400 MHz, CDCl_3 at 7.27 ppm) δ : 3.97 (d, 5.8 Hz, 4H), 3.70-3.60 (m, ca 1H), 3.19-2.99 (2 sets of peaks, 2H), 2.39-2.25 (m, 8H), 2.22 (singlet, 6H), 1.86-1.76 (m, 2H), 1.72-1.37 (m, ca 11H), 1.37-1.10 (60H), 0.91-0.85 (m, 18H).

5

EXAMPLE 13

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-DECYL-5-(DIBUTYLAMINO)PENTANAMIDO)NONADECANEDIOATE (COMPOUND I-31)

**Synthesis of Compound I-31**


10 A mixture of **4-3** (284 mg, 0.29 mmol), THF (10 mL), sodium iodide (5 mg) and dibutylamine (10 mmol, 1.29 g, 1.68 mL) in a pressure flask was stirred at 78 °C overnight. The reaction mixture was concentrated. The residue was taken up in a mixture of hexane, ethyl acetate and Et_3N (80:20:1) and was filtered through a pad of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a

15 brownish oil (the product and dibutylamine). The oil was diluted with hexane and washed with dilute aqueous HCl solution (0.5 M) twice, sat NaHCO_3 and brine and dried with sodium sulfate. The extract was concentrated under reduce pressure. The

crude product (309 mg) was purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product was obtained as a colorless oil (216 mg, 0.21 mmol, 72%). ¹HNMR (400 MHz, CDCl₃) δ: 4.50-4.35 (br, estimated 0.3H, due to slow isomerization about amide bond), 3.97, 3.96 (2 sets of doublets, 5.8 Hz, 4H), 3.61 (quintet-like, 7.0 Hz, 0.7H), 3.07-2.99 (m, 2H), 2.45-2.36 (m, 6H), 2.34-2.27 (m, 6H), 1.70-1.56 (m, 8H), 1.56-1.36 (m, 12H), 1.37-1.10 (70 H), 0.97-0.85 (m, 21H).

EXAMPLE 14

SYNTHESIS OF 10-(4-(DIMETHYLAMINO)-N-NONYLBUTANAMIDO)NONADECANE-1,19-DIYL BIS(2-BUTYLOCTANOATE) (COMPOUND I-32)


**Synthesis Compound I-32**

To a solution of 4-dimethylaminobutyric acid hydrochloride (2 equiv, 1.08 mmol, 181 mg) and 4-dimethylaminopyridine (3 equiv, DMAP, 1.62 mmol, 198 mg) in acetonitrile (10 mL) was added DCC (2.2 equiv, 1.18 mmol, 245 mg) and the mixture stirred at room temperature for 45 min. Then a solution of **14-1** (442 mg, 0.54 mmol) in CH₂Cl₂ (2 mL) was added. The resulting mixture was stirred overnight. More

DCC (140 mg) was added and the mixture was stirred for another day. The mixture was then concentrated under reduce pressure. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave yellow oil (381 mg). The crude product (381 mg) was purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product was obtained as colorless oil (345 mg, 0.38 mmol, 70%). ¹HNMR (400 MHz, CDCl₃ at 7.26 ppm) δ: 5.30-4.34 (br., 0.3H), 4.061, 4.056 (two sets of triplets, 6.7 Hz, 4H), 3.64 (quintet-like, 6.8 Hz, 0.7H), 3.07-3.01 (m, 2H), 2.35-2.26 (m, 6H), 2.214, 2.211 (two sets of singlet, 6H), 1.82 (sextet-like, 7.6 Hz, 2H), 1.65-1.48 (m, 10H), 1.48-1.37 (m, 8H), 1.37-1.02 (60H), 0.90-0.85 (m, 15H).

EXAMPLE 15

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-DECYL-3-(PYRROLIDIN-1-YL)PROPANAMIDO)NONADECANEDIOATE (COMPOUND I-33)


**Synthesis of Compound I-33**

A mixture of **10-1** (283 mg, 0.30 mmol), pyrrolidine (1.25 mL, 15 mmol) and THF (10 mL) in a pressure tube was heated at 64 °C overnight. The mixture

was cooled and concentrated under reduce pressure. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave yellow oil. The crude product (314 mg) was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product was obtained as colorless oil (113 mg, 0.12 mmol, 40%). ¹HNMR (400 MHz, CDCl₃ at 7.26) δ: 4.55-4.30 (br, estimated 0.3H, due to slow isomerization about amide bond), 3.97 (shouldered doublet, 5.8 Hz, 4H), 3.61 (quintet-like, 7.0 Hz, 0.7H), 3.06-3.00 (t-like, 2H), 2.81 (q-like, 7.6 Hz, 2H), 2.58-2.51(m, 6H), 2.292, 2.285 (2 sets of triplets, 7.5 Hz, 4H), 1.83-1.73 (m, 4H), 1.65-1.56 (m, 6H), 1.56-1.48 (m, 2H), 1.48-1.37 (m, 4H), 1.37-1.10 (66H), 0.91-0.85 (m, 15H).

EXAMPLE 16

SYNTHESIS OF BIS(2-BUTYLOCTYL) 10-(N-DECYL-5-(HEXYL(METHYL)AMINO)PENTANAMIDO)NONADECANEDIOATE (COMPOUND I-34)



5 Synthesis of 16-1

A mixture of **4-3** (200 mg, 0.20 mmol), hexylamine (20 mmol, 2 g), N,N-diisopropylethylamine (5 equiv., 1.0 mmol, 0.17 mL), and sodium iodide (10 mg) in acetonitrile (6 mL) was sealed and heated at 70 °C for 24 h. The reaction mixture was concentrated under reduced pressure (ca 30 mmHg) at 75 to 85 °C. TLC showed that

10 most of the excess hexylamine was removed. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of


silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a brown oil (192 mg) which was used for the next step without further purification.

Synthesis of Compound I-34

- 5 To a solution of **16-1** (192 mg, 0.19 mmol) in THF (5 mL) was added formaldehyde HCHO solution (500 mg, 37 wt. % solution in water) at RT. The resulting mixture was stirred for 30 min before introducing sodium triacetoxymethylborohydride (1.2 mmol, 243 mg). The resulting mixture was stirred at RT overnight. The reaction mixture was concentrated. The residue was taken in a mixture
- 10 of hexane and washed with dilute NaOH solution, saturated sodium bicarbonate solution and brine. After dried over sodium sulfate, the solution was concentrated to dryness (yellow oil). The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a yellow oil (220 mg). The
- 15 crude product (220 mg) was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product was obtained as colorless oil (113 mg, 0.13 mmol, 69%). ¹HNMR (400 MHz, CDCl₃ at 7.26 ppm) δ: 4.50-4.35 (br, estimated 0.3H, due to slow isomerization about amide bond), 3.97, 3.96 (2 sets of doublets, 5.8 Hz, 4H), 3.60 (quintet-like, 7.0 Hz, 0.7H),
- 20 3.06-2.98 (m, 2H), 2.36-2.26 (m, 10H), 2.191, 2.189 (2 sets of singlet, 3H), 1.70-1.56 (m, 8H), 1.56-1.36 (m, 10H), 1.37-1.10 (72 H), 0.91-0.85 (m, 18H).

EXAMPLE 17

SYNTHESIS OF BIS(2-HEXYLDECYL) 7-(N-DECYL-3-(DIMETHYLAMINO)PROPANAMIDO)TRIDECANEDIOATE (COMPOUND I-35)




5 Synthesis of Compound I-35

To a solution of 3-dimethylaminopropionic acid hydrochloride (2 equiv, 0.62 mmol, 95 mg) and 4-dimethylaminopyridine (3 equiv, DMAP, 0.93 mmol, 114 mg) in acetonitrile (10 mL) was added DCC (2.2 equiv, 0.68 mmol, 141 mg) and the mixture stirred at room temperature for 45 min. A solution of **8-2** (262 mg, 0.31 mmol) in CH_2Cl_2 (2 mL) was added and the resulting mixture was stirred over the weekend. TLC (chloroform/MeOH, 9:1) showed a major spot at the solvent front which could be

the elimination product, no starting material and a little desired product. The reaction mixture was concentrated. The possible elimination product (107 mg colorless oil) was isolated by column chromatography (hexane-EtOAc, 95:5) and was treated with a solution of dimethyl amine in THF (2M, 9 mL) at RT for 4 days. The mixture was concentrated. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a yellow oil. The crude product was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product was obtained as colorless oil (62 mg). ¹HNMR (400 MHz, CDCl₃ at 7.26) δ : 4.50-4.36 (br, estimated 0.3H, due to slow isomerization about amide bond), 3.97, 3.96 (two sets of doublets, 5.8 Hz, 4H), 3.61 (quintet-like, 7.0 Hz, 0.7H), 3.07-3.00 (m, 2H), 2.65 (q-like, 7.2 Hz, 2H), 2.53-2.41 (m, 2H), 2.31-2.26 (m, 4H), 2.26, 2.25 (2 sets of singlet, 6H), 1.66-1.56 (m, 6H), 1.56-1.48 (m, 2H), 1.48-1.37 (m, 4H), 1.37-1.10 (70H), 0.91-0.85 (m, 15H)

EXAMPLE 18

SYNTHESIS OF BIS(2-HEXYLDECYL) 7-(3-(DIMETHYLAMINO)-N-(6-((2-ETHYLHEXYL)OXY)-6-OXOHXYL)PROPANAMIDO)TRIDECANEDIOATE (COMPOUND I-36)



5 Synthesis of 18-2

To a solution of 3-bromopropionic acid (0.34 mmol, 52 mg) in CH₂Cl₂ (3 mL) and DMF (1 drop from a thin needle) was added oxalyl chloride (0.86 mmol, 109 mg, 74 uL) at RT. The mixture was stirred at RT overnight. The mixture was then concentrated under reduced pressure for 60 min at RT. The residual liquid/solid (yellow) was dissolved in 5 mL of CH₂Cl₂ and added to a solution of **18-1** (160 mg,


0.17 mmol) and triethylamine (0.86 mmol, 0.12 mL) and DMAP (1 mg) in CH₂Cl₂ (5 mL) at RT in 1 min. After addition, the mixture was stirred at RT for 3 h and then concentrated. The product was isolated by column chromatography on silica gel (hexane, EtOAc, and Et₃N, from 95:5 to 85:15). The desired product was obtained as a
5 colorless oil (99 mg, 0.09 mmol, 53%).

Synthesis of Compound I-36

To a pressure flask containing **18-2** (99 mg, 0.09 mmol) was added dimethylamine (2M in THF, 5 mL). The solution was stirred at 68 °C (oil bath temp)
10 for 2 days. The mixture was cooled and concentrated under reduce pressure. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a brown oil (94 mg). The product (94 mg) was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in
15 chloroform with a trace of Et₃N). The desired product was obtained as colorless oil (71 mg, 0.069 mmol, 76%). ¹HNMR (400 MHz, CDCl₃ at 7.26) δ: 4.53-4.35 (br, estimated 0.3H, due to slow isomerization about amide bond), 4.02-3.93 (m, 6H), 3.62 (quintet-like, 7.0 Hz, 0.7H), 3.07-3.01 (m, 2H), 2.65 (q-like, 7.6 Hz, 2H), 2.50-2.44 (m, 2H), 2.34-2.26 (m, 6H), 2.26, 2.25 (2 sets of singlet, 6H), 1.69-1.56-1.48 (m, estimated 11H,
20 overlapped with water peak), 1.49-1.37 (m, 4H), 1.37-1.10 (66H), 0.92-0.86 (m, 18H).

EXAMPLE 19

SYNTHESIS OF DI(TRIDECAN-7-YL) 10-(N-DECYL-4-(DIMETHYLAMINO)BUTANAMIDO)NONADECANEDIOATE (COMPOUND I-37)

5 **Synthesis of 19-2**

A solution of **19-1** (1 eq., 0.493 g, 0.70 mmol) and 1-decylamine (1.5 eq, 1.05 mmol, 165 mg, 0.21 mL) in DCE (10 mL) was stirred at RT for about 15 min. To the solution was added sodium triacetoxyborohydride (1.5 eq., 1.05 mmol, 222 mg) and AcOH (1.5 eq, 1.05 mmol, 63 mg, 0.059 mL). The mixture was stirred at RT for 2 days.

- 10 The reaction mixture was then concentrated. The residue was diluted with hexanes and washed with dilute NaOH, sat NaHCO_3 and brine. After the organic extract was dried over sodium sulfate, the solvent was removed under reduce pressure. The residue was

taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave the desired product as a colorless oil (582 mg, 0.69 mmol, 98%). The product was used for the next step without further purification.

5

Synthesis of Compound I-37


To a solution of 4-dimethylaminobutyric acid hydrochloride (1.71 mmol, 287 mg) and 4-dimethylaminopyridine (3 equiv, DMAP, 2.07 mmol, 253 mg) in acetonitrile (15 mL) was added DCC (2.2 equiv, 1.52 mmol, 313 mg) and the mixture stirred at room temperature for 45 min. A solution of **19-2** (582 mg, 0.69 mmol) in CH₂Cl₂ (3 mL) was added and the resulting mixture was stirred overnight. On the next day more DCC (200 mg) was added and stirring was continued for the week end (4 days). The mixture was then concentrated under reduced pressure. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a yellow oil. The product was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product was obtained as colorless oil.

10

15

EXAMPLE 20

SYNTHESIS OF BIS(2-HEXYLDECYL) 10-(N-DECYL-4-(DIMETHYLAMINO)BUTANAMIDO)NONADECANEDIOATE (COMPOUND I-38)



5 Synthesis of 20-2

A solution of the ketone **20-1** (0.92 g, 1.16 mmol) and 1-decylamine (2.03 mmol, 319 mg, 0.40 mL) in DCE (6 mL) was stirred at RT for 15 min, followed by addition of sodium triacetoxyborohydride (2.03 mmol, 429 mg) and AcOH (2.03 mmol, 121 mg, 0.115 mL). After the mixture was stirred at RT for 2 days, the reaction mixture was concentrated. The residue was diluted with hexanes and washed with dilute NaOH, saturated NaHCO₃ and brine. The organic phase was separated, dried over

sodium sulfate. The extract was filtered through a short column of silica gel and the column was washed with a mixture of hexane/EtOAc/Et₃N, 95:5:0 to 80:20:1). The desired product was obtained as a colorless oil (814 mg colorless oil, 0.87 mmol, 75% yield).

5


Synthesis of I-38

To a solution of 4-dimethylaminobutyric acid hydrochloride (2 equiv, 0.76 mmol, 127 mg), 4-dimethylaminopyridine (3 eq, 1.14 mmol, 139 mg) in CH₃CN (5 mL) was added DCC (2.2 equiv, 0.84 mmol, 172 mg) and the mixture stirred at room temperature for 45 min. A solution of **20-2** (350 mg, 0.38 mmol) in DCM (1 mL) was added and the resulting mixture was stirred overnight. On the next day, more DCC (50 mg) was added and stirred for another day. The mixture was then concentrated under reduce pressure. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave a colorless oil. The product was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform with a trace of Et₃N). The desired product was obtained as colorless oil (260 mg).

10

15

EXAMPLE 21

SYNTHESIS OF BIS(2-HEXYLDECYL) 10-(N-DECYL-4-(PYRROLIDIN-1-
YL)BUTANAMIDO)NONADECANEDIOATE (COMPOUND I-39)

5 Synthesis of 21-1

To a solution of 4-bromobutyric acid (1.00 mmol, 167mg) in DCM (3 mL) and DMF (one tiny drop) was added oxalyl chloride (3 eq, 3.00 mmol, 381 mg, 0.26 mL) at RT. The mixture was stirred at RT overnight. The reaction mixture was then concentrated in vacuo. The residual liquid/solid (slightly yellow) was dissolved in 5 mL of DCM and was added a solution of **20-2** (464 mg, 0.50 mmol) and triethylamine (0.42 mL) and DMAP (2 mg) in DCM (5 mL) at RT in 2 min. After addition, the resulting mixture was stirred at RT for 2.5 h. TLC (Hexane/Ethyl acetate = 9:1) showed two major spots. The reaction mixture was then concentrated at RT under reduced pressure. The residue was used for the next reaction without any purification.


Synthesis of I-39

The above residue was taken up in a mixture of pyrrolidine (2.20 mL, 26 mmol) and THF (15 mL). The mixture was transferred into a pressure flask and heated at 68 C overnight. The mixture was cooled and concentrated under reduced pressure. The residue was taken up in a mixture of hexane, ethyl acetate and Et₃N (80:20:1) and was filtered through a short column of silica gel and washed with the same solvent mixture. Concentration of the filtrate gave yellow oil/solid (359 mg). The crude product (359 mg) was further purified by flash dry column chromatography on silica gel (0 to 5% MeOH in chloroform). The desired product was obtained as colorless oil (165 mg).

The various embodiments described above can be combined to provide further embodiments. All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, including U.S. 62/791,566 and 62/890,469, are incorporated herein by reference, in their entirety. Aspects of the embodiments can be modified, if necessary to employ concepts of the various patents, applications and publications to provide yet further embodiments. These and other changes can be made to the embodiments in light of the above-detailed description. In general, in the following claims, the terms used should not be construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

CLAIMS

1. A compound having the following structure (I):



or a pharmaceutically acceptable salt, tautomer, or stereoisomer thereof, wherein:

R^1 is optionally substituted C_1 - C_{24} alkyl or optionally substituted C_2 - C_{24} alkenyl;

R^2 and R^3 are each independently optionally substituted C_1 - C_{36} alkyl;

R^4 and R^5 are each independently optionally substituted C_1 - C_6 alkyl, or R^4 and R^5 join, along with the nitrogen atom to which they are attached, to form an optionally substituted heterocyclyl ring or an optionally substituted heteroaryl ring;


L^1 , L^2 , and L^3 are each independently optionally substituted C_1 - C_{18} alkylene;

G^1 is a direct bond, $-(CH_2)_nO(C=O)-$, $-(CH_2)_n(C=O)O-$, or $-(C=O)-$;


G^2 and G^3 are each independently $-(C=O)O-$ or $-O(C=O)-$; and

n is an integer greater than 0.

2. The compound of claim 1, having the following structure (IA):




3. The compound of claim 1, having the following structure (IB):



(IB)

4. The compound of claim 1, wherein R¹ is optionally substituted C₆-C₁₈ alkyl or optionally substituted C₁₄-C₁₈ alkenyl.
5. The compound of claim 1, wherein G¹ is a direct bond, -(CH₂)_nO(C=O)-, or -(CH₂)_n(C=O)O-.
6. The compound of claim 5, wherein G¹ is a direct bond.
7. The compound of claim 5, wherein G¹ is -(CH₂)_n(C=O)O- and n is greater than 1.
8. The compound of claim 1, wherein L¹ is unsubstituted C₁-C₆ alkylene.
9. The compound of claim 1, wherein R² and R³ are both unsubstituted C₈-C₂₄ alkyl.
10. The compound of claim 1, wherein R² and R³ are optionally substituted branched C₁-C₃₆ alkyl.


11. The compound of claim 1, wherein R^2 and R^3 each independently have one of the following structures:



wherein:

R^6 and R^7 are each independently unsubstituted C_2 - C_{12} alkyl.

12. The compound of claim 11, wherein R^2 and R^3 each independently have one of the following structures:




13. The compound of claim 1, wherein L^2 and L^3 are each independently unsubstituted C_4 - C_{10} alkylene.

14. The compound of claim 1, wherein R^4 and R^5 are each independently unsubstituted C_1 - C_6 alkyl.


15. The compound of claim 14, wherein R^4 and R^5 are both methyl, ethyl or n-butyl.

16. The compound of claim 1, wherein R^4 and R^5 join, along with the nitrogen atom to which they are attached, to form an optionally substituted heterocyclyl ring.

17. The compound of claim 16, wherein the optionally substituted heterocyclyl ring has the following structure:





18. The compound of claim 1, wherein the compound has one of the following structures:














19. A composition comprising the compound of claim 1 and a therapeutic agent.
20. A lipid nanoparticle comprising the compound of claim 1.
21. The composition of claim 19, wherein the therapeutic agent comprises messenger RNA.
22. The lipid nanoparticle of claim 20, further comprising messenger RNA.