

PROCESS FOR PREPARING POLYAMIDE MICROCAPSULES

Technical Field

The present invention relates to a new process for the preparation of polyamide
5 microcapsules. Polyamide microcapsules obtainable by said process are also an object of
the invention. Perfuming compositions and consumer products comprising said
microcapsules, in particular perfumed consumer products in the form of home care or
personal care products, are also part of the invention.

10 **Background of the Invention**

One of the problems faced by the perfumery industry lies in the relatively rapid
loss of olfactive benefit provided by odoriferous compounds due to their volatility,
particularly that of “top-notes”. In order to tailor the release rates of volatiles, delivery
systems such as microcapsules containing a perfume are needed to protect and later
15 release the core payload when triggered. A key requirement from the industry regarding
these systems is to survive suspension in challenging bases without physically
dissociating or degrading. This is referred to as *stability* for the delivery system. For
instance, fragranced personal and household cleansers containing high levels of
aggressive surfactant detergents are very challenging for the stability of microcapsules.

20 Polyurea and polyurethane-based microcapsule slurry are widely used for example
in perfumery industry for instance as they provide a long lasting pleasant olfactory effect
after their applications on different substrates. Those microcapsules have been widely
disclosed in the prior art (see for example WO2007/004166 or EP 2300146 from the
Applicant).

25 Therefore, there is still a need to provide new microcapsules, while not
compromising on the performance of the microcapsules, in particular in terms of stability
in a challenging medium such as a consumer product base, as well as in delivering a good
performance in terms of hydrophobic material delivery, e.g. olfactive performance in the
case of perfuming ingredients.

The present invention is proposing a solution to the above-mentioned problem by providing a new process for the preparation of polyamide microcapsules.

Summary of the Invention

5 It has now been found that performing core-shell microcapsules encapsulating hydrophobic material, preferably hydrophobic active ingredient could be obtained by reacting an acyl chloride with at least one amino-compound during the interfacial polymerization. The process of the invention therefore provides a solution to the above-mentioned problems as it allows preparing microcapsules with the desired stability in
10 challenging bases.

In a first aspect, the present invention relates to a process for preparing a polyamide core-shell microcapsule slurry comprising the following steps:

- a) Dissolving at least one acyl chloride and at least one stabilizer in a hydrophobic material, preferably a perfume to form an oil phase;
 - 15 b) Dispersing the oil phase obtained in step a) into a water phase comprising optionally an amino compound A or a base to form an oil-in-water emulsion; and
 - c) Performing a curing step to form polyamide microcapsules in the form of a slurry.
- wherein at least an amino compound B is added in the water phase before the formation of the oil-in-water emulsion and/or in the oil-in water emulsion obtained after step b)

20 A second object of the invention is a polyamide core-shell microcapsule slurry obtainable by the process as defined above.

A perfuming composition comprising:

- (i) microcapsule slurry as defined above, wherein the hydrophobic material comprises a perfume,
 - 25 (ii) at least one ingredient selected from the group consisting of a perfumery carrier and a perfumery base
 - (iii) optionally at least one perfumery adjuvant,
- is another object of the invention.

30 Another object of the invention is a consumer product comprising:

- a personal care active base, and
- microcapsules as defined above or the perfuming composition as defined above, wherein the consumer product is in the form of a personal care composition.

5 Another object of the invention is a consumer product comprising:

- a home care or a fabric care active base, and
- microcapsules as defined above or the perfuming composition as defined above, wherein the consumer product is in the form of a home care or a fabric care composition.

10

Detailed description of the invention

Unless stated otherwise, percentages (%) are meant to designate a percentage by weight of a composition.

15 By “**active ingredient**”, it is meant a single compound or a combination of ingredients.

By “**perfume or flavour oil**”, it is meant a single perfuming or flavouring compound or a mixture of several perfuming or flavouring compounds.

By “**consumer product**” or “**end-product**” it is meant a manufactured product ready to be distributed, sold and used by a consumer.

20 For the sake of clarity, by the expression “**dispersion**” in the present invention it is meant a system in which particles are dispersed in a continuous phase of a different composition and it specifically includes a suspension or an emulsion.

25 A “**microcapsule**”, or the similar, in the present invention, it is meant that core-shell microcapsules have a particle size distribution in the micron range (e.g. a mean diameter ($d(v, 0.5)$) comprised between about 1 and 3000 microns) and comprise an external solid polymer-based shell and an internal continuous oil phase enclosed by the external shell.

By “**amino-compound**” it should be understood a compound having at least two reactive amine groups.

In the present invention, the wordings “**acyl chloride**” or “**acid chloride**” are used indifferently.

By “**polyamide microcapsules**”, it means that the microcapsule’s shell comprises a polyamide material. The wording “polyamide microcapsules” can also encompass a
5 shell made of a composite comprising a polyamide material and another material, for example a biopolymer.

It has been found that polyamide core-shell microcapsules with overall good performance in terms of stability in challenging bases could be obtained when a stabilizer is added in the oil phase during the process.

10

Process for preparing a polyamide microcapsule slurry

A first object of the invention is therefore a process for preparing a polyamide core-shell microcapsule slurry comprising the following steps:

- a) Dissolving at least one acyl chloride and at least one stabilizer in a hydrophobic
15 material, preferably a perfume to form an oil phase;
- b) Dispersing the oil phase obtained in step a) into a water phase comprising optionally an amino compound A or a base to form an oil-in-water emulsion; and
- c) Performing a curing step to form polyamide microcapsules in the form of a slurry.
wherein at least an amino compound B is added in the water phase before the formation
20 of the oil-in-water emulsion and/or in the oil-in water emulsion obtained after step b)

According to an embodiment, the process comprises the steps of:

- a) Dissolving at least one acyl chloride and at least one stabilizer in a hydrophobic material, preferably a perfume to form an oil phase;
- b) Dispersing the oil phase obtained in step a) into a water phase comprising optionally an
25 amino compound A or a base to form an oil-in-water emulsion;
- c) Adding to the oil-in-water emulsion obtained in step b) an amino compound B; and
- d) Performing a curing step to form polyamide microcapsules in the form of a slurry.

In one step of the process, an oil phase is formed by admixing at least one hydrophobic material with at least one acyl chloride and a stabilizer.

- 30 According to a particular embodiment, the acyl chloride is chosen in the group consisting of benzene-1,3,5-tricarbonyl chloride, benzene-1,2,4-tricarbonyl trichloride, benzene-

1,2,4,5-tetracarbonyl tetrachloride, cyclohexane-1,3,5-tricarbonyl trichloride, isophthaloyl dichloride, diglycolyl dichloride, succinic dichloride, and mixtures thereof.

The weight ratio between acyl chloride and the hydrophobic material is preferably comprised between 0.01 and 0.09, more preferably between 0.03 and 0.07.

5 The acyl chloride can be dissolved directly in the perfume oil or can be pre-dispersed in an inert solvent such as benzyl benzoate before mixing with the hydrophobic material, preferably a perfume oil.

According to a particular embodiment, a polyisocyanate having at least two isocyanate functional groups is added in the oil phase.

10 Suitable polyisocyanates used according to the invention include aromatic polyisocyanate, aliphatic polyisocyanate and mixtures thereof. Said polyisocyanate comprises at least 2, preferably at least 3 but may comprise up to 6, or even only 4, isocyanate functional groups. According to a particular embodiment, a triisocyanate (3 isocyanate functional group) is used.

15 According to one embodiment, said polyisocyanate is an aromatic polyisocyanate.

The term "aromatic polyisocyanate" is meant here as encompassing any polyisocyanate comprising an aromatic moiety. Preferably, it comprises a phenyl, a toluyl, a xylyl, a naphthyl or a diphenyl moiety, more preferably a toluyl or a xylyl moiety. Preferred aromatic polyisocyanates are biurets, polyisocyanurates and trimethylol propane adducts of diisocyanates, more preferably comprising one of the above-cited specific aromatic moieties. More preferably, the aromatic polyisocyanate is a polyisocyanurate of toluene diisocyanate (commercially available from Bayer under the tradename Desmodur[®] RC), a trimethylol propane-adduct of toluene diisocyanate (commercially available from Bayer under the tradename Desmodur[®] L75), a trimethylol propane-adduct of xylylene diisocyanate (commercially available from Mitsui Chemicals under the tradename Takenate[®] D-110N). In a most preferred embodiment, the aromatic polyisocyanate is a trimethylol propane-adduct of xylylene diisocyanate.

25 According to another embodiment, said polyisocyanate is an aliphatic polyisocyanate. The term "aliphatic polyisocyanate" is defined as a polyisocyanate which
30 does not comprise any aromatic moiety. Preferred aliphatic polyisocyanates are a trimer

of hexamethylene diisocyanate, a trimer of isophorone diisocyanate, a trimethylol propane-adduct of hexamethylene diisocyanate (available from Mitsui Chemicals) or a biuret of hexamethylene diisocyanate (commercially available from Bayer under the tradename Desmodur[®] N 100), among which a biuret of hexamethylene diisocyanate is
5 even more preferred.

According to another embodiment, the at least one polyisocyanate is in the form of a mixture of at least one aliphatic polyisocyanate and of at least one aromatic polyisocyanate, both comprising at least two or three isocyanate functional groups, such as a mixture of a biuret of hexamethylene diisocyanate with a trimethylol propane-adduct
10 of xylylene diisocyanate, a mixture of a biuret of hexamethylene diisocyanate with a polyisocyanurate of toluene diisocyanate and a mixture of a biuret of hexamethylene diisocyanate with a trimethylol propane-adduct of toluene diisocyanate. Most preferably, it is a mixture of a biuret of hexamethylene diisocyanate with a trimethylol propane-adduct of xylylene diisocyanate. Preferably, when used as a mixture the molar ratio
15 between the aliphatic polyisocyanate and the aromatic polyisocyanate is ranging from 80:20 to 10:90.

According to an embodiment, the at least one polyisocyanate used in the process of the invention is present in amounts representing from 0.1 to 15%, preferably from 0.5 to 10% and more preferably from 0.8 to 6%, and even more preferably between 1 and 3%
20 by weight based on the total amount of the oil phase.

Hydrophobic material

According to an embodiment, the hydrophobic material is a hydrophobic active ingredient. By “hydrophobic active ingredient”, it is meant any hydrophobic active ingredient –
25 single ingredient or a mixture of ingredients - which forms a two-phase dispersion when mixed with water. The hydrophobic active ingredient is liquid at about 20°C.

Hydrophobic active ingredients are preferably chosen from the group consisting of flavor, flavor ingredients, perfume, perfume ingredients, nutraceuticals, cosmetics, pest control agents, biocide actives and mixtures thereof.

According to a particular embodiment, the hydrophobic active ingredient comprises a mixture of a perfume with another ingredient selected from the group consisting of nutraceuticals, cosmetics, pest control agents and biocide actives.

According to a particular embodiment, the hydrophobic active ingredient
5 comprises a mixture of biocide actives with another ingredient selected from the group consisting of perfume, nutraceuticals, cosmetics, pest control agents.

According to a particular embodiment, the hydrophobic active ingredient comprises a mixture of pest control agents with another ingredient selected from the group consisting of perfume, nutraceuticals, cosmetics, biocide actives.

10 According to a particular embodiment, the hydrophobic active ingredient comprises a perfume.

According to a particular embodiment, the hydrophobic active ingredient consists of a perfume.

According to a particular embodiment, the hydrophobic active ingredient consists
15 of biocide actives.

According to a particular embodiment, the hydrophobic active ingredient consists of pest control agents.

By “perfume” (or also “perfume oil”) what is meant here is an ingredient or composition that is a liquid at about 20°C. According to any one of the above
20 embodiments said perfume oil can be a perfuming ingredient alone or a mixture of ingredients in the form of a perfuming composition. As a “perfuming ingredient” it is meant here a compound, which is used for the primary purpose of conferring or modulating an odour. In other words such an ingredient, to be considered as being a perfuming one, must be recognized by a person skilled in the art as being able to at least
25 impart or modify in a positive or pleasant way the odor of a composition, and not just as having an odor. For the purpose of the present invention, perfume oil also includes combination of perfuming ingredients with substances which together improve, enhance or modify the delivery of the perfuming ingredients, such as perfume precursors, emulsions or dispersions, as well as combinations which impart an additional benefit
30 beyond that of modifying or imparting an odor, such as long-lasting, blooming, malodour counteraction, antimicrobial effect, microbial stability, pest control.

The nature and type of the perfuming ingredients present in the oil phase do not warrant a more detailed description here, which in any case would not be exhaustive, the skilled person being able to select them on the basis of its general knowledge and according to intended use or application and the desired organoleptic effect. In general terms, these perfuming ingredients belong to chemical classes as varied as alcohols, aldehydes, ketones, esters, ethers, acetates, nitriles, terpenoids, nitrogenous or sulphurous heterocyclic compounds and essential oils, and said perfuming co-ingredients can be of natural or synthetic origin. Many of these co-ingredients are in any case listed in reference texts such as the book by S. Arctander, *Perfume and Flavor Chemicals*, 1969, Montclair, New Jersey, USA, or its more recent versions, or in other works of a similar nature, as well as in the abundant patent literature in the field of perfumery. It is also understood that said ingredients may also be compounds known to release in a controlled manner various types of perfuming compounds.

In particular one may cite perfuming ingredients which are commonly used in perfume formulations, such as:

- Aldehydic ingredients: decanal, dodecanal, 2-methyl-undecanal, 10-undecenal, octanal, nonanal and/or nonenal;
- Aromatic-herbal ingredients: eucalyptus oil, camphor, eucalyptol, 5-methyltricyclo[6.2.1.0~2,7~]undecan-4-one, 1-methoxy-3-hexanethiol, 2-ethyl-4,4-dimethyl-1,3-oxathiane, 2,2,7/8,9/10-Tetramethylspiro[5.5]undec-8-en-1-one, menthol and/or alpha-pinene;
- Balsamic ingredients: coumarin, ethylvanillin and/or vanillin;
- Citrus ingredients: dihydromyrcenol, citral, orange oil, linalyl acetate, citronellyl nitrile, orange terpenes, limonene, 1-p-menthen-8-yl acetate and/or 1,4(8)-p-menthadiene;
- Floral ingredients: methyl dihydrojasmonate, linalool, citronellol, phenylethanol, 3-(4-tert-butylphenyl)-2-methylpropanal, hexylcinnamic aldehyde, benzyl acetate, benzyl salicylate, tetrahydro-2-isobutyl-4-methyl-4(2H)-pyranol, beta ionone, methyl 2-(methylamino)benzoate, (E)-3-methyl-4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-one, (1E)-1-(2,6,6-trimethyl-2-cyclohexen-1-yl)-1-penten-3-one, 1-(2,6,6-trimethyl-1,3-cyclohexadien-1-yl)-2-buten-1-one, (2E)-1-(2,6,6-

- trimethyl-2-cyclohexen-1-yl)-2-buten-1-one, (2E)-1-[2,6,6-trimethyl-3-cyclohexen-1-yl]-2-buten-1-one, (2E)-1-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2-buten-1-one, 2,5-dimethyl-2-indanmethanol, 2,6,6-trimethyl-3-cyclohexene-1-carboxylate, 3-(4,4-dimethyl-1-cyclohexen-1-yl)propanal, hexyl salicylate, 3,7-dimethyl-1,6-nonadien-3-ol, 3-(4-isopropylphenyl)-2-methylpropanal, verdyl acetate, geraniol, p-menth-1-en-8-ol, 4-(1,1-dimethylethyl)-1-cyclohexyle acetate, 1,1-dimethyl-2-phenylethyl acetate, 4-cyclohexyl-2-methyl-2-butanol, amyl salicylate, high cis methyl dihydrojasmonate, 3-methyl-5-phenyl-1-pentanol, verdyl propionate, geranyl acetate, tetrahydro linalool, cis-7-p-menthanol, propyl (S)-2-(1,1-dimethylpropoxy)propanoate, 2-methoxynaphthalene, 2,2,2-trichloro-1-phenylethyl acetate, 4/3-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1-carbaldehyde, amylcinnamic aldehyde, 8-decen-5-olide, 4-phenyl-2-butanone, isononyl acetate, 4-(1,1-dimethylethyl)-1-cyclohexyl acetate, verdyl isobutyrate and/or mixture of methylionones isomers;
- 15 - Fruity ingredients: gamma-undecalactone, 2,2,5-trimethyl-5-pentylcyclopentanone, 2-methyl-4-propyl-1,3-oxathiane, 4-decanolide, ethyl 2-methyl-pentanoate, hexyl acetate, ethyl 2-methylbutanoate, gamma-nonolactone, allyl heptanoate, 2-phenoxyethyl isobutyrate, ethyl 2-methyl-1,3-dioxolane-2-acetate, 3-(3,3/1,1-dimethyl-5-indanyl)propanal, diethyl 1,4-cyclohexanedicarboxylate, 3-methyl-2-hexen-1-yl acetate, 1-[3,3-dimethylcyclohexyl]ethyl [3-ethyl-2-oxiranyl]acetate and/or diethyl 1,4-cyclohexane dicarboxylate;
- 20 - Green ingredients: 2-methyl-3-hexanone (E)-oxime, 2,4-dimethyl-3-cyclohexene-1-carbaldehyde, 2-tert-butyl-1-cyclohexyl acetate, styrallyl acetate, allyl (2-methylbutoxy)acetate, 4-methyl-3-decen-5-ol, diphenyl ether, (Z)-3-hexen-1-ol and/or 1-(5,5-dimethyl-1-cyclohexen-1-yl)-4-penten-1-one;
- 25 - Musk ingredients: 1,4-dioxa-5,17-cycloheptadecanedione, (Z)-4-cyclopentadecen-1-one, 3-methylcyclopentadecanone, 1-oxa-12-cyclohexadecen-2-one, 1-oxa-13-cyclohexadecen-2-one, (9Z)-9-cycloheptadecen-1-one, 2-{1S)-1-[(1R)-3,3-dimethylcyclohexyl]ethoxy}-2-oxoethyl propionate 3-methyl-5-cyclopentadecen-1-one, 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethyl-cyclopenta-g-2-benzopyrane, 30 (1S,1'R)-2-[1-(3',3'-dimethyl-1'-cyclohexyl)ethoxy]-2-methylpropyl propanoate,

oxacyclohexadecan-2-one and/or (1S,1'R)-[1-(3',3'-dimethyl-1'-cyclohexyl)ethoxycarbonyl]methyl propanoate, ;

- Woody ingredients: 1-[(1R,6SR)-2,2,6-trimethylcyclohexyl]-3-hexanol, 3,3-dimethyl-5-[(1R)-2,2,3-trimethyl-3-cyclopenten-1-yl]-4-penten-2-ol, 3,4'-dimethylspiro[oxirane-2,9'-tricyclo[6.2.1.0^{2,7}]undec[4]ene, (1-ethoxyethoxy)cyclododecane, 2,2,9,11-tetramethylspiro[5.5]undec-8-en-1-yl acetate, 1-(octahydro-2,3,8,8-tetramethyl-2-naphthalenyl)-1-ethanone, patchouli oil, terpenes fractions of patchouli oil, clearwood[®], (1'R,E)-2-ethyl-4-(2',2',3'-trimethyl-3'-cyclopenten-1'-yl)-2-buten-1-ol, 2-ethyl-4-(2,2,3-trimethyl-3-cyclopenten-1-yl)-2-buten-1-ol, methyl cedryl ketone, 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol, 1-(2,3,8,8-tetramethyl-1,2,3,4,6,7,8,8a-octahydronaphthalen-2-yl)ethan-1-one and/or isobornyl acetate;
- Other ingredients (e.g. amber, powdery spicy or watery): dodecahydro-3a,6,6,9a-tetramethyl-naphtho[2,1-b]furan and any of its stereoisomers, heliotropin, anisic aldehyde, eugenol, cinnamic aldehyde, clove oil, 3-(1,3-benzodioxol-5-yl)-2-methylpropanal, 7-methyl-2H-1,5-benzodioxepin-3(4H)-one, 2,5,5-trimethyl-1,2,3,4,4a,5,6,7-octahydro-2-naphthalenol, 1-phenylvinyl acetate, 6-methyl-7-oxa-1-thia-4-azaspiro[4.4]nonan and/or 3-(3-isopropyl-1-phenyl)butanal.

It is also understood that said ingredients may also be compounds known to release in a controlled manner various types of perfuming compounds also known as properfume or profragrance. Non-limiting examples of suitable properfume may include 4-(dodecylthio)-4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-2-butanone, 4-(dodecylthio)-4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2-butanone, trans-3-(dodecylthio)-1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-1-butanone, 2-phenylethyl oxo(phenyl)acetate or a mixture thereof.

The perfuming ingredients may be dissolved in a solvent of current use in the perfume industry. The solvent is preferably not an alcohol. Examples of such solvents are diethyl phthalate, isopropyl myristate, Abalyn[®] (rosin resins, available from Eastman), benzyl benzoate, ethyl citrate, limonene or other terpenes, or isoparaffins. Preferably, the solvent is very hydrophobic and highly sterically hindered, like for example Abalyn[®] or benzyl benzoate. Preferably the perfume comprises less than 30% of solvent. More preferably the perfume comprises less than 20% and even more preferably less than 10%

of solvent, all these percentages being defined by weight relative to the total weight of the perfume. Most preferably, the perfume is essentially free of solvent.

Preferred perfuming ingredients are those having a high steric hindrance and in particular those from one of the following groups:

- 5 - Group 1: perfuming ingredients comprising a cyclohexane, cyclohexene, cyclohexanone or cyclohexenone ring substituted with at least one linear or branched C₁ to C₄ alkyl or alkenyl substituent;
- Group 2: perfuming ingredients comprising a cyclopentane, cyclopentene, cyclopentanone or cyclopentenone ring substituted with at least one linear or
10 branched C₄ to C₈ alkyl or alkenyl substituent;
- Group 3: perfuming ingredients comprising a phenyl ring or perfuming ingredients comprising a cyclohexane, cyclohexene, cyclohexanone or cyclohexenone ring substituted with at least one linear or branched C₅ to C₈ alkyl or alkenyl substituent or with at least one phenyl substituent and optionally one or more linear or branched C₁
15 to C₃ alkyl or alkenyl substituents;
- Group 4: perfuming ingredients comprising at least two fused or linked C₅ and/or C₆ rings;
- Group 5: perfuming ingredients comprising a camphor-like ring structure;
- Group 6: perfuming ingredients comprising at least one C₇ to C₂₀ ring structure;
- 20 - Group 7: perfuming ingredients having a logP value above 3.5 and comprising at least one tert-butyl or at least one trichloromethyl substituent;

Examples of ingredients from each of these groups are:

- Group 1: 2,4-dimethyl-3-cyclohexene-1-carbaldehyde (origin: Firmenich SA, Geneva, Switzerland), isocyclocitral, menthone, isomenthone, Romascone[®] (methyl 2,2-
25 dimethyl-6-methylene-1-cyclohexanecarboxylate, origin: Firmenich SA, Geneva, Switzerland), nerone, terpineol, dihydroterpineol, terpenyl acetate, dihydroterpenyl acetate, dipentene, eucalyptol, hexylate, rose oxide, Perycorolle[®] ((S)-1,8-p-menthadiene-7-ol, origin: Firmenich SA, Geneva, Switzerland), 1-p-menthene-4-ol, (1RS,3RS,4SR)-3-p-mentanyl acetate, (1R,2S,4R)-4,6,6-trimethyl-
30 bicyclo[3,1,1]heptan-2-ol, Doremox[®] (tetrahydro-4-methyl-2-phenyl-2H-pyran, origin: Firmenich SA, Geneva, Switzerland), cyclohexyl acetate, cyclanol acetate,

- Fructalate[®] (1,4-cyclohexane diethyldicarboxylate, origin: Firmenich SA, Geneva, Switzerland), Koumalactone[®] ((3ARS,6SR,7ASR)-perhydro-3,6-dimethyl-benzo[B]furan-2-one, origin: Firmenich SA, Geneva, Switzerland), Natactone[®] ((6R)-perhydro-3,6-dimethyl-benzo[B]furan-2-one, origin: Firmenich SA, Geneva, Switzerland), 2,4,6-trimethyl-4-phenyl-1,3-dioxane, 2,4,6-trimethyl-3-cyclohexene-1-carbaldehyde;
- 5 - Group 2: (E)-3-methyl-5-(2,2,3-trimethyl-3-cyclopenten-1-yl)-4-penten-2-ol (origin: Givaudan SA, Vernier, Switzerland), (1'R,E)-2-ethyl-4-(2',2',3'-trimethyl-3'-cyclopenten-1'-yl)-2-buten-1-ol (origin: Firmenich SA, Geneva, Switzerland),
- 10 Polysantol[®] ((1'R,E)-3,3-dimethyl-5-(2',2',3'-trimethyl-3'-cyclopenten-1'-yl)-4-penten-2-ol, origin: Firmenich SA, Geneva, Switzerland), fleuramone, Hedione[®] HC (methyl-cis-3-oxo-2-pentyl-1-cyclopentane acetate, origin: Firmenich SA, Geneva, Switzerland), Veloutone[®] (2,2,5-Trimethyl-5-pentyl-1-cyclopentanone, origin: Firmenich SA, Geneva, Switzerland), Nirvanol[®] (3,3-dimethyl-5-(2,2,3-trimethyl-3-
- 15 cyclopenten-1-yl)-4-penten-2-ol, origin: Firmenich SA, Geneva, Switzerland), 3-methyl-5-(2,2,3-trimethyl-3-cyclopenten-1-yl)-2-pentanol (origin, Givaudan SA, Vernier, Switzerland);
- 20 - Group 3: damascones, Neobutenone[®] (1-(5,5-dimethyl-1-cyclohexen-1-yl)-4-penten-1-one, origin: Firmenich SA, Geneva, Switzerland), nectalactone ((1'R)-2-[2-(4'-methyl-3'-cyclohexen-1'-yl)propyl]cyclopentanone), alpha-ionone, beta-ionone, damascenone, Dynascone[®] (mixture of 1-(5,5-dimethyl-1-cyclohexen-1-yl)-4-penten-1-one and 1-(3,3-dimethyl-1-cyclohexen-1-yl)-4-penten-1-one, origin: Firmenich SA, Geneva, Switzerland), Dorinone[®] beta (1-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2-
- 25 buten-1-one, origin: Firmenich SA, Geneva, Switzerland), Romandolide[®] ((1S,1'R)-[1-(3',3'-Dimethyl-1'-cyclohexyl)ethoxycarbonyl]methyl propanoate, origin: Firmenich SA, Geneva, Switzerland), 2-tert-butyl-1-cyclohexyl acetate (origin: International Flavors and Fragrances, USA), Limbanol[®] (1-(2,2,3,6-tetramethyl-cyclohexyl)-3-hexanol, origin: Firmenich SA, Geneva, Switzerland), trans-1-(2,2,6-
- 30 trimethyl-1-cyclohexyl)-3-hexanol (origin: Firmenich SA, Geneva, Switzerland), (E)-3-methyl-4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-one, terpenyl isobutyrate, Lorysia[®] (4-(1,1-dimethylethyl)-1-cyclohexyl acetate, origin: Firmenich SA, Geneva,

- Switzerland), 8-methoxy-1-p-menthene, Helvetolide[®] ((1S,1'R)-2-[1-(3',3'-dimethyl-1'-cyclohexyl) ethoxy]-2-methylpropyl propanoate, origin: Firmenich SA, Geneva, Switzerland), para tert-butylcyclohexanone, menthenethiol, 1-methyl-4-(4-methyl-3-pentenyl)-3-cyclohexene-1-carbaldehyde, allyl cyclohexylpropionate, cyclohexyl salicylate, 2-methoxy-4-methylphenyl methyl carbonate, ethyl 2-methoxy-4-methylphenyl carbonate, 4-ethyl-2-methoxyphenyl methyl carbonate;
- 5
- Group 4: Methyl cedryl ketone (origin: International Flavors and Fragrances, USA), Verdylate, vetyverol, vetyverone, 1-(octahydro-2,3,8,8-tetramethyl-2-naphtalenyl)-1-ethanone (origin: International Flavors and Fragrances, USA), (5RS,9RS,10SR)-2,6,9,10-tetramethyl-1-oxaspiro[4.5]deca-3,6-diene and the (5RS,9SR,10RS) isomer, 10 2,6,9,10-tetramethyl-1-oxaspiro[4.5]deca-3,6-diene, 1,2,3,5,6,7-hexahydro-1,1,2,3,3-pentamethyl-4-indenone (origin: International Flavors and Fragrances, USA), Hivernal[®] (a mixture of 3-(3,3-dimethyl-5-indanyl)propanal and 3-(1,1-dimethyl-5-indanyl)propanal, origin: Firmenich SA, Geneva, Switzerland), Rhubofix[®] (3',4-dimethyl-tricyclo[6.2.1.0(2,7)]undec-4-ene-9-spiro-2'-oxirane, 15 origin: Firmenich SA, Geneva, Switzerland), 9/10-ethyldiene-3-oxatricyclo[6.2.1.0(2,7)]undecane, Polywood[®] (perhydro-5,5,8A-trimethyl-2-naphthalenyl acetate, origin: Firmenich SA, Geneva, Switzerland), octalynol, Cetalex[®] (dodecahydro-3a,6,6,9a-tetramethyl-naphtho[2,1-b]furan, origin: Firmenich SA, Geneva, Switzerland), tricyclo[5.2.1.0(2,6)]dec-3-en-8-yl acetate and 20 tricyclo[5.2.1.0(2,6)]dec-4-en-8-yl acetate as well as tricyclo[5.2.1.0(2,6)]dec-3-en-8-yl propanoate and tricyclo[5.2.1.0(2,6)]dec-4-en-8-yl propanoate, (+)-(1S,2S,3S)-2,6,6-trimethyl-bicyclo[3.1.1]heptane-3-spiro-2'-cyclohexen-4'-one;
 - Group 5: camphor, borneol, isobornyl acetate, 8-isopropyl-6-methyl-25 bicyclo[2.2.2]oct-5-ene-2-carbaldehyde, camphopinene, cedramber (8-methoxy-2,6,6,8-tetramethyl-tricyclo[5.3.1.0(1,5)]undecane, origin: Firmenich SA, Geneva, Switzerland), cedrene, cedrenol, cedrol, Florex[®] (mixture of 9-ethylidene-3-oxatricyclo[6.2.1.0(2,7)]undecan-4-one and 10-ethylidene-3-oxatricyclo[6.2.1.0(2,7)]undecan-4-one, origin: Firmenich SA, Geneva, Switzerland), 30 3-methoxy-7,7-dimethyl-10-methylene-bicyclo[4.3.1]decane (origin: Firmenich SA, Geneva, Switzerland);

- Group 6: Cedroxyde[®] (trimethyl-13-oxabicyclo-[10.1.0]-trideca-4,8-diene, origin: Firmenich SA, Geneva, Switzerland), Ambrettolide LG ((E)-9-hexadecen-16-olide, origin: Firmenich SA, Geneva, Switzerland), Habanolide[®] (pentadecenolide, origin: Firmenich SA, Geneva, Switzerland), muscenone (3-methyl-(4/5)-cyclopentadecenone, origin: Firmenich SA, Geneva, Switzerland), muscone (origin: Firmenich SA, Geneva, Switzerland), Exaltolide[®] (pentadecanolide, origin: Firmenich SA, Geneva, Switzerland), Exaltone[®] (cyclopentadecanone, origin: Firmenich SA, Geneva, Switzerland), (1-ethoxyethoxy)cyclododecane (origin: Firmenich SA, Geneva, Switzerland), Astrotone, 4,8-cyclododecadien-1-one;
- 10 - Group 7: Lilial[®] (origin: Givaudan SA, Vernier, Switzerland), rosinol.

Preferably, the perfume comprises at least 30%, preferably at least 50%, more preferably at least 60% of ingredients selected from Groups 1 to 7, as defined above. More preferably said perfume comprises at least 30%, preferably at least 50% of ingredients from Groups 3 to 7, as defined above. Most preferably said perfume

15 comprises at least 30%, preferably at least 50% of ingredients from Groups 3, 4, 6 or 7, as defined above.

According to another preferred embodiment, the perfume comprises at least 30%, preferably at least 50%, more preferably at least 60% of ingredients having a logP above 3, preferably above 3.5 and even more preferably above 3.75.

20 Preferably, the perfume used in the invention contains less than 10% of its own weight of primary alcohols, less than 15% of its own weight of secondary alcohols and less than 20% of its own weight of tertiary alcohols. Advantageously, the perfume used in the invention does not contain any primary alcohols and contains less than 15% of secondary and tertiary alcohols.

25 According to an embodiment, the oil phase (or the oil-based core) comprises:

- 25-100wt% of a perfume oil comprising at least 15wt% of high impact perfume raw materials having a Log T<-4, and
- 0-75wt% of a density balancing material having a density greater than 1.07 g/cm³.

30 “High impact perfume raw materials” should be understood as perfume raw materials having a LogT<-4. The odor threshold concentration of a chemical compound is

determined in part by its shape, polarity, partial charges and molecular mass. For convenience, the threshold concentration is presented as the common logarithm of the threshold concentration, i.e., Log [Threshold] ("LogT").

A "*density balancing material*" should be understood as a material having a density
5 greater than 1.07 g/cm³ and having preferably low or no odor.

The odor threshold concentration of a perfuming compound is determined by using a gas chromatograph ("GC"). Specifically, the gas chromatograph is calibrated to determine the exact volume of the perfume oil ingredient injected by the syringe, the precise split ratio, and the hydrocarbon response using a hydrocarbon standard of known concentration
10 and chain-length distribution. The air flow rate is accurately measured and, assuming the duration of a human inhalation to last 12 seconds, the sampled volume is calculated. Since the precise concentration at the detector at any point in time is known, the mass per volume inhaled is known and hence the concentration of the perfuming compound. To determine the threshold concentration, solutions are delivered to the sniff port at the back-calculated concentration. A panelist sniffs the GC effluent and identifies the retention
15 time when odor is noticed. The average across all panelists determines the odor threshold concentration of the perfuming compound. The determination of odor threshold is described in more detail in C. Vuilleumier et al., Multidimensional Visualization of Physical and Perceptual Data Leading to a Creative Approach in Fragrance Development, Perfume & Flavorist, Vol. 33, September,, 2008, pages 54-61.
20

The nature of high impact perfume raw materials having a Log T<-4 and density balancing material having a density greater than 1.07 g/cm³ are described in WO2018115250, the content of which are included by reference.

According to an embodiment, the high impact perfume raw materials having a
25 Log T<-4 are selected from the list in Table A below.

Table A: high impact perfume raw materials having a Log T<-4

Perfume raw materials (Log T<-4)
(+)-1-METHOXY-3-HEXANETHIOL
4-(4-HYDROXY-1-PHENYL)-2-BUTANONE

(+)-2-(4-METHYL-3-CYCLOHEXEN-1-YL)-2-PROPANETHIOL
2-METHOXY-4-(1-PROPENYL)-1-PHENYL ACETATE
PYRAZOBUTYLE
3-PROPYLPHENOL
1-(3-METHYL-1-BENZOFURAN-2-YL)ETHANONE
2-(3-PHENYLPROPYL)PYRIDINE
1-(3,3-DIMETHYL-1-CYCLOHEXEN-1-YL)-4-PENTEN-1-ONE (A) + 1-(5,5-DIMETHYL-1-CYCLOHEXEN-1-YL)-4-PENTEN-1-ONE (B)
1-(5,5-DIMETHYL-1-CYCLOHEXEN-1-YL)-4-PENTEN-1-ONE (3RS,3ARS,6SR,7ASR)-PERHYDRO-3,6-DIMETHYL- BENZO[B]FURAN-2-ONE (A) + (3SR,3ARS,6SR,7ASR)- PERHYDRO-3,6-DIMETHYL-BENZO[B]FURAN-2-ONE (B)
(+)-1-(5-ETHYL-5-METHYL-1-CYCLOHEXEN-1-YL)-4-PENTEN- 1-ONE
(1'S,3'R)-1-METHYL-2-[(1',2',2'-TRIMETHYLBICYCLO[3.1.0]HEX- 3'-YL)METHYL]CYCLOPROPYL}METHANOL
(+)-3-MERCAPTOHEXYL ACETATE
(2E)-1-(2,6,6-TRIMETHYL-1,3-CYCLOHEXADIEN-1-YL)-2- BUTEN-1-ONE
7-METHYL-2H-1,5-BENZODIOXEPIN-3(4H)-ONE
(2E,6Z)-2,6-NONADIEN-1-OL
(4Z)-4-DODECENAL
(+)-4-HYDROXY-2,5-DIMETHYL-3(2H)-FURANONE
METHYL 2,4-DIHYDROXY-3,6-DIMETHYLBENZOATE
3-METHYLINDOLE
(+)-PERHYDRO-4ALPHA,8ABETA-DIMETHYL-4A- NAPHTHALENOL
PATCHOULOL

2-METHOXY-4-(1-PROPENYL)PHENOL
(+)-5,6-DIHYDRO-4-METHYL-2-PHENYL-2H-PYRAN (A) + TETRAHYDRO-4-METHYLENE-2-PHENYL-2H-PYRAN (B) 4-METHYLENE-2-PHENYL-TETRAHYDRO-2H-PYRAN (A) + (+)- 4-METHYL-2-PHENYL-3,6-DIHYDRO-2H-PYRAN (B)
4-HYDROXY-3-METHOXYBENZALDEHYDE
NONYLENIC ALDEHYDE
2-METHOXY-4-PROPYLPHENOL
(2Z)-3-METHYL-5-PHENYL-2-PENTENENITRILE (A) + (2E)-3- METHYL-5-PHENYL-2-PENTENENITRILE (B)
1-(SPIRO[4.5]DEC-6-EN-7-YL)-4-PENTEN-1-ONE (A) + 1- (SPIRO[4.5]DEC-7-EN-7-YL)-4-PENTEN-1-ONE (B)
2-METHOXYNAPHTHALENE
(-)-(3AR,5AS,9AS,9BR)-3A,6,6,9A- TETRAMETHYLDODECAHYDRONAPHTHO[2,1-B]FURAN
5-NONANOLIDE
(3AR,5AS,9AS,9BR)-3A,6,6,9A- TETRAMETHYLDODECAHYDRONAPHTHO[2,1-B]FURAN
7-ISOPROPYL-2H,4H-1,5-BENZODIOXEPIN-3-ONE
COUMARIN
4-METHYLPHENYL ISOBUTYRATE
(2E)-1-(2,6,6-TRIMETHYL-1,3-CYCLOHEXADIEN-1-YL)-2-BUTEN-1-ONE
BETA,2,2,3-TETRAMETHYL-DELTA-METHYLENE-3- CYCLOPENTENE-1-BUTANOL
DELTA DAMASCONE ((2E)-1-[(1RS,2SR)-2,6,6-TRIMETHYL-3- CYCLOHEXEN-1-YL]-2-BUTEN-1-ONE)
(+)-3,6-DIHYDRO-4,6-DIMETHYL-2-PHENYL-2H-PYRAN
ANISALDEHYDE

PARACRESOL
3-ETHOXY-4-HYDROXYBENZALDEHYDE
METHYL 2-AMINOBENZOATE
ETHYL METHYLPHENYLGLYCIDATE
OCTALACTONE G
ETHYL 3-PHENYL-2-PROPENOATE
(-)-(2E)-2-ETHYL-4-[(1R)-2,2,3-TRIMETHYL-3-CYCLOPENTEN-1-YL]-2-BUTEN-1-OL
PARACRESYL ACETATE
DODECALACTONE
TRICYCLONE
(+)-(3R,5Z)-3-METHYL-5-CYCLOPENTADECEN-1-ONE
UNDECALACTONE
(1R,4R)-8-MERCAPTO-3-P-MENTHANONE
(3S,3AS,6R,7AR)-3,6-DIMETHYLHEXAHYDRO-1-BENZOFURAN-2(3H)-ONE
BÉTA IONONE
(+)-6-PENTYLTETRAHYDRO-2H-PYRAN-2-ONE
(3E,5Z)-1,3,5-UNDECATRIENE
10-UNDECENAL (A) + (9E)-9-UNDECENAL (B) + (9Z)-9-UNDECENAL (C)
(Z)-4-DECENAL
(+)-ETHYL 2-METHYLPENTANOATE
1,2-DIALLYLDISULFANE
(2Z)-2-TRIDECENENITRILE (A) + (3Z)-3-TRIDECENENITRILE (B) + (3E)-3-TRIDECENENITRILE (C) + (2E)-2-TRIDECENENITRILE (D)
(+)-2-ETHYL-4,4-DIMETHYL-1,3-OXATHIANE

(+)-(3R,5Z)-3-METHYL-5-CYCLOPENTADECEN-1-ONE
3-(4-TERT-BUTYLPHENYL)PROPANAL
ALLYL (CYCLOHEXYLOXY)ACETATE
METHYLNAPHTHYLKETONE
(+)-(4E)-3-METHYL-4-CYCLOPENTADECEN-1-ONE (A) + (+)-(5E)-3-METHYL-5-CYCLOPENTADECEN-1-ONE (B) + (+)-(5Z)-3-METHYL-5-CYCLOPENTADECEN-1-ONE (C)
CYCLOPROPYLMETHYL (3Z)-3-HEXENOATE (A) + CYCLOPROPYLMETHYL (3E)-3-HEXENOATE (B)
(4E)-4-METHYL-5-(4-METHYLPHENYL)-4-PENTENAL
(+)-1-(5-PROPYL-1,3-BENZODIOXOL-2-YL)ETHANONE
4-METHYL-2-PENTYLPYRIDINE
(+)-(E)-3-METHYL-4-(2,6,6-TRIMETHYL-2-CYCLOHEXEN-1-YL)-3-BUTEN-2-ONE
(3ARS,5ASR,9ASR,9BRS)-3A,6,6,9A-TETRAMETHYLDODECAHYDRONAPHTHO[2,1-B]FURAN
(2S,5R)-5-METHYL-2-(2-PROPANYL)CYCLOHEXANONE OXIME
6-HEXYLTETRAHYDRO-2H-PYRAN-2-ONE
(+)-3-(3-ISOPROPYL-1-PHENYL)BUTANAL
METHYL 2-((1RS,2RS)-3-OXO-2-PENTYLCYCLOPENTYL)ACETATE (A) + METHYL 2-((1RS,2SR)-3-OXO-2-PENTYLCYCLOPENTYL)ACETATE (B)
1-(2,6,6-TRIMETHYL-1-CYCLOHEX-2-ENYL)PENT-1-EN-3-ONE
INDOL
7-PROPYL-2H,4H-1,5-BENZODIOXEPIN-3-ONE
ETHYL PRALINE
(4-METHYLPHENOXY)ACETALDEHYDE
ETHYL TRICYCLO[5.2.1.0.(2,6)]DECANE-2-CARBOXYLATE
(+)-(1'S,2S,E)-3,3-DIMETHYL-5-(2',2',3'-TRIMETHYL-3'-

CYCLOPENTEN-1'-YL)-4-PENTEN-2-OL
(2R,4E)-3,3-DIMETHYL-5-[(1R)-2,2,3-TRIMETHYL-3-CYCLOPENTEN-1-YL]-4-PENTEN-2-OL (A) + (2S,4E)-3,3-DIMETHYL-5-[(1R)-2,2,3-TRIMETHYL-3-CYCLOPENTEN-1-YL]-4-PENTEN-2-OL (B)
8-ISOPROPYL-6-METHYL-BICYCLO[2.2.2]OCT-5-ENE-2-CARBALDEHYDE
METHYLNONYLACETALDEHYDE
4-FORMYL-2-METHOXYPHENYL 2-METHYLPROPANOATE
(E)-4-DECENAL
(+)-2-ETHYL-4-(2,2,3-TRIMETHYL-3-CYCLOPENTEN-1-YL)-2-BUTEN-1-OL
(1R,5R)-4,7,7-TRIMETHYL-6-THIABICYCLO[3.2.1]OCT-3-ENE (A) + (1R,4R,5R)-4,7,7-TRIMETHYL-6-THIABICYCLO[3.2.1]OCTANE (B)
(-)-(3R)-3,7-DIMETHYL-1,6-OCTADIEN-3-OL
(E)-3-PHENYL-2-PROPENENITRILE
4-METHOXYBENZYL ACETATE
(E)-3-METHYL-5-(2,2,3-TRIMETHYL-3-CYCLOPENTEN-1-YL)-4-PENTEN-2-OL
ALLYL (3-METHYLBUTOXY)ACETATE (A) + (+)-ALLYL (2-METHYLBUTOXY)ACETATE
(+)-(2E)-1-(2,6,6-TRIMETHYL-2-CYCLOHEXEN-1-YL)-2-BUTEN-1-ONE
(1E)-1-(2,6,6-TRIMETHYL-1-CYCLOHEXEN-1-YL)-1-PENTEN-3-ONE

According to an embodiment, perfume raw materials having a Log T<-4 are chosen in the group consisting of aldehydes, ketones, alcohols, phenols, esters lactones, ethers, epoxydes, nitriles and mixtures thereof.

According to an embodiment, perfume raw materials having a Log T<-4 comprise at least one compound chosen in the group consisting of alcohols, phenols, esters lactones, ethers, epoxydes, nitriles and mixtures thereof, preferably in amount comprised between 20 and 70% by weight based on the total weight of the perfume raw materials having a

5 Log T<-4.

According to an embodiment, perfume raw materials having a Log T<-4 comprise between 20 and 70% by weight of aldehydes, ketones, and mixtures thereof based on the total weight of the perfume raw materials having a Log T<-4.

The remaining perfume raw materials contained in the oil-based core may have

10 therefore a Log T>-4.

Non limiting examples of perfume raw materials having a Log T>-4 are listed in table B below.

Table B: perfume raw materials having a Log T>-4

Perfume raw materials (Log T>-4)
ETHYL 2-METHYLBUTYRATE
(E)-3-PHENYL-2-PROPENYL ACETATE
(+)-8-SEC-BUTYLQUINOLINE (A) + (+)-6-SEC-BUTYLQUINOLINE
(+)-3-(1,3-BENZODIOXOL-5-YL)-2- METHYLPROPANAL
VERDYLE PROPIONATE
1-(OCTAHYDRO-2,3,8,8- TETRAMETHYL-2-NAPHTALENYL)- 1-ETHANONE
METHYL 2-((1RS,2RS)-3-OXO-2- PENTYLCYCLOPENTYL)ACETATE
(+)-(E)-4-METHYL-3-DECEN-5-OL
2,4-DIMETHYL-3-CYCLOHEXENE-1-

CARBALDEHYDE
1,3,3-TRIMETHYL-2- OXABICYCLO[2.2.2]OCTANE
TETRAHYDRO-4-METHYL-2-(2- METHYL-1-PROPENYL)-2H-PYRAN
ALDEHYDE C 12
1-OXA-12-CYCLOHEXADECEN-2- ONE (A) + 1-OXA-13- CYCLOHEXADECEN-2-ONE (B)
(+)-3-(4-ISOPROPYLPHENYL)-2- METHYLPROPANAL
ALDEHYDE C 11 LENIQUE
(+)-2,6-DIMETHYL-7-OCTEN-2-OL
(+)-2,6-DIMETHYL-7-OCTEN-2-OL
ALLYL 3- CYCLOHEXYLPROPANOATE
(Z)-3-HEXENYL ACETATE
(2RS,5SR)-5-METHYL-2-(2- PROPANYL)CYCLOHEXANONE (A) + (2RS,5RS)-5-METHYL-2-(2- PROPANYL)CYCLOHEXANONE (B)
ALLYL HEPTANOATE
(1RS,2RS)-2-(2-METHYL-2- PROPANYL)CYCLOHEXYL ACETATE (A) + (1RS,2SR)-2-(2-METHYL-2- PROPANYL)CYCLOHEXYL ACETATE (B)
1,1-DIMETHYL-2-PHENYLETHYL BUTYRATE
GERANYL ACETATE (A) + NERYL

ACETATE (B)
(+)-1-PHENYLETHYL ACETATE
1,1-DIMETHYL-2-PHENYLETHYL ACETATE
3-METHYL-2-BUTENYL ACETATE
ETHYL 3-OXOBUTANOATE (A) <=> (2Z)-ETHYL 3-HYDROXY-2- BUTENOATE (B)
8-P-MENTHANOL
8-P-MENTHANYL ACETATE (A) + 1- P-MENTHANYL ACETATE (B)
(+)-2-(4-METHYL-3-CYCLOHEXEN-1- YL)-2-PROPANYL ACETATE
(+)-2-METHYLBUTYL BUTANOATE
2-{(1S)-1-[(1R)-3,3- DIMETHYLCYCLOHEXYL]ETHOXY}- 2-OXOETHYL PROPIONATE
3,5,6-TRIMETHYL-3-CYCLOHEXENE- 1-CARBALDEHYDE (A) + 2,4,6- TRIMETHYL-3-CYCLOHEXENE-1- CARBALDEHYDE (B)
2-CYCLOHEXYLETHYL ACETATE
ALDEHYDE C 8
ETHYL BUTANOATE
(+)-(3E)-4-(2,6,6-TRIMETHYL-2- CYCLOHEXEN-1-YL)-3-BUTEN-2- ONE (A) + (3E)-4-(2,6,6-TRIMETHYL- 1-CYCLOHEXEN-1-YL)-3-BUTEN-2- ONE (B);

1-[(1R,6SR)-2,2,6- TRIMETHYLCYCLOHEXYL]-3- HEXANOL
1,3,3-TRIMETHYL-2- OXABICYCLO[2.2.2]OCTANE
1,3,3-TRIMETHYL-2- OXABICYCLO[2.2.2]OCTANE
ETHYL HEXANOATE
UNDECANAL
ALDEHYDE C 10
2-PHENYLETHYL ACETATE
(1S,2S,4S)-1,7,7- TRIMETHYLBICYCLO[2.2.1]HEPTAN- 2-OL (A) + (1S,2R,4S)-1,7,7- TRIMETHYLBICYCLO[2.2.1]HEPTAN- 2-OL (B)
(+)-3,7-DIMETHYL-3-OCTANOL
1-METHYL-4-(2- PROPANYLIDENE)CYCLOHEXENE
(+)-(R)-4-(2-METHOXYPROPAN-2- YL)-1-METHYLCYCLOHEX-1-ENE
VERDYL ACETATE
(3R)-1-[(1R,6S)-2,2,6- TRIMETHYLCYCLOHEXYL]-3- HEXANOL (A) + (3S)-1-[(1R,6S)-2,2,6- TRIMETHYLCYCLOHEXYL]-3- HEXANOL (B) + (3R)-1-[(1S,6S)-2,2,6- TRIMETHYLCYCLOHEXYL]-3- HEXANOL (C)

(+)-(1S,1'R)-2-[1-(3',3'-DIMETHYL-1'-CYCLOHEXYL)ETHOXY]-2-METHYLPROPYL PROPANOATE

According to an embodiment, the oil phase (or the oil-based core) comprises 2-75wt% of a density balancing material having a density greater than 1.07 g/cm^3 and 25-98wt% of a perfume oil comprising at least 15wt% of high impact perfume raw materials having a Log T<-4.

The density of a component is defined as the ratio between its mass and its volume (g/cm^3).

Several methods are available to determine the density of a component.

One may refer for example to the ISO 298:1998 method to measure d20 densities of essential oils.

According to an embodiment, the density balancing material is chosen in the group consisting of benzyl salicylate, benzyl benzoate, cyclohexyl salicylate, benzyl phenylacetate, phenylethyl phenoxyacetate, triacetin, methyl and ethyl salicylate, benzyl cinnamate, and mixtures thereof.

According to a particular embodiment, the density balancing material is chosen in the group consisting of benzyl salicylate, benzyl benzoate, cyclohexyl salicylate and mixtures thereof.

According to a particular embodiment, the hydrophobic material is free of any active ingredient (such as perfume). According to this particular embodiment, it comprises, preferably consists of hydrophobic solvents, preferably chosen in the group consisting of isopropyl myristate, tryglycerides (e.g. Neobee® MCT oil, vegetable oils), D-limonene, silicone oil, mineral oil, and mixtures thereof with optionally hydrophilic solvents preferably chosen in the group consisting of 1,4 butanediol, benzyl alcohol, triethyl citrate, triacetin, benzyl acetate, ethyl acetate, propylene glycol (1,2-propanediol), 1,3-Propanediol, dipropylene glycol, glycerol, glycol ethers and mixtures thereof.

The term "biocide" refers to a chemical substance capable of killing living organisms (e.g. microorganisms) or reducing or preventing their growth and/or accumulation. Biocides are commonly used in medicine, agriculture, forestry, and in

industry where they prevent the fouling of, for example, water, agricultural products including seed, and oil pipelines. A biocide can be a pesticide, including a fungicide, herbicide, insecticide, algicide, molluscicide, miticide and rodenticide; and/or an antimicrobial such as a germicide, antibiotic, antibacterial, antiviral, antifungal, antiprotozoal and/or antiparasite.

As used herein, a "pest control agent" indicates a substance that serves to repel or attract pests, to decrease, inhibit or promote their growth, development or their activity. Pests refer to any living organism, whether animal, plant or fungus, which is invasive or troublesome to plants or animals, pests include insects notably arthropods, mites, spiders, fungi, weeds, bacteria and other microorganisms.

By "flavour ingredient or composition" it is meant here a flavouring ingredient or a mixture of flavouring ingredients, solvent or adjuvants of current use for the preparation of a flavouring formulation, i.e. a particular mixture of ingredients which is intended to be added to an edible composition or chewable product to impart, improve or modify its organoleptic properties, in particular its flavour and/or taste. Taste modulator as also encompassed in said definition. Flavouring ingredients are well known to a skilled person in the art and their nature does not warrant a detailed description here, which in any case would not be exhaustive, the skilled flavourist being able to select them on the basis of his general knowledge and according to the intended use or application and the organoleptic effect it is desired to achieve. Many of these flavouring ingredients are listed in reference texts such as in the book by S. Arctander, Perfume and Flavor Chemicals, 1969, Montclair, N.J., USA, or its more recent versions, or in other works of similar nature such as Fenaroli's Handbook of Flavor Ingredients, 1975, CRC Press or Synthetic Food Adjuncts, 1947, by M.B. Jacobs, can Nostrand Co., Inc. Solvents and adjuvants or current use for the preparation of a flavouring formulation are also well known in the art.

In a particular embodiment, the flavour is selected from the group consisting of terpenic flavours including citrus and mint oil, and sulfury flavours.

According to any one of the invention's embodiments, the hydrophobic material represents between about 10% and 60% w/w, or even between 15% and 45% w/w, by weight, relative to the total weight of the dispersion as obtained after step b).

According to a particular embodiment, the oil phase essentially consists of the acyl chloride, a perfume or flavor oil and a stabilizer.

According to the invention, a stabilizer is added in the oil phase to form later on the emulsion. According to an embodiment, the stabilizer is a colloidal stabilizer.

5 The colloidal stabilizer can be a molecular emulsifier (standard emulsion) or solid particles (Pickering emulsion).

By “stabilizer”, it is meant compounds capable to stabilize oil/water interface as an emulsion. According to a particular embodiment, the stabilizer is a biopolymer.

By “biopolymers” it is meant biomacromolecules produced by living organisms.
10 Biopolymers are characterized by molecular weight distributions ranging from 1,000 (1 thousand) to 1,000,000,000 (1 billion) Daltons. These macromolecules may be carbohydrates (sugar based) or proteins (amino-acid based) or a combination of both (gums) and can be linear or branched.

According to an embodiment, the stabilizer is chosen in the group consisting of
15 gum Arabic, modified starch, polyvinyl alcohol, PVP (polyvinylpyrrolidone), CMC (carboxymethylcellulose), anionic polysaccharides, acrylamide copolymer, inorganic particles, protein such as soy protein, rice protein, whey protein, white egg albumin, sodium caseinate, gelatin, bovine serum albumin, hydrolyzed soy protein, hydrolyzed sericin, Pseudocollagen, Silk protein, sericin powder, and mixtures thereof.

20 According to a particular embodiment, the stabilizer is a biopolymer chosen in the group consisting of consisting of protein such as whey protein, sodium caseinate, bovine serum albumin, and mixtures thereof.

When added in the oil phase, the stabilizer can be pre-dispersed in an inert solvent such as benzyl benzoate or can be mixed to the active ingredient, preferably comprising a
25 perfume oil.

The stabilizer and acyl chloride can be premixed and can be heated at a temperature between for example 10 and 80°C before mixing with the hydrophobic material, preferably comprising a perfume oil.

According to any one of the above embodiments of the present invention, the
30 dispersion comprises between about 0.01% and 3.0% of at least stabilizer, percentage being expressed on a w/w basis relative to the total weight of the dispersion as obtained

after step b). In still another aspect of the invention, the dispersion comprises between about 0.05% and 2.0%, preferably between about 0.05% and 1.0% of at least a colloid stabilizer. In still another aspect of the invention, the dispersion comprises between about 0.1% and 1.6%, preferably between about 0.1% and 0.8% of at least a colloid stabilizer.

- 5 In another step of the process according to the invention, the oil phase of step a) is dispersed into an aqueous solution comprising optionally an amino compound A or a base to form an oil-in-water emulsion.

 The mean droplet size of the emulsion is preferably comprised between 1 and 1000 microns, more preferably between 1 and 500 microns, and even more preferably
10 between 5 and 50 microns.

 When added, the amino compound A may be an amino-acid, preferably chosen in the group consisting of L-Lysine, L-Arginine, L-Histidine, L-Tryptophane, L-Serin, L-Glutamine, L-Threonine and mixtures thereof, preferably L-Lysine, L-Arginine, L-Histidine, L-Tryptophane and mixtures thereof, more preferably L-Lysine, L-Arginine, L-
15 Histidine and mixtures thereof. The amino-acid has preferably two nucleophilic groups.

 When added, the amino compound A may be chosen in the group consisting of L-Lysine, L-Lysine ethyl ester, guanidine carbonate, chitosan, 3-aminopropyltriethoxysilane, and mixtures thereof. According to a particular embodiment, the amino compound A is L-Lysine.

- 20 According to a particular embodiment, when added, the base is not an amino-compound and is chosen in the group consisting of sodium carbonate, sodium bicarbonate, sodium hydroxide and mixtures thereof.

 Base is preferably added in an amount comprised between 0.1% and 10% by weight based on the dispersion, more preferably between 0.5% and 5% by weight.

- 25 In another step of the process according to an embodiment, an amino compound B is added to the oil-in-water emulsion obtained in step b) to form a polyamide shell.

 As non-limiting examples, the amino-compound B is chosen in the group consisting of a xylylene diamine, 1,2-diaminocyclohexane, 1,4-diaminocyclohexane, L-lysine, L-Lysine ethyl ester, Jeffamine® (Polyetheramines), ethylene diamine, diethylene triamine,
30 spermine, spermidine, polyamidoamine (PAMAM), guanidine carbonate, chitosan, tris-(2-aminoethyl)amine, 3-aminopropyltriethoxysilane, L-arginine, an amine having a

disulfide bond such as cystamine, cystamine hydrochloride, cystine, cystine hydrochloride, cystine dialkyl ester, cystine dialkyl ester hydrochloride and mixtures thereof.

According to an embodiment, the amino-compound B is an amine having a disulfide bond and is chosen in the group consisting of cystamine, cystamine hydrochloride, cystine, cystine hydrochloride, cystine dialkyl ester, cystine dialkyl ester hydrochloride and mixtures thereof.

According to another embodiment, the amino-compound B is chosen in the group consisting of xylylene diamine, 1,2-diaminocyclohexane, 1,4-diaminocyclohexane, L-lysine, L-Lysine ethyl ester, Jeffamine® (Polyetheramines), ethylene diamine, diethylene triamine, spermine, spermidine, polyamidoamine (PAMAM), guanidine carbonate, chitosan, tris-(2-aminoethyl)amine, 3-aminopropyltriethoxysilane, L-arginine and mixtures thereof.

According to a particular embodiment, when amino compound A is present, the amino compound A and the amino compound B are the same.

According to another particular embodiment, when amino compound A is present, the amino compound A and the amino compound B are different.

According to an embodiment, when amino compound A is present, the weight ratio between the amino compound A and the amino compound B is comprised between 0.5 and 25, preferably between 1.3 and 10, more preferably between 1.3 and 7.

The amount of the amino compound B used is typically adjusted so that the molar ratio between the functional group NH_2 of the amino compound B and COCl of the acyl chloride is comprised between 0.01 and 7.5, preferably from 0.1 to 3.0.

When added, the amount of the amino compound A used is typically adjusted so that the molar ratio between the functional groups NH_2 of the amino compound A and the functional groups COCl of the acyl chloride is comprised between 0.2 and 3, preferably from 0.5 to 2.

According to an embodiment, a base is added to adjust the pH at the end of step c) or d). One may cite as non-limiting examples guanidine carbonate, sodium bicarbonate or triethanolamine.

Base is preferably added in an amount comprised between 0.1% and 10% based on the dispersion, more preferably between 0.5% and 5%.

This is followed by a curing step c) or d) which allows ending up with microcapsules in the form of a slurry. According to a preferred embodiment, to enhance the kinetics, said step is performed at a temperature comprised between 5 and 90°C, possibly under pressure, for 1 to 8 hours. More preferably it is performed at between 10 and 80°C for between 30 minutes and 5 hours.

According to a particular embodiment, no polyol is added at any step of the process.

10

Optional outer coating: According to a particular embodiment of the invention, during or at the end of step c) or d) one may also add to the invention's slurry a polymer selected from the group consisting of a non-ionic polysaccharide, a cationic polymer and mixtures thereof to form an outer coating to the microcapsule.

Non-ionic polysaccharide polymers are well known to a person skilled in the art and are described for instance in WO2012/007438 page 29, lines 1 to 25 and in WO2013/026657 page 2, lines 12 to 19 and page 4, lines 3 to 12. Preferred non-ionic polysaccharides are selected from the group consisting of locust bean gum, xyloglucan, guar gum, hydroxypropyl guar, hydroxypropyl cellulose and hydroxypropyl methyl cellulose.

20

Cationic polymers are well known to a person skilled in the art.. Preferred cationic polymers have cationic charge densities of at least 0.5 meq/g, more preferably at least about 1.5 meq/g, but also preferably less than about 7 meq/g, more preferably less than about 6.2 meq/g. The cationic charge density of the cationic polymers may be determined by the Kjeldahl method as described in the US Pharmacopoeia under chemical tests for Nitrogen determination. The preferred cationic polymers are chosen from those that contain units comprising primary, secondary, tertiary and/or quaternary amine groups that can either form part of the main polymer chain or can be borne by a side substituent directly connected thereto. The weight average (Mw) molecular weight of the cationic polymer is preferably between 10,000 and 3.5M Dalton, more preferably between 50,000 and 1.5M Dalton. According to a particular embodiment, one will use cationic polymers

30

based on acrylamide, methacrylamide, N-vinylpyrrolidone, quaternized N,N-dimethylaminomethacrylate, diallyldimethylammonium chloride, quaternized vinylimidazole (3-methyl-1-vinyl-1H-imidazol-3-ium chloride), vinylpyrrolidone, acrylamidopropyltrimonium chloride, cassia hydroxypropyltrimonium chloride, guar
5 hydroxypropyltrimonium chloride or polygalactomannan 2-hydroxypropyltrimethylammonium chloride ether, starch hydroxypropyltrimonium chloride and cellulose hydroxypropyltrimonium chloride. Preferably copolymers shall be selected from the group consisting of polyquaternium-5, polyquaternium-6, polyquaternium-7, polyquaternium-10, polyquaternium-11, polyquaternium-16,
10 polyquaternium-22, polyquaternium-28, polyquaternium-43, polyquaternium-44, polyquaternium-46, cassia hydroxypropyltrimonium chloride, guar hydroxypropyltrimonium chloride or polygalactomannan 2-hydroxypropyltrimethylammonium chloride ether, starch hydroxypropyltrimonium chloride and cellulose hydroxypropyltrimonium chloride. As specific examples of
15 commercially available products, one may cite Salcare[®] SC60 (cationic copolymer of acrylamidopropyltrimonium chloride and acrylamide, origin: BASF) or Luviquat[®], such as the PQ 11N, FC 550 or Style (polyquaternium-11 to 68 or quaternized copolymers of vinylpyrrolidone origin: BASF), or also the Jaguar[®] (C13S or C17, origin Rhodia).

According to any one of the above embodiments of the invention, there is added
20 an amount of polymer described above comprised between about 0% and 5% w/w, or even between about 0.1% and 2% w/w, percentage being expressed on a w/w basis relative to the total weight of the slurry as obtained after step c) or d). It is clearly understood by a person skilled in the art that only part of said added polymers will be incorporated into/deposited on the microcapsule shell.

25 Another object of the invention is a process for preparing a microcapsule powder comprising the steps as defined above and an additional step d) or e) consisting of submitting the slurry obtained in step c) or d) to a drying, like spray-drying, to provide the microcapsules as such, i.e. in a powdery form. It is understood that any standard method known by a person skilled in the art to perform such drying is also applicable. In
30 particular the slurry may be spray-dried preferably in the presence of a polymeric carrier material such as polyvinyl acetate, polyvinyl alcohol, dextrans, natural or modified starch,

vegetable gums, pectins, xanthans, alginates, carragenans or cellulose derivatives to provide microcapsules in a powder form.

According to a particular embodiment, the carrier material contains free perfume oil which can be the same or different from the perfume from the core of the
5 microcapsules.

Another object of the invention is a polyamide microcapsule slurry obtainable by the process as described above. Another object of the invention is a polyamide microcapsule powder obtained by drying the microcapsule slurry defined above.

10 **Perfuming composition/consumer products**

The microcapsules of the invention can be used in combination with active ingredients. An object of the invention is therefore a composition comprising:

- (i) microcapsules as defined above;
- (ii) an active ingredient, preferably chosen in the group consisting of a cosmetic
15 ingredient, skin caring ingredient, perfume ingredient, flavor ingredient, malodour counteracting ingredient, bactericide ingredient, fungicide ingredient, pharmaceutical or agrochemical ingredient, a sanitizing ingredient, an insect repellent or attractant, and mixtures thereof.

The microcapsules of the invention can be used for the preparation of perfuming
20 or flavouring compositions which are also an object of the invention.

The capsules of the invention show very good performance in terms of stability in challenging medium.

- Another object of the present invention is a perfuming composition comprising:
- 25 (i) microcapsules as defined above, wherein the oil comprises a perfume;
 - (ii) at least one ingredient selected from the group consisting of a perfumery carrier, a perfumery co-ingredient and mixtures thereof;
 - (iii) optionally at least one perfumery adjuvant.

As liquid perfumery carrier one may cite, as non-limiting examples, an
30 emulsifying system, i.e. a solvent and a surfactant system, or a solvent commonly used in perfumery. A detailed description of the nature and type of solvents commonly used in

perfumery cannot be exhaustive. However, one can cite as non-limiting examples solvents such as dipropyleneglycol, diethyl phthalate, isopropyl myristate, benzyl benzoate, 2-(2-ethoxyethoxy)-1-ethanol or ethyl citrate, which are the most commonly used. For the compositions which comprise both a perfumery carrier and a perfumery co-ingredient, other suitable perfumery carriers than those previously specified, can be also ethanol, water/ethanol mixtures, limonene or other terpenes, isoparaffins such as those known under the trademark Isopar[®] (origin: Exxon Chemical) or glycol ethers and glycol ether esters such as those known under the trademark Dowanol[®] (origin: Dow Chemical Company). By “perfumery co-ingredient” it is meant here a compound, which is used in a perfuming preparation or a composition to impart a hedonic effect and which is not a microcapsule as defined above. In other words such a co-ingredient, to be considered as being a perfuming one, must be recognized by a person skilled in the art as being able to at least impart or modify in a positive or pleasant way the odor of a composition, and not just as having an odor.

The nature and type of the perfuming co-ingredients present in the perfuming composition do not warrant a more detailed description here, which in any case would not be exhaustive, the skilled person being able to select them on the basis of his general knowledge and according to the intended use or application and the desired organoleptic effect. In general terms, these perfuming co-ingredients belong to chemical classes as varied as alcohols, lactones, aldehydes, ketones, esters, ethers, acetates, nitriles, terpenoids, nitrogenous or sulphurous heterocyclic compounds and essential oils, and said perfuming co-ingredients can be of natural or synthetic origin. Many of these co-ingredients are in any case listed in reference texts such as the book by S. Arctander, Perfume and Flavor Chemicals, 1969, Montclair, New Jersey, USA, or its more recent versions, or in other works of a similar nature, as well as in the abundant patent literature in the field of perfumery. It is also understood that said co-ingredients may also be compounds known to release in a controlled manner various types of perfuming compounds.

By “perfumery adjuvant” we mean here an ingredient capable of imparting additional added benefit such as a color, a particular light resistance, chemical stability, etc. A detailed description of the nature and type of adjuvant commonly used in

perfuming bases cannot be exhaustive, but it has to be mentioned that said ingredients are well known to a person skilled in the art.

Preferably, the perfuming composition according to the invention comprises between 0.01 and 30 % by weight of microcapsules as defined above.

5 The invention's microcapsules can advantageously be used in many application fields and used in consumer products. Microcapsules can be used in liquid form applicable to liquid consumer products as well as in powder form, applicable to powder consumer products.

 According to a particular embodiment, the consumer product as defined above is
10 liquid and comprises:

- a) from 2 to 65% by weight, relative to the total weight of the consumer product, of at least one surfactant;
- b) water or a water-miscible hydrophilic organic solvent; and
- c) microcapsule slurry as defined above,
- 15 d) optionally non-encapsulated perfume.

 According to a particular embodiment, the consumer product as defined above is in a powder form and comprises:

- a) from 2 to 65% by weight, relative to the total weight of the consumer product, of at least one surfactant;
- 20 b) microcapsule powder as defined above.
- c) optionally perfume powder that is different from the microcapsules defined above.

 In the case of microcapsules including a perfume oil-based core, the products of the invention, can in particular be of used in perfumed consumer products such as product belonging to fine fragrance or "functional" perfumery. Functional perfumery includes in
25 particular personal-care products including hair-care, body cleansing, skin care, hygiene-care as well as home-care products including laundry care and air care. Consequently, another object of the present invention consists of a perfumed consumer product comprising as a perfuming ingredient, the microcapsules defined above or a perfuming composition as defined above. The perfume element of said consumer product can be a

combination of perfume microcapsules as defined above and free or non-encapsulated perfume, as well as other types of perfume microcapsule than those here-disclosed.

In particular a liquid consumer product comprising:

- a) from 2 to 65% by weight, relative to the total weight of the consumer product, of at least one surfactant;
- b) water or a water-miscible hydrophilic organic solvent; and
- c) a perfuming composition as defined above is another object of the invention.

Also a powder consumer product comprising

- (a) from 2 to 65% by weight, relative to the total weight of the consumer product, of at least one surfactant; and
- (b) a perfuming composition as defined above is part of the invention.

The invention's microcapsules can therefore be added as such or as part of an invention's perfuming composition in a perfumed consumer product.

For the sake of clarity, it has to be mentioned that, by "perfumed consumer product" it is meant a consumer product which is expected to deliver among different benefits a perfuming effect to the surface to which it is applied (e.g. skin, hair, textile, paper, or home surface) or in the air (air-freshener, deodorizer etc). In other words, a perfumed consumer product according to the invention is a manufactured product which comprises a functional formulation also referred to as "base", together with benefit agents, among which an effective amount of microcapsules according to the invention.

The nature and type of the other constituents of the perfumed consumer product do not warrant a more detailed description here, which in any case would not be exhaustive, the skilled person being able to select them on the basis of his general knowledge and according to the nature and the desired effect of said product. Base formulations of consumer products in which the microcapsules of the invention can be incorporated can be found in the abundant literature relative to such products. These formulations do not warrant a detailed description here which would in any case not be exhaustive. The person skilled in the art of formulating such consumer products is perfectly able to select the suitable components on the basis of his general knowledge and of the available literature.

Non-limiting examples of suitable perfumed consumer product can be a perfume, such as a fine perfume, a cologne, an after-shave lotion, a body-splash; a fabric care product, such as a liquid or solid detergent, tablets and pods, a fabric softener, a dryer sheet, a fabric refresher, an ironing water, or a bleach; a personal-care product, such as a hair-care product (e.g. a shampoo, hair conditioner, a colouring preparation or a hair spray), a cosmetic preparation (e.g. a vanishing cream, body lotion or a deodorant or antiperspirant), or a skin-care product (e.g. a perfumed soap, shower or bath mousse, body wash, oil or gel, bath salts, or a hygiene product); an air care product, such as an air freshener or a “ready to use” powdered air freshener; or a home care product, such all-purpose cleaners, liquid or power or tablet dishwashing products, toilet cleaners or products for cleaning various surfaces, for example sprays & wipes intended for the treatment / refreshment of textiles or hard surfaces (floors, tiles, stone-floors etc.); a hygiene product such as sanitary napkins, diapers, toilet paper.

Another object of the invention is a consumer product comprising:

- 15- a personal care active base, and
 - microcapsules slurry or microcapsule powder as defined above or the perfuming composition as defined above,
- wherein the consumer product is in the form of a personal care composition.

Personal care active base in which the microcapsules of the invention can be incorporated can be found in the abundant literature relative to such products. These formulations do not warrant a detailed description here which would in any case not be exhaustive. The person skilled in the art of formulating such consumer products is perfectly able to select the suitable components on the basis of his general knowledge and of the available literature.

25 The personal care composition is preferably chosen in the group consisting of a hair-care product (e.g. a shampoo, hair conditioner, a colouring preparation or a hair spray), a cosmetic preparation (e.g. a vanishing cream, body lotion or a deodorant or antiperspirant), or a skin-care product (e.g. a perfumed soap, shower or bath mousse, body wash, oil or gel, bath salts, or a hygiene product).

30 Another object of the invention is a consumer product comprising:

- a home care or a fabric care active base, and

- microcapsules slurry or microcapsule powder as defined above or the perfuming composition as defined above,

wherein the consumer product is in the form of a home care or a fabric care composition.

Home care or fabric care active base in which the microcapsules of the invention can be incorporated can be found in the abundant literature relative to such products. These formulations do not warrant a detailed description here which would in any case not be exhaustive. The person skilled in the art of formulating such consumer products is perfectly able to select the suitable components on the basis of his general knowledge and of the available literature.

Preferably, the consumer product comprises from 0.1 to 15 wt%, more preferably between 0.2 and 5 wt% of the microcapsules of the present invention, these percentages being defined by weight relative to the total weight of the consumer product. Of course the above concentrations may be adapted according to the benefit effect desired in each product.

According to a particular embodiment, the consumer product in which the microcapsules are incorporated has a pH preferably lower than 4.5.

Fabric softener

An object of the invention is a consumer product in the form of a fabric softener composition comprising:

- a fabric softener active base; preferably chosen in the group consisting of dialkyl quaternary ammonium salts, dialkyl ester quaternary ammonium salts (esterquats), Hamburg esterquat (HEQ), TEAQ (triethanolamine quat), cationic guar, silicones and mixtures thereof, preferably in an amount comprised between 85 and 99.95% by weight based on the total weight of the composition,
- a microcapsule slurry as defined above, preferably in an amount comprised between 0.05 to 15 wt%, more preferably between 0.1 and 5 wt% by weight based on the total weight of the composition.

Liquid detergent

An object of the invention is a consumer product in the form of a liquid detergent composition comprising:

- a liquid detergent active base; preferably chosen in the group consisting of anionic surfactant such as alkylbenzenesulfonate (ABS), secondary alkyl sulfonate (SAS),
5 primary alcohol sulfate (PAS), lauryl ether sulfate (LES), methyl ester sulfonate (MES) and nonionic surfactant such as alkyl amines, alkanolamide, fatty alcohol poly(ethylene glycol) ether, fatty alcohol ethoxylate (FAE), ethylene oxide (EO) and propylene oxide (PO) copolymers, amine oxydes, alkyl polyglucosides, alkyl polyglucosamides, preferably in an amount comprised between 85 and 99.95% by
10 weight based on the total weight of the composition,
- a microcapsule slurry as defined above, preferably in an amount comprised between 0.05 to 15 wt%, more preferably between 0.1 and 5 wt% by weight based on the total weight of the composition.

15 **Solid detergent**

An object of the invention is a consumer product in the form of a solid detergent composition comprising:

- a solid detergent active base; preferably chosen in the group consisting of anionic surfactant such as alkylbenzenesulfonate (ABS), secondary alkyl sulfonate (SAS),
20 primary alcohol sulfate (PAS), lauryl ether sulfate (LES), methyl ester sulfonate (MES) and nonionic surfactant such as alkyl amines, alkanolamide, fatty alcohol poly(ethylene glycol) ether, fatty alcohol ethoxylate (FAE), ethylene oxide (EO) and propylene oxide (PO) copolymers, amine oxydes, alkyl polyglucosides, alkyl polyglucosamides, preferably in an amount comprised between 85 and 99.95% by
25 weight based on the total weight of the composition,
- a microcapsule slurry or microcapsule powder as defined above, preferably in an amount comprised between 0.05 to 15 wt%, more preferably between 0.1 and 5 wt% by weight based on the total weight of the composition.

Solid scent booster

An object of the invention is a consumer product in the form of a solid scent booster comprising:

- a solid carrier, preferably chosen in the group consisting of urea, sodium chloride, sodium sulphate, sodium acetate, zeolite, sodium carbonate, sodium bicarbonate, clay, talc, calcium carbonate, magnesium sulfate, gypsum, calcium sulfate, magnesium oxide, zinc oxide, titanium dioxide, calcium chloride, potassium chloride, magnesium chloride, zinc chloride, saccharides such as sucrose, mono-, di-, and polysaccharides and derivatives such as starch, cellulose, methyl cellulose, ethyl cellulose, propyl cellulose, polyols/sugar alcohols such as sorbitol, maltitol, xylitol, erythritol, and isomalt, PEG, PVP, citric acid or any water soluble solid acid, fatty alcohols or fatty acids and mixtures thereof.
- microcapsules powder as defined above, in a powdered form, preferably in an amount comprised between 0.05 to 15 wt%, more preferably between 0.1 and 5 wt% by weight based on the total weight of the composition.

Liquid scent booster

An object of the invention is a consumer product in the form of a liquid scent booster comprising:

- an aqueous phase,
- a surfactant system essentially consisting of one or more than one non-ionic surfactant, wherein the surfactant system has a mean HLB between 10 and 14, preferably chosen in the group consisting of ethoxylated aliphatic alcohols, POE/PPG (polyoxyethylene and polyoxypropylene) ethers, mono and polyglyceryl esters, sucrose ester compounds, polyoxyethylene hydroxylesters, alkyl polyglucosides, amine oxides and combinations thereof;
- a linker chosen in the group consisting of alcohols, salts and esters of carboxylic acids, salts and esters of hydroxyl carboxylic acids, fatty acids,

fatty acid salts, glycerol fatty acids, surfactant having an HLB less than 10 and mixtures thereof, and

- microcapsules slurry as defined above, in the form of a slurry, preferably in an amount comprised between 0.05 to 15 wt%, more preferably between 0.1 and 5 wt% by weight based on the total weight of the composition.

Shampoo/shower gel

An object of the invention is a consumer product in the form of a shampoo or a shower gel composition comprising:

- a shampoo or a shower gel active base; preferably chosen in the group consisting of sodium alkylether sulfate, ammonium alkylether sulfates, alkylamphoacetate, cocamidopropyl betaine, cocamide MEA, alkylglucosides and aminoacid based surfactants and mixtures thereof, preferably in an amount comprised between 85 and 99.95% by weight based on the total weight of the composition,
- a microcapsule slurry as defined above, preferably in an amount comprised between 0.05 to 15 wt%, more preferably between 0.1 and 5 wt% by weight based on the total weight of the composition.

Rinse-Off Conditioner

An object of the invention is a consumer product in the form of a rinse-off conditioner composition comprising:

- a rinse-off conditioner active base; preferably chosen in the group consisting of cetyltrimonium chloride, stearyl trimonium chloride, benzalkonium chloride, behentrimonium chloride and mixture thereof, preferably in an amount comprised between 85 and 99.95% by weight based on the total weight of the composition,
- a microcapsule slurry as defined above, preferably in an amount comprised between 0.05 to 15 wt%, more preferably between 0.1 and 5 wt% by weight based on the total weight of the composition.

Hair coloration

An object of the invention is a consumer product in the form of an oxidative hair coloring composition comprising:

- an oxidizing phase comprising an oxidizing agent and an alkaline phase comprising an alkakine agent, a dye precursor and a coupling compound; wherein
5 said dye precursor and said coupling compound form an oxidative hair dye in the presence of the oxidizing agent, preferably in an amount comprised between 85 and 99.95% by weight based on the total weight of the composition,
- a microcapsule slurry as defined above, preferably in an amount comprised
10 between 0.05 to 15 wt%, more preferably between 0.1 and 5 wt% by weight based on the total weight of the composition.

By “oxidative hair coloring composition”, it is meant a composition comprising two groups of colorless dye molecules: the dye precursor and the coupling agent. Upon reaction with each other through an oxidation process, they form a wide range of colored
15 molecules (dyes) that are then trapped into the hair due their size. In other words, the dye precursor and the coupling compound form an oxidative hair dye in the presence of the oxidizing agent.

“Dye precursor” and “oxidative dye precursor” are used indifferently in the present invention.

20 Dye precursors can be aromatic compounds derived from benzene substituted by at least two electron donor groups such as NH_2 and OH in *para* or *ortho* positions to confer the property of easy oxidation.

According to an embodiment, dye precursors are chosen in the group consisting of p-phenylene diamine, 2,5-diamino toluene, N,N-bis(2-hydroxymethyl)-p-phenylene
25 diamine, 4-aminophenol, 1,4-diamino-benzene, and mixtures thereof.

The primary dye precursors is used in combination with coupling agents. Coupling agents are preferably aromatic compounds derived from benzene and substituted by groups such as NH_2 and OH in the *meta* position and do not produce color
30 singly, but which modify the color, shade or intensity of the colors developed by the dye precursor.

According to an embodiment, the coupling agent is chosen in the group consisting of resorcinol, 2-methyl resorcinol, 4-chlororesorcinol, 2,5-diamino-toluene, 1,3-diamino-benzene, 2,4-diaminophenoxyethanol HCl, 2-amino-hydroxyethylaminoanisole sulfate, 4-amino-2-hydroxytoluene, and mixtures thereof.

- 5 The oxidative dye precursor is preferably used in an amount comprised between 0.001% and 5%, preferably between 0.1% and 4% by weight based on the total weight of the composition.

 The use of oxidative dye precursors and coupling agents in hair coloring formulation have been widely disclosed in the prior art and is well-known from the person skilled in the art. One may cite for example EP0946133A1, the content of which is
10 incorporated by reference.

 The alkaline phase comprises an alkaline agent, preferably chosen in the group consisting of ammonia hydroxide, ammonia carbonate, ethanolamine, potassium hydroxide, sodium borate, sodium carbonate, triethanolamine and mixtures thereof.

- 15 The alkaline agent is preferably used in an amount comprised between 1% and 10%, preferably between 3% and 9% by weight based on the total weight of the composition.

 According to the invention, the coupling agent and the dye precursor in an alkaline medium form an oxidative hair dye in the presence of the oxidizing agent.

- 20 The oxidizing agent will supply the necessary oxygen gas to develop color molecules and create a change in hair color.

 The oxidizing agent should be safe and effective for use in the compositions herein.

- Preferably, the oxidizing agents suitable for use herein will be soluble in the compositions according to the present invention when in liquid form and/or in the form
25 intended to be used.

 Preferably, oxidizing agents suitable for use herein will be water-soluble. Suitable oxidizing agents for use herein are selected from inorganic peroxygen oxidizing agents, preformed organic peroxyacid oxidizing agents and organic peroxide oxidizing agents or mixtures thereof.

The oxidizing agent is preferably used in an amount comprised between 5 and 30%, preferably between 5 and 25% by weight based on the total weight of the composition.

Components commonly used in cosmetic compositions may be added into the hair coloring composition as defined in the present invention. One may cite for example, 5 surfactants, cationic polymers, oily substances, silicone derivatives, free perfume, preservatives, ultraviolet absorbents, antioxidants, germicides, propellants, thickeners.

According to a particular embodiment, the hair coloring composition comprises one or more quaternary ammonium compounds, preferably chosen in the group consisting 10 of cetyltrimonium chloride, stearyl trimonium chloride, benzalkonium chloride, behentrimonium chloride and mixture thereof to confer hair conditioner benefits.

Perfuming composition

According to a particular embodiment, the consumer product is in the form of a 15 perfuming composition comprising:

- 0.1 to 30%, preferably 0.1 to 20% of microcapsules as defined previously,
- 0 to 40%, preferably 3-40% of perfume, and
- 20-90, preferably 40-90% of ethanol, by weight based on the total weight of the perfuming composition.

20 The invention will now be further described by way of examples. It will be appreciated that the invention as claimed is not intended to be limited in any way by these examples.

EXAMPLES

25

General protocol

1,3,5-benzene tricarbonyl chloride (BTC, 1.77 g, Table 1) was dissolved in benzyl benzoate (5 g). An emulsifier (0.95 g) was dispersed in benzyl benzoate (5 g) and was 30 optionally maintained under stirring at 60°C for one hour. Both solutions were mixed together, stirred at room temperature for 10 minutes, and then added to a perfume oil (25

g, Table 2a or Table 2b) at room temperature to form an oil phase. Oil phase was mixed with water (94.05 g), the latter comprising optionally an amino compound A (first amino-compound) or a base. Reaction mixture was stirred with an Ultra Turrax at 24,000 rpm for 30 s or 1 minute to afford an emulsion. An amino compound B (second amino-compound) (Table 3) was dissolved in water (5 g) and this solution was added dropwise to the emulsion. A solution of guanidine carbonate (30 wt% in water, 5 to 10 g) was optionally added to control pH value at about 8. The reaction mixture was stirred at 30°C for 4 h to afford a white dispersion.

10

Example 1**Preparation of microcapsules according to the process of the invention**

Microcapsules were prepared according to the general protocol described previously.

15 Materials

Table 1: List of ingredients used in the examples

Ingredients	Function
BTC ¹⁾	Acyl chloride
Isophthaloyl chloride ²⁾	Acyl chloride
mXDA ³⁾	Amino compound
L-lysine ⁴⁾	Amino compound
EDA ⁵⁾	Amino compound
Cystamine Dihydrochloride ⁶⁾	Amino compound
Diethylene triamine ⁷⁾	Amino compound
Spermine dihydrate ⁸⁾	Amino compound
Guanidine carbonate ⁹⁾	Amino compound
Cystine dtertbutyl ester ¹⁰⁾	Amino compound
Zein ¹¹⁾	Colloidal stabilizer
Gum arabic ¹²⁾	Colloidal stabilizer
Bovin Serum Albumin ¹³⁾	Colloidal stabilizer

White Egg ¹⁴⁾	Colloidal stabilizer
Bio Pro 2E063 ¹⁵⁾	Colloidal stabilizer
Bio Pur β -lactoglobulin ¹⁶⁾	Colloidal stabilizer
Sodium Caseinate ¹⁷⁾	Colloidal Stabilizer
Lactose ¹⁸⁾	Colloidal Stabilizer
Sodium bicarbonate ¹⁹⁾	Base
Sodium Carbonate ²⁰⁾	Base
Sodium Hydroxide ²¹⁾	Base

- 1) 1,3,5-benzene tricarbonyl chloride; origin : Aldrich, Switzerland
- 2) Isophthaloyl chloride²⁾ ; origin : Aldrich, Switzerland
- 3) m-xylylene diamine; origin : Aldrich, Switzerland
- 4) L-Lysine; origin : Aldrich, Switzerland
- 5) Ethylene diamine; origin : Aldrich, Switzerland
- 6) Cystamine Dihydrochloride; origin : Aldrich, Switzerland
- 7) Diethylene triamine; origin : Aldrich, Switzerland
- 8) Spermine dihydrate; origin : Aldrich, Switzerland
- 9) Guanidine carbonate; origin : Aldrich, Switzerland
- 10) Cystine ditertbutyl ester; origin: Aldrich, Switzerland
- 11) Zein; origin : Aldrich, Switzerland
- 12) Acacia gum Superstab AA, origin: Nexira, France
- 13) Bovin Serum Albumin; origin : Aldrich, Switzerland
- 14) White Egg
- 15) Bio Pro 2E063; origin : Agropur Inc. USA
- 16) Bio Pur β -lactoglobulin; origin : Danisco
- 17) Sodium Caseinate; origin : Aldrich, Switzerland
- 18) Lactose; origin : Aldrich, Switzerland
- 19) Sodium bicarbonate; origin : Aldrich, Switzerland
- 20) Sodium carbonate; origin : Aldrich, Switzerland
- 21) Sodium Hydroxide; origin : Aldrich, Switzerland

5 Perfume oil composition:**Table 2a:** Perfume oil A composition

Raw Materials	% in oil
2,4-Dimethyl-3-cyclohexene-1-carbaldehyde	3.30%
Allyl Heptanoate	5.50%
Allyl amyl glycolate	10.99%
Delta Damascone	1.65%
Verdyl acetate	20.30%
Hedione® ¹⁾	4.95%
Iso E Super ® ²⁾	16.49%
Ald. Hexylcinnamique	9.89%
Ethyl-2-methylvalerate	3.3%
Lilial	21.98%
(3Z)-3-hexen-1-yl butyrate	1.1%
Ambrox® ³⁾	0.55%
Total	100%

1) Methyl dihydrojasmonate, Firmenich SA, Geneva, Switzerland

2) 1-(octahydro-2,3,8,8-tetramethyl-2-naphtalenyl)-1-ethanone, International Flavors &

10 Fragrances, USA

3) (-)-(8R)-8,12-epoxy-13,14,15,16-tetranorlabdane, Firmenich SA, Geneva, Switzerland

Table 2b: Perfume oil B composition

Ingredient	Parts
Ethyl 2-methyl-pentanoate	3.2
Eucalyptol	7.8

Aldehyde C10	0.75
2,4-Dimethyl-3-cyclohexene- 1-carbaldehyde ¹⁾	0.75
Citronellyl nitrile	4.3
Isobornyl acetate	3.0
Verdox ²⁾	9.8
Citronellyl acetate	1.3
2-methylundecanal	3.0
Diphenyloxide	0.8
Aldehyde C12	1.3
Dicyclopentadiene acetate	9.85
Ionone beta	3.3
Undecalactone gamma	18.75
Hexyl salicylate	15.9
Benzyl salicylate	16.2

1) Origin: Firmenich SA, Geneva, Switzerland

2) Trademark from IFF; 2-tert-butyl-1-cyclohexyl acetate

Capsules A: Preparation of capsules A with Bovin Serum Albumin as stabilizer

5

Table 3: Capsule A compositions

Capsules	Base	Base (g)	Amine	Amine (g)	[NH ₂] /[COCl] (mol/mol)
A1	No base	0	Ethylene diamine	0.48	0.8
A2	No base	0	3-aminopropyl triethoxysilane	3.76	0.8
A3	Sodium carbonate	0.8	mXDA	1.5	1.5
A4	Sodium bicarbonate	0.8	mXDA	1	1
A5	Sodium bicarbonate	0.5	mXDA	1.4	1.5

Capsules B: Preparation of capsules B with Bovin Serum Albumin5 **Table 4:** Capsule B compositions

Capsules	1 st amino compound	(g)	2 nd amino compound	(g)
B1	L-Lysine	2.5	EDA	0.48
B2	L-Lysine	2.5	Diethylene triamine	0.82
B3	L-Lysine	2.5	m-XDA	1.08
B4	L-Lysine	2.5	Spermine dihydrate	1.9
B5	L-Lysine	2.5	Guanidine carbonate	0.72
B6	3-aminopropyl triethoxysilane	3.76	EDA	0.48
B7	3-aminopropyl triethoxysilane	3.76	Diethylene triamine	0.82
B8	L-Lysine	2.5	Cystamine dihydrochloride	1.8
B9	L-Lysine	2.5	L-cystine bis(tert-butyl)ester dihydrochloride	3.4
B10	3-aminopropyl triethoxysilane	3.76	Cystamine dihydrochloride	1.8
B11 **	L-Lysine	2.5	EDA	0.24
B12 **	3-aminopropyl triethoxysilane	3.76	Spermine dihydrate	1.9
B13 **	3-aminopropyl triethoxysilane	3.76	Ethylene diamine	0.48

** Protein and acid chloride mixed at 60°C for 1 hour

Capsules C: Preparation of capsules C with Sodium Caseinate5 **Table 5:** Capsule C compositions

Capsules	1 st amino compound	(g)	2 nd amino compound	(g)	Mass Colloidal Stabilizer
C1	L-Lysine	2.5	EDA	0.48	0.95
C2	L-Lysine	2.5	EDA	0.24	0.95
C3	L-Lysine	2.5	EDA	0.24	2
C4	L-Lysine	2.5	-	-	2
C5	L-Lysine	2.5	Cystamine Dihydrochloride	0.90	0.95

Capsules D: Preparation of capsules D with different stabilizers and combination of them in the oil phase

- 10 Capsules D were prepared in the presence of different proteins and diamines according to the protocol used to prepare capsules B in the presence of L-Lysine (2.5 g) as an amino compound A and ethylene diamine (0.48g or 0.24g) as an amino compound B.

Table 6: Capsules D compositions

15

Capsules	Protein	Protein (g)
D1	Gum Arabic	0.95
D2	Zein	0.95
D3	White Egg	0.95
D4	New Rice	0.95
D5	Bi Pro LE 063-18-420	0.95

D6	Bio Pure β -lacto globulin	0.95
D7	Sodium Caseinate / Bi Pro LE 063-18-420	2
D8	Sodium Caseinate / Lactose	2

Capsules E: Preparation of capsules E with a mixture of acyl chloride and caseinate in the oil phase

- 5 Capsules E were prepared in the presence of Caseinate and diamines according the protocol used to prepare capsules B in the presence of L-Lysine (2.5 g) as an amino compound A and ethylene diamine (0.48g or 0.24g) as an amino compound B.

Table 7: Capsules E compositions

10

Capsules	Acyl chloride	Molarity OCl mmol
E1	Isophthaloyl chloride/BTC	10/10
E8	Isophthaloyl chloride/BTC	5/15

Capsules F: Preparation of capsules F with only one amino compound and caseinate and mixture of proteins in the oil phase

15

Capsules F were prepared in the presence of protein of mixture of proteins according the protocol used to prepare capsules B with only the presence of L-Lysine (2.5 g) as an amino compound A in the water phase during the process of emulsification. No additional amino compound B is added after the emulsion.

20

Table 8: Capsules F compositions

Capsules	Protein	(g)
F1	Caseinate	2
F2	75% Caseinate/25%Whey protein	2

Example 2

Storage stability in a Fabric Softener composition

- 5 The storage stability of the capsules in fabric softener was evaluated. Capsule dispersion (0.27 g) of the present invention (with encapsulated perfume oil B) was diluted in the fabric softener composition described in Table 9 (29.73 g). The softener was stored for up to one month at 37°C. The amount of perfume having leaked out of the capsules was then measured by solvent extraction and GC-FID analysis (Table 10).

10

Table 9: Fabric Softener composition

Product	Wt %
Stepantex VL 90A	8.88
Calcium Chloride Sol. 10%	0.36
Proxel GXL	0.04
Perfume	1
Water	89.72
TOTAL	100

Table 10: Oil leakage of microcapsules in Fabric Softener composition

Capsules	Leakage 3 days (%)	Leakage 30 days (%)
A1	7	25
A3	14	25
A4	4	5

A5	8	8
A2	9.5	20.8
B8	10.8	16.6
B11	8.2	19.7
C1	22	32
C2	11	25
C3	13	23
C4	25	49
C5	24	33
D1	21	29
D4	22	37
D5	19	38
D6	9	22
D7	11	22
D8	16	27
E1	21	50
E2	20	48
F1	25	49
F2	20	56

One can conclude that microcapsules of the invention show a satisfactory stability in a challenging medium.

5

Example 3

Spray-dried microcapsules preparation

Emulsions A-E having the following ingredients are prepared.

10 **Table 11:** Composition of Emulsions A-E and composition of granulated powder A-E after spray-drying

Ingredients	Emulsion A	Emulsion B	Emulsion C	Emulsion D	Emulsion E
Modified starch ¹⁾	2.6%	2.6%	2.6%	12.5%	2%
Maltodextrin ²⁾	26.8%	22.8%	19.3%	0%	19.1%
Maltose ³⁾	0%	0%	0%	7.9%	0%
Citric Acid	0%	0%	0%	1%	0%
Tripotassium Citrate	0%	0%	0%	1.9%	0%
Microcapsules A-F	12.0%	24%	37.0%	8.9%	56.2%
Silica ⁴⁾	1.1%	1.1%	1.1%	0%	0%
Free Perfume C ⁵⁾	0%	0%	0%	11%	0%
Water	57.6%	49.6%	40.1%	56.9%	22.7%
	Granule A	Granule B	Granule C	Granule D	Granule E
Modified starch ¹⁾	7.5%	7.4%	7.2%	31.6%	4.9%
Maltodextrin ²⁾	77.4%	65.5%	53.8%	0%	44.7%
Maltose	0%	0%	0%	20.9%	

Citric Acid	0%	0%	0%	2.6%	0%
Tripotassium citrate	0%	0%	0%	4.9%	0%
Encapsulated perfume C	0%	0%	0%	28.1%	0%
Microcapsules I-L	12.%	24.1%	36.1%	9.8%	48.4%
Silica	3.0	3.0%	2.9%	2.0%	2%
Fragrance loading in powder after spray-drying	10.1%	20.1%	30%	35.8%	40.2%

- 1) Capsul™, Ingredion
- 2) Maltodextrin 10DE origin: Roquette
- 3) Maltose, Lehmann & Voss
- 5 4) Silica, Evonik
- 5) see table 12

Table 12: Composition of Perfume C

Component	%
ACÉTATE DE 4-(1,1-DIMÉTHYLÉTHYL)-1-CYCLOHEXYLE ¹⁾	14.50
LINALOL BJ	10.50
LILIAL® ²⁾	10.00
ISO E SUPER ³⁾	10.00
CITRONELLYL NITRILE	9.00

DIPHENYLOXYDE	6.50
ISOBORNYL ACETATE	6.00
BETA IONONE	6.00
TRICYCLO[5.2.1.0~2,6~]DEC-3-EN-8-YL ACETATE (A) + TRICYCLO[5.2.1.0~2,6~]DEC-4-EN-8-YL ACETATE (B) 4)	5.50
ETHER MT	4.00
HEDIONE® 5)	4.00
GERANIOL 60	3.00
CITRAL	2.50
ALDEHYDE C 10	2.50
ALLYL HEPTANOATE	2.50
ETHYL METHYL-2-BUTYRATE	1.50
GERANYL ACETATE	1.00
2,4-DIMETHYL-3-CYCLOHEXENE-1-CARBALDEHYDE 6)	1.00

1) Firmenich SA, Switzerland

2) 3-(4-tert-butylphenyl)-2-methylpropanal, Givaudan SA, Vernier, Switzerland

3) 1-(octahydro-2,3,8,8-tetramethyl-2-naphtalenyl)-1-ethanone, International Flavors &

5 Fragrances, USA

4) Firmenich SA, Switzerland

5) Methyl dihydrojasmonate, Firmenich SA, Switzerland

6) Firmenich SA, Switzerland

10 Components for the polymeric matrix (Maltodextrin and capsulTM, or capsulTM, citric acid and tripotassium citrate) are added in water at 45-50°C until complete dissolution.

For emulsion D, free perfume C is added to the aqueous phase.

Microcapsules slurry is added to the obtained mixture. Then, the resulting mixture is then mixed gently at 25°C (room temperature).

15 Granulated powder A-E are prepared by spray-drying Emulsion A-E using a Sodeva Spray Dryer (Origin France), with an air inlet temperature set to 215°C and a throughput

set to 500 ml per hour. The air outlet temperature is of 105°C. The emulsion before atomization is at ambient temperature.

5

Example 4

Liquid scent booster composition

A sufficient amount of microcapsule slurry A-F is weighed and mixed in a liquid scent booster (Table 13) to add the equivalent of 0.2% perfume.

10

Table 13: Liquid scent booster composition

Ingredients	Amount (% wt)					
	1	2	3	4	5	6
Water	71.20%	89.5%	78.8%	79.4%	70%	70%
Propylene glycol	20.30%	-	-	-	20%	20%
Polyethylene glycol ethers of decyl alcohol ¹⁾	4.00%	6%				
Polyethylene glycol ether of Lauryl Alcohol ²⁾					4.00%	4.00%
alkyl polyglucoside C8-C10 ³⁾			8.30%	7.7%		
Deceth-3 ¹⁾	1.50%					
Lauryl lactate				1%		
Lauric acid		1.5%	1.60%			
Glyceryl Caprylate					3.00%	3.00%
Fragrance	3.00%	3.0%	3.00%	3.00%	3.00%	0%

1) Deceth-8; trademark and origin : KLK Oleo

2) Laureth-9; ; trademark and origin

3) Plantacare 2000UP; trademark and origin : BASF

15

Different ringing gel compositions are prepared (compositions 1-6) according to the following protocol.

In a first step, the aqueous phase (water), the solvent (propylene glycol) if present and surfactants are mixed together at room temperature under agitation with magnetic
5 stirrer at 300 rpm for 5 min.

In a second step, the linker is dissolved in the hydrophobic active ingredient (fragrance) at room temperature under agitation with magnetic stirrer at 300 rpm. The resulting mixture is mixed for 5 min.

Then, the aqueous phase and the oil phase are mixed together at room temperature
10 for 5 min leading to the formation of a transparent or opalescent ringing gel.

Example 5

Liquid detergent composition

15 A sufficient amount of microcapsule slurry A-F is weighed and mixed in a liquid detergent (Table 14) to add the equivalent of 0.2% perfume.

Table 14: Liquid detergent composition

Ingredients	Concentration [wt%]
Sodium C ₁₄₋₁₇ Alkyl Sec Sulfonate ¹⁾	7
Fatty acids, C ₁₂₋₁₈ and C ₁₈ -unsaturated ²⁾	7.5
C _{12/14} fatty alcohol polyglycol ether with 7 mol EO ³⁾	17
Triethanolamine	7.5
Propylene Glycol	11
Citric acid	6.5
Potassium Hydroxyde	9.5
Properase L ⁴⁾	0.2

Puradax EG L ⁴⁾	0.2
Purastar ST L ⁴⁾	0.2
Acrylates/Steareth-20 Methacrylate structuring Crosspolymer ⁵⁾	6
Deionized Water	27.4

1) Hostapur SAS 60; Origin: Clariant

2) Edenor K 12-18; Origin: Cognis

3) Genapol LA 070; Origin: Clariant

4) Origin: Genencor International

5) Aculyn 88; Origin: Dow Chemical

Example 6

Powder detergent composition

- 10 A sufficient amount of granules A-E is weighed and mixed in a powder detergent composition (Table 15) to add the equivalent of 0.2% perfume.

Table 15: Powder detergent composition

Ingredients	Part
Anionic (Linear Alkyl Benzene Sulphonates)	20%
Nonionics (Alcohol Ethoxylates (5-9 ethylene oxide)	6%
Builders (zeolites, sodium carbonate)	25%
Silicates	6%

Sodium Sulphate	35%
Others (Enzymes, Polymers, Bleach)	7.5%
Spray-dried granule powder A-E	0.5%

Example 7Concentrated All Purpose Cleaner composition

5

A sufficient amount of microcapsule slurry A-F is weighed and mixed in a concentrated all-purpose cleaner composition (Table 16) to add the equivalent of 0.2% perfume.

Table 16: concentrated all-purpose cleaner composition

10

Ingredients	Amount (% wt)	Function
Ethoxylated Alcohol (C9-C11, 8EO) ⁽¹⁾	20	Non-ionic surfactant
Sodium Dodecyl Benzene Sulfonate ⁽²⁾	16	Anionic surfactant
Sodium Cumene Sulfonate ⁽³⁾	8	Hydrotrope
Methyl chloro isothiazolinone Methyl isothiazolinone 3.3:1 ⁽⁴⁾	0.8%	preservative
Water	55.9	solvent

1) Neodol 91-8 ®; trademark and origin : Shell Chemical

2) Biosoft D-40®; trademark and origin : Stepan Company

3) Stepanate SCS®; trademark and origin : Stepan Company

4) Kathon CG®; trademark and origin : Dow Chemical Company

All ingredients are mixed together and then the mixture was diluted with water to 100%.

Example 8

Solid scent booster composition

5

The following compositions are prepared.

Table 17: Salt-based solid scent booster compositions

Ingredients	Part
Sodium chloride	95
Spray-dried granule powder A-E	5

10

Table 18: Urea-based solid scent booster compositions

Ingredients	Part
Urea (beads)	86
Spray-dried granule powder A-E	8
Bentonite	3
Perfume	3

Example 9

Shampoo composition

15

A sufficient amount of microcapsule slurry A-F is weighed and mixed in a shampoo composition (Table 19) to add the equivalent of 0.2% perfume.

Table 19: Shampoo composition

	Ingredients	Concentration
--	-------------	---------------

20

		[wt%]
A	Water deionized	44.4
	Polyquaternium-10 ¹⁾	0.3
	Glycerin 85% ²⁾	1
	DMDM Hydantoin ³⁾	0.2
B	Sodium Laureth Sulfate ⁴⁾	28
	Cocamidopropyl Betaine ⁵⁾	3.2
	Disodium Cocoamphodiacetate ⁶⁾	4
	Ethoxy (20) Stearyl Alcohol ⁶⁾	1
C	Sodium Laureth Sulfate ⁴⁾	3
	Glyceryl Laureate ⁷⁾	0.2
D	Water deionized	1
	Sodium Methylparaben ⁸⁾	0.1
E	Sodium Chloride 10% aqueous sol.	15
	Citric acid 10% aqueous sol. till pH 5.5-6	q.s.
	Perfume	0.5
	T O T A L:	100

1) Ucare Polymer JR-400, Noveon

2) Schweizerhall

3) Glydant, Lonza

4) Texapon NSO IS, Cognis

5) Tego Betain F 50, Evonik

6) Amphotensid GB 2009, Zschimmer & Schwarz

7) Monomuls 90 L-12, Gruenau

8) Nipagin Monosodium, NIPA

Polyquaternium-10 is dispersed in water. The remaining ingredients of phase A are mixed separately by addition of one after the other while mixing well after each adjunction. Then this pre-mix is added to the Polyquaternium-10 dispersion and was mixed for 5 min. Then Phase B and the premixed Phase C (heat to melt Monomuls 90L-12 in Texapon NSO IS) are added. The mixture is mixed well. Then, Phase D and Phase E are added while agitating. The pH was adjusted with citric acid solution till pH: 5.5 – 6.0.

Example 10

Shampoo composition

10

A sufficient amount of microcapsule slurry A-F is weighed and mixed in a shampoo composition (Table 20) to add the equivalent of 0.2% perfume.

Table 20: Shampoo composition

	Ingredients	Concentration [wt%]
A	Water deionized	45.97
	Tetrasodium EDTA ¹⁾	0.05
	Guar Hydroxypropyltrimonium Chloride ²⁾	0.05
	Polyquaternium-10 ³⁾	0.075
B	NaOH 10% aqueous sol.	0.3
C	Ammonium Lauryl Sulfate ⁴⁾	34
	Ammonium Laureth Sulfate ⁵⁾	9.25
	Cocamidopropyl Betaine ⁶⁾	2
	Dimethicone (&) C12-13 Pareth-4 (&) C12-13 Pareth-23 (&) Salicylic Acid ⁷⁾	2.5
D	Cetyl Alcohol ⁸⁾	1.2
	Cocamide MEA ⁹⁾	1.5

	Glycol Distearate ¹⁰⁾	2
E	Methylchloroisothiazolinone & Methylisothiazolinone ¹¹⁾	0.1
	D-Panthenol 75% ¹²⁾	0.1
	Water deionized	0.3
F	Sodium Chloride 25% aqueous sol.	0.6
	T O T A L:	100

- 1) EDETA B Powder, BASF
- 2) Jaguar C14 S, Rhodia
- 3) Ucare Polymer JR-400, Noveon
- 4) Sulfetal LA B-E, Zschimmer & Schwarz
- 5) Zetesol LA, Zschimmer & Schwarz
- 6) Tego Betain F 50, Evonik
- 7) Xiameter MEM-1691, Dow Corning
- 8) Lanette 16, BASF
- 9) Comperlan 100, Cognis
- 10) Cutina AGS, Cognis
- 11) Kathon CG, Rohm & Haas
- 12) D-Panthenol, Roche

A premix comprising Guar Hydroxypropyltrimonium Chloride and Polyquaternium-10 are added to water and Tetrasodium EDTA while mixing. When the mixture is homogeneous, NaOH is added. Then, Phase C ingredients are added and the mixture was

5 heat to 75 °C. Phase D ingredients are added and mixed till homogeneous. The heating is stopped and temperature of the mixture is decreased to RT. At 45 °C, ingredients of Phase E while mixing final viscosity is adjusted with 25% NaCl solution and pH of 5.5-6 is adjusted with 10% NaOH solution.

Rinse-off hair composition

A sufficient amount of microcapsule slurry A-F is weighed and mixed in a rinse-off composition (Table 21) to add the equivalent of 0.2% perfume.

5

Table 21: rinse-off composition

	Ingredients	Concentration [wt%]
A	Water deionized	81.8
	Behentrimonium Chloride ¹⁾	2.5
	Hydroxyethylcellulose ²⁾	1.5
B	Cetearyl Alcohol ³⁾	4
	Glyceryl Stearate (and) PEG-100 Stearate ⁴⁾	2
	Behentrimonium Methosulfate (and) Cetyl alcohol (and) Butylene Glycol ⁵⁾	4
	Ethoxy (20) Stearyl Alcohol ⁶⁾	1
C	Amodimethicone (and) Trideceth-12 (and) Cetrimonium Chloride ⁷⁾	3
	Chlorhexidine Digluconate ⁸⁾ 20% aqueous solution	0.2
D	Citric acid 10% aqueous sol. till pH 3.5-4	q.s.
	T O T A L:	100

1) Genamin KDMP, Clariant

2) Tylose H10 Y G4, Shin Etsu

10

3) Lanette O, BASF

4) Arlacel 165, Croda

5) Incroquat Behenyl TMS-50-PA- (MH), Croda

6) Brij S20, Croda

7) Xiameter MEM-949, Dow Corning

15

8) Alfa Aesar

Ingredients of Phase A are mixed until an uniform mixture was obtained. Tylose is allowed to completely dissolve. Then the mixture is heated up to 70-75°C. Ingredients of Phase B are combined and melted at 70-75°C. Then ingredients of Phase B are added to Phase A with good agitation and the mixing is continued until cooled down to 60°C. Then, ingredients of Phase C are added while agitating and keeping mixing until the mixture cooled down to 40°C. The pH is adjusted with citric acid solution till pH: 3.5 – 4.0.

Example 12

Antiperspirant spray anhydrous composition

A sufficient amount of microcapsule slurry A-F is weighed and mixed in an antiperspirant spray anhydrous composition (Table 22) to add the equivalent of 0.2% perfume.

Table 22: antiperspirant spray anhydrous composition

Ingredient	Amount (wt %)
Cyclomethicone ¹⁾	53.51
Isopropyl miristate	9.04
Silica ²⁾	1.03
Quaternium-18-Hectorite ³⁾	3.36
Aluminium Chlorohydrate ⁴⁾	33.06

1) Dow Corning[®] 345 Fluid; trademark and origin: Dow Corning

2) Aerosil[®] 200 ; trademark and origin : Evonik

3) Bentone[®] 38; trademark and origin : Elementis Specialities

4) Micro Dry Ultrafine; origin : Reheis

Using a high speed stirrer, Silica and Quaternium-18-Hectorite are added to the Isopropyl miristate and Cyclomethicone mixture. Once completely swollen, Aluminium Chlorohydrate is added portion wise under stirring until the mixture was homogeneous

and without lumps. The aerosol cans are filled with 25 % Suspension of the suspension and 75 % of Propane/Butane (2,5 bar).

Example 13

5

Antiperspirant spray emulsion composition

A sufficient amount of microcapsule slurry A-F is weighed and mixed in antiperspirant spray emulsion composition (Table 23) to add the equivalent of 0.2% perfume.

10 **Table 23:** antiperspirant spray emulsion composition

Ingredient	Amount (wt %)
Polysorbate 65 ¹⁾ (Part A)	0.95
Polyglyceryl-2 dipolyhydroxystearate ²⁾ (Part A)	1.05
Cetyl PEG/PPG-10/1 Dimethicone ³⁾ (Part A)	2.75
Cyclomethicone ⁴⁾ (Part A)	16.4
Isopropylisostearate ⁵⁾ (Part A)	4.5
Phenoxyethanol ⁶⁾ (Part A)	0.5
Ethylhexylglycerin ⁷⁾ (Part A)	0.2
C12-15 Alkyl Benzoate ⁸⁾ (Part A)	5.65
Silica Silylate ⁹⁾ (Part A)	0.1
Sodium Methylparaben ¹⁰⁾ (Part B)	0.1

Aluminium Chlorohydrate ⁽¹¹⁾ (Part B)	20
Water (Part B)	44.47
Fragrance (Part C)	3.33

- 1) Tween 65; trademark and origin : CRODA
- 2) Dehymuls PGPH; trademark and origin : BASF
- 3) Abil EM-90; trademark and origin : BASF
- 4) Dow Corning 345 fluid; trademark and origin : Dow Corning
- 5 5) Crodamol ipis; trademark and origin : CRODA
- 6) Phenoxyethanol; trademark and origin : LANXESS
- 7) Sensiva sc 50; trademark and origin : KRAFT
- 8) Tegosoft TN; trademark and origin : Evonik
- 9) Aerosil R 812; trademark and origin : Evonik
- 10 10) Nipagin mna; trademark and origin : CLARIANT
- 11) Locron L; trademark and origin : CLARIANT

The ingredients of Part A and Part B are weighted separately. Ingredients of Part A are heated up to 60°C and ingredients of Part B are heated to 55 °C. Ingredients of Part B are poured small parts while continuous stirring into A. Mixture were stirred well until the room temperature was reached. Then, ingredients of part C are added. The emulsion is mixed and is introduced into the aerosol cans. The propellant is crimped and added.

Aerosol filling: 30% Emulsion: 70% Propane / Butane 2,5 bar

20

Example 14

Deodorant spray composition

A sufficient amount of microcapsule slurry A-F is weighed and mixed in antiperspirant deodorant spray composition (Table 24) to add the equivalent of 0.2% perfume.

25

Table 24: deodorant spray composition

Ingredient	Amount (wt %)
Ethanol 95 %	90.65
Triclosan ¹⁾	0.26
Isopropyl miristate	9.09

1) Irgasan[®] DP 300; trademark and origin : BASF

5

All the ingredients according to the sequence of the Table 24 are mixed and dissolved. Then the aerosol cans are filled, crimp and the propellant is added (Aerosol filling: 40% active solution 60% Propane / Butane 2.5 bar).

10

Example 15

Antiperspirant roll-on emulsion composition

A sufficient amount of microcapsule slurry A-F is weighed and mixed in antiperspirant roll-on emulsion composition (Table 25) to add the equivalent of 0.2% perfume.

15

Table 25: antiperspirant roll-on emulsion composition

Ingredient	Amount (wt %)
Steareth-2 ¹⁾ (Part A)	3.25
Steareth-21 ²⁾ (Part A)	0.75
PPG-15 Stearyl Ether ³⁾ (Part A)	4
WATER deionised (Part B)	51
Aluminum Chlorohydrate 50% aqueous solution ⁴⁾ (Part C)	40

Fragrance (Part D)	1
--------------------	---

- 1) BRIJ 72; origin : ICI
- 2) BRIJ 721; origin : ICI
- 3) ARLAMOL E; origin : UNIQEMA-CRODA
- 4) LOCRON L; origin : CLARIAN

5

Part A and B are heated separately to 75°C; Part A is added to part B under stirring and the mixture is homogenized for 10 minutes. Then, the mixture is cooled down under stirring; and part C is slowly added when the mixture reached 45°C and part D when the mixture reached at 35 °C while stirring. Then the mixture is cooled down to RT.

10

Example 16

Antiperspirant roll-on composition

- 15 A sufficient amount of microcapsule slurry A-F is weighed and mixed in antiperspirant roll-on composition (Table 26) to add the equivalent of 0.2% perfume.

Table 26: antiperspirant roll-on composition

Ingredient	QUANTITY
Water (Part A)	45
Aluminum Chlorohydrate 50% aqueous solution ¹⁾ (Part B)	20
Alcohol Denat. (Ethanol 96%) (Part B)	30
Cetareth-12 ²⁾ (Part C)	2
Cetareth-30 ³⁾ (Part C)	2
Fragrance (Part D)	1

- 1) LOCRON L; origin: CLARIANT
- 2) EUMULGIN B-1; origin : BASF
- 3) EUMULGIN B-3; origin : BASF

- 5 The ingredients of part B are mixed in the vessel then ingredient of part A is added. Then dissolved part C in part A and B. With perfume, 1 part of Cremophor RH40 for 1 part of perfume is added while mixing well

Example 17

10 Antiperspirant roll-on composition

A sufficient amount of microcapsule slurry A-F is weighed and mixed in antiperspirant roll-on emulsion composition (Table 27) to add the equivalent of 0.2% perfume.

15 **Table 27:** antiperspirant roll-on emulsion composition

Ingredient	Amount (wt %)
Water (Part A)	50.51
Hydroxyethylcellulose ¹⁾ (Part A)	0.71
Ethanol 95 % (Part B)	40.40
1,2-Propylene Glycol (Part B)	5.05
Triclosan ²⁾ (Part B)	0.30
PEG-40 Hydrogenated castor oil ³⁾ (Part C)	3.03

- 1) Natrosol[®] 250 H; trademark and origin: Ashland
- 2) Irgasan[®] DP 300; trademark and origin : BASF
- 3) Cremophor[®] RH 40; trademark and origin : BASF

- 20 Part A is prepared by sprinkling little by little the Hydroxyethylcellulose in the water whilst rapidly stirring with the turbine. Stirring is continued until the Hydroxyethylcellulose is entirely swollen and giving a limp gel. Then, Part B is poured little by little in Part A whilst continuing stirring until the whole is homogeneous. Part C is added.

Example 18Deodorant pump without alcohol formulation

- 5 A sufficient amount of microcapsule slurry A-F is weighed and mixed in the following composition (Table 28) to add the equivalent of 0.2% perfume.

Table 28: deodorant composition

Ingredients	Amount (wt %)
C12-15 Alkyl Lactate ¹⁾	5
Dimethicone ²⁾	91.6
Cetyl Lactate ³⁾	1
Octyldodecanol ⁴⁾	0.8
Triclosan ⁵⁾	0.1
PERFUME	1.5

- 10 1) Ceraphyl 41; trademark and origin ASHLAND
 2) DOW CORNING 200 FLUID 0.65cs; trademark and origin DOW CORNING CORPORATION
 3) Ceraphyl 28; trademark and origin ASHLAND
 4) Eutanol G; trademark and origin BASF
 15 5) Irgasan[®] DP 300; trademark and origin : BASF

All the ingredients of Table 28 are mixed according to the sequence of the table and the mixture is heated slightly to dissolve the Cetyl Lactate.

Example 19Deodorant pump with alcohol formulation

- 5 A sufficient amount of microcapsule slurry A-F is weighed and mixed in the following composition (Table 29) to add the equivalent of 0.2% perfume.

Table 29: deodorant composition

Ingredients	Amount (wt %)
Ethyl Alcohol (Part A)	60
PEG-6 Caprylic/Capric Glycerides ¹⁾ (Part A)	2
Water (Part A)	35.6
PEG-40 Hydrogenated Castor Oil ²⁾ (Part B)	0.4
PERFUME (Part B)	2

1) Softigen 767; trademark and origin CRODA

10 2) Cremophor[®] RH 40; trademark and origin : BASF

Ingredients from Part B are mixed together. Ingredients of Part A are dissolved according to the sequence of the Table and are poured into part B.

15

20

Example 20Talc formulation

A sufficient amount of granules A-F is weighed and mixed in introduced in a standard
 5 talc base: 100% talc, very slight characteristic odor, white powder, origin: LUZENAC to
 add the equivalent of 0.2% perfume.

Example 21Shower-gel Reference

10

A sufficient amount of microcapsule slurry A-F is weighed and mixed in the following
 composition (Table 30) to add the equivalent of 0.2% perfume.

15 **Table 30:** shower gel composition

Ingredients	Amount (% wt)	Function
WATER deionised	49.350	Solvent
Tetrasodium EDTA ¹⁾	0.050	Chelating agent
Acrylates Copolymer ²⁾	6.000	Thickener
Sodium C12-C15 Pareth Sulfate ³⁾	35.000	Surfactant
Sodium Hydroxide 20% aqueous solution	1.000	pH adjuster
Cocamidopropyl Betaine ⁴⁾	8.000	Surfactant
Methylchloroisothiazolinone and Methylisothiazolinone ⁵⁾	0.100	Preservative
Citric Acid (40%)	0.500	pH adjuster

1) EDETA B POWDER; trademark and origin: BASF

2) CARBOPOL AQUA SF-1 POLYMER; trademark and origin: NOVEON

3) ZETESOL AO 328 U; trademark and origin: ZSCHIMMER & SCHWARZ

4) TEGO-BETAIN F 50; trademark and origin: GOLDSCHMIDT

5) KATHON CG; trademark and origin: ROHM & HASS

Ingredients are mixed, pH is adjusted to 6-6.3 (Viscosity: 4500cPo +/-1500cPo

5 (Brookfield RV / Spindle#4 / 20RPM)).

Example 22

Shower-gel composition

- 10 A sufficient amount of microcapsule slurry A-F is weighed and mixed in the following composition (Table 31) to add the equivalent of 0.2% perfume.

Table 31: shower gel composition

Ingredients	Amount (% wt)	Function
WATER deionized	52.40	Solvent
Tetrasodium EDTA ¹⁾	0.10	Chelating agent
Sodium Benzoate	0.50	Preservative
Propylene Glycol	2.00	Solvent
Sodium C12-C15 Pareth Sulfate ²⁾	35.00	Surfactant
Cocamidopropyl Betaine ³⁾	8.00	Surfactant
Polyquaternium-7 ⁴⁾	0.20	Conditioning agent
Citric Acid (40%)	1.00	pH adjuster
Sodium Chloride	0.80	Viscosity adjuster

1) EDETA B POWDER; trademark and origin: BASF

15 2) ZETESOL AO 328 U; trademark and origin: ZSCHIMMER & SCHWARZ

3) TEGO-BETAIN F 50; trademark and origin: GOLDSCHMIDT

4) MERQUAT 550; trademark and origin: LUBRIZOL

Ingredients are mixed, pH is adjusted to 4.5 (Viscosity: 3000cPo +/-1500cPo (Brookfield RV / Spindle#4 / 20RPM)).

5

Example 23

Shower-gel composition

A sufficient amount of microcapsule slurry A-F is weighed and mixed in the following composition (Table 32) to add the equivalent of 0.2% perfume.

10

Table 32: shower gel composition

Ingredients	Amount (% wt)	Function
WATER deionized	50.950	Solvent
Tetrasodium EDTA ¹⁾	0.050	Chelating agent
Sodium Benzoate	0.500	Preservative
Glycerin 86%	3.500	Solvent
Sodium Laureth Sulfate ²⁾	27.000	Surfactant
Polyquaternium-7 ³⁾	1.000	Conditioning Agent
Coco-Betaine ⁴⁾	6.000	Surfactant
PEG-120 Methyl Glucose trioleate ⁵⁾	1.000	Thickener
Citric Acid (40%)	1.000	pH adjuster
Glycol Distearate & Laureth-4 & Cocamidopropyl Betaine ⁶⁾	3.000	Pearlizing agent
Sodium Chloride 20%	5.000	Viscosity adjuster
PEG-40 Hydrogenated Castor Oil ⁷⁾	1.000	Viscosity adjuster

1) EDETA B POWDER; trademark and origin: BASF

- 2) Texapon NSO IS; trademark and origin: COGNIS
- 3) MERQUAT 550; trademark and origin: LUBRIZOL
- 4) DEHYTON AB-30; trademark and origin: COGNIS
- 5) GLUCAMATE LT; trademark and origin: LUBRIZOL
- 5 6) EUPERLAN PK 3000 AM; trademark and origin: COGNIS
- 7) CREMOPHOR RH 40; trademark and origin: BASF

Ingredients are mixed, pH is adjusted to 4.5 (Viscosity: 4000cPo +/-1500cPo (Brookfield RV / Spindle#4 / 20RPM))

10

Example 24

Hand Dishwash

A sufficient amount of microcapsule slurry A-F is weighed and mixed in the following
15 composition (Table 33) to add the equivalent of 0.2% perfume.

Table 33: Hand dishwash composition

Ingredients	Amount (% wt)	Function
Linear alkylbenzene sulfonic acid ⁽¹⁾	20	Anionic surfactant
Diethanolamide ⁽²⁾	3.5	Foam booster
Sodium Hydroxide (50%) ⁽³⁾	3.4	pH Adjuster/neutralizer
Secondary alcohol ethoxolate ⁽⁴⁾	2.5	Non-ionic surfactant
Sodium xylene sulfonate	6.3	Hydrotrope
Water	64.3	Solvent

--	--	--

- 1) Biosoft S-118®; trademark and origin : Stepan Company
- 2) Ninol 40-CO®; trademark and origin : Stepan Company
- 3) Stepanate SXS®; trademark and origin : Stepan Company
- 4) Tergitol 15-S-9®; trademark and origin : Dow Chemical Company

5

Water with sodium hydroxide and diethanolamide are mixed. LAS is added. After the LAS is neutralized, the remaining ingredients are added. The pH was Checked (=7-8) and adjusted if necessary.

10

Example 25

Toothpaste formulation

A sufficient amount of microcapsule slurry M (corresponding to microcapsules A except that a menthol flavor is encapsulated) is weighed and mixed in the following composition (Table 34) to add the equivalent of 0.2% flavor.

15

Table 34: Toothpaste formulation

Ingredients	Amount (% wt)
Polyethylene glycol 400	2.0%
Xanthan Gum	0.6%
Sorbitol 70% Solution	50%
Sodium Fluoride	0.220%
Sodium Benzoate	0.2%
Water	15.230%
Hydrated Silica ¹⁾	22.0%
Hydrated Silica ²⁾	7.0%
Titanium Dioxide CI77891	0.5%
Sodium Lauryl Sulfate	1.250%
Flavor	1.2%

TOTAL	100%

1) Tixosil 73; trademark and origin :

2) Tixosil 43; trademark and origin :

Example 26

5

Dicalcium Phosphate based toothpaste formulation

A sufficient amount of microcapsule slurry M (corresponding to microcapsules A except that a menthol flavor is encapsulated) is weighed and mixed in the following composition (Table 35) to add the equivalent of 0.2% flavor.

10

Table 35: Toothpaste formulation

Ingredients	Amount (% wt)
Sodium carboxymethyl cellulose	1.2%
Flavor	1.2%
DI/Purified Water	Q.S to Final Wt.
Sodium Lauryl Sulfate	1.3%
Glycerine	20.0%
Sodium Saccharin	0.2%
Dicalcium phosphate dihydrate	36.0%
Methylparaben	0.2%
Silica ¹⁾	3.0%
TOTAL	100%

1) Aerosil®200; trademark and origin:

15

Example 27**Mouthwash alcohol free formulation**

A sufficient amount of microcapsule slurry M (corresponding to microcapsules A except
 5 that a menthol flavor is encapsulated) is weighed and mixed in the following composition
 (Table 36) to add the equivalent of 0.2% flavor.

Table 36: Mouthwash formulation

Ingredients	Amount (% wt)
Propylene Glycol	10%
Flavor	0.240%
DI/Purified Water	Q.S to Final Wt.
Poloxamer 407 NF	0.240%
Sodium Lauryl Sulfate	0.040%
Sorbitol 70% Solution	10.0%
Sodium Saccharin	0.030%
Glycerine	3.0%
Sodium Benzoate	0.100%
Sucralose	0.020%
Benzoic Acid	0.050%
TOTAL	100%

10

Example 28**Mouthwash formulation**

15 A sufficient amount of microcapsule slurry M (corresponding to microcapsules A except
 that a menthol flavor is encapsulated) is weighed and mixed in the following composition
 (Table 37) to add the equivalent of 0.2% flavor.

Table 37: Mouthwash formulation

Ingredients	Amount (% wt)
Ethyl Alcohol 190 Proof	15.0%
Flavor	0.240%
DI/Purified Water	Q.S to Final Wt.
Poloxamer 407 NF	0.240%
Sodium Lauryl Sulfate	0.040%
Sorbitol 70% Solution	10.0%
Sodium Saccharin	0.030%
Glycerine	3.0%
Sodium Benzoate	0.100%
Sucralose	0.020%
Benzoic Acid	0.050%
TOTAL	100%

5

10

15

CLAIMS

1. A process for preparing a polyamide core-shell microcapsule slurry comprising the following steps:

- 5 a) Dissolving at least one acyl chloride and at least one stabilizer in a hydrophobic material, preferably a perfume to form an oil phase;
- b) Dispersing the oil phase obtained in step a) into a water phase comprising an amino compound A to form an oil-in-water emulsion; and
- 10 c) Performing a curing step to form polyamide microcapsules in the form of a slurry,
- wherein at least an amino compound B is added in the water phase before the formation of the oil-in-water emulsion and/or in the oil-in water emulsion obtained after step b),
- wherein said amino compound A is an amino acid.

15

2. The process according to claim 1, wherein the acyl chloride is chosen in the group consisting of benzene-1,3,5-tricarbonyl chloride, benzene-1,2,4-tricarbonyl trichloride, benzene-1,2,4,5-tetracarbonyl tetrachloride, cyclohexane-1,3,5-tricarbonyl trichloride, isophthalyol dichloride, diglycolyl dichloride, succinic dichloride, and mixtures thereof.

20

3. The process according to any one of the preceding claims, wherein the amino acid is L-Lysine.

4. The process according to any one of the preceding claims, wherein the amino
25 compound B is chosen in the group consisting of cystamine, cystamine hydrochloride, cystine, cystine hydrochloride, cystine dialkyl ester, cystine dialkyl ester hydrochloride, a xylylene diamine, 1,2-diaminocyclohexane, 1,4-diaminocyclohexane, L-lysine, L-Lysine ethyl ester, polyetheramines, ethylene diamine, diethylene triamine, spermine, spermidine, polyamidoamine (PAMAM), guanidine carbonate, chitosan, tris-(2-
30 aminoethyl)amine, 3-aminopropyltriethoxysilane, L-arginine and mixtures thereof.

5. The process according to any one of the preceding claims, wherein amino-compound A and amino-compound B are different.

5 6. The process according to any one of the preceding claims, wherein the stabilizer is chosen in the group consisting of gum Arabic, modified starch, polyvinyl alcohol, polyvinylpyrrolidone, carboxymethylcellulose, anionic polysaccharides, acrylamide copolymer, inorganic particles, proteins and mixtures thereof, or wherein the stabilizer is
10 a protein chosen in the group consisting of soy protein, rice protein, whey protein, white egg albumin, sodium caseinate, gelatin, bovine serum albumin, hydrolyzed soy protein, hydrolyzed sericin, pseudocollagen, silk protein, sericin powder, and mixtures thereof.

15

7. The process according to any one of the preceding claims, wherein the molar ratio between the functional group NH_2 of the amino compound B and the functional group COCl of the acyl chloride is comprised between 0.01 and 7.5.

20 8. The process according to any one of the preceding claims, wherein the weight ratio between acyl chloride and the hydrophobic material is comprised between 0.01 and 0.09.

9. The process according to any one of the preceding claims, wherein the
25 hydrophobic material is a hydrophobic active ingredient, preferably wherein the hydrophobic active ingredient comprises a perfume.