optionally substituted by one or more groups selected from –OH, halogen, -CN, (C_1-C_6)alkyl, (C_1-C_6)alkoxy, (C_1-C_6)haloalkyl, (C_1-C_6)haloalkoxy and aryl(C_1-C_6)alkyl;

n is at each occurrence independently 0 or an integer from 1 to 3;

m is at each occurrence independently an integer from 1 to 3;

E is absent or is selected from -O-, -NR_7-, -NR_7-C(O)-, -C(O)-NR_7-, -OC(O)-, -C(O)-$(CH_2)_n$-O-; -NR_7-C(O)-$(CH_2)_n$-O-, -NR_7-C(O)-NR_7- and -S-;


G is arylene or heteroarylene, optionally substituted by one or more substituents selected from halogen atoms, -OH, oxo (=O), -SH, -NO_2, -CN, -$CON(R_6)_2$, -NH_2, -$NHCOR_6$, -CO_2R_6, (C_1-C_{10})alkylsulfanyl, (C_1-C_{10})alkylsulfinyl, (C_1-C_{10})alkylsulfonyl, (C_1-C_{10})alkyl, aryl, haloaryl, heteroaryl and (C_1-C_{10})alkoxy;

R_1 is selected from (C_3-C_8)cycloalkyl, (C_3-C_8)heterocycloalkyl, aryl, heteroaryl, aryl(C_1-C_6)alkyl, heteroaryl(C_1-C_6)alkyl and (C_3-C_8)cycloalkyl(C_1-C_6)alkyl, optionally substituted by one or more group selected independently from halogen, (C_1-C_8)alkyl, and (C_1-C_{10})alkoxy;

s is 0 or an integer from 1 to 3;

R_2 is a nitrogen containing group which is selected from:

- a group (a) which is -NR_3R_4 wherein R_3 and R_4 are independently hydrogen or (C_1-C_4) alkyl; and
- a group (b) of formula J1, J2, J3, J4 or J5



J1 **J2** **J3** **J4** **J5**

R_5 is a group of formula **K**