


This Sheet Has Been Replaced



PROCESS FOR THE MANUFACTURE OF 2-ALKOXYMETHYLACROLEIN

It is an object of this invention to provide a convenient and effective method for the manufacture of 2-alkoxymethylacrolein compounds useful in the preparation of pyridine and quinoline herbicide intermediates. There is also provided a process for the manufacture of 3-alkoxymethylquinolines from 2-alkoxymethylacrolein.

The present invention relates to a process for the manufacture of compounds of formula I




(I)

wherein R is C₁ to C₆ alkyl.

The formula I alkoxymethylacrolein compound is prepared by reacting an alcohol, ROH wherein R is as described for formula I, with at least one molar equivalent of acrolein in the presence of an acid, a catalytic amount of a trisubstituted amine and a solvent to form an intermediate and reacting said intermediate with at least one molar equivalent of formaldehyde in the presence of an acid, a catalytic amount of disubstituted amine and a solvent to obtain the desired formula I compound.

The invention further relates to a process for the manufacture of 2-alkoxymethylacrolein compounds


of formula I which comprises reacting a suitable compound of formula II



wherein R is as described for formula I and W is CHO or $\text{CH}(\text{OR}_1)_2$ and R_1 is C_1 to C_4 alkyl, with at least one molar equivalent of formaldehyde in the presence of an acid, a catalytic amount of a disubstituted amine and a solvent.

The invention also relates to the use of compounds of formula I in the manufacture of important quinoline and pyridine herbicidal intermediates.

The present invention provides a process for the manufacture of compounds of formula I. The compounds of formula I may be used to prepare 3-alkoxymethylquinolines and 5-alkoxymethylpyridine-2,3-dicarboxylates which are key intermediate compounds in the preparation of 2-(imidazolin-2-yl)nicotinate herbicides. For example, the formula I compounds of the present invention may be used to prepare 3-alkoxymethylquinolines of formula III



wherein

R is C_1 - C_6 alkyl;


R_4 is hydrogen or C_1 - C_4 alkyl; and

R_5 , R_6 and R_7 are each independently hydrogen or OR_4 .



3-Alkoxyethylquinolines may be prepared by reacting a substituted aniline of formula IV with at least one molar equivalent of a 2-alkoxyethylacrolein of formula I in the presence of an acid and a solvent and optionally in the presence of a substituted nitrobenzene of formula V, preferably at an elevated temperature, to form desired formula III compounds. The above reaction scheme is shown in flow diagram I.

FLOW DIAGRAM I




Acids suitable for use in the process used to prepare formula III compounds are strong organic and mineral acids such as sulfuric acid, phosphoric acid and hydrochloric acid. Solvents suitable for use in the process used to prepare formula III compounds are water and ROH alcohols wherein R is C_1-C_6 alkyl. Preferred solvents are ROH alcohols wherein R corresponds to the R of the formula I compound with methanol being a most preferred solvent. Reaction temperatures of from about $30^\circ C$ to $120^\circ C$, preferably about $50^\circ C$ to

110°C, are suitable for use in the process used to prepare formula III compounds.

The thus-obtained formula III 3-alkoxymethyl-quinoline compounds may be oxidized to form 5-alkoxymethylpyridine-2,3-dicarboxylic acids of formula VI. The above reaction scheme is shown in flow diagram II.



FLOW DIAGRAM II



Among the methods suitable to oxidize formula III compounds to formula VI compounds are nitric acid oxidation, base peroxide oxidation, base peroxide followed by sodium hypochlorite oxidation, chlorate catalyzed by vanadium oxidation, ozone oxidation and the like.



Advantageously, the formula I compounds of the present invention may also be reacted with an α -halo- β -keto ester in the presence of an ammonium salt to form the corresponding pyridine-2,3-dicarboxylate product as shown in flow diagram III.

Flow Diagram III




15 Further, 2-alkoxymethylacrolein compounds of formula I may be reacted with a dialkyl dihalomaleate in the presence of ammonia to form the desired pyridinedicarboxylate product as shown in flow diagram IV.

Flow Diagram IV



It has now been found, that the alkoxymethyl compounds of formula I may be effectively prepared in an efficient process from readily available starting materials. In accordance with the method of invention, an appropriate alcohol having the formula, ROH wherein R is C₁ to C₆ alkyl, may be reacted with at least one molar equivalent of acrolein in the presence of an acid, a catalytic amount of a trisubstituted amine and a solvent to form an intermediate and the intermediate may be reacted with at least one molar equivalent of formaldehyde in the presence of an acid, a disubstituted amine and a solvent to form the desired formula I alkoxymethylacrolein compound. The reaction is shown in flow diagram V.


Flow Diagram V



In general, the formation of the intermediate and the formula I product are temperature dependent, that is, increased reaction temperature increases the rate of formation. Convenient reaction times may be obtained by increasing the reaction temperature to about 20° to 110°C, preferably about 75° to 100°C. Suitable reaction solvents are water or mixtures of water and a water-miscible organic solvent. Acids suitable for use in the present process are strong

mineral acids, preferably polybasic acids such as sulfuric acid and phosphoric acid. Trisubstituted amines, $N(R_2)_3$, wherein R_2 is C_1-C_4 alkyl, C_1-C_4 alkanol and the like are suitable for use in the present invention. Disubstituted amines, $NH(R_3)_2$, wherein R_3 is C_1-C_6 alkyl, preferably dibutylamine, may be used in the inventive process. It is intended formaldehyde be used in any of its readily available forms and preferably as an aqueous solution of about 37% concentration.


Advantageously, compounds of formula I may also be prepared by reacting a compound of formula II



(II)

wherein R is C_1-C_6 alkyl and W is CHO or $CH(OR_1)_2$ and R_1 is C_1-C_4 alkyl, with at least one molar equivalent of formaldehyde in the presence of an acid, a catalytic amount of a disubstituted amine, $NH(R_3)_2$ wherein R_3 is C_1-C_4 alkyl, and a solvent. The reaction is shown in flow diagram VI.

Flow Diagram VI



Compounds of formula II may be prepared according to methods known in the art. The formaldehyde employed in the above process may be in any of its readily available forms and preferably as an


aqueous solution of about 37%. Acids suitable for use are strong mineral acids such as those mentioned hereinabove and preferably sulfuric acid or phosphoric acid. The reaction is temperature dependent, therefore convenient reaction times may be obtained by elevating the reaction temperature to about 20° to 110°C, preferably about 75° to 100°C.

In order to present a more clear understanding of the invention, the following examples are set forth. The examples are primarily for the purpose of demonstrating more specific details thereof and the invention is not to be limited thereby except as defined in the claims.

Unless otherwise noted, all parts are parts by weight. The term NMR designates nuclear magnetic resonance spectroscopy.

EXAMPLE 1

Preparation of 3-(Methoxymethyl)-8-quinolinol




A solution of methanol and sulfuric acid (10.0 g, 0.1 mol) is heated to 65°C, treated with o-aminophenol (4.36 g, 0.04 mol) and o-nitrophenol (2.78 g, 0.02 mol), heated to 70°C, treated with methoxymethacrolein (6.0 g, 0.06 mol) over 40 minutes

at 85° to 90°C and diluted with water. The aqueous mixture is adjusted to about pH 2 with 50% sodium hydroxide solution and filtered. The filtrate is adjusted to about pH 7 with 50% sodium hydroxide solution and extracted with chloroform. The combined organic extracts are concentrated in vacuo to give the title product as a solid which is identified by ¹H and ¹³CNMR spectral analyses.

Using essentially the same procedure, but substituting *o*-anisidine for *o*-aminophenol, 8-methoxy-3-(methoxymethyl)quinoline is obtained.

EXAMPLE 2


Preparation of 5-(Methoxymethyl)-2,3-pyridinedicarboxylic acid



A mixture of 3-(methoxymethyl)-8-quinolinol (0.92 g, 4.9 mmol) in water is treated with 50% sodium hydroxide solution (1.0 g real, 25.0 mmol), heated to 85°C and treated with 30% hydrogen peroxide solution (3.4 g real, 100.0 mmol) over 2 hours while maintaining the temperature between 65°-90°C. The reaction mixture (26.0 g) is then cooled to room temperature, assayed by HPLC and found to contain 3.15% of the title product (80% yield).

EXAMPLE 3


Preparation of Dimethyl 5-(methoxymethyl)-2,3-pyridine-dicarboxylate



10 Ozone is bubbled through a solution of
 3-(methoxymethyl)-8-quinolinol (0.145 g, 0.78 mmol) in
 90% acetic acid at room temperature. After 20 minutes,
 the reaction mixture is concentrated in vacuo and the
 resulting solid is dissolved in 95% ethanol and treated
 15 with excess diazomethane. After stirring at room
 temperature for 14 hours, the reaction mixture is
 treated with acetic acid and concentrated in vacuo to
 give a yellow oil. Flash chromatography of the oil
 using silica gel and a 4:1 to 2:1 hexanes/ethyl acetate
 20 solution gives the title product as a colorless oil,
 0.072 g (41% yield), which is identified by ¹H and
¹³CNMR spectral analyses.

EXAMPLE 4

Preparation of Methyl 3-formyl-5-(methoxymethyl)-picolinate



Ozone is bubbled through a solution of 8-methoxy-3-(methoxymethyl)quinoline (0.134 g, 0.66 mmol) in a 9:1 acetonitrile/water mixture at 0°C. After 10 minutes, the reaction mixture is concentrated in vacuo and the resulting oil is dissolved in methanol and treated with excess diazomethane. After stirring for 14 hours at room temperature, the reaction mixture is treated with acetic acid and concentrated in vacuo to obtain a red oil. Flash chromatography of the oil using silica gel and a 3:1 hexanes/ethyl acetate mixture gives the title product as a colorless oil which is identified by ¹HNMR spectral analysis.

Using essentially the same procedure, but substituting 90% acetic acid for the 9:1 acetonitrile/water mixture, the title product is obtained as a colorless oil which is identified by ¹HNMR spectral analysis.

EXAMPLE 5

Preparation of 2-methoxymethylacrolein from acrolein

A stirred mixture of acrolein (112g, 2.0 mole), methanol (310g, 9.08 mole), triethanolamine (7.5g, 0.05 mole) and 85% phosphoric acid (5.7g, 0.049 mole) in water is heated at reflux temperature for 9 hours, cooled to room temperature and filtered. The filtrate is diluted with water and treated with a 37% formaldehyde solution (162g, 2.0 mole formaldehyde), concentrated sulfuric acid (11.6g, 0.11 mole), and dibutylamine (27g, 0.21 mole), heated at reflux temperature for 4 hours, cooled to room temperature and extracted with methylene chloride. The extracts are combined and concentrated and the concentrate is vacuum distilled to give the title product, 88g (44% yield) bp 64-66°/70mm Hg, identified by NMR analysis.

EXAMPLE 6

Preparation of 2-ethoxymethylacrolein from acrolein

A stirred mixture of anhydrous ethanol (235.5g, 5.12 mole) and acrolein (77.3g, 1.38 mole) is treated with 0.7 ml concentrated HCl and NH_4Cl (6.0g, 0.11 mole) and heated to reflux temperature over a 3 hour period. The reaction mixture is heated at reflux temperatures for 18 hours in a flask fitted with a 1 Dean Stark trap. The trap is removed and the reaction mixture is vacuum distilled. The distillate is redistilled, and 42g is added to a stirred mixture of water, 0.8g of concentrated H_2SO_4 , hydroquinone (0.05g, 0.45 mmole) and dibutylamine (1.78g, 0.014 mole). A 37% formaldehyde solution (19.5g, 0.24 mole) is added to the reaction mixture simultaneously at $80^\circ\text{--}85^\circ\text{C}$. The reaction mixture is stirred for 6 hours at $80^\circ\text{--}85^\circ\text{C}$, cooled to room temperature and extracted with hexanes. The extracts are combined and fractionally distilled to yield the title product, identified by NMR analysis.

EXAMPLE 7

Preparation of 2-methoxymethylacrolein from 1,1,3-trimethoxypropane

To a mixture of 96% sulfuric acid (3.0g, 0.024 mole), dibutylamine (6.7g, 0.052 mole) and hydroquinone (1.6g, 0.013 mole) in water at 85°C , is added a mixture of 1,1,3-trimethoxypropane (120.6g, 0.90 mole) and 37% formaldehyde solution (84g, 1.1 mole formaldehyde) over a 1.25 hour period. The reaction mixture is heated at reflux temperature for 5 hours, cooled to room temperature and extracted with methylene chloride. The extracts are combined and fractionally distilled to give the title product, identified by NMR analysis.

EXAMPLE 8

Preparation of 2-methoxymethylacrolein from β -methoxymethylpropionaldehyde

A mixture of β -methoxymethylpropionaldehyde (44g, 0.43 mole), 37% formaldehyde (40.5g, 0.52 mole), dibutylamine (6.8g, 0.053 mole), 96% sulfuric acid (3.0g, 0.029 mole) and hydroquinone (0.6g, 0.055 mole) in water is heated at reflux temperature for 2 hours, cooled to room temperature and extracted with methylene chloride. The extracts are combined and fractionally distilled to give the title product, 20.5g (48% yield), identified by NMR analysis.


EXAMPLE 9

Preparation of 2-alkoxymethylacrolein from β -alkoxymethylpropionaldehyde

Using essentially the same procedure described in Example 4 and substituting the appropriate β -alkoxymethylpropionaldehyde, the following compounds are obtained: 2-butoxymethylacrolein, 1.4g (28% yield), identified by NMR analysis and 2-isopropoxymethylacrolein, 1.4g (22% yield), identified by NMR analysis.


I claim:

1. A process for the manufacture of a compound of formula I




wherein R is C₁ to C₆ alkyl, which is characterized by reacting ROH wherein R is C₁ to C₆ alkyl with at least one molar equivalent of acrolein in the presence of a mineral acid, a catalytic amount of a trisubstituted amine and a solvent, wherein the solvent is water or a mixture of water and a water-miscible organic solvent, to form an intermediate and reacting said intermediate with at least one molar equivalent of formaldehyde in the presence of a mineral acid, a catalytic amount of a disubstituted amine and a solvent, wherein the solvent is water or a mixture of water and a water-miscible organic solvent, to form the formula I compound.

2. A process for the manufacture of a compound of formula I



wherein R is C₁ to C₆ alkyl, which is characterized by reacting a compound of formula II






wherein R is C_1 to C_6 alkyl and W is CHO or $CH(OR_1)_2$ and R_1 is C_1 - C_4 alkyl, with at least one molar equivalent of formaldehyde in the presence of a mineral acid, a catalytic amount of a disubstituted amine and a solvent, wherein the solvent is water or a mixture of water and a water-miscible organic solvent.

3. The process according to claim 1 or 2, wherein the reaction temperature is about 20°C to 110°C ; the acid is selected from the group consisting of sulfuric acid, phosphoric acid, hydrochloric acid and hydrobromic acid; the trisubstituted amine is selected from the group consisting of tri(C_1 - C_4 -alkyl)amine and tri(C_1 - C_4 alkanol)amine and wherein the disubstituted amine is di(C_1 - C_4 alkyl)amine.

4. The process according to claim 1 or 2 wherein the trisubstituted amine is triethylamine or triethanolamine and wherein the disubstituted amine is dibutylamine.

5. A compound having the structural formula



wherein

R is C_1 - C_6 alkyl;


R_4 is hydrogen or C_1 - C_4 alkyl; and

R_5 , R_6 and R_7 are each independently hydrogen or OR_4 .



6. The compound according to claim 5 selected from the group consisting of 3-(methoxymethyl)-8-quinolinol and 8-methoxy-3-(methoxymethyl)quinoline.

7. A process for the preparation of a compound of formula III




wherein


R is C₁-C₆alkyl;

R₄ is hydrogen or C₁-C₄alkyl; and

R₅, R₆ and R₇ are each independently hydrogen or OR₄, which is characterized by reacting a compound of formula IV



wherein R₄, R₅, R₆ and R₇ are as described above with at least one molar equivalent of a compound of formula I




(I)






wherein R is as described above in the presence of an acid and a solvent and optionally in the presence of a compound of formula V



wherein R_4 , R_5 , R_6 and R_7 are as described above to form the formula III compound.

8. The process according to claim 7 wherein the formula V compound is present; the reaction temperature is about 30°C to 120°C and the acid is selected from the group consisting of sulfuric acid, phosphoric acid and hydrochloric acid.

9. A process for the preparation of a compound of formula III



wherein


R is $\text{C}_1\text{-C}_6$ alkyl;

R_4 is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; and

R_5 , R_6 and R_7 are each independently hydrogen or OR_4 , which is characterized by reacting ROH wherein R is C_1 to C_6 alkyl with at least one molar equivalent of




acrolein in the presence of a mineral acid, a catalytic amount of a trisubstituted amine of the formula $N(R_2)_3$, wherein R_2 is C_1-C_4 alkyl or C_1-C_4 alkanol, and a first solvent wherein the first solvent is water or a mixture of water and a water-miscible organic solvent, to form an intermediate and reacting said intermediate with at least one molar equivalent of formaldehyde in the presence of a mineral acid, a catalytic amount of a disubstituted amine of the formula $NH(R_3)_2$, wherein R_3 is C_1-C_6 alkyl, and a second solvent wherein the second solvent is water or a mixture of water and a water-miscible organic solvent to form a compound of formula I



(I)

wherein R is as described above, reacting at least one molar equivalent of the compound of formula I with a compound of formula IV




(IV)

wherein R_4 , R_5 , R_6 , and R_7 are as described above, in the presence of an acid and a third solvent and optionally in the presence of a compound of formula V



This Sheet Has Been Replaced



wherein R_4 , R_5 , R_6 and R_7 are as described above to form the formula III compound.

10. The process according to claim 9 wherein the formula V compound is present; the reaction temperature is about $30^{\circ}C$ to $120^{\circ}C$; the acid is selected from the group consisting of sulfuric acid, phosphoric acid and hydrochloric acid; and wherein the third solvent is selected from the group consisting of water and a C_1-C_6 alcohol.



PROCESS FOR THE MANUFACTURE OF 2-ALKOXYMETHYLACROLEIN


It is an object of this invention to provide a convenient and effective method for the manufacture of 2-alkoxymethylacrolein compounds useful in the preparation of pyridine and quinoline herbicide intermediates. There is also provided a process for the manufacture of 3-alkoxymethylquinolines from 2-alkoxymethylacrolein.

Although there are literature procedures which have been used successfully to prepare certain substituted acrolein compounds, there is, as yet, no known method reported for the manufacture of 2-alkoxy methylacrolein.

Alpha, beta-unsaturated aldehydes such as 2-alkoxymethylacrolein are useful in the preparation of substituted pyridine-2,3-dicarboxylates, which are key intermediates in the manufacture of a new class of imidazolinone herbicides. The use of α,β -unsaturated aldehydes in the preparation of pyridine-2,3-dicarboxylates is described in U.S. Pat. Nos. 4,723,011, 4,948,896 and 5,008,392.

The thus-obtained pyridinedicarboxylate compounds may be converted to the corresponding herbicidal 2-(imidazolin-2-yl)nicotinic acids, esters and salts described in U.S. Patent No. 4,798,619.

The present invention relates to a process for the manufacture of compounds of formula I



wherein R is C₁ to C₆ alkyl.

The formula I alkoxymethylacrolein compound is prepared by reacting an alcohol, ROH wherein R is as described for formula I, with at least one molar equivalent of acrolein in the presence of an acid, a catalytic amount of a trisubstituted amine and a

This is the replacement sheet
Date: - 6 JAN 1999

REPLACEMENT SHEETS		SG
☒	Description	: paged 1, 1, 3 to paged 1, 1, 3
☒	Claim(s)	: paged 1, 1, 15 to paged 1, 1, 15
☐	Drawing(s)	: paged to paged
☐	Abstract	: paged to paged
DATE OF RECEIPT 6/1/99		

solvent to form an intermediate and reacting said intermediate with at least one molar equivalent of formaldehyde in the presence of an acid, a catalytic amount of disubstituted amine and a solvent to obtain the desired formula I compound.


The invention further relates to a process for the manufacture of 2-alkoxymethylacrolein compounds

This is the replacement sheet

Date: - 6 JAN 1999

3-Alkoxymethylquinolines may be prepared by reacting a substituted aniline of formula IV with at least one molar equivalent of a 2-alkoxymethylacrolein of formula I in the presence of an acid and a solvent and optionally in the presence of a substituted nitrobenzene of formula V, preferably at an elevated temperature, to form desired formula III compounds. The above reaction scheme is shown in flow diagram I.

FLOW DIAGRAM I



Acids suitable for use in the process used to prepare formula III compounds are strong organic and mineral acids such as sulfuric acid, phosphoric acid and hydrochloric acid. Solvents suitable for use in the process used to prepare formula III compounds are water and ROH alcohols wherein R is $\text{C}_1\text{-C}_6$ alkyl. Preferred solvents are ROH alcohols wherein R corresponds to the R of the formula I compound with methanol being a most preferred solvent. Reaction temperatures of from 30°C to 120°C , preferably 50°C to

This is the replacement sheet


Date: 6 JAN 1999

wherein R is C_1 to C_6 alkyl and W is CHO or $CH(OR_1)_2$ and R_1 is C_1 - C_4 alkyl, with at least one molar equivalent of formaldehyde in the presence of a mineral acid, a catalytic amount of a disubstituted amine and a solvent, wherein the solvent is water or a mixture of water and a water-miscible organic solvent.

3. The process according to claim 1 or 2, wherein the reaction temperature is 20°C to 110°C ; the acid is selected from the group consisting of sulfuric acid, phosphoric acid, hydrochloric acid and hydrobromic acid; the trisubstituted amine is selected from the group consisting of $\text{tri}(C_1$ - C_4 -alkyl)amine and $\text{tri}(C_1$ - C_4 alkanol)amine and wherein the disubstituted amine is $\text{di}(C_1$ - C_4 alkyl)amine.

4. The process according to claim 1 or 2 wherein the trisubstituted amine is triethylamine or triethanolamine and wherein the disubstituted amine is dibutylamine.

5. A compound having the structural formula



wherein

R is C_1 - C_6 alkyl;

R_4 is hydrogen or C_1 - C_4 alkyl; and

R_5 , R_6 and R_7 are each independently hydrogen or OR_4 .




This is the replacement sheet

- 6 JAN 1999

Date:


wherein R is as described above in the presence of an acid and a solvent and optionally in the presence of a compound of formula V



wherein R_4 , R_5 , R_6 and R_7 are as described above to form the formula III compound.

8. The process according to claim 7 wherein the formula V compound is present; the reaction temperature is $30^{\circ}C$ to $120^{\circ}C$ and the acid is selected from the group consisting of sulfuric acid, phosphoric acid and hydrochloric acid.

9. A process for the preparation of a compound of formula III



wherein

R is C_1 - C_6 alkyl;


R_4 is hydrogen or C_1 - C_4 alkyl; and

R_5 , R_6 and R_7 are each independently hydrogen or OR_4 , which is characterized by reacting ROH wherein R is C_1 to C_6 alkyl with at least one molar equivalent of



This is the replacement sheet

Date: - 6 JAN 1999



wherein R₄, R₅, R₆ and R₇ are as described above to form the formula III compound.

10. The process according to claim 9 wherein the formula V compound is present; the reaction temperature is 30°C to 120°C; the acid is selected from the group consisting of sulfuric acid, phosphoric acid and hydrochloric acid; and wherein the third solvent is selected from the group consisting of water and a C₁-C₆ alcohol.



This is the replacement sheet

Date: - 6 JAN 1999

PROCESS FOR THE MANUFACTURE OF 2-ALKOXYMETHYLACROLEIN**ABSTRACT OF THE INVENTION**

There is provided a process for the manufacture of 2-alkoxymethylacrolein compounds via the reaction of an appropriate alcohol and acrolein in the presence of an acid and a trisubstituted amine to form an intermediate and the subsequent reaction of the intermediate with formaldehyde in the presence of an acid and a disubstituted amine. There is also provided a process for the manufacture of 3-alkoxymethylquinolines from 2-alkoxymethylacrolein.

REPLACED SHEETS		SG
☒	Description : paged 13 to paged 17	
☒	Claim(s) : paged 15, 17, 18 to paged 17	
☐	Drawing(s) : paged to paged	
☐	Abstract : paged to paged	
DATE OF RECEIPT 22 / 2 / 96		