

AUG 08 2001

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

STRITENSEC

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
26 July 2001 (26.07.2001)

PCT

(10) International Publication Number
WO 01/53273 A1(51) International Patent Classification⁷: C07D 239/96,
403/04, 471/04, 471/06, 498/06, 403/14, 519/00, 487/04,
487/10, 413/04, 417/04, 409/04, A61K 31/505, A61P
31/04

(21) International Application Number: PCT/US00/33656

(22) International Filing Date:
12 December 2000 (12.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/178,252 24 January 2000 (24.01.2000) US
60/241,267 18 October 2000 (18.10.2000) USDrive, Ann Arbor, MI 48103 (US). SINGH, Rajeshwar
[CA/CA]; 7927 22nd Avenue, Edmonton, Alberta T6K
1Z2 (CA). STIER, Michael, Andrew [US/US]; 4886
Club Place, Ypsilanti, MI 48197 (US). TRAN, Tuan,
Phong [US/US]; 43517 Argonne Court, Canton, MI 48188
(US). WATSON, Brian, Morgan [US/US]; 3816 Brian
Place, Carmel, IN 46033 (US). YIP, Judy [CA/CA]; #306,
4607-20 Avenue, Edmonton, Alberta T6L 5W7 (CA).(74) Agents: FEDERMAN, Evan, J.; Warner-Lambert Com-
pany, 201 Tabor Road, Morris Plains, NJ 07950 et al. (US).(81) Designated States (*national*): AE, AG, AL, AU, BA, BB,
BG, BR, BZ, CA, CN, CR, CU, CZ, DM, DZ, EE, GD, GE,
HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV,
MA, MG, MK, MN, MX, MZ, NO, NZ, PL, RO, SG, SI,
SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA.(71) Applicant (*for all designated States except US*):
WARNER-LAMBERT COMPANY [US/US]; 201
Tabor Road, Morris Plains, NJ 07950 (US).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): BIRD, Paul
[CA/CA]; 11407-73 Avenue, Edmonton, Alberta T6G
0E2 (CA). ELLSWORTH, Edmund, Lee [US/US]; 6899
Kestrel Court, Brighton, MI 48116 (US). NGUYEN, Dai,
Quoc [CA/CA]; 4928-13 Avenue, Edmonton, Alberta T6L
6J9 (CA). SANCHEZ, Joseph, Peter [US/US]; 22372
Antler Drive, Novi, MI 48050 (US). SHOWALTER,
Howard, Daniel, Hollis [US/US]; 3578 Lamplighter(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 3-AMINOQUINAZOLIN-2,4-DIONE ANTIBACTERIAL AGENTS



(I)

(57) Abstract: Antibacterial 3-aminoquinazolin-2,4-diones have formula (I)
wherein: R₁ and R₃ include alkyl, alkenyl, alkynyl, cycloalkyl, aryl, hetero-
cyclic, and heteroaryl; R₅, R₆, and R₈ include H, alkyl, alkoxy, halo, NO₂, CN,
NH₂, alkyl and dialkylamino; R₇ includes hydrogen, alkyl, cycloalkyl, hetero-
cyclic, fused heterocyclic, aryl and fused aryl; J and K are C or N; and pharma-
ceutically acceptable salts thereof.

WO 01/53273 A1

-1-

3-AMINOQUINAZOLIN-2,4-DIONE ANTIBACTERIAL AGENTS


Field of the Invention

This invention relates to 3-aminoquinazolin-2,4-diones that have potent in vitro and in vivo antibacterial activity. In addition, this invention relates to a method of inhibiting both wild-type and quinolone resistant mutants of DNA gyrase. This invention further relates to a method of treating patients having antibiotic resistance and in need of treatment for bacterial infections.

BACKGROUND OF THE INVENTION

Antibiotic resistance is a worldwide problem with catastrophic potential. The American Society of Microbiology Task Force recently issued a report defining the resistance problem and calling for new antibacterial agents with novel structures or mechanisms to offer alternatives to existing therapeutic choices (*Southern Med. J.*, 1995;88:797).


The quinolones are a class of antibacterials that are widely used throughout the world. The quinolones are potent inhibitors of gram positive and gram negative bacteria, and may be administered orally or intravenously. The quinolones are routinely used even though they have side effects (*J. Antimicrob. Chemother.*, 1994;33:685), and significant resistance has been frequently noted (Gootz, *Medicinal Research*, 1996;Rev. 16:433). One of the most widely used quinolones is ciprofloxacin, which has the following chemical structure:



The quinolones have a distinct structure activity relationship (SAR) which is defined by several thousands of analogs prepared over the last 30 years


-2-

(S. Mitsuhashi, ed. *Progress in Drug Research*, 1992;38:11-147). In the quinolone SAR (referring to the formula below representing substituted quinolones), it is well-established that the N₁ group, in combination with the C₃-carboxyl and the C₄-carbonyl, are essential for activity, and that any substituents at C₂ detract from activity (*J. Antimicrob. Chemother.*, 1994;33:685 and Gootz, supra., 1996). It is also well-established that R₆ is ideally fluorine, and that R₇ is a nitrogen containing heterocycle. R₁ is ideally a small alkyl, a cycloalkyl, or a phenyl group.



The quinolones inhibit bacterial growth by inhibition of DNA gyrase and topoisomerase IV (Gootz, supra., 1996). The gyrase interaction appears to rely on the N₁/C₄-carbonyl/C₃-carboxyl relationship.


Attempts to design novel quinolone mimics have focused on the N₁/C₄-carbonyl/C₃-carboxyl relationship. Certain tricyclic isothiazole analogs were designed to keep an all-planar relationship, and to have the NH of the isothiazole ring be as acidic as the quinolone C₃-carboxyl group (Chu, *Drugs Exptl. Clin. Res.*, 1990;16:215).



While maintaining excellent antibacterial activity, these compounds also show antitumor and mammalian topoisomerase activity (*Drugs of the Future*, 1992;17:1101). These activities are undesired in an antibacterial agent.

-3-

Several publications (US Patent No. 5,283,248; *J. Med. Chem.*, 1992;35:1358; *Antimicrob. Agents Chemother.*, 1995;39:163) disclose tricyclic compounds (such as the structure below) having antibacterial activity and DNA gyrase inhibiting activity.




5

For such compounds, the relationship of the N₁ to the C₄-carbonyl is skewed. Compounds of this type are ineffective against bacteria that are quinolone resistant.

Other tricyclic analogs of the following structure are disclosed as quinolone mimics (JP 4,091,090 3/92; Interscience Conference on Antimicrobial Agents and Chemotherapy 1991, Abstract 1494).

10





While the ideal quinolone N₁-C₄-carbonyl relationship is maintained, the C₂ region, where substitution is undesirable, is part of a fused-ring system.

15

These compounds are not active against quinolone resistant bacteria.


WO 96/04288 describes a series of benzoheterocycles of the formulas



-4-


where X, Y, and Z are hydrogen bond acceptor and donator groups. Among the compounds depicted are N-hydroxy-quinazoline-2,4-diones, where R₁-R₄ can represent hydroxy, amino, nitro, a variety of alkyls, esters, and amides. In all cases, the substituent on N₁ is hydrogen. None of the substituents R₁-R₄ are nitrogen containing heterocycles. No antibacterial activity is disclosed for these compounds.

United States Patent No. 5,155,110 describes N₁-aryl-N-hydroxy-quinazoline-2,4-diones of the formula



where R may be halo, cyano, hydroxy, alkoxy, and substituted amino. Amino heterocycles are not included in R, and no antibacterial activity is reported for such compounds.


Compounds taught in International Publication No. WO 95/19346 as AMPA, NMDA, and kinase receptor antagonists have the formula



where R substituents are defined as alkyl or phenyl, and R₁ and R₂ are defined as H, alkyl, alkoxy, alkenyl, halogen, NO₂, (un)substituted NH₂, CN, etc. Again, no antibacterial activity is disclosed for these compounds.

Kornet and coworkers (Kornet M.J. et al., *J. Heterocycl. Chem.*, 1984;21:1533-1535) disclosed anticonvulsant compounds having the formula


-5-



where R is defined as H, Me, or Cl; R₁ and R₂ as H or Me; and R₃ as Me or t-butyl. Alternatively, R₂ and R₃, together with the nitrogen atom to which they are attached, can form a heterocyclic ring such as piperidinyl, morpholinyl, etc.


5 These compounds are not reported to have antibacterial activity.

A number of publications teach compounds having diuretic and sedative activities and with the generic structure as follows (ES 80/489675, JP 80/33323, BE 80/199792, FR 78/33438, JP 70/97932; Baronnet R., Callendret R., Blanchard L., Foussard-Blanpin O., Bretaudeau J., *Eur. J. Med.-Chim. Ther.*,
 10 1983;18:241-247).



These are compounds in which NR₁R₂ can form a heterocycle, or R₁ and R₂ independently can represent H or a small alkyl. R₃ and R₄ can represent H, halogen, sulphonamide, alkyl, or alkoxy. No antibacterial activity is reported for
 15 these compounds.

International Publication No. WO 99/21840 describes 3-hydroxy-quinazoline-2,4-diones as antibacterial agents.



-6-


Compounds of this type are quinolone mimics where the 3-position exocyclic carboxylic acid of the quinolones is replaced by an oxo at the 2-position and a hydroxy at the N-3-position. Compounds of this type are designed to keep an all-planar relationship and to have the N-OH remain acidic, as in the quinolone CO₂H (Chu, *Drugs Exptl. Clin. Res.*, 1990;16:215). These compounds inhibit DNA gyrase and topoisomerase IV, and demonstrate antibacterial activity in vitro. However, such compounds are not effective against quinolone resistant bacteria.

The need continues to find new antibacterial agents that have the potency of the quinolones, but which are active against quinolone resistant bacterial strains. The present invention provides such compounds, which are characterized as 3-aminoquinazolin-2,4-diones. An object of this invention is to provide these new compounds, and a method for treating bacterial infections in mammals by administering these compounds.

SUMMARY OF THE INVENTION

Disclosed herein is a 3-aminoquinazolin-2, 4-diones having antibacterial activity. The compounds may inhibit DNA gyrase, but are not cross-resistant with quinolone mutants that are no longer sensitive to agents such as ciprofloxacin.

In a first aspect of the present invention, there is provided a compound of Formula I



or a pharmaceutically acceptable salt thereof wherein:

R₁ and R₃ independently are H,

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,

C₂-C₁₂ alkynyl and substituted alkynyl,

C₃-C₁₂ cycloalkyl and substituted cycloalkyl,

-7-

aryl and substituted aryl,
heterocyclic and substituted heterocyclic,
or heteroaryl and substituted heteroaryl;

R₅, R₆, and R₈ independently are H,

5 (O)_nC₁-C₁₂ alkyl and substituted alkyl;
(O)_nC₂-C₁₂ alkenyl and substituted alkenyl;
(O)_nC₂-C₁₂ alkynyl and substituted alkynyl;

halo,

NO₂,

10 CN,

NR'R'' wherein n is 0 or 1;

R' and R'' independently are H,

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,

15 C₂-C₁₂ alkynyl and substituted alkynyl,

CO-C₁-C₁₂ alkyl and substituted alkyl,


or R' and R'' taken together with the nitrogen to which they are attached
form a 3- to 7-membered ring containing from 1 to 3 heteroatoms
selected from N, O, and S, said ring being unsubstituted or
20 substituted with up to 4 substituent groups;

R₁ and R₈ can be taken together with the atoms to which they are attached to
form a cyclic ring having from 1 to 3 heteroatoms selected from N, O, and
S, and optionally substituted by R' and R'';


R₇ is selected from

25

7a



7b




J and K independently are C or N, provided that when J or K is N, R_6 or R_8 is absent at that position.

-8-

J and K independently are C or N, provided that when J or K is N, R₆ or R₈ is absent at that position.

R₇ is referred to herein as the "side chain" of the 3-aminoquinazolin-2,4-dione nucleus. The R₇ side chain can be any group known in the quinolone art as a suitable substituent at the 7-position of the quinolone ring.


In a second aspect of the present invention, there is provided a compound of Formula II



II

wherein R₁, R₃, R₆, R₇, and R₈ are as defined in the first aspect.

In a third aspect of the present invention, there is provided a compound of Formula III




III

wherein R₃, R₆, R₇, and R' are as defined in the first aspect.

In a fourth aspect of the present invention, there is provided a compound of Formula IV


-9-



IV

wherein R_1 , R_3 , R_6 , and R_7 , are as defined in the first aspect.

In a fifth aspect of the present invention, there is provided a compound of Formula V



V

or a pharmaceutically acceptable salt thereof, wherein

R_1 and R_3 independently are H,

C_1 - C_{12} alkyl and substituted alkyl,

C_2 - C_{12} alkenyl and substituted alkenyl,

C_2 - C_{12} alkynyl and substituted alkynyl,

C_3 - C_{12} cycloalkyl and substituted cycloalkyl,

aryl and substituted aryl,

heterocyclic and substituted heterocyclic,

or heteroaryl and substituted heteroaryl;

R_6 and R_8 independently are H,

$(O)_n C_1$ - C_{12} alkyl and substituted alkyl,

$(O)_n C_2$ - C_{12} alkenyl and substituted alkenyl,

$(O)_n C_2$ - C_{12} alkynyl and substituted alkynyl,

wherein n is 0 or 1,

halo,

NO_2 ,

CN,

$NR'R''$;

R' and R''

9a

taken together with the nitrogen to which they are attached form a 3- to 7-membered ring containing from 1 to 3 heteroatoms selected from N, O, and S, said ring being unsubstituted or substituted with up to 4 substituent groups.


5

An especially preferred

embodiment are compounds of Formula V wherein R' and R'' are taken together with the nitrogen to which they are attached to complete a ring which is optionally substituted with groups such as alkyl and substituted alkyl, and amino, alkylamino, dialkylamino, and aryl amino.

10


In a sixth aspect of the present invention, there is provided a compound of Formula VI



VI

wherein R₁, R₃, R₆, R', and R'' are as defined in the fifth aspect.

In a seventh aspect of the present invention, there is provided a compound of Formula VII



VII

15

wherein R₁, R₃, R₇, and R₈ are as defined in the first aspect.

In a ninth aspect of the present invention, there is provided a compound selected from:

3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

5 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-piperazin-1-yl-1*H*-quinazoline-2,4-dione;

3-Amino-7-[3-(aminomethyl)-3-methylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(6-amino-3-aza-bicyclo[3.1.0]hex-3-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazolin-2,4-dione;

10 3-Amino-7-((*S*)-3-N-methylaminopyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-((*S*)-3-aminopyrrolidin-1-yl)-8-chloro-1-(2,4-difluoroanilino)-6-fluoro-1*H*-quinazoline-2,4-dione;

15 3-Amino-7-[(*S*)-3-aminopyrrolidin-1-yl]-1-cyclopropylamino-6,8-difluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(*S*)-3-aminopyrrolidin-1-yl]-1-cyclopropylamino-5,8-difluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-((*S*)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-5,6,8-trifluoro-1*H*-quinazoline-2,4-dione;

20 3-Amino-7-(3-aminopyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

7-((*S*)-3-Aminopyrrolidin-1-yl)-1-cyclopropyl-3,5-diamino-6,8-difluoro-1*H*-quinazoline-2,4-dione;

25 3-Amino-7-(3-hydroxymethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminoethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-((*S*)-7-amino-5-azaspiro[2.4]hept-5-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

30 3-Amino-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

-11-

3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(pyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

5 3-Amino-7-(3-aminopyrrolidin-1-yl)-8-chloro-6-fluoro-1-isopropyl-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-*sec*-butyl-8-chloro-6-fluoro-1*H*-quinazoline-2,4-dione;

10 3-Amino-7-[3-aminopyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1*H*-quinazoline-2,4-dione;

3-Amino-7-[3-aminopyrrolidin-1-yl]-8-chloro-1-cyclobutylamino-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-fluoro-1*H*-quinazoline-2,4-dione;

15 3-Amino-7-(3-aminopyrrolidin-1-yl)-6-chloro-1-cyclopropyl-8-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-((*S*)-3-aminopyrrolidin-1-yl)-1-cyclopropylmethyl-8-fluoro-1*H*-quinazoline-2,4-dione;

20 3-Amino-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropylmethyl-8-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-((*S*)-3-aminopyrrolidin-1-yl)-1-ethyl-8-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-1-ethyl-6-fluoro-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione;

25 3-Amino-7-((*S*)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

30 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-fluoro-6-methoxy-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-ethoxy-6-fluoro-1H-quinazoline-2,4-dione;

5 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-8-carbonitrile;

3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(3-[1,2,3-triazol]-1-yl)pyrrolidin-1-yl)-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(S)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(S)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

5-Amino-9-((S)-3-aminopyrrolidin-1-yl)-8-fluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione;

20 5-Amino-9-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-8-fluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione;

2-Amino-8-((S)-3-aminopyrrolidin-1-yl)-9-fluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione,;

2-Amino-8-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-9-fluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione;

25 3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

3-Amino-7-(6-amino-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione,;

30 3-Amino-7-(3-aminomethyl-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione,;

3-Amino-1-cyclopropyl-6-fluoro-7-(octahydropyrrolo[3,4-c]pyridin-2-yl)-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

-13-

3-Amino-1-cyclopropyl-6-fluoro-7-(octahydropyrrolo[3,4-b]pyridin-2-yl)-
1H-pyrido[2,3-d]pyrimidine-2,4-dione;

3-Amino-7-[(R)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropyl-
6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

5 3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-1H-
pyrido[2,3-d]pyrimidine-2,4-dione;

trans-3-Amino-7-(3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl)-1-
cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

10 3-Amino-1-cyclopropyl-6-fluoro-7-[(R)-3-((R)-1-methylamino-
ethyl)pyrrolidin-1-yl]-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-
fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-aminopropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-
methyl-1H-quinazoline-2,4-dione;

15 3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-[(R)-3-((S)-1-
methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione;

3-Amino-7-[7-(1-aminoethyl)-5-azaspiro[2.4]hept-5-yl]-1-cyclopropyl-6-
fluoro-8-methyl-1H-quinazoline-2,4-dione;

20 1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-
tetrahydroquinazolin-7-yl)piperidine-3-carboxylic acid amide;

3-Amino-[7-trans-3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl]-8-
chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-{3-[(2,2,2-trifluoro-
ethylamino)methyl]pyrrolidin-1-yl}-1H-quinazoline-2,4-dione;

25 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(5-methylhexahydro-
pyrrolo[3,4-c]pyrrol-2-yl)-1H-quinazoline-2,4-dione;

3-Amino-8-chloro-1-cyclopropyl-7-(2,7-diazaspiro[4.4]non-2-yl)-6-fluoro-
1H-quinazoline-2,4-dione;

30 3-Amino-7-(3-aminomethyl-3-benzylpyrrolidin-1-yl)-8-chloro-1-
cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-8-chloro-1-
cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

-14-

3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(3-hydroxyimino-pyrrolidin-1-yl)-1*H*-quinazoline-2,4-dione;

3-Amino-7-[trans-3-amino-4-trifluoromethylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

5 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-[(*R*)-3-(*S*)-1-methylaminoethyl]pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione;

3-Amino-7-(trans-3-amino-4-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

10 3-Amino-7-[trans-3-amino-4-(4-hydroxyphenyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

N-[1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-ylmethyl]methanesulfonamide;

3-Amino-7-(3-aminomethylpiperidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

15 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-[3-(isopropylamino-methyl)pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-6,8-dichloro-1-cyclopropyl-1*H*-quinazoline-2,4-dione;

20 *N*-[1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-ylmethyl]methanesulfonamide;

3-Amino-7-[(*R*)-3-(*S*)-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

25 3-Amino-7-[3-(1-aminoethyl)-3-methoxypyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

3-Amino-7-[3-(1-aminoethyl)-3-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1*H*-quinazoline-2,4-dione;

30 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-5-methyl-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(*R*)-3-(*S*)-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-5-methyl-1*H*-quinazoline-2,4-dione;

-15-

3-Amino-7-(3-aminomethyl-3-methoxymethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyl-3-fluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

5 3-Amino-7-(trans-3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[3-(1-aminoethyl)piperidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-[3-(aminoethyl)piperidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[4-(1-aminoethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[4-(1-aminoethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-(3-amino-3-phenylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[3-(1-aminoethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-(3-aminomethyl-3-phenylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(7-aminomethyl-5-azaspiro[2,4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(7-aminomethyl-5-azaspiro[2,4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-(3-aminomethyl-3-hydroxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-(3-amino-4-methoxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-amino-4-methoxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-16-

3-Amino-7-(3-amino-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyl-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-ethyl-6-fluoro-1H-quinazoline-2,4-dione;

1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid trifluoroacetate;

10 1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid trifluoroacetate;

3-Amino-7-(1-aminomethyl-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(1-aminomethyl-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-(octahydropyrrolo [3,4-b]pyridin-6-yl)-1H-quinazoline-2,4-dione;

3-Amino-7-(trans-3-aminomethyl-4-methylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(trans-3-aminomethyl-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-(trans-3-aminomethyl-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(trans-3-amino-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-(3-aminomethylmorpholin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-17-

3-Amino-7-[3-(1-aminoethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-amino-3-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

5 3-Amino-7-(3-amino-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione;

10 3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)-pyrrolidin-1-yl]-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)-pyrrolidin-1-yl]-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-1-cyclopropyl-6-fluoro-5-methyl-7-[(R)-3-((S)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione;

(S)-1-[(R)-1-(3-Amino-1-cyclopropyl-7-[3-(1-ethylaminoethyl)pyrrolidin-1-yl]-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-1-cyclopropyl-7-[(R)-3-((S)-1-ethylaminoethyl)pyrrolidin-1-yl]-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(6-amino-1-methyl-3-azabicyclo[3.2.0]hept-3-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(4-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-(3-aminomethylazetidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

cis-3-Amino-7-(3-aminomethyl-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 trans-3-Amino-7-(3-aminomethyl-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(1-amino-5-azaspiro[2.4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-18-

3-Amino-7-(3-aminomethyloctahydroisoindol-2-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyloctahydroisoindol-2-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(R)-3-((S)-1-aminoethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(6-amino-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-7-[3-(isopropylaminomethyl)pyrrolidin-1-yl]-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-7-[(4aS,7aS)-octahydropyrrolo[3,4-b]pyridin-6-yl]-1H-quinazoline-2,4-dione;

15 3-Amino-7-(3-amino-4-fluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyl-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-7-(4-aminopiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminopiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-7-[(R)-3-((S)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione;

25 3-Amino-7-(3-aminoazetidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(3R,4S)-3-((S)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-(R)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4S)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(3R,4R)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(R)-4-(R)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((S)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-(S)-1-amino-2-methoxyethyl]-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(R)-3-(R)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(R)-3-((S)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(R)-4-((R)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((S)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

-21-

3-Amino-7-[(R)-4-((S)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(R)-3-((S)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(R)-3-((R)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(R)-4-((R)-1-amino-2-ethoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-ethoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

-22-

3-Amino-7-[(R)-3-((S)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

{(R)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(R)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(S)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

- {(S)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-
 1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;
 {(R)-2-Amino-2-[(3R,4S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-
 2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;
 5 {(R)-2-Amino-2-[(3R,4S)-1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-
 1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;
 {(S)-2-Amino-2-[(3R,4R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-
 2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;
 {(S)-2-Amino-2-[(3R,4R)-1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-
 10 1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;
 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-
 1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]ethyl}urea;
 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-
 tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]ethyl}urea;
 15 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-
 1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}urea;
 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-
 tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}urea;
 3-Amino-7-[(3R,4S)-3-((S)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-
 20 cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
 3-Amino-7-[(3R,4S)-3-((S)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-
 cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
 3-Amino-7-[(3R,4R)-3-(R)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-
 cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
 25 3-Amino-7-[(3R,4R)-3-((R)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-
 cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
 3-Amino-7-[(3R,4S)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-
 1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
 3-Amino-7-[(3R,4S)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-
 30 1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
 3-Amino-7-[(3R,4R)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-
 1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

- 3-Amino-7-[(3R,4R)-3-(1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
- 5 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 10 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 15 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(R)-4-((R)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
- 20 3-Amino-7-[(R)-4-((R)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(R)-4-((S)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
- 25 3-Amino-7-[(R)-4-((S)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(R)-3-((R)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;
- 3-Amino-7-[(R)-3-((R)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;
- 30 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-25-

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(R)-3-((S)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(R)-4-((R)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((S)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(R)-4-((S)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(R)-3-((S)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-ethoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-ethoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(R)-3-((S)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

-27-

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

20 {(R)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(R)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

25 {(S)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(S)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(R)-2-Amino-2-[(3R,4S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

30 {(R)-2-Amino-2-[(3R,4S)-1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

-28-

- {(S)-2-Amino-2-[(3R,4R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;
- {(S)-2-Amino-2-[(3R,4R)-1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;
- 5 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]ethyl}urea;
- {2-Amino-2-[(1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]ethyl}urea;
- {2-Amino-2-[(1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}urea;
- 10 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}urea;
- 3-Amino-7-[3-(1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 15 3-Amino-7-[3-(1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 20 3-Amino-7-[4-(1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Amino-7-[3-(1-amino-2-phenoxyethyl)-pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 25 3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 3-Amino-7-[4-(1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;
- 30 3-Amino-7-[3-(1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

5 3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4,4-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

10 3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[4,3-d]pyrimidin-7-yl)pyrrolidin-3-yl]ethylurea;

15 {2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[4,3-d]pyrimidin-7-yl)-4-methylpyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[4,3-d]pyrimidin-7-yl)-4-fluoropyrrolidin-3-yl]ethylurea;

20 {2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[4,3-d]pyrimidin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethylurea;

3-Amino-7-[3-(1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

25 3-Amino-7-[3-(1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

30 3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

-30-

3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

5 3-Amino-7-[4-(1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[4-(1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

10 3-Amino-7-[3-(1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

15 3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

20 3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[4-(1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[4-(1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

25 3-Amino-7-[3-(1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

30 3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

5 3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4,4-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4,4-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

10 3-Amino-7-[3-(1-amino-2-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

15 3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

20 3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)pyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)pyrrolidin-3-yl]ethylurea;

25 {2-Amino-2-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4-methylpyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4-methylpyrrolidin-3-yl]ethylurea;

30 {2-Amino-2-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4-fluoropyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4-fluoropyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethylurea;

5 {2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethylurea;

3-Amino-7-(3-aminomethylphenyl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylphenyl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

10 3-Amino-7-(3-aminomethylphenyl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylphenyl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-(3-aminomethylphenyl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylphenyl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(2-aminomethylthiazol-4-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

20 3-Amino-7-(2-aminomethylthiazol-4-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(2-aminomethylthiazol-4-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

25 3-Amino-7-(2-aminomethylthiazol-4-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(2-aminomethylthiazol-4-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(2-aminomethylthiazol-4-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

5 3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(5-aminomethylthiophen-3-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(5-aminomethylthiophen-3-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

15 3-Amino-7-(5-aminomethylthiophen-3-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(5-aminomethylthiophen-3-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-(5-aminomethylthiophen-3-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(5-aminomethylthiophen-3-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

25 3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

30 3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione

-34-

3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

5 3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

10 3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

20 3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-(1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-5,8-dimethyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-(3-(1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-5,8-dimethyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-(1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-5-methyl-1H-quinazoline-2,4-dione;

-35-

3-Amino-7-(3-(1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-8-methoxy-5-methyl-1H-quinazoline-2,4-dione;

3,8-Diamino-7-[3-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-5-methyl-1H-quinazoline-2,4-dione; and

3,8-Diamino-7-[3-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-5-methyl-1H-quinazoline-2,4-dione.

In a tenth aspect of the present invention, there is provided a pharmaceutical composition comprising a compound as defined in the first aspect admixed with a carrier, diluent, or excipient.

In an eleventh aspect of the present invention, there is provided a use of an antibacterial effective amount of a compound as defined in the first aspect in the manufacture of a medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof.

In a twelfth aspect of the present invention, there is provided a use of an effective amount of a compound as defined in the first aspect in the manufacture of a medicament for inhibiting a wild-type mutant of DNA gyrase in a mammal comprising administering to the mammal in need thereof.

In a thirteenth aspect of the present invention, there is provided a use of an effective amount of a compound as defined in the first aspect in the manufacture of a medicament for inhibiting a quinolone resistant mutant of DNA gyrase in a mammal comprising administering to the mammal in need thereof.

35a

A further embodiment may be a pharmaceutical composition comprising a compound of Formula I together with an excipient, carrier, or diluent. Preferred compositions will have a compound of Formulas II, III, IV, V, VI, or VII mixed with an excipient, carrier, or diluent.

Still another embodiment may be a method of treating bacterial infections in mammals comprising administering an antibacterial effective amount of a compound of Formula I. Preferred methods comprise administering a compound of Formulas II, III, IV, V, VI, or VII.

Another embodiment may relate to a method of preventing or eliminating bacterial growth on surfaces comprising applying to the surface an antibacterially effective amount of a compound of Formulas I-VII.

DETAILED DESCRIPTION

All references cited herein, including patents, are incorporated herein by reference.

The terms used in this specification and the appended claims have the following meanings. The term "halo" means chloro, bromo, fluoro, and iodo.

The term "alkyl" means a straight or branched hydrocarbon radical having from 1 to 12 carbon atoms and includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, 2-isopropylheptyl, 3-n-butyloctyl, n-nonyl, n-decyl, undecyl, and dodecyl.

Preferred alkyl groups are C₁-C₆ alkyl such as methyl, isopropyl, neopentyl, and 1-methylpentyl.

-36-

The term "C₂-C₁₂ alkenyl" means a straight or branched hydrocarbon radical having from 1 to 3 double bonds. Examples include ethenyl, 2-propen-1-yl, 1,3-butadien-1-yl, 3-hexen-1-yl, 5-octen-2-yl, 2-isopropyl-3,5-octadien-1-yl, cis-3-hexen-1-yl, and trans-2-hepten-1-yl. Preferred alkenyl groups include C₂-C₆ alkenyls such as ethenyl, 2-propen-1-yl, 2-buten-1-yl, and 3-penten-1-yl.

The term "C₂-C₁₂ alkynyl" means a straight or branched hydrocarbon radical having from 1 to 3 triple-bonds. Examples include ethynyl, propynyl, 3-butyn-1-yl, 4-hexyn-1-yl, 5-octyn-3-yl, 4,6-octadiyn-1-yl, 4-decyn-1-yl, and 1,1-dimethyl-5-decyn-1-yl. Preferred alkynyl groups are C₂-C₆ alkynyls such as ethynyl, propynyl, 3-butyn-1-yl, and 5-hexyn-1-yl.

The alkyl, alkenyl, and alkynyl groups can be substituted with 1 to 3 groups selected from halo, hydroxy, cyano, C₁-C₆ alkoxy, nitro, amino, C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, carboxy, C₁-C₆ alkoxycarbonyl, aminocarbonyl, halo-dihalo and trihalomethyl, thiol, alkylsulfanyl, alkylsulfinyl, and aminosulfonyl. Examples of substituted alkyl groups include fluoromethyl, tribromomethyl, hydroxymethyl, 3-methoxypropyl, 3-carboxypentyl, 3,5-dibromo-6-aminocarbonyldecyl, and 4-ethylsulfinyloctyl. Examples of substituted alkenyl groups include 2-bromoethenyl, 1-amino-2-propen-1-yl, 3-hydroxypent-2-en-1-yl, 4-methoxycarbonyl-hex-2-en-1-yl, and 2-nitro-3-bromo-4-iodo-oct-5-en-1-yl. Typical substituted alkynyl groups include 2-hydroxyethynyl, 3-dimethylamino-hex-5-yn-1-yl, and 2-cyano-hept-3-yn-1-yl.

The term "cycloalkyl" means a hydrocarbon ring which contains from 3 to 12 carbon atoms, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, decalinyl, norpinanyl, and adamantyl. Where possible, the cycloalkyl group may contain double bonds, for example, 3-cyclohexen-1-yl. The cycloalkyl ring may be unsubstituted or substituted by 1 to 3 substituents selected from alkyl, alkoxy, thioalkoxy, hydroxy, thiol, nitro, halogen, amino, alkyl and dialkylamino, formyl, carboxyl, CN, -NH-CO-R', -CO-NHR', -CO₂R', -COR', aryl, or heteroaryl, wherein alkyl, aryl, and heteroaryl are as defined herein. Examples of substituted cycloalkyl groups include fluorocyclopropyl, 2-iodocyclobutyl, 2,3-dimethylcyclopentyl, 2,2-dimethoxycyclohexyl, and 3-phenylcyclopentyl.

-37-

The term "heterocyclic" means a cyclic or fused bicyclic or polycyclic ring system having from 3 to 12 ring atoms, with from 1 to 5 being heteroatoms selected from N, O, and S. The heterocyclic groups can be substituted with 1 to 3 of the substituents listed above for the cycloalkyl groups. Examples of

5 heterocyclic groups include cyclic ethers (oxiranes) such as ethyleneoxide, tetrahydrofuran, dioxane, and substituted cyclic ethers, wherein the substituents are those described above for the alkyl and cycloalkyl groups. Typical substituted cyclic ethers include propyleneoxide, phenyloxirane (styrene oxide), cis-2-butene-oxide (2,3-dimethyloxirane), 3-chlorotetrahydrofuran, 2,6-dimethyl-1,4-dioxane,

10 and the like. Heterocycles containing nitrogen are groups such as pyrrolidine, piperidine, piperazine, tetrahydrotriazine, tetrahydropyrazole, and substituted groups such as 3-aminopyrrolidine, 4-methylpiperazin-1-yl, and the like. Typical sulfur containing heterocycles include tetrahydrothiophene, dihydro-1,3-dithiol-2-yl, and hexahydrothiepin-4-yl. Other commonly employed heterocycles include

15 dihydro-oxathiol-4-yl, tetrahydro-oxazolyl, tetrahydro-oxadiazolyl, tetrahydro-dioxazolyl, tetrahydro-oxathiazolyl, hexahydrotriazinyl, tetrahydro-oxazinyl, morpholinyl, thiomorpholinyl, tetrahydropyrimidinyl, dioxolinyl, octahydrobenzofuranyl, octahydrobenzimidazolyl, and octahydrobenzothiazolyl. For heterocycles containing sulfur, the oxidized sulfur heterocycles containing SO or SO₂ groups are also included. Examples include the sulfoxide and sulfone forms of tetrahydrothiophene.

20

The term "aryl" means a cyclic or polycyclic aromatic ring having from 5 to 12 carbon atoms, and being unsubstituted or substituted with up to 3 of the substituent groups recited above for alkyl, alkenyl, and alkynyl. Examples of aryl

25 groups include phenyl, 2,6-dichlorophenyl, 3-methoxyphenyl, naphthyl, 4-thionaphthyl, tetralinyl, anthracinyl, phenanthrenyl, benzonaphthenyl, fluorenyl, 2-acetamidofluoren-9-yl, and 4'-bromobiphenyl.

The term "heteroaryl" means an aromatic cyclic or polycyclic ring system having from 1 to 4 heteroatoms selected from N, O, and S. Typical heteroaryl

30 groups include 2- or 3-thienyl, 2- or 3-furanyl, 2- or 3-pyrrolyl, 2-, 4-, or 5-imidazolyl, 3-, 4-, or 5-pyrazolyl, 2-, 4-, or 5-thiazolyl, 3-, 4-, or 5-isothiazolyl, 2-, 4-, or 5-oxazolyl, 3-, 4-, or 5-isoxazolyl, 3- or 5-1,2,4-triazolyl, 4- or

-38-


5-1,2,3-triazolyl, tetrazolyl, 2-, 3-, or 4-pyridinyl, 3-, 4-, or 5-pyridazinyl, 2-pyrazinyl, 2-, 4-, or 5-pyrimidinyl, 2-, 3-, 4-, 5-, 6-, 7-, or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, or 8-isoquinolinyl, 2-, 3-, 4-, 5-, 6-, or 7-indolyl, 2-, 3-, 4-, 5-, 6-, or 7-benzo[b]thienyl, 2-, 4-, 5-, 6-, or 7-benzoxazolyl, 2-, 4-, 5-, 6-, or 7-benzimidazolyl, 2-, 4-, 5-, 6-, or 7-benzothiazolyl. The heteroaryl groups may be unsubstituted or substituted by 1 to 3 substituents selected from those described above for alkyl, alkenyl, and alkynyl, for example, cyanothienyl and formylpyrrolyl.

10 Preferred aromatic fused heterocyclic rings of from 8 to 10 atoms include but are not limited to 2-, 3-, 4-, 5-, 6-, 7-, or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, or 8-isoquinolinyl, 2-, 3-, 4-, 5-, 6-, or 7-indolyl, 2-, 3-, 4-, 5-, 6-, or 7-benzo[b]thienyl, 2-, 4-, 5-, 6-, or 7-benzoxazolyl, 2-, 4-, 5-, 6-, or 7-benzimidazolyl, 2-, 4-, 5-, 6-, or 7-benzothiazolyl.


15 Preferred compounds of the invention have Formula I wherein R₇ is a heterocyclic or heteroaryl group such as those described above. Such heterocyclic and heteroaryl groups will be referred to herein generically as " $\bigcirc\text{N}^{\sim\sim}$ ".

Further examples of typical heterocycles, fused heterocycles, and heteroaryl groups that are preferred as R₇ substituents are listed below in Table A.

Table A



-40-



The foregoing groups are especially preferred as R₇ side chains on the quinazoline ring system. The heterocycles and heteroaryl groups such as those depicted above are known in the quinolone art and are prepared by literature methods such as *J. Med. Chem.*, 1992;35:1764; *J. Med. Chem.*, 1996;39:3070; *Synlett*, 1996:1097; and *J. Med. Chem.*, 1986;29:445. Any of the primary or

-41-

secondary amines shown as substituents on the heteroaryl or heterocyclic groups may be substituted by alkyl, such as methyl, ethyl, isopropyl, and the like.

Some of the compounds of Formula I are capable of forming pharmaceutically acceptable acid addition and/or base salts. All of these forms are within the scope of the present invention and are prepared by art recognized methods. For example, an acid addition salt is prepared by contacting the free base form of the invention compound with a sufficient amount of the desired acid to produce the salt in the conventional manner. Pharmaceutically acceptable acid addition salts of the compounds of Formula I include salts derived from inorganic acids such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydriodic, hydrofluoric, phosphorous, and the like, as well as the salts derived from organic acids, such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc. Such salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, acetate, trifluoroacetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinate suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, phthalate, benzensulfonate, toluenesulfonate, phenylacetate, citrate, lactate, tartrate, methanesulfonate, and the like. Also contemplated are salts of amino acids such as arginate and the like, as well as gluconate and galacturonate. All such salts are readily prepared, for example as described by Berge S.M. et al., "Pharmaceutical Salts," *Journal of Pharmaceutical Science*, 1977;66:1-19.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines, when compounds of Formula I have an acidic group such as carboxy. The base addition salts of acidic invention compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. Typical bases include sodium hydroxide, calcium oxide, diethylamine, and pyridine. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are

N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge S.M., supra., 1977).

Certain of the compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

Certain of the compounds of the present invention possess one or more chiral centers. Each center may exist in the R or S configuration. The present invention includes all diastereomeric, enantiomeric, and epimeric forms, as well as the appropriate mixtures thereof. Additionally, the compounds of the present invention may exist as geometric isomers. The present invention includes all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers, as well as the mixtures thereof.

Structures of representative compounds of the invention are shown below in Table 1.

TABLE 1







TABLE 1 (cont)




-43-




5



6




9




11

5











18



21

TABLE 1 (cont)



5





Table 1 (cont)



The compounds of Formula I can be prepared utilizing standard synthetic methodology well-known to those skilled in the art of organic chemistry. Many of the compounds have reactive functional groups that generally need to be derivatized with a protecting group in order to avoid unwanted side reactions. For example, functional groups such as alcohols, acid groups, and amines generally are protected while a reaction is carried out at a different site in the molecule, and then the protecting group is subsequently removed following the desired chemical transformation. The use of such protecting groups is standard in organic chemistry synthesis, as described, for example, in T.W. Green and P.G. Wuts, *Protective Groups in Organic Synthesis*, 2nd edition, New York City: John Wiley & Sons, 1991. Thus, for example, protecting groups such as the following may be utilized to protect suitable amino, hydroxyl, and other groups of related reactivity:

carboxylic acyl groups, such as formyl, acetyl, trifluoroacetyl; alkoxycarbonyl groups, such as ethoxycarbonyl, *t*-butoxycarbonyl (BOC), β,β,β -trichloroethoxycarbonyl (TCEC), β -iodoethoxycarbonyl; aryloxycarbonyl groups, such as benzyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, phenoxycarbonyl; trialkyl silyl groups, such as trimethylsilyl and *t*-butyldimethylsilyl (TBDMS); and groups such as trityl, tetrahydropyranyl, vinyloxycarbonyl, *o*-nitrophenylsulfonyl, diphenylphosphinyl, *p*-toluenesulfonyl, and benzyl may all be utilized. The protecting group may be removed, after completion of the synthetic reaction of interest, by procedures known to those skilled in the art. For example, a BOC group may be removed by acidolysis, a trityl group by acidolysis or

-46-

hydrogenolysis, TBDMS by treatment with fluoride ions, and TCEC by treatment with zinc.

Illustrations of typical preparations of compounds of the present invention having Formula I are shown in Schemes 1-22. Typical heterocyclic and aromatic side chains defined by R₇ in Formula I are prepared as described in Schemes A1-A8. All of the 3-aminoquinazoline-2,4-diones of the invention may be prepared from appropriately substituted benzoic acid starting materials. Protecting groups (referred to in the schemes as "Pro") may be used when appropriate throughout many of the schemes. Although specifically noted in certain schemes, the appropriate use and choice of protecting groups is well-known by those skilled in the art, and is not limited to the specific illustrations shown below. It should also be understood that such protecting groups not only serve to protect chemically reactive sites, but also to enhance solubility or otherwise change physical properties of the underlying invention compound. A number of general reactions such as oxidations and reductions are not shown in detail in the schemes, but can be carried out by standard methods well-known to those skilled in the art. In general, the starting materials used in the following schemes are obtained from commercial sources, or are readily prepared by standard methods. All cited published articles, patents, books, and the like are incorporated herein by reference.

As shown in Scheme 1, a difluoro substituted benzoic acid is reacted with oxalyl chloride or an equivalent acylating reagent (such as an acid anhydride), and the acid halide or anhydride is reacted with an alcohol (ZOH) to afford the respective ester (Z is C₁-C₆ alkyl such as methyl, ethyl, isopropyl, etc.). The ester is reacted with an amine, for example, a heterocyclic amine, to produce the desired 4-heterocyclic phenyl derivative. Alternatively, carbocycles and aryls may also be introduced at this 4-position using palladium catalyzed couplings of tin or boronate carbocycles and aryls, with starting materials containing a Br, I, or triflate at the 4-position as described by Suzuki A., *Pure Appl. Chem.*, 1994;66(2):213-222 and Stille J.K., *Angew. Chem.* 1986;98(6):504-519.

-47-

Reaction of the 2-fluoro benzoic acid analog with a primary amine R_1NH_2 affords the corresponding 2-amino benzoic acid ester. The ester group is readily hydrolyzed by reaction with an acid such as hydrochloric acid or a base such as sodium hydroxide to give the corresponding polysubstituted 2-amino benzoic acid. The acid is then coupled to an appropriately protected hydrazine to provide the corresponding amide, which in turn is cyclized to generate the quinazoline-2,4-dione by treatment with, for example, phosgene. At this point, the protecting group may be removed to give the free amine A of Formula I, which can represent a final product of the invention, and which can be further derivatized, if desired, for instance by alkylation with R_3Cl .


Alternatively, the quinazoline-2,4-dione may also be metallated at low temperatures with bases such as, for example, NaH, KH, and lithium diisopropylamine, and then alkylated or acylated to provide the corresponding N-protected-mono-substituted (R_3) amine. The protecting group (Pro) of the N-protected-mono-substituted amine is removed by conventional methods such as hydrogenation, treatment with acid or base, or metal catalysis to afford invention compound C.

If R_5 is a leaving group such as F, it is activated towards displacement with a nucleophile $HY-Pro'$ where Y is N or O and Pro' is a protecting group such as trimethylsilyl. Other R_5 groups such as chlorine, bromine, or sulfonyl are also good leaving groups. The displacement generally is carried out in a solvent such as ethanol, DMSO, DMF, THF, and at a temperature of about $0^\circ C$ to $120^\circ C$.

The protecting groups (Pro) may be selectively removed by hydrogenation, acid or base treatment, metal catalysis, or such other standard methods. When Pro is benzyl, the benzyl may be removed with Pd/C and hydrogen. A t-butyl carbamate group may be removed by alcoholic HCl, TFA, or TFA in dichloromethane, ethyl acetate or diethyl ether. Allylic carbamates may be removed by $PhSiH_3$ and Pd catalyst. Solvents such as alcohol, THF, alcohol/THF, alcohol/THF/DMF, diethyl ether, etc. are generally employed in such protecting group cleavage reactions.


-48-

Scheme 1



-49-

Scheme 1 (cont)



In Scheme 2, an *ortho*-aminobenzoic acid is utilized as the starting material, and is alkylated on the amino group. For example, when R_1 is cyclopropyl, the alkylation is carried out according to the method of Gillaspay (*Tetrahedron Letters*, 1995:7399) to provide the cyclopropyl amine. When R_1 is phenyl or substituted phenyl, the respective amine is prepared from the *ortho*-fluorobenzoic acid using a base, such as, lithium diisopropylamide or Li-hexamethyl disilazide, and the appropriate aryl amine ($R_1\text{NH}_2$). When R_1 is any alkyl group, such as *t*-butyl or isopropyl, the R_1 can be introduced by reacting

-50-

the amine and ortho halo benzoic acid with a Cu catalyst such as copper, bronze, or cupric acetate in the presence of a base such as potassium acetate, triethylamine, or pyridine.


5 The resulting R₁-substituted amino benzoic acid is then coupled to an appropriately protected hydrazine to provide the corresponding benzamide using methods described in the literature. The corresponding amide can be further reacted, if R₇ is a leaving group such as fluoro, with various heterocyclic amines (e.g., piperidine or pyrrolidine) to form the desired 4-heterocyclic benzamide derivative. Alternatively, carbocycles and aryls (e.g., cyclobutyl or phenyl) may
10 also be introduced at this 4-position using palladium catalyzed couplings of tin or boronate carbocycles and aryls, if the starting material contains a Br, I, or triflate at the 4-position.

The 4-substituted benzamide derivative is then cyclized to generate the quinazoline-2,4-dione by reaction with carbonyldiimidazole (CDI), phosgene,
15 triphosgene or the like in ethereal solvents such as diethyl ether, chlorinated hydrocarbons such as dichloromethane, or aromatic hydrocarbons such as toluene, in the presence of a base such as triethylamine or NaHCO₃.

Alternatively, the corresponding amide is first cyclized and then the R₇ halo group is displaced by reaction with a carbocyclic amine C_nNH_2 to afford the
20 same product. Deprotection by conventional methods provides the invention compound A.

-51-

Scheme 2



-52-

Scheme 3 illustrates alkylation at the 1-position of a 3-aminoquinazolin-2,4-dione to provide invention compounds wherein R_1 is alkyl. A 2-aminobenzoic acid is reacted with an N,N-diprotected hydrazine such as dibenzyl hydrazine (H_2N-NBn_2) to provide the corresponding N-protected amide. This intermediate


5 can then be reacted with phosgene or phosgene/base in an ethereal solvent, or with a phosgene equivalent such as triphosgene in a chlorinated hydrocarbon such as dichloromethane, to give the quinazoline-2,4-dione. The alkylation of the quinazoline-2,4-dione to provide a 1-alkylated-quinazoline-2,4-dione is accomplished by reaction with an alkyl halide as described by Bouzard, *supra.*,

10 1990. Typically, such reactions are carried out in THF, ether, DMSO, an alkanol, or DMF, and in the presence of a base. Typical alkyl halides (R_1X where X is halo) include ethyl iodide, ethyl bromide, cyclopropyl iodide, n-decyl bromide, and the like. Typical bases include sodium hydride, potassium carbonate, and the like. Conversion of the 1-alkylated-quinazoline-2,4-dione to other invention

15 compounds (and removal of protecting groups such as benzyl Bn) can be carried out according to Scheme 1, for example, to give the corresponding 1-alkyl ($R_1 =$ alkyl) 3-amino (NH_2) compounds such as A.

-53-

Scheme 3



Displacement of leaving groups located at the 7-position of the quinazolin (e.g., $\text{R}_7 = \text{halo}$) as shown in Schemes 1 and 2 is not limited to nitrogen

- 5 heterocycles. Other nucleophiles (Nu) such as CH_3O^- , N_3^- , $\text{R}'\text{R}''\text{NH}$, $\text{R}'\text{-NH}_2$, and $\text{R}'\text{S}^-$ (where R' and R'' are defined above) also displace a leaving group such as F, Cl, or NO_2 at the 7-position as shown in Scheme 4. When the leaving group is a triflate or higher halide, organo tin reagents or organoboronates may be used with palladium catalysts to deliver a carbon nucleophile. The methodology shown
- 10 in Scheme 4 follows that of Stille et al., *Angew. Chem. Int. Ed. Eng.*, 1986;25:508 and is exemplified by Mitchell (*Synthesis*, 1992:803).

Chemical reaction scheme showing the deprotection of a 6-substituted-2,3-dihydroquinolin-4(1H)-one derivative.

The starting material is a 6-substituted-2,3-dihydroquinolin-4(1H)-one derivative. It features a benzene ring with substituents R_5 , R_6 , R_7 , and R_8 . The heterocyclic core is a 2,3-dihydroquinolin-4(1H)-one, with an N-protecting group (Pro) and an N1 substituent (R_1).

The reaction proceeds via a horizontal arrow to an intermediate and then a vertical arrow labeled "Deprotection" to the final product.

The final product is a 6-substituted-2,3-dihydroquinolin-4(1H)-one derivative, where the N-protecting group (Pro) has been removed, replaced by an NH_2 group.

5

10


15

-55-

using a nucleophilic amine such as diethylamine, pyrrolidine, methylpiperazine, and the like to give the corresponding amino substituted analog. This analog is then reacted with R_1NH_2 to displace the second leaving group (e.g., the 2-chloro group). The resulting 2,6-disubstituted-pyridylamide is then reacted with CDI, phosgene, or other phosgene equivalents to form the cyclized quinazoline product. Amination at the 3-position NH is achieved by reaction with any number of aminating reagents such as ammonia and N-alkylamines, for example, as described by Kloetzer (*Sci. Pharm.*, 1984;52:46-50) to give the corresponding invention compound.

10

Scheme 5




Tricyclic compounds (i.e., where R_1 and R_8 in Formula I are taken together with the atoms to which they are attached to form a ring) can be prepared

-56-

according to Schemes 6 and 7. Schemes 6 and 7 differ in the introduction of the R_1 substitutions in Structures B and C wherein $R_1 = R_1'$ and R_1 is as defined above for Formula I and R_1' is alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, and the like. In Scheme 6, the *ortho*-fluoro nitro compound (other leaving groups such as chlorine, bromine, and sulfonyl may also be employed in place of fluoro) is displaced by the α -nucleophile substituted ester (where Q is a nucleophile Nu such as methyl or methoxy). The nitro group is then reduced using, for example, Raney Ni, H_2 over Pd/C, or an active metal in acid such as iron or tin in HCl or acetic acid. The newly formed amine readily cyclizes with the ester (other acid analogs may be employed such as thioesters, amides, and the like). The cyclized product is then reduced with hydride reducing agents such as $LiAlH_4$ and the like to produce the dihydroquinoline derivative, which in turn is reacted with chloral hydrate and then an acid to form the dione ring. The dione ring is subsequently opened using, for example, sodium hydroxide and hydrogen peroxide to give the benzoic acid. The 3-aminoquinazolinedione ring is then prepared using the chemistry described in Schemes 2, 3, or 5 to give the invention compound B, where R_3' is alkyl, alkenyl, cycloalkyl, aryl, and heteroaryl.

-57-


Scheme 6



In a similar series of reactions depicted in Scheme 7, the *ortho*-fluoro nitro compound is reacted with an α -nucleophile substituted ketone, and the resulting product is likewise reduced. In this sequence, the resulting aniline forms a cyclic imine, which is further reduced with H_2 on Pd/C or by chemical hydride reducing agents such as sodium borohydride or sodium cyanoborohydride to give the dihydroquinoline. Such reductive aminations are well-known in the art and are typically performed in THF, alcohol, water alcohol mixtures, or in water DMF mixtures. The remaining steps to produce C follow those of Scheme 6. When the C-7 substituent is a leaving group (e.g., $R_7 = \text{halo}$), compounds B and C may be further reacted with nucleophiles (such as pyrrolidine or piperidine) to give compounds of Formula I as in the previous schemes.

-58-

Scheme 7



In Scheme 8, the target tricyclic compounds as C are prepared in a slightly different manner. In this case, the nucleophile Q (such as hydroxymethyl) is attached to the phenyl ring of the starting aniline, and a leaving group L (such as halo) is attached alpha to the ketone reactant. The nucleophile may be activated with bases such as sodium hydride or potassium hydride, triethylamine or 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU), or sodium, potassium, or cesium carbonate to displace the leaving group L. In this sequence, the resulting aniline forms a cyclic imine, which is further reduced with H₂ on Pd/C or by chemical hydride reducing agents such as sodium borohydride or sodium cyanoborohydride to give a dihydroquinoline. Such reductive aminations are well-known in the art and are typically performed in THF, alcohol, water alcohol mixtures, or in water DMF mixtures. The dihydroquinoline intermediate is reacted with chloral hydrate, and then an acid to form the dione ring. The dione ring is subsequently opened by reaction with a base, for example sodium hydroxide and hydrogen peroxide, to give the benzoic acid. The 3-aminoquinazolin-4(1H)-one ring is then prepared by first forming an ester on the benzoic acid as in Scheme 2, followed by reaction of the

-59-

benzoic acid ester with chlorosulfonylisocyanate or the like at temperatures of 0°C and below, followed by treatment with a base such as triethylamine or diisopropylethylamine. Amination can then be carried out in the same manner as described in Scheme 5 to provide compounds of structure C. When R₇ is a


5 leaving group such as Cl or F, the invention compounds of structures B (in Scheme 6) and D (in Scheme 8) can be prepared by coupling to the R₇ side chain, e.g., various heterocyclic amines C_nNH to produce the desired derivatives.

Alternatively, carbocycles and aryls may also be introduced as R₇ side chains using palladium catalyzed couplings with tin or boronate carbocycles and aryls, for

10 example when R₇ is a Br, I, or triflate.

-60-

Scheme 8



It should be noted from Schemes 6, 7, and 8 that R_1' and R_3' will form chiral centers, giving *R* and *S* enantiomers and diastereomers. Such enantiomers or diastereomers may be separated, if desired, at any stage by chiral HPLC, or by other conventional resolution techniques. Resolution of the precursor benzoic acid


-61-

may also be accomplished by fractional crystallization using mandelic acid, tartaric acid, or other chiral, optically pure acid bearing resolving agents. Chiral amides may also be prepared from the benzoic acids using chiral amines such as camphorsulfonamide, benzylamine, or the like. The isomers can then be separated and the chiral amide hydrolyzed as desired.

Scheme 9 illustrates synthesis of compounds of Formula I wherein one or both of R_6 and R_8 are halo via halogenation. The halogenation is carried out on a 2-aminobenzoic acid where one or both of R_6 and R_8 is hydrogen. If both R_6 and R_8 in the benzoic acid are H, then halogenation can be accomplished at both positions selectively or simultaneously. Thus, for example, chlorination at R_6 or R_8 is achieved by reaction of the benzoic acid with N-chlorosuccinamide, *t*-butylhypochlorite, chlorine gas, and the like. Similarly, bromination at R_6 and R_8 can also be accomplished by reaction of the benzoic acid with Br_2 , N-bromosuccinimide, and the like. Such halogenations are well-known in the art. Halogenation provides the respective mono- or dihalo compound. The halogenated benzoic acid can be further reacted as shown in Schemes 1 and 2 to provide the quinazoline-2,4-diones of Formula I.

-62-

Scheme 9




In Scheme 10, compounds where R_8 is H are halogenated as described in Scheme 9 to provide the corresponding 3-halo-2-aminobenzoic acid ($Y = \text{halo}$). This intermediate can then be diazotized by reaction of the 2-amino group with sodium nitrite or t-butyl nitrite, which is then converted to a 2-halobenzoic acid in the presence of an appropriate sodium, potassium, or copper salt such as sodium iodide, potassium chloride, and the like. The resulting 2,3-dihalobenzoic acid

-63-

(where X and Y both are halo) is then converted to the 3-halo-2-aminobenzoic acid by reaction with an amine R_1NH_2 in the presence of a copper catalyst. The 3-halo-2-aminobenzoic acid is converted to an amide and cyclized to the corresponding 8-halo-3-amino-quinazolin-2,4-dione, as illustrated in Scheme 2.

5

Scheme 10




Compounds of Formula I wherein R_6 and/or R_8 is halo such as chloro or bromo are readily dehalogenated by reaction with metal catalysts under hydrogen pressure (Scheme 11). Suitable catalysts include the many variations of Pd on carbon, Raney nickel, or other reagents that are well-known to effect such dehalogenation.

10

-64-

Scheme 11




Compounds of Formula I wherein R_6 and/or R_8 are hydrogen can be halogenated to give the mono- or dihalo compound (e.g., R_6 or $R_8 = \text{Cl or Br}$).

- 5 The invention compounds are preferably prepared by first halogenating a benzoic acid derivative, and then cyclizing the halogenated benzoate (Scheme 12). If both R_6 and R_8 are H, then halogenation can be accomplished at both positions selectively or simultaneously. Halogenations can be carried out as described above for Scheme 10. The resulting compound is then converted to the
- 10 2-substituted-aminobenzoic acid as depicted in Scheme 1, which is subsequently cyclized and aminated.

-65-


Scheme 12



Invention compounds of Formula I can also be prepared as shown in Scheme 13. Substituted benzoic acids can be converted into esters (where Z is an ester forming group such as alkyl or benzyl) by a number of methods known by those skilled in the art. The ester is reacted with an isocyanate such as trimethylsilylisocyanate, chlorosulfanyl isocyanate, and chlorocarbonyl isocyanate, followed by treatment with a base such as triethylamine, sodium t-butoxide or the like, to provide a quinazoline-2,4-dione. The quinazoline-2,4-dione can be aminated by reaction with an aminating agent such as ammonia as described in Scheme 5 to generate the corresponding 3-aminoquinazoline-2,4-dione. This compound may be further reacted, when R₇ is a leaving group such as fluoro, with various heterocyclic amines. Again, carbocycles and aryls may also be introduced at R₇ if R₇ is a Br, I, or triflate, using palladium catalyzed couplings of tin or boronate carbocycles and aryls. Removal of any protecting groups (Pro) by normal means provides invention compounds such as A, as described above in Scheme 1.

-66-


Scheme 13



- Invention compounds of Formula I having a cyano substituent can be prepared as shown in Scheme 14. For example, a benzoic acid wherein R_8 is
- 5 hydrogen can be metallated with a strong base such as lithium hexamethyldisilazane or lithium diisopropylamine. The resulting metallated intermediate can then be quenched with dimethylformamide or an equivalent to provide an aldehyde. The aldehyde can be converted to an oxime by reaction with an alkoxy amine (other electrophiles such as alkyl halides, activated amides, esters and halide sources such as 1,2-dichloro-tetrafluoroethane may also be employed
- 10 to provide other benzoic acid derivatives that can be used as defined in Schemes 1, 2, 13, and 15). The oxime is converted into a cyano group under the cyclization conditions required to form the quinazolinedione ring system (e.g., reaction with phosgene) as described in Schemes 1, 12, and 13. Deprotection
- 15 provides invention compounds where R_8 is $-\text{CN}$.

-67-

Scheme 14




Another alternative for preparing invention compounds is illustrated in Scheme 15. Appropriately substituted benzoic acids can be converted into benzamides by any number of methods as described above. A benzamide can then be treated with oxalyl chloride in a chlorinated solvent such as dichloroethane or an equivalent to provide an isocyanate. The isocyanate is reacted with a substituted primary amine to give a benzoyl substituted urea. This intermediate can be cyclized to form a quinazolin-2(1H)-one ring system by reaction with sodium hydride, potassium hexamethyldisilazane or other non-nucleophilic bases, generally in a solvent such as tetrahydrofuran (THF)/dimethylformamide, THF with 18-crown-6, THF/dioxane, THF/glyme, THF/diglyme, dimethoxyethane/toluene or an equivalent. The quinazolin-2(1H)-one can then be aminated by reaction with any number of aminating reagents as described by Kloetzer (*Sci. Pharm.*, 1984;52:46-50). The resulting 3-aminoquinazolin-2(1H)-one ring system can then be

-68-


readily coupled with an appropriately substituted heterocyclic amine (such as those noted above in Table A) by reaction in the presence of a base such as triethyl amine, diisopropylethyl amine, tetramethyl guanidine, and the like in solvents such as dimethyl sulfoxide, dimethylformamide, dimethylacetamide, sulfolane or the equivalent. Any protecting group associated with the heterocyclic amine side chain is then removed by methods known to those skilled in the art to provide invention compounds having Structure A.

Scheme 15



Tricyclic compounds (i.e., where R₁ and R₈, together with the atoms to which they are attached, form a carbocyclic ring in invention compounds of Formula I) can be prepared according to Scheme 16 where a palladium mediated carbon monoxide insertion on a quinoline (X = Br, I, or triflate) under well-precedented conditions gives rise to an ester. The quinoline ring can then be hydrogenated to provide a tetrahydroquinoline by standard hydrogenation conditions. The remainder of Scheme 16 follows that of Scheme 8 to provide invention compounds such as G, substituted with a displaceable substituent at R₇ (e.g., halo). These compounds may be further reacted with nucleophiles such as amines from Table A to give compounds of Formula I.


Scheme 16



Compounds of Formula I where K is N may be prepared by the routes shown in Schemes 1, 2, 3, 4, and 6, or by routes such as the one illustrated in Scheme 17, which follows closely the chemistry shown in Scheme 15. The leaving groups (e.g. halo) *ortho* and *para* to the carboxyl group of the pyridyl starting material are highly activated. The two leaving groups *ortho* to the pyridine nitrogen are generally chlorine, but fluorine, alkylthiol, and sulfoxides such as methylsulfoxide are also good leaving groups for such compounds. The urea intermediate readily cyclizes to the 7-halo-quinazolinone. Amination at the 3-position gives an invention compound that is an important intermediate due to the good leaving group at the 7-position ($R_7 = \text{halo}$), which is readily displaced by reaction with an amine NH_2 .

-70-


Scheme 17



5 Scheme 18 illustrates the synthesis of benzoic acid starting materials that
 have a bromo for R_7 . A difluoro substituted benzoic acid is converted to an ester
 by first reaction with oxalyl chloride or an equivalent reagent (including acid
 anhydride), and the acid halide or anhydride is reacted with an alkanol to afford
 the respective ester (Z =methyl, ethyl, isopropyl, etc.). The ester is then reacted
 with 4-methoxybenzylamine or an equivalent to produce the desired 4-substituted
 10 benzoic acid derivative. This intermediate is reacted with triethylsilane and
 trifluoroacetic acid in a chlorinated solvent such as dichloromethane or an
 equivalent to provide an aniline derivative. Alternatively, one skilled in the art
 might also employ transition metal catalysis. The resulting aniline is then
 subjected to diazotization and converted to a bromide by treatment with cuprous
 15 bromide. The ester is then hydrolyzed by well-known methods to the desired
 benzoic acid, which can be used as a starting material as illustrated in Scheme 20
 below.


-71-

Scheme 18



In Scheme 19, 4-bromo-2-fluorobenzoic acid derivatives can be selectively
 5 chlorinated by treatment with chlorine gas in chlorosulfonic acid at a temperature
 of 40°C-100°C.

Scheme 19



Scheme 20 illustrates use of 4-bromobenzoic acids as starting materials to
 10 make invention compounds wherein R_7 is aryl. The 4-bromobenzoic acids are
 converted into benzamides by any number of methods known in the art. A
 benzamide is reacted with oxalyl chloride in a chlorinated solvent such as
 dichloroethane or an equivalent to provide an isocyanate. The isocyanate is
 reacted with a substituted primary amine to give a benzoyl substituted urea. This
 15 intermediate can be cyclized to form a quinazolinedione ring system by reaction
 with sodium hydride, potassium hexamethyldisilazane or other non-nucleophilic
 bases in tetrahydrofuran/dimethylformamide, tetrahydrofuran with 18-crown-6,
 toluene/dioxane, tetrahydrofuran/glyme, tetrahydrofuran/diglyme, glyme/toluene


-72-

or an equivalent. The quinazolinedione can then be aminated with a number of aminating reagents as described in Scheme 15. The resulting 3-aminoquinazolinedione ring system can then be readily coupled with a stannane or boronic acid derivative of a substituted aryl such as phenyl or substituted aromatic heterocycle (Ar).

Alternatively, the 3-position amine can be protected with a protecting group such as a tert-butyl carbamate or trifluoroacetamide. Protecting groups of these types are well known in the art. The 3-derivatized aminoquinazolinedione ring system can then be coupled via palladium catalysis with an aromatic (Ar) stannane or boronic acid.

Each of the outlined routes, upon deprotection by standard procedures, provides invention compounds of Formula I where R₇ is aryl such as phenyl or substituted phenyl, or heteroaryl such as pyridyl or substituted pyridyl.


Scheme 20



-73-


Compounds of Formula I wherein R₇ is a heteroaryl group (such as those in Table A) are alternatively prepared as illustrated in Scheme 21, where R₇ is a substituted thiazole. A 3-protected 3-amino-7-bromoquinazolidinedione from Scheme 20 is reacted with 1-tributyl-ethoxyvinyltin in the presence of a palladium catalyst. The resulting 7-substituted adduct is reacted with a brominating reagent such as n-bromosuccinimide and the like in tetrahydrofuran/H₂O to provide an α-bromoketone at the quinazolidinedione 7-position. Reaction of this intermediate with a thioamide (or thiourea) in a polar solvent such as dimethyl formamide, dimethylacetamide, or ethanol, and at an elevated temperature of about 80°C to 120°C, effects cyclization to form a thiazolyl group. Deprotection by known methods can then be applied to provide invention compounds of Formula I wherein R₇ is the heteroaryl, optionally substituted with T, which is H, alkyl, substituted alkyl, NH₂, NH-alkyl and N-dialkyl.

Scheme 21



Alternatively, 7-aromatic or heterocyclic aromatic compounds of Formula I are prepared as shown in Scheme 22. A 4-bromo-2-fluorobenzoic acid (Schemes 18 or 19) is first esterified (Z is alkyl or benzyl), and the ester is reacted with a substituted primary amine (R_1NH_2) in dimethylsulfoxide (DMSO) at elevated temperature of about $100^\circ C$ to provide an anthranilic ester. The resulting aniline is then reacted with an appropriately protected hydrazine to provide the corresponding amide, for example as described in Scheme 1. The resulting amido aniline is then cyclized by reaction with phosgene, CDI, or the like to generate the quinazoline-2,4-dione. The cyclization typically is carried out in a solvent, such as ethereal solvents, chlorinated hydrocarbons such as chloroform, or aromatic hydrocarbons such as toluene, and in the presence of a base such as triethylamine or $NaHCO_3$. The quinazolin-2,4-dione ring system is then further modified as described in Scheme 20 to provide the desired 7-aryl (Ar) compound of Formula I.

Scheme 22



As noted above, R_7 in Formula I is referred to as the "side chain" of the 3-aminoquinazolin-2,4-dione nucleus. The R_7 side chains are any of those typically found on the quinolone antibiotics. Most of the R_7 side chain reactants that are required to make the Formula I compounds are readily available from commercial sources. Typical R_7 side chains are shown in Table A above.


-75-

The synthesis of the R₇ side chains can be accomplished by standard synthetic methods, for example as shown in the following schemes or as found in the literature. Those of ordinary skill in the art will be able to make any of the starting materials required to prepare the invention compounds, although most are available from commercial sources. The following schemes are provided to illustrate the synthesis of typical R₇ side chain reactants.

Scheme A1 illustrates the synthesis of typical pyrrolidines, which are preferred side chains (R₇) for invention compounds of Formula I. In Scheme A1, appropriately activated enones can undergo [3+2]cycloadditions under the conditions described by Tsuge et al. (Recent advances in azomethine ylid chemistry. In: *Advances in Heterocyclic Chemistry* [Katritzky A., ed.] San Diego: Academic Press, 231-349). These reactions are carried out in a chlorinated hydrocarbon solvent such as dichloromethane, chloroform, or dichloroethane and the like, and in the presence of a catalytic acid such as trifluoacetic acid, to provide substituted pyrrolidines (wherein S₁, S₂, S₃, and S₄ independently are alkyl, substituted alkyl, aryl, amino, alkyl and dialkylamino). These intermediates can then be treated with hydroxylamine or any O-alkylated or arylated hydroxylamine under a variety of conditions known to those skilled in the art to provide oximated substituted pyrrolidines. The oximes are reduced to amines with lithium aluminum hydride, diisobutylaluminum hydride, borane, or by selective catalytic hydrogenation. The resulting primary amines can then be protected using a number of methods as described in "Protecting Groups in Organic Synthesis" by Theodora Green (supra). The benzylic pyrrolidine can then be deprotected by hydrogenation, and the resulting pyrrolidine used in the preparation of 7-cyclic amino substituted 3-aminoquinazolinediones of Formula I as shown in the schemes above.

-76-


Scheme A1



In Scheme A2, enoates (or vinylogous nitriles) may also be used in [3+2]cycloadditions to provide 3- and 4-substituted pyrrolidines. The ester (or nitrile) functionality can then be treated directly with alkyl lithium reagents at low temperature (0°C) in ether to provide S₁ substituted ketones, or the ester (Z is alkyl or benzyl) can be hydrolyzed under conditions usually employed by those skilled in the art to provide carboxylic acids. This intermediate can then be transformed into an acid chloride with oxalyl chloride and catalytic dimethyl formamide in solvents such as dichloromethane or chloroform or into an activated amide (Singh J., Satyamurthi N., Aidhen I., Singh J., *Prakt. Chem. [Weinheim, Ger.]*, 2000;342(4):340-347). An acid chloride can be converted into a ketone using organocopper reagents (Lipshutz B.H., Sengupta S., *Org. React. [N.Y.]*, 1992;41:135-631), and Weinreb amides can be converted into ketones according to the methods described by Weinreb (*Tetrahedron Lett.*, 1981;22(39):3815-3818) using organo Grignard reagents. The resulting ketones can then be converted into pyrrolidines (optionally substituted with S₂, S₃, and S₄) as described in Scheme A1. As noted above, pyrrolidines and substituted pyrrolidines are preferred R₇ groups in Formula I.

-77-


Scheme A2



In Scheme A3, 3-carboxylic acid N-benzyl substituted pyrrolidines (Scheme A2) can be converted into amides by a number of methods well-known to those who practice the art of organic synthesis. The amides can then be treated with lithium aluminum hydride in an ethereal solvent such as diethyl ether or tetrahydrofuran or the like to provide amines that can be protected as described in Scheme A1. Hydrogenation with palladium catalysis provides an appropriately substituted pyrrolidine (S₂, S₃, S₄ are independently alkyl, lower alkyl, aryl, etc.).

10

Scheme A3




In Scheme A4, a 3-carboxypyrrolidinone (Culbertson T.P., Domagala J.M., Nichols J.B., Priebe S., Skeece R.W., *J. Med. Chem.*,

-78-

1987;30(10):1711-1715) can be converted into an acid chloride or Weinreb amide in the same fashion as that defined in Scheme A3. The acid chlorides are reacted with an organocopper reagent or an organo Grignard reagent (for a Weinreb amide) to provide a ketone that can be manipulated as described in Scheme A1 to provide an appropriately substituted pyrrolidine with a variety of substitutions at S₁ (alkyl, lower alkyl, substituted alkyl, aryl). The use of S-methylbenzyl (or R-methylbenzyl) as a protecting group for the pyrrolidine nitrogen allows for the separation of enantiomers and diastereomers at any step in the reaction sequence.

Scheme A4



10

In Scheme A5, carboxybenzyl protected 3-pyrrolidinone is reacted with a Wittig salt in the presence of sodium hydride or an equivalent in dimethylsulfoxide or any polar aprotic solvent to give an olefin. The resulting olefin is oxidized with 3-chloroperoxybenzoic acid in dichloromethane or other suitable chlorinated hydrocarbon to provide an epoxide. The epoxide is reacted with ammonium hydroxide in methanol, ethanol, dimethylformamide, dimethylacetamide or the like to generate an aminoalcohol. Methylation of the tertiary hydroxyl group can then be carried out by formation of a Schiff-base, followed by alkoxide formation with sodium hydride and treatment with iodomethane. Hydrolysis of the Schiff-base under standard conditions liberates the amine. Reaction of the amine with di-*tert*-butyldicarbonate in methanol, followed


15

20

-79-

by hydrogenation via palladium catalysis provides R₇ side chains (S₁ is alkyl, lower alkyl, and aryl) that can be used in the synthesis of compounds of Formula I.

Scheme A5




5

In Scheme A6, the aminoalcohol intermediate described in Scheme A5 is reacted with di-*tert*-butyldicarbonate in methanol to produce the N-protected intermediate. Other amine protection strategies familiar to those skilled in the art can also be utilized. Hydrogenation under palladium catalysis provides substituted pyrrolidine R₇ side chains that are used in the synthesis of compounds of Formula I.

10


Scheme A6



-80-

In Scheme A7, an epoxide intermediate (Scheme A5) is selectively opened with hydrofluoric acid in pyridine, and the resulting secondary alcohol is reacted with methanesulfonylchloride, or an equivalent sulfonating reagent, in dichloromethane or another chlorinated hydrocarbon and triethylamine. Other appropriate bases include diisopropylethylamine, pyridine, collidine, *n*-methylmorpholine, and the like. The mesylate is then displaced by reaction with sodium azide, and the azide is reduced with Raney nickel under hydrogen pressure in methanol. Such transformations are well-precedented in the art. The resulting amine is then treated with di-*tert*-butyldicarbonate in methanol as shown in Schemes A4-A6, and the product is hydrogenated to provide the R7 side chain as shown above.

Scheme A7




The synthesis of 3-substituted piperidines is demonstrated in Scheme A8. Esterification and alkylation of 3-piperidine carboxylic acid is simultaneously accomplished by treatment with a benzylbromide in a polar solvent such as dimethylformamide or dimethylsulfoxide and a base such as potassium carbonate (K₂CO₃) or an equivalent. The ester is then hydrolyzed, and the resulting carboxylic acid is converted to a methyl ketone by treatment with methyl lithium at low temperature (0°C). This intermediate is then converted to an oxime by a

-81-

number of methods known by those skilled in the art. The oxime is reduced with lithium aluminum hydride in tetrahydrofuran to provide an amine. The amine is reacted with di-*tert*-butyldicarbonate in dichloromethane and triethylamine as a base. As shown in Schemes A4-A6, the product is then hydrogenated to provide the R-7 side chain.

Scheme A8



It should be recognized from Schemes A1-A8 that substitutions on ring systems such as a pyrrolidine, piperazine, or piperidine ring will form chiral centers giving *R* and *S* enantiomers and diastereomers. Such enantiomers or diastereomers may be separated, if desired, by chiral HPLC at any stage. Resolution of any of the intermediates may be performed with techniques of fractional crystallization using mandelic acid, tartaric acid, or other chiral, optically pure acid resolving agents. Chiral benzylic amines can be used in the preparation of starting materials for the above schemes, and chiral amides may

-82-


also be prepared using chiral acids, such as mandelic acid and the like. The isomers can then be separated and the chiral amine can be hydrogenated, or the chiral amide can be hydrolyzed.

The synthesis of invention compounds of Formula I is further illustrated by the following detailed examples. The examples are not to be construed as limiting the invention in scope or spirit to the specific procedures described. The starting materials and various intermediates utilized in the following examples may be obtained from commercial sources, or are readily prepared from commercially available organic compounds, using well-known synthetic methods.

EXAMPLE 1

Synthesis of Benzoates


(a) *2,3,4,5-Tetrafluorobenzoic acid ethyl ester*



A solution of 2,3,4,5-tetrafluorobenzoic acid (4.68 g, 24.1 mmol) in dichloromethane (50 mL) at 0°C is treated with oxalyl chloride (11.5 mL, 132 mmol) followed by *N,N*-dimethylformamide (4 drops). The mixture is stirred at 0°C for 5 minutes and then at room temperature for 1.5 hours. The mixture is concentrated to dryness and subsequently co-evaporated from benzene. The resulting acid chloride is diluted with dichloromethane (50 mL), cooled to 0°C, and anhydrous ethanol (15.0 mL, 256 mmol) is added dropwise. The resulting solution is stirred at room temperature for 4.5 hours, then poured into saturated NaHCO₃ and extracted with chloroform. The combined organic extracts are washed with brine, dried over Na₂SO₄, filtered, and concentrated under vacuum to afford the title compound (5.37 g). ¹H NMR (CDCl₃): δ 7.67–7.55 (m, 1H), 4.42 (q, 2H), 1.41 (t, 3H).


-83-

(b) 2,3,4,6-Tetrafluorobenzoic acid ethyl ester



2,3,4,6-Tetrafluorobenzoic acid (4.8 g, 24.7 mmol) in dichloromethane (80 mL) is cooled to 0°C under a nitrogen atmosphere and treated with oxalyl chloride (11.2 mL, 128 mmol) followed by anhydrous *N,N*-dimethylformamide (2 drops). The mixture is warmed to room temperature and stirred for 2 hours. The solution is co-evaporated with benzene to yield an oil that is taken up in dichloromethane (80 mL), cooled to 0°C under an inert atmosphere, and treated with anhydrous ethanol (15 mL, 258 mmol). After 5 hours at room temperature, the solution is poured into saturated NaHCO₃ and extracted with chloroform. The organic phase is washed with brine, dried over Na₂SO₄, filtered, and concentrated under vacuum to provide the title compound (3.3 g). This material is used without further purification. ¹H NMR (CDCl₃): δ 6.93-6.75 (m, 1H), 4.42 (q, 2H), 1.39 (t, 3H).

(c) 3-Chloro-2,4,5-trifluorobenzoic acid ethyl ester




To a solution of 3-chloro-2,4,5-trifluorobenzoic acid (Example 20, 4.90 g, 23.3 mmol) in dichloromethane (50 mL) is added oxalyl chloride (5.1 mL, 58.2 mmol) and *N,N*-dimethylformamide (1 drop). After 45 minutes, the reaction mixture is concentrated under vacuum. The resulting residue is dissolved in dichloromethane (50 mL) and treated with ethanol (10 mL, 172 mmol). After 30 minutes, the reaction mixture is diluted with dichloromethane and washed with saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄.

-84-


filtered, and the filtrate is concentrated to afford the title compound (5.59 g). ^1H NMR (CDCl_3): δ 7.76-7.68 (m, 1 H), 4.40 (q, 2 H), 1.39 (t, 3 H).

(d) *2,3,4,5,6-Pentafluorobenzoic acid benzyl ester*



5 A solution of 2,3,4,5,6-pentafluorobenzoic acid (15.25 g, 71.9 mmol),
4-dimethylaminopyridine (4.4 g, 36.0 mmol) and benzyl alcohol (7.4 mL,
71.5 mmol) in dichloromethane (150 mL) is cooled to 0°C under an inert
atmosphere. 1-[3-(Dimethylamino)propyl]-3-ethyl carbodiimide hydrochloride
10 (16.7 g, 87.1 mmol) is added and the mixture stirred at 0°C for 2 hours. The
mixture is warmed to ambient temperature, stirred for 24 hours, and then poured
into brine. The mixture is extracted with chloroform and the organic phase washed
with NaHCO_3 , dried over Na_2SO_4 , filtered and concentrated under vacuum to
yield the title compound as a white solid (16.7 g). This material is used without
further purification. ^1H NMR (CDCl_3): δ 7.49-7.27 (m, 5 H), 5.40 (s, 2 H).

15 (e) *2,4,5-Trifluorobenzoic acid ethyl ester*




To a solution of 2,4,5-trifluorobenzoic acid (10.85 g, 61.6 mmol) in
dichloromethane (100 mL) is added oxalyl chloride (13.5 mL, 154 mmol) and
N,N-dimethylformamide (1 drop). After 45 minutes, the reaction mixture is
20 concentrated under vacuum. The resulting residue is dissolved in dichloromethane
(100 mL), and to this solution is added ethanol (18 mL, 310 mmol). After
30 minutes, the reaction mixture is diluted with dichloromethane and washed with
saturated NaHCO_3 , water, and brine. The organic layer is dried over MgSO_4 ,
filtered, and the filtrate is concentrated to afford the title compound as an oil

-85-


(12.23 g). ^1H NMR (CDCl_3): δ 7.85-7.73 (m, 1 H), 7.06-6.94 (m, 1 H), 4.38 (q, 2 H), 1.38 (t, 3 H).

(f) *2,4,5-Trifluoro-3-methoxybenzoic acid ethyl ester*



5 A solution of 2,4,5-trifluoro-3-methoxybenzoic acid (2.513 g, 12.19 mmol) in dichloromethane (50 mL) at 0°C is treated with oxalyl chloride (5.4 mL, 61.9 mmol) followed by *N,N*-dimethylformamide (4 drops). The mixture is stirred at 0°C for 5 minutes and then at room temperature for 2.5 hours. The mixture is concentrated to near dryness and then co-evaporated from benzene. The resulting acid chloride is diluted with dichloromethane (50 mL), cooled to 0°C , and then anhydrous ethanol (7.30 mL, 124 mmol) is added dropwise. The resulting solution is stirred at room temperature for 4.5 hours, poured into saturated NaHCO_3 solution, and extracted with dichloromethane. The combined organic extracts are washed with brine, dried over Na_2SO_4 , filtered, and concentrated under vacuum to afford the title compound as a light yellow liquid (2.82 g). ^1H NMR (CDCl_3): δ 7.47 (ddd, 1 H), 4.39 (q, 2 H), 4.05 (s, 3 H), 1.40 (t, 3 H).

(g) *2,3,4-Trifluorobenzoic acid ethyl ester*




20 To a solution of 2,3,4-trifluorobenzoic acid (10.85 g, 61.6 mmol) in dichloromethane (100 mL) is added oxalyl chloride (13.5 mL, 154 mmol) and *N,N*-dimethylformamide (3 drops). After 45 minutes, the reaction mixture is concentrated under vacuum. The resulting residue is dissolved in dichloromethane

-86-


(100 mL) and to this solution is added ethanol (18 mL, 310 mmol). After 30 minutes, the reaction mixture is diluted with dichloromethane and washed with saturated NaHCO_3 , water, and brine. The organic layer is dried over MgSO_4 , filtered, and the filtrate is concentrated to afford the title compound as an oil (12.23 g). ^1H NMR (CDCl_3): δ 7.93-7.67 (m, 1 H), 7.09-6.96 (m, 1 H), 4.39 (q, 2 H), 1.40 (t, 3 H).

(h) 2,3,4-Trifluoro-5-methoxybenzoic acid ethyl ester



A solution of 2,3,4-trifluoro-5-methoxybenzoic acid (Hayashi et al., *Bull. Chem. Soc. Jap.*, 1972;45(9):2909-2914, 3.443 g, 16.7 mmol) in dichloromethane (100 mL) at 0°C is treated with oxalyl chloride (7 mL, 80 mmol) followed by *N,N*-dimethylformamide (4 drops). The mixture is stirred at room temperature for 3.5 hours, concentrated to near dryness, and co-evaporated with benzene. The acid chloride is then diluted with dichloromethane, cooled to 0°C , and treated with absolute ethanol (10 mL, 170 mmol). The resulting solution is stirred at room temperature for 20 hours, poured into saturated NaHCO_3 solution, and extracted with dichloromethane. The organic extracts are washed with brine, dried over Na_2SO_4 , filtered, and concentrated under vacuum to afford the title compound (3.785 g). ^1H NMR (200 MHz, CDCl_3): δ 7.36-7.29 (m, 1H), 4.42 (q, 2H), 3.93 (s, 3H), 1.44 (t, 3H).


(i) 2,4,5-Trifluoro-3-methylbenzoic acid ethyl ester



-87-

A solution of 3-methyl-2,4,5-trifluorobenzoic acid (Shimizu T., Asai T., Kumai S., Jpn. Kokai Tokkyo Koho (1997) Japanese Appl. JP 95-219069, 4.3 g, 22.6 mmol) in dichloromethane (100 mL) at 0°C is treated with oxalyl chloride (4.2 g, 33 mmol) followed by *N,N*-dimethylformamide (2 drops). The mixture is stirred at room temperature for 3 hours, concentrated and dried in vacuo. The acid chloride is diluted with dichloromethane, cooled to 0°C, and treated with absolute ethanol (10 mL, 170 mmol). The resulting solution is stirred at room temperature for 20 hours, poured into saturated NaHCO₃ solution and extracted with dichloromethane. The organic extracts are washed with brine, dried over Na₂SO₄, filtered, and concentrated under vacuum to afford the title compound (3.8 g). ¹H NMR (200 MHz, CDCl₃): δ 7.67-7.26 (m, 1H), 4.44 (q, 2H), 2.27 (dd, 3H), 1.42 (t, 3H).

(j) *3-(Benzyloxyiminomethyl)-2,4,5-trifluorobenzoic acid ethyl ester*



2,4,5-Trifluoro-3-formylbenzoic acid (Horiuchi N., Yonezawa T., Chiba K., Yoshida H., PCT Int. Appl. [1999] WO 97-JP2918), (5.70 g, 27.9 mmol), triethylamine (11 mL, 78.9 mmol), O-benzylhydroxylamine hydrochloride (4.47 g, 28.0 mmol) and anhydrous tetrahydrofuran (150 mL) are stirred at room temperature for 50 hours. The mixture is poured into brine and the pH adjusted to 5.5-6.0 using 1.0 M hydrochloric acid. The mixture is extracted with ethyl acetate. The organic phase is dried over Na₂SO₄, filtered, and concentrated in vacuo to provide 3-(benzyloxyiminomethyl)-2,4,5-trifluorobenzoic acid (8.63 g), which is used without further purification. ¹H NMR (200 MHz, CDCl₃): δ 9.82-9.18 (bs, 1H), 8.24 (s, 1H), 7.94-7.68 (m, 1H), 7.55-7.23 (m, 5H), 5.23 (d, 2H). MS (EI, M-1) m/z 308.

-88-

3-(Benzyloxyiminomethyl)-2,4,5-trifluorobenzoic acid (0.117 g, 0.378 mmol), 4-dimethylaminopyridine (0.023 g, 0.189 mmol), anhydrous ethanol (0.030 mL, 0.517 mmol) are mixed in dichloromethane (6 mL), cooled to 0°C, and treated with 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.082 g, 0.427 mmol). After 2 hours at 0°C and room temperature for 21 hours, the solvents are evaporated. The residue is dissolved in chloroform, washed with saturated NaHCO₃ and brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. Purification by chromatography (1:8 ethyl acetate/hexanes) provided the title compound (0.068 g). ¹H NMR (200 MHz, CDCl₃): δ 8.27 (s, 1H), 7.87-7.66 (m, 1H), 7.49-7.28 (m, 5H), 5.27 (s, 2H), 4.48-4.29 (q, 2H), 1.46-1.31 (t, 3H).


EXAMPLE 2

Synthesis of 4-Heterocyclic Benzoates

General Procedure

A mixture of 4-fluoro substituted benzoic acid ester (1 eq.), substituted pyrrolidine (1.3-2.5 eq.), and triethylamine (2-10 eq.) in *N,N*-dimethylacetamide or acetonitrile is heated to 50°C-100°C for 15 to 60 hours. The solution is then concentrated to near dryness, poured into saturated NaHCO₃ solution, and extracted with ethyl acetate. The combined organic extracts are washed with brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. The resulting residue is purified by column chromatography (ethyl acetate/hexanes) to afford 4-pyrrolidinyl substituted benzoic acid esters.

(a) *4-[(S)-3-tert-Butoxycarbonylamino]pyrrolidin-1-yl]-2,3,5-trifluorobenzoic acid ethyl ester*




-89-

A solution of 2,3,4,5-tetrafluorobenzoic acid ethyl ester (Example 1a, 4.03 g, 18.1 mmol), (S)-pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (4.02 g, 21.6 mmol) (Sanchez J.P., Domagala J.M., Heifetz C.L., Priebe S.R., Sesnie J.A., Trehan A.K., *J. Med. Chem.*, 1992;35(10):1764-1773), and triethylamine (18.0 mL, 129 mmol) in acetonitrile (50 mL) is heated at 50°C under a nitrogen atmosphere for 4.5 hours. The yellow solution is concentrated in vacuo and poured into a saturated NaHCO₃ solution. The mixture is then extracted with chloroform, and the combined organic extracts are washed with brine, dried with Na₂SO₄, filtered, and concentrated in vacuo. The resulting residue is then purified by flash column chromatography (1:7 ethyl acetate/hexanes) to afford the title compound (5.24 g). ¹H NMR (CDCl₃): δ 7.35 (ddd, 1H), 4.70-4.65 (bs, 1H), 4.42-4.20 (m, 3H), 3.90-3.62 (m, 3H), 3.58-3.42 (m, 1H), 2.28-2.03 (m, 1H), 1.99-1.80 (m, 1H), 1.46 (s, 9H), 1.37 (t, 3H).

(b) 4-[(S)-3-*tert*-Butoxycarbonylamino

pyrrolidin-1-yl]-2,3,6-trifluorobenzoic acid ethyl ester




A solution of 2,3,4,6-tetrafluorobenzoic acid ethyl ester (Example 1b, 5.1 g, 22.9 mmol), triethylamine (22 mL, 157 mmol), and (S)-pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (5.2 g, 27.9 mmol) in acetonitrile (60 mL) is stirred at room temperature for 64 hours. The mixture is concentrated in vacuo and the residue dissolved in chloroform, washed with NaHCO₃ and then brine, dried over Na₂SO₄, filtered, and concentrated under vacuum to afford a solid. The solid is purified by column chromatography (1:1:7 ethyl acetate/chloroform/hexanes) to afford the title compound (6.21 g). ¹H NMR (CDCl₃): δ 6.07 (ddd, 1H),

-90-

4.78-4.63 (bd, 1H), 4.49-4.22 (m, 3H), 3.80-3.68 (m, 1H), 3.66-3.28 (m, 3H),
2.33-2.13 (m, 1H), 2.04-1.83 (m, 1H), 1.45 (s, 9H), 1.36 (t, 3H).

(c) *4-[(S)-3-tert-Butoxycarbonylaminopyrrolidin-1-yl]-3-chloro-2,5-difluorobenzoic acid ethyl ester*




5

To a solution of 3-chloro-2,4,5-trifluorobenzoic acid ethyl ester (Example 1c, 5.50 g, 23.05 mmol) in acetonitrile (25 mL) is added triethylamine (6.43 mL, 46.1 mmol) and (S)-pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (4.72 g, 25.4 mmol). After 48 hours, the mixture is diluted with ethyl acetate and washed with saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate is concentrated to afford the title compound (9.34 g). MS CI: *m/z* 405 (MH⁺).

10

(d) *4-[(S)-3-(tert-Butoxycarbonylamino)pyrrolidin-1-yl]-2,3,5,6-tetrafluorobenzoic acid benzyl ester*



15


A solution of 2,3,4,5,6-pentafluorobenzoic acid benzyl ester (Example 1d, 8.05 g, 26.6 mmol), triethylamine (26 mL, 186 mmol), and (S)-pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (5.6 g, 30 mmol) in acetonitrile (150 mL) is stirred at room temperature for 24 hours. The solvent is removed under vacuum and the residue dissolved in chloroform, washed with NaHCO₃ and then brine, dried over Na₂SO₄, filtered, and concentrated in vacuo to afford a solid. The solid is then purified by column chromatography (1:1:6 ethyl acetate/chloroform/hexanes) to

20

-91-

yield the title compound as a solid (7.92 g). ^1H NMR (CDCl_3): δ 7.48-7.30 (m, 5H), 5.35 (s, 2H), 4.77-4.50 (m, 1H), 4.37-4.19 (m, 1H), 4.01-3.63 (m, 3H), 3.59-3.46 (m, 1H), 2.29-2.09 (m, 1H), 2.01-1.82 (m, 1H), 1.45 (s, 9H).

(e) **4-(Pyrrolidin-1-yl)-2,5-difluorobenzoic acid ethyl ester**




5

To a solution of 2,4,5-trifluorobenzoic acid ethyl ester (Example 1e, 4.32 g, 21.2 mmol) in acetonitrile (25 mL) is added triethylamine (5.9 mL, 42.3 mmol) and pyrrolidine (2.2 mL, 25.4 mmol). After 24 hours, the reaction mixture is diluted with ethyl acetate and washed with saturated NaHCO_3 , water, and brine. The organic layer is dried over MgSO_4 , filtered, and the filtrate is concentrated to afford the title compound as an oil (5.25 g). MS CI: m/z 256 (MH^+).

10

(f) **4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2,5-difluoro-3-methoxybenzoic acid ethyl ester**



15


A solution of 2,4,5-trifluoro-3-methoxybenzoic acid ethyl ester (Example 1f, 2.821 g, 12.05 mmol), pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (3.310 g, 17.77 mmol), and triethylamine (9.13 g, 12.6 mL, 90.23 mmol) in acetonitrile (50 mL) is heated at 40°C under nitrogen for 2 days. The yellow solution is concentrated to near dryness and poured into saturated NaHCO_3 solution and extracted with dichloromethane. The combined organic extracts are washed with brine, dried over Na_2SO_4 , filtered, and concentrated under vacuum. The resulting residue is purified by flash column chromatography

20

-92-

(1:7 ethyl acetate/hexanes) to afford the title compound as a yellow solid (3.743 g). MS EI: m/z 401 (MH^+).

(g) **4-[3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl]-3-chloro-2,5-difluorobenzoic acid ethyl ester**




5

To a solution of 3-chloro-2,4,5-trifluorobenzoic acid ethyl ester (Example 1c, 5.50 g, 23.05 mmol) in acetonitrile (25 mL) is added triethylamine (6.43 mL, 46.1 mmol) and pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (4.72 g, 25.4 mmol). After 48 hours, the mixture is diluted with ethyl acetate and washed with saturated $NaHCO_3$, water, and brine. The organic layer is dried over $MgSO_4$, filtered, and the filtrate is concentrated to afford the title compound (9.2 g). MS CI: m/z 405 (MH^+).

10

(h) **4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2,3-difluorobenzoic acid ethyl ester**



15


A solution of 2,3,4-trifluorobenzoic acid ethyl ester (Example 1g, 3.70 g, 18.1 mmol), pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (4.02 g, 21.6 mmol), and triethylamine (18.0 mL, 129 mmol) in acetonitrile (50 mL) is heated at 50°C under nitrogen atmosphere for 4.5 hours. The solution is concentrated in vacuo and poured into a saturated $NaHCO_3$ solution. The mixture is then extracted with chloroform, and the combined organic extracts are washed with brine, dried with Na_2SO_4 , filtered, and the mixture concentrated in vacuo. The resulting residue is

20

-93-

then purified by column chromatography (1:7 ethyl acetate/hexanes) to afford the title compound (4.70 g). ^1H NMR (CDCl_3): δ 7.59-7.49 (m, 1H), 6.35-6.26 (m, 1H), 4.88-4.84 (bd, 1H), 4.39-4.28 (m, 3H), 3.81-3.50 (m, 3H), 3.48-3.34 (m, 1H), 2.32-2.15 (m, 1H), 2.02-1.76 (m, 1H), 1.45 (s, 9H), 1.40 (t, 3H).

- 5 (i) 4-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-2,5-difluorobenzoic acid ethyl ester



To a solution of 2,4,5-trifluorobenzoic acid ethyl ester (Example 1e, 14.8 g, 72.4 mmol) in acetonitrile (225 mL) is added triethylamine (65 mL, 466 mmol) and pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (16.07 g, 86.2 mmol). After 45 hours, the reaction mixture is diluted with ethyl acetate and washed with saturated NaHCO_3 , water, and brine. The organic layer is dried over MgSO_4 , filtered, and the filtrate is concentrated to afford an oil. The resulting residue is purified by flash chromatography (1:5 ethyl acetate/hexanes) to afford the title compound. MS (ES, $\text{M}+1$) m/z 371, mp 98-100°C.

The following compounds are synthesized according to general procedures described in Examples 2a-i above.

- (k) 4-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-2,5-difluoro-3-methylbenzoic acid ethyl ester (MS ES, MH^+) m/z 385) using 2,4,5-trifluoro-3-methylbenzoic acid ethyl ester (Example 1i) and pyrrolidin-3-ylcarbamic acid *tert*-butyl ester.

- (l) 4-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-2,5-difluoro-3-methylbenzoic acid ethyl ester (MS ES, MH^+) m/z 399) using 2,4,5-trifluoro-3-methylbenzoic acid ethyl ester (Example 1i) and 3-(*tert*-butoxycarbonylaminomethyl)pyrrolidine (Rogers D.H., Saunders J., Williams J.P., DE 19955794).

-94-

(m) 4-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-2,3-difluoro-5-methoxybenzoic acid ethyl ester (MS ES, MH^+) m/z 401) using 2,3,4-trifluoro-5-methoxybenzoic acid ethyl ester (Example 1h) and pyrrolidin-3-ylcarbamic acid *tert*-butyl ester.

5 (n) 4-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-2,5-difluoro-3-methoxybenzoic acid ethyl ester (MS ES, MH^+) m/z 415) using 2,4,5-trifluoro-3-methoxybenzoic acid ethyl ester (Example 1f) and 3-(*tert*-butoxycarbonylaminomethyl)pyrrolidine (Rogers D.H., Saunders J., Williams J.P., DE 19955794).

10 (o) 4-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-3-ethoxy-2,5-difluorobenzoic acid ethyl ester. (^1H NMR (200 MHz, $CDCl_3$): δ 7.31 (dd, 1H), 4.71 (bs, 1H), 4.34-3.4 (m, 9H), 2.23-1.81 (m, 2H), 1.45 (s, 9H), 1.37 (t, 3H), 1.33 (t, 3H); ((MS ES, MH^+) m/z 415.) using 3-ethoxy-2,5-difluorobenzoic acid ethyl ester (Sanchez J.P., Gogliotti R.D., Domagala J.M., Gracheck S.J., Huband M.D.,
15 Sesnie J.A., Cohen M.A., Shapiro M.A., *J. Med. Chem.*, 1995;38(22):4478-4487) and pyrrolidin-3-ylcarbamic acid *tert*-butyl ester.

(p) 3-(Benzyloxyiminomethyl)-4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2,5-difluorobenzoic acid ethyl ester (MS ES, MH^+) m/z 504) using 3-(benzyloxyiminomethyl)-2,4,5-trifluorobenzoic acid
20 ethyl ester (Example 1j) and pyrrolidin-3-ylcarbamic acid *tert*-butyl ester.

(q) 3-Chloro-2,5-difluoro-4-(3-[1,2,3]-triazol-1-ylpyrrolidin-1-yl)benzoic acid ethyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.79 (s, 1H), 7.65 (s, 1H), 7.41-7.51 (m, 1H), 5.40-5.30 (m, 1H), 4.34-4.24 (q, 2H), 4.13-3.45 (m, 4H), 2.65-2.50 (m, 1H), 2.43-2.28 (m, 1H), 1.34-1.27 (t, 3H)) using 3-chloro-
25 2,4,5-trifluorobenzoic acid ethyl ester (Example 1c) and 1-pyrrolidin-3-yl-1H-[1,2,3]triazole (Singh R., Singh I.P., Thomas G., Singh M.P., Micetich R.G., Fahti-Afshar R., Doerksen T.R., PCT Int. Appl., 1992, WO 9210492 A1).

-95-


EXAMPLE 3.

Synthesis of 2-Substituted-Amino Benzoates

General Procedure

A solution of 4-(pyrrolidin-1-yl)-2-fluorobenzoic acid ethyl esters (Example 2) and cyclopropylamine (10-15 eq.) in dimethyl sulfoxide and triethylamine (2-5 eq.) is heated in a sealed glass tube for 3 days at 140°C. Compressed air is blown into the mixture to remove excess cyclopropylamine. The resulting solution is concentrated under vacuum and purified by column chromatography (ethyl acetate/hexanes) to afford 4-(substituted pyrrolidin-1-yl)-2-cyclopropylamino-benzoic acid esters.


(a) *4-[(S)-3-tert-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,5-difluorobenzoic acid ethyl ester*



A solution of 4-[(S)-3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-2,3,5-trifluorobenzoic acid ethyl ester (Example 2a, 1.95 g, 5.01 mmol) and cyclopropylamine (14.0 mL, 201 mmol) in dimethyl sulfoxide (5 mL) is heated in a sealed glass tube for 44.5 hours at 100°C. Compressed air is blown into the black solution to remove excess cyclopropylamine. The solution is concentrated under high vacuum and purified by flash column chromatography (1:9 ethyl acetate/hexanes) to afford the title compound (1.77 g). ¹H NMR (CDCl₃): δ 7.32 (dd, 1H), 4.81-4.67 (bs, 1H), 4.35-4.16 (m, 3H), 3.95-3.40 (m, 4H), 2.92-2.80 (m, 1H), 2.25-2.08 (m, 1H), 1.97-1.77 (m, 1H), 1.46 (s, 9H), 1.34 (t, 3H), 0.72-0.63 (m, 2H), 0.57-0.49 (m, 2H).


-96-

(b) *4-[(S)-3-tert-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,6-difluorobenzoic acid ethyl ester*



A solution of 4-[(S)-3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-2,3,6-trifluorobenzoic acid ethyl ester (Example 2b, 5.4 g, 13.9 mmol),
 5 cyclopropylamine (40 mL, 577 mmol), and dimethyl sulfoxide (28 mL) in a sealed glass tube is heated at 100°C for 27 hours. The excess amine is removed by blowing in compressed air before the solution is concentrated under high vacuum. The residue is then purified by column chromatography (1:1:8 ethyl acetate/
 10 chloroform/hexanes) to afford the title compound (4.80 g). ¹H NMR (CDCl₃):
 δ 7.42 (bs, 1H), 5.69 (ddd, 1H), 4.76-4.61 (m, 1H), 4.37-4.19 (m, 3H),
 3.81-3.66 (m, 1H), 3.65-3.43 (m, 2H), 3.39-3.28 (m, 1H), 2.98-2.82 (m, 1H),
 2.30-2.09 (m, 1H), 1.98-1.82 (m, 1H), 1.45 (s, 9H), 1.34 (t, 3H), 0.72-0.68 (m,
 2H), 0.67-0.44 (m, 2H).

(c) *4-[(S)-3-tert-Butoxycarbonylamino]pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoic acid ethyl ester*




A solution of 4-[(S)-3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-3-chloro-2,5-difluorobenzoic acid ethyl ester (Example 2c, 9.34 g, 23.1 mmol),
 20 cyclopropylamine (16 mL, 228 mmol), and dimethyl sulfoxide (30 mL) is heated in a sealed tube at 110°C for 3 days. After cooling to room temperature, the reaction mixture is diluted with ethyl acetate and washed with saturated NaHCO₃,

-97-


water, and brine. The organic layer is dried over MgSO_4 , filtered, and the filtrate is concentrated to afford a brown residue. The residue is then purified via flash column chromatography (1:1 ethyl acetate/hexanes) to afford the title compound (4.41 g). MS CI: m/z 442 (MH^+).

- 5 (d) **4-[(S)-3-*tert*-Butoxycarbonylamino**
pyrrolidin-1-yl]-2-cyclopropylamino-
3,5,6-trifluorobenzoic acid benzyl ester



A solution of 4-[(S)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-
 2,3,5,6-tetrafluorobenzoic acid benzyl ester (Example 2d, 2.5 g, 5.3 mmol),
 10 cyclopropylamine (40 mL, 577 mmol), and dimethyl sulfoxide (20 mL) in a sealed
 glass tube is heated at 80°C for 24 hours. The excess amine is removed by
 blowing in compressed air, and the solution is then concentrated under high
 vacuum. The residue is purified by column chromatography (1:1:7 ethyl acetate/
 chloroform/hexanes) to afford the title compound as a solid (2.31 g). ^1H NMR
 15 (CDCl_3): δ 7.47-7.27 (m, 5H), 7.12 (bs, 1H), 5.30 (s, 2H), 4.78-4.62 (m, 1H),
 4.35-4.17 (m, 1H), 3.96-3.56 (m, 3H), 3.54-3.39 (m, 1H), 2.91-2.73 (m, 1H),
 2.24-2.02 (m, 1H), 1.96-1.77 (m, 1H), 1.45 (s, 9H), 0.69-0.56 (m, 2H),
 0.53-0.40 (m, 2H).

- (e) **2-Cyclopropylamino-5-fluoro-4-pyrrolidin-1-ylbenzoic acid ethyl ester**




20 4-(Pyrrolidin-1-yl)-2,5-difluorobenzoic acid ethyl ester (Example 2e,
 5.25 g, 20.6 mmol) and cyclopropylamine (30 mL, 428 mmol) are stirred in

-98-


dimethyl sulfoxide (30 mL) at 110°C for 3 days in a sealed tube. After cooling to room temperature, the reaction mixture is diluted with ethyl acetate and washed with saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate concentrated to afford a brown residue. The residue is purified via flash column chromatography (1:9 hexanes/dichloromethane) to afford the title compound as a solid (5.39 g). MS CI: m/z 293 (MH⁺).

(f) **4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-5-fluoro-2-isopropylaminobenzoic acid ethyl ester**



4-[3-(*tert*-Butoxycarbonylamino)-pyrrolidin-1-yl]-2,5-difluorobenzoic acid ethyl ester (Example 2j, 5.4 g, 13.9 mmol), isopropylamine (40 mL, 469 mmol), and dimethyl sulfoxide (25 mL) are heated at 100°C for 4 days in a sealed glass tube. The excess amine is removed by blowing in compressed air. The residue is filtered and concentrated under vacuum to a thick oil. Purification by column chromatography (1:1:7 ethyl acetate/chloroform/hexanes) to provide the title compound as a solid (1.6 g). MS: m/z 410 (MH⁺).

(g) **4-[(*S*)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-*sec*-butylamino-3-chloro-5-fluorobenzoic acid ethyl ester**




4-[(*S*)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-3-chloro-2,5-difluorobenzoic acid ethyl ester (Example 2c, 1.95 g, 4.81 mmol), *sec*-butylamine (30 mL, 296 mmol), and dimethyl sulfoxide (20 mL) are heated in a sealed glass tube at 110°C for 24 hours. The reaction mixture is evaporated, and purification

-99-

by column chromatography (1:8, ethyl acetate/hexanes) provides the title compound as a yellow syrup (3.69 g). MS EI: m/z 458 (MH^+).


(h) *4-[3-tert-butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoic acid ethyl ester*



A solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2,5-difluoro-3-methoxybenzoic acid ethyl ester (Example 2f, 3.087 g, 7.71 mmol) and cyclopropylamine (12.0 mL, 172 mmol) in dimethyl sulfoxide (5 mL) is heated in a sealed glass tube for 3 days at 110°C. Compressed air is blown into the black solution to remove excess cyclopropylamine. The solution is concentrated under vacuum and purified by flash column chromatography (1:5 ethyl acetate/hexanes) to afford the title compound as a yellow solid (0.875 g). ^1H NMR ($CDCl_3$):

δ 7.33 (d, 1H), 7.15 (bs, 1H), 4.78-4.65 (m, 1H), 4.33-4.15 (m, 3H), 3.92-3.50 (m, 3H), 3.55 (s, 3H), 3.45-3.34 (m, 1H), 3.02-2.89 (m, 1H), 2.28-2.09 (m, 1H), 1.93-1.77 (m, 1H), 1.46 (s, 9H), 1.34 (t, 3H), 0.7-0.59 (m, 2H), 0.49-0.42 (m, 2H).

(i) *4-[3-tert-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclobutylamino-5-fluorobenzoic acid ethyl ester*




A solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2,5-difluorobenzoic acid ethyl ester (2j, 3.059 g, 8.26 mmol) and cyclobutylamine (15.0 mL, 175 mmol) in dimethyl sulfoxide (20 mL) is heated in a sealed vessel for 4 days at 110°C. Compressed air is blown into the black solution to remove

-100-


excess cyclobutylamine. The solution is concentrated under vacuum and purified by flash column chromatography (1:5 ethyl acetate/hexanes) to afford the title compound as a yellow solid (1.715 g). ¹H NMR (CDCl₃): δ 7.67 (bs, 1H), 7.48 (d, 1H), 5.54 (d, 1H), 4.77-4.64 (bs, 1H), 4.38-4.18 (m, 3H), 3.97-3.85 (m, 1H), 3.80-3.42 (m, 3H), 3.41-3.30 (m, 1H), 2.50-2.32 (m, 2H), 2.28-2.12 (m, 1H), 2.05-1.70 (m, 5H), 1.46 (s, 9H), 1.35 (t, 3H).

(j) **4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3-fluorobenzoic acid ethyl ester**



A solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2,3-difluorobenzoic acid ethyl ester (Example 2h, 1.85 g, 5.01 mmol) and cyclopropylamine (14.0 mL, 201 mmol) in *N,N*-dimethylacetamide (5 mL) is heated in a sealed glass tube for 48 hours at 100°C. The solution is then concentrated under high vacuum and purified by column chromatography (1:9 ethyl acetate/hexanes) to afford the title compound (1.57 g). ¹H NMR (CDCl₃): δ 7.59 (bs, 1H), 7.57 (dd, 1H), 6.00 (dd, 1H), 4.82-4.68 (bd, 1H), 4.39-4.18 (m, 3H), 3.80-3.49 (m, 3H), 3.41-3.30 (m, 1H), 3.00-2.88 (m, 1H), 2.31-2.11 (m, 1H), 2.00-1.85 (m, 1H), 1.45 (s, 9H), 1.37 (t, 3H), 0.73-0.50 (m, 4H).

(k) **4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-5-chloro-2-cyclopropylamino-3-fluorobenzoic acid ethyl ester**



-101-

A solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-5-chloro-2,3-difluorobenzoic acid ethyl ester (Example 21, 3.75 g, 9.3 mmol) and cyclopropylamine (14.0 mL, 201 mmol) in *N,N*-dimethylacetamide (5 mL) is heated in a sealed glass tube for 48 hours at 100°C. The solution is then concentrated under high vacuum and purified by column chromatography (1:9 ethyl acetate/hexanes) to afford the title compound (2.0 g). ¹H NMR (CDCl₃): δ 7.64 (d, 1H), 7.56 (bs, 1H), 4.98-4.90 (bd, 1H), 4.40-4.20 (m, 3H), 3.78-3.63 (m, 4H), 2.97-2.83 (m, 1H), 2.35-2.21 (m, 1H), 1.93-1.81 (m, 1H), 1.46 (s, 9H), 1.38 (t, 3H), 0.75-0.61 (m, 2H), 0.57-0.50 (m, 2H).

The following compounds are synthesized according to general procedures Examples 3a-k described above.

(l) 4-(3-*tert*-Butoxycarbonylamino)pyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-3-methylbenzoic acid ethyl ester ([MS ES, MH⁺] *m/z* 422) using 4-(3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl)-2,5-difluoro-3-methylbenzoic acid ethyl ester (Example 2k).

(m) 4-[3-(*tert*-Butoxycarbonylamino)methyl]pyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-3-methylbenzoic acid ethyl ester. ([MS ES, M+1] *m/z* 436) using 4-[3-(*tert*-butoxycarbonylamino)methyl]-pyrrolidin-1-yl)-2,5-difluoro-3-methylbenzoic acid ethyl ester (Example 2l).

(n) 4-(3-*tert*-Butoxycarbonylamino)pyrrolidin-1-yl)-2-cyclopropylamino-3-fluoro-5-methoxybenzoic acid ethyl ester. ([MS ES, M+1] *m/z* 439) using 4-(3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl)-2,3-difluoro-5-methoxybenzoic acid ethyl ester (Example 2m).

(o) 4-[3-(*tert*-Butoxycarbonylamino)methyl]pyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-3-methoxybenzoic acid ethyl ester ([MS ES, MH⁺] *m/z* 453) using 4-[3-(*tert*-butoxycarbonylamino)methyl]-pyrrolidin-1-yl)-2,5-difluoro-3-methoxybenzoic acid ethyl ester (Example 2n).

(p) 4-(3-*tert*-Butoxycarbonylamino)pyrrolidin-1-yl)-2-cyclopropylamino-3-ethoxy-5-fluorobenzoic acid ethyl ester ([MS ES, MH⁺]

-102-

m/z 452). using 4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-3-ethoxy-2,5-difluorobenzoic acid ethyl ester (Example 2o).

(q) 3-(Benzyloxyiminomethyl)-4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-5-fluorobenzoic acid ethyl ([MS ES, MH⁺] m/z 541) using 3-(benzyloxyiminomethyl)-4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2,5-difluorobenzoic acid ethyl ester (Example 2p).

(r) 3-Chloro-2-cyclopropylamino-5-fluoro-4-(3-[1,2,3]-triazol-1-yl-pyrrolidin-1-yl)benzoic acid ethyl ester (¹H NMR (200 MHz, CDCl₃): δ 8.00 (s, 1H), 7.73 (s, 1H), 7.52 (d, 1H), 7.19 (bs, 1H), 5.53-5.39 (m, 1H), 4.36-4.21 (q, 2H), 4.14-4.01 (m, 1H), 3.98-3.83 (m, 1H), 2.71-2.58 (m, 1H), 2.43-2.29 (m, 1H), 1.41-1.28 (t, 3H), 0.73-0.59 (m, 2H), 0.52-0.37 (m, 2H)) using 3-chloro-2,5-difluoro-4-(3-[1,2,3]-triazol-1-yl-pyrrolidin-1-yl)benzoic acid ethyl ester (Example 2q)


EXAMPLE 4

Ester Hydrolysis

General Procedure


A solution of 4-(substituted-pyrrolidin-1-yl)-2-cyclopropylaminobenzoic acid ester (Example 3) and aqueous sodium hydroxide (10-20 eq., 1.0N) in tetrahydrofuran and methanol is heated for 16 to 24 hours at 70°C. The solution is cooled, diluted with water, and partially concentrated under vacuum. The resulting solution is acidified to pH 6 with aqueous 1.0N hydrochloric acid, then the pH is adjusted to pH 7-8 with 5% NaHCO₃ solution and extracted with ethyl acetate. The combined organic extracts are dried over Na₂SO₄, filtered, and concentrated in vacuo to afford 4-(substituted-pyrrolidin-1-yl)-2-cyclopropylaminobenzoic acid.

(a) 4-[(S)-3-*tert*-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,5-difluorobenzoic acid



A solution of 4-[(S)-3-*tert*-butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,5-difluorobenzoic acid ethyl ester (Example 3a, 1.40 g, 3.29 mmol) and aqueous sodium hydroxide (1.33 N, 15.0 mL, 20.0 mmol) in tetrahydrofuran (20 mL) and methanol (50 mL) is heated for 3 hours at 50°C. The solution is cooled, diluted with water, and partially concentrated in vacuo. The resulting solution is acidified to pH 6 with aqueous 1N hydrochloric acid, and then the pH is adjusted to pH 7-8 with 5% NaHCO₃ and extracted with chloroform. The combined organic extracts are dried over Na₂SO₄, filtered, and concentrated to give the title compound as a pale yellow solid (1.31 g). ¹H NMR (DMSO-*d*₆): δ 12.65 (bs, 1H), 7.48 (bs, 1H), 7.27-7.14 (m, 2H), 4.08-3.92 (m, 1H), 3.85-3.45 (m, 3H), 3.45-3.29 (m, 1H), 2.86-2.74 (m, 1H), 2.10-1.91 (m, 1H), 1.89-1.72 (m, 1H), 1.39 (s, 9H), 0.71-0.60 (m, 2H), 0.47-0.38 (m, 2H).

(b) 4-[(S)-3-*tert*-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,6-difluorobenzoic acid




A solution of 4-[(S)-3-*tert*-butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,6-difluorobenzoic acid ethyl ester (Example 3b, 1.45 g, 3.40 mmol) and aqueous sodium hydroxide (1.33N, 17.0 mL, 22.6 mmol) in tetrahydrofuran (20 mL) and methanol (50 mL) is heated for 21 hours at 50°C.

-104-

The solution is cooled, diluted with water, and then partially concentrated in vacuo. The resulting solution is acidified to pH 6 with aqueous 1N hydrochloric acid, and then the pH is adjusted to pH 7-8 with 5% NaHCO₃ and extracted with ethyl acetate three times. The combined organic extracts are dried over Na₂SO₄, filtered, and concentrated under vacuum to give the title compound as a pale green solid (1.33 g). ¹H NMR (DMSO-*d*₆): δ 7.21 (bd, 1H), 5.83 (dd, 1H), 4.12-3.98 (m, 1H), 3.75-3.20 (m, 5H), 2.88-2.78 (m, 1H), 2.14-1.96 (m, 1H), 1.90-1.73 (m, 1H), 1.39 (s, 9H), 0.69-0.59 (m, 2H), 0.45-0.36 (m, 2H).


(c) **4-[(S)-3-*tert*-Butoxycarbonylamino**pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluoro-benzoic acid



To a solution of 4-[(S)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoic acid ethyl ester (Example 3c, 2.90 g, 6.56 mmol) in methanol (30 mL) and tetrahydrofuran (30 mL) is added a 1N solution of sodium hydroxide (50 mL). The reaction mixture is stirred at 80°C for 20 hours, cooled to room temperature, and diluted with ethyl acetate. The organic layer is washed with 1N hydrochloric acid, water, and brine. The organic layer is dried over MgSO₄, filtered, and filtrate concentrated to the title compound as a solid (2.74 g). MS CI: *m/z* 414 (MH⁺).


-105-

(d) 4-[(S)-3-*tert*-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,5,6-trifluorobenzoic acid



A solution of benzyl 4-[(S)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3,5,6-trifluorobenzoate (Example 3d, 3.68 g, 7.31 mmol), 1.0 M NaOH (75 mL), methanol (50 mL), and tetrahydrofuran (50 mL) is heated at 50°C for 24 hours. The mixture is partially concentrated under vacuum and then extracted with dichloromethane. The aqueous phase is brought to pH 7.5 using 1.0 M HCl and extracted with ethyl acetate. The combined ethyl acetate extracts are dried over Na₂SO₄, filtered, and concentrated under vacuum to afford 4-[(S)-3-*tert*-butoxycarbonylamino-pyrrolidin-1-yl]-2-cyclopropylamino-3,5,6-trifluorobenzoic acid as a solid (2.9 g). This material is used without further purification. ¹H NMR (DMSO-d₆): δ 7.19 (bd, 1H), 4.12-3.95 (m, 1H), 3.83-3.52 (m, 3H), 3.48-3.33 (m, 1H), 2.83-2.69 (m, 1H), 2.10-1.94 (m, 1H), 1.89-1.72 (m, 1H), 1.49 (s, 9H), 0.70-0.56 (m, 2H), 0.44-0.33 (m, 2H).

(e) 2-Cyclopropylamino-5-fluoro-4-pyrrolidin-1-ylbenzoic acid




To a solution of 4-(pyrrolidin-1-yl)-2-cyclopropylamino-5-fluorobenzoic acid ethyl ester (Example 3e, 3.10 g, 10.4 mmol) in methanol (50 mL) and tetrahydrofuran (20 mL) is added 1N solution of sodium hydroxide (35 mL). The reaction mixture is stirred at 80°C for 20 hours, cooled to room temperature, and diluted with ethyl acetate. The organic layer is washed with 1N hydrochloric acid,

-106-


water, and brine. The organic layer is dried over MgSO_4 , filtered, and the filtrate concentrated to afford the title compound as a solid (2.47 g). MS CI: m/z 263 (M^+).

5 (f) 4-[3-*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-5-fluoro-2-isopropylaminobenzoic acid



10 A solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-5-fluoro-2-isopropylaminobenzoic acid ethyl ester (Example 3f, 1.5 g, 3.40 mmol), aqueous 1 M sodium hydroxide (50 mL), tetrahydrofuran (40 mL), and methanol (40 mL) are heated for 25 hours at 70°C. The solution is cooled, diluted with water, and then partially concentrated. The resulting solution is acidified to pH 7-8 with aqueous 1.0 M hydrochloric acid and extracted with ethyl acetate. The combined organic extracts are dried over Na_2SO_4 , filtered, and concentrated under vacuum to give the title compound as a solid (1.25 g). MS EI: m/z 380 (M^+).

15 (g) 4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-*sec*-butylamino-3-chloro-5-fluorobenzoic acid




20 A solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-*sec*-butylamino-3-chloro-5-fluorobenzoic acid ethyl ester (Example 3g, 3.57 g, 7.79 mmol), 1.0N sodium hydroxide (50 mL), and methanol (80 mL) are heated at 80°C for 16.5 hours. The mixture is partially evaporated, diluted with water (150 mL), and the pH adjusted to 7.0-8.0 using 1.0 M hydrochloric acid. The

-107-


mixture is extracted with ethyl acetate, dried over Na_2SO_4 , and concentrated under vacuum to give the title compound as an off-white solid (3.26 g) which is used without further purification. MS EI: m/z 430 (M^+).

5 (h) 4-[3-*tert*-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoic acid



A solution of 4-[3-*tert*-butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoic acid ethyl ester (Example 3h, 0.173 g, 0.395 mmol) and aqueous sodium hydroxide (1.33N, 2.0 mL, 2.66 mmol) in tetrahydrofuran (10 mL) and methanol (30 mL) is heated for 22 hours at 70°C. The solution is cooled, diluted with water, and partially concentrated under vacuum. The resulting solution is acidified to pH 6 with aqueous 1N hydrochloric acid, and then the pH is adjusted to pH 7-8 with 5% NaHCO_3 solution and extracted with ethyl acetate. The combined organic extracts are dried over Na_2SO_4 , filtered, and concentrated in vacuo to afford the title compound as a yellow solid (0.160 g). ^1H NMR (CDCl_3): δ 7.46 (d, 1H), 4.80-4.69 (bs, 1H), 4.33-4.18 (m, 1H), 3.90-3.45 (m, 3H), 3.62 (s, 3H), 3.43-3.32 (m, 1H), 2.91-2.80 (m, 1H), 2.28-2.10 (m, 1H), 1.96-1.78 (m, 1H), 1.46 (s, 9H), 0.63-0.52 (m, 4H).


20 (i) 4-[3-*tert*-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclobutylamino-5-fluorobenzoic acid



-108-

A solution of 4-[3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclobutylamino-5-fluorobenzoic acid ethyl ester (Example 3i, 1.51 g, 3.58 mmol) and aqueous sodium hydroxide (1.33N, 20.0 mL, 26.6 mmol) in tetrahydrofuran (15 mL) and methanol (40 mL) is refluxed for 19 hours. The solution is cooled, diluted with water, and partially concentrated under vacuum. The resulting solution is acidified to pH 6 with aqueous 1.0N hydrochloric acid, and then the pH is adjusted to pH 7-8 with 5% NaHCO₃ solution and extracted with ethyl acetate. The combined organic extracts are dried over Na₂SO₄, filtered, and concentrated under vacuum to afford the title compound as a yellow solid (1.353 g). MS EI: *m/z* 394 (MH⁺).


(i) **4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3-fluorobenzoic acid**



A solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3-fluorobenzoic acid ethyl ester (Example 3j, 1.34 g, 3.29 mmol) and aqueous sodium hydroxide (2N, 20 mL) in tetrahydrofuran (20 mL) and methanol (20 mL) is refluxed for 1 hour. The solution is partially concentrated in vacuo, then acidified to pH 6 and extracted with chloroform. The combined organic extracts are dried over Na₂SO₄, filtered, and concentrated to give the title compound (1.40 g). ¹H NMR (CDCl₃): δ 7.59 (dd, 1H), 6.06 (dd, 1H), 4.80-4.70 (bd, 1H), 4.38-4.22 (m, 1H), 3.79-3.50 (m, 3H), 3.47-3.35 (m, 1H), 2.98-2.87 (m, 1H), 2.30-2.14 (m, 1H), 1.97-1.84 (m, 1H), 1.46 (s, 9H), 0.69-0.54 (m, 4H).

-109-

(k) 4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-5-chloro-2-cyclopropylamino-3-fluorobenzoic acid



5 A solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-5-chloro-2-cyclopropylamino-3-fluorobenzoic acid ethyl ester (Example 3k, 2.00 g, 4.50 mmol) and aqueous sodium hydroxide (2.0N, 20 mL) in tetrahydrofuran (20 mL), and methanol (20 mL) is refluxed for 1 hour. The solution is partially concentrated in vacuo, acidified to pH 6, and extracted with chloroform. The combined organic extracts are dried over Na₂SO₄, filtered, and concentrated to
10 give the title compound (1.70 g). ¹H NMR (CDCl₃): δ 7.71 (d, 1H), 4.91-4.82 (bd, 1H), 4.40-4.25 (m, 1H), 3.77-3.69 (m, 2H), 3.53-3.34 (m, 2H), 3.00-2.88 (m, 1H), 2.24-2.15 (m, 1H), 1.90-1.80 (m, 1H), 1.46 (s, 9H), 0.72-0.54 (m, 4H).

15 The following compounds are synthesized according to general procedures of Examples 4a-k.

(l) 4-(3-*tert*-Butoxycarbonylamino)pyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-3-methylbenzoic acid ([MS ES, MH⁺] m/z 394.) using 4-(3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-3-methylbenzoic acid ethyl ester (Example 3l).

20 (m) 4-[3-(*tert*-Butoxycarbonylamino)methyl]pyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-3-methylbenzoic acid ([MS ES, M+1] m/z 408) using 4-[3-(*tert*-butoxycarbonylamino)methyl]pyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-3-methylbenzoic acid ethyl ester (Example 3m).

25 (n) 4-(3-*tert*-Butoxycarbonylamino)pyrrolidin-1-yl)-2-cyclopropylamino-3-fluoro-5-methoxybenzoic acid (¹H NMR (200 MHz, CDCl₃): δ 12.50 (bs, 1H), 7.18-7.09 (m, 1H), 7.07 (d, 1H), 4.07-3.90 (m, 1H),

-110-

3.80-3.50 (m, 2H), 3.66 (s, 3H), 3.42-3.29 (m, 2H), 2.88-2.71 (m, 1H), 2.10-1.91 (m, 1H), 1.86-1.67 (m, 1H), 1.39 (s, 9H), 0.70-0.57 (m, 2H), 0.46-0.32 (m, 2H)) using 4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-3-fluoro-5-methoxy-benzoic acid ethyl ester (Example 3n).

5 (o) 4-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoic acid. ([MS ES, M+1] m/z 424) using 4-[3-(*tert*-butoxycarbonylaminomethyl)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoic acid ethyl ester (Example 3o)

(p) 4-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-3-ethoxy-5-fluorobenzoic acid ([MS ES, MH⁺] m/z 424) using 4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-3-ethoxy-5-fluorobenzoic acid ethyl ester (Example 3p).

(q) 3-(Benzyloxyiminomethyl)-4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-5-fluorobenzoic acid (MS ES: m/z 513 (MH⁺)) using 3-(benzyloxyiminomethyl)-4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-5-fluorobenzoic acid ethyl ester (Example 3q)

(r) 3-Chloro-2-cyclopropylamino-5-fluoro-4-(3-[1,2,3-triazol]-1-yl-pyrrolidin-1-yl)benzoic acid (¹H NMR (200 MHz, CDCl₃): δ 8.50 (bs, 1H), 8.03 (s, 1H), 7.72 (s, 1H), 7.60 (d, 1H), 7.54-5.39 (m, 1H), 4.13-3.97 (m, 1H), 3.92-3.85 (m, 1H), 3.68-3.63 (m, 1H), 3.54-3.42 (m, 1H), 3.10-3.00 (m, 1H), 2.78-2.60 (m, 1H), 2.39-2.33 (m, 1H), 0.70-0.58 (m, 2H), 0.58-0.46 (m, 2H)) using 3-chloro-2-cyclopropylamino-5-fluoro-4-(3-[1,2,3-triazol]-1-yl-pyrrolidin-1-yl)benzoic acid ethyl ester (Example 3r).


25

EXAMPLE 5

Synthesis of N-Substituted Anthranilic Acids


(a) 2-Cyclopropylamino-4,5-difluorobenzoic acid

-111-



To a solution of 4,5-difluoroanthranilic acid (1.13 g, 6.53 mmol) in anhydrous methanol (40 mL) is added molecular sieves (3 Å), acetic acid (3.70 mL, 65.3 mmol) and [(1-ethoxycyclopropyl)oxy]trimethylsilane (5.25 mL, 26.11 mmol). After 30 minutes, sodium cyanoborohydride (2.08 g, 32.64 mmol) is added, and the reaction mixture is heated to reflux. After 16 hours, the reaction is cooled to room temperature, filtered, washed with methanol, and the combined filtrate concentrated under vacuum to afford a viscous oil. The resulting oil is dissolved in ethyl acetate and washed with 1.0 M hydrochloric acid, water, and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate is concentrated under vacuum to afford as a beige solid (1.32 g). MS CI: *m/z* 212 (M⁺).

(b) *5-Chloro-2-cyclopropylamino-4-fluorobenzoic acid.*




To a solution of 4,5-difluoroanthranilic acid (1.20 g, 6.37 mmol) in anhydrous methanol (50 mL) is added molecular sieves (3 Å), acetic acid (3.70 mL, 65.3 mmol), and [(1-ethoxycyclopropyl)oxy]trimethylsilane (5.12 mL, 25.5 mmol). After 30 minutes, sodium cyanoborohydride (2.03 g, 31.9 mmol) is added, and the reaction mixture is heated to reflux. After 16 hours, the reaction is cooled to room temperature, filtered, washed with methanol, and the combined filtrate concentrated under vacuum to afford a viscous oil. The resulting oil is dissolved in ethyl acetate and washed with 1.0 M hydrochloric acid, water, and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate is

-112-


concentrated under vacuum to afford as a beige solid (1.78 g). MS CI: m/e 228 (M^+).

(c) *2-(2,4-Difluoroanilino)-4,5-difluorobenzoic acid*



5 Lithium diisopropylamide is generated at -5°C by combining
diisopropylamine (7.2 mL, 51 mmol) and *n*-butyllithium (33 mL, 53 mmol) in
anhydrous tetrahydrofuran (150 mL) under an inert atmosphere. After 0.5 hour,
the solution is cooled to -78°C and 2,4-difluoroaniline (3.46 mL, 34 mmol) is
added and stirred for 2 hours. 2,4,5-Trifluorobenzoic acid (3.0 g, 17 mmol) is
10 added, and the mixture is subsequently allowed to warm to room temperature over
18 hours. A saturated solution of hydrogen chloride/dioxane (10 mL) is added,
and after 1 hour, the mixture is concentrated to a solid. The solid is dissolved in
chloroform and washed with 1.0 M hydrochloric acid, water, and brine. The
solution is dried, concentrated in vacuo, and purified by column chromatography
15 (3:1 ethyl acetate/hexanes) to give 2-(2,4-difluoroanilino)-4,5-difluorobenzoic
acid as a solid (2.18 g). ^1H NMR (CDCl_3): δ 8.95 (bs, 1H), 7.80-7.75 (m, 2H),
7.50-6.85 (m, 3H), 6.62-6.40 (m, 1H). MS EI: m/z 286 (M^+).

(d) *3-Chloro-2-cyclopropylamino-4,5-difluorobenzoic acid*




-113-

In a sealed tube a mixture of 2-bromo-3-chloro-4,5-difluorobenzoic acid (Example 20, 7.96 g, 29.3 mmol), cyclopropyl amine (4.20 mL, 58.7 mmol), potassium acetate (5.77 g, 58.6 mmol), cupric acetate monohydrate (0.50 g, 2.5 mmol), and triethylamine (4.9 mL, 35.19 mmol) in isopropyl alcohol is stirred at 80°C. After 16 hours, the reaction mixture is concentrated under vacuum, and the resulting residue is dissolved in ethyl acetate. The organic layer is washed with 1.0 M hydrochloric acid, water, and brine. The organic layer is dried over MgSO₄ and filtered. The filtrate is concentrated under vacuum and purified via flash column chromatography (5% isopropyl alcohol/1% formic acid/94% dichloromethane) to afford the title compound (4.62 g). MS CI: m/z 248 (MH⁺).

EXAMPLE 6

*Synthesis of Hydrazinecarboxylic *t*-Butyl Esters*


(a) *(2-Cyclopropylamino-4,5-difluorobenzoyl)hydrazinecarboxylic acid tert-butyl ester*



To a solution of 2-cyclopropylamino-4,5-difluorobenzoic acid (Example 5) (1.32 g, 6.19 mmol) and *tert*-butyl carbazate (1.30 g, 9.75 mmol) in dichloromethane (30 mL) is added 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (1.87 g, 9.75 mmol). After 16 hours, the reaction mixture is diluted with dichloromethane and washed with saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄ and filtered. The filtrate is concentrated under vacuum and purified via flash column chromatography (1:2 ethyl acetate/hexanes) to afford N₂-(2-cyclopropylamino-4,5-difluorobenzoyl)hydrazinecarboxylic acid *tert*-butyl ester as a solid (1.65 g). MS EI: m/z 328 (M⁺).


-114-

(b) **[2-(2,4-Difluoroanilino)-4,5-difluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester**



To a solution of 2-(2,4-difluoroanilino)-4,5-difluorobenzoic acid
 5 (Example 5, 2.18 g, 7.6 mmol) and *tert*-butyl carbazate (1.57 g, 11.8 mmol) in
 dichloromethane (30 mL) is added 1-ethyl-3-(3-dimethylaminopropyl)-
 carbodiimide (2.25 g, 11.77 mmol). After stirring for 16 hours at room
 temperature, the reaction mixture is diluted with dichloromethane and then
 washed with saturated NaHCO₃, water, and brine. The organic layer is dried over
 10 Na₂SO₄, concentrated under vacuum, and purified via column chromatography
 (1:3 ethyl acetate/hexanes) to afford [2-(2,4-difluoroanilino)-4,5-difluorobenzoyl]-
 hydrazinecarboxylic acid *tert*-butyl ester as a solid (2.20 g). MS EI: *m/z* 400 (M⁺).

(c) **(3-Chloro-2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester**




To a solution of 3-chloro-2-cyclopropylamino-4,5-difluorobenzoic acid
 15 (Example 5d, 2.09 g, 8.45 mmol) in dichloromethane (30 mL) is added *tert*-butyl
 carbazate (1.67 g, 12.7 mmol) and 1-ethyl-3-(3-dimethylaminopropyl)-
 carbodiimide (2.43 g, 12.7 mmol). After 16 hours, the reaction mixture is diluted

-115-

with dichloromethane and washed with saturated NaHCO_3 , water, and brine. The organic layer is dried over MgSO_4 and filtered. The filtrate is concentrated under vacuum and purified via flash column chromatography (1:2 ethyl acetate/hexanes) to afford the title compound (1.93 g). MS EI: m/z 360 (M^+).

- 5 (d) (5-Chloro-2-cyclopropylamino-4-fluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester



- 10 To a solution of 5-chloro-2-cyclopropylamino-4-fluorobenzoic acid (Example 5, 1.48 g, 6.46 mmol) in dichloromethane (50 mL) is added *tert*-butyl carbazate (1.28 g, 9.69 mmol) and 1-ethyl-3-(3-dimethylaminopropyl)carbo-
- 15 diimide (1.86 g, 9.69 mmol). After 16 hours, the reaction mixture is diluted with dichloromethane and washed with saturated NaHCO_3 , water, and brine. The organic layer is dried over MgSO_4 and filtered. The filtrate is concentrated under vacuum and purified via flash column chromatography (1:2 ethyl acetate/hexanes) to afford (3-chloro-2-cyclopropylamino-4,5-difluorobenzoyl)hydrazinecarboxylic acid *tert*-butyl ester as a solid (1.04g). MS CI: m/e 344 (M^++1).

EXAMPLE 7

Synthesis of 4-Heterocyclic-benzoylhydrazinecarboxylic acid t-butyl esters

General Procedure A

- 20 A solution of a substituted benzoylhydrazinecarboxylic acid *tert*-butyl ester from Example 6 (where $\text{R}_7 = \text{F}$), heterocyclic amine (2 eq.), and triethylamine (10 eq.) is stirred in dimethyl sulfoxide or *N,N*-dimethylformamide (6 mL) at 120°C in a sealed tube for 16 hours. After cooling to room temperature, the reaction mixture is diluted with ethyl acetate and washed with saturated
- 25 NaHCO_3 , water, and brine. The organic layer is dried over MgSO_4 , filtered, and

-116-

the filtrate concentrated to afford a brown residue. Trituration with diethyl ether or purification via flash column chromatography (ethyl acetate/hexanes) yields the product.

General Procedure B

- 5 To a suspension of a benzoic acid of Example 4 (3.30 mmol) and *tert*-butyl carbazate (1.5 eq.) in anhydrous dichloromethane is added
1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (1.5 eq.). The mixture is stirred at room temperature for 20 hours, poured into saturated NaHCO₃, and extracted with dichloromethane. The combined organic extracts are
10 washed with brine, dried over Na₂SO₄, filtered, and concentrated in vacuo. Purification by flash column chromatography (ethyl acetate/hexanes) afford the title compounds.

The following compounds are synthesized according to General Procedures A or B above:

- 15 (a) *N'*-{4-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS CI: *m/z* 508 (MH⁺)) using (2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6a) and 3-(*tert*-butoxycarbonylaminomethyl)pyrrolidine (General Method A).
- 20 (b) *N'*-{4-[4-(*tert*-Butoxycarbonylpiperazin-1-yl)-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (MS CI: *m/z* 494 (MH⁺)) using (2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6a) and 4-*tert*-butoxycarbonylpiperazine (General Method A).
- 25 (c) *N'*-{4-[3-(*tert*-Butoxycarbonylaminomethyl)-3-methylpyrrolidin-1-yl]-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS CI: *m/z* 522 (MH⁺)) using (2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6a) and 3-(*tert*-butoxycarbonylaminomethyl)-3-methylpyrrolidine (Sanchez J.P., Bridges A.J.,
30 Bucsh R., Domagala J.M., Gogliotti R.D., Hagen S.E., Heifetz C.L.,

Joannides E.T., Sesnie J.C., et al. *J. Med. Chem.*, 1992;35(2):361-367) (General Method A).

(d) $N^{\prime}$ -[4-(6-*tert*-Butoxycarbonylamino-3-azabicyclo[3.1.0]hex-3-yl)-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/z 506 (MH^{+})) using (2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6a) and (3-azabicyclo[3.1.0]hex-6-yl)carbamic acid *tert*-butyl ester (Brighty K.E., 1991, EP 413455) (General Method A).

(e) $N^{\prime}$ -{4-[(S)-3-(*tert*-Butoxycarbonyl-*N*-methylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/z 508 (MH^{+})) using (2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6a) and (S)-3-(*tert*-butoxycarbonyl-*N*-methylamino)pyrrolidine (Chu D.T., Li Q., Cooper C.S., Fung A.K.L., Lee C.M., Plattner J.J., PCT Int. Appl., 1995:255. WO 9510519) (General Method A).

(f) $N^{\prime}$ -{4-[(S)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-(2,4-difluoroanilino)-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/z 566 (MH^{+})) using [2-(2,4-difluoroanilino)-4,5-difluorobenzoyl]-hydrazinecarboxylic acid *tert*-butyl ester (Example 6b) and (S)-pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (General Method A).

(g) $N^{\prime}$ -{4-[(S)-3-*tert*-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,5-difluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester MS CI: m/z 538 (MH^{+}) using 4-[(S)-3-*tert*-butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,5-difluorobenzoic (Example 4a) (General Method B).

(h) $N^{\prime}$ -{4-[(S)-3-*tert*-Butoxycarbonylamino]pyrrolidin-1-yl]-2-cyclopropylamino-3,6-difluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (^1H NMR ($CDCl_3$): δ 8.08 (bs, 2H), 6.50 (bs, 1H), 5.70 (dd, 1H), 4.76-4.68 (m, 1H), 4.37-4.23 (m, 1H), 3.78-3.68 (m, 1H), 3.65-3.42 (m, 2H), 3.40-3.30 (m, 1H), 2.96-2.83 (m, 1H), 2.28-2.14 (m, 1H), 2.00-1.84 (m, 1H), 1.49 (s, 9H), 1.46 (s, 9H), 0.68-0.61 (m, 2H), 0.58-0.48 (m, 2H)) using 4-[(S)-3-*tert*-

butoxycarbonylaminopyrrolidin-1-yl]-2-cyclopropylamino-3,6-difluorobenzoic acid (Example 4b) (General Method B).

(i) N' -{4-[(S)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3,5,6-trifluorobenzoyl}hydrazine carboxylic acid *tert*-butyl ester (^1H NMR (CDCl_3): δ 8.04 (bd, 1H), 7.55 (bs, 1H), 6.52 (bs, 1H), 4.79-4.67 (m, 1H), 4.37-4.19 (m, 1H), 3.97-3.55 (m, 3H), 3.53-3.40 (m, 1H), 2.90-2.75 (m, 1H), 2.29-2.06 (m, 1H), 1.96-1.75 (m, 1H), 1.49 (s, 9H), 1.46 (s, 9H), 0.69-0.57 (m, 2H), 0.54-0.43 (m, 2H)) using 4-[(S)-3-*tert*-butoxycarbonylaminopyrrolidin-1-yl]-2-cyclopropylamino-3,5,6-trifluorobenzoic acid (Example 4d) (General Method B).

(j) N' -{4-[(S)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic *tert*-butyl ester (MS CI: m/z 528 (MH^+)) using 4-[(S)-3-*tert*-butoxycarbonylaminopyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoic acid (Example 4c) (General Method B).

(k) N' -[(S)-4-(7-*tert*-Butoxycarbonylamino-5-azaspiro[2.4]hept-5-yl)-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/z 520 (MH^+)) using (2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6a) and (5-azaspiro[2,4]hept-7-yl)carbamic acid *tert*-butyl ester (General Method A).

(l) N' -{4-[(R)-3-(1-*tert*-Butoxycarbonylamino-1-methylethyl)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/z 570 (MH^+)) using (3-chloro-2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6c) and [(R)-3-(1-*tert*-butoxycarbonylamino-1-methylethyl)pyrrolidine (Fedij V., Lenoir E.A., III, Suto M.J., Zeller J.R., Wemple J., *Tetrahedron: Asymmetry*, 1994;5[7]:1131-1134) (General Method A).

(m) N' -[4-(Pyrrolidin-1-yl)-2-cyclopropylamino-5-fluorobenzoyl]-hydrazinecarboxylic *tert*-butyl ester (MS CI: m/z 379 (MH^+)) using

2-cyclopropylamino-5-fluoro-4-pyrrolidin-1-ylbenzoic acid (Example 4e)
(General Method B).

(n) N' -{4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-5-fluoro-2-isopropylaminobenzoyl}hydrazine carboxylic acid *tert*-butyl ester (MS EI: m/z 496 (MH^+)) using 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-5-fluoro-2-isopropylaminobenzoic acid (Example 4f) (General Method B).

(o) N' -{4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-*sec*-butylamino-3-chloro-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS EI: m/z 544 (MH^+)) using 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-*sec*-butylamino-3-chloro-5-fluorobenzoic acid (Example 4g) (General Method B).

(p) N' -{4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS EI: m/z 524 (MH^+)) using 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoic acid (Example 4h) (General Method B).

(q) N' -{4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclobutylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS EI: m/z 394 (M^+)) using 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclobutylamino-5-fluorobenzoic acid (Example 4i) (General Method B).

(r) N' -{4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS EI: m/z 494 (MH^+)) using 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3-fluorobenzoic acid (Example 4j) (General Method B).

(s) N' -{4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-5-chloro-2-cyclopropylamino-3-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (^1H NMR ($CDCl_3$): δ 8.77 (bs, 1H), 7.35 (d, 1H), 6.82 (bs, 1H), 6.55 (bs, 1H), 5.01-4.97 (m, 1H), 4.29 (bs, 1H), 3.74-3.57 (m, 2H), 3.41-3.24 (m, 2H), 2.84-2.75 (m, 1H), 2.33-2.16 (m, 1H), 1.92-1.72 (m, 1H), 1.47 (s, 9H), 1.45 (s, 9H), 0.69-0.62 (m, 2H), 0.60-0.50 (m, 2H)) using 4-[3-(*tert*-

-120-

butoxycarbonylamino)pyrrolidin-1-yl]-5-chloro-2-cyclopropylamino-3-fluorobenzoic acid (Example 4k) (General Method B).

(t) N' -[4-(3-*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methylbenzoyl]hydrazinecarboxylic acid *tert*-butyl ester ([MS ES, MH^+] m/z 508) using 4-(3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methylbenzoic acid (Example 4l) (General Method B).

(u) N' -{4-[3-(*tert*-Butoxycarbonylamino)methyl]pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methylbenzoyl}hydrazinecarboxylic acid *tert*-butyl ester ([MS ES, MH^+] m/z 522) using 4-[3-(*tert*-butoxycarbonylamino)methyl]-pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methylbenzoic acid (Example 4m) (General Method B).

(v) N' -[4-(3-*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3-fluoro-5-methoxybenzoyl]hydrazinecarboxylic acid *tert*-butyl ester. ([MS ES, MH^+] m/z 525) using 4-(3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3-fluoro-5-methoxybenzoic acid (Example 4n) (General Method B).

(w) N' -{4-[3-(*tert*-Butoxycarbonylamino)methyl]pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoyl}hydrazinecarboxylic acid *tert*-butyl ester ([MS ES, MH_2^{+2}] m/z 539) using 4-[3-(*tert*-butoxycarbonylamino)methyl]pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoic acid (Example 4o) (General Method B).

(x) N' -[3-(Benzyloxyiminomethyl)-4-(3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester. ([MS ES, MH^+] m/z 627) using 3-(benzyloxyiminomethyl)-4-(3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluorobenzoic acid (Example 4q) (General Method B)

(y) N' -[3-Chloro-2-cyclopropylamino-5-fluoro-4-(3-[1,2,3-triazol]-1-yl)pyrrolidin-1-yl]benzoyl]hydrazinecarboxylic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 9.90 (bs, 1H), 7.97 (s, 1H), 7.72 (s, 1H), 7.52 (d, 1H), 7.17

-121-

(bs, 1H), 5.51-5.41 (m, 1H), 4.59 (bs, 1H), 4.06-3.95 (m, 1H), 3.87-3.74 (m, 1H), 3.61-3.55 (m, 1H), 3.48-3.36 (m, 1H), 2.97-2.90 (m, 1H), 2.77-2.59 (m, 1H) 2.42-2.31 (m, 1H), 1.46 (s, 9H), 0.68-0.50 (m, 4H)) using 3-chloro-2-cyclopropylamino-5-fluoro-4-(3-[1,2,3-triazol]-1-yl)pyrrolidin-1-yl)benzoic acid (Example 4r) (General Method B).

(z) *N*-(1-{4-[3-(*tert*-Butoxycarbonylaminomethyl)-piperidin-1-yl]-2-cyclopropylamino-5-fluorophenyl}methanoyl)hydrazinecarboxylic acid *tert*-butyl ester (MS CI: *m/e* 522 (MH⁺)) from piperidin-3-ylmethylcarbamic acid *tert*-butyl ester (Hilpert K., Ackermann J., Banner D.W., Gast A., Gubernator K., Hadvary P., Labler L., Mueller K., Schmid G., et al., *J. Med. Chem.*, 1994;37[23]:3889-3901) and (2-Cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6a) (General Method A).

(aa) *N*-[1-(4-{3-[(*tert*-Butoxycarbonylisopropylamino)-methyl]pyrrolidin-1-yl}-2-cyclopropylamino-5-fluorophenyl)methanoyl]-hydrazinecarboxylic acid *tert*-butyl ester (MS CI: *m/e* 550 (MH⁺)) using isopropylpyrrolidin-3-ylmethylamine (Mich T.F., Culbertson T.P., US 4550103) and (2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6a) (General Method B).

(bb) *N'*-(1-{4-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-5-chloro-2-cyclopropylaminophenyl}methanoyl)hydrazinecarboxylic acid *tert*-butyl ester (MS CI: *m/e* 524 [M⁺+1]) using (3-chloro-2-cyclopropylamino-4,5-difluorobenzoyl)hydrazinecarboxylic acid *tert*-butyl ester (Example 6c) and 3-(*tert*-butoxycarbonylaminomethyl)pyrrolidine (General Method A).

(cc) *N'*-[4-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-3-ethoxy-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (MS ES: *m/z* 537 (MH⁺)) from (4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-3-ethoxy-5-fluorobenzoic acid (Example 4p) (General Method A).

EXAMPLE 8

Synthesis of (2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid tert-butyl esters

General Method A

5 To a solution of a benzoylhydrazinecarboxylic acid *tert*-butyl ester (from Example 7 (0.435 mmol) in tetrahydrofuran (15 mL) is added potassium carbonate (0.300 g, 2.18 mol) and triphosgene (0.167 g, 0.566 mmol). The reaction mixture is refluxed for 90 minutes, cooled to room temperature, and diluted with ethyl acetate. The organic layer is washed with water and brine, then dried over
10 MgSO₄, and filtered. The filtrate is concentrated under vacuum and purified by flash column chromatography (ethyl acetate/hexanes) to afford the title compounds.

General Method B

A solution of (2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid
15 *tert*-butyl ester (such as, for example, Example 14), pyrrolidine side chain (2 eq.), and triethylamine (10 eq.) are stirred in dimethyl sulfoxide or acetonitrile (3 mL) at 80°C for 48 hours. After cooling to room temperature, the reaction mixture is diluted with ethyl acetate and washed with saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄, filtered, and concentrated. The resulting
20 residue is purified via flash column chromatography (ethyl acetate/hexanes) to afford the title compounds.

General Method C

To a solution of a C-6- and/or C-8-halogenated-(2,4-dioxo-1,4-dihydro-
2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester in ethanol and tetrahydrofuran is
25 added triethylamine and 20% palladium on carbon. Hydrogen is introduced to the reaction mixture at high pressure for several days, then the reaction mixture is filtered through Celite, washed with ethanol, and the combined filtrate concentrated in vacuo and purified via flash column chromatography (ethyl acetate/hexanes) to afford the title compounds.

-123-

General Method D

5 A suspension of benzoylhydrazinecarboxylic acid *tert*-butyl ester (from Example 7) and triethylamine (10 eq.) are stirred in anhydrous tetrahydrofuran for 5 minutes. Phosgene (20% solution in toluene, 2 eq.) is added in one portion to the suspension. The reaction mixture is heated at 60°C for 4.5 hours, cooled to room temperature, poured into saturated NaHCO₃ solution, and extracted with chloroform. The combined organic extracts are washed with water, brine, and then dried over Na₂SO₄, filtered, and concentrated under vacuum. The resulting residue is purified by flash column chromatography (1:1 ethyl acetate/hexanes) to afford the title compounds.

10

General Method E

An appropriately substituted pyrrolidine (1.5-2 eq.), a 3-dibenzylamino-6, 7-difluoro-1-substituted-1H-quinazoline-2,4-dione (Example 16) (1 eq.), and triethylamine (10 eq.) are combined in acetonitrile and heated to reflux for 15 72 hours. The solution is cooled, concentrated, and re-dissolved in chloroform. The organic solution is washed with 1.0 N hydrochloric acid, saturated NaHCO₃, brine, and dried over magnesium sulfate. The solution is concentrated to give the title compounds.

General Method F

20 A solution of (2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (such as, for example, Example 14), pyrrolidine side chain (2 eq.), and triethylamine (10 eq.) are stirred in dimethyl sulfoxide or acetonitrile (3 mL) at 80°C for 20 to 40 hours. After cooling to room temperature, the reaction mixture is diluted with ethyl acetate and washed with saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄, filtered, and concentrated. The resulting residue is redissolved in methylene chloride and then treated with di-*tert*-butyl dicarbonate (2 eq.). After 1 hour, the reaction mixture is concentrated and the product purified via flash column chromatography (ethylacetate/hexanes) to afford the title compounds.

25

The following compounds are synthesized according to General Procedures A-F of Example 8:

5 (a) {7-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (MS CI: *m/z* 568 (MH⁺)) from *N'*-{4-[3-(*tert*-butoxycarbonylaminomethyl)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 18a) (General Method A)

10 (b) 7-[4-(*tert*-Butoxycarbonylpiperazin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (MS CI: *m/z* 554 (MH⁺)) from *N'*-4-[4-(*tert*-butoxycarbonylpiperazin-1-yl)-3-chloro-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (Example 18b) (General Method A).

15 (c) {7-[3-(*tert*-Butoxycarbonylaminomethyl)-3-methylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS CI: *m/z* 582 (MH⁺)) from *N'*-{4-[3-(*tert*-butoxycarbonylaminomethyl)-3-methylpyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 18c) (General Method A).

20 (d) [7-(6-*tert*-Butoxycarbonylamino-3-azabicyclo[3.1.0]hex-3-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]-carbamic acid *tert*-butyl ester (MS CI: *m/z* 566 (MH⁺)) from *N'*-[4-(6-*tert*-Butoxycarbonylamino-3-azabicyclo[3.1.0]hex-3-yl)-3-chloro-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (Example 18d) (General Method A).

25 (e) {7-[(*S*)-3-(*tert*-Butoxycarbonyl-N-methylamino)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (MS CI: *m/z* 568 (MH⁺)) from *N'*-{4-[(*S*)-3-(*tert*-butoxycarbonyl-N-methylamino)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluoro-benzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 18e) (General Method A).

30

-125-

(f) {7-[(S)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-8-chloro-1-(2,4-difluoroanilino)-6-fluoro-1,4-dihydro-2,4-dioxo-2*H*-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (^1H NMR (CDCl_3): δ 7.45-7.38 (m, 1H), 6.97-6.81 (m, 1H), 6.73-6.66 (m, 1H), 6.54-6.42 (m, 1H), 4.90-4.81 (bs, 1H), 4.31-4.21 (bs, 1H), 3.86-3.36 (m, 4H), 2.32-2.04 (m, 1H), 1.96-1.84 (m, 1H), 1.61 (s, 9H), 1.45 (s, 9H)) from *N'*-{4-[(S)-3-(*tert*-butoxycarbonyl-N-amino)-pyrrolidin-1-yl]-3-chloro-2-(2,4-difluoroanilino)-5-fluorobenzoyl}-hydrazinecarboxylic acid *tert*-butyl ester (Example 18f) (General Method A).

(g) 7-[(S)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropylamino-6,8-difluoro-1,4-dihydro-2,4-dioxo-2*H*-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (^1H NMR (CDCl_3): δ 7.54 (dd, 1H), 6.67 (bs, 1H), 4.79-4.68 (bs, 1H), 4.39-4.23 (m, 1H), 4.10-3.65 (m, 3H), 3.58-3.45 (m, 1H), 3.37-3.25 (m, 1H), 2.30-2.11 (m, 1H), 2.03-1.86 (m, 1H), 1.50 (s, 9H), 1.47 (s, 9H), 1.20-1.12 (m, 2H), 0.83-0.74 (bs, 2H)) from *N'*-{4-[(S)-3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3,5-difluorobenzoyl}-hydrazinecarboxylic acid *tert*-butyl ester (Example 5g) (General Method A).

(h) {7-[(S)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropylamino-5,8-difluoro-1,4-dihydro-2,4-dioxo-2*H*-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (MS CI: m/z 538 (M^+)) from *N'*-{4-[(S)-3-*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3,6-difluorobenzoyl}-hydrazinecarboxylic acid *tert*-butyl ester (Example 5h) (General Method A).

(i) {7-[(S)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-1,4-dihydro-2,4-dioxo-5,6,8-trifluoro-2*H*-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (MS CI: m/z 556 (MH^+)) from *N'*-{4-[(S)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-3,5,6-trifluorobenzoyl}hydrazine carboxylic acid *tert*-butyl ester (Example 5i) (General Method A)

(j) {7-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS CI: m/z 552 (MH^+)) from *N'*-{4-[(S)-3-(*tert*-

butoxycarbonylamino)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic *tert*-butyl ester (Example 5j) (General Method A).

(k) [7-(3-Hydroxymethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (MS CI: *m/z* 467 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and 3-Hydroxymethylpyrrolidine (Roussi G., Zhang J., *Tetrahedron Lett.*, 1988;29(28):3481-3482) (General Method B).

(l) {7-[3-(*tert*-Butoxycarbonylaminoethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS CI: *m/z* 580 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and 3-(*tert*-butoxycarbonylaminoethyl)pyrrolidine (Antonsson K.T., Bylund R.E., Gustafsson N.D., Nilsson N.O.I. WO 9429336) (General Method B).

(m) [7-(7-*tert*-Butoxycarbonylamino-5-azaspiro[2.4]hept-5-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (MS CI: *m/z* 580 (MH^+)) from *N'*-[(*S*)-4-(7-*tert*-butoxycarbonylamino-5-azaspiro[2.4]hept-5-yl)-3-chloro-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (Example 18g) (General Method A).

(n) {7-[(*R*)-3-(1-*tert*-Butoxycarbonylamino-1-methylethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS CI: *m/z* 596 (MH^+)) from *N'*-{4-[(*R*)-3-(1-*tert*-butoxycarbonylamino-1-methylethyl)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 7l) (General Method A).

(o) {7-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS CI: *m/z* 534 (MH^+)) from {7-[3-(*tert*-butoxycarbonylaminomethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-

fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8a) (General Method C).

(p) [7-(Pyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (MS CI: *m/z* 437 (MH⁺)) using *N*'-[4-(Pyrrolidin-1-yl)-3-chloro-2-cyclopropylamino-5-fluoro-
5 benzoyl]hydrazinecarboxylic *tert*-butyl ester (Example 18h) (General Method A).

(q) {7-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-8-chloro-6-fluoro-1-isopropyl-2,4-dioxo-6-fluoro-2*H*-quinazolin-3-yl}carbamic acid *tert*-
10 butyl ester. (MS EI: *m/z* 530 (M⁺)) using *N*'-{4-[3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-3-chloro-5-fluoro-2-isopropylaminobenzoyl}-
10 hydrazinecarboxylic acid *tert*-butyl ester (Example 18i) (General Method A).

(r) {7-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-1-*sec*-butyl-8-chloro-6-fluoro-2,4-dioxo-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester
15 (MS EI: *m/z* 570 (MH⁺)) using *N*'-{4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-*sec*-butylamino-3-chloro-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-
15 butyl ester (Example 7o) (General Method A).

(s) {7-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid
20 *tert*-butyl ester (MS EI: *m/z* 550 (MH⁺)) using *N*'-{4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-
20 3-methoxybenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 7p) (General Method D).

(t) {7-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-8-chloro-1-cyclobutylamino-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic
25 acid *tert*-butyl ester (MS EI: *m/z* 568 (MH⁺)) using *N*'-{4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclobutylamino-5-fluorobenzoyl}-
25 hydrazinecarboxylic acid *tert*-butyl ester (Example 7q) (General Method A).

(u) {7-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-8-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester
30 (MS EI: *m/z* 520 (MH⁺)) using *N*'-{4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-

1-yl]-2-cyclopropylamino-3-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 7r) (General Method D).

(v) {7-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-6-chloro-1-cyclopropyl-8-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (¹H NMR (CDCl₃): δ 7.36 (d, 1H), 6.64 (bs, 1H), 4.86-4.80 (bd, 1H), 4.31-4.25 (m, 1H), 3.81-3.66 (m, 2H), 3.51-3.31 (m, 2H), 3.02-2.97 (m, 1H), 2.32-2.22 (m, 1H), 1.95-1.85 (m, 1H), 1.63 (s, 9H), 1.46 (s, 9H), 0.77-0.73 (m, 2H), 0.63-0.57 (bs, 2H)) using *N'*-(4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-5-chloro-2-cyclopropylamino-3-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 7s) (General Method D).

(w) [(*S*)-1-(1-Cyclopropylmethyl-3-dibenzylamino-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]carbamic acid *tert*-butyl ester (MS CI, *m/z* 614 (MH⁺)) using 3-dibenzylamino-6,7-difluoro-1-cyclopropylmethyl-1*H*-quinazoline-2,4-dione (Example 17a) and (*S*)-pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (General Method E).

(x) {1-[(*R*)-1-(Cyclopropylmethyl-3-dibenzylamino-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]-1-methylethyl}-carbamic acid *tert*-butyl ester (MS CI: *m/z* 656 (MH⁺)) using 3-dibenzylamino-6,7-difluoro-1-cyclopropylmethyl-1*H*-quinazoline-2,4-dione (Example 17a) and (*R*)-3-(1-*tert*-butoxycarbonylamino-1-methylethyl)pyrrolidine (General Method E).

(y) [(*S*)-1-(3-Dibenzylamino-1-ethyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]carbamic acid *tert*-butyl ester (mp 104-106°C, MS CI: *m/z* 588 (MH⁺)) using 3-dibenzylamino-6,7-difluoro-1-ethyl-1*H*-quinazoline-2,4-dione (Example 17b) and (*S*)-pyrrolidin-3-ylcarbamic acid *tert*-butyl ester (General Method E).

(z) [(*R*)-1-(3-Dibenzylamino-1-ethyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-1-methyl-1-pyrrolidin-3-ylethyl]carbamic acid *tert*-butyl ester (MS CI: *m/z* 630 (MH⁺)) using 3-dibenzylamino-6,7-difluoro-1-ethyl-1*H*-quinazoline-2,4-dione (Example 17b) and (*R*)-(1-*tert*-butoxycarbonylamino-1-methylethyl)pyrrolidine (General Method E).

(aa) [7-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester ([MS ES, MH^+] m/z 534) using *N'*-[4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-3-methylbenzoyl]-hydrazinecarboxylic acid *tert*-butyl ester (Example 7t) (General Method D).

(bb) {7-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (MS ES, MH^+) m/z 548)) using *N'*-{4-[3-(*tert*-butoxycarbonylaminomethyl)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methylbenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 7u) (General Method D).

(cc) [7-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-1-cyclopropyl-8-fluoro-6-methoxy-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester ([MS ES, MH_2^+] m/z 551) using *N'*-[4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-3-fluoro-5-methoxybenzoyl]-hydrazinecarboxylic acid *tert*-butyl ester (Example 7v) (General Method D).

(dd) {7-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}-carbamic acid *tert*-butyl ester ([MS ES, MH_2^{+2}] m/z 565) using *N'*-{4-[3-(*tert*-butoxycarbonylaminomethyl)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluoro-3-methoxybenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 7w) (General Method D).

(ee) [7-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-1-cyclopropyl-8-ethoxy-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester ([MS ES, MH_2^{+2}] m/z 565) from *N'*-[4-(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-3-ethoxy-5-fluorobenzoyl]-hydrazinecarboxylic acid *tert*-butyl ester (Example 7cc) (General Method A).

(ff) [7-(3-*tert*-Butoxycarbonylaminopyrrolidin-1-yl)-8-cyano-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester ([MS ES, MH^+] m/z 545) from *N'*-[3-(benzyloxyiminomethyl)-4-

-130-

(3-*tert*-butoxycarbonylaminopyrrolidin-1-yl)-2-cyclopropylamino-5-fluoro-benzoyl]hydrazinecarboxylic acid *tert*-butyl ester (Example 7x) (General Method A).

(gg) [8-Chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-7-(3-[1,2,3-triazol]-1-yl-pyrrolidin-1-yl)-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (¹H NMR (200 MHz, CDCl₃): δ 7.96 (s, 1H), 7.74 (s, 1H), 7.29 (d, 1H), 6.26 (bs, 1H), 5.51-5.39 (m, 1H), 4.17-4.07 (m, 1H), 3.99-3.88 (m, 1H), 3.71-3.66 (m, 1H), 3.57-3.45 (m, 1H), 3.15-3.05 (m, 1H), 2.74-2.60 (m, 1H), 2.44-2.32 (m, 1H), 1.64 (s, 9H), 0.76-0.49 (m, 4H)) from N'-[3-chloro-2-cyclopropylamino-5-fluoro-4-(3-[1,2,3-triazol]-1-yl-pyrrolidin-1-yl)benzoyl]hydrazinecarboxylic acid *tert*-butyl ester (Example 7y) (General Method A).

(hh) [7-(3-Carbamoylpiperidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (MS CI: *m/z* 496 (MH⁺)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and 3-carbamoyl-piperidine (General Method B).

(ii) {7-[(*trans*-3-(*tert*-Butoxycarbonylaminomethyl)-4-trifluoromethylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS CI: *m/z* 635 (M⁺)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and *trans*-3-(*tert*-butoxycarbonylaminomethyl)-4-trifluoromethylpyrrolidine (Li Q., Wang W., Berst K.B., Claiborne A., Hasvold L., Raye K., Tufano M., et al., *J. Med. Chem. Lett.*, 1998;8[15]:1953-1958) (General Method B).

(jj) 8-Chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-7-{3-[(2,2,2-trifluoroethylamino)methyl]pyrrolidin-1-yl}-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (MS CI: *m/z* 548 (M⁺)) using (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and {3-[(2,2,2-trifluoroethylamino)methyl]pyrrolidin-1-yl}carbamic acid *tert*-butyl ester (Domagala J.M., Mich T.F., Sanchez J.P. US 5097032) (General Method B).

(kk) [8-Chloro-1-cyclopropyl-6-fluoro-7-(5-methylhexahydro-pyrrolo[3,4-*c*]pyrrol-2-yl)-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (MS CI: m/z 494 (MH^+)) using (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and 5-methyloctahydropyrrolo[3,4-*c*]pyrrole (Ohnmacht C.J. Jr., Draper C.W., Dedinas R.F., Loftus P., Wong J.J., *J. Heterocycl. Chem.*, 1983;20[2]:321-329) (General Method B).

(ll) [8-Chloro-1-cyclopropyl-7-(2,7-diazaspiro[4.4]non-2-yl)-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (MS CI: m/z 494 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and (2,7-diaza-spiro[4.4]non-2-yl)carbamic acid *tert*-butyl ester (Culbertson T.P., Sanchez J.P., Gambino L., Sesnie J.A., *J. Med. Chem.*, 1990;33(8):2270-2275) (General Method B).

(mm) (7-[3-Benzyl-3-(*tert*-butoxycarbonylaminoethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)-carbamic acid *tert*-butyl ester (MS CI: m/z 658 (MH^+)) using (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and (3-benzylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A7o) (General Method B).

(nn) {7-[(*R*)-3-((*S*)-1-*tert*-Butoxycarbonylaminoethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}-carbamic acid *tert*-butyl ester ((MS CI: m/z 582 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and (*R*)-(3-yl)pyrrolidin-3-ylethyl)carbamic acid *tert*-butyl ester (Johnson D.R., Szotek D.L., Domagala J.M., Stickney T.M., Michel A., Kampf J.W., *J. Heterocycl. Chem.*, 1992;29[6]:1481-1488) (General Method B).

(oo) [8-Chloro-1-cyclopropyl-6-fluoro-7-(3-hydroxyiminopyrrolidin-yl)-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (MS CI: m/z 469 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-

1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and pyrrolidin-3-one oxime (Example A7) (General Method B).

(pp) [7-*trans*-(3-*tert*-Butoxycarbonylamino-4-trifluoromethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]-carbamic acid *tert*-butyl ester (MS CI: *m/z* 622 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and *trans*-(4-trifluoromethylpyrrolidin-3-yl)carbamic acid *tert*-butyl ester (Fukui H., Shibata T., Naito T. Nakano J., Maejima T., Senda H., Iwatani W., et al., *Bioorg. Med. Chem. Lett.*, 1998;8[20]:2833-2838) (General Method B).

(qq) (7-{(R)-3-(S)-1-(*tert*-Butoxycarbonylmethylamino) ethyl}pyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (MS CI: *m/z* 596 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and (R)-3-((S)-1-methylaminoethyl)pyrrolidine (*J. Het. Chem.*, 1992;29:1481) (General Method F).

(rr) *trans*-[7-((3-*tert*-Butoxycarbonylamino-4-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (MS CI: *m/z* 630 (MH^+)) using (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and *trans*-4-phenylpyrrolidin-3-ylamine carbamic acid *tert*-butyl ester (Bucsh R.A., Domagala J.M., Laborde E., Sesnie J.C., *J. Med. Chem.*, 1993;36[26]:4139-4151) (General Method B).

(ss) *trans*-{7-[3-*tert*-Butoxycarbonylamino-4-(4-hydroxyphenyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (MS: *m/z* 646 (MH^+)) using (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and *trans*-4-(4-hydroxyphenyl)pyrrolidin-3-ylamine carbamic acid *tert*-butyl ester (Bucsh R.A., Domagala J.M., Laborde E., Sesnie J.C., *J. Med. Chem.*, 1993;36[26]:4139-4151) (General Procedure B).

-133-

(tt) {7-[3-(*tert*-Butoxycarbonylaminomethyl)piperidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS CI: m/z 580 (M^+)) using (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and (piperidin-3-ylmethyl)carbamic acid *tert*-butyl ester (General Method B).

(uu) (7-{3-[(*tert*-Butoxycarbonylisopropylamino)methyl]pyrrolidin-1-yl}-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)-carbamic acid *tert*-butyl ester (MS CI: m/z 608 (M^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and 3-(isopropylaminomethyl)pyrrolidine (Domagala et al., *J. Med. Chem.*, 1994;3889-3901) (General Method F).

(vv) {7-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-6,8-dichloro-1-cyclopropyl-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS CI: m/z 584 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and (pyrrolidine-3-ylmethyl)carbamic acid *tert*-butyl ester (General Method B).

(ww) [7-(3-*tert*-Butoxycarbonylamino-3-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$) δ 7.68 (d, 1H), 7.50-7.29 (m, 5H), 6.72 (s, 1H), 5.24 (s, 1H), 4.18-3.85 (m, 3H), 3.78-3.52 (m, 2H), 2.70 (s, 1H), 2.58-2.39 (m, 1H), 1.62-0.62 (m, 22H)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and (3-phenylpyrrolidin-3-yl)carbamic acid *tert*-butyl ester (Example A30) using General Method B.

(xx) {7-[3-(*tert*-Butoxycarbonylaminomethyl)-4-methylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (MS ES: m/z 582 (MH^+)) from (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14) and trans-(4-methylpyrrolidin-3-yl)carbamic acid

-134-

tert-butyl ester (Kuniyoshi M., Seigo S., Keiji H., Takayoshi I., Eur. Pat. Appl., 1987, EP 208210) using general procedure F.

EXAMPLE 9

Deprotection of Quinazolin-2,4-diones

5 General Method A

Hydrogen chloride gas is bubbled into diethyl ether (or dichloromethane or dichloroethane for 15 minutes. The resulting solution is then cooled to 0°C and added to a solution of a 2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 6). The reaction mixture is slowly warmed to room
10 temperature. After 30 hours, the precipitate is filtered, washed with ether (or dichloromethane or dichloroethane) and hexanes, and dried under vacuum to afford the hydrochloride salt of the desired deprotected amine as a solid.

General Method B

A compound is dissolved in trifluoroacetic acid and stirred at 0°C for 3 to
15 8 hours. The acid is removed by blowing compressed air over the solution or concentrate in vacuo. The residue is twice co-evaporated with ethanol and dried in vacuo to yield the trifluoroacetate salt of the title compound as a solid.

General Method C

A 3-dibenzylamino-6-fluoro-1-substituted-2,4-dioxo-1,2,3,4-tetrahydro-
20 quinazolin-7-yl]pyrrolidin-3-yl]carbamic acid *tert*-butyl ester and 20% Pd/C (cat.) are combined in methanol and shaken under 50 psi of hydrogen for 22.5 hours. The suspension is filtered through Celite and concentrated. The residue is purified via chromatography (SiO₂, CHCl₃) to give a solid. The solid is then dissolved in methylene chloride, cooled to 0°C, and HCl gas is passed through the solution for
25 10 minutes. The suspension is stirred for 2 hours and concentrated to give the title compound.

-135-

General Method D

Hydrogen chloride gas is bubbled through a solution of substrate in diethyl ether, and the solution stirred for 2 hours. The solvent is removed under reduced pressure to afford the desired title compound as a solid.

5 General Method E

A solution of substrate in ethanol is treated with a solution of ethanol saturated with gaseous hydrogen chloride. The reaction mixture is stirred at room temperature for 18 hours, and the solvent is removed in vacuo. The residue is dissolved in water, filtered through a fiber glass pad to clarify, and lyophilized.
10 The solid residue is triturated with ether, filtered, washed with ether and dried in vacuo.

General Method F

A solution of 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b), 1.2 eq. of heterocyclic amine side chain,
15 and 1:3 *N,N*-diisopropylethylamine/acetonitrile is heated at 50°C for 18 hours. The reaction mixture is diluted with water, cooled to 0°C, and the solid is removed by filtration, washed with 50% aqueous acetonitrile, and dried in vacuo to give a solid that is dissolved in ethanol and treated with a solution of ethanol saturated with gaseous hydrogen chloride. The mixture is then stirred at room temperature
20 for 18 hours. The solvent is removed in vacuo and the residue is triturated with ether. The solid is removed by filtration, washed with ether and dried in vacuo.

The following compounds are synthesized according to General Procedures A-F of Example 9:

(a) 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (Compound 1)
25 (mp 153-155°C, MS CI: *m/z* 368 (MH⁺)) from 7-[3-(*tert*-butoxycarbonylamino-methyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8a) using General Method A.

(b) 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-piperazin-1-yl-1H-quinazoline-2,4-dione hydrochloride (**Compound 2**) (mp 156-159°C, MS CI: m/z 354 (MH^+)) from 7-[4-(*tert*-butoxycarbonylpiperazin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8b) using General Method A.

(c) 3-Amino-7-[3-(aminomethyl)-3-methylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 3**) (mp 166-168°C, MS CI: m/z 382 (MH^+)) from {7-[3-(*tert*-butoxycarbonylamino-methyl)-3-methylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8c) using General Method A.

(d) 3-Amino-7-(6-amino-3-aza-bicyclo[3.1.0]hex-3-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazolin-2,4-dione hydrochloride (**Compound 4**) (mp 111-114°C, MS CI: m/z 366 (MH^+)) from [7-(6-*tert*-butoxycarbonylamino-3-azabicyclo[3.1.0]hex-3-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8d) using General Method A.

(e) 3-Amino-7-((S)-3-N-methylaminopyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 5**) (mp 118-120°C, MS CI: m/z 368 (MH^+)) from {7-[(S)-3-(*tert*-butoxycarbonyl-N-methylamino)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8e) using General Method A.

(f) 3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-8-chloro-1-(2,4-difluoroanilino)-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 6**) (mp 184-186°C, MS CI: m/z 427(MH^+)) from {7-[(S)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-8-chloro-1-(2,4-difluoroanilino)-6-fluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8f) using General Method A.

(g) 3-Amino-7-[(S)-3-aminopyrrolidin-1-yl]-1-cyclopropylamino-6,8-difluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 7**) (mp 288°C

-137-

(dec.), MS EI: m/z 338 (M^+) from 7-[(S)-3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-1-cyclopropylamino-6,8-difluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8g) using General Method A.

5 (h) 3-Amino-7-[(S)-3-aminopyrrolidin-1-yl]-1-cyclopropylamino-5,8-difluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 8**) (mp 338°C (dec), MS EI: m/z 338 (M^+)) from {7-[(S)-3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-1-cyclopropylamino-5,8-difluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8h) using General
10 Method A

(i) 3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-5,6,8-trifluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 9**) (mp 271°C (dec.), MS CI: m/z 356 (MH^+)) from {7-[(S)-3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-1-cyclopropyl-1,4-dihydro-2,4-dioxo-5,6,8-trifluoro-2H-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (Example 8i) using General
15 Method A.

(j) 3-Amino-7-(3-aminopyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 10**) (mp 134-136°C, MS CI: m/z 354 (MH^+)) from {7-[3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8a) using General
20 Method A.

(k) 7-((S)-3-Aminopyrrolidin-1-yl)-1-cyclopropyl-3,5-diamino-6,8-difluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 11**) (mp 215-218°C (dec.), MS EI: 352 (M^+)) from {5-amino-7-[(S)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-6,8-difluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 12) using
25 Method A.

(l) 3-Amino-7-(3-hydroxymethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 12**) (mp 78-80°C, MS CI: m/z 369 (MH^+)) from [7-(3-hydroxymethylpyrrolidin-1-yl)-
30

-138-

8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8k) using Method A.

(m) 3-Amino-7-(3-aminoethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 13**)

5 (mp 158-160°C, MS CI: m/z 382 (MH^+)) from {7-[3-(*tert*-butoxycarbonylaminoethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8l) using Method A.

(n) 3-Amino-7-((*S*)-7-amino-5-azaspiro[2.4]hept-5-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 14**)

10 (mp 177-179°C, MS CI: m/z 380 (MH^+)) from [7-(7-*tert*-butoxycarbonylamino-5-azaspiro[2.4]hept-5-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8m) using General Method A.

(o) 3-Amino-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 15**) (mp 117-120°C, MS CI: m/z 396 (MH^+)) from {7-[(*R*)-3-(1-*tert*-butoxycarbonylamino-1-methylethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8n) using General Method A.

20 (p) 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 16**) (mp 170-172°C, MS CI: m/z 334 (MH^+)) from {7-[3-(*tert*-butoxycarbonylamino-methyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8o) using General Method A.

25 (q) 3-Amino-7-(pyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 17**) (mp 96-98°C, MS CI: m/z 339 (MH^+)) from [7-(pyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8p) using General Method A.

30

- (r) 3-Amino-7-(3-aminopyrrolidin-1-yl)-8-chloro-6-fluoro-1-isopropyl-1*H*-quinazoline-2,4-dione trifluoroacetate (**Compound 18**) (mp 151°C, MS EI: m/z 356 (MH^+)) from {7-[3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-8-chloro-6-fluoro-1-isopropyl-2,4-dioxo-6-fluoro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8q) using General Method B.
- (s) 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-*sec*-butyl-8-chloro-6-fluoro-1*H*-quinazoline-2,4-dione trifluoroacetate (**Compound 19**) (mp 115°C) from {7-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-1-*sec*-butyl-8-chloro-6-fluoro-2,4-dioxo-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8r) using General Method B.
- (t) 3-Amino-7-[3-aminopyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 20**) (mp 216-218°C, MS EI: m/z 350 (MH^+)) from {7-[3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8s) using Method A.
- (u) 3-Amino-7-[3-aminopyrrolidin-1-yl]-8-chloro-1-cyclobutylamino-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 21**) (mp 205-208°C (dec.), MS EI: m/z 368 (MH^+)) from {7-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-8-chloro-1-cyclobutylamino-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8t) using General Method A.
- (v) 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-fluoro-1*H*-quinazoline-2,4-dione trifluoroacetate (**Compound 22**) (mp 210-212°C, MS EI: m/z 320 (MH^+)) from {7-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-8-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8u) using Method A.
- (w) 3-Amino-7-(3-aminopyrrolidin-1-yl)-6-chloro-1-cyclopropyl-8-fluoro-1*H*-quinazoline-2,4-dione trifluoroacetate (**Compound 23**) (mp 135-137°C, MS EI: m/z 354 (MH^+)) from {7-[3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-6-chloro-1-cyclopropyl-8-fluoro-2,4-dioxo-1,4-dihydro-2*H*-

quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8v) using General Method B.

(x) 3-Amino-7-((*S*)-3-aminopyrrolidin-1-yl)-1-cyclopropylmethyl-8-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 24**) (mp >250°C; MS CI: *m/z* 334 (MH^+)) from [(*S*)-1-(1-cyclopropylmethyl-3-dibenzylamino-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 8w) using General Method C.

(y) 3-Amino-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropylmethyl-8-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 25**) (mp >250°C, MS CI: *m/z* 376 (MH^+)) from {1-[(*R*)-1-(cyclopropylmethyl-3-dibenzylamino-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]-1-methylethyl} carbamic acid *tert*-butyl ester (Example 8x) using General Method C.

(z) 3-Amino-7-((*S*)-3-aminopyrrolidin-1-yl)-1-ethyl-8-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 26**) (mp 248-251°C, MS CI: *m/z* 308 (MH^+)) from [(*S*)-1-(3-dibenzylamino-1-ethyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]-carbamic acid *tert*-butyl ester (Example 8y) using General Method C.

(aa) 3-Amino-1-ethyl-6-fluoro-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 27**) (mp >250°C, MS CI: *m/z* 350 (MH^+)) from [(*R*)-1-(3-dibenzylamino-1-ethyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-1-methyl-1-pyrrolidin-3-ylethyl]carbamic acid *tert*-butyl ester (Example 8z) using Method C.

(bb) 3-Amino-7-((*S*)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 28**) (mp >250°C, MS CI: *m/z* 320 (MH^+)) from 3-amino-{7-[(*S*)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione (Example 25) using General Method A.

(cc) 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione trifluoroacetate (**Compound 29**) (mp 110-112°C, MS ES, MH^+) *m/z* 334) from [7-(3-*tert*-butoxycarbonylamino-

-141-

pyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8aa) using General Method B.

(dd) 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione trifluoroacetate (**Compound 30**) (mp 82-84°C, MS ES, m/z 348 (MH^+)) from {7-[3-(*tert*-butoxycarbonylaminomethyl)-pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8bb) using General Method B.

(ee) 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-fluoro-6-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 31**) (mp 208-210°C, MS ES, m/z 350(MH^+)) from [7-(3-*tert*-butoxycarbonylamino-pyrrolidin-1-yl)-1-cyclopropyl-8-fluoro-6-methoxy-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8cc) using General Method A.

(ff) 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 32**) (mp 180-182°C, MS ES, m/z 364 (MH^+)) from {7-[3-(*tert*-butoxycarbonylaminomethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}-carbamic acid *tert*-butyl ester (Example 8dd) using General Method A.

(gg) 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-ethoxy-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 33**) (mp 220°C, MS ES, m/z 364 (MH^+)) from [7-(3-*tert*-butoxycarbonylamino-pyrrolidin-1-yl)-1-cyclopropyl-8-ethoxy-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8ee) using General Method A.

(hh) 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-8-carbonitrile hydrochloride (**Compound 34**) (mp 277°C, MS ES, m/z 345 (MH^+)) from [7-(3-*tert*-butoxycarbonylamino-pyrrolidin-1-yl)-8-cyano-1-cyclopropyl-6-fluoro-2,4-dioxo-

-142-

1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8ff) using General Method A.

(ii) 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(3-[1,2,3-triazol]-1-yl-pyrrolidin-1-yl)-1H-quinazoline-2,4-dione trifluoroacetate (**Compound 35**)
5 (mp 145-147°C, MS ES, *m/z* 406 (MH⁺)) from [8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-7-(3-[1,2,3-triazol]-1-yl-pyrrolidin-1-yl)-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8gg) using General Method B.

(jj) 3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride
10 (**Compound 36**), (mp 215-217°C) from {(R)-1-[(S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}-carbamic acid *tert*-butyl ester (Example 28a) using General Method E.

(kk) 3-Amino-7-[(S)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride
15 (**Compound 37**) (mp 221-223°C) from {(S)-1-[(S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}-carbamic acid *tert*-butyl ester (Example 28b) using General Method A.

(ll) 3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride
20 (**Compound 38**) (mp 197-199°C) from {(R)-1-[(S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28c) using General Method E.

(mm) 3-Amino-7-[(S)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride
25 (**Compound 39**) (mp 192-194°C) from {(S)-1-[(S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28d) using General Method E.

(nn) 5-Amino-9-((S)-3-aminopyrrolidin-1-yl)-8-fluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione hydrochloride (**Compound 40**)
30 (mp 202-204°C) from [(S)-1-(5-amino-8-fluoro-3-methyl-4,6-dioxo-2,3,5,6-

tetrahydro-4H-1-oxa-3a,5-diazaphenalen-9-yl]pyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28e) using General Method E.

(oo) 5-Amino-9-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-8-fluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenylene-4,6-dione hydrochloride
5 (Compound 41) (mp 192-194°C) from {(S)-1-[(R)-1-(5-amino-8-fluoro-3-methyl-4,6-dioxo-2,3,5,6-tetrahydro-4H-1-oxa-3a,5-diazaphenalen-9-yl)-pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28f) using General Method E.

(pp) 2-Amino-8-((S)-3-aminopyrrolidin-1-yl)-9-fluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione, hydrochloride (Compound 42)
10 (mp 202-204°C) from [(S)-1-(2-amino-9-fluoro-5-methyl-1,3-dioxo-2,3,6,7-tetrahydro-1H,5H-pyrido[3,2,1-ij]quinazolin-8-yl)pyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28g) using General Method E.

(qq) 2-Amino-8-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-9-fluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione hydrochloride
15 (Compound 43) (mp 197-199°C) from {(R)-3-[(S)-1-(2-amino-9-fluoro-5-methyl-1,3-dioxo-2,3,6,7-tetrahydro-1H,5H-pyrido[3,2,1-ij]quinazolin-8-yl)-pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28h) using General Method E.

(rr) 3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione hydrochloride (Compound 44)
20 (mp 213-215°C) from [1-(3-Amino-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydro-pyrido[2,3-d]pyrimidin-7-yl)pyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28i) using General Method E.

(ss) 3-Amino-7-(6-amino-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione, hydrochloride (Compound 45)
25 (mp 268-270°C) from 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b) and 3-azabicyclo[3.1.0]hex-6-ylcarbamic acid *tert*-butyl ester using General Method F.

(tt) 3-Amino-7-(3-aminomethyl-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione, hydrochloride
30

(Compound 46) (mp 187-189°C) from 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b) and [1-methyl-(3-methylpyrrolidin-3-yl)]methylcarbamic acid *tert*-butyl ester (*J. Med. Chem.*, 1992;361-367) using General Method F.

5 (uu) 3-Amino-1-cyclopropyl-6-fluoro-7-(octahydropyrrolo[3,4-c]pyridin-2-yl)-1H-pyrido[2,3-d]pyrimidine-2,4-dione hydrochloride
(Compound 47) (mp 200-203°C) from 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b) and (3aS,7aR)-octahydropyrrolo[3,4-c]pyridine-5-carboxylic acid *tert*-butyl ester (Example A27)
10 using General Method F.

(vv) 3-Amino-1-cyclopropyl-6-fluoro-7-(octahydropyrrolo[3,4-b]pyridin-2-yl)-1H-pyrido[2,3-d]pyrimidine-2,4-dione hydrochloride
(Compound 48) (mp 200-202°C) from 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b) and octahydropyrrolo[3,4-b]pyridine-5-carboxylic acid *tert*-butyl ester using General Method F.
15

(ww) 3-Amino-7-[(R)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione hydrochloride
(Compound 49) (mp >260°C) from 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b) and (R)-3-(1-amino-1-methylethyl)-pyrrolidine using General Method F.
20

(xx) 3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione, dihydrochloride (Compound 50)
(mp 152-154°C) from 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b) and (3-aminomethylpiperidin-1-yl)carbamic acid *tert*-butyl ester (Hilpert K. et al., *J. Med. Chem.*, 1994;37[23]:3889-3901) using General Method F.
25

(yy) *trans*-3-Amino-7-(3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione hydrochloride
(Compound 51) (mp 214-216°C) from 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b) and 3-aminomethyl-4-trifluoromethylpyrrolidine (Li Q., Wang W., Berst K.B., Claiborne A.,
30

Hasvold L., Raye K., Tufano M., et al., *Bioorg. Med. Chem. Lett.*, 1998;8:1953-1958) using General Method F.

(zz) 3-Amino-1-cyclopropyl-6-fluoro-7-[(R)-3-((R)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-pyrido[2,3-d]pyrimidine-2,4-dione hydrochloride
5 (Compound 52) (mp 137-139°C) from 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b) and (R)-3-((R)-1-methylaminoethyl)pyrrolidine using General Method F.

(aaa) 3-Amino-7-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione hydrochloride
10 (Compound 53) (mp >260°C) from {(S)-1-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-7-yl)pyrrolidin-3-yl]-ethyl}carbamic acid *tert*-butyl ester (Example 28j) using General Method E.

(bbb) 3-Amino-7-[3-(1-aminopropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (Compound 54) (MS
15 CI: m/z 376 (MH^+)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]propyl}carbamic acid *tert*-butyl ester (Example 28ooo) using General Method E.

(ccc) 3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-[(R)-3-((S)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione hydrochloride
20 (Compound 55) (mp 153-5°C) from {(S)-1-[(R)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]-ethyl}methylcarbamic acid *tert*-butyl ester (Example 28ppp) using General Method E.

(ddd) 3-Amino-7-[7-(1-aminoethyl)-5-azaspiro[2.4]hept-5-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride
25 (Compound 56) (mp 178-80°C, MS CI: m/z 388 (MH^+)) from {1-[5-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-5-aza-spiro[2.4]hept-7-yl]-ethyl}carbamic acid *tert*-butyl ester (Example 28qqq) using General Method E.

(eee) 1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidine-3-carboxylic acid amide (Compound 58)
30

-146-

(mp 141-144°C, MS CI: m/z 396 (MH^+)) from [7-(3-carbamoylpiperidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]-carbamic acid *tert*-butyl ester using General Method A (Example 8hh).

(fff) 3-Amino-[7-*trans*-3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride
(Compound 59) (mp 171-173°C, MS CI: m/z 436 (MH^+)) from {7-[*trans*-3-(*tert*-butoxycarbonylaminomethyl)-4-trifluoromethylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 8ii) using General Method E.

(ggg) 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-{3-[(2,2,2-trifluoroethylamino)methyl]pyrrolidin-1-yl}-1*H*-quinazoline-2,4-dione hydrochloride
(Compound 60) (mp 163-164°C, MS CI: m/z 450 (MH^+)) from 8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-7-{3-[(2,2,2-trifluoroethylamino)methyl]pyrrolidin-1-yl}-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester
(Example 8jj) using General Method E.

(hhh) 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(5-methylhexahydropyrrolo[3,4-*c*]pyrrol-2-yl)-1*H*-quinazoline-2,4-dione hydrochloride
(Compound 61) (mp 133-135°C, MS CI: m/z 394 (MH^+)) from [8-chloro-1-cyclopropyl-6-fluoro-7-(5-methylhexahydropyrrolo[3,4-*c*]pyrrol-2-yl)-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8kk) using General Method E.

(iii) 3-Amino-8-chloro-1-cyclopropyl-7-(2,7-diazaspiro[4.4]non-2-yl)-6-fluoro-1*H*-quinazoline-2,4-dione (Compound 62) (mp 147-149°C, MS CI: m/z 394 (MH^+)) from [8-chloro-1-cyclopropyl-7-(2,7-diazaspiro[4.4]non-2-yl)-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8li) using General Method E.

(jjj) 3-Amino-7-(3-aminomethyl-3-benzylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (Compound 63) (mp 148-150°C, MS CI: m/z 458 (MH^+)) from (7-[3-benzyl-3-(*tert*-butoxycarbonylaminomethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-

-147-

2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester
(Example 8mm) using General Method E.

(kkk) 3-Amino-7-[(*R*)-3-((*S*)-1-aminoethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 64**)

5 (mp 164-166°C, MS CI: *m/z* 382 (MH^+)) from {7-[(*R*)-3-((*S*)-1-*tert*-butoxycarbonylaminoethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}carbamic acid *tert*-butyl ester
(Example 8nn) using General Method E.

10 (lll) 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(3-hydroxyimino-pyrrolidin-1-yl)-1*H*-quinazoline-2,4-dione (**Compound 65**) (mp 147-9°C, MS CI: *m/z* 368 (MH^+)) from) [8-chloro-1-cyclopropyl-6-fluoro-7-(3-hydroxyimino-pyrrolidin-yl)-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8oo) using General Method E.

15 (mmm) 3-Amino-7-[trans-3-amino-4-trifluoromethylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 66**) (mp 181-183°C, MS CI: *m/z* 422 (MH^+)) from [7-(trans-3-*tert*-butoxycarbonylamino-4-trifluoromethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 8pp) using General Method E.

20 (nnn) 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-[(*R*)-3-(*S*)-1-methylaminoethyl]pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 67**) (mp 122-124°C, MS CI: *m/z* 396 (MH^+)) from (7-[(*R*)-3-(*S*)-1-(*tert*-butoxycarbonylmethylamino)ethyl]pyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl)carbamic acid *tert*-butyl ester
25 (Example 8qq) using General Method E.

(ooo) 3-Amino-7-(trans-3-amino-4-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (**Compound 68**)
(mp 174-175°C, MS CI: *m/z* 430 (MH^+)) from [7-(trans-3-*tert*-butoxycarbonylamino-4-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl]carbamic acid *tert*-butyl ester
30 (Example 8rr) using General Method E.

(ppp) 3-Amino-7-[trans-3-amino-4-(4-hydroxyphenyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride

(Compound 69) (mp >200°C, MS CI: *m/z* 446 (MH⁺)) using 7-[trans-3-*tert*-butoxycarbonylamino-4-(4-hydroxyphenyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8ss) and General Method E.

(qqq) *N*-[1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-ylmethyl]methanesulfonamide hydrochloride (Compound 70) (mp 109-112°C, MS CI: *m/e* 446 (MH⁺)) from {8-chloro-1-cyclopropyl-6-fluoro-7-[3-(methanesulfonylamino-methyl)-pyrrolidin-1-yl]-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 36) using General Method E.

(rrr) 3-Amino-7-(3-aminomethylpiperidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione hydrochloride (Compound 71) (mp 131-133°C, MS CI: *m/z* 382 (MH⁺)) from {7-[3-(*tert*-butoxycarbonylamino-methyl)piperidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester (Example 8tt) using General Method E.

(sss) 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-[3-(isopropylamino-methyl)pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione hydrochloride (Compound 72) (mp 125-127°C, MS CI: *m/z* 410 (MH⁺)) from (7-{3-[(*tert*-butoxycarbonyl-isopropylamino)methyl]pyrrolidin-1-yl}-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl) carbamic acid *tert*-butyl ester (Example 8uu) using General Method E.

(ttt) 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-6,8-dichloro-1-cyclopropyl-1*H*-quinazoline-2,4-dione hydrochloride (Compound 73) (mp 147-149°C, MS CI: *m/z* 384 (MH⁺)) from {7-[3-(*tert*-butoxycarbonylamino-methyl)-pyrrolidin-1-yl]-6,8-dichloro-1-cyclopropyl-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl} carbamic acid *tert*-butyl ester using General Method E.

(uuu) *N*-[1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-ylmethyl]methanesulfonamide

(Compound 74) (mp 109-112°C, MS CI: m/z 446 (MH^+)) from {8-chloro-1-cyclopropyl-6-fluoro-7-[3-(methanesulfonylaminomethyl)pyrrolidin-1-yl]-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Example 36) using General Method E.

(vvv) 3-Amino-7-[(*R*)-3-(*S*)-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride

(Compound 75) (mp 194-196°C, ^1H NMR (200 MHz, DMSO- d_6): δ 8.25 (bs, 3H), 7.47 (d, 1H), 4.68 (bs, 3H), 3.59-3.23 (m, 7H), 2.49 (s, 3H), 2.02-1.92 (m, 1H), 1.80-1.60 (m, 1H), 1.30 (d, 3H), 1.12-1.08 (m, 2H), 0.62-0.50 (m, 2H); MS ES: m/z 362 (MH^+)) from {1-[1-[(*R*)-3-(*S*)-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28k) using General Method A.

(www) 3-Amino-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (Compound 76) (mp 200-202°C, MS ES: m/z 376 (MH^+)) from {1-[(*R*)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]-1-methylethyl}carbamic acid *tert*-butyl ester (Example 28l) using Method A.

(xxx) 3-Amino-7-[3-(1-aminoethyl)-3-methoxypyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (Compound 77) (mp 190-192°C, ^1H NMR (200 MHz, DMSO- d_6): δ 8.07 (bs, 3H), 7.49 (d, 1H), 3.89-3.10 (m, 12H), 2.42 (s, 3H), 2.39-2.08 (m, 2H), 1.29 (d, 3H), 1.07-1.03 (m, 2H), 0.62-0.52 (m, 2H); MS ES: m/z 392 (MH^+)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methoxypyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28m) using General Method A.

(yyy) 3-Amino-7-[3-(1-aminoethyl)-3-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride

-150-

(**Compound 78**) (mp 190-192°C, MS ES: m/z 396 (MH⁺)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-fluoropyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28o) using General Method A.

5 (zzzz) 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-5-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 79**) (mp 238-240°C, MS ES: m/z 334 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-5-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28p) using General Method A.

10 (aaaa) 3-Amino-7-[(R)-3-(S)-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-5-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 80**) (mp 245-247°C, MS ES: m/z 362 (MH⁺)) from {1-(S)-[1-(R)-(3-amino-1-cyclopropyl-6-fluoro-5-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28q) using
15 General Method A.

 (bbbb) 3-Amino-7-(3-aminomethyl-3-methoxymethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 81**) (mp 172-177°C, MS ES: m/z 392 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methoxymethylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester
20 (Example 28r) using General Method A.

 (cccc) 3-Amino-7-(3-aminomethyl-3-fluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (**Compound 82**) (mp 160-163°C, MS ES: m/z 380 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-fluoromethylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28s) using General Method A.
25

 (dddd) 3-Amino-7-(trans-3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 83**) (mp 205-207°C, MS ES: m/z 432 (MH⁺)) from [trans-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-
30

-151-

4-trifluoromethylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester

(Example 28t) using General Method A.

(eeee) 3-Amino-7-[3-(1-aminoethyl)piperidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 84**)

5 (mp 198°C, MS ES: m/z 392 (MH^+)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28u) using General Method A.

(ffff) 3-Amino-7-[3-(aminoethyl)piperidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 85**) (mp 200°C,

10 ^1H NMR (200 MHz, DMSO- d_6): δ 8.25 (bs, 3H), 7.52 (d, 1H), 6.36 (bs, 3H), 3.54-2.80 (m, 6H), 2.48 (s, 3H), 2.02-1.15 (m, 8H), 1.02-0.50 (m, 4H); MS ES: m/z 376(MH^+)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28v) using General Method A.

15 (gggg) 3-Amino-7-[4-(1-aminoethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 86**) (mp 230°C, MS ES: m/z 406 (MH^+)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28w) using General Method A.

20 (hhhh) 3-Amino-7-[4-(1-aminoethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 87**) (mp 246°C, MS ES: m/z 390 (MH^+)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28x) using General Method A.

25 (iiii) 3-Amino-7-(3-amino-3-phenylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 88**) (mp 236°C, MS ES: m/z 410 (MH^+)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-phenylpyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28z) using General Method A.

30

-152-

(jjjj) 3-Amino-7-[3-(1-aminoethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride
(Compound 89) (mp 240°C, MS ES: m/z 376 (MH⁺)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28y) using
5 General Method A.

(kkkk) 3-Amino-7-(3-aminomethyl-3-phenylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride
(Compound 90) (mp 227-231°C, MS ES: m/z 424 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)3-phenylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28aa) using
10 General Method A.

(llll) 3-Amino-7-(7-aminomethyl-5-azaspiro[2,4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride
(Compound 91) (mp 167°C, MS ES: m/z 374 (MH⁺)) from [5-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-5-azaspiro[2.4]hept-7-ylmethyl]carbamic acid *tert*-butyl ester (Example 28bb) using
15 General Method A.

(mmmm) 3-Amino-7-(7-aminomethyl-5-azaspiro[2,4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride
(Compound 92) (mp 178°C, MS ES: m/z 390 (MH⁺)) from [5-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-5-azaspiro[2.4]hept-7-ylmethyl]carbamic acid *tert*-butyl ester (Example 28cc) using
20 General Method A.

(nnnn) 3-Amino-7-(3-aminomethyl-3-hydroxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride
(Compound 93) (mp 202°C, MS ES: m/z 380 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-hydroxypyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28dd)
25 using General Method A.
30

-153-

(oooo) 3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 95**)

(mp 219-221°C, MS ES: m/z 362 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28ee) using General Method A.

(pppp) 3-Amino-7-(3-amino-4-methoxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 96**)

(mp >250°C, MS ES: m/z 364 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methoxypyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28ff) using General Method A.

(qqqq) 3-Amino-7-(3-amino-4-methoxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 97**)

(mp >250°C, MS ES: m/z 380 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methoxypyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28gg) using General Method A.

(rrrr) 3-Amino-7-(3-amino-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 98**)

(mp >250°C, MS ES: m/z 368 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28hh) using General Method A.

(ssss) 3-Amino-7-(3-aminomethyl-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride

(**Compound 99**) (mp 210°C, MS ES: m/z 362 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28ii) using General Method A.

(tttt) 3-Amino-7-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-ethyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride

(**Compound 100**) (mp 155-157°C, MS ES: m/z 376 (MH⁺)) from {(S)-1-[(R)-1-(3-amino-1-cyclopropyl-8-ethyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-

-154-

7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28jj) using General Method A.

(uuuu) 1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid trifluoroacetate

5 (Compound 101) (mp 252°C (decomp), MS ES: m/z 379 (MH^+)) from 1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid *tert*-butyl ester (Example 28kk) using General Method B.

10 (vvvv) 1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid trifluoroacetate

(Compound 102) (mp 226°C (decomp), MS ES: m/z 363 (MH^+)) from 1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid *tert*-butyl ester (Example 28ll) using General Method A.

15 (www) 3-Amino-7-(1-aminomethyl-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride

(Compound 103) (mp 183-185°C, MS ES: m/z 376 (MH^+)) from [3-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-azabicyclo[3.1.0]hex-1-ylmethyl]yl]carbamic acid *tert*-butyl ester (Example 28mm) using General Method A.

20 (xxxx) 3-Amino-7-(1-aminomethyl-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride

(Compound 104) (mp 189-194°C, MS ES: m/z 360 (MH^+)) from [3-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-azabicyclo[3.1.0]hex-1-ylmethyl]yl]carbamic acid *tert*-butyl ester (Example 28nn) using General Method A.

25 (yyyy) 3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride

(Compound 105) (mp 175-179°C, MS ES: m/z 378 (MH^+)) from {(R)-1-[(S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-

30

-155-

tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28oo) using General Method A.

(zzzzz) 3-Amino-7-[(*S*)-3-((*R*)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (Compound 106) (mp 172-176°C, MS (ES: m/z 362 (MH^+)) from {(*R*)-1-[(*S*)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28pp) using General Method A.

(aaaaa) 3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-(octahydropyrrolo[3,4-*b*]pyridin-6-yl)-1H-quinazoline-2,4-dione hydrochloride (Compound 107) (mp 220°C, MS ES: m/z 374 (MH^+)) from 6-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)octahydropyrrolo[3,4-*b*]pyridine-1-carboxylic acid *tert*-butyl ester (Example 28qq) using General Method A.

(bbbbb) 3-Amino-7-(*trans*-3-aminomethyl-4-methylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (Compound 108) (mp 215-217°C, MS ES: m/z 382 (MH^+)) from {7-[*trans*-3-(*tert*-butoxycarbonyl-aminomethyl)-4-methylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester (Compound 109) (Example 8xx) using General Method A.

(ccccc) 3-Amino-7-(*trans*-3-aminomethyl-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (Compound 110) (mp 189°C, MS ES: m/z 378 (MH^+)) from [*trans*-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28rr) using General Method A.

(dddd) 3-Amino-7-(*trans*-3-aminomethyl-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (Compound 111) (mp 220°C, MS ES: m/z 362 (MH^+)) from [*trans*-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-

-156-

methylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28ss) using General Method A.

(eeee) 3-Amino-7-(trans-3-amino-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride
5 (Compound 112) (mp 209°C, MS ES: m/z 364 (MH^+)) from [trans-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]carbamic acid *tert*-butyl ester (Example 28tt) using General Method A.

(ffff) 3-Amino-7-(3-aminomethylmorpholin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (Compound 113)
10 (mp 195°C, MS ES: m/z 380 (MH^+)) from [4-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)morpholin-2-ylmethyl]carbamic acid *tert*-butyl ester (Example 28uu) using General Method A.

(gggg) 3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (Compound 114)
15 (mp 201°C, MS ES: m/z 378 (MH^+)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidin-3-ylmethyl]-carbamic acid *tert*-butyl ester (Example 28vv) using General Method A.

(hhhh) 3-Amino-7-[3-(1-aminoethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride
20 (Compound 115) (mp 204.5°C, MS ES: m/z 392 (MH^+)) from {1-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28ww) using General Method A.

(iiii) 3-Amino-7-(3-amino-3-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione hydrochloride (Compound 116)
25 (mp 198-200°C, MS ES: m/z 430 (MH^+)) from [7-(3-*tert*-butoxycarbonylamino-3-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl]carbamic acid *tert*-butyl ester (Example 6ww) using General
30 Method A.

- (jjjj) 3-Amino-7-(3-amino-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 117**) (mp 237-240°C, MS ES: m/z 348 (MH⁺)) from [1-(1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methylpyrrolidin-3-yl]]carbamic acid *tert*-butyl ester (Example 28xx) using General Method A.
- 5 (kkkk) 3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione hydrochloride (**Compound 118**) (mp 178-180°C, MS ES: m/z 319 (MH⁺)) from 3-amino-1-cyclopropyl-6-fluoro-8-methyl-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione (Example 28yy) using General Method A.
- 10 (llll) 3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione hydrochloride (**Compound 119**) (mp 92-94°C, MS ES: m/z 335 (MH⁺)) from 3-amino-1-cyclopropyl-6-fluoro-8-methoxy-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione (Example 28zz) using General Method A.
- 15 (mmmm) 3-Amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)pyrrolidin-1-yl]-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 120**) (mp 116-120°C, MS ES: m/z 377 (MH⁺)) from 3-amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)pyrrolidin-1-yl]-8-methyl-1H-quinazoline-2,4-dione (Example 28aaa) using General Method E.
- 20 (nnnn) 3-Amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)pyrrolidin-1-yl]-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 121**) (mp 134°C (decomp), MS ES: m/z 393 (MH⁺)) from 3-amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)pyrrolidin-1-yl]-8-methoxy-1H-quinazoline-2,4-dione (Example 28bbb) using General Method E.
- 25 (oooo) 3-Amino-1-cyclopropyl-6-fluoro-5-methyl-7-[(R)-3-((S)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione hydrochloride (**Compound 122**) (mp 182-185°C, MS ES: m/z 376 (MH⁺)) from {(S)-1-[(R)-3-amino-1-cyclopropyl-6-fluoro-5-methyl-7-[(R)-3-((S)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione (Example 28ccc) using General Method E.
- 30 (pppp) (S)-1-[(R)-1-(3-Amino-1-cyclopropyl-7-[3-(1-ethylaminoethyl)pyrrolidin-1-yl]-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione

-158-

hydrochloride (**Compound 123**) (mp 185-190°C, MS ES: m/z 406 (MH⁺)) from (S)-1-[(R)-1-(3-amino-1-cyclopropyl-7-[3-(1-ethylaminoethyl)pyrrolidin-1-yl]-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 28ddd) using General Method E.

5 (qqqq) 3-Amino-1-cyclopropyl-7-[(R)-3-((S)-1-ethylaminoethyl)pyrrolidin-1-yl]-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (**Compound 124**) (mp 210-215°C, MS ES: m/z 390 (MH⁺)) from 3-amino-1-cyclopropyl-7-[(R)-3-((S)-1-ethylaminoethyl)pyrrolidin-1-yl]-6-fluoro-8-methyl-1H-quinazoline-2,4-dione (Example 28eee) using General Method E.

10 (rrrr) 3-Amino-7-(6-amino-1-methyl-3-azabicyclo[3.2.0]hept-3-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 125**) (mp >190°C, MS ES: m/z 390 (MH⁺)) from 3-amino-7-(6-amino-1-methyl-3-azabicyclo[3.2.0]hept-3-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 28fff) using General Method E.

15 (ssss) 3-Amino-7-(4-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 126**) (mp 201°C, MS ES: m/z 378 (MH⁺)) from 3-amino-7-(4-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 28hhh) using General Method E.

20 (tttt) 3-Amino-7-(3-aminomethylazetidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 127**) (MS ES: m/z 350 (MH⁺)) from 3-amino-7-(3-aminomethylazetidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 28ggg) using General Method E.

25 (uuuu) cis-3-Amino-7-(3-aminomethyl-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (**Compound 128**) (mp 169-179°C, MS ES: m/z 382 (MH⁺)) from cis-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28iii) using General Method A.

30

-159-

(vvvvv) trans-3-Amino-7-(3-aminomethyl-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (Compound 129) (mp 144-147°C, MS ES: m/z 382 (MH⁺)) from trans-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Example 28jjj) using General Method A.

(wwwww) 3-Amino-7-(1-amino-5-azaspiro[2.4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (Compound 130) (mp >250°C, MS APCI: m/z 376 (MH⁺)) from [1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-(1-amino-5-azaspiro[2.4]hept-5-yl)]carbamic acid *tert*-butyl ester (Example 28kkk) using General Method A.

(xxxxx) 3-Amino-7-(3a-aminomethyloctahydroisoindol-2-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (Compound 131) (mp >250°C, MS APCI: m/z 418 (MH⁺)) from 1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-aminomethyloctahydroisoindol-2-yl]carbamic acid *tert*-butyl ester (Example 28lll) using General Method A.

(yyyyy) 3-Amino-7-(3a-aminomethyloctahydroisoindol-2-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride (Compound 132) (mp >250°C, MS APCI: m/z 402 (MH⁺)) from 1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-aminomethyloctahydroisoindol-2-yl]carbamic acid *tert*-butyl ester (Example 28mmm) using General Method A.

(zzzzz) 3-Amino-7-[(R)-3-((S)-1-aminoethylpyrrolidin-1-yl)]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione hydrochloride (Compound 133) (mp 228-230°C, MS APCI: m/z 378 (MH⁺)) from {(S)-1-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (Example 28nnn) using General Method E.


-160-

(aaaaaa) 3-Amino-7-(6-amino-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione hydrochloride

(Compound 134) (mp 180-182°C, MS ES: m/z 346(MH^+)) from [3-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-azabicyclo[3.1.0]hex-6-yl]carbamic acid tert-butyl ester (Example 28n) using General Method A.

EXAMPLE 10

Synthesis of {7-[(S)-3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-5-benzylamino-1-cyclopropyl-6,8-difluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl}carbamic acid tert-butyl ester




A solution of {7-[(S)-3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-1,4-dihydro-2,4-dioxo-5,6,8-trifluoro-2H-quinazolin-3-yl}carbamic acid tert-butyl ester from Example 8 (0.51 g, 0.914 mmol), triethylamine (1.3 mL, 9.3 mmol), and benzylamine (0.50 mL, 4.6 mmol) in dimethyl sulfoxide (7.5 mL) is heated at 100°C for 16 hours in a sealed glass tube. The mixture is concentrated under high vacuum and the residue purified by column chromatography (1:2 ethyl acetate/hexanes) to afford {7-[(S)-3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-5-benzylamino-1-cyclopropyl-6,8-difluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl}carbamic acid tert-butyl ester as a solid (0.51 g). ^1H NMR ($CDCl_3$): δ 8.50 (bs, 1H), 7.37-7.18 (m, 5H), 6.56 (bs, 1H), 4.76-4.62 (bd, 1H), 4.59-4.48 (m, 2H), 4.34-4.18 (m, 1H), 3.93-3.49 (m, 3H), 3.48-3.20 (m, 2H), 2.24-2.04 (m, 1H),

-161-

1.96-1.74 (m, 1H), 1.49 (s, 9H), 1.46 (s, 9H), 1.13-1.00 (m, 2H), 0.75-0.61 (m, 2H).

EXAMPLE 11

Synthesis of {7-[(S)-3-(tert-Butoxycarbonyl-N-methylamino)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}-methylcarbamic acid tert-butyl ester




To a solution of {7-[(S)-3-(tert-butoxycarbonyl-N-methylamino)-pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester from Example 8 (0.238 g, 0.421 mmol) in *N,N*-dimethylformamide (5 mL) at -78°C under nitrogen atmosphere is added sodium hydride (60% dispersion in mineral oil, 0.025 g, 0.631 mmol). After 30 minutes, iodomethane (0.052 mL, 0.842 mmol) is added, and the reaction mixture is warmed to room temperature. After 1 hour, the reaction mixture is diluted with ethyl acetate and washed with saturated ammonium chloride solution, water, and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate concentrated. The resulting residue is purified by flash column chromatography (1:1 ethyl acetate/hexanes) to afford the title compound (0.118 g). ¹H NMR (CDCl₃): δ 7.63 (dd, 1 H), 4.78 (bs, 1 H), 3.87-3.45 (m, 6 H), 2.62-2.44 (m, 6 H), 2.31-2.12 (m, 2 H), 1.49-1.47 (m, 18 H), 1.23-1.07 (m, 2 H), 0.75-0.53 (m, 2H). MS CI: *m/z* 582 (MH⁺).

EXAMPLE 12

*Synthesis of {5-Amino-7-[(S)-3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-6,8-difluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl}carbamic acid *tert*-butyl ester*

-162-




5 {7-[(S)-3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-5-benzylamino-1-cyclopropyl-6,8-difluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl} carbamic acid *tert*-butyl ester from Example 10 (0.435 g, 0.676 mmol) in tetrahydrofuran (20 mL) is hydrogenated at room temperature and atmospheric pressure over 20% palladium hydroxide on carbon (0.114 g) for 27 hours. The mixture is filtered through Celite, the solid is washed with chloroform, and the combined filtrates concentrated under vacuum. Purification by column chromatography (2:3 ethyl acetate/hexanes) gives {5-amino-7-[(S)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-6,8-difluoro-1,4-dihydro-2,4-dioxo-2H-quinazolin-3-yl}-carbamic acid *tert*-butyl ester as a solid (0.284 g). ¹H NMR (CDCl₃): δ 6.61 (bs, 1H), 5.94 (bs, 2H), 4.82-4.69 (bd, 1H), 4.40-4.21 (m, 1H), 4.00-3.59 (m, 3H), 3.55-3.41 (m, 1H), 3.35-3.21 (m, 1H), 2.32-2.08 (m, 1H), 2.01-1.80 (m, 1H), 1.50 (s, 9H), 1.46 (s, 9H), 1.17-1.01 (m, 2H), 0.77-0.63 (m, 2H).

15

EXAMPLE 13

Synthesis of 8-Chloro-1-cyclopropyl-6-fluoro-3-methylamino-7-[(S)-3-methylaminopyrrolidin-1-yl]-1H-quinazoline-2,4-dione




20 Hydrogen chloride gas is bubbled into diethyl ether (30 mL) for 15 minutes. The resulting solution is then cooled to 0°C and added to {7-[(S)-

-163-

3-(*tert*-butoxycarbonylmethylamino)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,4-dihydro-2*H*-quinazolin-3-yl}methyl-carbamic acid *tert*-butyl ester from Example 11 (0.118 g, 0.202 mmol). The reaction mixture is slowly warmed to room temperature. After 30 hours, the precipitate is filtered, washed with ether and hexanes, and dried under vacuum to afford the hydrochloride salt of the title compound as a solid (0.0632 g, mp 77-79°C). MS Cl: m/z 382 (MH^+).

EXAMPLE 14

Synthesis of (8-Chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid tert-butyl ester




To a solution of (3-chloro-2-cyclopropylamino-4,5-difluorobenzoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 6c) (1.93 g, 5.34 mmol) in tetrahydrofuran (50 mL) is added potassium carbonate (3.69 g, 26.7 mol) and triphosgene (2.06 g, 6.95 mmol). The reaction mixture is refluxed for 90 minutes, cooled to room temperature, and diluted with ethyl acetate. The organic layer is washed with water and brine, then dried over $MgSO_4$ and filtered. The filtrate is concentrated under vacuum and purified via flash column chromatography (1:2 ethyl acetate/hexanes) to afford the title compound (1.27 g). MS EI: m/z 386 (M^+).

EXAMPLE 15


Synthesis of 2-Amino-4, 5-difluorobenzoic acid, 2,2-dibenzylhydrazide

-164-




4,5-Difluoroanthranilic acid (4.73 g, 27.3 mmol) and N,N-dibenzylhydrazine (8.69 g, 42 mmol) are combined in 200 mL of methylene chloride. 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDAC) (7.85 g, 41 mmol) is then added to this solution, and the mixture is stirred for 18 hours at 25°C. The solution is washed with saturated NaHCO₃, brine, and dried over magnesium sulfate. The solution is concentrated to give 15.8 g of a dark oil and purified via chromatography (SiO₂, CHCl₃) to give 3.4 g of the title compound as a solid. MS CI: *m/z* 368 (MH⁺).

EXAMPLE 16


Synthesis of 3-Dibenzylamino-6, 7-difluoro-1H-quinazoline-2, 4-dione

2-Amino-4,5-difluorobenzoic acid, 2,2-dibenzylhydrazide (Example 15) (0.81 g, 2.2 mmol) and triphosgene (0.33 g, 1.1 mmol) are combined in 100 mL of methylene chloride and stirred at 25°C for 20 hours. The solution is poured into 200 mL of saturated NaHCO₃, the layers are separated, and the aqueous layer is washed three times with ethyl acetate. The combined organic layers are dried over magnesium sulfate and then concentrated to give 0.86 g of the title compound as a solid. MS CI: *m/z* 394 (MH⁺).

EXAMPLE 17

*Synthesis of 3-Dibenzylamino-6,7-difluoro-1-substituted-1H-quinazoline-2,4-dione**(a) 3-Dibenzylamino-6,7-difluoro-1-cyclopropylmethyl-1H-quinazoline-2,4-dione*

A solution of 3-dibenzylamino-6,7-difluoro-1H-quinazoline-2,4-dione (Example 16) (0.43 g, 1.1 mmol) in 10 mL of *N,N*-dimethylformamide is added to a suspension of sodium hydride (0.05 g, 1.3 mmol) in 10 mL of *N,N*-dimethylformamide and stirred for 30 minutes. Bromomethylcyclopropane (0.16 mL, 1.6 mmol) is added, and the mixture is stirred at 25°C for 18 hours. The reaction is quenched with 1 mL of water and concentrated in vacuo. The residue is dissolved in chloroform, washed with water, brine, and dried over magnesium sulfate. The solution is concentrated and purified via chromatography (SiO_2 , CHCl_3) to give 0.33 g of the title compound as a solid. MS CI: m/z 448 (MH^+).

(b) 3-Dibenzylamino-6,7-difluoro-1-ethyl-1H-quinazoline-2,4-dione

A solution of 3-dibenzylamino-6,7-difluoro-1H-quinazoline-2,4-dione (Example 16) (0.43 g, 1.1 mmol) in 10 mL of *N,N*-dimethylformamide is added to a suspension of sodium hydride (0.05 g, 1.3 mmol) in 10 mL of *N,N*-dimethylformamide and stirred for 30 minutes. Ethyl iodide (0.13 mL, 1.6 mmol)

-166-

is added, and the mixture is stirred at 25°C for 18 hours. The reaction is quenched with 1 mL of water and concentrated in vacuo. The residue is dissolved in chloroform, washed with water, brine, and dried over magnesium sulfate. The solution is concentrated and purified via chromatography (SiO₂, CHCl₃) to give
5 0.34 g of the title compound as a solid, mp 133-135°C, MS CI, *m/z* 422 (MH⁺).

EXAMPLE 18

Chlorination

General Chlorination Method

To a solution of a benzoylhydrazinecarboxylic acid *tert*-butyl ester
10 (Example 7) in acetic acid (5 mL) is added *N*-chlorosuccinimide (1.2 eq.). After 1 hour, the reaction mixture is diluted with ethyl acetate and washed with saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄ and filtered. The filtrate is concentrated under vacuum and purified via flash column chromatography (ethyl acetate/hexanes) to afford the corresponding chloro-
15 benzoylhydrazinecarboxylic acid *tert*-butyl ester.

The following compounds are synthesized according to the General Chlorination Method above:

- (a) *N'*{4-[3-(*tert*-Butoxycarbonylaminomethyl)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS CI: *m/z* 542 (MH⁺)) from *N'*{4-[3-(*tert*-butoxycarbonylaminomethyl)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluorobenzoyl}-hydrazinecarboxylic acid *tert*-butyl ester (Example 7a).
20
- (b) *N'*-4-[4-(*tert*-Butoxycarbonylpiperazin-1-yl)-3-chloro-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (MS
25 CI: *m/z* 528 (MH⁺)) from *N'*-4-[4-(*tert*-butoxycarbonylpiperazin-1-yl)-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (Example 7b).
- (c) *N'*{4-[3-(*tert*-Butoxycarbonylaminomethyl)-3-methylpyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid

-167-

tert-butyl ester (MS CI: m/z 556 (MH^+)) from N' -{4-[3-(*tert*-butoxycarbonylaminoethyl)-3-methylpyrrolidin-1-yl]-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 7c).

(d) N' -[4-(6-*tert*-Butoxycarbonylamino-3-azabicyclo[3.1.0]hex-3-yl)-3-chloro-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/z 540 (MH^+)) from N' -[4-(6-*tert*-butoxycarbonylamino-3-azabicyclo[3.1.0]hex-3-yl)-2-cyclopropylamino-5-fluorobenzoyl]-hydrazinecarboxylic acid *tert*-butyl ester (Example 7d).

(e) N' -{4-[(*S*)-3-(*tert*-Butoxycarbonyl-*N*-methylamino)pyrrolidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/z 542 (MH^+)) from N' -{4-[(*S*)-3-(*tert*-butoxycarbonyl-*N*-methylamino)pyrrolidin-1-yl]-2-cyclopropylamino-5-fluorobenzoyl}-hydrazinecarboxylic acid *tert*-butyl ester (Example 7e).

(f) N' -{4-[(*S*)-3-(*tert*-Butoxycarbonyl-*N*-amino)pyrrolidin-1-yl]-3-chloro-2-(2,4-difluoroanilino)-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester ($CDCl_3$): δ 9.70 (bs, 1H), 7.76 (d, 1H), 6.93-6.82 (m, 1H), 6.70-6.62 (m, 2H), 6.48-6.34 (m, 2H), 4.91-4.87 (bs, 1H), 4.29-4.20 (m, 1H), 3.78-3.60 (m, 2H), 3.50-3.28 (m, 2H), 2.33-2.20 (m, 1H), 1.94-1.66 (m, 1H), 1.44 (s, 9H), 1.42 (s, 9H), (MS CI: m/z 601 (MH^+)) from N' -{4-[(*S*)-3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2-(2,4-difluoroanilino)-5-fluorobenzoyl}-hydrazinecarboxylic acid *tert*-butyl ester (Example 7f).

(g) N' -[4-((*S*)-7-*tert*-Butoxycarbonylamino-5-azaspiro[2.4]hept-5-yl)-3-chloro-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (MS CI: 554 (MH^+)) from N' -[(*S*)-4-(7-*tert*-butoxycarbonylamino-5-azaspiro[2.4]hept-5-yl)-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic acid *tert*-butyl ester (Example 7k).

(h) N' -[4-(Pyrrolidin-1-yl)-3-chloro-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic *tert*-butyl ester (MS CI: m/z 413 (MH^+)) from N' -[4-(pyrrolidin-1-yl)-2-cyclopropylamino-5-fluorobenzoyl]hydrazinecarboxylic *tert*-butyl ester (Example 7m).

-168-

(i) N' -{4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-3-chloro-5-fluoro-2-isopropylaminobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS: m/z 530 (MH^+)) from N' -{4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-5-fluoro-2-isopropylaminobenzoyl}hydrazine carboxylic acid *tert*-butyl ester (Example 7n).


(j) N' -{4-[3-(*tert*-Butoxycarbonylamino)pyrrolidin-1-yl]-2-cyclobutylamino-3-chloro-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (MS EI: m/e 542 (MH^+)) from N' -{4-[3-(*tert*-butoxycarbonylamino)-pyrrolidin-1-yl]-2-cyclobutylamino-5-fluorobenzoyl}hydrazinecarboxylic acid *tert*-butyl ester (Example 7q).

(k) N' -(1-{4-[3-(*tert*-Butoxycarbonylamino-methyl)piperidin-1-yl]-3-chloro-2-cyclopropylamino-5-fluorophenyl}methanoyl)hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/e 556 (MH^+)) from N' -(1-{4-[3-(*tert*-butoxycarbonylamino-methyl)piperidin-1-yl]-2-cyclopropylamino-5-fluorophenyl}methanoyl)hydrazinecarboxylic acid *tert*-butyl ester (Example 7z).


(l) N' -[1-(4-{3-[(*tert*-Butoxycarbonylisopropylamino)methyl]-pyrrolidin-1-yl}-3-chloro-2-cyclopropylamino-5-fluorophenyl)methanoyl]-hydrazine-carboxylic acid *tert*-butyl ester (MS CI: m/e 584 (MH^+)) from N' -[1-(4-{3-[(*tert*-butoxycarbonylisopropylamino)methyl]pyrrolidin-1-yl}-2-cyclopropylamino-5-fluorophenyl)methanoyl]hydrazinecarboxylic acid *tert*-butyl ester (Example 7aa).

(m) N' -(1-{4-[3-(*tert*-Butoxycarbonylamino-methyl)pyrrolidin-1-yl]-3,5-dichloro-2-cyclopropylaminophenyl}methanoyl)hydrazinecarboxylic acid *tert*-butyl ester (MS CI: m/e 558 (MH^+)) from N' -(1-{4-[3-(*tert*-butoxycarbonylamino-methyl)pyrrolidin-1-yl]-5-chloro-2-cyclopropylaminophenyl}methanoyl)-hydrazinecarboxylic acid *tert*-butyl ester (Example 7bb).

EXAMPLE 19

*Chlorination of Substituted Benzoic Acids**(a) Synthesis of 3-Chloro-4,5-difluoroanthranilic acid*


5 To a solution of 4,5-difluoroanthranilic acid (2.45 g, 14.2 mmol) in dichloromethane (25 mL) is added acetic acid (10 mL) and hypochlorous acid *tert*-butyl ester (1.75 mL, 15.6 mmol). After 2 hours, the reaction mixture is diluted with ethyl acetate and washed with water and brine. The organic layer is dried over MgSO₄, filtered, and concentrated. The resulting residue is purified via
10 flash column chromatography (5% isopropyl alcohol/1% formic acid/94% dichloromethane) to afford 3-chloro-4,5-difluoroanthranilic acid as an oil (2.97 g). MS CI: *m/z* 206 (M⁺).

(b) 5-Chloro-4-fluoroanthranilic acid

15 To a solution of 4-fluoroanthranilic acid (0.882 g, 5.69 mmol) in methylene chloride (50 mL) and acetic acid (10 mL) is added *tert*-butyl hypochlorite (0.71 mL, 6.25 mmol) solution in dichloromethane (1 mL) dropwise over 1 minute. After 90 minutes, the reaction mixture is washed with water and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate
20 concentrated to afford 5-chloro-4-fluoroanthranilic acid (1.20 g). MS CI: *m/e* 188 (M⁺).

-170-


EXAMPLE 20

Synthesis of 2-Bromo-3-chloro-4,5-difluorobenzoic acid

A solution of cuprous bromide (10.96 g, 49.2 mmol) in acetonitrile
5 (100 mL) is cooled to 0°C and then *tert*-butyl nitrite (7.31 mL, 61.5 mmol) and
3-chloro-4,5-difluoroanthranilic acid (Example 19) (8.51 g, 41.0 mmol) are added.
The mixture is slowly warmed to room temperature, and after 20 hours, the
solvent is removed under vacuum. The resulting residue is dissolved in ethyl
acetate and washed with 1.0 M hydrochloric acid, water, and brine. The organic
10 layer is dried over MgSO₄ and filtered. The filtrate is concentrated under vacuum
and purified via flash column chromatography (5% isopropyl alcohol/1% formic
acid/94% dichloromethane) to afford 2-bromo-3-chloro-4,5-difluorobenzoic acid
as a solid (7.96 g). MS CI: *m/z* 271 (MH⁺).

EXAMPLE 21

15 *Synthesis of 4-[3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-2,3-*
difluorobenzoic acid ethyl ester




To a solution of 4-[3-(*tert*-butoxycarbonylamino)pyrrolidin-1-yl]-2,3-
20 difluorobenzoic acid ethyl ester (Example 2h) (3.50 g, 9.50 mmol) in
dichloromethane (80 mL) is added acetic acid (8 mL) and *tert*-butyl hypochlorite
(1.75 mL, 15.6 mmol). After 20 minutes, the reaction mixture is diluted with ethyl
acetate and washed with water and brine. The organic layer is dried over sodium

-171-

sulfate, filtered, and the filtrate concentrated. The resulting residue is purified by column chromatography (1:4 ethyl acetate/hexanes) to afford the title compound (3.75 g) as an oil. ^1H NMR (CDCl_3): δ 7.69 (dd, 1H), 4.85-4.76 (bd, 1H), 4.50-4.20 (m, 3H), 3.90-3.74 (m, 2H), 3.62-3.36 (m, 2H), 2.40-2.20 (m, 1H), 1.91-1.84 (m, 1H), 1.45 (s, 9H), 1.40 (s, 9H).

EXAMPLE 22


Synthesis of 2-cyclopropylamino-4,5-difluorobenzoic acid methyl ester

To a solution of 2-cyclopropylamino-4,5-difluorobenzoic acid (Example 5a) (6.0 g, 28.1 mmol), dichloromethane (100 mL), and methanol (20 mL) is added (trimethylsilyl)diazomethane (2.9 M solution in hexane) dropwise until the evolution of gas stops. The solution is stirred at room temperature for 0.5 hour and 88% formic acid is added dropwise until the evolution of gas again stops. Water is added, and the solution is extracted with dichloromethane, dried over MgSO_4 , filtered, and the solvent removed under reduced pressure. The residue is purified by silica gel column chromatography using hexane/EtOAc (9:1) as the eluent to afford a 5.4 g of the title compound as a clear oil. MS CI: m/z 228 (MH^+).

EXAMPLE 23


(a) Synthesis of 1-Cyclopropyl-6,7-difluoro-1H-quinazoline-2,4-dione

-172-



To a solution of 2-cyclopropylamino-4,5-difluorobenzoic acid ethyl ester (Example 22) (5.0 g, 22.0 mmol) in dry dichloromethane (120 mL) under a N₂ atmosphere is added chlorosulfanyl isocyanate (3.11 g, 22 mmol). The solution is reacted at room temperature for 4 hours, and the solvent is removed under reduced pressure. The residue is cooled to -20°C, and a cold brine solution (100 mL) buffered with NaHCO₃ is added. The solution is warmed to room temperature for 1 hour. The volume is reduced by half with a stream of air, and the solid is collected by filtration. The dry solid is added to a solution of triethylamine (5 mL) in THF (250 mL) and refluxed overnight. The reaction is cooled, the solvent removed under reduced pressure, dissolved in water (100 mL), and acidified to pH 1-2 with 1.0 M hydrochloric acid. The resulting precipitate is collected via filtration to afford 2.0 g of the title compound. MS CI: *m/z* 239 (MH⁺).

(b) *1-Cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione*




A solution of 1-cyclopropyl-3-(3-methoxy-2,4,5-trifluorobenzoyl)urea (Example 27b, 0.94 g, 3.26 mmol) in tetrahydrofuran (20 mL) and *N,N*-dimethylformamide (5 mL) is treated with sodium hydride (0.275 g, 6.8 mmol, 60% in mineral oil dispersion) and heated at reflux for 16 hours. The mixture is cooled, treated with saturated NH₄Cl and extracted with dichloromethane. The

-173-

organic layer is then washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. Flash column chromatography (1:3 ethyl acetate/hexanes) afforded the title compound. MS (EI, $M+1$) m/z 269.


(c) *Synthesis of 8,9-Difluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diaza-phenalene-4,6-dione*



A solution of 1.5 g (6.2 mmol) of 7,8-difluoro-3-methyl-3,4-dihydro-2H-1,4-benzoxazine-5-carboxylic acid methyl ester (Example 32a) in 25 mL of dichloromethane is cooled to -20°C and treated dropwise with 0.92 g (6.5 mmol) of chlorosulfonyl isocyanate. The reaction is stirred at room temperature for 18 hours and the solvent removed in vacuo. The residue is treated with a saturated sodium bicarbonate solution and extracted with dichloromethane. The organic layer is dried (MgSO_4), filtered, and concentrated in vacuo. The residue is dissolved in 50 mL of tetrahydrofuran, and the resulting solution is treated with 5.0 g (5.0 mmol) of triethylamine. After heating at reflux for 4 hours, the solvent is removed in vacuo, and the residue is triturated with 5 mL of dichloromethane/ethyl acetate (80:20). The insoluble material is removed by filtration, washed with the above solvent mixture (2 mL) to give 0.25 g of the title compound, mp $259-261^{\circ}\text{C}$. The filtrate is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane/ethyl acetate (80:20) to give 0.38 g of starting material. Continued elution afforded an additional 0.2 g of the title compound. ^1H NMR (400 MHz, CDCl_3): 11.78 (s, 1H), 7.50 (m, 1H), 4.68 (m, 1H), 4.52 (d, 1H), 4.17 (d, 1H), 1.23 (d, 3H).


-174-

(d) *Synthesis of 8,9-Difluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione*



A solution of 1.07 g (4.4 mmol) of 5,6-difluoro-2-methyl-1,2,3,4-tetrahydro-quinoline-8-carboxylic acid methyl ester (Example 32) in 25 mL of dichloromethane is cooled to -20°C and treated dropwise with 0.85 g (6.0 mmol) of chlorosulfonyl isocyanate. The reaction is stirred at -20°C for 1 hour, treated with 1.0 g (12 mmol) of solid sodium acetate, and the solvent is removed in vacuo. The residue is cooled in an ice bath and triturated with a saturated solution of sodium acetate in brine. The precipitate is removed by filtration, washed with the sodium acetate solution and dried in vacuo. The dried precipitate is washed with ether to remove starting material and dried in vacuo. The dried solid is suspended in 40 mL of dry tetrahydrofuran, cooled to 5°C and treated portionwise with 1.2 g (12 mmol) of sodium tert-butoxide. After the addition is complete, the reaction mixture is stirred at 5°C for 30 minutes, the bath is removed, and the reaction is stirred to room temperature over 1 hour. The solvent is removed in vacuo; the residue is dissolved in water, cooled to 5°C and acidified with formic acid. The resulting precipitate is removed by filtration, washed with water and dried in vacuo affording 0.8 g of the title compound, mp 212-214°C.

(e) *1-Cyclopropyl-6-fluoro-7-methylsulfanyl-1H-pyrido[2,3-d]pyrimidine-2,4-dione*



-175-

A solution of 4.1 g (15.2 mmol) of ethyl 2-cyclopropylamino-5-fluoro-6-methylsulfanylnicotinate (Example 34) in 125 mL of dichloromethane is cooled to -20°C and treated dropwise with 5.24 g (37 mmol) of chlorosulfonyl isocyanate. The reaction is stirred at -20°C for 3 hours and allowed to come to room temperature overnight. The mixture is recooled to -20°C and treated with 6.8 g (80 mmol) of solid sodium acetate. After stirring for 1 hour without cooling, the solvent is removed in vacuo without heating. The residue is triturated with a saturated solution of sodium acetate in brine and extracted with ethyl acetate (3 × 75 mL). The combined organic layers are dried (MgSO₄), filtered and evaporated in vacuo. The residue is suspended in 100 mL of tetrahydrofuran and treated with 3.6 g (32 mmol) of solid sodium tert-butoxide and stirred at room temperature for 36 hours. The solvent is removed in vacuo, the residue triturated with 100 mL of 0.1 M formic acid and extracted with ethyl acetate (3 × 100 mL). The combined organic layers are washed with water, dried (MgSO₄), filtered and evaporated in vacuo to give a residue which is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane to give 0.75 g of the title compound, mp 220-222°C.

General Procedure A

Sodium hydride (3 eq., 60% mineral oil dispersion) is added portionwise to a solution of 1-cyclopropyl-3-benzoylurea (Example 27) in 20:1 tetrahydrofuran/ *N,N*-dimethylformamide at -25°C. The mixture is stirred at room temperature for 30 minutes, then refluxed overnight. It is poured into ice water, and the solution is acidified with 10% hydrochloric acid and extracted with ethyl acetate. The extract is washed with brine, dried, and concentrated. The residue is purified by silica gel column chromatography (ethyl acetate/hexanes) to afford 8-substituted 1-cyclopropyl-6,7-difluoro-1H-quinazoline-2,4-diones.

The following compounds are synthesized according to General Procedure A. of Example 23:

(e) 1-Cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione
(¹H NMR (200 MHz, CDCl₃): δ 8.61 (bs, 1H), 7.83 (dd, 1H), 3.37-3.35 (m, 1H),

-176-

2.60 (dd, 3H), 1.31-1.12 (m, 2H), 0.71-0.63 (m, 2H) from 1-cyclopropyl-3-(2,4,5-trifluoro-3-methylbenzoyl)urea (Example 27c).


(f) 1-Cyclopropyl-6,7-difluoro-5-methyl-1H-quinazoline-2,4-dione (¹H NMR (200 MHz, CDCl₃): δ 8.37 (bs, 1H), 7.96 (dd, 1H), 2.81-2.74 (m, 1H), 2.61 (d, 3H), 1.26-1.14 (m, 2H), 0.81-0.73 (m, 2H)) from 1-cyclopropyl-3-(3,4,6-trifluoro-2-methylbenzoyl)urea (Example 27d).

(g) 1-cyclopropyl-6,7-difluoro-8-ethyl-1H-quinazoline-2,4-dione (¹H NMR (200 MHz, CDCl₃): δ 9.41 (bs, 1H), 7.86-7.77 (m, 1H), 3.40-3.20 (m, 3H), 1.30-1.10 (m, 5H), 0.75-0.67 (m, 2H)) from 1-cyclopropyl-3-(3-ethyl-2,4,5-trifluorobenzoyl)urea (Example 27e).

(i) 7-Chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (mp 214-216°C) from 1-cyclopropyl-3-(2,6-dichloro-5-fluoro-pyridine-3-carbonyl)urea (Example 27a).

EXAMPLE 24

(a) *Synthesis of 3-Amino-1-cyclopropyl-6,7-difluoro-1H-quinazoline-2,4-dione*




To a solution of 1-cyclopropyl-6,7-difluoro-1H-quinazoline-2,4-dione (Example 23a, 2.0 g, 8.92 mmol) in dioxane (7.5 mL) and dimethylformamide (7.5 mL) is added NaH (60% dispersion in oil, 428 mg, 10.7 mmol). The solution is heated to 60°C for 10 minutes and cooled to room temperature. To the cooled solution is added O-(2,4-dinitrophenyl)hydroxylamine (1.77 g, 8.92 mmol), and the solution is heated to 80°C for 30 minutes. The resulting red solution is cooled to room temperature, poured over crushed ice, and extracted with EtOAc. The combined organic layers are dried over MgSO₄, filtered, and the solvent removed

-177-

under reduced pressure. The resulting solid is triturated with Et₂O and air dried to afford 1.44 g of the title compound. MS CI: m/z 254 (MH⁺).

(b) *3-Amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione*




5

A solution of 3.83 g (15 mmol) of 7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 23i) in 50 mL of dimethylformamide-dioxane (1:1) is cooled to 0°C and treated portionwise with 0.8 g (20 mmol) of 60% sodium hydride/mineral oil. The reaction is stirred to room temperature for 30 minutes, re-cooled to 0°C and 3.2 g (16 mmol) of 2,4-dinitrophenylhydroxylamine is added all at once. After stirring at room temperature overnight, the dioxane is removed in vacuo and the residue is diluted to 300 mL with ice and water and stirred at 5°C for 1 hour. The precipitate is removed by filtration, washed with water, ether and dried in vacuo to give 2.65 g of the title compound, mp 192-194°C. A second crop, (0.6 g) could be isolated by evaporating the filtrate and triturating the residue with petroleum ether.

15

(c) *3-Amino-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydro-pyrido[2,3-d]pyrimidine-7-thiosulfonic acid*



20

A solution of 0.47 g (1.57 mmol) of 1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydro-pyrido[2,3-d]pyrimidine-7-thiosulfonic acid (Example 35) in 20 mL of dimethylformamide/dioxane (1:1) is treated portionwise with 0.08 g (2.0 mmol) of 60% sodium hydride/mineral oil. The reaction is stirred at room

-178-

temperature for 1 hour and 0.32 g (1.6 mmol) of 2,4-dinitrophenylhydroxylamine is added all at once. After stirring at room temperature overnight, the dioxane is removed in vacuo and the residue is diluted to 100 mL with ice and water and stirred at 5°C for 1 hour. The aqueous solution is extracted with ethyl acetate (3 × 50 mL) with the combined organic layers being washed with water, dried (MgSO₄), filtered and evaporated in vacuo to give 0.4 g of the title compound, mp 167-169°C.

General Procedure A (Example 24)

To a solution of a 1-cyclopropyl-6,7-difluoro-1H-quinazoline-2,4-dione (Example 23) in 1:1 dry tetrahydrofuran or dioxane and dry *N,N*-dimethylformamide is added portionwise sodium hydride (1.1 eq., 60% mineral oil dispersion) at room temperature. After stirring at 50°C for 20 to 30 minutes, the resulting solution is cooled to room temperature and 2,4-dinitrophenylhydroxylamine (4 eq.) is added. The mixture is heated at 60°C to 80°C for 30 minutes, cooled, and the dioxane removed reduced pressure. The residue is poured into ice water and extracted with ethyl acetate. The organic layers are combined, washed with brine, dried with Na₂SO₄, and concentrated in vacuo. The residue is purified by flash column chromatography (ethyl acetate/hexanes) to afford a 3-amino-1-cyclopropyl-6,7-difluoro-1H-quinazoline-2,4-dione.

The compounds are synthesized according to General Procedure A of Example 24:

(d) 3-Amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (MS EI: *m/z* 284 (MH⁺)) from 1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 23b).

(e) 3-Amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (¹H NMR (200 MHz, CDCl₃): δ 7.86 (dd, 1H), 5.20 (bs, 2H), 3.48-3.38 (m, 1H), 2.61 (dd, 3H), 1.29-1.17 (m, 2H), 0.73-0.62 (m, 2H)) from 1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 23e).

-179-

(f) 3-Amino-1-cyclopropyl-6,7-difluoro-5-methyl-1H-quinazoline-2,4-dione (¹H NMR (200 MHz, CDCl₃): δ 7.45 (dd, 1H), 5.28 (bs, 2H), 2.94-2.83 (m, 1H), 2.77 (d, 3H), 1.41-1.31 (m, 2H), 0.98-0.93 (m, 2H)) from 1-cyclopropyl-6,7-difluoro-5-methyl-1H-quinazoline-2,4-dione (Example 23f).


(g) 3-Amino-1-cyclopropyl-6,7-difluoro-8-ethyl-1H-quinazoline-2,4-dione (¹H NMR (200 MHz, CDCl₃): δ 7.84-7.75 (t, 1H), 5.37 (bs, 2H), 3.50-3.40 (m, 1H), 3.37-3.21 (m, 2H), 1.29-1.15 (m, 5H), 0.75-0.66 (m, 2H)) from 1-cyclopropyl-6,7-difluoro-8-ethyl-1H-quinazoline-2,4-dione (Example 23g).

(h) 5-Amino-8,9-difluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione (mp 171-173°C) from 8,9-difluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione (Example 23c).

(i) 2-Amino-8,9-difluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione, mp 140-142°C, ¹H NMR (400 MHz, CDCl₃): 7.90 (m, 1H), 5.30 (bs, 2H), 5.07 (m, 1H), 3.08 (m, 1H), 2.85 (m, 1H), 2.15 (m, 1H), 1.95 (m, 1H), 1.31 (d, 3H)) from 8,9-difluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione (Example 23d).

EXAMPLE 25

Synthesis of {3-Amino-7-[(S)-3-(tert-butoxycarbonylamino)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione



To a solution of 3-amino-1-cyclopropyl-6,7-difluoro-1H-quinazoline-2,4-dione (Example 24a) (50.0 mg, 0.20 mmol) and triethylamine (74 mg, 0.74 mmol) in acetonitrile (3 mL) is added (S)-pyrrolidin-3-yl carbamic acid *tert*-butyl ester (74 mg, 0.40 mmol). The solution is refluxed for 18 hours, the solvent is removed under reduced pressure, and the residue triturated with water. The solid is collected by filtration and air dried to afford 74 mg of the title compound. MS CI: *m/z* 420 (MH⁺).

-180-

EXAMPLE 26

Synthesis of Benzamides

General Procedure A

A solution of an appropriately substituted benzoic acid, oxalyl chloride (1.5 eq.), and dichloromethane is treated with *N,N*-dimethylformamide (2 drops), the solution is stirred at room temperature for 2 hours. The mixture is concentrated under reduced pressure, and the residue is dissolved in dry THF and slowly added to a solution of ammonia in diethyl ether at -70°C. The mixture is then allowed to warm to ambient temperature and stirred for 30 minutes. The mixture is then filtered and the resulting solid dissolved in ethyl acetate, washed with water, and dried over Na₂SO₄. The solvent is then concentrated in vacuo to provide the desired amide.

The following compounds are synthesized according to the General Procedure A method above:


(a) 3,4,6-Trifluoro-2-methylbenzamide. (¹H NMR (200 MHz, CDCl₃): δ 6.89-6.77 (m, 1H), 6.41 (bs, 1H), 5.95 (bs, 1H), 2.39 (d, 3H)) from 3,4,6-trifluorobenzoic acid (Hagen S. et al., *Heterocycl. Chem.*, 1990;27[6]:1609-1616).

(b) 3-Ethyl-2,4,5-trifluorobenzamide, (¹H NMR (200 MHz, CDCl₃): δ 7.78-7.60 (m, 1H), 6.61 (bs, 2H), 2.67 (q, 2H), 1.15 (t, 3H)) from 3-ethyl-2,4,5-trifluorobenzoic acid (Takemura, PCT Int. Appl., 1996, WO 9623782).

EXAMPLE 27

Synthesis of 1-cyclopropyl-3-benzoyl substituted ureas.

(a) 1-Cyclopropyl-3-(2,6-dichloro-5-fluoro-pyridine-3-carbonyl)urea



-181-

A solution of 5.8 g (27.8 mmol) of 2,6-dichloro-5-fluoronicotinamide (*Chem. Pharm. Bull.*, 1987;35:2280) in 60 mL of dichloromethane is treated dropwise with 5.1 g (40 mmol) of oxalyl chloride, and the reaction mixture is heated at reflux for 18 hours. The solvent is removed in vacuo, and the residue is dissolved in 50 mL of dichloromethane, which is also removed in vacuo. The residue is dissolved in 50 mL of dichloromethane, cooled to -20°C and treated dropwise with 2.28 g (40 mmol) of cyclopropylamine. The reaction mixture is stirred at -20° to -10°C for ½ hour, then allowed to come to room temperature over 2 hours. The solvent is removed in vacuo and the residue is triturated with 25 mL of dichloromethane, which is also removed in vacuo to give 8.0 g of the title compound, mp 171-173°C.

General Procedure A

To a solution of substituted benzamide (Example 26) in 1,2-dichloroethane is added oxalyl chloride (1.5 eq.) and the mixture refluxed for 16 hours. The solvent is then removed under reduced pressure to obtain a crude isocyanate. The isocyanate is then taken up into dioxane, cooled to 0°C and treated with cyclopropylamine (1.5 eq.) in dioxane. The mixture is then warmed to ambient temperature and stirred for 3 hours. The mixture is then concentrated under reduced pressure, dissolved in ethyl acetate, washed with water, dried over Na₂SO₄ and evaporated under reduced pressure. The resulting residue is then purified by flash column chromatography (ethyl acetate/hexanes) to afford the 1-cyclopropyl-3-benzoyl urea.

The following compounds are synthesized according to the General Procedure A method above:

- (b) 1-Cyclopropyl-3-(2,4,5-trifluoro-3-methoxy-benzoyl)urea (MS ES (MH⁺): m/z 289) from 2,4,5-trifluoro-3-methoxybenzamide (Masuzawa K., Suzue S., Hirai K., Ishizaki T., *Eur. Pat. Appl.*, 1987, EP 230295).
- (c) 1-Cyclopropyl-3-(2,4,5-trifluoro-3-methylbenzoyl)urea (¹H NMR (200 MHz, CDCl₃): δ 8.69 (d, 1H), 8.48 (bs, 1H), 7.75-7.62 (m, 1H), 2.83-2.71 (m, 1H), 2.30 (dd, 3H), 0.87-0.78 (m, 2H), 0.67-0.59 (m, 2H)) from 2,4,5-

-182-


trifluoro-3-methylbenzamide (Masuzawa K., Suzue S., Hirai K., Ishizaki T., Eur. Pat. Appl., 1987, EP 237955).

(d) 1-Cyclopropyl-3-(3,4,6-trifluoro-2-methylbenzoyl)urea (^1H NMR (200 MHz, CDCl_3): δ 8.99 (bs, 1H), 8.37 (bs, 1H), 6.93-6.80 (m, 1H), 2.74-2.65 (m, 1H), 2.35 (d, 3H), 0.84-0.71 (m, 2H), 0.66-0.58 (m, 2H)) from 3,4,6-trifluoro-2-methylbenzamide (Example 26a).

(e) 1-Cyclopropyl-3-(3-ethyl-2,4,5-trifluorobenzoyl)urea (^1H NMR (200 MHz, CDCl_3): δ 8.76-8.69 (bd, 1H), 8.50 (bs, 1H), 7.76-7.63 (m, 1H), 2.84-2.72 (m, 3H), 1.27-1.20 (m, 3H), 0.91-0.81 (m, 2H), 0.71-0.61 (m, 2H)) from 3-ethyl-2,4,5-trifluorobenzamide (Example 26b).

EXAMPLE 28

(a) *Synthesis of {(R)-1-[(S)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}-carbamic acid tert-butyl ester*




15

A solution of 0.57 g (2.0 mmol) of 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d), 0.64 g (3.0 mmol) of ((S)-(R)-1-pyrrolidin-3-ylethyl)carbamic acid *tert*-butyl ester (*J. Het. Chem.*, 1992;29:1481), 0.81 g (8.0 mmol) of triethylamine and 10 mL of dimethyl sulfoxide is heated at 110°C for 4 hours. The reaction mixture is cooled to room temperature, poured into 150 mL of ice and water, and extracted with ethyl acetate (2 x 125 mL). The combined organic layers are washed with water, dried (MgSO_4), filtered and evaporated in vacuo to give 1.1 g of crude product. Chromatography on flash grade silica gel (230-400 mesh) eluting with dichloromethane/ethanol (9:1) provided 0.92 g of the title compound, mp 96-98°C.

25


-183-

(b) *Synthesis of {(S)-1-[(S)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}-carbamic acid tert-butyl ester*



5 A solution of 0.57 g (2.0 mmol) of 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d), 0.64 g (3.0 mmol) of ((S)-(S)-1-pyrrolidin-3-ylethyl)carbamic acid *tert*-butyl ester (*J. Het. Chem.*, 1992;29:1481), 0.81 g (8.0 mmol) of triethylamine, and 10 mL of dimethyl
10 sulfoxide is heated at 110°C for 4 hours. The reaction mixture is cooled to room temperature, poured into 150 mL of ice and water and extracted with ethyl acetate (2 × 125 mL). The combined organic layers are washed with water, dried (MgSO₄), filtered and evaporated in vacuo to give 1.08 g of crude product. Chromatography on flash grade silica gel (230-400 mesh) eluting with dichloromethane/ethanol (9:1) provided 0.9 g of the title compound, mp 87-89°C.

15 (c) *Synthesis of {(R)-1-[(S)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]-ethyl}-carbamic acid tert-butyl ester*




20 A solution of 0.5 g (1.9 mmol) of 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e), 0.64 g (3.0 mmol) of ((S)-(R)-1-pyrrolidin-3-ylethyl)carbamic acid *tert*-butyl ester, 0.81 g (8.0 mmol) of triethylamine and 10 mL of dimethyl sulfoxide is heated at 110°C for 24 hours. After thin layer chromatography showed incomplete reaction, an additional 0.43 g (2.0 mmol) of pyrrolidine derivative and 0.81 g (8.0 mmol) of triethylamine are

-184-

added and the reaction is heated at 120°C for 18 hours. The reaction mixture is cooled to room temperature, poured into 200 mL of ice and water and extracted with ethyl acetate (2 × 125 mL). The combined organic layers are washed with water, dried (MgSO₄), filtered and evaporated in vacuo to give 1.1 g of crude product. Chromatography on flash grade silica gel (230-400 mesh) eluting with ethyl acetate provided 0.4 g of the title compound, mp 86-88°C.


(d) *Synthesis of {(S)-1-[(S)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid tert-butyl ester*



A solution of 0.5 g (1.9 mmol) of 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e), 0.64 g (3.0 mmol) of ((S)-1-pyrrolidin-3-ylethyl)carbamic acid *tert*-butyl ester, 0.81 g (8.0 mmol) of triethylamine and 10 mL of dimethyl sulfoxide is heated at 110°C for 24 hours. After thin layer chromatography showed incomplete reaction, an additional 0.43 g (2.0 mmol) of pyrrolidine derivative and 0.81 g (8.0 mmol) of triethylamine are added and the reaction is heated at 120°C for 18 hours. The reaction mixture is cooled to room temperature, poured into 200 mL of ice and water and extracted with ethyl acetate (2 × 125 mL). The combined organic layers are washed with water, dried (MgSO₄), filtered and evaporated in vacuo to give 1.3 g of crude product. Chromatography on flash grade silica gel (230-400 mesh) eluting with 400 mL of dichloromethane/ethyl acetate (8:2), 600 mL of (6:4) and 1 L of (4:6) provided 0.46 g of the title compound, mp 84-86°C.


(e) *Synthesis of [(S)-1-(5-Amino-8-fluoro-3-methyl-4,6-dioxo-2,3,5,6-tetrahydro-4H-1-oxa-3a,5-diazaphenalen-9-yl)pyrrolidin-3-yl]carbamic acid tert-butyl ester*

-185-



A solution of 0.2 g (0.75 mmol) of 5-amino-8,9-difluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione (Example 24h), 0.58 g (2.0 mmol) of (S)-3-pyrrolidinylcarbamic acid *tert*-butyl ester (*J. Med. Chem.*, 1992;35:1764), 0.5 g (0.5 mmol) of triethylamine and 20 mL of acetonitrile is heated at reflux for 18 hours. The solvent is removed in vacuo and the residue is partitioned between dichloromethane-water. The organic layer is washed with water, dried (MgSO₄) and concentrated in vacuo. The residue is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane/ ethanol (95:5) to give 0.34 g of the title compound, mp 114-116°C.

(f) *{(S)-1-[(R)-1-(5-Amino-8-fluoro-3-methyl-4,6-dioxo-2,3,5,6-tetrahydro-4H-1-oxa-3a,5-diazaphenalene-9-yl)pyrrolidin-3-yl]ethyl}carbamic acid tert-butyl ester*




A solution of 0.11 g (0.41 mmol) of 5-amino-8,9-difluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione (Example 24h), 0.26 g (1.3 mmol) of ((R)-(S)-1-pyrrolidin-3-ylethyl)carbamic acid *tert*-butyl ester (*J. Het. Chem.*, 1992;29:1481), 0.25 g (2.5 mmol) of triethylamine and 10 mL of acetonitrile is heated at reflux for 18 hours. The solvent is removed in vacuo and the residue is partitioned between ethyl acetate/water. The organic layer is washed with water, dried (MgSO₄) and concentrated in vacuo. The residue is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane/ethanol (95:5) to give 0.14 g of the title compound. ¹H NMR (400 MHz, CDCl₃): 7.43 (d, 1H),

-186-


5.24 (bs, 2H), 4.82 (m, 1H), 4.56 (m, 1H), 4.33 (d, 1H), 3.99 (m, 1H), 3.86 (m, 1H), 3.73 (m, 1H), 3.59 (m, 2H), 2.18 (m, 1H), 3.01 (m, 1H), 1.67 (m, 1H), 1.44 (s, 9H), 1.40 (m, 3H), 1.25 (m, 1H), 1.20 (d, 3H).

(g) *[(S)-1-(2-Amino-9-fluoro-5-methyl-1,3-dioxo-2,3,6,7-tetrahydro-1H,5H-pyrido[3,2,1-ij]quinazolin-8-yl) pyrrolidin-3-yl]carbamic acid tert-butyl ester*



A solution of 0.12 g (0.45 mmol) of 5-amino-8,9-difluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione (Example 24i), 0.34 g (1.8 mmol) of (S)-3-pyrrolidinylcarbamic acid *tert*-butyl ester (*J. Med. Chem.*, 1992;35:1764), 0.2 g (2.0 mmol) of triethylamine and 7 mL of dimethyl sulfoxide is heated at 110°C for 18 hours. The reaction is cooled to room temperature, diluted with 80 mL of ice and water and stirred at 5°C for 1 hour. The resulting precipitate is removed by filtration, washed with water and dried in vacuo. The solid is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane/ethanol (9:1) to give 0.14 g of the title compound, mp 98-100°C.

(h) *{(R)-3-[(S)-1-(2-Amino-9-fluoro-5-methyl-1,3-dioxo-2,3,6,7-tetrahydro-1H,5H-pyrido[3,2,1-ij]quinazolin-8-yl)pyrrolidin-3-yl}ethylcarbamic acid tert-butyl ester*




A solution of 0.12 g (0.45 mmol) of 5-amino-8,9-difluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione (Example 24i), 0.39 g (1.8 mmol) of ((R)-3-[(S)-1-(2-amino-9-fluoro-5-methyl-1,3-dioxo-2,3,6,7-tetrahydro-1H,5H-pyrido[3,2,1-ij]quinazolin-8-yl)pyrrolidin-3-yl]ethyl)carbamic acid *tert*-butyl ester (*J. Het. Chem.*, 1992;29:1481), in 5 mL of dimethyl sulfoxide is heated at 110°C for

-187-


18 hours. The reaction is cooled to room temperature, diluted with 80 mL of ice and water and stirred at 5°C for 1 hour. The resulting precipitate is removed by filtration, washed with water and dried in vacuo. The solid is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane/ethanol (9:1) to give 0.13 g of the title compound, mp 93-95°C.

(i) *[1-(3-Amino-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-7-yl)pyrrolidin-3-yl]carbamic acid tert-butyl ester*



A solution of 0.11 g (0.4 mmol) of 3-amino-7-chloro-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 24b), 0.105 g (0.56 mmol) of (S)-3-pyrrolidinylcarbamic acid tert-butyl ester (*J. Med. Chem.*, 1992;35:1764), 1.2 mL of N, N-diisopropylethylamine and 3 mL of acetonitrile is heated at 50°C for 18 hours. The reaction mixture is diluted with 9 mL of water, cooled to 0°C and the solid is removed by filtration, washed with 50% aqueous acetonitrile and dried in vacuo to give 0.14 g of the title compound, mp 108-110°C.

(j) *{{(S)-1-[(R)-1-(3-Amino-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid tert-butyl ester*



A solution of 0.12 g (0.38 mmol) of 3-amino-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydro-pyrido[2,3-d]pyrimidine-7-thiosulfonic acid (Compound 24c) in 10 mL of acetonitrile is treated with 0.24 g (1.14 mmol) of ((R)-S)-1-pyrrolidin-3-ylethyl)carbamic acid tert-butyl ester (*J. Het. Chem.*,

-188-

1992;29:1481), 0.25 g (2.5 mmol) of triethylamine and stirred at room temperature for 18 hours then 50°C for 1 hour. The solvent is removed in vacuo and the residue is partitioned between ethyl acetate (50 mL) and water (25 mL). The organic layer is washed with water, dried (MgSO₄), filtered and evaporated
5 in vacuo to give 0.28 g. Chromatography on flash grade silica gel (230-400 mesh) column (2 × 12 cm) eluting with dichloromethane/ethanol (95:5) provided 0.1 g of the title compound, mp 175-177°C.

General Procedure A

A solution of Example 24, 1-2.5 eq of heterocyclic amine side chain, and
10 3-3.5 eq. 1,8-diazabicyclo-5,4,0-undecen-7-ene (DBU) or triethylamine are stirred in dimethyl sulfoxide (1 mL) at 130°C for 3 to 48 hours. After cooling to room temperature, the reaction mixture is diluted with ethyl acetate and washed with saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate concentrated. The resulting residue is purified via flash
15 column chromatography (isopropanol/dichloromethane) to afford the product.

General Procedure B

A solution of Example 24 and heterocyclic diamine side chain (1.5-3.0 eq.) is stirred in dimethyl sulfoxide at 130°C for 3 hours. After cooling to room temperature the reaction mixture is diluted with ethyl acetate and washed with
20 saturated NaHCO₃, water, and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate concentrated. The resulting residue is redissolved in dichloromethane and while stirring, di-*tert*-butyl dicarbonate (0.41 g, 1.87 mmol) is added. After 1 hour, the reaction mixture is concentrated and the product is purified via flash column chromatography (1:1 EtOAc/hexanes) to afford the
25 product.

General procedure C

A solution of Example 24, heterocyclic diamine side chain (1.5-3.0 eq.) and tetramethyl guanidine (1-5.0 eq.) is stirred in dimethyl sulfoxide at (80°C-130°C) for 36 to 24 hours. After cooling to room temperature the reaction
30 mixture is diluted with water and saturated NH₄Cl. The solid is washed with

water and dried to give the crude solid. The product is purified via flash column chromatography (SiO₂) using (CHCl₃/MeOH or ethyl acetate/hexanes) to afford the product.

The following compounds are synthesized according to the General Procedures A, B, and C of Example 28:

(k) {1-[(R)-1-((S)-3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 462 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (R)-((S)-1-pyrrolidin-3-yl)ethylcarbamic acid *tert*-butyl ester (Kimura Y., Atarashi S., Takahashi M., Hayakkawa I., *Chem. Pharm. Bull.*, 1994;42:1442) using General Method A.

(l) {1-[(S)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]-1-methylethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 476 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and [(S)-(pyrrolidin-3-yl)-1-methylethyl]carbamic acid *tert*-butyl ester using General Method A.

(m) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methoxypyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 492 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and [1-(3-methoxypyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (Example A5b) using General Method A.

(n) [3-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-azabicyclo[3.1.0]hex-6-yl]carbamic acid *tert*-butyl ester (MS ES: m/z 446 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and 3-azabicyclo[3.1.0]hex-6-ylcarbamic acid *tert*-butyl ester (Brighty K.E., 1991, EP 413455) using General Method A.

(o) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-fluoropyrrolidin-3-yl]ethyl}carbamic acid

tert-butyl ester (MS ES: m/z 496 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (4-fluoropyrrolidin-3-yl)ethylcarbamic acid *tert*-butyl ester (Example A5c) using General Method A.

5 (p) [1-(3-Amino-1-cyclopropyl-6-fluoro-5-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]carbamic acid *tert*-butyl ester (MS ES: m/z 434 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-5-methyl-1H-quinazoline-2,4-dione (Example 24f) and (pyrrolidin-3-yl)carbamic acid *tert*-butyl ester using General Method A.

10 (q) {1-(R)-[1-(S)-(3-Amino-1-cyclopropyl-6-fluoro-5-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 462 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-5-methyl-1H-quinazoline-2,4-dione (Example 24f) and ((R)-((S)-1-pyrrolidin-3-ylethyl)amine using General Method A.

15 (r) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methoxymethylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 493 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (3-methoxymethylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A5d) using General Method A.

20 (s) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-fluoromethylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.60 (d, 1H), 5.18 (bs, 1H), 4.97 (bs, 2H), 4.61-4.10 (m, 4H), 3.72-3.19 (m, 5H), 2.46 (s, 3H), 2.00-1.71 (m, 2H), 25 1.45 (s, 9H), 1.21-1.10 (m, 2H), 0.68-0.56 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and 1-(3-fluoromethylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Li Q., Wang W., Berst K.B., Claiborne A., Hasvold L., Raye K., Tufano M., et al., *Bioorg. Med. Chem. Lett.*, 1998;8:1953-1958) using General Method A.

30 (t) [trans-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-trifluoromethylpyrrolidin-3-

-191-

ylmethyl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, CDCl_3): δ 7.56 (d, 1H), 5.17 (s, 2H), 4.75 (bs, 1H), 3.97-3.68 (m, 3H), 3.57 (s, 3H), 3.51-3.13 (m, 4H), 2.98-2.60 (m, 2H), 1.45 (s, 9H), 1.20-1.08 (m, 2H), 0.71-0.55 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and [trans-(4-trifluoromethylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (Li Q., Wang W., Berst K.B., Claiborne A., Hasvold L., Raye K., Tufano M., et al., *Bioorg. Med. Chem. Lett.*, 1998;8:1953-1958) using General Method A.

(u) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS (ES, $M+1$) m/z 492 ($M\text{H}^+$)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (1-piperidin-3-ylethyl)carbamic acid *tert*-butyl ester (Example A5e) using General Method A.

(v) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 476 ($M\text{H}^+$)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and 1-(piperidin-3-yl)ethylcarbamic acid *tert*-butyl ester (Example A5e) using General Method A.

(w) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 506 ($M\text{H}^+$)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and [1-(4,4-dimethylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (Example A5f) using General Method A.

(x) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 490 ($M\text{H}^+$)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and [1-(4,4-dimethylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (Example A5f) using General Method A.

- (y) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl} carbamic acid tert-butyl ester (MS ES: m/z 476 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and [1-(4-methylpyrrolidin-3-yl)ethyl]carbamic acid tert-butyl ester (Example A5g) using General Method A.
- (z) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-phenylpyrrolidin-3-yl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.61 (d, 1H), 7.50-7.12 (m, 5H), 5.19 (bs, 2H), 4.07-3.70 (m, 3H), 3.62-3.34 (m, 2H), 2.68-2.52 (m, 1H), 2.45 (s, 3H), 1.68 (m, 2H), 1.50-0.95 (m, 11H), 0.62 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (3-phenylpyrrolidin-3-yl)carbamic acid *tert*-butyl ester (Example A30) (using General Method A).
- (aa) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-phenylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.68 (d, 1H), 7.50-7.29 (m, 5H), 6.72 (s, 1H), 5.24 (s, 2H), 4.18-3.85 (m, 3H), 3.78-3.52 (m, 2H), 2.70 (s, 1H), 2.58-2.39 (m, 4H), 1.62-0.62 (m, 15H) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (3-phenylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A5p) using General Method A.
- (bb) [5-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-5-azaspiro[2.4]hept-7-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 474 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (5-azaspiro[2.4]hept-7-ylmethyl)carbamic acid *tert*-butyl ester (Example A5h) using General Method A.
- (cc) [5-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-5-azaspiro[2.4]hept-7-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 490 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and 5-azaspiro[2.4]hept-7-ylmethyl)carbamic acid *tert*-butyl ester (Example A5h) using General Method A.

-193-

(dd) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-hydroxypyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 480 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (3-hydroxypyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A5i) using General Method A.

(ee) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidin-3-ylmethyl]carbamic acid *tert*-butyl ester (REF) (MS ES: m/z 462 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (piperidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Hilpert K., Ackermann J., Banner D.W., Gast A., Gubernator K., Hadvary P., Labler L., et al., *J. Med. Chem.*, 1994;37[23]:3889-3901) using General Method A.

(ff) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methoxypyrrolidin-3-yl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.60-7.54 (d, 1H), 5.16 (s, 2H), 4.82 (s, 1H), 4.20-3.85 (m, 5H), 3.42 (s, 3H), 3.23 (t, 2H), 2.41 (s, 3H), 1.47 (s, 9H), 1.18-1.10 (m, 2H), 0.61 (s, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and 4-(methoxypyrrolidin-3-yl)carbamic acid *tert*-butyl ester (Li Q., Cooper C.S., Anthony K.L., Lee C.M., Plattner J.J., Ma Z., Wang W., 1996, WO 9639407) using Method A.

(gg) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methoxypyrrolidin-3-yl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.21 (d, 1H), 5.23-5.14 (m, 3H), 4.33-3.22 (m, 13H), 1.40 (s, 9H), 1.18 (m, 2H), 0.60 (s, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and 4-(methoxypyrrolidin-3-yl)carbamic acid *tert*-butyl ester (Li Q., Cooper C.S., Anthony K.L., Lee C.M., Plattner J.J., Ma Z., Wang W., 1996, WO 9639407) using Method A.

(hh) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $DMSO-d_6$): δ 7.40-7.34 (d, 1H), 5.30-4.80 (m, 3H) 4.50-

3.10 (m, 10H), 1.47 (s, 9H), 1.30-0.50 (m, 4H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and 4-(fluoropyrrolidin-3-yl)carbamic acid *tert*-butyl ester (Li Q., Wang W., Berst K.B., Claiborne A., Hasvold L., Raye K., Tufano M., et al., *Bioorg. & Med. Chem. Lett.*, 1998;8:1953-1958) using General Method A.

(ii) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 462 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (3-methylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (*J. Med. Chem.*, 1992:361-367) using General Method A.

(jj) {(S)-1-[(R)-1-(3-Amino-1-cyclopropyl-8-ethyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 476 (MH^+)) from 3-amino-1-cyclopropyl-8-ethyl-6,7-difluoro-1H-quinazoline-2,4-dione (Example 24g) and ((R)-[(S)-1-pyrrolidin-3-yl-ethyl]ethyl)carbamic acid *tert*-butyl ester using General Method A.

(kk) 1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.51 (d, 1H), 5.15 (s, 2H), 3.88-3.58 (m, 4H), 3.50 (s, 3H), 3.42-3.30 (m, 1H), 3.14-3.02 (m, 1H), 2.25-2.17 (m, 2H), 1.48 (s, 9H), 1.14-1.05 (m, 2H), 0.68-0.61 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and pyrrolidine-3-carboxylic acid *tert*-butyl ester (Hayakawa I., Tanaka Y., EP 101829) using General Method A.

(ll) 1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.61 (d, 1H), 5.16 (s, 2H), 3.70-3.40 (m, 5H), 3.17-3.03 (m, 1H), 2.43 (s, 3H), 2.29-2.19 (m, 2H), 1.47 (s, 9H), 1.22-1.12 (m, 2H), 0.67-0.57 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and pyrrolidine-3-carboxylic acid *tert*-butyl ester (Hayakawa I., Tanaka Y., EP 101829) using General Method A.

-195-

(mm) [3-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-azabicyclo[3.1.0]hex-1-ylmethyl]yl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, CDCl_3): δ 7.53 (d, 1H), 5.31 (bs, 2H), 4.73 (bs, 1H), 3.82-3.01 (m, 11H), 1.45 (s, 9H), 1.42-1.34 (m, 1H), 1.14-1.10 (m, 3H), 0.71-0.63 (m, 2H)), from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (3-azabicyclo[3.1.0]hex-1-ylmethyl)carbamic acid *tert*-butyl ester (Example A5j) using General Method A.

(nn) [3-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-azabicyclo[3.1.0]hex-1-ylmethyl]yl]carbamic acid *tert*-butyl ester (MS ES: m/z 460) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (3-azabicyclo[3.1.0]hex-1-ylmethyl)carbamic acid *tert*-butyl ester (Example A5j) using General Method A.

(oo) {(R)-1-[(S)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 478 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (R)-1-((S)-pyrrolidine-3-yl)ethyl)carbamic acid *tert*-butyl ester (Schroeder M.C., Kiely J.S., Johnson D.R., Szotek D.L., Domagala J.M., Stickney T.M., Kampf J.W., *J. Heterocyclic Chem.*, 1992;29:1481) using General Method A.

(pp) {(R)-1-[(S)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester ^1H NMR (200 MHz, CDCl_3): δ 7.51-7.44 (d, 1H), 5.17 (bs, 2H), 4.62 (bs, 0.5H), 4.58 (bs, 0.5H), 3.79-3.74 (m, 3H), 3.61-3.53 (m, 2H), 3.49 (s, 3H), 3.41-3.32 (m, 1H), 2.22-2.06 (m, 2H), 1.87-1.67 (m, 1H), 1.46 (s, 9H), 1.23-1.19 (d, 3H), 1.08-1.02 (m, 2H) 0.76-0.62 (m, 2H). (MS ES: m/z 462) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (R)-1-((S)-pyrrolidine-3-yl)ethyl)carbamic acid *tert*-butyl ester using General Method A.

(qq) 6-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)octahydropyrrolo[3,4-b]pyridine-1-carboxylic acid *tert*-butyl ester (^1H NMR (200 MHz, CDCl_3): δ 7.61-7.55 (d, 1H), 5.16 (bs, 2H), 4.52

(bs, 0.5H), 4.48 (bs, 0.5H), 4.95-3.38 (m, 6H), 2.40 (s, 3H), 2.36-1.83(m, 3H), 1.46 (s, 9H), 1.21-1.17 (m, 5H), 0.65-0.57 (m, 2H), from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and octahydropyrrolo[3,4-b]pyridine-5-carboxylic acid *tert*-butyl ester using General Method A.

(rr) [(trans)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 478 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (trans-4-methylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Kuniyoshi M., Seigo S., Keiji H., Takayoshi I., Eur. Pat. Appl. EP 208210, 1987) using General Method A.

(ss) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 462 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (4-methylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Kuniyoshi M., Seigo S., Keiji H., Takayoshi I., Eur. Pat. Appl. EP 208210, 1987) using General Method A.

(tt) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]carbamic acid *tert*-butyl ester (MS ES: m/z 464 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (4-methylpyrrolidin-3-yl)carbamic acid *tert*-butyl ester (Di Cesare P., Bouzard D., Essiz M., Jacquet J.P., Ledoussai B., Kiechet J.R., Remuzon P. et al., *J. Med. Chem.*, 1992,35:4205) using General Method A.

(uu) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)morpholin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 480 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and 1-(morpholin-3-ylmethyl)carbamic acid *tert*-butyl ester (Araki K., Kuroda T., Uemori S., Moriguchi A., Ikeda Y.,

-197-

Hirayama F., Yokoyama Y. et al., *J. Med. Chem.*, 1993,36:1356) using General Method A.

(vv) [1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 478 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and 1-piperidin-3-ylmethylcarbamic acid *tert*-butyl ester (Hilpert K., et al., *J. Med. Chem.*, 1994;37(23):3889-3901) using General Method A.

(ww) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS ES: m/z 492 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and [1-(4-methylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (A5g) using General Method A.

(xx) [1-(1-Cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-3-methylpyrrolidin-3-yl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.61 (d, 1H), 5.15 (s, 2H), 4.74 (s, 1H), 3.71-3.37 (m, 6H), 2.45-2.20 (m, 4H), 1.45 (s, 12H), 1.21-1.08 (m, 2H), 0.61 (d, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and 3-methylpyrrolidin-3-ylcarbamic acid *tert*-butyl ester (Yoshida T., Yamamoto Y., Orita H., Kakiuchi M., Takahashi Y., Itakura M., Kado N. et al., *Chem. Pharm. Bull.*, 1996;44[7]:1376-1386) using General Method A

(yy) 3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione (^1H NMR (200MHz, $CDCl_3$): δ 7.61-7.54 (d, 1H), 5.15 (bs, 2H), 3.51-3.40 (m, 5H), 2.39 (s, 3H), 2.01-1.95 (m, 4H), 1.21-1.11 (m, 2H), 0.67-0.62 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and pyrrolidine using General Method A.

(zz) 3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione (^1H NMR (200MHz, $CDCl_3$): δ 7.49-7.42 (d, 1H), 5.13 (bs, 2H), 3.66-3.54 (m, 4H), 3.50 (s, 3H), 3.42-3.31 (m, 1H), 2.02-1.92 (m, 4H),

-198-

1.17–1.07 (m, 2H), 0.69–0.61 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and pyrrolidine using General Method A.

(aaa) 3-Amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxyethyl-1-methyl)-pyrrolidin-1-yl]-8-methyl-1H-quinazoline-2,4-dione (¹H NMR (200 MHz, CDCl₃): δ 7.59 (d, 1H), 5.15 (s, 2H), 3.68–3.53 (m, 2H), 3.44–3.31 (m, 3H), 2.42 (s, 3H), 2.00–1.82 (m, 2H), 1.58 (s, 1H), 1.28 (s, 6H), 1.20–1.10 (m, 2H), 1.00–0.80 (m, 1H), 0.66–0.60 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and 1-methyl-1-pyrrolidin-3-yl-ethanol (Example A5n) using General Method A.

(bbb) 3-Amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)-pyrrolidin-1-yl]-8-methoxy-1H-quinazoline-2,4-dione (¹H NMR (200 MHz, CDCl₃): δ 7.50 (d, 1H), 5.14 (s, 2H), 3.86–3.70 (m, 2H), 3.49 (s, 3H), 3.60–3.30 (m, 3H), 2.40–2.22 (m, 1H), 2.00–1.80 (m, 2H), 1.30 (s, 6H), 1.09–0.85 (m, 2H), 0.75–0.57 (m, 2H)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and 1-methyl-1-pyrrolidin-3-yl-ethanol (Example A5n) using General Method A.

(ccc) 1-[(R)-1-(3-(S)-Amino-1-cyclopropyl-6-fluoro-5-methyl-7-[3-(1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione (MS ES: m/z 376 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-5-methyl-1H-quinazoline-2,4-dione (Example 24e) and methyl-((R)-((S)-1-pyrrolidin-3-yl)ethyl)amine using General Method A.

(ddd) 1-[(R)-1-(3-(S)-Amino-1-cyclopropyl-7-[3-(1-ethylaminoethyl)-pyrrolidin-1-yl]-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione (MS ES: m/z 406 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and ethyl-((R)-((S)-1-pyrrolidin-3-yl)ethyl)amine (Domagala J.M., Hagen S.E., Joannides T., Kiely J.S., Laborde E., Schroeder M.C., Sesnie J.A., et al., *J. Med. Chem.*, 1993;36[7]:871–882) using General Method A.

(eee) 3-Amino-1-cyclopropyl-7-[(R)-3-((S)-1-ethylaminoethyl)-pyrrolidin-1-yl]-6-fluoro-8-methyl-1H-quinazoline-2,4-dione (MS ES: m/z 390

-199-

(MH⁺) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and ethyl-((R)-((S)-1-pyrrolidin-3-yl)ethyl)amine using General Method A.

5 (fff) 3-Amino-7-(6-amino-1-methyl-3-azabicyclo[3.2.0]hept-3-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione (MS ES: m/z 390 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and 1-methyl-3-azabicyclo[3.2.0]hept-6-ylamine (Kim C.S., Kim J.W., Lee J.M., Youn Y.S., Shin Y.J., Lee K.H., Kim J.H. WO 94/15933) using General Method A.

10 (ggg) 3-Amino-7-(3-aminomethylazetidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione (MS ES: m/z 350 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and azetidin-3-ylmethylamine (Okada T., Ezumi K., Yamakawa M., Sato H., Tsuji T., Tsushima T., Motokawa K., Komatsu Y., *Chem. Pharm. Bull.*, 1993;41:126-131) using General Method A.

15 (hhh) 3-Amino-7-(4-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione (MS ES: m/z 378 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and piperidin-4-ylmethylamine using General Method A.

20 (iii) [(cis)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 482 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (cis-4-fluoropyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester [A5k] using General Method A.

25 (jii) [(trans)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-ylmethyl]carbamic acid *tert*-butyl ester (MS ES: m/z 482 (MH⁺)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (trans-4-methylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester [A5l] using General Method A.

30

-200-

(kkk) [1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-(1-amino-5-azaspiro[2.4]hept-5-yl)]carbamic acid *tert*-butyl ester (MS APCI: m/z 476 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 14d) and 1-amino-5-azaspiro[2.4]hept-5-yl)]carbamic acid *tert*-butyl ester (Li Q., Chu D.T.W., Claiborne A., Cooper C.S., Lee C.M., Raye K., Berst K.B., et al., *J. Med. Chem.*, 1996;39[16]:3070-3088) using General Method C.

(lll) [1-(3-Amino-7-(3a-aminomethyloctahydroisindol-2-yl)-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)]carbamic acid *tert*-butyl ester (MS APCI: m/z 518 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 24d) and (3a-aminomethyloctahydro-isindol-2-yl)carbamic acid *tert*-butyl ester (Ma Z., Chu D.T.W., Cooper C.S., Li Q., Fung A.K.L., Wang S., Shen L.L., et al., *J. Med. Chem.*, 1999;42(20):4202-4213) using General Method C.

(mmm) 3-Amino-7-(3a-aminomethyloctahydroisindol-2-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione (MS APCI: m/z 502 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and 3a-aminomethyloctahydroisindol-2-ylcarbamic acid-*tert*-butyl ester (Ma Z., Chu D.T.W., Cooper C.S., Li Q., Fung A.K.L., Wang S., Shen L.L., et al., *J. Med. Chem.*, 1999;42(20):4202-4213) using General Method C.

(nnn) {(S)-1-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}carbamic acid *tert*-butyl ester (MS APCI: m/z 476 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methoxy-1H-quinazoline-2,4-dione (Example 14d) and ((R)-(S)-1-pyrrolidin-3-yl)ethylcarbamic acid *tert*-butyl ester (*J. Het. Chem.*, 1992;29:1481) using General Method C.

(ooo) {1-[1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydro-quinazolin-7-yl)-pyrrolidin-3-yl]-propyl}-carbamic acid *tert*-butyl ester (MS: m/e 476 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and (1-pyrrolidin-3-yl-propyl)-

-201-


carbamic acid *tert*-butyl ester (Kimura Y., Atarashi S., Takahashi M., Hayakkawa I., *Chem. Pharm. Bull.*, 1994;42:1442) using General Method A.

(ppp) {(S)-1-[(R)-1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydro-quinazolin-7-yl)pyrrolidin-3-yl]-ethyl}-methyl-carbamic acid dimethylethyl ester (MS: *m/e* 476 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and methyl-(R)-((S)-1-pyrrolidin-3-yl)ethylamine (Plummer J.S., Emery L.A., Stier M.A., Suto M.J., *Tetrahedron Lett.*, 1993;34(47):7529-7532) using General Method A.

(qqq) {1-[5-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-5-aza-spiro[2.4]hept-7-yl]ethyl}carbamic acid *tert*-butyl ester (MS CI: *m/e* 488 (MH^+)) from 3-amino-1-cyclopropyl-6,7-difluoro-8-methyl-1H-quinazoline-2,4-dione (Example 24e) and [1-(5-aza-spiro[2.4]hept-7-yl)ethyl]carbamic acid *tert*-butyl ester (Example A5a) using General Method A.

EXAMPLE 29

(7,8-Difluoro-3-methyl-2,3-dihydro-1,4-benzoxazin-4-yl)-oxoacetaldehyde oxime



A solution of 8.27 g (50 mmol) of chloral hydrate in 125 mL of water is treated with 112.8 g (0.35 mmol) of sodium sulfate decahydrate. The mixture is stirred and treated with a solution of 7,8-difluoro-3-methyl-3,4-dihydro-2H-benz[1,4]oxazine (Chem. Pharm. Bull., 1984;32:4907) hydrochloride (prepared by dissolving 8.26 g (44.6 mmol) of the free base in 50 mmol of concentrated hydrochloric acid in 70 mL of 50% aqueous ethanol). The reaction is stirred and a solution of 9.8 g (140 mmol) of hydroxylamine hydrochloride in 50 mL of water is added. After heating at 75 to 80°C for 3 hours, the reaction is stirred at room


-202-

temperature overnight. The resulting precipitate is removed by filtration, washed with water, and the wet filter cake is dissolved in dichloromethane. The organic solution is washed with water, dried (MgSO₄), decolorized with charcoal, filtered and concentrated in vacuo to give 10.5 g of the title compound, mp 188-190°C.

5

EXAMPLE 30


Synthesis of 6,7-Difluoro-3-methyl-3,4-dihydro-5-oxa-2a-aza-acenaphthylene-1,2-dione



10 A solution of 45 mL of 98% sulfuric acid in 18 mL of water is heated to 50°C to 60°C and treated portionwise over 30 minutes with 10.5 g (41 mmol) of (7,8-difluoro-3-methyl-2,3-dihydro-1,4-benzoxazin-4-yl)oxoacetaldehyde oxime (Example 29). After the addition is complete, the reaction is heated to 80°C for 15 minutes and poured onto 200 mL of ice and water. The mixture is stirred until the ice melts and the solid is removed by filtration, washed with water and the wet
15 filter cake is dissolved in dichloromethane. The organic solution is washed with water (2 × 200 mL), dried (MgSO₄), filtered and concentrated in vacuo affording 7.1 g of the title compound, mp 169-171°C.

EXAMPLE 31

20 *Synthesis of 7,8-Difluoro-3-methyl-3,4-dihydro-2H-1,4-benzoxazine-5-carboxylic acid*




-203-

A suspension of 6.9 g (28.8 mmol) of 6,7-difluoro-3-methyl-3,4-dihydro-5-oxa-2a-aza-acenaphthylene-1,2-dione (Example 30) in 250 mL of 4.5% aqueous sodium hydroxide is treated dropwise with 16.4 mL of 30% hydrogen peroxide over 1 hour. The reaction is filtered to remove a trace of insoluble material and the filtrate is acidified to pH 4.5 with acetic acid. After cooling to 5°C, the precipitate is removed by filtration, washed with water and dried in vacuo to give 5.5 g of the title compound, mp 200-202°C.

EXAMPLE 32

Synthesis of Anthranilic Esters


(a) *7,8-Difluoro-3-methyl-3,4-dihydro-2H-1,4-benzoxazine-5-carboxylic acid methyl ester*



A solution of 2.1 g (9.1 mmol) of 7,8-difluoro-3-methyl-3,4-dihydro-2H-1,4-benzoxazine-5-carboxylic acid (Example 31) in 50 mL of methanol is cooled to 0°C and saturated with hydrogen chloride gas. The reaction is stirred at room temperature for 18 hours, recooled to 0°C and resaturated with hydrogen chloride gas. After stirring an additional 24 hours at room temperature, the solvent is removed in vacuo and the residue is partitioned between dichloromethane/5% aqueous sodium bicarbonate (200:100 mL). The organic layer is separated, washed with 5% aqueous sodium bicarbonate, water, dried (MgSO₄), filtered, and concentrated in vacuo affording 1.9 g of the title compound. ¹H NMR (400 MHz, CDCl₃): 7.27 (bs, 1H), 7.24 (m, 1H), 4.26 (m, 1H), 3.82 (s, 3H), 3.75 (m, 1H), 3.62 (m, 1H), 1.22 (d, 3H).

(b) *Synthesis of 5,6-Difluoro-2-methyl-1,2,3,4-tetrahydro-quinoline-8-carboxylic acid methyl ester*


-204-



A solution of 8.0 g (33.7 mmol) of 5,6-difluoro-2-methylquinoline-8-carboxylic acid methyl ester (Example 33) in 100 mL of glacial acetic acid is treated with 2.15 g of platinum on carbon (0.8 g of active catalyst + 62.8% water) and then shaken in a hydrogen atmosphere at temperatures of 24°C to 29°C and pressures of 23.9 to 46.6 psi for 3 hours. The catalyst is removed by filtration and the solvent is removed in vacuo at 50°C. The residue is triturated with water (50 mL) and extracted with dichloromethane (2 × 200 mL). The combined organic layers are stirred with 5% sodium bicarbonate solution, washed with water, dried (MgSO₄), filtered and evaporated in vacuo. The residue is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane to give 5.6 g of the title compound as a light yellow oil. ¹H NMR (400 MHz, CDCl₃): 7.66 (bs, 1H), 7.49 (m, 1H), 3.82 (s, 3H), 3.46 (m, 1H), 2.88 (m, 1H), 2.67 (m, 1H), 1.97 (m, 1H), 1.27 (d, 3H).

15

EXAMPLE 33

Synthesis of 5,6-Difluoro-2-methylquinoline-8-carboxylic acid methyl ester


In a stainless steel reactor, a solution of 9.53 g (36.9 mmol) of 8-bromo-5,6-difluoro-2-methylquinoline (*Chem. Pharm. Bull.*, 1996;44:642), 6.8 g (67.6 mmol) of triethylamine, 0.67 g (3.0 mmol) of palladium (II) acetate, 1.28 g (3.1 mmol) of 1,3-bis(diphenyl-phosphino)propane in 90 mL of dimethylformamide and 65 mL of methanol is flushed with nitrogen gas and pressurized to 550 psi with carbon monoxide. The reaction is rocked and heated to

-205-

75°C for 24 hours, cooled to room temperature and evaporated in vacuo. The residue is triturated with water (50 mL) and extracted with dichloromethane (2 × 200 mL). The combined organic layers are washed with water (2 × 50 mL), dried (MgSO₄), filtered and evaporated in vacuo. The residue is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane (200 mL), then dichloromethane/ethyl acetate (90:10) to give 8.0 g of the title compound, mp 99-101°C.

EXAMPLE 34

Synthesis of 2-cyclopropylamino-5-fluoro-6-methylsulfanylnicotinic acid ethyl ester




A solution of 8.7 g (34.8 mmol) of 2-chloro-5-fluoro-6-methylsulfanylnicotinic acid ethyl ester (*J. Med. Chem.*, 1993;36:2676) and 5.7 g (100 mmol) of cyclopropylamine in 100 mL of acetonitrile is heated at reflux for 18 hours. After TLC showed the presence of unreacted starting material, 4.12 g (72 mmol) of cyclopropylamine is added and the reaction mixture is refluxed for 48 hours. The solvent is removed in vacuo and the residue is partitioned between dichloromethane (250 mL) and water (100 mL). The organic layer is washed with water, dried (MgSO₄), filtered, and concentrated in vacuo. The residue is chromatographed over flash grade silica gel (230-400 mesh) eluting with dichloromethane to give 6.5 g of the title compound, mp 67-69°C.

EXAMPLE 35

Synthesis of 1-Cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydropyrido[2,3-d]pyrimidine-7-thiosulfonic acid


-206-



A solution of 0.65 g (2.4 mmol) of 1-cyclopropyl-6-fluoro-7-methylsulfonyl-1H-pyrido[2,3-d]pyrimidine-2,4-dione (Example 23e) in 20 mL of formic acid is treated with 1.83 g (19.5 mmol) of urea-hydrogen peroxide and the reaction mixture is stirred at room temperature overnight. The resulting precipitate is removed by filtration, washed with water, ethyl acetate and dried in vacuo to give 0.63 g of the title compound, mp 300-302°C.

EXAMPLE 36

Synthesis of {8-Chloro-1-cyclopropyl-6-fluoro-7-[3-(methanesulfonylamino-methyl)pyrrolidin-1-yl]-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl}carbamic acid tert-butyl ester



A solution of (8-chloro-1-cyclopropyl-6,7-difluoro-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl)carbamic acid *tert*-butyl ester (Example 14, 0.37 g, 0.97 mmol) in acetonitrile (20 mL) is added pyrrolidin-3-yl-methylamine (0.115 g, 1.16 mmol) and triethylamine (2.7 mL, 19.3 mmol). After refluxing for 20 hours, the reaction mixture is cooled to room temperature and diluted with ethyl acetate. The organic layer is washed with saturated sodium bicarbonate, water, and brine. The organic layer is dried over MgSO₄, filtered, and the filtrate concentrated. The resulting residue is redissolved in methylene chloride (10 mL) at 0°C and while stirring, triethylamine (0.33 mL, 2.34 mmol) and methanesulfonyl chloride (0.072 mL, 0.94 mmol) is added. After 2 hours, the reaction mixture is diluted with ethyl acetate and washed with saturated sodium


-207-

bicarbonate, water, and brine. The organic layer is dried over MgSO_4 , filtered, and the filtrate concentrated and the product is purified via flash column chromatography (1:1 EtOAc/hexanes) to afford {8-chloro-1-cyclopropyl-6-fluoro-7-[3-(methanesulfonylaminomethyl)pyrrolidin-1-yl]-2,4-dioxo-1,4-dihydro-2H-quinazolin-3-yl} carbamic acid *tert*-butyl ester (0.043 g). MS CI: m/e 446 (MH^+ -Boc).

Amine Side Chain Synthesis

EXAMPLE A1


(a) *5-Benzyl-5-azaspiro[2.4]heptan-1-carboxylic acid ethyl ester*



A solution of cyclopropylidene acetic acid ethyl ester (192 g, 1520 mmol [Salaun J., Bennani F., Compain J.C., Fadel A., Ollivier J.J., *Org. Chem.*, 1980;45(21):4129-3415]) and N-(methoxymethyl)-N-(trimethylsilylmethyl)-benzylamine (572 g, 2410 mmol [Gerlach K., Hoffmann H.M.R., Wartchow R.J., *Chem. Soc., Perkin Trans. 11*, 1998(22):3867-3872]) in dichloromethane (5.5 L) was treated with a solution of 1.0N trifluoroacetic acid in dichloromethane (80 mL) and the reaction is stirred at ambient temperature for 30 minutes. The solvent is then evaporated at reduced pressure and the product purified by Kugelrohr distillation (277 g). MS CI: m/z 260 (MH^+).

EXAMPLE A2


(a) *Synthesis of 1-(5-Benzyl-5-azaspiro[2.4]hept-7-yl)ethanone*



-208-

A solution (-78°C) of 5-benzyl-5-azaspiro[2.4]heptane-7-carboxylic acid ethyl ester (Example A1a, 8.15 g, 31.4 mmol) in diethyl ether (200 mL) under a nitrogen atmosphere is treated with methyllithium (1.4 M in diethyl ether, 22.4 mL, 31.4 mmol) over a period of 10 minutes. After 1 hour, reaction mixture is warmed to -10°C and quenched with saturated aqueous ammonium chloride. The mixture is poured into a separatory funnel and the organic layer is washed with saturated aqueous ammonium chloride, water, and brine. The organic layer is then dried with MgSO₄, filtered, and the filtrate concentrated. The resulting residue is purified via flash column chromatography (1% triethylamine/7% isopropanol/92% CH₂Cl₂) to afford the title compound (3.93 g). MS CI: *m/z* 230 (MH⁺).

EXAMPLE A3

(a) *Synthesis of 1-(5-Benzyl-5-azaspiro[2.4]hept-7-yl)ethanone oxime*

1-(5-benzyl-5-azaspiro[2.4]hept-7-yl)ethanone (Example A2a, 3.93 g, 17.1 mmol) and hydroxylamine hydrochloride (1.78 g, 25.7 mmol) are stirred in pyridine (10 mL) at 90°C. After 20 hours, the reaction mixture is diluted with ethyl acetate and the organic layer is washed with saturated NaHCO₃, water, and brine. The organic layer is dried with MgSO₄, filtered, and the filtrate concentrated to afford 1-(5-benzyl-5-azaspiro[2.4]hept-7-yl)ethanone oxime as an oil (3.07 g). MS CI: *m/z* 245 (MH⁺).

General Method A

A solution of substrate, hydroxylamine hydrochloride (1.2 eq.), sodium acetate (0.9 eq.) in ethanol is stirred at room temperature for 2 hours. The solvent is removed in vacuo and the residue is partitioned between ethyl acetate and saturated sodium chloride solution. The aqueous is re-extracted with ethyl acetate

-209-

and the combined extracts are dried over Na_2SO_4 , filtered and concentrated under vacuum to afford the title compound.

The compounds below were prepared according to General Method A.


(b) 1-(1-Benzylpiperidin-3-yl)ethanone oxime (^1H NMR (200 MHz, CDCl_3): δ 7.48-7.28 (m, 5H), 3.92-3.68 (m, 2H), 3.28-3.0 (m, 4H), 2.88-2.62 (m, 2H), 2.38-1.72 (m, 3H), 1.3-1.18 (m, 3H)) from 1-(1-benzylpiperidin-3-yl)ethanone (Example A23).

(c) 1-(1-Benzyl-4,4-dimethylpyrrolidin-3-yl)ethanone oxime (^1H NMR (200 MHz, CDCl_3): δ 7.68-7.30 (m, 5H), 6.60-5.80 (bs, 1H), 3.80-2.38 (m, 6H), 1.90 (s, 3H), 1.38-0.96 (m, 7H)) from 1-(1-benzyl-4,4-dimethylpyrrolidin-3-yl)ethanone (Example A18b).

(d) 1-(1-Benzyl-4-methylpyrrolidin-3-yl)ethanone oxime (^1H NMR (200 MHz, CDCl_3): δ 7.41-7.15 (m, 5H), 3.78-3.44 (m, 3H), 2.98-2.80 (m, 1H), 2.75-2.39 (m, 3H), 2.35-2.08 (m, 2H), 1.90 (m, 3H), 1.18-0.95 (m, 3H)) from 1-(1-benzyl-4-methylpyrrolidin-3-yl)ethanone (Example A23b).

EXAMPLE A4

(a) *Synthesis of 1-(5-Benzyl-5-azaspiro[2.4]hept-7-yl)-ethyl]carbamic acid tert-butyl ester*



To a solution of 1-(5-benzyl-5-azaspiro[2.4]hept-7-yl)ethanone oxime (Example A3a, 2.90 g, 11.9 mmol) in methanol (100 mL) is added Raney-Nickel (2 g). Hydrogen is introduced to the reaction mixture at high pressure (48 psi) for 20 hours and the reaction mixture is filtered through celite, washed with methanol, and the combined filtrates concentrated in vacuo. The resulting residue is redissolved in dichloromethane (25 mL) and treated with di-*tert*-butyl dicarbonate (3.24 g, 14.9 mmol). After 1 hour, the mixture is concentrated and the product purified via flash column chromatography (1% triethylamine/7% isopropanol/92%

-210-

dichloromethane) to afford [1-(5-benzyl-5-azaspiro[2.4]hept-7-yl)-ethyl]carbamic acid *tert*-butyl ester (1.47 g). MS CI: *m/z* 331 (MH^+).

General Method A

Di-*tert*-butyl dicarbonate (0.69 g, 0.3 mmol) is added to a solution of
5 amine in methanol (4 mL) at 0°C. After stirring at room temperature for 2 hours, the mixture is concentrated and the residue is purified by chromatography (SiO₂, ethyl acetate/hexanes) to afford the title carbamic esters.

General Method B

An ice-cold solution of substrate in 2:1 methanol and water is treated with
10 di-*tert*-butyl dicarbonate (1.25 eq.). The resulting solution is stirred at room temperature for 18 hours and extracted with dichloromethane. The organic extracts are washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum. The residue is purified by flash column chromatography (ethyl acetate/hexanes) to afford the title compounds.

15 General Method C

To a solution of substrate in dichloromethane is added di-*tert*-butyl
dicarbonate (2 eq.) and triethylamine (2 eq.). The resulting mixture is stirred at room temperature for 18 hours and diluted with water. The organic extract is washed with water, brine, dried over Na₂SO₄, filtered and concentrated in vacuo.
20 The residue is then purified by column chromatography to afford the title compound.

Examples A4b-A4m are prepared according to General Methods A, B, and C.

(b) 3-(1-*tert*-Butoxycarbonylaminoethyl)-3-methoxypyrrolidine-1-
25 carboxylic acid benzyl ester (¹H NMR (200 MHz, CDCl₃): δ 7.37-7.29 (m, 5H), 5.13 (s, 2H), 4.70 (bs, 1H), 3.91-3.22 (m, 8H), 2.03-1.75 (m, 2H), 1.43 (s, 9H), 1.16-1.11 (m, 3H)) from 3-(1-aminoethyl)-3-methoxypyrrolidine-1-carboxylic acid benzyl ester (Example A13a) using General Method A.

(c) 3-(1-*tert*-Butoxycarbonylaminoethyl)-3-fluoropyrrolidine-1-
30 carboxylic acid benzyl ester (¹H NMR (200 MHz, CDCl₃): δ 7.35 (s, 5H), 5.13 (s,

-211-

2H), 4.79 (d, 1H), 3.97-3.26 (m, 5H), 2.17-1.75 (m, 2H), 1.44 (s, 9H), 1.24 (d, 3H)) from 3-(1-aminoethyl)-3-fluoropyrrolidine-1-carboxylic acid benzyl ester (Example A13b) using General Method A.

(d) (1-Benzyl-3-methoxymethylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (¹H NMR (200 MHz, CDCl₃): δ 7.35-7.17 (m, 5H), 5.78 (bs, 1H), 3.56 (s, 2H), 3.32 (s, 3H), 3.27 (s, 2H), 3.21-3.12 (m, 2H), 2.71-2.62 (m, 1H), 2.54-2.30 (m, 3H), 1.78-1.54 (m, 2H), 1.46 (s, 9H)) from (1-benzyl-3-methoxymethylpyrrolidin-3-yl)methylamine (Example A13c) using General Method B.

(e) [1-(1-Benzylpiperidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (¹H NMR (200 MHz, CDCl₃): δ 7.3 (s, 5H), 4.46-4.30 (m, 1H), 3.62-3.42 (m, 4H), 2.94-2.62 (m, 3H), 2.0-1.0 (m, 8H), 1.44 (s, 9H)) from 1-(1-benzylpiperidin-3-yl)ethylamine (Example A13d) using General Method C.

(f) [1-(1-Benzyl-4,4-dimethylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (¹H NMR (200 MHz, CDCl₃): **Isomer-1**: δ 7.36-7.22 (m, 5H), 3.82-3.38 (m, 2H), 2.98-1.78 (m, 5H), 1.52-1.0 (m, 19H); **Isomer-2**: δ 7.35-7.25 (m, 5H), 4.35-4.18 (m, 1H), 3.70-3.45 (m, 2H), 2.90-1.54 (m, 6H), 1.43 (s, 9H), 1.18-0.92 (m, 9H)) from 1-(1-Benzyl-4,4-dimethylpyrrolidin-3-yl)ethylamine (Example A13e) using General Method C.

(g) [1-(1-Benzyl-4-methylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (¹H NMR (200 MHz, CDCl₃): δ 7.38-7.13 (m, 5H), 3.71-3.32 (m, 4H), 3.08-1.53 (m, 6H), 1.52-1.31 (m, 9H), 1.14-0.94 (m, 6H). MS ES: *m/z* 319 (MH⁺)) from 1-(1-benzyl-4-methylpyrrolidin-3-yl)ethylamine (Example A13f) using General Method C.

(h) (5-Benzyl-5-azaspiro[2,4]hept-7-ylmethyl)carbamic acid *tert*-butyl ester (MS ES: *m/z* 317 (MH⁺)) from 1-(5-benzyl-5-azaspiro[2,4]hept-7-yl)methylamine (Example A13g) using General Method C.

(i) (1-Benzyl-3-hydroxypyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (MS ES: *m/z* 307 (MH⁺)) from 3-aminomethyl-1-benzylpyrrolidin-3-ol

-212-

(Grohe K., Schriewer M., Haller I., Metzger K., Endermann R., Zeiler H., EP 0326916) using General Method C.

(i) [3-(1-Phenylethyl)-3-azabicyclo[3.1.0]hex-1-ylmethylcarbamic acid *tert*-butyl ester (MS ES: m/z 317 (MH^+)) from 1-[3-(1-phenylethyl)-3-azabicyclo[3.1.0]hex-1-yl]methylaniline (Example A13h) using General Method C.

(k) (1,3-Dibenzylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (MS EI+: m/z 381 (MH^+)) using 1-(1,3-dibenzylpyrrolidin-3-yl)methylaniline (Example A13i) using General Method C.

(l) (1-Benzyl-4-fluoropyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.30 (s, 5H), 5.04 (m, 1H), 5.04 (m, 1H), 4.77 (m, 1H), 3.63 (s, 2H), 3.20 (t, 2H), 2.86-3.00 (m, 2H), 2.76 (d, 1H), 2.20-2.60 (m, 2H), 1.45 (s, 9H)) from (1-benzyl-4-fluoropyrrolidin-3-yl)methylaniline (Bouzard D., Di Cesare P., Essiz M., Jacquet J.P., Kiechel J.R., Remuzon P., Weber A., et al., *J. Med. Chem.*, 1990;33:1344-1352) using General Method C.

(m) *cis*-(1-Benzyl-4-fluoropyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.30 (s, 5H), 5.04 (m, 1H), 4.77 (m, 1H), 3.63 (s, 2H), 3.20 (t, 2H), 3.00-2.86 (m, 2H), 2.76 (d, 1H), 2.60-2.20 (m, 2H), 1.45 (s, 9H)) from *cis*-(1-benzyl-4-fluoro-pyrrolidin-3-yl)methylaniline (Matsumoto J., Nakano J., Chiba K., Minamida A., Nishimura Y., Jpn. Kokai Tokkyo Koho, 1987;11 pp., JP 62072660, A2 19870403) using General Method C.

(n) *trans*-(1-Benzyl-4-fluoropyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 7.30 (s, 5H), 5.25 (m, 1H), 4.77 (m, 1H), 3.85 (s, 2H), 3.55 (t, 2H), 3.40-2.40 (m, 5H), 1.47 (s, 9H) from *trans*-(1-benzyl-4-fluoropyrrolidin-3-yl)methylaniline (Matsumoto J., Nakano J., Chiba K., Minamida A., Nishimura Y., Jpn. Kokai Tokkyo Koho, 1987;11 pp., JP 62072660, A2 19870403) using General Method C.


(o) (1-Benzyl-3-phenylpyrrolidine-3-ylmethyl)carbamic acid *tert*-butyl ester (MS ES: m/z 367 (MH^+)) from (1-benzyl-3-phenylpyrrolidine-3-

-213-

yl)methylamine ester (Hagen S., Domagala J.M., Heifetz C.L., Sanchez J.P., Solomon M., *J. Med. Chem.*, 1990;33:849-854) using General Method C.

EXAMPLE A5

(a) *Synthesis of [1-(5-Azaspiro[2.4]hept-7-yl)ethyl]carbamic acid tert-butyl ester*



A solution of [1-(5-benzyl-5-azaspiro[2.4]hept-7-yl)ethyl]carbamic acid *tert*-butyl ester (Example A4a, 1.47 g) in methanol (30 mL) in a Parr shaker is treated with 10% palladium on carbon (0.5 g) and hydrogen introduced (50 psi) for 24 hours. The reaction mixture is filtered through celite, washed with methanol, and the combined filtrate concentrated in vacuo to afford the title compound. MS CI: m/z 241 (MH^+).

General Method A

To a solution of substrate in methanol is added ammonium formate (5 eq.) and 10% palladium on charcoal (0.75 wt. vol. eq.). After 2 hours at reflux, the reaction mixture is cooled, diluted with dichloromethane and filtered. The filtrate is concentrated to afford the title compound.

Examples A5b-A5o are prepared using General Method A.

(b) [1-(3-Methoxypyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 5.30 (bs, 1H), 5.03 (dd, 1H), 3.92 (bs, 1H), 3.28-2.82 (m, 7H), 1.97-1.72 (m, 2H), 1.44 (s, 9H), 1.19-1.14 (m, 3H)) from 3-(1-*tert*-butoxycarbonylaminoethyl)-3-methoxypyrrolidine-1-carboxylic acid benzyl ester (Example A4b) using General Method A.

(c) [1-(3-Fluoropyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, $CDCl_3$): δ 4.89 (d, 1H), 3.95-3.61 (m, 1H), 3.21-2.70 (m, 4H), 2.11-1.53 (m, 1H), 1.44 (s, 9H), 1.26 (d, 3H)) from 3-(1-*tert*-

-214-

butoxycarbonylaminoethyl)-3-fluoropyrrolidine-1-carboxylic acid benzyl ester (Example A4c) using General Method A.

(d) (3-Methoxymethylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, CDCl_3): δ 5.25 (bs, 1H), 3.35 (s, 3H), 3.30 (s, 2H), 3.27-3.15 (m, 2H), 3.09-2.94 (m, 2H), 2.78 (s, 2H), 1.68-1.56 (m, 2H), 1.45 (s, 9H). (MS ES: m/z 245 (MH^+)) from (1-benzyl-3-methoxymethylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A4d) using General Method A.

(e) (1-Piperidin-3-ylethyl)carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, CDCl_3): δ 4.52-4.32 (m, 1H), 3.68-3.18 (m, 4H), 2.74-2.42 (m, 1H), 2.02-1.2 (m, 5H), 1.44 (s, 9H), 1.30-1.04 (m, 3H)) from [1-(1-benzylpiperidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (Example A4e) using General Method A.

(f) [1-(4,4-Dimethylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, CDCl_3): 6.28-5.85 (bs, 1H), 4.75 (d, 1H), 3.92-1.80 (m, 6H), 1.43 (s, 9H), 1.30-1.0 (9H)) from [1-(1-benzyl-4,4-dimethylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (Example A4f).

(g) [1-(4-Methylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, CDCl_3): δ 4.88-4.40 (m, 1H), 4.02-3.38 (m, 4H), 3.25-2.75 (m, 2H), 2.46-1.88 (m, 2H), 1.44 (s, 9H), 1.31-1.04 (m, 6H)) from [1-(1-benzyl-4-methylpyrrolidin-3-yl)ethyl]carbamic acid *tert*-butyl ester (Example A4g).

(h) (5-Azaspiro[2,4]hept-7-ylmethyl)carbamic acid *tert*-butyl ester (MS ES: m/z 227 (MH^+)) from (5-benzyl-5-azaspiro[2,4]hept-7-ylmethyl)carbamic acid *tert*-butyl ester (Example A4h).

(i) (3-Hydroxypyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (MS ES: m/z 217 (MH^+)) using (1-benzyl-3-hydroxypyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A4i).

(j) (3-Azabicyclo[3.1.0]hex-1-ylmethyl)carbamic acid *tert*-butyl ester (MS ES: m/z 213 (MH^+)) using [3-(1-phenylethyl)-3-azabicyclo[3.1.0]hex-1-ylmethyl]carbamic acid *tert*-butyl ester (Example A4j).

(k) *cis*-(4-Fluoropyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (^1H NMR (200 MHz, CDCl_3): δ 5.60-5.00 (m, 4H), 3.60-3.10 (m, 4H), 3.00 (t,

-215-

1H), 2.75-2.30 (m, 1H), 1.44 (s, 9H)) using *cis*-(1-benzyl-4-fluoropyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A4I).

(l) *trans*-(4-Fluoropyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (¹H NMR (200 MHz, CDCl₃): δ 6.60-5.70 (m, 2H), 5.25 (bs, 0.5H), 4.98 (bs, 0.5H), 3.70-2.90 (m, 5H) 2.90-2.50 (m, 1H), 1.44 (s, 9H)) using *trans*-(1-benzyl-4-fluoropyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A4m).

(m) 3-Hydroxy-4-hydroxymethylpyrrolidine (¹H NMR (200 MHz, DMSO-d₆): δ 5.50-4.30 (bs, 3H), 4.30-1.50 (m, 8H)) using 1-benzyl-3-hydroxy-4-hydroxymethylpyrrolidine (Jaeger E., Biel J.H., *J. Org. Chem.*, 1965;30:740-744).


(n) 1-Methyl-1-pyrrolidin-3-yl-ethanol (MS ES: *m/z* 130 (MH⁺)) from 2-(1-benzylpyrrolidin-3-yl)propan-2-ol (Example A25).

(o) (3-benzylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (mp 183-185°C) from (1,3-dibenzylpyrrolidin-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A4K).

(p) A5p (3-Phenylpyrrolidine-3-ylmethyl)carbamic acid *tert*-butyl ester (MS APCI: *m/z* 277 (MH⁺)) from (1-benzyl-3-phenylpyrrolidine-3-ylmethyl)carbamic acid *tert*-butyl ester (Example A4o).

EXAMPLE A6


Synthesis of 1-Benzylpyrrolidin-3-one oxime



To a solution of 1-benzylpyrrolidin-3-one (1.0 g, 5.71 mmol) in pyridine (5 mL) is added hydroxylamine hydrochloride (0.59 g, 8.57 mmol). After stirring at 90°C for 5 hours, the reaction mixture is diluted with ethyl acetate and washed with saturated sodium bicarbonate, water, and brine. The organic layer is dried over MgSO₄, filtered, and filtrate concentrated to afford 1-benzylpyrrolidin-3-one oxime (1.11 g) as an oil. MS CI: *m/z* 191 (MH⁺).


-216-

EXAMPLE A7

Synthesis of Pyrrolidin-3-one oxime

5 In a Parr shaker, 1-benzylpyrrolidin-3-one oxime (1.5 g) in methanol (20 mL) is treated with 20% palladium on carbon (0.5 g) under Hydrogen pressure (50 psi) for 24 hours. The mixture is then filtered through Celite, washed with methanol, and the combined filtrate concentrated in vacuo to afford the title compound as an oil (1.2 g). MS CI: m/e 101 ($M^{+}+1$).

EXAMPLE A8


10 *Synthesis of 3-Ethylidenepyrrolidine-1-carboxylic acid benzyl ester*

Under an inert atmosphere, sodium-hydride (0.160 g, 7 mmol) and ethyltriphenylphosphonium bromide (2 g, 5.4 mmol) are mixed in dry dimethyl sulfoxide (10 mL) and stirred for 40 minutes. 3-Oxopyrrolidine-1-carboxylic acid benzyl ester (1.18 g, 5.4 mmol (*J. Med. Chem.*, 1992;35:1392)) in dry dimethyl sulfoxide (5 mL) is added to the reaction and the mixture is stirred at 70°C for 3 hours. The reaction is cooled, diluted with cold water and extracted with ethyl acetate. The organic phase is washed with brine, dried over Na_2SO_4 and evaporated. The residue is purified by column chromatography (65:35 hexanes/ethyl acetate) to afford the title compound (0.650 g) as a viscous oil. ^1H NMR (200 MHz, CDCl_3): δ 7.37-7.28 (m, 5H), 5.40-5.32 (m, 1H), 5.15 (s, 2H), 3.97 (bs, 2H), 3.60-3.44 (m, 2H), 2.49 (bs, 2H), 1.65-1.60 (m, 3H).

EXAMPLE A9

25 *Synthesis of 2-Methyl-1-oxa-5-azaspiro[2,4]heptane-5-carboxylic acid benzyl ester*


-217-



A solution of 3-ethylidenepyrrolidine-1-carboxylic acid benzyl ester (Example A8, 0.65 g, 2.8 mmol) and 3-chloroperoxybenzoic acid (0.86 g, 5 mmol) in dichloromethane (15 mL) is stirred at room temperature for 3 hours. The excess peracid is decomposed by the addition of 10% sodium sulfite. The organic layer is washed with 5% sodium bicarbonate solution, dried over Na₂SO₄ and evaporated. The residue is purified by column chromatography (1:1 hexanes/ethyl acetate) to afford the title compound (0.310 g). ¹H NMR (200 MHz, CDCl₃): δ 7.36-7.26 (m, 5H), 5.14 (s, 2H), 3.73-3.55 (m, 3H), 3.34 (d, 1H), 3.20-3.13 (m, 1H), 2.26-2.11 (m, 1H), 1.90-1.62 (m, 1H), 1.33 (d, 3H).

EXAMPLE A10

Synthesis of 3-(1-Aminoethyl)-3-hydroxypyrrolidine-1-carboxylic acid benzyl ester



A solution of 2-methyl-1-oxa-5-azaspiro[2,4]heptane-5-carboxylic acid benzyl ester (Example A9, 0.31 g, 1.25 mmol) in methanol (10 mL) and ammonium hydroxide (7 mL) is heated in a sealed tube at 100°C for 16 hours. After cooling, the reaction mixture is dissolved ethyl acetate, washed with water, dried over Na₂SO₄ and evaporated to afford the title compound (0.340 g). ¹H NMR (200 MHz, CDCl₃): δ 7.33-7.26 (m, 5H), 5.28 (bs, 2H), 5.07(s, 2H), 3.56-3.16(m, 5H), 2.03-1.76 (m, 2H), 1.15 (d, 3H).

-218-

EXAMPLE A11

Synthesis of 3-Hydroxy-3-{1-[(2-hydroxybenzylidene)amino]ethyl}pyrrolidine-1-carboxylic acid benzyl ester

A solution of 3-(1-aminoethyl)-3-hydroxypyrrolidine-1-carboxylic acid benzyl ester (Example A10, 0.34 g, 0.93 mmol) and salicylaldehyde (0.147 g, 1.2 mmol) in dry ethanol (5 mL) is refluxed for 3 hours. After evaporation of the solvent, the residue is purified by column chromatography (30:70:0.01 ethyl acetate/hexanes/ammonium hydroxide) to afford the title compound (0.12 g).
¹H NMR (200 MHz, CDCl₃): δ 8.39 (s, 1H), 7.41-7.25 (m, 7H), 6.97-6.85 (m, 2H), 5.09 (s, 2H), 3.65-3.34 (m, 5H), 2.01-1.87 (m, 2H), 1.37-1.32 (m, 3H).

EXAMPLE A12


Synthesis of 3-Methoxy-3-{1-[(2-methoxybenzylidene)amino]ethyl}pyrrolidine-1-carboxylic acid benzyl ester.

To a stirred solution of 3-hydroxy-3-{1-[(2-hydroxybenzylidene)amino]ethyl}pyrrolidine -1-carboxylic acid benzyl ester (Example A11, 0.120 g, 0.33 mmol) in tetrahydrofuran (5 mL) at 0°C is added 0.024 g (1 mmol) of sodium hydride. The mixture is stirred at room temperature for 2 hours and methyl iodide (3 mL) is added and stirring was continued overnight. After evaporation, ethyl acetate and water are added to the residue. The organic layer is separated, dried over Na₂SO₄ and concentrated to afford the title compound (0.125 g). ¹H NMR (200 MHz, CDCl₃): δ 8.70 (s, 1H), 7.95 (d, 1H), 7.41-7.32 (m, 6H), 7.01-6.88 (m, 2H), 5.09 (s, 2H), 3.86 (s, 3H), 3.72-3.22 (m, 8H), 2.33-1.73 (m, 2H), 1.27-1.24 (m, 3H).

EXAMPLE A13


(a) *Synthesis of 3-(1-Aminoethyl)-3-methoxypyrrolidine-1-carboxylic acid benzyl ester*

-219-



A solution of 3-methoxy-3-{1-[(2-methoxybenzylidene)amino]ethyl}pyrrolidine-1-carboxylic acid benzyl ester (Example A12, 0.120 g, 0.3 mmol) in 3 M hydrogen chloride in methanol (1 mL) is heated at 40°C overnight. The reaction is basified, extracted with ethyl acetate and the organic layer is dried over Na₂SO₄. After concentration, the residue is purified by column chromatography (20:80 methanol/dichloromethane) to afford the title compound (0.060 g). ¹H NMR (200 MHz, CDCl₃): δ 7.38-7.29 (m, 5H), 5.12 (s, 2H), 3.64-3.19 (m, 8H), 2.47 (bs, 2H), 2.07-1.80 (m, 2H), 1.16-1.10 (m, 3H).


(b) *3-(1-Aminoethyl)-3-fluoropyrrolidine-1-carboxylic acid benzyl ester*



A solution of 3-(1-azidoethyl)-3-fluoropyrrolidine-1-carboxylic acid benzyl ester (Example A16, 0.16 g, 0.5 mmol) in methanol (10 mL) is treated with Raney nickel (100 mg, wet weight) and shaken in a hydrogen atmosphere at room temperature and at 55 psi for 17 hours. The catalyst is removed by filtration through Celite, and the solvent is removed in vacuo to afford the title compound (0.12 g). ¹H NMR (200 MHz, CDCl₃): δ 7.30 (s, 5H), 5.12 (s, 2H), 3.84-3.22 (m, 5H), 2.23-1.79 (m, 4H), 1.20 (d, 3H).

(c) *1-(1-Benzyl-3-methoxymethylpyrrolidin-3-yl)methylamine*

-220-



Hydrazine hydrate (85% solution in water, 0.372 g, 6.32 mmol) is added to a solution of 2-(1-benzyl-3-methoxymethylpyrrolidin-3-ylmethyl)isoindole-1,3-dione (Example A20, 1.581 g, 4.34 mmol) in ethanol (50 mL). The resulting mixture is refluxed for 4.5 hours, cooled and acidified with concentrated hydrochloric acid. The precipitate is removed by filtration and the filtrate is concentrated in vacuo. The residue is dissolved in ethanol/water (2:1) and the insoluble material is removed by filtration. The filtrate is basified to pH 10 with 1N sodium hydroxide solution and extracted with dichloromethane. The combined organic extracts are dried over Na₂SO₄, filtered and concentrated under vacuum to afford the title compound (0.910 g). ¹H NMR (200 MHz, CDCl₃): δ 7.41-7.15 (m, 5H), 3.57 (s, 2H), 3.33 (s, 3H), 3.30 (s, 2H), 2.71 (s, 2H), 2.64-2.51 (m, 2H), 2.42-2.24 (m, 2H), 1.68-1.52 (m, 2H), 1.38-1.10 (m, 2H).

General Method A

Lithium aluminium hydride (2 eq.) is slowly added to a solution of oxime, nitrile, or amide derivative in dry tetrahydrofuran. After the addition is complete, the mixture is refluxed for 2 to 6 hours, cooled in an ice bath and quenched with saturated sodium sulfate solution. The solid is removed by filtration and the filtrate is concentrated in vacuo to afford the title compound.

Examples A13d-i are prepared according to General Method A.

(d) 1-(1-Benzylpiperidin-3-yl)ethylamine (¹H NMR (200 MHz, DMSO-d₆): δ 7.38-7.18 (m, 5H), 3.52-3.22 (m, 2H), 2.82-2.58 (m, 3H), 1.92-0.75 (m, 10H)) from 1-(1-benzylpiperidin-3-yl)ethanone oxime (Example A3b).

(e) 1-(1-Benzyl-4,4-dimethylpyrrolidin-3-yl)ethylamine (¹H NMR (200 MHz, CDCl₃): δ 7.36-7.22 (m, 5H), 3.66-3.42 (m, 2H), 2.94-2.10 (m, 5H), 1.32-0.88 (m, 10H)) from 1-(1-benzyl-4,4-dimethylpyrrolidin-3-yl)ethanone oxime (Example A3c).

-221-

(f) 1-(1-Benzyl-4-methylpyrrolidin-3-yl)ethylamine (^1H NMR (200 MHz, CDCl_3): δ 7.38-7.11 (m, 5H), 3.63-3.12 (m, 2H), 2.78-2.27 (m, 5H), 2.23-1.18 (m, 4H), 1.10-0.78 (m, 6H)) from 1-(1-benzyl-4-methylpyrrolidin-3-yl)ethanone oxime (Example A3d).


5 (g) 1-(5-Benzyl-5-azaspiro[2,4]hept-7-yl)methylamine (^1H NMR (CDCl_3): δ 7.36-7.21 (m, 5H), 3.68-3.53 (m, 2H), 3.1 (dd, 1H), 2.61-2.38 (m, 5H), 2.1-1.92 (m, 1H), 1.29 (bs, 2H), 0.72-0.32 (m, 4H)) from 5-benzyl-5-azaspiro[2,4]heptane-7-carboxylic acid amide (Example A24).

(h) 1-[3-(1-Phenylethyl)-3-azabicyclo[3.1.0]hex-1-yl]methylamine
10 (MS ES: m/z 217 (MH^+)) from 3-(1-phenylethyl)-3-azabicyclo[3.1.0]hexane-1-carbonitrile (Takemura M., Kimura Y., Kawakami K., EP 0807630).

(i) 1,3-Dibenzyl-3-pyrrolidinemethanamine (MS EI: m/z 281 (MH^+)) from 1,3-dibenzyl-3-pyrrolidine carboxamide (Example A29).

EXAMPLE A14

15 *Synthesis of 3-Fluoro-3-(1-hydroxyethyl)pyrrolidine-1-carboxylic acid benzyl ester*



A solution of 0.247 g, (1 mmol) of 2-methyl-1-oxa-5-azaspiro[2,4]-heptane-5-carboxylic acid benzyl ester (Example A9) in dichloromethane (3 mL)
20 is added to a polypropylene flask containing 0.5 mL of hydrogen fluoride-pyridine at -40°C . The reaction is allowed to come to room temperature and stirring is continued for 3 hours. The reaction mixture is poured into an aqueous ice-cold ammonium hydroxide solution (30 mL) and extracted with dichloromethane. The organic layer is dried, filtered and concentrated to afford the title compound

-222-

(0.267 g). ^1H NMR (200 MHz, CDCl_3): δ 7.35 (s, 5H), 5.13 (s, 2H), 3.96-3.34 (m, 5H), 2.20-1.98 (m, 3H), 1.33-1.26 (m, 3H).

EXAMPLE A15

Synthesis of 3-Fluoro-3-(1-methanesulfonyloxyethyl)pyrrolidine-1-carboxylic acid benzyl ester

To a -10°C solution of 3-fluoro-3-(1-hydroxyethyl)pyrrolidine-1-carboxylic acid benzyl ester (Example A14, 0.267 g, 1.00 mmol), triethylamine (0.4 g, 4 mmol) and dichloromethane (10 mL) is added methanesulfonyl chloride (0.229 g, 2.00 mmol). The mixture is stirred at room temperature for 2 hours and concentrated in vacuo. The residue is partitioned between ethyl acetate and water. The organic layer is dried over Na_2SO_4 and concentrated to afford the title compound (0.330 g). ^1H NMR (200 MHz, CDCl_3): δ 7.36 (s, 5H), 5.14 (s, 2H), 4.99-4.81 (m, 1H), 3.76-3.37 (m, 4H), 3.06 (s, 3H), 2.25-1.96 (m, 2H), 1.52-1.47 (m, 3H).

EXAMPLE A16


Synthesis of 3-(1-Azidoethyl)-3-fluoropyrrolidine-1-carboxylic acid benzyl ester

To a solution of 3-fluoro-3-(1-methanesulfonyloxyethyl)pyrrolidine-1-carboxylic acid benzyl ester (Example A15, 0.25 g, 0.72 mmol) in *N,N*-dimethylformamide (5 mL) is added sodium azide (0.13 g, 2 mmol). The reaction mixture is heated at 120°C overnight. The mixture is diluted with water and extracted with ethyl acetate. The extracts are washed with brine, dried over Na_2SO_4 and concentrated to afford the title compound (0.160 g). ^1H NMR (200 MHz, CDCl_3): δ 7.28 (s, 5H), 5.08 (s, 2H), 3.80-3.25 (m, 5H), 2.13-1.75 (m, 2H), 1.40 (d, 3H).

Example A17

Synthesis of 2-Methoxymethylacrylic acid ethyl ester

-223-



A solution of ethyl α -(bromomethyl)acrylate (0.99 g, 5.1 mmol [Villieras J., Rambaud M., *Synthesis*, 1982:924-926]) in methanol (10 mL) is added to an ice-cold solution of sodium methoxide (0.33 g, 6.1 mmol) in methanol (50 mL). The suspension is refluxed for 21 hours, cooled, diluted with water and extracted with dichloromethane. The organic extracts are dried over Na_2SO_4 , filtered and concentrated under vacuum. The resulting residue is purified by flash column chromatography (1:5 ethyl acetate/hexanes) to afford the title compound (0.211 g). ^1H NMR (200 MHz, CDCl_3): δ 6.33-6.26 (m, 1H), 5.88-5.80 (m, 1H), 4.23 (q, 2H), 4.19-4.10 (m, 2H), 3.41 (s, 3H), 1.31 (t, 3H).

EXAMPLE A18

General Method A

A solution of substrate and *N*-(methoxymethyl)-*N*-(trimethylsilylmethyl)benzylamine (1.3 eq.) in dichloromethane at 0°C is treated with trifluoroacetic acid (0.07 eq). The mixture is stirred at room temperature for 3 hours, diluted with dichloromethane and washed successively with a saturated NaHCO_3 solution and brine. The organic extracts are dried over Na_2SO_4 , filtered, and concentrated under vacuum to afford the title compounds.

The examples below (Example A18a-e) were prepared according to General Method A.

(a) 1-Benzyl-3-methoxymethylpyrrolidine-3-carboxylic acid ethyl ester (^1H NMR (200 MHz, CDCl_3): δ 7.45-7.16 (m, 5H), 4.19 (q, 2H), 3.78-3.67 (m, 2H), 3.55 (s, 2H), 3.31 (s, 3H), 3.11-2.92 (m, 1H), 2.89-2.49 (m, 2H), 2.42-2.21 (m, 2H), 1.99-1.78 (m, 1H), 1.26 (t, 3H)) from 2-methoxymethylacrylic acid ethyl ester (Example A17).

-224-

(b) 1-(1-Benzyl-4,4-dimethylpyrrolidin-3-yl)ethanone (^1H NMR (200 MHz, CDCl_3): δ 7.36-7.22 (m, 5H), 3.60 (d, 2H), 3.05-2.20 (m, 5H), 2.15 (s, 3H), 1.30 (s, 3H), 0.96 (s, 3H)) from mesityl oxide.


5 (c) 1-Benzyl-4-methylpyrrolidine-3-carboxylic acid ethyl ester (^1H NMR (200 MHz, CDCl_3): δ 7.44-7.13 (m, 5H), 4.21-4.04 (q, 1H), 3.71-3.49 (m, 3H), 2.94-2.71 (m, 3H), 2.61-2.41 (m, 2H), 2.28-2.04 (m, 1H), 1.32-1.18 (t, 3H), 1.16-1.04 (d, 3H). MS ES: m/z 248 (MH^+)) from ethyl crotonate.

(d) Pyrrolidine-3-carboxylic acid *tert*-butyl ester (MS ES: m/z 172 (MH^+)) from *tert*-butylacrylate.

10 (e) Ethyl 1,3-dibenzyl-3-pyrrolidinecarboxylic acid (MS EI+: m/z 324 (MH^+)) from ethyl-2-phenylmethylacrylic acid (Vassiliou S., Mucha A., Cuniasse P., Georgiadis D., Lucet-Levannier K., Beau F., Kannan R., et al., *J. Med. Chem.*, 1999;42(14):2610-2620).

EXAMPLE A19

15 *Synthesis of (1-Benzyl-3-methoxymethylpyrrolidin-3-yl)methanol*



A solution of 1-benzyl-3-methoxymethylpyrrolidine-3-carboxylic acid ethyl ester (Example A18a, 2.30 g, 8.3 mmol) in anhydrous tetrahydrofuran (10 mL) is added dropwise to an ice-cold suspension of lithium aluminum hydride (0.821 g, 22 mmol) in tetrahydrofuran (50 mL). The resulting mixture is stirred at room temperature for 18 hours, cooled and quenched by the sequential dropwise addition of 1N sodium hydroxide and water. The solids are removed by filtration, re-suspended in hot tetrahydrofuran and filtered. The combined filtrates are concentrated in vacuo and the residue is partitioned between dichloromethane and water. The organic extracts are dried over Na_2SO_4 , filtered and concentrated under vacuum to afford the title compound (1.71 g). ^1H NMR (200 MHz,


20

25

-225-


CDCl₃): δ 7.41-7.17 (m, 5H), 3.84-3.43 (m, 4H), 3.34 (s, 2H), 3.32 (s, 3H), 2.85-2.60 (m, 2H), 2.54-2.14 (m, 3H), 1.98-1.50 (m, 2H).

EXAMPLE A20

2-(1-Benzyl-3-methoxymethylpyrrolidin-3-ylmethoxy)isoindole-1,3-dione

5 A solution of (1-benzyl-3-methoxymethylpyrrolidin-3-yl)methanol (Example A19, 1.71 g, 7.27 mmol), triphenylphosphine (2.16 g, 8.22 mmol), phthalimide (1.14 g, 7.73 mmol), and anhydrous tetrahydrofuran is cooled to 0°C and treated with diethyl azodicarboxylate (1.62 g, 9.31 mmol). The resulting
10 mixture is stirred at room temperature for 18 hours and concentrated under vacuum. The residue is purified by flash column chromatography (1:5 ethyl acetate/hexanes) to afford the title compound, (0.58 g) as a solid. ¹H NMR (200 MHz, CDCl₃): δ 7.91-7.80 (m, 2H), 7.78-7.66 (m, 2H), 7.39-7.11 (m, 5H), 3.95-3.72 (m, 2H), 3.65-3.40 (m, 2H), 3.31 (s, 2H), 3.27 (s, 3H), 2.72-2.34 (m,
15 4H), 2.02-1.82 (m, 1H), 1.76-1.56 (m, 1H). MS ES: m/z 365 (MH⁺).

EXAMPLE A21


Synthesis of 1-Benzylpiperidine-3-carboxylic acid benzyl ester

20 To a solution added of nipecotic acid (8.0 g, 61.9 mmol) in *N,N*-dimethylformamide (80 mL) is benzyl bromide (31.8 g, 186 mmol) and potassium carbonate (42.8 g, 310 mmol). The mixture is heated at 70°C for 18 hours and

-226-

diluted with ethyl acetate and water. The organic extract is washed with 1.0 M hydrochloric acid, water and brine, dried over Na₂SO₄, filtered and concentrated in vacuo. The residue is purified by column chromatography (1:7 ethyl acetate/hexanes) to afford the title compound (8.6 g). ¹H NMR (200 MHz, DMSO-d₆): δ 7.27-7.42 (m, 10H), 5.07 (s, 2H), 3.45 (s, 2H), 2.88-2.49 (m, 3H), 2.32-1.32 (m, 6H).

EXAMPLE A22

Synthesis of 1-Benzylpiperidine-3-carboxylic acid

To a solution 3.0 g (9.7 mmol) of 1-benzylpiperidine-3-carboxylic acid benzyl ester (Example A21) in methanol (20 mL) was added 1N sodium hydroxide (20 mL), and the reaction is stirred at room temperature for 18 hours. The methanol is removed in vacuo and the basic aqueous solution is acidified to pH 4 with 1.0 M hydrochloric acid. After washing with ethyl acetate, the aqueous acid layer is lyophilized. The solid is suspended in methanol (10 mL), stirred for 10 minutes and filtered. The filtrate is evaporated in vacuo to afford the title compound (2.0 g). ¹H NMR (200 MHz, DMSO-d₆): δ 7.38-7.23 (m, 5H), 3.58 (s, 2H), 2.98-2.40 (m, 3H), 2.30-1.25 (m, 6H).

EXAMPLE A23

General procedure A

Methyl lithium (1.1 eq.) is added dropwise to a -70°C solution of substrate in tetrahydrofuran (15 mL). The resulting mixture is stirred at -70°C for 1 hour, and then warmed to room temperature for 18 hours. The reaction is diluted with ice water and ether and the organic layer is separated. The aqueous layer is

-227-

extracted with ether and the combined extracts are dried with Na₂SO₄ and concentrated in vacuo to afford the title compound.


The following compounds are prepared according to general procedure A.

5 (a) 1-(1-Benzylpiperidin-3-yl)ethanone (¹H NMR (200 MHz, DMSO-d₆): δ 7.38-7.22 (m, 5H), 3.45 (s, 2H), 2.85-2.48 (m, 3H), 2.05 (s, 3H), 2.16-1.06 (m, 6H)) from 1-benzylpiperidine-3-carboxylic acid (Example A22).

10 (b) 1-(1-Benzyl-4-methylpyrrolidin-3-yl)ethanone (¹H NMR (200 MHz, CDCl₃): δ 7.42-7.18 (m, 5H), 3.71-3.45 (m, 3H), 2.89-2.33 (m, 5H), 2.16 (s, 3H), 1.12 (d, 3H)) from 1-benzyl-4-methylpyrrolidine-3-carboxylic acid ethyl ester (Example A18c).

EXAMPLE A24

Synthesis of 5-Benzyl-5-azaspiro[2,4]heptane-7-carboxylic acid amide



15 5-Benzyl-5-azaspiro[2,4]heptane-7-carboxylic acid amide (3.32 g) is prepared from 5-benzyl-5-azaspiro[2,4]heptane-7-carboxylic acid (Example A28b, 5.0 g, 21.64 mmol) in dichloromethane by conversion to the acid chloride using oxalyl chloride (5.0 mL, 65 mmol) and *N,N*-dimethylformamide (3 drops). The mixture is allowed to stir for 2 hours then concentrate in vacuo. The residue is redissolved in dichloromethane and the mixture is again
20 concentrated in vacuo. The residue is dissolved in ether and treated with a solution of ammonia in ether. After 1 hour, the mixture is concentrated in vacuo to provide the title compound. ¹H NMR (CDCl₃): δ 7.6 (bs, 1H), 7.38-7.23 (m, 5H), 5.42 (bs, 1H), 3.75-3.57 (m, 2H), 3.19 (d, 1H), 2.83-2.64 (m, 2H), 2.4 (d, 2H), 0.96-0.58 (m, 4H).


-228-

EXAMPLE A25

Synthesis of 2(1-Benzylpyrrolidin-3-yl)propan-2-ol

To a 5°C solution of 1-benzylpyrrolidine-3-carboxylic acid ethyl ester (0.467 g, 2.0 mmol) in tetrahydrofuran (25 mL) is added an etherial solution of methylmagnesium bromide (3.3 mL, 10 mmol). The reaction mixture is stirred at room temperature for 2 hours, cooled to 0°C and saturated aqueous ammonium chloride (10 mL) is added followed by water (10 mL). The mixture is extracted with ethyl acetate (2 × 100 mL) and the combined organic extracts are washed with brine, dried over Na₂SO₄, filtered and concentrated under vacuum to afford the title compound (0.43 g). ¹H NMR (200 MHz, CDCl₃): δ 7.30-7.22 (m, 5H), 3.59 (s, 2H), 3.0 (bs, 1H), 2.80-2.66 (m, 2H), 2.48-2.30 (m, 2H), 2.20-2.05 (m, 1H), 1.94-1.80 (m, 2H), 1.19 (s, 3H), 1.16 (s, 3H).

EXAMPLE A26


Synthesis of 2-Benzyloctahydropyrrolo[3,4-c]pyridine-5-carboxylic acid tert-butyl ester

2-Benzyl-2,3,4,5,6,7,8,9-octahydropyrrolo[3,4-c]pyridine (21.6 g, 100 mmol [Himmeler T., Petersen U., Bremm K-D., Endermann R., Stegemann M., Wetzstein H-G., 1995, US 5578604] is dissolved in 1000 mL of dichloromethane containing triethylamine (16.7 mL, 120 mmol). Then di-*t*-butyldicarbonate (22.6 g, 120 mmol) is added and the mixture is stirred at room temperature for 18 hours. The organic layer is washed with saturated NaHCO₃, brine, dried, and concentrated to provide 30 g of the title compound as an oil; MS EI⁺: 318 (MH⁺).

EXAMPLE A27


Synthesis of octahydropyrrolo[3,4-c]pyridine-5-carboxylic acid tert-butyl ester

-229-



2-Benzyloctahydropyrrolo[3,4-c]pyridine-5-carboxylic acid *tert*-butyl ester (Example A26, 24.98 g, 78.8 mmol) in 400 mL of methanol is treated with 2.5 g of 20% Pd/C and shaken for 20.5 hours under 50 PSI of hydrogen. The solution is filtered and concentrated to afford 17.8 g of the title compound as a solid. MS APCI: m/z 228 (MH^+).

EXAMPLE A28

(a) *Synthesis of 1,3-Dibenzylpyrrolidine-3-carboxylic acid*


To a solution of 1,3-dibenzylpyrrolidine-3-carboxylic acid ethyl ester (Example A18e, 13.1 g, 41 mmol) in 250 mL of methanol is added 2N NaOH (46 mL, 92 mmol) and the reaction is heated to reflux for 24 hours. The solution is cooled, concentrated, redissolved in water and extracted with ether. The pH of the aqueous layer is adjusted to 7 and the precipitated solid is collected, to give 9.3 g of the title compound, MS EI^+ : m/z 296 (MH^+), mp 209-210°C.

(b) 5-Benzyl-5-azaspiro[2,4]heptane-7-carboxylic acid (MS ES: m/z 232) from 5-benzyl-7-ethoxycarbonyl-5-azaspiro[2,4]heptane (A1) using the method of Example A28a

EXAMPLE A29

20 *Synthesis of 1,3-Dibenzylpyrrolidine-3-carboxylic acid amide*


-230-



A mixture of 1,3-dibenzylpyrrolidine-3-carboxylic acid (Example A28, 4.0 g, 13.5 mmol), 1,1'-carbonyldiimidazole (2.42 g, 14.9 mmol), and triethylamine (2.08 mL, 14.9 mmol) in tetrahydrofuran (100 mL) is heated at reflux for 1.5 hours. The solution is cooled, poured into NH₄OH (300 mL), and stirred for 18 hours. The mixture is extracted with chloroform, the organic layer is washed with water, dried, and concentrated to afford 4.13 g of the title compound as an oil; MS EI⁺: m/z 295 (MH⁺).

EXAMPLE A30

Synthesis of (3-Phenylpyrrolidin-3-yl)carbamic acid tert-butyl ester



(1-Benzyl-3-phenylpyrrolidin-3-yl)carbamate (Hagen S.E., Domagala J.M., Heifetz C.L., Sanchez J.P., Solomon M., *J. Med. Chem.*, 1990;33(2):849-854) (2.79 g, 7.9 mmol) in 100 mL of methanol is treated with 1.0 g of 20% Pd/C and shaken for 17 hours under 50 psi of hydrogen. The solution is filtered and concentrated to afford 2.07 g of the title compound as a foam; MS EI⁺: m/z 263 (MH⁺).

The compounds of the present invention are potent inhibitors of bacterial growth both in animals and on surfaces, and also are inhibitors of bacterial enzymes. The invention compounds have been evaluated in a number of standard

-231-

in vitro and in vivo assays routinely used by those in the antibacterial art to measure antibacterial activity of test compounds.

Antibacterial Assay



5 The compounds of the present invention are tested against an assortment of Gram negative and Gram positive organisms using standard microtitration techniques (Cohen et al., *Antimicrob. Agents Chemother.*, 1985;28:766; Heifetz et al., *Antimicrob. Agents Chemother.*, 1974;6:124). The results of the evaluation of representative 3-aminoquinazolin-2,4-diones of Formula I are shown in Table 2 and compared to two 3-hydroxyquinazolin-2,4-diones.

DNA Gyrase Assay

10 The effects of invention compounds on the activity of DNA gyrase is determined in the supercoiling inhibition assay, following reaction conditions recommended by the enzyme supplier (Lucent, Ltd., Leicester, UK). Reactions are performed in buffer G (35 mM Tris-HCl (pH 7.5), 24 mM KCl, 4 mM MgCl₂,
15 2 mM DTT, 1.8 mM spermidine, 1 mM ATP, 0.1 mg/mL bovine serum albumin). Relaxed plasmid pBR322 (0.25 µg, Lucent, Ltd., Leicester, UK) is reacted with 1 U *E. coli* gyrase (Lucent, Ltd., Leicester, UK), in the absence or presence of test compounds, for 30 minutes at 37°C. Reactions are stopped by the addition of SDS and proteinase K to respective final concentrations of 1% and 0.5 mg/mL. After an
20 additional 30 minutes at 37°C, one-tenth volume of 10X loading buffer (0.3% bromophenol blue, 16% Ficoll, 10 mM Na₂HPO₄) is added, and reactions are loaded onto agarose gels and electrophoresed. The concentration of representative invention compounds of Formula I inhibiting 50% of the supercoiling activity of DNA gyrase is given as an IC₅₀ and recorded in Table 2.

-232-


TABLE 2. Antibacterial and *E. coli* gyrase Activities

Compound	Minimum Inhibitory Concentrations $\mu\text{g/mL}$						<i>E. coli</i>
Number or	Gram Negatives			Gram Positives			gyrase
Structure	<i>E. coli</i>	<i>E. coli</i>	<i>E. coli</i>	<i>E.</i>	<i>S.</i>	<i>S.</i>	IC ₅₀
	MC4100	B90	Tol C	<i>faecalis</i>	<i>aureus</i>	<i>pyogenes</i>	(μM)
				RB1	29213	C203	
1	2.0	1.0	0.25	1.0	0.5	0.13	1.0
2	16.0	4.0	2.0	>64	>64	>64	26.0
7	2.0	0.5	0.25	8.0	8.0	4.0	1.3
9	2.0	1.0	0.5	32.0	16.0	16.0	3.0
11	2.0	0.5	0.25	8.0	8.0	4.0	11.0
12	16.0	0.25	0.13	4.0	2.0	8.0	2.1
14	4.0	0.25	0.5	8.0	8.0	4.0	2.4
	1.0	0.25	0.25	4.0	4.0	2.0	0.9
	8.0	2.0	1.0	64.0	32.0	32.0	5.0

The in vivo activity of invention compounds is evaluated according to the procedure of Miller et al., *Proc. Soc. Exp. Biol. Med.*, 1944;57:261. The median protective dose (PD₅₀) is determined in mice given lethal systemic infections.

Two 3-aminoquinazolin-2,4-diones of the invention are compared to a 3-hydroxyquinazolin-2,3-dione, and the results are presented in Table 3. The results establish the unexpected superiority of the 3-amino invention compounds compared to similar 3-hydroxy analogs.

TABLE 3. In Vivo Median Protective Dose (PD₅₀) in Mice

Compound Number or Structure	Organism	PD ₅₀ (mg/kg)
1	<i>S. pyogenes</i>	12
10	<i>E. coli</i>	72
	<i>S. pyogenes</i>	>100

Cross Resistance Antibacterial Assay

The compounds of the present invention are tested against an assortment of ciprofloxacin resistant *E. coli* and *S. aureus* organisms described below using standard microtitration techniques (Cohen, et al., *Antimicrob. Agents Chemother.*, 1985;28:766; Heifetz, et al., *Antimicrob. Agents Chemother.*, 1974;6:124). The results of the evaluation are shown in Table 4 and compared to two 3-hydroxyquinazolin-2,4-diones.

E. coli and *S. aureus* organisms:

E. coli JL4: *E. coli* W3110 + Tol C⁻ (Tol C pump deficient strain with *E. coli* W3110 background)

E. coli LLM-1: Isogenic to *E. coli* JL4 with a D87G mutation (aspartic acid to glycine at position 87) in the QRDR (quinolone resistance determining region)



S. aureus UC-76: Typical sensitive laboratory strain (Wild type)

S. aureus 2552: Isogenic to *S. aureus* UC-76, with upregulated *norA* pump

S. aureus 2554: Isogenic to *S. aureus* 2552, with point mutation at position 80 of *grlA*.subunit

S. aureus 2558: Isogenic to *S. aureus* 2554, with point mutation at position 84 of *gyrA* subunit

TABLE 4. Antibacterial Activities Against Ciprofloxacin Resistant Strains

Compound Number or Structure	Minimum Inhibitory Concentrations $\mu\text{g/mL}$					
	<i>E. coli</i> JL4	<i>E. coli</i> LLM1	<i>S. aureus</i> UC-76	<i>S. aureus</i> 2552	<i>S. aureus</i> 2554	<i>S. aureus</i> 2558
1	0.13	0.5	0.13	1	1	1
3	0.5	1	1	2	2	4
12	0.06	0.25	1	1	1	2
	0.25	4	4	32	32	>64
	0.5	2	4	>64	>64	>64

The quinolone mimics provided by this invention display Gram-negative and Gram-positive activity. The compounds also show inhibition of bacterial DNA gyrase/activity. The compounds demonstrate in vivo protective activity in mice and are not highly cytotoxic to mammalian cells, thus indicating selectivity for bacteria. The compounds of Formula I are thus useful for treating and preventing bacterial disease and growth in both animals and on surfaces. The compounds can be applied to wood and metal surfaces, for instance, to eliminate bacterial growth. The compounds can be administered to animals such as humans, horses, dogs, sheep, and cattle to prevent and treat bacterial infections.

The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms for convenient administration to animals. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of the present invention can be administered by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. It will be obvious to those skilled in the art that the

-235-

following dosage forms may comprise as the active component, either a compound of Formula I or a corresponding pharmaceutically acceptable salt of a compound of Formula I. The formulations typically will comprise from about 1 to about 95 percent by weight of the active invention compound. The compounds
5 can be dissolved or suspended in liquids, such as ethanol, and applied to wood or metal surfaces to prevent and/or kill bacterial growth.

For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets,
10 suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

15 In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium
20 carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers,
25 is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is
30 dispersed homogeneously therein, as by stirring. The molten homogenous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

-236-

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water propylene glycol solutions. For parenteral injection liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

5 Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents as desired.

 Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or,
10 synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

 Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These
15 preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

 The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is divided into unit doses containing appropriate quantities of
20 the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

25 The quantity of invention compound in a unit dose preparation may be varied or adjusted from 0.1 mg to 100 mg, preferably from 0.5 mg to 100 mg, according to the particular application and the potency of the active component as determined by a skilled physician. The composition can, if desired, also contain other compatible therapeutic agents.

30 In therapeutic use as agents for the treatment of infections caused by a bacteria, the compounds utilized in the pharmaceutical method of this invention are administered at the initial dosage of about 0.01 mg to about 500 mg/kg daily.

-237-

A daily dose range of about 0.01 mg to about 100 mg/kg is preferred. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the particular compound being employed. Determination of the proper dosage for a particular situation is within the skill of the medical art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day, if desired.

The following additional examples illustrate typical formulations for use according to the invention.

EXAMPLE 37

1. Tablet Formulation

Ingredient	Amount (mg)
Compound No. 10	50
Cornstarch (mix)	15
Cornstarch (paste)	5
Lactose	50
Calcium stearate	2
Dicalcium phosphate	28
	150

Compound 10 is blended to uniformity with the cornstarch mix, lactose, and dicalcium phosphate. The cornstarch paste is prepared as a 10% aqueous paste, and it is blended to uniformity with the other mixture. The wet granulation is passed through a standard 8-mesh screen. The wet granules are dried and then blended to uniformity with the calcium stearate and pressed into a tablet. Such tablets are administered orally to patients suffering from bacterial infections at the rate of from one to about four times a day.

-238-

2. Capsule Formulation

Ingredients	Amount (mg)
Compound No. 25	120
Lactose USP	150
Starch	100
Magnesium stearate	5
	150

The ingredients are blended to uniformity and packed into a soft gelatin capsule. The capsules are administered at the rate of one to two per day to a patient infected with bacteria resistant to commercially available quinolone antibiotics.

3. Liquid Preparation

Ingredient	Amount
Compound No. 18	500 mg
Water	1000 mL

5 The compound is dissolved in water, and the solution is sprayed onto wood surfaces and in shower stalls to control and eliminate bacterial growth.


10 The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

-239-

CLAIMS

What is claimed is:

1. A compound of Formula I



I

5

or a pharmaceutically acceptable salt thereof wherein:

R₁ and R₃ independently are H,

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,

C₂-C₁₂ alkynyl and substituted alkynyl,

10

C₃-C₁₂ cycloalkyl and substituted cycloalkyl,

aryl and substituted aryl,

heterocyclic and substituted heterocyclic,

or heteroaryl and substituted heteroaryl;

R₅, R₆, and R₈ independently are H,

15

(O)_nC₁-C₁₂ alkyl and substituted alkyl,

(O)_nC₂-C₁₂ alkenyl and substituted alkenyl,

(O)_nC₂-C₁₂ alkynyl and substituted alkynyl,

wherein n is 0 or 1,

halo,

20

NO₂,

CN,

NR'R'';

R' and R'' independently are H,

C₁-C₁₂ alkyl and substituted alkyl,

-240-

C₂-C₁₂ alkenyl and substituted alkenyl,

C₂-C₁₂ alkynyl and substituted alkynyl,

CO-C₁-C₁₂ alkyl and substituted alkyl,

or R' and R'' taken together with the nitrogen to which they are

5 attached form a 3- to 7-membered ring containing from 1 to
3 heteroatoms selected from N, O, and S, said ring being
unsubstituted or substituted with up to 4 substituent groups;

R₁ and R₈ can be taken together with the atoms to which they are attached

10 to form a cyclic ring having from 1 to 3 heteroatoms selected from
N, O, and S, and optionally substituted by R' and R'';


R₇ is selected from

15


20

25

240a




240b



J and K independently are C or N, provided that when J or K is N, R₆ or R_g is absent at that position.

-241-

2. A compound of Formula II



II

or a pharmaceutically acceptable salt thereof,
wherein:

R₁ and R₃ independently are H,

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,

C₂-C₁₂ alkynyl and substituted alkynyl,

C₃-C₁₂ cycloalkyl and substituted cycloalkyl,

aryl and substituted aryl,

heterocyclic and substituted heterocyclic,

or heteroaryl and substituted heteroaryl;

R₆ and R₈ independently are H,

(O)_nC₁-C₁₂ alkyl and substituted alkyl,

(O)_nC₂-C₁₂ alkenyl and substituted alkenyl

(O)_nC₂-C₁₂ alkynyl and substituted alkynyl,

wherein n is 0 or 1,


halo,

NO₂,


CN,

NR'R'';

R₇ is selected from




242a



242b

(3.) A compound of Formula III



III

15

wherein

R₃ is H,

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,

C₂-C₁₂ alkynyl and substituted alkynyl,

20

C₃-C₁₂ cycloalkyl and substituted cycloalkyl,

aryl and substituted aryl,

heterocyclic and substituted heterocyclic,

or heteroaryl and substituted heteroaryl;

R₆ is H,

25

(O)_nC₁-C₁₂ alkyl and substituted alkyl,

-243-

(O)_nC₂-C₁₂ alkenyl and substituted alkenyl,

(O)_nC₂-C₁₂ alkynyl and substituted alkynyl,

wherein n is 0 or 1,

halo,

5


NO₂,

CN,


NR'R'';

R₇ is selected from

243a



243b



243c


R' is H,

25

C₁-C₁₂ alkyl and substituted alkyl,C₂-C₁₂ alkenyl and substituted alkenyl,C₂-C₁₂ alkynyl and substituted alkynyl, orCO-C₁-C₁₂ alkyl and substituted alkyl.

-244-

4. A compound of Formula IV



IV

or a pharmaceutically acceptable salt thereof,
wherein:

5 R_1 and R_3 independently are H,

C_1 - C_{12} alkyl and substituted alkyl,

C_2 - C_{12} alkenyl and substituted alkenyl,

C_2 - C_{12} alkynyl and substituted alkynyl,

C_3 - C_{12} cycloalkyl and substituted cycloalkyl,

10 aryl and substituted aryl,

heterocyclic and substituted heterocyclic,

or heteroaryl and substituted heteroaryl;

R_6 is H,

$(O)_n C_1$ - C_{12} alkyl and substituted alkyl,

15 $(O)_n C_2$ - C_{12} alkenyl and substituted alkenyl,

$(O)_n C_2$ - C_{12} alkynyl and substituted alkynyl,

wherein n is 0 or 1,

halo,


NO_2 ,

20 CN,


$NR'R''$;

R_7 is selected from

25



245a



245b

R' and R'' independently are H;

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,

15

C₂-C₁₂ alkynyl and substituted alkynyl,

CO-C₁-C₁₂ alkyl and substituted alkyl,

or R' and R'' taken together with the nitrogen to which they are


attached form a 3- to 7-membered ring containing from 1 to

3 heteroatoms selected from N, O, and S, said ring being

20

unsubstituted or substituted with up to 4 substituent groups.

5. A compound of Formula V



V

or a pharmaceutically acceptable salt thereof,

wherein:

25

R₁ and R₃ independently are H,

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,
C₂-C₁₂ alkynyl and substituted alkynyl,
C₃-C₁₂ cycloalkyl and substituted cycloalkyl,
aryl and substituted aryl,
5 heterocyclic and substituted heterocyclic,
or heteroaryl and substituted heteroaryl;

R₆ and R₈ independently are H,

(O)_nC₁-C₁₂ alkyl and substituted alkyl,
(O)_nC₂-C₁₂ alkenyl and substituted alkenyl,
10 (O)_nC₂-C₁₂ alkynyl and substituted alkynyl,

wherein n is 0 or 1,

halo,

NO₂,

CN,


15 NR'R'';

R' and R''

taken together with the nitrogen to which they are
attached form a 3- to 7-membered ring containing from 1 to
3 heteroatoms selected from N, O, and S, said ring being
unsubstituted or substituted with up to 4 substituent groups.

6. A compound of Formula VI

-247-



VI

or a pharmaceutically acceptable salt thereof,

wherein:

R₁ and R₃ independently are H,

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,

C₂-C₁₂ alkynyl and substituted alkynyl,

C₃-C₁₂ cycloalkyl and substituted cycloalkyl,

aryl and substituted aryl,

heterocyclic and substituted heterocyclic,

or heteroaryl and substituted heteroaryl;

R₆ is H,

(O)_nC₁-C₁₂ alkyl and substituted alkyl,

(O)_nC₂-C₁₂ alkenyl and substituted alkenyl,

(O)_nC₂-C₁₂ alkynyl and substituted alkynyl,

wherein n is 0 or 1,

halo,

NO₂,

CN,


NR'R'';

R' and R''

-248-

taken together with the nitrogen to which they are attached form a 3- to 7-membered ring containing from 1 to 3 heteroatoms selected from N, O, and S, said ring being unsubstituted or substituted with up to 4 substituent groups.

5 7. A compound of Formula VII




VII

or a pharmaceutically acceptable salt thereof,
wherein:


R₁ and R₃ independently are H,

- 10 C₁-C₁₂ alkyl and substituted alkyl,
 C₂-C₁₂ alkenyl and substituted alkenyl,
 C₂-C₁₂ alkynyl and substituted alkynyl,
 C₃-C₁₂ cycloalkyl and substituted cycloalkyl,
 aryl and substituted aryl,
15 heterocyclic and substituted heterocyclic,
 or heteroaryl and substituted heteroaryl;

R₇ is selected from



249a.



R₈ is H,

(O)_nC₁-C₁₂ alkyl and substituted alkyl,

10

(O)_nC₂-C₁₂ alkenyl and substituted alkenyl,

(O)_nC₂-C₁₂ alkynyl and substituted alkynyl,

wherein n is 0 or 1,

halo,

NO₂,

15

CN,

NR'R'';

wherein R' and R'' independently are H,

C₁-C₁₂ alkyl and substituted alkyl,

C₂-C₁₂ alkenyl and substituted alkenyl,

20

C₂-C₁₂ alkynyl and substituted alkynyl,

CO-C₁-C₁₂ alkyl and substituted alkyl,

or R' and R'' taken together with the nitrogen to which they are
attached form a 3- to 7-membered ring containing from 1 to
3 heteroatoms selected from N, O, and S, said ring being
25 unsubstituted or substituted with up to 4 substituent groups;

R₁ and R₈ can be taken together with the atoms to which they are attached
to form a cyclic ring having from 1 to 3 heteroatoms selected from
N, O, and S, and optionally substituted by R' and R'';

R₇ is hydrogen,

30

C₁-C₁₂ alkyl and substituted alkyl,

-250-

C₂-C₁₂ alkenyl and substituted alkenyl,

C₂-C₁₂ alkynyl and substituted alkynyl,

halo,

NO₂,

5

CN,

NR'R'',

COOR',

aryl,

fused aryl,

10

heterocyclic,

fused heterocyclic,

bicyclic heterocyclic, or

spiro heterocyclic,

wherein all ring groups can be substituted; and

15

J and K independently are C or N, provided that when J or K is N,

R₆ or R₈ is absent at that position.

8. A compound of Claim 1 wherein R₇ is pyrrolidinyl, substituted pyrrolidinyl, piperazinyl, substituted piperazinyl, piperidinyl or substituted piperidinyl.
9. A compound selected from:
- 5 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;
- 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-piperazin-1-yl-1*H*-quinazoline-2,4-dione;
- 10 3-Amino-7-[3-(aminomethyl)-3-methylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;
- 3-Amino-7-(6-amino-3-aza-bicyclo[3.1.0]hex-3-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazolin-2,4-dione;
- 3-Amino-7-((*S*)-3-*N*-methylaminopyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-8-chloro-1-(2,4-difluoroanilino)-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(S)-3-aminopyrrolidin-1-yl]-1-cyclopropylamino-6,8-difluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(S)-3-aminopyrrolidin-1-yl]-1-cyclopropylamino-5,8-difluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-5,6,8-trifluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

7-((*S*)-3-Aminopyrrolidin-1-yl)-1-cyclopropyl-3,5-diamino-6,8-difluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-hydroxymethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminoethylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

-254-

3-Amino-7-((S)-7-amino-5-azaspiro[2.4]hept-5-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

5 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(pyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

10 3-Amino-7-(3-aminopyrrolidin-1-yl)-8-chloro-6-fluoro-1-isopropyl-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-*sec*-butyl-8-chloro-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-[3-aminopyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1*H*-quinazoline-2,4-dione;

15 3-Amino-7-[3-aminopyrrolidin-1-yl]-8-chloro-1-cyclobutylamino-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-fluoro-1*H*-quinazoline-2,4-dione;

20 3-Amino-7-(3-aminopyrrolidin-1-yl)-6-chloro-1-cyclopropyl-8-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-1-cyclopropylmethyl-8-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropylmethyl-8-fluoro-1*H*-quinazoline-2,4-dione;

25 3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-1-ethyl-8-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-1-ethyl-6-fluoro-7-[(R)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione;

30 3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

-255-

3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-fluoro-6-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-8-ethoxy-6-fluoro-1H-quinazoline-2,4-dione;

10 3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-8-carbonitrile;

3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(3-[1,2,3-triazol]-1-yl)-pyrrolidin-1-yl)-1H-quinazoline-2,4-dione;

3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(S)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(S)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

5-Amino-9-((S)-3-aminopyrrolidin-1-yl)-8-fluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione;

5-Amino-9-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-8-fluoro-3-methyl-2,3-dihydro-1-oxa-3a,5-diazaphenalene-4,6-dione;

25 2-Amino-8-((S)-3-aminopyrrolidin-1-yl)-9-fluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione,;

2-Amino-8-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-9-fluoro-5-methyl-6,7-dihydro-5H-pyrido[3,2,1-ij]quinazoline-1,3-dione;

30 3-Amino-7-((S)-3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

3-Amino-7-(6-amino-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione,;

-256-

3-Amino-7-(3-aminomethyl-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione,;

3-Amino-1-cyclopropyl-6-fluoro-7-(octahydropyrrolo[3,4-c]pyridin-2-yl)-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

5 3-Amino-1-cyclopropyl-6-fluoro-7-(octahydropyrrolo[3,4-b]pyridin-2-yl)-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

3-Amino-7-[(R)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

10 3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

trans-3-Amino-7-(3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-7-[(R)-3-((R)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

15 3-Amino-7-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-1H-pyrido[2,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-aminopropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-[(R)-3-((S)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione;

3-Amino-7-[7-(1-aminoethyl)-5-azaspiro[2.4]hept-5-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)piperidine-3-carboxylic acid amide;

25 3-Amino-[7-trans-3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-{3-[(2,2,2-trifluoroethylamino)methyl]pyrrolidin-1-yl}-1H-quinazoline-2,4-dione;

30 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(5-methylhexahydropyrrolo[3,4-c]pyrrol-2-yl)-1H-quinazoline-2,4-dione;

3-Amino-8-chloro-1-cyclopropyl-7-(2,7-diazaspiro[4.4]non-2-yl)-6-fluoro-1H-quinazoline-2,4-dione;

-257-

3-Amino-7-(3-aminomethyl-3-benzylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(*R*)-3-((*S*)-1-aminoethyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

5 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-(3-hydroxyimino-pyrrolidin-1-yl)-1*H*-quinazoline-2,4-dione;

3-Amino-7-[trans-3-amino-4-trifluoromethylpyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

10 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-[(*R*)-3-(*S*)-1-methylaminoethyl]pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione;

3-Amino-7-(trans-3-amino-4-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

3-Amino-7-[trans-3-amino-4-(4-hydroxyphenyl)pyrrolidin-1-yl]-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

15 *N*-[1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-ylmethyl]methanesulfonamide;

3-Amino-7-(3-aminomethylpiperidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1*H*-quinazoline-2,4-dione;

20 3-Amino-8-chloro-1-cyclopropyl-6-fluoro-7-[3-(isopropylaminomethyl)pyrrolidin-1-yl]-1*H*-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylpyrrolidin-1-yl)-6,8-dichloro-1-cyclopropyl-1*H*-quinazoline-2,4-dione;

N-[1-(3-Amino-8-chloro-1-cyclopropyl-6-fluoro-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-ylmethyl]methanesulfonamide;

25 3-Amino-7-[(*R*)-3-(*S*)-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

3-Amino-7-[(*R*)-3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

30 3-Amino-7-[3-(1-aminoethyl)-3-methoxypyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1*H*-quinazoline-2,4-dione;

3-Amino-7-[3-(1-aminoethyl)-3-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1*H*-quinazoline-2,4-dione;

-258-

3-Amino-7-(3-aminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-5-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-(S)-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-5-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-(3-aminomethyl-3-methoxymethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyl-3-fluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

10 3-Amino-7-(trans-3-aminomethyl-4-trifluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[3-(1-aminoethyl)piperidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[3-(aminoethyl)piperidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[4-(1-aminoethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[4-(1-aminoethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-(3-amino-3-phenylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[3-(1-aminoethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyl-3-phenylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-(7-aminomethyl-5-azaspiro[2,4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(7-aminomethyl-5-azaspiro[2,4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-(3-aminomethyl-3-hydroxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

-259-

3-Amino-7-(3-amino-4-methoxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-amino-4-methoxypyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-(3-amino-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyl-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(R)-3-((S)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-ethyl-6-fluoro-1H-quinazoline-2,4-dione;

1-(3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid trifluoroacetate;

1-(3-Amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidine-3-carboxylic acid trifluoroacetate;

15 3-Amino-7-(1-aminomethyl-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(1-aminomethyl-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(S)-3-((R)-1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-(octahydropyrrolo[3,4-b]pyridin-6-yl)-1H-quinazoline-2,4-dione;

25 3-Amino-7-(trans-3-aminomethyl-4-methylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(trans-3-aminomethyl-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-(trans-3-aminomethyl-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(trans-3-amino-4-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-260-

3-Amino-7-(3-aminomethylmorpholin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[3-(1-aminoethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-amino-3-phenylpyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

10 3-Amino-7-(3-amino-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-8-methyl-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-7-pyrrolidin-1-yl-1H-quinazoline-2,4-dione;

15 3-Amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)-pyrrolidin-1-yl]-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-7-[3-(1-hydroxy-1-methylethyl)-pyrrolidin-1-yl]-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-1-cyclopropyl-6-fluoro-5-methyl-7-[(R)-3-((S)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione;

(S)-1-[(R)-1-(3-Amino-1-cyclopropyl-7-[3-(1-ethylaminoethyl)pyrrolidin-1-yl]-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-1-cyclopropyl-7-[(R)-3-((S)-1-ethylaminoethyl)pyrrolidin-1-yl]-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-(6-amino-1-methyl-3-azabicyclo[3.2.0]hept-3-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-(4-aminomethylpiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylazetidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-261-

cis-3-Amino-7-(3-aminomethyl-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

trans-3-Amino-7-(3-aminomethyl-4-fluoropyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-(1-amino-5-azaspiro[2.4]hept-5-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyloctahydroisoindol-2-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-(3-aminomethyloctahydroisoindol-2-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-aminoethylpyrrolidin-1-yl)]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(6-amino-3-azabicyclo[3.1.0]hex-3-yl)-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[3-(1-amino-1-methylethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-1-cyclopropyl-6-fluoro-7-[3-(isopropylaminomethyl)pyrrolidin-1-yl]-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-7-[(4aS,7aS)-octahydropyrrolo[3,4-b]pyridin-6-yl]-1H-quinazoline-2,4-dione;

3-Amino-7-(3-amino-4-fluoromethylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethyl-3-methylpyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-(4-aminopiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminopiperidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-1-cyclopropyl-6-fluoro-8-methoxy-7-[(R)-3-((S)-1-methylaminoethyl)pyrrolidin-1-yl]-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminoazetidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-262-

3-Amino-7-[(3R,4S)-3-((S)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4R)-3-(R)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(3R,4S)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4R)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-(-1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

-263-

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-(R)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((S)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-(S)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-(R)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(R)-3-((S)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(R)-4-((R)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((S)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(R)-4-((S)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(R)-3-((S)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

-265-

3-Amino-7-[(R)-3-((R)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(R)-4-((R)-1-amino-2-ethoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-ethoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(R)-3-((S)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

-266-

3-Amino-7-[(R)-3-((S)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-
1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-
methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-
2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-
methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-
2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-3-methoxypropyl)pyrrolidin-1-
yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-3-methoxypropyl)pyrrolidin-1-
yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-
methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-
2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-
methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-quinazoline-
2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-quinazoline-2,4-dione;

{(R)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-
methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-
yl]ethyl}urea;

{(R)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-8-methyl-2,4-
dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

-267-

{(S)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

5 {(S)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(R)-2-Amino-2-[(3R,4S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

10 {(R)-2-Amino-2-[(3R,4S)-1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

{(S)-2-Amino-2-[(3R,4R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

15 {(S)-2-Amino-2-[(3R,4R)-1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

{2-Amino-2-[(1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]ethyl}urea;

20 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]ethyl}urea;

{2-Amino-2-[(1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}urea;

25 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}urea;

3-Amino-7-[(3R,4S)-3-((S)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(3R,4S)-3-((S)-1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

-268-

3-Amino-7-[(3R,4R)-3-(R)-1-aminoethyl]-4-fluoropyrrolidin-1-yl]-
1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-(R)-1-aminoethyl]-4-fluoropyrrolidin-1-
yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4S)-3-(1-amino-1-methylethyl)-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-
2,4-dione;

3-Amino-7-[(3R,4S)-3-(1-amino-1-methylethyl)-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-
10 dione;

3-Amino-7-[(3R,4R)-3-(1-amino-1-methylethyl)-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-
2,4-dione;

3-Amino-7-[(3R,4R)-3-(1-amino-1-methylethyl)-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-
15 dione;

3-Amino-7-[(3R,4S)-3-(R)-1-amino-2-methoxyethyl]-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-
2,4-dione;

20 3-Amino-7-[(3R,4S)-3-(R)-1-amino-2-methoxyethyl]-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-
dione;

3-Amino-7-[(3R,4R)-3-(S)-1-amino-2-methoxyethyl]-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-
25 2,4-dione;

3-Amino-7-[(3R,4R)-3-(S)-1-amino-2-methoxyethyl]-4-
fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-
dione;

3-Amino-7-[(3R,4S)-3-(R)-1-amino-2-methoxyethyl]-4-
methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-
30 quinazoline-2,4-dione;

-269-

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(R)-4-((R)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-(R)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(R)-4-((S)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(R)-4-(S)-1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(R)-3-(R)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

-270-

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(R)-3-((S)-1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(3R,4S)-3-((S)-1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(R)-4-((R)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(R)-4-((S)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-271-

3-Amino-7-[(R)-4-((S)-1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(R)-3-((S)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(R)-3-((R)-1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(3R,4S)-3-((R)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-4-((R)-1-amino-2-ethoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(R)-4-((R)-1-amino-2-ethoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-272-

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

5 3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((S)-1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

10 3-Amino-7-[(R)-3-((S)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((S)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

15 3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

20 3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

25 3-Amino-7-[(3R,4S)-3-((S)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(R)-3-((R)-1-amino-3-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

30 3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-273-

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

3-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methoxy-1H-quinazoline-2,4-dione;

-Amino-7-[(3R,4R)-3-((R)-1-amino-3-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methoxy-1H-quinazoline-2,4-dione;

{(R)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(R)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(S)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(S)-2-Amino-2-[(R)-1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)pyrrolidin-3-yl]ethyl}urea;

{(R)-2-Amino-2-[(3R,4S)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

{(R)-2-Amino-2-[(3R,4S)-1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

{(S)-2-Amino-2-[(3R,4R)-1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

{(S)-2-Amino-2-[(3R,4R)-1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-methylpyrrolidin-3-yl]ethyl}urea;

-274-

{2-Amino-2-[(1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]ethyl}urea;

5 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4-fluoropyrrolidin-3-yl]ethyl}urea;

{2-Amino-2-[(1-(3-amino-1-cyclopropyl-6-fluoro-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}urea;

10 {2-Amino-2-[(1-(3-amino-1-cyclopropyl-8-methoxy-2,4-dioxo-1,2,3,4-tetrahydroquinazolin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethyl}urea;

3-Amino-7-[3-(1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

15 3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

20 3-Amino-7-[4-(1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)-pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

25 3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[4-(1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

30 3-Amino-7-[3-(1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

-275-

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4,4-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

5 3-Amino-7-[3-(1-amino-2-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

10 3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[4,3-d]pyrimidine-2,4-dione;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[4,3-d]pyrimidin-7-yl)pyrrolidin-3-yl]ethylurea;

15 {2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[4,3-d]pyrimidin-7-yl)-4-methylpyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[4,3-d]pyrimidin-7-yl)-4-fluoropyrrolidin-3-yl]ethylurea;

20 {2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[4,3-d]pyrimidin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethylurea;

3-Amino-7-[3-(1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

25 3-Amino-7-[3-(1-aminoethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-1-methylethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

30 3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

-276-

3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-fluoropyrrolidin-1-yl]-
1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-
yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-
dione;

3-Amino-7-[3-(1-amino-2-methoxyethyl)-4-methylpyrrolidin-1-
yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[4-(1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-
1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-
dione;

3-Amino-7-[4-(1-amino-2-methoxyethyl)-3,3-dimethylpyrrolidin-
1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-
cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)pyrrolidin-1-yl]-1-
cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-
1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-methylpyrrolidin-1-yl]-
1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-
1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-phenoxyethyl)-4-fluoropyrrolidin-1-yl]-
1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[4-(1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-
1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-
dione;

3-Amino-7-[4-(1-amino-2-phenoxyethyl)-3,3-dimethylpyrrolidin-
1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-
cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

-277-

3-Amino-7-[3-(1-amino-2-ethoxyethyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

5 3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

10 3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4,4-dimethylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

15 3-Amino-7-[3-(1-amino-2-ethoxyethyl)-4,4-dimethylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)pyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

20 3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-methylpyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

25 3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

3-Amino-7-[3-(1-amino-2-methoxypropyl)-4-fluoropyrrolidin-1-yl]-1-cyclopropyl-8-methyl-1H-pyrido[3,2-d]pyrimidine-2,4-dione;

30 {2-Amino-2-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)pyrrolidin-3-yl]ethylurea;

-278-

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)pyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4-methylpyrrolidin-3-yl]ethylurea;

5

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4-methylpyrrolidin-3-yl]ethylurea;

10

{2-Amino-2-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4-fluoropyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4-fluoropyrrolidin-3-yl]ethylurea;

15

{2-Amino-2-[1-(3-amino-1-cyclopropyl-6-fluoro-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethylurea;

{2-Amino-2-[1-(3-amino-1-cyclopropyl-8-methyl-2,4-dioxo-1,2,3,4-tetrahydropyrido[3,2-d]pyrimidin-7-yl)-4,4-dimethylpyrrolidin-3-yl]ethylurea;

20

3-Amino-7-(3-aminomethylphenyl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylphenyl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

25

3-Amino-7-(3-aminomethylphenyl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylphenyl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

30

3-Amino-7-(3-aminomethylphenyl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-aminomethylphenyl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

-279-

3-Amino-7-(2-aminomethylthiazol-4-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(2-aminomethylthiazol-4-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

5 3-Amino-7-(2-aminomethylthiazol-4-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(2-aminomethylthiazol-4-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-(2-aminomethylthiazol-4-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(2-aminomethylthiazol-4-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

15 3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

20 3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-[2-(1-aminoethyl)thiazol-4-yl]-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

25 3-Amino-7-(5-aminomethylthiophen-3-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(5-aminomethylthiophen-3-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

30 3-Amino-7-(5-aminomethylthiophen-3-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(5-aminomethylthiophen-3-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

-280-

3-Amino-7-(5-aminomethylthiophen-3-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(5-aminomethylthiophen-3-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

10 3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione

15 3-Amino-7-(4-aminomethyl-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

20 3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

25 3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-aminoethyl)-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

30 3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

-281-

3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-6-fluoro-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-chloro-1-cyclopropyl-1H-quinazoline-2,4-dione;

5 3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-methyl-1-cyclopropyl-1H-quinazoline-2,4-dione;

3-Amino-7-(4-(1-amino-1-methylethyl)-3,3-difluoropyrrolidin-1-yl)-8-methoxy-1-cyclopropyl-1H-quinazoline-2,4-dione;

10 3-Amino-7-(3-(1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-5,8-dimethyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-(1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-5,8-dimethyl-1H-quinazoline-2,4-dione;

3-Amino-7-(3-(1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methoxy-5-methyl-1H-quinazoline-2,4-dione;

15 3-Amino-7-(3-(1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-8-methoxy-5-methyl-1H-quinazoline-2,4-dione;

3,8-Diamino-7-[3-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-6-fluoro-5-methyl-1H-quinazoline-2,4-dione; and

20 3,8-Diamino-7-[3-(1-aminoethyl)pyrrolidin-1-yl]-1-cyclopropyl-5-methyl-1H-quinazoline-2,4-dione.

10. A pharmaceutical composition comprising a compound of Claim 1 admixed with a carrier, diluent, or excipient.

11. A pharmaceutical composition comprising a compound of Claim 2 admixed with a carrier, diluent, or excipient.

25 12. A pharmaceutical composition comprising a compound of Claim 3 admixed with a carrier, diluent, or excipient.

13. A pharmaceutical composition comprising a compound of Claim 4 admixed with a carrier, diluent, or excipient.

-282-

14. A pharmaceutical composition comprising a compound of Claim 5 admixed with a carrier, diluent, or excipient.
15. A pharmaceutical composition comprising a compound of Claim 6 admixed with a carrier, diluent, or excipient.
16. A pharmaceutical composition comprising a compound of Claim 7 admixed with a carrier, diluent, or excipient.
17. A pharmaceutical composition comprising a compound of Claim admixed with a carrier, diluent, or excipient.
18. A use of an antibacterial effective amount of a compound of Claim 1 in the manufacture of a medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof.
19. A use of an antibacterial effective amount of a compound of Claim 2 in the manufacture of a medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof.
20. A use of an antibacterial effective amount of a compound of Claim 3 in the manufacture of a medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof.
21. A use of an antibacterial effective amount of a compound of Claim 4 in the manufacture of a medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof.
22. A use of an antibacterial effective amount of a compound of Claim 5 in the manufacture of a medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof.

-283-

23. A use of an antibacterial effective amount of a compound of Claim 6 in the manufacture of medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof
24. A use of an antibacterial effective amount of a compound of Claim 7 in the manufacture of medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof.
25. A use of an antibacterial effective amount of a compound of Claim 9 in the manufacture of a medicament for treating a bacterial infection in a mammal comprising administering to the mammal in need thereof.
26. A use of an effective amount of a compound of Claim 1 in the manufacture of a medicament for inhibiting a wild-type mutant of DNA gyrase in a mammal comprising administering to the mammal in need thereof
27. A use of an effective amount of a compound of Claim 1 in the manufacture of a medicament for inhibiting a quinolone resistant mutant of DNA gyrase in a mammal comprising administering to the mammal in need thereof.