


exchange of GDP with GTP in eIF2 is catalyzed by its dedicated guanine nucleotide exchange factor (GEF) eIF2B. eIF2B exists as a decamer (B1₂B2₂B3₂B4₂B5₂) or dimer of two pentamers in cells (Gordiyenko, Y. et al, *Nat Commun* (2014) 5:3902; Wortham, N.C. et al, *FASEB J* (2014) 28:2225-2237). Molecules such as ISRIB interact with and stabilize the eIF2B dimer conformation, thereby enhancing intrinsic GEF activity and making cells less sensitive to the cellular effects of phosphorylation of eIF2 α (Sidrauski, C. et al, *eLife* (2015) e07314; Sekine, Y. et al, *Science* (2015) 348:1027-1030). As such, small molecule therapeutics that can modulate eIF2B activity may have the potential to attenuate the PERK branch of the UPR and the overall ISR, and therefore may be used in the prevention and/or treatment of various diseases, such as a neurodegenerative disease, a leukodystrophy, cancer, an inflammatory disease, a musculoskeletal disease, or a metabolic disease.

SUMMARY

In one aspect, the present disclosure refers to a compound of Formula (I):



Formula (I)

or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, *N*-oxide, or stereoisomer thereof, wherein:

—— D is a bridged bicyclic cycloalkyl, bridged bicyclic heterocyclyl, or cubanyl, wherein each bridged bicyclic cycloalkyl, bridged bicyclic heterocyclyl, or cubanyl is optionally substituted on one or more available carbons with 1-4 R^X; and wherein if the bridged bicyclic heterocyclyl contains a substitutable nitrogen moiety, the substitutable nitrogen may be optionally substituted by R^{N1};

—— L¹ is a bond, C₁-C₆ alkylene, 2-7 membered heteroalkylene, NR^{N2}, or O, wherein C₁-C₆ alkylene or 2-7 membered heteroalkylene is optionally substituted with 1-5 R^{L1};

—— L² is a bond, C₁-C₆ alkylene, or 2-7 membered heteroalkylene, wherein C₁-C₆ alkylene or 2-7 membered heteroalkylene is optionally substituted with 1-5 R^{L2};

—— R¹ is hydrogen or C₁-C₆ alkyl;

—— R² is hydrogen or C₁-C₆ alkyl;

—— W is a 8-10 membered, partially unsaturated, fused bicyclic ring moiety comprising a 5-6 membered heterocyclyl fused to a phenyl or 5-6 membered heteroaryl; wherein the heterocyclyl may be optionally substituted on one or more available saturated carbons with

~~1-4 R^{W1}; and wherein the phenyl or heteroaryl may optionally be substituted on one or more available unsaturated carbons with 1-4 R^{W2}; and wherein if the heterocyclyl contains a substitutable nitrogen moiety, the substitutable nitrogen may optionally be substituted with R^{N3}; and wherein W is attached to L² through an available saturated carbon or nitrogen atom within the heterocyclyl;~~

~~——— A is phenyl or 5-6-membered heteroaryl, wherein phenyl or 5-6-membered heteroaryl is optionally substituted on one or more available carbons with 1-5 R^Y; and wherein if the 5-6-membered heteroaryl contains a substitutable nitrogen moiety, the substitutable nitrogen may be optionally substituted by R^{N4};~~

~~——— each R^{L1} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, oxo, halo, cyano, —OR^A, —NR^BR^C, —NR^BC(O)R^D, —C(O)NR^BR^C, —C(O)R^D, —C(O)OH, —C(O)OR^D, —SR^E, —S(O)R^D, and —S(O)₂R^D;~~

~~——— each R^{L2} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, oxo, halo, cyano, —OR^A, —NR^BR^C, —NR^BC(O)R^D, —C(O)NR^BR^C, —C(O)R^D, —C(O)OH, —C(O)OR^D, —SR^E, —S(O)R^D, and —S(O)₂R^D;~~

~~——— R^{N1} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl, halo-C₂-C₆ alkyl, amino-C₂-C₆ alkyl, cyano-C₂-C₆ alkyl, —C(O)NR^BR^C, —C(O)R^D, —C(O)OR^D, and —S(O)₂R^D;~~

~~——— R^{N2} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl, halo-C₂-C₆ alkyl, amino-C₂-C₆ alkyl, cyano-C₂-C₆ alkyl, —C(O)NR^BR^C, —C(O)R^D, —C(O)OR^D, and —S(O)₂R^D;~~

~~——— R^{N3} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkenyl, —C(O)-C₁-C₆ alkyl, —C(O)-C₁-C₆ cycloalkyl, C₁-C₆ alkyl-CO₂H, C₁-C₆ alkyl-CO₂-C₁-C₆ alkyl, —C(O)-C₁-C₃ alkyl-O-C₁-C₃ alkyl-O-C₁-C₃ alkyl, —C(O)-phenyl, —C(O)-heteroaryl, —C(O)-heterocyclyl, —S-C₁-C₆ alkyl, —S(O)₂-C₁-C₆ alkyl, —S(O)₂-phenyl, —S(O)₂-heteroaryl, —C(O)NR^BR^C and —C(O)OR^D;~~

~~——— wherein C₁-C₆ alkyl, C₁-C₆ alkenyl, C(O)-C₁-C₆ alkyl, C(O)-C₁-C₆ cycloalkyl, C₁-C₆ alkyl-CO₂H, C₁-C₆ alkyl-CO₂-C₁-C₆ alkyl, —C(O)-heterocyclyl, —S-C₁-C₆ alkyl and —S(O)₂-C₁-C₆ alkyl may optionally be substituted by one or more substituents each independently selected from the group consisting of fluoro, hydroxyl, C₁-C₆ alkoxy, C₁-C₆ alkyl (optionally substituted by one, two or three fluorine atoms) and S(O)_w-C₁₋₆ alkyl (wherein w is 0, 1 or 2); and~~

~~wherein C(O) phenyl, C(O) heteroaryl, S(O)₂ phenyl and S(O)₂ heteroaryl may optionally be substituted by one or more substituents each independently selected from the group consisting of halogen, hydroxyl, C₁-C₆ alkyl (optionally substituted by one, two or three fluorine atoms), C₁-C₆ alkoxy (optionally substituted by one, two or three fluorine atoms), S(O₂)NR^BR^C and SO₂F;~~

~~R^{N4} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy C₂-C₆ alkyl, halo C₂-C₆ alkyl, amino C₂-C₆ alkyl, cyano C₂-C₆ alkyl, C(O)NR^BR^C, C(O)R^D, C(O)OR^D, and S(O)₂R^D;~~

~~each R^{W1} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl (optionally substituted by CO₂H), hydroxy C₁-C₆ alkyl, hydroxy C₂-C₆ alkyl O, halo C₁-C₆ alkyl, amino C₁-C₆ alkyl, cyano C₁-C₆ alkyl, oxo, halo, cyano, OR^A, NR^BR^C, NR^BR^{CC}, NR^BC(O)R^D, C(O)NR^BR^C, C(O)R^D, C(O)OH, C(O)OR^D, SR^E, S(O)R^D, and S(O)₂R^D;~~

~~each R^{W2} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy C₁-C₆ alkyl, hydroxy C₂-C₆ alkyl O, halo C₁-C₆ alkyl, halo C₁-C₆ alkoxy, amino C₁-C₆ alkyl, cyano C₁-C₆ alkyl, halo, cyano, OR^A, NR^BR^C, NR^BC(O)R^D, C(O)NR^BR^C, C(O)R^D, C(O)OH, C(O)OR^D, S(R^F)_m, S(O)R^D, and S(O)₂R^D; or~~

~~2 R^{W2} groups on adjacent atoms, together with the atoms to which they are attached, form a 3-7 membered fused cycloalkyl, 3-7 membered fused heterocyclyl, fused aryl, or 5-6 membered fused heteroaryl, each of which is optionally substituted with 1-5 R^X;~~

~~each R^X is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy C₁-C₆ alkyl, halo C₁-C₆ alkyl, amino C₁-C₆ alkyl, cyano C₁-C₆ alkyl, oxo, halo, cyano, OR^A, NR^BR^C, NR^BC(O)R^D, C(O)NR^BR^C, C(O)R^D, C(O)OH, C(O)OR^D, SR^E, S(O)R^D, and S(O)₂R^D;~~

~~each R^Y is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy C₁-C₆ alkyl, halo C₁-C₆ alkyl, halo C₁-C₆ alkoxy, amino C₁-C₆ alkyl, cyano C₁-C₆ alkyl, halo, cyano, OR^A, NR^BR^C, NR^BC(O)R^D, C(O)NR^BR^C, C(O)R^D, C(O)OH, C(O)OR^D, S(R^F)_m, S(O)R^D, S(O)₂R^D, and G⁺; or~~

~~2 R^Y groups on adjacent atoms, together with the atoms to which they are attached form a 3-7 membered fused cycloalkyl, 3-7 membered fused heterocyclyl, fused aryl, or 5-6 membered fused heteroaryl, each of which is optionally substituted with 1-5 R^X;~~

~~each G⁺ is independently 3-7 membered cycloalkyl, 3-7 membered heterocyclyl, aryl, or 5-6 membered heteroaryl, wherein each 3-7 membered cycloalkyl, 3-7 membered heterocyclyl, aryl, or 5-6 membered heteroaryl is optionally substituted with 1-3 R^Z;~~

each R^Z is independently selected from the group consisting of C₁-C₆-alkyl, hydroxy-C₁-C₆-alkyl, halo-C₁-C₆-alkyl, halo, cyano, OR^A , $NR^B R^C$, $NR^B C(O)R^D$, $C(O)NR^B R^C$, $C(O)R^D$, $C(O)OH$, $C(O)OR^D$, and $S(O)_2 R^D$;

R^A is, at each occurrence, independently hydrogen, C₁-C₆-alkyl, halo-C₁-C₆-alkyl, $C(O)NR^B R^C$, $C(O)R^D$, or $C(O)OR^D$;

each of R^B and R^C is independently hydrogen or C₁-C₆-alkyl; or

R^B and R^C together with the atom to which they are attached form a 3-7 membered heterocyclyl ring optionally substituted with 1-3 R^Z ;

each R^{CC} is independently selected from the group consisting of C₁-C₆-alkyl-OH, C₁-C₆-alkyl-CO₂H and C₁-C₆-alkyl-CO₂-C₁-C₆-alkyl;

each R^D is independently C₁-C₆-alkyl or halo-C₁-C₆-alkyl;

each R^E is independently hydrogen, C₁-C₆-alkyl, or halo-C₁-C₆-alkyl;

each R^F is independently hydrogen, C₁-C₆-alkyl, or halo; and

m is 1 when R^F is hydrogen or C₁-C₆-alkyl, 3 when R^F is C₁-C₆-alkyl, or 5 when R^F is halo.

In another aspect, the present disclosure refers to a compound selected from any compound set forth in Table 1 or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, *N*-oxide, or stereoisomer thereof.

In yet another aspect, the present disclosure refers to a pharmaceutically acceptable composition comprising a compound disclosed herein and a pharmaceutically acceptable carrier.

In a further aspect, the present disclosure refers to a use of a composition in the manufacture of a medicament for treating a neurodegenerative disease, a leukodystrophy, a cancer, an inflammatory disease, an autoimmune disease, a viral infection, a skin disease, a fibrotic disease, a hemoglobin disease, a kidney disease, a hearing loss condition, an ocular disease, a musculoskeletal disease, a metabolic disease, or a mitochondrial disease in a subject, wherein the composition comprises a compound disclosed herein.

In another aspect, the present disclosure refers to a use of a composition in the manufacture of a medicament for treating a disease related to a modulation of eIF2B activity or levels, eIF2 α activity or levels, or the activity or levels of a component of the eIF2 pathway or the ISR pathway, wherein the composition comprises a compound disclosed herein.

In yet another aspect, the present disclosure refers to a use of a composition in combination with an immunotherapeutic agent, in the manufacture of a medicament for treating cancer in a subject, wherein the composition comprises a compound disclosed herein.


~~In a further aspect, the present disclosure refers to a composition for use in treating a neurodegenerative disease, a leukodystrophy, a cancer, an inflammatory disease, an autoimmune disease, a viral infection, a skin disease, a fibrotic disease, a hemoglobin disease, a kidney disease, a hearing loss condition, an ocular disease, a musculoskeletal disease, a metabolic disease, or a mitochondrial disease in a subject; or~~

~~——— for use in treating a disease related to a modulation of eIF2B activity or levels, eIF2 α activity or levels, or the activity or levels of a component of the eIF2 pathway or the ISR pathway; or~~

~~——— for use in treating cancer in a subject in combination with an immunotherapeutic agent, wherein the composition comprises a compound disclosed herein.~~

~~The present disclosure is directed, at least in part, to compounds, compositions, and methods for the modulation of eIF2B (e.g., activation of eIF2B) and the attenuation of the ISR signaling pathway. In some examples, disclosed herein is an eIF2B modulator (e.g., an eIF2B activator) comprising a compound of Formula (I) or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, N-oxide, or stereoisomer thereof. In other examples, disclosed herein are methods of using a compound of Formula (I) or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, N-oxide, or stereoisomer thereof for the treatment of a disease or disorder, e.g., a neurodegenerative disease, a leukodystrophy, cancer, an inflammatory disease, a musculoskeletal disease, a metabolic disease, or a disease or disorder associated with impaired function of eIF2B or components in the ISR pathway (e.g., eIF2 pathway).~~

~~For example, disclosed herein is a compound of Formula (I):~~



~~Formula (I)~~


~~or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, N-oxide, or stereoisomer thereof, wherein:~~

~~——— D is a bridged bicyclic cycloalkyl, bridged bicyclic heterocyclyl, or cubanyl, wherein each bridged bicyclic cycloalkyl, bridged bicyclic heterocyclyl, or cubanyl is optionally substituted on one or more available carbons with 1-4 R^X; and wherein if the bridged bicyclic~~

exchange of GDP with GTP in eIF2 is catalyzed by its dedicated guanine nucleotide exchange factor (GEF) eIF2B. eIF2B exists as a decamer (B₁₂ B₂₂ B₃₂ B₄₂ B₅₂) or dimer of two pentamers in cells (Gordiyenko, Y. et al, *Nat Commun* (2014) 5:3902; Wortham, N.C. et al, *FASEB J* (2014) 28:2225-2237). Molecules such as ISRIB interact with and stabilize the eIF2B dimer conformation, thereby enhancing intrinsic GEF activity and making cells less sensitive to the cellular effects of phosphorylation of eIF2 α (Sidrauski, C. et al, *eLife* (2015) e07314; Sekine, Y. et al, *Science* (2015) 348:1027-1030). As such, small molecule therapeutics that can modulate eIF2B activity may have the potential to attenuate the PERK branch of the UPR and the overall ISR, and therefore may be used in the prevention and/or treatment of various diseases, such as a neurodegenerative disease, a leukodystrophy, cancer, an inflammatory disease, a musculoskeletal disease, or a metabolic disease.

SUMMARY


In one aspect of the present disclosure, there is provided a compound of Formula (I):



Formula (I)

or a pharmaceutically acceptable salt thereof, wherein:

D is selected from the group consisting of:



L¹ is a bond, C₁-C₆ alkylene, 2-7 membered heteroalkylene, -NR^{N2}-, or -O-, wherein C₁-C₆ alkylene or 2-7 membered heteroalkylene is optionally substituted with 1-5 R^{L1};

L² is a bond, C₁-C₆ alkylene, or 2-7 membered heteroalkylene, wherein C₁-C₆ alkylene or 2-7 membered heteroalkylene is optionally substituted with 1-5 R^{L2};

R¹ is hydrogen or C₁-C₆ alkyl;

R² is hydrogen or C₁-C₆ alkyl;

W is a 8-10 membered, partially unsaturated, fused bicyclic ring moiety comprising a 5-6 membered heterocyclyl fused to a phenyl or 5-6-membered heteroaryl; wherein the heterocyclyl may be optionally substituted on one or more available saturated carbons with 1-4 R^{W1}; and wherein the phenyl or heteroaryl may optionally be substituted on one or more available unsaturated carbons with 1-4 R^{W2}; and wherein if the heterocyclyl contains a substitutable nitrogen moiety, the substitutable nitrogen may optionally be substituted with R^{N3}; and wherein W is attached to L² through an available saturated carbon or nitrogen atom within the heterocyclyl;

A is phenyl or 5-6-membered heteroaryl, wherein phenyl or 5-6-membered heteroaryl is optionally substituted on one or more available carbons with 1-5 R^Y; and wherein if the 5-6-membered heteroaryl contains a substitutable nitrogen moiety, the substitutable nitrogen may be optionally substituted by R^{N4};

each R^{L1} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, oxo, halo, cyano, -OR^A, -NR^BR^C, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -SR^E, -S(O)R^D, and -S(O)₂R^D;

each R^{L2} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, oxo, halo, cyano, -OR^A, -NR^BR^C, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -SR^E, -S(O)R^D, and -S(O)₂R^D;

R^{N1} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl, halo-C₂-C₆ alkyl, amino-C₂-C₆ alkyl, cyano-C₂-C₆ alkyl, -C(O)NR^BR^C, -C(O)R^D, -C(O)OR^D, and -S(O)₂R^D;

R^{N2} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl, halo-C₂-C₆ alkyl, amino-C₂-C₆ alkyl, cyano-C₂-C₆ alkyl, -C(O)NR^BR^C, -C(O)R^D, -C(O)OR^D, and -S(O)₂R^D;

R^{N3} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₆ alkenyl, -C(O)-C₁-C₆ alkyl, -C(O)-C₁-C₆ cycloalkyl, C₁-C₆ alkyl-CO₂H, C₁-C₆ alkyl-CO₂-C₁-C₆ alkyl, -

C(O)-C₁-C₃ alkyl-O-C₁-C₃ alkyl-O-C₁-C₃ alkyl, -C(O)-phenyl, -C(O)-heteroaryl, -C(O)-heterocyclyl, -S-C₁-C₆ alkyl, -S(O)₂-C₁-C₆ alkyl, -S(O)₂-phenyl, -S(O)₂-heteroaryl, -C(O)NR^BR^C and -C(O)OR^D;

wherein C₁-C₆ alkyl, C₁-C₆ alkenyl, C(O)-C₁-C₆ alkyl, -C(O)-C₁-C₆ cycloalkyl, C₁-C₆ alkyl-CO₂H, C₁-C₆ alkyl-CO₂-C₁-C₆ alkyl, -C(O)-heterocyclyl, -S-C₁-C₆ alkyl and -S(O)₂-C₁-C₆ alkyl may optionally be substituted by one or more substituents each independently selected from the group consisting of fluoro, hydroxyl, C₁-C₆ alkoxy, C₁-C₆ alkyl (optionally substituted by one, two or three fluorine atoms) and S(O)_wC₁₋₆ alkyl (wherein w is 0, 1 or 2); and

wherein -C(O)-phenyl, -C(O)-heteroaryl, -S(O)₂-phenyl and -S(O)₂-heteroaryl may optionally be substituted by one or more substituents each independently selected from the group consisting of halogen, hydroxyl, C₁-C₆ alkyl (optionally substituted by one, two or three fluorine atoms), C₁-C₆ alkoxy (optionally substituted by one, two or three fluorine atoms), S(O)₂NR^BR^C and SO₂F;

R^{N4} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl, halo-C₂-C₆ alkyl, amino-C₂-C₆ alkyl, cyano-C₂-C₆ alkyl, -C(O)NR^BR^C, -C(O)R^D, -C(O)OR^D, and -S(O)₂R^D;

each R^{W1} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl (optionally substituted by -CO₂H), hydroxy-C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl-O-, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, oxo, halo, cyano, -OR^A, -NR^BR^C, -NR^BR^{CC}, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -SR^E, -S(O)R^D, and -S(O)₂R^D;

each R^{W2} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl-O-, halo-C₁-C₆ alkyl, halo-C₁-C₆ alkoxy, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, halo, cyano, -OR^A, -NR^BR^C, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -S(R^F)_m, -S(O)R^D, and -S(O)₂R^D; or

2 R^{W2} groups on adjacent atoms, together with the atoms to which they are attached, form a 3-7-membered fused cycloalkyl, 3-7-membered fused heterocyclyl, fused aryl, or 5-6 membered fused heteroaryl, each of which is optionally substituted with 1-5 R^X;

each R^X is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, oxo, halo, cyano, -OR^A, -NR^BR^C, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -SR^E, -S(O)R^D, and -S(O)₂R^D;

each R^Y is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, halo-C₁-C₆ alkoxy, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, halo, cyano, $-OR^A$, $-NR^B R^C$, $-NR^B C(O)R^D$, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OH$, $-C(O)OR^D$, $-S(R^F)_m$, $-S(O)R^D$, $-S(O)_2R^D$, and G^1 ; or

2 R^Y groups on adjacent atoms, together with the atoms to which they are attached form a 3-7-membered fused cycloalkyl, 3-7-membered fused heterocyclyl, fused aryl, or 5-6 membered fused heteroaryl, each of which is optionally substituted with 1-5 R^X ;

each G^1 is independently 3-7-membered cycloalkyl, 3-7-membered heterocyclyl, aryl, or 5-6-membered heteroaryl, wherein each 3-7-membered cycloalkyl, 3-7-membered heterocyclyl, aryl, or 5-6-membered heteroaryl is optionally substituted with 1-3 R^Z ;

each R^Z is independently selected from the group consisting of C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, halo, cyano, $-OR^A$, $-NR^B R^C$, $-NR^B C(O)R^D$, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OH$, $-C(O)OR^D$, and $-S(O)_2R^D$;

R^A is, at each occurrence, independently hydrogen, C₁-C₆ alkyl, halo-C₁-C₆ alkyl, $-C(O)NR^B R^C$, $-C(O)R^D$, or $-C(O)OR^D$;

each of R^B and R^C is independently hydrogen or C₁-C₆ alkyl; or

R^B and R^C together with the atom to which they are attached form a 3-7-membered heterocyclyl ring optionally substituted with 1-3 R^Z ;

each R^{CC} is independently selected from the group consisting of C₁-C₆ alkyl-OH, C₁-C₆ alkyl-CO₂H and C₁-C₆ alkyl-CO₂-C₁-C₆ alkyl;









each R^D is independently C₁-C₆ alkyl or halo-C₁-C₆ alkyl;









each R^E is independently hydrogen, C₁-C₆ alkyl, or halo-C₁-C₆ alkyl;









each R^F is independently hydrogen, C₁-C₆ alkyl, or halo; and

m is 1 when R^F is hydrogen or C₁-C₆ alkyl, 3 when R^F is C₁-C₆ alkyl, or 5 when R^F is halo.











In another aspect of the present disclosure, there is provided a compound, wherein the compound is selected from the group consisting of:
























- 2g -




- 2h -









,




,




,




,




,











,



















,











,

















- 2p -



or a pharmaceutically acceptable salt thereof.

The present disclosure is directed, at least in part, to compounds, compositions, and methods for the modulation of eIF2B (e.g., activation of eIF2B) and the attenuation of the ISR signaling pathway. In some examples, disclosed herein is an eIF2B modulator (e.g., an eIF2B activator) comprising a compound of Formula (I) or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, N-oxide, or stereoisomer thereof. In other examples, disclosed herein are methods of using a compound of Formula (I) or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, N-oxide, or stereoisomer thereof for the treatment of a disease or disorder, e.g., a neurodegenerative disease, a leukodystrophy, cancer, an inflammatory disease, a musculoskeletal disease, a metabolic disease, or a disease or disorder associated with impaired function of eIF2B or components in the ISR pathway (e.g., eIF2 pathway).

For example, disclosed herein is a compound of Formula (I):



Formula (I)


or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, *N*-oxide, or stereoisomer thereof, wherein:

D is a bridged bicyclic cycloalkyl, bridged bicyclic heterocyclyl, or cubanyl, wherein each bridged bicyclic cycloalkyl, bridged bicyclic heterocyclyl, or cubanyl is optionally substituted on one or more available carbons with 1-4 R^X; and wherein if the bridged bicyclic

CLAIMS

We claim:


1. A compound of Formula (I):



Formula (I)

or a pharmaceutically acceptable salt, ~~solvent, hydrate, tautomer, N-oxide, or stereoisomer~~ thereof, wherein:

D is ~~a bridged bicyclic cycloalkyl, bridged bicyclic heterocyclyl, or cubanyl, wherein each bridged bicyclic cycloalkyl, bridged bicyclic heterocyclyl, or cubanyl is optionally substituted on one or more available carbons with 1-4 R^x; and wherein if the bridged bicyclic heterocyclyl contains a substitutable nitrogen moiety, the substitutable nitrogen may be optionally substituted by R^{N1}; selected from the group consisting of:~~



L¹ is a bond, C₁-C₆ alkylene, 2-7 membered heteroalkylene, -NR^{N2}-, or -O-, wherein C₁-C₆ alkylene or 2-7 membered heteroalkylene is optionally substituted with 1-5 R^{L1};

L² is a bond, C₁-C₆ alkylene, or 2-7 membered heteroalkylene, wherein C₁-C₆ alkylene or 2-7 membered heteroalkylene is optionally substituted with 1-5 R^{L2};

R¹ is hydrogen or C₁-C₆ alkyl;

R² is hydrogen or C₁-C₆ alkyl;

W is a 8-10 membered, partially unsaturated, fused bicyclic ring moiety comprising a 5-6 membered heterocyclyl fused to a phenyl or 5-6-membered heteroaryl; wherein the heterocyclyl may be optionally substituted on one or more available saturated carbons with 1-4 R^{W1}; and wherein the phenyl or heteroaryl may optionally be substituted on one or more available unsaturated carbons with 1-4 R^{W2}; and wherein if the heterocyclyl contains a

substitutable nitrogen moiety, the substitutable nitrogen may optionally be substituted with R^{N3} ; and wherein W is attached to L^2 through an available saturated carbon or nitrogen atom within the heterocyclyl;

A is phenyl or 5-6-membered heteroaryl, wherein phenyl or 5-6-membered heteroaryl is optionally substituted on one or more available carbons with 1-5 R^Y ; and wherein if the 5-6-membered heteroaryl contains a substitutable nitrogen moiety, the substitutable nitrogen may be optionally substituted by R^{N4} ;

each R^{L1} is independently selected from the group consisting of hydrogen, C_1 - C_6 alkyl, hydroxy- C_1 - C_6 alkyl, halo- C_1 - C_6 alkyl, amino- C_1 - C_6 alkyl, cyano- C_1 - C_6 alkyl, oxo, halo, cyano, $-OR^A$, $-NR^B R^C$, $-NR^B C(O)R^D$, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OH$, $-C(O)OR^D$, $-SR^E$, $-S(O)R^D$, and $-S(O)_2 R^D$;

each R^{L2} is independently selected from the group consisting of hydrogen, C_1 - C_6 alkyl, hydroxy- C_1 - C_6 alkyl, halo- C_1 - C_6 alkyl, amino- C_1 - C_6 alkyl, cyano- C_1 - C_6 alkyl, oxo, halo, cyano, $-OR^A$, $-NR^B R^C$, $-NR^B C(O)R^D$, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OH$, $-C(O)OR^D$, $-SR^E$, $-S(O)R^D$, and $-S(O)_2 R^D$;

R^{N1} is selected from the group consisting of hydrogen, C_1 - C_6 alkyl, hydroxy- C_2 - C_6 alkyl, halo- C_2 - C_6 alkyl, amino- C_2 - C_6 alkyl, cyano- C_2 - C_6 alkyl, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OR^D$, and $-S(O)_2 R^D$;

R^{N2} is selected from the group consisting of hydrogen, C_1 - C_6 alkyl, hydroxy- C_2 - C_6 alkyl, halo- C_2 - C_6 alkyl, amino- C_2 - C_6 alkyl, cyano- C_2 - C_6 alkyl, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OR^D$, and $-S(O)_2 R^D$;

R^{N3} is selected from the group consisting of hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkenyl, $-C(O)-C_1$ - C_6 alkyl, $-C(O)-C_1$ - C_6 cycloalkyl, C_1 - C_6 alkyl- CO_2H , C_1 - C_6 alkyl- CO_2 - C_1 - C_6 alkyl, $-C(O)-C_1$ - C_3 alkyl-O- C_1 - C_3 alkyl-O- C_1 - C_3 alkyl, $-C(O)$ -phenyl, $-C(O)$ -heteroaryl, $-C(O)$ -heterocyclyl, $-S-C_1$ - C_6 alkyl, $-S(O)_2-C_1$ - C_6 alkyl, $-S(O)_2$ -phenyl, $-S(O)_2$ -heteroaryl, $-C(O)NR^B R^C$ and $-C(O)OR^D$;

wherein C_1 - C_6 alkyl, C_1 - C_6 alkenyl, $C(O)-C_1$ - C_6 alkyl, $-C(O)-C_1$ - C_6 cycloalkyl, C_1 - C_6 alkyl- CO_2H , C_1 - C_6 alkyl- CO_2 - C_1 - C_6 alkyl, $-C(O)$ -heterocyclyl, $-S-C_1$ - C_6 alkyl and $-S(O)_2-C_1$ - C_6 alkyl may optionally be substituted by one or more substituents each independently selected from the group consisting of fluoro, hydroxyl, C_1 - C_6 alkoxy, C_1 - C_6 alkyl (optionally substituted by one, two or three fluorine atoms) and $S(O)_w C_{1-6}$ alkyl (wherein w is 0, 1 or 2); and

wherein $-C(O)$ -phenyl, $-C(O)$ -heteroaryl, $-S(O)_2$ -phenyl and $-S(O)_2$ -heteroaryl may optionally be substituted by one or more substituents each independently

selected from the group consisting of halogen, hydroxyl, C₁-C₆ alkyl (optionally substituted by one, two or three fluorine atoms), C₁-C₆ alkoxy (optionally substituted by one, two or three fluorine atoms), S(O₂)NR^BR^C and SO₂F;

R^{N4} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl, halo-C₂-C₆ alkyl, amino-C₂-C₆ alkyl, cyano-C₂-C₆ alkyl, -C(O)NR^BR^C, -C(O)R^D, -C(O)OR^D, and -S(O)₂R^D;

each R^{W1} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl (optionally substituted by -CO₂H), hydroxy-C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl-O-, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, oxo, halo, cyano, -OR^A, -NR^BR^C, -NR^BR^{CC}, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -SR^E, -S(O)R^D, and -S(O)₂R^D;

each R^{W2} is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl-O-, halo-C₁-C₆ alkyl, halo-C₁-C₆ alkoxy, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, halo, cyano, -OR^A, -NR^BR^C, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -S(R^F)_m, -S(O)R^D, and -S(O)₂R^D; or

2 R^{W2} groups on adjacent atoms, together with the atoms to which they are attached, form a 3-7-membered fused cycloalkyl, 3-7-membered fused heterocyclyl, fused aryl, or 5-6 membered fused heteroaryl, each of which is optionally substituted with 1-5 R^X;

each R^X is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, oxo, halo, cyano, -OR^A, -NR^BR^C, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -SR^E, -S(O)R^D, and -S(O)₂R^D;

each R^Y is independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, halo-C₁-C₆ alkoxy, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, halo, cyano, -OR^A, -NR^BR^C, -NR^BC(O)R^D, -C(O)NR^BR^C, -C(O)R^D, -C(O)OH, -C(O)OR^D, -S(R^F)_m, -S(O)R^D, -S(O)₂R^D, and G¹; or

2 R^Y groups on adjacent atoms, together with the atoms to which they are attached form a 3-7-membered fused cycloalkyl, 3-7-membered fused heterocyclyl, fused aryl, or 5-6 membered fused heteroaryl, each of which is optionally substituted with 1-5 R^X;

each G¹ is independently 3-7-membered cycloalkyl, 3-7-membered heterocyclyl, aryl, or 5-6-membered heteroaryl, wherein each 3-7-membered cycloalkyl, 3-7-membered heterocyclyl, aryl, or 5-6-membered heteroaryl is optionally substituted with 1-3 R^Z;

each R^Z is independently selected from the group consisting of C_1 - C_6 alkyl, hydroxy- C_1 - C_6 alkyl, halo- C_1 - C_6 alkyl, halo, cyano, $-OR^A$, $-NR^B R^C$, $-NR^B C(O)R^D$, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OH$, $-C(O)OR^D$, and $-S(O)_2 R^D$;

R^A is, at each occurrence, independently hydrogen, C_1 - C_6 alkyl, halo- C_1 - C_6 alkyl, $-C(O)NR^B R^C$, $-C(O)R^D$, or $-C(O)OR^D$;

each of R^B and R^C is independently hydrogen or C_1 - C_6 alkyl; or

R^B and R^C together with the atom to which they are attached form a 3-7-membered heterocyclyl ring optionally substituted with 1-3 R^Z ;

each R^{CC} is independently selected from the group consisting of C_1 - C_6 alkyl-OH, C_1 - C_6 alkyl-CO₂H and C_1 - C_6 alkyl-CO₂- C_1 - C_6 alkyl;


each R^D is independently C_1 - C_6 alkyl or halo- C_1 - C_6 alkyl;

each R^E is independently hydrogen, C_1 - C_6 alkyl, or halo- C_1 - C_6 alkyl;



each R^F is independently hydrogen, C_1 - C_6 alkyl, or halo; and

m is 1 when R^F is hydrogen or C_1 - C_6 alkyl, 3 when R^F is C_1 - C_6 alkyl, or 5 when R^F is halo.


~~2. The compound of claim 1, wherein D is a bridged bicyclic cycloalkyl or heterocyclyl (e.g., bridged bicyclic 5-8 membered cycloalkyl or heterocyclyl) optionally substituted with 1-4 R^X .~~





42. The compound of ~~any one of claims 1-3~~ claim 1, wherein D is



53. The compound of ~~any one of claims 1-4~~ claim 1 or 2, wherein D is



64. The compound of ~~any one of claims 1-4~~ claim 1 or 2, wherein D is



75. The compound of claim 64, wherein each R^x is independently selected from the group consisting of oxo, -OH, -C(O)OH, -C(O)OR^D, halo, and hydroxy-C₁-C₆ alkyl.


86. The compound of any one of claims 1-75, wherein L¹ is a bond, 2-7 membered heteroalkylene, -NR^{N2}-, or -O-, wherein 2-7 membered heteroalkylene is optionally substituted by 1-5 R^{L1}.

97. The compound of any one of claims 1-86, wherein L¹ is selected from a bond, CH₂O-*, CH₂OCH₂-, -NCH₃-, -NH-, or -O-, wherein “-*” indicates the attachment point to A.

108. The compound of any one of claims 1-97, wherein R¹ is hydrogen or -CH₃.

~~1419~~. The compound of any one of claims 1-~~1408~~, wherein R^2 is hydrogen or $-CH_3$.

~~14210~~. The compound of any one of claims 1-~~1419~~, wherein A is selected from the group consisting of:



~~14311~~. The compound of any one of claims 1-~~14210~~, wherein each R^Y is independently selected from the group consisting of hydrogen, chloro, fluoro, $-CHF_2$, $-CF_3$, $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, $-OCH_3$, $-OCHF_2$, $-OCF_3$, $-OCH_2CF_3$, $-OCH(CH_3)_2$, and $-CN$, or 2 R^Y on adjacent carbons, together with the atoms to which they are attached, form a 1,3-dioxolanyl ring, which is optionally substituted with 1-2 R^X .


~~14412~~. The compound of claim ~~14311~~, wherein each R^X is independently fluoro.

~~14513~~. The compound of any one of claims 1-~~14412~~, wherein L^2 is a bond or C_1 - C_6 alkylene, wherein C_1 - C_6 alkylene is optionally substituted by 1-5 R^{L2} .

~~14614~~. The compound of any one of claims 1-~~14513~~, wherein L^2 is selected from a bond or CH_2- *, wherein “-*” indicates the attachment point to W.

~~14715~~. The compound of any one of claims 1-~~14614~~, wherein L^2 is a bond.

~~14816~~. The compound of any one of claims 1-~~14715~~, wherein W is represented by Formula (W-a):



Formula (W-a)

wherein:

T^1 is nitrogen or $C(R^{W2})$;

T^2 is nitrogen or $C(R^{W2})$;

T^3 is nitrogen or $C(R^{W2})$;

T^4 is nitrogen or $C(R^{W2})$;

wherein no more than two of T^1 , T^2 , T^3 , and T^4 may be nitrogen;

U^1 is selected from the group consisting of a bond, $-O-$, $-NR^{N3}-$, and $-S(O)_w-$ (wherein w is 0, 1, or 2);

V^1 is selected from the group consisting of $+O-^{\#}$, $+C(R^{V11}R^{V12})-^{\#}$, $+C(R^{V11}R^{V12})-C(O)-^{\#}$, $+C(R^{V11}R^{V12})-C(R^{V13}R^{V14})-^{\#}$, $+C(R^{V15}R^{V16})-O-^{\#}$, $+C(R^{V15}R^{V16})-NR^{N3}-^{\#}$, $+C(O)-NR^{N3}-^{\#}$, $+NR^{N3}-^{\#}$, $+O-C(R^{V15}R^{V16})-^{\#}$, $+NR^{N3}-C(R^{V15}R^{V16})-^{\#}$, $+NR^{N3}-C(O)-^{\#}$, $+C(O)-O-^{\#}$, $+O-C(O)-^{\#}$, $+C(R^{V15}R^{V16})-S(O)_w-^{\#}$, and $+S(O)_w-C(R^{V15}R^{V16})-^{\#}$ (wherein w is 0, 1, or 2); wherein the “+” and “-” indicate the attachment points of V^1 as indicated in Formula (W-a);

wherein if V^1 is $+O-^{\#}$, $+NR^{N3}-^{\#}$, or $+C(R^{V11}R^{V12})-^{\#}$; U^1 is not a bond;


R^{V11} and R^{V12} are each independently selected from the group consisting of hydrogen, C_1 - C_6 alkyl, hydroxy- C_1 - C_6 alkyl, halo- C_1 - C_6 alkyl, amino- C_1 - C_6 alkyl, cyano- C_1 - C_6 alkyl, halo, cyano, $-OR^A$, $-NR^B R^C$, $-NR^B R^{CC}$, $-NR^B C(O)R^D$, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OH$, $-C(O)OR^D$, $-SR^E$, $-S(O)R^D$, and $-S(O)_2R^D$;

R^{V13} and R^{V14} are each independently selected from the group consisting of hydrogen, C_1 - C_6 alkyl, hydroxy- C_1 - C_6 alkyl, halo- C_1 - C_6 alkyl, amino- C_1 - C_6 alkyl, cyano- C_1 - C_6 alkyl, halo, cyano, $-OR^A$, $-NR^B R^C$, $-NR^B R^{CC}$, $-NR^B C(O)R^D$, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OH$, $-C(O)OR^D$, $-SR^E$, $-S(O)R^D$, and $-S(O)_2R^D$;


R^{V15} and R^{V16} are each independently selected from the group consisting of hydrogen, C_1 - C_6 alkyl, hydroxy- C_2 - C_6 alkyl, halo- C_2 - C_6 alkyl, amino- C_2 - C_6 alkyl, cyano- C_2 - C_6 alkyl, $-C(O)NR^B R^C$, $-C(O)R^D$, $-C(O)OH$, and $-C(O)OR^D$; and

R^{W1} is selected from the group consisting of hydrogen and C_1 - C_6 alkyl.


~~19~~17. The compound of claim ~~18~~16, wherein W is represented by Formula (W-a-1), Formula (W-a-2), Formula (W-a-3), Formula (W-a-4), or Formula (W-a-5):




Formula (W-a-1),




Formula (W-a-2),



Formula (W-a-3),




Formula (W-a-4), or



Formula (W-a-5).

~~20~~18. The compound of claim ~~19~~17, wherein W is represented by Formula (W-a-1):



Formula (W-a-1).

2419. The compound of any one of claims 48-2016-18, wherein U^1 is selected from the group consisting of a bond, $-O-$, and $-NR^{N3}-$; and V^1 is selected from the group consisting of $+O-^{\#}$, $+C(R^{V11}R^{V12})-^{\#}$, $+C(R^{V11}R^{V12})-C(R^{V13}R^{V14})-^{\#}$, $+C(R^{V15}R^{V16})-O-^{\#}$, $+C(R^{V11}R^{V12})-C(O)-^{\#}$, $+O-C(R^{V15}R^{V16})-^{\#}$, $+C(R^{V15}R^{V16})-NR^{N3}-^{\#}$, and $+C(O)-NR^{N3}-^{\#}$; wherein “+” and “-” indicate the attachment points of V^1 as indicated in Formula (W-a); and wherein if V^1 is $+O-^{\#}$ or $+C(R^{V11}R^{V12})-^{\#}$, U^1 is not a bond.

2220. The compound of any one of claims 48-2416-19, wherein each of R^{V11} , R^{V12} , R^{V13} , and R^{V14} is independently selected from the group consisting of hydrogen, halo, C_1 - C_3 alkyl, cyano, $-OR^A$, $-NR^BR^C$, and $-NR^BR^{CC}$.

2321. The compound of any one of claims 48-2216-20, wherein each of R^{V11} , R^{V12} , R^{V13} , and R^{V14} is independently selected from the group consisting of hydrogen, hydroxyl and $-NR^BR^{CC}$.

2422. The compound of any one of claims 48-2316-21, wherein each of R^{V15} and R^{V16} is independently selected from the group consisting of hydrogen and C_1 - C_3 alkyl.


2523. The compound of any one of claims 48-2416-22, wherein each of R^{V15} and R^{V16} is hydrogen.

2624. The compound of any one of claims 48-2516-23, wherein U^1 is selected from the group consisting of a bond, $-O-$, $-NH-$ and $-NCH_3-$; and V^1 is selected from the group consisting of $+O-^{\#}$, $+CH_2-^{\#}$, $+CH_2-CH_2-^{\#}$, $+CH_2-C(O)-^{\#}$, $+CH_2-O-^{\#}$, $+O-CH_2-^{\#}$, $+CH_2-NH-^{\#}$, $+CH_2-NCH_3-^{\#}$, $+C(O)-NH-^{\#}$, and $+C(O)-NCH_3-^{\#}$, and wherein “+” and “-” indicate the attachment points of V^1 as indicated in Formula (W-a).


2725. The compound of any one of claims ~~18-26~~16-24, wherein R^{W1} is selected from the group consisting of hydrogen, hydroxyl, $-\text{CH}_3$, $-\text{NH}(\text{CH}_2)_2\text{OH}$, $-\text{NH}(\text{CH}_2)_2\text{CO}_2\text{H}$, and $-\text{NH}(\text{CH}_2)_2\text{CO}_2\text{CH}_3$.

2826. The compound of any one of claims ~~18-27~~16-25, wherein W is a benzo[d][1,3]dioxole, 3,4-dihydro-2H-benzo[b][1,4]oxazine, chromane, chroman-4-one, 2H-benzo[b][1,4]oxazin-3(4H)-one, 2,3-dihydrobenzo[b][1,4]dioxine, indoline, or 2,3-dihydrobenzofuran moiety; wherein each of which is attached to L^2 through a saturated carbon atom, and wherein each of which is optionally substituted on one or more available unsaturated carbons with 1-4 R^{W2} , and wherein each R^{W2} is independently selected from the group consisting of C_1 - C_6 alkyl, halo- C_1 - C_6 alkyl, halo, oxo, cyano, and $-\text{OR}^A$.

2927. The compound of any one of claims ~~18-28~~16-26, wherein W is selected from the group consisting of:



3028. The compound of any one of claims 1-~~18~~15, wherein W is represented by Formula (W-b):



Formula (W-b)

wherein:

T^5 is nitrogen or $\text{C}(\text{R}^{W2})$;

T^6 is nitrogen or $\text{C}(\text{R}^{W2})$;

T⁷ is nitrogen or C(R^{W2});

T⁸ is nitrogen or C(R^{W2});

wherein no more than two of T⁵, T⁶, T⁷, and T⁸ may be nitrogen;

V² is selected from the group consisting of *C(R^{V21}R^{V22})-#, *C(R^{V21}R^{V22})-C(R^{V23}R^{V24})-#, *C(R^{V21}R^{V22})-C(R^{V23}R^{V24})-C(R^{V23}R^{V24})-#, *C(R^{V21}R^{V22})-C(R^{V21}R^{V22})-O-#, *C(R^{V21}R^{V22})-C(R^{V21}R^{V22})-NR^{N3}-#, *C(R^{V21}R^{V22})-NR^{N3}-#, *C(O)-C(R^{V23}R^{V24})-#, *C(O)-C(R^{V23}R^{V24})-C(R^{V23}R^{V24})-#, *C(O)-NR^{N3}-#, and *C(O)-O-#, wherein “*“ and “-#” indicate the attachment points of V² as indicated in Formula (W-b);

U² is selected from the group consisting of a bond, *C(O)-+, and *C(R^{U21}R^{U22})-+, wherein “*“ and “-+” indicate the attachment points of U² as indicated in Formula (W-b);


wherein if V² is *C(R^{V21}R^{V22})-#, U² is not a bond;

R^{U21} and R^{U22} are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl, halo-C₂-C₆ alkyl, amino-C₂-C₆ alkyl, cyano-C₂-C₆ alkyl, -C(O)NR^{BR^C}, -C(O)R^D, -C(O)OH, -C(O)OR^D, C₁-C₆ alkyl-C(O)OH, and C₁-C₆ alkyl-C(O)OR^D;


R^{V21} and R^{V22} are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₂-C₆ alkyl, halo-C₂-C₆ alkyl, amino-C₂-C₆ alkyl, cyano-C₂-C₆ alkyl, -C(O)NR^{BR^C}, -C(O)R^D, -C(O)OH, and -C(O)OR^D; and

R^{V23} and R^{V24} are each independently selected from the group consisting of hydrogen, C₁-C₆ alkyl, hydroxy-C₁-C₆ alkyl, halo-C₁-C₆ alkyl, amino-C₁-C₆ alkyl, cyano-C₁-C₆ alkyl, halo, cyano, -OR^A, -NR^{BR^C}, -NR^{BC}(O)R^D, -C(O)NR^{BR^C}, -C(O)R^D, -C(O)OH, -C(O)OR^D, -SR^E, -S(O)R^D, and -S(O)₂R^D.


~~3429~~. The compound of claim ~~3028~~, wherein W is represented by Formula (W-b-1), Formula (W-b-2), Formula (W-b-3), Formula (W-b-4), or Formula (W-b-5):




Formula (W-b-1),




Formula (W-b-2),



Formula (W-b-3),




Formula (W-b-4), or



Formula (W-b-5).

3230. The compound of claim 3128, wherein W is represented by Formula (W-b-1):



Formula (W-b-1).

3331. The compound of any one of claims 30-3228-30, wherein V^2 is selected from the group consisting of $^*C(R^{V21}R^{V22})-^{\#}$, $^*C(R^{V21}R^{V22})-C(R^{V23}R^{V24})-^{\#}$, $^*C(O)-C(R^{V23}R^{V24})-^{\#}$, and $^*C(R^{V21}R^{V22})-C(R^{V23}R^{V24})-C(R^{V23}R^{V24})-^{\#}$; wherein “ * ” and “ $^{\#}$ ” indicate the attachment points of V^2 as indicated in Formula (W-b).

3432. The compound of any one of claims 30-3328-31, wherein each of R^{V21} and R^{V22} is independently selected from the group consisting of hydrogen and C_1 - C_3 alkyl.

3533. The compound of any one of claims 30-3428-32, wherein each of R^{V21} and R^{V22} is hydrogen.


~~36~~34. The compound of any one of claims ~~30-35~~28-33, wherein each of R^{V23} and R^{V24} is independently selected from the group consisting of hydrogen, halo, C_1 - C_3 alkyl, cyano, $-OR^A$, and $-NR^B R^C$.


~~37~~35. The compound of any one of claims ~~30-36~~28-34, wherein each of R^{V23} and R^{V24} is hydrogen.

~~38~~36. The compound of any one of claims ~~30-37~~28-35, wherein U^2 is selected from the group consisting of a bond, $^*-C(O)^-+$, $^*-CH_2^+$, and $^*-CH(CH_2CO_2H)^-+$, wherein “ * ” and “ $^-+$ ” indicate the attachment points of U^2 as indicated in Formula (W-b); and V^2 is selected from the group consisting of $^*-CH_2\#$, $^*-CH_2-CH_2\#$, $^*-C(O)-CH_2\#$, $^*-C(O)-NH\#$, $^*-CH_2-NH\#$, and $^*-CH_2-CH_2-CH_2\#$; wherein “ * ” and “ $\#$ ” indicate the attachment points of V^2 as indicated in Formula (W-b).

~~39~~37. The compound of any one of claims ~~30-38~~28-36, wherein W is an indoline, indolin-2-one, isoindoline, isoindolin-1-one, 1,2,3,4-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, quinazoline-2,4(1*H*,3*H*)-dione, or 2,3-dihydroquinazolin-4(1*H*)-one moiety; wherein each of which is attached to L^2 through a nitrogen atom, and wherein each of which is optionally substituted on one or more available unsaturated carbon atoms with 1-4 R^{W2} , and wherein each R^{W2} is independently selected from the group consisting of C_1 - C_6 alkyl, halo- C_1 - C_6 alkyl, hydroxy- C_2 - C_6 alkyl-O-, halo, cyano, and $-OR^A$.


~~40~~38. The compound of any one of claims ~~30-39~~28-37, wherein W is selected from the group consisting of:





and ; wherein R^{N3} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, and hydroxy-C₂-C₆ alkyl.

~~4439~~. The compound of any one of claims 1-~~4038~~, wherein the compound of Formula (I) is a compound of Formula (I-a):



Formula (I-a)

or a pharmaceutically acceptable salt, ~~solvate, hydrate, tautomer, N oxide, or stereoisomer~~ thereof, wherein:

D is bicyclo[1.1.1]pentanyl or bicyclo[2.2.2]octanyl, each of which is optionally substituted with 1-4 R^X groups;

L^1 is selected from the group consisting of a bond, CH_2O^* , $CH_2OCH_2^*$, $-NCH_3-$, $-NH-$, and $-O-$, wherein “ $-^*$ ” indicates the attachment point to A;

L^2 is a bond;

R^1 is selected from the group consisting of hydrogen and $-CH_3$;

R^2 is selected from the group consisting of hydrogen and $-CH_3$;

A is phenyl, pyrazine or pyridyl, each of which is optionally substituted with 1-5 R^Y groups;

W is a benzo[d][1,3]dioxole, 3,4-dihydro-2*H*-benzo[b][1,4]oxazine, chromane, 2*H*-benzo[b][1,4]oxazin-3(4*H*)-one, 2,3-dihydrobenzo[b][1,4]dioxine, indoline, or 2,3-dihydrobenzofuran moiety; wherein each of which is attached to L^2 through a saturated carbon atom, and wherein each of which is optionally substituted on one or more available unsaturated carbon atoms with 1-4 R^{W2} groups; and wherein 3,4-dihydro-2*H*-benzo[b][1,4]oxazine, 2*H*-benzo[b][1,4]oxazin-3(4*H*)-one, and indoline may be optionally substituted on an available nitrogen atom with hydrogen or CH_3 ;

each R^{W2} is independently selected from the group consisting of hydrogen, chloro, fluoro, $-CHF_2$, $-CF_3$, $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, $-OCH_3$, $-OCHF_2$, $-OCF_3$, $-OCH_2CF_3$, $-OCH(CH_3)_2$, and $-CN$; or


2 R^{W2} groups on adjacent carbons, together with the atoms to which they are attached form a 1,3-dioxolanyl ring, which is optionally substituted with 1-2 fluorine atoms;

each R^X is independently fluoro, oxo, $-OH$, $-OCH_3$, $-C(O)OH$, or $-C(O)OCH_3$; and

each R^Y is independently chloro, fluoro, $-CHF_2$, $-CF_3$, $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, $-OCH_3$, $-OCHF_2$, $-OCF_3$, $-OCH_2CF_3$, $-OCH(CH_3)_2$, or $-CN$; or

2 R^Y groups on adjacent atoms, together with the atoms to which they are attached form a 1,3-dioxolanyl ring, which is optionally substituted with 1-2 fluorine atoms.

~~4240.~~ The compound of any one of claims 1-~~4038~~, wherein the compound of Formula (I) is a compound of Formula (I-b):



Formula (I-b)

or a pharmaceutically acceptable salt, ~~solvate, hydrate, tautomer, N-oxide, or stereoisomer~~ thereof, wherein:

D is bicyclo[1.1.1]pentanyl or bicyclo[2.2.2]octanyl, each of which is optionally substituted with 1-4 R^X groups;

L^1 is selected from the group consisting of a bond, CH_2O^* , $CH_2OCH_2^*$, $-NCH_3-$, $-NH-$, and $-O-$, wherein “ * ” indicates the attachment point to A;

L^2 is CH_2^* , wherein “ * ” indicates the attachment point to W;

R^1 is selected from the group consisting of hydrogen and $-CH_3$;

R^2 is selected from the group consisting of hydrogen and $-CH_3$;

A is phenyl or pyridyl, each of which is optionally substituted with 1-5 R^Y groups;

W is an indoline moiety; wherein indoline is attached to L^2 through a nitrogen atom, and wherein indoline is optionally substituted on one or more available unsaturated carbon atoms with 1-4 R^{W2} groups;

each R^{W2} is independently selected from the group consisting of hydrogen, chloro, fluoro, $-CHF_2$, $-CF_3$, $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, $-OCH_3$, $-OCHF_2$, $-OCF_3$, $-OCH_2CF_3$, $-OCH(CH_3)_2$, and $-CN$; or


2 R^{W2} groups on adjacent carbons, together with the atoms to which they are attached form a 1,3-dioxolanyl ring, which is optionally substituted with 1-2 fluorine atoms;

each R^X is independently fluoro, oxo, $-OH$, $-OCH_3$, $-C(O)OH$, or $-C(O)OCH_3$; and


each R^Y is independently chloro, fluoro, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$, $-\text{OCH}_3$, $-\text{OCHF}_2$, $-\text{OCF}_3$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCH}(\text{CH}_3)_2$, or $-\text{CN}$; or

2 R^Y groups on adjacent atoms, together with the atoms to which they are attached form a 1,3-dioxolanyl ring, which is optionally substituted with 1-2 fluorine atoms.


4341. The compound of any one of claims 1-4240, wherein the compound of Formula (I) is a compound of Formula (I-c-1), Formula (I-c-2), Formula (I-c-3), Formula (I-c-4), Formula (I-c-5), Formula (I-c-6), Formula (I-c-7), Formula (I-c-8), Formula (I-c-9) or Formula (I-c-10):




Formula (I-c-1)




Formula (I-c-2)




Formula (I-c-3)




Formula (I-c-4)




Formula (I-c-5)




Formula (I-c-6)




Formula (I-c-7)



Formula (I-c-8)




Formula (I-c-9)




Formula (I-c-10)

or a pharmaceutically acceptable salt, ~~solvate, hydrate, tautomer, N-oxide, or stereoisomer~~ thereof.


4442. The compound of any one of claims 1-4240, wherein the compound of Formula (I) is a compound of Formula (I-d-1), Formula (I-d-2), Formula (I-d-3), Formula (I-d-4), Formula (I-d-5), Formula (I-d-6), Formula (I-d-7), Formula (I-d-8), Formula (I-d-9) or Formula (I-d-10):




Formula (I-d-1)




Formula (I-d-2)




Formula (I-d-3)




Formula (I-d-4)




Formula (I-d-5)




Formula (I-d-6)




Formula (I-d-7)



Formula (I-d-8)











Formula (I-d-9)



















Formula (I-d-10)











or a pharmaceutically acceptable salt, ~~solvate, hydrate, tautomer, N-oxide, or stereoisomer~~ thereof.









4543. A compound, ~~wherein the compound is~~ selected from ~~any compound set forth in Table 1 or a pharmaceutically acceptable salt, solvate, hydrate, tautomer, N-oxide, or stereoisomer thereof~~ the group consisting of:


























































































or a pharmaceutically acceptable salt thereof.

46.—~~A pharmaceutically acceptable composition comprising a compound of any one of claims 1-45 and a pharmaceutically acceptable carrier.~~

47.—~~Use of a composition in the manufacture of a medicament for treating a neurodegenerative disease, a leukodystrophy, a cancer, an inflammatory disease, an autoimmune disease, a viral infection, a skin disease, a fibrotic disease, a hemoglobin disease, a kidney disease, a hearing loss condition, an ocular disease, a musculoskeletal disease, a metabolic disease, or a mitochondrial disease in a subject, wherein the composition comprises a compound of any one of claims 1-45.~~

48.—~~The use of claim 47, wherein:~~

~~the neurodegenerative disease comprises a leukodystrophy, a leukoencephalopathy, a hypomyelinating or demyelinating disease, an intellectual disability syndrome, a cognitive impairment, a glial cell dysfunction, a brain injury, vanishing white matter disease, childhood ataxia with CNS hypo-myelination, Alzheimer's disease, amyotrophic lateral sclerosis, Creutzfeldt-Jakob disease, frontotemporal dementia, Gerstmann-Straussler-Scheinker disease, Huntington's disease, dementia, kuru, multiple sclerosis, Parkinson's disease, or a prion disease;~~

~~the cancer comprises pancreatic cancer, breast cancer, multiple myeloma, or a cancer of the secretory cells;~~

~~the inflammatory disease comprises postoperative cognitive dysfunction, arthritis, systemic lupus erythematosus (SLE), myasthenia gravis, diabetes), Guillain-Barre syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, ankylosing spondylitis, psoriasis, Sjogren's syndrome, vasculitis, glomerulonephritis, auto-immune thyroiditis, Behcet's disease, Crohn's disease, ulcerative colitis, bullous pemphigoid, sarcoidosis, ichthyosis, Graves' ophthalmopathy, inflammatory bowel disease, Addison's disease, vitiligo, acne vulgaris, celiac disease, chronic prostatitis, pelvic inflammatory disease, reperfusion injury, sarcoidosis, transplant rejection, interstitial cystitis, atherosclerosis, or atopic dermatitis;~~

~~the musculoskeletal disease comprises muscular dystrophy, multiple sclerosis, amyotrophic lateral sclerosis, primary lateral sclerosis, progressive muscular atrophy, progressive bulbar palsy, pseudobulbar palsy, spinal muscular atrophy, progressive spinobulbar muscular atrophy, spinal cord spasticity, spinal muscle atrophy, myasthenia gravis, neuralgia, fibromyalgia, Machado-Joseph disease, cramp-fasciculation syndrome,~~

Freidrich's ataxia, a muscle wasting disorder), an inclusion body myopathy, motor neuron disease, or paralysis;

the metabolic disease comprises non-alcoholic steatohepatitis (NASH), non-alcoholic fatty liver disease (NAFLD), liver fibrosis, obesity, heart disease, atherosclerosis, arthritis, cystinosis, diabetes, phenylketonuria, proliferative retinopathy, or Kearns-Sayre disease;

the mitochondrial disease is associated with or is a result of mitochondrial dysfunction, one or more mitochondrial protein mutations, or one or more mitochondrial DNA mutations or is a mitochondrial myopathy, selected from the group consisting of Barth syndrome, chronic progressive external ophthalmoplegia (cPEO), Kearns-Sayre syndrome (KSS), Leigh syndrome (*e.g.*, MILS, or maternally inherited Leigh syndrome), mitochondrial DNA depletion syndromes (MDDS, *e.g.*, Alpers syndrome), mitochondrial encephalomyopathy (*e.g.*, mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes (MELAS)), mitochondrial neurogastrointestinal encephalomyopathy (MNGIE), myoclonus epilepsy with ragged red fibers (MERRF), neuropathy, ataxia, retinitis pigmentosa (NARP), Leber's hereditary optic neuropathy (LHON and Pearson syndrome;

the autoimmune disease is selected from the group consisting of Achalasia, Addison's disease, Adult Still's disease, Agammaglobulinemia, Alopecia areata, Amyloidosis, Ankylosing spondylitis, Anti-GBM/Anti-TBM nephritis, Antiphospholipid syndrome, Autoimmune angioedema, Autoimmune dysautonomia, Autoimmune encephalomyelitis, Autoimmune hepatitis, Autoimmune inner ear disease (AIED), Autoimmune myocarditis, Autoimmune oophoritis, Autoimmune orchitis, Autoimmune pancreatitis, Autoimmune retinopathy, Autoimmune urticaria, Axonal & neuronal neuropathy (AMAN), Baló disease, Behcet's disease, Benign mucosal pemphigoid, Bullous pemphigoid, Castleman disease (CD), Celiac disease, Chagas disease, Chronic inflammatory demyelinating polyneuropathy (CIDP), Chronic recurrent multifocal osteomyelitis (CRMO), Churg-Strauss Syndrome (CSS) or Eosinophilic Granulomatosis (EGPA), Cicatricial pemphigoid, Cogan's syndrome, Cold agglutinin disease, Congenital heart block, Cocksackie myocarditis, CREST syndrome, Crohn's disease, Dermatitis herpetiformis, Dermatomyositis, Devic's disease (neuromyelitis optica), Discoid lupus, Dressler's syndrome, Endometriosis, Eosinophilic esophagitis (EoE), Eosinophilic fasciitis, Erythema nodosum, Essential mixed cryoglobulinemia, Evans syndrome, Fibromyalgia, Fibrosing alveolitis, Giant cell arteritis (temporal arteritis), Giant cell myocarditis, Glomerulonephritis, Goodpasture's syndrome, Granulomatosis with Polyangiitis, Graves' disease, Guillain-Barre syndrome, Hashimoto's thyroiditis, Hemolytic anemia, Henoch-Schonlein purpura (HSP), Herpes gestationis or pemphigoid gestationis

(PG), Hidradenitis Suppurativa (HS) (Acne Inversa), Hypogammaglobulinemia, IgA Nephropathy, IgG4 related sclerosing disease, Immune thrombocytopenic purpura (ITP), Inclusion body myositis (IBM), Interstitial cystitis (IC), Juvenile arthritis, Juvenile diabetes (Type 1 diabetes), Juvenile myositis (JM), Kawasaki disease, Lambert Eaton syndrome, Leukocytoclastic vasculitis, Lichen planus, Lichen sclerosus, Ligneous conjunctivitis, Linear IgA disease (LAD), Lupus, Lyme disease chronic, Meniere's disease, Microscopic polyangiitis (MPA), Mixed connective tissue disease (MCTD), Mooren's ulcer, Mucha-Habermann disease, Multifocal Motor Neuropathy (MMN) or MMNCB, Multiple sclerosis, Myasthenia gravis, Myositis, Narcolepsy, Neonatal Lupus, Neuromyelitis optica, Neutropenia, Ocular cicatricial pemphigoid, Optic neuritis, Palindromic rheumatism (PR), PANDAS, Paraneoplastic cerebellar degeneration (PCD), Paroxysmal nocturnal hemoglobinuria (PNH), Parry Romberg syndrome, Pars planitis (peripheral uveitis), Parsonnage Turner syndrome, Pemphigus, Peripheral neuropathy, Perivenous encephalomyelitis, Pernicious anemia (PA), POEMS syndrome, Polyarteritis nodosa, Polyglandular syndrome type I, Polyglandular syndrome type II, Polyglandular syndrome type III, Polymyalgia rheumatica, Polymyositis, Postmyocardial infarction syndrome, Postpericardiotomy syndrome, Primary biliary cirrhosis, Primary sclerosing cholangitis, Progesterone dermatitis, Psoriasis, Psoriatic arthritis, Pure red cell aplasia (PRCA), Pyoderma gangrenosum, Raynaud's phenomenon, Reactive Arthritis, Reflex sympathetic dystrophy, Relapsing polychondritis, Restless legs syndrome (RLS), Retroperitoneal fibrosis, Rheumatic fever, Rheumatoid arthritis, Sarcoidosis, Schmidt syndrome, Scleritis, Scleroderma, Sjögren's syndrome, Sperm & testicular autoimmunity, Stiff person syndrome (SPS), Subacute bacterial endocarditis (SBE), Susac's syndrome, Sympathetic ophthalmia (SO), Takayasu's arteritis, Temporal arteritis/Giant cell arteritis, Thrombocytopenic purpura (TTP), Tolosa-Hunt syndrome (THS), Transverse myelitis, Type 1 diabetes, Ulcerative colitis (UC), Undifferentiated connective tissue disease (UCTD), Uveitis, Vasculitis, Vitiligo, Vogt-Koyanagi-Harada Disease, and Wegener's granulomatosis (or Granulomatosis with Polyangiitis (GPA));

the viral infection is selected from the group consisting of influenza, human immunodeficiency virus (HIV) and herpes;

the skin disease is selected from the group consisting of acne, alopecia areata, basal cell carcinoma, Bowen's disease, congenital erythropoietic porphyria, contact dermatitis, Darier's disease, disseminated superficial actinic porokeratosis, dystrophic epidermolysis bullosa, eczema (atopic eczema), extra-mammary Paget's disease, epidermolysis bullosa

simplex, erythropoietic protoporphyria, fungal infections of nails, Hailey-Hailey disease, herpes simplex, hidradenitis suppurativa, hirsutism, hyperhidrosis, ichthyosis, impetigo, keloids, keratosis pilaris, lichen planus, lichen sclerosus, melanoma, melasma, mucous membrane pemphigoid, pemphigoid, pemphigus vulgaris, pityriasis lichenoides, pityriasis rubra pilaris, plantar warts (verrucae), polymorphic light eruption, psoriasis, plaque psoriasis, pyoderma gangrenosum, rosacea, scabies, scleroderma, shingles, squamous cell carcinoma, sweet's syndrome, urticaria and angioedema and vitiligo;

the fibrotic disease is selected from the group consisting of adhesive capsulitis, arterial stiffness, arthrofibrosis, atrial fibrosis, cardiac fibrosis, cirrhosis, congenital hepatic fibrosis, Crohn's disease, cystic fibrosis, Dupuytren's contracture, endomyocardial fibrosis, glial scar, hepatitis C, hypertrophic cardiomyopathy, hypersensitivity pneumonitis, idiopathic pulmonary fibrosis, idiopathic interstitial pneumonia, interstitial lung disease, keloid, mediastinal fibrosis, myelofibrosis, nephrogenic systemic fibrosis, non-alcoholic fatty liver disease, old myocardial infarction, Peyronie's disease, pneumoconiosis, pneumonitis, progressive massive fibrosis, pulmonary fibrosis, radiation-induced lung injury, retroperitoneal fibrosis, scleroderma/systemic sclerosis, silicosis and ventricular remodeling;

the hemoglobin disease is selected from the group consisting of "dominant" β -thalassemia, acquired (toxic) methemoglobinemia, carboxyhemoglobinemia, congenital Heinz body hemolytic anemia, HbH disease, HbS/ β -thalassemia, HbE/ β -thalassemia, HbSC disease, homozygous α^+ -thalassemia (phenotype of α^0 -thalassemia), Hydrops fetalis with Hb Bart's, sickle cell anemia/disease, sickle cell trait, sickle β -thalassemia disease, α^+ -thalassemia, α^0 -thalassemia, α -Thalassemia associated with myelodysplastic syndromes, α -Thalassemia with mental retardation syndrome (ATR), β^0 -Thalassemia, β^+ -Thalassemia, δ -Thalassemia, γ -Thalassemia, β -Thalassemia major, β -Thalassemia intermedia, $\delta\beta$ -Thalassemia, and $\epsilon\gamma\delta\beta$ -Thalassemia;

the kidney disease is selected from the group consisting of Abderhalden-Kaufmann-Lignac syndrome (Nephropathic Cystinosis), Abdominal Compartment Syndrome, Acetaminophen-induced Nephrotoxicity, Acute Kidney Failure/Acute Kidney Injury, Acute Lobar Nephronia, Acute Phosphate Nephropathy, Acute Tubular Necrosis, Adenine Phosphoribosyltransferase Deficiency, Adenovirus Nephritis, Alagille Syndrome, Alport Syndrome, Amyloidosis, ANCA Vasculitis Related to Endocarditis and Other Infections, Angiomyolipoma, Analgesic Nephropathy, Anorexia Nervosa and Kidney Disease, Angiotensin Antibodies and Focal Segmental Glomerulosclerosis, Antiphospholipid Syndrome, Anti-TNF- α Therapy-related Glomerulonephritis, APOL1 Mutations, Apparent

Mineralocorticoid Excess Syndrome, Aristolochic Acid Nephropathy, Chinese Herbal Nephropathy, Balkan Endemic Nephropathy, Arteriovenous Malformations and Fistulas of the Urologic Tract, Autosomal Dominant Hypocalcemia, Bardet-Biedl Syndrome, Bartter Syndrome, Bath Salts and Acute Kidney Injury, Beer Potomania, Beeturia, β -Thalassemia Renal Disease, Bile Cast Nephropathy, BK Polyoma Virus Nephropathy in the Native Kidney, Bladder Rupture, Bladder Sphincter Dyssynergia, Bladder Tamponade, Border-Crossers' Nephropathy, Bourbon Virus and Acute Kidney Injury, Burnt Sugarcane Harvesting and Acute Renal Dysfunction, Byetta and Renal Failure, C1q Nephropathy, C3 Glomerulopathy, C3 Glomerulopathy with Monoclonal Gammopathy, C4 Glomerulopathy, Calcineurin Inhibitor Nephrotoxicity, Callilepis Laureola Poisoning, Cannabinoid Hyperemesis Acute Renal Failure, Cardiorenal syndrome, Carfilzomib-Induced Renal Injury, CFHR5 nephropathy, Charcot-Marie-Tooth Disease with Glomerulopathy, Chinese Herbal Medicines and Nephrotoxicity, Cherry Concentrate and Acute Kidney Injury, Cholesterol Emboli, Churg-Strauss syndrome, Chyluria, Ciliopathy, Cocaine and the Kidney, Cold Diuresis, Colistin Nephrotoxicity, Collagenofibrotic Glomerulopathy, Collapsing Glomerulopathy, Collapsing Glomerulopathy Related to CMV, Combination Antiretroviral (cART) Related Nephropathy, Congenital Anomalies of the Kidney and Urinary Tract (CAKUT), Congenital Nephrotic Syndrome, Congestive Renal Failure, Conorenal syndrome (Mainzer-Saldino Syndrome or Saldino-Mainzer Disease), Contrast Nephropathy, Copper Sulphate Intoxication, Cortical Necrosis, Crizotinib-related Acute Kidney Injury, Cryocrystalglobulinemia, Cryoglobulinemia, Crystalglobulin-Induced Nephropathy, Crystal-Induced Acute Kidney injury, Crystal-Storing Histiocytosis, Cystic Kidney Disease, Acquired, Cystinuria, Dasatinib-Induced Nephrotic-Range Proteinuria, Dense Deposit Disease (MPGN Type 2), Dent Disease (X-linked Recessive Nephrolithiasis), DHA Crystalline Nephropathy, Dialysis Disequilibrium Syndrome, Diabetes and Diabetic Kidney Disease, Diabetes Insipidus, Dietary Supplements and Renal Failure, Diffuse Mesangial Sclerosis, Diuresis, Djenkol Bean Poisoning (Djenkolism), Down Syndrome and Kidney Disease, Drugs of Abuse and Kidney Disease, Duplicated Ureter, EAST syndrome, Ebola and the Kidney, Ectopic Kidney, Ectopic Ureter, Edema, Swelling, Erdheim-Chester Disease, Fabry's Disease, Familial Hypocalciuric Hypercalcemia, Fanconi Syndrome, Fraser syndrome, Fibronectin Glomerulopathy, Fibrillary Glomerulonephritis and Immunotactoid Glomerulopathy, Fraley syndrome, Fluid Overload, Hypervolemia, Focal Segmental Glomerulosclerosis, Focal Sclerosis, Focal Glomerulosclerosis, Galloway-Mowat syndrome, Giant Cell (Temporal) Arteritis with Kidney Involvement, Gestational Hypertension;

Gitelman Syndrome, Glomerular Diseases, Glomerular Tubular Reflux, Glycosuria, Goodpasture Syndrome, Green Smoothie Cleanse Nephropathy, HANAC Syndrome, Harvoni (Ledipasvir with Sofosbuvir) Induced Renal Injury, Hair Dye Ingestion and Acute Kidney Injury, Hantavirus Infection Podocytopathy, Heat Stress Nephropathy, Hematuria (Blood in Urine), Hemolytic Uremic Syndrome (HUS), Atypical Hemolytic Uremic Syndrome (aHUS), Hemophagocytic Syndrome, Hemorrhagic Cystitis, Hemorrhagic Fever with Renal Syndrome (HFRS, Hantavirus Renal Disease, Korean Hemorrhagic Fever, Epidemic Hemorrhagic Fever, Nephropathis Epidemica), Hemosiderinuria, Hemosiderosis related to Paroxysmal Nocturnal Hemoglobinuria and Hemolytic Anemia, Hepatic Glomerulopathy, Hepatic Veno-Occlusive Disease, Sinusoidal Obstruction Syndrome, Hepatitis C Associated Renal Disease, Hepatocyte Nuclear Factor 1 β Associated Kidney Disease, Hepatorenal Syndrome, Herbal Supplements and Kidney Disease, High Altitude Renal Syndrome, High Blood Pressure and Kidney Disease, HIV Associated Immune Complex Kidney Disease (HIVICK), HIV Associated Nephropathy (HIVAN), HNF1B related Autosomal Dominant Tubulointerstitial Kidney Disease, Horseshoe Kidney (Renal Fusion), Hunner's Ulcer, Hydroxychloroquine-induced Renal Phospholipidosis, Hyperaldosteronism, Hypercalcemia, Hyperkalemia, Hypermagnesemia, Hyponatremia, Hyperoxaluria, Hyperphosphatemia, Hypocalcemia, Hypocomplementemic Urticarial Vasculitic Syndrome, Hypokalemia, Hypokalemia-induced renal dysfunction, Hypokalemic Periodic Paralysis, Hypomagnesemia, Hyponatremia, Hypophosphatemia, Hypophosphatemia in Users of Cannabis, Hypertension, Hypertension, Monogenic, Iced Tea Nephropathy, Ifosfamide Nephrotoxicity, IgA Nephropathy, IgG4 Nephropathy, Immersion Diuresis, Immune Checkpoint Therapy Related Interstitial Nephritis, Infliximab Related Renal Disease, Interstitial Cystitis, Painful Bladder Syndrome (Questionnaire), Interstitial Nephritis, Interstitial Nephritis, Karyomegalic, Ivemark's syndrome, JC Virus Nephropathy, Joubert Syndrome, Ketamine Associated Bladder Dysfunction, Kidney Stones, Nephrolithiasis, Kombucha Tea Toxicity, Lead Nephropathy and Lead Related Nephrotoxicity, Lecithin Cholesterol Acyltransferase Deficiency (LCAT Deficiency), Leptospirosis Renal Disease, Light Chain Deposition Disease, Monoclonal Immunoglobulin Deposition Disease, Light Chain Proximal Tubulopathy, Liddle Syndrome, Lightwood Albright Syndrome, Lipoprotein Glomerulopathy, Lithium Nephrotoxicity, LMX1B Mutations Cause Hereditary FSGS, Loin Pain Hematuria, Lupus, Systemic Lupus Erythematosus, Lupus Kidney Disease, Lupus Nephritis, Lupus Nephritis with Antineutrophil Cytoplasmic Antibody Seropositivity, Lupus Podocytopathy, Lyme Disease Associated Glomerulonephritis, Lysinuric Protein Intolerance, Lysozyme Nephropathy, Malarial

Nephropathy, Malignancy-Associated Renal Disease, Malignant Hypertension, Malakoplakia, McKittrick-Wheelock Syndrome, MDMA (Molly; Ecstasy; 3,4-Methylenedioxymethamphetamine) and Kidney Failure, Meatal Stenosis, Medullary Cystic Kidney Disease, Urolodulin-Associated Nephropathy, Juvenile Hyperuricemic Nephropathy Type 1, Medullary Sponge Kidney, Megaureter, Melamine Toxicity and the Kidney, MELAS Syndrome, Membranoproliferative Glomerulonephritis, Membranous Nephropathy, Membranous-like Glomerulopathy with Masked IgG-Kappa Deposits, MesoAmerican Nephropathy, Metabolic Acidosis, Metabolic Alkalosis, Methotrexate-related Renal Failure, Microscopic Polyangiitis, Milk-alkalai syndrome, Minimal Change Disease, Monoclonal Gammopathy of Renal Significance, Dysproteinemia, Mouthwash Toxicity, MUC1 Nephropathy, Multicystic dysplastic kidney, Multiple Myeloma, Myeloproliferative Neoplasms and Glomerulopathy, Nail-patella Syndrome, NARP Syndrome, Nephrocalcinosis, Nephrogenic Systemic Fibrosis, Nephroptosis (Floating Kidney, Renal Ptois), Nephrotic Syndrome, Neurogenic Bladder, 9/11 and Kidney Disease, Nodular Glomerulosclerosis, Non-Gonococcal Urethritis, Nutcracker syndrome, Oligomeganephronia, Orofaciodigital Syndrome, Orotic Aciduria, Orthostatic Hypotension, Orthostatic Proteinuria, Osmotic Diuresis, Osmotic Nephrosis, Ovarian Hyperstimulation Syndrome, Oxalate Nephropathy, Page Kidney, Papillary Necrosis, Papillorrenal Syndrome (Renal-Coloboma Syndrome, Isolated Renal Hypoplasia), PARN Mutations and Kidney Disease, Parvovirus B19 and the Kidney, The Peritoneal Renal Syndrome, POEMS Syndrome, Posterior Urethral Valve, Podocyte Infolding Glomerulopathy, Post-infectious Glomerulonephritis, Post-streptococcal Glomerulonephritis, Post-infectious Glomerulonephritis, Atypical, Post-Infectious Glomerulonephritis (IgA-Dominant), Mimicking IgA-Nephropathy, Polyarteritis Nodosa, Polycystic Kidney Disease, Posterior Urethral Valves, Post-Obstructive Diuresis, Preeclampsia, Propofol infusion syndrome, Proliferative Glomerulonephritis with Monoclonal IgG Deposits (Nasr Disease), Propolis (Honeybee Resin) Related Renal Failure, Proteinuria (Protein in Urine), Pseudohyperaldosteronism, Pseudohypobicarbonatemia, Pseudohypoparathyroidism, Pulmonary Renal Syndrome, Pyelonephritis (Kidney Infection), Pyonephrosis, Pyridium and Kidney Failure, Radiation Nephropathy, Ranolazine and the Kidney, Refeeding syndrome, Reflux Nephropathy, Rapidly Progressive Glomerulonephritis, Renal Abscess, Peripnephric Abscess, Renal Agenesis, Renal Arcuate Vein Microthrombi-Associated Acute Kidney Injury, Renal Artery Aneurysm, Renal Artery Dissection, Spontaneous, Renal Artery Stenosis, Renal Cell Cancer, Renal Cyst, Renal Hypouricemia with Exercise-induced Acute Renal Failure, Renal Infarction, Renal Osteodystrophy, Renal

~~Tubular Acidosis, Renin Mutations and Autosomal Dominant Tubulointerstitial Kidney Disease, Renin Secreting Tumors (Juxtaglomerular Cell Tumor), Reset Osmostat, Retrocaval Ureter, Retroperitoneal Fibrosis, Rhabdomyolysis, Rhabdomyolysis related to Bariatric Sugery, Rheumatoid Arthritis Associated Renal Disease, Sarcoidosis Renal Disease, Salt Wasting, Renal and Cerebral, Schistosomiasis and Glomerular Disease, Schimke immuno-osseous dysplasia, Scleroderma Renal Crisis, Serpentine Fibula Polycystic Kidney Syndrome, Exner Syndrome, Sickle Cell Nephropathy, Silica Exposure and Chronic Kidney Disease, Sri Lankan Farmers' Kidney Disease, Sjögren's Syndrome and Renal Disease, Synthetic Cannabinoid Use and Acute Kidney Injury, Kidney Disease Following Hematopoietic Cell Transplantation, Kidney Disease Related to Stem Cell Transplantation, TAFRO Syndrome, Tea and Toast Hyponatremia, Tenofovir Induced Nephrotoxicity, Thin Basement Membrane Disease, Benign Familial Hematuria, Thrombotic Microangiopathy Associated with Monoclonal Gammopathy, Trench Nephritis, Trigonitis, Tuberculosis, Genitourinary, Tuberous Sclerosis, Tubular Dysgenesis, Immune Complex Tubulointerstitial Nephritis Due to Autoantibodies to the Proximal Tubule Brush Border, Tumor Lysis Syndrome, Uremia, Uremic Optic Neuropathy, Ureteritis Cystica, Ureterocele, Urethral Caruncle, Urethral Stricture, Urinary Incontinence, Urinary Tract Infection, Urinary Tract Obstruction, Urogenital Fistula, Uromodulin-Associated Kidney Disease, Vancomycin-Associated Cast Nephropathy, Vasomotor Nephropathy, Vesicointestinal Fistula, Vesicoureteral Reflux, VGEF Inhibition and Renal Thrombotic Microangiopathy, Volatile Anesthetics and Acute Kidney Injury, Von Hippel Lindau Disease, Waldenstrom's Macroglobulinemic Glomerulonephritis, Warfarin Related Nephropathy, Wasp Stings and Acute Kidney Injury, Wegener's Granulomatosis, Granulomatosis with Polyangiitis, West Nile Virus and Chronic Kidney Disease, Wunderlich syndrome, Zellweger Syndrome, or Cerebrohepatorenal Syndrome;~~

~~the hearing loss condition is selected from the group consisting of mitochondrial nonsyndromic hearing loss and deafness, hair cell death, age-related hearing loss, noise-induced hearing loss, genetic or inherited hearing loss, hearing loss experienced as a result of ototoxic exposure, hearing loss resulting from disease, and hearing loss resulting from trauma; or~~

~~the ocular disease cataracts, glaucoma, endoplasmic reticulum (ER) stress, autophagy deficiency, age-related macular degeneration (AMD), or diabetic retinopathy.~~

49. — The use of claim 47 or 48, wherein the neurodegenerative disease comprises vanishing white matter disease.

50. — The use of any one of claims 47-49, wherein the composition further comprises a second agent for treating a neurodegenerative disease, a leukodystrophy, a cancer, an inflammatory disease, an autoimmune disease, a viral infection, a skin disease, a fibrotic disease, a hemoglobin disease, a kidney disease, a hearing loss condition, an ocular disease, a musculoskeletal disease, a metabolic disease, a mitochondrial disease, or a disease or disorder associated with impaired function of eIF2B, eIF2 α , or a component of the eIF2 pathway or ISR pathway.

51. — Use of a composition in the manufacture of a medicament for treating a disease related to a modulation of eIF2B activity or levels, eIF2 α activity or levels, or the activity or levels of a component of the eIF2 pathway or the ISR pathway, wherein the composition comprises a compound of any one of claims 1-45.

52. — Use of a composition in combination with an immunotherapeutic agent, in the manufacture of a medicament for treating cancer in a subject, wherein the composition comprises a compound of any one of claims 1-45.

53. — A composition for use in treating a neurodegenerative disease, a leukodystrophy, a cancer, an inflammatory disease, an autoimmune disease, a viral infection, a skin disease, a fibrotic disease, a hemoglobin disease, a kidney disease, a hearing loss condition, an ocular disease, a musculoskeletal disease, a metabolic disease, or a mitochondrial disease in a subject; or

——— for use in treating a disease related to a modulation of eIF2B activity or levels, eIF2 α activity or levels, or the activity or levels of a component of the eIF2 pathway or the ISR pathway; or

——— for use in treating cancer in a subject in combination with an immunotherapeutic agent, wherein the composition comprises a compound of any one of claims 1-45.