

Description**FIELD OF THE INVENTION**

- 5 The invention relates to novel chiral 2,5-disubstituted phospholanes and a method for their preparation. The compounds, when complexed with transition metals, are efficient catalyst precursors for the enantioselective hydrogenation of unsaturated substrates.

BACKGROUND OF THE INVENTION

- 10 The development of novel catalytic systems exhibiting unique reactivity and high enantioselectivity requires the synthesis of chiral ligands for transition metals. Generally, some of the most successful chiral ligands have been chelating phosphines possessing a C_2 symmetry axis. The synthesis of these phosphines in optically pure form often involves tedious synthetic routes that are limited to only one
15 antipode or require a resolution step.

- One synthetic route is through a diol intermediate. S. Masamune et al., Journal of Organic Chemistry, Vol. 54, p. 1755 (1989) teaches use of Baker's yeast for the reduction of 2,5-hexanedione to the corresponding (S,S)-diol, followed by reaction with methanesulfonyl chloride and ring closure with benzylamine to form the optically pure (2R,5R)-2,5-dimethylpyrrolidine. Wilson et al., Synlett, pp. 199-200, April
20 (1990) disclose similar use of a diol intermediate formed via a Baker's yeast reduction of a diketone in the preparation of a 2,5-dimethylphospholane (Compound 10). The phospholane is prepared by reacting the diol with methanesulfonyl chloride followed by phenylphosphine in the presence of potassium hydroxide. However, enzymatic reductions generally provide only one enantiomer of the desired product, and can have limitations such as high substrate specificity, low product yields, or involved isolation procedures.

- 25 In addition, many of the chiral phosphines known in the art have at least two aryl substituents on the phosphorus, rendering that center relatively electron poor. The mechanism of asymmetric induction using these phosphines has been linked to the proper conformational relationship between the phenyl groups on the phosphorus centers.

- More recently, chiral phosphines having relatively electron-rich phosphorus centers have been reported.
30 Brunner et al., Journal of Organometallic Chemistry, Vol. 328, PP. 71-80 (1987) teach 3,4-disubstituted phospholanes derived from tartaric acid having chloro, methoxy, or dimethylamino substituents. These were complexed with manganese and rhodium and used as catalysts in the hydrogenation of alpha-N-acetamidocinnamic acid. Relatively low optical yields of (S)-N-acetylphenylalanine of from 6.6% enantiomeric excess to 16.8% enantiomeric excess were obtained.

- 35 A need exists for transition metal complexes providing high levels of stereochemical control and asymmetric induction in stoichiometric and catalytic transformations. A need also exists for efficient synthetic routes for the preparation of chiral ligands having a high degree of enantiomeric purity for transition metal catalysts.

- 40 It is therefore an object of the present invention to provide novel phospholane compounds as chiral ligands for transition metals.

It is a further object of the present invention to provide transition metal catalysts which provide high levels of stereochemical control of reactions.

It is a further object of the present invention to provide transition metal catalysts which result in high levels of asymmetric induction in hydrogenation reactions.


- 45 It is a further object of the present invention to provide efficient synthetic routes for the preparation of these phospholane compounds.

50

55

SUMMARY OF THE INVENTION

The present invention comprises phospholane compounds represented by the following formula I




wherein:

R is a C₁-C₆ alkyl, trifluoromethyl, phenyl, substituted phenyl, aralkyl, or ring substituted aralkyl; and n is an integer from 1 to 12.

The present invention further comprises transition metal complexes of compounds of formula I.


A second aspect of the present invention further comprises phospholane compounds represented by the following formula II



wherein R and n are as defined above for formula I, and A is CCH₃, CH, N or P.

The present invention further comprises transition metal complexes of compounds of formula II.

A third aspect of the present invention comprises phospholane compounds represented by the following formula III



wherein:

R is an alkyl group of two to six carbon atoms, trifluoromethyl, phenyl, substituted phenyl, aralkyl, or ring substituted aralkyl.

A fourth aspect of the present invention comprises a process for the preparation of compounds of formula I as described above wherein a phenyl substituted phospholane of formula III is reacted with lithium and either 1) a dihalo compound of formula X-(CH₂)_n-X wherein X is halogen, and n is an integer from 1 to 12, or 2) a compound of formula R'O(CH₂)_nOR' wherein n is an integer from 1 to 12, and R'O or OR' is methanesulfonate, trifluoromethanesulfonate or p-toluenesulfonate.


A fifth aspect of the present invention comprises a process for the preparation of compounds of formula II wherein a compound of formula III as defined above is reacted with lithium and a trihalo compound of formula A[(CH₂)_nX]₃ wherein X is halogen, n is defined as above, and A is CCH₃, CH, N or P.

DETAILED DESCRIPTION OF THE INVENTION


The present invention provides chiral phospholane substituted alkanes useful as ligands for transition metals in asymmetric catalysis. The peralkylated nature of these compounds renders the phosphorus center electron rich. Transition metal complexes containing these ligands demonstrate a high level of enantioselective control and asymmetric induction in the catalyzed hydrogenation of unsaturated substrates. It has been found that the close proximity of the chirality to the metal center of the complex results in an increase of the asymmetric induction achieved.

This invention also provides an efficient stereospecific process for the preparation of the chiral phospholanes. The availability of optically active 1,4-diols with a high degree of enantiomeric purity permits preparation of optically active phospholane compounds with a high degree of enantiomeric purity.

The present invention comprises novel phospholane substituted alkane compounds of formulae I and II



I



II

wherein for each of formulae I and II:

R is a C₁-C₆ alkyl, trifluoromethyl, phenyl, substituted phenyl, aralkyl, or ring-substituted aralkyl; and

n is an integer from 1 to 12; and

for formula II, A is CCH₃, CH, N or P.

The present invention further comprises transition metal complexes of these compounds.

Preferred are compounds of formulae I and II wherein n is 1 to 3. Most preferred are those compounds of formula I and II wherein R is methyl and n is 1 to 3.

Examples of such compounds include, but are not limited to, 1,2-bis[(2R,5R)-2,5-dimethylphospholano]ethane; 1,3-bis[(2R,5R)-2,5-dimethylphospholano]propane; tris[[(2S,5S)-2,5-dimethylphospholano)methyl]methane; tris[2-[(2R,5R)-2,5-dimethylphospholano]ethyl]amine; or 1,1,1-tris[2-[(2R,5R)-2,5-dimethylphospholano]ethyl]ethane.

The above phospholane compounds of formulae I and II can be complexed with any of the transition metals of Groups 3 through 12 of the periodic table, plus the lanthanides and actinides. Due to the electron-rich nature of the phospholane compounds they generally coordinate best with the transition metals of Groups 4 through 10. Such complexes are formed by methods known in the art. Preferred transition metal complexes of the present invention are those comprising the above described preferred compounds complexed with rhodium.


The phospholane compounds of formulae I and II of the present invention are useful as transition metal ligands in asymmetric catalysis. The use of these ligands in transition metal catalysts results in a high level of enantioselective and stereochemical control in the catalyzed hydrogenation of unsaturated substrates.

By a high level of enantioselectivity is meant a hydrogenation that yields a product of greater than or equal to about 80%, preferably, greater than or equal to about 90% enantiomeric excess (abbreviated ee).

Enantiomeric excess is defined as the ratio (%R - %S)/(%R + %S), where %R is the percentage of R enantiomer and %S is the percentage of S enantiomer in a sample of optically active compound.

For the purpose of this application, by a "compound with a high degree of enantiomeric purity", or a "compound of high enantiomeric purity" is meant a compound that exhibits optical activity to the extent of greater than or equal to about 90%, preferably, greater than or equal to about 95% enantiomeric excess (abbreviated ee).

A further aspect of the present invention comprises the phospholane compounds of formula III



wherein R is an alkyl group having 2 to 6 carbon atoms, trifluoromethyl, phenyl, substituted phenyl, aralkyl, or ring substituted aralkyl.


Preferred compounds of formula III are those wherein R is an alkyl group of 2 to 6 carbon atoms.

15 The phospholane compounds of formula III of the present invention are useful as intermediates in the preparation of the compounds of formulae I and II.

Another aspect of the present invention comprises processes for the preparation of compounds of formulae I, II and III. The phospholane compounds are prepared by processes of the present invention which provide high yields and high substrate stereo-selectivity. The processes are summarized in reaction

20 Scheme I hereinafter.

SCHEME I



The first step introduces the desired chirality and utilizes a Ru(BINAP) [Ru-(R)-(+)] or (S)-(-)-2,2'-bis-(diphenylphosphino)-1,1'-binaphthyl] catalyst as taught in Noyori et al., J. Amer. Chem. Soc., Vol. 110, p. 629 (1988), which is herein incorporated by reference, for the asymmetric reduction of a β -keto ester to the corresponding β -hydroxy ester. Hydrolysis with a strong base such as KOH provides the free carboxylic acid, which is then subjected to electrochemical Kolbe-coupling to afford a chiral diol. In the Kolbe-coupling reaction a β -hydroxy carboxylic acid of formula $\text{R}^1\text{R}^2\text{C}(\text{OH})\text{CH}_2\text{COOH}$, wherein R^1 and R^2 are each independently hydrogen, lower alkyl, phenyl, substituted phenyl, aralkyl or ring-substituted aralkyl, or R^1 and R^2 are joined together to form a 4-, 5- or 6-membered ring, is dissolved or suspended in a lower alcohol solvent, together with a catalytic amount of a corresponding alkali metal alkoxide. Electrical current is then passed through the solution or suspensions and the chiral diol product isolated by methods known in

the art. The chiral diol is reacted with an alkylsulfonyl chloride, preferably methanesulfonyl chloride, in the presence of a tertiary amine such as triethylamine to form the bis(alkylsulfonate) derivative of the diol. Dilithium phenylphosphide is then added to obtain the chiral 2,5-disubstituted-1-phenylphospholane of formula III.

5 The preparation of phospholane compounds of formulae I and II using the compounds of formula III requires two additional steps. Treatment of a phenylphospholane of formula III with lithium with a clean metallic surface results in selective cleavage of the phenyl group and yields a mixture of 2,5-disubstituted lithium phosphide and phenyllithium. This reaction is conducted in tetrahydrofuran or an equivalent solvent. It is conducted at a temperature range of from about 0 °C to about 40 °C, preferably from about 20 °C to
10 about 25 °C. This reaction is conducted in the absence of oxygen and water under an inert atmosphere at a pressure of about 1 atm. Preferably the inert atmosphere is argon. Agitation is required since it is a heterogeneous reaction with lithium metal. The overall reaction time can range from about 5 to about 30 hours, and typically is from about 10 to about 20 hours.

The resulting mixture is then reacted directly with a compound of formula $R^1O(CH_2)_nOR^1$ wherein OR^1 or R^1O is methanesulfonate, trifluoromethanesulfonate, or p-toluenesulfonate, or with a dihalo alkane of
15 formula $X-(CH_2)_n-X$ wherein X is halogen, preferably chloro or bromo, to obtain the desired chiral chelating bis(phospholanes) of formula I. Reacting the mixture with a compound of formula $A[(CH_2)_nX]_3$ wherein X is halogen, preferably chloro or bromo; A is CCH_3 , CH, N or P; and n is 1 to 12; yields the desired tris-
20 (phospholanes) of formula II. These reactions are conducted at a temperature range of from about -78 °C to about 40 °C, preferably at from about 0 °C to about 25 °C, in tetrahydrofuran solvent. An inert atmosphere is employed, preferably argon or nitrogen at about 1 atm. pressure. The reaction mixture is agitated. The overall reaction time for these reactions is from about 0.5 to about 2 hours, typically from about 0.5 to about 1 hour. The desired product is isolated using methods well known in the art such as distillation, crystallization, evaporation of solvent, filtration, chromatography and the like.

25 The following examples illustrate the present invention but are not intended to limit it in any manner.

GENERAL PROCEDURES

30 All reactions and manipulations were performed in a nitrogen-filled Vacuum Atmospheres Dri-Lab glovebox or using standard Schlenk (inert atmosphere) techniques. Benzene, toluene, diethyl ether (Et_2O), tetrahydrofuran (THF), glyme, hexane, and pentane were distilled from sodium benzo-phenone ketyl under nitrogen. Acetonitrile (CH_3CN) and methylene chloride (CH_2Cl_2) were distilled from CaH_2 . Methanol ($MeOH$) was distilled from $Mg(OMe)_2$.

Melting points were determined using a Mel-Temp apparatus in capillaries sealed under nitrogen and
35 were uncorrected. HPLC analyses were performed using a Hewlett Packard Model HP 1090 LC interfaced to a HP 9000 Series 300 computer workstation. Optical Rotations were obtained using a Perkin Elmer Model Polarimeter. NMR spectra were obtained on Nicolet NT-360 wide-bore (360 MHz 1H , 145 MHz ^{31}P), Nicolet NMC-300 wide-bore (300 MHz 1H , 120.5 MHz ^{31}P , 75.5 Mz ^{13}C) and Nicolet QM-300 narrow-bore (300 MHz 1H) spectrometers. ^{13}C and ^{31}P NMR chemical shifts were positive downfield (and negative upfield) from
40 external Me_4Si and 85% H_3PO_4 , respectively. IR spectra were recorded on a Nicolet 5DXB FT-IR spectrometer. Elemental analyses were performed by Oneida Research Services, Inc., Whitesboro, NY, or Schwarzkopf Microanalytical Laboratory, Inc., Woodside, NY.

The precursor chiral β -hydroxy esters used in the following examples of diol synthesis were prepared as described by Noyori et al., J. Amer. Chem. Soc., 109, 5856 (1987) which is herein incorporated by
45 reference. The asymmetric reduction of β -keto esters to the β -hydroxy esters was conducted using a ruthenium catalyst bearing the chiral phosphine ligand BINAP, (R)-(+)- or (S)-(-)-2,2'-bis(diphenylphosphine)-1,1'-binaphthyl, (both enantiomers commercially available from Strem Chemicals, 7 Mulliken Way, Dexter Industrial Park, P.O. Box 108, Newburyport, MA 01950).

50 EXAMPLE 1

a) Preparation of chiral β -hydroxy acids. A mixture of methyl (3R)-3-hydroxypentanoate (290 g, 2.2 mol) in water (200 mL) and ethanol (200 mL) was cooled to 0 °C. To this cold solution was added a solution of
55 KOH (185 g, 3.3 mol) in water (1 L). The reaction was then allowed to stir at 25 °C for 48 h. The resulting solution was concentrated to ca. 500 mL and acidified (conc. HCl) until pH = 1 was reached. The precipitated salts were filtered and the filtrate subjected to continuous liquid/liquid extraction with diethyl ether (1 L) for 24 h. The diethyl ether was removed on a rotovap to afford the product β -hydroxy acid as a colorless oil (250 g, 97%). The crude product was sufficiently pure to use in the next step (Kolbe-

coupling).

b) Preparation of (2R,5R)-2,5-hexanediol. A 100 mL reaction vessel was charged with (3R)-3-hydroxybutyric acid (1.0 g, 9.6 mmol), methanol (30 mL) and sodium methoxide (1.0 mL of a 0.5 N solution in methanol, 0.05 mmol), and then was cooled to 0 °C. Using a Pt foil anode (5 cm²), a Pt screen cathode (5 cm²), and a 50 V/40 amp power supply, a constant current (current density 0.25 A/cm²) was applied until 1388 coulombs (1.5 F/mol) were passed. The reaction and gas evolution (H₂ and CO₂) proceeded normally until ca. 1.0 F/mol current were passed, after which the resistance was observed to increase. The colorless solution was concentrated on a rotovap. Chromatography on SiO₂ (70% ethyl acetate/hexane) afforded the product as a colorless crystalline solid (0.36 g, 64%). mp 53-54 °C; [α]_D²⁵ = -37.6° (c 1, CHCl₃); ¹H NMR (CD₂Cl₂) δ 1.15 (d, J_{HH} = 6.2 Hz, 6H CH₃), 1.50 (m, 4H, CH₂), 2.95 (br, 2H, OH), 3.75 (m, 2H, CH); ¹³C NMR (CD₂Cl₂) δ 23.6, 35.9, 68.1.

c) Preparation of (2R,5R)-2,5-hexanediol bis(methanesulfonate). To a solution of (2R,5R)-2,5-hexanediol (8.9 g, 0.075 mol) in CH₂Cl₂ (200 mL) was added triethylamine (26.2 mL, 0.188 mol). The solution was cooled to 0 °C, and methanesulfonyl chloride (12.82 mL, 0.166 mol) in CH₂Cl₂ (30 mL) was added dropwise over 30 min. Upon complete addition, the mixture containing precipitated salts was allowed to stir at 0 °C for 30 min, and then at 25 °C for 30 min. The mixture was then poured into 1N HCl (250 mL) at 0 °C. After shaking, the layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2 x 50 mL). The combined organic layers were washed successively with 1N HCl (50 mL), saturated NaHCO₃, and brine. After drying (MgSO₄), the solution was concentrated on a rotovap to a pale yellow oil (18.2 g, 88%). The crude product thus obtained was sufficiently pure to be used in further reactions. ¹H NMR (CDCl₃) δ 1.41 (d, J_{HH} = 6.3 Hz, 6H, CH₃), 1.78 (m, 4H, CH₂), 3.0 (s, 6H, CH₃), 4.85 (m, 2H, CH).

d) Preparation of (2R,5R)-2,5-dimethyl-1-phenylphospholane. To a slurry of Li₂PPh-THF (20.3 g, 0.105 mol) in THF (300 mL) at -78 °C was added dropwise a solution of (2S,5S)-2,5-hexanediol bis(methanesulfonate) (26.0 g, 0.095 mol) in THF (50 mL). Upon complete addition, the orange mixture was allowed to stir at -78 °C for 1 h. The reaction was then slowly warmed to 25 °C and stirring was continued for 16 h. The resulting pale yellow mixture was filtered through a coarse frit, and concentrated to a semi-solid. Extraction with pentane (100 mL) and filtration, followed by concentration *in vacuo* yields a pale yellow oil. Distillation afforded the product as a colorless oil (13.9 g, 76%); bp 61-64 °C (0.2 torr); [α]_D²⁵ = -49.0 ± 2° (c 1, hexane); ¹H NMR (C₆D₆) δ 0.70 (dd, J_{HH} = 7.2 Hz, J_{PH} = 10.6 Hz, 3H, CH₃), 1.1-1.3 (m, 2H, CH₂), 1.20 (dd, J_{HH} = 7.2 Hz, J_{PH} = 18.8 Hz, 3H, CH₃), 1.65 (m, 1H, CH), 2.0 (m, 2H, CH₂), 2.45 (m, 1H, CH); ³¹P NMR (C₆D₆) δ 10.0; ¹³C NMR (C₆D₆) δ 15.43 (CH₃), 21.23 (d, J_{PC} = 34.2 Hz, CH₃), 32.25 (d, J_{PC} = 10.0 Hz, CH), 35.62 (d, J_{PC} = 13.1 Hz, CH), 37.17 (CH₂), 37.24 (d, J_{PC} = 3.6 Hz, CH₂), 128.11, 128.30, 134.51 (d, J_{PC} = 19.0 Hz, *ortho*), 137.67 (d, J_{PC} = 28.1 Hz, *ipso* Ph); HRMS (EI, direct insert): m/z 192.1068 (M⁺, exact mass calcd for C₁₂H₁₇P: 192.1068), 177.0839 (M-CH₃), 150.0559 (M-C₃H₆), 135.0367 (M-C₄H₉), 108.0127 (C₆H₅P fragment).

e) (2S,5S)-2,5-dimethyl-1-phenylphospholane [α]_D²⁵ = +51.6 ± 2° (c 1, hexane). Other spectroscopic properties were identical to Example 1 d).

f) Preparation of 1,2-Bis((2R,5R)-2,5-dimethylphospholane)ethane. To phospholane of Example 1 d) (6.0 g, 0.031 mol) in THF (200 mL) at 25 °C under Ar was added clean Li ribbon (0.54 g, 0.078 mol), and the reaction was allowed to stir for 10 h. To the resulting brown/green mixture was added dropwise a solution of ethylene glycol di-p-tosylate (6.90 g, 0.018 mol) in THF (100 mL). After stirring for 1 h, the mixture was filtered (coarse frit) and MeOH (2 mL) was added to the filtrate which turned pale yellow. The reaction was allowed to stir for 30 min and then filtered. The filtrate was concentrated to dryness *in vacuo*, and the resulting solids were extracted with pentane (200 mL) and filtered. The pentane filtrate was concentrated to a yellow oil which was distilled to afford the product as a colorless oil (2.10 g, 52%); bp 64-67 °C (0.06 torr); [α]_D²⁵ = +222 ± 6° (c 1, hexane); ¹H NMR (C₆D₆) δ 0.98 (dd, J_{HH} = 7.2 Hz, J_{PH} = 9.1 Hz, 3H, CH₃), 1.0-1.35 (m, 5H, CH₂), 1.22 (dd, J_{HH} = 7.2 Hz, J_{PH} = 17.3 Hz, 3H, CH₃), 1.55 (m, 2H, CH, CH₂), 1.70 (m, 5H, CH, CH₂), 1.90 (m, 4H, CH, CH₂); ³¹P NMR (C₆D₆) δ 3.2; ¹³C NMR (C₆D₆) δ 14.6 (CH₃), 20.71 (d, J_{PC} = 6.0 Hz, CH₃), 21.48 (dd, J_{PC} = 15.5 Hz, bridge CH₂), 34.44 (dd, J_{PC} = 5.8, 5.9 Hz, ring CH), 37.02 (ring CH₂), 37.42 (ring CH₂), 38.32 (dd, J_{PC} = 5.5 Hz, ring CH); HRMS (EI, direct insert): m/z 258.1670 (M⁺, exact mass calcd for C₁₄H₂₈P₂: 258.1667), 230.1344 (M-C₂H₄), 175.0785 (M-C₆H₁₁), 144.1072 (M-C₆H₁₁P), 116.0748 (M-C₈H₁₅P).

g) Optical Purity of Phosphines. The phosphines of Example 1 d), e) and f) were ascertained to be optically pure (within the limits of detection) by reacting each with (R)-[dimethyl-(α -methylbenzyl)-aminato-C,N]palladium (II) chloride dimer and monitoring the ³¹P NMR spectrum. Comparisons were made with the spectrum of the opposite phosphine enantiomer.

EXAMPLE 2

Preparation of 1,3-Bis((2R,5R)-2,5-dimethylphospholano)propane. To phospholane of Example 1 d) (6.0 g, 0.031 mol) in THF (200 mL) at 25 °C under Ar was added clean Li ribbon (0.54 g, 0.078 mol), and the reaction was allowed to stir for 10 h. To the resulting brown/green mixture was added dropwise a solution of 1,3-dichloropropane (2.11 g, 0.018 mol) in THF (25 mL). The reaction decolorized towards the end of the addition, and after stirring for 30 min, MeOH (2 mL) was added. This mixture was allowed to stir 10 min, then was filtered, and the filtrate concentrated *in vacuo*. The resulting oil was extracted with pentane (125 mL), filtered, and the pentane layer was concentrated to a yellow oil. Distillation afforded the product as a colorless oil (3.2 g, 75%); bp 98-101 °C (0.08 torr); $[\alpha]_D^{25} = +279 \pm 6$ (c 1, hexane); $^1\text{H NMR}$ (C_6D_6) δ 1.0 (dd, $J_{\text{HH}} = 7.1$ Hz, $J_{\text{PH}} = 9.6$ Hz, 3H, CH_3), 1.05 (m, 2H, CH_2), 1.20 (dd, $J_{\text{HH}} = 7.1$ Hz, $J_{\text{PH}} = 17.5$ Hz, 3H, CH_3), 1.20 (m, 2H, CH_2), 1.40 (m, 2H, CH_2), 1.55 (m, 4H, CH_2), 1.70 (m, 4H, CH, CH_2), 1.90 (m, 4H, CH, CH_2); $^{31}\text{P NMR}$ (C_6D_6) δ -2.85; $^{13}\text{C NMR}$ (C_6D_6) δ 14.6 (CH_3), 21.45 (d, $J_{\text{PC}} = 30.8$ Hz, CH_3), 24.34 (t, $J_{\text{PC}} = 18.9$ Hz, bridge central CH_2), 25.70 (dd, $J_{\text{PC}} = 11.3$, 22.3 Hz, bridge CH_2), 34.05 (d, $J_{\text{PC}} = 12.1$ Hz, ring CH), 37.10 (d, $J_{\text{PC}} = 3.6$ Hz, ring CH_2), 37.51 (ring CH_2), 38.30 (d, $J_{\text{PC}} = 11.5$ Hz, ring CH); HRMS (EI, direct insert): m/z 272.1816 (M^+ exact mass calcd for $\text{C}_{15}\text{H}_{30}\text{P}_2$: 272.1823), 229.1283 ($M - \text{C}_3\text{H}_7$), 188.0831 ($M - \text{C}_6\text{H}_{12}$), 157.1139 ($M - \text{C}_6\text{H}_{12}\text{P}$), 130.0900 ($M - \text{C}_8\text{H}_{15}\text{P}$), 116.0742 ($\text{C}_6\text{H}_{13}\text{P}$ fragment).

Optical Purity. The phosphine of Example 2 was ascertained to be optically pure (within the limits of detection) by reacting it with (R)-[dimethyl-(α -methylbenzyl)-aminato-C,N]palladium (II) chloride dimer and monitoring the $^{31}\text{P NMR}$ spectrum. Comparisons were made with the spectrum of the opposite phosphine enantiomer.

EXAMPLE 3

a) Rhodium complex $[(\text{COD})\text{Rh}(1,3\text{-bis}((2\text{R},5\text{R})\text{-2,5-dimethylphospholano})\text{propane})]^+\text{PF}_6^-$. To a mixture of $[(\text{COD})\text{RhCl}]_2$ (0.44 g, 0.89 mmol, COD = 1,5-cyclooctadiene) and NaPF_6 (0.40 g, 2.4 mmol) in THF (20 mL) at 25 °C was added dropwise a solution of 1,3-bis((2R, 5R)-2,5-dimethylphospholano)propane (0.50 g, 1.8 mmol) in THF (5 mL). The solution turned orange from yellow upon the phosphine addition. The reaction was allowed to stir for 30 min, and then was concentrated to a volume of ca. 5 mL. The slow addition of Et_2O (30 mL) to the solution produced an orange precipitate which was filtered, washed with Et_2O , and briefly dried. The solids were dissolved in CH_2Cl_2 (5 mL), filtered, and Et_2O (30 mL) was added slowly to the orange filtrate to provide the product as an orange microcrystalline solid (0.86 g, 75%); $^1\text{H NMR}$ (CD_2Cl_2) δ 1.15 (dd, $J_{\text{HH}} = 6.9$ Hz, $J_{\text{PH}} = 14.8$ Hz, 6H, CH_3), 1.50 (dd, $J_{\text{HH}} = 7.2$ Hz, $J_{\text{PH}} = 18.7$ Hz, 6H, CH_3), 1.3-1.6 (m, 6H, CH_2), 1.80 (m, 2H, CH, CH_2), 2.10 (m, 4H, CH, CH_2), 2.20-2.60 (m, 12H, CH_2 , CH), 4.80 (m (br), 2H, COD-CH), 5.15 (m (br), 2H, COD-CH); $^{31}\text{P NMR}$ (CD_2Cl_2) δ 27.7 (d, $J_{\text{RHP}} = 139.6$ Hz), -145 (sept., PF_6); Anal. Calcd for $\text{C}_{23}\text{H}_{42}\text{F}_6\text{P}_3\text{Rh}$: C, 43.96; H, 6.74. Found: C, 44.19; H, 6.43.

b) Rhodium complex $[(\text{COD})\text{Rh}((2\text{R},5\text{R})\text{-2,5-dimethyl-1-phenylphospholane})_2]^+\text{SbF}_6^-$. This complex was prepared in a manner analogous to that of Example 3 a). $^1\text{H NMR}$ (CD_2Cl_2) δ 0.70 (dd, $J_{\text{HH}} = 6.9$ Hz, $J_{\text{PH}} = 13.8$ Hz, 6H, CH_3), 1.00 (m, 2H, CH_2), 1.40 (m, 2H, CH_2), 1.62 (dd, $J_{\text{HH}} = 7.2$ Hz, $J_{\text{PH}} = 18.6$ Hz, 6H, CH_3), 1.90 (m, 2H, CH_2 , CH), 2.20 (m, 6H, CH_2 , CH), 2.40 (m, 8H, COD- CH_2), 4.90 (br, 2H, COD-CH), 5.34 (br, 2H, COD-CH), 7.0 (m, 4H, Ph), 7.30 (m, 4H, Ph), 7.40 (m, 2H, Ph); $^{31}\text{P NMR}$ (CD_2Cl_2) δ 43.8 (d, $J_{\text{RHP}} = 143.4$ Hz). Anal. calcd. for $\text{C}_{32}\text{H}_{46}\text{P}_2\text{F}_6\text{SbRh}$: C, 46.23; H, 5.58; P, 7.45. Found: C, 46.26; H, 5.47; P, 7.43.

X-ray Crystallographic Analysis. Crystal data for $\text{C}_{32}\text{H}_{46}\text{F}_6\text{P}_2\text{RhSb}$: monoclinic, P21 (No. 4), $a = 12.184(4)$ Å, $b = 12.734(2)$ Å, $c = 10.771(1)$ Å, $\beta = 96.83(1)^\circ$, $T = -100^\circ\text{C}$, $V = 1659.3$ Å³, Mo K α radiation, $\mu_{\text{calcd}} = 14.58$ cm⁻¹, $d_{\text{calcd}} = 1.664$ g cm⁻³, $Z = 2$, FW = 831.33.

Suitable crystals of the rhodium complex were obtained by slow vapor diffusion of Et_2O into a CH_2Cl_2 solution at 25 °C. An orange needle of dimensions 0.16 x 0.32 x 0.38 mm was mounted in a nitrogen-filled thin-walled glass capillary, and data was collected on a Syntex R3 diffractometer at -100 °C. The unit cell dimensions were determined by least squares refinement of 49 reflections. The crystal stability was monitored throughout the data collection by measuring the intensity of three standard reflections every 180 data points. The data were adjusted for a 2% decrease in intensity over the course of data acquisition. Lorentzian, polarization and absorption (azimuthal) corrections were applied, due to the relatively large absorption coefficient $\mu(\text{Mo}) = 14.58$ cm⁻¹.

The structure was solved using direct methods (SHELXS). All hydrogen atoms were determined from difference Fourier maps and idealized (C-H = 0.95 Å). Anisotropic refinement was carried out by full-matrix least squares on F.

Neutral atom scattering factors and anomalous scattering terms for P, Rh, and Sb were obtained from the *International Tables for X-ray Crystallography*, Vol. IV, herein incorporated by reference. Non-hydrogen atoms were refined anisotropically, and the hydrogens were refined isotropically for a total of 378 parameters. The refinement converged at $R = 0.035$, $R_w = 0.041$, and $EOF = 1.61$ for 6845 unique reflections with $I > 3.0\sigma(I)$. Fractional coordinates and isotropic thermal parameters for all non-H atoms were provided in Table S-1, while anisotropic thermal parameters were given in Table S-2. A complete listing of bond lengths and angles were also supplied.

The structure consisted of discrete molecules in which the usual square planar geometry of Rh(I) was somewhat distorted. The molecule had C_2 symmetry, but the 1,5-cyclooctadiene ring was rotated out of the P-Rh-P plane by 17.1 degrees. This distortion was the result of interactions with the methyl groups of the phosphine ligands. The Rh-C distances also were disparate with C1 and C5 being closer than C2 and C6. The rather large thermal motion of the phenyl rings and the fluorine atoms were self-evident and limited the quality of the structure. Attempts to refine the hydrogen atoms were unsatisfactory and these atoms were idealized and fixed with isotropic thermal parameters one higher than their associated carbon atoms. A full hemisphere of data was used in the refinement of the structure; the enantiomorphic structure refined to higher values $R = 0.039$ and $R_w = 0.048$.

c) Rhodium complex [(COD)Rh-Bis((2R,5R)-2,5-dimethylphospholano)ethane] $^+$ SbF $_6^-$. This complex was prepared in a manner analogous to that of Example 3 a). 1H NMR (CD_2Cl_2) δ 1.20 (dd, $J_{HH} = 6.9$ Hz, $J_{PH} = 14.4$ Hz, 6H, CH_3), 1.40 (dd, $J_{HH} = 7.1$ Hz, $J_{PH} = 17.7$ Hz, 6H, CH_3), 1.3-1.6 (m, 6H, CH_2), 1.95 (m, 2H, CH, CH_2), 2.10-2.60 (m, 10H, CH_2 , CH), 4.95 (br, 2H, COD-CH), 5.40 (br, 2H, COD-CH); ^{31}P NMR (CD_2Cl_2) δ 76.7 (d, $J_{RhP} = 146.3$ Hz). Calcd. for $C_{22}H_{40}F_6P_2SbRh$: C, 37.47; H, 5.72. Found: C, 37.64; H, 5.37.

X-ray Crystallographic Analysis. Suitable crystals of the rhodium complex were obtained by slow crystallization from a CH_2Cl_2 /hexane (2/1) solution at 25°C. An orange needle of dimensions 0.10 x 0.12 x 0.71 mm was mounted in a nitrogen-filled thin-walled glass capillary, and data was collected on a Syntex R3 diffractometer at -100°C. The unit cell dimensions were determined by least squares refinement of 49 reflections. The crystal stability was monitored throughout the data collection by measuring the intensity of three standard reflections every 182 data points. The data were adjusted for a 4% decrease in intensity over the course of data acquisition. Lorentzian, polarization and absorption (azimuthal) corrections were applied, due to the relatively large absorption coefficient $\mu(Mo) = 17.72$ cm^{-1} .

The structure was solved using direct methods (SHELXS). All hydrogen atoms were determined from difference Fourier maps and idealized (C-H = 0.95 Å). Anisotropic refinement was carried out by full-matrix least squares on F. Neutral atom scattering factors and anomalous scattering terms for P, Rh, and Sb were obtained from the *International Tables for X-ray Crystallography*, Vol. IV, herein incorporated by reference. Non-hydrogen atoms were refined anisotropically, and the hydrogens were refined isotropically for a total of 289 parameters. The refinement converged at $R = 0.038$, $R_w = 0.036$, and $EOF = 1.07$. Fractional coordinates and isotropic thermal parameters for all non-H atoms were provided in Table S-1, while anisotropic thermal parameters were given in Table S-2. A complete listing of bond lengths and angles were also supplied.

Like the Rh complex of Example 3 a), this structure consisted of discrete molecules in which the usual square planar geometry was distorted to an even greater extent. The complex had essentially C_2 symmetry with the 1,5-cyclooctadiene ligand rotated well out of the P-Rh-P plane by 24.3 degrees.

EXAMPLE 4

Hydrogenation Procedure. In a dry box, a 100 mL Fisher-Porter tube was charged with substrate (1.26 mmol), dry degassed MeOH or THF (20 mL), and the catalyst of Example 3 a), b) or c) (0.2 mol %). After two freeze-pump-thaw cycles, the tube was pressurized to an initial pressure of 10 psig with H_2 (Matheson, 99.998%). The reactions were allowed to stir at 25°C for 3-12 h. Hydrogen uptake was monitored and complete reaction was indicated by GC and NMR analyses. Reactions were worked-up as previously described. Enantiomeric excesses were determined by HPLC (methyl acetamidophenylalanine, Chiralcel OB, 5% IPA/Hexane) or shift reagent (dimethyl methylsuccinate, (+)-Eu(hfc) $_3$) analyses. The resulting data is listed in Table I.

TABLE I

Substrate	Enantiomeric Excess
Methyl acetamidocinnamate	85%
Dimethyl itaconate	91%

EXAMPLE 5

Preparation of Tris(((2S,5S)-2,5-dimethylphospholano)methyl)methane. To (2S,5S)-2,5-dimethyl-1-phenylphospholane (6.07 g, 0.032 mol) in THF (200 mL) at 25 °C under Ar was added clean Li ribbon (0.55 g, 0.079 mmol), and the reaction was allowed to stir for 15 h. To the resulting brown-orange mixture was added dropwise a solution of 1,3-dichloro-2-(chloromethyl)propane (1.70 g, 10.5 mmol) in THF (15 mL) at 25 °C. The reaction remained brown throughout the addition, and after stirring for 30 min, MeOH (3 mL) was added. The resulting colorless mixture was allowed to stir 15 min, then was filtered through a celite pad, and the filtrate concentrated *in vacuo*. The resulting solids were extracted with pentane (125 mL), filtered, and the pentane layer was concentrated to ca. 15 mL. Rapid filtration afforded the product as a colorless crystalline solid (1.0 g). The filtrate was then concentrated to a pale yellow solid which was dissolved in a minimum amount of Et₂O (3 mL). To this solution was added MeOH (15 mL) and the mixture was cooled to -20 °C for 12 h. The resulting white crystals were filtered, washed with cold MeOH and dried *in vacuo* (1.65 g). Combined yield 2.65 g (63%): $[\alpha]_D^{25} = -329 \pm 6^\circ$ (c 1, hexane); ¹H NMR (C₆D₆) δ 1.0-1.2 (m, 3H, CH, CH₂), 1.07 (dd, J_{HH} = 7.2 Hz, J_{PH} = 9.8 Hz, 9H, CH₃), 1.29 (dd, J_{HH} = 7.0 Hz, J_{PH} = 17.6 Hz, 9H, CH₃), 1.40 (m, 3H, CH, CH₂), 1.60 (m, 3H, CH₂), 1.80 (m, 3H, CH, CH₂), 1.9-2.2 (m, 13H, CH, CH₂); ³¹P NMR (C₆D₆) δ -8.0; ¹³C NMR (C₆D₆) δ 14.89 (CH₃), 21.51 (d, J_{PC} = 30.7 Hz, CH₃), 31.85 (dt, J_{PC} = 8.3, 22.1 Hz, bridge P-CH₂), 32.67 (q, J_{PC} = 14.9 Hz, bridge CH), 34.12 (d, J_{PC} = 11.6 Hz, ring CH), 37.30 (d, J_{PC} = 3.8 Hz, ring CH₂), 37.47 (ring CH₂), 38.49 (d, J_{PC} = 11.3 Hz, ring CH); HRMS (EI, direct insert): m/z 400.2583 (M⁺ exact mass calcd for C₂₂H₄₃P₃: 400.2578), 357.2021 (M-C₃H₇), 315.1557 (M-C₆H₁₃), 285.1896 (M-C₆H₁₂P), 273.1104 (M-C₉H₁₉), 232.0678 (M-C₁₂H₂₄), 201.0955 (M-C₁₂H₂₄P).

EXAMPLE 6

Preparation of Tris(2-((2R,5R)-2,5-dimethylphospholano)ethyl)amine. To (2R,5R)-2,5-dimethyl-1-phenylphospholane (3.0 g, 15.6 mmol) in THF (100 mL) at 25 °C under Ar was added clean Li ribbon (0.27 g, 39.0 mmol), and the reaction was allowed to stir for 15 h. To the resulting brown/orange mixture was added dropwise a solution of tris(2-chloroethyl)amine (1.06 g, 5.2 mmol) in THF (15 mL) at 25 °C. The reaction remained brown throughout the addition, and after stirring for 30 min, MeOH (3 mL) was added. The resulting colorless mixture was allowed to stir 15 min, then was filtered through a celite pad, and the filtrate concentrated *in vacuo*. The resulting solids were extracted with pentane (125 mL), filtered, and the pentane layer was concentrated to ca. 15 mL. Rapid filtration afforded the product as a colorless crystalline solid (0.6 g). The filtrate was then concentrated to a pale yellow solid which was dissolved in a minimum amount of Et₂O (3 mL). To this solution was added MeOH (10 mL) and the mixture was cooled to -20 °C for 12 h. The resulting white crystals were filtered, washed with cold MeOH and dried *in vacuo* (1.13 g). Combined yield 1.73 g (75%): $[\alpha]_D^{25} = +167 \pm 2^\circ$ (c 1, hexane); ¹H NMR (C₆D₆) δ 1.0-1.2 (m, 3H, CH, CH₂), 1.11 (dd, J_{HH} = 7.2 Hz, J_{PH} = 9.8 Hz, 9H, CH₃), 1.32 (dd, J_{HH} = 7.0 Hz, J_{PH} = 17.7 Hz, 9H, CH₃), 1.30-1.45 (m, 3H, CH, CH₂), 1.55 (m, 3H, CH₂), 1.70-2.15 (m, 15H, CH, CH₂), 2.80 (m, 6H, NCH₂); ³¹P NMR (C₆D₆) δ -3.4.


EXAMPLE 7

Preparation of 1,1,1-Tris(2-((2R,5R)-2,5-dimethylphospholano)ethyl)ethane. To (2R,5R)-2,5-dimethyl-1-phenylphospholane (0.53 g, 2.76 mmol) in THF (10 mL) at 25 °C under Ar was added clean Li ribbon (0.048 g, 6.9 mmol), and the reaction was allowed to stir for 15 h. To the resulting brown/orange mixture was added dropwise a solution of 1,1,1-tris(2-chloroethyl)ethane (0.2 g, 0.92 mmol) in THF (5 mL) at 25 °C. The reaction remained brown throughout the addition, and after stirring for 30 min, MeOH (1 mL) was added. The resulting colorless mixture was allowed to stir 15 min, then was filtered through a celite pad, and the filtrate concentrated *in vacuo*. The resulting solids/oil were extracted with pentane (50 mL), filtered, and the pentane layer was concentrated to yield the product as a colorless, viscous oil (0.257 g, 61%): ¹H NMR (C₆D₆) δ 0.84 (s, 3H, CH₃), 1.11 (dd, J_{HH} = 6.9 Hz, J_{PH} = 9.8 Hz, 9H, CH₃), 1.20 (dd, J_{HH} = 7.2 Hz, J_{PH} =

17.6 Hz, 9H, CH₃), 1.00-1.50 (m, 18H, CH, CH₂), 1.80 (m, 3H, CH₂), 1.90-2.15 (m, 9H, CH, CH₂); ³¹P NMR (C₆D₆) δ 0.3; HRMS (EI, direct insert): m/z 456.3170 (M⁺ exact mass calcd for C₂₆H₅₁P₃: 456.3204), 371.2206 (M-C₆H₁₃), 341.2645 (M-C₆H₁₂P), 257.1612 (M-C₁₂H₂₄P).

5 Claims

1. A compound represented by the following formula I:




wherein:

R is a C₁-C₆ alkyl, trifluoromethyl, phenyl, substituted phenyl, aralkyl or ring-substituted aralkyl;
and

n is an integer having a value from 1 to 12.

2. The compound of Claim 1 which exhibits optical activity of greater than or equal to 90%.
3. A compound of Claim 1 or Claim 2 wherein R is methyl.
4. A compound of Claim 1, 2 or 3 wherein n is 1 to 3.
5. A compound of Claim 1 which is 1,2-bis-(2,5-dimethyl-phospholano)ethane.
6. A compound of Claim 1 which is 1,3-bis(2,5-dimethylphospholano)propane.
7. A complex comprising one or more transition metals, lanthanides or actinides and one or more compounds of any one of Claims 1 to 6 as ligands.
8. The complex of Claim 7 wherein the transition metal is rhodium.
9. The transition metal complex of Claim 8 which is [(COD)Rh-bis[(2R,5R)-2,5-dimethylphospholano]ethane]⁺SbF₆⁻.
10. The transition metal complex of Claim 8 which is [(COD)Rh-1,3-bis[(2R,5R)-2,5-dimethylphospholano]propane]⁺PF₆⁻.
11. A compound represented by the following formula II:



wherein:

R is a C₁-C₆ alkyl, trifluoromethyl, phenyl, substituted phenyl, aralkyl or ring-substituted aralkyl;


A is CCH₃, CH, N, or P; and

n is an integer having a value of from 1 to 12.

17.6 Hz, 9H, CH₃), 1.00-1.50 (m, 18H, CH, CH₂), 1.80 (m, 3H, CH₂), 1.90-2.15 (m, 9H, CH, CH₂); ³¹P NMR (C₆D₆) δ 0.3; HRMS (EI, direct insert): m/z 456.3170 (M⁺ exact mass calcd for C₂₆H₅₁P₃: 456.3204), 371.2206 (M-C₆H₁₃), 341.2645 (M-C₆H₁₂P), 257.1612 (M-C₁₂H₂₄P).

5 Claims

1. A compound represented by the following formula I:




wherein:

R is a C₁-C₆ alkyl, trifluoromethyl, phenyl, substituted phenyl, aralkyl or ring-substituted aralkyl; and

n is an integer having a value from 1 to 12.

2. The compound of Claim 1 which exhibits optical activity of greater than or equal to 90%.
3. A compound of Claim 1 or Claim 2 wherein R is methyl.
4. A compound of Claim 1, 2 or 3 wherein n is 1 to 3.
5. A compound of Claim 1 which is 1,2-bis-(2,5-dimethyl-phospholano)ethane.
6. A compound of Claim 1 which is 1,3-bis(2,5-dimethylphospholano)propane.
7. A complex comprising one or more transition metals, lanthanides or actinides and one or more compounds of any one of Claims 1 to 6 as ligands.
8. The complex of Claim 7 wherein the transition metal is rhodium.
9. The transition metal complex of Claim 8 which is [(COD)Rh-bis[(2R,5R)-2,5-dimethylphospholano]-ethane]⁺SbF₆⁻.
10. The transition metal complex of Claim 8 which is [(COD)Rh-1,3-bis[(2R,5R)-2,5-dimethylphospholano]-propane]⁺PF₆⁻.
11. A compound represented by the following formula II:



wherein:


R is a C₁-C₆ alkyl, trifluoromethyl, phenyl, substituted phenyl, aralkyl or ring-substituted aralkyl;

A is CCH₃, CH, N, or P; and


n is an integer having a value of from 1 to 12.



12. A compound of Claim 11 which exhibits optical activity of greater than or equal to 90%.
13. A compound of Claim 11 or Claim 12 wherein R is methyl.
14. A compound of Claim 11, 12 or 13 wherein n is 1 to 3.
15. A compound of Claim 12 which is tris[[(2S,5S)-2,5-dimethylphospholano]methyl]methane.
16. A compound of Claim 12 which is tris(2-[(2R,5R)-2,5-dimethylphospholano]ethyl)amine.
17. A compound of Claim 12 which is 1,1,1-tris(2-[(2R,5R)-2,5-dimethylphospholano]ethyl)ethane.
18. A complex comprising one or more transition metals, lanthanides, or actinides and one or more compounds of any one of Claims 11 to 17 as ligands.
19. A complex of Claim 18 comprising a transition metal and one or more compounds of Claim 12 as ligands.
20. A compound represented by the following formula III:



- wherein R is an alkyl group of two to six carbon atoms, trifluoromethyl, phenyl, substituted phenyl, aralkyl or ring substituted aralkyl.
21. A compound of Claim 20 which exhibits optical activity of greater than or equal to 90%.
22. The transition metal complex which is $[(\text{COD})\text{Rh}((2R,5R)\text{-}2,5\text{-dimethyl-1-phenylphospholane})_2]^+\text{SbF}_6^-$.
23. A process for the preparation of a compound of formula I



wherein:


R is a $\text{C}_1\text{-C}_6$ alkyl, trifluoromethyl, phenyl, substituted phenyl, aralkyl or ring-substituted aralkyl;

and

n is an integer of 1 to 12;

comprising reacting a phenyl substituted phospholane of formula III





10 wherein R is defined as in formula I, with lithium and a compound of formula




15 wherein X is halogen, OR^1 or R^1O are methanesulfonate, trifluoromethanesulfonate or p-toluenesulfonate, and n is an integer from 1 to 12 to yield the desired compound of formula I.

24. The process of Claim 23 conducted in tetrahydrofuran as a solvent.

25. The process of Claim 23 or Claim 24 conducted at a temperature of from about $0^\circ C$ to about $40^\circ C$.

26. The process of Claim 23, 24 or 25 conducted in an inert atmosphere.

27. A process for the preparation of a compound of formula II




35 wherein:

R is a C_1-C_6 alkyl, trifluoromethyl, phenyl, substituted phenyl, aralkyl or ring-substituted aralkyl;

n is an integer having a value of from 1 to 12; and

A is CCH_3 , CH, N or P;

comprising reacting a phenyl substituted phospholane of formula III



45 wherein R is as defined in formula II, with lithium and a trihalo compound of formula



50 wherein X is halogen and A is as defined in formula II, to yield the desired compound of formula II.

55 28. The process of Claim 27 conducted in tetrahydrofuran as a solvent.


29. The process of Claim 27 or Claim 28 conducted at a temperature of from about $-78^\circ C$ to about $40^\circ C$.



30. The process of Claim 27, 28 or 29 conducted in an inert atmosphere.

Patentansprüche

1. Verbindung, die durch die folgende Formel I dargestellt wird:




worin:

R C₁-C₆-Alkyl, Trifluormethyl, Phenyl, substituiertes Phenyl, Aralkyl oder ringsubstituiertes Aralkyl ist und

n eine ganze Zahl mit einem Wert von 1 bis 12 ist.

2. Verbindung gemäß Anspruch 1, die eine optische Aktivität von größer oder gleich 90% zeigt.
3. Verbindung gemäß Anspruch 1 oder 2, worin R Methyl ist.
4. Verbindung gemäß Anspruch 1, 2 oder 3, worin n 1 bis 3 ist.
5. Verbindung gemäß Anspruch 1, bei der es sich um 1,2-Bis(2,5-dimethylphospholano)ethan handelt.
6. Verbindung gemäß Anspruch 1, bei der es sich um 1,3-Bis(2,5-dimethylphospholano)propan handelt.
7. Komplex, der ein oder mehrere Übergangsmetalle, Lanthanide oder Actinide sowie eine oder mehrere Verbindungen gemäß einem der Ansprüche 1 bis 6 als Liganden umfaßt.
8. Komplex gemäß Anspruch 7, worin das Übergangsmetall Rhodium ist.
9. Übergangsmetallkomplex gemäß Anspruch 8, bei dem es sich um [(COD)Rh-bis[(2R,5R)-2,5-dimethylphospholano]ethan]⁺SbF₆⁻ handelt.
10. Übergangsmetallkomplex gemäß Anspruch 8, bei dem es sich um [(COD)Rh-1,3-bis[(2R,5R)-2,5-dimethylphospholano]propan]⁺PF₆⁻ handelt.
11. Verbindung, die durch die folgende Formel II dargestellt wird:



worin:

R C₁-C₆-Alkyl, Trifluormethyl, Phenyl, substituiertes Phenyl, Aralkyl oder ringsubstituiertes Aralkyl ist;

A CCH₃, CH, N oder P ist und

n eine ganze Zahl mit einem Wert von 1 bis 12 ist.

12. Verbindung gemäß Anspruch 11, die eine optische Aktivität von größer oder gleich 90% zeigt.



13. Verbindung gemäß Anspruch 11 oder 12, worin R Methyl ist.

14. Verbindung gemäß Anspruch 11, 12 oder 13, worin n 1 bis 3 ist.

15. Verbindung gemäß Anspruch 12, bei der es sich um Tris[[(2S,5S)-2,5-dimethylphospholano)methyl]-methan handelt.


16. Verbindung gemäß Anspruch 12, bei der es sich um Tris(2-[(2R,5R)-2,5-dimethylphospholano]ethyl)-amin handelt.

17. Verbindung gemäß Anspruch 12, bei der es sich um 1,1,1-Tris(2-[(2R,5R)-2,5-dimethylphospholano]ethyl)ethan handelt.

18. Komplex, der ein oder mehrere Übergangsmetalle, Lanthanide oder Actinide sowie eine oder mehrere Verbindungen gemäß einem der Ansprüche 11 bis 17 als Liganden umfaßt.

19. Komplex gemäß Anspruch 18, der ein Übergangsmetall und eine oder mehrere Verbindungen gemäß Anspruch 12 als Liganden umfaßt.

20. Verbindung, die durch die folgende Formel III dargestellt wird:




worin R eine Alkylgruppe mit zwei bis sechs Kohlenstoffatomen, Trifluormethyl, Phenyl, substituiertes Phenyl, Aralkyl oder ringsubstituiertes Aralkyl ist.

21. Verbindung gemäß Anspruch 20, die eine optische Aktivität von größer oder gleich 90% zeigt.

22. Übergangsmetallkomplex, bei dem es sich um $[(\text{COD})\text{Rh}((2R,5R)\text{-}2,5\text{-dimethyl-1-phenylphospholan})_2]^+ \text{SbF}_6^-$ handelt.

23. Verfahren zur Herstellung einer Verbindung der Formel I




worin:

R C₁-C₆-Alkyl, Trifluormethyl, Phenyl, substituiertes Phenyl, Aralkyl oder ringsubstituiertes Aralkyl ist und

n eine ganze Zahl mit einem Wert von 1 bis 12 ist;

umfassend die Umsetzung eines phenylsubstituierten Phospholans der Formel III





III

worin R wie in Formel I definiert ist, mit Lithium und einer Verbindung der Formel




worin X Halogen ist, OR^1 oder R^1O Methansulfonat, Trifluormethansulfonat oder *p*-Toluolsulfonat sind und n eine ganze Zahl von 1 bis 12 ist, wobei man die gewünschte Verbindung der Formel I erhält.

24. Verfahren gemäß Anspruch 23, das in Tetrahydrofuran als Lösungsmittel durchgeführt wird.

25. Verfahren gemäß Anspruch 23 oder 24, das bei einer Temperatur von etwa $0^\circ C$ bis etwa $40^\circ C$ durchgeführt wird.

26. Verfahren gemäß Anspruch 23, 24 oder 25, das in einer inerten Atmosphäre durchgeführt wird.

27. Verfahren zur Herstellung einer Verbindung der Formel II



II


worin:

R C_1 - C_6 -Alkyl, Trifluormethyl, Phenyl, substituiertes Phenyl, Aralkyl oder ringsubstituiertes Aralkyl ist;

n eine ganze Zahl mit einem Wert von 1 bis 12 ist und


A CCH_3 , CH, N oder P ist;

umfassend die Umsetzung eines phenylsubstituierten Phospholans der Formel III



III

worin R wie in Formel II definiert ist, mit Lithium und einer Trihalogenverbindung der Formel



worin X Halogen ist und A wie in Formel II definiert ist, wobei man die gewünschte Verbindung der Formel II erhält.

28. Verfahren gemäß Anspruch 27, das in Tetrahydrofuran als Lösungsmittel durchgeführt wird.


29. Verfahren gemäß Anspruch 27 oder 28, das bei einer Temperatur von etwa $-78^\circ C$ bis etwa $40^\circ C$ durchgeführt wird.



30. Verfahren gemäß Anspruch 27, 28 oder 29, das in einer inerten Atmosphäre durchgeführt wird.

Revendications

1. Un composé représenté par la formule I suivante :




dans laquelle :

R est un groupe alkyle en C₁-C₆, trifluorométhyle, phényle, phényle substitué, aralkyle ou aralkyle substitué sur le noyau ; et

n est un nombre entier ayant une valeur de 1 à 12.

2. Le composé de la revendication 1, qui présente une activité optique supérieure ou égale à 90 %.
3. Un composé de la revendication 1 ou de la revendication 2, dans lequel R est le groupe méthyle.
4. Un composé de la revendication 1, 2 ou 3, dans lequel n est de 1 à 3.
5. Un composé de la revendication 1, qui est le 1,2-bis(2,5-diméthylphospholano)éthane.
6. Un composé de la revendication 1, qui est le 1,3-bis(2,5-diméthylphospholano)propane.
7. Un complexe comprenant un ou plusieurs métaux de transition, lanthanides ou actinides et un ou plusieurs composés de l'une quelconque des revendications 1 à 6 comme ligands.
8. Le complexe de la revendication 7, dans lequel le métal de transition est le rhodium.
9. Le complexe de métal de transition de la revendication 8, qui est le [(COD)Rh-bis[(2R,5R)-2,5-diméthylphospholano]éthane]⁺SbF₆⁻.
10. Le complexe de métal de transition de la revendication 8, qui est le [(COD)Rh-1,3-bis[(2R,5R)-2,5-diméthylphospholano]propane]⁺PF₆⁻.
11. Un composé représenté par la formule II suivante :



dans laquelle :


R est un groupe alkyle en C₁-C₆, trifluorométhyle, phényle, phényle substitué, aralkyle ou aralkyle substitué sur le noyau ;

A est CCH₃, CH, N ou P ; et


n est un nombre entier ayant une valeur de 1 à 12.



12. Un composé de la revendication 11, qui présente une activité optique supérieure ou égale à 90 %.
13. Un composé de la revendication 11 ou de la revendication 12, dans lequel R est le groupe méthyle.
14. Un composé de la revendication 11, 12 ou 13, dans lequel n est de 1 à 3.
15. Un composé de la revendication 12, qui est le tris[[(2S,5S)-2,5-diméthylphospholano]méthyl]méthane.
16. Un composé de la revendication 12, qui est la tris(2-[(2R,5R)-2,5-diméthylphospholano]éthyl)amine.
17. Un composé de la revendication 12, qui est le 1,1,1-tris(2-[(2R,5R)-2,5-diméthylphospholano]éthyl)-éthane.
18. Un complexe comprenant un ou plusieurs métaux de transition, lanthanides ou actinides et un ou plusieurs composés de l'une quelconque des revendications 11 à 17 comme ligands.
19. Un complexe de la revendication 18, comprenant un métal de transition et un ou plusieurs composés de la revendication 12 comme ligands.
20. Un composé représenté par la formule III suivante :



- dans laquelle R est un groupe alkyle de deux à six atomes de carbone, trifluorométhyle, phényle, phényle substitué, aralkyle ou aralkyle substitué sur le noyau.
21. Un composé de la revendication 20, qui présente une activité optique supérieure ou égale à 90 %.
22. Le complexe de métal de transition qui est le [(COD)Rh((2R,5R)-2,5-diméthyl-1-phénylphospholane)₂]⁺SbF₆⁻.
23. Un procédé pour la préparation d'un composé de formule I



- dans laquelle :
- R est un groupe alkyle en C₁-C₆, trifluorométhyle, phényle, phényle substitué, aralkyle ou aralkyle substitué sur le noyau ; et
- n est un nombre entier de 1 à 12 ;
- comprenant la réaction d'un phospholane substitué par un groupe phényle de formule III





10 où R est tel que défini dans la formule I, avec du lithium et un composé de formule




15 où X est un halogène, OR' ou $R'O$ est un radical méthanesulfonate, trifluorométhanesulfonate ou *p*-toluènesulfonate, et n est un nombre entier de 1 à 12, pour produire le composé désiré de formule I.

24. Le procédé de la revendication 23, conduit dans du tétrahydrofuranne comme solvant.

25. Le procédé de la revendication 23 ou de la revendication 24, conduit à une température d'environ 0 °C à environ 40 °C.

26. Le procédé de la revendication 23, 24 ou 25, conduit dans une atmosphère inerte.

27. Un procédé pour la préparation d'un composé de formule II




dans laquelle :

35 R est un groupe alkyle en C_1-C_6 , trifluorométhyle, phényle, phényle substitué, aralkyle ou aralkyle substitué sur le noyau ;

n est un nombre entier ayant une valeur de 1 à 12 ; et

A est CCH_3 , CH, N ou P ;

40 comprenant la réaction d'un phospholane substitué par un groupe phényle de formule III



50 où R est tel que défini dans la formule II, avec du lithium et un composé trihalogéné de formule



55 où X est un halogène et A est tel que défini dans la formule II, pour produire le composé désiré de formule II.

28. Le procédé de la revendication 27, conduit dans du tétrahydrofuranne comme solvant.



29. Le procédé de la revendication 27 ou de la revendication 28, conduit à une température d'environ -78°C à environ 40°C .

30. Le procédé de la revendication 27, 28 ou 29, conduit dans une atmosphère inerte.

5

10

15

20

25

30

35

40

45

50

55

