

REPLACED SHEETS 50

☒ Description : paged 1 to paged 12

☒ Claim(s) : paged 13 to paged 16


☐ Drawing(s) : paged to paged

☒ Abstract : paged 17 to paged 18


DATE OF RECEIPT

The present invention relates to novel 1-[[[(di-alkylamino)alkyl]amino]-4-substituted-thioxanthen-9-ones, to pharmaceutical compositions containing the thioxanthenones, to methods of treating tumors with the thioxanthenones and to methods of treating cancer in mammals with the compositions containing the thioxanthenones.

Collins and Rosi U.S. Patent 3,745,172 disclose as an intermediate in the synthesis of antifungal and antibacterial agents:



and as an anthelmintic and antibacterial agent:




Blanz and French [J. Med. Chem. 6, 185-191 (1963)] disclose the synthesis of a series of thioxanthenones related to lucanthone and the results of the testing of

This Sheet Has Been Replaced




the compounds against a leukemia and two solid tumors.
Among the compounds disclosed are



wherein R is methyl, methoxy, and ethoxy.

- 5 Yarinsky and Freele [Journal of Tropical Medicine
and Hygiene 73, 23-27 (1970)] disclose




as an antischistosomal agent.

- 10 Palmer et al. [J. Med. Chem. 31, 707-712 (1988)]
disclose N-[2-(dimethylamino)ethyl]-9-oxo-9H-thio-
xanthene-4-carboxamide monohydrochloride which was
tested in vitro versus murine leukemia (L1210) and
in vivo versus P388 leukemia cells and was found to be
unlikely to be worth pursuing as a potential antitumor
15 agent.

This Sheet Has Been Replaced



Archer, U.S. Patent 4,539,412 discloses compounds of the formula:



wherein, for X=S:

- 5 R^1 and R^2 are individually selected from one of lower-alkyls, and jointly selected from one of pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl and N-substituted piperazinyl; and

R^3 is H or hydroxy.


- 10 wherein, for X=O:

R^1 and R^2 are individually selected from one of lower-alkyls, and jointly selected from one of pyrrolidinyl, piperidyl, morpholinyl, piperazinyl and N-substituted piperazinyl; and

- 15 R^3 is hydroxy.

The compounds are said to be useful as antitumor agents.

Archer *et al.* [J. Med. Chem. 31, 254-260 (1988)] disclose compounds of the general formula:



- 20 wherein $R=C_2H_5$, $R'=H$, $R''=CH_2OH$ (example 1);
 $R=CH_3$, $R'=OH$, $R''=CH_2OH$ (example 2);
 $R=C_2H_5$, $R'=H$, $R''=CH_2OCONHCH_3$ (example 4);
 $R=CH_3$, $R'=OH$, $R''=CH_2OCONHCH_3$ (example 5);

This Sheet Has Been Replaced



$R=C_2H_5$, $R'=H$, $R''=CH_2OCONHC_6H_5$ (example 23);

$R=C_2H_5$; $R'=H$, $R''=CH_2OCONHC_3H_7$ (example 24);

$R=C_2H_5$, $R'=H$, $R''=CH_2OCONHC_4H_9$ (example 25);


$R=C_2H_5$, $R'=H$, $R''=CH_2OCOCH_3$ (example 26);

5 $R=C_2H_5$, $R'=H$, $R''=CH_2OCOCH_3$ (example 27) or

$R=C_2H_5$, $R'=H$, $R''=CH_2OCOC_6H_3(NO_2)_2$ (example 28)

and the results of the testing of these compounds for antitumor and/or antischistosomicidal activity.

According to the present invention there is
10 provided a compound of the formula I



wherein: (1) n is 2 or 3;

R^1 and R^2 are independently lower-alkyl;

15 Q is a residue chosen from the group consisting of
 CH_2NHR^3 , $CH_2N(R^4)SO_2R^7$, CH_2NHCHO , $CH=N-Ar$,
 $C(O)NR^5R^6$, $CH_2N(R^4)C(O)R^7$, $CH_2N(C_2H_5)CHO$,
 $CH_2N(R^4)P(O)(O-lower-alkyl)_2$, $CH_2N=CH-N(R^9)(R^{10})$,
 $CH_2N(R^4)C(O)CF_3$ and $CH_2N(R^4)C(O)OR^7$;

R^3 is hydrogen or lower-alkyl;

20 R^4 is hydrogen, lower-alkyl or Ar;

R^5 is hydrogen, lower-alkyl or Ar;

R^6 is hydrogen or lower-alkyl;

R^7 is lower-alkyl, or Ar;

R^8 is hydroxy;

25 Ar is phenyl or phenyl substituted with methyl, methoxy, hydroxy, halogen or nitro; and

R^9 and R^{10} are independently lower-alkyl; or



(2) Q is a residue chosen from the group consisting of $\text{CH}_2\text{N}(\text{R}^4)\text{SO}_2\text{R}^7$, $\text{CH}=\text{N}-\text{Ar}$, $\text{C}(\text{O})\text{NR}^5\text{R}^6$, $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{R}^7$, $\text{CH}_2\text{N}(\text{R}^4)\text{P}(\text{O})(\text{O-lower-alkyl})_2$, $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{CF}_3$ and $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{OR}^7$; R^8 is hydrogen, lower-alkyl, lower-alkoxy, or hydroxy; Ar is phenyl substituted by hydroxy; and n, R^1 , R^2 , R^4 , R^5 , R^6 , and R^7 are as defined hereinabove in part (1) with the proviso that one or more of R^4 , R^5 , or R^7 is Ar; or

(3) Q is a residue chosen from the group consisting of $\text{CH}_2\text{N}=\text{CH}-\text{N}(\text{R}^9)(\text{R}^{10})$, $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{CF}_3$ and $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{OR}^7$; R^8 is hydrogen, lower-alkyl, lower-alkoxy, or hydroxy; and n, R^1 , R^2 , R^4 , R^7 , Ar, R^9 and R^{10} are as defined hereinabove in part (1); or a pharmaceutically acceptable acid-addition salt or solvate thereof;.

The compounds are useful for the treatment of tumors in mammals.

Lower-alkyl as used herein describes linear, branched or cyclic hydrocarbons containing four or fewer carbon atoms. Halogen describes bromine, chlorine or fluorine. The term lower-alkoxy means linear or branched alkyloxy substituents having from one to about four carbon atoms, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, and the like.

Preferred compounds within the scope of part (1) are those wherein Q is a residue chosen from the group consisting of CH_2NHR^3 , $\text{CH}_2\text{N}(\text{R}^4)\text{SO}_2\text{R}^7$, CH_2NHCHO , $\text{CH}=\text{N}-\text{Ar}$, $\text{C}(\text{O})\text{NR}^5\text{R}^6$, $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{R}^7$, $\text{CH}_2\text{N}(\text{C}_2\text{H}_5)\text{CHO}$ and $\text{CH}_2\text{N}(\text{R}^4)\text{P}(\text{O})(\text{O-lower-alkyl})_2$ and R^4 is hydrogen or lower alkyl, especially when Ar is phenyl or phenyl substituted with methyl, methoxy, halogen or nitro. Particularly preferred are those wherein n is 2, both R^1 and R^2 are ethyl, R^3 is hydrogen or methyl and R^7 is lower-alkyl and especially 1-[[2-(diethylamino)ethyl]-amino]-4-[(methylamino)methyl]-7-hydroxythioxanthen-

This Sheet Has Been Replaced



9-one and N-[[1-(2-(diethylamino)ethyl)amino]-7-hydroxy-9-oxothioxanthen-4-yl]methyl]methanesulfonamide.

Preferred compounds within the scope of part (3) are those wherein R^8 is hydrogen, lower-alkyl, or lower-alkoxy; R^4 is hydrogen and Ar is phenyl or phenyl substituted with methyl, methoxy, halogen or nitro. Particularly preferred are those wherein n is 2 and both of R^1 and R^2 are ethyl and R^8 is hydrogen or methoxy, especially wherein Q is $CH_2N=CH-N(R^9)(R^{10})$ and R^9 and R^{10} are both methyl or wherein Q is $CH_2N(R^4)C(O)OR^7$, especially wherein R^7 is lower-alkyl. Most preferred compounds are methyl N-[[1-[[2-(diethylamino)ethyl)-amino]-9-oxothioxanthen-4-yl]methyl]carbamate and methyl N-[[1-[[2-(diethylamino)ethyl)amino]-7-methoxy-9-oxo-
thioxanthen-4-yl]methyl]carbamate.


The invention further relates to compositions for treating tumors and cancer in mammals which comprise compounds of formula I together with pharmaceutically acceptable excipients or diluents.

The invention further relates to a method of use of a compound of formula I for treating tumors in mammals which comprises administering to the mammal a compound of formula I.


The invention further relates to a method of use of a compound of formula I for treating cancer in a mammal which comprises administering to the mammal a composition of a compound of formula I together with pharmaceutically acceptable excipients or diluents.

The synthesis of compounds of the invention may be outlined as shown in Schemes A, B and C:

This Sheet Has Been Replaced




SCHEME A



This Sheet Has Been Replaced



Scheme B



The compounds of formula III (formula I wherein Q is -CH=N-Ar) may be synthesized by heating an aldehyde of formula II together with about one equivalent of the appropriate aniline in an inert, azeotroping solvent, preferably xylene or toluene at reflux.

The compounds of formula IV may be prepared from an aldehyde of formula II by heating at 150° to 185°C in the presence of from about 5 to about 17 equivalents of formic acid in formamide, N-alkylformamide or N-Ar-formamide as a solvent. The conditions are well known in the art for the Leuckart reaction. The compounds of formula V are then obtained by acid hydrolysis of the formamide.



The compounds of formula VI (formula I wherein Q is $-\text{CH}_2\text{N}(\text{R}^4)\text{SO}_2\text{R}^7$) may be synthesized by sulfonylation of the amine V with a slight excess of a lower-alkyl-sulfonyl, or arylsulfonyl chloride in (a) a suitable solvent, preferably pyridine, at 0° to 50°C ; or

(b) a suitable solvent, such as dichloromethane, in the presence of an excess of a base, such as triethylamine, at a temperature in the range of about 0°C up to about room temperature.

Alternatively, the compounds of formula VI, wherein R^4 is lower-alkyl, may be synthesized by sulfonylation of the amine V, wherein R^4 is hydrogen, as described above, followed by treatment of the resulting sulfonamide VI, R^4 is hydrogen, with an excess of a base, preferably sodium hydride, in a suitable solvent, such as tetrahydrofuran or N,N-dimethylformamide, followed by treatment with an excess of an appropriate lower-alkyl halide at a temperature in the range of about 0°C up to the boiling point of the solvent used.

The carboxamides of formula IX may be synthesized by reacting the aldehyde II with a 5-6 fold excess of hydroxylamine hydrochloride in pyridine, optionally containing a cosolvent, followed by dehydration of the oxime (VII) by treating with an excess of acetic anhydride and heating in an inert, high-boiling solvent such as xylene, and, finally, partial hydrolysis of the nitrile (VIII) in concentrated sulfuric acid. In the cases where it is desired that R^5 and R^6 be other than hydrogen, the amide IX may be further hydrolyzed to the corresponding acid in 16% alcoholic KOH and the acid may be condensed with the appropriate amine using procedures well known in the art.


The compounds of formula X (formula I wherein Q is $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{R}^7$) may be synthesized by acylation of the amine V with an excess of a lower-alkyl acid chloride, or aryl acid chloride in a suitable solvent,



preferably pyridine, optionally in the presence of a cosolvent, preferably dichloromethane, at a temperature in the range from about 0°C to about 50°C.

The compounds of formula XI (formula I
5 wherein Q is $\text{CH}_2\text{N}(\text{R}^4)\text{P}(\text{O})(\text{O-lower-alkyl})_2$) may be synthesized by treatment of the amine V with an excess of an appropriate di-lower-alkyl phosphorochloridate in a suitable solvent, such as dichloromethane, in the presence of an excess of a base, preferably triethyl-
10 amine, at a temperature in the range of from about 0°C up to about 50°C.

Scheme C



This Sheet Has Been Replaced



The compounds of formula XII (formula I wherein Q is $\text{CH}_2\text{N}=\text{CHN}(\text{R}^9)(\text{R}^{10})$) may be synthesized by the treatment of the amine of formula V with an excess of a dimethyl acetal of formula $(\text{MeO})_2\text{CHN}(\text{R}^9)(\text{R}^{10})$, at a temperature of about 60°C .

The compounds of formula XIII (formula I wherein Q is $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{CF}_3$) may be synthesized by treatment of the amine V, in a suitable solvent, preferably dichloromethane, with an excess of a solution of a trifluoroacetyl halide of formula $\text{CF}_3\text{C}(\text{O})\text{X}$ wherein X is halogen, preferably trifluoroacetyl chloride, in a suitable solvent, preferably toluene, at a temperature of about 0°C .

The compounds of formula XIV (formula I wherein Q is $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{OR}^7$) may be synthesized by treatment of the amine V with an excess of an appropriate haloformate of formula $\text{R}^7\text{OC}(\text{O})\text{X}$ wherein X is halogen, preferably chlorine, in the presence of an excess of a base, preferably triethylamine, in a suitable solvent, such as dichloromethane or chloroform, at a temperature in the range of about 0°C up to about room temperature.

The aldehyde II is available by the method disclosed in U.S. Patent 3,294,803, by MnO_2 oxidation of the alcohol obtained by the method of U.S. Patent 3,711,512, or by the methods described hereinafter in the examples.

Simple chemical transformations which are conventional and well known to those skilled in the art of chemistry can be used for effecting changes in the functional groups of the compounds of the invention, for example, dealkylation of lower-alkyl aryl ethers to produce the corresponding phenol derivatives.

The compounds of formula I are useful both in the free base form and in the form of acid-addition salts, and both forms are within the purview of the invention. The acid-addition salts are in some cases a more



convenient form for use, and in practice the use of the salt form inherently amounts to the use of the base form. The acids which can be used to prepare the acid-addition salts include preferably those which produce, when combined with the free base, medicinally acceptable salts, that is, salts whose anions are relatively innocuous to the animal organism in medicinal doses of the salts so that the beneficial properties inherent in the free base are not vitiated by side effects ascribable to the anions. In practising the present invention it is convenient to form the hydrochloride, fumarate, toluenesulfonate, methanesulfonate, or maleate salts. However, other appropriate medicinally acceptable salts within the scope of the invention are those derived from other mineral acids and organic acids. The acid-addition salts of the basic compounds are prepared either by dissolving the free base in aqueous alcohol solution containing the appropriate acid and isolating the salt by evaporating the solution, or by reacting the free base and an acid in an organic solvent, in which case the salt separates directly, is precipitated with a second organic solvent, or can be obtained by concentration of the solution. Although medicinally acceptable salts of the basic compounds are preferred, all acid-addition salts are within the scope of the present invention. All acid-addition salts are useful as sources of the free base form even if the particular salt per se is desired only as an intermediate product, as, for example, when the salt is formed only for purposes of purification or identification, or when it is used as an intermediate in preparing a medicinally acceptable salt by ion exchange procedures.

The structures of the compounds of the invention were established by the mode of synthesis and by one or more of elemental analysis, and infrared, ultraviolet



and nuclear magnetic resonance spectroscopy. The course of the reactions and the identity and homogeneity of the products were assessed by thin layer chromatography (TLC) or gas-liquid chromatography (GLC). The melting points are given in degrees C and are uncorrected. The starting materials are either commercially available or may be prepared by procedures well known in the art.

The invention will now be illustrated with reference to the following examples but is in no way to be construed as limited thereto.

Example 1

1-[[2-(Diethylamino)ethyl]amino]-4-(N-phenylformimidoyl)thioxanthen-9-one

(I: $R^1=R^2=Et$; $Q=CH=N-C_6H_5$; $R^8=H$; $n=2$)

A mixture of 17.7 g (50 mmol) 1-[[2-(diethylamino)ethyl]amino]-9-oxo-thioxanthen-4-carboxyaldehyde and 15.1 g (150 mmol) aniline in 100 ml toluene was refluxed for 8 hours with a Dean-Stark trap. TLC on alumina with chloroform/hexane/isopropylamine 10:10:2 indicated incomplete reaction. The toluene was distilled off, 25 ml aniline was added and the mixture refluxed for 4 hours. 50 ml xylene was added and the reaction refluxed again for 3 hours. The solvent and excess aniline were removed in vacuo and the residue recrystallized from benzene to yield 19.9 g of crude product. This was recrystallized from approximately 1.5 l hexane to yield 15.8 g (86%) of product, m.p. 125-126°.

Example 2

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]formamide

(I: $R^1=R^2=Et$; $Q=CH_2NHCHO$; $R^8=H$; $n=2$)

A solution of 35.4 g (0.1 mol) 1-[[2-(diethylamino)ethyl]amino]-9-oxo-thioxanthen-4-carboxyaldehyde, 420 ml formamide and 50 ml (1 mol) formic acid was



heated at 160° for 1 hour. The reaction was cooled, poured into 2 l water and made basic with about 50 ml 35% sodium hydroxide solution. The gummy precipitate was filtered off and dried in vacuo. The dried

5 precipitate was dissolved in about 1.5 l hot ethyl acetate, treated with charcoal, and crystallized by cooling. The product was filtered off, washed with ethyl acetate and dried to provide 29.0 g (75%) of product, m.p. 154-155°.

10 Example 3

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-methylformamide

(IV: R¹=R²=Et; R⁴=Me; R⁸=H; n=2)

By a procedure analogous to that of Example 2,
15 24.6 g of the N-methylformamide was prepared from 35.4 g (0.1 mol) of 1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-carboxyaldehyde, 394 g of N-methylformamide and 50 ml formic acid. The product was recrystallized from 150 ml acetone to a m.p. of
20 127-130°.

Example 4

4-(Aminomethyl)-1-[[2-(diethylamino)ethyl]amino]-thioxanthen-9-one

(I: R¹=R²=Et; Q=CH₂NH₂; R⁸=H; n=2)

25 A solution of 24.4 g (64 mmol) of the formamide of Example 2 in 240 ml 2N hydrochloric acid was heated on a steam bath for 1 hour. The reaction was cooled to room temperature, made basic with 35% aqueous sodium hydroxide, and the resulting yellow precipitate
30 collected by filtration. The product was dissolved in benzene, treated with charcoal, dried with magnesium sulfate, filtered and azeotroped to remove traces of water. The dried residue was crystallized from methanol and isopropanol by the addition of ethereal hydrogen
35 chloride. The resulting solid was recrystallized in



several crops from methanol to yield 10.6 g of product, m.p. 270-272°, as the dihydrochloride salt.

Example 5

1-[[2-(Diethylamino)ethyl]amino]-4-[(methylamino)-methyl]thioxanthen-9-one

(I: $R^1=R^2=Et$; $Q=CH_2NHCH_3$; $R^8=H$; $n=2$)

By a process precisely analogous to that of Example 4, 10.5 g of the methylamine was obtained as the dihydrochloride hemihydrate from 14.6 g (37 mmol) of the N-methylformamide of Example 3 and 150 ml 2N hydrochloric acid. The product melted at 241-243°.

Example 6

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]methanesulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2NHSO_2CH_3$; $R^8=H$; $n=2$)

A solution of 10.65 g (30 mmol) of the free base of the amine of Example 4 in 100 ml pyridine was cooled in an ice bath and 4 g (35 mmol) of methanesulfonyl-chloride was added in one portion. The mixture was stirred for 2 hours at room temperature and poured into 750 ml water containing 2 g sodium hydroxide. The dark yellow precipitate was collected, washed with water and dried in vacuo overnight. A second crop was obtained by adding excess sodium hydroxide to the filtrate and filtering the resulting solid. The combined precipitates after drying were recrystallized from benzene to yield 6.4 g of the methanesulfonamide, m.p. 169-170°.

Example 7

1-[[2'-(Diethylamino)ethyl]amino]-9-oxothioxanthene-4-carboxamide

(I: $R^1=R^2=Et$; $Q=CONH_2$; $R^8=H$; $n=2$)

A suspension of 74g (0.23 mol) 1-[[2-(diethyl-amino)ethyl]amino]-9-oxo-thioxanthen-4-carboxaldehyde and 74 g (1.06 mol) hydroxylamine hydrochloride in



400 ml pyridine and 400 ml ethanol was refluxed 0.5 hour and 70 ml water was added to provide a homogeneous solution. The solution was heated for a further 2 hours and allowed to sit at room temperature 14 hours. The resulting crystalline oxime was filtered off to provide a quantitative yield, mp 215-218°.

123g of the oxime was heated briefly on a steambath in 180 ml acetic anhydride to achieve solution. The solution was cooled, 100 ml 1.8 M HCl in ether was added and the resulting suspension was diluted with 500 ml ether. The suspension was allowed to sit 14 hours at 0° and filtered. The residue (123g, mp 109-112°) was slurried in 250 ml xylene and refluxed 20 min. The mixture was cooled and 71.3g of the nitrile was filtered off, mp 265°.

10g nitrile was stirred in 200 ml conc. H₂SO₄ at room temperature for 3 days. The reaction was neutralized with conc. NH₄OH and the residue filtered off. The residue was digested in warm EtOAc/EtOH, filtered and the product crystallized from the chilled solution, mp 241-243°. It was dissolved in ethanol and one equivalent of HCl in ethanol was added. 6g of the amide hydrochloride was obtained, mp 271-272°.

Example 8

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]N-methylmethanesulfonamide

(I: R¹=R²=Et; Q=CH₂N(CH₃)SO₂CH₃; R⁸=H; n=2)

A solution of 1.5 g (3.5 mmol) methanesulfonamide of Example 6 in THF (60 ml) was cooled to 0°C in an ice-bath and NaH 0.16 g (4.0 mmol) was added. The reaction mixture was warmed to room temperature, stirred for 10 minutes, then methyl iodide 0.25 ml (4.0 mmol) was added. The reaction mixture was stirred at room temperature for 24 hours and the solvent was removed in vacuo. The residue was purified by column chromatography on silica eluting with chloroform (100%)



then 1% isopropylamine/chloroform to afford 1.15 g (74%) of the N-methylmethanesulfonamide as a yellow powder, m.p. 175-177°C. The free base was also treated with methanesulfonic acid in methanol to afford the methanesulfonate salt, m.p. 194-195.5°C (labelled Example 8a hereinafter).

Example 9

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]phenylsulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2NHSO_2Ph$; $R^8=H$; $n=2$)

Following a procedure substantially similar to that described in Example 6, 2.4 g (57%) of the phenylsulfonamide was obtained as the methanesulfonic acid salt from 2.54 g (7.15 mmol) of the free base of the amine of Example 4, pyridine (50 ml) and benzenesulfonyl chloride (1.1 ml, 8.62 mmol), followed by treatment with methanesulfonic acid in methanol. The product was recrystallized from ethanol.

Example 10

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]acetamide

(I: $R^1=R^2=Et$; $Q=CH_2NHC(O)CH_3$; $R^8=H$; $n=2$)

Following a procedure substantially similar to that described in Example 6, 2.3 g (52%) of the acetamide was obtained as an orange solid from 4.15 g (11.7 mmol) of the free base of the amine of Example 4, pyridine (60 ml) and acetyl chloride (0.82 ml, 11.53 mmol). The product was recrystallized from acetone and melted at 182-183°C.

Example 11

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]benzamide

(I: $R^1=R^2=Et$; $Q=CH_2NHC(O)Ph$; $R^8=H$; $n=2$)

Following a procedure substantially similar to that described in Example 6, 1.02 g (68%) of the benzamide was obtained as a yellow powder from 1.17 g



(3.29 mmol) of the free base of the amine of Example 4, pyridine (25 ml) and benzoyl chloride (0.42 ml, 3.62 mmol). The product was purified by column chromatography on silica eluting with chloroform (100%) to 1% isopropylamine/chloroform, followed by recrystallization from ethyl acetate. The product melted at 161-163°C.

Example 12

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]diethyl phosphoramidate

(I: $R^1=R^2=Et$; $Q=CH_2NHP(O)(OEt)_2$; $R^8=H$; $n=2$)

A solution of 2.28 g (6.41 mmol) of the free base of the amine of Example 4, CH_2Cl_2 (50 ml), and triethylamine (2 ml) at 0°C was treated with diethyl phosphorochloridate (1.0 ml, 6.9 mmol). The reaction mixture was stirred at 0°C for 2 hours, then at room temperature for 1 hour. The solvent was removed *in vacuo* and the residue was purified by column chromatography on silica eluting with ethyl acetate (100%), then 5% methanol/-ethyl acetate and finally methanol/isopropylamine/ethyl acetate (5/5/90) to afford 2.28 g (72%) of the diethyl phosphoramidate as a yellow solid, m.p. 108-110°C when recrystallized from ethyl acetate.

Example 13

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-ethylformamide

(IV: $R^1=R^2=Et$; $R^4=Et$; $R^8=H$; $n=2$)

A solution of 2.0 g (5.6 mmol) of 1-[[2-(diethylamino)ethyl]amino]-9-oxo-thioxanthen-4-carboxaldehyde, N-ethylformamide (24.0 ml) and formic acid (3.0 ml, 79.5 mmol) was heated at 170°C for 4 hours. The reaction mixture was cooled, poured into water and made basic with 10% sodium hydroxide. A solid was obtained which was collected by filtration and washed with water. The solid residue was taken up in chloroform/water, and the organic layer was separated and dried over Na_2SO_4 .



The solvent was removed in vacuo and the residue was purified by radial chromatography eluting with isopropylamine/methanol/ethyl acetate (0.5/1/98.5) to afford 1.32 g (57%) of the N-ethylformamide as an orange solid, m.p. 75-77°C.

Example 14

1-[[2-(Diethylamino)ethyl]amino]-4-[(ethylamino)-methyl]thioxanthen-9-one

(I: $R^1=R^2=Et$; $Q=CH_2NHC_2H_5$; $R^8=H$; $n=2$)

By a process substantially similar to that described in Example 4, 1.29 g (92%) of the ethylamine was obtained as the dihydrochloride from 1.3 g (3.2 mmol) of the N-ethylformamide of Example 13 and 10.8 ml 2N hydrochloric acid. The product was recrystallized from ethanol/tetrahydrofuran and melted at 160°C (dec.).

Example 15

1-[[2-(Diethylamino)ethyl]amino]-4-(dimethylamino-methyleneaminomethyl)thioxanthen-9-one trihydrochloride

(I: $R^1=R^2=Et$; $Q=CH_2N=CHN(Me)_2$; $R^8=H$; $n=2$)

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]formamide (3g) was diluted with 50 ml 2N HCl and the solution was heated on a steam bath for 90 min. The mixture was cooled, basified to pH 10 with 35% sodium hydroxide solution, and extracted into chloroform. The organic layer was separated, filtered through K_2CO_3 , and concentrated in vacuo and the resulting crude product was allowed to react with dimethylformamide dimethyl acetal overnight at 60°C. Excess DMF-dimethyl acetal was removed in vacuo and the desired title compound was purified by flash chromatography (silica gel; chloroform/*i*PrNH₂/MeOH (98:1:1). This product was dissolved in 2.5 M HCl/EtOH (100 ml), cooled in an ice-bath, filtered, and dried to afford 2.38 g of 1-[[2-(diethylamino)ethyl]amino]-



4-(dimethylaminomethyleneaminomethyl)-thioxanthen-9-one trihydrochloride as an orange solid, m.p. 258-260°C.

Example 16

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]trifluoroacetamide

(I: $R^1=R^2=Et$; $Q=CH_2NHC(O)CF_3$; $R^8=H$; $n=2$)

A solution of 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]thioxanthen-9-one (2.91 g; 8.19 mmol) in 80 ml methylene chloride at 0°C was treated with trifluoroacetyl chloride (14.75 ml 0.61 M solution in toluene; 9.0 mmol) and the reaction mixture was stirred at 0°C for 90 min. The mixture was concentrated in vacuo and the residue was purified by flash chromatography (silica gel; EtOAc (100%), then 2% isopropylamine/EtOAc) and then recrystallization from ethyl acetate to afford 2.52 g (68%) of the product as the free base, m.p. 189-190°C (Example 16). The free base was dissolved in methanol and treated with methanesulfonic acid (0.55 g, 5.72 mmol) to afford the methanesulfonate salt, m.p. 152-154°C after recrystallization from acetone (Example 16a).

Example 17

(a)

A mixture of thiosolicyclic acid (50.14 g, 0.33 mol) and cupric acetate (5.0 g) in DMSO (500 ml) was brought to reflux and potassium carbonate (54.3 g) was added in portions. 3-Bromochlorobenzene (42 ml, 0.36 mol) was then added via syringe and the mixture was refluxed for 3 hours. The reaction mixture was poured into water, treated with charcoal and filtered through celite. The filtrate was acidified with concentrated HCl and the precipitate which formed was collected by filtration, washed with water and dried in vacuo at 60°C to afford 75.01 g (86%) of 2-[(3-chlorophenyl)thio]-benzoic acid.



(b)

To a stirred solution of concentrated H_2SO_4 at 0°C was added 2-[(3-chlorophenyl)thio]benzoic acid (75.00 g, 0.28 mol) in portions over 1 hour. The mixture was stirred for 2 hours, poured into concentrated NH_4OH (500 ml) in water (2.5 l) and the precipitate which formed was collected by filtration, washed with water and dried in vacuo at 60°C to afford 65.9 g (95%) of a mixture of 1-chloro and 3-chlorothioxanthen-9-one.

(c)

A mixture of 1-chloro and 3-chlorothioxanthen-9-one (14.01 g, 56.8 mmol), pyridine (20 ml) and diethylaminopropylamine (5.13 g, 39.4 mmol) was refluxed until the reaction was complete. The heat was removed, the solvent was removed in vacuo and the residue was taken up in chloroform and purified by column chromatography on silica eluting with chloroform to remove the unreacted 3-chloroisomer and then 5% isopropylamine/chloroform to afford 5.10 g (54%) of 1-[[3-(diethylamino)propyl]amino]thioxanthen-9-one as an orange gum.

(d)

A mixture of 1-[[3-(diethylamino)propyl]amino]-thioxanthen-9-one (5.10 g, 15.0 mmol), formalin (160 ml) and 5N acetic acid (0.8 ml) was heated to 90°C 16 hours, additional 5N acetic acid (0.20 ml) was added, followed by formalin (50 ml) and the mixture was heated at 90°C for approximately 57 hours. The mixture was diluted with water, basified with 5N NaOH and extracted with chloroform. The organic layer was dried over Na_2SO_4 and passed through a silica column eluting with 2% methanol/chloroform and then isopropylamine/methanol/-chloroform (2/2/96) to afford 3.82 g (69%) of 1-[[3-(diethylamino)propyl]amino]-4-(hydroxymethyl)thioxanthen-9-one as an orange/brown gum.

This Sheet Has Been Replaced



(e)

1-[[3-(Diethylamino)propyl]amino]-9-oxothio-
xanthen-4-carboxaldehyde

(II: $R^1=R^2=Et$; $R^8=H$; $n=3$)

5 A mixture of 1-[[3-(diethylamino)propyl]amino]-
4-(hydroxymethyl)thioxanthen-9-one (3.82 g), toluene
(60 ml) and manganese oxide (7.5 g) was refluxed for
6.5 hours. The mixture was cooled to room temperature,
filtered through celite, and the filtrate was
10 concentrated in vacuo to afford 3.3 g (87%) of the
product as a brown oil.

(f)

1-[[3-(Diethylamino)propyl]amino]-4-(methylaminomethyl)
thioxanthen-9-one dihydrochloride 3/2 hydrate

15 (I: $R^1=R^2=Et$; $Q=CH_2NHMe$; $R^8=H$; $n=3$)

A solution of 1-[[3-(diethylamino)propyl]amino]-
9-oxothioxanthen-4-carboxaldehyde (3.3 g, 8.96 mmol) and
3 g of formic acid in 50 ml N-methylformamide was
allowed to reflux for 2 h. The mixture was basified
20 with 5 ml 5N sodium hydroxide solution and extracted
into chloroform (3x150ml). The organic layer was dried
over sodium sulfate, concentrated in vacuo, and a crude
oil was dissolved in a 3N aqueous HCl solution (50 ml)
and heated on a steam bath for 3 h. The above mixture
25 was cooled, basified with 30 ml 35% NaOH, extracted into
chloroform (3x150ml), the organic layer dried over
sodium sulfate and concentrated in vacuo to afford a
brown oil. The brown oil was purified by flash
chromatography (silica gel; 5% triethylamine/Et₂O, then
30 5% Et₃N/EtOAc, then triethylamine/methanol/EtOAc
(5:5:90)) to afford 1.1 g of 1-[[3-(diethylamino)-
propyl]amino]-4-(methylaminomethyl)thioxanthen-9-one, as
a clear orange gum. The above gum was converted into
the corresponding dihydrochloride by treatment with
35 6N HCl in ether to afford 1.04 g of the dihydro-
chloride·3/2 hydrate as a yellow powder, m.p. 222-224°C.



Example 18

(a)

A mixture of 1-chloro and 3-chlorothioxanthen-9-one (20 g), pyridine (40 ml) and N,N-dimethylethylene diamine (11 ml) was refluxed for approximately 22 hours and then the solvent was removed in vacuo. The residue was combined with the crude product (3.51 g) obtained from three similar experimental runs and the mixture was purified by column chromatography on silica eluting with chloroform, then 1% isopropylamine/chloroform and finally 2% isopropylamine/chloroform to afford 13.4 g of 1-[[2-(dimethylamino)ethyl]amino]thioxanthen-9-one.

(b)

A mixture of 1-[[2-(dimethylamino)ethyl]amino]-thioxanthen-9-one (13 g, 0.044 mol), 37% formalin (390 ml) and 5N acetic acid (6.5 ml) was heated to 100°C for 8.5 hours, allowed to stand over the weekend and then was heated at 100°C for several more hours. The mixture was poured into ice water, basified with 35% NaOH and extracted with chloroform. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column chromatography on silica eluting in succession with; 25% CHCl₃/hexane; 50% CHCl₃/hexane; 75% CHCl₃/hexane; 100% CHCl₃; 0.5% isopropylamine/chloroform; 1% isopropylamine/chloroform; 2% isopropylamine/chloroform; then 2% isopropylamine/2% MeOH/CHCl₃ to afford 9.2 g (64%) of 1-[[2-(dimethylamino)ethyl]amino]-4-(hydroxymethyl)thioxanthen-9-one.

(c)

1-[[2-(dimethylamino)ethyl]amino]-9-oxothioxanthen-4-carboxaldehyde

(Formula II: R¹=R²=Me; R⁸=H; n=2).

A mixture of 1-[[2-(dimethylamino)ethyl]amino]-4-(hydroxymethyl)thioxanthen-9-one (9.2 g, 0.028 mol) in toluene (322 ml) was heated to about 60°C and then



manganese oxide (MnO_2 , 16 g) was added and the mixture was heated at 60°C for 1 hour. The mixture was filtered and the filtrate was concentrated in vacuo to afford 7.9 g (87%) of the product.

5 (d)

N-[[1-[[2-(Dimethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]formamide

(I: $\text{R}^1=\text{R}^2=\text{Me}$; $\text{Q}=\text{CH}_2\text{NHCHO}$; $\text{R}^8=\text{H}$; $n=2$)

A mixture of 1-[[2-(dimethylamino)ethyl]amino]-
10 9-oxothioxanthen-4-carboxaldehyde (4.75 g), formamide (66.5 ml) and formic acid (7.6 ml) was heated at 170°C for 4 hours. The mixture was poured into ice water (250 ml), basified with 35% NaOH and extracted with chloroform. The organic layer was washed with water
15 (2X), then brine (1X) and the solvent was dried over Na_2SO_4 and concentrated in vacuo to afford 6.2 g of the product.

(e)

4-(Aminomethyl)-1-[[2-(dimethylamino)ethyl]amino]-
20 9-oxothioxanthen-9-one dihydrochloride-1/2 hydrate

(I: $\text{R}^1=\text{R}^2=\text{Me}$; $\text{Q}=\text{CH}_2\text{NH}_2$; $\text{R}^8=\text{H}$; $n=2$)

A mixture of N-[[1-[[2-(dimethylamino)ethyl]-
amino]-9-oxothioxanthen-4-yl]methyl]formamide (6.2 g)
and 2N HCl (52 ml) was heated to 100°C for 1.5-2 hours.
25 The reaction mixture was poured into ice water, basified with 35% NaOH, and extracted with chloroform. The organic layer was washed with water (2X), then brine (1X), dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by column chromatography on silica
30 eluting with ethyl acetate, then 0.5% triethylamine/-EtOAc, then 2% triethylamine/EtOAc, then CHCl_3 /1-2% isopropylamine and finally CHCl_3 /1-2% isopropylamine/-2% MeOH to afford 3.3 g (58%) of the product as the free base. A portion of the free base (1.25 g) was dissolved
35 in methanol and treated with concentrated HCl (3.3 ml)



in MeOH (6 ml) to afford 1.2 g of the product as the dihydrochloride·1/2 hydrate, m.p. 213°C (dec.).

(f)

N-[[1-[[2-(Dimethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]methanesulfonamide methanesulfonate

(I: $R^1=R^2=Me$; $Q=CH_2NHSO_2Me$; $R^8=H$; $n=2$)

4-(Aminomethyl)-1-[[2-(dimethylamino)]ethyl]-amino]-9-oxothioxanthen-9-one (2 g, 6 mmol) in 30 ml dry pyridine under nitrogen was stirred at room temperature until the solution was complete. The solution was chilled in an ice-bath and 0.52 ml (6.7 mmol) methanesulfonyl chloride in chilled pyridine was added dropwise and the mixture was stirred for 1 h at room temperature. The reaction mixture was poured into 500 ml water containing 0.51 g sodium hydroxide, extracted into chloroform, the organic layer was washed with water (2x) and brine, and dried over anhydrous sodium sulfate. The mixture was filtered and concentrated in vacuo. The residue (2.5 g) was stirred in ether, filtered, and dried to afford 2 g of the product, m.p. 126-127°C. The free base was dissolved in MeOH, and treated with methanesulfonic acid (0.48 g) to afford 2.0 g (67%) of the product as the methanesulfonate salt, m.p. 168°C (dec.).

Example 19

(a)

Following a procedure similar to that described in Example 17(c), there was prepared 6.83 g of 1-[[3-(dimethylamino)propyl]amino]thioxanthen-9-one, from a mixture of 1-chloro and 3-chlorothioxanthen-9-one (15.15 g, 61.4 mmol), pyridine (20 ml) and dimethylaminopropylamine (6.01 g, 58.7 mmol).

(b)

Following a procedure similar to that described in Example 17(d), there was obtained 6.74 g (90%) of



product was dissolved in chloroform and filtered through silica gel (chloroform; 1% isopropylamine/chloroform) to afford 2.38 g of the desired amine as an orange gum. The product was converted into the corresponding hydrochloride salt by dissolution in MeOH and treatment with concentrated HCl to afford 0.98 g of the dihydrochloride monohydrate, m.p. 228-229°C.

Example 20

(a)

N-[[1-[[2-(dimethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-methylformamide

[Formula IV: $R^1=R^2=Me$; $R^4=Me$; $R^8=H$; $n=2$)

A mixture of 1-[[2-(dimethylamino)ethyl]amino]-9-oxothioxanthen-4-carboxaldehyde (4.75 g, 0.15 mol), N-methylformamide (48 ml) and formic acid (3.9 ml) was heated at 170°C for 4.5 hours and then was allowed to stand at room temperature for approximately 64 hours. The reaction mixture was poured into water (250 ml), basified with 35% NaOH, extracted with $CHCl_3$ (3X). The organic layer was separated, washed with water (2X) and then brine (1X) and was dried over Na_2SO_4 . The solvent was removed in vacuo to afford 5.75 g of the product.

(b)

1-[[2-(Dimethylamino)ethyl]amino]-4-[(methylamino)-methyl]thioxanthen-9-one dihydrochloride-5/4 hydrate

(I: $R^1=R^2=Me$; $Q=CH_2NHMe$; $R^8=H$; $n=2$)

By a process similar to that of Example 19(e), 1.8 g 1-[[2-(dimethylamino)ethyl]amino]-4-[(methylamino)methyl]thioxanthen-9-one was obtained from 5.7 g (15.4 mmol) of the corresponding N-methylformamide of Example 20(a) and 50 ml 2 N HCl, after purification of the free base by flash chromatography (silica gel; chloroform; then 0.5% isopropylamine/chloroform; then 1% isopropylamine/chloroform). The free base was converted into the corresponding dihydrochloride



5/4 hydrate salt by treating with conc. HCl in methanol to afford 1.8 g (30%) of the product, m.p. 177°C (dec.).

Example 21

(a)

5 N-[[1-[[3-(dimethylamino)propyl]amino]-9-oxothioxanthen-4-yl]methyl]formamide

(Formula I: $R^1=R^2=Me$; $Q=CH_2NHCHO$; $R^8=H$; $n=3$)

10 A solution of 1-[[3-(dimethylamino)propyl]amino]-9-oxothioxanthen-4-carboxaldehyde (3.6 g; 10.57 mmol) in 50 ml formamide containing 3.6 g of formic acid was refluxed for 1.5 h and then was allowed to stand at room temperature overnight. The reaction mixture was diluted with water (400ml), basified with 3 ml 5N NaOH solution, stirred rapidly for 30 min, and the precipitated solid
15 was filtered, washed with water, and dried, yielding 3.1 g (79%) of the product as a yellow powder.

(b)

4-(aminomethyl)-1-[[3-(dimethylamino)propyl]amino]-thioxanthen-9-one

20 (Formula I: $R^1=R^2=Me$; $Q=CH_2NH_2$; $R^8=H$; $n=3$)

A solution of the formamide of Example 21(a) (2.98 g; 8.07 mmol) in 40 ml 3N HCl was heated on a steam bath for 4 h, allowed to cool to room temperature, chilled in ice, and neutralized to pH 8 with 5 N NaOH
25 solution. The resulting heterogenous mixture was extracted into chloroform (5x100ml) and the organic layer was dried over sodium sulfate and filtered through a pad of silica gel (first 5% triethylamine/ether, then 1-5% isopropylamine/chloroform) to afford 2.3 g (83%) of
30 the product as a bright yellow oil.

This Sheet Has Been Replaced



(c)

N-[[1-[[3-(Dimethylamino)propyl]amino]-9-oxothioxanthen-4-yl]methyl]methanesulfonamide methanesulfonate-1/2 hydrate

5 (I: $R^1=R^2=Me$; $Q=CH_2NHSO_2Me$; $R^8=H$; $n=3$)

To an ice cold solution of the amine of Example 21(b) (2.2 g; 6.44 mmol) in pyridine was added methanesulfonyl chloride (0.51 ml; 6.59 mmol) and the resulting mixture was stirred at room temperature overnight. The mixture was diluted with chloroform and passed through a large plug of silica gel eluting with 5% triethylamine/EtOAc affording 1.32 g of N-[[1-[[3-(dimethylamino)propyl]amino]-9-oxothioxanthen-4-yl]methyl]methanesulfonamide as a yellow powder. The free base was dissolved in methanol (10 ml) and treated with methanesulfonic acid (0.31 g, 1 eq.) in methanol to afford 1.38 g of the methanesulfonate-1/2 hydrate salt as an orange solid, m.p. $>107^\circ C$.

Example 22

20 N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-methylethanesulfonamide methanesulfonate

(I: $R^1=R^2=Et$; $Q=CH_2N(CH_3)SO_2Et$; $R^8=H$; $n=2$)

A solution of 2.03 g (5.49 mmol) of 1-[[2-(diethylamino)ethyl]amino]-4-[(methylamino)methyl]-thioxanthen-9-one (prepared by the method described in Example 5) and triethylamine in 45 ml methylene chloride was cooled to $0^\circ C$ and treated with ethanesulfonyl chloride (0.74 g, 5.76 mmol). After 15 min at $0^\circ C$, the reaction mixture was stirred at room temperature for 72 h. The mixture was concentrated in vacuo, the residue dissolved in chloroform and purified by passing through a pad of silica gel (chloroform; then 1% triethylamine/chloroform) affording 2.43 g (96%) of N-[[1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-methylethanesulfonamide. The sulfonamide was recrystallized from ethyl acetate and treated with



methanesulfonic acid in isopropanol to afford the product as the methanesulfonate salt, m.p. 159-161°C.

Example 23

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl](p-methoxy)benzenesulfonamide methane-
sulfonate

(I: $R^1=R^2=Et$; $Q=CH_2NHSO_2C_6H_4-p-OMe$; $R^8=H$; $n=2$)

A solution of 1.40 g (3.94 mmol) of 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]thioxanthen-9-one (prepared by the method described in Example 4) in 30 ml chloroform containing 1.5 ml triethylamine was cooled to 0°C and treated with p-methoxybenzenesulfonyl chloride (0.83 g, 4.02 mmol). After 10 min at 0°C the reaction mixture was stirred at room temperature for 2 h. Chloroform was removed in vacuo, the residue dissolved in 100 ml methylene chloride containing 1 ml triethylamine and treated with additional p-methoxybenzenesulfonyl chloride (0.85 g) with stirring at room temperature. The mixture was concentrated in vacuo, the residue was purified by passing through a pad of silica gel (1% triethylamine/chloroform) affording 1.57 g of N-[[1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-p-methoxybenzenesulfonamide. The sulfonamide was treated with methanesulfonic acid (0.3 g) in isopropanol/isopropyl acetate/methanol to afford 1.07 g of the methanesulfonate salt, m.p. 133-137°C.

Example 24

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]ethanesulfonamide methanesulfonate

(I: $R^1=R^2=Et$; $Q=CH_2NHSO_2Et$; $R^8=H$; $n=2$)

A solution of 2.5 g of 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]thioxanthen-9-one (prepared by the method described in Example 4) in 30 ml pyridine was cooled in an ice bath for 15 min and 0.95 g of ethanesulfonyl chloride in 5 ml pyridine was added rapidly



dropwise and the reaction mixture was stirred at room temperature for 1 h. The mixture was poured into 75 ml water containing 0.75 g NaOH, extracted into chloroform, the organic layer was washed with water (2x) and brine, and dried over sodium sulfate. The mixture was concentrated in vacuo and the residue was stirred in ether and dried (40°C/0.1mm) to afford 1.7 g of N-[[1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]ethanesulfonamide, m.p. 105°C (dec.). The sulfonamide was dissolved in methanol and treated with methanesulfonic acid in methanol to afford 1.61 g (42%) of the methanesulfonate salt, m.p. 135°C (dec.).

Example 25

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-ethylmethanesulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2N(Et)SO_2Me$; $R^8=H$; $n=2$)

A solution of 2.10 g (5.48 mmol) of 1-[[2-(diethylamino)ethyl]amino]-4-[(ethylamino)methyl]-thioxanthen-9-one (prepared by the method of Example 14) in 30 ml methylene chloride was cooled to 0°C and treated with 2 ml triethylamine and methanesulfonyl chloride (0.7 ml) and the reaction mixture was stirred at room temperature for 6 h. The solvent was removed in vacuo, the residue dissolved in chloroform, and the solution was purified by passing through a pad of silica gel (eluting with chloroform followed by 2% triethylamine/chloroform). The yellow solid isolated was recrystallized from ethyl acetate and dried to afford 1.11 g (44%) of the product as a yellow powder, m.p. 172-176°C.



Example 26

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-3,4-dichlorobenzenesulfonamidemethanesulfonate-1/2 hydrate

5 (I: $R^1=R^2=Et$; $Q=CH_2NHSO_2C_6H_3$ -3,4-dichloro; $R^8=H$; $n=2$)

To a solution of 3,4-dichlorobenzenesulfonyl chloride (1.84 g, 7.5 mmol) in 35 ml dry pyridine was added 2.5 g (7 mmol) 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]thioxanthen-9-one (prepared by the method described in Example 4) under nitrogen and the reaction mixture was stirred at room temperature for 15 min. and then was allowed to stand for approximately 72 hours. The reaction mixture was poured into 75 ml water containing 0.75 g NaOH, and extracted into 15 chloroform. The organic layer was washed with water (2x) and brine, and dried over sodium sulfate. Chloroform was removed in vacuo, the residue was recrystallized from ethanol to afford 1.24 g of N-[[1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-20 4-yl]methyl]-3,4-dichlorobenzenesulfonamide, m.p. 95°C (dec.). The free base was dissolved in methanol and treated with methanesulfonic acid in methanol to afford the methanesulfonate-1/2 hydrate, m.p. 55°C (dec.).

Example 27

25 N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-2-fluorobenzenesulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2NHSO_2C_6H_4$ -2-F; $R^8=H$; $n=2$)

A solution of 1.36 g (3.83 mmol) 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]thioxanthen-9-one (prepared by the method described in Example 4) in 25 ml methylene chloride containing 1 ml triethylamine was cooled to 0°C and treated with 2-fluorobenzene-sulfonyl chloride (0.84 g; 4.32 mmol) and the reaction mixture was stirred for several hours. The solvent was removed in vacuo, the residue dissolved in chloroform, and purified by flash chromatography (silica gel:



chloroform followed by 1% triethylamine/chloroform). The solvent was removed in vacuo and the product was recrystallized from ethyl acetate as an orange powder, 1.08 g (55%), m.p. 125-127°C.

5 Example 28

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-propylmethanesulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2N(C_3H_7)SO_2Me$; $R^8=H$; $n=2$)

The oil from 0.2 g of 60% dispersion of sodium
10 hydride in mineral oil was removed by triturating with pentane (4x). Dry DMF (40 ml) was added under nitrogen to sodium hydride with stirring, and 2 g N-[[1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-methanesulfonamide (Example 6) was then added to the
15 reaction mixture while stirring under nitrogen and the mixture was heated to 50°C for 2 h. The above mixture was chilled in an ice bath for 15 min, 0.87 g of propyl iodide in a small volume of DMF was added, and the mixture was allowed to stir at room temperature
20 overnight. The mixture was stirred with 35 ml water, filtered, and the residue was washed with water and dried (50°C/0.1mm/ P_2O_5) to afford 2.17 g of the product, m.p. 142-143°C.

Example 29

25 N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-methyl-benzenesulfonamide methane-sulfonate

(I: $R^1=R^2=Et$; $Q=CH_2N(Me)SO_2C_6H_5$; $R^8=H$; $n=2$)

A solution of 5.32 g (14.4 mmol) of 1-[[2-(diethylamino)ethyl]amino]-4-[(methylamino)methyl]-
30 thioxanthen-9-one (prepared by the method described in Example 5) in 100 ml methylene chloride was cooled to 0°C and treated with triethylamine (5 ml) and benzenesulfonyl chloride (2 ml; 15.67 mmol) and the
35 reaction mixture was stirred for 2 h. The mixture was concentrated in vacuo, and the residue was purified by



passing through a pad of silica gel (eluting with
chloroform; then 1/2%-1% isopropylamine/chloroform)
yielding 6.24 g of a yellow gum. The product was
dissolved in ethyl acetate and the solvent removed
5 in vacuo affording 6.06 g (83%) of N-[[1-[[2-(diethyl-
amino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-
N-methylbenzenesulfonamide. The sulfonamide (2.5 g) was
suspended in isopropanol and treated with methane-
sulfonic acid (0.51 g) to yield 2.63 g of the
10 methanesulfonate salt, m.p. 171-174°C.

Example 30

(a)

To a mixture of m-anisic acid (250 g, 1.67 mol) in
acetic acid (1 l) was added bromine (85 ml) and then
15 water (1 l). The mixture was heated to reflux, cooled
in an ice bath and the product which precipitated was
collected by filtration and washed with water to afford
305.7 g (79%) of 2-bromo-5-methoxybenzoic acid,
m.p. 154-156°C.

20 (b)

To a mixture of 3-chlorothiophenol (20 g,
0.138 mol), and cupric acetate (1.8 g) and DMF (200 ml)
was added K₂CO₃ (23 g). The mixture was heated to 150°C
for 15-20 minutes, then 2-bromo-5-methoxybenzoic acid
25 (35.8 g, 0.155 mol) was added in portions. The mixture
was heated overnight, poured into water (600 ml),
filtered and the filtrate was treated with charcoal,
filtered and diluted with HCl. The resulting
precipitate was collected by filtration, washed with
30 water, and dried at 50°C in vacuo over P₂O₅ to afford
27.6 g 2-[(3-chlorophenyl)thio]-5-methoxybenzoic acid.

(c)

To cooled sulfuric acid (89 ml) under nitrogen was
added 2-[(3-chlorophenyl)thio]-5-methoxybenzoic acid



(27 g, 0.092 mol) in portions over 1.5-2 hours. The mixture was stirred at ambient temperature overnight, poured into water (900 ml) containing conc. NH_4OH (218 ml) and ice. The solid which precipitated was collected by filtration and dried at 50°C in vacuo over P_2O_5 to afford 21 g (42%) of a mixture of 1-chloro and 3-chloro-7-methoxy-thioxanthen-9-one.

(d)

A mixture of 1-chloro and 3-chloro-7-methoxy-thioxanthen-9-one (20.7 g), pyridine (69 ml) and diethylaminoethylamine (16.1 g, 0.138 mol) was heated at 115°C under N_2 for 20 hours. The solvent was removed in vacuo and the residue was purified by column chromatography on silica eluting with CHCl_3 (100%) and then 1% isopropylamine/chloroform to afford 11.22 g of 1-[[2-(diethylamino)ethyl]amino]-7-methoxythioxanthen-9-one.

(e)

A mixture of 1-[[2-(diethylamino)ethyl]amino]-7-methoxythioxanthen-9-one (11.2 g, 0.031 mol), 37% formaldehyde (277 mL) and 5N acetic acid (4.6 ml) was heated at 100°C for 3 hours. The reaction mixture was cooled, filtered, and the filtrate was poured into ice-water (600 ml) and made basic with 35% NaOH . The mixture was extracted with chloroform (3X), washed with brine and dried over Na_2SO_4 . The solvent was removed in vacuo and the residue was purified by column chromatography on silica eluting with 25% CHCl_3 /hexane, then 50% CHCl_3 /hexane, then 75% CHCl_3 /hexane, then 0.5% isopropylamine/ CHCl_3 to afford 8.8 g (73%) of 1-[[2-(diethylamino)ethyl]amino]-4-(hydroxymethyl)-7-methoxythioxanthen-9-one.



(f)

1-[[2-(diethylamino)ethyl]amino]-7-methoxy-9-oxothio-
xanthen-4-carboxaldehyde

(Formula II: $R^1=R^2=Et$; $R^8=7-OCH_3$; $n=2$)

5 A solution of 1-[[2-(diethylamino)ethyl]amino]-
4-(hydroxymethyl)-7-methoxythioxanthen-9-one (8.8 g,
0.023 mol) in toluene (268 ml) was heated to 60°C under
nitrogen and then MnO_2 (13.2 g) was added. The mixture
was heated overnight, filtered, and the filtrate was
10 concentrated in vacuo to afford 7.05 g (81%) of the
product.

(g)

N-[[1-[[2-(diethylamino)ethyl]amino]-7-methoxy-9-oxo-
thioxanthen-4-yl]methyl]-N-methylformamide

15 (Formula IV: $R^1=R^2=Et$; $R^4=Me$; $R^8=7-OCH_3$; $n=2$)

A solution of 1-[[2-(diethylamino)ethyl]amino]-
7-methoxy-9-oxothioxanthen-4-carboxaldehyde (3 g,
7.8 mmol) and 1.5 ml formic acid in 25.5 ml N-methyl-
formamide was heated to 170°C for 8 h with stirring
20 under nitrogen. The reaction mixture was poured into
160 ml ice/water, basified with 35% of NaOH solution,
and extracted into chloroform (3x). The organic layer
was washed with water (2x) and brine, dried over sodium
sulfate, and the solvent was removed in vacuo affording
25 3 g (89.9 %) of the product.

(h)

1-[[2-(Diethylamino)ethyl]amino]-4-[(methylamino)-
methyl]-7-methoxythioxanthen-9-one

(I: $R^1=R^2=Et$; $Q=CH_2NHMe$; $R^8=7-OMe$; $n=2$)

30 The N-methylformamide of Example 30(g) (3.0 g) in
a 2N aqueous HCl solution (24 ml) was heated at 100°C
for 2 h under nitrogen with stirring. The above mixture
was cooled, poured into 125 ml ice/water, basified with
35% NaOH solution, and was extracted into chloroform and
5 washed with water (2X), then brine. The organic layer
was dried over sodium sulfate and concentrated in vacuo



to afford 3.1 g of the crude product. The product was trituated in ether and the filtrate was purified by several flash chromatography columns (silica gel; eluting with 50% hexane/chloroform, then chloroform, and then 0.25-0.5% isopropylamine/chloroform (column 1); chloroform, then 1% isopropylamine/1% MeOH/CHCl₃ (column 2); and CHCl₃, then 0.5% isopropylamine/CHCl₃ (column 3)) affording 0.746 g of the product.

Example 31

(a)

N-[[1-[[2-(Diethylamino)ethyl]amino]-7-methoxy-9-oxo-thioxanthen-4-yl]methyl]formamide

(I: R¹=R²=Et; Q=CH₂NHCHO; R⁸=7-OCH₃; n=2)

A mixture of 1-[[2-(diethylamino)ethyl]amino]-7-methoxy-9-oxothioxanthen-4-carboxaldehyde (3.6 g, 0.0094 mol), formamide (48 ml) and formic acid (6 ml) was heated to 170°C under N₂ for 8 hours. The mixture was poured into ice-water, basified with 35% NaOH and extracted with chloroform. The organic layer was separated, washed with water (2X), then brine and was dried over Na₂SO₄ and concentrated in vacuo to afford 3.88 g of the product.

(b)

4-(Aminomethyl)-1-[[2-(diethylamino)ethyl]amino]-7-methoxythioxanthen-9-one

(I: R¹=R²=Et; Q=CH₂NH₂; R⁸=7-OCH₃; n=2)

A mixture of the formamide of Example 31(a) (3.88 g) and 2N HCl (32 ml) was heated at 100°C for 2 h under nitrogen with stirring. The above mixture was cooled, poured into water, basified with 10% NaOH solution, and extracted into chloroform and washed with water, then brine. The organic layer dried over sodium sulfate and concentrated in vacuo to afford 3.6 g of the crude product. The product was dissolved in chloroform and purified by flash chromatography (silica gel; eluting with hexane/chloroform (50:50) and then 1% isopropylamine in hexane/chloroform (50:50)) affording



1.75 g of the desired product.

(c)

N-[[1-[[2-(Diethylamino)ethyl]amino]-7-methoxy-9-oxothioxanthen-4-yl]methyl]methanesulfonamide

5 (I: $R^1=R^2=Et$; $Q=CH_2NHSO_2Me$; $R^8=7-OMe$; $n=2$)

To a solution of 1.75 g (0.0045 mol) of the amine of Example 31(b) in 22.5 ml pyridine cooled in an ice bath under nitrogen with stirring was added dropwise 0.39 ml (0.005 mol) methanesulfonyl chloride in a small volume of pyridine and the resulting reaction mixture was stirred at room temperature for 2 h. The mixture was poured into 375 ml water containing 0.38 g NaOH, extracted into chloroform, and the organic layer was washed with water and brine. The chloroform layer was dried over sodium sulfate, the solvent removed in vacuo, and the residue was dried in vacuo to afford 1.61 g (77%) of the product, m.p. 144°C (dec.).

Example 32

(a)

20 To a mixture of 3-chlorothiophenol (20 g, 0.138 mol), cupric acetate (1.75 g) and DMF (199 ml) under N_2 was added in portions K_2CO_3 (23 g). The mixture was heated to 150°C and then 2,5-dibromobenzoic acid (43.5 g) was added. The mixture was heated overnight, poured into water (600 ml), filtered, the filtrate was treated with charcoal and filtered again. The filtrate was acidified with conc. HCl, extracted with $CHCl_3$, the organic layer was washed with brine and dried over Na_2SO_4 . The solvent was removed in vacuo to afford 28.9 g of 2-[(3-chlorophenyl)thio]-5-bromobenzoic acid.

(b)

35 A mixture of 2-[(3-chlorophenyl)thio]-5-bromobenzoic acid (28.4 g) and conc. sulfuric acid (80 ml) was stirred at 0°C and then at room temperature overnight. The mixture was poured into ice-water



(850 ml) containing conc. NH_4OH (199 ml) and the product which precipitated was collected by filtration and dried at 50°C in vacuo to afford 15.0 g of a mixture of 1-chloro and 3-chloro-7-bromothioxanthen-9-one.

5 (c)

A mixture of 1-chloro and 3-chloro-7-bromothioxanthen-9-one (13.6 g), pyridine (108 ml) and N,N-diethylethylenediamine (16.3 ml) was heated to 115°C for 20 hours. The solvent was removed in vacuo, and the
10 residue was purified by column chromatography on silica eluting with CHCl_3 (100%), then 1% isopropyl-amine/ CHCl_3 to afford 9.3 g of 1-[[2-(diethylamino)-ethyl]amino]-7-bromothioxanthen-9-one.

(d)

15 A mixture of 1-[[2-(diethylamino)ethyl]amino]-7-bromothioxanthen-9-one (9.3 g, 22.9 mmol), 203 ml 37% formaldehyde solution, and 3.4 ml 5N acetic acid solution was heated to 100°C under nitrogen overnight. The mixture was cooled to room temperature and the solid
20 formed was removed by filtration. The filtrate was diluted with water, basified with 35% NaOH solution and extracted into chloroform. The organic layer was washed with brine, dried over sodium sulfate, and the solvent was removed in vacuo to afford 10 g of oil. The above
25 oil in methylene chloride was filtered, the solvent concentrated in vacuo, and the crude hydroxymethyl analog was purified by flash chromatography (silica gel; eluting with 25% chloroform/hexane, then chloroform/-hexane (1:1), then 25% chloroform/hexane, then CHCl_3
30 (100%), and then 0.5-1% isopropylamine/chloroform) to afford 3.2 g of 1-[[2-(diethylamino)ethyl]amino]-4-(hydroxymethyl)-7-bromothioxanthen-9-one.

This Sheet Has Been Replaced



(e)

1-[[2-(Diethylamino)ethyl]amino]-7-bromo-9-oxothioxanthen-4-carboxaldehyde

(II: $R^1=R^2=Et$; $R^8=7-Br$; $n=2$)

5 A mixture of 3.2 g (7.34 mmol) of the alcohol of Example 32(d) and 4.3 g MnO_2 in 85 ml toluene was heated at 60°C for 1 h under nitrogen. The mixture was filtered, washed with $CHCl_3$, and the combined filtrate was concentrated in vacuo affording 3 g of a yellow
10 solid. The yellow solid was triturated in ether, filtered, and dried to afford 2.7 g (94.3%) of the product, m.p. 145-146°C.

(f)

15 N-[[1-[[2-(Diethylamino)ethyl]amino]-7-bromo-9-oxothioxanthen-4-yl]methyl]formamide

(I: $R^1=R^2=Et$; $Q=CH_2NHCHO$; $R^8=7-Br$; $n=2$)

A mixture of 2.7 g (6.2 mmol) 1-[[2-(diethylamino)ethyl]amino]-7-bromo-9-oxothioxanthen-4-carboxaldehyde, 31.7 ml formamide, and 3.6 ml formic
20 acid was heated to 170°C under nitrogen and with stirring for 8 h and the mixture was allowed to stand at room temperature for 72 h. The mixture was poured into 150 ml ice/water, basified with 35% NaOH solution, and the solid product was filtered and washed with water.
25 The solid product was dissolved in chloroform, washed with brine, dried over sodium sulfate, and the solvent was concentrated in vacuo to afford 2.85 g of the desired formamide, as a yellow/orange solid, m.p. 132°C (dec.).

30 (g)

1-[[2-(Diethylamino)ethyl]amino]-4-(aminomethyl)-7-bromothioxanthen-9-one

(I: $R^1=R^2=Et$; $Q=CH_2NH_2$; $R^8=7-Br$; $n=2$)

A mixture of 2.85 g (6.6 mmol) of the above
35 formamide (Example 32(f)) in 26 ml 2 N HCl solution was heated to 100°C under nitrogen for 2h and the mixture was allowed to stand at room temperature overnight. ~~The~~



reaction mixture was poured into 200 ml ice/water, basified with 35% NaOH solution, and extracted into chloroform. The organic layer was washed with water and brine, dried over sodium sulfate, and concentrated in vacuo to yield 2.67 g of a dark oil. The dark oil was purified by flash chromatography (silica gel; 1250 ml hexane/chloroform (1:1), and then 1% isopropylamine in hexane/chloroform (1:1)) to afford 1.87 g (70%) of the product, m.p. 79-82°C.

Example 33

N-[[1-[[2-(Diethylamino)ethyl]amino]-7-bromo-9-oxothioxanthen-4-yl]methyl]methanesulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2NHSO_2Me$; $R^8=7-Br$; $n=2$)

1-[[2-(Diethylamino)ethyl]amino]-4-(aminomethyl)-7-bromothioxanthen-9-one (1 g, 2.3 mmol) in 11.5 ml dry pyridine under nitrogen was stirred in an ice bath for 15 min and 0.2 ml (2.6 mmol) methanesulfonyl chloride in chilled pyridine was added dropwise and the mixture was stirred at room temperature. The reaction mixture was poured into 200 ml water, added 0.19 g sodium hydroxide in ice/water, and extracted into chloroform. The organic layer was washed with water (2x) and brine, and dried over anhydrous sodium sulfate. The mixture was filtered, concentrated in vacuo and the residue was stirred in ether, filtered, and dried to afford 1.02g of the product, m.p. 134-139°C.

Example 34

Methyl N-[[1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]carbamate

(I: $R^1=R^2=Et$; $Q=CH_2NHCOOMe$; $R^8=H$; $n=2$)

To a solution of 2.94 g (8.27 mmol) of 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]thioxanthen-9-one (prepared by the process described in Example 4) in 50 ml methylene chloride containing 5 ml triethylamine chilled to 0°C was added 0.7 ml (9.06 mmol) of methyl chloroformate and the mixture was stirred for



2.5 h. The solvent was removed in vacuo, the residue was suspended in chloroform and was purified by flash chromatography (silica gel; chloroform, then 1% isopropylamine/chloroform) affording 2.36 g (69%) of the product, as a yellow solid, m.p. 129-131°C.

Example 35

1-[[2-(Diethylamino)ethyl]amino]-4-[(methylamino)-methyl]-7-hydroxy-thioxanthen-9-one

(I: $R^1=R^2=Et$; $Q=CH_2NHMe$; $R^8=7-OH$; $n=2$)

10 A solution of 1.6 g (4 mmol) 1-[[2-(diethylamino)-ethyl]amino]-4-[(methylamino)methyl]-7-methoxythioxanthen-9-one (prepared by the process described in Example 30(h)) in 10 ml 48% HBr solution was heated to 110°C for 5 h. After cooling, the reaction mixture was
15 neutralized with saturated sodium bicarbonate and extracted into chloroform (3x100ml). A dark gum, insoluble in water or chloroform, was dissolved in methanol and combined with chloroform solution. The solvent was concentrated in vacuo to afford 1.67 g of a
20 dark orange solid. The orange solid product was purified by flash chromatography (silica gel; isopropylamine/methanol/chloroform (1:1:98) followed by a second silica column eluting with isopropylamine/-MeOH/ $CHCl_3$ (2:2:96)) affording 0.56 g (36%) of
25 the product, m.p. 167-169°C.

Example 36

Methyl N-[[1-[[2-(diethylamino)ethyl]amino]-7-methoxy-9-oxothioxanthen-4-yl]-methyl]carbamate

(I: $R^1=R^2=Et$; $Q=CH_2NHCOOMe$; $R^8=7-OMe$; $n=2$)

30 To a solution of 1.55 g of 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]-7-methoxythioxanthen-9-one (prepared as in Example 31(b)) in 40 ml chloroform containing 2 ml triethylamine chilled to 0°C was added 0.45 ml methyl chloroformate and the mixture was stirred
35 at room temperature for several hours. The solvent was removed in vacuo, the residue was purified by flash chromatography (silica gel; eluting with chloroform,



then 1% triethylamine in chloroform/hexane (1:1))
affording 1.2 g of the product which was recrystallized
from ethyl acetate to yield 0.79 g of a bright yellow
solid, m.p. 131-132°C.

5 Example 37

N-[[1-(2-Diethylamino)ethyl]amino]-7-hydroxy-9-oxothio-
xanthen-4-yl]methyl]methanesulfonamide.3/4 hydrate

(I: $R^1=R^2=Et$, $Q=CH_2NHSO_2CH_3$; $R^8=7-OH$; $n=2$)

To a solution of N-[[1-(2-(diethylamino)ethyl)-
10 amino]-7-methoxy-9-oxothioxanthen-4-yl]methyl]methane-
sulfonamide (0.5 g) (prepared as in Example 31 (c)) in
 CH_2Cl_2 (45 ml) at $-78^\circ C$ was added 1N BBr_3 in CH_2Cl_2
(1.75 ml). The mixture was warmed to room temperature,
stirred overnight, and then poured into ice-water
15 (250 ml) containing 35% NaOH (8 ml). The mixture was
acidified with dilute HCl, then basified with solid
 Na_2CO_3 and then was extracted with ethyl acetate. The
organic layer was separated, washed with brine, dried
over Na_2SO_4 and concentrated in vacuo. The residue was
20 purified by column chromatography on silica eluting with
5% MeOH/EtOAc to afford 0.28 g (58%) of product,
m.p. $78^\circ C$ (dec.).

Example 38

25 4-(Aminomethyl)-1-[[2-(diethylamino)ethyl]amino]-
6-methylthioxanthen-9-one


(I: $R^1=R^2=Et$; $Q=CH_2NH_2$; $R^8=6-CH_3$; $n=2$)

Following a procedure similar to that described in
Example 30, parts (a)-(f), but substituting 4-methyl-
benzoic acid for 3-methoxybenzoic acid in part (a), it
30 is contemplated that there can be prepared 1-[[2-(di-
ethylamino)ethyl]amino]-6-methyl-9-oxothioxanthen-
4-carboxaldehyde. It is further contemplated that the
latter can be converted into 4-(aminomethyl)-1-[[2-(di-
ethylamino)ethyl]amino]-6-methylthioxanthen-9-one by
35 following procedures similar to those described in
Example 31, parts (a)-(b).



Examples 39-41

Following a procedure similar to that described in Example 34, but substituting, if necessary, an appropriate amine of the formula V for 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]thioxanthen-9-one it is contemplated that the following compounds of formula I can be prepared:



Example No.	R ¹	R ²	n	Q	R ⁸
39	C ₂ H ₅	C ₂ H ₅	2	CH ₂ NHC(O)OC ₆ H ₅	H
40	C ₂ H ₅	C ₂ H ₅	2	CH ₂ N(CH ₃)C(O)O-4-CH ₃ -C ₆ H ₄	7-OCH ₃
41	C ₂ H ₅	C ₂ H ₅	2	CH ₂ NHC(O)O-2-NO ₂ -C ₆ H ₄	6-CH ₃

Example 42

1-[[2-Diethylamino)ethyl]amino]-4-[(methylamino)methyl]-6-methylthioxanthen-9-one

(I: R¹=R²=Et; Q=CH₂NHCH₃; R⁸=6-CH₃; n=2)

Following a procedure similar to that described in Example 30, parts (g)-(h) but substituting 1-[[2-(diethylamino)ethyl]amino]-6-methyl-9-oxothioxanthen-4-carboxaldehyde for 1-[[2-(diethylamino)ethyl]amino]-7-methoxy-9-oxo-thioxanthen-4-carboxaldehyde, the above product can be prepared.



Example 43

N-[[1-[[2-(Diethylamino)ethyl]amino]-6-methyl-9-oxothioxanthen-4-yl]methyl]methanesulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2NHSO_2CH_3$; $R^8=6-CH_3$; $n=2$)

Following a procedure similar to that described in Example 6, it is contemplated that the above product can be prepared from 4-(amino-methyl)-1-[[2-(diethylamino)-ethyl]amino]-6-methylthioxanthen-9-one, pyridine and methanesulfonyl chloride.

Example 44

N-[[1-[[2-(Diethylamino)ethyl]amino]-7-methoxy-9-oxothioxanthen-4-yl]methyl]phenylsulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2NHSO_2Ph$; $R^8=7-CH_3O$; $n=2$)

Following a procedure similar to that described in Example 31, part (c), it is contemplated that the above product can be prepared from 4-(aminomethyl)-1-[[2-(diethylamino)ethyl]amino]-7-methoxythioxanthen-9-one, pyridine and benzenesulfonyl chloride.

Example 45

(a)

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-phenylformamide

(IV: $R^1=R^2=Et$; $R^4=Ph$; $R^8=H$; $n=2$)

A mixture of 1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-carboxaldehyde (3.40 g, 9.59 mmol), formanilide (31 g) and formic acid (3 ml) were heated to 160°C for 2 hours. The mixture was cooled, poured into water (200 ml) and extracted with ether (3 x 150 ml). The aqueous layer was basified with 5N NaOH, extracted with $CHCl_3$ (3 x 150 ml) and the combined organic extracts were dried over Na_2SO_4 and filtered through florisil eluting with $CHCl_3$ (100%), then 5% isopropylamine/ $CHCl_3$. The solvent was removed in vacuo and the residue was purified by column chromatography on silica eluting with 0.5% to 1% isopropylamine/ $CHCl_3$ and then by



recrystallization from benzene/hexane (2x) to afford 1.88 g of the desired product.

(b)

1-[[2-(Diethylamino)ethyl]amino]-4-[(phenylamino)methyl]thioxanthen-9-one

(V: $R^1=R^2=Et$; $R^4=Ph$; $R^8=H$; $n=2$)

A mixture of the N-phenylformamide of Example 45(a) (1.70 g, 3.70 mmol) and 3N HCl (100 ml) was heated on a steam bath for 1 hour. The mixture was cooled, basified with 5N NaOH, extracted with $CHCl_3$ (3 x 100 ml) and the combined organic layers were dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by column chromatography on silica eluting with ether (100%), then 5% isopropylamine/ether; followed by a second silica column eluting with 1% isopropylamine/-ether to afford 0.80 g of the desired product, m.p. 133-135°C.

(c)

N-[[1-[[2-(Diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]-N-phenylmethanesulfonamide

(I: $R^1=R^2=Et$; $Q=CH_2N(Ph)SO_2Me$; $R^8=H$; $n=2$)

Following a procedure similar to that described in Example 22 it is contemplated that the above product can be prepared from 1-[[2-(diethylamino)ethyl]amino]-4-[(phenylamino)methyl]thioxanthen-9-one of Example 45(b), methanesulfonyl chloride, methylene chloride and triethylamine.

It is contemplated that other members of the genus I may be made in a fashion analogous to that of Examples 1-14, substituting the appropriate 1-[[2-(di-alkylamino)ethyl]amino]- or 1-[[3-(dialkylamino)propyl]amino]-9-oxothioxanthen-4-carboxaldehyde for 1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-carboxaldehyde, or in a fashion analogous to that of Examples 15-45 but utilizing an appropriate 1-[[2-(di-alkylamino)alkyl]amino]-9-oxothioxanthen-



4-carboxaldehyde. Many of the aldehydes and their precursors are described in U.S. Patent 3,294,803.

Representative examples of the invention were
5 tested for antitumor activity in mice according to the following procedure:

The animals were pooled, implanted subcutaneously with 30 to 60 mg tumor fragments by 12-gauge trocar, and again pooled before unselective distribution to the
10 various treatment and control groups. For early-stage treatment, chemotherapy was started within 1 to 5 days after tumor implantation while the number of cells was relatively small (10^7 to 10^8 cells). For advanced-stage treatment, chemotherapy was delayed until the tumor
15 became relatively large (200 to 300 mg in size). A 300 mg tumor contains approximately 3×10^8 total cells. Tumors within a given advanced-stage trial were within a 2.5-fold size range for 90% of the animals. Tumors were measured with a caliper weekly (or twice weekly for the
20 more rapidly growing tumors). Mice were sacrificed when their tumors reached 1500 mg (i.e., before they could cause the animal discomfort). Tumor weights were estimated from two-dimensional measurements.

The treatment and control groups were measured
25 when the control group tumors reached approximately 700 to 1200 mg in size (median of group). The median tumor weight of each group was determined (including zeros). The T/C value (weight of treated tumors over the weight of control tumors) in % is an indication of
30 antitumor effectiveness: A T/C equal to or less than 42% is considered significant antitumor activity by the Drug Evaluation Branch of the Division of Cancer Treatment (NCI). A T/C value $<10\%$ is considered to indicate highly significant antitumor activity. A body weight
35 loss nadir (mean of group) of greater than 20% or



greater than 20% drug-deaths is considered to indicate an excessively toxic dosage.

The results are shown in Table I for pancreatic ductal adenocarcinoma #03 and in Table 2 for colon
5 adenocarcinoma #38.

This Sheet Has Been Replaced



TABLE 1

Example #	T/C (%)	Weight Loss (g)*	Drug Deaths	Total Dose (mg/kg) i.v. or p.o.
1	0	1.6	0	1739
2	0	2.0	0	576
2	7	1.6	0	144
4	0	0.4	0	570
5	0	1.6	0	222
6	0	1.6	0	124
7	0	3.2	1/5	400
8	0	0.8	0	304
9	8	1.6	0	1395
10	0	2.4	0	540
11	0	0.8	0	855
12	36	3.2	0	1298
13	0	2.4	0	431
14	0	1.2	0	448
15	0	2.0	0	390
16(a)	21	+0.8	0	1171
17(f)	17	2.0	0	1060
18(e)	82	1.0	0	128
18(f)	0	5.2	2/5	256
19(e)	4	3.4	0	610
20(b)	10	0.4	0	383
21(c)	0	4.5	0	208
22	0	3.2	0	465
23	20	2.4	0	1212
24	0	2.3	0	203
25	0	4.6	0	288
26	13	2.2	0	654
27	5	+0.8	0	2594
28	0	2.8	0	552
29	6	2.4	0	2403

This Sheet Has Been Replaced



TABLE 1 (cont'd)

<u>Example #</u>	<u>T/C (%)</u>	<u>Weight Loss (g)*</u>	<u>Drug Deaths</u>	<u>Total Dose (mg/kg) i.v. or p.o.</u>
31(c)	0	5.6	0	248
30(h)	0	1.4	0	880
33	28	2.0	0	1281
34	0	3.6	0	248
36	0	0.4	0	155
37	7	0	0	32

5 *Average body weight was 25 g.

This Sheet Has Been Replaced



The compound of Example 6 was tested by bolus intravenous injection against a number of other tumors as shown in Table 4.

5 The compound of Example 8(a) was tested against a number of tumors as shown in Table 5.

The compound of Example 36 was tested against a number of tumors as shown in Table 6.

10 Representative compounds of the invention were tested against Mammary Adenocarcinoma 16/C/RP as shown in Table 7.

Representative compounds of the invention were tested against P388/adriamycin resistant leukemia as shown in Table 8.

This Sheet Has Been Replaced



TABLE 3

Tumor	Total Dosage mg/kg	Schedule	Wt. Loss at Nadir g/mouse	T/C Value in %	Tumor Free	Days of Observation
Human Adeno Squamous Lung #125	960	16 hr infusion days 7,20	-0.4	16%	1/4	33
			-0.4	16%	0/4	132
	600		-0.4	15%	0/5	33
			-0.4	15%	0/5	132
Mammary 16/C	720	4 hr infusion days 1,4	-2.0	18%	0/7	13
	454		-1.2	16%	0/7	13
Mammary 16/C/Adr*	960	3 hr infusion days 1,4	-2.8	23%	0/7	27
Colon Adenocarcinoma #38	732	3-15 bolus	-2.0	0%	1/5	100
	477		-2.8	9%	0/5	100
Colon Adenocarcinoma #38	960	4 hr infusion days 6,13	-2.9	11%	2/7	20
			-2.9	11%	1/7	100
	600		-2.0	26%	0/7	20
			-2.0	26%	0/7	100
Colon #51	720	3 hr infusion days 3,7	-1.1	8%	0/5	50
	454		-2.8	12%	0/7	50
Panc 03	222	3 hr infusion	-1.6	0	1/4	108

*adriamycin resistant


This Sheet Has Been Replaced

TABLE 4

<u>Tumor</u>	<u>Total Dosage mg/kg</u>	<u>Schedule</u>	<u>Wt. Loss at Nadir g/mouse</u>	<u>T/C Value in %</u>	<u>Tumor Free</u>	<u>Days of Observation</u>
Colon Adenocarcinoma #38	82	days 6-9	-1.7	1	2/7	132
	51	bolus	-1.2	18	1/7	132
Mammary 16/C	82	days 1-4	-3.0	6	0/5	23
	51	bolus	-1.2	13	0/5	23
Colon 51/A	96	days 3-6,9,11	-3.0	14	0/6	32
	66	bolus	-2.3	28	0/6	32
Panc 02	82	days 1-4	-2.0	23	0/5	28
	51	bolus	-1.2	42	0/5	28
Panc 03	124	days 3-4, 6-14	-1.6	0	3/5	245
	79	bolus	-1.0	0	1/5	245



This Sheet Has Been Replaced

Table 5

Tumor	Total Dosage mg/kg	Drug Route	Schedule	Wt. Loss at Nadir g/mouse	T/C Value in %	Drug Deaths	Tumor Free	(Days of Observation)
Colon Adenocarcinoma #38	340	i.p.	days 3-10	-2.0	0	0	5/5	168
	196	i.p.	days 3-10	-0.0	0	0	1/5	168
	112	i.p.	days 3-10	+0.4	3	0	- -	168
	550	p.o.	days 3-7	-2.8	0	0	5/5	168
	275	p.o.	days 3-7	-0.4	0	0	3/5	168
Colon 51/A	500	p.o.	days 3-7	-4.6	0	4/5	0/5	31
	250	p.o.	days 3-7	-1.6	16	0	0/5	31
Mammary 16/C/Adr*	340	i.v.→p.o.	i.v. 2X/day days 1-4; p.o. day 5	-7.1	32	5/6	0/6	22
	284	i.v.→p.o.	i.v. 2X/day days 1-4; p.o. days 5-6	-2.3	61	3/6	0/6	22
	183	i.v.→p.o.	i.v. 2X/day days 1-4; p.o. days 5-6	-4.3	39	0/4	0/4	22

*Anti-fungal resistant

This Sheet Has Been Replaced


Table 6

Tumor	Total Dosage mg/kg	Drug Route(a)	Schedule	Wt. Loss at Nadir g/mouse	T/C Value in %	Drug Deaths	Tumor Free	Days of Observation
Panc 03	420	i.v.	days 3-6	-5.6	0	3/5	2/5	45
	240	i.v.	days 3-6	-0.4	0	0	5/5	45
	155	i.v.	days 3-6	-0.4	0	0	5/5	45
Colon Adenocarcinoma #38	330	i.v.	days 3-7	-3.2	0	0	5/5	33
	220	i.v.	days 3-7	-0.8	0	0	5/5	33
	147	i.v.	days 3-7	0	0	0	5/5	33
Mammary 16/C	363	i.v.	days 1-4	-2.4	2	2/5	2/5	21
	242	i.v.	days 1-4	-1.2	8	1/5	0/5	21
	161	i.v.	days 1-4	-0.4	9	0	0/5	21
Human MX-1 Mammary	297	i.v.	days 1-6	-2.4	39	0	0/5	26
	132	i.v.	days 1-6	-0.8	27	0	0/5	26

(a): Note some injections were changed to p.o. after 2-4 days of injections due to tail vein damage.

This Sheet Has Been Replaced



Table 7

Example No.	Total Dosage mg/kg	Drug Route	Schedule	Wt. Loss at Nadir g/mouse	T/C Value in %	Drug Deaths	Tumor Free	Days of Observation
30(H)	900	i.v. infusion	infusion days 1,4	0	29	1/6	0/5	63
	600	i.v. infusion	infusion days 1,4	-0.7	23	0/5	0/5	63
	400	i.v. infusion	infusion days 1,4	-1.0	46	0/5	0/5	63
31(C)	224	i.v.	2X/day, days 3-6	-4.4	0	0	1/5	28
	156.8	i.v.	2X/day, days 3-6	-1.6	5	0	0/5	28
	109.6	i.v.	2X/day, days 3-6	-1.6	6	0	1/5	28

This Sheet Has Been Replaced

Table 8

Example No.	# of P388/Adr cells implanted i.v. on day 0	Total Dosage mg/kg	Drug Route	Schedule	Wt. Loss at Nadir g/mouse	Drug Deaths	% ILS	Log ₁₀ Tumor cell kill	Tumor Cell	Days Observa
10	10 ⁵	520	i.v. 3 hour infusion	days 1,4	-3.75	3/7	54	3.5	0/7	129
	10 ⁵	320	i.v. 3 hour infusion	days 1,4	-1.75	0/5	23	1.5	0/5	129
	10 ⁵	200	i.v. 3 hour infusion	days 1,4	-1.5	0/7	8	0.5	0/7	129
13	10 ⁵	450	i.v. 3 hour infusion	days 1,4	-3.0	2/7	38	2.5	0/7	129
	10 ⁵	300	i.v. 3 hour infusion	days 1,4	-1.75	0/7	15	1.0	0/7	129
	10 ⁵	150	i.v. 3 hour infusion	days 1,4	-1.75	0/8	8	0.5	0/8	129
30(H)	10 ⁶	450	i.v. 3 hour infusion	day 1	-2.3	0/7	62	5.3	0/7	43



This Sheet Has Been Replaced

Table 8 (cont'd)

Example No.	# of P388/Adr cells implanted i.v. on day 0	Total Dosage mg/kg	Drug Route	Schedule	Wt. Loss at Nadir g/mouse	Drug Deaths	% ILS	Log ₁₀ Tumor cell kill	Tumor Cell	Days Observa
	10 ⁶	280	i.v. 3 hour infusion	day 1	-1.7	0/7	69	6.0	2/7	43
31(C)	10 ⁶	248	i.v.→p.o.*	days 1,4	-3.0	1/6	0	0	0/6	43
	10 ⁶	112	i.v.→p.o.*	days 1,4	-2.7	0/6	0	0	0/6	43
34	10 ⁶	320	i.v. 3.5 hour infusion	day 1	-3.6	5/5	toxic	LD ₁₀₀	0/5	23
		198	i.v. 3.5 hour infusion	day 1	-1.2	0/5	41.7	3.3	0/5	23
		123	i.v. 3.5 hour infusion	day 1	-1.2	0/5	8.3	0.7	0/5	23
		76	i.v. 3.5 hour infusion	day 1	-0.4	0/5	0	0	0/5	23

*: p.o. injection day 4 due to tail vein damage.

%ILS = the percentage increase in lifespan.

This Sheet Has Been Replaced



The pharmaceutical compositions of the present invention include one or more of the compounds of this invention formulated into compositions together with one or more non-toxic physiologically acceptable carriers, adjuvants or vehicles which are collectively referred to herein as carriers, for parenteral injection, for oral administration in solid or liquid form, for rectal or topical administration, and the like.

The compositions can be administered to humans and animals either orally, rectally, parenterally (intravenously, intramuscularly or subcutaneously), intracisternally, intravaginally, intraperitoneally, locally (powders, ointments or drops), or as a buccal or nasal spray.

Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propylene glycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride and the like. Prolonged absorption of the



injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

If desired, and for more effective distribution, the compounds can be incorporated into slow release or targeted delivery systems such as polymer matrices, liposomes, and microspheres. They may be sterilized, for example, by filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, e.g. starches, lactose, sucrose, glucose, mannitol and silicic acid, (b) binders, e.g. carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia, (c) humectants, e.g. glycerol, (d) disintegrating agents, e.g. agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates and sodium carbonate, (e) solution retarders, e.g. paraffin, (f) absorption accelerators, e.g. quaternary ammonium compounds, (g) wetting agents, e.g. cetyl alcohol and glycerol monostearate, (h) adsorbents, e.g. kaolin and bentonite, and (i) lubricants, e.g. talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate or mixtures thereof. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents.

The percentage of active component in the composition and method for treating tumors or cancer can be varied so that a suitable dosage is obtained. The

dosage administered to a particular patient is variable depending upon the clinician's judgement using as the criteria: the route of administration, the duration of treatment, the size and condition of the patient, the
5 potency of the active component, and the patient's response thereto. An effective dosage amount of active component can thus readily be determined by the clinician considering all criteria and utilizing his best judgement on the patient's behalf.

This Sheet Has Been Replaced






5

CLAIMS:

1. A compound of formula



wherein: (1) n is 2 or 3;

R¹ and R² are independently lower-alkyl;

Q is a residue chosen from the group consisting of CH₂NHR³, CH₂N(R⁴)SO₂R⁷, CH₂NHCHO, CH=N-Ar, C(O)NR⁵R⁶, CH₂N(R⁴)C(O)R⁷, CH₂N(C₂H₅)CHO, CH₂N(R⁴)P(O)(O-lower-alkyl)₂, CH₂N=CH-N(R⁹)(R¹⁰), CH₂N(R⁴)C(O)CF₃ and CH₂N(R⁴)C(O)OR⁷;

R³ is hydrogen or lower-alkyl;

R⁴ is hydrogen, lower-alkyl or Ar;

R⁵ is hydrogen, lower-alkyl or Ar;

R⁶ is hydrogen or lower-alkyl;

R⁷ is lower-alkyl, or Ar;

R⁸ is hydroxy;

Ar is phenyl or phenyl substituted with methyl, methoxy, hydroxy, halogen or nitro; and

R⁹ and R¹⁰ are independently lower-alkyl; or

(2) Q is a residue chosen from the group consisting of CH₂N(R⁴)SO₂R⁷, CH=N-Ar, C(O)NR⁵R⁶, CH₂N(R⁴)C(O)R⁷, CH₂N(R⁴)P(O)(O-lower-alkyl)₂, CH₂N(R⁴)C(O)CF₃ and CH₂N(R⁴)C(O)OR⁷; R⁸ is hydrogen, lower-alkyl, lower-alkoxy, or hydroxy; Ar is phenyl substituted by hydroxy; and n, R¹, R², R⁴, R⁵, R⁶, and R⁷ are as defined hereinabove in part (1) with the proviso that one or more of R⁴, R⁵, or R⁷ is Ar; or

(3) Q is a residue chosen from the group consisting of CH₂N=CH-N(R⁹)(R¹⁰), CH₂N(R⁴)C(O)CF₃ and





$\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{OR}^7$; R^8 is hydrogen, lower-alkyl, lower-alkoxy, or hydroxy; and n , R^1 , R^2 , R^4 , R^7 , Ar , R^9 and R^{10} are as defined hereinabove in part (1); or a pharmaceutically acceptable acid-addition salt or solvate thereof;.

2. A compound as claimed in claim 1 wherein:

n is 2 or 3;

R^1 and R^2 are independently lower-alkyl;

Q is a residue chosen from the group consisting of

CH_2NHR^3 , $\text{CH}_2\text{N}(\text{R}^4)\text{SO}_2\text{R}^7$, CH_2NHCHO , $\text{CH}=\text{N}-\text{Ar}$,
 $\text{C}(\text{O})\text{NR}^5\text{R}^6$, $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{R}^7$, $\text{CH}_2\text{N}(\text{C}_2\text{H}_5)\text{CHO}$,
 $\text{CH}_2\text{N}(\text{R}^4)\text{P}(\text{O})(\text{O-lower-alkyl})_2$, $\text{CH}_2\text{N}=\text{CH}-\text{N}(\text{R}^9)(\text{R}^{10})$,
 $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{CF}_3$ and $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{OR}^7$;

R^3 is hydrogen or lower-alkyl;

R^4 is hydrogen, lower-alkyl or Ar ;

R^5 is hydrogen, lower-alkyl or Ar ;

R^6 is hydrogen or lower-alkyl;

R^7 is lower-alkyl, or Ar ;

R^8 is hydroxy;

Ar is phenyl or phenyl substituted with methyl, methoxy, hydroxy, halogen or nitro; and

R^9 and R^{10} are independently lower-alkyl;

or a pharmaceutically acceptable acid-addition salt or solvate thereof.

3. A compound as claimed in claim 2 wherein:

Q is a residue chosen from the group consisting of

CH_2NHR^3 , $\text{CH}_2\text{N}(\text{R}^4)\text{SO}_2\text{R}^7$, CH_2NHCHO , $\text{CH}=\text{N}-\text{Ar}$,
 $\text{C}(\text{O})\text{NR}^5\text{R}^6$, $\text{CH}_2\text{N}(\text{R}^4)\text{C}(\text{O})\text{R}^7$, $\text{CH}_2\text{N}(\text{C}_2\text{H}_5)\text{CHO}$, and
 $\text{CH}_2\text{N}(\text{R}^4)\text{P}(\text{O})(\text{O-lower-alkyl})_2$;

and R^4 is hydrogen or lower alkyl.

4. A compound as claimed in claim 3 wherein

Ar is phenyl or phenyl substituted with methyl, methoxy, halogen or nitro.

This Sheet Has Been Replaced



<



5. A compound as claimed in claim 4 wherein n is 2, both R^1 and R^2 are ethyl, R^3 is hydrogen or methyl and R^7 is lower-alkyl.

6. A compound as claimed in claim 5 selected from the group consisting of
1-[[2-(diethylamino)ethyl]amino]-4-[(methylamino)-methyl]-7-hydroxy-thioxanthen-9-one and
N-[[1-(2-(diethylamino)ethyl)amino]-7-hydroxy-9-oxothioxanthen-4-yl]methyl]methanesulfonamide.

7. A compound as claimed in claim 1 wherein
 n is 2 or 3;
 R^1 and R^2 are independently lower-alkyl;
 Q is a residue chosen from the group consisting of
 $CH_2N(R^4)SO_2R^7$, $CH=N-Ar$, $C(O)NR^5R^6$, $CH_2N(R^4)-C(O)R^7$, $CH_2N(R^4)P(O)(O-lower-alkyl)_2$,
 $CH_2N(R^4)C(O)CF_3$ and $CH_2N(R^4)C(O)OR^7$;
 R^4 is hydrogen, lower-alkyl or Ar ;
 R^5 is hydrogen, lower-alkyl or Ar ;
 R^6 is hydrogen or lower-alkyl;
 R^7 is lower-alkyl, or Ar ;
 R^8 is hydrogen, lower-alkyl, lower-alkoxy, or hydroxy; and

Ar is phenyl substituted by hydroxy; with the proviso that one or more of R^4 , R^5 or R^7 is Ar ;
or a pharmaceutically acceptable acid-addition salt or solvate thereof.

8. A compound as claimed in claim 1 wherein:
 n is 2 or 3;
 R^1 and R^2 are independently lower-alkyl;
 Q is a residue chosen from the group consisting of
 $CH_2N=CH-N(R^9)(R^{10})$, $CH_2N(R^4)C(O)CF_3$ and
 $CH_2N(R^4)C(O)OR^7$;
 R^4 is hydrogen, lower-alkyl or Ar ;
 R^7 is lower-alkyl, or Ar ;
 R^8 is hydrogen, lower-alkyl, lower-alkoxy, or hydroxy;

This Sheet Has Been Replaced





Ar is phenyl or phenyl substituted with methyl, methoxy, hydroxy, halogen or nitro; and R^9 and R^{10} are independently lower-alkyl; or a pharmaceutically acceptable acid-addition salt or solvate thereof.

9. A compound as claimed in claim 8 wherein R^8 is hydrogen, lower-alkyl, or lower-alkoxy; R^4 is hydrogen and Ar is phenyl or phenyl substituted with methyl, methoxy, halogen, or nitro.

10. A compound as claimed in claim 9 wherein n is 2 and both of R^1 and R^2 are ethyl and R^8 is hydrogen or methoxy.

11. A compound as claimed in claim 10 wherein Q is $\text{CH}_2\text{N}=\text{CH}-\text{N}(\text{R}^9)(\text{R}^{10})$ and R^9 and R^{10} are both methyl.

12. A compound as claimed in claim 10 wherein Q is $\text{CH}_2\text{NHC}(\text{O})\text{OR}^7$.

13. A compound as claimed in claim 12 wherein R^7 is lower-alkyl.

14. A compound as claimed in claim 13 selected from the group consisting of methyl N-[[1-[[2-(diethylamino)ethyl]amino]-9-oxothioxanthen-4-yl]methyl]carbamate and methyl N-[[1-[[2-(diethylamino)ethyl]amino]-7-methoxy-9-oxothioxanthen-4-yl]methyl]carbamate.

15. A pharmaceutical composition which comprises a compound as claimed in any one of claims 1 to 14 and a pharmaceutically acceptable carrier or diluent.

16. The use of a compound as claimed in any one of claims 1 to 14 for the preparation of a medicament for treating a tumor in a mammal.

17. The use of a composition as claimed in claim 15 for the preparation of a medicament for treating cancer in a mammal.

