

CD20 BINDING MOLECULES AND USES THEREOF**CROSS REFERENCE**

[0001] This application claims benefit of priority of International Patent Application No. PCT/CN2020/102463 filed on July 16, 2020, the content of which is incorporated herein by reference in its entirety.

SEQUENCE LISTING

[0002] This application incorporates by reference a Sequence Listing submitted with this application as a text file, entitled 14651-026-228_SEQ_LISTING.txt, created on July 9, 2021, and is 236,191 bytes in size.

1. FIELD

[0003] The present disclosure relates to anti-CD20 single domain antibodies, chimeric antigen receptors, engineered immune effector cells, and methods of use thereof. The present disclosure further relates to activation and expansion of cells for therapeutic uses, especially to chimeric antigen receptor-based T cell immunotherapies.

2. BACKGROUND

[0004] CD20 is a surface antigen expressed at certain stages of B-cell differentiation. Targeting the CD20-positive B cells with therapeutic monoclonal antibodies (mAbs) has been an effectual strategy in the treatment of hematologic malignancies such as non-Hodgkin's lymphoma (NHL) and chronic lymphocytic leukemia (CLL). Initial success with Rituximab (RTX) has encouraged the creation and development of more effective CD20 based therapeutics. However, treatment with conventional mAbs has not been adequate to overcome the problems such as refractory/relapsed diseases (Shanehbandi *et al.*, *Current Cancer Drug Targets*, 17(5): 423-444 (2007)).

[0005] Chimeric antigen receptor T (CAR-T) cell therapy is an emerging and effective cancer immunotherapy. Especially in hematological malignancies, CAR-T cells have achieved exciting results. Early clinical studies of CD20 CAR-T cells have been conducted, which indicate that CD20 CAR-T cells have encouraging therapeutic potential in B-cell NHLs (Brudno and Kochenderfer, *Nature Reviews Clinical Oncology*, 15: 31-46 (2018)). However, the application of CAR-T cells is obviously hampered by adverse effects, such as cytokines release syndrome and on-target off-tumor toxicity (Yu *et al.*, *Molecular Cancer*, 18 (1): 125 (2019)).

[0006] Thus, improved CD20-binding molecules and engineered CD20-targeting cells are needed. For example, there is a need to develop stable and small-sized CD20 binding molecules for use in more effective or efficient CAR-T therapies.

3. SUMMARY

[0007] In one aspect, provided herein is an anti-CD20 single domain antibody (sdAb) comprising

(i) a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 2; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3; (ii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 5; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 6; (iii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 7; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8; (iv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 9; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 10; (v) a CDR1 comprising the amino acid sequence of SEQ ID NO: 11 or 196; a CDR2 comprising the amino acid sequence of SEQ ID NO: 12; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8; (vi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 13; a CDR2 comprising the amino acid sequence of SEQ ID NO: 9; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 10; (vii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 14 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 15; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16; (viii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 17; a CDR2 comprising the amino acid sequence of SEQ ID NO: 18; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19; (ix) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 21; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16; (x) a CDR1 comprising the amino acid sequence of SEQ ID NO: 22; a CDR2 comprising the amino acid sequence of SEQ ID NO: 23; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19; (xi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 24 or 198; a CDR2 comprising the amino acid sequence of SEQ ID NO: 25; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 26; (xii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 27; a CDR2 comprising the amino acid sequence of SEQ ID NO: 28; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 29; (xiii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 14 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 30; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16; (xiv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 17; a CDR2 comprising the amino acid sequence of SEQ ID NO: 31; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19; (xv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 32; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3; (xvi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 6; (xvii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a

CDR2 comprising the amino acid sequence of SEQ ID NO: 34; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16; (xviii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 22; a CDR2 comprising the amino acid sequence of SEQ ID NO: 35; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19; (xix) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 36; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16; (xx) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 37; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16; (xxi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 38; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16; (xxii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 39; (xxiii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 40; (xxiv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 165 or 199; a CDR2 comprising the amino acid sequence of SEQ ID NO: 166; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 167; (xxv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 168; a CDR2 comprising the amino acid sequence of SEQ ID NO: 169; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 170; (xxvi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 171 or 200; a CDR2 comprising the amino acid sequence of SEQ ID NO: 172; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 173; (xxvii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 174; a CDR2 comprising the amino acid sequence of SEQ ID NO: 175; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 176; (xxviii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 177 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 178; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 179; or (xxix) a CDR1 comprising the amino acid sequence of SEQ ID NO: 180; a CDR2 comprising the amino acid sequence of SEQ ID NO: 181; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 182.

[0008] In another aspect, provided herein is an anti-CD20 single domain antibody (sdAb) comprising: (i) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 41; (ii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 42; (iii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 43; (iv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3,

respectively, as set forth in SEQ ID NO: 44; (v) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 45; (vi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 46; (vii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 47; (viii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 48; (ix) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 49; (x) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 50; (xi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 51; (xii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 52; (xiii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 53; (xiv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 54; (xv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 55; (xvi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 56; (xvii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 57; (xviii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 58; (xix) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 59; (xx) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 60; (xxi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 61; (xxii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 62; or (xxiii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 63; (xxiv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 183; (xxv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 184; or (xxvi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 185.

[0009] In some embodiments, the CDR1, CDR2 or CDR3 are determined according to the Kabat numbering scheme, the IMGT numbering scheme, the AbM numbering scheme, the Chothia numbering scheme, the Contact numbering scheme, or a combination thereof.

[0010] In some embodiments, the anti-CD20 sdAb provided herein further comprises one or more FR regions as set forth in SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 and/or SEQ ID NO: 185.

[0011] In some embodiments, the anti-CD20 sdAb provided herein comprises the amino acid sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 or SEQ ID NO: 185.

[0012] In other embodiments, the anti-CD20 sdAb provided herein comprises or consists of an amino acid sequence having at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or more sequence identity with the sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 or SEQ ID NO: 185.

[0013] In some embodiments, the anti-CD20 sdAb is a camelid sdAb. In some embodiments, the anti-CD20 sdAb is a humanized sdAb. In some embodiments, the anti-CD20 sdAb is a VHH domain. In some embodiments, the anti-CD20 sdAb is a humanized VHH domain.

[0014] In some embodiments, the anti-CD20 sdAb is genetically fused or chemically conjugated to an agent.

[0015] In another aspect, provided herein is a chimeric antigen receptor (CAR), comprising (a) an extracellular antigen binding domain comprising the anti-CD20 sdAb provided herein; (b) a transmembrane domain; and (c) an intracellular signaling domain. In some embodiments, the extracellular antigen binding domain further comprises one or more additional antigen binding domain(s). In some embodiments, the extracellular antigen binding domain further comprises one additional antigen binding domain. In other embodiments, the extracellular antigen binding domain further comprises two additional antigen binding domains. In some embodiments, the

one or more additional antigen binding domain(s) bind to one or more antigen(s) selected from a group consisting of CD20, CD19, CD22, CD33, CD38, BCMA, CS1, ROR1, GPC3, CD123, IL-13R, CD138, c-Met, EGFRvIII, GD-2, NY-ESO-1, MAGE A3, and glycolipid F77. In some embodiments, the CD20 CAR is monospecific. In some embodiments, the CD20 CAR is monovalent. In some embodiments, the CD20 CAR is multivalent. In some embodiments, the extracellular antigen binding domain comprises at least two anti-CD20 sdAbs provided herein.

[0016] In some embodiments, provided herein is a multivalent (such as bivalent and trivalent) CAR comprising a polypeptide comprising: (a) an extracellular antigen binding domain comprising two or more single domain antibodies (sdAbs) specifically binding to CD20; (b) a transmembrane domain; and (c) an intracellular signaling domain. In some embodiments, the extracellular antigen binding domain comprises two single domain antibodies (sdAbs) specifically binding to CD20 provided herein. In other embodiments, the extracellular antigen binding domain comprises three single domain antibodies (sdAbs) specifically binding to CD20 provided herein.

[0017] In some embodiments according to any one of the CARs described above, the transmembrane domain is derived from a molecule selected from a group consisting of CD8 α , CD4, CD28, CD137, CD80, CD86, CD152, and PD1. In some embodiments, the transmembrane domain is derived from CD8 α .

[0018] In some embodiments according to any one of the CARs described above, the intracellular signaling domain comprises a primary intracellular signaling domain of an immune effector cell. In some embodiments, the primary intracellular signaling domain is derived from CD3 ζ .

[0019] In other embodiments according to any one of the CARs described above, the intracellular signaling domain further comprises a co-stimulatory signaling domain. In some embodiments, the co-stimulatory signaling domain is derived from a co-stimulatory molecule selected from the group consisting of CD27, CD28, CD137, OX40, CD30, CD40, CD3, LFA-1, CD2, CD7, LIGHT, NKG2C, B7-H3, ligands of CD83 and combinations thereof. In some embodiments, the co-stimulatory signaling domain is derived from CD137.

[0020] In some embodiments, the CAR provided herein further comprises a hinge domain located between the C-terminus of the extracellular antigen binding domain and the N-terminus of the transmembrane domain. In some embodiments, the hinge domain is derived from CD8 α .

[0021] In some embodiments, the CAR provided herein further comprises a signal peptide located at the N-terminus of the polypeptide. In some embodiments, the signal peptide is derived from CD8 α .

[0022] In certain embodiments, provided herein is a chimeric antigen receptor (CAR), comprising (i) an amino acid sequence selected from the group consisting of SEQ ID NO: 81,

SEQ ID NO: 82, SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, SEQ ID NO: 95, SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 99, SEQ ID NO: 100, SEQ ID NO: 101, SEQ ID NO: 102, SEQ ID NO: 103, SEQ ID NO: 104, SEQ ID NO: 105, SEQ ID NO: 106, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191; or (ii) an amino acid sequence having at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or more sequence identity with the sequence of SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, SEQ ID NO: 95, SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 99, SEQ ID NO: 100, SEQ ID NO: 101, SEQ ID NO: 102, SEQ ID NO: 103, SEQ ID NO: 104, SEQ ID NO: 105, SEQ ID NO: 106, SEQ ID NO: 189, SEQ ID NO: 190, or SEQ ID NO: 191.

[0023] In another aspect, provided herein is an isolated nucleic acid comprising a nucleic acid sequence encoding the anti-CD20 sdAb provided herein. In another aspect, provided herein is a vector comprising a nucleic acid encoding the anti-CD20 sdAb provided herein.

[0024] In yet another aspect, provided herein is an isolated nucleic acid comprising a nucleic acid sequence encoding the CAR provided herein. In another aspect, provided herein is a vector comprising a nucleic acid encoding the CAR provided herein.

[0025] In yet another aspect, provided herein is an engineered immune effector cell, comprising the CAR, the isolated nucleic acid, or the vector provided herein. In some embodiments, the engineered immune effector cell is a T cell or B cell.

[0026] In another aspect, provided herein is a pharmaceutical composition, comprising the anti-CD20 sdAb, the engineered immune effector cell, or the vector provided herein, and a pharmaceutically acceptable excipient.

[0027] In yet another aspect, provided herein is a method of treating a disease or disorder in a subject, comprising administering to the subject an effective amount of the anti-CD20 sdAb, the engineered immune effector cell, or the pharmaceutical composition provided herein. In some embodiments, the disease or disorder is a B cell associated disease or disorder and/or CD20 associated disease or disorder. In some embodiments, the disease or disorder is cancer. In other embodiments, the disease or disorder is a B cell malignancy. In some embodiments, the B cell malignancy is a B cell leukemia or B cell lymphoma. In some embodiments, the disease or disorder is selected from a group consisting of marginal zone lymphoma (*e.g.*, splenic marginal zone lymphoma), diffuse large B cell lymphoma (DLBCL), mantle cell lymphoma (MCL), primary central nervous system (CNS) lymphoma, primary mediastinal B cell lymphoma (PMBL), small lymphocytic lymphoma (SLL), B cell prolymphocytic leukemia (B-PLL),

follicular lymphoma (FL), burkitt lymphoma, primary intraocular lymphoma, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), hairy cell leukemia (HCL), precursor B lymphoblastic leukemia, non-hodgkin lymphoma (NHL), high-grade B-cell lymphoma (HGBL), and multiple myeloma (MM). In other embodiments, the disease or disorder is an autoimmune and/or inflammatory disease. In some embodiments, the autoimmune and/or inflammatory disease is associated with inappropriate or enhanced B cell numbers and/or activation.

4. BRIEF DESCRIPTION OF THE FIGURES

[0028] **FIGs.1A-1B** show the exemplary transduction efficiency of VHH-based CAR-T cells (**FIG.1A**) and scFv-based CAR-T cells (**FIG.1B**). UnT refers to T cells un-transduced with CAR.

[0029] **FIGs.2A-2F** show *in vitro* cytotoxicity of VHH-based CAR-T cells compared to that of scFv-based CAR-T cells against CD20 positive cell lines (**FIGs.2A-2B** and **FIGs.2E-2F**) or CD20 negative cell lines (**FIGs.2C-2D**).

[0030] **FIGs.3A-3D** show *in vitro* cytotoxicity of mono-, bi- and tri-VHH CAR-T cells compared to that of CD20 scFv CAR-T cells against CD20 positive cell lines (**FIGs.3A-3B**) or CD20 negative cell lines (**FIGs.3C-3D**).

[0031] **FIGs.4A-4B** show *in vitro* cytotoxicity of bi-VHH CAR-T cells with different length of peptide linkers against Raji.Luc cells.

[0032] **FIGs.5A-5B** show IFN- γ release level of VHH-based CAR-T cells compared to that of scFv-based CAR-T cells after co-culture with Raji.Luc or K562-CD20.Luc cells at different E:T ratios for 24 hours.

[0033] **FIG.6** shows IFN- γ release level of mono-, bi- and tri-VHH CAR-T cells compared to that of CD20 scFv CAR-T cells after co-culture with Raji.Luc or K562.Luc cells for 24 hours.

[0034] **FIGs.7A-7D** show the proliferation capacity and CAR positive ratio of the exemplary CD20 VHH CAR-T cells compared to that of CD20 scFv CAR-T cells with or without antigen specific stimulation.

[0035] **FIG.8** shows the results of exhaustion markers assessment of exemplary CD20 VHH CAR-T cells and CD20 scFv CAR-T cells.

[0036] **FIGs.9A-9D** show the results of memory markers assessment of exemplary CD20 VHH CAR-T cells and CD20 scFv CAR-T cells.

[0037] **FIGs.10A-10F** show *in vivo* anti-tumor and re-challenging efficacy of VHH-273 CAR-T cells and VHH-496 CAR-T cells in a Raji xenograft NCG mouse model. Mice were assessed on a regular basis to monitor tumor growth by bioluminescence imaging (**FIGs.10A-10B**), and body weight (**FIG.10C**), as well as CAR copies in mouse peripheral blood (**FIG.10D**). Then, mice at least 5 weeks tumor-free with undetectable CAR copy number (copies/ng of genomic DNA) in

PBMC were selected for re-challenging assay. The Raji tumors were monitored weekly by bioluminescence imaging (FIG.10E) and the mice body weight was measured (FIG.10F).

[0038] FIGs.11A-11F show *in vivo* anti-tumor and re-challenging efficacy of mono-, bi- and tri-VHH CAR-T cells in a Raji xenograft NCG mouse model. Mice were assessed on a regular basis to monitor tumor growth by bioluminescence imaging (FIGs.11A-11B), and body weight (FIG.11C), as well as CAR copies in mouse peripheral blood (FIGs.11D-11E; FIG.11D is adopted from FIG.11E and the scale of Y-axis is adjusted to enhance the resolution of copies/ng genomic DNA). Then, mice at least 6 weeks tumor-free were selected for re-challenging assay. The Raji tumors were monitored weekly by bioluminescence imaging (FIG.11F).

[0039] FIGs.12A-12E show the results from studies assessing binding affinities of anti-CD20 VHH-huIgG1Fc mAbs. MFI=mean fluorescence intensity.

[0040] FIGs.13A-13K show the cytotoxicity of the exemplary humanized CD20 VHH CAR-T cells against Raji.Luc cells, K652-CD20.Luc cells, K562.Luc cells and Nalm.6.Luc at different effector cells to target cells ratios (E:T) of 15:1, 10:1, 5:1 or 2.5:1. FIGs.13A-13C show the cytotoxicity of huVHH-746 CAR-T cells, huVHH-750 CAR-T cells, and huVHH-753 CAR-T cells; FIGs.13D-13G show the cytotoxicity of huVHH-253 CAR-T cells and huVHH-256 CAR-T cells; FIGs.13H-13K show the cytotoxicity of huVHH-260 CAR-T cells.

5. DETAILED DESCRIPTION

[0041] The present disclosure is based in part on the novel single domain antibodies (*e.g.*, VHH domains) that bind to CD20, chimeric antigen receptors or engineered cells comprising same, and improved properties thereof.

5.1. Definitions

[0042] Techniques and procedures described or referenced herein include those that are generally well understood and/or commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized methodologies described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (3d ed. 2001); *Current Protocols in Molecular Biology* (Ausubel et al. eds., 2003); *Therapeutic Monoclonal Antibodies: From Bench to Clinic* (An ed. 2009); *Monoclonal Antibodies: Methods and Protocols* (Albitar ed. 2010); and *Antibody Engineering* Vols 1 and 2 (Kontermann and Dübel eds., 2d ed. 2010). Unless otherwise defined herein, technical and scientific terms used in the present description have the meanings that are commonly understood by those of ordinary skill in the art. For purposes of interpreting this specification, the following description of terms will apply and whenever appropriate, terms used in the singular will also include the plural and vice versa. In the event that any description of a term set forth conflicts with any document incorporated herein by reference, the description of the term set forth below shall control.

[0043] The term “antibody,” “immunoglobulin,” or “Ig” is used interchangeably herein, and is

used in the broadest sense and specifically covers, for example, monoclonal antibodies (including agonist, antagonist, neutralizing antibodies, full length or intact monoclonal antibodies), antibody compositions with polyepitopic or monoepitopic specificity, polyclonal or monovalent antibodies, multivalent antibodies, multispecific antibodies (*e.g.*, bispecific antibodies so long as they exhibit the desired biological activity), formed from at least two intact antibodies, single chain antibodies, and fragments thereof (*e.g.*, domain antibodies), as described below. An antibody can be human, humanized, chimeric and/or affinity matured, as well as an antibody from other species, for example, mouse, rabbit, llama, *etc.* The term “antibody” is intended to include a polypeptide product of B cells within the immunoglobulin class of polypeptides that is able to bind to a specific molecular antigen and is composed of two identical pairs of polypeptide chains, wherein each pair has one heavy chain (about 50-70 kDa) and one light chain (about 25 kDa), each amino-terminal portion of each chain includes a variable region of about 100 to about 130 or more amino acids, and each carboxy-terminal portion of each chain includes a constant region. *See, e.g., Antibody Engineering* (Borrebæck ed., 2d ed. 1995); and Kuby, *Immunology* (3d ed. 1997). Antibodies also include, but are not limited to, synthetic antibodies, recombinantly produced antibodies, single domain antibodies including from Camelidae species (*e.g.*, llama or alpaca) or their humanized variants, intrabodies, anti-idiotypic (anti-Id) antibodies, and functional fragments (*e.g.*, antigen-binding fragments) of any of the above, which refers to a portion of an antibody heavy or light chain polypeptide that retains some or all of the binding activity of the antibody from which the fragment was derived. Non-limiting examples of functional fragments (*e.g.*, antigen-binding fragments) include single-chain Fvs (scFv) (*e.g.*, including monospecific, bispecific, *etc.*), Fab fragments, F(ab') fragments, F(ab)₂ fragments, F(ab')₂ fragments, disulfide-linked Fvs (dsFv), Fd fragments, Fv fragments, diabody, triabody, tetrabody, and minibody. In particular, antibodies provided herein include immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, for example, antigen-binding domains or molecules that contain an antigen-binding site that binds to an antigen (*e.g.*, one or more CDRs of an antibody). Such antibody fragments can be found in, for example, Harlow and Lane, *Antibodies: A Laboratory Manual* (1989); *Mol. Biology and Biotechnology: A Comprehensive Desk Reference* (Myers ed., 1995); Huston *et al.*, 1993, *Cell Biophysics* 22:189-224; Plückthun and Skerra, 1989, *Meth. Enzymol.* 178:497-515; and Day, *Advanced Immunochemistry* (2d ed. 1990). The antibodies provided herein can be of any class (*e.g.*, IgG, IgE, IgM, IgD, and IgA) or any subclass (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2) of immunoglobulin molecule. Antibodies may be agonistic antibodies or antagonistic antibodies. Antibodies may be neither agonistic nor antagonistic.

[0044] An “antigen” is a structure to which an antibody can selectively bind. A target antigen may be a polypeptide, carbohydrate, nucleic acid, lipid, hapten, or other naturally occurring or

synthetic compound. In some embodiments, the target antigen is a polypeptide. In certain embodiments, an antigen is associated with a cell, for example, is present on or in a cell.

[0045] An “intact” antibody is one comprising an antigen-binding site as well as a CL and at least heavy chain constant regions, CH1, CH2 and CH3. The constant regions may include human constant regions or amino acid sequence variants thereof. In certain embodiments, an intact antibody has one or more effector functions.

[0046] “Single-chain Fv” also abbreviated as “sFv” or “scFv” are antibody fragments that comprise the VH and VL antibody domains connected into a single polypeptide chain. Preferably, the sFv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the sFv to form the desired structure for antigen binding. For a review of the sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

[0047] The term “heavy chain-only antibody” or “HCAb” refers to a functional antibody, which comprises heavy chains, but lacks the light chains usually found in 4-chain antibodies. Camelid animals (such as camels, llamas, or alpacas) are known to produce HCAs.

[0048] “Single domain antibody” or “sdAb” as used herein refers to a single monomeric variable antibody domain and which is capable of antigen binding (*e.g.*, single domain antibodies that bind to CD20). Single domain antibodies include VHH domains as described herein. Examples of single domain antibodies include, but are not limited to, antibodies naturally devoid of light chains such as those from Camelidae species (*e.g.*, llama), single domain antibodies derived from conventional 4-chain antibodies, engineered antibodies and single domain scaffolds other than those derived from antibodies. Single domain antibodies may be derived from any species including, but not limited to mouse, human, camel, llama, goat, rabbit, and bovine. For example, a single domain antibody can be derived from antibodies raised in Camelidae species, for example in camel, llama, dromedary, alpaca and guanaco, as described herein. Other species besides Camelidae may produce heavy chain antibodies naturally devoid of light chain; VHHs derived from such other species are within the scope of the disclosure. In some embodiments, the single domain antibody (*e.g.*, VHH) provided herein has a structure of FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4. Single domain antibodies may be genetically fused or chemically conjugated to another molecule (*e.g.*, an agent) as described herein. Single domain antibodies may be part of a bigger binding molecule (*e.g.*, a multispecific antibody or a chimeric antigen receptor).

[0049] The terms “binds” or “binding” refer to an interaction between molecules including, for example, to form a complex. Interactions can be, for example, non-covalent interactions including hydrogen bonds, ionic bonds, hydrophobic interactions, and/or van der Waals interactions. A complex can also include the binding of two or more molecules held together by

covalent or non-covalent bonds, interactions, or forces. The strength of the total non-covalent interactions between a single antigen-binding site on an antibody and a single epitope of a target molecule, such as an antigen, is the affinity of the antibody or functional fragment for that epitope. The ratio of dissociation rate (k_{off}) to association rate (k_{on}) of a binding molecule (*e.g.*, an antibody) to a monovalent antigen ($k_{\text{off}}/k_{\text{on}}$) is the dissociation constant K_D , which is inversely related to affinity. The lower the K_D value, the higher the affinity of the antibody. The value of K_D varies for different complexes of antibody and antigen and depends on both k_{on} and k_{off} . The dissociation constant K_D for an antibody provided herein can be determined using any method provided herein or any other method well known to those skilled in the art. The affinity at one binding site does not always reflect the true strength of the interaction between an antibody and an antigen. When complex antigens containing multiple, repeating antigenic determinants, such as a polyvalent antigen, come in contact with antibodies containing multiple binding sites, the interaction of antibody with antigen at one site will increase the probability of a reaction at a second site. The strength of such multiple interactions between a multivalent antibody and antigen is called the avidity.

[0050] In connection with the binding molecules described herein terms such as “bind to,” “that specifically bind to,” and analogous terms are also used interchangeably herein and refer to binding molecules of antigen binding domains that specifically bind to an antigen, such as a polypeptide. A binding molecule or antigen binding domain that binds to or specifically binds to an antigen can be identified, for example, by immunoassays, Octet[®], Biacore[®], or other techniques known to those of skill in the art. In some embodiments, a binding molecule or antigen binding domain binds to or specifically binds to an antigen when it binds to an antigen with higher affinity than to any cross-reactive antigen as determined using experimental techniques, such as radioimmunoassay (RIA) and enzyme linked immunosorbent assay (ELISA). Typically, a specific or selective reaction will be at least twice background signal or noise and may be more than 10 times background. See, *e.g.*, Fundamental Immunology 332-36 (Paul ed., 2d ed. 1989) for a discussion regarding binding specificity. In certain embodiments, the extent of binding of a binding molecule or antigen binding domain to a “non-target” protein is less than about 10% of the binding of the binding molecule or antigen binding domain to its particular target antigen, for example, as determined by fluorescence activated cell sorting (FACS) analysis or RIA. A binding molecule or antigen binding domain that binds to an antigen includes one that is capable of binding the antigen with sufficient affinity such that the binding molecule is useful, for example, as a therapeutic and/or diagnostic agent in targeting the antigen. In certain embodiments, a binding molecule or antigen binding domain that binds to an antigen has a dissociation constant (K_D) of less than or equal to 1 μM , 800 nM, 600 nM, 550 nM, 500 nM, 300 nM, 250 nM, 100 nM, 50 nM, 10 nM, 5 nM, 4 nM, 3 nM, 2 nM, 1 nM, 0.9 nM, 0.8 nM, 0.7 nM,

0.6 nM, 0.5 nM, 0.4 nM, 0.3 nM, 0.2 nM, or 0.1 nM. In certain embodiments, a binding molecule or antigen binding domain binds to an epitope of an antigen that is conserved among the antigen from different species.

[0051] In certain embodiments, the binding molecules or antigen binding domains can comprise “chimeric” sequences in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (*see* U.S. Pat. No. 4,816,567; and Morrison *et al.*, 1984, Proc. Natl. Acad. Sci. USA 81:6851-55). Chimeric sequences may include humanized sequences.

[0052] In certain embodiments, the binding molecules or antigen binding domains can comprise portions of “humanized” forms of nonhuman (*e.g.*, camelid, murine, non-human primate) antibodies that include sequences from human immunoglobulins (*e.g.*, recipient antibody) in which the native CDR residues are replaced by residues from the corresponding CDR of a nonhuman species (*e.g.*, donor antibody) such as camelid, mouse, rat, rabbit, or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, one or more FR region residues of the human immunoglobulin sequences are replaced by corresponding nonhuman residues. Furthermore, humanized antibodies can comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. A humanized antibody heavy or light chain can comprise substantially all of at least one or more variable regions, in which all or substantially all of the CDRs correspond to those of a nonhuman immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. In certain embodiments, the humanized antibody will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see, Jones *et al.*, Nature 321:522-25 (1986); Riechmann *et al.*, Nature 332:323-29 (1988); Presta, Curr. Op. Struct. Biol. 2:593-96 (1992); Carter *et al.*, Proc. Natl. Acad. Sci. USA 89:4285-89 (1992); U.S. Pat. Nos. 6,800,738; 6,719,971; 6,639,055; 6,407,213; and 6,054,297.

[0053] In certain embodiments, the binding molecules or antigen binding domains can comprise portions of a “fully human antibody” or “human antibody,” wherein the terms are used interchangeably herein and refer to an antibody that comprises a human variable region and, for example, a human constant region. The binding molecules may comprise a single domain antibody sequence. In specific embodiments, the terms refer to an antibody that comprises a variable region and constant region of human origin. “Fully human” antibodies, in certain

embodiments, can also encompass antibodies which bind polypeptides and are encoded by nucleic acid sequences which are naturally occurring somatic variants of human germline immunoglobulin nucleic acid sequence. The term “fully human antibody” includes antibodies having variable and constant regions corresponding to human germline immunoglobulin sequences as described by Kabat *et al.* (See Kabat *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). A “human antibody” is one that possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. Human antibodies can be produced using various techniques known in the art, including phage-display libraries (Hoogenboom and Winter, J. Mol. Biol. 227:381 (1991); Marks *et al.*, J. Mol. Biol. 222:581 (1991)) and yeast display libraries (Chao *et al.*, Nature Protocols 1: 755-68 (2006)). Also available for the preparation of human monoclonal antibodies are methods described in Cole *et al.*, Monoclonal Antibodies and Cancer Therapy 77 (1985); Boerner *et al.*, J. Immunol. 147(1):86-95 (1991); and van Dijk and van de Winkel, Curr. Opin. Pharmacol. 5: 368-74 (2001). Human antibodies can be prepared by administering the antigen to a transgenic animal that has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled, *e.g.*, mice (*see, e.g.*, Jakobovits, Curr. Opin. Biotechnol. 6(5):561-66 (1995); Brüggemann and Taussing, Curr. Opin. Biotechnol. 8(4):455-58 (1997); and U.S. Pat. Nos. 6,075,181 and 6,150,584 regarding XENOMOUSE™ technology). See also, for example, Li *et al.*, Proc. Natl. Acad. Sci. USA 103:3557-62 (2006) regarding human antibodies generated via a human B-cell hybridoma technology.

[0054] In certain embodiments, the binding molecules or antigen binding domains can comprise portions of a “recombinant human antibody,” wherein the phrase includes human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell, antibodies isolated from a recombinant, combinatorial human antibody library, antibodies isolated from an animal (*e.g.*, a mouse or cow) that is transgenic and/or transchromosomal for human immunoglobulin genes (*see, e.g.*, Taylor, L. D. *et al.*, *Nucl. Acids Res.* 20:6287-6295 (1992)) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies can have variable and constant regions derived from human germline immunoglobulin sequences (See Kabat, E. A. *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). In certain embodiments, however, such recombinant human antibodies are subjected to *in vitro*

mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

[0055] In certain embodiments, the binding molecules or antigen binding domains can comprise a portion of a “monoclonal antibody,” wherein the term as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *e.g.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts or well-known post-translational modifications such as amino acid isomerization or deamidation, methionine oxidation or asparagine or glutamine deamidation, each monoclonal antibody will typically recognize a single epitope on the antigen. In specific embodiments, a “monoclonal antibody,” as used herein, is an antibody produced by a single hybridoma or other cell. The term “monoclonal” is not limited to any particular method for making the antibody. For example, the monoclonal antibodies useful in the present disclosure may be prepared by the hybridoma methodology first described by Kohler *et al.*, *Nature* 256:495 (1975), or may be made using recombinant DNA methods in bacterial or eukaryotic animal or plant cells (*see, e.g.*, U.S. Pat. No. 4,816,567). The “monoclonal antibodies” may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature* 352:624-28 (1991) and Marks *et al.*, *J. Mol. Biol.* 222:581-97 (1991), for example. Other methods for the preparation of clonal cell lines and of monoclonal antibodies expressed thereby are well known in the art. *See, e.g.*, Short Protocols in Molecular Biology (Ausubel *et al.* eds., 5th ed. 2002).

[0056] A typical 4-chain antibody unit is a heterotetrameric glycoprotein composed of two identical light (L) chains and two identical heavy (H) chains. In the case of IgGs, the 4-chain unit is generally about 150,000 daltons. Each L chain is linked to an H chain by one covalent disulfide bond, while the two H chains are linked to each other by one or more disulfide bonds depending on the H chain isotype. Each H and L chain also has regularly spaced intrachain disulfide bridges. Each H chain has at the N-terminus, a variable domain (VH) followed by three constant domains (CH) for each of the α and γ chains and four CH domains for μ and ϵ isotypes. Each L chain has at the N-terminus, a variable domain (VL) followed by a constant domain (CL) at its other end. The VL is aligned with the VH, and the CL is aligned with the first constant domain of the heavy chain (CH1). Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains. The pairing of a VH and VL together forms a single antigen-binding site. For the structure and properties of the different classes of antibodies, *see, for example*, Basic and Clinical Immunology 71 (Stites *et al.* eds., 8th ed. 1994); and Immunobiology (Janeway *et al.* eds., 5th ed. 2001).

[0057] The term “Fab” or “Fab region” refers to an antibody region that binds to antigens. A conventional IgG usually comprises two Fab regions, each residing on one of the two arms of the Y-shaped IgG structure. Each Fab region is typically composed of one variable region and one constant region of each of the heavy and the light chain. More specifically, the variable region and the constant region of the heavy chain in a Fab region are VH and CH1 regions, and the variable region and the constant region of the light chain in a Fab region are VL and CL regions. The VH, CH1, VL, and CL in a Fab region can be arranged in various ways to confer an antigen binding capability according to the present disclosure. For example, VH and CH1 regions can be on one polypeptide, and VL and CL regions can be on a separate polypeptide, similarly to a Fab region of a conventional IgG. Alternatively, VH, CH1, VL and CL regions can all be on the same polypeptide and oriented in different orders as described in more detail the sections below.

[0058] The term “variable region,” “variable domain,” “V region,” or “V domain” refers to a portion of the light or heavy chains of an antibody that is generally located at the amino-terminal of the light or heavy chain and has a length of about 120 to 130 amino acids in the heavy chain and about 100 to 110 amino acids in the light chain, and are used in the binding and specificity of each particular antibody for its particular antigen. The variable region of the heavy chain may be referred to as “VH.” The variable region of the light chain may be referred to as “VL.” The term “variable” refers to the fact that certain segments of the variable regions differ extensively in sequence among antibodies. The V region mediates antigen binding and defines specificity of a particular antibody for its particular antigen. However, the variability is not evenly distributed across the 110-amino acid span of the variable regions. Instead, the V regions consist of less variable (*e.g.*, relatively invariant) stretches called framework regions (FRs) of about 15-30 amino acids separated by shorter regions of greater variability (*e.g.*, extreme variability) called “hypervariable regions” that are each about 9-12 amino acids long. The variable regions of heavy and light chains each comprise four FRs, largely adopting a β sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases form part of, the β sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site of antibodies (*see, e.g., Kabat et al., Sequences of Proteins of Immunological Interest* (5th ed. 1991)). The constant regions are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity (ADCC) and complement dependent cytotoxicity (CDC). The variable regions differ extensively in sequence between different antibodies. In specific embodiments, the variable region is a human variable region.

[0059] The term “variable region residue numbering according to Kabat” or “amino acid position numbering as in Kabat”, and variations thereof, refer to the numbering system used for heavy

chain variable regions or light chain variable regions of the compilation of antibodies in Kabat *et al.*, *supra*. Using this numbering system, the actual linear amino acid sequence may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, an FR or CDR of the variable domain. For example, a heavy chain variable domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 and three inserted residues (*e.g.*, residues 82a, 82b, and 82c, *etc.* according to Kabat) after residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a “standard” Kabat numbered sequence. The Kabat numbering system is generally used when referring to a residue in the variable domain (approximately residues 1-107 of the light chain and residues 1-113 of the heavy chain) (*e.g.*, Kabat *et al.*, *supra*). The “EU numbering system” or “EU index” is generally used when referring to a residue in an immunoglobulin heavy chain constant region (*e.g.*, the EU index reported in Kabat *et al.*, *supra*). The “EU index as in Kabat” refers to the residue numbering of the human IgG 1 EU antibody. Other numbering systems have been described, for example, by AbM, Chothia, Contact, IMGT, and AHon

[0060] The term “heavy chain” when used in reference to an antibody refers to a polypeptide chain of about 50-70 kDa, wherein the amino-terminal portion includes a variable region of about 120 to 130 or more amino acids, and a carboxy-terminal portion includes a constant region. The constant region can be one of five distinct types, (*e.g.*, isotypes) referred to as alpha (α), delta (δ), epsilon (ϵ), gamma (γ), and mu (μ), based on the amino acid sequence of the heavy chain constant region. The distinct heavy chains differ in size: α , δ , and γ contain approximately 450 amino acids, while μ and ϵ contain approximately 550 amino acids. When combined with a light chain, these distinct types of heavy chains give rise to five well known classes (*e.g.*, isotypes) of antibodies, IgA, IgD, IgE, IgG, and IgM, respectively, including four subclasses of IgG, namely IgG1, IgG2, IgG3, and IgG4.

[0061] The term “light chain” when used in reference to an antibody refers to a polypeptide chain of about 25 kDa, wherein the amino-terminal portion includes a variable region of about 100 to about 110 or more amino acids, and a carboxy-terminal portion includes a constant region. The approximate length of a light chain is 211 to 217 amino acids. There are two distinct types, referred to as kappa (κ) or lambda (λ) based on the amino acid sequence of the constant domains.

[0062] As used herein, the terms “hypervariable region,” “HVR,” “Complementarity Determining Region,” and “CDR” are used interchangeably. A “CDR” refers to one of three hypervariable regions (H1, H2 or H3) within the non-framework region of the immunoglobulin (Ig or antibody) VH β -sheet framework, or one of three hypervariable regions (L1, L2 or L3) within the non-framework region of the antibody VL β -sheet framework. Accordingly, CDRs are variable region sequences interspersed within the framework region sequences.

[0063] CDR regions are well known to those skilled in the art and have been defined by well-known numbering systems. For example, the Kabat Complementarity Determining Regions (CDRs) are based on sequence variability and are the most commonly used (*see, e.g., Kabat et al., supra*). Chothia refers instead to the location of the structural loops (*see, e.g., Chothia and Lesk, J. Mol. Biol. 196:901-17 (1987)*). The end of the Chothia CDR-H1 loop when numbered using the Kabat numbering convention varies between H32 and H34 depending on the length of the loop (this is because the Kabat numbering scheme places the insertions at H35A and H35B; if neither 35A nor 35B is present, the loop ends at 32; if only 35A is present, the loop ends at 33; if both 35A and 35B are present, the loop ends at 34). The AbM hypervariable regions represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular's AbM antibody modeling software (*see, e.g., Antibody Engineering Vol. 2 (Kontermann and Dübel eds., 2d ed. 2010)*). The "contact" hypervariable regions are based on an analysis of the available complex crystal structures. Another universal numbering system that has been developed and widely adopted is ImMunoGeneTics (IMGT) Information System[®] (Lafranc *et al.*, Dev. Comp. Immunol. 27(1):55-77 (2003)). IMGT is an integrated information system specializing in immunoglobulins (IG), T-cell receptors (TCR), and major histocompatibility complex (MHC) of human and other vertebrates. Herein, the CDRs are referred to in terms of both the amino acid sequence and the location within the light or heavy chain. As the "location" of the CDRs within the structure of the immunoglobulin variable domain is conserved between species and present in structures called loops, by using numbering systems that align variable domain sequences according to structural features, CDR and framework residues are readily identified. This information can be used in grafting and replacement of CDR residues from immunoglobulins of one species into an acceptor framework from, typically, a human antibody. An additional numbering system (AHon) has been developed by Honegger and Plückthun, J. Mol. Biol. 309: 657-70 (2001). Correspondence between the numbering system, including, for example, the Kabat numbering and the IMGT unique numbering system, is well known to one skilled in the art (*see, e.g., Kabat, supra; Chothia and Lesk, supra; Martin, supra; Lefranc et al., supra*). The residues from each of these hypervariable regions or CDRs are exemplified in Table 1 below.

Table 1. Exemplary CDRs According to Various Numbering Systems

Loop	Kabat	AbM	Chothia	Contact	IMGT
CDR L1	L24--L34	L24--L34	L26--L32 or L24--L34	L30--L36	L27--L38
CDR L2	L50--L56	L50--L56	L50--L52 or L50--L56	L46--L55	L56--L65
CDR L3	L89--L97	L89--L97	L91--L96 or L89--L97	L89--L96	L105-L117
CDR H1	H31--H35B	H26--H35B	H26--H32..34	H30--H35B	H27--H38

Loop	Kabat	AbM	Chothia	Contact	IMGT
	(Kabat Numbering)				
CDR H1	H31--H35 (Chothia Numbering)	H26--H35	H26--H32	H30--H35	
CDR H2	H50--H65	H50--H58	H53--H55 or H52--H56	H47--H58	H56--H65
CDR H3	H95--H102	H95--H102	H96--H101 or H95--H102	H93--H101	H105-H117

[0064] The boundaries of a given CDR may vary depending on the scheme used for identification. Thus, unless otherwise specified, the terms “CDR” and “complementary determining region” of a given antibody or region thereof, such as a variable region, as well as individual CDRs (*e.g.*, CDR-H1, CDR-H2) of the antibody or region thereof, should be understood to encompass the complementary determining region as defined by any of the known schemes described herein above. In some instances, the scheme for identification of a particular CDR or CDRs is specified, such as the CDR as defined by the IMGT, Kabat, Chothia, or Contact method. In some instances, one or more positions according to the Kabat numbering may not be occupied in the actual sequence, or the actual sequence may contain more amino acid residues than the number allowed for by the Kabat numbering. *See, e.g.*, Deschacht *et al.*, 2010. J Immunol 184: 5696-704 for an exemplary numbering for VHH domains according to Kabat. In other cases, the particular amino acid sequence of a CDR is given. It should be noted CDR regions may also be defined by a combination of various numbering systems, *e.g.*, a combination of Kabat and Chothia numbering systems, or a combination of Kabat and IMGT numbering systems. Therefore, the term such as “a CDR as set forth in a specific VH or VHH” includes any CDR1 as defined by the exemplary CDR numbering systems described above, but is not limited thereby. Once a variable region (*e.g.*, a VHH, VH or VL) is given, those skilled in the art would understand that CDRs within the region can be defined by different numbering systems or combinations thereof.

[0065] Hypervariable regions may comprise “extended hypervariable regions” as follows: 24-36 or 24-34 (L1), 46-56 or 50-56 (L2), and 89-97 or 89-96 (L3) in the VL, and 26-35 or 26-35A (H1), 50-65 or 49-65 (H2), and 93-102, 94-102, or 95-102 (H3) in the VH.

[0066] The term “constant region” or “constant domain” refers to a carboxy terminal portion of the light and heavy chain which is not directly involved in binding of the antibody to antigen but exhibits various effector function, such as interaction with the Fc receptor. The term refers to the portion of an immunoglobulin molecule having a more conserved amino acid sequence relative to the other portion of the immunoglobulin, the variable region, which contains the antigen binding site. The constant region may contain the CH1, CH2, and CH3 regions of the heavy

chain and the CL region of the light chain.

[0067] The term “framework” or “FR” refers to those variable region residues flanking the CDRs. FR residues are present, for example, in chimeric, humanized, human, domain antibodies (*e.g.*, single domain antibodies), diabodies, linear antibodies, and bispecific antibodies. FR residues are those variable domain residues other than the hypervariable region residues or CDR residues.

[0068] The term “Fc region” herein is used to define a C-terminal region of an immunoglobulin heavy chain, including, for example, native sequence Fc regions, recombinant Fc regions, and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is often defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue. A “functional Fc region” possesses an “effector function” of a native sequence Fc region. Exemplary “effector functions” include C1q binding; CDC; Fc receptor binding; ADCC; phagocytosis; downregulation of cell surface receptors (*e.g.*, B cell receptor), *etc.* Such effector functions generally require the Fc region to be combined with a binding region or binding domain (*e.g.*, an antibody variable region or domain) and can be assessed using various assays known to those skilled in the art. A “variant Fc region” comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification (*e.g.*, substituting, addition, or deletion). In certain embodiments, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, for example, from about one to about ten amino acid substitutions, or from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of a parent polypeptide. The variant Fc region herein can possess at least about 80% homology with a native sequence Fc region and/or with an Fc region of a parent polypeptide, or at least about 90% homology therewith, for example, at least about 95% homology therewith.

[0069] As used herein, an “epitope” is a term in the art and refers to a localized region of an antigen to which a binding molecule (*e.g.*, an antibody comprising a single domain antibody sequence) can specifically bind. An epitope can be a linear epitope or a conformational, non-linear, or discontinuous epitope. In the case of a polypeptide antigen, for example, an epitope can be contiguous amino acids of the polypeptide (a “linear” epitope) or an epitope can comprise

amino acids from two or more non-contiguous regions of the polypeptide (a “conformational,” “non-linear” or “discontinuous” epitope). It will be appreciated by one of skill in the art that, in general, a linear epitope may or may not be dependent on secondary, tertiary, or quaternary structure. For example, in some embodiments, a binding molecule binds to a group of amino acids regardless of whether they are folded in a natural three dimensional protein structure. In other embodiments, a binding molecule requires amino acid residues making up the epitope to exhibit a particular conformation (*e.g.*, bend, twist, turn or fold) in order to recognize and bind the epitope.

[0070] A “blocking” antibody or an “antagonist” antibody is one that inhibits or reduces a biological activity of the antigen it binds. In some embodiments, blocking antibodies or antagonist antibodies substantially or completely inhibit the biological activity of the antigen.

[0071] An “agonist” or activating antibody is one that enhances or initiates signaling by the antigen to which it binds. In some embodiments, agonist antibodies cause or activate signaling without the presence of the natural ligand.

[0072] “Percent (%) amino acid sequence identity” and “homology” with respect to a peptide, polypeptide or antibody sequence are defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific peptide or polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or MEGALIGN™ (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared.

[0073] The term “specificity” refers to selective recognition of an antigen binding protein (such as a CAR or an sdAb) for a particular epitope of an antigen. Natural antibodies, for example, are monospecific. The term “multispecific” as used herein denotes that an antigen binding protein (such as a CAR or an sdAb) has two or more antigen-binding sites of which at least two bind different antigens. “Bispecific” as used herein denotes that an antigen binding protein (such as a CAR or an sdAb) has two different antigen-binding specificities. The term “monospecific” CARs as used herein denotes an antigen binding protein (such as a CAR or an sdAb) that has one or more binding sites each of which bind the same antigen.

[0074] The term “valent” as used herein denotes the presence of a specified number of binding sites in an antigen binding protein (such as a CAR or an sdAb). A natural antibody for example or a full length antibody has two binding sites and is bivalent. As such, the terms “trivalent”,

"tetravalent", "pentavalent" and "hexavalent" denote the presence of two binding site, three binding sites, four binding sites, five binding sites, and six binding sites, respectively, in an antigen binding protein (such as a CAR or an sdAb).

[0075] "Chimeric antigen receptor" or "CAR" as used herein refers to genetically engineered receptors, which can be used to graft one or more antigen specificity onto immune effector cells, such as T cells. Some CARs are also known as "artificial T-cell receptors," "chimeric T cell receptors," or "chimeric immune receptors." In some embodiments, the CAR comprises an extracellular antigen binding domain specific for one or more antigens (such as tumor antigens), a transmembrane domain, and an intracellular signaling domain of a T cell and/or other receptors. "CAR-T cell" refers to a T cell that expresses a CAR.

[0076] The terms "polypeptide" and "peptide" and "protein" are used interchangeably herein and refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid, including but not limited to, unnatural amino acids, as well as other modifications known in the art. It is understood that, because the polypeptides of this disclosure may be based upon antibodies or other members of the immunoglobulin superfamily, in certain embodiments, a "polypeptide" can occur as a single chain or as two or more associated chains.

[0077] "Polynucleotide" or "nucleic acid," as used interchangeably herein, refers to polymers of nucleotides of any length and includes DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase or by a synthetic reaction. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and their analogs. "Oligonucleotide," as used herein, refers to short, generally single-stranded, synthetic polynucleotides that are generally, but not necessarily, fewer than about 200 nucleotides in length. The terms "oligonucleotide" and "polynucleotide" are not mutually exclusive. The description above for polynucleotides is equally and fully applicable to oligonucleotides. A cell that produces a binding molecule of the present disclosure may include a parent hybridoma cell, as well as bacterial and eukaryotic host cells into which nucleic acids encoding the antibodies have been introduced. Unless specified otherwise, the left-hand end of any single-stranded polynucleotide sequence disclosed herein is the 5' end; the left-hand direction of double-stranded polynucleotide sequences is referred to as the 5' direction. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription

direction; sequence regions on the DNA strand having the same sequence as the RNA transcript that are 5' to the 5' end of the RNA transcript are referred to as "upstream sequences"; sequence regions on the DNA strand having the same sequence as the RNA transcript that are 3' to the 3' end of the RNA transcript are referred to as "downstream sequences."

[0078] An "isolated nucleic acid" is a nucleic acid, for example, an RNA, DNA, or a mixed nucleic acids, which is substantially separated from other genome DNA sequences as well as proteins or complexes such as ribosomes and polymerases, which naturally accompany a native sequence. An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid molecule. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. In a specific embodiment, one or more nucleic acid molecules encoding a single domain antibody or an antibody as described herein are isolated or purified. The term embraces nucleic acid sequences that have been removed from their naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogues or analogues biologically synthesized by heterologous systems. A substantially pure molecule may include isolated forms of the molecule. Specifically, an "isolated" nucleic acid molecule encoding a CAR or an sdAb described herein is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the environment in which it was produced.

[0079] The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

[0080] As used herein, the term "operatively linked," and similar phrases (*e.g.*, genetically fused), when used in reference to nucleic acids or amino acids, refer to the operational linkage of nucleic acid sequences or amino acid sequence, respectively, placed in functional relationships with each other. For example, an operatively linked promoter, enhancer elements, open reading frame, 5' and 3' UTR, and terminator sequences result in the accurate production of a nucleic acid molecule (*e.g.*, RNA). In some embodiments, operatively linked nucleic acid elements result in the transcription of an open reading frame and ultimately the production of a polypeptide (*i.e.*, expression of the open reading frame). As another example, an operatively linked peptide is one in which the functional domains are placed with appropriate distance from each other to impart the intended function of each domain.

[0081] The term “vector” refers to a substance that is used to carry or include a nucleic acid sequence, including for example, a nucleic acid sequence encoding a binding molecule (*e.g.*, an antibody) as described herein, in order to introduce a nucleic acid sequence into a host cell. Vectors applicable for use include, for example, expression vectors, plasmids, phage vectors, viral vectors, episomes, and artificial chromosomes, which can include selection sequences or markers operable for stable integration into a host cell’s chromosome. Additionally, the vectors can include one or more selectable marker genes and appropriate expression control sequences. Selectable marker genes that can be included, for example, provide resistance to antibiotics or toxins, complement auxotrophic deficiencies, or supply critical nutrients not in the culture media. Expression control sequences can include constitutive and inducible promoters, transcription enhancers, transcription terminators, and the like, which are well known in the art. When two or more nucleic acid molecules are to be co-expressed (*e.g.*, both an antibody heavy and light chain or an antibody VH and VL), both nucleic acid molecules can be inserted, for example, into a single expression vector or in separate expression vectors. For single vector expression, the encoding nucleic acids can be operationally linked to one common expression control sequence or linked to different expression control sequences, such as one inducible promoter and one constitutive promoter. The introduction of nucleic acid molecules into a host cell can be confirmed using methods well known in the art. Such methods include, for example, nucleic acid analysis such as Northern blots or polymerase chain reaction (PCR) amplification of mRNA, immunoblotting for expression of gene products, or other suitable analytical methods to test the expression of an introduced nucleic acid sequence or its corresponding gene product. It is understood by those skilled in the art that the nucleic acid molecules are expressed in a sufficient amount to produce a desired product and it is further understood that expression levels can be optimized to obtain sufficient expression using methods well known in the art.

[0082] The term “host” as used herein refers to an animal, such as a mammal (*e.g.*, a human).

[0083] The term “host cell” as used herein refers to a particular subject cell that may be transfected with a nucleic acid molecule and the progeny or potential progeny of such a cell. Progeny of such a cell may not be identical to the parent cell transfected with the nucleic acid molecule due to mutations or environmental influences that may occur in succeeding generations or integration of the nucleic acid molecule into the host cell genome.

[0084] As used herein, the term “autologous” is meant to refer to any material derived from the same individual to whom it is later to be re-introduced into the individual.

[0085] “Allogeneic” refers to a graft derived from a different individual of the same species.

[0086] The term “transfected” or “transformed” or “transduced” as used herein refers to a process by which exogenous nucleic acid is transferred or introduced into the host cell. A “transfected” or “transformed” or “transduced” cell is one which has been transfected,

transformed or transduced with exogenous nucleic acid. The cell includes the primary subject cell and its progeny.

[0087] The term “pharmaceutically acceptable” as used herein means being approved by a regulatory agency of the Federal or a state government, or listed in United States Pharmacopeia, European Pharmacopeia, or other generally recognized Pharmacopeia for use in animals, and more particularly in humans.

[0088] “Excipient” means a pharmaceutically-acceptable material, composition, or vehicle, such as a liquid or solid filler, diluent, solvent, or encapsulating material. Excipients include, for example, encapsulating materials or additives such as absorption accelerators, antioxidants, binders, buffers, carriers, coating agents, coloring agents, diluents, disintegrating agents, emulsifiers, extenders, fillers, flavoring agents, humectants, lubricants, perfumes, preservatives, propellants, releasing agents, sterilizing agents, sweeteners, solubilizers, wetting agents and mixtures thereof. The term “excipient” can also refer to a diluent, adjuvant (*e.g.*, Freund’s adjuvant (complete or incomplete) or vehicle.

[0089] In some embodiments, excipients are pharmaceutically acceptable excipients. Examples of pharmaceutically acceptable excipients include buffers, such as phosphate, citrate, and other organic acids; antioxidants, including ascorbic acid; low molecular weight (*e.g.*, fewer than about 10 amino acid residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers, such as polyvinylpyrrolidone; amino acids, such as glycine, glutamine, asparagine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates, including glucose, mannose, or dextrans; chelating agents, such as EDTA; sugar alcohols, such as mannitol or sorbitol; salt-forming counterions, such as sodium; and/or nonionic surfactants, such as TWEENTM, polyethylene glycol (PEG), and PLURONICSTM. Other examples of pharmaceutically acceptable excipients are described in Remington and Gennaro, Remington’s Pharmaceutical Sciences (18th ed. 1990).

[0090] In one embodiment, each component is “pharmaceutically acceptable” in the sense of being compatible with the other ingredients of a pharmaceutical formulation, and suitable for use in contact with the tissue or organ of humans and animals without excessive toxicity, irritation, allergic response, immunogenicity, or other problems or complications, commensurate with a reasonable benefit/risk ratio. *See, e.g.*, Lippincott Williams & Wilkins: Philadelphia, PA, 2005; Handbook of Pharmaceutical Excipients, 6th ed.; Rowe et al., Eds.; The Pharmaceutical Press and the American Pharmaceutical Association: 2009; Handbook of Pharmaceutical Additives, 3rd ed.; Ash and Ash Eds.; Gower Publishing Company: 2007; Pharmaceutical Preformulation and Formulation, 2nd ed.; Gibson Ed.; CRC Press LLC: Boca Raton, FL, 2009. In some embodiments, pharmaceutically acceptable excipients are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. In some embodiments, a

pharmaceutically acceptable excipient is an aqueous pH buffered solution.

[0091] In some embodiments, excipients are sterile liquids, such as water and oils, including those of petroleum, animal, vegetable, or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil, and the like. Water is an exemplary excipient when a composition (*e.g.*, a pharmaceutical composition) is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid excipients, particularly for injectable solutions. An excipient can also include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol, and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents.

Compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations, and the like. Oral compositions, including formulations, can include standard excipients such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, *etc.*

[0092] Compositions, including pharmaceutical compounds, may contain a binding molecule (*e.g.*, an antibody), for example, in isolated or purified form, together with a suitable amount of excipients.

[0093] The term “effective amount” or “therapeutically effective amount” as used herein refers to the amount of a single domain antibody or a therapeutic molecule comprising an agent and the single domain antibody or pharmaceutical composition provided herein which is sufficient to result in the desired outcome.

[0094] The terms “subject” and “patient” may be used interchangeably. As used herein, in certain embodiments, a subject is a mammal, such as a non-primate or a primate (*e.g.*, human). In specific embodiments, the subject is a human. In one embodiment, the subject is a mammal, *e.g.*, a human, diagnosed with a disease or disorder. In another embodiment, the subject is a mammal, *e.g.*, a human, at risk of developing a disease or disorder.

[0095] “Administer” or “administration” refers to the act of injecting or otherwise physically delivering a substance as it exists outside the body into a patient, such as by mucosal, intradermal, intravenous, intramuscular delivery, and/or any other method of physical delivery described herein or known in the art.

[0096] As used herein, the terms “treat,” “treatment” and “treating” refer to the reduction or amelioration of the progression, severity, and/or duration of a disease or condition resulting from the administration of one or more therapies. Treating may be determined by assessing whether there has been a decrease, alleviation and/or mitigation of one or more symptoms associated with the underlying disorder such that an improvement is observed with the patient, despite that the patient may still be afflicted with the underlying disorder. The term “treating” includes both

managing and ameliorating the disease. The terms “manage,” “managing,” and “management” refer to the beneficial effects that a subject derives from a therapy which does not necessarily result in a cure of the disease.

[0097] The terms “prevent,” “preventing,” and “prevention” refer to reducing the likelihood of the onset (or recurrence) of a disease, disorder, condition, or associated symptom(s) (*e.g.*, diabetes or a cancer).

[0098] As used herein, “delaying” the development of cancer means to defer, hinder, slow, retard, stabilize, and/or postpone development of the disease. This delay can be of varying lengths of time, depending on the history of the disease and/or individual being treated. As is evident to one skilled in the art, a sufficient or significant delay can, in effect, encompass prevention, in that the individual does not develop the disease. A method that “delays” development of cancer is a method that reduces probability of disease development in a given time frame and/or reduces the extent of the disease in a given time frame, when compared to not using the method. Such comparisons are typically based on clinical studies, using a statistically significant number of individuals. Cancer development can be detectable using standard methods, including, but not limited to, computerized axial tomography (CAT Scan), Magnetic Resonance Imaging (MRI), abdominal ultrasound, clotting tests, arteriography, or biopsy. Development may also refer to cancer progression that may be initially undetectable and includes occurrence, recurrence, and onset.

[0099] “B cell associated disease or disorder” as used herein refers to a disease or disorder mediated by B cells or conferred by abnormal B cell functions (such as dysregulation of B-cell function). “B cell associated disease or disorder” as used herein includes but not limited to a B cell malignancy such as a B cell leukemia or B cell lymphoma. It also includes marginal zone lymphoma (*e.g.*, splenic marginal zone lymphoma), diffuse large B cell lymphoma (DLBCL), mantle cell lymphoma (MCL), primary central nervous system (CNS) lymphoma, primary mediastinal B cell lymphoma (PMBL), small lymphocytic lymphoma (SLL), B cell prolymphocytic leukemia (B-PLL), follicular lymphoma (FL), burkitt lymphoma, primary intraocular lymphoma, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), hairy cell leukemia (HCL), precursor B lymphoblastic leukemia, non-hodgkin lymphoma (NHL), high-grade B-cell lymphoma (HGBL), and multiple myeloma (MM). “B cell associated disease or disorder” also includes certain autoimmune and/or inflammatory disease, such as those associated with inappropriate or enhanced B cell numbers and/or activation.

[00100] “CD20 associated disease or disorder” as used herein refers to a disease or disorder that comprises a cell or tissue in which CD20 is expressed. In some embodiments, CD20 associated disease or disorder comprises a cell on which CD20 is abnormally expressed. In other embodiments, CD20 associated disease or disorder comprises a cell in or on which CD20 is

deficient.

[00101] The terms “about” and “approximately” mean within 20%, within 15%, within 10%, within 9%, within 8%, within 7%, within 6%, within 5%, within 4%, within 3%, within 2%, within 1%, or less of a given value or range.

[00102] As used in the present disclosure and claims, the singular forms “a”, “an” and “the” include plural forms unless the context clearly dictates otherwise.

[00103] It is understood that wherever embodiments are described herein with the term “comprising” otherwise analogous embodiments described in terms of “consisting of” and/or “consisting essentially of” are also provided. It is also understood that wherever embodiments are described herein with the phrase “consisting essentially of” otherwise analogous embodiments described in terms of “consisting of” are also provided.

[00104] The term “between” as used in a phrase as such “between A and B” or “between A-B” refers to a range including both A and B.

[00105] The term “and/or” as used in a phrase such as “A and/or B” herein is intended to include both A and B; A or B; A (alone); and B (alone). Likewise, the term “and/or” as used in a phrase such as “A, B, and/or C” is intended to encompass each of the following embodiments: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

5.2. Single Domain Antibodies

5.2.1. Single Domain Antibodies that Bind to CD20

[00106] In one aspect, provided herein are single domain antibodies (*e.g.*, VHH domains) capable of binding to CD20.

[00107] In some embodiments, the single domain antibodies (*e.g.*, VHH domains) provided herein bind to human CD20. In some embodiments, the anti-CD20 single domain antibody provided herein modulates one or more CD20 activities. In some embodiments, the anti-CD20 single domain antibody provided herein is an antagonist antibody.

[00108] In some embodiments, the anti-CD20 single domain antibody provided herein binds to CD20 (*e.g.*, human CD20) with a dissociation constant (K_D) of $\leq 1\mu\text{M}$, $\leq 100\text{ nM}$, $\leq 10\text{ nM}$, $\leq 1\text{ nM}$, $\leq 0.1\text{ nM}$, $\leq 0.01\text{ nM}$, or $\leq 0.001\text{ nM}$ (*e.g.* 10^{-8} M or less, *e.g.* from 10^{-8} M to 10^{-13} M , *e.g.*, from 10^{-9} M to 10^{-13} M). A variety of methods of measuring binding affinity are known in the art, any of which can be used for purposes of the present disclosure, including by RIA, for example, performed with the Fab version of an antibody of interest and its antigen (Chen et al., 1999, J. Mol Biol 293:865-81); by biolayer interferometry (BLI) or surface plasmon resonance (SPR) assays by Octet®, using, for example, an Octet®Red96 system, or by Biacore®, using, for example, a Biacore®TM-2000 or a Biacore®TM-3000. An “on-rate” or “rate of association” or “association rate” or “kon” may also be determined with the same biolayer interferometry (BLI)

or surface plasmon resonance (SPR) techniques described above using, for example, the Octet®Red96, the Biacore®TM-2000, or the Biacore®TM-3000 system.

[00109] In some embodiments, the anti-CD20 single domain antibodies provided herein are VHH domains. Exemplary VHH domains provided herein are generated as described below in Section 6, and these VHH domains are referred to as VHH-273, VHH-283, VHH-313, VHH-440, VHH-466, VHH-496, VHH-653, huVHH-253, huVHH-256, huVHH-260, huVHH-746, huVHH-750, huVHH-753, huVHH-836, huVHH-840, huVHH-843, huVHH-846, 2082H1, 2082H2, 2082H3, 2082H4, 2082H5, 2082H6, VHH-623, VHH-640, and/or VHH-657 as also shown in Table 2 below.

[00110] Thus, in some embodiments, the single domain antibody provided herein comprises one or more CDR sequences of any one of VHH-273, VHH-283, VHH-313, VHH-440, VHH-466, VHH-496, VHH-653, huVHH-253, huVHH-256, huVHH-260, huVHH-746, huVHH-750, huVHH-753, huVHH-836, huVHH-840, huVHH-843, huVHH-846, 2082H1, 2082H2, 2082H3, 2082H4, 2082H5, 2082H6, VHH-623, VHH-640, and/or VHH-657. In some embodiments, provided herein is a single domain antibody that binds to CD20 comprising the following structure: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4, wherein the CDR sequences are selected from those in VHH-273, VHH-283, VHH-313, VHH-440, VHH-466, VHH-496, VHH-653, huVHH-253, huVHH-256, huVHH-260, huVHH-746, huVHH-750, huVHH-753, huVHH-836, huVHH-840, huVHH-843, huVHH-846, 2082H1, 2082H2, 2082H3, 2082H4, 2082H5, 2082H6, VHH-623, VHH-640, and/or VHH-657.

Table 2. Exemplary Single Domain Antibodies

Clone ID	VHH Domain	CDR1	CDR2	CDR3
VHH-273	SEQ ID NO: 41	SEQ ID NO: 1 or 195	SEQ ID NO: 2	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 5	SEQ ID NO: 6
VHH-283	SEQ ID NO: 42	SEQ ID NO: 1 or 195	SEQ ID NO: 7	SEQ ID NO: 8
		SEQ ID NO: 4	SEQ ID NO: 9	SEQ ID NO: 10
VHH-313	SEQ ID NO: 43	SEQ ID NO: 11 or 196	SEQ ID NO: 12	SEQ ID NO: 8
		SEQ ID NO: 13	SEQ ID NO: 9	SEQ ID NO: 10
VHH-440	SEQ ID NO: 44	SEQ ID NO: 14 or 197	SEQ ID NO: 15	SEQ ID NO: 16
		SEQ ID NO: 17	SEQ ID NO: 18	SEQ ID NO: 19
VHH-466	SEQ ID NO: 45	SEQ ID NO: 20 or 197	SEQ ID NO: 21	SEQ ID NO: 16
		SEQ ID NO: 22	SEQ ID NO: 23	SEQ ID NO: 19

Clone ID	VHH Domain	CDR1	CDR2	CDR3
VHH-496	SEQ ID NO: 46	SEQ ID NO: 24 or 198	SEQ ID NO: 25	SEQ ID NO: 26
		SEQ ID NO: 27	SEQ ID NO: 28	SEQ ID NO: 29
VHH-653	SEQ ID NO: 47	SEQ ID NO: 14 or 197	SEQ ID NO: 30	SEQ ID NO: 16
		SEQ ID NO: 17	SEQ ID NO: 31	SEQ ID NO: 19
huVHH-253	SEQ ID NO: 48	SEQ ID NO: 1 or 195	SEQ ID NO: 32	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 33	SEQ ID NO: 6
huVHH-256	SEQ ID NO: 49	SEQ ID NO: 1 or 195	SEQ ID NO: 2	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 33	SEQ ID NO: 6
huVHH-260	SEQ ID NO: 50	SEQ ID NO: 11 or 196	SEQ ID NO: 12	SEQ ID NO: 8
		SEQ ID NO: 13	SEQ ID NO: 9	SEQ ID NO: 10
huVHH-746	SEQ ID NO: 51	SEQ ID NO: 24 or 198	SEQ ID NO: 25	SEQ ID NO: 26
		SEQ ID NO: 27	SEQ ID NO: 28	SEQ ID NO: 29
huVHH-750	SEQ ID NO: 52	SEQ ID NO: 24 or 198	SEQ ID NO: 25	SEQ ID NO: 26
		SEQ ID NO: 27	SEQ ID NO: 28	SEQ ID NO: 29
huVHH-753	SEQ ID NO: 53	SEQ ID NO: 24 or 198	SEQ ID NO: 25	SEQ ID NO: 26
		SEQ ID NO: 27	SEQ ID NO: 28	SEQ ID NO: 29
huVHH-836	SEQ ID NO: 54	SEQ ID NO: 20 or 197	SEQ ID NO: 34	SEQ ID NO: 16
		SEQ ID NO: 22	SEQ ID NO: 35	SEQ ID NO: 19
huVHH-840	SEQ ID NO: 55	SEQ ID NO: 20 or 197	SEQ ID NO: 36	SEQ ID NO: 16
		SEQ ID NO: 22	SEQ ID NO: 23	SEQ ID NO: 19
huVHH-843	SEQ ID NO: 56	SEQ ID NO: 20 or 197	SEQ ID NO: 37	SEQ ID NO: 16
		SEQ ID NO: 22	SEQ ID NO: 23	SEQ ID NO: 19
huVHH-846	SEQ ID NO: 57	SEQ ID NO: 20 or 197	SEQ ID NO: 38	SEQ ID NO: 16
		SEQ ID NO: 22	SEQ ID NO: 23	SEQ ID NO: 19

Clone ID	VHH Domain	CDR1	CDR2	CDR3
2082H1	SEQ ID NO: 58	SEQ ID NO: 1 or 195	SEQ ID NO: 2	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 33	SEQ ID NO: 6
2082H2	SEQ ID NO: 59	SEQ ID NO: 1 or 195	SEQ ID NO: 2	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 33	SEQ ID NO: 39
2082H3	SEQ ID NO: 60	SEQ ID NO: 1 or 195	SEQ ID NO: 2	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 33	SEQ ID NO: 6
2082H4	SEQ ID NO: 61	SEQ ID NO: 1 or 195	SEQ ID NO: 2	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 33	SEQ ID NO: 40
2082H5	SEQ ID NO: 62	SEQ ID NO: 1 or 195	SEQ ID NO: 2	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 33	SEQ ID NO: 39
2082H6	SEQ ID NO: 63	SEQ ID NO: 1 or 195	SEQ ID NO: 2	SEQ ID NO: 3
		SEQ ID NO: 4	SEQ ID NO: 33	SEQ ID NO: 6
VHH-623	SEQ ID NO: 183	SEQ ID NO: 165 or 199	SEQ ID NO: 166	SEQ ID NO: 167
		SEQ ID NO: 168	SEQ ID NO: 169	SEQ ID NO: 170
VHH-640	SEQ ID NO: 184	SEQ ID NO: 171 or 200	SEQ ID NO: 172	SEQ ID NO: 173
		SEQ ID NO: 174	SEQ ID NO: 175	SEQ ID NO: 176
VHH-657	SEQ ID NO: 185	SEQ ID NO: 177 or 197	SEQ ID NO: 178	SEQ ID NO: 179
		SEQ ID NO: 180	SEQ ID NO: 181	SEQ ID NO: 182

[00111] In some embodiments, there is provided an anti-CD20 single domain antibody comprising one, two, or all three CDRs of the amino acid sequence of SEQ ID NO: 41. In some embodiments, there is provided an anti-CD20 single domain antibody comprising one, two, or all three CDRs of the amino acid sequence of SEQ ID NO: 42. In some embodiments, there is provided an anti-CD20 single domain antibody comprising one, two, or all three CDRs of the amino acid sequence of SEQ ID NO: 43. In some embodiments, there is provided an anti-CD20 single domain antibody comprising one, two, or all three CDRs of the amino acid sequence of SEQ ID NO: 44. In some embodiments, there is provided an anti-CD20 single domain antibody comprising one, two, or all three CDRs of the amino acid sequence of SEQ ID NO: 45. In some

[illegible]

three CDRs of the amino acid sequence of SEQ ID NO: 185. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00112] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 41. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 41. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 41 (*e.g.*, SEQ ID NO: 3 or 6). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 41. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 41. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 41. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 41. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00113] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 42. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 42. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 42 (*e.g.*, SEQ ID NO: 8 or 10). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 42. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 42. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ

ID NO: 42. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 42. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00114] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 43. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 43. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 43 (*e.g.*, SEQ ID NO: 8 or 10). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 43. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 43. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 43. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 43. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00115] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 44. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 44. In other embodiments, the single domain antibody has a CDR3 having an amino acid

sequence of the CDR3 as set forth in SEQ ID NO: 44 (*e.g.*, SEQ ID NO: 16 or 19). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 44. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 44. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 44. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 44. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00116] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 45. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 45. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 45 (*e.g.*, SEQ ID NO: 16 or 19). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 45. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 45. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 45. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 45. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody

comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00117] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 46. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 46. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 46 (*e.g.*, SEQ ID NO: 26 or 29). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 46. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 46. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 46. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 46. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00118] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 47. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 47. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 47 (*e.g.*, SEQ ID NO: 16 or 19). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 47. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 47. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 47. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 47. CDR sequences can be determined according to well-known numbering systems. In

some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00119] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 48. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 48. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 48 (*e.g.*, SEQ ID NO: 3 or 6). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 48. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 48. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 48. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 48. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00120] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 49. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 49. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 49 (*e.g.*, SEQ ID NO: 3 or 6). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 49. In some embodiments, the single

domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 49. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 49. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 49. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00121] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 50. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 50. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 50 (*e.g.*, SEQ ID NO: 8 or 10). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 50. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 50. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 50. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 50. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00122] In some embodiments, the single domain antibody has a CDR1 having an amino acid

sequence of the CDR1 as set forth in SEQ ID NO: 51. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 51. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 51 (*e.g.*, SEQ ID NO: 26 or 29). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 51. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 51. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 51. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 51. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00123] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 52. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 52. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 52 (*e.g.*, SEQ ID NO: 26 or 29). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 52. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 52. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 52. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 52. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In

other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00124] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 53. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 53. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 53 (*e.g.*, SEQ ID NO: 26 or 29). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 53. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 53. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 53. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 53. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00125] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 54. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 54. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 54 (*e.g.*, SEQ ID NO: 16 or 19). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 54. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 54. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ

ID NO: 54. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 54. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00126] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 55. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 55. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 55 (*e.g.*, SEQ ID NO: 16 or 19). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 55. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 55. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 55. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 55. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00127] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 56. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 56. In other embodiments, the single domain antibody has a CDR3 having an amino acid

sequence of the CDR3 as set forth in SEQ ID NO: 56 (*e.g.*, SEQ ID NO: 16 or 19). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 56. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 56. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 56. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 56. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00128] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 57. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 57. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 57 (*e.g.*, SEQ ID NO: 16 or 19). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 57. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 57. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 57. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 57. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises

an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00129] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 58. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 58. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 58 (*e.g.*, SEQ ID NO: 3 or 6). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 58. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 58. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 58. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 58. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00130] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 59. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 59. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 59 (*e.g.*, SEQ ID NO: 3 or 39). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 59. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 59. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 59. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 59. CDR sequences can be determined according to well-known numbering systems. In

some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00131] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 60. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 60. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 60 (*e.g.*, SEQ ID NO: 3 or 6). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 60. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 60. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 60. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 60. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00132] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 61. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 61. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 61 (*e.g.*, SEQ ID NO: 3 or 40). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 61. In some embodiments, the single

domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 61. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 61. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 61. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00133] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 62. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 62. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 62 (*e.g.*, SEQ ID NO: 3 or 39). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 62. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 62. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 62. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 62. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00134] In some embodiments, the single domain antibody has a CDR1 having an amino acid

sequence of the CDR1 as set forth in SEQ ID NO: 63. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 63. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 63 (*e.g.*, SEQ ID NO: 3 or 6). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 63. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 63. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 63. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 63. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00135] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 183. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 183. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 183 (*e.g.*, SEQ ID NO: 167 or 170). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 183. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 183. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 183. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 183. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In

other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, e.g., a human immunoglobulin framework or a human consensus framework.

[00136] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 184. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 184. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 184 (e.g., SEQ ID NO: 173 or 176). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 184. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 184. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ ID NO: 184. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 184. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, e.g., a human immunoglobulin framework or a human consensus framework.

[00137] In some embodiments, the single domain antibody has a CDR1 having an amino acid sequence of the CDR1 as set forth in SEQ ID NO: 185. In some embodiments, the single domain antibody has a CDR2 having an amino acid sequence of the CDR2 as set forth in SEQ ID NO: 185. In other embodiments, the single domain antibody has a CDR3 having an amino acid sequence of the CDR3 as set forth in SEQ ID NO: 185 (e.g., SEQ ID NO: 179 or 182). In some embodiments, the single domain antibody has a CDR1 and a CDR2 having amino acid sequences of the CDR1 and the CDR2 as set forth in SEQ ID NO: 185. In some embodiments, the single domain antibody has a CDR1 and a CDR3 having amino acid sequences of the CDR1 and the CDR3 as set forth in SEQ ID NO: 185. In some embodiments, the single domain antibody has a CDR2 and a CDR3 having amino acid sequences of the CDR2 and the CDR3 as set forth in SEQ

ID NO: 185. In some embodiments, the single domain antibody has a CDR1, a CDR2, and a CDR3 having amino acid sequences of the CDR1, the CDR2, and the CDR3 as set forth in SEQ ID NO: 185. CDR sequences can be determined according to well-known numbering systems. In some embodiments, the CDRs are according to IMGT numbering. In some embodiments, the CDRs are according to Kabat numbering. In some embodiments, the CDRs are according to AbM numbering. In other embodiments, the CDRs are according to Chothia numbering. In other embodiments, the CDRs are according to Contact numbering. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, e.g., a human immunoglobulin framework or a human consensus framework.

[00138] In some embodiments, provided herein is a single domain antibody that binds to CD20 comprising the following structure: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4, wherein (i) the CDR1 comprises an amino acid sequence of SEQ ID NO: 1 or 195, SEQ ID NO: 4, SEQ ID NO: 11 or 196, SEQ ID NO: 13, SEQ ID NO: 14 or 197, SEQ ID NO: 17, SEQ ID NO: 20 or 197, SEQ ID NO: 22, SEQ ID NO: 24 or 198, SEQ ID NO: 27; SEQ ID NO: 165 or 199, SEQ ID NO: 168, SEQ ID NO: 171 or 200, SEQ ID NO: 174, SEQ ID NO: 177 or 197, or SEQ ID NO: 180; (ii) the CDR2 comprises an amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 166, SEQ ID NO: 169, SEQ ID NO: 172, SEQ ID NO: 175, SEQ ID NO: 178, or SEQ ID NO: 181; and/or (iii) the CDR3 comprises an amino acid sequence of SEQ ID NO: 3, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 19, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 39, SEQ ID NO: 40, SEQ ID NO: 167, SEQ ID NO: 170, SEQ ID NO: 173, SEQ ID NO: 176, SEQ ID NO: 179 or SEQ ID NO: 182. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, e.g., a human immunoglobulin framework or a human consensus framework.

[00139] In other embodiments, provided herein is a single domain antibody that binds to CD20 comprising the following structure: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4, wherein (i) the CDR1 comprises an amino acid sequence having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO: 1 or 195, SEQ ID NO: 4, SEQ ID NO: 11 or 196, SEQ ID NO: 13, SEQ ID NO: 14 or 197, SEQ ID NO: 17, SEQ ID NO: 20 or 197, SEQ ID NO: 22, SEQ ID NO: 24 or 198, SEQ ID NO: 27,

SEQ ID NO: 165 or 199, SEQ ID NO: 168, SEQ ID NO: 171 or 200, SEQ ID NO: 174, SEQ ID NO: 177 or 197, or SEQ ID NO: 180; (ii) the CDR2 comprises an amino acid sequence having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO: 2, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 12, SEQ ID NO: 15, SEQ ID NO: 18, SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 166, SEQ ID NO: 169, SEQ ID NO: 172, SEQ ID NO: 175, SEQ ID NO: 178, or SEQ ID NO: 181; and/or (iii) the CDR3 comprises an amino acid sequence having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO: 3, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 16, SEQ ID NO: 19, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 39, SEQ ID NO: 40, SEQ ID NO: 167, SEQ ID NO: 170, SEQ ID NO: 173, SEQ ID NO: 176, SEQ ID NO: 179 or SEQ ID NO: 182. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00140] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 2; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 1; a CDR2 comprising the amino acid sequence of SEQ ID NO: 2; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 2; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00141] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 5; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 6. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments,

the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00142] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 7; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 1; a CDR2 comprising the amino acid sequence of SEQ ID NO: 7; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 7; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00143] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 9; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 10. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00144] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 11 or 196; a CDR2 comprising the amino acid sequence of SEQ ID NO: 12; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 11; a CDR2 comprising the amino acid sequence of SEQ ID NO: 12; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 196; a CDR2 comprising the amino acid sequence of SEQ ID NO: 12; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human

consensus framework.

[00145] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 13; a CDR2 comprising the amino acid sequence of SEQ ID NO: 9; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 10. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00146] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 14 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 15; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 14; a CDR2 comprising the amino acid sequence of SEQ ID NO: 15; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 15; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00147] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 17; a CDR2 comprising the amino acid sequence of SEQ ID NO: 18; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00148] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 21; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20; a CDR2 comprising the amino acid sequence of SEQ ID NO: 21; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein

is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 21; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00149] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 22; a CDR2 comprising the amino acid sequence of SEQ ID NO: 23; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00150] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 24 or 198; a CDR2 comprising the amino acid sequence of SEQ ID NO: 25; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 26. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 24; a CDR2 comprising the amino acid sequence of SEQ ID NO: 25; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 26. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 198; a CDR2 comprising the amino acid sequence of SEQ ID NO: 25; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 26. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00151] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 27; a CDR2 comprising the amino acid sequence of SEQ ID NO: 28; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 29. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00152] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb)

comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 14 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 30; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 14; a CDR2 comprising the amino acid sequence of SEQ ID NO: 30; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 30; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00153] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 17; a CDR2 comprising the amino acid sequence of SEQ ID NO: 31; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00154] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 32; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 1; a CDR2 comprising the amino acid sequence of SEQ ID NO: 32; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 32; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00155] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising

the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 6. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00156] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 34; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20; a CDR2 comprising the amino acid sequence of SEQ ID NO: 34; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 34; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00157] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 22; a CDR2 comprising the amino acid sequence of SEQ ID NO: 35; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00158] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 36; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20; a CDR2 comprising the amino acid sequence of SEQ ID NO: 36; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 36; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, the

anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00159] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 37; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20; a CDR2 comprising the amino acid sequence of SEQ ID NO: 37; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 37; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00160] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 38; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 20; a CDR2 comprising the amino acid sequence of SEQ ID NO: 38; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 38; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00161] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 39. In some embodiments, the anti-CD20 single domain antibody is camelid. In

some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00162] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 40. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00163] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 165 or 199; a CDR2 comprising the amino acid sequence of SEQ ID NO: 166; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 167. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 165; a CDR2 comprising the amino acid sequence of SEQ ID NO: 166; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 167. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 199; a CDR2 comprising the amino acid sequence of SEQ ID NO: 166; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 167. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00164] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 168; a CDR2 comprising the amino acid sequence of SEQ ID NO: 169; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 170. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00165] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 171 or 200; a CDR2 comprising the amino acid sequence of SEQ ID NO: 172; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 173. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ

ID NO: 171; a CDR2 comprising the amino acid sequence of SEQ ID NO: 172; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 173. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 200; a CDR2 comprising the amino acid sequence of SEQ ID NO: 172; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 173. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00166] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 174; a CDR2 comprising the amino acid sequence of SEQ ID NO: 175; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 176. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00167] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 177 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 178; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 179. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 177; a CDR2 comprising the amino acid sequence of SEQ ID NO: 178; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 179. In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 178; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 179. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework, *e.g.*, a human immunoglobulin framework or a human consensus framework.

[00168] In some embodiments, provided herein is an anti-CD20 single domain antibody (sdAb) comprising a CDR1 comprising the amino acid sequence of SEQ ID NO: 180; a CDR2 comprising the amino acid sequence of SEQ ID NO: 181; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 182. In some embodiments, the anti-CD20 single domain antibody is camelid. In some embodiments, the anti-CD20 single domain antibody is humanized. In some embodiments, the anti-CD20 single domain antibody comprises an acceptor human framework,

e.g., a human immunoglobulin framework or a human consensus framework.

[00169] In some embodiments, the single domain antibody further comprises one or more framework region(s) of VHH-273, VHH-283, VHH-313, VHH-440, VHH-466, VHH-496, VHH-653, huVHH-253, huVHH-256, huVHH-260, huVHH-746, huVHH-750, huVHH-753, huVHH-836, huVHH-840, huVHH-843, huVHH-846, 2082H1, 2082H2, 2082H3, 2082H4, 2082H5, 2082H6, VHH-623, VHH-640, and/or VHH-657. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 41. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 42. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 43. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 44. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 45. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 46. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 47. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 48. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 49. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 50. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 51. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 52. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 53. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 54. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 55. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 56. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 57. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain

comprising the sequence of SEQ ID NO: 58. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 59. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 60. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 61. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 62. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 63. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 183. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 184. In some embodiments, the single domain antibody comprises one or more framework(s) derived from a VHH domain comprising the sequence of SEQ ID NO: 185.

[00170] In some embodiments, the single domain antibody provided herein is a humanized single domain antibody. In some embodiments, humanized single domain antibodies can be generated using the method exemplified in the Section 6 below or the methods described in the section below.

[00171] Framework regions described herein are determined based upon the boundaries of the CDR numbering system. In other words, if the CDRs are determined by, *e.g.*, Kabat, IMGT, or Chothia, then the framework regions are the amino acid residues surrounding the CDRs in the variable region in the format, from the N-terminus to C-terminus: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4. For example, FR1 is defined as the amino acid residues N-terminal to the CDR1 amino acid residues as defined by, *e.g.*, the Kabat numbering system, the IMGT numbering system, or the Chothia numbering system, FR2 is defined as the amino acid residues between CDR1 and CDR2 amino acid residues as defined by, *e.g.*, the Kabat numbering system, the IMGT numbering system, or the Chothia numbering system, FR3 is defined as the amino acid residues between CDR2 and CDR3 amino acid residues as defined by, *e.g.*, the Kabat numbering system, the IMGT numbering system, or the Chothia numbering system, and FR4 is defined as the amino acid residues C-terminal to the CDR3 amino acid residues as defined by, *e.g.*, the Kabat numbering system, the IMGT numbering system, or the Chothia numbering system.

[00172] In some embodiments, there is provided an isolated anti-CD20 single domain antibody comprising a VHH domain having the amino acid sequence of SEQ ID NO: 41. In some embodiments, there is provided a polypeptide comprising the amino acid sequence of SEQ ID NO: 41. In some embodiments, there is provided an isolated anti-CD20 single domain antibody

[illegible]

[illegible]

comprising a VHH domain having the amino acid sequence of SEQ ID NO: 185. In some embodiments, there is provided a polypeptide comprising the amino acid sequence of SEQ ID NO: 185.

[00173] In certain embodiments, an antibody described herein or an antigen-binding fragment thereof comprises amino acid sequences with certain percent identity relative to any one of antibodies VHH-273, VHH-283, VHH-313, VHH-440, VHH-466, VHH-496, VHH-653, huVHH-253, huVHH-256, huVHH-260, huVHH-746, huVHH-750, huVHH-753, huVHH-836, huVHH-840, huVHH-843, huVHH-846, 2082H1, 2082H2, 2082H3, 2082H4, 2082H5, 2082H6, VHH-623, VHH-640, and VHH-657.

[00174] The determination of percent identity between two sequences (*e.g.*, amino acid sequences or nucleic acid sequences) can be accomplished using a mathematical algorithm. A non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, Proc. Natl. Acad. Sci. U.S.A. 87:2264 2268 (1990), modified as in Karlin and Altschul, Proc. Natl. Acad. Sci. U.S.A. 90:5873 5877 (1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul *et al.*, J. Mol. Biol. 215:403 (1990). BLAST nucleotide searches can be performed with the NBLAST nucleotide program parameters set, *e.g.*, for score=100, word length=12 to obtain nucleotide sequences homologous to a nucleic acid molecules described herein. BLAST protein searches can be performed with the XBLAST program parameters set, *e.g.*, to score 50, word length=3 to obtain amino acid sequences homologous to a protein molecule described herein. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, Nucleic Acids Res. 25:3389 3402 (1997). Alternatively, PSI BLAST can be used to perform an iterated search which detects distant relationships between molecules (*Id.*). When utilizing BLAST, Gapped BLAST, and PSI Blast programs, the default parameters of the respective programs (*e.g.*, of XBLAST and NBLAST) can be used (see, *e.g.*, National Center for Biotechnology Information (NCBI) on the worldwide web, ncbi.nlm.nih.gov). Another non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS 4:11-17 (1998). Such an algorithm is incorporated in the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, typically only exact matches are counted.

[00175] In some embodiments, there is provided an anti-CD20 single domain antibody comprising a VHH domain having at least about any one of 75%, 80%, 85%, 86%, 87%, 88%,

SEQ ID NO: 63, wherein the single domain antibody binds to CD20. In certain embodiments, the single domain antibody described herein comprises a VHH domain having at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 183, wherein the single domain antibody binds to CD20. In certain embodiments, the single domain antibody described herein comprises a VHH domain having at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 184, wherein the single domain antibody binds to CD20. In certain embodiments, the single domain antibody described herein comprises a VHH domain having at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 185, wherein the single domain antibody binds to CD20.

[00177] In some embodiments, functional epitopes can be mapped, *e.g.*, by combinatorial alanine scanning, to identify amino acids in the CD20 protein that are necessary for interaction with anti-CD20 single domain antibodies provided herein. In some embodiments, conformational and crystal structure of anti-CD20 single domain antibody bound to CD20 may be employed to identify the epitopes. In some embodiments, the present disclosure provides an antibody that specifically binds to the same epitope as any of the anti-CD20 single domain antibodies provided herein. For example, in some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 41. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 42. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 43. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 44. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 45. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 46. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 47. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 48. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20

single domain antibody comprising the amino acid sequence of SEQ ID NO: 49. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 50. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 51. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 52. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 53. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 54. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 55. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 56. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 57. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 58. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 59. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 60. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 61. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 62. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 63. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 183. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 184. In some embodiments, an antibody is provided that binds to the same epitope as an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 185.

[00178] In some embodiments, provided herein is an anti-CD20 antibody, or antigen binding fragment thereof, that specifically binds to CD20 competitively with any one of the anti-CD20 single domain antibodies described herein. In some embodiments, competitive binding may be

[illegible]

domain antibody comprising the amino acid sequence of SEQ ID NO: 58. In some embodiments, an antibody is provided that specifically binds to CD20 competitively with an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 59. In some embodiments, an antibody is provided that specifically binds to CD20 competitively with an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 60. In some embodiments, an antibody is provided that specifically binds to CD20 competitively with an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 61. In some embodiments, an antibody is provided that specifically binds to CD20 competitively with an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 62. In some embodiments, an antibody is provided that specifically binds to CD20 competitively with an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 63. In some embodiments, an antibody is provided that specifically binds to CD20 competitively with an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 183. In some embodiments, an antibody is provided that specifically binds to CD20 competitively with an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 184. In some embodiments, an antibody is provided that specifically binds to CD20 competitively with an anti-CD20 single domain antibody comprising the amino acid sequence of SEQ ID NO: 185.

[00179] In some embodiments, provided herein is a CD20 binding protein comprising any one of the anti-CD20 single domain antibodies described above. In some embodiments, the CD20 binding protein is a monoclonal antibody, including a camelid, chimeric, humanized or human antibody. In some embodiments, the anti-CD20 antibody is an antibody fragment, *e.g.*, a VHH fragment. In some embodiments, the anti-CD20 antibody is a full-length heavy-chain only antibody comprising an Fc region of any antibody class or isotype, such as IgG1 or IgG4. In some embodiments, the Fc region has reduced or minimized effector function. In some embodiments, the CD20 binding protein is a fusion protein comprising the anti-CD20 single domain antibody provided herein. In other embodiments, the CD20 binding protein is a multispecific antibody comprising the anti-CD20 single domain antibody provided herein. Other exemplary CD20 binding molecules are described in more detail in the following sections.

[00180] In some embodiments, the anti-CD20 antibody (such as anti-CD20 single domain antibody) or antigen binding protein according to any of the above embodiments may incorporate any of the features, singly or in combination, as described in Sections 5.2.2 to 5.2.7 below.

5.2.2. Humanized Single Domain Antibodies

[00181] The single domain antibodies described herein include humanized single domain antibodies. General strategies to humanize single domain antibodies from Camelidae species have been described (see, *e.g.*, Vincke *et al.*, J. Biol. Chem., 284(5):3273-3284 (2009)) and may be useful for producing humanized VHH domains as disclosed herein. The design of humanized

single domain antibodies from Camelidae species may include the hallmark residues in the VHH, such as residues 11, 37, 44, 45 and 47 (residue numbering according to Kabat) (Muyldermans, *Reviews Mol Biotech* 74:277-302 (2001)).

[00182] Humanized antibodies, such as the humanized single domain antibodies disclosed herein can also be produced using a variety of techniques known in the art, including but not limited to, CDR-grafting (European Patent No. EP 239,400; International publication No. WO 91/09967; and U.S. Patent Nos. 5,225,539, 5,530,101, and 5,585,089), veneering or resurfacing (European Patent Nos. EP 592,106 and EP 519,596; Padlan, *Molecular Immunology* 28(4/5):489-498 (1991); Studnicka *et al.*, *Protein Engineering* 7(6):805-814 (1994); and Roguska *et al.*, *PNAS* 91:969-973 (1994)), chain shuffling (U.S. Patent No. 5,565,332), and techniques disclosed in, *e.g.*, U.S. Pat. No. 6,407,213, U.S. Pat. No. 5,766,886, WO 9317105, Tan *et al.*, *J. Immunol.* 169:1119-25 (2002), Caldas *et al.*, *Protein Eng.* 13(5):353-60 (2000), Morea *et al.*, *Methods* 20(3):267-79 (2000), Baca *et al.*, *J. Biol. Chem.* 272(16):10678-84 (1997), Roguska *et al.*, *Protein Eng.* 9(10):895-904 (1996), Couto *et al.*, *Cancer Res.* 55 (23 Supp):5973s- 5977s (1995), Couto *et al.*, *Cancer Res.* 55(8):1717-22 (1995), Sandhu JS, *Gene* 150(2):409-10 (1994), and Pedersen *et al.*, *J. Mol. Biol.* 235(3):959-73 (1994). *See also* U.S. Patent Pub. No. US 2005/0042664 A1 (Feb. 24, 2005), each of which is incorporated by reference herein in its entirety.

[00183] In some embodiments, single domain antibodies provided herein can be humanized single domain antibodies that bind to CD20, including human CD20. For example, humanized single chain antibodies of the present disclosure may comprise one or more CDRs set forth in SEQ ID NOs: 41-63 and 183-185. Various methods for humanizing non-human antibodies are known in the art. For example, a humanized antibody can have one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as “import” residues, which are typically taken from an “import” variable domain. Humanization may be performed, for example, following the method of Jones *et al.*, *Nature* 321:522-25 (1986); Riechmann *et al.*, *Nature* 332:323-27 (1988); and Verhoeven *et al.*, *Science* 239:1534-36 (1988)), by substituting hypervariable region sequences for the corresponding sequences of a human antibody. In a specific embodiment, humanization of the single domain antibody provided herein is performed as described in Section 6 below.

[00184] In some cases, the humanized antibodies are constructed by CDR grafting, in which the amino acid sequences of the CDRs of the parent non-human antibody are grafted onto a human antibody framework. For example, Padlan *et al.* determined that only about one third of the residues in the CDRs actually contact the antigen, and termed these the “specificity determining residues,” or SDRs (Padlan *et al.*, *FASEB J.* 9:133-39 (1995)). In the technique of SDR grafting, only the SDR residues are grafted onto the human antibody framework (*see, e.g.*, Kashmiri *et al.*,

Methods 36:25-34 (2005)).

[00185] The choice of human variable domains to be used in making the humanized antibodies can be important to reduce antigenicity. For example, according to the so-called “best-fit” method, the sequence of the variable domain of a non-human antibody is screened against the entire library of known human variable-domain sequences. The human sequence that is closest to that of the non-human antibody may be selected as the human framework for the humanized antibody (Sims *et al.*, J. Immunol. 151:2296-308 (1993); and Chothia *et al.*, J. Mol. Biol. 196:901-17 (1987)). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, Proc. Natl. Acad. Sci. USA 89:4285-89 (1992); and Presta *et al.*, J. Immunol. 151:2623-32 (1993)). In some cases, the framework is derived from the consensus sequences of the most abundant human subclasses, V_L6 subgroup I (V_L6I) and V_H subgroup III (V_HIII). In another method, human germline genes are used as the source of the framework regions.

[00186] In an alternative paradigm based on comparison of CDRs, called superhumanization, FR homology is irrelevant. The method consists of comparison of the non-human sequence with the functional human germline gene repertoire. Those genes encoding the same or closely related canonical structures to the murine sequences are then selected. Next, within the genes sharing the canonical structures with the non-human antibody, those with highest homology within the CDRs are chosen as FR donors. Finally, the non-human CDRs are grafted onto these FRs (*see, e.g.*, Tan *et al.*, J. Immunol. 169:1119-25 (2002)).

[00187] It is further generally desirable that antibodies be humanized with retention of their affinity for the antigen and other favorable biological properties. To achieve this goal, according to one method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. These include, for example, WAM (Whitelegg and Rees, Protein Eng. 13:819-24 (2002)), Modeller (Sali and Blundell, J. Mol. Biol. 234:779-815 (1993)), and Swiss PDB Viewer (Guex and Peitsch, Electrophoresis 18:2714-23 (1997)). Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *e.g.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are

directly and most substantially involved in influencing antigen binding.

[00188] Another method for antibody humanization is based on a metric of antibody humanness termed Human String Content (HSC). This method compares the mouse sequence with the repertoire of human germline genes, and the differences are scored as HSC. The target sequence is then humanized by maximizing its HSC rather than using a global identity measure to generate multiple diverse humanized variants (Lazar *et al.*, Mol. Immunol. 44:1986-98 (2007)).

[00189] In addition to the methods described above, empirical methods may be used to generate and select humanized antibodies. These methods include those that are based upon the generation of large libraries of humanized variants and selection of the best clones using enrichment technologies or high throughput screening techniques. Antibody variants may be isolated from phage, ribosome, and yeast display libraries as well as by bacterial colony screening (*see, e.g.*, Hoogenboom, Nat. Biotechnol. 23:1105-16 (2005); Dufner *et al.*, Trends Biotechnol. 24:523-29 (2006); Feldhaus *et al.*, Nat. Biotechnol. 21:163-70 (2003); and Schlapschy *et al.*, Protein Eng. Des. Sel. 17:847-60 (2004)).

[00190] In the FR library approach, a collection of residue variants are introduced at specific positions in the FR followed by screening of the library to select the FR that best supports the grafted CDR. The residues to be substituted may include some or all of the “Vernier” residues identified as potentially contributing to CDR structure (*see, e.g.*, Foote and Winter, J. Mol. Biol. 224:487-99 (1992)), or from the more limited set of target residues identified by Baca *et al.* J. Biol. Chem. 272:10678-84 (1997).

[00191] In FR shuffling, whole FRs are combined with the non-human CDRs instead of creating combinatorial libraries of selected residue variants (*see, e.g.*, Dall’Acqua *et al.*, Methods 36:43-60 (2005)). A one-step FR shuffling process may be used. Such a process has been shown to be efficient, as the resulting antibodies exhibited improved biochemical and physicochemical properties including enhanced expression, increased affinity, and thermal stability (*see, e.g.*, Damschroder *et al.*, Mol. Immunol. 44:3049-60 (2007)).

[00192] The “humaneering” method is based on experimental identification of essential minimum specificity determinants (MSDs) and is based on sequential replacement of non-human fragments into libraries of human FRs and assessment of binding. This methodology typically results in epitope retention and identification of antibodies from multiple subclasses with distinct human V-segment CDRs.

[00193] The “human engineering” method involves altering a non-human antibody or antibody fragment by making specific changes to the amino acid sequence of the antibody so as to produce a modified antibody with reduced immunogenicity in a human that nonetheless retains the desirable binding properties of the original non-human antibodies. Generally, the technique involves classifying amino acid residues of a non-human antibody as “low risk,” “moderate risk,”

or “high risk” residues. The classification is performed using a global risk/reward calculation that evaluates the predicted benefits of making particular substitution (*e.g.*, for immunogenicity in humans) against the risk that the substitution will affect the resulting antibody’s folding. The particular human amino acid residue to be substituted at a given position (*e.g.*, low or moderate risk) of a non-human antibody sequence can be selected by aligning an amino acid sequence from the non-human antibody’s variable regions with the corresponding region of a specific or consensus human antibody sequence. The amino acid residues at low or moderate risk positions in the non-human sequence can be substituted for the corresponding residues in the human antibody sequence according to the alignment. Techniques for making human engineered proteins are described in greater detail in Studnicka *et al.*, Protein Engineering 7:805-14 (1994); U.S. Pat. Nos. 5,766,886; 5,770,196; 5,821,123; and 5,869,619; and PCT Publication WO 93/11794.

[00194] A composite human antibody can be generated using, for example, Composite Human Antibody™ technology (Antitope Ltd., Cambridge, United Kingdom). To generate composite human antibodies, variable region sequences are designed from fragments of multiple human antibody variable region sequences in a manner that avoids T cell epitopes, thereby minimizing the immunogenicity of the resulting antibody.

[00195] A deimmunized antibody is an antibody in which T-cell epitopes have been removed. Methods for making deimmunized antibodies have been described. *See, e.g.*, Jones *et al.*, Methods Mol Biol. 525:405-23 (2009), xiv, and De Groot *et al.*, Cell. Immunol. 244:148-153(2006)). Deimmunized antibodies comprise T-cell epitope-depleted variable regions and human constant regions. Briefly, variable regions of an antibody are cloned and T-cell epitopes are subsequently identified by testing overlapping peptides derived from the variable regions of the antibody in a T cell proliferation assay. T cell epitopes are identified via *in silico* methods to identify peptide binding to human MHC class II. Mutations are introduced in the variable regions to abrogate binding to human MHC class II. Mutated variable regions are then utilized to generate the deimmunized antibody.

5.2.3. Single Domain Antibody Variants

[00196] In some embodiments, amino acid sequence modification(s) of the single domain antibodies that bind to CD20 described herein are contemplated. For example, it may be desirable to optimize the binding affinity and/or other biological properties of the antibody, including but not limited to specificity, thermostability, expression level, effector functions, glycosylation, reduced immunogenicity, or solubility. Thus, in addition to the single domain antibodies that bind to CD20 described herein, it is contemplated that variants of the single domain antibodies that bind to CD20 described herein can be prepared. For example, single domain antibody variants can be prepared by introducing appropriate nucleotide changes into the

encoding DNA, and/or by synthesis of the desired antibody or polypeptide. Those skilled in the art who appreciate that amino acid changes may alter post-translational processes of the single domain antibody.

Chemical Modifications

[00197] In some embodiments, the single domain antibodies provided herein are chemically modified, for example, by the covalent attachment of any type of molecule to the single domain antibody. The antibody derivatives may include antibodies that have been chemically modified, for example, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, or conjugation to one or more immunoglobulin domains (*e.g.*, Fc or a portion of an Fc). Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to, specific chemical cleavage, acetylation, formulation, metabolic synthesis of tunicamycin, *etc.* Additionally, the antibody may contain one or more non-classical amino acids.

[00198] In some embodiments, an antibody provided herein is altered to increase or decrease the extent to which the antibody is glycosylated. Addition or deletion of glycosylation sites to an antibody may be conveniently accomplished by altering the amino acid sequence such that one or more glycosylation sites is created or removed.

[00199] When the single domain antibody provided herein is fused to an Fc region, the carbohydrate attached thereto may be altered. Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 of the CH2 domain of the Fc region. *See, e.g.*, Wright et al. *TIBTECH* 15:26-32 (1997). The oligosaccharide may include various carbohydrates, *e.g.*, mannose, N-acetyl glucosamine (GlcNAc), galactose, and sialic acid, as well as a fucose attached to a GlcNAc in the “stem” of the biantennary oligosaccharide structure. In some embodiments, modifications of the oligosaccharide in the binding molecules provided herein may be made in order to create variants with certain improved properties.

[00200] In other embodiments, when the single domain antibody provided herein is fused to an Fc region, antibody variants provided herein may have a carbohydrate structure that lacks fucose attached (directly or indirectly) to said Fc region. For example, the amount of fucose in such antibody may be from 1% to 80%, from 1% to 65%, from 5% to 65% or from 20% to 40%. The amount of fucose is determined by calculating the average amount of fucose within the sugar chain at Asn297, relative to the sum of all glycostructures attached to Asn 297 (*e.g.*, complex, hybrid and high mannose structures) as measured by MALDI-TOF mass spectrometry, as described in WO 2008/077546, for example. Asn297 refers to the asparagine residue located at about position 297 in the Fc region (EU numbering of Fc region residues); however, Asn297 may also be located about ± 3 amino acids upstream or downstream of position 297, *i.e.*, between

positions 294 and 300, due to minor sequence variations in antibodies. Such fucosylation variants may have improved ADCC function. *See, e.g.*, US Patent Publication Nos. US 2003/0157108 and US 2004/0093621. Examples of publications related to “defucosylated” or “fucose-deficient” antibody variants include: US 2003/0157108; WO 2000/61739; WO 2001/29246; US 2003/0115614; US 2002/0164328; US 2004/0093621; US 2004/0132140; US 2004/0110704; US 2004/0110282; US 2004/0109865; WO 2003/085119; WO 2003/084570; WO 2005/035586; WO 2005/035778; WO2005/053742; WO2002/031140; Okazaki et al. *J. Mol. Biol.* 336:1239-1249 (2004); Yamane-Ohnuki et al. *Biotech. Bioeng.* 87: 614 (2004). Examples of cell lines capable of producing defucosylated antibodies include Lec13 CHO cells deficient in protein fucosylation (Ripka et al. *Arch. Biochem. Biophys.* 249:533-545 (1986); US Patent Application No. US 2003/0157108; and WO 2004/056312, especially at Example 11), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, *FUT8*, knockout CHO cells (*see, e.g.*, Yamane-Ohnuki et al. *Biotech. Bioeng.* 87: 614 (2004); Kanda, Y. et al., *Biotechnol. Bioeng.*, 94(4):680-688 (2006); and WO2003/085107).

[00201] The binding molecules comprising a single domain antibody provided herein are further provided with bisected oligosaccharides, *e.g.*, in which a biantennary oligosaccharide attached to the Fc region is bisected by GlcNAc. Such variants may have reduced fucosylation and/or improved ADCC function. Examples of such variants are described, *e.g.*, in WO 2003/011878 (Jean-Mairet et al.); US Patent No. 6,602,684 (Umana et al.); and US 2005/0123546 (Umana et al.). Variants with at least one galactose residue in the oligosaccharide attached to the Fc region are also provided. Such variants may have improved CDC function. Such variants are described, *e.g.*, in WO 1997/30087; WO 1998/58964; and WO 1999/22764.

[00202] In molecules that comprise the present single domain antibody and an Fc region, one or more amino acid modifications may be introduced into the Fc region, thereby generating an Fc region variant. The Fc region variant may comprise a human Fc region sequence (*e.g.*, a human IgG1, IgG2, IgG3 or IgG4 Fc region) comprising an amino acid modification (*e.g.* a substitution) at one or more amino acid positions.

[00203] In some embodiments, the present application contemplates variants that possesses some but not all effector functions, which make it a desirable candidate for applications in which the half life of the binding molecule *in vivo* is important yet certain effector functions (such as complement and ADCC) are unnecessary or deleterious. *In vitro* and/or *in vivo* cytotoxicity assays can be conducted to confirm the reduction/depletion of CDC and/or ADCC activities. For example, Fc receptor (FcR) binding assays can be conducted to ensure that the binding molecule lacks FcγR binding (hence likely lacking ADCC activity), but retains FcRn binding ability. Non-limiting examples of *in vitro* assays to assess ADCC activity of a molecule of interest is described in U.S. Patent No. 5,500,362 (*see, e.g.* Hellstrom, I. et al. *Proc. Nat'l Acad. Sci. USA*

83:7059-7063 (1986)) and Hellstrom, I et al., *Proc. Nat'l Acad. Sci. USA* 82:1499-1502 (1985); 5,821,337 (*see* Bruggemann, M. et al., *J. Exp. Med.* 166:1351-1361 (1987)). Alternatively, non-radioactive assays methods may be employed (*see*, for example, ACTITM non-radioactive cytotoxicity assay for flow cytometry (CellTechnology, Inc. Mountain View, CA; and CytoTox 96® non-radioactive cytotoxicity assay (Promega, Madison, WI). Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, *e.g.*, in an animal model such as that disclosed in Clynes et al. *Proc. Nat'l Acad. Sci. USA* 95:652-656 (1998). C1q binding assays may also be carried out to confirm that the antibody is unable to bind C1q and hence lacks CDC activity. *See, e.g.*, C1q and C3c binding ELISA in WO 2006/029879 and WO 2005/100402. To assess complement activation, a CDC assay may be performed (*see*, for example, Gazzano-Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996); Cragg, M.S. et al., *Blood* 101:1045-1052 (2003); and Cragg, M.S. and M.J. Glennie, *Blood* 103:2738-2743 (2004)). FcRn binding and *in vivo* clearance/half life determinations can also be performed using methods known in the art (*see, e.g.*, Petkova, S.B. et al., *Int'l. Immunol.* 18(12):1759-1769 (2006)).

[00204] Binding molecules with reduced effector function include those with substitution of one or more of Fc region residues 238, 265, 269, 270, 297, 327 and 329 (U.S. Patent No. 6,737,056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called "DANA" Fc mutant with substitution of residues 265 and 297 to alanine (US Patent No. 7,332,581).

[00205] Certain variants with improved or diminished binding to FcRs are described. (*See, e.g.*, U.S. Patent No. 6,737,056; WO 2004/056312, and Shields et al., *J. Biol. Chem.* 9(2): 6591-6604 (2001).)

[00206] In some embodiments, a variant comprises an Fc region with one or more amino acid substitutions which improve ADCC, *e.g.*, substitutions at positions 298, 333, and/or 334 of the Fc region (EU numbering of residues). In some embodiments, alterations are made in the Fc region that result in altered (*i.e.*, either improved or diminished) C1q binding and/or Complement Dependent Cytotoxicity (CDC), *e.g.*, as described in US Patent No. 6,194,551, WO 99/51642, and Idusogie et al. *J. Immunol.* 164: 4178-4184 (2000).

[00207] Binding molecules with increased half lives and improved binding to the neonatal Fc receptor (FcRn), which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., *J. Immunol.* 117:587 (1976) and Kim et al., *J. Immunol.* 24:249 (1994)), are described in US2005/0014934A1 (Hinton et al.). Those molecules comprise an Fc region with one or more substitutions therein which improve binding of the Fc region to FcRn. Such Fc variants include those with substitutions at one or more of Fc region residues: 238, 256, 265, 272, 286, 303, 305,

307, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 413, 424 or 434, *e.g.*, substitution of Fc region residue 434 (US Patent No. 7,371,826). See also Duncan & Winter, *Nature* 322:738-40 (1988); U.S. Patent No. 5,648,260; U.S. Patent No. 5,624,821; and WO 94/29351 concerning other examples of Fc region variants.

[00208] In some embodiments, it may be desirable to create cysteine engineered antibodies, in which one or more residues of an antibody are substituted with cysteine residues. In some embodiments, the substituted residues occur at accessible sites of the antibody. By substituting those residues with cysteine, reactive thiol groups are thereby positioned at accessible sites of the antibody and may be used to conjugate the antibody to other moieties, such as drug moieties or linker-drug moieties, to create an immunoconjugate, as described further herein.

Substitutions, Deletions, or Insertions

[00209] Variations may be a substitution, deletion, or insertion of one or more codons encoding the single domain antibody or polypeptide that results in a change in the amino acid sequence as compared with the original antibody or polypeptide. Sites of interest for substitutional mutagenesis include the CDRs and FRs.

[00210] Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, *e.g.*, conservative amino acid replacements. Standard techniques known to those of skill in the art can be used to introduce mutations in the nucleotide sequence encoding a molecule provided herein, including, for example, site-directed mutagenesis and PCR-mediated mutagenesis which results in amino acid substitutions. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. In certain embodiments, the substitution, deletion, or insertion includes fewer than 25 amino acid substitutions, fewer than 20 amino acid substitutions, fewer than 15 amino acid substitutions, fewer than 10 amino acid substitutions, fewer than 5 amino acid substitutions, fewer than 4 amino acid substitutions, fewer than 3 amino acid substitutions, or fewer than 2 amino acid substitutions relative to the original molecule. In a specific embodiment, the substitution is a conservative amino acid substitution made at one or more predicted non-essential amino acid residues. The variation allowed may be determined by systematically making insertions, deletions, or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the parental antibodies.

[00211] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing multiple residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue.

[00212] Single domain antibodies generated by conservative amino acid substitutions are included in the present disclosure. In a conservative amino acid substitution, an amino acid

residue is replaced with an amino acid residue having a side chain with a similar charge. As described above, families of amino acid residues having side chains with similar charges have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Alternatively, mutations can be introduced randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for biological activity to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed and the activity of the protein can be determined. Conservative (*e.g.*, within an amino acid group with similar properties and/or side chains) substitutions may be made, so as to maintain or not significantly change the properties. Exemplary substitutions are shown in Table 3 below.

Table 3. Amino Acid Substitutions

Original Residue	Exemplary Substitutions	Original Residue	Exemplary Substitutions
Ala (A)	Val; Leu; Ile	Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe
Arg (R)	Lys; Gln; Asn	Lys (K)	Arg; Gln; Asn
Asn (N)	Gln; His; Asp, Lys; Arg	Met (M)	Leu; Phe; Ile
Asp (D)	Glu; Asn	Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr
Cys (C)	Ser; Ala	Pro (P)	Ala
Gln (Q)	Asn; Glu	Ser (S)	Thr
Glu (E)	Asp; Gln	Thr (T)	Val; Ser
Gly (G)	Ala	Trp (W)	Tyr; Phe
His (H)	Asn; Gln; Lys; Arg	Tyr (Y)	Trp; Phe; Thr; Ser
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine

[00213] Amino acids may be grouped according to similarities in the properties of their side chains (*see, e.g.*, Lehninger, Biochemistry 73-75 (2d ed. 1975)): (1) non-polar: Ala (A), Val (V), Leu (L), Ile (I), Pro (P), Phe (F), Trp (W), Met (M); (2) uncharged polar: Gly (G), Ser (S), Thr (T), Cys (C), Tyr (Y), Asn (N), Gln (Q); (3) acidic: Asp (D), Glu (E); and (4) basic: Lys (K), Arg (R), His(H). Alternatively, naturally occurring residues may be divided into groups based on common side-chain properties: (1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile; (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln; (3) acidic: Asp, Glu; (4) basic: His, Lys, Arg; (5) residues that influence chain orientation: Gly, Pro; and (6) aromatic: Trp, Tyr, Phe. For example, any cysteine residue not involved in maintaining the proper conformation of the single domain

antibody also may be substituted, for example, with another amino acid, such as alanine or serine, to improve the oxidative stability of the molecule and to prevent aberrant crosslinking. Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

[00214] One type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (*e.g.*, a humanized or human antibody). Generally, the resulting variant(s) selected for further study will have modifications (*e.g.*, improvements) in certain biological properties (*e.g.*, increased affinity, reduced immunogenicity) relative to the parent antibody and/or will have substantially retained certain biological properties of the parent antibody. An exemplary substitutional variant is an affinity matured antibody, which may be conveniently generated, *e.g.*, using phage display-based affinity maturation techniques such as those described herein. Briefly, one or more CDR residues are mutated and the variant antibodies displayed on phage and screened for a particular biological activity (*e.g.* binding affinity).

[00215] Alterations (*e.g.*, substitutions) may be made in CDRs, *e.g.*, to improve antibody affinity. Such alterations may be made in CDR “hotspots,” *i.e.*, residues encoded by codons that undergo mutation at high frequency during the somatic maturation process (*see, e.g.*, Chowdhury, *Methods Mol. Biol.* 207:179-196 (2008)), and/or SDRs (a-CDRs), with the resulting variant antibody or fragment thereof being tested for binding affinity. Affinity maturation by constructing and reselecting from secondary libraries has been described, *e.g.*, in Hoogenboom et al. in *Methods in Molecular Biology* 178:1-37 (O’Brien et al., ed., Human Press, Totowa, NJ, (2001).) In some embodiments of affinity maturation, diversity is introduced into the variable genes chosen for maturation by any of a variety of methods (*e.g.*, error-prone PCR, chain shuffling, or oligonucleotide-directed mutagenesis). A secondary library is then created. The library is then screened to identify any antibody variants with the desired affinity. Another method to introduce diversity involves CDR-directed approaches, in which several CDR residues (*e.g.*, 4-6 residues at a time) are randomized. CDR residues involved in antigen binding may be specifically identified, *e.g.*, using alanine scanning mutagenesis or modeling. More detailed description regarding affinity maturation is provided in the section below.

[00216] In some embodiments, substitutions, insertions, or deletions may occur within one or more CDRs so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative alterations (*e.g.*, conservative substitutions as provided herein) that do not substantially reduce binding affinity may be made in CDRs. In some embodiments of the variant VHH sequences provided herein, each CDR either is unaltered, or contains no more than one, two or three amino acid substitutions.

[00217] A useful method for identification of residues or regions of an antibody that may be

targeted for mutagenesis is called “alanine scanning mutagenesis” as described by Cunningham and Wells, *Science*, 244:1081-1085 (1989). In this method, a residue or group of target residues (*e.g.*, charged residues such as Arg, Asp, His, Lys, and Glu) are identified and replaced by a neutral or negatively charged amino acid (*e.g.*, alanine or polyalanine) to determine whether the interaction of the antibody with antigen is affected. Further substitutions may be introduced at the amino acid locations demonstrating functional sensitivity to the initial substitutions. Alternatively, or additionally, a crystal structure of an antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues may be targeted or eliminated as candidates for substitution. Variants may be screened to determine whether they contain the desired properties.

[00218] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (*e.g.*, for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

[00219] The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis (*see, e.g.*, Carter, *Biochem J.* 237:1-7 (1986); and Zoller *et al.*, *Nucl. Acids Res.* 10:6487-500 (1982)), cassette mutagenesis (*see, e.g.*, Wells *et al.*, *Gene* 34:315-23 (1985)), or other known techniques can be performed on the cloned DNA to produce the single domain antibody variant DNA.

5.2.4. *In vitro* Affinity Maturation

[00220] In some embodiments, antibody variants having an improved property such as affinity, stability, or expression level as compared to a parent antibody may be prepared by *in vitro* affinity maturation. Like the natural prototype, *in vitro* affinity maturation is based on the principles of mutation and selection. Libraries of antibodies are displayed on the surface of an organism (*e.g.*, phage, bacteria, yeast, or mammalian cell) or in association (*e.g.*, covalently or non-covalently) with their encoding mRNA or DNA. Affinity selection of the displayed antibodies allows isolation of organisms or complexes carrying the genetic information encoding the antibodies. Two or three rounds of mutation and selection using display methods such as phage display usually results in antibody fragments with affinities in the low nanomolar range. Affinity matured antibodies can have nanomolar or even picomolar affinities for the target antigen.

[00221] Phage display is a widespread method for display and selection of antibodies. The antibodies are displayed on the surface of Fd or M13 bacteriophages as fusions to the

bacteriophage coat protein. Selection involves exposure to antigen to allow phage-displayed antibodies to bind their targets, a process referred to as “panning.” Phage bound to antigen are recovered and used to infect bacteria to produce phage for further rounds of selection. For review, see, for example, Hoogenboom, *Methods. Mol. Biol.* 178:1-37 (2002); and Bradbury and Marks, *J. Immunol. Methods* 290:29-49 (2004).

[00222] In a yeast display system (*see, e.g.,* Boder *et al.*, *Nat. Biotech.* 15:553–57 (1997); and Chao *et al.*, *Nat. Protocols* 1:755-68 (2006)), the antibody may be fused to the adhesion subunit of the yeast agglutinin protein Aga2p, which attaches to the yeast cell wall through disulfide bonds to Aga1p. Display of a protein via Aga2p projects the protein away from the cell surface, minimizing potential interactions with other molecules on the yeast cell wall. Magnetic separation and flow cytometry are used to screen the library to select for antibodies with improved affinity or stability. Binding to a soluble antigen of interest is determined by labeling of yeast with biotinylated antigen and a secondary reagent such as streptavidin conjugated to a fluorophore. Variations in surface expression of the antibody can be measured through immunofluorescence labeling of either the hemagglutinin or c-Myc epitope tag flanking the single chain antibody (*e.g.,* scFv). Expression has been shown to correlate with the stability of the displayed protein, and thus antibodies can be selected for improved stability as well as affinity (*see, e.g.,* Shusta *et al.*, *J. Mol. Biol.* 292:949-56 (1999)). An additional advantage of yeast display is that displayed proteins are folded in the endoplasmic reticulum of the eukaryotic yeast cells, taking advantage of endoplasmic reticulum chaperones and quality-control machinery. Once maturation is complete, antibody affinity can be conveniently “titrated” while displayed on the surface of the yeast, eliminating the need for expression and purification of each clone. A theoretical limitation of yeast surface display is the potentially smaller functional library size than that of other display methods; however, a recent approach uses the yeast cells’ mating system to create combinatorial diversity estimated to be 10^{14} in size (*see, e.g.,* U.S. Pat. Publication 2003/0186374; and Blaise *et al.*, *Gene* 342:211–18 (2004)).

[00223] In ribosome display, antibody-ribosome-mRNA (ARM) complexes are generated for selection in a cell-free system. The DNA library coding for a particular library of antibodies is genetically fused to a spacer sequence lacking a stop codon. This spacer sequence, when translated, is still attached to the peptidyl tRNA and occupies the ribosomal tunnel, and thus allows the protein of interest to protrude out of the ribosome and fold. The resulting complex of mRNA, ribosome, and protein can bind to surface-bound ligand, allowing simultaneous isolation of the antibody and its encoding mRNA through affinity capture with the ligand. The ribosome-bound mRNA is then reverse transcribed back into cDNA, which can then undergo mutagenesis and be used in the next round of selection (*see, e.g.,* Fukuda *et al.*, *Nucleic Acids Res.* 34:e127 (2006)). In mRNA display, a covalent bond between antibody and mRNA is established using

puromycin as an adaptor molecule (Wilson *et al.*, Proc. Natl. Acad. Sci. USA 98:3750-55 (2001)).

[00224] As these methods are performed entirely *in vitro*, they provide two main advantages over other selection technologies. First, the diversity of the library is not limited by the transformation efficiency of bacterial cells, but only by the number of ribosomes and different mRNA molecules present in the test tube. Second, random mutations can be introduced easily after each selection round, for example, by non-proofreading polymerases, as no library must be transformed after any diversification step.

[00225] In some embodiments, mammalian display systems may be used.

[00226] Diversity may also be introduced into the CDRs of the antibody libraries in a targeted manner or via random introduction. The former approach includes sequentially targeting all the CDRs of an antibody via a high or low level of mutagenesis or targeting isolated hot spots of somatic hypermutations (*see, e.g.*, Ho *et al.*, J. Biol. Chem. 280:607-17 (2005)) or residues suspected of affecting affinity on experimental basis or structural reasons. Diversity may also be introduced by replacement of regions that are naturally diverse via DNA shuffling or similar techniques (*see, e.g.*, Lu *et al.*, J. Biol. Chem. 278:43496-507 (2003); U.S. Pat. Nos. 5,565,332 and 6,989,250). Alternative techniques target hypervariable loops extending into framework-region residues (*see, e.g.*, Bond *et al.*, J. Mol. Biol. 348:699-709 (2005)) employ loop deletions and insertions in CDRs or use hybridization-based diversification (*see, e.g.*, U.S. Pat. Publication No. 2004/0005709). Additional methods of generating diversity in CDRs are disclosed, for example, in U.S. Pat. No. 7,985,840. Further methods that can be used to generate antibody libraries and/or antibody affinity maturation are disclosed, *e.g.*, in U.S. Patent Nos. 8,685,897 and 8,603,930, and U.S. Publ. Nos. 2014/0170705, 2014/0094392, 2012/0028301, 2011/0183855, and 2009/0075378, each of which are incorporated herein by reference.

[00227] Screening of the libraries can be accomplished by various techniques known in the art. For example, single domain antibodies can be immobilized onto solid supports, columns, pins, or cellulose/poly (vinylidene fluoride) membranes/other filters, expressed on host cells affixed to adsorption plates or used in cell sorting, or conjugated to biotin for capture with streptavidin-coated beads or used in any other method for panning display libraries.

[00228] For review of *in vitro* affinity maturation methods, *see, e.g.*, Hoogenboom, Nature Biotechnology 23:1105-16 (2005); Quiroz and Sinclair, Revista Ingenieria Biomedica 4:39-51 (2010); and references therein.

5.2.5. Modifications of Single Domain Antibodies

[00229] Covalent modifications of single domain antibodies are included within the scope of the present disclosure. Covalent modifications include reacting targeted amino acid residues of a single domain antibody with an organic derivatizing agent that is capable of reacting with

selected side chains or the N- or C- terminal residues of the single domain antibody. Other modifications include deamidation of glutaminyl and asparaginyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains (*see, e.g.*, Creighton, Proteins: Structure and Molecular Properties 79-86 (1983)), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

[00230] Other types of covalent modification of the single domain antibody included within the scope of this present disclosure include altering the native glycosylation pattern of the antibody or polypeptide as described above (*see, e.g.*, Beck *et al.*, Curr. Pharm. Biotechnol. 9:482-501 (2008); and Walsh, Drug Discov. Today 15:773-80 (2010)), and linking the antibody to one of a variety of nonproteinaceous polymers, *e.g.*, polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth, for example, in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192; or 4,179,337. The single domain antibody that binds to CD20 of the disclosure may also be genetically fused or conjugated to one or more immunoglobulin constant regions or portions thereof (*e.g.*, Fc) to extend half-life and/or to impart known Fc-mediated effector functions.

[00231] The single chain antibody that binds to CD20 of the present disclosure may also be modified to form chimeric molecules comprising the single chain antibody that binds to CD20 fused to another, heterologous polypeptide or amino acid sequence, for example, an epitope tag (*see, e.g.*, Terpe, Appl. Microbiol. Biotechnol. 60:523-33 (2003)) or the Fc region of an IgG molecule (*see, e.g.*, Aruffo, Antibody Fusion Proteins 221-42 (Chamow and Ashkenazi eds., 1999)). The single chain antibody that binds to CD20 may also be used to generate CD20 binding chimeric antigen receptor (CAR), as described in more detail below.

[00232] Also provided herein are fusion proteins comprising the single chain antibody that binds to CD20 of the disclosure and a heterologous polypeptide. In some embodiments, the heterologous polypeptide to which the antibody is genetically fused or chemically conjugated is useful for targeting the antibody to cells having cell surface-expressed CD20.

[00233] Also provided herein are panels of antibodies that bind to a CD20 antigen. In specific embodiments, the panels of antibodies have different association rates, different dissociation rates, different affinities for a CD20 antigen, and/or different specificities for a CD20 antigen. In some embodiments, the panels comprise or consist of about 10 to about 1000 antibodies or more. Panels of antibodies can be used, for example, in 96-well or 384-well plates, for assays such as ELISAs.

5.2.6. Preparation of Single Domain Antibodies

[00234] Methods of preparing single domain antibodies have been described. *See, e.g.*, Els

Pardon et al, *Nature Protocol*, 9(3): 674 (2014). Single domain antibodies (such as VHs) may be obtained using methods known in the art such as by immunizing a *Camelid* species (such as camel or llama) and obtaining hybridomas therefrom, or by cloning a library of single domain antibodies using molecular biology techniques known in the art and subsequent selection by ELISA with individual clones of unselected libraries or by using phage display.

[00235] Single domain antibodies provided herein may be produced by culturing cells transformed or transfected with a vector containing a single domain antibody-encoding nucleic acids. Polynucleotide sequences encoding polypeptide components of the antibody of the present disclosure can be obtained using standard recombinant techniques. Desired polynucleotide sequences may be isolated and sequenced from antibody producing cells such as hybridomas cells or B cells. Alternatively, polynucleotides can be synthesized using nucleotide synthesizer or PCR techniques. Once obtained, sequences encoding the polypeptides are inserted into a recombinant vector capable of replicating and expressing heterologous polynucleotides in host cells. Many vectors that are available and known in the art can be used for the purpose of the present disclosure. Selection of an appropriate vector will depend mainly on the size of the nucleic acids to be inserted into the vector and the particular host cell to be transformed with the vector. Host cells suitable for expressing antibodies of the present disclosure include prokaryotes such as Archaeobacteria and Eubacteria, including Gram-negative or Gram-positive organisms, eukaryotic microbes such as filamentous fungi or yeast, invertebrate cells such as insect or plant cells, and vertebrate cells such as mammalian host cell lines. Host cells are transformed with the above-described expression vectors and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. Antibodies produced by the host cells are purified using standard protein purification methods as known in the art.

[00236] Methods for antibody production including vector construction, expression, and purification are further described in Plückthun *et al.*, Antibody Engineering: Producing antibodies in Escherichia coli: From PCR to fermentation 203-52 (McCafferty *et al.* eds., 1996); Kwong and Rader, *E. coli Expression and Purification of Fab Antibody Fragments*, in Current Protocols in Protein Science (2009); Tachibana and Takekoshi, *Production of Antibody Fab Fragments in Escherichia coli*, in Antibody Expression and Production (Al-Rubeai ed., 2011); and Therapeutic Monoclonal Antibodies: From Bench to Clinic (An ed., 2009).

[00237] It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare anti-CD20 single domain antibodies. For instance, the appropriate amino acid sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques (*see, e.g.*, Stewart *et al.*, Solid-Phase Peptide Synthesis (1969); and Merrifield, J. Am. Chem. Soc. 85:2149-54 (1963)). *In vitro* protein synthesis may be performed

using manual techniques or by automation. Various portions of the anti-CD20 antibody may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the desired anti-CD20 antibody. Alternatively, antibodies may be purified from cells or bodily fluids, such as milk, of a transgenic animal engineered to express the antibody, as disclosed, for example, in U.S. Pat. Nos. 5,545,807 and 5,827,690.

[00238] Specifically, the single domain antibodies, or other CD20 binders provided herein, can be generated by immunizing llamas, performing single B-cell sorting, undertaking V-gene extraction, cloning the CD20 binders, such as VHH-Fc fusions, and then performing small scale expression and purification. Additional screening of the single domain antibodies and other molecules that bind to CD20 can be performed, including one or more of selecting for ELISA-positive, BLI-positive, and K_D less than 100 nM. These selection criteria can be combined as described in Section 6 below. Additionally, individual VHH binders (and other molecules that bind to CD20) can be assayed for their ability to bind to cells expressing CD20. Such assay can be performed using FACS analysis with cells expressing CD20, and measuring the mean fluorescence intensity (MFI) of fluorescently-labeled VHH molecules. Various aspects mentioned above are described in more details below.

Polyclonal Antibodies

[00239] Polyclonal antibodies are generally raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen to a protein that is immunogenic in the species to be immunized, *e.g.*, keyhole limpet hemocyanin (KLH), serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor, using a bifunctional or derivatizing agent, *e.g.*, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N}=\text{C}=\text{NR}$, where R and R^1 are independently lower alkyl groups. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

[00240] For example, the animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, *e.g.*, 100 μg or 5 μg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later, the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to fourteen days later, the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitable to

enhance the immune response.

Monoclonal Antibodies

[00241] Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translational modifications (*e.g.*, isomerizations, amidations) that may be present in minor amounts. Thus, the modifier “monoclonal” indicates the character of the antibody as not being a mixture of discrete antibodies.

[00242] For example, the monoclonal antibodies may be made using the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Pat. No. 4,816,567).

[00243] In the hybridoma method, an appropriate host animal is immunized to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 59-103 (Academic Press, 1986).

[00244] The immunizing agent will typically include the antigenic protein or a fusion variant thereof. Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press (1986), pp. 59-103. Immortalized cell lines are usually transformed mammalian cells. The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, parental myeloma cells. Preferred immortalized myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium.

[00245] Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. The culture medium in which the hybridoma cells are cultured can be assayed for the presence of monoclonal antibodies directed against the desired antigen. Such techniques and assays are known in the art. For example, binding affinity may be determined by the Scatchard analysis of Munson *et al.*, *Anal. Biochem.*, 107:220 (1980).

[00246] After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution procedures and grown by standard methods (Goding, *supra*). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown *in vivo* as tumors in a mammal.

[00247] The monoclonal antibodies secreted by the subclones are suitably separated from the culture medium, ascites fluid, or serum by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

[00248] Monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567, and as described above. DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, in order to synthesize monoclonal antibodies in such recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra *et al.*, *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Plickthun, *Immunol. Revs.* 130:151-188 (1992).

[00249] In a further embodiment, antibodies can be isolated from antibody phage libraries generated using the techniques described in McCafferty *et al.*, *Nature*, 348:552-554 (1990). Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991). Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.*, *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse *et al.*, *Nucl. Acids Res.*, 21:2265-2266 (1993)). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

[00250] The DNA also may be modified, for example, by substituting the coding sequence (U.S. Pat. No. 4,816,567; Morrison, *et al.*, *Proc. Natl Acad. Sci. USA*, 81:6851 (1984)), or by covalently joining to the coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such non-immunoglobulin polypeptides can be substituted to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different antigen.

[00251] Chimeric or hybrid antibodies also may be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide-exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate.

Recombinant Production in Prokaryotic Cells

[00252] Polynucleic acid sequences encoding the antibodies of the present disclosure can be obtained using standard recombinant techniques. Desired polynucleic acid sequences may be isolated and sequenced from antibody producing cells such as hybridoma cells. Alternatively, polynucleotides can be synthesized using nucleotide synthesizer or PCR techniques. Once obtained, sequences encoding the polypeptides are inserted into a recombinant vector capable of replicating and expressing heterologous polynucleotides in prokaryotic hosts. Many vectors that are available and known in the art can be used for the purpose of the present disclosure. Selection of an appropriate vector will depend mainly on the size of the nucleic acids to be inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components, depending on its function (amplification or expression of heterologous polynucleotide, or both) and its compatibility with the particular host cell in which it resides. The vector components generally include, but are not limited to, an origin of replication, a selection marker gene, a promoter, a ribosome binding site (RBS), a signal sequence, the heterologous nucleic acid insert and a transcription termination sequence.

[00253] In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, a plasmid derived from an *E. coli* species. Examples of pBR322 derivatives used for expression of particular antibodies are described in detail in Carter *et al.*, U.S. Pat. No. 5,648,237.

[00254] In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, bacteriophage such as GEM™-11 may be utilized in making a recombinant vector which can be used to transform susceptible host cells such as *E. coli* LE392.

[00255] The expression vector of the present application may comprise two or more promoter-cistron pairs, encoding each of the polypeptide components. A promoter is an untranslated regulatory sequence located upstream (5') to a cistron that modulates its expression. Prokaryotic promoters typically fall into two classes, inducible and constitutive. Inducible promoter is a promoter that initiates increased levels of transcription of the cistron under its control in response to changes in the culture condition, *e.g.* the presence or absence of a nutrient or a change in temperature.

[00256] A large number of promoters recognized by a variety of potential host cells are well known. The selected promoter can be operably linked to cistron DNA encoding the present antibody by removing the promoter from the source DNA via restriction enzyme digestion and inserting the isolated promoter sequence into the vector of the present application. Both the

native promoter sequence and many heterologous promoters may be used to direct amplification and/or expression of the target genes. In some embodiments, heterologous promoters are utilized, as they generally permit greater transcription and higher yields of expressed target gene as compared to the native target polypeptide promoter.

[00257] Promoters suitable for use with prokaryotic hosts include the PhoA promoter, the -galactamase and lactose promoter systems, a tryptophan (trp) promoter system and hybrid promoters such as the tac or the trc promoter. However, other promoters that are functional in bacteria (such as other known bacterial or phage promoters) are suitable as well. Their nucleic acid sequences have been published, thereby enabling a skilled worker operably to ligate them to cistrons encoding the target peptide (Siebenlist *et al. Cell* 20: 269 (1980)) using linkers or adaptors to supply any required restriction sites.

[00258] In one aspect, each cistron within the recombinant vector comprises a secretion signal sequence component that directs translocation of the expressed polypeptides across a membrane. In general, the signal sequence may be a component of the vector, or it may be a part of the target polypeptide DNA that is inserted into the vector. The signal sequence selected for the purpose of this invention should be one that is recognized and processed (*i.e.* cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the signal sequences native to the heterologous polypeptides, the signal sequence can be substituted by a prokaryotic signal sequence selected, for example, from the group consisting of the alkaline phosphatase, penicillinase, Ipp, or heat-stable enterotoxin II (STII) leaders, LamB, PhoE, PelB, OmpA and MBP.

[00259] In some embodiments, the production of the antibodies according to the present disclosure can occur in the cytoplasm of the host cell, and therefore does not require the presence of secretion signal sequences within each cistron. Certain host strains (*e.g.*, the *E. coli* trxB⁻ strains) provide cytoplasm conditions that are favorable for disulfide bond formation, thereby permitting proper folding and assembly of expressed protein subunits.

[00260] Prokaryotic host cells suitable for expressing the antibodies of the present disclosure include Archaeobacteria and Eubacteria, such as Gram-negative or Gram-positive organisms. Examples of useful bacteria include *Escherichia* (*e.g.*, *E. coli*), *Bacilli* (*e.g.*, *B. subtilis*), Enterobacteria, *Pseudomonas* species (*e.g.*, *P. aeruginosa*), *Salmonella typhimurium*, *Serratia marcescans*, *Klebsiella*, *Proteus*, *Shigella*, *Rhizobia*, *Vitreoscilla*, or *Paracoccus*. In some embodiments, gram-negative cells are used. In one embodiment, *E. coli* cells are used as hosts. Examples of *E. coli* strains include strain W3110 (Bachmann, *Cellular and Molecular Biology*, vol. 2 (Washington, D.C.: American Society for Microbiology, 1987), pp. 1190-1219; ATCC Deposit No. 27,325) and derivatives thereof, including strain 33D3 having genotype W3110 AfhuA (AtonA) ptr3 lac Iq lacL8 AompT A(nmpc-fepE) degP41 kan^R (U.S. Pat. No. 5,639,635).

Other strains and derivatives thereof, such as *E. coli* 294 (ATCC 31,446), *E. coli* B, *E. coli* 1776 (ATCC 31,537) and *E. coli* RV308 (ATCC 31,608) are also suitable. These examples are illustrative rather than limiting. Methods for constructing derivatives of any of the above-mentioned bacteria having defined genotypes are known in the art and described in, for example, Bass *et al.*, *Proteins*, 8:309-314 (1990). It is generally necessary to select the appropriate bacteria taking into consideration replicability of the replicon in the cells of a bacterium. For example, *E. coli*, *Serratia*, or *Salmonella* species can be suitably used as the host when well known plasmids such as pBR322, pBR325, pACYC177, or pKN410 are used to supply the replicon.

[00261] Typically the host cell should secrete minimal amounts of proteolytic enzymes, and additional protease inhibitors may desirably be incorporated in the cell culture.

[00262] Host cells are transformed with the above-described expression vectors and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. Transformation means introducing DNA into the prokaryotic host so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integrant. Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride is generally used for bacterial cells that contain substantial cell-wall barriers. Another method for transformation employs polyethylene glycol/DMSO. Yet another technique used is electroporation.

[00263] Prokaryotic cells used to produce the antibodies of the present application are grown in media known in the art and suitable for culture of the selected host cells. Examples of suitable media include luria broth (LB) plus necessary nutrient supplements. In some embodiments, the media also contains a selection agent, chosen based on the construction of the expression vector, to selectively permit growth of prokaryotic cells containing the expression vector. For example, ampicillin is added to media for growth of cells expressing ampicillin resistant gene.

[00264] Any necessary supplements besides carbon, nitrogen, and inorganic phosphate sources may also be included at appropriate concentrations introduced alone or as a mixture with another supplement or medium such as a complex nitrogen source. Optionally the culture medium may contain one or more reducing agents selected from the group consisting of glutathione, cysteine, cystamine, thioglycollate, dithioerythritol and dithiothreitol. The prokaryotic host cells are cultured at suitable temperatures and pHs.

[00265] If an inducible promoter is used in the expression vector of the present application, protein expression is induced under conditions suitable for the activation of the promoter. In one aspect of the present application, PhoA promoters are used for controlling transcription of the polypeptides. Accordingly, the transformed host cells are cultured in a phosphate-limiting medium for induction. Preferably, the phosphate-limiting medium is the C.R.A.P medium (see,

e.g., Simmons *et al.*, *J. Immunol. Methods* 263:133-147 (2002)). A variety of other inducers may be used, according to the vector construct employed, as is known in the art.

[00266] The expressed antibodies of the present disclosure are secreted into and recovered from the periplasm of the host cells. Protein recovery typically involves disrupting the microorganism, generally by such means as osmotic shock, sonication or lysis. Once cells are disrupted, cell debris or whole cells may be removed by centrifugation or filtration. The proteins may be further purified, for example, by affinity resin chromatography. Alternatively, proteins can be transported into the culture media and isolated therein. Cells may be removed from the culture and the culture supernatant being filtered and concentrated for further purification of the proteins produced. The expressed polypeptides can be further isolated and identified using commonly known methods such as polyacrylamide gel electrophoresis (PAGE) and Western blot assay.

[00267] Alternatively, protein production is conducted in large quantity by a fermentation process. Various large-scale fed-batch fermentation procedures are available for production of recombinant proteins. To improve the production yield and quality of the antibodies of the present disclosure, various fermentation conditions can be modified. For example, the chaperone proteins have been demonstrated to facilitate the proper folding and solubility of heterologous proteins produced in bacterial host cells. Chen *et al.* *J Bio Chem* 274:19601-19605 (1999); U.S. Pat. No. 6,083,715; U.S. Pat. No. 6,027,888; Bothmann and Pluckthun, *J. Biol. Chem.* 275:17100-17105 (2000); Ramm and Pluckthun, *J. Biol. Chem.* 275:17106-17113 (2000); Arie *et al.*, *Mol. Microbiol.* 39:199-210 (2001).

[00268] To minimize proteolysis of expressed heterologous proteins (especially those that are proteolytically sensitive), certain host strains deficient for proteolytic enzymes can be used for the present invention, as described in, for example, U.S. Pat. No. 5,264,365; U.S. Pat. No. 5,508,192; Hara *et al.*, *Microbial Drug Resistance*, 2:63-72 (1996). *E. coli* strains deficient for proteolytic enzymes and transformed with plasmids overexpressing one or more chaperone proteins may be used as host cells in the expression system encoding the antibodies of the present application.

[00269] The antibodies produced herein can be further purified to obtain preparations that are substantially homogeneous for further assays and uses. Standard protein purification methods known in the art can be employed. The following procedures are exemplary of suitable purification procedures: fractionation on immunoaffinity or ion-exchange columns, ethanol precipitation, reverse phase HPLC, chromatography on silica or on a cation-exchange resin such as DEAE, chromatofocusing, SDS-PAGE, ammonium sulfate precipitation, and gel filtration using, for example, Sephadex G-75. Protein A immobilized on a solid phase for example can be used in some embodiments for immunoaffinity purification of binding molecules of the present disclosure. The solid phase to which Protein A is immobilized is preferably a column comprising

a glass or silica surface, more preferably a controlled pore glass column or a silicic acid column. In some embodiments, the column has been coated with a reagent, such as glycerol, in an attempt to prevent nonspecific adherence of contaminants. The solid phase is then washed to remove contaminants non-specifically bound to the solid phase. Finally the antibodies of interest is recovered from the solid phase by elution.

Recombinant Production in Eukaryotic Cells

[00270] For eukaryotic expression, the vector components generally include, but are not limited to, one or more of the following, a signal sequence, an origin of replication, one or more marker genes, and enhancer element, a promoter, and a transcription termination sequence.

[00271] A vector for use in a eukaryotic host may also an insert that encodes a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. The heterologous signal sequence selected preferably is one that is recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell. In mammalian cell expression, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available. The DNA for such precursor region can be ligated in reading frame to DNA encoding the antibodies of the present application.

[00272] Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

[00273] Expression and cloning vectors may contain a selection gene, also termed a selectable marker. Selection genes may encode proteins that confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline; complement auxotrophic deficiencies; or supply critical nutrients not available from complex media.

[00274] One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin, mycophenolic acid and hygromycin.

[00275] Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up nucleic acid encoding the antibodies of the present application. For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An exemplary appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity. Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with the polypeptide encoding-DNA sequences, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such

as an aminoglycosidic antibiotic.

[00276] Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the nucleic acid encoding the desired polypeptide sequences. Eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of the transcription of many genes may be included. The 3' end of most eukaryotic may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences may be inserted into eukaryotic expression vectors.

[00277] Polypeptide transcription from vectors in mammalian host cells can be controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, *e.g.*, the actin promoter or an immunoglobulin promoter, from heat-shock promoters, provided such promoters are compatible with the host cell systems.

[00278] Transcription of a DNA encoding the antibodies of the present disclosure by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. *See also* Yaniv, *Nature* 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the polypeptide encoding sequence, but is preferably located at a site 5' from the promoter.

[00279] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the polypeptide-encoding mRNA. One useful transcription termination component is the bovine growth hormone polyadenylation region.

[00280] Suitable host cells for cloning or expressing the DNA in the vectors herein include higher eukaryote cells described herein, including vertebrate host cells. Propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10);

Chinese hamster ovary cells/-DHFR (CHO, Urlaub *et al.*, *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TR1 cells (Mather *et al.*, *Annals N.Y. Acad. Sci.* 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

[00281] Host cells can be transformed with the above-described expression or cloning vectors for antibodies production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

[00282] The host cells used to produce the antibodies of the present application may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ((MEM), (Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham *et al.*, *Meth. Enz.* 58:44 (1979), Barnes *et al.*, *Anal. Biochem.* 102:255 (1980), U.S. Pat. No. 4,767,704; 4,657,866; 4,927,762; 4,560,655; or 5,122,469; WO 90/03430; WO 87/00195; or U.S. Pat. Re. 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as GENTAMYCIN™ drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

[00283] When using recombinant techniques, the antibodies can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. If the antibody is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, are removed, for example, by centrifugation or ultrafiltration. Where the antibody is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of

adventitious contaminants.

[00284] The protein composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly (styrene-divinyl) benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™ chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered. Following any preliminary purification step(s), the mixture comprising the antibody of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography.

5.2.7. Binding Molecules Comprising the Single Domain Antibodies

[00285] In another aspect, provided herein is a binding molecule comprising a single domain antibody (*e.g.*, a VHH domain against CD20) provided herein. In addition to chimeric antigen receptors (CARs) provided herein as described in Section 5.3 below, in some embodiments, a single domain antibody against CD20 provided herein is part of other binding molecules. Exemplary binding molecules of the present disclosure are described herein.

Fusion Protein

[00286] In various embodiments, the single domain antibody provided herein can be genetically fused or chemically conjugated to another agent, for example, protein-based entities. The single domain antibody may be chemically-conjugated to the agent, or otherwise non-covalently conjugated to the agent. The agent can be a peptide or antibody (or a fragment thereof).

[00287] Thus, in some embodiments, provided herein are single domain antibodies (*e.g.*, VHH domains) that are recombinantly fused or chemically conjugated (covalent or non-covalent conjugations) to a heterologous protein or polypeptide (or fragment thereof, for example, to a polypeptide of about 10, about 20, about 30, about 40, about 50, about 60, about 70, about 80, about 90, about 100, about 150, about 200, about 250, about 300, about 350, about 400, about 450 or about 500 amino acids, or over 500 amino acids) to generate fusion proteins, as well as uses thereof. In particular, provided herein are fusion proteins comprising an antigen-binding fragment of the single domain antibody provided herein (*e.g.*, CDR1, CDR2, and/or CDR3) and a heterologous protein, polypeptide, or peptide.

[00288] Moreover, antibodies provided herein can be fused to marker or “tag” sequences, such as a peptide, to facilitate purification. In specific embodiments, the marker or tag amino acid

sequence is a hexa-histidine peptide, hemagglutinin (“HA”) tag, and “FLAG” tag.

[00289] Methods for fusing or conjugating moieties (including polypeptides) to antibodies are known (see, *e.g.*, Arnon et al., *Monoclonal Antibodies for Immunotargeting of Drugs in Cancer Therapy*, in *Monoclonal Antibodies and Cancer Therapy* 243-56 (Reisfeld et al. eds., 1985); Hellstrom et al., *Antibodies for Drug Delivery*, in *Controlled Drug Delivery* 623-53 (Robinson et al. eds., 2d ed. 1987); Thorpe, *Antibody Carriers of Cytotoxic Agents in Cancer Therapy: A Review*, in *Monoclonal Antibodies: Biological and Clinical Applications* 475-506 (Pinchera et al. eds., 1985); Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody in Cancer Therapy, in *Monoclonal Antibodies for Cancer Detection and Therapy* 303-16 (Baldwin et al. eds., 1985); Thorpe et al., *Immunol. Rev.* 62:119-58 (1982); U.S. Pat. Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,723,125; 5,783,181; 5,908,626; 5,844,095; and 5,112,946; EP 307,434; EP 367,166; EP 394,827; PCT publications WO 91/06570, WO 96/04388, WO 96/22024, WO 97/34631, and WO 99/04813; Ashkenazi et al., *Proc. Natl. Acad. Sci. USA*, 88: 10535-39 (1991); Traunecker et al., *Nature*, 331:84-86 (1988); Zheng et al., *J. Immunol.* 154:5590-600 (1995); and Vil et al., *Proc. Natl. Acad. Sci. USA* 89:11337-41 (1992)).

[00290] Fusion proteins may be generated, for example, through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as “DNA shuffling”). DNA shuffling may be employed to alter the activities of the single domain antibodies as provided herein, including, for example, antibodies with higher affinities and lower dissociation rates (see, *e.g.*, U.S. Pat. Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458; Patten et al., *Curr. Opin. Biotechnol.* 8:724-33 (1997); Harayama, *Trends Biotechnol.* 16(2):76-82 (1998); Hansson et al., *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo and Blasco, *Biotechniques* 24(2):308-13 (1998)). Antibodies, or the encoded antibodies, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion, or other methods prior to recombination. A polynucleotide encoding an antibody provided herein may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

[00291] In some embodiments, a single domain antibody provided herein (*e.g.*, VHH domain) is conjugated to a second antibody to form an antibody heteroconjugate.

[00292] In various embodiments, the single domain antibody is genetically fused to the agent. Genetic fusion may be accomplished by placing a linker (*e.g.*, a polypeptide) between the single domain antibody and the agent. The linker may be a flexible linker.

[00293] In various embodiments, the single domain antibody is genetically conjugated to a therapeutic molecule, with a hinge region linking the single domain antibody to the therapeutic molecule.

[00294] Also provided herein are methods for making the various fusion proteins provided herein. The various methods described in Section 5.2.6 above may also be utilized to make the fusion proteins provided herein.

[00295] In a specific embodiment, the fusion protein provided herein is recombinantly expressed. Recombinant expression of a fusion protein provided herein may require construction of an expression vector containing a polynucleotide that encodes the protein or a fragment thereof. Once a polynucleotide encoding a protein provided herein or a fragment thereof has been obtained, the vector for the production of the molecule may be produced by recombinant DNA technology using techniques well-known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. Also provided are replicable vectors comprising a nucleotide sequence encoding a fusion protein provided herein, or a fragment thereof, or a CDR, operably linked to a promoter.

[00296] The expression vector can be transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce a fusion protein provided herein. Thus, also provided herein are host cells containing a polynucleotide encoding a fusion protein provided herein or fragments thereof operably linked to a heterologous promoter.

[00297] A variety of host-expression vector systems may be utilized to express the fusion protein provided herein. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express a fusion protein provided herein *in situ*. These include but are not limited to microorganisms such as bacteria (*e.g.*, *E. coli* and *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing coding sequences; yeast (*e.g.*, *Saccharomyces Pichia*) transformed with recombinant yeast expression vectors containing coding sequences; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing coding sequences; plant cell systems infected with recombinant virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV, tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing coding sequences; or mammalian cell systems (*e.g.*, COS, CHO, BHK, 293, NS0, and 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter). Bacterial cells such as *Escherichia*

coli, or, eukaryotic cells, especially for the expression of whole recombinant antibody molecule, can be used for the expression of a recombinant fusion protein. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies or variants thereof. In a specific embodiment, the expression of nucleotide sequences encoding the fusion proteins provided herein is regulated by a constitutive promoter, inducible promoter or tissue specific promoter.

[00298] In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the fusion protein being expressed. For example, when a large quantity of such a fusion protein is to be produced, for the generation of pharmaceutical compositions of a fusion protein, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited to, the *E. coli* expression vector pUR278 (Ruther *et al.*, EMBO 12:1791 (1983)), in which the coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, Nucleic Acids Res. 13:3101-3109 (1985); Van Heeke & Schuster, J. Biol. Chem. 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

[00299] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region EI or E3) will result in a recombinant virus that is viable and capable of expressing the fusion protein in infected hosts (*e.g.*, see Logan & Shenk, Proc. Natl. Acad. Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for efficient translation of inserted coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, *etc.* (see, *e.g.*, Bittner *et al.*, Methods

in Enzymol. 153:51-544 (1987)).

[00300] In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (*e.g.*, glycosylation) and processing (*e.g.*, cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, Hela, COS, MDCK, 293, 3T3, W138, BT483, Hs578T, HTB2, BT2O and T47D, NS0 (a murine myeloma cell line that does not endogenously produce any immunoglobulin chains), CRL7O3O and HsS78Bst cells.

[00301] For long-term, high-yield production of recombinant proteins, stable expression can be utilized. For example, cell lines which stably express the fusion proteins may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, *etc.*), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the fusion protein. Such engineered cell lines may be particularly useful in screening and evaluation of compositions that interact directly or indirectly with the binding molecule.

[00302] A number of selection systems may be used, including but not limited to, the herpes simplex virus thymidine kinase (Wigler *et al.*, Cell 11:223 (1977)), hypoxanthineguanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy *et al.*, Cell 22:8-17 (1980)) genes can be employed in tk-, hgp^rt- or ap^rt-cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: *dhfr*, which confers resistance to methotrexate (Wigler *et al.*, Natl. Acad. Sci. USA 77:357 (1980); O'Hare *et al.*, Proc. Natl. Acad. Sci. USA 78:1527 (1981)); *gpt*, which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 (Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993);

Mulligan, Science 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, TIB TECH 11(5):155-2 15 (1993)); and *hygro*, which confers resistance to hygromycin (Santerre *et al.*, Gene 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel *et al.* (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli *et al.* (eds.), Current Protocols in Human Genetics, John Wiley & Sons, NY (1994); Colberre-Garapin *et al.*, J. Mol. Biol. 150:1 (1981), which are incorporated by reference herein in their entireties.

[00303] The expression level of a fusion protein can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol. 3 (Academic Press, New York, 1987)). When a marker in the vector system expressing a fusion protein is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the fusion protein gene, production of the fusion protein will also increase (Crouse *et al.*, Mol. Cell. Biol. 3:257 (1983)).

[00304] The host cell may be co-transfected with multiple expression vectors provided herein. The vectors may contain identical selectable markers which enable equal expression of respective encoding polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing multiple polypeptides. The coding sequences may comprise cDNA or genomic DNA.

[00305] Once a fusion protein provided herein has been produced by recombinant expression, it may be purified by any method known in the art for purification of a polypeptide (*e.g.*, an immunoglobulin molecule), for example, by chromatography (*e.g.*, ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, sizing column chromatography, and Kappa select affinity chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. Further, the fusion protein molecules provided herein can be fused to heterologous polypeptide sequences described herein or otherwise known in the art to facilitate purification.

Immunoconjugates

[00306] In some embodiments, the present disclosure also provides immunoconjugates comprising any of the antibodies (such as anti-CD20 single domain antibodies) described herein conjugated to one or more cytotoxic agents, such as chemotherapeutic agents or drugs, growth inhibitory agents, toxins (*e.g.*, protein toxins, enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof), or radioactive isotopes.

[00307] In some embodiments, an immunoconjugate is an antibody-drug conjugate (ADC) in

which an antibody is conjugated to one or more drugs, including but not limited to a maytansinoid (see U.S. Patent Nos. 5,208,020, 5,416,064 and European Patent EP 0 425 235 B1); an auristatin such as monomethylauristatin drug moieties DE and DF (MMAE and MMAF) (see U.S. Patent Nos. 5,635,483 and 5,780,588, and 7,498,298); a dolastatin; a calicheamicin or derivative thereof (see U.S. Patent Nos. 5,712,374, 5,714,586, 5,739,116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, and 5,877,296; Hinman et al., *Cancer Res.* 53:3336-3342 (1993); and Lode et al., *Cancer Res.* 58:2925-2928 (1998)); an anthracycline such as daunomycin or doxorubicin (see Kratz et al., *Current Med. Chem.* 13:477-523 (2006); Jeffrey et al., *Bioorganic & Med. Chem. Letters* 16:358-362 (2006); Torgov et al., *Bioconj. Chem.* 16:717-721 (2005); Nagy et al., *Proc. Natl. Acad. Sci. USA* 97:829-834 (2000); Dubowchik et al., *Bioorg. & Med. Chem. Letters* 12:1529-1532 (2002); King et al., *J. Med. Chem.* 45:4336-4343 (2002); and U.S. Patent No. 6,630,579); methotrexate; vindesine; a taxane such as docetaxel, paclitaxel, larotaxel, tesetaxel, and ortataxel; a trichothecene; and CC1065.

[00308] In some embodiments, an immunoconjugate comprises an antibody as described herein conjugated to an enzymatically active toxin or fragment thereof, including but not limited to diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes.

[00309] In some embodiments, an immunoconjugate comprises an antibody as described herein conjugated to a radioactive atom to form a radioconjugate. A variety of radioactive isotopes are available for the production of radioconjugates. Examples include At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³², Pb²¹² and radioactive isotopes of Lu. When the radioconjugate is used for detection, it may comprise a radioactive atom for scintigraphic studies, for example tc99m or I123, or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, mri), such as iodine-123 again, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

[00310] Conjugates of an antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a

ricin immunotoxin can be prepared as described in Vitetta et al., *Science* 238:1098 (1987).

Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

[00311] The linker may be a “cleavable linker” facilitating release of the conjugated agent in the cell, but non-cleavable linkers are also contemplated herein. Linkers for use in the conjugates of the present disclosure include, without limitation, acid labile linkers (*e.g.*, hydrazone linkers), disulfide-containing linkers, peptidase-sensitive linkers (*e.g.*, peptide linkers comprising amino acids, for example, valine and/or citrulline such as citrulline-valine or phenylalanine-lysine), photolabile linkers, dimethyl linkers, thioether linkers, or hydrophilic linkers designed to evade multidrug transporter-mediated resistance.

[00312] The immunoconjugates or ADCs herein contemplate, but are not limited to such conjugates prepared with cross-linker reagents including, but not limited to, BMPS, EMCS, GMBS, HBVS, LC-SMCC, MBS, MPBH, SBAP, SIA, SIAB, SMCC, SMPB, SMPH, sulfo-EMCS, sulfo-GMBS, sulfo-KMUS, sulfo-MBS, sulfo-SIAB, sulfo-SMCC, and sulfo-SMPB, and SVSB (succinimidyl-(4-vinylsulfone)benzoate) which are commercially available (*e.g.*, from Pierce Biotechnology, Inc., Rockford, IL., U.S.A).

[00313] In other embodiments, antibodies provided herein are conjugated or recombinantly fused, *e.g.*, to a diagnostic molecule. Such diagnosis and detection can be accomplished, for example, by coupling the antibody to detectable substances including, but not limited to, various enzymes, such as, but not limited to, horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic groups, such as, but not limited to, streptavidin/biotin or avidin/biotin; fluorescent materials, such as, but not limited to, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride, or phycoerythrin; luminescent materials, such as, but not limited to, luminol; bioluminescent materials, such as, but not limited to, luciferase, luciferin, or aequorin; chemiluminescent material, such as, 225Ac γ -emitting, Auger-emitting, β -emitting, an alpha-emitting or positron-emitting radioactive isotope.

5.3. Chimeric antigen receptors

[00314] In another aspect, provided herein is a chimeric antigen receptor (CAR) comprising an extracellular antigen binding domain comprising at least one single domain antibody (*e.g.*, VHH) provided herein that binds to CD20. Exemplary CARs comprising the present VHH domains (*i.e.*, VHH-based CARs) are illustrated and compared with conventional CARs comprising scFvs (*i.e.*, scFv-based CARs) as described in Section 6 below.

[00315] In some embodiments, the chimeric antigen receptor (CAR) provided herein comprises a polypeptide comprising: (a) an extracellular antigen binding domain comprising at least one

single domain antibody (sdAb) specifically binding to CD20 as provided herein, and optionally one or more additional binding domain(s); (b) a transmembrane domain; and (c) an intracellular signaling domain. Each components and additional regions are described in more detail below.

5.3.1. Extracellular antigen binding domain

[00316] The extracellular antigen binding domain of the CARs described herein comprises one or more (such as any one of 1, 2, 3, 4, 5, 6 or more) single domain antibodies. The single domain antibodies can be fused to each other directly via peptide bonds, or via peptide linkers.

Single domain antibodies

[00317] The CARs of the present disclosure comprise an extracellular antigen binding domain comprising one or more single domain antibodies. The sdAbs may be of the same or different origins, and of the same or different sizes. Exemplary sdAbs include, but are not limited to, heavy chain variable domains from heavy-chain only antibodies (*e.g.*, VHH or V_{NAR}), binding molecules naturally devoid of light chains, single domains (such as V_H or V_L) derived from conventional 4-chain antibodies, humanized heavy-chain only antibodies, human single domain antibodies produced by transgenic mice or rats expressing human heavy chain segments, and engineered domains and single domain scaffolds other than those derived from antibodies. Any sdAbs known in the art or developed by the present disclosure, including the single domain antibodies described above in the present disclosure, may be used to construct the CARs described herein. The sdAbs may be derived from any species including, but not limited to mouse, rat, human, camel, llama, lamprey, fish, shark, goat, rabbit, and bovine. Single domain antibodies contemplated herein also include naturally occurring single domain antibody molecules from species other than Camelidae and sharks.

[00318] In some embodiments, the sdAb is derived from a naturally occurring single domain antigen binding molecule known as heavy chain antibody devoid of light chains (also referred herein as “heavy chain only antibodies”). Such single domain molecules are disclosed in WO 94/04678 and Hamers-Casterman, C. *et al.*, *Nature* 363:446-448 (1993), for example. For clarity reasons, the variable domain derived from a heavy chain molecule naturally devoid of light chain is known herein as a VHH to distinguish it from the conventional V_H of four chain immunoglobulins. Such a VHH molecule can be derived from antibodies raised in *Camelidae* species, for example, camel, llama, vicuna, dromedary, alpaca and guanaco. Other species besides Camelidae may produce heavy chain molecules naturally devoid of light chain, and such VHHs are within the scope of the present disclosure. In addition, humanized versions of VHHs as well as other modifications and variants are also contemplated and within the scope of the present disclosure.

[00319] VHH molecules from Camelids are about 10 times smaller than IgG molecules. They are single polypeptides and can be very stable, resisting extreme pH and temperature conditions.

Moreover, they can be resistant to the action of proteases which is not the case for conventional 4-chain antibodies. Furthermore, *in vitro* expression of VHHs produces high yield, properly folded functional VHHs. In addition, antibodies generated in Camelids can recognize epitopes other than those recognized by antibodies generated *in vitro* through the use of antibody libraries or via immunization of mammals other than Camelids (see, for example, WO9749805). As such, multispecific or multivalent CARs comprising one or more VHH domains may interact more efficiently with targets than multispecific or multivalent CARs comprising antigen binding fragments derived from conventional 4-chain antibodies. Since VHHs are known to bind into “unusual” epitopes such as cavities or grooves, the affinity of CARs comprising such VHHs may be more suitable for therapeutic treatment than conventional multispecific polypeptides.

[00320] In some embodiments, the sdAb is derived from a variable region of the immunoglobulin found in cartilaginous fish. For example, the sdAb can be derived from the immunoglobulin isotype known as Novel Antigen Receptor (NAR) found in the serum of shark. Methods of producing single domain molecules derived from a variable region of NAR ("IgNARs") are described in WO 03/014161 and Streltsov, *Protein Sci.* 14:2901-2909 (2005).

[00321] In some embodiments, the sdAb is recombinant, CDR-grafted, humanized, camelized, de-immunized and/or *in vitro* generated (*e.g.*, selected by phage display). In some embodiments, the amino acid sequence of the framework regions may be altered by “camelization” of specific amino acid residues in the framework regions. Camelization refers to the replacing or substitution of one or more amino acid residues in the amino acid sequence of a (naturally occurring) V_H domain from a conventional 4-chain antibody by one or more of the amino acid residues that occur at the corresponding position(s) in a VHH domain of a heavy chain antibody. This can be performed in a manner known in the field, which will be clear to the skilled person. Such “camelizing” substitutions are preferably inserted at amino acid positions that form and/or are present at the V_H-V_L interface, and/or at the so-called Camelidae hallmark residues, as defined herein (see for example WO 94/04678, Davies and Riechmann FEBS Letters 339: 285-290 (1994); Davies and Riechmann, *Protein Engineering* 9 (6): 531-537 (1996); Riechmann, *J. Mol. Biol.* 259: 957-969 (1996); and Riechmann and Muyldermans, *J. Immunol. Meth.* 231: 25-38 (1999)).

[00322] In some embodiments, the sdAb is a human single domain antibody produced by transgenic mice or rats expressing human heavy chain segments. *See, e.g.*, US20090307787, U.S. Pat. No. 8,754,287, US20150289489, US20100122358, and WO2004049794. In some embodiments, the sdAb is affinity matured.

[00323] In some embodiments, naturally occurring VHH domains against a particular antigen or target, can be obtained from (naïve or immune) libraries of Camelid VHH sequences. Such methods may or may not involve screening such a library using said antigen or target, or at least

one part, fragment, antigenic determinant or epitope thereof using one or more screening techniques known in the field. Such libraries and techniques are for example described in WO 99/37681, WO 01/90190, WO 03/025020 and WO 03/035694. Alternatively, improved synthetic or semi-synthetic libraries derived from (naïve or immune) VHH libraries may be used, such as VHH libraries obtained from (naïve or immune) VHH libraries by techniques such as random mutagenesis and/or CDR shuffling, as for example described in WO 00/43507.

[00324] In some embodiments, the single domain antibodies are generated from conventional four-chain antibodies. See, for example, EP 0 368 684; Ward et al., *Nature*, 341 (6242): 544-6 (1989); Holt et al., *Trends Biotechnol.*, 21(11):484-490 (2003); WO 06/030220; and WO 06/003388.

[00325] In some embodiments, the extracellular antigen binding domain provided herein comprises at least one binding domain, and the at least one binding domain comprises a single domain antibody that binds to CD20 as provided herein, *e.g.*, the anti-CD20 single domain antibodies described in Section 5.2 above.

[00326] In some embodiments, provided herein is a CAR comprising a polypeptide comprising: (a) an extracellular antigen binding domain comprising an anti-CD20 sdAb; (b) a transmembrane domain; and (c) an intracellular signaling domain, wherein the anti-CD20 sdAb is an anti-CD20 sdAb as described in Section 5.2 above, including, *e.g.*, the VHH domains in Table 2 and those having one, two or all three CDRs in any of those VHH domains in Table 2. In some embodiments, the anti-CD20 sdAb is camelid, chimeric, human, or humanized.

[00327] In some embodiments, provided herein is a CAR comprising a polypeptide comprising: (a) an extracellular antigen binding domain comprising an anti-CD20 sdAb; (b) a transmembrane domain; and (c) an intracellular signaling domain, wherein the anti-CD20 sdAb comprises the amino acid sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184, or SEQ ID NO: 185. In other embodiments, provided herein is a CAR comprising a polypeptide comprising: (a) an extracellular antigen binding domain comprising an anti-CD20 sdAb; (b) a transmembrane domain; and (c) an intracellular signaling domain, wherein the anti-CD20 sdAb comprises an amino acid sequence having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184, or SEQ ID NO: 185.

NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184, or SEQ ID NO: 185.

[00328] In other embodiments, the extracellular antigen binding domain comprises two or more antigen binding domains. Among these two or more antigen binding domains, at least one is a VHH that binds to CD20 as provided herein. In some embodiments, the one or more additional binding domain(s) is/are also VHH(s) that bind(s) to CD20. In other embodiments, the one or more additional binding domain(s) bind(s) to one or more additional different antigen(s), *e.g.*, 1, 2, 3, 4 or more additional single domain antibody binding regions (sdAbs) targeting one or more additional different antigen(s).

[00329] In some embodiments, provided herein is a multivalent (such as bivalent and trivalent) CAR comprising a polypeptide comprising: (a) an extracellular antigen binding domain comprising two or more single domain antibodies (sdAbs) specifically binding to CD20; (b) a transmembrane domain; and (c) an intracellular signaling domain. In some embodiments, the extracellular antigen binding domain comprises two single domain antibodies (sdAbs) specifically binding to CD20 provided herein. In other embodiments, the extracellular antigen binding domain comprises three single domain antibodies (sdAbs) specifically binding to CD20 provided herein. In some embodiments, the two or more anti-CD20 sdAbs are selected from those anti-CD20 sdAbs described in Section 5.2 above, including, *e.g.*, the VHH domains in Table 2 and those having one, two or all three CDRs in any of those VHH domains in Table 2. In some embodiments, the anti-CD20 sdAb is camelid, chimeric, human, or humanized. In some embodiments, the two or more anti-CD20 sdAbs are each independently selected from anti-CD20 sdAbs comprising an amino acid sequence having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184, and SEQ ID NO: 185.

[00330] In other embodiments, provided herein is a multispecific (such as bispecific and trispecific) CAR comprising a polypeptide comprising: (a) an extracellular antigen binding domain comprising a first single domain antibody (sdAb) specifically binding to CD20; (b) a transmembrane domain; and (c) an intracellular signaling domain. In some embodiments, the CAR further comprises a second single domain antibody (sdAb) specifically binding to a second antigen (such as a second tumor antigen). In some embodiments, the CAR further comprises a second single domain antibody (sdAb) specifically binding to a second antigen (such as a second

tumor antigen); and a third single domain antibody (sdAb) specifically binding to a third antigen (such as a third tumor antigen).

[00331] In some embodiments, the additional antigen(s) targeted by the CARs of the present disclosure are cell surface molecules. The single domain antibodies may be chosen to recognize an antigen that acts as a cell surface marker on target cells associated with a special disease state. In some embodiments, the antigen is a tumor antigen. In some embodiments, the tumor antigen is associated with a B cell malignancy. Tumors express a number of proteins that can serve as a target antigen for an immune response, particularly T cell mediated immune responses. The antigens targeted by the CAR may be antigens on a single diseased cell or antigens that are expressed on different cells that each contribute to the disease. The antigens targeted by the CAR may be directly or indirectly involved in the diseases.

[00332] Tumor antigens are proteins that are produced by tumor cells that can elicit an immune response, particularly T-cell mediated immune responses. The selection of the additional targeted antigen of the present disclosure will depend on the particular type of cancer to be treated. Exemplary tumor antigens include, but not limited to, a glioma-associated antigen, carcinoembryonic antigen (CEA), β -human chorionic gonadotropin, alphafetoprotein (AFP), lectin-reactive AFP, thyroglobulin, RAGE-1, MN-CAIX, human telomerase reverse transcriptase, RU1, RU2 (AS), intestinal carboxyl esterase, mut hsp70-2, M-CSF, prostase, prostate-specific antigen (PSA), PAP, NY-ESO-1, LAGE-1a, p53, prostein, PSMA, HER2/neu, survivin and telomerase, prostate-carcinoma tumor antigen-1 (PCTA-1), MAGE, ELF2M, neutrophil elastase, ephrinB2, CD19, insulin growth factor (IGF)-I, IGF-II, IGF-I receptor and mesothelin.

[00333] In some embodiments, the tumor antigen comprises one or more antigenic cancer epitopes associated with a malignant tumor. Malignant tumors express a number of proteins that can serve as target antigens for an immune attack. These molecules include, but are not limited to, tissue-specific antigens such as MART-1, tyrosinase and gp100 in melanoma and prostatic acid phosphatase (PAP) and prostate-specific antigen (PSA) in prostate cancer. Other target molecules belong to the group of transformation-related molecules such as the oncogene HER2/Neu/ErbB-2. Yet another group of target antigens are onco-fetal antigens such as carcinoembryonic antigen (CEA). In B-cell lymphoma the tumor-specific idiotype immunoglobulin constitutes a truly tumor-specific immunoglobulin antigen that is unique to the individual tumor. In addition to CD20, B-cell differentiation antigens such as CD22 and CD37 are other candidates for target antigens in B-cell lymphoma.

[00334] In some embodiments, the tumor antigen is a tumor-specific antigen (TSA) or a tumor-associated antigen (TAA). A TSA is unique to tumor cells and does not occur on other cells in the body. A TAA associated antigen is not unique to a tumor cell, and instead is also expressed

on a normal cell under conditions that fail to induce a state of immunologic tolerance to the antigen. The expression of the antigen on the tumor may occur under conditions that enable the immune system to respond to the antigen. TAAs may be antigens that are expressed on normal cells during fetal development, when the immune system is immature, and unable to respond or they may be antigens that are normally present at extremely low levels on normal cells, but which are expressed at much higher levels on tumor cells.

[00335] Non-limiting examples of TSA or TAA antigens include: differentiation antigens such as MART-1/MelanA (MART-I), gp 100 (Pmel 17), tyrosinase, TRP-1, TRP-2 and tumor-specific multilineage antigens such as MAGE-1, MAGE-3, BAGE, GAGE-1, GAGE-2, p15; overexpressed embryonic antigens such as CEA; overexpressed oncogenes and mutated tumor-suppressor genes such as p53, Ras, HER2/neu; unique tumor antigens resulting from chromosomal translocations; such as BCR-ABL, E2A-PRL, H4-RET, IGH-IGK, MYL-RAR; and viral antigens, such as the Epstein Barr virus antigens EBVA and the human papillomavirus (HPV) antigens E6 and E7.

[00336] Other large, protein-based antigens include TSP-180, MAGE-4, MAGE-5, MAGE-6, RAGE, NY-ESO, p185erbB2, p180erbB-3, c-met, nm-23HI, PSA, TAG-72, CA 19-9, CA 72-4, CAM 17.1, NuMa, K-ras, beta-Catenin, CDK4, Mum-1, p 15, p 16, 43-9F, 5T4, 791Tgp72, alpha-fetoprotein, beta-HCG, BCA225, BTAA, CA 125, CA 15-3\CA 27.29\BCAA, CA 195, CA 242, CA-50, CAM43, CD68\P1, CO-029, FGF-5, G250, Ga733\EpCAM, HTgp-175, M344, MA-50, MG7-Ag, MOV18, NB/70K, NY-CO- 1, RCAS 1, SDCCAG16, TA-90\Mac-2 binding protein\cyclophilin C-associated protein, TAAL6, TAG72, TLP, and TPS.

[00337] In some more specific embodiments, the one or more additional antigen(s) is selected from a group consisting of CD20, CD19, CD33, CD38, BCMA, CS1, ROR1, GPC3, CD123, IL-13R, CD138, c-Met, EGFRvIII, GD-2, NY-ESO-1, MAGE A3, and glycolipid F77.

[00338] In a specific embodiment, the CAR provided herein comprises a VHH that binds to CD20 and a VHH that binds to CD19. In another specific embodiment, the CAR provided herein comprises a VHH that binds to CD20 and a VHH that binds to CD22. In some embodiments, the sdAb provided herein is camelid, chimeric, human, or humanized.

[00339] In addition to the one or more antigen binding domain(s) provided herein, the CAR provided herein may further comprise one or more of the following: a linker (*e.g.*, a peptide linker), a transmembrane domain, a hinge region, a signal peptide, an intracellular signaling domain, a co-stimulatory signaling domain, each of which is described in more detail below.

[00340] For example, in some embodiments, the intracellular signaling domain comprises a primary intracellular signaling domain of an immune effector cell (such as T cell). In some embodiments, the primary intracellular signaling domain is derived from CD3ζ. In some embodiments, the intracellular signaling domain comprises a co-stimulatory signaling domain. In

some embodiments, the co-stimulatory signaling domain is derived from a co-stimulatory molecule selected from the group consisting of CD27, CD28, CD137, OX40, CD30, CD40, CD3, LFA-1, CD2, CD7, LIGHT, NKG2C, B7-H3, ligands of CD83 and combinations thereof. In some embodiments, the co-stimulatory signaling domain is derived from CD137. In some embodiments, the CD20 CAR further comprises a hinge domain (such as a CD8 α hinge domain) located between the C-terminus of the extracellular antigen binding domain and the N-terminus of the transmembrane domain. In some embodiments, the CD20 CAR further comprises a signal peptide (such as a CD8 α signal peptide) located at the N-terminus of the polypeptide. In some embodiments, the polypeptide comprises from the N-terminus to the C-terminus: a CD8 α signal peptide, the extracellular antigen-binding domain, a CD8 α hinge domain, a CD8 α transmembrane domain, a co-stimulatory signaling domain derived from CD137, and a primary intracellular signaling domain derived from CD3 ζ . In some embodiments, the CD20 CAR is monospecific. In some embodiments, the CD20 CAR is monovalent. In other embodiments, the CD20 CAR is multivalent.

Peptide linkers

[00341] The various single domain antibodies in the multispecific or multivalent CARs described herein may be fused to each other via peptide linkers. In some embodiments, the single domain antibodies are directly fused to each other without any peptide linkers. The peptide linkers connecting different single domain antibodies (*e.g.*, VHH) may be the same or different. Different domains of the CARs may also be fused to each other via peptide linkers.

[00342] Each peptide linker in a CAR may have the same or different length and/or sequence depending on the structural and/or functional features of the single domain antibodies and/or the various domains. Each peptide linker may be selected and optimized independently. The length, the degree of flexibility and/or other properties of the peptide linker(s) used in the CARs may have some influence on properties, including but not limited to the affinity, specificity or avidity for one or more particular antigens or epitopes. For example, longer peptide linkers may be selected to ensure that two adjacent domains do not sterically interfere with one another. In some embodiments, a short peptide linker may be disposed between the transmembrane domain and the intracellular signaling domain of a CAR. In some embodiment, a peptide linker comprises flexible residues (such as glycine and serine) so that the adjacent domains are free to move relative to each other. For example, a glycine-serine doublet can be a suitable peptide linker.

[00343] The peptide linker can be of any suitable length. In some embodiments, the peptide linker is at least about any of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 50, 75, 100 or more amino acids long. In some embodiments, the peptide linker is no more than about any of 100, 75, 50, 40, 35, 30, 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5 or fewer amino acids long. In some embodiments, the length of the peptide linker is any of

about 1 amino acid to about 10 amino acids, about 1 amino acids to about 20 amino acids, about 1 amino acid to about 30 amino acids, about 5 amino acids to about 15 amino acids, about 10 amino acids to about 25 amino acids, about 5 amino acids to about 30 amino acids, about 10 amino acids to about 30 amino acids long, about 30 amino acids to about 50 amino acids, about 50 amino acids to about 100 amino acids, or about 1 amino acid to about 100 amino acids.

[00344] The peptide linker may have a naturally occurring sequence, or a non-naturally occurring sequence. For example, a sequence derived from the hinge region of heavy chain only antibodies may be used as the linker. *See*, for example, WO1996/34103. In some embodiments, the peptide linker is a flexible linker. Exemplary flexible linkers include but not limited to glycine polymers (G)_n, glycine-serine polymers (including, for example, (GS)_n, (GSGGS)_n, (GGGS)_n, and (GGGGS)_n, where n is an integer of at least one), glycine-alanine polymers, alanine-serine polymers, and other flexible linkers known in the art. Exemplary peptide linkers are listed in the table below. In a specific embodiment, the peptide linker that connects two or more anti-CD20 VHH domains provided herein is (GGGGS)_n (SEQ ID NO: 138), wherein n is optionally 1, 2, 3, 4, or 5.

Table 4. Exemplary Peptide Linkers

Sequences	SEQ ID NO
(GS) _n , n is an integer including, <i>e.g.</i> , 1, 2, 3, 4, 5, and 6.	SEQ ID NO: 134
(GSGGS) _n , n is an integer including, <i>e.g.</i> , 1, 2, 3, 4, 5, and 6.	SEQ ID NO: 135
(GGGS) _n , n is an integer including, <i>e.g.</i> , 1, 2, 3, 4, 5, and 6.	SEQ ID NO: 136
GGGSGGGSGGGGGSGSGGGSGGGSGGGGS	SEQ ID NO: 137
(GGGGS) _n , n is an integer including, <i>e.g.</i> , 1, 2, 3, 4, 5, and 6.	SEQ ID NO: 138
DGGGS	SEQ ID NO: 139
TGEKP	SEQ ID NO: 140
GGRR	SEQ ID NO: 141
GGGSGGGSGGGGGSGSGGGGS	SEQ ID NO: 142
EGKSSGSGSESKVD	SEQ ID NO: 143
KESGSVSSEQLAQFRS	SEQ ID NO: 144
GGRRGGGS	SEQ ID NO: 145
LRQRDGERP	SEQ ID NO: 146
LRQKDGGGSERP	SEQ ID NO: 147
LRQKDGGSGGGSERP	SEQ ID NO: 148
GSTSGSGKPGSGEGST	SEQ ID NO: 149
GSTSGSGKSSEKKG	SEQ ID NO: 150
KESGSVSSEQLAQFRSLD	SEQ ID NO: 151

[00345] Other linkers known in the art, for example, as described in WO2016014789, WO2015158671, WO2016102965, US20150299317, WO2018067992, US7741465, Colcher *et*

al., *J. Nat. Cancer Inst.* 82:1191-1197 (1990), and Bird *et al.*, *Science* 242:423-426 (1988) may also be included in the CARs provided herein, the disclosure of each of which is incorporated herein by reference.

5.3.2. Transmembrane domain

[00346] The CARs of the present disclosure comprise a transmembrane domain that can be directly or indirectly fused to the extracellular antigen binding domain. The transmembrane domain may be derived either from a natural or from a synthetic source. As used herein, a “transmembrane domain” refers to any protein structure that is thermodynamically stable in a cell membrane, preferably an eukaryotic cell membrane. Transmembrane domains compatible for use in the CARs described herein may be obtained from a naturally occurring protein. Alternatively, it can be a synthetic, non-naturally occurring protein segment, *e.g.*, a hydrophobic protein segment that is thermodynamically stable in a cell membrane.

[00347] Transmembrane domains are classified based on the three dimensional structure of the transmembrane domain. For example, transmembrane domains may form an alpha helix, a complex of more than one alpha helix, a beta-barrel, or any other stable structure capable of spanning the phospholipid bilayer of a cell. Furthermore, transmembrane domains may also or alternatively be classified based on the transmembrane domain topology, including the number of passes that the transmembrane domain makes across the membrane and the orientation of the protein. For example, single-pass membrane proteins cross the cell membrane once, and multi-pass membrane proteins cross the cell membrane at least twice (*e.g.*, 2, 3, 4, 5, 6, 7 or more times). Membrane proteins may be defined as Type I, Type II or Type III depending upon the topology of their termini and membrane-passing segment(s) relative to the inside and outside of the cell. Type I membrane proteins have a single membrane-spanning region and are oriented such that the N-terminus of the protein is present on the extracellular side of the lipid bilayer of the cell and the C-terminus of the protein is present on the cytoplasmic side. Type II membrane proteins also have a single membrane-spanning region but are oriented such that the C-terminus of the protein is present on the extracellular side of the lipid bilayer of the cell and the N-terminus of the protein is present on the cytoplasmic side. Type III membrane proteins have multiple membrane- spanning segments and may be further sub-classified based on the number of transmembrane segments and the location of N- and C-termini.

[00348] In some embodiments, the transmembrane domain of the CAR described herein is derived from a Type I single-pass membrane protein. In some embodiments, transmembrane domains from multi-pass membrane proteins may also be compatible for use in the CARs described herein. Multi-pass membrane proteins may comprise a complex (at least 2, 3, 4, 5, 6, 7 or more) alpha helices or a beta sheet structure. In some embodiments, the N-terminus and the C-terminus of a multi-pass membrane protein are present on opposing sides of the lipid bilayer,

e.g., the N-terminus of the protein is present on the cytoplasmic side of the lipid bilayer and the C-terminus of the protein is present on the extracellular side.

[00349] In some embodiments, the transmembrane domain of the CAR comprises a transmembrane domain chosen from the transmembrane domain of an alpha, beta or zeta chain of a T-cell receptor, CD28, CD3 epsilon, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, CD154, KIRDS2, OX40, CD2, CD27, LFA-1 (CD11a, CD18), ICOS (CD278), 4-1BB (CD137), GITR, CD40, BAFFR, HVEM (LIGHTR), SLAMF7, NKp80 (KLRFl), CD160, CD19, IL-2R beta, IL-2R gamma, IL-7R a, ITGA1, VLA1, CD49a, ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD11d, ITGAE, CD103, ITGAL, CD11a, LFA-1, ITGAM, CD11b, ITGAX, CD11c, ITGB1, CD29, ITGB2, CD18, LFA-1, ITGB7, TNFR2, DNAM1 (CD226), SLAMF4 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRT AM, Ly9 (CD229), CD160 (BY55), PSGL1, CD100 (SEMA4D), SLAMF6 (NTB-A, Ly108), SLAM (SLAMF1, CD150, IPO-3), BLAME (SLAMF8), SELPLG (CD162), LTBR, PAG/Cbp, NKp44, NKp30, NKp46, NKG2D, and/or NKG2C. In some embodiments, the transmembrane domain is derived from a molecule selected from the group consisting of CD8 α , CD4, CD28, CD137, CD80, CD86, CD152 and PD1.

[00350] In some specific embodiments, the transmembrane domain is derived from CD8 α . In some embodiments, the transmembrane domain is a transmembrane domain of CD8 α comprising the amino acid sequence of SEQ ID NO: 154.

[00351] Transmembrane domains for use in the CARs described herein can also comprise at least a portion of a synthetic, non-naturally occurring protein segment. In some embodiments, the transmembrane domain is a synthetic, non-naturally occurring alpha helix or beta sheet. In some embodiments, the protein segment is at least approximately 20 amino acids, *e.g.*, at least 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, or more amino acids. Examples of synthetic transmembrane domains are known in the art, for example in U.S. Patent No. 7,052,906 and PCT Publication No. WO 2000/032776, the relevant disclosures of which are incorporated by reference herein.

[00352] The transmembrane domain provided herein may comprise a transmembrane region and a cytoplasmic region located at the C-terminal side of the transmembrane domain. The cytoplasmic region of the transmembrane domain may comprise three or more amino acids and, in some embodiments, helps to orient the transmembrane domain in the lipid bilayer. In some embodiments, one or more cysteine residues are present in the transmembrane region of the transmembrane domain. In some embodiments, one or more cysteine residues are present in the cytoplasmic region of the transmembrane domain. In some embodiments, the cytoplasmic region of the transmembrane domain comprises positively charged amino acids. In some embodiments, the cytoplasmic region of the transmembrane domain comprises the amino acids arginine, serine,

and lysine.

[00353] In some embodiments, the transmembrane region of the transmembrane domain comprises hydrophobic amino acid residues. In some embodiments, the transmembrane domain of the CAR provided herein comprises an artificial hydrophobic sequence. For example, a triplet of phenylalanine, tryptophan and valine may be present at the C terminus of the transmembrane domain. In some embodiments, the transmembrane region comprises mostly hydrophobic amino acid residues, such as alanine, leucine, isoleucine, methionine, phenylalanine, tryptophan, or valine. In some embodiments, the transmembrane region is hydrophobic. In some embodiments, the transmembrane region comprises a poly-leucine-alanine sequence. The hydrophathy, or hydrophobic or hydrophilic characteristics of a protein or protein segment, can be assessed by any method known in the art, for example the Kyte and Doolittle hydrophathy analysis.

5.3.3. Intracellular signaling domain

[00354] The CARs of the present disclosure comprise an intracellular signaling domain. The intracellular signaling domain is responsible for activation of at least one of the normal effector functions of the immune effector cell expressing the CARs. The term “effector function” refers to a specialized function of a cell. Effector function of a T cell, for example, may be cytolytic activity or helper activity including the secretion of cytokines. Thus the term “cytoplasmic signaling domain” refers to the portion of a protein which transduces the effector function signal and directs the cell to perform a specialized function. While usually the entire cytoplasmic signaling domain can be employed, in many cases it is not necessary to use the entire chain. To the extent that a truncated portion of the cytoplasmic signaling domain is used, such truncated portion may be used in place of the intact chain as long as it transduces the effector function signal. The term cytoplasmic signaling domain is thus meant to include any truncated portion of the cytoplasmic signaling domain sufficient to transduce the effector function signal.

[00355] In some embodiments, the intracellular signaling domain comprises a primary intracellular signaling domain of an immune effector cell. In some embodiments, the CAR comprises an intracellular signaling domain consisting essentially of a primary intracellular signaling domain of an immune effector cell. “Primary intracellular signaling domain” refers to cytoplasmic signaling sequence that acts in a stimulatory manner to induce immune effector functions. In some embodiments, the primary intracellular signaling domain contains a signaling motif known as immunoreceptor tyrosine-based activation motif, or ITAM. An “ITAM,” as used herein, is a conserved protein motif that is generally present in the tail portion of signaling molecules expressed in many immune cells. The motif may comprises two repeats of the amino acid sequence YxxL/I separated by 6-8 amino acids, wherein each x is independently any amino acid, producing the conserved motif YxxL/Ix(6-8)YxxL/I. ITAMs within signaling molecules are important for signal transduction within the cell, which is mediated at least in part by

phosphorylation of tyrosine residues in the ITAM following activation of the signaling molecule. ITAMs may also function as docking sites for other proteins involved in signaling pathways. Exemplary ITAM-containing primary cytoplasmic signaling sequences include those derived from CD3 ζ , FcR gamma (FCER1G), FcR beta (Fc Epsilon Rib), CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b, and CD66d.

[00356] In some embodiments, the primary intracellular signaling domain is derived from CD3 ζ . In some embodiments, the intracellular signaling domain consists of the cytoplasmic signaling domain of CD3 ζ . In some embodiments, the primary intracellular signaling domain is a cytoplasmic signaling domain of wild-type CD3 ζ . In some embodiments, the primary intracellular signaling domain of CD3 ζ comprises the amino acid sequence of SEQ ID NO: 156. In some embodiments, the primary intracellular signaling domain of wild-type CD3 ζ . In some embodiments, the primary intracellular signaling domain is a functional mutant of the cytoplasmic signaling domain of CD3 ζ containing one or more mutations, such as Q65K.

5.3.4. Co-stimulatory signaling domain

[00357] Many immune effector cells require co-stimulation, in addition to stimulation of an antigen-specific signal, to promote cell proliferation, differentiation and survival, as well as to activate effector functions of the cell. In some embodiments, the CAR comprises at least one co-stimulatory signaling domain. The term "co-stimulatory signaling domain," as used herein, refers to at least a portion of a protein that mediates signal transduction within a cell to induce an immune response such as an effector function. The co-stimulatory signaling domain of the chimeric receptor described herein can be a cytoplasmic signaling domain from a co-stimulatory protein, which transduces a signal and modulates responses mediated by immune cells, such as T cells, NK cells, macrophages, neutrophils, or eosinophils. "Co-stimulatory signaling domain" can be the cytoplasmic portion of a co-stimulatory molecule. The term "co-stimulatory molecule" refers to a cognate binding partner on an immune cell (such as T cell) that specifically binds with a co-stimulatory ligand, thereby mediating a co-stimulatory response by the immune cell, such as, but not limited to, proliferation and survival.

[00358] In some embodiments, the intracellular signaling domain comprises a single co-stimulatory signaling domain. In some embodiments, the intracellular signaling domain comprises two or more (such as about any of 2, 3, 4, or more) co-stimulatory signaling domains. In some embodiments, the intracellular signaling domain comprises two or more of the same co-stimulatory signaling domains. In some embodiments, the intracellular signaling domain comprises two or more co-stimulatory signaling domains from different co-stimulatory proteins, such as any two or more co-stimulatory proteins described herein. In some embodiments, the intracellular signaling domain comprises a primary intracellular signaling domain (such as

cytoplasmic signaling domain of CD3 ζ) and one or more co-stimulatory signaling domains. In some embodiments, the one or more co-stimulatory signaling domains and the primary intracellular signaling domain (such as cytoplasmic signaling domain of CD3 ζ) are fused to each other via optional peptide linkers. The primary intracellular signaling domain, and the one or more co-stimulatory signaling domains may be arranged in any suitable order. In some embodiments, the one or more co-stimulatory signaling domains are located between the transmembrane domain and the primary intracellular signaling domain (such as cytoplasmic signaling domain of CD3 ζ). Multiple co-stimulatory signaling domains may provide additive or synergistic stimulatory effects.

[00359] Activation of a co-stimulatory signaling domain in a host cell (*e.g.*, an immune cell) may induce the cell to increase or decrease the production and secretion of cytokines, phagocytic properties, proliferation, differentiation, survival, and/or cytotoxicity. The co-stimulatory signaling domain of any co-stimulatory molecule may be compatible for use in the CARs described herein. The type(s) of co-stimulatory signaling domain is selected based on factors such as the type of the immune effector cells in which the effector molecules would be expressed (*e.g.*, T cells, NK cells, macrophages, neutrophils, or eosinophils) and the desired immune effector function (*e.g.*, ADCC effect). Examples of co-stimulatory signaling domains for use in the CARs can be the cytoplasmic signaling domain of co-stimulatory proteins, including, without limitation, members of the B7/CD28 family (*e.g.*, B7-1/CD80, B7-2/CD86, B7-H1/PD-L1, B7-H2, B7-H3, B7-H4, B7-H6, B7-H7, BTLA/CD272, CD28, CTLA-4, Gi24/VISTA/B7-H5, ICOS/CD278, PD-1, PD-L2/B7-DC, and PDCD6); members of the TNF superfamily (*e.g.*, 4-1BB/TNFSF9/CD137, 4-1BB Ligand/TNFSF9, BAFF/BLyS/TNFSF13B, BAFF R/TNFRSF13C, CD27/TNFRSF7, CD27 Ligand/TNFSF7, CD30/TNFRSF8, CD30 Ligand/TNFSF8, CD40/TNFRSF5, CD40 Ligand/TNFSF5, DR3/TNFRSF25, GITR/TNFRSF18, GITR Ligand/TNFSF18, HVEM/TNFRSF14, LIGHT/TNFSF14, Lymphotoxin-alpha/TNF-beta, OX40/TNFRSF4, OX40 Ligand/TNFSF4, RELT/TNFRSF19L, TACI/TNFRSF13B, TL1A/TNFSF15, TNF-alpha, and TNF RII/TNFRSF1B); members of the SLAM family (*e.g.*, 2B4/CD244/SLAMF4, BLAME/SLAMF8, CD2, CD2F-10/SLAMF9, CD48/SLAMF2, CD58/LFA-3, CD84/SLAMF5, CD229/SLAMF3, CRACC/SLAMF7, NTB-A/SLAMF6, and SLAM/CD150); and any other co-stimulatory molecules, such as CD2, CD7, CD53, CD82/Kai-1, CD90/Thy1, CD96, CD160, CD200, CD300a/LMIR1, HLA Class I, HLA-DR, Ikaros, Integrin alpha 4/CD49d, Integrin alpha 4 beta 1, Integrin alpha 4 beta 7/LPAM-1, LAG-3, TCL1A, TCL1B, CRTAM, DAP12, Dectin-1/CLEC7A, DPPIV/CD26, EphB6, TIM-1/KIM-1/HAVCR, TIM-4, TSLP, TSLP R, lymphocyte function associated antigen-1 (LFA-1), and NKG2C.

[00360] In some embodiments, the one or more co-stimulatory signaling domains are selected

from the group consisting of CD27, CD28, CD137, OX40, CD30, CD40, CD3, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, B7-H3 and ligands that specially bind to CD83.

[00361] In some embodiments, the intracellular signaling domain in the CAR of the present disclosure comprises a co-stimulatory signaling domain derived from CD137 (*i.e.*, 4-1BB). In some embodiments, the intracellular signaling domain comprises a cytoplasmic signaling domain of CD3 ζ and a co-stimulatory signaling domain of CD137. In some embodiments, the intracellular signaling domain comprises a co-stimulatory signaling domain of CD137 comprising the amino acid sequence of SEQ ID NO: 155.

[00362] Also within the scope of the present disclosure are variants of any of the co-stimulatory signaling domains described herein, such that the co-stimulatory signaling domain is capable of modulating the immune response of the immune cell. In some embodiments, the co-stimulatory signaling domains comprises up to 10 amino acid residue variations (*e.g.*, 1, 2, 3, 4, 5, or 8) as compared to a wild-type counterpart. Such co-stimulatory signaling domains comprising one or more amino acid variations may be referred to as variants. Mutation of amino acid residues of the co-stimulatory signaling domain may result in an increase in signaling transduction and enhanced stimulation of immune responses relative to co-stimulatory signaling domains that do not comprise the mutation. Mutation of amino acid residues of the co-stimulatory signaling domain may result in a decrease in signaling transduction and reduced stimulation of immune responses relative to co-stimulatory signaling domains that do not comprise the mutation.

5.3.5. Hinge region

[00363] The CARs of the present disclosure may comprise a hinge domain that is located between the extracellular antigen binding domain and the transmembrane domain. A hinge domain is an amino acid segment that is generally found between two domains of a protein and may allow for flexibility of the protein and movement of one or both of the domains relative to one another. Any amino acid sequence that provides such flexibility and movement of the extracellular antigen binding domain relative to the transmembrane domain of the effector molecule can be used.

[00364] The hinge domain may contain about 10-100 amino acids, *e.g.*, about any one of 15-75 amino acids, 20-50 amino acids, or 30-60 amino acids. In some embodiments, the hinge domain may be at least about any one of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, or 75 amino acids in length.

[00365] In some embodiments, the hinge domain is a hinge domain of a naturally occurring protein. Hinge domains of any protein known in the art to comprise a hinge domain are compatible for use in the chimeric receptors described herein. In some embodiments, the hinge domain is at least a portion of a hinge domain of a naturally occurring protein and confers

flexibility to the chimeric receptor. In some embodiments, the hinge domain is derived from CD8 α . In some embodiments, the hinge domain is a portion of the hinge domain of CD8 α , *e.g.*, a fragment containing at least 15 (*e.g.*, 20, 25, 30, 35, or 40) consecutive amino acids of the hinge domain of CD8 α . In some embodiments, the hinge domain of CD8 α comprises the amino acid sequence of SEQ ID NO: 153.

[00366] Hinge domains of antibodies, such as an IgG, IgA, IgM, IgE, or IgD antibodies, are also compatible for use in the pH-dependent chimeric receptor systems described herein. In some embodiments, the hinge domain is the hinge domain that joins the constant domains CH1 and CH2 of an antibody. In some embodiments, the hinge domain is of an antibody and comprises the hinge domain of the antibody and one or more constant regions of the antibody. In some embodiments, the hinge domain comprises the hinge domain of an antibody and the CH3 constant region of the antibody. In some embodiments, the hinge domain comprises the hinge domain of an antibody and the CH2 and CH3 constant regions of the antibody. In some embodiments, the antibody is an IgG, IgA, IgM, IgE, or IgD antibody. In some embodiments, the antibody is an IgG antibody. In some embodiments, the antibody is an IgG1, IgG2, IgG3, or IgG4 antibody. In some embodiments, the hinge region comprises the hinge region and the CH2 and CH3 constant regions of an IgG1 antibody. In some embodiments, the hinge region comprises the hinge region and the CH3 constant region of an IgG1 antibody.

[00367] Non-naturally occurring peptides may also be used as hinge domains for the chimeric receptors described herein. In some embodiments, the hinge domain between the C-terminus of the extracellular ligand-binding domain of an Fc receptor and the N-terminus of the transmembrane domain is a peptide linker, such as a (GxS) $_n$ linker, wherein x and n, independently can be an integer between 3 and 12, including 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or more.

5.3.6. Signal peptide

[00368] The CARs of the present disclosure may comprise a signal peptide (also known as a signal sequence) at the N-terminus of the polypeptide. In general, signal peptides are peptide sequences that target a polypeptide to the desired site in a cell. In some embodiments, the signal peptide targets the effector molecule to the secretory pathway of the cell and will allow for integration and anchoring of the effector molecule into the lipid bilayer. Signal peptides including signal sequences of naturally occurring proteins or synthetic, non-naturally occurring signal sequences, which are compatible for use in the CARs described herein will be evident to one of skill in the art. In some embodiments, the signal peptide is derived from a molecule selected from the group consisting of CD8 α , GM-CSF receptor α , and IgG1 heavy chain. In some embodiments, the signal peptide is derived from CD8 α . In some embodiments, the signal peptide of CD8 α comprises the amino acid sequence of SEQ ID NO: 152.

5.3.7. Exemplary CARs

[00369] Exemplary monovalent CARs are generated as shown in Section 6 below, such as VHH-273 CAR, VHH-283 CAR, VHH-313 CAR, VHH-440 CAR, VHH-466 CAR, VHH-496 CAR, VHH-653 CAR, huVHH-253 CAR, huVHH-256 CAR, huVHH-260 CAR, huVHH-746 CAR, huVHH-750 CAR, huVHH-753 CAR, huVHH-836 CAR, huVHH-840 CAR, huVHH-843 CAR, huVHH-846 CAR, and VHH-623 CAR, VHH-640 CAR, and/or VHH-657 CAR.

[00370] In addition, exemplary multivalent CARs, such as bivalent and trivalent CARs, are generated as shown in Section 6, including Bi-VHH1 CAR, Bi-VHH2 CAR, Bi-VHH3 CAR, Bi-VHH4 (G4S)₁ CAR, Bi-VHH4 (G4S)₂ CAR, Bi-VHH4 (G4S)₃ CAR, Bi-VHH4 (G4S)₄ CAR, Bi-VHH4 (G4S)₅ CAR, and Tri-VHH CAR.

[00371] In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 81. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 82. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 83. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 84. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 85. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 86. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 87. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 88. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 89. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 90. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 91. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 92. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 93. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 94. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 95. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 96. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 97. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 98. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 99. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 100. In some

embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 101. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 102. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 103. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 104. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 105. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 106. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 189. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 190. In some embodiments, provided herein is a CAR comprising or consisting of the amino acid sequence of SEQ ID NO: 191.

[00372] In certain embodiments, the CAR provided herein comprises amino acid sequences with certain percent identity relative to any one of the CARs exemplified in the Section 6 below.

[00373] In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 81. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 82. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 83. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 84. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 85. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 86. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 87. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%,

97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 88. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 89. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 90. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 91. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 92. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 93. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 94. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 95. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 96. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 97. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 98. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 99. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 100. In some embodiments, provided herein is a CD20 CAR

comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 101. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 102. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 103. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 104. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 105. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 106. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 189. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 190. In some embodiments, provided herein is a CD20 CAR comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 191.

[00374] In some embodiments, provided herein is an isolated nucleic acid encoding any of the CD20 CARs provided herein. More detailed description regarding nucleic acid sequences and vectors are provided below.

5.4. Engineered immune effector cells

[00375] In yet another aspect, provided herein are host cells (such as immune effector cells) comprising any one of the CARs described herein.

[00376] Thus, in some embodiments, provided herein is an engineered immune effector cell (such as T cell) comprising a CAR which comprises a polypeptide comprising: (a) an extracellular antigen binding domain comprising one or more anti-CD20 sdAb(s); (b) a

transmembrane domain; and (c) an intracellular signaling domain, wherein the anti-CD20 sdAb is an anti-CD20 sdAb as described in Section 5.2 above, including, *e.g.*, the VHH domains in Table 2 and those having one, two or all three CDRs in any of those VHH domains in Table 2. In some embodiments, the anti-CD20 sdAb is camelid, chimeric, human, or humanized. In some embodiments, the transmembrane domain is selected from the group consisting of CD8 α , CD4, CD28, CD137, CD80, CD86, CD152 and PD1. In some embodiments, the intracellular signaling domain comprises a primary intracellular signaling domain of an immune effector cell (such as T cell). In some embodiments, the primary intracellular signaling domain is derived from CD3 ζ . In some embodiments, the intracellular signaling domain comprises a co-stimulatory signaling domain. In some embodiments, the co-stimulatory signaling domain is derived from a co-stimulatory molecule selected from the group consisting of CD27, CD28, CD137, OX40, CD30, CD40, CD3, LFA-1, CD2, CD7, LIGHT, NKG2C, B7-H3, ligands of CD83 and combinations thereof. In some embodiments, the CAR further comprises a hinge domain (such as a CD8 α hinge domain) located between the C-terminus of the extracellular antigen binding domain and the N-terminus of the transmembrane domain. In some embodiments, the CAR further comprises a signal peptide (such as a CD8 α signal peptide) located at the N-terminus of the polypeptide. In some embodiments, the polypeptide comprises from the N-terminus to the C-terminus: a CD8 α signal peptide, the extracellular antigen binding domain, a CD8 α hinge domain, a CD8 α transmembrane domain, a co-stimulatory signaling domain derived from CD137, and a primary intracellular signaling domain derived from CD3 ζ .

[00377] In some embodiments, provided herein is an engineered immune effector cell (such as T cell) comprising a CAR which comprises a polypeptide comprising: (a) an extracellular antigen binding domain comprising one or more anti-CD20 sdAb(s); (b) a transmembrane domain; and (c) an intracellular signaling domain, wherein the anti-CD20 sdAb comprises the amino acid sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184, or SEQ ID NO: 185. In some embodiments, provided herein is an engineered immune effector cell (such as T cell) comprising a CAR which comprises a polypeptide comprising: (a) an extracellular antigen binding domain comprising one or more anti-CD20 sdAb(s); (b) a transmembrane domain; and (c) an intracellular signaling domain, wherein the anti-CD20 sdAb comprises an amino acid sequence having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 41, SEQ

ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184, or SEQ ID NO: 185. In some embodiments, the transmembrane domain is selected from the group consisting of CD8 α , CD4, CD28, CD137, CD80, CD86, CD152 and PD1. In some embodiments, the intracellular signaling domain comprises a primary intracellular signaling domain of an immune effector cell (such as T cell). In some embodiments, the primary intracellular signaling domain is derived from CD3 ζ . In some embodiments, the intracellular signaling domain comprises a co-stimulatory signaling domain. In some embodiments, the co-stimulatory signaling domain is derived from a co-stimulatory molecule selected from the group consisting of CD27, CD28, CD137, OX40, CD30, CD40, CD3, LFA-1, CD2, CD7, LIGHT, NKG2C, B7-H3, ligands of CD83 and combinations thereof. In some embodiments, the CAR further comprises a hinge domain (such as a CD8 α hinge domain) located between the C-terminus of the extracellular antigen binding domain and the N-terminus of the transmembrane domain. In some embodiments, the CAR further comprises a signal peptide (such as a CD8 α signal peptide) located at the N-terminus of the polypeptide. In some embodiments, the polypeptide comprises from the N-terminus to the C-terminus: a CD8 α signal peptide, the extracellular antigen binding domain, a CD8 α hinge domain, a CD8 α transmembrane domain, a co-stimulatory signaling domain derived from CD137, and a primary intracellular signaling domain derived from CD3 ζ .

[00378] In some embodiments, provided herein is an engineered immune effector cell (such as T cell) comprising a CAR which comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, SEQ ID NO: 95, SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 99, SEQ ID NO: 100, SEQ ID NO: 101, SEQ ID NO: 102, SEQ ID NO: 103, SEQ ID NO: 104, SEQ ID NO: 105, SEQ ID NO: 106, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191. In some embodiments, provided herein is an engineered immune effector cell (such as T cell) comprising a CAR which comprises a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, SEQ ID NO: 95, SEQ

ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 99, SEQ ID NO: 100, SEQ ID NO: 101, SEQ ID NO: 102, SEQ ID NO: 103, SEQ ID NO: 104, SEQ ID NO: 105, SEQ ID NO: 106, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191.

[00379] In other embodiments, there is provided an engineered immune effector cell (such as T cell) comprising a multispecific (such as bispecific or trispecific) chimeric antigen receptor (CAR) comprising a polypeptide comprising: (a) an extracellular antigen binding domain comprising a first single domain antibody (sdAb) specifically binding to CD20 and one or more additional antigen binding domain(s); (b) a transmembrane domain; and (c) an intracellular signaling domain. In some embodiments, the additional antigen binding domain binds to an antigen selected from the group consisting of CD22, CD19, CD33, CD38, BCMA, CS1, ROR1, GPC3, CD123, IL-13R, CD138, c-Met, EGFRvIII, GD-2, NY-ESO-1, MAGE A3, and glycolipid F77. In some embodiments, the first sdAb and/or the additional sdAb is camelid, chimeric, human, or humanized. In some embodiments, the first single domain antibody and the additional single domain antibody are fused to each other via a peptide bond or a peptide linker. In some embodiments, the peptide linker is no more than about 50 (such as no more than about any one of 35, 25, 20, 15, 10, or 5) amino acids long. In some embodiments, the transmembrane domain is selected from the group consisting of CD8 α , CD4, CD28, CD137, CD80, CD86, CD152 and PD1. In some embodiments, the intracellular signaling domain comprises a primary intracellular signaling domain of an immune effector cell (such as T cell). In some embodiments, the primary intracellular signaling domain is derived from CD3 ζ . In some embodiments, the intracellular signaling domain comprises a co-stimulatory signaling domain. In some embodiments, the co-stimulatory signaling domain is derived from a co-stimulatory molecule selected from the group consisting of CD27, CD28, CD137, OX40, CD30, CD40, CD3, LFA-1, CD2, CD7, LIGHT, NKG2C, B7-H3, ligands of CD83 and combinations thereof. In some embodiments, the multispecific CAR further comprises a hinge domain (such as a CD8 α hinge domain) located between the C-terminus of the extracellular antigen binding domain and the N-terminus of the transmembrane domain. In some embodiments, the multispecific CAR further comprises a signal peptide (such as a CD8 α signal peptide) located at the N-terminus of the polypeptide. In some embodiments, the polypeptide comprises from the N-terminus to the C-terminus: a CD8 α signal peptide, the extracellular antigen binding domain, a CD8 α hinge domain, a CD8 α transmembrane domain, a co-stimulatory signaling domain derived from CD137, and a primary intracellular signaling domain derived from CD3 ζ .

[00380] In some embodiments, the engineered immune effector cell is a T cell, an NK cell, a peripheral blood mononuclear cell (PBMC), a hematopoietic stem cell, a pluripotent stem cell, or an embryonic stem cell. In some embodiments, the engineered immune effector cell is

autologous. In some embodiments, the engineered immune effector cell is allogenic.

[00381] Also provided are engineered immune effector cells comprising (or expressing) two or more different CARs. Any two or more of the CARs described herein may be expressed in combination. The CARs may target different antigens, thereby providing synergistic or additive effects. The two or more CARs may be encoded on the same vector or different vectors.

[00382] The engineered immune effector cell may further express one or more therapeutic proteins and/or immunomodulators, such as immune checkpoint inhibitors. *See, e.g.*, International Patent Application NOs. PCT/CN2016/073489 and PCT/CN2016/087855, which are incorporated herein by reference in their entirety.

5.4.1. Vectors

[00383] The present disclosure provides vectors for cloning and expressing any one of the CARs described herein. In some embodiments, the vector is suitable for replication and integration in eukaryotic cells, such as mammalian cells. In some embodiments, the vector is a viral vector. Examples of viral vectors include, but are not limited to, adenoviral vectors, adeno-associated virus vectors, lentiviral vector, retroviral vectors, vaccinia vector, herpes simplex viral vector, and derivatives thereof. Viral vector technology is well known in the art and is described, for example, in Sambrook *et al.* (2001, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York), and in other virology and molecular biology manuals.

[00384] A number of viral based systems have been developed for gene transfer into mammalian cells. For example, retroviruses provide a convenient platform for gene delivery systems. The heterologous nucleic acid can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to the engineered mammalian cell *in vitro* or *ex vivo*. A number of retroviral systems are known in the art. In some embodiments, adenovirus vectors are used. A number of adenovirus vectors are known in the art. In some embodiments, lentivirus vectors are used. In some embodiments, self-inactivating lentiviral vectors are used. For example, self-inactivating lentiviral vectors carrying the immunomodulator (such as immune checkpoint inhibitor) coding sequence and/or self-inactivating lentiviral vectors carrying chimeric antigen receptors can be packaged with protocols known in the art. The resulting lentiviral vectors can be used to transduce a mammalian cell (such as primary human T cells) using methods known in the art. Vectors derived from retroviruses such as lentivirus are suitable tools to achieve long-term gene transfer, because they allow long-term, stable integration of a transgene and its propagation in progeny cells. Lentiviral vectors also have low immunogenicity, and can transduce non-proliferating cells.

[00385] In some embodiments, the vector comprises any one of the nucleic acids encoding a CAR described herein. The nucleic acid can be cloned into the vector using any known molecular

cloning methods in the art, including, for example, using restriction endonuclease sites and one or more selectable markers. In some embodiments, the nucleic acid is operably linked to a promoter. Varieties of promoters have been explored for gene expression in mammalian cells, and any of the promoters known in the art may be used in the present disclosure. Promoters may be roughly categorized as constitutive promoters or regulated promoters, such as inducible promoters.

[00386] In some embodiments, the nucleic acid encoding the CAR is operably linked to a constitutive promoter. Constitutive promoters allow heterologous genes (also referred to as transgenes) to be expressed constitutively in the host cells. Exemplary constitutive promoters contemplated herein include, but are not limited to, Cytomegalovirus (CMV) promoters, human elongation factors-1 alpha (hEF1 α), ubiquitin C promoter (UbiC), phosphoglycerokinase promoter (PGK), simian virus 40 early promoter (SV40), and chicken β -Actin promoter coupled with CMV early enhancer (CAGG). The efficiencies of such constitutive promoters on driving transgene expression have been widely compared in a huge number of studies. For example, Michael C. Milone *et al* compared the efficiencies of CMV, hEF1 α , UbiC and PGK to drive chimeric antigen receptor expression in primary human T cells, and concluded that hEF1 α promoter not only induced the highest level of transgene expression, but was also optimally maintained in the CD4 and CD8 human T cells (Molecular Therapy, 17(8): 1453-1464 (2009)). In some embodiments, the nucleic acid encoding the CAR is operably linked to a hEF1 α promoter.

[00387] In some embodiments, the nucleic acid encoding the CAR is operably linked to an inducible promoter. Inducible promoters belong to the category of regulated promoters. The inducible promoter can be induced by one or more conditions, such as a physical condition, microenvironment of the engineered immune effector cell, or the physiological state of the engineered immune effector cell, an inducer (*i.e.*, an inducing agent), or a combination thereof.

[00388] In some embodiments, the inducing condition does not induce the expression of endogenous genes in the engineered mammalian cell, and/or in the subject that receives the pharmaceutical composition. In some embodiments, the inducing condition is selected from the group consisting of: inducer, irradiation (such as ionizing radiation, light), temperature (such as heat), redox state, tumor environment, and the activation state of the engineered mammalian cell.

[00389] In some embodiments, the vector also contains a selectable marker gene or a reporter gene to select cells expressing the CAR from the population of host cells transfected through lentiviral vectors. Both selectable markers and reporter genes may be flanked by appropriate regulatory sequences to enable expression in the host cells. For example, the vector may contain transcription and translation terminators, initiation sequences, and promoters useful for regulation of the expression of the nucleic acid sequences.

[00390] In some embodiments, the vector comprises more than one nucleic acid encoding CARs. In some embodiments, the vector comprises a nucleic acid comprising a first nucleic acid sequence encoding a first CAR and a second nucleic acid sequence encoding a second CAR, wherein the first nucleic acid is operably linked to the second nucleic acid via a third nucleic acid sequence encoding a self-cleaving peptide. In some embodiments, the self-cleaving peptide is selected from the group consisting of T2A, P2A and F2A.

5.4.2. Immune effector cells

[00391] “Immune effector cells” are immune cells that can perform immune effector functions. In some embodiments, the immune effector cells express at least FcγRIII and perform ADCC effector function. Examples of immune effector cells which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells, neutrophils, and eosinophils.

[00392] In some embodiments, the immune effector cells are T cells. In some embodiments, the T cells are CD4⁺/CD8⁻, CD4⁻/CD8⁺, CD4⁺/CD8⁺, CD4⁻/CD8⁻, or combinations thereof. In some embodiments, the T cells produce IL-2, TFN, and/or TNF upon expressing the CAR and binding to the target cells, such as CD20⁺ tumor cells. In some embodiments, the CD8⁺ T cells lyse antigen-specific target cells upon expressing the CAR and binding to the target cells.

[00393] In some embodiments, the immune effector cells are NK cells. In other embodiments, the immune effector cells can be established cell lines, for example, NK-92 cells.

[00394] In some embodiments, the immune effector cells are differentiated from a stem cell, such as a hematopoietic stem cell, a pluripotent stem cell, an iPS, or an embryonic stem cell.

[00395] The engineered immune effector cells are prepared by introducing the CARs into the immune effector cells, such as T cells. In some embodiments, the CAR is introduced to the immune effector cells by transfecting any one of the isolated nucleic acids or any one of the vectors described above. In some embodiments, the CAR is introduced to the immune effector cells by inserting proteins into the cell membrane while passing cells through a microfluidic system, such as CELL SQUEEZE[®] (*see, e.g.*, U.S. Patent Application Publication No. 20140287509).

[00396] Methods of introducing vectors or isolated nucleic acids into a mammalian cell are known in the art. The vectors described can be transferred into an immune effector cell by physical, chemical, or biological methods.

[00397] Physical methods for introducing the vector into an immune effector cell include calcium phosphate precipitation, lipofection, particle bombardment, microinjection, electroporation, and the like. Methods for producing cells comprising vectors and/or exogenous nucleic acids are well-known in the art. *See, e.g.*, Sambrook *et al.* (2001) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York. In some embodiments, the

vector is introduced into the cell by electroporation.

[00398] Biological methods for introducing the vector into an immune effector cell include the use of DNA and RNA vectors. Viral vectors have become the most widely used method for inserting genes into mammalian, *e.g.*, human cells.

[00399] Chemical means for introducing the vector into an immune effector cell include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. An exemplary colloidal system for use as a delivery vehicle *in vitro* is a liposome (*e.g.*, an artificial membrane vesicle).

[00400] In some embodiments, RNA molecules encoding any of the CARs described herein may be prepared by a conventional method (*e.g.*, *in vitro* transcription) and then introduced into the immune effector cells via known methods such as mRNA electroporation. *See, e.g.*, Rabinovich *et al.*, Human Gene Therapy 17:1027-1035 (2006).

[00401] In some embodiments, the transduced or transfected immune effector cell is propagated *ex vivo* after introduction of the vector or isolated nucleic acid. In some embodiments, the transduced or transfected immune effector cell is cultured to propagate for at least about any of 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, or 14 days. In some embodiments, the transduced or transfected immune effector cell is further evaluated or screened to select the engineered mammalian cell.

[00402] Reporter genes may be used for identifying potentially transfected cells and for evaluating the functionality of regulatory sequences. In general, a reporter gene is a gene that is not present in or expressed by the recipient organism or tissue and that encodes a polypeptide whose expression is manifested by some easily detectable property, *e.g.*, enzymatic activity. Expression of the reporter gene is assayed at a suitable time after the DNA has been introduced into the recipient cells. Suitable reporter genes may include genes encoding luciferase, beta-galactosidase, chloramphenicol acetyl transferase, secreted alkaline phosphatase, or the green fluorescent protein gene (*e.g.*, Ui-Tei *et al.* FEBS Letters 479: 79-82 (2000)). Suitable expression systems are well known and may be prepared using known techniques or obtained commercially. Other methods to confirm the presence of the nucleic acid encoding the CARs in the engineered immune effector cell, include, for example, molecular biological assays well known to those of skill in the art, such as Southern and Northern blotting, RT-PCR and PCR; biochemical assays, such as detecting the presence or absence of a particular peptide, *e.g.*, by immunological methods (such as ELISAs and Western blots).

5.4.3. Sources of T Cells

[00403] In some embodiments, prior to expansion and genetic modification of the T cells, a source of T cells is obtained from a subject. T cells can be obtained from a number of sources,

including peripheral blood mononuclear cells, bone marrow, lymph node tissue, cord blood, thymus tissue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. In some embodiments, any number of T cell lines available in the art, may be used. In some embodiments, T cells can be obtained from a unit of blood collected from a subject using any number of techniques known to the skilled artisan, such as Ficoll™ separation. In some embodiments, cells from the circulating blood of an individual are obtained by apheresis. The apheresis product typically contains lymphocytes, including T cells, monocytes, granulocytes, B cells, other nucleated white blood cells, red blood cells, and platelets. In some embodiments, the cells collected by apheresis may be washed to remove the plasma fraction and to place the cells in an appropriate buffer or media for subsequent processing steps. In some embodiments, the cells are washed with phosphate buffered saline (PBS). In some embodiments, the wash solution lacks calcium and may lack magnesium or may lack many if not all divalent cations. Initial activation steps in the absence of calcium may lead to magnified activation. As those of ordinary skill in the art would readily appreciate a washing step may be accomplished by methods known to those in the art, such as by using a semi-automated “flow-through” centrifuge (for example, the Cobe 2991 cell processor, the Baxter CytoMate, or the Haemonetics Cell Saver 5) according to the manufacturer's instructions. After washing, the cells may be resuspended in a variety of biocompatible buffers, such as, for example, Ca^{2+} -free, Mg^{2+} -free PBS, PlasmaLyte A, or other saline solution with or without buffer. Alternatively, the undesirable components of the apheresis sample may be removed and the cells directly resuspended in culture media.

[00404] In some embodiments, T cells are isolated from peripheral blood lymphocytes by lysing the red blood cells and depleting the monocytes, for example, by centrifugation through a PERCOLL™ gradient or by counterflow centrifugal elutriation. A specific subpopulation of T cells, such as CD3^+ , CD28^+ , CD4^+ , CD8^+ , CD45RA^+ , and CD45RO^+ T cells, can be further isolated by positive or negative selection techniques. For example, in some embodiments, T cells are isolated by incubation with anti-CD3/anti-CD28 (*i.e.*, 3×28)-conjugated beads, such as DYNABEADS® M-450 CD3/CD28 T, for a time period sufficient for positive selection of the desired T cells. In some embodiments, the time period is about 30 minutes. In a further embodiment, the time period ranges from 30 minutes to 36 hours or longer and all integer values there between. In a further embodiment, the time period is at least 1, 2, 3, 4, 5, or 6 hours. In some embodiments, the time period is 10 to 24 hours. In some embodiments, the incubation time period is 24 hours. For isolation of T cells from patients with leukemia, use of longer incubation times, such as 24 hours, can increase cell yield. Longer incubation times may be used to isolate T cells in any situation where there are few T cells as compared to other cell types, such in isolating tumor infiltrating lymphocytes (TIL) from tumor tissue or from immune-compromised individuals. Further, use of longer incubation times can increase the efficiency of capture of

CD8+ T cells. Thus, in some embodiments, by simply shortening or lengthening the time T cells are allowed to bind to the CD3/CD28 beads and/or by increasing or decreasing the ratio of beads to T cells, subpopulations of T cells can be preferentially selected for or against at culture initiation or at other time points during the process. Additionally, by increasing or decreasing the ratio of anti-CD3 and/or anti-CD28 antibodies on the beads or other surface, subpopulations of T cells can be preferentially selected for or against at culture initiation or at other desired time points. The skilled artisan would recognize that multiple rounds of selection can also be used. In some embodiments, it may be desirable to perform the selection procedure and use the “unselected” cells in the activation and expansion process. “Unselected” cells can also be subjected to further rounds of selection.

[00405] Enrichment of a T cell population by negative selection can be accomplished with a combination of antibodies directed to surface markers unique to the negatively selected cells. One method is cell sorting and/or selection via negative magnetic immunoadherence or flow cytometry that uses a cocktail of monoclonal antibodies directed to cell surface markers present on the cells negatively selected. For example, to enrich for CD4+ cells by negative selection, a monoclonal antibody cocktail typically includes antibodies to CD14, CD20, CD11b, CD16, HLA-DR, and CD8. In certain embodiments, it may be desirable to enrich for or positively select for regulatory T cells which typically express CD4+, CD25+, CD62Lhi, GITR+, and FoxP3+. Alternatively, in certain embodiments, T regulatory cells are depleted by anti-CD25 conjugated beads or other similar method of selection.

[00406] For isolation of a desired population of cells by positive or negative selection, the concentration of cells and surface (*e.g.*, particles such as beads) can be varied. In certain embodiments, it may be desirable to significantly decrease the volume in which beads and cells are mixed together (*i.e.*, increase the concentration of cells), to ensure maximum contact of cells and beads. For example, in one embodiment, a concentration of 2 billion cells/ml is used. In one embodiment, a concentration of 1 billion cells/ml is used. In a further embodiment, greater than 100 million cells/ml is used. In a further embodiment, a concentration of cells of 10, 15, 20, 25, 30, 35, 40, 45, or 50 million cells/ml is used. In yet another embodiment, a concentration of cells from 75, 80, 85, 90, 95, or 100 million cells/ml is used. In further embodiments, concentrations of 125 or 150 million cells/ml can be used. Using high concentrations may result in increased cell yield, cell activation, and cell expansion. Further, use of high cell concentrations may allow more efficient capture of cells that may weakly express target antigens of interest, such as CD28-negative T cells, or from samples where there are many tumor cells present (*i.e.*, leukemic blood, tumor tissue, etc.). Such populations of cells may have therapeutic value and would be desirable to obtain. In some embodiments, using high concentration of cells allows more efficient selection of CD8+ T cells that normally have weaker CD28 expression.

[00407] In some embodiments, it may be desirable to use lower concentrations of cells. By significantly diluting the mixture of T cells and surface (e.g., particles such as beads), interactions between the particles and cells is minimized. This selects for cells that express high amounts of desired antigens to be bound to the particles. For example, CD4⁺ T cells express higher levels of CD28 and are more efficiently captured than CD8⁺ T cells in dilute concentrations. In some embodiments, the concentration of cells used is 5×10^6 /ml. In some embodiments, the concentration used can be from about 1×10^5 /ml to 1×10^6 /ml, and any integer value in between.

[00408] In some embodiments, the cells may be incubated on a rotator for varying lengths of time at varying speeds at either 2–10°C, or at room temperature.

[00409] T cells for stimulation can also be frozen after a washing step. Without being bound by theory, the freeze and subsequent thaw step may provide a more uniform product by removing granulocytes and to some extent monocytes in the cell population. After the washing step that removes plasma and platelets, the cells may be suspended in a freezing solution. While many freezing solutions and parameters are known in the art and will be useful in this context, one method involves using PBS containing 20% DMSO and 8% human serum albumin, or culture media containing 10% dextran 40 and 5% dextrose, 20% human serum albumin and 7.5% DMSO, or 31.25% plasmalyte-A, 31.25% dextrose 5%, 0.45% NaCl, 10% dextran 40 and 5% dextrose, 20% human serum albumin, and 7.5% DMSO or other suitable cell freezing media containing for example, Hespan and PlasmaLyte A. The cells then are frozen to –80°C at a rate of 1° per minute and stored in the vapor phase of a liquid nitrogen storage tank. Other methods of controlled freezing may be used as well as uncontrolled freezing immediately at –20°C or in liquid nitrogen.

[00410] In some embodiments, cryopreserved cells are thawed and washed as described herein and allowed to rest for one hour at room temperature prior to activation.

[00411] Also contemplated in the present disclosure is the collection of blood samples or apheresis product from a subject at a time period prior to when the expanded cells as described herein might be needed. As such, the source of the cells to be expanded can be collected at any time point necessary, and desired cells, such as T cells, isolated and frozen for later use in T cell therapy for any number of diseases or conditions that would benefit from T cell therapy, such as those described herein. In one embodiment, a blood sample or an apheresis is taken from a generally healthy subject. In certain embodiments, a blood sample or an apheresis is taken from a generally healthy subject who is at risk of developing a disease, but who has not yet developed a disease, and the cells of interest are isolated and frozen for later use. In certain embodiments, the T cells may be expanded, frozen, and used at a later time. In certain embodiments, samples are collected from a patient shortly after diagnosis of a particular disease as described herein but

prior to any treatments. In a further embodiment, the cells are isolated from a blood sample or an apheresis from a subject prior to any number of relevant treatment modalities, including but not limited to treatment with agents such as natalizumab, efalizumab, antiviral agents, chemotherapy, radiation, immunosuppressive agents, such as cyclosporin, azathioprine, methotrexate, mycophenolate, and FK506, antibodies, or other immunoablative agents such as CAMPATH, anti-CD3 antibodies, cytoxan, fludarabine, cyclosporin, FK506, rapamycin, mycophenolic acid, steroids, FR901228, and irradiation. These drugs inhibit either the calcium dependent phosphatase calcineurin (cyclosporine and FK506) or inhibit the p70S6 kinase that is important for growth factor induced signaling (rapamycin) (Liu et al., Cell 66:807-815 (1991); Henderson et al., Immun 73:316-321 (1991); Bierer et al., Curr. Opin. Immun. 5:763-773 (1993)). In a further embodiment, the cells are isolated for a patient and frozen for later use in conjunction with (*e.g.*, before, simultaneously or following) bone marrow or stem cell transplantation, T cell ablative therapy using either chemotherapy agents such as, fludarabine, external-beam radiation therapy (XRT), cyclophosphamide, or antibodies such as OKT3 or CAMPATH. In another embodiment, the cells are isolated prior to and can be frozen for later use for treatment following B-cell ablative therapy such as agents that react with CD20, *e.g.*, Rituxan.

[00412] In some embodiments, T cells are obtained from a patient directly following treatment. In this regard, it has been observed that following certain cancer treatments, in particular treatments with drugs that damage the immune system, shortly after treatment during the period when patients would normally be recovering from the treatment, the quality of T cells obtained may be optimal or improved for their ability to expand *ex vivo*. Likewise, following *ex vivo* manipulation using the methods described herein, these cells may be in a preferred state for enhanced engraftment and *in vivo* expansion. Thus, it is contemplated within the context of the present disclosure to collect blood cells, including T cells, dendritic cells, or other cells of the hematopoietic lineage, during this recovery phase. Further, in certain embodiments, mobilization (for example, mobilization with GM-CSF) and conditioning regimens can be used to create a condition in a subject wherein repopulation, recirculation, regeneration, and/or expansion of particular cell types is favored, especially during a defined window of time following therapy. Illustrative cell types include T cells, B cells, dendritic cells, and other cells of the immune system.

5.4.4. Activation and Expansion of T Cells

[00413] In some embodiments, prior to or after genetic modification of the T cells with the CARs described herein, the T cells can be activated and expanded generally using methods as described, for example, in U.S. Pat. Nos. 6,352,694; 6,534,055; 6,905,680; 6,692,964; 5,858,358; 6,887,466; 6,905,681; 7,144,575; 7,067,318; 7,172,869; 7,232,566; 7,175,843; 5,883,223; 6,905,874; 6,797,514; 6,867,041; and U.S. Patent Application Publication No. 20060121005.

[00414] Generally, T cells can be expanded by contact with a surface having attached thereto an agent that stimulates a CD3/TCR complex associated signal and a ligand that stimulates a co-stimulatory molecule on the surface of the T cells. In particular, T cell populations may be stimulated as described herein, such as by contact with an anti-CD3 antibody, or antigen-binding fragment thereof, or an anti-CD2 antibody immobilized on a surface, or by contact with a protein kinase C activator (*e.g.*, bryostatin) in conjunction with a calcium ionophore. For co-stimulation of an accessory molecule on the surface of the T cells, a ligand that binds the accessory molecule is used. For example, a population of T cells can be contacted with an anti-CD3 antibody and an anti-CD28 antibody, under conditions appropriate for stimulating proliferation of the T cells. To stimulate proliferation of either CD4⁺ T cells or CD8⁺ T cells, an anti-CD3 antibody and an anti-CD28 antibody. Examples of an anti-CD3 antibody include UCHT1, OKT3, HIT3a (BioLegend, San Diego, US) can be used as can other methods commonly known in the art (Graves J, et al., J.Immunol. 146:2102 (1991); Li B, et al., Immunology 116:487 (2005); Rivollier A, et al., Blood 104:4029 (2004)). Examples of an anti-CD28 antibody include 9.3, B-T3, XR-CD28 (Diacclone, Besancon, France) can be used as can other methods commonly known in the art (Berg et al., Transplant Proc. 30(8):3975-3977 (1998); Haanen et al., J. Exp. Med. 190(9):13191328 (1999); Garland et al., J. Immunol Meth. 227(1-2):53-63 (1999)).

[00415] In some embodiments, the primary stimulatory signal and the co-stimulatory signal for the T cell may be provided by different protocols. For example, the agents providing each signal may be in solution or coupled to a surface. When coupled to a surface, the agents may be coupled to the same surface (*i.e.*, in “cis” formation) or to separate surfaces (*i.e.*, in “trans” formation). Alternatively, one agent may be coupled to a surface and the other agent in solution. In one embodiment, the agent providing the co-stimulatory signal is bound to a cell surface and the agent providing the primary activation signal is in solution or coupled to a surface. In certain embodiments, both agents can be in solution. In another embodiment, the agents may be in soluble form, and then cross-linked to a surface, such as a cell expressing Fc receptors or an antibody or other binding agent which will bind to the agents. In this regard, see for example, U.S. Patent Application Publication Nos. 20040101519 and 20060034810 for artificial antigen presenting cells (aAPCs) that are contemplated for use in activating and expanding T cells in certain embodiments in the present disclosure.

[00416] In some embodiments, the T cells, are combined with agent-coated beads, the beads and the cells are subsequently separated, and then the cells are cultured. In an alternative embodiment, prior to culture, the agent-coated beads and cells are not separated but are cultured together. In a further embodiment, the beads and cells are first concentrated by application of a force, such as a magnetic force, resulting in increased ligation of cell surface markers, thereby inducing cell stimulation.

[00417] By way of example, cell surface proteins may be ligated by allowing paramagnetic beads to which anti-CD3 and anti-CD28 are attached (3×28 beads) to contact the T cells. In one embodiment, the cells (for example, 10^4 to 4×10^8 T cells) and beads (for example, anti-CD3/CD28 MACSiBead particles) at a recommended titer of 1:100 are combined in a buffer, preferably PBS (without divalent cations such as, calcium and magnesium). Those of ordinary skill in the art can readily appreciate any cell concentration may be used. For example, the target cell may be very rare in the sample and comprise only 0.01% of the sample or the entire sample (*i.e.*, 100%) may comprise the target cell of interest. Accordingly, any cell number is within the context of the present disclosure. In certain embodiments, it may be desirable to significantly decrease the volume in which particles and cells are mixed together (*i.e.*, increase the concentration of cells), to ensure maximum contact of cells and particles. For example, in one embodiment, a concentration of about 2 billion cells/mL is used. In another embodiment, greater than 100 million cells/mL is used. In a further embodiment, a concentration of cells of 10, 15, 20, 25, 30, 35, 40, 45, or 50 million cells/mL is used. In yet another embodiment, a concentration of cells from 75, 80, 85, 90, 95, or 100 million cells/mL is used. In further embodiments, concentrations of 125 or 150 million cells/mL can be used. Using high concentrations may result in increased cell yield, cell activation, and cell expansion. Further, use of high cell concentrations may allow more efficient capture of cells that may weakly express target antigens of interest, such as CD28-negative T cells. Such populations of cells may have therapeutic value and would be desirable to obtain in certain embodiments. For example, using high concentration of cells allows more efficient selection of CD8⁺ T cells that normally have weaker CD28 expression.

[00418] In some embodiments, the mixture may be cultured for several hours (about 3 hours) to about 14 days or any hourly integer value in between. In another embodiment, the mixture may be cultured for 21 days. In one embodiment, the beads and the T cells are cultured together for about eight days. In another embodiment, the beads and T cells are cultured together for 2-3 days. Several cycles of stimulation may also be desired such that culture time of T cells can be 60 days or more. Conditions appropriate for T cell culture include an appropriate media (*e.g.*, Minimal Essential Media or RPMI Media 1640 or, X-vivo 15, (Lonza)) that may contain factors necessary for proliferation and viability, including serum (*e.g.*, fetal bovine or human serum), interleukin-2 (IL-2), insulin, IFN- γ , IL-4, IL-7, GM-CSF, IL-10, IL-12, IL-15, TGF β , and TNF- α or any other additives for the growth of cells known to the skilled artisan. Other additives for the growth of cells include, but are not limited to, surfactant, plasmanate, and reducing agents such as N-acetyl-cysteine and 2-mercaptoethanol. Media can include RPMI 1640, AIM-V, DMEM, MEM, α -MEM, F-12, X-Vivo 15, and X-Vivo 20, optimizer, with added amino acids, sodium pyruvate, and vitamins, either serum-free or supplemented with an appropriate amount of serum (or plasma) or a defined set of hormones, and/or an amount of cytokine(s) sufficient for the

growth and expansion of T cells. Antibiotics, *e.g.*, penicillin and streptomycin, are included only in experimental cultures, not in cultures of cells that are to be infused into a subject. The target cells are maintained under conditions necessary to support growth, for example, an appropriate temperature (*e.g.*, 37°C) and atmosphere (*e.g.*, air plus 5% CO₂). T cells that have been exposed to varied stimulation times may exhibit different characteristics. For example, typical blood or apheresed peripheral blood mononuclear cell products have a helper T cell population (TH, CD4+) that is greater than the cytotoxic or suppressor T cell population (TC, CD8). *Ex vivo* expansion of T cells by stimulating CD3 and CD28 receptors produces a population of T cells that prior to about days 8-9 consists predominately of TH cells, while after about days 8-9, the population of T cells comprises an increasingly greater population of TC cells. Accordingly, depending on the purpose of treatment, infusing a subject with a T cell population comprising predominately of TH cells may be advantageous. Similarly, if an antigen-specific subset of TC cells has been isolated it may be beneficial to expand this subset to a greater degree.

[00419] Further, in addition to CD4 and CD8 markers, other phenotypic markers vary significantly, but in large part, reproducibly during the course of the cell expansion process. Thus, such reproducibility enables the ability to tailor an activated T cell product for specific purposes.

5.5. Polynucleotides

[00420] In certain embodiments, the disclosure provides polynucleotides that encode the single domain antibodies that bind to CD20 and fusion proteins comprising the single domain antibodies that bind to CD20 described herein. The polynucleotides of the disclosure can be in the form of RNA or in the form of DNA. DNA includes cDNA, genomic DNA, and synthetic DNA; and can be double-stranded or single-stranded, and if single stranded can be the coding strand or non-coding (anti-sense) strand. In some embodiments, the polynucleotide is in the form of cDNA. In some embodiments, the polynucleotide is a synthetic polynucleotide. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 41, such as a nucleic acid having SEQ ID NO: 64. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 42, such as a nucleic acid having SEQ ID NO: 65. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 43, such as a nucleic acid having SEQ ID NO: 66. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 44, such as a nucleic acid having SEQ ID NO: 67. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of

[illegible]

NO: 61. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 62. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 63. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 183, such as a nucleic acid having SEQ ID NO: 186. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 184, such as a nucleic acid having SEQ ID NO: 187. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the single domain antibody having the sequence of SEQ ID NO: 185, such as a nucleic acid having SEQ ID NO: 188.

[00421] In certain embodiments, the disclosure provides polynucleotides that encode the CD20 CAR provided herein. The polynucleotides of the disclosure can be in the form of RNA or in the form of DNA. DNA includes cDNA, genomic DNA, and synthetic DNA; and can be double-stranded or single-stranded, and if single stranded can be the coding strand or non-coding (anti-sense) strand. In some embodiments, the polynucleotide is in the form of cDNA. In some embodiments, the polynucleotide is a synthetic polynucleotide. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 81, such as a nucleic acid having SEQ ID NO: 108. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 82, such as a nucleic acid having SEQ ID NO: 109. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 83, such as a nucleic acid having SEQ ID NO: 110. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 84, such as a nucleic acid having SEQ ID NO: 111. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 85, such as a nucleic acid having SEQ ID NO: 112. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 86, such as a nucleic acid having SEQ ID NO: 113. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 87, such as a nucleic acid having SEQ ID NO: 114. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 88, such as a nucleic acid having SEQ ID NO: 115. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 89, such as a nucleic acid having SEQ ID NO: 116. In

exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 90, such as a nucleic acid having SEQ ID NO: 117. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 91, such as a nucleic acid having SEQ ID NO: 118. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 92, such as a nucleic acid having SEQ ID NO: 119. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 93, such as a nucleic acid having SEQ ID NO: 120. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 94, such as a nucleic acid having SEQ ID NO: 121. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 95, such as a nucleic acid having SEQ ID NO: 122. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 96, such as a nucleic acid having SEQ ID NO: 123. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 97, such as a nucleic acid having SEQ ID NO: 124. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 98, such as a nucleic acid having SEQ ID NO: 125. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 99, such as a nucleic acid having SEQ ID NO: 126. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 100, such as a nucleic acid having SEQ ID NO: 127. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 101, such as a nucleic acid having SEQ ID NO: 128. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 102, such as a nucleic acid having SEQ ID NO: 129. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 103, such as a nucleic acid having SEQ ID NO: 130. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 104, such as a nucleic acid having SEQ ID NO: 131. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 105, such as a nucleic acid having SEQ ID NO: 132. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 106, such as a

nucleic acid having SEQ ID NO: 133. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 189, such as a nucleic acid having SEQ ID NO: 192. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 190, such as a nucleic acid having SEQ ID NO: 193. In exemplary embodiments, the nucleic acid molecule provided herein comprises a sequence that encodes the CAR having the sequence of SEQ ID NO: 191, such as a nucleic acid having SEQ ID NO: 194.

[00422] The present disclosure further relates to variants of the polynucleotides described herein, wherein the variant encodes, for example, fragments, analogs, and/or derivatives of the single domain antibody or CAR that binds CD20 of the disclosure. In certain embodiments, the present disclosure provides a polynucleotide comprising a polynucleotide having a nucleotide sequence at least about 75% identical, at least about 80% identical, at least about 85% identical, at least about 90% identical, at least about 95% identical, and in some embodiments, at least about 96%, 97%, 98% or 99% identical to a polynucleotide encoding the single domain antibody or CAR that binds CD20 of the disclosure. As used herein, the phrase “a polynucleotide having a nucleotide sequence at least, for example, 95% “identical” to a reference nucleotide sequence” is intended to mean that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence can include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence can be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence can be inserted into the reference sequence. These mutations of the reference sequence can occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

[00423] The polynucleotide variants can contain alterations in the coding regions, non-coding regions, or both. In some embodiments, a polynucleotide variant contains alterations which produce silent substitutions, additions, or deletions, but does not alter the properties or activities of the encoded polypeptide. In some embodiments, a polynucleotide variant comprises silent substitutions that results in no change to the amino acid sequence of the polypeptide (due to the degeneracy of the genetic code). Polynucleotide variants can be produced for a variety of reasons, for example, to optimize codon expression for a particular host (*i.e.*, change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*). In some embodiments, a polynucleotide variant comprises at least one silent mutation in a non-coding or a coding region of the sequence.

[00424] In some embodiments, a polynucleotide variant is produced to modulate or alter expression (or expression levels) of the encoded polypeptide. In some embodiments, a polynucleotide variant is produced to increase expression of the encoded polypeptide. In some embodiments, a polynucleotide variant is produced to decrease expression of the encoded polypeptide. In some embodiments, a polynucleotide variant has increased expression of the encoded polypeptide as compared to a parental polynucleotide sequence. In some embodiments, a polynucleotide variant has decreased expression of the encoded polypeptide as compared to a parental polynucleotide sequence.

[00425] Also provided are vectors comprising the nucleic acid molecules described herein. In an embodiment, the nucleic acid molecules can be incorporated into a recombinant expression vector. The present disclosure provides recombinant expression vectors comprising any of the nucleic acids of the disclosure. As used herein, the term “recombinant expression vector” means a genetically-modified oligonucleotide or polynucleotide construct that permits the expression of an mRNA, protein, polypeptide, or peptide by a host cell, when the construct comprises a nucleotide sequence encoding the mRNA, protein, polypeptide, or peptide, and the vector is contacted with the cell under conditions sufficient to have the mRNA, protein, polypeptide, or peptide expressed within the cell. The vectors described herein are not naturally-occurring as a whole; however, parts of the vectors can be naturally-occurring. The described recombinant expression vectors can comprise any type of nucleotides, including, but not limited to DNA and RNA, which can be single-stranded or double-stranded, synthesized or obtained in part from natural sources, and which can contain natural, non-natural or altered nucleotides. The recombinant expression vectors can comprise naturally-occurring or non-naturally-occurring internucleotide linkages, or both types of linkages. The non-naturally occurring or altered nucleotides or internucleotide linkages do not hinder the transcription or replication of the vector.

[00426] In an embodiment, the recombinant expression vector of the disclosure can be any suitable recombinant expression vector, and can be used to transform or transfect any suitable host. Suitable vectors include those designed for propagation and expansion or for expression or both, such as plasmids and viruses. The vector can be selected from the group consisting of the pUC series (Fermentas Life Sciences, Glen Burnie, Md.), the pBluescript series (Stratagene, LaJolla, Calif.), the pET series (Novagen, Madison, Wis.), the pGEX series (Pharmacia Biotech, Uppsala, Sweden), and the pEX series (Clontech, Palo Alto, Calif.). Bacteriophage vectors, such as λ GT10, λ GT11, λ EMBL4, and λ NM1149, λ ZapII (Stratagene) can be used. Examples of plant expression vectors include pBI01, pBI01.2, pBI121, pBI101.3, and pBIN19 (Clontech). Examples of animal expression vectors include pEUK-Cl, pMAM, and pMAMneo (Clontech). The recombinant expression vector may be a viral vector, *e.g.*, a retroviral vector, *e.g.*, a gamma retroviral vector.

[00427] In an embodiment, the recombinant expression vectors are prepared using standard recombinant DNA techniques described in, for example, Sambrook et al., supra, and Ausubel et al., supra. Constructs of expression vectors, which are circular or linear, can be prepared to contain a replication system functional in a prokaryotic or eukaryotic host cell. Replication systems can be derived, *e.g.*, from ColE1, SV40, 2 μ plasmid, λ , bovine papilloma virus, and the like.

[00428] The recombinant expression vector may comprise regulatory sequences, such as transcription and translation initiation and termination codons, which are specific to the type of host (*e.g.*, bacterium, plant, fungus, or animal) into which the vector is to be introduced, as appropriate, and taking into consideration whether the vector is DNA- or RNA-based.

[00429] The recombinant expression vector can include one or more marker genes, which allow for selection of transformed or transfected hosts. Marker genes include biocide resistance, *e.g.*, resistance to antibiotics, heavy metals, etc., complementation in an auxotrophic host to provide prototrophy, and the like. Suitable marker genes for the described expression vectors include, for instance, neomycin/G418 resistance genes, histidinol x resistance genes, histidinol resistance genes, tetracycline resistance genes, and ampicillin resistance genes.

[00430] The recombinant expression vector can comprise a native or normative promoter operably linked to the nucleotide sequence of the disclosure. The selection of promoters, *e.g.*, strong, weak, tissue-specific, inducible and developmental-specific, is within the ordinary skill of the artisan. Similarly, the combining of a nucleotide sequence with a promoter is also within the skill of the artisan. The promoter can be a non-viral promoter or a viral promoter, *e.g.*, a cytomegalovirus (CMV) promoter, an RSV promoter, an SV40 promoter, or a promoter found in the long-terminal repeat of the murine stem cell virus.

[00431] The recombinant expression vectors can be designed for either transient expression, for stable expression, or for both. Also, the recombinant expression vectors can be made for constitutive expression or for inducible expression.

[00432] Further, the recombinant expression vectors can be made to include a suicide gene. As used herein, the term "suicide gene" refers to a gene that causes the cell expressing the suicide gene to die. The suicide gene can be a gene that confers sensitivity to an agent, *e.g.*, a drug, upon the cell in which the gene is expressed, and causes the cell to die when the cell is contacted with or exposed to the agent. Suicide genes are known in the art and include, for example, the Herpes Simplex Virus (HSV) thymidine kinase (TK) gene, cytosine deaminase, purine nucleoside phosphorylase, and nitroreductase.

[00433] In certain embodiments, a polynucleotide is isolated. In certain embodiments, a polynucleotide is substantially pure.

[00434] Also provided are host cells comprising the nucleic acid molecules described herein.

The host cell may be any cell that contains a heterologous nucleic acid. The heterologous nucleic acid can be a vector (e.g., an expression vector). For example, a host cell can be a cell from any organism that is selected, modified, transformed, grown, used or manipulated in any way, for the production of a substance by the cell, for example the expression by the cell of a gene, a DNA or RNA sequence, a protein or an enzyme. An appropriate host may be determined. For example, the host cell may be selected based on the vector backbone and the desired result. By way of example, a plasmid or cosmid can be introduced into a prokaryote host cell for replication of several types of vectors. Bacterial cells such as, but not limited to DH5 α , JM109, and KCB, SURE[®] Competent Cells, and SOLOPACK Gold Cells, can be used as host cells for vector replication and/or expression. Additionally, bacterial cells such as *E. coli* LE392 could be used as host cells for phage viruses. Eukaryotic cells that can be used as host cells include, but are not limited to yeast (e.g., YPH499, YPH500 and YPH501), insects and mammals. Examples of mammalian eukaryotic host cells for replication and/or expression of a vector include, but are not limited to, HeLa, NIH3T3, Jurkat, 293, COS, Saos, PC12, SP2/0 (American Type Culture Collection (ATCC), Manassas, VA, CRL-1581), NS0 (European Collection of Cell Cultures (ECACC), Salisbury, Wiltshire, UK, ECACC No. 85110503), FO (ATCC CRL-1646) and Ag653 (ATCC CRL-1580) murine cell lines. An exemplary human myeloma cell line is U266 (ATCC CRL-TIB-196). Other useful cell lines include those derived from Chinese Hamster Ovary (CHO) cells such as CHO-K1SV (Lonza Biologics, Walkersville, MD), CHO-K1 (ATCC CRL-61) or DG44.

5.6. Pharmaceutical Compositions

[00435] In one aspect, the present disclosure further provides pharmaceutical compositions comprising a single domain antibody, a binding molecule or therapeutic molecule comprising a single domain antibody, or an engineered immune effector cell of the present disclosure. In some embodiments, a pharmaceutical composition comprises a therapeutically effective amount of the single domain antibody, the binding molecule or therapeutic molecule comprising the single domain antibody, or the engineered immune effector cell of the present disclosure and a pharmaceutically acceptable excipient.

[00436] In some embodiments, provided herein is a pharmaceutical composition comprising a therapeutically effective amount of the single domain antibody provided herein and a pharmaceutically acceptable excipient.

[00437] In some embodiments, provided herein is a pharmaceutical composition comprising a therapeutically effective amount of the therapeutic molecule (such as a fusion protein, immunoconjugate, and a multispecific binding molecule) comprising the single domain antibody provided herein and a pharmaceutically acceptable excipient.

[00438] In other embodiments, provided herein is a pharmaceutical composition comprising a therapeutically effective amount of CAR comprising the single domain antibody provided herein and a pharmaceutically acceptable excipient.

[00439] In other embodiments, provided herein is a pharmaceutical composition comprising a therapeutically effective amount of engineered immune effector cells provided herein and a pharmaceutically acceptable excipient.

[00440] In other embodiments, provided herein is a pharmaceutical composition comprising a therapeutically effective amount of a nucleic acid provided herein, *e.g.*, in a vector, and a pharmaceutically acceptable excipient, *e.g.*, suitable for gene therapy.

[00441] In a specific embodiment, the term “excipient” can also refer to a diluent, adjuvant (*e.g.*, Freund's adjuvant (complete or incomplete), carrier or vehicle. Pharmaceutical excipients can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid excipients. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. Examples of suitable pharmaceutical excipients are described in Remington's Pharmaceutical Sciences (1990) Mack Publishing Co., Easton, PA. Such compositions will contain a prophylactically or therapeutically effective amount of the active ingredient provided herein, such as in purified form, together with a suitable amount of excipient so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

[00442] In some embodiments, the choice of excipient is determined in part by the particular cell, binding molecule, and/or antibody, and/or by the method of administration. Accordingly, there are a variety of suitable formulations.

[00443] Typically, acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers, antioxidants including ascorbic acid, methionine, Vitamin E, sodium metabisulfite; preservatives, isotonicifiers, stabilizers, metal complexes (*e.g.* Zn-protein complexes); chelating agents such as EDTA and/or non-ionic surfactants.

[00444] Buffers may be used to control the pH in a range which optimizes the therapeutic effectiveness, especially if stability is pH dependent. Suitable buffering agents for use with the present disclosure include both organic and inorganic acids and salts thereof. For example,

citrate, phosphate, succinate, tartrate, fumarate, gluconate, oxalate, lactate, acetate. Additionally, buffers may comprise histidine and trimethylamine salts such as Tris.

[00445] Preservatives may be added to retard microbial growth. Suitable preservatives for use with the present disclosure include octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium halides (*e.g.*, chloride, bromide, iodide), benzethonium chloride; thimerosal, phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol, 3-pentanol, and m-cresol.

[00446] Tonicity agents, sometimes known as “stabilizers” can be present to adjust or maintain the tonicity of liquid in a composition. When used with large, charged biomolecules such as proteins and antibodies, they are often termed “stabilizers” because they can interact with the charged groups of the amino acid side chains, thereby lessening the potential for inter and intramolecular interactions. Exemplary tonicity agents include polyhydric sugar alcohols, trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol and mannitol.

[00447] Additional exemplary excipients include: (1) bulking agents, (2) solubility enhancers, (3) stabilizers and (4) agents preventing denaturation or adherence to the container wall. Such excipients include: polyhydric sugar alcohols (enumerated above); amino acids such as alanine, glycine, glutamine, asparagine, histidine, arginine, lysine, ornithine, leucine, 2-phenylalanine, glutamic acid, threonine, etc.; organic sugars or sugar alcohols such as sucrose, lactose, lactitol, trehalose, stachyose, mannose, sorbose, xylose, ribose, ribitol, myoinositol, myoinositol, galactose, galactitol, glycerol, cyclitols (*e.g.*, inositol), polyethylene glycol; sulfur containing reducing agents, such as urea, glutathione, thiocetic acid, sodium thioglycolate, thioglycerol, α -monothioglycerol and sodium thio sulfate; low molecular weight proteins such as human serum albumin, bovine serum albumin, gelatin or other immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; monosaccharides (*e.g.*, xylose, mannose, fructose, glucose; disaccharides (*e.g.*, lactose, maltose, sucrose); trisaccharides such as raffinose; and polysaccharides such as dextrin or dextran.

[00448] Non-ionic surfactants or detergents (also known as “wetting agents”) may be present to help solubilize the therapeutic agent as well as to protect the therapeutic protein against agitation-induced aggregation, which also permits the formulation to be exposed to shear surface stress without causing denaturation of the active therapeutic protein or antibody. Suitable non-ionic surfactants include, *e.g.*, polysorbates (20, 40, 60, 65, 80, etc.), polyoxamers (184, 188, etc.), PLURONIC® polyols, TRITON®, polyoxyethylene sorbitan monoethers (TWEEN®-20, TWEEN®-80, etc.), lauromacrogol 400, polyoxyl 40 stearate, polyoxyethylene hydrogenated castor oil 10, 50 and 60, glycerol monostearate, sucrose fatty acid ester, methyl cellulose and carboxymethyl cellulose. Anionic detergents that can be used include sodium lauryl sulfate, dioctyle sodium sulfosuccinate and dioctyl sodium sulfonate. Cationic detergents include

benzalkonium chloride or benzethonium chloride.

[00449] In order for the pharmaceutical compositions to be used for *in vivo* administration, they are preferably sterile. The pharmaceutical composition may be rendered sterile by filtration through sterile filtration membranes. The pharmaceutical compositions herein generally can be placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[00450] The route of administration is in accordance with known and accepted methods, such as by single or multiple bolus or infusion over a long period of time in a suitable manner, *e.g.*, injection or infusion by subcutaneous, intravenous, intraperitoneal, intramuscular, intraarterial, intralesional or intraarticular routes, topical administration, inhalation or by sustained release or extended-release means.

[00451] In another embodiment, a pharmaceutical composition can be provided as a controlled release or sustained release system. In one embodiment, a pump may be used to achieve controlled or sustained release (*see, e.g.*, Sefton, Crit. Ref. Biomed. Eng. 14:201-40 (1987); Buchwald *et al.*, Surgery 88:507-16 (1980); and Saudek *et al.*, N. Engl. J. Med. 321:569-74 (1989)). In another embodiment, polymeric materials can be used to achieve controlled or sustained release of a prophylactic or therapeutic agent (*e.g.*, a fusion protein as described herein) or a composition provided herein (*see, e.g.*, Medical Applications of Controlled Release (Langer and Wise eds., 1974); Controlled Drug Bioavailability, Drug Product Design and Performance (Smolen and Ball eds., 1984); Ranger and Peppas, J. Macromol. Sci. Rev. Macromol. Chem. 23:61-126 (1983); Levy *et al.*, Science 228:190-92 (1985); During *et al.*, Ann. Neurol. 25:351-56 (1989); Howard *et al.*, J. Neurosurg. 71:105-12 (1989); U.S. Pat. Nos. 5,679,377; 5,916,597; 5,912,015; 5,989,463; and 5,128,326; PCT Publication Nos. WO 99/15154 and WO 99/20253). Examples of polymers used in sustained release formulations include, but are not limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl alcohol), polyacrylamide, poly(ethylene glycol), polylactides (PLA), poly(lactide-co-glycolides) (PLGA), and polyorthoesters. In one embodiment, the polymer used in a sustained release formulation is inert, free of leachable impurities, stable on storage, sterile, and biodegradable. In yet another embodiment, a controlled or sustained release system can be placed in proximity of a particular target tissue, for example, the nasal passages or lungs, thus requiring only a fraction of the systemic dose (*see, e.g.*, Goodson, Medical Applications of Controlled Release Vol. 2, 115-38 (1984)). Controlled release systems are discussed, for example, by Langer, Science 249:1527-33 (1990). Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more agents as described herein (*see, e.g.*, U.S. Pat. No. 4,526,938, PCT publication Nos.

WO 91/05548 and WO 96/20698, Ning *et al.*, Radiotherapy & Oncology 39:179-89 (1996); Song *et al.*, PDA J. of Pharma. Sci. & Tech. 50:372-97 (1995); Cleek *et al.*, Pro. Int'l. Symp. Control. Rel. Bioact. Mater. 24:853-54 (1997); and Lam *et al.*, Proc. Int'l. Symp. Control Rel. Bioact. Mater. 24:759-60 (1997)).

[00452] The pharmaceutical compositions described herein may also contain more than one active compound or agent as necessary for the particular indication being treated. Alternatively, or in addition, the composition may comprise a cytotoxic agent, chemotherapeutic agent, cytokine, immunosuppressive agent, or growth inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[00453] The active ingredients may also be entrapped in microcapsules prepared, for example, by coascervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 18th edition.

[00454] Various compositions and delivery systems are known and can be used with the therapeutic agents provided herein, including, but not limited to, encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the single domain antibody or therapeutic molecule provided herein, construction of a nucleic acid as part of a retroviral or other vector, *etc.*

[00455] In some embodiments, the pharmaceutical composition provided herein contains the binding molecules and/or cells in amounts effective to treat or prevent the disease or disorder, such as a therapeutically effective or prophylactically effective amount. Therapeutic or prophylactic efficacy in some embodiments is monitored by periodic assessment of treated subjects. For repeated administrations over several days or longer, depending on the condition, the treatment is repeated until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful and can be determined.

5.7. Methods and Uses

[00456] In another aspect, provided herein are methods for using and uses of the CD20 binding molecules provided herein, including the anti-CD20 VHH, chimeric antigen receptors (CARs), and/or engineered cells expressing the recombinant receptors.

5.7.1. Therapeutic Methods and Uses

[00457] Such methods and uses include therapeutic methods and uses, for example, involving administration of the molecules, cells, or compositions containing the same, to a subject having a disease, condition, or disorder expressing or associated with CD20 expression, and/or in which cells or tissues express CD20. In some embodiments, the molecule, cell, and/or composition is

administered in an effective amount to effect treatment of the disease or disorder. Uses include uses of the antibodies and cells in such methods and treatments, and in the preparation of a medicament in order to carry out such therapeutic methods. In some embodiments, the methods are carried out by administering the antibodies or cells, or compositions comprising the same, to the subject having or suspected of having the disease or condition. In some embodiments, the methods thereby treat the disease or disorder in the subject.

[00458] In some embodiments, the treatment provided herein cause complete or partial amelioration or reduction of a disease or disorder, or a symptom, adverse effect or outcome, or phenotype associated therewith. Desirable effects of treatment include, but are not limited to, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastasis, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. The terms include, but do not imply, complete curing of a disease or complete elimination of any symptom or effect(s) on all symptoms or outcomes.

[00459] As used herein, in some embodiments, the treatment provided herein delay development of a disease or disorder, *e.g.*, defer, hinder, slow, retard, stabilize, suppress and/or postpone development of the disease (such as cancer). This delay can be of varying lengths of time, depending on the history of the disease and/or individual being treated. As is evident to one skilled in the art, a sufficient or significant delay can, in effect, encompass prevention, in that the individual does not develop the disease or disorder. For example, a late stage cancer, such as development of metastasis, may be delayed. In other embodiments, the method or the use provided herein prevents a disease or disorder.

[00460] In some embodiments, the disease or disorder is a CD20 associated disease or disorder. In some embodiments, the disease or disorder is a B cell associated disease or disorder. In some embodiments, the disease or disorder is a B cell malignancy. In some embodiments, the B cell malignancy is a B cell leukemia or B cell lymphoma. In a specific embodiment, the disease or disorder is marginal zone lymphoma (*e.g.*, splenic marginal zone lymphoma). In a specific embodiment, the disease or disorder is diffuse large B cell lymphoma (DLBCL). In another specific embodiment, the disease or disorder is mantle cell lymphoma (MCL). In another specific embodiment, the disease or disorder is primary central nervous system (CNS) lymphoma. In another specific embodiment, the disease or disorder is primary mediastinal B cell lymphoma (PMBL). In another specific embodiment, the disease or disorder is small lymphocytic lymphoma (SLL). In another specific embodiment, the disease or disorder is B cell prolymphocytic leukemia (B-PLL). In another specific embodiment, the disease or disorder is follicular lymphoma (FL). In another specific embodiment, the disease or disorder is burkitt lymphoma. In another specific embodiment, the disease or disorder is primary intraocular

lymphoma. In another specific embodiment, the disease or disorder is chronic lymphocytic leukemia (CLL). In another specific embodiment, the disease or disorder is acute lymphoblastic leukemia (ALL). In another specific embodiment, the disease or disorder is hairy cell leukemia (HCL). In another specific embodiment, the disease or disorder is precursor B lymphoblastic leukemia. In another specific embodiment, the disease or disorder is non-hodgkin lymphoma (NHL). In another specific embodiment, the disease or disorder is high-grade B-cell lymphoma (HGBL). In another specific embodiment, the disease or disorder is multiple myeloma (MM). In other embodiments, the disease or disorder is a relapsed or refractory B cell malignancy, such as relapsed or refractory ALL (R/R ALL).

[00461] In other embodiments, the disease or disorder is an autoimmune and inflammatory disease, including, *e.g.*, those associated with inappropriate or enhanced B cell numbers and/or activation.

[00462] In some embodiments, the methods include adoptive cell therapy, whereby genetically engineered cells expressing the provided CD20-targeted CARs are administered to a subject. Such administration can promote activation of the cells (*e.g.*, T cell activation) in a CD20-targeted manner, such that the cells of the disease or disorder are targeted for destruction.

[00463] In some embodiments, the methods include administration of the cells or a composition containing the cells to a subject, tissue, or cell, such as one having, at risk for, or suspected of having the disease or disorder. In some embodiments, the cells, populations, and compositions are administered to a subject having the particular disease or disorder to be treated, *e.g.*, via adoptive cell therapy, such as adoptive T cell therapy. In some embodiments, the cells or compositions are administered to the subject, such as a subject having or at risk for the disease or disorder. In some embodiments, the methods thereby treat, *e.g.*, ameliorate one or more symptom of the disease or disorder, such as by lessening tumor burden in a CD20-expressing cancer.

[00464] Methods for administration of cells for adoptive cell therapy are known, as described, *e.g.*, in US Patent Application Publication No. 2003/0170238; U.S. Pat. No. 4,690,915; Rosenberg, *Nat Rev Clin Oncol.* 8 (10):577-85 (2011); Themeli et al., *Nat Biotechnol.* 31(10): 928-933 (2013); Tsukahara et al., *Biochem Biophys Res Commun* 438(1): 84-9 (2013); and Davila et al., *PLoS ONE* 8(4): e61338 (2013). These methods may be used in connection with the methods and compositions provided herein.

[00465] In some embodiments, the cell therapy (*e.g.*, adoptive T cell therapy) is carried out by autologous transfer, in which the cells are isolated and/or otherwise prepared from the subject who is to receive the cell therapy, or from a sample derived from such a subject. Thus, in some aspects, the cells are derived from a subject in need of a treatment and the cells, following isolation and processing are administered to the same subject. In other embodiments, the cell therapy (*e.g.*, adoptive T cell therapy) is carried out by allogeneic transfer, in which the cells are

isolated and/or otherwise prepared from a subject other than a subject who is to receive or who ultimately receives the cell therapy, *e.g.*, a first subject. In such embodiments, the cells then are administered to a different subject, *e.g.*, a second subject, of the same species. In some embodiments, the first and second subjects are genetically identical. In some embodiments, the first and second subjects are genetically similar. In some embodiments, the second subject expresses the same HLA class or supertype as the first subject.

[00466] In some embodiments, the subject, to whom the cells, cell populations, or compositions are administered is a primate, such as a human. The subject can be male or female and can be any suitable age, including infant, juvenile, adolescent, adult, and geriatric subjects. In some examples, the subject is a validated animal model for disease, adoptive cell therapy, and/or for assessing toxic outcomes.

[00467] The CD20-binding molecules, such as VHHs and chimeric receptors containing the VHHs and cells expressing the same, can be administered by any suitable means, for example, by injection, *e.g.*, intravenous or subcutaneous injections, intraocular injection, periocular injection, subretinal injection, intravitreal injection, trans-septal injection, subcleral injection, intrachoroidal injection, intracameral injection, subconjunctival injection, subconjunctival injection, sub-Tenon's injection, retrobulbar injection, peribulbar injection, or posterior juxtасleral delivery. In some embodiments, they are administered by parenteral, intrapulmonary, and intranasal, and, if desired for local treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration.

[00468] The amount of a prophylactic or therapeutic agent provided herein that will be effective in the prevention and/or treatment of a disease or condition can be determined by standard clinical techniques. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems. For the prevention or treatment of disease, the appropriate dosage of the binding molecule or cell may depend on the type of disease or disorder to be treated, the type of binding molecule, the severity and course of the disease or disorder, whether the therapeutic agent is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the agent, and the discretion of the attending physician. The compositions, molecules and cells are in some embodiments suitably administered to the patient at one time or over a series of treatments.

[00469] For example, depending on the type and severity of the disease, dosages of antibodies may include about 10 ug/kg to 100 mg/kg or more. Multiple doses may be administered intermittently. An initial higher loading dose, followed by one or more lower doses may be administered. In some embodiments, wherein the pharmaceutical composition comprises any one of the single domain antibodies described herein, the pharmaceutical composition is administered at a dosage of about 10 ng/kg up to about 100 mg/kg of body weight of the individual or more

per day, for example, at about 1 mg/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature (*see, e.g.*, U.S. Pat. Nos. 4,657,760; 5,206,344; and 5,225,212).

[00470] In the context of genetically engineered cells containing the binding molecules, in some embodiments, a subject may be administered the range of about one million to about 100 billion cells and/or that amount of cells per kilogram of body weight. In some embodiments, wherein the pharmaceutical composition comprises any one of the engineered immune cells described herein, the pharmaceutical composition is administered at a dosage of at least about any of 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , or 10^9 cells/kg of body weight of the individual. Dosages may vary depending on attributes particular to the disease or disorder and/or patient and/or other treatments.

[00471] In some embodiments, the pharmaceutical composition is administered for a single time. In some embodiments, the pharmaceutical composition is administered for multiple times (such as any of 2, 3, 4, 5, 6, or more times). In some embodiments, the pharmaceutical composition is administered once or multiple times during a dosing cycle. A dosing cycle can be, *e.g.*, 1, 2, 3, 4, 5 or more week(s), or 1, 2, 3, 4, 5, or more month(s). The optimal dosage and treatment regime for a particular patient can be determined by one skilled in the art of medicine by monitoring the patient for signs of disease and adjusting the treatment accordingly.

[00472] In some embodiments, the cells or antibodies are administered as part of a combination treatment, such as simultaneously with or sequentially with, in any order, another therapeutic intervention, such as another antibody or engineered cell or receptor or agent, such as a cytotoxic or therapeutic agent.

[00473] In some embodiments, the cells or antibodies are co-administered with one or more additional therapeutic agents or in connection with another therapeutic intervention, either simultaneously or sequentially in any order. In some contexts, the cells are co-administered with another therapy sufficiently close in time such that the cell populations enhance the effect of one or more additional therapeutic agents, or vice versa. In some embodiments, the cells or antibodies are administered prior to the one or more additional therapeutic agents. In some embodiments, the cells or antibodies are administered after to the one or more additional therapeutic agents.

[00474] In certain embodiments, once the cells are administered to a mammal (*e.g.*, a human), the biological activity of the engineered cell populations and/or antibodies is measured by any of a number of known methods. Parameters to assess include specific binding of an engineered or natural T cell or other immune cell to antigen, *in vivo*, *e.g.*, by imaging, or *ex vivo*, *e.g.*, by ELISA or flow cytometry. In certain embodiments, the ability of the engineered cells to destroy target cells can be measured using any suitable method known in the art, such as cytotoxicity assays described in, for example, Kochenderfer et al., J. Immunotherapy, 32(7): 689-702 (2009), and Herman et al. J. Immunological Methods, 285(1): 25-40 (2004). In certain embodiments, the

biological activity of the cells also can be measured by assaying expression and/or secretion of certain cytokines, such as CD107a, IFN γ , IL-2, and TNF. In some aspects the biological activity is measured by assessing clinical outcome, such as reduction in tumor burden or load.

[00475] In some specific embodiments, provided herein is a method for treating a disease or disorder in a subject comprising administering to the subject a binding molecule comprising a single domain antibody that binds to CD20 as described in Section 5.2 above, including, *e.g.*, those with CDRs in Table 2, those comprising the amino acid sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 or SEQ ID NO: 185; and those comprising an amino acid sequence having at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184, or SEQ ID NO: 185. In some embodiments, the disease or disorder is a CD20 associated disease or disorder. In some embodiments, the disease or disorder is a B cell associated disease or disorder. In some embodiments, the disease or disorder is a B cell malignancy. In some embodiments, the B cell malignancy is a B cell leukemia or B cell lymphoma. In a specific embodiment, the disease or disorder is marginal zone lymphoma (*e.g.*, splenic marginal zone lymphoma). In a specific embodiment, the disease or disorder is diffuse large B cell lymphoma (DLBCL). In another specific embodiment, the disease or disorder is mantle cell lymphoma (MCL). In another specific embodiment, the disease or disorder is primary central nervous system (CNS) lymphoma. In another specific embodiment, the disease or disorder is primary mediastinal B cell lymphoma (PMBL). In another specific embodiment, the disease or disorder is small lymphocytic lymphoma (SLL). In another specific embodiment, the disease or disorder is B cell prolymphocytic leukemia (B-PLL). In another specific embodiment, the disease or disorder is follicular lymphoma (FL). In another specific embodiment, the disease or disorder is burkitt lymphoma. In another specific embodiment, the disease or disorder is primary intraocular lymphoma. In another specific embodiment, the disease or disorder is chronic lymphocytic leukemia (CLL). In another specific embodiment, the disease or disorder is acute lymphoblastic leukemia (ALL). In another specific embodiment, the disease or disorder is hairy cell leukemia (HCL). In another specific embodiment, the disease or disorder is precursor B lymphoblastic

leukemia. In another specific embodiment, the disease or disorder is non-hodgkin lymphoma (NHL). In another specific embodiment, the disease or disorder is high-grade B-cell lymphoma (HGBL). In another specific embodiment, the disease or disorder is multiple myeloma (MM). In other embodiments, the disease or disorder is a relapsed or refractory B cell malignancy, such as relapsed or refractory ALL (R/R ALL). In other embodiments, the disease or disorder is an autoimmune and inflammatory disease, including, *e.g.*, those associated with inappropriate or enhanced B cell numbers and/or activation.

[00476] In other embodiments, provided herein is a method for treating a disease or disorder comprising administering to the subject an engineered immune effector cell (such as T cell) as provided in Section 5.4, including, *e.g.*, the cells comprising a CAR provided in Section 5.3. In some embodiments, the engineered immune cell administered to the subject comprises a CAR comprising a polypeptide comprising: (a) an extracellular antigen binding domain comprising one or more anti-CD20 sdAb(s); (b) a transmembrane domain; and (c) an intracellular signaling domain, wherein the anti-CD20 sdAb is as described in Section 5.2 above, including *e.g.*, those with CDRs in Table 2, those comprising the amino acid sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 or SEQ ID NO: 185; and those comprising an amino acid sequence having at least 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 or SEQ ID NO: 185. In some embodiments, the engineered immune cell administered to the subject comprises a CAR comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, SEQ ID NO: 95, SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 99, SEQ ID NO: 100, SEQ ID NO: 101, SEQ ID NO: 102, SEQ ID NO: 103, SEQ ID NO: 104, SEQ ID NO: 105, SEQ ID NO: 106, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191; or comprising a polypeptide having at least 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID

NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, SEQ ID NO: 95, SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 99, SEQ ID NO: 100, SEQ ID NO: 101, SEQ ID NO: 102, SEQ ID NO: 103, SEQ ID NO: 104, SEQ ID NO: 105, SEQ ID NO: 106, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191. In some embodiments, the disease or disorder is a CD20 associated disease or disorder. In some embodiments, the disease or disorder is a B cell associated disease or disorder. In some embodiments, the disease or disorder is a B cell malignancy. In some embodiments, the B cell malignancy is a B cell leukemia or B cell lymphoma. In a specific embodiment, the disease or disorder is marginal zone lymphoma (*e.g.*, splenic marginal zone lymphoma). In a specific embodiment, the disease or disorder is diffuse large B cell lymphoma (DLBCL). In another specific embodiment, the disease or disorder is mantle cell lymphoma (MCL). In another specific embodiment, the disease or disorder is primary central nervous system (CNS) lymphoma. In another specific embodiment, the disease or disorder is primary mediastinal B cell lymphoma (PMBL). In another specific embodiment, the disease or disorder is small lymphocytic lymphoma (SLL). In another specific embodiment, the disease or disorder is B cell prolymphocytic leukemia (B-PLL). In another specific embodiment, the disease or disorder is follicular lymphoma (FL). In another specific embodiment, the disease or disorder is burkitt lymphoma. In another specific embodiment, the disease or disorder is primary intraocular lymphoma. In another specific embodiment, the disease or disorder is chronic lymphocytic leukemia (CLL). In another specific embodiment, the disease or disorder is acute lymphoblastic leukemia (ALL). In another specific embodiment, the disease or disorder is hairy cell leukemia (HCL). In another specific embodiment, the disease or disorder is precursor B lymphoblastic leukemia. In another specific embodiment, the disease or disorder is non-hodgkin lymphoma (NHL). In another specific embodiment, the disease or disorder is high-grade B-cell lymphoma (HGBL). In another specific embodiment, the disease or disorder is multiple myeloma (MM). In other embodiments, the disease or disorder is a relapsed or refractory B cell malignancy, such as relapsed or refractory ALL (R/R ALL). In other embodiments, the disease or disorder is an autoimmune and inflammatory disease, including, *e.g.*, those associated with inappropriate or enhanced B cell numbers and/or activation.

5.7.2. Diagnostic and Detection Methods and Uses

[00477] In another aspect, provided herein are methods involving use of the binding molecules provided herein, *e.g.*, VHHs that binds CD20 and molecules (such as conjugates and complexes) containing such VHHs, for detection, prognosis, diagnosis, staging, determining binding of a particular treatment to one or more tissues or cell types, and/or informing treatment decisions in a subject, such as by the detection of CD20 and/or the presence of an epitope thereof recognized by

the antibody.

[00478] In some embodiments, an anti-CD20 antibody (such as any one of the anti-CD20 single domain antibodies described herein) for use in a method of diagnosis or detection is provided. In a further aspect, a method of detecting the presence of CD20 in a biological sample is provided. In certain embodiments, the method comprises detecting the presence of CD20 protein in a biological sample. In certain embodiments, CD20 is human CD20. In some embodiments, the methods are diagnostic and/or prognostic methods in association with a CD20-expressing disease or disorder. The methods in some embodiments include incubating and/or probing a biological sample with the antibody and/or administering the antibody to a subject. In certain embodiments, a biological sample includes a cell or tissue or portion thereof, such as tumor or cancer tissue or biopsy or section thereof. In certain embodiments, the contacting is under conditions permissive for binding of the anti-CD20 antibody to CD20 present in the sample. In some embodiments, the methods further include detecting whether a complex is formed between the anti-CD20 antibody and CD20 in the sample, such as detecting the presence or absence or level of such binding. Such a method may be an *in vitro* or *in vivo* method. In one embodiment, an anti-CD20 antibody is used to select subjects eligible for therapy with an anti-CD20 antibody or engineered antigen receptor, *e.g.*, where CD20 is a biomarker for selection of patients.

[00479] In some embodiments, a sample, such as a cell, tissue sample, lysate, composition, or other sample derived therefrom is contacted with the anti-CD20 antibody and binding or formation of a complex between the antibody and the sample (*e.g.*, CD20 in the sample) is determined or detected. When binding in the test sample is demonstrated or detected as compared to a reference cell of the same tissue type, it may indicate the presence of an associated disease or disorder, and/or that a therapeutic containing the antibody will specifically bind to a tissue or cell that is the same as or is of the same type as the tissue or cell or other biological material from which the sample is derived. In some embodiments, the sample is from human tissues and may be from diseased and/or normal tissue, *e.g.*, from a subject having the disease or disorder to be treated and/or from a subject of the same species as such subject but that does not have the disease or disorder to be treated. In some cases, the normal tissue or cell is from a subject having the disease or disorder to be treated but is not itself a diseased cell or tissue, such as a normal tissue from the same or a different organ than a cancer that is present in a given subject.

[00480] Various methods known in the art for detecting specific antibody-antigen binding can be used. Exemplary immunoassays include fluorescence polarization immunoassay (FPIA), fluorescence immunoassay (FIA), enzyme immunoassay (EIA), nephelometric inhibition immunoassay (NIA), enzyme linked immunosorbent assay (ELISA), and radioimmunoassay (RIA). An indicator moiety, or label group, can be used so as to meet the needs of various uses of the method which are often dictated by the availability of assay equipment and compatible

immunoassay procedures. Exemplary labels include radionuclides (*e.g.* ^{125}I , ^{131}I , ^{35}S , ^3H , or ^{32}P and/or chromium (^{51}Cr), cobalt (^{57}Co), fluorine (^{18}F), gadolinium (^{153}Gd , ^{159}Gd), germanium (^{68}Ge), holmium (^{166}Ho), indium (^{115}In , ^{113}In , ^{112}In , ^{111}In), iodine (^{125}I , ^{123}I , ^{121}I), lanthanum (^{140}La), lutetium (^{177}Lu), manganese (^{54}Mn), molybdenum (^{99}Mo), palladium (^{103}Pd), phosphorous (^{32}P), praseodymium (^{142}Pr), promethium (^{149}Pm), rhenium (^{186}Re , ^{188}Re), rhodium (^{105}Rh), ruthenium (^{97}Ru), samarium (^{153}Sm), scandium (^{47}Sc), selenium (^{75}Se), (^{85}Sr), sulphur (^{35}S), technetium (^{99}Tc), thallium (^{201}Tl), tin (^{113}Sn , ^{117}Sn), tritium (^3H), xenon (^{133}Xe), ytterbium (^{169}Yb , ^{175}Yb), yttrium (^{90}Y), enzymes (*e.g.*, alkaline phosphatase, horseradish peroxidase, luciferase, or β -galactosidase), fluorescent moieties or proteins (*e.g.*, fluorescein, rhodamine, phycoerythrin, GFP, or BFP), or luminescent moieties (*e.g.*, QdotTM nanoparticles supplied by the Quantum Dot Corporation, Palo Alto, Calif.). Various general techniques to be used in performing the various immunoassays noted above are known.

[00481] In certain embodiments, labeled antibodies (such as anti-CD20 single domain antibodies) are provided. Labels include, but are not limited to, labels or moieties that are detected directly (such as fluorescent, chromophoric, electron-dense, chemiluminescent, and radioactive labels), as well as moieties, such as enzymes or ligands, that are detected indirectly, *e.g.*, through an enzymatic reaction or molecular interaction. In other embodiments, antibodies are not labeled, and the presence thereof can be detected using a labeled antibody which binds to any of the antibodies.

5.8. Kits and Articles of Manufacture

[00482] Further provided are kits, unit dosages, and articles of manufacture comprising any of the single domain antibodies, the chimeric antigen receptors, or the engineered immune effector cells described herein. In some embodiments, a kit is provided which contains any one of the pharmaceutical compositions described herein and preferably provides instructions for its use.

[00483] The kits of the present application are in suitable packaging. Suitable packaging includes, but is not limited to, vials, bottles, jars, flexible packaging (*e.g.*, sealed Mylar or plastic bags), and the like. Kits may optionally provide additional components such as buffers and interpretative information. The present application thus also provides articles of manufacture, which include vials (such as sealed vials), bottles, jars, flexible packaging, and the like.

[00484] The article of manufacture can comprise a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, *etc.* The containers may be formed from a variety of materials such as glass or plastic. Generally, the container holds a composition which is effective for treating a disease or disorder (such as cancer) described herein, and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The label or package insert indicates that the composition is used for treating the particular

condition in an individual. The label or package insert will further comprise instructions for administering the composition to the individual. The label may indicate directions for reconstitution and/or use. The container holding the pharmaceutical composition may be a multi-use vial, which allows for repeat administrations (e.g. from 2-6 administrations) of the reconstituted formulation. Package insert refers to instructions customarily included in commercial packages of therapeutic products that contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products. Additionally, the article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

[00485] The kits or article of manufacture may include multiple unit doses of the pharmaceutical composition and instructions for use, packaged in quantities sufficient for storage and use in pharmacies, for example, hospital pharmacies and compounding pharmacies.

[00486] For the sake of conciseness, certain abbreviations are used herein. One example is the single letter abbreviation to represent amino acid residues. The amino acids and their corresponding three letter and single letter abbreviations are as follows:

Amino acid	Three letter	One letter	Amino acid	Three letter	One letter
alanine	Ala	(A)	leucine	Leu	(L)
arginine	Arg	(R)	lysine	Lys	(K)
asparagine	Asn	(N)	methionine	Met	(M)
aspartic acid	Asp	(D)	phenylalanine	Phe	(F)
cysteine	Cys	(C)	proline	Pro	(P)
glutamic acid	Glu	(E)	serine	Ser	(S)
glutamine	Gln	(Q)	threonine	Thr	(T)
glycine	Gly	(G)	tryptophan	Trp	(W)
histidine	His	(H)	tyrosine	Tyr	(Y)
isoleucine	Ile	(I)	valine	Val	(V)

[00487] The disclosure is generally disclosed herein using affirmative language to describe the numerous embodiments. The disclosure also specifically includes embodiments in which particular subject matter is excluded, in full or in part, such as substances or materials, method steps and conditions, protocols, procedures, assays or analysis. Thus, even though the disclosure is generally not expressed herein in terms of what the disclosure does not include, aspects that are not expressly included in the disclosure are nevertheless disclosed herein.

[00488] A number of embodiments of the disclosure have been described. Nevertheless, it will

be understood that various modifications may be made without departing from the spirit and scope of the disclosure. Accordingly, the following examples are intended to illustrate but not limit the scope of disclosure described in the claims.

6. EXAMPLES

[00489] The following is a description of various methods and materials used in the studies, and are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present disclosure, and are not intended to limit the scope of what the inventors regard as their disclosure nor are they intended to represent that the experiments below were performed and are all of the experiments that may be performed. It is to be understood that exemplary descriptions written in the present tense were not necessarily performed, but rather that the descriptions can be performed to generate the data and the like associated with the teachings of the present disclosure. Efforts have been made to ensure accuracy with respect to numbers used (*e.g.*, amounts, percentages, *etc.*), but some experimental errors and deviations should be accounted for.

6.1. Example 1—Preparation of Anti-CD20 VHH

[00490] To develop VHH with high binding affinity to CD20 antigen, camels were immunized with human CD20 protein. A phage-display library was then constructed to screen VHH leads. Unique clones were picked based on specific binding and were ranked according to the VHH complementarity determining region (CDR), especially CDR3 which enlarges antigen recognition repertoire and binding.

6.1.1. Cell Line Construction

[00491] Human membrane spanning 4-domains A1 (MS4A1) encoding cDNA (NM_021950.3) was synthesized into plasmid PLVX-puro (Clontech, #632164) by GenScript and the plasmid was named as PLVX-CD20. High-titer recombinant lentiviruses were produced in virus packaging system and were transduced to K562 cells (ATCC, #CRL-24), CHO cells (ATCC, #CCL-61) and Dubca cells (ATCC, #CRL2276). After 3 rounds of selection with puromycin-containing cell culture medium, the surface expression of human CD20 on the transduced cell lines was validated by flow cytometry using PE conjugated anti-human CD20 specific monoclonal antibody (MILTENYI, #130098084). Briefly, 2×10^5 of transduced cells or parental cells were incubated with PE conjugated anti-human CD20 antibody at 4°C for 30 mins, followed by three time washes, and were re-suspended in 200 μ L of DPBS with 0.5% FBS for FACS analysis on Attune NXT flow cytometry (Thermo Fisher) to detect the expression level of human CD20 antigen. The mean fluorescence intensity (MFI) of CHO-CD20 was 1067.44 folds higher than that of un-transduced CHO cells, and the mean fluorescence intensity (MFI) of Dubca-CD20 was 1621.21 folds higher than that of un-transduced Dubca cells.

6.1.2. Animal Immunization and Immune Response Testing

[00492] One adult male alpaca (*Vicugna pacos*) was immunized five times (days 1, 21, 35, 49, and 63) subcutaneously with human CD20 protein. Complete Freund's adjuvant, incomplete Freund's adjuvant and no adjuvant were used for the first, the second to the forth, and the fifth immunization, respectively. A pre-immune 10 mL serum sample and serum sample during the vaccination procedure were collected to monitor the antigen specific humoral response. Binding of alpaca immunoglobulins presented in the diluted pre-immune samples (Pre) and the final immune serum sample (TB) was scored for differential staining of CD20 transfected versus non-transfected cells via flow cytometry. FITC conjugated anti-llama IgG heavy and light chain antibody (BETHYL, #A160100F) was used as the detection antibody. The antibody in the final Test-bleeding serum (TB) showed significant increase of binding to CHO-CD20 cells, starting from 0.16% binding in pre-immunization to 86.8% binding in TB. The data suggested the animal immunization was successful.

[00493] At the same time, one adult male camel (*Camelus bactrian*) was immunized subcutaneously with human CD20 protein for five times with two week intervals. Blood was collected on the pre-immune day (Pre) and final immunization day (TB). Immune response of the camel was evaluated by ELISA, in which the serum samples were tested for binding to immobilized antigen. A series of diluted sera were added to the plate. HRP-conjugated anti-llama IgG secondary antibody (BETHYL, #A160100P) was added as the second antibody for detection. The ELISA results suggested that a robust immune response was induced upon CD20 antigen injection to the animal, with the serum titer reached >1:729 k, indicting the animal immunization was also successful.

[00494] Three to five days after the final immunization, 100 mL of blood was collected from the jugular vein as production bleed. Peripheral blood lymphocytes (PBLs) were isolated from the blood according to the procedure of lymphoprep.

6.1.3. Antibody Phage Library Construction

[00495] Total RNAs were extracted from the isolated lymphocytes according to the manual of TRIzol® Reagent (Thermofisher, #15596026). After quality testing, total RNAs were reverse transcribed into cDNA using oligo(dT)₂₀ primer according to the manual of PrimeScript™ 1st Strand cDNA Synthesis Kit (Takara, #6110A). Nested PCR primers (as described in CN105555310B) were designed for the amplification of VHH fragments, with two *Sfi*I restriction sites introduced. The VHH fragments were amplified using a two-step polymerase chain reaction (PCR), and the PCR products were digested with *Sfi*I and gel purified, and then inserted into phagemid vector-pFL249, which were electro-transferred into *E.coli* cells to generate the phage display VHH immune library.

[00496] A small portion of the transformed cells were diluted and streaked on 2×YT plates supplemented with 100 µg/mL ampicillin. The colonies were counted to calculate the library size.

Positive clones were randomly picked and sequenced to assess the quality of the library. The rest of the transformed cells were streaked onto 245-mm square 2×YT-agar dishes supplemented with 100 µg/mL ampicillin and 2% glucose. Lawns of colonies were scraped off the dishes. A small aliquot of the cells were used for library plasmids isolation. The rest were supplemented with glycerol and stored at -80°C as stock.

6.1.4. Phage-display Panning

[00497] After infection with helper phage, recombinant phage particles which display VHH doamins on the surface as gene III fusion proteins were produced. Phage particles were prepared according to standard methods and stored after filter sterilization at 4°C for further research. Before each round of panning, VHH-displaying phage particles were rescued and amplified by adding helper phage.

[00498] Phage libraries were used with different panning strategies. Based on protein level, in the first and second rounds, biotinylated human CD20 (biotinylating in house according to manufacturer instructions using Sulfo-NHS-LC-Biotin Kit) was incubated with the phage libraries and subsequently captured on Streptavidin Dynabeads (Invitogen). Followed by extensive washing, bound phages were eluted with triethylamine.

[00499] Based on cellular level, in the first selection, CHO-CD20 cells, Dubca-CD20 cells or Raji cells (ATCC, #CCL-86) expressing human CD20 antigen were incubated with the phage libraries. After extensive washing, bound phages were eluted from the cells by resuspension in 1 mg/mL trypsin. In a second selection, phage libraries were incubated with CHO-CD20 cells. After extensive washing, bound phages were eluted as previously described.

6.1.5. ELISA Screening

[00500] Individual library clones were inoculated and induced for expression in 96-deep-well plates. ELISA screening was performed to isolate VHH clones which recognize human CD20 antigen specifically.

[00501] Single clones were randomly picked from the output library, and cultured in a 96-deep-well plate. When the OD600 of bacteria culture reached as 0.6~0.8, IPTG was added to induce the expression for overnight. The bacteria were collected by centrifugation and seeded in a microwell plate.

[00502] Exemplary anti-CD20 VHH domains of the disclosure (*i.e.*, VHH-273, VHH-283, VHH-313, VHH-440, VHH-466, VHH-496, VHH-653, VHH-623, VHH-640 and VHH-657) were selected and sequenced. The CDR (*e.g.*, as defined by Kabat or IMGT numbering scheme) and VHH sequences are summarized in Table 2 and the Sequence Listing provided herein.

6.2. Example 2—Constructions and Immune Cells Expression of VHH Chimeric Receptor Polypeptides

6.2.1. Construction of CD20 VHH CARs

[00503] A nucleic acid sequence encoding a CAR backbone polypeptide comprising from the N-terminus to the C-terminus: a CD8 α hinge domain, a CD8 α transmembrane domain, a CD137 cytoplasmic domain, and a CD3 ζ cytoplasmic domain was chemically synthesized and cloned into a pre-modified lentiviral vector downstream and operably linked to a hEF1 α promoter. Multi-cloning sites (MCS) in the vector allowed insertion of a nucleic acid sequence comprising a Kozak sequence (GCCGCCACC (SEQ ID NO: 160)) operably linked to a nucleic acid sequence encoding a CD8 α signal peptide fused to the N-terminus of VHH fragment(s), and the upstream was operably linked to the CAR backbone sequence.

[00504] To construct a monovalent VHH-based CAR using the CAR backbone vector, the nucleic acid sequence encoding the VHH domain was operably linked to the 3' of the nucleic acid sequence encoding the CD8 α signal peptide. The fusion nucleic acid sequence was chemically synthesized and cloned into the CAR backbone via the *EcoRI* (5' -GAATTC-3' (SEQ ID NO: 161)) and *SpeI* (5' -ACTAGT-3' (SEQ ID NO: 162)) restriction sites by molecular cloning techniques known in the art. Exemplary monospecific and monovalent CD20 VHH CAR constructs are listed in Table 5. Anti-CD20 scFv (Leu16 scFv) (SEQ ID NO: 157) construct was also cloned into the CAR backbone to serve as a positive control and is listed in Table 5 too.

Table 5. Exemplary Monospecific and Monovalent CD20 CAR Constructs

Exemplary CAR Code	Amino Acid Sequence	Signal Peptide	Extracellular antigen binding domain	Hinge & TM	Co-stimulator y signaling domain	Primary intracellular signaling domain
VHH-273 CAR	SEQ ID NO: 81	CD8 α	VHH-273	CD8 α	CD137	CD3 ζ
VHH-283 CAR	SEQ ID NO: 82	CD8 α	VHH-283	CD8 α	CD137	CD3 ζ
VHH-313 CAR	SEQ ID NO: 83	CD8 α	VHH-313	CD8 α	CD137	CD3 ζ
VHH-440 CAR	SEQ ID NO: 84	CD8 α	VHH-440	CD8 α	CD137	CD3 ζ
VHH-466 CAR	SEQ ID NO: 85	CD8 α	VHH-466	CD8 α	CD137	CD3 ζ
VHH-496 CAR	SEQ ID NO: 86	CD8 α	VHH-496	CD8 α	CD137	CD3 ζ
VHH-653 CAR	SEQ ID NO: 87	CD8 α	VHH-653	CD8 α	CD137	CD3 ζ
CD20 scFv CAR	SEQ ID NO: 107	CD8 α	Leu16 scFv	CD8 α	CD137	CD3 ζ
VHH-623 CAR	SEQ ID NO: 189	CD8 α	VHH-623	CD8 α	CD137	CD3 ζ
VHH-640 CAR	SEQ ID NO: 190	CD8 α	VHH-640	CD8 α	CD137	CD3 ζ
VHH-657 CAR	SEQ ID NO: 191	CD8 α	VHH-657	CD8 α	CD137	CD3 ζ

[00505] To construct a multivalent VHH-based CAR, a nucleic acid sequence encoding multiple copies of a VHH, or multiple different VHHS fused to each other via peptide linkers was cloned into the CAR backbone vector. These constructs were prepared by fusing two or three anti-CD20 VHH domains with a series of Glycine-Serine peptide linkers followed by directly synthesizing this fusion sequence in combination with a Kozak-CD8 α signal peptide nucleic acid sequence and cloning into the CAR backbone via *EcoRI* and *SpeI* restriction sites. Exemplary

monospecific and multivalent CD20 VHH CAR constructs are listed in Table 6.

Table 6. Exemplary Monospecific and Multivalent CD20 CAR Constructs

Exemplary CAR Code	Amino Acid	Signal Peptide	Extracellular antigen binding domain					Hinge & TM	CT SD ²	PI SD ³
			VHH #1	linker #1 ¹	VHH #2	linker #2 ¹	VHH #3			
Bi-VHH1 CAR	SEQ ID NO: 98	CD8 α	VHH-273	X2	VHH-496	NA	NA	CD8 α	CD137	CD3 ζ
Bi-VHH2 CAR	SEQ ID NO: 99	CD8 α	VHH-283	X2	VHH-496	NA	NA	CD8 α	CD137	CD3 ζ
Bi-VHH3 CAR	SEQ ID NO: 100	CD8 α	VHH-313	X2	VHH-440	NA	NA	CD8 α	CD137	CD3 ζ
Bi-VHH4 (G4S) ₁ CAR	SEQ ID NO: 101	CD8 α	VHH-273	X1	VHH-653	NA	NA	CD8 α	CD137	CD3 ζ
Bi-VHH4 (G4S) ₂ CAR	SEQ ID NO: 102	CD8 α	VHH-273	X2	VHH-653	NA	NA	CD8 α	CD137	CD3 ζ
Bi-VHH4 (G4S) ₃ CAR	SEQ ID NO: 103	CD8 α	VHH-273	X3	VHH-653	NA	NA	CD8 α	CD137	CD3 ζ
Bi-VHH4 (G4S) ₄ CAR	SEQ ID NO: 104	CD8 α	VHH-273	X4	VHH-653	NA	NA	CD8 α	CD137	CD3 ζ
Bi-VHH4 (G4S) ₅ CAR	SEQ ID NO: 105	CD8 α	VHH-273	X5	VHH-653	NA	NA	CD8 α	CD137	CD3 ζ
Tri-VHH CAR	SEQ ID NO: 106	CD8 α	VHH-273	X2	VHH-496	X2	VHH-440	CD8 α	CD137	CD3 ζ

Note:

¹The linker used in the constructs in Table 6 is (GGGGS)_n, n is an integer including, e.g., 1, 2, 3, 4, 5, and 6 (SEQ ID NO: 138), wherein in linker X1, n=1; in linker X2, n=2; in linker X3, n=3; in linker X4, n=4; and in linker X5, n=5.

²CT SD represents co-stimulatory signaling domain.

³PI SD represents primary intracellular signaling domain.

[00506] The nucleic acid sequences of monovalent and multivalent CD20 VHH CARs described above are shown in SEQ ID NOs: 108-133 and 192-194 in the Sequence Listing. In these exemplary CAR constructs, the sequences of the CD8 α signal peptide, the CD8 α hinge domain, the CD8 α transmembrane domain, the CD137 cytoplasmic domain, and the CD3 ζ cytoplasmic domain are shown in SEQ ID NO: 152, SEQ ID NO: 153, SEQ ID NO: 154, SEQ ID NO: 155, and SEQ ID NO: 156, respectively.

6.2.2. Packaging of Lentivirus Vector

[00507] The lentivirus packaging plasmids mixture containing pMDLg.pRRE (Addgene, #12251), pRSV-REV (Addgene, #12253) and pMD2.G (Addgene, #12259) was pre-mixed with the vectors expressing CAR constructs at a pre-optimized ratio with polyetherimide (PEI). The transfection mixture was then added dropwise to the HEK293T cells and mixed gently, followed by medium replacement post 6~8 hours. The virus-containing supernatants were collected at 48 hours and 72 hours, then centrifuged at 3000 g for 10 mins at 4°C. Post lentivirus concentration, the supernatants were carefully discarded and the virus pellets were re-suspended with D10

medium (DMEM, 10% FBS, 1 mM Sodium Pyruvate and 2 mM L-Glutamine). The harvested virus was aliquoted and stored at -80°C immediately. The virus titer was assessed and determined by CHO mammalian cells transduction efficiency. The LV titers of CD20 mono-/bi-/tri-VHH-based CAR reached within a range of $1 \times 10^8 \sim 4.5 \times 10^8$.

6.2.3. T cell Isolation and Activation

[00508] Human PBMCs were collected from healthy donors. Human T cells were purified from PBMCs using Miltenyi Pan T cell isolation kit (Cat. #130-096-535), according to manufacturer's protocol as described below. The cell number was counted and the cell suspension was centrifuged at 300 g for 10 mins at 4°C. The supernatant was then aspirated off and the cell pellets were re-suspended in 40 µL of the buffer per 10^7 total cells. 10 µL of Pan T Cell Biotin-Antibody Cocktail was added per 10^7 total cells, mixed thoroughly and incubated for 5 mins at 4°C. 30 µL of the buffer was then added per 10^7 total cells. 20 µL of Pan T Cell MicroBead Cocktail was added per 10^7 cells. The cell suspension mixture was mixed thoroughly and incubated for an additional 10 mins at 4°C. A minimum volume (vol.) of 500 µL was required for magnetic separation. In magnetic separation, an LS column was placed in the magnetic field of a suitable MACS Separator. The LS column was rinsed with 3 mL of buffer. The cell suspension was then applied onto the column, and flow-through was collected containing the unlabeled cells which represented the enriched T cell fractions. Additional T cells were collected by washing the column with 3 mL of buffer and collecting unlabeled cells that passed through. These unlabeled cells again represented the enriched T cells and were combined with the flow-through from the previous step. The pooled enriched T cells were then centrifuged and re-suspended with T cell culture medium (RPMI1640, 10% heat-inactivated fetal bovine serum (FBS) and 300 IU/mL of IL-2).

[00509] The isolated T cells were subsequently pre-activated for 48 hours with human T Cell TransAct™ (Miltenyi, Cat. #130-111-160) according to manufacturer's protocol. The freshly isolated T cells were activated by the addition of anti-CD3/CD28 MACSiBead particles (Miltenyi, Cat. #130-111-160) in T cell culture medium according to manufacturer's protocol.

6.2.4. Generation of CD20 VHH CAR-T Cells

[00510] Based on the preliminary results of CD20 CAR-T cells generated by RNA electroporation and screened by *in vitro* assays, mono-, bi- and tri-VHH CAR-T cells were selected, designed and generated by lentivirus transduction for the efficacy analysis in human primary T cells. Additionally, CD20 scFv CAR-T cells were generated as the positive control. Activated T cells were cultured at 0.5×10^6 cells in 0.5 mL medium per well of a 24-well plate. After 24 hours, when T cells were blasting, 0.5 mL of non-concentrated, or smaller volumes of concentrated viral supernatant was added. T cells were transduced at a multiplicity of infection (MOI) of 10 by centrifugation at 1200 g for 1.5 hours at 32°C. The transduced cells were then

transferred to the cell culture incubator for transgene expression under suitable conditions. T cells began to divide in a logarithmic growth pattern, which was monitored by measuring the cell number (viable cells/per mL) and viability (%). The T cell culture was replenished with fresh medium every two days. As the T cells began to rest down after approximately 6~7 days, they were ready to be harvested and cryopreserved for later analysis.

[00511] Before cryopreserving, the percentages of cells transduced (expressing VHH domain or scFv domain on T cell surface) were determined by flow cytometric analysis. The T cells were stained with LIVE/DEAD™ Fixable Dead Cell Stain Kits (Invitrogen, Cat. #L34976), VHH-based CAR-T cells were stained with Goat anti-Llama IgG FITC Conjugate (Bethyl, Cat. #A160-100F), and scFv-based CAR-T cells were stained with FITC-labeled Recombinant Protein L (Acro, Cat. #RPL-PF141) at 4°C for 30 mins, followed by three-time washes, and were re-suspended in 200 µL of DPBS with 0.5% FBS for FACS analysis on a NovoCyt Flow Cytometer (ACEA Biosciences).

[00512] The transduced T cells showed different CAR expression levels with mono-, bi- and tri-VHH domains, indicating the transduction efficiency (percent cells transduced) was related with the size and structure of CAR. Mono-VHH CAR-T cells showed significantly higher transduction efficiency than bi- or tri-VHH CAR-T cells, and were comparable to scFv CAR-T cells. CD20 VHH CAR-T cells expanded about 20~30 folds in a week. The cell counts and viability (82% ~ 96%) of the CD20 VHH CAR-T cell cultures indicated that there was no detectable negative effect of the VHH(s) on the T cells ability to proliferate and expand when compared to the un-transduced T cells (UnT), as shown in Table 7.

Table 7. Viability and Expansion of CD20 CAR-T Cells

Exemplary CAR-T cell Code	CAR Positive Ratio (CAR+ %)	Cell Culture (Day 7)	
		Viability (%)	Expansion fFold
VHH-273 CAR-T cells	47.1	82.3	17.7
VHH-313 CAR-T cells	44.7	81.9	20.6
VHH-466 CAR-T cells	29.8	86.6	32.8
VHH-496 CAR-T cells	42.4	82.6	16.7
Bi-VHH1 CAR-T cells	24.4	95.5	33.3
Bi-VHH2 CAR-T cells	23.3	90.4	28.9
Bi-VHH3 CAR-T cells	26.2	93.9	29.9
Tri-VHH CAR-T cells	6.9	87.1	19.5
CD20 scFv CAR-T cells	41.8	81.7	25.4
UnT	0	83.4	32.7

6.3. Example 3—Characterization of Immune Cells Expressing Chimeric Receptor Polypeptides *In Vitro*

6.3.1. Expression of CD19, CD20 and CD22 Antigens on Target Cell Surface

[00513] To evaluate the expression levels of CD19, CD20 and CD22 on the assessed target cell surface, 5×10^5 of cells per well were incubated with PE-labeled anti-CD19, anti-CD20 and anti-CD22 mAbs, respectively (BioLegend, Cat. #302208, Cat. #302306 & Cat. #302506, respectively), and assessed by flow cytometry with QUANTI-BRITE PE beads (BD Bioscience, Cat. # 340495). The assays and data analysis were performed in accordance with the manufacturer's instructions. "Receptor Number per Cell" indicates the approximate absolute number of molecules per cell on each of the indicated cell lines and is shown in Table 8.

Table 8. CD19, CD20 and CD22 Receptor Number per Target Cell

Cell line	CD19 receptor number per cell	CD20 receptor number per cell	CD22 receptor number per cell
K562.Luc	<10	<10	<50
K562-CD19.Luc	189789	NA	NA
K562-CD20.Luc	<10	28303	NA
K562-CD22.Luc	<10	NA	93339
Raji.Luc	77641	132614	34854
Daudi.Luc	49158	247729	39586
Nalm.6.Luc	32773	<10	10441

6.3.2. Efficacy Evaluation of CD20 VHH CAR-T Cells

[00514] To assess the cytotoxicity of CD20 VHH CAR-T cells against tumor cells, the cells generated as described above were counted and co-cultured with antigen specific cancer cells to read out the killing potency. The control CD20 scFv CAR-T cells were used in all assays to compare assay variation and/or act as an internal control. The un-transduced T cells (UnT) were used as the non-targeting T cells control. CAR-T cell killing assays were conducted towards CD20 positive cell lines - human lymphoma cell line Raji (ATCC, #CCL-86) and K562-CD20; and CD20 negative cell lines - K562-CD19 and K562 (ATCC, #CCL-243). All cell lines were engineered in-house to express firefly luciferase as a reporter for cell viability/killing. The transduced cells were selected with puromycin and refreshed by the selection culture medium (Eagle's Minimum Essential Medium supplemented with 10% FBS and 2 μ g/mL puromycin) in every 2-3 days. Post three rounds of selections, the selected cell clones were harvested and preserved for further use. The cytotoxicity of CD20 VHH CAR-T cells was measured at the effector to target cell ratio (E:T) of 15:1, 10:1 or 5:1 for 24 hours. Assays were initiated by mixing the respective number of T cells with a constant number of target cells. The remaining luciferase activity per well was assessed by ONE-Glo luciferase assay (Promega, Cat. #E6110), to quantify the remaining viable target cells per well.

[00515] Hundreds of monovalent VHH CAR-T cells targeting CD20 were constructed and screened by *in vitro* cytotoxicity assay. Exemplary data are shown in **FIGs.1A-1B** and **FIGs.2A-2F**. The T cells were stained with LIVE/DEAD™ Fixable Dead Cell Stain Kits (Invitrogen, Cat. #L34976), VHH-based CAR-T cells were stained with Goat anti-Llama IgG FITC Conjugate

(Bethyl, Cat. #A160-100F), and scFv-based CAR-T cells were stained with FITC-labeled Recombinant Protein L (Acro, Cat. #RPL-PF141). CAR⁺ expression level (%) of the exemplary VHH CAR-T cells reached around 30% at the harvest day (**FIG.1A**). All of the exemplary CD20 VHH CAR-T cells exhibited higher cytotoxicity than positive control (*i.e.*, CD20 scFv CAR-T cells) against Raji.Luc cells (E:T=10:1) (**FIG.2A, and FIGs.2E-2F**), and were comparable to CD20 scFv CAR-T cells against K562-CD20.Luc cells (**FIG.2B, and FIGs.2E-2F**). No significant cytotoxicity effect was detected against CD20 negative cell lines by the CD20 VHH CAR-T cells as compared to UnT control (**FIGs.2C-2D**).

[00516] To assess the potency of multivalent VHH CAR-T cells, mono-, bi- and tri-VHH CAR-T cells were constructed and generated, and the CAR-T cells' cytotoxicity were assessed. The exemplary data showed that multivalent CD20 VHH CAR-T cells exhibit different levels of cytotoxicity against Raji.Luc and K562-CD20.Luc cells (**FIGs.3A-3B**). In the 24 hours of *in vitro* killing assay, mono-VHH CAR-T cells and bi-VHH CAR-T cells showed higher killing than Tri-VHH CAR-T cells against Raji.Luc cells. None of the CD20 VHH CAR-T cells showed cytotoxicity effect against CD20-negative cell lines, as compared to UnT control (**FIGs.3C-3D**).

[00517] To assess the effect of (G4S) linker(s) between the VHHs, two exemplary VHHs linked with (G4S)_n (n=1~5) (SEQ ID NO: 138) peptides were constructed and five of the Bi-VHH4 CAR-T cells were generated (Table 6). The exemplary data showed that the shorter the (G4S) linker between two VHH domains, the more potent the bi-VHH CAR-T cells were in killing target cells- Raji.Luc cells, and the bi-VHH CAR-T cells with the linker as (G4S)₁ showed the highest cytotoxicity against Raji.Luc at E:T ratios of 20:1 and 10:1 (**FIGs.4A-4B**).

[00518] The observation above indicates that the CD20 VHH CARs provided herein induce T cell activation via specific recognizing CD20 receptor expressing cells, activate T cell endogenous signaling pathway, induce activation of cytotoxic T lymphocytes (CTL) and enhance anti-tumor responses.

6.3.3. IFN- γ Release Evaluation of CD20 VHH CAR-T Cells

[00519] To measure cytokine production of CD20 VHH CAR-T cells in response to CD20-expressing cells, CAR-T cells were co-cultured with the CD20 positive cell lines - Raji.Luc or K562-CD20.Luc and CD20 negative cell line - K562.Luc at an E:T ratio of 15:1, 10:1 or 5:1 for 24 hours, after which the media was harvested for the cytokine analysis using the human IFN- γ kit (Cisbio, Cat. #62HIFNGPEG) for cytokine quantification, and the absorbance of each well (triplicate per teste article) was read by a multimode microplate reader (Tecan Spark).

[00520] The data showed that the exemplary cells VHH-273 CAR-T cells and VHH-496 CAR-T cells produced less IFN- γ than CD20 scFv CAR-T cells when co-cultured with Raji.Luc cells (**FIG.5A**), and produced comparable IFN- γ to CD20 scFv CAR-T cells when co-cultured with K562-CD20.Luc cells (**FIG.5B**). Mono-VHH CAR-T cells, bi-VHH CAR-T cells and tri-VHH

CAR-T cells showed different levels of IFN- γ release against Raji.Luc cells (**FIG.6**). In contrast, IFN- γ release was not detectable or extremely low in cultures containing either un-transduced T cells or negative control cells - K562.Luc, confirming that CD20-specificity of the CAR-T cells was required for reactivity to CD20 expressing cells (**FIG.6**).

6.3.4. CD20 VHH CAR-T Cells Proliferation Capacity

[00521] To assess the proliferation capacity of CD20 VHH CAR-T cells, T cells were stimulated with irradiated K562-CD20 cells or K562 cells at an E:T ratio of 2:1 and co-cultured for 40 days (triplicates per test article). In every four days, T cells were counted and stained with LIVE/DEAD™ Fixable Dead Cell Stain Kit (Invitrogen, Cat. #L34976), CD3+ and CAR+ were assessed by FACS. CD3 receptors were stained with FITC-anti-human CD3 (Biolegend, Cat. #300306). VHH-based CAR-T cells were stained with Goat anti-Llama IgG FITC Conjugate (Bethyl, Cat. #A160-100F), and scFv-based CAR-T cells were stained with FITC-labeled Recombinant Protein L (ACRO, Cat. #RPL-PF141) at 4°C, for 30 mins, followed by three-time washes, and were re-suspended in 200 μ L of DPBS with 0.5% FBS for FACS analysis on a NovoCyte Flow Cytometer (ACEA Biosciences). The FACS data was analyzed by Novoexpress software.

[00522] In the expansion stimulated by CD20 specific antigen (co-cultured with K562-CD20 cells), VHH-273 CAR-T cells showed dramatically enhanced CD3+ T cells and CAR+ T cells proliferation, compared to CD20 scFv CAR-T cells from Day 20 to Day 40, and showed around 1.6 ~ 2 folds higher than CD20 scFv CAR-T cells in CD3+ or CAR+ T cells proliferation from Day 28 to Day 40 (**FIGs.7A-7B**). Enhanced CD3+ T cells or CAR+ T cells proliferation was not observed in co-cultures containing negative control cells - K562 (**FIGs.7A-7B**). Both of CD20 VHH CAR-T cells and CD20 scFv CAR-T cells demonstrated that the percentages of CAR+ expression were increased in a time-dependent manner in the first 20 days and plateaued post 20 days (**FIG.7C**). In addition, CD20 VHH CAR-T cells exhibited significantly higher percentage of CAR+ expression than CD20 scFv CAR-T cells (**FIG.7C**). CD20 antigen stimulated CD20 VHH CAR-T cells or CD20 scFv CAR-T cells maintained significantly higher viability in 40-days proliferation assay than the respective non-stimulated CD20 VHH CAR-T cells or CD20 scFv CAR-T cells (**FIG.7D**). The data indicates that CD20 VHH CAR-T cells induce robust and persistent T cells activation and proliferation in a tumor associated antigen dependent manner.

6.3.5. CD20 VHH CAR-T Cells Exhaustion Markers and Memory Markers Assessment

[00523] To assess the exhaustion markers and memory markers of CD20 VHH CAR-T cells, T cells were stimulated with irradiated K562-CD20 cells at an E:T ratio of 2:1 and co-cultured for 28 days (triplicates per test article). In every four days, T cells were counted with a live/dead staining, and CAR+, T cell exhaustion markers and T cell memory markers were assessed by a

multi-color FACS staining. T cell exhaustion markers were co-stained with fluorochrome conjugated anti-human CD279 (PD-1), anti-human CD223 (LAG-3) and anti-human CD366 (Tim-3). T cell memory markers were co-stained with fluorochrome conjugated anti-human CD45RA and anti-human CD197 (CCR7). FACS analysis was ran with a BD LSRFortessa Flow Cytometer (BD Biosciences).

[00524] The FACS data of exhaustion markers assessment showed that the exhaustion markers of both VHH-273 CAR-T cells and CD20 scFv CAR-T cells were peaked at Day 4 post antigen specific stimulation and dropped dramatically afterward. Post Day 8, the expression levels of exhaustion markers of both CD20 VHH CAR-T cells and CD20 scFv CAR-T cells were kept low till Day 28 (**FIG.8**). The data suggests that T cell exhaustion is driven by strong T cell activation only at early stage when exposed to tumor associated antigen and normally the expression returns to baseline after activation.

[00525] The FACS data of memory markers assessment showed that VHH-273 CAR-T cells and CD20 scFv CAR-T cells demonstrated similar time-dependent pattern of central memory T cells (T_{CM}) and effector memory T cells (T_{EM}) post antigen-specific stimulation. In addition, CAR-T cells initially differentiated into T_{CM}, which peaked at Day 4, then gradually differentiated to T_{EM} or T_{EMRA}, which peaked at Day 8 or Day 16 respectively (**FIGs.9A-9C**). T_{CM} and T_{EM}/T_{EMRA} of UnT were maintained at the baseline (**FIGs.9A-9C**). The engineered second generation CAR-T cells direct T cells to their fates by balancing effector and memory programs.

6.4. Example 4—*In Vivo* Efficacy of CD20 VHH CAR-T Cells in Tumor

Xenograft Mice

[00526] Anti-tumor activity of CD20 VHH CAR-T cells was assessed *in vivo* in a Raji xenograft NCG mouse model and CD20 scFv CAR-T cells and un-transduced cells (UnT) were evaluated as either positive or negative control.

[00527] Cell line: Raji (ATCC #CCL-86) is the lymphoblast-like cell line, established by Pulvertaft in 1963 from a Burkitt's lymphoma of the 11-year-old male. Raji cells were grown in RPMI medium containing 10% fetal bovine serum. This cell line grows in suspension in tissue culture flasks. This cell line persists and expands in mice when implanted intravenously. The Raji cells had been modified to express luciferase, so that tumor cell growth could also be monitored by imaging the mice. The Raji model endogenously expresses high levels of CD19, CD20 and CD22 and thus, can be used to test the *in vivo* efficacy of CD20-directed engineered T cells.

[00528] Mice: 5-6 weeks old NCG (NOD-Prkdcem26Cd52Il2rgem26Cd22/Nju) female mice were received from Model Animal Research Center of Nanjing University, with similar weight (around 20 g). Animals were allowed to acclimate in the animal facility for 7 days prior to experimentation. Animals were handled in accordance with ACUC regulations and guidelines.

6.4.1. CD20 Mono-VHH CAR-T Cells Efficacy and Re-challenging *In Vivo* Tests

6.4.1.1. CD20 Mono-VHH CAR-T Cells Efficacy *In Vivo* Test

[00529] To create the tumor xenograft, NCG Mice were injected intravenously with Raji.Luc. The mice were treated with the T cells post 4 days with Raji.Luc tumor cell implantation. The mice were injected intravenously via the tail vein with 400 μ L of the T cells for a dose of 1×10^6 T cells per mouse. The four mice per group were treated with either CD20 mono-VHH CAR-T cells (VHH-273 CAR-T cells or VHH-496 CAR-T cells) or CD20 scFv CAR-T cells, and 400 μ L of HBSS alone and un-transduced T cells (UnT) were used as controls. All the T cells were prepared from the same donor in parallel. Animal health status was monitored twice per week, including body weight measurement. Tumors were monitored weekly by bioluminescence imaging (BLI) until animals achieved endpoint.

[00530] The mean bioluminescence for all treatment groups is plotted in **FIGs.10A-10F**. The HBSS treatment group (vehicle), which did not receive any T cells, demonstrated baseline Raji tumor growth kinetics in intravenously injected NCG mice. The UnT treatment group received non-transduced T cells as negative control for the engineered T cells. Both the HBSS and the UnT treatment groups demonstrated continuous aggressive tumor progression throughout this study and were euthanized on Day 16. CD20 mono-VHH CAR-T cells significantly inhibited tumor growth when compared to UnT treatment and showed complete tumor inhibition during the 35 days of the *in vivo* efficacy study, as indicated from the mean bioluminescence and the image of bioluminescence (**FIGs.10A-10B**). In addition, the twice per week monitored mice health status was normal and the body weights were increased in CD20 mono-VHH CAR-T cells treatment groups throughout the 35 days of *in vivo* study (**FIG.10C**).

6.4.1.2. PK Study - Detection Copies of CARs in Mouse Peripheral Blood

[00531] To assess pharmacokinetics of CD20 VHH CAR-T cells and to monitor circulating CAR-T cells proliferation and persistence in mouse, the genomic DNA of mouse peripheral blood was extracted with QIAamp-DNA-mini Blood Kit (Qiagen, Cat. #51106) and the concentration was measured by Nanodrop One (ThermoFisher, Cat. #ND-ONE-W). The genomic DNA (of 3 mice per group) was diluted at the concentration of 50 ng/ μ L with ddH₂O and mixed with PCR reaction reagents, including forward and reverse primer sets binding onto CAR backbone. The mixture was then assessed with digital PCR and ran on the QX200 Droplet Reader (Bio Rad, Cat. #1864003). The copies/ng genomic DNA of VHH-273 CAR was 2 folds higher than that of CD20 scFv CAR observed at peak level (Day 14) (**FIG.10D**).

6.4.1.3. Raji.Luc Tumor Cells Re-challenging *In Vivo* Study

[00532] To investigate CAR-T cells efficacy on those at least 5 weeks tumor-free mice under

the additional tumor burdens, those mice with undetectable CAR copy number (copies/ng of genomic DNA) in PBMC were selected for re-challenging assay. The selected mice were injected intravenously with half amount of Raji.Luc tumor cells. Animal health status was monitored twice per week, including body weight measurement. The Raji tumors were monitored weekly by bioluminescence imaging (BLI) until animals achieved endpoint.

[00533] The naïve mice (not implanted with Raji.Luc cells or treated with CAR-T cells previously) injected with Raji.Luc as control demonstrated continuous aggressive tumor progression and were euthanized on Day 19. VHH-273 CAR-T cells showed complete tumor inhibition for 28 days post re-challenging. Both CD20 VHH CAR-T cells and CD20 scFv CAR-T cells treatment groups relapsed around Day 12, as indicated from the mean bioluminescence (**FIG.10E**). In addition, the twice per week monitored mice health status and body weight under VHH-273 CAR-T cells treatment were normal throughout the 28 days re-challenging *in vivo* study (**FIG.10F**).

6.4.2. CD20 Mono-VHH, Bi-VHH and Tri-VHH CAR-T Cells Efficacy and Re-challenging *In Vivo* Tests

6.4.2.1. CD20 Mono-VHH, Bi-VHH and Tri-VHH CAR-T Cells

Efficacy *In Vivo* Test

[00534] To create the tumor xenograft, NCG Mice were injected intravenously with Raji.Luc. The mice were treated with the T cells post 4 days with Raji.Luc tumor cell implantation. The mice were injected intravenously via the tail vein with 400 μ L of the T cells for a dose of 1×10^6 T cells per mouse. The six mice per group were treated with CD20 mono-VHH CAR-T cells (VHH-313 CAR-T cells), bi-VHH CAR-T cells (Bi-VHH3 CAR-T cells) or Tri-VHH CAR-T cells, and CD20 scFv CAR-T cells, 400 μ L of HBSS alone and un-transduced T cells (UnT) were used as the controls. All the T cells were prepared from the same donor in parallel. Animal health status was monitored twice per week, including body weight measurement. Tumors were monitored weekly by bioluminescence imaging (BLI) until animals achieved endpoint.

[00535] The HBSS treatment group (vehicle), which did not receive any T cells, demonstrated baseline Raji tumor growth kinetics in intravenously injected NCG mice. The UnT treatment group received non-transduced T cells as negative control for the engineered T cells. Both the HBSS and the UnT treatment groups demonstrated continuous aggressive tumor progression throughout this study and were euthanized on Day 14. VHH-313 CAR-T cells, Bi-VHH3 CAR-T cells and Tri-VHH CAR-T cells all significantly inhibited tumor growth when compared to UnT treatment and showed complete tumor inhibition during 42 days of the *in vivo* efficacy study, as indicated from the mean bioluminescence and the image of bioluminescence (**FIGs.11A-11B**). Tri-VHH CAR-T cells were slightly more efficacious than mono-VHH CAR-T cells and bi-VHH CAR-T cells in Raji tumor inhibition. CD20 scFv CAR-T cells treatment group relapsed at Day

28 (FIGs.11A-11B). In addition, the twice per week monitored mice health status was normal and body weights were increased in CD20 mono-VHH, bi-VHH and tri-VHH CAR-T cells treatment groups throughout the *in vivo* study (FIG.11C).

6.4.2.2. PK Study - Detection Copies of CARs in Mouse Peripheral Blood

[00536] To assess pharmacokinetics of CD20 mono-VHH, bi-VHH and tri-VHH CAR-T cells and to monitor circulating CAR-T cells proliferation and persistence in mouse, the genomic DNA of mouse peripheral blood was extracted with QIAamp-DNA-mini Blood Kit (Qiagen, Cat. #51106) and the concentration was measured by Nanodrop One (ThermoFisher, Cat. #ND-ONE-W). The genomic DNA (of 3 mice per group) was diluted at the concentration of 50 ng/μL with ddH₂O and mixed with PCR reaction reagents, including forward and reverse primer sets binding onto CAR backbone. The mixture was then assessed with digital PCR and ran on the QX200 Droplet Reader (Bio Rad, Cat. #1864003). Both VHH-313 CAR copies and Bi-VHH3 CAR copies peaked at Day 7 and CD20 scFv CAR copies peaked at Day 21 (FIG.11D, which is adopted from FIG.11E and the scale of Y-axis is adjusted to enhance the resolution of copies/ng genomic DNA). In addition, the Tri-VHH CAR-T cells demonstrated higher expansion capability and peaked at Day 21, and then gradually dropped till the endpoint (Day 35) (FIG.11E).

6.4.2.3. Raji.Luc Tumor Cells Re-challenging *In Vivo* Study

[00537] To investigate CD20 mono-VHH, bi-VHH and tri-VHH CAR-T cells efficacy on those at least 6 weeks tumor-free mice under the additional tumor burdens, the mice were selected for re-challenging assay. The selected mice were injected intravenously with half amount of Raji.Luc cells. Animal health status was monitored weekly, including body weight measurement. The Raji tumors were monitored weekly by bioluminescence imaging (BLI) until animals achieved endpoint.

[00538] CD20 scFv CAR-T cells treatment group showed continued remissions and relapses, VHH-313 CAR-T cells or Bi-VHH3 CAR-T cells treated mice relapsed around Day 11 to Day 18, and Tri-VHH CAR-T cells treated mice showed significant tumor inhibition for 32 days post Raji.Luc cells re-challenging, as indicated by the mean bioluminescence (FIG.11F).

6.5. Example 5—Characterization of Anti-CD20 VHH-huIgG1Fc Monoclonal Antibody (mAb) Binding and On-/off-target Activity *In Vitro*

6.5.1. Anti-CD20 VHH-huIgG1Fc mAb On-target Binding to CD20

Receptor Expressing Cells and EC₅₀

[00539] Each of three exemplary of anti-CD20 VHH sequences (clone ID: VHH-273, VHH-466 and VHH-496) with human IgG1 Fc fragment sequence (SEQ ID NO: 159) were cloned into a mammalian expression vector – pcDNA3.4, to facilitate the recombinant protein expression. The DNA codons were further optimized for optimal expression in mammalian host

cell – Expi293F. The antibodies were harvested from the supernatant of cell culture, one-step purified by MabSelect SuRe LX and sterilized via a 0.2 µm filter. The purified antibody concentrations were determined by A280 and reached 2~6 mg/mL with ~90% purity. Anti-CD20 scFv-huIgG1Fc mAb with amino acid sequence of Rituximab (C2B8) scFv shown in SEQ ID NO: 158 and anti-CD20 Fab-huIgG1Fc mAb with amino acid sequences of Leu16 VH-CH1 and VL-CL shown in SEQ ID NO: 163 and SEQ ID NO: 164 were used as positive controls for the cell surface binding assay. CD20 positive cell lines - Raji, K562-CD20 and negative cell line - K562 were re-suspended in complete culture medium, cells concentration was diluted to 1×10^6 cells/mL and the staining was performed in 2×10^5 cells per well. The mAbs were serially diluted from maximal concentration of 2000 nM (3-fold reduction, 12 pts) and added according to the experimental plan and protocol. The mAbs and cells were co-incubated for 1 hour at 4°C. Then, the cells were washed with 200 µL of DPBS + 0.5% FBS and spun at 300 g for 5 mins, 4°C. The cells were stained with the detection antibody - PE-conjugated mouse anti-human IgG1 Fc (BioLegend, Cat. # 409304, 1:100) for 40 mins at 4°C. The cells were then washed again and re-suspended with 200 µL of DPBS + 0.5% FBS for flow cytometric analysis on a NovoCyte Flow Cytometer (ACEA Biosciences). The FACS data was analyzed by Novoexpress software, and the MFI (median fluorescent intensity) was analyzed by GraphPad PRISM version 6.0.

[00540] The cell surface binding data showed that anti-CD20 VHH-huIgG1Fc mAbs specifically bound to CD20 positive cells - Raji cells and K562-CD20 cells in a dose-dependent manner, and significantly stronger binding than anti-CD20 scFv-huIgG1Fc mAb on both target cells (**FIGs.12A-12B**). Anti-CD20 VHH-huIgG1Fc mAbs showed a comparable binding to anti-CD20 Fab-huIgG1Fc mAb on target cells and stronger binding to Raji at higher mAb concentration (**FIG.12D**). None of the antibodies showed significant binding to CD20 negative cell line - K562 (**FIG.12C and FIG.12E**). EC₅₀ of anti-CD20 VHH-huIgG1Fc mAbs vs. positive control mAbs are shown in Table 9 and Table 10. The term “EC₅₀”, also known as half maximal effective concentration, refers to the concentration of an antibody which induces a response halfway between the baseline and maximum after a specified exposure time.

Table 9. EC₅₀ of Anti-CD20 VHH-huIgG1Fc mAbs and Anti-CD20 scFv-huIgG1Fc mAb Binding with Raji or K562-CD20 Cells

mAb	EC ₅₀ (nM) On-target Raji cells	EC ₅₀ (nM) On-target K562-CD20 cells
VHH-273-huIgG1Fc mAb	9.8	4.9
VHH-466-huIgG1Fc mAb	8.9	4.7
VHH-496-huIgG1Fc mAb	13.5	5.1
VHH-440-huIgG1Fc mAb	11.12	6.24
Anti-CD20 scFv-huIgG1Fc mAb	75.3	41.1

Table 10. EC₅₀ of Anti-CD20 VHH-huIgG1Fc mAbs and Anti-CD20 Fab-huIgG1Fc mAb Binding with Raji Cells

mAb	EC ₅₀ (nM) On-target Raji cells
VHH-273-huIgG1Fc mAb	14.28
VHH-466-huIgG1Fc mAb	9.12
VHH-496-huIgG1Fc mAb	16.00
Anti-CD20 Fab-huIgG1Fc mAb	13.51

6.5.2. Anti-CD20 VHH-huIgG1Fc mAb Off-target Binding Assessment

[00541] To validate off-target binding, anti-CD20 VHH-huIgG1Fc mAbs were assessed with various human cell lines. The exemplary tested cell lines are listed in Table 11. The mAbs were incubated with 1×10^5 cells per well for 1 hour at 4°C. Then, the cells were washed with 200 μ L of DPBS and 0.5% FBS and spun at 300 g for 5 mins at 4°C. The cells were stained with the detection antibody - PE-conjugated mouse anti-human IgG1 Fc (BioLegend, Cat. # 409304, 1:100) for 30 mins at 4°C. The cells were then washed again and re-suspended with 200 μ L of DPBS + 0.5% FBS for flow cytometric analysis on a NovoCyt Flow Cytometer (ACEA Biosciences) and the FACS histogram data was analyzed with Novoexpress software. As for the above three exemplary anti-CD20 VHH-huIgG1Fc mAbs (VHH clone ID: VHH-273, VHH-466 and VHH-496), non-specific binding to off-target cells was not observed at the concentration yielding maximum binding to Raji cell (Table 11).

Table 11. Anti-CD20 VHH-huIgG1Fc mAbs Binding to Off-target Cell Lines

Cell Line	ATCC ID	Primary Site	CD20 expression level	Anti-CD20 VHH-huIgG1Fc mAb Binding
Raji	CCL-86	Lymphoid	+	+
Daudi	CCL-213	Lymphoid	+	+
Nalm.6	CRL-3273	Lymphoid	-	-
RPMI-8226	CCL-155	Lymphoid	very low	very weak
Jurkat	TIB-152	Lymphoid	-	-
K562	CCL-243	Bone marrow	-	-
HL-60	CCL-240	Peripheral blood	-	-
THP-1	TIB-85	Peripheral blood	-	-
U87-MG	HTB-14	Brain/Neuron	-	-
IMR-32	CCL-127	Brain/Neuron	-	-
FaDu	HTB-43	Pharynx	-	-
A-253	HTB-41	Salivary gland	-	-
SK-BR-3	HTB-30	Breast	-	-
A549	CCL-185	Lung	-	-
NCI-H446	HTB-171	Lung	-	-
HepG2	HB-8065	Liver	-	-
NCI-N87	CRL-5822	Stomach	-	-
HEK-293T	632180	Kidney	-	-
PANC-1	CRL-1469	Pancreas	-	-
HCT 116	CCL-247	Colorectal	-	-

SK-OV-3	HTB-77	Ovary	-	-
Hela	CCL-2	Cervix	-	-
BeWo	CCL-98	Placenta	-	-
A375	CRL-1619	Skin	-	-
U-2 OS	HTB-96	Bone	-	-

6.6. Example 6—Generation and Characterization of Humanized CD20 VHH CARs

6.6.1. Humanization of Anti-CD20 VHH Antibodies

[00542] To reduce the immunogenicity in human, camelid VHH antibodies were humanized, since much of the immune response occurs against the non-human antibody constant region. When different framework regions are combined with the camelid CDRs, chimeric human and camelid antibodies specific for the same antigen can elicit different effector functions, extending their therapeutic benefits. Mono-specific camelid VHHs were humanized by using sequence-based approaches and framework shuffling to most homologous human germline sequence or related scaffold. The non-compatibility of camelid CDRs being supported by non-native human framework scaffold and elimination of key conformational residues were resolved by *in silico* CDR-grafting, homology structural modeling (tertiary conformation & fold), sequence alignment, structure based-back mutation design and reintroduction of key conformational residues from camelid VHH antibody. The antibody humanization process may not only eliminate steric clashes but also restore function in relation to binding affinity to its antigen.

[00543] Universal humanized VHH framework h-NbBcII10FGLA (Protein Data Bank, PDB code: 3EAK, <https://www.rcsb.org/structure/3EAK>) designed by Cécile Vincke et al. was adopted for humanization design based on sequence homology. The homologous modeling of camelid VHH was performed using the modeling software MODELLER. According to alignment with human germline gene, IGHV3-64*04 was chosen as one human acceptor for anti-CD20 VHH domains. Relative solvent accessibility of the amino acids was calculated according to the three-dimensional structure of the protein. If one of the amino acids of VHH was exposed to a solvent, it would be replaced with the original amino acid. The exemplary humanized VHH domains (*e.g.*, huVHH-253, huVHH-256, huVHH-260, huVHH-746, huVHH-750, huVHH-753, huVHH-836, huVHH-840, huVHH-843, huVHH-846, 2082H1, 2082H2, 2082H3, 2082H4, 2082H5 and 2082H6) generated herein are shown in Table 2, and the corresponding sequences are provided in the Sequence Listing provided herein.

6.6.2. Characterization of Humanized CD20 VHH CAR-T Cells

[00544] Exemplary humanized CD20 VHH CARs (see Table 12) were generated using the method described above.

Table 12. Exemplary Humanized CD20 VHH CAR Constructs

Exemplary humanized CAR Code	Amino Acid Sequence	Signal Peptide	Extracellular antigen binding domain	Hinge & TM	Co-stimulatory signaling domain	Primary intracellular signaling domain
huVHH-253 CAR	SEQ ID NO: 88	CD8 α	huVHH-253	CD8 α	CD137	CD3 ζ
huVHH-256 CAR	SEQ ID NO: 89	CD8 α	huVHH-256	CD8 α	CD137	CD3 ζ
huVHH-260 CAR	SEQ ID NO: 90	CD8 α	huVHH-260	CD8 α	CD137	CD3 ζ
huVHH-746 CAR	SEQ ID NO: 91	CD8 α	huVHH-746	CD8 α	CD137	CD3 ζ
huVHH-750 CAR	SEQ ID NO: 92	CD8 α	huVHH-750	CD8 α	CD137	CD3 ζ
huVHH-753 CAR	SEQ ID NO: 93	CD8 α	huVHH-753	CD8 α	CD137	CD3 ζ
huVHH-836 CAR	SEQ ID NO: 94	CD8 α	huVHH-836	CD8 α	CD137	CD3 ζ
huVHH-840 CAR	SEQ ID NO: 95	CD8 α	huVHH-840	CD8 α	CD137	CD3 ζ
huVHH-843 CAR	SEQ ID NO: 96	CD8 α	huVHH-843	CD8 α	CD137	CD3 ζ
huVHH-846 CAR	SEQ ID NO: 97	CD8 α	huVHH-846	CD8 α	CD137	CD3 ζ

[00545] The humanized CD20 VHH CAR-T cells were generated by lentivirus transduction in human primary T cells and were assessed by *in vitro* efficacy study according to the standard method. The transduced T cells showed different CAR expression levels (%) with humanized CD20 VHH CAR. Exemplary humanized CD20 VHH CAR-T cells' viability was about 90%~96%, CAR+ (%) was about 12%~43% and the expansion folds were within 35~74 in a 7-day culture, indicating that there was no detectable negative effect of humanized VHH on the T cells' capability to proliferate and expand when compared to the un-transduced T cells (UnT).

[00546] Humanized CD20 VHH CAR-T cells generated as described above were counted and co-cultured with antigen specific cancer cell lines to assess the killing potency, parental camelid CD20 VHH CAR-T cells and CD20 scFv CAR-T cells were used as control and the un-transduced T cells (UnT) were used as non-targeting T cells control. Humanized CD20 VHH CAR-T cell killing assays were conducted towards CD20 positive cell lines - Raji (ATCC #CCL-86), K562-CD20 and negative cell line - K562 (ATCC #CCL-243). All the cell lines were engineered in-house to express firefly luciferase as a reporter for cell viability/killing. The transduced cells were selected with puromycin and refreshed by the selection culture medium (Eagle's Minimum Essential Medium supplemented with 10% FBS and 2 μ g/mL of puromycin) in every 2-3 days. Post three rounds of selection, the selected cell clones were harvested and preserved for further use. The cytotoxicity of humanized CD20 VHH CAR-T cells was measured at an effector cells to target cells ratio (E:T) of 15:1, 10:1, 5:1 or 2.5:1 for 24 hours. Assays were initiated by mixing the respective number of T cells with a constant number of target cells. The

remaining luciferase activity per well was assessed by ONE-Glo luciferase assay (Promega, Cat. #E6110) to quantify the remaining viable target cells per well.

[00547] The results showed that the humanized CD20 VHH CAR-T cells (including huVHH-746 CAR-T cells, huVHH-750 CAR-T cells, huVHH-753 CAR-T cells, huVHH-253 CAR-T cells, huVHH-256 CAR-T cells, and huVHH-260 CAR-T cells) induced lysis of antigen specific target cells and exhibited higher potency or maintained potency against the target cells (**FIGs.13A-13B, FIGs.13D-13E and FIGs.13H-13I**). No significant cytotoxicity effect was detected against negative cell line - K562.Luc or Nalm.6.Luc by humanized CD20 VHH CAR-T cells as compared to UnT control (**FIG.13C, FIGs.13F-13G and FIGs.13J-13K**).

[00548] The teachings of all patents, published applications and references cited herein are incorporated by reference in their entirety.

[00549] While example embodiments have been particularly shown and described, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the embodiments encompassed by the appended claims.

[00550] From the foregoing, it will be appreciated that, although specific embodiments have been described herein for the purpose of illustration, various modifications may be made without deviating from the spirit and scope of what is provided herein. All of the references referred to above are incorporated herein by reference in their entireties.

WHAT IS CLAIMED:

1. An anti-CD20 single domain antibody (sdAb) comprising:
 - (i) a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 2; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3;
 - (ii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 5; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 6;
 - (iii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 7; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8;
 - (iv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 9; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 10;
 - (v) a CDR1 comprising the amino acid sequence of SEQ ID NO: 11 or 196; a CDR2 comprising the amino acid sequence of SEQ ID NO: 12; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 8;
 - (vi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 13; a CDR2 comprising the amino acid sequence of SEQ ID NO: 9; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 10;
 - (vii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 14 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 15; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16;
 - (viii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 17; a CDR2 comprising the amino acid sequence of SEQ ID NO: 18; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19;
 - (ix) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 21; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16;
 - (x) a CDR1 comprising the amino acid sequence of SEQ ID NO: 22; a CDR2 comprising the amino acid sequence of SEQ ID NO: 23; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19;
 - (xi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 24 or 198; a CDR2 comprising the amino acid sequence of SEQ ID NO: 25; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 26;

- (xii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 27; a CDR2 comprising the amino acid sequence of SEQ ID NO: 28; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 29;
- (xiii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 14 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 30; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16;
- (xiv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 17; a CDR2 comprising the amino acid sequence of SEQ ID NO: 31; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19;
- (xv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 1 or 195; a CDR2 comprising the amino acid sequence of SEQ ID NO: 32; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 3;
- (xvi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 6;
- (xvii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 34; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16;
- (xviii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 22; a CDR2 comprising the amino acid sequence of SEQ ID NO: 35; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 19;
- (xix) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 36; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16;
- (xx) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 37; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16;
- (xxi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 20 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 38; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 16;
- (xxii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 39;
- (xxiii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 4; a CDR2 comprising the amino acid sequence of SEQ ID NO: 33; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 40;

(xxiv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 165 or 199; a CDR2 comprising the amino acid sequence of SEQ ID NO: 166; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 167;

(xxv) a CDR1 comprising the amino acid sequence of SEQ ID NO: 168; a CDR2 comprising the amino acid sequence of SEQ ID NO: 169; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 170;

(xxvi) a CDR1 comprising the amino acid sequence of SEQ ID NO: 171 or 200; a CDR2 comprising the amino acid sequence of SEQ ID NO: 172; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 173;

(xxvii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 174; a CDR2 comprising the amino acid sequence of SEQ ID NO: 175; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 176;

(xxviii) a CDR1 comprising the amino acid sequence of SEQ ID NO: 177 or 197; a CDR2 comprising the amino acid sequence of SEQ ID NO: 178; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 179; or

(xxix) a CDR1 comprising the amino acid sequence of SEQ ID NO: 180; a CDR2 comprising the amino acid sequence of SEQ ID NO: 181; and a CDR3 comprising the amino acid sequence of SEQ ID NO: 182.

2. An anti-CD20 single domain antibody (sdAb) comprising:

(i) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 41;

(ii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 42;

(iii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 43;

(iv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 44;

(v) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 45;

(vi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 46;

(vii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 47;

- (viii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 48;
- (ix) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 49;
- (x) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 50;
- (xi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 51;
- (xii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 52;
- (xiii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 53;
- (xiv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 54;
- (xv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 55;
- (xvi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 56;
- (xvii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 57;
- (xviii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 58;
- (xix) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 59;
- (xx) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 60;
- (xxi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 61;
- (xxii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 62;
- (xxiii) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 63;
- (xxiv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 183;
- (xxv) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 184; or

(xxvi) a CDR1, a CDR2, and a CDR3 having the amino acid sequences of the CDR1, CDR2, and CDR3, respectively, as set forth in SEQ ID NO: 185.

3. The anti-CD20 sdAb of claim 2, wherein the CDR1, CDR2 or CDR3 are determined according to the Kabat numbering scheme, the IMGT numbering scheme, the AbM numbering scheme, the Chothia numbering scheme, the Contact numbering scheme, or a combination thereof.

4. The anti-CD20 sdAb of any one of claims 1 to 3, further comprising one or more FR regions as set forth in SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 and/or SEQ ID NO: 185.

5. The anti-CD20 sdAb of any one of claims 1 to 4, comprising the amino acid sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 or SEQ ID NO: 185.

6. The anti-CD20 sdAb of any one of claims 1 to 4, wherein anti-CD20 sdAb comprises or consists of an amino acid sequence having at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or more sequence identity with the sequence of SEQ ID NO: 41, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 49, SEQ ID NO: 50, SEQ ID NO: 51, SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54, SEQ ID NO: 55, SEQ ID NO: 56, SEQ ID NO: 57, SEQ ID NO: 58, SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 61, SEQ ID NO: 62, SEQ ID NO: 63, SEQ ID NO: 183, SEQ ID NO: 184 or SEQ ID NO: 185.

7. The anti-CD20 sdAb of claim 1 or claim 2, wherein anti-CD20 sdAb is a camelid sdAb.

8. The anti-CD20 sdAb of claim 1 or claim 2, wherein anti-CD20 sdAb is a humanized sdAb.

9. The anti-CD20 sdAb of any one of claims 1 to 8, wherein the anti-CD20 sdAb is genetically fused or chemically conjugated to an agent.

10. A chimeric antigen receptor (CAR), comprising:

(a) an extracellular antigen binding domain comprising the anti-CD20 sdAb of any

one of claims 1 to 9;

- (b) a transmembrane domain; and
- (c) an intracellular signaling domain.

11. The CAR of claim 10, wherein the extracellular antigen binding domain further comprises one or more additional antigen binding domain(s).
12. The CAR of claim 11, wherein the extracellular antigen binding domain further comprises one additional antigen binding domain; or wherein the extracellular antigen binding domain further comprises two additional antigen binding domains.
13. The CAR of any one of claims 11 to 12, wherein the one or more additional antigen binding domain(s) bind to one or more antigen(s) selected from a group consisting of CD20, CD19, CD22, CD33, CD38, BCMA, CS1, ROR1, GPC3, CD123, IL-13R, CD138, c-Met, EGFRvIII, GD-2, NY-ESO-1, MAGE A3, and glycolipid F77.
14. The CAR of any one of claims 11 to 13, wherein the extracellular antigen binding domain comprises two or three of the anti-CD20 sdAbs of any one of claims 1 to 9.
15. The CAR of any one of claims 10 to 14, wherein the transmembrane domain is derived from a molecule selected from a group consisting of CD8 α , CD4, CD28, CD137, CD80, CD86, CD152, and PD1.
16. The CAR of claim 15, wherein the transmembrane domain is derived from CD8 α .
17. The CAR of any one of claims 10 to 16, wherein the intracellular signaling domain comprises a primary intracellular signaling domain of an immune effector cell.
18. The CAR of claim 17, wherein the primary intracellular signaling domain is derived from CD3 ζ .
19. The CAR of claim 17 or claim 18, wherein the intracellular signaling domain further comprises a co-stimulatory signaling domain.
20. The CAR of claim 19, wherein the co-stimulatory signaling domain is derived from a co-stimulatory molecule selected from the group consisting of CD27, CD28, CD137, OX40, CD30, CD40, CD3, LFA-1, CD2, CD7, LIGHT, NKG2C, B7-H3, ligands of CD83 and combinations thereof.
21. The CAR of claim 20, wherein the co-stimulatory signaling domain is derived from CD137.
22. The CAR of any one of claims 10 to 21, further comprising a hinge domain located between the C-terminus of the extracellular antigen binding domain and the N-terminus of the

transmembrane domain.

23. The CAR of claim 22, wherein the hinge domain is derived from CD8 α .
24. The CAR of any one of claims 10 to 23, further comprising a signal peptide located at the N-terminus of the polypeptide.
25. The CAR of claim 24, wherein the signal peptide is derived from CD8 α .
26. A chimeric antigen receptor (CAR), comprising (i) an amino acid sequence selected from the group consisting of SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, SEQ ID NO: 95, SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 99, SEQ ID NO: 100, SEQ ID NO: 101, SEQ ID NO: 102, SEQ ID NO: 103, SEQ ID NO: 104, SEQ ID NO: 105, SEQ ID NO: 106, SEQ ID NO: 189, SEQ ID NO: 190, and SEQ ID NO: 191; or (ii) an amino acid sequence having at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or more sequence identity with the sequence of SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 89, SEQ ID NO: 90, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, SEQ ID NO: 94, SEQ ID NO: 95, SEQ ID NO: 96, SEQ ID NO: 97, SEQ ID NO: 98, SEQ ID NO: 99, SEQ ID NO: 100, SEQ ID NO: 101, SEQ ID NO: 102, SEQ ID NO: 103, SEQ ID NO: 104, SEQ ID NO: 105, SEQ ID NO: 106, SEQ ID NO: 189, SEQ ID NO: 190, or SEQ ID NO: 191.
27. An isolated nucleic acid comprising a nucleic acid sequence encoding the anti-CD20 sdAb of any one of claims 1 to 9.
28. A vector comprising the isolated nucleic acid of claim 27.
29. An isolated nucleic acid comprising a nucleic acid sequence encoding the CAR of any one of claims 10 to 26.
30. A vector comprising the isolated nucleic acid of claim 29.
31. An engineered immune effector cell, comprising the CAR of any one of claims 10 to 26, the isolated nucleic acid of claim 29, or the vector of claim 30.
32. The engineered immune effector cell of claim 31, wherein the immune effector cell is a T cell or B cell.
33. A pharmaceutical composition, comprising the anti-CD20 sdAb of any one of claims 1 to 9, the engineered immune effector cell of claim 31 or claim 32, or the vector of claim 28 or claim 30, and a pharmaceutically acceptable excipient.

34. A method of treating a disease or disorder in a subject, comprising administering to the subject an effective amount of the anti-CD20 sdAb of any one of claims 1 to 9, the engineered immune effector cell of claim 31 or claim 32, or the pharmaceutical composition of claim 33.
35. The method of claim 34, wherein the disease or disorder is a B cell associated disease or disorder and/or CD20 associated disease or disorder.
36. The method of claim 35, wherein the disease or disorder is cancer.
37. The method of claim 36, wherein the disease or disorder is a B cell malignancy.
38. The method of claim 37, wherein the B cell malignancy is a B cell leukemia or B cell lymphoma.
39. The method of claim 34, wherein the disease or disorder is selected from a group consisting of marginal zone lymphoma (*e.g.*, splenic marginal zone lymphoma), diffuse large B cell lymphoma (DLBCL), mantle cell lymphoma (MCL), primary central nervous system (CNS) lymphoma, primary mediastinal B cell lymphoma (PMBL), small lymphocytic lymphoma (SLL), B cell prolymphocytic leukemia (B-PLL), follicular lymphoma (FL), burkitt lymphoma, primary intraocular lymphoma, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), hairy cell leukemia (HCL), precursor B lymphoblastic leukemia, non-hodgkin lymphoma (NHL), high-grade B-cell lymphoma (HGBL), and multiple myeloma (MM).
40. The method of claim 34, wherein the disease or disorder is an autoimmune and/or inflammatory disease.
41. The method of claim 40, wherein the autoimmune and/or inflammatory disease is associated with inappropriate or enhanced B cell numbers and/or activation.